



# SCOPE 2.0: A model to simulate vegetated land surface fluxes and satellite signals

Peiqi Yang<sup>1</sup>, Egor Prikaziuk<sup>1</sup>, Wout Verhoef<sup>1</sup>, and Christiaan van der Tol<sup>1</sup>

<sup>1</sup>University of Twente, Faculty ITC, P.O. Box 217, 7500 AE Enschede, The Netherlands

**Correspondence:** Christiaan van der Tol (c.vandertol@utwente.nl)

**Abstract.** The Soil Canopy Observation of Photosynthesis and Energy fluxes (SCOPE) model aims at linking satellite observations in the visible, infrared and thermal domains with land surface processes in a physically based manner, and quantifying the micro-climate in the canopy. It simulates radiative transfer in the soil, leaves and vegetation canopies, as well as photosynthesis and non-radiative heat dissipation through convection and mechanical turbulence. Since the first publication 11 years ago, SCOPE has been applied in remote sensing studies of solar-induced chlorophyll fluorescence (SIF), energy balance fluxes, gross primary productivity (GPP) and directional thermal signals. Here we present a thoroughly revised version, SCOPE 2.0, which features a number of new elements: (1) It enables the definition of layers consisting of leaves with different properties, thus enabling the simulation of vegetation with an understory or with a vertical gradient in leaf chlorophyll concentration; (2) It enables the simulation of soil reflectance; (3) It includes the simulation of leaf and canopy reflectance changes induced by the xanthophyll cycle; and (4) The computation speed has been reduced by 90% compared to earlier versions due to a fundamental optimization of the model. These new features improve the capability of the model to represent complex canopies and to explore the response of remote sensing signals to vegetation physiology. The improvements in the computational efficiency make it possible to use SCOPE 2.0 routinely for the simulation of satellite data and land surface fluxes. It also strengthens the operability for numerical retrieval of land surface products from satellite or airborne data.

15

## 1 Introduction

Vegetation, as a dynamic component of the earth system, affects the climate via its influence on the exchange of energy and matter between the land surface and the atmosphere. Quantification of this exchange is relevant for a wide range of applications including weather prediction, climate projections, agriculture, and ecological and hydrological studies.

20 Earth observation with satellites can be used to monitor key characteristics of vegetation that are responsible for the surface-atmosphere exchanges, and identify changes therein. The most commonly used remote sensing indicator of vegetation biophysical and biochemical properties is reflectance (Ollinger, 2011). For example, the MODIS (Moderate Resolution Imaging Spectroradiometer) normalized vegetation reflectance index (NDVI), and the MERIS (MEdium Resolution Imaging Spec-



trometer) Terrestrial Chlorophyll Index (MTCI) have been empirically correlated with canopy leaf area index (LAI) and leaf  
25 chlorophyll content, respectively (Huete et al., 2002). Recent developments of sun-induced chlorophyll fluorescence (SIF)  
have offered an additional way to monitor vegetation (Mohammed et al., 2019). SIF has been successfully used to estimate  
gross primary production (GPP) (Guanter et al., 2014; Ryu et al., 2019) and stress detection (Ač et al., 2015; Rossini et al.,  
2015). Additional to reflectance and SIF, thermal signals provide insights into the physical processes of surface energy and  
water balance, such as demonstrated by the mapping of evapotranspiration globally with satellite thermal radiance (Nemani  
30 and Running, 1989; Allen et al., 2007).

Combined radiative transfer and plant physiological modelling is a promising way to investigate the exchanges of energy,  
water and carbon among soil, vegetation and atmosphere, and to develop remote sensing techniques for monitoring of vegeta-  
tion functioning. Many factors affect the signals observed from remote sensing, including sun-observation geometry, vegetation  
canopy structure and composition of the Earth's surface and atmosphere. Physically consistent exploitation of remote sensing  
35 data, therefore, requires the modelling of radiative transfer in the soil-vegetation-atmosphere system. Radiative transfer models  
(RTMs) describe the relationship between vegetation characteristics and remote sensing observations obtained under varying  
sun-observer geometry. However, for a complete understanding of the role of vegetation in the energy budget of the Earth's  
surface, radiative transfer modelling is not sufficient. One also needs to model non-radiative processes of energy dissipation via  
photosynthesis, phase transitions of water, heat storage and turbulent heat exchange between the surface and the atmosphere.  
40 This enables investigations beyond the monitoring of vegetation biophysical and biochemical properties, towards monitoring  
of fluxes.

The Soil-Canopy-Observation of Photosynthesis and Energy fluxes (SCOPE) model simulates the radiative transfer of in-  
cident light and thermal and fluorescence radiation emitted by soil and plants, component temperatures, photosynthesis and  
turbulent heat exchange (Van der Tol et al., 2009). In SCOPE, the radiative transfer and the non-radiative energy fluxes are  
45 computed in an assemblage of leaves and soil. The energy balance is maintained at all levels of spatial aggregation. Maintaining  
an energy budget is necessary for the simulation of thermal radiation, which depends on the within-canopy temperature  
distribution. To obtain this distribution, stomatal aperture, latent and sensible heat fluxes of individual elements have to be  
resolved together with the radiative fluxes in the vegetation canopy.

SCOPE has been applied in a wide range of studies. Thanks to the coupling of photosynthesis and radiative transfer of  
50 fluorescence in the SCOPE model, it has been used as a convenient tool for in-depth process-based studies to unravel the  
relationship between fluorescence and photosynthesis (Damm et al., 2015; Verrelst et al., 2016; Migliavacca et al., 2017).  
Besides, it has also been used for simulating directional anisotropy of satellite-measured surface temperatures (Duffour et al.,  
2015), for predicting evapotranspiration (Galleguillos et al., 2011), and as a benchmark for other simple radiative transfer  
models (Bian et al., 2020). Contemporary simulations of satellite observations and plant physiological processes make SCOPE  
55 a useful tool to monitor dynamic vegetation response to environmental conditions (Zhang et al., 2014; Pacheco-Labrador et al.,  
2019).

Since the original publication, SCOPE been extended with new features:

1. The soil reflectance model BSM (Verhoef et al., 2018; Yang et al., 2020b) has been introduced.



2. The radiative transfer of fluorescence has been improved (Van der Tol et al., 2019b).
- 60 3. Changes in reflectance due to xanthophyll pigment changes have been included (Vilfan et al., 2018).
4. The RTMs in the SCOPE have been adapted for multi-layer canopies (Yang et al., 2017).

These new features have not been described together. In the new version of SCOPE (SCOPE 2.0) presented here, these improvements are coherently incorporated. Moreover, the model has been optimized in many ways to improve the computational efficiency, and the options to provide data input formats have been extended. We present a description of the basic functionality  
65 of the model followed by several recent developments.

## 2 General description of SCOPE

### 2.1 Starting points

SCOPE is designed to simulate photosynthetic, hydrological, and radiative transfer processes at the vegetated land surface. For these purposes, it combines several RTMs with a leaf biochemical model and an aerodynamic resistance scheme. These  
70 models provide simulations of emanating hyperspectral radiance and net radiation  $R_n$  (via radiative transfer processes), photosynthesis rates (via photosynthetic processes), and sensible heat flux  $H$ , latent heat flux  $\lambda E$  and ground heat flux  $G$  (via micro-meteorological processes), for both individual elements of the land surface (e.g., soil and leaves) and the whole vegetation stand. In order to meet the requirements of broad applicability, the models are as much as possible physically based. The model has been developed further since the first publication Van der Tol et al. (2009), but the main model structure and  
75 functionality remain. The SCOPE model described in this section refers to the first-release version in 2009.

The central idea of SCOPE is the modelling of interactions between radiative and non-radiative fluxes among elements of the vegetation canopy. Remote sensing signals, such as reflectance, fluorescence and thermal radiance, are outputs of these interactions. The modelling of radiative fluxes facilitates the simulation of optical properties (i.e., reflectance, transmittance and absorptance) of soil, leaves and canopies. This is complemented with the modelling of non-radiative fluxes in vegetation  
80 canopies, respecting energy conservation at all levels of spatial aggregation from the photosystem to the whole stand. The energy budget is determined by both the radiative transfer of incident and emitted (thermal) radiation, and the exchange of (latent) heat with the atmosphere. The surface temperature is resolved as the outcome of this balance.

### 2.2 Model domain and representation

In the spatial domain, the typical representation of land surfaces in SCOPE is a vegetation layer consisting of leaves bounded  
85 underneath by a soil surface. The representation of the vegetation layer is one-dimensional in the sense that fluxes in the vertical ( $z$ ) direction are considered only. This implies that even if the model is applied pixel-by-pixel in a spatial grid, the horizontal interactions between neighbouring pixels are not considered. Thus, typical 3-D effects in the vegetation, such as boundary



effects at the edge of fields or forests, or effects of topography and horizontal heat advection are not included. Radiative transfer is based on turbid medium representations of both leaves and canopies.

90 In the temporal domain, SCOPE assumes steady-state conditions. This means that a simulation with SCOPE outputs the energy and spectrally resolved radiation budgets of the surface for a single set of surface and weather characteristics at one moment in time. The lack of memory of state variables in time also means that storage of carbon and water are not considered, and similarly, vegetation growth is not simulated. A complete run of SCOPE may consist of many simulations, either for one location as a function of time, or for different locations or surface types, but the simulations in such a sequence are  
95 treated independently without interactions, and thus the order of model simulations is arbitrary. One exception is the (optional) modelling of the soil heat budget with a thermal inertia approach, which is described in section 3.4.

In the spectral domain, SCOPE simulates visible to thermal infrared radiance from 0.4 to 50 $\mu$ m within and above vegetation canopies. The spectral resolutions in the spectral regions from 0.4 to 2.5 $\mu$ m, from 2.5 to 15 $\mu$ m, and from 15 to 50  $\mu$ m are 1 nm, 100 nm, and 1000 nm, respectively. It also covers the fluorescence emission spectral region from 640 to 850 nm with a  
100 resolution of 1 nm. It is noted that the spectral resolutions in these regions are easily adapted to simulation requirements and spectral input data.

### 2.3 Structure of the model

The model code at the highest hierarchical level, SCOPE, calls sub-models which operate in series. The main sub-models are listed in Table 1. Besides the listed sub-models for radiative transfer and energy balance, SCOPE requires functions for input,  
105 output and some supporting functions (such as Planck's equation). Therefore, all the functions used in SCOPE can be organized in four types: (1) RTMs, (2) modules for energy balance, (3) input-output functions, and (4) supporting functions.

#### 2.3.1 RTMs

SCOPE includes seven RTMs, which together simulate the spectrally resolved radiance emanating from the vegetation: one for the soil (BSM, only available in SCOPE 2.0, Verhoef et al., 2018; Yang et al., 2020b), one for the leaf (Fluspect, Vilfan  
110 et al., 2016, 2018), and five for the whole stand, i.e. the combined system of soil and foliage. They include one RTM for incident radiation from the sun and the sky (RTMo), two for thermal radiation emitted by the soil and vegetation (RTMt\_sb and RTMt\_planck), one for chlorophyll fluorescence, RTMf (Van der Tol et al., 2009, 2019b), and one for the dynamic modulations of leaf reflectance and transmittance due to pigment changes in the xanthophyll cycle (RTMz, only available in SCOPE v1.70 or later, Vilfan et al., 2018).

115 Four types of fluxes are involved in the radiative transfer processes, namely a direct solar flux, two hemispherical (semi-isotropic) diffuse fluxes (up- and downward) and a flux in the direction of viewing. Following the Kubelka-Munk theory, the radiative transfer in the vertical direction is expressed with a set of linear differential equations (Verhoef, 1984). These equations are solved either with analytical or numerical approaches.



**Table 1.** Main sub-models in SCOPE.

sub-models	main functions	main input	main output
BSM	simulating soil reflectance	soil moisture, brightness and two spectral shape related parameters	anisotropic soil reflectance
Fluspect	leaf RTM	leaf biophysical properties	leaf reflectance, transmittance and fluorescence emission matrices
RTMo	RTM for incident radiation	canopy structure, leaf reflectance, transmittance and soil reflectance	canopy reflectance, radiation absorbed by each leaf
RTMf	RTM for fluorescence fluxes	canopy structure, leaf reflectance, transmittance, soil reflectance and fluorescence emission matrices	fluorescence of each leaf and of the whole canopy
RTMt_sb/RTMt_planck	RTM for thermal fluxes	leaf temperature, incoming thermal radiation, emissivity of soil and leaves	thermal emission of each leaf and of the whole canopy
RTMz	RTM for fluxes induced by the xanthophyll cycle	leaf absorbed radiation, canopy structure, leaf reflectance, transmittance, soil reflectance	dynamic modulations of canopy reflectance
biochemical	biochemical model for photosystem energy partitioning	leaf absorbed radiation, leaf temperature, photosynthetic parameters	photosynthesis rate, fluorescence emission efficiency and heat dissipation
ebal	energy balance module	leaf absorbed radiation, leaf temperature	sensible and latent heat fluxes

### 2.3.2 Energy balance module

120 The energy balance module in SCOPE minimizes the energy balance closure error  $e_{ebal}$ :

$$e_{ebal} = R_n - H + \lambda E - G \quad (1)$$

for all leaf and soil elements by iteratively updating their temperature. In this equation,  $R_n$  is the net radiation,  $H$  the sensible heat flux,  $\lambda E$  the latent heat flux, and  $G$  the ground heat flux (i.e., zero for leaf elements), all in  $\text{Wm}^{-2}$ . In the energy balance, chemical conversions (e.g., photosynthesis and respiration) and fluorescence are neglected.

125 The net radiation is obtained after spectral integration of the radiative transfer modules for incident radiation (RTMo) and internally generated thermal radiation (RTMt). The radiative transfer for incident radiation is computed before the energy balance closure loop, while the internally generated thermal radiation is calculated within this loop, because of its dependence on leaf and soil temperatures. The sensible and latent heat fluxes are calculated with an aerodynamic resistance scheme, where resistances for the leaf and soil boundary layer, the vegetation layer and the atmospheric surface layer apply. The aerodynamic  
 130 resistances depend on atmospheric stability, wind speed and surface roughness (Van der Tol et al., 2009). For latent heat flux of leaves, a stomatal resistance is calculated with a combined photosynthesis and stomatal model (Van der Tol et al., 2014), while for the soil, a surface resistance applies which is either a pre-defined input, or an empirical function of soil moisture.



### 2.3.3 Leaf biochemical model

The biochemical model simulates the energy partitioning into fluorescence, heat or photochemistry in photosystems (Maxwell  
135 and Johnson, 2000; Van der Tol et al., 2014). It is based a conventional photosynthesis model of Collatz et al. (1991) for C3  
and Collatz et al. (1992) for C4 vegetation, in which photosynthetic rates (or photosynthetic light use efficiency) are simulated  
as a function of leaf temperature, ambient radiation levels, intercellular CO<sub>2</sub> concentrations  $C_i$ , and other leaf physiological  
parameters (e.g., photosynthetic pathways, maximum carboxylation rate  $V_{c_{mo}}$ ).


Van der Tol et al. (2014) established empirical relationships between fluorescence emission efficiency and photosynthetic  
140 light use efficiency under various environmental conditions by using active fluorescence measurements. With these relation-  
ships, the fraction of the absorbed radiation by a leaf emitted as fluorescence and dissipated as heat can be simulated.

### 2.3.4 Interactions among the sub-models

Fig. 1 is a schematic overview of the SCOPE model structure, which also shows the connections among the sub-models. A full  
list of input parameters is provided in Table 2. A simulation with SCOPE starts with calculating soil reflectance (BSM), the  
145 leaf reflectance and transmittance and fluorescence emission excitation matrices (Fluspect). These simulations of soil and leaf  
optical properties, together with canopy structure and irradiance, are the input of canopy RTMs. The sub-model for radiative  
transfer of solar and sky radiation (RTMo) takes leaf optical properties, and soil reflectance as input and simulates canopy  
reflectance and radiation fluxes including the net absorbed solar radiation by soil and leaves. RTMf takes the leaf fluorescence  
emission excitation matrices and the radiation fluxes as input, and simulates canopy fluorescence.

150 The radiative transfer of emitted thermal radiation relies on the temperatures of soil and leaves, which are not known a priori.  
For this reason, the thermal radiative transfer model is carried out in the energy balance closure loop, as described in section  
2.3.2. For the purpose of computational efficiency, the radiative transfer of emitted thermal radiation is carried out in broadband  
by using RTMt\_sb. The letters 'sb' denote the use of the Stefan–Boltzmann law to describe the spectrally integrated radiance  
from a leaf or soil in terms of its temperature. Leaf temperature is also used together with the radiation absorbed by leaf  
155 chlorophyll pigments and other leaf physiological parameters, to simulate photosystem energy partitioning in the biochemical  
model (Van der Tol et al., 2014). The energy balance residual is used to update the initial estimate of the temperature of each  
element.

After energy balance closure, the thermal radiation fluxes are simulated spectrally resolved in observation direction by using  
RTMt\_planck, where 'planck' denotes the use of Planck's law to describe the spectrally resolved radiance from a leaf or  
160 soil in terms of its temperature. The radiative transfer of the emitted fluorescence is simulated with RTMf. This module uses  
the radiative fluxes interacting with leaves as simulated with RTMo, and the fluorescence emission matrices simulated with  
Fluspect, to simulate leaves' fluorescence emission, which is aggregated to canopy fluorescence signals. Finally, the effect of  
(small) changes in reflectance and transmittance due to the illumination and temperature dependent xanthophyll epoxidation  
state are simulated with RTMz.



**Figure 1.** Schematic overview of the SCOPE model structure. For a complete list of input data, see Table 2

## 165 2.4 Model inputs and outputs

### 2.4.1 Input variables for soil, vegetation and meteorology

The inputs of the SCOPE model comprise soil, leaf and canopy properties as well as sun-observer geometry and meteorological conditions. Besides the intermediate variables, which are passed between the sub-models, the main input variables of SCOPE are given in Table 2.

170 Leaf biophysical and biochemical parameters characterize leaf pigment, water and dry matter contents, which determine leaf optical properties. Canopy structural parameters describe the arrangement of the leaves in the canopy. Sun-observer geometry is determined by sun and observer's zenith angles and their absolute azimuth difference. Both the canopy structural parameters and sun-observer geometry strongly affect remote sensing signals observed above the canopy. It is worth noting that viewing angles only affect the directional signals observed at the top of a canopy, such as reflectance, fluorescence and thermal radiation,



175 but they have no impact on the radiative fluxes inside the canopy, canopy photosynthesis and respiration and non-radiative fluxes, such as latent and sensible heat.

The meteorological inputs for SCOPE include the typical synoptic weather variables of air temperature, humidity, wind speed, air pressure and the concentrations of oxygen and carbon dioxide. All these inputs are required close to the Earth surface. The height above the surface of the terrain for which they are specified must be given in the input, as input  $z$ , typically  
180 2.5 times the vegetation height. Thus,  $z$  is not the height of the terrain above sea level, but rather the height above the terrain at in surface layer, where the wind profile is logarithmic. The value of  $z$  must be given in the input as it is used to calculate the aerodynamic roughness of the surface.

#### 2.4.2 Input irradiance for the atmosphere boundary condition

In addition to the variables listed in Table 2, SCOPE requires the radiative properties of the atmosphere as the upper boundary  
185 condition. They can be provided in two different ways. The first option is to provide irradiance at the bottom of atmosphere (BOA) in the form of a file with two columns representing the spectra of direct solar irradiance  $E_{sun}$ , and diffuse sky irradiance  $E_{sky}$  [ $\text{Wm}^{-2}\mu\text{m}^{-1}\text{sr}^{-1}$ ], respectively. These spectra could either be measured in the field or generated with an atmospheric RTM (e.g. MODTRAN, Berk et al., 1999). Using an atmospheric RTM has the disadvantage that  $E_{sky}$  may not be accurate, because  $E_{sky}$  also depends on the surface (canopy) reflectance in the surroundings, which may not be known a priori in the  
190 atmospheric radiative transfer simulation. Therefore, if the surface reflectance assumed in the atmospheric radiative transfer simulation largely differs from the canopy reflectance produced by SCOPE, errors in  $E_{sky}$  occur. The second and preferred option is using an atmospheric RTM to generate some optical properties of the atmosphere rather than the direct output of  $E_{sky}$  and  $E_{sun}$ . The optical properties should include the following coefficients.

- $E_s \cos(\theta_s)$ , the product of the solar irradiance at TOA and the cosine of the solar zenith angle. This product is the  
195 irradiance at TOA projected on the surface [ $\text{Wm}^{-2}\mu\text{m}^{-1}\text{sr}^{-1}$ ].
- $\rho_{dd}$ , the diffuse reflectance of the atmosphere (i.e. the spherical albedo).
- $\tau_{ss}$ , the direct atmospheric transmittance in the direction of the sun.
- $\tau_{sd}$ , the diffuse atmospheric transmittance for solar incidence.
- $L_a$ , the thermal emission by atmosphere at BOA, towards the surface ( $\text{Wm}^{-2}\mu\text{m}^{-1}\text{sr}^{-1}$ ).

200 The coefficients listed can be extracted from MODTRAN simulations by using the T18 system, which is described in detail in Verhoef et al. (2018). A database of the optical coefficients for several typical atmospheric conditions is provided together with the SCOPE model. With these coefficients, SCOPE can simulate the BOA direct and diffuse irradiance spectra in the module RTMo with consideration of surface-atmosphere interactions. The BOA irradiances  $E_{sun}$  and  $E_{sky}$  are calculated in



**Table 2.** Main input variables of SCOPE

symbol	abbreviation	unit	default value	sub-model	description
$C_{ab}$	Cab	$[\mu\text{gcm}^{-2}]$	80	Fluspect	leaf chlorophyll concentration
$C_{ca}$	Cca	$[\mu\text{gcm}^{-2}]$	20	Fluspect	leaf carotenoid concentration
$C_w$	Cw	[cm]	0.01	Fluspect	equivalent water thickness in leaves
$C_s$	Cs	[]	0	Fluspect	leaf senescence parameters
$C_{dm}$	Cdm	$[\text{g cm}^{-2}]$	0.012	Fluspect	leaf dry matter content
$C_{ant}$	Cant	$[\mu\text{g cm}^{-2}]$	0	Fluspect	Anthocyanin content
$N$	N	[]	1.4	Fluspect	leaf structure parameter
$L$	LAI	[]	3	canopy RTMs	projected leaf area per unit ground area
$h_c$	hc	[m]	2	canopy RTMs	vegetation height
$LIDFa$	LIDFa	[]	-0.35	canopy RTMs	parameter for the mean leaf zenith angle
$LIDFb$	LIDFb	[]	-0.15	canopy RTMs	bimodality of leaf angle distribution
$\theta_s$	tts	[deg]	30	canopy RTMs	solar zenith angle
$\theta_o$	tto	[deg]	0	canopy RTMs	viewing zenith angle
$\psi$	psi	[deg]	0	canopy RTMs	absolute azimuth difference
$R_{si}$	Rin	$[\text{Wm}^{-2}]$	600	canopy RTMs	shortwave irradiance
$R_{li}$	Rli	$[\text{Wm}^{-2}]$	300	canopy RTMs	longwave irradiance
$p_a$	p	[hPa]	970	energy balance	air pressure
$T_a$	T	[°C]	20	energy balance	air temperature
$u$	u	$[\text{ms}^{-1}]$	2	energy balance	wind speed
$e_a$	ea	[hPa]	15	energy balance	vapour pressure
$z$	z	[m]	2	energy balance	measurement height
$\Theta$	SMC	[]	25	BSM, energy balance	surface volumetric soil moisture content
$B$	BSMBrightness	[]	0.5	BSM	soil brightness
$\varphi$	BSMlat	[deg]	25	BSM	soil 'latitude' parameter (not geographical)
$\lambda$	BSMlon	[deg]	45	BSM	soil 'longitude' parameter (not geographical)
$C_a$	Ca	[ppm]	380	biochemical model	atmospheric CO <sub>2</sub> concentration
$V_{cmo}$	Vcmo	$[\mu\text{mol m}^{-2}]$	70	biochemical model	carboxylation capacity at 25 deg C
$m$	m	[]	12	biochemical model	Ball-Berry stomatal parameter



the function RTMo as:

$$E_{sun} = E_s \cos(\theta_s) \overline{\tau_{ss}}$$
$$E_{sky} = \frac{E_s \cos(\theta_s) (\overline{\tau_{sd}} + \overline{\tau_{ss} \rho_{dd} r_{sd}}) + \pi((1 - r_{dd}) L_s \overline{\rho_{dd}} + L_a)}{1 - \overline{\rho_{dd} r_{dd}}} \quad (2)$$

where  $r_{sd}$  and  $r_{dd}$  are the surface reflectance for direct and diffuse incoming radiation, respectively and  $L_s$  the thermal emission by the (vegetated) surface ( $\text{Wm}^{-2} \mu\text{m}^{-1} \text{sr}^{-1}$ ). All of them are simulated with SCOPE. The overbars denote the spectral averaging to the SCOPE resolution (1 nm in the VNIR). Note that  $\tau_{ss}$  and  $\rho_{dd}$  are aggregated to the SCOPE resolution separately, but also the product  $\tau_{ss} \rho_{dd}$ , in order to accommodate spectral correlation effects in the finite bands. The coupling with the atmosphere is described in detail in Verhoef et al. (2018) and Yang et al. (2020b).

Finally, SCOPE offers the possibility to provide additional values for the spectrally integrated irradiance (direct solar radiation  $E_{sun}$  plus  $E_{sky}$ ) over the ranges from 0.4 to  $2.5 \mu\text{m}$ , and 2.5 to  $50 \mu\text{m}$ . These are the input fields  $R_{si}$  and  $R_{li}$ , respectively. However, it is not necessary to specify these inputs, because the broadband irradiances  $R_{si}$  and  $R_{li}$  are already calculated internally as the integral of the irradiance spectra. If the values for these two inputs are specified, then the solar and sky irradiance spectra  $E_{sun}$  plus  $E_{sky}$  are linearly scaled (each by the same factor so that the ratio  $E_{sun}/E_{sky}$  remains unaltered) in the two spectral regions separately, to match the values provided for  $R_{si}$  and  $R_{li}$ . This option can be useful if time series of synoptic weather data are used as input, and if it is computationally not feasible to carry out atmospheric radiative transfer simulations for every time step separately. For coupled surface-atmosphere simulations this is not recommended, because of obvious inconsistencies between SCOPE and the atmospheric model. In that case, the input fields for  $R_{si}$  and  $R_{li}$  must be left blank.


### 2.4.3 Model outputs

In Table 3, the main outputs of SCOPE are listed. The general output of SCOPE includes 1) spectral simulations of radiance in the viewing direction and upward flux for the whole upper hemisphere from optical to thermal domain including fluorescence 2) radiation budget, such as incoming and outgoing radiation for shortwave from 0.5 to  $2.5 \mu\text{m}$  and longwave from 2.5 to  $50 \mu\text{m}$  3) fluxes such as sensible heat, latent heat and the ground heat flux for canopy, soil and the combined system and 4) canopy absorption, such as absorbed PAR by chlorophyll. Most of the stored outputs of SCOPE are for the whole canopy, although similar variables of leaves are also computed internally in SCOPE.



**Table 3.** SCOPE outputs

output	description	unit
<b>spectral simulation</b>		
Eout_spectrum	hemispherical leaving irradiance	$[\text{Wm}^{-2}\mu\text{m}^{-1}]$
Lo_spectrum	radiance in the viewing direction	$[\text{Wm}^{-2}\mu\text{m}^{-1}\text{sr}^{-1}]$
fluorescence	fluorescence radiance in the viewing direction	$[\text{Wm}^{-2}\mu\text{m}^{-1}\text{sr}^{-1}]$
fluorescence_hemis	hemispheric leaving fluorescence irradiance	$[\text{Wm}^{-2}\mu\text{m}^{-1}]$
reflectance	TOC reflectance in the viewing direction	[]
<b>vegetation</b>		
aPAR	PAR absorbed by the vegetation	$[\mu\text{mol m}^{-2}\text{s}^{-1}]$
aPARbyCab	PAR absorbed by chlorophyll	$[\mu\text{mol m}^{-2}\text{s}^{-1}]$
aPARbyCab_en	PAR energy absorbed by chlorophyll	$[\text{Wm}^{-2}]$
Photosynthesis	canopy photosynthesis rate	$[\mu\text{mol m}^{-2}\text{s}^{-1}]$
LST	black-body radiometric land surface temperature	[K]
<b>fluxes</b>		
Rnctot	Net radiation of canopy	$[\text{Wm}^{-2}]$
lEctot	Latent heat flux of canopy	$[\text{Wm}^{-2}]$
Hctot	Sensible heat flux of canopy	$[\text{Wm}^{-2}]$
Actot	Net photosynthesis of canopy	$[\text{Wm}^{-2}]$
Tcave	Average canopy temperature	[°C]
Rnstot	Net radiation of soil	$[\text{Wm}^{-2}]$
lEstot	Latent heat flux of soil	$[\text{Wm}^{-2}]$
Hstot	Sensible heat flux of soil	$[\text{Wm}^{-2}]$
Gtot	Soil heat flux	$[\text{Wm}^{-2}]$
Tsave	Average soil temperature	[°C]
Rntot	Total net radiation	$[\text{Wm}^{-2}]$
lEtot	Total latent heat flux	$[\text{Wm}^{-2}]$
Htot	Total sensible heat flux	$[\text{Wm}^{-2}]$
<b>radiation</b>		
ShortIn	Incoming shortwave radiation	$[\text{Wm}^{-2}]$
LongIn	Incoming longwave radiation	$[\text{Wm}^{-2}]$
HemisOutShort	hemispherical outgoing shortwave radiation	$[\text{Wm}^{-2}]$
HemisOutLong	hemispherical outgoing longwave radiation	$[\text{Wm}^{-2}]$
Lo	radiance in observation direction	$[\text{Wm}^{-2}\text{sr}^{-1}]$
Lot	thermal radiance in observation direction	$[\text{Wm}^{-2}\text{sr}^{-1}]$
Lote	emitted radiance in observation direction	$[\text{Wm}^{-2}\text{sr}^{-1}]$



**Figure 2.** Reflectance simulations with the BSM model. The black curves in each panels are the same simulation.

### 3 Major improvements of SCOPE 2.0 compared with SCOPE

#### 3.1 Implementation of the BSM soil reflectance model

230 In the first published version of SCOPE, the soil reflectance spectrum was an input variable. The users should either provide a measured soil spectrum or select one from the soil reflectance library incorporated in the SCOPE model. In SCOPE 2.0, we provide the users the option to simulate soil reflectance thanks to the implementation of a soil reflectance model.

The Brightness-Shape-Moisture (BSM) model simulates the isotropic soil reflectance. This model is based on an empirical reflectance model of dry soil (Verhoef et al., 2018; Jiang and Fang, 2019) and incorporates the effects of soil moisture by using  
235 the water film coating approach (Ångström, 1925; Yang et al., 2020b). To simulate reflectance of dry soil, the model requires soil brightness ( $B$ ) and two spectral-shape related parameters ( $\varphi$  and  $\lambda$ ) as inputs. Soil moisture is necessary for simulating wet soil reflectance.

Fig. 2 shows the effects of the four parameters on soil reflectance. It is evident that soil brightness only affects the ‘intensity’ of soil reflectance and the ‘shape’ of soil reflectance is controlled by  $\varphi$  and  $\lambda$ . Soil moisture affects reflectance intensity over  
240 all wavelengths, but reflectance at the water absorption bands are more sensitive to soil moisture. Soil moisture effects on reflectance are considerably similar to the effects of soil brightness, and the soil is dark when it is wet, as explained in Lekner and Dorf (1988).




### 3.2 Inclusion of dynamic reflectance induced by the xanthophyll cycle

A new feature in SCOPE 2.0 is modelling the photochemical reflectance dynamics induced by the xanthophyll cycle at both  
245 leaf and canopy levels. In the original leaf RTM Fluspect (Vilfan et al., 2016), leaf optical properties are determined by leaf  
biophysical properties. However, in natural conditions, the xanthophyll cycle that is involved in photo-protection mechanisms  
under excess light can provoke a change in reflectance and transmittance as the composition of the pigment pool is regulated.  
Changes in the de-epoxidation state (DEPS) of xanthophyll cycle pigments (e.g., violaxanthin and zeaxanthin) can be observed  
as changes in the leaf absorption of light with wavelengths between 500 to 570 nm. These spectral changes can be a good  
250 remote sensing indicator of the photosynthetic efficiency. The photochemical reflectance index (PRI,  $\frac{R_{570}-R_{531}}{R_{570}+R_{531}}$ ) proposed by  
Gamon et al. (1992) is a example of a measure for the effects of xanthophyll cycle pigments on the reflectance. It takes changes  
in reflectance at 531 nm to estimate DEPS with reflectance at 570 nm as a reference to correct changes in reflectance induced  
by other factors, such as sun-observer geometry.

Vilfan et al. (2018) incorporated the effects of the xanthophyll cycle on leaf optical properties in Fluspect and developed the  
255 Fluspect-CX model. The main idea of Fluspect-CX is to use *in vivo* specific absorption coefficients for two extreme states of  
carotenoids, representing the two extremes of the xanthophyll de-epoxidation. A ‘photochemical reflectance parameter’ (Cx)  
is employed to describe the intermediate states as a linear mixture of these two states. Cx controls the specific absorption  
coefficient of carotenoids in a leaf, and thus affects leaf reflectance and transmittance.

The propagation of changes in leaf reflectance and transmittance induced by the xanthophyll cycle to TOC reflectance  
260 is carried out with RTMz, which is largely similar to RTMf in the sense that both the xanthophyll cycle and fluorescence  
emission lead to small changes in (apparent) reflectance but for different spectral regions (i.e., 500-570 nm and 640-850 nm,  
respectively). RTMf and RTMz take fluorescence emission efficiency and Cx (simulated from the leaf biochemical model),  
respectively, as inputs, of which the magnitudes vary among individual leaves due to their ambient light intensities, temperature,  
etc. Fig. 3 depicts an example of the effects of Cx on the leaf and canopy reflectance as simulated by SCOPE 2.0 with the default  
265 model inputs (Table 2). Although the effects on canopy reflectance seem small, it could be helpful to monitor the variation in  
DEPS.

Fig. 4 compares simulations of PRI in a day with SCOPE and SCOPE 2.0. In these illustrative simulations, the default model  
inputs are used except for the incoming radiation and solar zenith angles. The values of incoming radiation and solar zenith  
angles ( $\theta_s$ ) are assigned according to the field measurements presented in Yang et al. (2020a) (i.e., on day 232 of the dataset in  
270 the referred paper). The comparison demonstrates that the inclusion of dynamic reflectance induced by the xanthophyll cycle  
has a clear impact on the simulation of diurnal changes in PRI. In SCOPE, diurnal variation of PRI is mainly regulated by  
sun-observer geometry, since leaf biophysical properties and canopy structure are kept unchanged in a day. Because the BRDF  
effects on reflectance at 531 nm and 570 nm are similar, they cancel out in PRI, and the diurnal variation of PRI simulated with  
SCOPE is small. Compared with SCOPE, SCOPE 2.0 considers the changes in leaf pigment pool induced by the xanthophyll  
275 cycle in response to the variation of incoming radiation besides the BRDF effects. The excessive incoming radiation in the




**Figure 3.** The effects of the xanthophyll cycle on leaf and canopy visible reflectance simulated with SCOPE 2.0 using the default model inputs (Table 2).  $C_x$  is a factor for the de-epoxidation state (DEPS) of xanthophyll cycle pigments.

midday leads to larger  $C_x$  values than in the morning and afternoon, and higher afternoon than morning temperatures to higher afternoon  $C_x$ , and thus more significant diurnal variation of PRI.

### 3.3 Adaption of the RTMs for multi-layer canopies

The original SCOPE model assumes that vegetation canopies are vertically homogeneous and horizontally infinite, as its radiative transfer routines are based on the classical 1-D SAIL model (Verhoef, 1984). The vertical heterogeneity of leaf biophysical and biochemical properties may have a large effect on the bi-directional reflectance, fluorescence and photosynthesis of vegetation canopies. To allow simulations of vertical heterogeneous canopies, Yang et al. (2017) modified the RTMs in SCOPE and developed a new branch of SCOPE, called mSCOPE. SCOPE 2.0 incorporates the essence of mSCOPE on radiative transfer modelling and adapts the capability to simulate reflectance, fluorescence and photosynthesis of vertically heterogeneous canopies.

RTMs in SCOPE 2.0 remain structurally the same with the original SCOPE. However, a more general solution of the radiative transfer problems is used. Compared to the classic SAIL analytical solution, SCOPE 2.0 (and mSCOPE) employs the adding method to solve the radiative transfer problems. The application of the adding method for TOC reflectance simulation is given in Verhoef (1985). Yang et al. (2017) extended this method to calculate the radiative flux profiles in the canopy. The procedure is summarized as follows: 1) divide the vertical layer into  $n$  thin homogeneous layers; 2) start from the bottom homogeneous layer, calculate the surface reflectance of the combined system of the bottom surface (e.g., soil) and this layer; 3) add a new homogeneous vegetation layer above the surface of the previous system in step 2, and calculate the surface reflectance of the new system; 4) repeat step 3 until all homogeneous layers are added; 5) Once the surface reflectance at each vertical level is obtained, the fluxes profile can be computed from top to bottom, given the incident fluxes at top of the canopy. For the radiative




**Figure 4.** The effects of the xanthophyll cycle on diurnal PRI variation by comparing simulations from SCOPE and SCOPE 2.0. The default model inputs are used except for the incoming radiation and solar zenith angles, which are values in a representative sunny day (two grey lines). Note that SCOPE does not include the effects of the xanthophyll cycle on TOC reflectance while SCOPE 2.0 does.

295 transfer of fluorescence and thermal radiation, the emission from leaves and soil should be included as extra radiation sources besides the incident fluxes at top of the canopy. In SCOPE 2.0, the value of  $n$  is set as 10 times LAI rather than a fixed value of 60 in mSCOPE, because this ensures the LAI of one elementary layer is small enough (i.e.,  $iLAI < 0.1$ ), and the use of less elementary layers improves the computational efficiencies of the RTMs.

### 3.4 An alternative way to estimate the ground heat flux

300 In SCOPE the ground heat flux is calculated for the sunlit and shaded soil. In the original SCOPE model, this was either a constant fraction of 0.35 of the net radiation on the soil, or calculated with the force restore method of Bhumralkar (1975). SCOPE 2.0 offers an alternative way to estimate the ground heat flux as a function of the soil temperature time series with the method of Wang and Bras (1999). The ground heat flux is determined by the gradient of soil temperature in the profile underneath the soil surface. The subsurface is outside the model domain of SCOPE, and therefore the soil temperature gradient



**Figure 5.** Representations of canopies in SCOPE and in SCOPE 2.0. In SCOPE, a canopy consists of leaves with identical biochemical and biophysical properties, and the leaf angle distribution is vertically invariant, while in SCOPE 2.0, both leaf properties and leaf angles can vary vertically.

305 is not simulated. However, this vertical gradient may equivalently be expressed the by the half-order time-derivative of the surface temperature (Wang and Bras, 1999). This enables the estimation of  $G$  from the time-history of the surface temperature:

$$G(t) = \Gamma / \sqrt{\pi} \int_{t_0}^t \frac{T(s)}{t-s} ds \quad (3)$$

where  $T$  is the soil temperature at time  $s$ ,  $\Gamma$  [ $\text{J m}^{-2} \text{s}^{-1/2} \text{K}^{-1}$ ] is the thermal inertia of the soil, calculated from physical properties of the soil:

$$310 \quad \Gamma = \sqrt{c_s \cdot \rho_s \cdot \lambda_s} \quad (4)$$

where  $c_s$  is the volumetric heat capacity of the soil [ $\text{J kg}^{-1} \text{K}^{-1}$ ],  $\rho_s$  the soil bulk density [ $\text{kg m}^{-3}$ ], and  $\lambda_s$  [ $\text{J m}^{-1} \text{s}^{-1} \text{K}^{-1}$ ] the heat conductivity of the soil. In SCOPE 2.0 a solution derived for a discrete time series of temperatures by Bennett et al. (2008) (Eq. A3 therein) was adopted:

$$G(t) = 2\Gamma / \sqrt{\pi} \sum_{i=1}^{12} \frac{T_{i+1} - T_i}{s_{i+1} - s_i} (\sqrt{t - s_i} - \sqrt{t - s_{i+1}}) \quad (5)$$

315 This approach is only meaningful if consecutive simulations are carried out in a time series, in which the diurnal variation of temperature is reproduced (at least one simulation per three-hour time step). The approximation of  $G = 0.35R_{ns}$  should be used for cases in which the state of the soil heat reservoir cannot be known, for example, if simulations are carried out for pixels in a satellite image taken at a single moment in time.



### 3.5 Improvements in energy balance closure

320 The energy balance loop starts by simulating the radiative transfer of internally generated radiation with initial estimates of component temperatures, followed by the calculation of aerodynamic and stomatal resistances (and photosynthesis), and the fluxes  $H$ ,  $\lambda E$  and  $G$ . Finally, new estimates of the component temperatures are calculated from the value of the energy balance closure error ( $\Delta E$ ) per leaf and soil element. Newton's method is used to estimate the new temperatures, which are the starting point for the next iteration in the loop.

$$325 \quad T_{new} = T_{old} + W \cdot \frac{e_{bal}}{\delta e_{bal}/\delta T} \quad (6)$$

where  $\delta e_{bal}/\delta T$  is the first derivative of the energy balance closure error to temperature, and  $W$  is a weighting for the step size. The derivative is estimated analytically:

$$T_{new} = T_{old} + W \frac{e_{bal}}{\rho \cdot c_p / r_a + \rho \cdot \lambda \cdot M_{H_2O} / M_{air} / p \cdot s / (r_a + r_s) + 4\varepsilon\sigma(T_{old} + 273.15)^3} \quad (7)$$

The derivative is estimated independently for all leaf and soil elements. In the estimate, it is assumed that the incident irradiance on the leaves (or soil) does not change. This is an approximation. The internally (in the canopy) generated incident irradiance depends on the temperature of the neighbouring leaves, which is updated in the next iteration step as well. Further, it is assumed that the resistances  $r_a$  and  $r_s$  do not change between iteration steps. This is an approximation as well, as both depend on leaf and soil temperature. Although these interactions cannot be resolved analytically, Eq. 7 is a sufficiently accurate approximation of the first derivative to obtain rapid energy balance convergence. Iterations continue until the maximum absolute closure error of all leaf and soil elements is less than  $1 \text{ Wm}^{-2}$ , and this is usually achieved in less than 10 iteration steps. If energy balance closure is not achieved after 10 steps, then the weighting coefficient  $W$  is gradually decreased from 1 (i.e. smaller update steps) to avoid the updated temperatures bouncing around the solution.

In earlier versions of SCOPE, a similar equation to Equation 7 has been used to update temperature in the energy balance loop. However, the partial derivative of latent heat flux to temperature was not included in the equation. The improvement in SCOPE 2.0 has substantially reduced the number of required iterations due to a more complete estimate of the derivative.

### 3.6 Angular aggregation of sunlit leaves


In the energy balance routine, the number of sunlit leaf elements that are considered is 13 leaf zenith times 36 leaf azimuth times  $10 \times \text{LAI}$  layers, while the number of shaded leaf elements is  $10 \times \text{LAI}$  since leaf angles play no role for the interactions between a leaf and radiative fluxes when a leaf is illuminated by isotropic diffuse light. Solving the energy budget for all these elements separately means that closure of energy balance should be achieved for each element, and this is computationally demanding. SCOPE 2.0 offers the possibility to simulate the non-radiative energy fluxes, photosynthesis and gas exchange for all inclination and azimuth angles of the sunlit leaves combined (the 'lite' option). This involves an aggregation (weighted averaging) of net radiation over all leaf angles, before entering the energy balance loop. One effective leaf for the  $13 \times 36$  sunlit leaf classes is used for each layer. The resulting number of elements is  $10 \times \text{LAI}$  for the sunlit leaves, and  $10 \times \text{LAI}$  for the shaded leaves. This significantly reduces the computation time of the energy balance routine.



The consequence of this internal aggregation is that the all sunlit leaves in a layer will have an identical temperature, gas exchange, photosynthesis rate, chlorophyll fluorescence emission efficiency, and latent and sensible heat flux, independent of their inclination towards the sun. Figs. 6 and 7 present examples for the effects of the angular aggregation on the profiles of leaf temperature and photosynthesis simulations, respectively. In these simulations, the default model inputs are used (Table 2).

355 Due to the simplifications in the energy balance and biochemical part in the lite mode, the layer average temperatures become slightly higher for both sunlit and shaded leaves (Fig. 6). A slight difference in photosynthetic production between the lite on and lite off modes can be found for sunlit leaves, but the difference for shaded leaves is negligible (Fig. 6). The photosynthetic production simulation for the whole canopy changes by about  $0.7 \mu\text{mol m}^{-2}\text{s}^{-1}$  (4%) when the lite mode is activated. The differences in leaf temperature and photosynthesis are apparently affected by the incoming radiation, leaf biochemistry, canopy


360 structure and other model inputs. The implementation of the lite mode might be helpful for estimation of the added value of consideration of various leaf orientations in a canopy in comparison of the simpler one-big-leaf or two-big-leaf models (Dai et al., 2004; Luo et al., 2018).



**Figure 6.** Layer average kinetic temperatures of the leaves in the vegetation canopy simulated by SCOPE 2.0 with the ‘lite’ representation on (red) and off (blue) of the vegetation, for sunlit (solid lines) and shaded (dash lines) leaves.


With the lite option switched on, the emitted thermal and fluorescence radiation is calculated for layer-average temperature and emission efficiency, respectively, albeit separately for the sunlit and shaded portions. The aggregated layer properties will propagate into the simulation of fluorescence and surface brightness temperature ( $T_b$ ) as observed above the canopy. Fig. 8 presents an example for the effects of the angular aggregation on fluorescence and  $T_b$  simulation with the default model inputs.

365 It is worth noting that the RTMs are all carried out with the original representation of the canopy, thus with  $13 \times 36$  leaf orientations per layer. This means that the lite mode has no influence on reflectance, net radiation in the optical domain, and



**Figure 7.** Layer photosynthesis per LAI in the vegetation canopy simulated by SCOPE 2.0 with the ‘lite’ representation on (red) and off (blue) of the vegetation, for sunlit (solid lines) and shaded (dash lines) leaves.

absorbed photosynthetically active radiation (APAR) by leaves. Moreover, the directionality and hotspot is still simulated (Fig. 370 8).



**Figure 8.** (a) Top-of-canopy fluorescence at 740 nm ( $F_{740}$ ) and (b) surface brightness temperature ( $T_b$ ) simulated by SCOPE 2.0 with the ‘lite’ representation on (red) and off (blue) of the vegetation versus viewing angles in the principle plane.



### 3.7 Improvements in the computational efficiency

In SCOPE 2.0, substantial reductions in computation have been achieved compared to SCOPE 1.70 (Table 4). In a test case of 100 scenarios run by SCOPE 2.0 using a regular PC, the computation time is 12.8% (lite option off) or 7% (lite option on) of the same 100 scenarios run by SCOPE 1.70. The reduction of computation time is due to (in order of decreasing contribution) (1) a more efficient energy balance closure, (2) more efficient saving of output (initially as binary files, later converted to CSV), (3) the overall reduction of the number of layers (from 60 to  $10 \times \text{LAI}$ ) and (4) introducing the mSCOPE radiative transfer equations, which allows for better re-use of earlier calculated quantities. A further reduction in computation time can be achieved when switching off the temperature correction of biochemical parameters (such as  $V_{cmo}$ ) with the option 'tempcor', due to a more rapid convergence of the energy balance loop (not shown).

**Table 4.** Breakdown of computation time to the most computationally intensive modules in SCOPE 1.70 and SCOPE 2.0 expressed as seconds per 100 simulations.

module	computation time (s/100 simulation)		
	SCOPE1.70	SCOPE2.0	
		lite off	lite on
SCOPE self time	0.87	0.51	0.51
BSM	/	0.28	0.28
Fluspect	3.5	1.29	1.26
RTMo	5.75	2.53	2.53
RTMt_planck	33.89	3.69	0.91
RTMf	14.1	0.58	0.63
RTMz	/	0.62	0.64
importdata	0.6	0.03	0.03
ebal	78.8	7.06	1.26
output	32.87	2.3	2.3
the rest	7.22	3.82	3.5
<b>total</b>	<b>177.6</b>	<b>22.7</b>	<b>13.85</b>

### 3.8 Additional outputs

In addition to the output of the original SCOPE model, more model output parameters are produced and stored in SCOPE 2.0, considering users' needs. In Table 5, the outputs available in SCOPE 2.0 but not in the original SCOPE model are presented. Nevertheless, it is worth noting that all the outputs produced in SCOPE 2.0 can also be computed in the original SCOPE with little effort, although they are not stored as outputs.



385 In the original SCOPE model, TOC reflectance spectral simulation in the viewing direction is provided as an output. It is computed as

$$R = \frac{\pi L_o}{E_{sun} + E_{sky}} \quad (8)$$

where  $L_o$  is the radiance in the viewing direction excluding fluorescence contribution, and  $E_{sun}$  and  $E_{sky}$  the incoming direct solar and diffuse sky irradiance.

390 In practice, many users do not have measurements of  $E_{sun}$  and  $E_{sky}$  or atmospheric properties available for inputs but the fraction of diffuse light ( $f_{sky}$ ). Therefore, we provide the directional reflectance factors of the surface as outputs:  $r_{so}$ ,  $r_{sd}$ ,  $r_{dd}$  and  $r_{do}$ . The two-letter subscripts indicate the incident and outgoing fluxes types:  $d$  referring to the diffuse fluxes,  $s$  referring to the direct solar flux and  $o$  referring to the flux in the viewing direction. These four reflectance factors are independent of the incoming irradiance but are optical properties of the soil-vegetation system. The canopy reflectance in the viewing direction  
395 can be estimated as

$$R = (1 - f_{sky})r_{so} + f_{sky}r_{do} \quad (9)$$

Furthermore, the radiance in the viewing direction including the fluorescence contribution is provided, which allows computing the apparent reflectance of a vegetation canopy besides the true reflectance.

We include several fluorescence variables as outputs to help to better interpret fluorescence signals in SCOPE 2.0, besides  
400 fluorescence at top of canopy. Because fluorescence produced by all photosystems is considered to have a more direct relationship with canopy GPP (Yang and Van der Tol, 2018; Van der Tol et al., 2019a), we include it in the outputs. This allows us to compute an important variable: the fluorescence scattering coefficient, which is defined as:

$$\sigma_F = \pi L_o^F / E_F; \quad (10)$$

where  $E_F$  is the total emitted fluorescence irradiance by all photosystems, calculated as the canopy integration of the product  
405 of absorbed photosynthetically active radiation by chlorophyll, the fluorescence yield, and the (constant) spectral shape of chlorophyll fluorescence. The coefficient  $\sigma_F$  is sometimes referred to as the 'escape probability' in the literature. It can be used to correct the fluorescence for both sun-observation geometry and reabsorption of fluorescence in the canopy in order to estimate a canopy-effective fluorescence yield (Yang et al., 2020a).

The biochemical model quantifies the energy partitioning into different pathways and computes their light use efficiencies  
410 at leaf scale. The energy partitioning concept is applied to the whole canopy. By taking the weighted average values of the efficiencies of individual leaves, we obtain canopy electron transport rate and non-photochemical quenching (NPQ), which describes the effective photosynthetic light use efficiency and the effective efficiency of the heat dissipation pathway of the canopy (Maxwell and Johnson, 2000). These variables are direct indicators of the physiological status of the whole canopy.



**Table 5.** Additional outputs in SCOPE 2.0

output	description	unit
<b>spectral simulation</b>		
Lo_spectrum_inclF	radiance in the viewing direction including fluorescence	$[\text{Wm}^{-2}\mu\text{m}^{-1}\text{sr}^{-1}]$
rso, rsd, rdd, rdo	four canopy reflectance factors	[]
sigmaF	fluorescence scattering coefficient	[]
<b>fluorescence_scalars</b>		
LFtot	spectrally integrated observed fluorescence	$[\text{Wm}^{-2}\text{sr}^{-1}]$
EFtot	spectrally and hemispherically integrated fluorescence	$[\text{Wm}^{-2}]$
EFtot_RC	spectrally and hemispherically integrated fluorescence corrected for reabsorption	$[\text{Wm}^{-2}]$
<b>vegetation</b>		
Electron_transport	canopy electron transport rate	$[\mu\text{mol m}^{-2}\text{s}^{-1}]$
NPQ_energy	energy dissipated as non-photochemical quenching	$[\text{Wm}^{-2}]$

#### 4 Conclusions

415 We presented a significantly improved version of the Soil-Canopy Observation of Photosynthesis and Energy fluxes (SCOPE) model. SCOPE 2.0 simulates the energy balance fluxes of net radiation, sensible and latent heat flux, ground heat flux and photosynthesis, as well as hyperspectral radiance in the optical and thermal domain including the contribution of fluorescence.

The improved computational efficiency and model stability make the model a suitable tool for routine estimation of fluxes and satellite signals in homogeneous vegetation canopies with an understory and overstory, or multi-layer structure. The new  
420 features also include the simulation of a subtle change in the reflectance due to the xanthophyll cycle dynamics in the range of 500-600 nm, allowing better investigating vegetation physiology under various weather conditions.

*Author contributions.* CT and PY designed the model general structure of SCOPE2.0. WV and PY developed the BSM model and the multi-layer vegetation parameterization. CT, PY and EP wrote the model code and performed the simulations. EP and CT wrote the online model documentation. PY and CT prepared the manuscript with contributions from all co-authors.

425 *Code availability.* The source code is freely available to users on Github (<https://github.com/Christiaanvandertol/SCOPE2>). SCOPE is written in Matlab and is compatible with versions of 2013a and later. A compiled version is available for the Matlab Runtime Compiler 2019a.

*Competing interests.* The authors declare that there is no competing interests.



*Acknowledgements.* CT was funded by ESA under Contract No.4000122680/17/NL/MP, CNN2. PY was funded by the Netherlands Organization for Scientific Research, grant ALWGO.2017.018. EP has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 721995. Many users contributed with their feedback and suggestions. Particular thanks to: Ari Kornfeld, Christian Frankenberg, Joe Berry, Albert Olivos, Jerome Démarty, Federico Magnani, Jose Moreno, Yves Goulas and Marco Celesti.



## References

- Ač, A., Malenovský, Z., Olejníčková, J., Gallé, A., Rascher, U., and Mohammed, G.: Meta-analysis assessing potential of steady-state chlorophyll fluorescence for remote sensing detection of plant water, temperature and nitrogen stress, *Remote Sensing of Environment*, 168, 420–436, 2015.
- Allen, R. G., Tasumi, M., and Trezza, R.: Satellite-based energy balance for mapping evapotranspiration with internalized calibration (METRIC)—Model, *Journal of irrigation and drainage engineering*, 133, 380–394, 2007.
- Bennett, W. B., Wang, J., and Bras, R. L.: Estimation of global ground heat flux, *Journal of Hydrometeorology*, 9, 744–759, 2008.
- 435 Berk, A., Anderson, G. P., Bernstein, L. S., Acharya, P. K., Dothe, H., Matthew, M. W., Adler-Golden, S. M., Chetwynd Jr, J. H., Richtsmeier, S. C., Pukall, B., et al.: MODTRAN4 radiative transfer modeling for atmospheric correction, in: *Optical spectroscopic techniques and instrumentation for atmospheric and space research III*, vol. 3756, pp. 348–353, International Society for Optics and Photonics, 1999.
- 440 Bhumralkar, C. M.: Numerical experiments on the computation of ground surface temperature in an atmospheric general circulation model, *Journal of Applied Meteorology*, 14, 1246–1258, 1975.
- 445 Bian, Z., Roujean, J.-L., Lagouarde, J.-P., Cao, B., Li, H., Du, Y., Liu, Q., Xiao, Q., and Liu, Q.: A semi-empirical approach for modeling the vegetation thermal infrared directional anisotropy of canopies based on using vegetation indices, *ISPRS Journal of Photogrammetry and Remote Sensing*, 160, 136–148, 2020.
- Collatz, G. J., Ball, J. T., Grivet, C., and Berry, J. A.: Physiological and environmental regulation of stomatal conductance, photosynthesis and transpiration: a model that includes a laminar boundary layer, *Agricultural and Forest meteorology*, 54, 107–136, 1991.
- 450 Collatz, G. J., Ribas-Carbo, M., and Berry, J.: Coupled photosynthesis-stomatal conductance model for leaves of C4 plants, *Functional Plant Biology*, 19, 519–538, 1992.
- Dai, Y., Dickinson, R. E., and Wang, Y.-P.: A two-big-leaf model for canopy temperature, photosynthesis, and stomatal conductance, *Journal of Climate*, 17, 2281–2299, 2004.
- Damm, A., Guanter, L., Paul-Limoges, E., Van der Tol, C., Hueini, A., Buchmann, N., Eugster, W., Ammann, C., and Schaepman, M. E.: Far-red sun-induced chlorophyll fluorescence shows ecosystem-specific relationships to gross primary production: An assessment based on observational and modeling approaches, *Remote Sensing of Environment*, 166, 91–105, 2015.
- 455 Duffour, C., Olioso, A., Demarty, J., Van der Tol, C., and Lagouarde, J.-P.: An evaluation of SCOPE: A tool to simulate the directional anisotropy of satellite-measured surface temperatures, *Remote sensing of environment*, 158, 362–375, 2015.
- Galleguillos, M., Jacob, F., Prévot, L., French, A., and Lagacherie, P.: Comparison of two temperature differencing methods to estimate daily evapotranspiration over a Mediterranean vineyard watershed from ASTER data, *Remote Sensing of Environment*, 115, 1326–1340, 2011.
- 460 Gamon, J., Penuelas, J., and Field, C.: A narrow-waveband spectral index that tracks diurnal changes in photosynthetic efficiency, *Remote Sensing of environment*, 41, 35–44, 1992.
- Guanter, L., Zhang, Y., Jung, M., Joiner, J., Voigt, M., Berry, J. A., Frankenberg, C., Huete, A. R., Zarco Tejada, P., Lee, J.-E., et al.: Global and time-resolved monitoring of crop photosynthesis with chlorophyll fluorescence, *Proceedings of the National Academy of Sciences*, 111, E1327–E1333, 2014.
- 465 Huete, A., Didan, K., Miura, T., Rodriguez, E. P., Gao, X., and Ferreira, L. G.: Overview of the radiometric and biophysical performance of the MODIS vegetation indices, *Remote Sensing of Environment*, 83, 195–213, 2002.
- Jiang, C. and Fang, H.: GSV: a general model for hyperspectral soil reflectance simulation, *International Journal of Applied Earth Observation and Geoinformation*, 83, 101 932, 2019.



- 470 Lekner, J. and Dorf, M. C.: Why some things are darker when wet, *Applied Optics*, 27, 1278–1280, 1988.
- Luo, X., Chen, J. M., Liu, J., Black, T. A., Croft, H., Staebler, R., He, L., Arain, M. A., Chen, B., Mo, G., et al.: Comparison of big-leaf, two-big-leaf, and two-leaf upscaling schemes for evapotranspiration estimation using coupled carbon-water modeling, *Journal of Geophysical Research: Biogeosciences*, 123, 207–225, 2018.
- Maxwell, K. and Johnson, G. N.: Chlorophyll fluorescence—a practical guide, *Journal of experimental botany*, 51, 659–668, 2000.
- 475 Migliavacca, M., Perez-Priego, O., Rossini, M., El-Madany, T. S., Moreno, G., Van der Tol, C., Rascher, U., Berninger, A., Bessenbacher, V., Burkart, A., et al.: Plant functional traits and canopy structure control the relationship between photosynthetic CO<sub>2</sub> uptake and far-red sun-induced fluorescence in a Mediterranean grassland under different nutrient availability, *New Phytologist*, 214, 1078–1091, 2017.
- Mohammed, G. H., Colombo, R., Middleton, E. M., Rascher, U., Van der Tol, C., Nedbal, L., Goulas, Y., Pérez-Priego, O., Damm, A., Meroni, M., et al.: Remote sensing of solar-induced chlorophyll fluorescence (SIF) in vegetation: 50 years of progress, *Remote Sensing of Environment*, 231, 111 177, 2019.
- 480 Nemani, R. R. and Running, S. W.: Estimation of regional surface resistance to evapotranspiration from NDVI and thermal-IR AVHRR data, *Journal of Applied meteorology*, 28, 276–284, 1989.
- Ollinger, S. V.: Sources of variability in canopy reflectance and the convergent properties of plants, *New Phytologist*, 189, 375–394, 2011.
- Pacheco-Labrador, J., Perez-Priego, O., El-Madany, T. S., Julitta, T., Rossini, M., Guan, J., Moreno, G., Carvalhais, N., Martín, M. P., Gonzalez-Cascon, R., et al.: Multiple-constraint inversion of SCOPE. Evaluating the potential of GPP and SIF for the retrieval of plant functional traits, *Remote Sensing of Environment*, 234, 111 362, 2019.
- Ångström, A.: The Albedo of Various Surfaces of Ground, *Geografiska Annaler*, 7, 323–342, 1925.
- Rossini, M., Nedbal, L., Guanter, L., Ač, A., Alonso, L., Burkart, A., Cogliati, S., Colombo, R., Damm, A., Drusch, M., et al.: Red and far red Sun-induced chlorophyll fluorescence as a measure of plant photosynthesis, *Geophysical Research Letters*, 42, 1632–1639, 2015.
- 490 Ryu, Y., Berry, J. A., and Baldocchi, D. D.: What is global photosynthesis? History, uncertainties and opportunities, *Remote Sensing of Environment*, 223, 95–114, 2019.
- Van der Tol, C., Verhoef, W., Timmermans, J., Verhoef, A., and Su, Z.: An integrated model of soil-canopy spectral radiances, photosynthesis, fluorescence, temperature and energy balance., *Biogeosciences*, 6, 2009.
- Van der Tol, C., Berry, J., Campbell, P., and Rascher, U.: Models of fluorescence and photosynthesis for interpreting measurements of solar-induced chlorophyll fluorescence, *J Geophys Res-Biogeophys*, 119, 2312–2327, 2014.
- 495 Van der Tol, C., Vilfan, N., Dauwe, D., Cendrero-Mateo, M. P., and Yang, P.: The scattering and re-absorption of red and near-infrared chlorophyll fluorescence in the models Fluspect and SCOPE, *Remote sensing of environment*, 232, 111 292, 2019a.
- Van der Tol, C., Vilfan, N., Dauwe, D., Cendrero-Mateo, M. P., and Yang, P.: The scattering and re-absorption of red and near-infrared chlorophyll fluorescence in the models Fluspect and SCOPE, *Remote sensing of environment*, 232, 111 292, 2019b.
- 500 Verhoef, W.: Light scattering by leaf layers with application to canopy reflectance modeling: the SAIL model, *Remote Sens Environ*, 16, 125–141, 1984.
- Verhoef, W.: Earth observation modeling based on layer scattering matrices, *Remote Sensing of Environment*, 17, 165–178, 1985.
- Verhoef, W., Van der Tol, C., and Middleton, E. M.: Hyperspectral radiative transfer modeling to explore the combined retrieval of biophysical parameters and canopy fluorescence from FLEX–Sentinel-3 tandem mission multi-sensor data, *Remote sensing of environment*, 204, 942–963, 2018.
- 505



- Verrelst, J., Van der Tol, C., Magnani, F., Sabater, N., Rivera, J. P., Mohammed, G., and Moreno, J.: Evaluating the predictive power of sun-induced chlorophyll fluorescence to estimate net photosynthesis of vegetation canopies: A SCOPE modeling study, *Remote Sensing of Environment*, 176, 139–151, 2016.
- 510 Vilfan, N., Van der Tol, C., Muller, O., Rascher, U., and Verhoef, W.: Fluspect-B: A model for leaf fluorescence, reflectance and transmittance spectra, *Remote Sens Environ*, 186, 596–615, 2016.
- Vilfan, N., Van der Tol, C., Yang, P., Wyber, R., Malenovský, Z., Robinson, S. A., and Verhoef, W.: Extending Fluspect to simulate xanthophyll driven leaf reflectance dynamics, *Remote Sens Environ*, 211, 345–356, 2018.
- Wang, J. and Bras, R.: Ground heat flux estimated from surface soil temperature, *Journal of hydrology*, 216, 214–226, 1999.
- 515 Yang, P. and Van der Tol, C.: Linking canopy scattering of far-red sun-induced chlorophyll fluorescence with reflectance, *Remote Sensing of Environment*, 209, 456–467, 2018.
- Yang, P., Verhoef, W., and Van der Tol, C.: The mSCOPE model: A simple adaptation to the SCOPE model to describe reflectance, fluorescence and photosynthesis of vertically heterogeneous canopies, *Remote sensing of environment*, 201, 1–11, 2017.
- 520 Yang, P., Van der Tol, C., Campbell, P. K., and Middleton, E. M.: Fluorescence Correction Vegetation Index (FCVI): A physically based reflectance index to separate physiological and non-physiological information in far-red sun-induced chlorophyll fluorescence, *Remote sensing of environment*, 240, 111 676, 2020a.
- Yang, P., Van der Tol, C., Yin, T., and Verhoef, W.: The SPART model: A soil-plant-atmosphere radiative transfer model for satellite measurements in the solar spectrum, *Remote Sensing of Environment*, 247, 111 870, 2020b.
- 525 Zhang, Y., Guanter, L., Berry, J. A., Joiner, J., Van der Tol, C., Huete, A., Gitelson, A., Voigt, M., and Köhler, P.: Estimation of vegetation photosynthetic capacity from space-based measurements of chlorophyll fluorescence for terrestrial biosphere models, *Global change biology*, 20, 3727–3742, 2014.