

Ice Algae Model Intercomparison Project phase 2 (IAMIP2)

Hakase Hayashida^{1,2}, Meibing Jin^{3,4,5}, Nadja S. Steiner^{6,7}, Neil C. Swart⁷, Eiji Watanabe⁸, Russell Fiedler⁹, Andrew McC. Hogg^{2,10}, Andrew E. Kiss^{2,10}, Richard J. Matear^{2,9}, Peter G. Strutton^{1,2}

¹Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, TAS, Australia

5 ²Australian Research Council Centre of Excellence for Climate Extremes, Australia

³School of Marine Sciences, Nanjing University of Information Science and Technology, Nanjing, China

⁴Southern Laboratory of Ocean Science and Engineering, Zhuhai, China

⁵International Arctic Research Center, University of Alaska Fairbanks, Fairbanks, AK, USA

⁶Fisheries and Oceans Canada, Institute of Ocean Sciences, Sidney, BC, Canada

10 ⁷Canadian Centre for Climate Modelling and Analysis, Environment and Climate Change Canada, Victoria, BC, Canada

⁸Japan Agency for Marine-Earth Science and Technology, Yokosuka, Kanagawa, Japan

⁹CSIRO Oceans and Atmosphere, Hobart, TAS, Australia

¹⁰Research School of Earth Sciences, Australian National University, Canberra, ACT, Australia

15 *Correspondence to:* Hakase Hayashida (hakase.hayashida@utas.edu.au)

Abstract. Ice algae play a fundamental role in shaping [sea-ice-associated](#) ecosystems and biogeochemistry. This role can be investigated by field observations, however the influence of ice algae at the regional and global scales remains unclear due to limited spatial and temporal coverage of observations, and because ice algae are typically not included in current Earth System Models. To address this knowledge gap, we introduce a new model intercomparison project (MIP), referred to here as the Ice Algae Model Intercomparison Project phase 2 (IAMIP2). IAMIP2 is built upon the experience from its previous phase, and expands its scope to global coverage (both Arctic and Antarctic) and centennial timescales (spanning the mid-twentieth century to the end of the twenty-first century). Participating models are three-dimensional regional and global coupled sea ice–ocean models that incorporate sea-ice ecosystem components. These models are driven by the same initial conditions and atmospheric forcing datasets by incorporating and expanding the protocols of the Ocean Model Intercomparison Project, an endorsed MIP of the Coupled Model Intercomparison Project phase 6 (CMIP6). Doing so provides more robust estimates of model bias and uncertainty, and consequently advances the science of polar marine ecosystems and biogeochemistry. A diagnostic protocol is designed to enhance the reusability of the model data products of IAMIP2. Lastly, the limitations and strengths of IAMIP2 are discussed in the context of prospective research outcomes.

Deleted: polar marine

1 Introduction

30 [Together with pelagic phytoplankton, microalgae](#) that colonize sea ice are the foundation of polar marine food webs. Understanding the susceptibility of ice algae to climate change is therefore essential to comprehending the climatic impacts on higher trophic levels, such as fish, seals, whales, penguins, polar bears, and humans (Cavan et al., 2019; Darnis et al.,

Deleted: Microalgae

Deleted: , referred to here as ice algae,

2012). Vernal blooms of ice algae also directly influence lower trophic levels, such as phytoplankton, zooplankton, and krill, by drawing down near-surface nutrients, reducing light transmission through sea ice, seeding subsequent pelagic blooms, and providing a food source for pelagic and benthic grazers (Leu et al., 2015). Ice algae also regulate ocean biogeochemistry through the biological carbon pump (Mortenson et al., 2020; Watanabe et al., 2015) and polar climates via the production of the climate-active trace gas dimethyl sulfide (Levasseur, 2013) and via the modification of sea-ice albedo (Zeebe et al., 1996).

Field observations of ice algae abundance and distribution are scarce, mainly due to logistical and methodological challenges in these remote and cold environments (Miller et al., 2015), even though technological advancements in recent years have enabled field sampling at much larger scales (Castellani et al., 2020; Cimoli et al., 2020; Lange et al., 2017). Satellite observations are incapable of detecting algae in sea ice. Therefore, the role of ice algae in polar marine ecosystems and biogeochemistry at the regional and global scales remains unclear. One approach to address this knowledge gap is numerical modelling, which simulates ice algae abundance and distribution across the sea-ice domain by incorporating a numerical model of the sea-ice ecosystem into a regional or global three-dimensional coupled sea ice–ocean general circulation model (Vancoppenolle and Tedesco, 2016).

An initial model intercomparison effort was made recently to understand the similarities and differences in simulated ice algae abundance and distribution among the existing three-dimensional models (IAMIP1 hereafter; Watanabe et al., 2019). This model intercomparison investigated the seasonal-to-decadal variability in ice-algal primary productivity in four regions across the Arctic during 1980-2009 simulated by five participating models. The conclusions were: (1) the decadal trend is unclear despite the ongoing reduction of Arctic sea ice; (2) the vernal bloom shifts to an earlier onset and briefer duration over the period of the simulations; and (3) the choice of the maximum growth rate is a key source of the inter-model spread in the simulated ice-algal primary productivity.

Polar regions, especially the Arctic, are warming faster than the rest of the globe, which results in the reduction of both Arctic and Antarctic sea ice (Smith et al., 2019). However, as demonstrated by the previous model intercomparison study (Watanabe et al., 2019), the transient response of ice algae to the anticipated loss of sea ice throughout the twenty-first century may not necessarily be linear. Using a one-dimensional sea-ice biogeochemistry model, Tedesco et al. (2019) found such a non-linear projected response of ice-algal primary productivity to global warming, that also has strong latitudinal dependence. Further investigation using three-dimensional models will provide a comprehensive view of the climate change impacts on polar marine ecosystems and biogeochemistry at the regional-to-global scale, and determine whether ice algae exert any influence on global climate. This knowledge will help to clarify whether ice algae should be incorporated into the next generation of Earth System Models (ESMs), such as those participating in the Coupled Model Intercomparison Project phase 6 (CMIP6; Eyring et al., 2016).

Deleted: Satellite observations are incapable of detecting algae in sea ice. Therefore, the role of ice algae in polar marine ecosystems and biogeochemistry at the regional and global scales remains unknown.

Deleted: (Watanabe et al., 2019)

This paper introduces a new model intercomparison effort for ice algae, referred to here as the Ice Algae Model Intercomparison Project phase 2 (IAMIP2). [IAMIP2 improves on the previous effort \(IAMIP1\) in its experimental design, which is based on the experimental protocols of CMIP6.](#) A consistent experimental design allows us to provide more robust estimates of model bias and uncertainty, and consequently advance the science of polar marine ecosystems and biogeochemistry. The scope of IAMIP2 is global, covering both the Arctic and Antarctic, and centennial timescales, spanning the mid-twentieth century to the end of the twenty-first century. IAMIP2 has five main objectives: (1) evaluation of systematic bias in the existing sea-ice ecosystem models; (2) mechanistic understanding of the past changes in ice algae abundance and distribution; (3) assessment of projected changes and their uncertainty in ice algae abundance and distribution; (4) regional-to-global impacts on marine ecosystems and biogeochemistry; and (5) open-access distribution of the model output for sea-ice research and education.

Deleted: IAMIP2 improves on the previous effort (IAMIP1 hereafter; Watanabe et al., 2019) in its experimental design, which is based on the experimental protocols of CMIP6....

2 Participating models

To date, six three-dimensional coupled sea ice–ocean models have been committed to participating in IAMIP2 (Table 1). [These models represent either a global or regional configuration of the sea-ice and ocean components of the ESMs or global coupled models \(GCMs\) participating in CMIP6.](#) A key difference from these parent model components is that the IAMIP2 models include sea-ice ecosystem components that simulate the biological sources and sinks of ice algae biomass and nutrient concentrations in the bottom ice layer (Fig. 1). [In each model, the sea-ice ecosystem component is coupled to the sea-ice component to account for physical processes that regulate the budgets of ice algae biomass and nutrient concentrations, such as sea-ice growth and melt.](#) The sea-ice ecosystem component is also coupled to ocean dynamics and ecosystem components to simulate the tracer exchange at the sea ice–ocean interface. Three of the IAMIP2 models are global configurations, while the other three are pan-Arctic regional configurations in which the tracer states in the ocean are prescribed at the lateral boundaries. The IAMIP2 models are driven by applying a common atmospheric forcing dataset at the surface boundary. We welcome participation of additional models.

Deleted: All of these

Deleted: sea

Deleted: skeletal

2.1 ACCESS-OM2

ACCESS-OM2 refers to the sea ice–ocean model of the Australian Community Climate Earth System Simulator (ACCESS), and is described in detail in Kiss et al. (2020). In brief, ACCESS-OM2 consists of an ocean dynamics component based on the Modular Ocean Model (MOM) version 5.1 and a sea-ice dynamics component based on the Los Alamos sea ice model (CICE) version 5.1.2. [Notably, these sea ice–ocean physical model components are identical to those adopted for ACCESS-CM2, the GCM of ACCESS contributing to CMIP6 \(Bi et al., 2020\).](#)

Deleted: Notably, these sea ice–ocean physical model components are identical to those adopted for ACCESS-CM2, the GCM of ACCESS contributing to CMIP6 (Bi et al., 2020)...

For IAMIP2, ocean and sea-ice ecosystem components are added to ACCESS-OM2. The ocean ecosystem component is the
115 Whole Ocean Model of Biogeochemistry and Trophic-dynamics (WOMBAT), which consists of nitrate, iron, phytoplankton,
zooplankton, detritus, dissolved oxygen, dissolved inorganic carbon, total alkalinity, and calcium carbonate (Ziehn et al.,
2020). The sea-ice ecosystem component is the Biogeochemistry of CICE (Jeffery et al., 2016), which is based on Jin et al.
(2006). For IAMIP2, ACCESS-OM2 simulates ice algae biomass and nitrate concentration in the bottom-ice layer, which are
120 coupled to phytoplankton biomass and nitrate concentration in the ocean surface layer, respectively. The horizontal
resolution of ACCESS-OM2 is nominally 1° (360 × 300) and the vertical resolution in the ocean surface layer is 2.3 m (Kiss
et al., 2020).

Deleted: skeletal

Deleted: respectively

2.2 CESM-IARC

CESM-IARC refers to the 1° global sea ice–ocean configuration of the Community Earth System Model (CESM) version 1
with a sub-grid-scale brine rejection parameterization that improves ocean mixing under sea ice (Jin et al., 2018). The ocean
125 dynamic component of CESM-IARC is the Parallel Ocean Program (POP) version 2 and the sea-ice dynamic component is
CICE version 5.1.2. The vertical resolution of the ocean surface layer is 10 m. The ocean ecosystem component consists of
multiple nutrients (nitrate, ammonium, iron, silicate, and phosphate), phytoplankton functional groups (diatoms, flagellates,
and diazotrophs), a single zooplankton group, dissolved organic matter (nitrogen, carbon, iron, and phosphorus), dissolved
oxygen, dissolved inorganic carbon, and total alkalinity (Moore et al., 2013). The sea-ice ecosystem component consists of
130 nitrate, ammonium, silicate, and ice algae (Jin et al., 2006).

2.3 RASM

RASM refers to the 1/12° sea ice–ocean configuration of the Regional Arctic System Model, and consists of the same
physical and ecosystem model components as CESM-IARC. The spatial coverage of RASM is pan-Arctic with lateral
boundaries located approximately along the 40 °N Atlantic sector and along the 30 °N Pacific sector (Jin et al., 2018). The
135 vertical resolution of the ocean surface layer is 5 m.

2.4 CanNEMO

CanNEMO is the ocean component of CanESM5 (Swart et al., 2019), which is modified from version 3.4.1 of the Nucleus
for European Modelling of the Ocean (NEMO; Madec and the NEMO team, 2012), and includes the Louvain-la-Neuve sea
Ice Model version 2 (LIM2; Bouillon et al., 2009; Fichefet and Maqueda, 1997). The modifications include the addition of a
140 lee wave mixing scheme based on Saenko et al. (2012) and an update to the mesoscale eddy mixing length-scale (Saenko
et al., 2018), as well as various parameter settings, as described in Swart et al. (2019). CanNEMO is configured on the ORCA1
tripolar grid, with a nominal grid spacing of 1°, refining to 1/3° within 20° of the equator. There are 45 vertical levels,
ranging from 6 m near the surface to 250 m in the abyss. In the version used here, ocean biogeochemistry is represented by
the Canadian Ocean Ecosystem (CanOE) model. CanOE contains two classes of phytoplankton, zooplankton and detritus,

with variable elemental (C/N/Fe) ratios in phytoplankton and fixed ratios for zooplankton and detritus; as well as prognostic carbonate chemistry, and cycles of iron, calcium carbonate and parameterized nitrogen fixation and denitrification (Swart et al., 2019).

150

For IAMIP2, CanNEMO–CanOE is coupled to the Canadian Sea-Ice Biogeochemistry (CSIB) model, which consists of nitrate, ammonium, and ice algae (Hayashida et al., 2019). Modelled ice algae have a higher sensitivity to low-light conditions relative to modelled phytoplankton (Mortenson et al., 2017). When modelled ice algae are released from sea ice into seawater, they contribute partly to the large detritus pool and the seeding of large phytoplankton.

155 2.5 NEMO-NAA

NEMO-NAA is a pan-Arctic regional sea ice–ocean model based on NEMO version 3.4 coupled to LIM2 (Hu and Myers, 2013). The version used here incorporates several modifications that improve the simulation of physical and biogeochemical processes in the Arctic (Hayashida et al., 2019; in revision). Specifically, these modifications include light penetration through snow and sea ice, the vertical resolution of the ocean model, river runoff of biogeochemical tracers, removal of iron dependency of phytoplankton and zooplankton, and changes to sea-ice model parameters. The horizontal grid resolution is 1/4°, which ranges from 10 km near the North American coastline to 14.5 km along the northern Eurasian coastline. The vertical resolution ranges from 1 m for the upper surface layer to 255 m in the deep ocean. Similar to CanNEMO, the ecosystem component of NEMO-NAA is CSIB–CanOE.

160

2.6 COCO–Arctic NEMURO

COCO–Arctic NEMURO refers to a pan-Arctic regional sea ice–ocean model developed at the Japan Agency for Marine-Earth Science and Technology (JAMSTEC; Watanabe et al., 2019). The physical model component is the Center for Climate System Research Ocean Component Model (COCO) version 4.9 (Hasumi, 2006). The sea-ice component of COCO accounts for seven-category distributions of sub-grid snow depth and ice thickness with a one-layer thermodynamic formulation (Bitz and Lipscomb, 1999; Bitz et al., 2001; Lipscomb, 2001) and the elastic–viscous–plastic rheology (Hunke and Dukowicz, 1997). The model domain covers the entire Arctic Ocean with boundaries at the North Atlantic (45 °N) and at the Bering Strait on the Pacific side. The Bering Strait throughflow across the Pacific model boundary is prescribed by idealized seasonal cycles of temperature, salinity, and current velocity based on Woodgate et al. (2005). The model adopts the spherical coordinate system rotated by 90°, which sets the singular points (the North and South Poles of the model grid) at the equator. The horizontal resolution of COCO–Arctic NEMURO is 1/4°, and there are 28 vertical levels in the ocean with variable resolutions from 2 m in the uppermost layer to 500 m below 1,000 m depth. The ecosystem model component is the Arctic and North Pacific Ecosystem Model for Understanding Regional Oceanography (Arctic NEMURO; Watanabe et al., 2015). The ocean ecosystem model component consists of 3 nutrients (nitrate, ammonium, and silicate), 5 plankton types (diatoms, flagellates, microzooplankton, copepod, and predator zooplankton), dissolved organic nitrogen, particulate organic

170

175

Deleted: (EVP)

180 nitrogen, and opal (Kishi et al., 2007). The sea-ice ecosystem model component consists of ice algae, ice-related fauna, and
particulate organic matter (Watanabe et al., 2015).

3 Experiments

The experimental design of IAMIP2 is developed based on the experience from IAMIP1 and the Ocean Model
Intercomparison Project (OMIP; Griffies et al., 2016; Orr et al., 2017), which is an endorsed Model Intercomparison Project
185 (MIP) of CMIP6. The OMIP protocol is useful because OMIP is based on sea ice–ocean models driven by common
atmospheric forcing fields at their surface boundaries. Applying common forcing eliminates uncertainties due to atmospheric
processes and feedbacks, and allows us to focus on differences in the sea-ice, ocean, and biogeochemistry components.

Four numerical experiments are planned for IAMIP2: historical, projection, exclusion, and control as described in detail
190 below and shown in Fig. 2. The IAMIP2 models are expected to conduct all these experiments.

Deleted: of

3.1 Historical

The historical experiment is designed to simulate changes in ice algae abundance and distribution since the mid-twentieth
century, the period for which we have realistic surface-atmospheric conditions based on observations. This experiment spans
the 61 years from January 1, 1958, to December 31, 2018, and uses the Japanese 55-year atmospheric reanalysis for driving
195 sea ice–ocean models version 1.4.0, referred to here as JRA55-do (Tsujino et al., 2018). For reference, version 1.4.0 is also
used for the second phase of OMIP (Tsujino et al., 2020). JRA55-do is regarded as a successor of the Coordinated Ocean-ice
Reference Experiment version 2 forcing dataset (CORE2; Large & Yeager, 2009), and has finer spatial and temporal
resolutions than CORE2. The surface atmospheric variables of JRA55-do are air temperature at 10 m, specific humidity at 10
m, eastward wind at 10 m, northward wind at 10 m, sea level pressure, downward shortwave [radiation](#), downward longwave
200 [radiation](#), rainfall flux, snowfall flux, river runoff, and calving flux (Fig. 1). The original spatial and temporal resolutions of
these variables are respectively $\sim 0.5^\circ$ and 3 hourly except for river runoff and calving flux, which are provided at 0.25° and
daily (Tsujino et al., 2018). However, in reality, the daily calving flux is provided by linear interpolation of monthly data for
Greenland and it is temporally constant for Antarctica (Tsujino et al., 2018). Although JRA55-do provides calving flux, none
of the IAMIP2 models have iceberg components, and therefore it is released into the ocean as meltwater, adopting the
205 approach before version 1.4.0 (Tsujino et al., 2018). JRA55-do is interpolated to the model grid either prior to or during the
experiment.

All IAMIP2 models are initialized from rest ([three-dimensional oceanic velocity fields](#) and [two-dimensional sea level fields](#)
[are all set to zero in the first time step](#)) and with ocean temperature, salinity, dissolved oxygen, and nutrients (nitrate,
210 phosphate, and silicate) from the World Ocean Atlas version 2 (WOA13v2; Garcia et al., 2013; Locarnini et al., 2013;

Deleted: zero

Zweng et al., 2013). More specifically, these fields are the version of WOA13v2 provided for OMIP (Griffies et al., 2016; Orr et al., 2017). There is no recommended protocol for the initialization of sea-ice and other biogeochemical fields. As discussed in Section 2.3 of Griffies et al. (2016), the restoring of sea-surface salinity is necessary to reduce drift in sea ice-ocean models over decadal timescales. However, there is no best practice for salinity restoring because it depends on model details. Therefore, the restoring procedure is left to the discretion of the participating groups, although it is recommended to choose a weak restoring as much as possible to minimize the impact on variability.

220 Three of the IAMIP2 models are based on pan-Arctic regional configurations that require lateral boundary conditions. How to prescribe these conditions is left to the discretion of the participating groups.

Although carbonate chemistry is not the primary focus of IAMIP2, we recommend that modelled total alkalinity and dissolved inorganic carbon should be initialized with the Global Ocean Data Analysis Project version 2 (GLODAPv2; 225 Lauvset et al., 2016) and prescribe the monthly global-mean atmospheric carbon dioxide concentration for the air-sea carbon flux (Meinshausen et al., 2017). These procedures are consistent with OMIP (Orr et al., 2017), and allow us to expand the use of the IAMIP2 product for future research.

3.2 Projection

230 The projection experiments are designed to simulate the projected changes in ice algae abundance and distribution throughout the twenty-first century under two of the greenhouse gas emission scenarios for CMIP6, known as the Shared Socioeconomic Pathways 1-2.6 and 5-8.5 (SSP1-2.6 and SSP5-8.5; O'Neill et al., 2016). SSP1-2.6 is a low emission scenario that informs the Paris Agreement goal of limiting the global warming to below 2 °C of the pre-industrial level. SSP5-8.5 is the highest emission scenario of CMIP6. Therefore, conducting these projections allows us to assess and compare between the impacts of strong mitigation and fossil-fueled development (O'Neill et al., 2016). Each of these 235 experiments spans 86 years from 2015 to 2100. The IAMIP2 models are initialized from states at the end of 2014 in the historical experiment. The atmospheric carbon dioxide concentrations are set to their monthly global-mean values prescribed for its respective SSP (Meinshausen et al., 2020).

3.2.1 Selection of the projected atmospheric forcing dataset

240 The IAMIP2 models are driven by the atmospheric output of selected CMIP6 models that provide the atmospheric forcing fields at the nominal spatial resolution of 100 km and at the temporal resolutions needed for simulating high-frequency (e.g., daily) variability (Holdsworth and Myers, 2015; Lebeaupin Brossier et al., 2012). To date, there are four CMIP6 models that satisfy these criteria: CMCC-CM2-SR5, CMCC-ESM2, EC-Earth3, and MRI-ESM2-0. Specifically, these models provide the simulated atmospheric forcing variables over 2015-2100 under both SSP1-2.6 and SSP5-8.5 at the temporal resolutions identical to those of JRA55-do except for sea level pressure, river runoff, and calving flux. Sea level pressure fields are

Deleted: experiment is

Deleted: highest

Deleted: scenario

Deleted: Pathway

Deleted: . This experiment spans 86 years from 2015 to 2100. The IAMIP2 models are initialized from states at the end of 2014 in the historical experiment. The atmospheric carbon dioxide concentrations are set to their monthly global-mean values prescribed for SSP5-8.5 (Meinshausen et al., 2020).
3.2.1 Selection of the projected atmospheric forcing dataset
The IAMIP2 models are driven by the atmospheric output of selected CMIP6 models that provide the atmospheric forcing fields at the ...

Deleted: variability.

Deleted: three

Deleted: this criterion: ACCESS

Deleted: the

Deleted: scenario

provided 6 hourly, as opposed to 3 hourly in JRA55-do. River runoff fields are provided at monthly, as opposed to daily in JRA55-do. Calving flux fields are not provided at all under [any of the SSP scenarios](#). To partly overcome this limitation, we prescribe a monthly climatology of the calving flux of JRA55-do over 1958-2018 for the projection experiment. This will provide interannually-invariant calving flux for Greenland, and constant calving flux for Antarctica as noted in Section 3.1.

Deleted: SSP5-8.5 scenario

Among the [four](#) CMIP6 models, we choose the output of [CMCC-ESM2](#) and EC-Earth3 for the atmospheric forcing for the projection [experiments](#) because [CMCC-ESM2](#) and EC-Earth3 [provide](#) overall the most realistic atmospheric conditions (in best agreement with JRA55-do) [for the South and North polar oceans, respectively](#). Specifically, we compare the global and polar surface air temperature and major climate modes derived from JRA55-do with those simulated by the following [26](#) CMIP6 models over 1958-2100 (Figures 3 and 4): ACCESS-CM2, ACCESS-ESM1-5, AWI-CM-1-1-MR, BCC-CSM2-MR, CESM2-WACCM, CIESM, [CMCC-CM2-SR5](#), [CMCC-ESM2](#), CanESM5, EC-Earth3, EC-Earth3-Veg, FGOALS-f3-L, FGOALS-g3, FIO-ESM-2-0, GFDL-ESM4, INM-CM4-8, INM-CM5-0, IPSL-CM6A-LR, KACE-1-0-G, MIROC6, MPI-ESM1-2-HR, MPI-ESM1-2-LR, MRI-ESM2-0, NESM3, NorESM2-LM, and NorESM2-MM. Although the objective here is to compare among the [four](#) candidates for the projected atmospheric forcing, we include the remaining [22](#) CMIP6 models that have provided the output at monthly resolution as well as the multi-model mean product of the [26](#) models for reference. These model products belong to Variant Label *r1i1p1f1* of Experiment IDs *historical* and *ssp585*, and [they](#) were downloaded from <https://esgf-node.llnl.gov/> (accessed on July [15, 2021](#)).

Deleted: three

Deleted: experiment

Deleted: provides

Deleted:).

Deleted: 24

Deleted: three

Deleted: 19

Deleted: 24

Deleted: 1, 2020

Formatted: English (UK)

3.2.2 Comparison of surface air temperature [over the ocean](#)

Surface air temperatures are averaged over the ocean grid cells globally, north of 60 °N, and south of 60 °S (Figure 3). In terms of global averages, [EC-Earth3](#) simulates surface air temperature [closest to](#) JRA55-do prior to 2000 (Figure 3a). After 2000, EC-Earth3 simulates warmer surface air temperature that agrees well with [CMCC-CM2-SR5](#) but [deviates from](#) JRA55-do. [At the beginning of the SSP projections \(2015\), the global average of CMCC-ESM2 matches well with JRA55-do.](#) [MRI-ESM2-0 consistently underestimates the global averages.](#) Compared with the multi-model averages, EC-Earth3 [projects](#) greater global warming [especially in the latter half of the 21st century](#), whereas MRI-ESM2-0 [projecton is close to the multi-model averages](#) (Figure 3b).

Deleted: the three candidates simulate roughly 1 °C colder

Deleted: than

Deleted: and ACCESS-CM2 project

Deleted: throughout

Deleted: projects lesser warming in the late 21st century

In terms of Arctic averages, MRI-ESM2-0 simulates surface air temperature that is in closest agreement with JRA55-do, but EC-Earth3 [compares well](#) later in the 2010s (Figure 3c). [In 2015, EC-Earth3 is closest to JRA55-do among the four candidates.](#) Polar amplification is evident in all models, but is more so in EC-Earth3 [\(>15 °C warming by the end of the 21st century relative to the 1958-2014 average;](#) Figure 3d).

Deleted: again

Deleted: and ACCESS-CM2 (>10

Deleted: their

Deleted: averages

In the Antarctic, the average air temperatures of CMCC-CM2-SR5 and CMCC-ESM2 agree by far the best with JRA55-do compared to the other two candidates (Figure 3e). EC-Earth3 simulates roughly 4 °C warmer than JRA55-do, whereas MRI-ESM2-0 simulates approximately 3 °C colder surface air over the South polar ocean. MRI-ESM2-0 exhibits greater interannual variability throughout the 21st century and the greatest Antarctic warming among the four candidates (approximately 5 °C warming by 2100 relative to the 1958-2014 average; Figure 3f).

3.2.3 Comparison of major climate modes of variability

Three major modes of climate variability considered here are the El Niño–Southern Oscillation (ENSO), the Northern Annular Mode (NAM), and the Southern Annular Mode (SAM). ENSO is the dominant mode of climate variability for the globe, which can be characterised by an index called the Equatorial Southern Oscillation Index (EQSOI; Bell and Halpert, 1998). EQSOI is the standardised sea level pressure difference between two regions over Indonesia (80 °W-130 °W, 5 °S-5 °N) and the eastern equatorial Pacific (90 °E-140 °E, 5 °S-5 °N). Because the CMIP6 models have their own ENSO variability, EQSOIs of these models are not expected to be in phase with that of JRA55-do (this is applicable also to other modes of climate variability, such as NAM and SAM discussed below). Instead, we compare the magnitude of the sea level pressure difference, which determines the strength of the easterly winds along the equator (Figure 4a). Furthermore, we compare the power spectra of EQSOI over 1958-2014 (Figure 4b).

CMCC-CM2-SR5 and CMCC-ESM2 simulate the sea level pressure difference for EQSOI that is in closer agreement with JRA55-do than the other two candidates whose sea level pressure difference is about 100 Pa lower than JRA55-do (Figure 4a). JRA55-do shows two maxima in the spectrum of EQSOI at periods between 3 and 6 years (Figure 4b). MRI-ESM2-0 captures the one of these maxima at period of about 3-4 years, while CMCC-ESM2 exhibits the other maximum at about 5-6 years. EC-Earth3 shows a maximum at period of 4-5 years. In contrast, the EQSOI signal of CMCC-CM2-SR5 is peaked at a longer period of 7-8 years.

NAM is the dominant mode of climate variability in the Northern Hemisphere, which can be characterised by the sea level pressure difference between the zonal means at 35 °N and 65 °N (Li and Wang, 2003). JRA55-do overall exhibits smaller sea level pressure difference than any of the four candidates or the multi-model averages (Figure 4c). MRI-ESM2-0 simulates somewhat larger sea level pressure difference than the other three candidates that are close to the multi-model averages. The power spectra of NAM are noisier at higher-frequency (<1 year) and the peaks are spread more broadly than those of EQSOI among the candidates and JRA55-do (Figure 4d). JRA55-do shows peaks at about 0.5, 1, and 10 years, indicating the importance of seasonal, interannual, and interdecadal variability of NAM, respectively. The four candidates also exhibit peaks within this seasonal-to-interdecadal time scale, but differ in terms of the exact locations of these peaks. The strongest signal of NAM is present at period of roughly 0.5-1 year (MRI-ESM2-0), 3 years (CMCC-CM2-SR5 and CMCC-ESM2), and 4-5 years (EC-Earth3).

Deleted: temperature

Deleted: EC-Earth3 agrees

Deleted: rest of the

Deleted: Besides EC-Earth3, only two other CMIP6 models simulate EC-Earth3 simulates

Deleted: temperature that is within 2 °C of JRA55-do.

Deleted: and highly-variable

Deleted: than ACCESS-CM2 and EC-Earth3 (

Deleted: (EQSOI; Bell and Halpert, 1998)

Deleted: ACCESS

Deleted: simulates

Deleted: EC-Earth3 and MRI-ESM2-0 (Figure 4a). These

Deleted: models simulate about 100 Pa lower

Deleted: . The three models agree with

Deleted: simulating strong signals

Deleted: >2

Deleted: shows a distinct maximum in

Deleted: spectrum

Deleted: .

Deleted: EC-Earth3 simulates closer

Deleted: to JRA55-do and

Deleted: mean than ACCESS-CM2 and MRI-ESM2-0

Deleted: three models

Deleted: power spectra of ACCESS-CM2 and EC-Earth3 are strongest at around 5 years, whereas the ...

Deleted: in MRI-ESM2-0

Deleted: less than

Deleted: .

375 SAM is the dominant mode of climate variability in the Southern Hemisphere, which can be characterised by the sea level
 pressure difference between the zonal means at 40 °S and 65 °S (Gong and Wang, 1999). ~~EC-Earth3 and MRI-ESM2-0~~
 compare equally well with JRA55-do in simulating the sea level pressure difference for SAM, which is lower than the multi-
 model mean of the 26 CMIP6 models (Figure 4e). ~~In contrast, CMCC-CM2-SR5 and CMCC-ESM2 simulate about 500 Pa~~
 380 ~~greater sea level pressure difference than JRA55-do.~~ Similar to NAM, the power spectra of SAM are distributed across the
 broad time scales, but the peaks occur at periods of ~~about 1 year or greater~~ (Figure 4f). The only exception is EC-Earth3, in
 which the strongest signal is present at period of slightly less than 1 year. ~~CMCC-CM2-SR5 agrees well~~ with JRA55-do in
 simulating ~~the absolute maximum at period of 1-2 years.~~ Consistent among the ~~four candidates~~ and JRA55-do, SAM exhibits
 appreciable variability also at lower frequencies, as indicated by the presence of local maxima at periods of 3-6 years and
 beyond 10 years. Notably, this low-frequency variability is the strongest signature of SAM simulated by MRI-ESM2-0.

385 3.2.4 Recommendations for the projected atmospheric forcing dataset

In summary, all ~~four~~ candidates for the projected atmospheric forcing dataset simulate reasonable historical atmospheric
 conditions compared to JRA55-do. ~~An~~ exception is the substantially colder surface air temperature over the Antarctic
 oceanic region, (Figure 3e), for which ~~CMCC-CM2-SR5 and CMCC-ESM2~~ are superior to the other two models. ~~In contrast,~~
 390 ~~the latter two models outperform the former in simulating surface air temperature over the Arctic oceanic region (Figure 3c).~~
~~Therefore, there is no single candidate that does a better job than the others at both poles.~~ For this reason, we adopt the
 output of ~~two models, CMCC-ESM2-0 and EC-Earth3,~~ as the standard atmospheric forcing dataset for the projection
 experiments of IAMIP2. ~~With two SSP scenarios, there are four combinations (CMCC-ESM2-0 ssp126, CMCC-ESM2-0~~
 395 ~~ssp585, EC-Earth3 ssp126, and EC-Earth3 ssp585; Table 2).~~ We encourage the IAMIP2 participants to perform all the four
 projections. ~~However, if computational resources are limited to undertake only one projection, we request one of the SSP5-~~
 8.5 projections, either ~~CMCC-ESM2-0 ssp858 (for Antarctic-focused models) or EC-Earth3 ssp585 (for Arctic-focused~~
 models). ~~Depending~~ on computational resources, additional projection experiments using the output of the other two
 candidates as well as other CMIP6 models may be considered if they provide the output at high temporal resolution. Such
 additional experiments will allow assessment of the sensitivity of the IAMIP2 models to projected atmospheric conditions.

3.3 Exclusion

400 The exclusion experiment is designed to simulate ocean biogeochemistry in the absence of ice algae, which is the case for all
 CMIP6 models. This experiment is done in the same set up as the ~~historical~~ experiment except that the sea-ice ecosystem
 component is excluded. Comparing the results of the ~~historical~~ and exclusion experiments allows us to quantify the impacts
 of ice algae on polar marine lower-trophic-level ecosystems and ~~the~~ biological carbon pump. Doing so assesses the
 significance of incorporating sea-ice ecosystem components into the next-generation ESMs.

Deleted: The three candidates

Deleted: 24

Deleted: >

Deleted: ACCESS

Deleted: and EC-Earth3 agree

Deleted: strong SAM signals

Deleted: periods

Deleted: around

Deleted: , whereas such signals are lacking in MRI-ESM2-0.

Deleted: three models

Deleted: three

Deleted: The only

Deleted: ,

Deleted: EC-Earth3 is

Deleted: under the SSP5-8.5 scenario

Deleted: experiment

Deleted: depending

Deleted:

Deleted: projection

Deleted: projection

Deleted: under global warming

3.4 Control

The control experiment is designed to diagnose artificial model drifts and to distinguish anthropogenic effects from natural variability. This experiment spans 143 years by repeating the annual cycle of JRA55-do from May 1, 1990 to April 30, 1991, during which major climate modes were all in neutral phases (Stewart et al., 2020).

430 4 Diagnostics

The following guidelines are developed for IAMIP2 in order to implement the Findability, Accessibility, Interoperability, and Reusability (FAIR) data principles (Wilkinson et al., 2016) and ensure that the IAMIP2 product reaches the end users (Objective 5). Specifically, we make the product discoverable from the website <http://cosima.org.au/index.php/working-groups/iamp2> which provides a list of hyperlinks to the IAMIP2 product archived by individual modelling groups.

435

All model diagnostics are saved as daily averages in the NetCDF format, and where applicable, their names must follow the CMIP6 naming conventions (Table 2). Daily temporal resolution is needed to quantify the bloom phenology (Watanabe et al., 2019) and to provide the ocean and sea-ice climate data for driving sea-ice biogeochemical models (Lavoie et al., 2010; Tedesco et al., 2019). To limit data storage needs, only two-dimensional fields are saved, and to preserve the spatial details in regional models, they are stored on the [models' native grids](#). To perform interpolation for analysis, an additional file containing longitude, latitude, and grid-cell area needs to be provided for each model. [The output should be stored using a common format for directory and file names \(Figure 5\)](#).

440

[A few examples of potential use of these diagnostics for investigating various roles of ice algae are illustrated here. Their ecological role as the foundation of the polar marine food web and their relative importance can be quantified by comparing the biomass \(*phycbi* and *phycos*; Table 2\) and primary productivity between ice algae and phytoplankton \(*intppbi* and *intpp*\). The latter quantities can also be used to quantify the biogeochemical role of ice algae in carbon fixation and their contribution to the biological carbon pump can be assessed using particulate organic carbon export \(*epc100*\). A combination of physical \(*siconc*, *paros*, *sst*, and *mlots2t*\) and biogeochemical diagnostics \(*no3os* and *phvos*\) can be used to estimate dimethyl sulfide concentration \(e.g., Bock et al., 2021; Galí et al., 2019\) as well as its emission using the wind speed which is available as the atmospheric forcing fields \(*uas* and *vas*; Figure 1\).](#)

445

450

5 Discussion

Although the experimental design of IAMIP2 advances from that of IAMIP1 (Watanabe et al., 2019) in many aspects, it has its own limitations which are discussed here to help the interpretation of results and for consideration [by prospective end users](#). First, the vertical extent of the sea-ice ecosystem component of the currently committed selection of IAMIP2 models

455

Deleted: model's

Deleted: grid

Deleted: of

460 is restricted to the bottom-ice layer, which likely underestimates depth-integrated ice algae biomass and primary production
because ice algae are present throughout the sea-ice column, especially in the Antarctic. Although simulating ice algae
throughout the sea-ice column is desirable and has been conducted under one-dimensional settings (Duarte et al., 2015;
Pogson et al., 2011) as well as in a recent study using an ESM (Jeffery et al., 2020), its implementation into high-resolution
three-dimensional models is computationally expensive. We anticipate that this limitation has a negligible effect on the
estimation of depth-integrated biomass and primary production in landfast sea ice (Meiners et al., 2018), but it could
465 underestimate these quantities substantially for pack ice as demonstrated by field observations (Meiners et al., 2012). We
consider this a necessary compromise in order to progress IAMIP2 with currently-available computational resources.

Deleted: skeletal

Deleted: .

Deleted: Nevertheless, we

Deleted: as demonstrated by field observations

Deleted: .

470 For projections of sea ice-ocean models, a few previous studies have prescribed a synthetic atmospheric forcing dataset
instead of applying the raw output of climate models in order to ensure that there is no undesirable step change in the
atmospheric conditions at the beginning of projections (Naughten et al., 2018) and that the high-frequency climate variability
is simulated realistically (Zhang et al., 2017). In contrast to the approach taken by these previous studies, the IAMIP2
models are driven by the raw output of CMIP6 model projections, which is made possible thanks to four CMIP6 models so
far that have provided their atmospheric output available at the temporal resolutions needed to simulate the high-frequency
climate variability (Holdsworth and Myers, 2015; Lebeaupin Brossier et al., 2012). Our approach is advantageous over that
475 of the previous studies in that the IAMIP2 model projections account for projected changes in both the high- and low-
frequency climate variability. As demonstrated in Section 3.2, there appears to be no single CMIP6 model that outperforms
the other models considered here in simulating surface air temperature in both marine polar regions. Some models are better
at simulating Arctic surface air temperature but not the Antarctic, and vice versa. Considering this finding, we aim to
perform multiple projections by prescribing the atmospheric output of two CMIP6 models (EC-Earth3 and CMCC-ESM2)
480 under two emission scenarios (SSP1-2.6 and SSP5-8.5). To a certain extent, these projections will allow us to investigate the
uncertainty in the projected changes in ice algae abundance and distribution due to the uncertainty in the projected
atmospheric conditions under the same climate change scenarios. Furthermore, the implications of climate change scenarios
for polar marine ecosystems and biogeochemistry can be addressed by comparison of the SSP1-2.6 and SSP5-8.5
projections.

Deleted: three

Deleted: .

Deleted: EC-Earth3 simulates the atmospheric conditions most comparable to JRA55-do especially in terms of Antarctic surface air temperature, and therefore its projected atmospheric output is adopted as the default atmospheric forcing dataset for the IAMIP2 projection experiment. However, it would be worthwhile investigating there

Deleted: among the CMIP6 models

Deleted: Similarly

Deleted: performing additional experiments based on other SSP scenarios (O'Neill et al., 2016). Given the commitment to undertake the experiments outlined in Section 3, such experiments are not the top priority of IAMIP2, but may be considered in the future, or alternatively can be investigated individually by the participating modelling groups...

485 One important source of biases in IAMIP2 projections by regional models is the lateral boundary conditions for nutrients. As
a limiting factor for primary production in sea ice, prescribing the temporally-varying nutrient boundary conditions are
desirable. However, this is not an easy task given the large uncertainty in projected changes in nutrients especially in polar
regions (e.g., Lannuzel et al., 2020).

490 IAMIP2 is an ongoing international effort aiming primarily to understand the role of ice algae in polar marine ecosystems
and biogeochemistry at the regional and hemispheric scales. This paper describes the design of IAMIP2 which is built upon

the experience from IAMIP1 (Watanabe et al., 2019) and by keeping up to date with the CMIP6 protocols (Griffies et al., 515 2016; O'Neill et al., 2016; Orr et al., 2017). Six three-dimensional regional and global coupled sea ice-ocean models are currently committed to participating in IAMIP2. These models are driven by the same initial conditions and atmospheric forcing dataset to assess systematic biases in these models in terms of simulating ice algae abundance and distribution. Five numerical experiments are designed to understand the past changes since the mid-twentieth century by applying realistic atmospheric forcing (JRA55-do) as well as the projected changes throughout the twenty-first century by applying the high-520 temporal-resolution output of selected CMIP6 model projections under the SSP1-2.6 and SSP5-8.5 scenarios. Two other experiments are used to separate the anthropogenic effect from natural variability and quantify the large-scale impacts of incorporating ice algae into regional and global models on polar marine ecosystems and biogeochemistry. The model data products of IAMIP2 are expected to meet the FAIR data principles and intended to be used for future research and as educational tools. In conclusion, IAMIP2 is expected to advance the science of polar marine ecosystems and 525 biogeochemistry.

Deleted: Two

Deleted: projection

Deleted: scenario

Data and code availability

The model output of IAMIP2 will be made available by individual modelling groups, which will be discoverable from the website <http://cosima.org.au/index.php/working-groups/iamip2>. The atmospheric forcing datasets for the four projection experiments are available on the National Computational Infrastructure (NCI) National Research Data Collection (Hayashida, 2021). 530

Deleted: .

Author contributions

HH conceived IAMIP2 and developed the details with MJ, NSS, NCS, and EW. AH, AK, RM, and PS supervised the ACCESS-OM2 contribution to IAMIP2. RF and HH developed the model code for enabling sea ice-ocean coupling of biogeochemistry in ACCESS-OM2. HH prepared the projected atmospheric forcing datasets, performed the analysis, and 535 wrote the paper with input from all co-authors.

Competing interests

All authors declare that they have no conflict of interest.

Acknowledgements

HH thanks the members of the Consortium for Ocean-Sea Ice Modelling in Australia (COSIMA) for discussion on atmospheric forcing dataset for future projections. This dataset is made available through technical support from Paola 540

Deleted: HH, AH, AK, RM, and PS acknowledge

[Petrelli from the CMS team](#) of the Australian Research Council Centre of Excellence for Climate Extremes ([CLEX](#)). [HH, AH, AK, RM, and PS](#) acknowledge support from [CLEX](#) (CE170100023). This study was undertaken with the assistance of resources and services from the [NCI](#), which is supported by the Australian Government. Analyses are performed using the Climate Data Operator (CDO) software (Schulzweida, 2019) and Python. Normalised power spectra are computed using SciPy (Virtanen et al., 2020). EW is supported by the Grant-in-Aid for Scientific Research of Japan Society for the Promotion of Science (JSPS) (KAKENHI 18H03368) and the Arctic Challenge for Sustainability II (ArCS II) project (JPMXD1420318865) in Japan. NSS and NCS acknowledge funding through the Departments of Environment and Climate Change Canada and Fisheries and Oceans Canada. This work is a contribution to the Biogeochemical Exchange processes at Sea Ice Interfaces (BEPSII) network.

Deleted: National Computational Infrastructure (

Deleted:),

555 References

Bell, G.D., and Halpert, M.S. (1998). Climate Assessment for 1997. *Bull. Am. Meteorol. Soc.* *79*, S1–S50.

Bi, D., Dix, M., Marsland, S., O'Farrell, S., Sullivan, A., Bodman, R., Law, R., Harman, I., Srbinovsky, J., and Rashid, H. (2020). Configuration and spin-up of ACCESS-CM2, the new generation Australian Community Climate and Earth System Simulator Coupled Model. *J. South. Hemisphere Earth Syst. Sci.* Submitt.

560 Bitz, C.M., and Lipscomb, W.H. (1999). An energy-conserving thermodynamic model of sea ice. *J. Geophys. Res. Oceans* *104*, 15669–15677.

Bitz, C.M., Holland, M.M., Weaver, A.J., and Eby, M. (2001). Simulating the ice-thickness distribution in a coupled climate model. *J. Geophys. Res. Oceans* *106*, 2441–2463.

565 [Bock, J., Michou, M., Nabat, P., Abe, M., Mulcahy, J.P., Olivié, D.J.L., Schwinger, J., Suntharalingam, P., Tjiputra, J., van Hulst, M., et al. \(2021\). Evaluation of ocean dimethylsulfide concentration and emission in CMIP6 models. *Biogeosciences* *18*, 3823–3860.](#)

Bouillon, S., Maqueda, M.A.M., Legat, V., and Fichefet, T. (2009). An elastic–viscous–plastic sea ice model formulated on Arakawa B and C grids. *Ocean Model.* *27*, 174–184.

570 [Castellani, G., Schaafsma, F.L., Arndt, S., Lange, B.A., Peeken, I., Ehrlich, J., David, C., Ricker, R., Krumpen, T., Hendricks, S., et al. \(2020\). Large-Scale Variability of Physical and Biological Sea-Ice Properties in Polar Oceans. *Front. Mar. Sci.* *7*.](#)

Cavan, E.L., Belcher, A., Atkinson, A., Hill, S.L., Kawaguchi, S., McCormack, S., Meyer, B., Nicol, S., Ratnarajah, L., Schmidt, K., et al. (2019). The importance of Antarctic krill in biogeochemical cycles. *Nat. Commun.* *10*, 4742.

575 [Cimoli, E., Lucieer, V., Meiners, K.M., Chennu, A., Castrisios, K., Ryan, K.G., Lund-Hansen, L.C., Martin, A., Kennedy, F., and Lucieer, A. \(2020\). Mapping the in situ microspatial distribution of ice algal biomass through hyperspectral imaging of sea-ice cores. *Sci. Rep.* *10*, 21848.](#)

- 580 Darnis, G., Robert, D., Pomerleau, C., Link, H., Archambault, P., Nelson, R.J., Geoffroy, M., Tremblay, J.-É., Lovejoy, C.,
Ferguson, S.H., et al. (2012). Current state and trends in Canadian Arctic marine ecosystems: II. Heterotrophic food web,
pelagic-benthic coupling, and biodiversity. *Clim. Change* *115*, 179–205.
- Duarte, P., Assmy, P., Hop, H., Spreen, G., Gerland, S., and Hudson, S.R. (2015). The importance of vertical resolution in
sea ice algae production models. *J. Mar. Syst.* *145*, 69–90.
- 585 Eyring, V., Bony, S., Meehl, G.A., Senior, C.A., Stevens, B., Stouffer, R.J., and Taylor, K.E. (2016). Overview of the
Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization. *Geosci. Model Dev.* *9*,
1937–1958.
- Fichefet, T., and Maqueda, M.A.M. (1997). Sensitivity of a global sea ice model to the treatment of ice thermodynamics and
dynamics. *J. Geophys. Res. Oceans* *102*, 12609–12646.
- 590 [Galí, M., Devred, E., Babin, M., and Levasseur, M. \(2019\). Decadal increase in Arctic dimethylsulfide emission. *Proc. Natl. Acad. Sci.* 201904378.](#)
- Garcia, H.E., Locarnini, R.A., Boyer, T.P., Antonov, J.I., Baranova, O.K., Zweng, M.M., Reagan, J.R., Johnson, D.R.,
Mishonov, A.V., and Levitus, S. (2013). World ocean atlas 2013. Volume 4, Dissolved inorganic nutrients (phosphate,
nitrate, silicate).
- Gong, D., and Wang, S. (1999). Definition of Antarctic Oscillation index. *Geophys. Res. Lett.* *26*, 459–462.
- 595 Griffies, S.M., Danabasoglu, G., Durack, P.J., Adcroft, A.J., Balaji, V., Böning, C.W., Chassignet, E.P., Curchitser, E.,
Deshayes, J., Drange, H., et al. (2016). OMIP contribution to CMIP6: experimental and diagnostic protocol for the physical
component of the Ocean Model Intercomparison Project. *Geosci. Model Dev.* *9*, 3231–3296.
- Hasumi, H. (2006). CCSR Ocean Component Model (COCO) version 4.0 (University of Tokyo).
- 600 [Hayashida, H. \(2021\). Atmospheric forcing dataset for the Ice Algae Model Intercomparison Project phase 2 \(IAMIP2\) projection experiment v1.0. NCI Natl. Res. Data Collect.](#)
- [Hayashida, H.](#), Christian, J.R., Holdsworth, A.M., Hu, X., Monahan, A.H., Mortenson, E., Myers, P.G., Riche, O.G.J., Sou,
T., and Steiner, N.S. (2019). CSIB v1 (Canadian Sea-ice Biogeochemistry): a sea-ice biogeochemical model for the NEMO
community ocean modelling framework. *Geosci. Model Dev.* *12*, 1965–1990.
- 605 [Holdsworth, A.M., and Myers, P.G. \(2015\). The Influence of High-Frequency Atmospheric Forcing on the Circulation and Deep Convection of the Labrador Sea. *J. Clim.* *28*, 4980–4996.](#)
- Hu, X., and Myers, P.G. (2013). A Lagrangian view of Pacific water inflow pathways in the Arctic Ocean during model
spin-up. *Ocean Model.* *71*, 66–80.
- Hunke, E.C., and Dukowicz, J.K. (1997). An elastic-viscous-plastic model for sea ice dynamics. *J. Phys. Oceanogr.* *27*,
1849–1867.
- 610 Jeffery, N., Elliott, S.M., Hunke, E.C., Lipscomb, W.H., and Turner, A.K. (2016). Biogeochemistry of Cice: The Los
Alamos Sea Ice Model Documentation and Software User's Manual Zbgc_colpkg Modifications to Version 5 (Los Alamos
National Lab. (LANL), Los Alamos, NM (United States)).

- Jeffery, N., Maltrud, M.E., Hunke, E.C., Wang, S., Wolfe, J., Turner, A.K., Burrows, S.M., Shi, X., Lipscomb, W.H., Maslowski, W., et al. (2020). Investigating controls on sea ice algal production using E3SMv1.1-BGC. *Ann. Glaciol.* 1–22.
- 615 Jin, M., Deal, C.J., Wang, J., Shin, K.-H., Tanaka, N., Whitledge, T.E., Lee, S.H., and Gradinger, R.R. (2006). Controls of the landfast ice–ocean ecosystem offshore Barrow, Alaska. *Ann. Glaciol.* 44, 63–72.
- Jin, M., Deal, C., Maslowski, W., Matrai, P., Roberts, A., Osinski, R., Lee, Y.J., Frants, M., Elliott, S., Jeffery, N., et al. (2018). Effects of Model Resolution and Ocean Mixing on Forced Ice-Ocean Physical and Biogeochemical Simulations Using Global and Regional System Models. *J. Geophys. Res. Oceans* 123, 358–377.
- 620 Kishi, M.J., Kashiwai, M., Ware, D.M., Megrey, B.A., Eslinger, D.L., Werner, F.E., Noguchi-Aita, M., Azumaya, T., Fujii, M., Hashimoto, S., et al. (2007). NEMURO—a lower trophic level model for the North Pacific marine ecosystem. *Ecol. Model.* 202, 12–25.
- Kiss, A.E., Hogg, A.M., Hannah, N., Boeira Dias, F., Brassington, G.B., Chamberlain, M.A., Chapman, C., Dobrohotoff, P., Domingues, C.M., Duran, E.R., et al. (2020). ACCESS-OM2 v1.0: a global ocean–sea ice model at three resolutions. *Geosci. Model Dev.* 13, 401–442.
- 625 [Lange, B.A., Katlein, C., Castellani, G., Fernández-Méndez, M., Nicolaus, M., Peeken, I., and Flores, H. \(2017\). Characterizing Spatial Variability of Ice Algal Chlorophyll a and Net Primary Production between Sea Ice Habitats Using Horizontal Profiling Platforms. *Front. Mar. Sci.* 4.](#)
- [Lannuzel, D., Tedesco, L., van Leeuwe, M., Campbell, K., Flores, H., Delille, B., Miller, L., Stefels, J., Assmy, P., Bowman, J., et al. \(2020\). The future of Arctic sea-ice biogeochemistry and ice-associated ecosystems. *Nat. Clim. Change* 1–10.](#)
- 630 Large, W.G., and Yeager, S.G. (2009). The global climatology of an interannually varying air–sea flux data set. *Clim. Dyn.* 33, 341–364.
- Lauvset, S.K., Key, R.M., Olsen, A., van Heuven, S., Velo, A., Lin, X., Schirnick, C., Kozyr, A., Tanhua, T., Hoppema, M., et al. (2016). A new global interior ocean mapped climatology: the 1-deg. by 1-deg. GLODAP version 2. *Earth Syst. Sci. Data* 8, 325–340.
- 635 Lavoie, D., Denman, K.L., and Macdonald, R.W. (2010). Effects of future climate change on primary productivity and export fluxes in the Beaufort Sea. *J. Geophys. Res. Oceans* 115, C04018.
- [Lebeaupin Brossier, C., Béranger, K., and Drobinski, P. \(2012\). Sensitivity of the northwestern Mediterranean Sea coastal and thermohaline circulations simulated by the 1/12°-resolution ocean model NEMO-MED12 to the spatial and temporal resolution of atmospheric forcing. *Ocean Model.* 43–44, 94–107.](#)
- 640 Leu, E., Mundy, C.J., Assmy, P., Campbell, K., Gabrielsen, T.M., Gosselin, M., Juul-Pedersen, T., and Gradinger, R. (2015). Arctic spring awakening – Steering principles behind the phenology of vernal ice algal blooms. *Prog. Oceanogr.* 139, 151–170.
- Lavasaur, M. (2013). Impact of Arctic meltdown on the microbial cycling of sulphur. *Nat. Geosci.* 6, 691–700.
- 645 Li, J., and Wang, J.X.L. (2003). A modified zonal index and its physical sense. *Geophys. Res. Lett.* 30.
- Lipscomb, W.H. (2001). Remapping the thickness distribution in sea ice models. *J. Geophys. Res. Oceans* 106, 13989–14000.

- Locarnini, R.A., Mishonov, A.V., Antonov, J.I., Boyer, T.P., Garcia, H.E., Baranova, O.K., Zweng, M.M., Paver, C.R., Reagan, J.R., and Johnson, D.R. (2013). World ocean atlas 2013. Volume 1, Temperature.
- 650 Madec, G., and the NEMO team (2012). NEMO ocean engine, version3.4 (Institut Pierre-Simon Laplace Note du Pole de Modélisation).
- Meiners, K.M., Vancoppenolle, M., [Thanassekos, S.](#), [Dieckmann, G.S.](#), [Thomas, D.N.](#), [Tison, J.-L.](#), [Arrigo, K.R.](#), [Garrison, D.L.](#), [McMinn, A.](#), [Lannuzel, D.](#), et al. (2012). Chlorophyll a in Antarctic sea ice from historical ice core data. *Geophys. Res. Lett.* *39*, L21602.
- 655 [Meiners, K.M.](#), [Vancoppenolle, M.](#), Carnat, G., Castellani, G., Delille, B., Delille, D., Dieckmann, G.S., Flores, H., Fripiat, F., Grotti, M., et al. (2018). Chlorophyll-a in Antarctic Landfast Sea Ice: A First Synthesis of Historical Ice Core Data. *J. Geophys. Res. Oceans* *123*, 8444–8459.
- Meinshausen, M., Vogel, E., Nauels, A., Lorbacher, K., Meinshausen, N., Etheridge, D.M., Fraser, P.J., Montzka, S.A., Rayner, P.J., Trudinger, C.M., et al. (2017). Historical greenhouse gas concentrations for climate modelling (CMIP6). *Geosci. Model Dev.* *10*, 2057–2116.
- 660 Meinshausen, M., Nicholls, Z.R.J., Lewis, J., Gidden, M.J., Vogel, E., Freund, M., Beyerle, U., Gessner, C., Nauels, A., Bauer, N., et al. (2020). The shared socio-economic pathway (SSP) greenhouse gas concentrations and their extensions to 2500. *Geosci. Model Dev.* *13*, 3571–3605.
- Miller, L.A., Fripiat, F., Else, B.G.T., Bowman, J.S., Brown, K.A., Collins, R.E., Ewert, M., Fransson, A., Gosselin, M., Lannuzel, D., et al. (2015). Methods for biogeochemical studies of sea ice: The state of the art, caveats, and recommendations. *Elem. Sci. Anthr.* *3*.
- Moore, J.K., Lindsay, K., Doney, S.C., Long, M.C., and Misumi, K. (2013). Marine Ecosystem Dynamics and Biogeochemical Cycling in the Community Earth System Model [CESM1(BGC)]: Comparison of the 1990s with the 2090s under the RCP4.5 and RCP8.5 Scenarios. *J. Clim.* *26*, 9291–9312.
- 670 Mortenson, E., Hayashida, H., Steiner, N., Monahan, A., Blais, M., Gale, M.A., Galindo, V., Gosselin, M., Hu, X., Lavoie, D., et al. (2017). A model-based analysis of physical and biological controls on ice algal and pelagic primary production in Resolute Passage. *Elem Sci Anth* *5*.
- Mortenson, E., Steiner, N., Monahan, A.H., Hayashida, H., Sou, T., and Shao, A. (2020). Modeled Impacts of Sea Ice Exchange Processes on Arctic Ocean Carbon Uptake and Acidification (1980–2015). *J. Geophys. Res. Oceans* *125*, e2019JC015782.
- 675 Naughten, K.A., Meissner, K.J., Galton-Fenzi, B.K., England, M.H., Timmermann, R., and Hellmer, H.H. (2018). Future Projections of Antarctic Ice Shelf Melting Based on CMIP5 Scenarios. *J. Clim.* *31*, 5243–5261.
- O'Neill, B.C., Tebaldi, C., Vuuren, D.P. van, Eyring, V., Friedlingstein, P., Hurtt, G., Knutti, R., Kriegler, E., Lamarque, J.-F., Lowe, J., et al. (2016). The Scenario Model Intercomparison Project (ScenarioMIP) for CMIP6. *Geosci. Model Dev.* *9*, 3461–3482.
- 680 Orr, J.C., Najjar, R.G., Aumont, O., Bopp, L., Bullister, J.L., Danabasoglu, G., Doney, S.C., Dunne, J.P., Dutay, J.-C., Graven, H., et al. (2017). Biogeochemical protocols and diagnostics for the CMIP6 Ocean Model Intercomparison Project (OMIP). *Geosci. Model Dev.* *10*, 2169–2199.

- Pogson, L., Tremblay, B., Lavoie, D., Michel, C., and Vancoppenolle, M. (2011). Development and validation of a one-dimensional snow-ice algae model against observations in Resolute Passage, Canadian Arctic Archipelago. *J. Geophys. Res.* *116*, C04010.
- Saenko, O.A., Zhai, X., Merryfield, W.J., and Lee, W.G. (2012). The Combined Effect of Tidally and Eddy-Driven Diapycnal Mixing on the Large-Scale Ocean Circulation. *J. Phys. Oceanogr.* *42*, 526–538.
- Saenko, O.A., Yang, D., and Gregory, J.M. (2018). Impact of Mesoscale Eddy Transfer on Heat Uptake in an Eddy-Parameterizing Ocean Model. *J. Clim.* *31*, 8589–8606.
- Schulzweida, U. (2019). CDO User Guide.
- Smith, D.M., Screen, J.A., Deser, C., Cohen, J., Fyfe, J.C., García-Serrano, J., Jung, T., Kattsov, V., Matei, D., Msadek, R., et al. (2019). The Polar Amplification Model Intercomparison Project (PAMIP) contribution to CMIP6: investigating the causes and consequences of polar amplification. *Geosci. Model Dev.* *12*, 1139–1164.
- 695 Stewart, K.D., Kim, W.M., Urakawa, S., Hogg, A.McC., Yeager, S., Tsujino, H., Nakano, H., Kiss, A.E., and Danabasoglu, G. (2020). JRA55-do-based repeat year forcing datasets for driving ocean–sea-ice models. *Ocean Model.* *147*, 101557.
- Swart, N.C., Cole, J.N.S., Kharin, V.V., Lazare, M., Scinocca, J.F., Gillett, N.P., Anstey, J., Arora, V., Christian, J.R., Hanna, S., et al. (2019). The Canadian Earth System Model version 5 (CanESM5.0.3). *Geosci. Model Dev.* *12*, 4823–4873.
- Tedesco, L., Vichi, M., and Scoccimarro, E. (2019). Sea-ice algal phenology in a warmer Arctic. *Sci. Adv.* *5*, eaav4830.
- 700 Tsujino, H., Urakawa, S., Nakano, H., Small, R.J., Kim, W.M., Yeager, S.G., Danabasoglu, G., Suzuki, T., Bamber, J.L., Bentsen, M., et al. (2018). JRA-55 based surface dataset for driving ocean–sea-ice models (JRA55-do). *Ocean Model.* *130*, 79–139.
- Tsujino, H., Urakawa, L.S., Griffies, S.M., Danabasoglu, G., Adcroft, A.J., Amaral, A.E., Arsouze, T., Bentsen, M., Bernardello, R., Böning, C.W., et al. (2020). Evaluation of global ocean–sea-ice model simulations based on the experimental protocols of the Ocean Model Intercomparison Project phase 2 (OMIP-2). *Geosci. Model Dev.* *13*, 3643–3708.
- 705 Vancoppenolle, M., and Tedesco, L. (2016). Numerical models of sea ice biogeochemistry. In *Sea Ice*, (Wiley-Blackwell), pp. 492–515.
- Virtanen, P., Gommers, R., Oliphant, T.E., Haberland, M., Reddy, T., Cournapeau, D., Burovski, E., Peterson, P., Weckesser, W., Bright, J., et al. (2020). SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat. Methods* *17*, 261–272.
- 710 Watanabe, E., Onodera, J., Harada, N., Aita, M.N., Ishida, A., and Kishi, M.J. (2015). Wind-driven interannual variability of sea ice algal production in the western Arctic Chukchi Borderland. *Biogeosciences* *12*, 6147–6168.
- Watanabe, E., Jin, M., Hayashida, H., Zhang, J., and Steiner, N. (2019). Multi-Model Intercomparison of the Pan-Arctic Ice-Algal Productivity on Seasonal, Interannual, and Decadal Timescales. *J. Geophys. Res. Oceans* *124*, 9053–9084.
- 715 Welch, P. (1967). The use of fast Fourier transform for the estimation of power spectra: A method based on time averaging over short, modified periodograms. *IEEE Trans. Audio Electroacoustics* *15*, 70–73.

Wilkinson, M.D., Dumontier, M., Aalbersberg, I.J., Appleton, G., Axton, M., Baak, A., Blomberg, N., Boiten, J.-W., da Silva Santos, L.B., Bourne, P.E., et al. (2016). The FAIR Guiding Principles for scientific data management and stewardship. *Sci. Data* 3, 1–9.

720 Woodgate, R.A., Aagaard, K., and Weingartner, T.J. (2005). Monthly temperature, salinity, and transport variability of the Bering Strait through flow. *Geophys. Res. Lett.* 32.

Zeebe, R.E., Eicken, H., Robinson, D.H., Wolf-Gladrow, D., and Dieckmann, G.S. (1996). Modeling the heating and melting of sea ice through light absorption by microalgae. *J. Geophys. Res. Oceans* 101, 1163–1181.

725 Zhang, X., Church, J.A., Monselesan, D., and McInnes, K.L. (2017). Sea level projections for the Australian region in the 21st century. *Geophys. Res. Lett.* 44, 8481–8491.

Ziehn, T., Chamberlain, M.A., Law, R.M., Lenton, A., Bodman, R.W., Dix, M., Stevens, L., Wang, Y.-P., and Srinovsky, J. (2020). The Australian Earth System Model: ACCESS-ESM1.5. *J. South. Hemisphere Earth Syst. Sci.*

Zweng, M.M., Reagan, J.R., Antonov, J.I., Locarnini, R.A., Mishonov, A.V., Boyer, T.P., Garcia, H.E., Baranova, O.K., Johnson, D.R., and Seidov, D. (2013). *World ocean atlas 2013. Volume 2, Salinity.*

730




Figure 1: Global (left) and regional (right) configurations of the three-dimensional coupled sea ice–ocean physical–biogeochemical models participating in IAMIP2. The surface-atmospheric forcing dataset consists of air temperature at 10 m (tas), specific humidity at

735 10 m (huss), eastward wind at 10 m (uas), northward wind at 10 m (vas), sea level pressure (psl), downward shortwave radiation (rsds), downward longwave radiation (rlds), rainfall flux (prra), snowfall flux (prsn), river runoff (friver), and calving flux (lcalvlf).





Figure 2: Timeline of the numerical experiments of IAMIP2. Years begin on January 1. Each experiment starts on January 1 and ends on December 31. For example, the historical experiment starts from January 1, 1958, and ends on December 31, 2018.





Figure 3: Comparisons of historical and projected global and polar warming simulated by 26 CMIP6 models over the historical period (1958-2014) and the projection period under the SSP5-8.5 scenario (2015-2100). Time series of annual mean surface air temperature averaged over the ocean grid cells of (a) the globe, (c) the Northern Hemisphere north of 60°N, and (e) the Southern Hemisphere south of 60°S. Projected changes in annual mean surface air temperature averaged over the ocean grid cells of (b) the globe,



Deleted:

Deleted: 24

750

(d) the Northern Hemisphere north of 60 °N, and (f) the Southern Hemisphere south of 60 °S, relative to their averages during 1958-2014. Blue, purple, green, and red denote 4 of the 26 models that provide the atmospheric output at the temporal resolutions needed for the projection experiments of IAMIP2. Gray denotes the remaining 22 models. Black denotes the multi-model mean of the 26 models. Orange denotes the JRA55-do dataset used for the atmospheric forcing for the historical experiment of IAMIP2.



Deleted: 3

Deleted: 24

Deleted: experiment

Deleted: 19

Deleted: 24



Deleted:

Deleted: 24

760 **Figure 4: Comparisons of major climate modes simulated by 26 CMIP6 models over the historical period (1958-2014) and the projection period under the SSP5-8.5 scenario (2015-2100).** Time series of annual mean sea level pressure difference between two regions representative of (a) EQSOI, (c) NAM, and (e) SAM. Normalised power spectra of (b) EQSOI, (d) NAM, and (f) SAM derived from time series of standardised monthly mean anomaly of sea level pressure difference over 1958-2014 based on Welch's method

(Welch, 1967). Blue, purple, green, and red denote 4 of the 24 models that provide the atmospheric output at the temporal resolutions needed for the projection experiment of IAMIP2. Gray denotes the remaining 22 models. Black denotes the multi-model mean of the 26 models. Orange denotes the JRA55-do dataset used for the atmospheric forcing for the historical experiment of IAMIP2.

Deleted: 3
 Deleted: 19
 Deleted: 24
 Deleted:



Figure 5: An example file tree diagram for the IAMIP2 output. Blue denotes the directories and white denotes files. A grid file (e.g., ACCESS-OM2_grid.nc) must be provided and placed under the model directory (e.g., IAMIP2/ACCESS-OM2/). Each diagnostic file (e.g., siconc_day_IAMIP2_ACCESS-OM2_ssp585_EC-Earth3_2015.nc) contains the output of a variable for a year.

770

775 Table 1: List of participating models for IAMIP2. For ocean and sea-ice ecosystem components, the letters denote nutrients (N), phytoplankton (P), zooplankton (Z), and detritus (D), which are followed by the numbers indicating the complexity. For example, N1 denotes one type of modelled nutrients (e.g., nitrate), whereas N2 indicates two types of modelled nutrients (e.g., nitrate and iron). The vertical resolution is rounded to the nearest integer.

| Model | ACCESS-OM2 | CESM-IARC | RASM | CanNEMO | NEMO-NAA | COCO-Arctic NEMURO |
|------------------------|---------------------------|-----------|------------|----------|------------|--------------------|
| Relevant CMIP6 model | ACCESS-CM2, ACCESS-ESM1.5 | CESM2 | CESM2 | CanESM5 | CanESM5 | MIROC6 |
| Ocean dynamics | MOM5.1 | POP2 | POP2 | OPA | OPA | COCO4.9 |
| Sea-ice dynamics | CICE5.1.2 | CICE5.1.2 | CICE5.1.2 | LIM2 | LIM2 | COCO4.9 |
| Ocean ecosystem | N1P1Z1D1 | N4P3Z1D1 | N4P3Z1D1 | N3P2Z2D2 | N3P2Z2D2 | N3P2Z3D2 |
| Sea-ice ecosystem | N1P1 | N4P1 | N4P1 | N2P1 | N2P1 | N3P1D1 |
| Spatial domain | Global | Global | Pan-Arctic | Global | Pan-Arctic | Pan-Arctic |
| Horizontal resolution | 1° | 1° | 1/12° | 1° | 1/4° | 1/4° |
| Vertical resolution of | 2 m | 10 m | 5 m | 6 m | 1 m | 2 m |

| | | | | | | |
|--------------------------------------|---|--------------------|--------------------|--|--|-------------------------|
| the surface ocean layer | | | | | | |
| Bottom-ice ecosystem layer thickness | 0.03 m | 0.03 m | 0.03 m | 0.03 m | 0.03 m | 0.02 m |
| Reference | (Jeffery et al., 2016; Kiss et al., 2020; Ziehn et al., 2020) | (Jin et al., 2018) | (Jin et al., 2018) | (Hayashida et al., 2019; Swart et al., 2019) | (Hayashida et al., 2019; Hu and Myers, 2013) | (Watanabe et al., 2015) |

785 **Table 2: List of model diagnostics for IAMIP2.** Where applicable, variable names and units follow the CMIP6 convention (<https://earthsystemcog.org/projects/wip/CMIP6DataRequest>). Tiers refer to mandatory (Tier 1) and optional (Tier 2).

| Variable name | Long name | Units | Tier |
|---------------|---|-------------------------------------|------|
| siconc | Percentage of grid cell covered by sea ice | % | 1 |
| sithick | Actual sea ice thickness (sea-ice volume divided by sea-ice-covered area) | m | 1 |
| sisnthick | Actual snow thickness (snow volume divided by snow-covered area) | m | 1 |
| intppbi | Vertically integrated primary organic carbon production by bottom ice algae | mol m ⁻² s ⁻¹ | 1 |
| intpp | Vertically integrated primary organic carbon production by phytoplankton | mol m ⁻² s ⁻¹ | 1 |
| epc100 | Downward flux of particulate organic carbon at 100 m | mol m ⁻² s ⁻¹ | 2 |
| phycbi | Bottom ice algae carbon concentration | mol m ⁻³ | 1 |
| phycos | Sea surface phytoplankton carbon concentration | mol m ⁻³ | 1 |
| no3bi | Bottom ice dissolved nitrate concentration | mol m ⁻³ | 1 |
| no3os | Sea surface dissolved nitrate concentration | mol m ⁻³ | 1 |
| parbi | Downwelling photosynthetic radiance flux at bottom ice | W m ⁻² | 2 |
| paros | Downwelling photosynthetic radiance flux at sea surface | W m ⁻² | 2 |
| sst | Sea surface temperature | °C | 1 |
| sss | Sea surface salinity | PSU | 1 |
| mlots2t | Ocean mixed layer thickness defined by sigma-t | m | 2 |
| dissicos | Sea surface dissolved inorganic carbon concentration | mol m ⁻³ | 2 |
| talkos | Sea surface total alkalinity | mol m ⁻³ | 2 |
| fgco2 | Sea surface downward flux of carbon dioxide | kg m ⁻² s ⁻¹ | 2 |