



Anoxic iron and sulphur cycling in the cGENIE.muffin Earth system model (v0.9.16)

Sebastiaan J. van de Velde^{1,*}, Dominik Hülse¹, Christopher T. Reinhard², and Andy Ridgwell¹

¹Department of Earth and Planetary Sciences, University of California, Riverside, CA 92521, USA

²School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA 30332, USA

*Current address: Bgeosys, Geoscience, Environment & Society, Université Libre de Bruxelles, Brussels, Belgium;
Operational Directorate Natural Environment, Royal Belgian Institute of Natural Sciences, Brussels, Belgium

Correspondence: sebastiaan.van.de.velde@ulb.ac.be

Abstract. The coupled biogeochemical cycles of iron and sulphur are central to the long-term biogeochemical evolution of Earth's oceans. For instance, before the development of a persistently oxygenated deep ocean, the ocean interior likely alternated between states buffered by reduced sulphur ('euxinic') vs. buffered by reduced iron ('ferruginous'), with important implications for the cycles and hence bioavailability of dissolved iron (and phosphate). Even after atmospheric oxygen concentrations rose to modern-like values, the ocean continued, episodically, to develop regions of euxinic or ferruginous conditions, such as associated with past key intervals of organic carbon deposition (e.g. during the Cretaceous) as well as extinction events (e.g. at the Permian/Triassic boundary). A better understanding of the cycling of iron and sulphur in an anoxic ocean, how geochemical patterns in the ocean relate to the available spatially heterogeneous geological observations, and quantification of the feedback strengths between nutrient cycling, biological productivity, and ocean redox, requires a spatially-resolved representation of ocean circulation together with an extended set of (bio)geochemical reactions.

Here, we extend the 'muffin' release of the intermediate-complexity Earth system model cGENIE, to now include an anoxic iron and sulphur cycle, enabling the model to simulate ferruginous and euxinic redox states as well as the precipitation of reduced iron and sulphur minerals (pyrite, siderite, greenalite) and attendant iron and sulphur isotope signatures, which we describe in full. While we cannot make direct model comparison with present-day (oxic) ocean observations, we use an idealized ocean configuration to explore model sensitivity across a selection of key parameters. We also present the spatial patterns of concentrations and $\delta^{56}\text{Fe}$ isotope signatures of both dissolved and solid-phase Fe species in an anoxic ocean as an example application. Our sensitivity analyses show how the first-order results of the model are relatively robust against the choice default kinetic parameters within the Fe-S system, and that simulated concentrations and reaction rates are comparable to those observed in process analogues for ancient oceans (i.e., anoxic lakes). Future model developments will address sedimentary recycling and benthic iron fluxes back to the water column, together with the coupling of nutrient (in particular phosphate) cycling to the iron cycle.



1 Introduction

The biogeochemical cycles of iron and sulphur are tightly coupled in the marine environment and play fundamental roles in the evolution and functioning of the Earth System (Raiswell and Canfield, 2012). In their main oxidised states, both sulphur (in the form of sulphate; SO_4^{2-}) and iron (in the form of iron (oxyhydr)oxides; $FeOOH$) are important electron acceptors in the oxidation of organic matter in anoxic environments such as marine sediments or oxygen-deficient water columns (e.g., Black Sea, stratified lakes) (Thamdrup, 2000; Crowe et al., 2008; Raiswell and Canfield, 2012). In metabolising organic matter, microbial reduction of SO_4^{2-} and $FeOOH$ produce reduced sulphide (H_2S) and ferrous iron (Fe^{2+}), respectively. When present at the same location, H_2S and Fe^{2+} combine into iron monosulphides and eventually pyrite (FeS_2) (Rickard, 1997, 2006), the burial of which couples the short-term, surface, cycles of Fe and S with their long-term, geological, cycles (Berner, 1989). Depending on the relative ocean inventory of H_2S vs. Fe^{2+} , the precipitation of FeS_2 can lead to an anoxic water body becoming either iron-rich ('ferruginous') or sulphide-rich ('euxinic') (Canfield, 1998; Poulton and Canfield, 2011) – $H_2S:Fe^{2+}$ ratios greater than 2:1 (the $S:Fe$ ratio in FeS_2) promoting euxinic conditions, and lower ratios promoting ferruginous conditions (Poulton and Canfield, 2011). For most of Earth's history, the ocean interior is thought to have been predominantly anoxic (Fig. 1; Lyons et al., 2014), which implies that reduced forms of iron and sulphur would have dominated the marine redox landscape (Poulton and Canfield, 2011; Raiswell and Canfield, 2012).

Whether an anoxic water body becomes ferruginous or euxinic can have significant impacts on the availability of nutrients, with ferruginous conditions potentially leading to phosphate limitation, and euxinic conditions potentially leading to depletion of key biological trace elements (Van Cappellen and Ingall, 1996; Bjerrum and Canfield, 2002; Reinhard et al., 2013, 2017; Wallmann et al., 2019). Furthermore, before the advent of oxygenic photosynthesis, the productivity of marine ecosystems was likely, at least partly, fuelled by the oxidation of H_2S or Fe^{2+} to their oxidised counterparts (Kharecha et al., 2005; Canfield et al., 2006; Ozaki et al., 2018; Thompson et al., 2019). As a result, the long-term evolution of the Earth system and the structure of marine ecosystems are closely tied to the evolution of the biogeochemical cycles of iron and sulphur. The ability to simulate in models the evolution of these cycles as well as geochemical distributions in the ocean hence becomes key to better understanding the early evolution of microbial ecosystems.

During the early stages of Earth's history (i.e., Archean to mid-Proterozoic), the ocean was rich in Fe^{2+} (Fig. 1), which enabled the extensive deposition of banded iron formations (BIFs) and ferruginous shales during those Eons (Bekker et al., 2010; Planavsky et al., 2011; Sperling et al., 2015; Konhauser et al., 2017). BIFs are rare in sediments deposited after 1.8 billion years ago, which was initially hypothesised to reflect a transition to oxygen-rich bottom waters that prevented the build-up of soluble iron by removing it as insoluble iron oxides (Cloud, 1972; Holland, 1984). In a seminal paper, Canfield (1998) suggested that the abundance of atmospheric oxygen during the Proterozoic (which was at most $\sim 10\%$ of today; Canfield and Teske, 1996; Lyons et al., 2014) was too low to oxygenate the deeper waters of the ocean. Instead, he proposed that the disappearance of BIFs was driven by an increase of oceanic sulphate concentrations, allowing sulphide (produced by microbial sulphate reduction) to remove reduced iron from solution by the precipitation of FeS_2 (Canfield, 1998). Hence, this hypothesis implied that the deep ocean was euxinic for most of the Proterozoic. Since then, a large wealth of geochemical proxy data has




Figure 1. First-order evolution of Earth's ocean redox landscape. Insets show conceptual models of the spatial redox structure of the ocean at certain points in Earth's history. Based on (Poulton et al., 2010; Poulton and Canfield, 2011; Raiswell and Canfield, 2012). After (van de Velde et al., in review). See text for details.

been collected, aided by the development of a sequential Fe extraction scheme that helps to differentiate between oxic, euxinic and ferruginous conditions by determining seven operationally defined iron mineral pools (Poulton et al., 2005; Poulton and Canfield, 2011). These paleo-reconstructions of Archean and Proterozoic deposits helped to further refine the spatial and temporal history of ocean redox. In contrast to the earlier view, the Proterozoic Eon appears to have been dominated by ferruginous conditions (Canfield et al., 2008; Planavsky et al., 2011; Poulton and Canfield, 2011; Guilbaud et al., 2015), likely interspersed with euxinic excursions along the shelf (Canfield et al., 2008; Poulton et al., 2010; Raiswell and Canfield, 2012) (Fig. 1). However, to date these redox landscapes have been largely qualitative, since field-based observations are restricted by the limited number of available unaltered deposits of a certain point in time. Furthermore, sample acquisition for a depth transect is an enormously labour-intensive task (see e.g. Poulton et al., 2010). Because of these limitations, both the spatial redox pattern and the exact nature of ferruginous versus euxinic conditions (i.e. the concentrations of Fe^{2+} or H_2S) are still relatively unconstrained. The uncertainty with respect to the ocean redox conditions propagates into hypotheses on nutrient availability and ecosystem evolution (see, e.g.; Reinhard et al., 2017).

Whether the ocean interior is ferruginous or euxinic can significantly impact the supply of essential nutrients to surface ocean environments, and thus nutrient availability for primary producers (Van Cappellen and Ingall, 1996; Bjerrum and Canfield,



2002; Guilbaud et al., 2020). For example, iron oxides ($FeOOH$) can efficiently scavenge phosphate (PO_4^{3-}) from a water column, which has been suggested to limit oceanic phosphate availability in the past (Berner, 1973; Van Cappellen and Ingall, 1996; Bjerrum and Canfield, 2002; Jones et al., 2015). In contrast, a euxinic water column could potentially induce trace metal limitations by titrating out essential trace elements like iron or molybdenum (Reinhard et al., 2013; Wallmann et al., 2019).

5 Indeed, lower nutrient availability - specifically phosphate - has been suggested to limit primary productivity for the majority of the Proterozoic (Bjerrum and Canfield, 2002; Reinhard et al., 2017; Ozaki et al., 2019). At the same time, nutrient availability is thought to have been critical in shaping and driving eukaryote evolution and proliferation (Brocks et al., 2017; Reinhard et al., 2020b). However, many of the most significant innovations in the history of Earth's biosphere have likely occurred in specific sites of the ocean that did not reflect the average state of the ocean as a whole (Nisbet and Sleep, 2001). Hence, the

10 ability to reconstruct iron and sulphur cycling in a spatially explicit way is critical for exploring the relationships between biospheric evolution and changes in ocean redox.

None of the currently available suite of global models that explicitly represent biogeochemical cycling under low-oxygen marine environmental conditions include an extensive treatment of biogeochemical iron cycling (Ozaki et al., 2011; Laakso and Schrag, 2014; Hülse et al., 2017; Lenton et al., 2018; Reinhard et al., 2020a). Consequently, many essential interactions

15 between the iron cycle and other elemental cycles (e.g. sulphur burial via pyrite precipitation) are abstracted and parameterised using techniques that may or may not be mechanistically robust across a range of scenarios. Moreover, most of the ocean models used to simulate the early stages of Earth's history are essentially box- or one-dimensional ocean models (Ozaki et al., 2011; Laakso and Schrag, 2014; Lenton et al., 2018) and are unable to resolve the spatial patterns necessary to start contrasting with the geological record (although some two- or three-box models attempt to resolve between the coastal zone and the open

20 ocean; Laakso and Schrag, 2019; Thompson et al., 2019; Alcott et al., 2019). Indeed, previous simulations of past oceans with three-dimensional ocean models have already indicated the importance of spatial patterns in ocean redox (e.g. Olson et al., 2013), specifically when considering habitability for complex eukaryotic life (Reinhard et al., 2016, 2020b).

In this paper, we present the development of a coupled anoxic oceanic iron and sulphur cycle, embedded within the 'muffin' release of the carbon-centric Grid ENabled Integrated Earth system model, 'cGENIE'. The aim is to extend the functionality

25 of cGENIE into regimes in which the ocean interior is pervasively anoxic, including much of Earth's Precambrian history and periods of significant perturbation to ocean redox during the Paleozoic and Mesozoic (during so-called 'Ocean Anoxic Events'; OAEs), where coupled iron-sulphur cycling dominated biogeochemical interactions in the ocean interior. Our extension explicitly models the formation, burial, and isotopic compositions of key mineral phases used in paleoenvironmental reconstructions — specifically iron oxide ($FeOOH$), siderite ($FeCO_3$), greenalite ($Fe_3Si_2O_5(OH)_4$) and pyrite (FeS_2) (Poulton et al., 2005; Poulton and Canfield, 2011) — which allows quantitative comparison with available data (e.g., Rouxel et al., 2005; Heard and Dauphas, 2020). In the next section (Section 2), we briefly discuss the cGENIE model framework, focusing on the features that are most relevant for our purpose of modeling pervasively anoxic oceans. Section 3 describes the included iron-sulphur reactions and the reasoning behind the chosen parameterisations. In Section 4 we discuss an example configuration and a series of sensitivity experiments to the chosen parameterisation. Finally, in Section 5 we discuss model limitations and

35 potential future developments.



2 The cGENIE.muffin Earth system model framework

cGENIE is an earth system model of intermediate complexity (EMIC) which comprises a modular framework that incorporates different components of the Earth system, including ocean circulation and biogeochemical cycling, ocean-atmosphere and ocean-sediment exchange, and the long-term (geological) cycle carbon and various solid-Earth derived tracers (Ridgwell et al., 2007; Ridgwell and Hargreaves, 2007; Colbourn et al., 2013; Adloff et al., 2020). Here, we use the current 'muffin' release that encompasses a range of developments and/or additions in the representation of: temperature-dependent metabolic processes in the ocean (Crichton et al., 2020), ocean-atmosphere cycling of methane (Reinhard et al., 2020a), marine ecosystems (Ward et al., 2018), organic matter preservation and burial in marine sediments (Hülse et al., 2018), and geological cycles of weathering-relevant trace-metals and isotopes (Adloff et al., 2020).

The climate component in cGENIE - C-GOLDSTEIN - has a reduced physics (frictional geostrophic) 2-D energy-moisture balance model of the atmosphere coupled to a 3-D ocean circulation model and a dynamic-thermodynamic sea-ice model (see Edwards and Marsh, 2005; Marsh et al., 2011, for full descriptions). In addition to the simplified atmosphere, to further facilitate the simulation of a relatively large number of interacting gaseous, dissolved, and solid tracers across atmosphere, ocean, and marine sediment (and land surface), cGENIE employs a much reduced spatial and temporal resolution relative to most high-resolution ocean general circulation models. While this precludes exploration of very detailed spatial patterns, it does provide a flexibility which is not available in more high-resolution models, and the relatively short run time of cGENIE (around 1 day per 10,000 model years on a single CPU core) allows us to run many different model experiments and carry out comprehensive parameter sweeps and sensitivity analysis (and hence parameter tuning and model calibration). This aligns with our ultimate aim here which is to explore ocean biogeochemistry during periods of the Mesozoic (>65 million years ago), Paleozoic (>250 million years ago), and Precambrian (>540 million years ago). Most of these changes likely occurred on timescales exceeding 10,000 years, and at present we have virtually no information with respect to seasonality or detailed spatial variability for many of these intervals (see, e.g., Poulton et al., 2010; Guilbaud et al., 2015). In addition, key boundary conditions and parameter values for these periods of Earth's history are often poorly constrained, necessitating large model ensembles in order to adequately assess the robustness of any given result.

2.1 Continental configuration and climatology

For the purpose of this study – implementing and characterising the coupled cycling of iron and sulphur in an anoxic ocean – we adopt a deliberately idealised model configuration. We configure the ocean model on a 18x18 equal-area horizontal grid with 16 logarithmically spaced z-coordinate levels (Fig. 2). The horizontal grid is uniform in longitude (20° resolution) and uniform in the sine of latitude (~ 3.2° at the equator to 19.2° near the poles) (Fig. 2a). The layer thickness in the vertical grid increases from 80.8 m at the surface to 765 m at the deepest layer (Fig. 2b). We adopt a 'Ridge World' set-up, with a thin strip of land connecting North and South poles (Fig. 2c), following Ferreira et al. (2010), which creates a single ocean basin with no circumpolar current (a little akin to the plate configuration prevailing during the late Permian). We apply idealised boundary




Figure 2. Schematic depiction of the cGENIE grid used for our simulations. The grid is 18x18x16, uniform in longitude and uniform in the sine of latitude. The layer thickness in the vertical grid increases from 80.8 m at the surface to 765 m at the deepest layer. (a) Top view of the cGENIE grid with indication of the layer numbers. (b) Side-view of the cGENIE grid with indication of the layer numbers. (c) Schematic picture of the continental configuration. 'Ridge World' has 1 continent which runs from pole to pole and extends from 0 to 20 °E (i.e., the width of a longitudinal grid-cell). The ocean is 5000 m everywhere.

conditions of zonally-averaged wind stress and speed, plus a zonally-averaged planetary albedo, all following Vervoort et al. (in review). The solar constant is set to modern (1368.0 W m^{-2}).

The physics parameters controlling the model climatology all follow the 16-level ocean based configuration assessed by Cao et al. (2009) (Table S1 in their manuscript), with the exception of the parameterisation controlling ocean mixing. When using the standard implementation of isoneutral diffusion and eddy-induced advection (Edwards and Marsh, 2005; Marsh et al., 2011), sharp vertical redox gradients simulated under extreme redox and high dissolved iron conditions resulted in unacceptably negative tracer concentrations at depth, particularly for dissolved iron. We hence disabled this parameterisation, reverting to the original un-adjusted horizontal+vertical diffusion physics configuration of Edwards and Shepherd (2002). We tested the modern cGENIE configuration of Cao et al. (2009) for both ocean mixing parameterisations, and found only a minimal difference in the large scale ocean circulation (i.e., a $\sim 2 \text{ Sv}$ stronger Atlantic meridional overturning circulation in the un-adjusted parameterisation).



2.2 The biological carbon pump

In this paper, we adopt a representation of biological export from the surface ocean driven by an implicit (i.e. unresolved) biological community with a highly parameterised uptake of nutrients in the photic zone. The description of the basic scheme can be found in Ridgwell et al. (2007), although we use the specific configuration (including iron co-limitation) as characterised by Tagliabue et al. (2016). The governing equations are summarised below.

Biological productivity in the euphotic zone (taken to be the surface layer in the ocean model) is controlled by dissolved phosphate (PO_4^{3-}) and dissolved iron (Fe^{3+}) availability, the fractional ice coverage of each grid cell (A), mean ambient light and temperature. With the exception of the parameterisation of the temperature term, the equation for photosynthetic nutrient uptake follows Doney et al. (2006):

$$\Lambda = F_I \cdot F_T \cdot \min(F_{PO_4^{3-}}, F_{Fe^{3+}}) \cdot (1 - A) \cdot \frac{\min([PO_4^{3-}], red_{P/Fe} \cdot [Fe^{3+}])}{\tau_{bio}} \quad (1)$$

Rates of photosynthetic nutrient uptake are scaled to ambient dissolved nutrient concentrations (PO_4^{3-} and Fe^{3+}), according to an optimal uptake timescale ($\tau_{bio}=63.4$ days; Meyer et al., 2016), and converting $[Fe^{3+}]$ to the equivalent dissolved phosphate concentration via the $Fe : P$ Redfield ratio ($red_{P/Fe}$). The various limitation terms are:

$$F_I = \frac{I}{I + \kappa_I} \quad (2)$$

15

$$F_{PO_4^{3-}} = \frac{[PO_4^{3-}]}{[PO_4^{3-}] + \kappa_{PO_4^{3-}}} \quad (3)$$

$$F_{Fe^{3+}} = \frac{[Fe^{3+}]}{[Fe^{3+}] + \kappa_{Fe^{3+}}} \quad (4)$$

where shortwave irradiance I is averaged over the entire mixed layer, and is assumed to decay exponentially from the sea surface with a length scale of 20 m, as per Doney et al. (2006). The κ terms in each equation represent half-saturation constants for each limiting component ($\kappa_I=40 \text{ W m}^{-2}$, $\kappa_{PO_4^{3-}}=0.1 \text{ } \mu\text{M}$, $\kappa_{Fe^{3+}}=0.1 \text{ nM}$) and are as used in Tagliabue et al. (2016). The influence of temperature on biological export is parameterised as:

$$F_T = k_{T0} \cdot \exp\left(\frac{T}{k_{eT}}\right) \quad (5)$$

where k_{T0} (0.59) is a scaling constant, k_{eT} (15.8°C) the e -folding temperature, and T is the *in-situ* temperature (°C). The scaling constants give rise to an approximately factor two change per temperature change of 10°C (Reinhard et al., 2020a).

A proportion ($\nu = 0.66$) of PO_4^{3-} taken up by biota is partitioned in dissolved organic phosphorus (DOP) while the remainder – as particulate organic phosphorus (POP) – is exported vertically out of the surface ocean. Because the biological configuration used here does not resolve explicit standing plankton biomass, the export flux of POP is always equal to the rate



of PO_4^{3-} uptake:

$$F_{z=h_e}^{POP} = \int_{h_e}^0 \rho(1 - \nu) \Lambda dz \quad (6)$$

where ρ is the density of seawater and h_e the thickness of the euphotic zone (=80.84 m). The particulate organic carbon (POC) export flux is calculated using a fixed Redfield ratio as:

$$5 \quad F_{z=h_e}^{POC} = 106 F_{z=h_e}^{POP} \quad (7)$$

After export from the surface grid cells, POC is remineralised instantaneously throughout the water column following a Martin -type curve (Martin et al., 1987), with a specified decay constant b (=0.7) (Fig. 3a):

$$F_z^{POC} = F_{z=h_e} (z/h_e)^{-b} \quad (8)$$

The b -value of 0.7 is slightly higher than the global average of 0.6 estimated based on modern observations by Henson et al. (2011), but leads to a better reconstruction of the distribution of $CaCO_3$ in deep-sea sediments (Wilson, pers. comm.).

3 Oceanic iron and sulphur cycling

The motivation behind this paper to provide a tool that can aid understanding of the key interactions between the biological carbon pump, the oxygenation of the atmosphere and ocean, and the marine biogeochemical cycles of iron (Fe) and sulphur (S). In taking the first step towards this end, we construct a parsimonious model of ocean biogeochemical cycling based on a simplified speciation scheme for both Fe and S, and only consider a relatively limited number of potential redox states. In this section we briefly describe the general conceptual cycle of Fe and S, as illustrated in Fig. 3b, and in the following subsections discuss the assumptions, reaction equations, and parameters for each cycle. The model equations that describe the biogeochemical reactions (summarised in Table 1) are given in Table 2 and the kinetic constants, their units and default values can be found in Table 3.

Redox cycling in the ocean (and sediments) is driven by the mineralisation of POC, which is produced in the photic zone by photosynthesis, and subsequently sinks through the water column (Ridgwell et al., 2007). The sinking flux of POC follows a 'Martin'-type decay (Fig. 3a), which determines the mineralisation rate (R_{min}) at each depth layer in the water column, and thus R_{min} at depth z is calculated as,

$$R_{min}(z) = F_{POC}(z-1) - F_{POC}(z) \quad (9)$$

Note that mineralisation only occurs below the surface layers; $z > 1$ in Fig. 2. The mineralisation of POC is coupled to the reduction of a terminal electron acceptor (TEA). These TEAs are used according to decreasing energy yield (Froelich et al., 1979), and relative consumption rates are scaled with TEA concentration and the local abundance of inhibitory substances (i.e. a more energy-yielding TEA). Our mineralisation scheme includes aerobic respiration (AR, R1 in Table 1), dissimilatory




Figure 3. (a) Example of the 'Martin' curve for vertical organic matter fluxes in the ocean. (b) Conceptual description of the iron-sulphur cycle implemented in the model. For simplicity, only FeOOH (oxidised iron) and Fe²⁺ (reduced iron) are depicted, but the model includes different iron species in both the oxic and anoxic states. Interaction between the methane cycle and sulphur cycle is omitted for clarity. See text for details.

iron reduction (DIR, R2 in Table 1), dissimilatory sulphate reduction (DSR, R3 in Table 1), and methanogenesis (MG, R4 in Table 1). Of these, AR and DSR existed in the original cGENIE biogeochemical framework (Ridgwell et al., 2007), while methanogenesis has been more recently added (Reinhard et al., 2020a). In the current paper, we ignore nitrate reduction (Monteiro et al., 2012).

- 5 The rate of TEA consumption is represented by a Michaelis-Menten type relationship with respect to TEA concentration that allows for a non-linear closure of the system:

$$\begin{aligned}
 f_{AR} &= \frac{[O_2]}{K_{0,O_2} + [O_2]} \\
 f_{DIR} &= \frac{[FeOOH]}{K_{0,FeOOH} + [FeOOH]} \frac{K_{i,O_2}}{K_{i,O_2} + [O_2]} \\
 f_{DSR} &= \frac{[SO_4^{2-}]}{K_{0,SO_4^{2-}} + [SO_4^{2-}]} \frac{K_{i,FeOOH}}{K_{i,FeOOH} + [FeOOH]} \frac{K_{i,O_2}}{K_{i,O_2} + [O_2]} \\
 f_{MG} &= \frac{K_{i,SO_4^{2-}}}{K_{i,SO_4^{2-}} + [SO_4^{2-}]} \frac{K_{i,FeOOH}}{K_{i,FeOOH} + [FeOOH]} \frac{K_{i,O_2}}{K_{i,O_2} + [O_2]}
 \end{aligned} \tag{10}$$

where K_0 are the half-saturation constants for the four primary redox reactions and K_i are the inhibition constants that act on the less energetic redox reaction (Table 3).

- 10 The consequence of this scheme is that in the oxic zone of the water column, aerobic respiration is responsible for nearly all of the POC mineralisation. When oxygen starts to become depleted, DIR, and subsequently DSR, become the dominant mineralisation pathways. When both iron oxides and sulphate are exhausted, MG represents the final mineralisation pathway. While the occurrence and parametrisation of DSR and MG in the water column are relatively straightforward, the possibility of DIR in the water column is more complex and is discussed in more detail in section 3.1.2. It should be noted that a ('Martin'-



Table 1. List of biogeochemical reactions included in reduced Fe-S scheme within the cGENIE model. A distinction is made between mineralisation ("Primary redox reactions"), re-oxidation of reduced products ("Secondary redox reactions") and precipitation reactions. The associated kinetic expressions are listed in table 2. Note that we do not include here any details for reactions involving the methane cycle, though they are included in the simulations described here. For more information about the parametrisation and reactions of the methane cycle in cGENIE, we refer the reader to Reinhard et al. (2020a).

Reaction number	Reaction name	Abbreviation	Equation
<i>Primary redox reactions</i>			
R1	Aerobic respiration	AR	$CH_2O + O_2 \rightarrow CO_2 + H_2O$
R2	Dissimilatory iron reduction ¹	DIR	$CH_2O + 4FeOOH + 7H^+ \rightarrow HCO_3^- + 4Fe^{2+} + 6H_2O$
R3	Dissimilatory sulphate reduction	DSR	$CH_2O + \frac{1}{2}SO_4^{2-} + H^+ \rightarrow CO_2 + \frac{1}{2}\Sigma H_2S + H_2O$
R4	Methanogenesis	MG	$CH_2O \rightarrow \frac{1}{2}CO_2 + \frac{1}{2}CH_4$
<i>Secondary redox reactions</i>			
R5	Ferrous iron oxidation	FIO	$Fe^{2+} + \frac{1}{4}O_2 + H^+ \rightarrow Fe^{3+} + \frac{1}{2}H_2O$
R6	Canonical sulphide oxidation	CSO	$\Sigma H_2S + 2O_2 \rightarrow SO_4^{2-} + 2H^+$
R7	Sulphide-mediated iron reduction ¹	SMI _{d/s}	$\Sigma H_2S + 8Fe^{3+}/FeOOH + 4H_2O \rightarrow SO_4^{2-} + 8Fe^{2+} + 10H^+$
<i>Precipitation reactions</i>			
R8	Iron oxide precipitation	IrP	$Fe^{3+} + 2H_2O \rightarrow FeOOH + 3H^+$
R9	Pyrite precipitation ²	PyP	$FeS_p + \frac{3}{4}\Sigma H_2S + \frac{1}{4}SO_4^{2-} + \frac{1}{2}H^+ \rightarrow FeS_2 + H_2O$
R10	Siderite precipitation	SiP	$Fe^{2+} + CO_3^{2-} \rightarrow FeCO_3$
R11	Greenalite precipitation	GrP	$3Fe^{2+} + 2SiO_2(aq) + 5H_2O \leftrightarrow Fe_3Si_2O_5(OH)_4 + 6H^+$

¹ DIR can only occur with solid phase $FeOOH$, whereas SMI can occur with dissolved Fe^{3+} and solid phase $FeOOH$. See text for more details. ² The calculation of FeS_p occurs implicitly from the concentrations of Fe^{2+} and ΣH_2S and the solubility of FeS_{aq} , see section 3.1.3 for more details.

type) decay curve of organic matter flux with depth is prescribed in the model, such that regardless of the relative availability of different electron acceptors, in any one depth interval in the ocean, exactly the same proportion of organic matter will be degraded. We avoid the alternative here – a fully kinetic set of equations where each electron acceptor is associated with a different rate of degradation – partly because of the additional set of poorly constrained (kinetic rate constant) parameters that would be required, and partly because to implement such a scheme effectively, requires knowledge about the composition of settling organic matter and how its relative reactivity changes with time (Ridgwell, 2011; LaRowe and Van Cappellen, 2011).

As a consequence of organic matter remineralisation in the model, DIR produces ferrous iron (Fe^{2+}), which is re-oxidised when it comes into contact with oxygen (e.g., via upwelling of the reduced compound or downwelling of oxygen) via ferrous iron oxidation (FIO, R5 in Table 1) (Millero et al., 1987a). Similarly, any reduced sulphide (because we do not consider sulphide speciation explicitly, reduced sulphide is defined as $\Sigma H_2S = H_2S + HS^-$) that comes into contact with oxygen is



Table 2. List of kinetic rate expressions for the reactions included in the cGENIE model. All expressions are based on standard kinetic formulations in biogeochemical models (see text for details). The values of the kinetic constants are listed in Table 3.

Reaction number	Reaction name	Kinetic expression
R1	Aerobic respiration	$f_{AR}R_{min}$
R2	Dissimilatory iron reduction	$f_{DIR}R_{min}$
R3	Dissimilatory sulphate reduction	$f_{DSR}R_{min}$
R4	Methanogenesis	$f_{MG}R_{min}$
R5	Ferrous iron oxidation	$k_{FIO}[\Sigma Fe^{2+}][O_2]$
R6	Canonical sulphide oxidation	$k_{CSO}[\Sigma H_2S][O_2]^2$
R7	Sulphide-mediated iron reduction ¹	$k_{SMI,d/s}[\Sigma H_2S]^{0.5}[Fe^{3+}]$
R8	Iron oxide precipitation ²	$k_{scav}[Fe^{3+}]F_{POC}(z)$
R9	Pyrite precipitation	$k_{PyP}[FeS_p][\Sigma H_2S]$
R10	Siderite precipitation	$k_{AFC} e^{(b_{AFC} \log_{10}(IAP_{siderite}))}$
R11	Greenalite precipitation	$k_{greenalite} e^{(b_{greenalite} \log_{10}(SI_{greenalite}))}$

¹Two different kinetic constants are used for the reactions between Fe^{3+} and sulphide ($k_{SMI,d}$) and $FeOOH$ and sulphide ($k_{SMI,s}$), see text for details. ² The formation of iron oxide minerals is parametrised according to Parekh et al. (2004), and is explained in more detail in section 3.1.1.

re-oxidised via canonical sulphur oxidation (CSO, R6 in Table 1) (Millero et al., 1987b). Additionally, H_2S is oxidised by reaction with oxidised iron via sulphide-mediated iron reduction (SMI, R7 in Table 1) (Fig. 3b; Canfield, 1992; Poulton et al., 2004a; Mikucki et al., 2009). The oxidised form of iron, ferric iron (Fe^{3+}), will precipitate out as iron oxide ($FeOOH$) minerals (IrP, R8 in Table 1).

- 5 When Fe^{2+} and H_2S are simultaneously present in the water column (in sufficient concentrations), they can form the mineral pyrite (FeS_2) via pyrite precipitation (PyP, R9 in Table 1) (Rickard, 1997). Alternatively, when Fe^{2+} accumulates past its saturation state with bicarbonate, siderite ($FeCO_3$) forms via siderite precipitation (SiP, R10 in Table 1) (Jimenez-Lopez et al., 2004; Jiang and Tosca, 2019). Finally, greenalite ($Fe_3Si_2O_5(OH)_4$) precipitates when Fe^{2+} and dissolved silica (SiO_2) are saturated with respect to greenalite precipitation (GrP, R11 in Table 1) (Tosca et al., 2015; Rasmussen et al., 2015).
- 10 Pyrite, siderite and greenalite are subsequently buried in the sediment, together with any solid $FeOOH$ that has not reacted (the half-life of iron oxides in euxinic waters is in the order of 10s - 100s of days, which is comparable to the residence time of a particle in the ocean; Poulton et al., 2004a) (Fig. 3b). These four solid iron phases are the main burial phases for reactive iron, and form the basis of the Fe-speciation proxy used to reconstruct local redox conditions in past oceans (Poulton and Canfield, 2011).



3.1 The iron cycle

The cGENIE model already included the representation of a simplified iron cycle designed to account for iron limitation of biological productivity at the ocean surface (Tagliabue et al., 2016). However, because its initial use was to model the present-day oceanic iron cycle, it does not contain an anoxic iron cycle (as 95% of the today's ocean volume is oxic; Diaz and Rosenberg, 2008), nor does it contain a coupling between the iron and sulphur cycles (e.g., via the precipitation of FeS_2). The absence of an anoxic iron cycle limits the suitability of cGENIE for simulating low-oxygen worlds in which the ocean interior is pervasively anoxic and iron and sulphur cycling would have dominated ocean biogeochemistry (Poulton and Canfield, 2011; Raiswell and Canfield, 2012). In addition to summarising the existing oxic cycle, we expand the model to include key processes operating in anoxic environments, as follows.

10 3.1.1 The oxic iron cycle

In oxygenated waters, iron is predominately present in its oxidised form (Fe^{3+}). The oxic iron cycle in cGENIE follows Parekh et al. (2004), and has been recently updated with a revised set of parameters, calibrated based on the present day iron cycle (Tagliabue et al., 2016). Briefly; total dissolved Fe_{total}^{3+} consists of free Fe_{free}^{3+} and ligand-bound Fe_{ligand}^{3+} , and only the free fraction can be scavenged (and thus form $FeOOH$), or taken up by biology to fuel primary productivity (see section 2.2). The scavenging rate of iron is a function of the concentration of free Fe^{3+} together with the magnitude of the POC flux from the grid cell, i.e.,

$$R_{IrP} = k_{scav}[Fe_{free}^{3+}]F_{POC}(z) \quad (11)$$

where k_{scav} ($=1.43 \times 10^{-6} \text{ mol}^{-1} \text{ m}^2 \text{ h}^{-1}$) is a scavenging constant calibrated to the modern day distribution of Fe (Tagliabue et al., 2016). We assume that the complexation reaction is always in equilibrium, which allows for the calculation of the amount of Fe_{free}^{3+} at each time step by the conservation equations

$$\begin{aligned} [Fe_{total}^{3+}] &= [Fe_{ligand}^{3+}] + [Fe_{free}^{3+}] \\ [L_{total}] &= [Fe_{ligand}^{3+}] + [L_{free}] \\ K_{sp}^{FeL} &= \frac{[Fe_{ligand}^{3+}]}{[Fe_{free}^{3+}][L_{free}]} \end{aligned} \quad (12)$$

where K_{sp}^{FeL} is the stability constant of the $L - Fe^{3+}$ complex ($1.0 \times 10^{11} \text{ M}^{-1}$; Table 3), and L_{total} is the total amount of ligand.

Oxidised iron is very insoluble in seawater and will rapidly form particulate oxides ($FeOOH$), driving down equilibrium dissolved Fe^{3+} abundance to picomolar values without stabilisation by ligands (Liu and Millero, 2002). Thus, Fe_{free}^{3+} will form colloidal or nanoparticulate iron oxides – $FeOOH$ – which can then adsorb on other particles or aggregate to bigger particles and sink through the water column (Raiswell and Canfield, 2012). In cGENIE, the pool of Fe_{free}^{3+} can be thought of as a mix of purely dissolved Fe^{3+} and a range of colloidal and nanoparticulate Fe^{3+} phases that do not settle efficiently




through the water column. The Fe_{free}^{3+} pool can then further react or scavenge onto particles that allow it to more effectively settle through the water column.

In our new formulation of the Fe cycle in cGENIE, Fe_{free}^{3+} (that is not stabilised by ligands) is scavenged by settling POC (using a fixed scavenging rate; Eq. 11) and sinks as a 'marine snow' particle through the water column. This scavenging mechanism is commonly used in reactive-transport models of ferruginous lakes or the modern ocean (see e.g. Taillefert and Gaillard, 2002; Tagliabue et al., 2016). Marine snow formation is what likely happens today, where relatively high production of organic matter drives the transport of chemically heterogeneous aggregates (containing iron oxides or barite; Dehairs et al., 1990; Balzano et al., 2009). When scavenged on POC, $FeOOH$ can be used for DIR when oxygen becomes depleted (see Section 3.1.2).

Note that we also provide the option in the model for Fe_{free}^{3+} to precipitate as a pure $FeOOH$ phase, without being associated with POC. In this case, dissimilatory iron reduction does not occur in the water column since both POC and $FeOOH$ are not associated within the same particle. This situation would be more appropriate in the case of high amounts of iron oxidation, and lower production of organic matter (which was potentially the case in the Archean; Thompson et al., 2019). This configuration is by default implemented using a numerical 'cut-off', where all Fe_{free}^{3+} that passes the solubility threshold of 0.5 nM (Liu and Millero, 2002) is considered particulate and sinks through the water column.

To close the oceanic iron cycle, we have introduced a new dissolved iron species – Fe^{2+} – any reduced Fe^{2+} that is mixed into the oxic zone is being rapidly oxidised to Fe_{total}^{3+} via ferrous iron oxidation (FIO),



where the rate equation can be expressed as

$$R_{FIO} = k_{FIO}[Fe^{2+}][O_2] \quad (14)$$

and where k_{FIO} is a reaction rate constant ($k_{FIO} = 0.115 \times 10^6 M^{-1}h^{-1}$) (Table 2; Millero et al., 1987a).

In summary, we have extended the basic representation of Parekh et al. (2004), that considered only a generic free 'dissolved' iron (that could be scavenged) and a ligand-bound iron phase, to now distinguish between the oxidation states of iron (and the tracers Fe^{2+} and Fe_{total}^{3+}), with Fe_{total}^{3+} being split into free (and scavengeable) and ligand-bound phase.

3.1.2 The anoxic iron cycle

In an anoxic water column, oxidised forms of iron can be reduced to Fe^{2+} either via dissimilatory iron reduction or via sulphur-mediated iron reduction. We describe the two processes individually as follows.

Dissimilatory iron reduction in the water column

30

The majority of oxidised iron exists in particulate form due to the low solubility of Fe^{3+} , which poses a challenge for microorganisms performing dissimilatory iron reduction in the water column. Iron reducers need physical contact with the iron mineral



to reduce iron (Gorby, 2006), which is difficult when both organic matter and $FeOOH$ are in particulate form and are dispersed separately throughout the aqueous medium. Even in a sediment column, where all particles are packed closely together, DIR generally requires sediment homogenisation by burrowing fauna to become volumetrically important (Thamdrup, 2000; van de Velde and Meysman, 2016). Indeed, in a sulphate-rich, ancient marine brine, Fe^{3+} was found to be the terminal electron
5 acceptor, but sulphur ultimately acted as a redox shuttle between organic matter and iron oxides (Mikucki et al., 2009). Studies of anoxic Lake Pavin suggest that most of the iron reduction occurs in the sediment, rather than the water column (Michard et al., 1994; Cosmidis et al., 2014), although the same studies indicated that manganese oxide reduction (which has a very similar mechanism and inhibition concentration as DIR; Lovley, 1991; Thamdrup, 2000) does occur in the water column. However, there is direct evidence for iron reduction occurring coupled to organic matter mineralisation in the water column in a range of
10 other anoxic lakes, such as Lake Sammamish (Washington, USA; Balistrieri et al., 1992), Lake Cadagno (Switzerland; Berg et al., 2016), Lake Matano (Indonesia; Crowe et al., 2008), and Paul Lake (Michigan, USA; Taillefert and Gaillard, 2002). These studies suggested that DIR was coupled to either the oxidation of dissolved organic matter (Crowe et al., 2008), or reduction of iron in aggregates with organic matter. The latter has been experimentally shown to occur (Balzano et al., 2009). We consider this to be strong evidence that DIR can be coupled to POC oxidation in a water column, especially in the ocean where the
15 residence time of particles is considerably longer than in a much shallower lake environment.

To model DIR in the water column, we use the mathematical expression of DIR in marine sediments, where $FeOOH$ is a common electron acceptor for organic matter mineralisation (Thamdrup, 2000). Dissimilatory iron reduction has an overall reaction stoichiometry of,



20 where it is implicitly assumed that every iron oxide particle consists solely of Fe^{3+} , rather than a mixture of redox states as is sometimes encountered at the interface of ferruginous lakes (Zegeye et al., 2012). This assumption greatly simplifies the reaction scheme and parameter set. Dissimilatory iron reduction generally becomes limited when concentrations of $FeOOH$ drop below $\sim 30 \mu mol cm^{-3}$ (Van Cappellen and Wang, 1996; Thamdrup, 2000). This limitation is expressed as $\frac{[FeOOH]}{K_{0,FeOOH} + [FeOOH]}$ in Eq. 10 and follows the conventional limitation-inhibition scheme (Soetaert et al., 1996), where the parameter $K_{0,FeOOH}$ expresses the concentration at which DIR occurs at half of its maximum rate. In marine sediments, DIR
25 generally occurs before sulphate reduction since it is a more energy-yielding electron acceptor (Thamdrup, 2000), although the wide range of reactivities for different iron oxide minerals often leads to an overlap of DIR and DSR zones (Postma and Jakobsen, 1996). This limitation is expressed as $\frac{[K_{i,FeOOH}]}{K_{i,FeOOH} + [FeOOH]}$ in Eq. 10, where $K_{i,FeOOH}$ expresses the concentration above which $FeOOH$ inhibits other mineralisation pathways (DSR and MG). In early diagenetic models, this concentration is
30 generally assumed to be identical to the limitation parameter $K_{0,FeOOH}$ (Van Cappellen and Wang, 1996; Soetaert et al., 1996; van de Velde and Meysman, 2016). As a baseline value, we assume that $K_{0,FeOOH}$ and $K_{i,FeOOH}$ both equal $10^{-3} mol kg^{-1}$ ($\sim 1 \mu mol cm^{-3}$, comparable to the inhibition concentration in marine sediments), but as explained above, these parameter values are subject to a high degree of uncertainty. Therefore, we include parameters $K_{0,FeOOH}$ and $K_{i,FeOOH}$ in our model



sensitivity testing (section 4.3).

Sulphur-mediated iron reduction

- 5 Oxidised iron in the ocean can also be reduced via sulphur-mediated iron reduction (SMI), which follows the stoichiometry:



where the reaction rate can be expressed as (Poulton et al., 2004a):

$$R_{SMI} = k_{SMI,d/s} [\Sigma H_2S]^{0.5} [Fe^{3+}] \quad (17)$$

- Note that $[Fe^{3+}]$ in Eq. 17 can represent dissolved Fe_{total}^{3+} or solid $FeOOH$. We assume that the dissolved form is highly
10 reactive with sulphide (a reaction time of ~ 5 minutes, which is comparable to the reactivity of freshly precipitated hydrous
ferric oxide) and has a reaction rate constant of $k_{SMI,d} = 2.64 \times 10^2 M^{-0.5} h^{-1}$ (Poulton et al., 2004a). The solid form of
 $FeOOH$ can represent a number of different oxidised iron minerals which are reactive towards sulphide on timescales ranging
from hours to hundreds of days (Poulton et al., 2004a). For our baseline simulations, we assume that all $FeOOH$ precipitates
as lepidocrocite, which has a reactivity constant of $k_{SMI,s} = 1.98 \times 10^0 M^{-0.5} h^{-1}$ (Poulton et al., 2004a). Lepidocrocite is
15 less crystalline than the other (non-hydrous) iron oxides and precipitates when the rate of Fe^{3+} supply is low relative to the rate
of precipitation, as is generally the case in natural systems (Crosby et al., 1983). We discuss the model sensitivity to choices of
 $k_{SMI,s}$ in Section 4.3. The reduction of Fe^{3+} produces Fe^{2+} , which can either be re-oxidised when it comes in contact with
 O_2 (see section 3.1.1) or can form reduced minerals.

3.1.3 Reduced iron mineral formation

- 20 Reduced Fe^{2+} can form complexes with a number of ligands that are common in seawater, including Cl^- , SO_4^{2-} and bicar-
bonate (Millero et al., 1995). Under anoxic conditions, however, the most important ligand is free sulphide (Rickard, 2006),
which is produced by sulphate reduction. Together, dissolved Fe_{free}^{2+} and the aqueous iron-sulphide complex (FeS_{aq}) make
up ~ 100 % of the total dissolved iron pool in sulphidic-anoxic seawater (Fig. 4a). The thermodynamic equilibrium can be
calculated as

$$25 \quad K_{sp}^{FeS_{aq}} = \frac{[Fe^{2+}][\Sigma H_2S]}{[FeS_{aq}]} = 10^{-5.08} M \quad (18)$$

which was obtained from the visualMINTEQ database (Gustafsson, 2019), and is based on stability constants calculated by
Luther et al. (1996).

- Since dissolved Fe_{free}^{2+} and the FeS_{aq} complex are the dominant dissolved forms of reduced ferrous iron in natural waters
(in anoxic seawater devoid of sulphide, dissolved Fe_{free}^{2+} still represents ~ 80 % of the dissolved Fe_{total}^{2+} pool; Fig. 4b), we
30 choose to only consider those two species. We do not explicitly model the FeS_{aq} complex, but calculate the thermodynamic
equilibrium before each reaction proceeds, implicitly assuming that it is reached much faster than any of the kinetic reactions




Figure 4. Relative importance of different dissolved Fe complexes, obtained by visualMINTEQ (Gustafsson, 2019). Concentrations of other constituents are chosen to represent average seawater, with dissolved organic carbon (DOC) concentration from Sharp et al. (1995).

in our reaction set (as is commonly done for the complexation of Fe^{3+} ; see section 3.1.1). This is important, since FeS_{aq} forms particulate FeS_p , which is the precursor of pyrite (FeS_2) (Rickard, 1997, 2006). Furthermore, siderite or greenalite can only precipitate from Fe^{2+}_{free} (Tucker, 2001; Tosca et al., 2015), and their precipitation rates are consequently dependent on the dissolved Fe^{2+}_{free} concentration.

- 5 Before each precipitation reaction, we calculate the concentration of FeS_{aq} , Fe^{2+}_{free} and H_2S_{free} using equation 18. We then compare the FeS_{aq} concentration to the solubility threshold ($10^{-5.7}$ M; Rickard, 2006). If FeS_{aq} surpasses the solubility value, some fraction of it precipitates instantaneously as particulate iron monosulfide (FeS_p) (Suits and Wilkin, 1998). Note that here we consider FeS_p solubility to be independent of pH, which should be valid for seawater pH above 7 (Rickard, 2006). Pyrite can then form locally via reaction between FeS_p and free ΣH_2S , with the production of free hydrogen gas (as S^{2-} in
- 10 ΣH_2S and FeS_p has to be oxidised to S^{1-} in FeS_2 ; Rickard, 1997),



Since hydrogen-gas is highly reactive it is almost instantaneously oxidised, with the reduction of an electron acceptor, like O_2 , SO_4^{2-} or CO_2 . Given that pyrite precipitates under sulphidic conditions, oxygen is absent, and sulphate is the most likely electron acceptor



To preserve the redox balance while avoiding the need to model an additional state variable (H_2), we can combine reactions 19 and 20 to



- By using equation 21 we implicitly assume that the hydrogen gas produced during pyrite precipitation immediately reacts with
- 20 SO_4^{2-} . Since pyrite formation is not an equilibrium reaction (pyrite minerals are stable in seawater), the reaction of pyrite is



described as a kinetic reaction with a second-order dependency on $[FeS_p]$ and $[\Sigma H_2S]$, with $k_{PyP} = 0.3708 \times 10^0 M^{-1} h^{-1}$ (Rickard, 1997). The rate equation for FeS_2 precipitation can then be written as

$$R_{PyP} = k_{PyP} [FeS_p] [\Sigma H_2S] \quad (22)$$

Any Fe^{2+} that is not complexed with sulphide (Fe_{free}^{2+}) can form siderite ($FeCO_3$), an iron-carbonate mineral (Tucker, 2001).



Recent experiments by Jiang and Tosca (2019) have shown that the precipitation of iron-carbonate phases is controlled by the formation of amorphous Fe carbonate (AFC), which has the stoichiometric formula $FeCO_3(OH)_{1/2}$. Since the thermodynamic data required to calculate the mineral solubility product of AFC is currently lacking, Jiang and Tosca (2019) have determined the rate of AFC precipitation based on the ion activity product (IAP),

$$IAP_{AFC} = \gamma_{Fe^{2+}} [Fe^{2+}] \gamma_{CO_3^{2-}} [CO_3^{2-}] (\gamma_{OH^-} [OH^-])^{0.5} \quad (24)$$

where γ represents the activity coefficient of a given ion in seawater (Table 3). The rate-limiting step in the precipitation reaction is the spontaneous nucleation of AFC, and hence the empirically derived rate of AFC precipitation follows an exponential function that describes the nucleation rate from water (Steeffel and Van Cappellen, 1990; Jiang and Tosca, 2019),

$$R_{SiP} = k_{AFC} e^{b_{AFC} \log_{10} IAP_{AFC}} \quad (25)$$

where k_{AFC} is $1.963 \times 10^{-14} M h^{-1}$ and b_{AFC} equals 9.042 (Table 3).

Another reduced iron mineral potentially important for anoxic, Fe-rich environments is the iron-silicate mineral greenalite ($Fe_3Si_2O_5(OH)_4$) (Tosca et al., 2015):



The precipitation rate of greenalite is dependent on its degree of supersaturation (i.e. its saturation index), which can be calculated based on its IAP,

$$IAP_{greenalite} = \frac{(\gamma_{Fe^{2+}} [Fe^{2+}])^3 (\gamma_{SiO_2} [SiO_2])^2}{(\gamma_{H^+} [H^+])^6} \quad (27)$$

where γ represents the activity coefficient of an ion in seawater (Table 3), and $K_{sp}^{greenalite}$ its solubility product (Rasmussen et al., 2015)

$$SI_{greenalite} = \log_{10} \left(\frac{IAP}{K_{sp}^{greenalite}} \right) \quad (28)$$

where $K_{sp}^{greenalite}$ equals $3.98 \times 10^{27} M^{-1}$ (Tosca et al., 2015). The rate equation follows an exponential function that describes the nucleation rate from water (as in siderite, see above; Rasmussen et al., 2015; Jiang and Tosca, 2019)

$$R_{GrP} = k_{greenalite} e^{b_{greenalite} \log_{10} SI_{greenalite}} \quad (29)$$

where $k_{greenalite}$ is $6.996 \times 10^{-13} M h^{-1}$ and $b_{greenalite}$ equals 1.856 (Table 3).



3.2 The sulphur cycle

The oxidised form of dissolved sulphur in the ocean is sulphate (SO_4^{2-}). Under anaerobic conditions, sulphate is used as terminal electron acceptor during organic matter remineralisation:



5 The produced sulphide can then be re-oxidised when it comes into contact with oxygen via canonical sulphide oxidation (CSO)



The rate of CSO in cGENIE is dependent on the concentration of the electron donor (ΣH_2S) and acceptor (O_2) and a second order rate constant $k_{CSO} = 0.625 \times 10^6 M^{-2} h^{-1}$ (Zhang and Millero, 1993):

$$10 \quad R_{CSO} = k_{CSO}[\Sigma H_2S][O_2]^2 \quad (32)$$

Alternatively, ΣH_2S can be re-oxidised with Fe^{3+} (either Fe_{total}^{3+} or solid $FeOOH$) during sulphide-mediated iron reduction (as described in reaction 16).

In cGENIE, the eventual sink for sulphur is the precipitation of pyrite (reaction 19) (ignoring sulphide reacting with organic matter – see Hülse et al. (2019)). We do not currently include precipitation of gypsum ($CaSO_4$) in our model description (Fig. 3) – a sink likely less important on a globally integrated basis during Precambrian time, or during any other period in which ocean $[SO_4^{2-}]$ was relatively low (Grotzinger and Kasting, 1993; Crowe et al., 2014; Fakhraee et al., 2019). Planned future developments to cGENIE will incorporate an explicit gypsum cycle.

3.3 Isotope geochemistry

A particularly important application of an anoxic Fe-S cycle in cGENIE is exploring ocean redox landscapes during Precambrian time, when ocean biogeochemical cycling was dominated by iron and sulphur (Raiswell and Canfield, 2012). As noted above, one way of comparing our model output to available data is the explicit simulation of the burial phases of iron, which allows comparison to the often used Fe-proxy (Poulton and Canfield, 2011). A different and independent constraint potentially exists in the form of Fe or S isotopes (see e.g. Beard et al., 1999; Gomes and Johnston, 2017; van de Velde et al., 2018).

Any chemical element with multiple stable isotopes (Fe for example has four stable forms ^{54}Fe , ^{56}Fe , ^{57}Fe and ^{58}Fe) can potentially be used to track physicochemical processes that act to partition stable isotopes according to thermodynamic or kinetic principles. For Fe the most abundant isotopes are ^{54}Fe and ^{56}Fe , and deviations in the $^{56}Fe/^{54}Fe$ ratio of Fe-bearing aqueous and mineral phases from that of a reference material can be described using conventional δ -notation:

$$\delta^{56}Fe = \left(\frac{\left(\frac{^{56}Fe}{^{54}Fe} \right)}{\left(\frac{^{56}Fe}{^{54}Fe} \right)_{ref}} - 1.0 \right) \times 1000 \quad (33)$$



Table 3. List of kinetic constants for the reactions included in the cGENIE model.

Reactivity constants	Symbol	Unit	Value	Reference
Limitation constant oxygen reduction	K_{0,O_2}	M	8.0×10^{-6}	1
Inhibition constant oxygen reduction	K_{i,O_2}	M	8.0×10^{-6}	1
Limitation constant DIR	$K_{0,FeOOH}$	M	1.0×10^{-2}	2
Inhibition constant DIR	$K_{i,FeOOH}$	M	1.0×10^{-2}	2
Limitation constant DSR	$K_{0,SO_4^{2-}}$	M	5.0×10^{-4}	3
Inhibition constant DSR	$K_{i,SO_4^{2-}}$	M	1.0×10^{-3}	1
Canonical sulphide oxidation	k_{CSO}	$M^{-2}h^{-1}$	0.625×10^6	4
Ferrous iron oxidation	k_{FIO}	$M^{-1}h^{-1}$	0.115×10^6	5
Sulphide-mediated iron reduction (dissolved)	k_{SMId}	$M^{-0.5}h^{-1}$	2.64×10^2	6
Sulphide-mediated iron reduction (solid)	k_{SMIs}	$M^{-0.5}h^{-1}$	1.98×10^0	6
Fe_{free}^{3+} Scavenging constant	k_{scav}	$mol^{-1}m^2h^{-1}$	1.43×10^{-6}	7
$L - Fe^{3+}$ complex stability constant	K_{sp}^{FeL}	M^{-1}	1.0×10^{11}	7
Solubility product FeS_{aq}	$K_{sp}^{FeS_{aq}}$	M	8.32×10^{-6}	8
Kinetic constant pyrite precipitation	k_{PyP}	$M^{-1}h^{-1}$	0.3708×10^0	9
Kinetic constant siderite precipitation	k_{AFC}	Mh^{-1}	1.963×10^{-14}	10
Kinetic exponent siderite precipitation	b_{AFC}	-	9.042×10^0	10
Solubility product greenalite	$K_{sp}^{greenalite}$	M^{-1}	3.98×10^{27}	11
Kinetic constant greenalite precipitation	$k_{greenalite}$	Mh^{-1}	6.996×10^{-13}	12
Kinetic exponent greenalite precipitation	$b_{greenalite}$	-	1.856×10^0	12
<i>Activity coefficients</i>				
Activity constant H^+	γ_{H^+}	-	0.73	13
Activity constant OH^-	γ_{OH^-}	-	0.69	14
Activity constant CO_3^{2-}	$\gamma_{CO_3^{2-}}$	-	1.17	15
Activity constant Fe^{2+}	$\gamma_{Fe^{2+}}$	-	0.23	14
Activity constant SiO_2	γ_{SiO_2}	-	1.13	14

¹(Ridgwell et al., 2007),²(Thamdrup, 2000),³(Olson et al., 2016),⁴(Zhang and Millero, 1993),⁵(Millero et al., 1987a),⁶(Poulton et al., 2004a),⁷(Ridgwell and DeAth, in prep.),⁸(Luther et al., 1996),⁹(Rickard, 1997),¹⁰(Jiang and Tosca, 2019),¹¹(Tosca et al., 2015),¹²(Rasmussen et al., 2015),¹³(Marion et al., 2011),¹⁴(Following the Davies equation),¹⁵(Johnson, 1982)

where $(\frac{^{56}Fe}{^{54}Fe})_{ref}$ is the isotope ratio of a standard reference material (IRMM-14). Any geochemical reaction, be it biotic (mediated by micro-organisms) or abiotic, can induce isotopic fractionation between co-occurring Fe- or S-bearing phases.



To model the isotopic signatures of Fe and S , we track the concentrations of the 'bulk' pools (C_i) and the isotope-specific pools ($^{56}C_i$ for example, in the case of Fe). The isotopic signature of an Fe species C_i is then calculated as

$$\delta^{56}Fe_{C_i} = \left(\frac{\left(\frac{^{56}C_i}{C_i - ^{56}C_i} \right)}{\left(\frac{^{56}Fe}{^{54}Fe} \right)_{ref}} - 1.0 \right) \times 1000 \quad (34)$$

Each individual reaction R_k is assigned a fractionation factor $^{56}\epsilon_{R_k}$ (in ‰; Table 4), which relates to $^{56}\alpha_{R_k}$ as

$$^{56}\alpha_{R_k} = 1 + \frac{^{56}\epsilon_{R_k}}{1000} \quad (35)$$

Fractionations are then implemented by calculating the rate for the ^{56}Fe pool from the bulk reaction rate R_k

$$^{56}R_k = \frac{^{56}\alpha_{R_k} \ ^{56}r_{C_i}}{1 + ^{56}\alpha_{R_k} \ ^{56}r_{C_i}} R_k \quad (36)$$

where

$$^{56}r_{C_i} = \frac{^{56}C_i}{C_i - ^{56}C_i} \quad (37)$$

10 In this way, we assign a fractionation factor to each of the reactions considered in our model, and are able to track the isotopic signature of each Fe and S species. This allows us to simulate the $\delta^{56}Fe$ and $\delta^{34}S$ values of dissolved species and solid mineral phases, both of which can potentially be compared to observations from the geological record.

The obvious limitation of assigning a constant fractionation factor to each reaction is that this is incapable of fully capturing natural isotopic variability. For instance, different microbial strains of sulphur reducers (or iron oxidisers) express different
 15 fractionation factors, even though the overall reaction remains the same (Gomes and Johnston, 2017; Pellerin et al., 2019). This is reflected in the often broad range of isotopic fractionation factors found in the literature for a given process (Table 4). Other factors influencing microbial fractionation are local environmental conditions, such as electron donor type/availability (Wing and Halevy, 2014; Pellerin et al., 2018), or evolutionary adaptation (Pellerin et al., 2015). Aside from biologically mediated
 20 transformations, kinetic effects associated with abiotic aqueous reactions and precipitation of solid phases also affect the fractionation that is eventually recorded in the end-product. All these factors can make interpretation of isotope fractionations very complex in natural settings, and it is thus highly unlikely that any particular model simulation will be able to exactly reproduce observed isotope records. Nevertheless, the scheme employed here should be able to discern first-order observations from the geologic record, and in some cases could potentially be used to rule out particular end-member hypotheses for ocean chemistry.

25 4 Model testing

For contrasting with observations, we are severely limited because the modern ocean is largely well-oxygenated, with only a few oxygen-minimum zones near highly productive margins such as the Indian ocean and the Peruvian margin (Keeling et al., 2010). However, even in these regions dissolved oxygen concentrations rarely reach zero, and the development of ferruginous



Table 4. Isotope fractionation factors

Reaction	Reactant	Product	Fractionation factor	Literature min	Literature max	References
$^{34}\epsilon_{R-P}$ (<i>S</i> fractionation)						
Dissimilatory sulphate reduction	SO_4^{2-}	H_2S	-30.0‰	-70.0‰	-2.0‰	[1]-[3]
Canonical sulphide oxidation	H_2S	SO_4^{2-}	-10.0‰	-18.0‰	+12.5‰	[4],[5]
Sulphide mediated iron reduction	H_2S	SO_4^{2-}	-1.8‰	-3.6‰	0.0‰	[6],[7]
Pyrite precipitation	H_2S	FeS_2	0.0‰	-0.4‰	1.2‰	[8],[9]
$^{56}\epsilon_{R-P}$ (<i>Fe</i> fractionation)						
Dissimilatory iron reduction	Fe^{3+}	Fe^{2+}	-1.3‰	-2.95‰	-1.3‰	[10]-[14]
Sulphide mediated iron reduction	Fe^{3+}	Fe^{2+}	-1.3‰	-1.3‰	-1.3‰	[10]-[14]
Ferrous iron oxidation	Fe^{2+}	Fe^{3+}	0.8‰	0.4‰	1.1‰	[13],[15]
Iron oxide precipitation ¹	Fe^{3+}	$FeOOH$	0.0‰	-	-	-
Pyrite precipitation	Fe^{2+}	FeS_2	-2.2‰	-2.9‰	-1.5‰	[15],[16]
Siderite precipitation	Fe^{2+}	$FeCO_3$	-0.3‰	-0.6‰	0.0‰	[12],[17]
Greenalite precipitation ²	Fe^{2+}	$Fe_3Si_2O_5(OH)_4$	0.0‰	-	-	-

[1] (Kaplan and Rittenberg, 1964), [2] (Detmers et al., 2001), [3] (Sim et al., 2011), [4] (Gomes and Johnston, 2017), [5] (Pellerin et al., 2019), [6] (Poser et al., 2014), [7] (Fry et al., 1988), [8] (Botzcher et al., 1998), [9] (Price and Shieh, 1979), [10] (Beard et al., 1999), [11] (Beard et al., 2003), [12] (Johnson et al., 2004), [13] (Bullen et al., 2001), [14] (Crosby et al., 2007), [15] (Rolison et al., 2018), [16] (Guilbaud et al., 2011), [17] (Wiesli et al., 2004)

¹ The isotopic fractionation for iron oxide precipitation is driven by the oxidation reaction (FIO). ² The isotope fraction factor for greenalite precipitation is currently unknown.

or euxinic conditions is essentially absent. Only highly restricted basins such as the Cariaco basin or the Black and Baltic Seas can develop euxinic conditions, but these conditions arise as a result of local circulation within silled or enclosed basins, and are not likely to be representative of an anoxic open ocean setting. As a result, we lack observations to which we can directly calibrate our model. Hence, instead of calibrating our model results on a large observational dataset from the modern oceans, we will show the spatial concentration and isotope features of a hypothetical anoxic ocean (section 4.2), and – where possible – compare our predicted reaction rates to rates obtained from ancient ocean analogues. Subsequently, we broadly illustrate the sensitivity of the model output to the newly introduced parameters of the iron-sulphur cycling (section 4.3). As discussed below, our model is largely robust across a range of values for key parameters, and predicts reaction and process rates that are comparable to those obtained experimentally or observed in modern analogue environments.

10 4.1 Model configuration

For testing and characterizing the Fe-S cycle model developments, we configure cGENIE with a single continent that runs from pole to pole (Fig. 2c; Fig. 5a). Each model experiment is initialised from a homogeneous and static ocean, with an imposed constant atmospheric O_2 concentration of 0.1 PAL (present atmospheric level, i.e. 21,000 ppm) and an atmospheric CO_2 concentration of ~ 16 PAL (5337.6 ppm). These atmospheric boundary conditions are chosen as broadly plausible for the



Precambrian Earth system and deliberately preclude the formation of sea-ice in order to simplify the interpretation of the results. The model is run in 'closed' configuration for all elements (notably C and P), in which the ocean-atmosphere inventory for each element is always conserved. An exception is made for Fe and S , which enter the surface ocean in the form of Fe^{2+} and SO_4^{2-} , and exit the ocean as $FeOOH$, FeS_2 , $FeCO_3$ or $Fe_3Si_2O_5(OH)_4$. Fluxes of Fe and S are chosen to be in balance with respect to FeS_2 burial (S:Fe=2:1), and to represent the best estimate of present-day weathering fluxes to the ocean (excluding reprocessing in inner shelf sediment settings) ($F_{Fe^{2+}} = 1.3 \times 10^{12} \text{ mol yr}^{-1}$, $F_{SO_4^{2-}} = 2.6 \times 10^{12} \text{ mol yr}^{-1}$; Poulton and Raiswell, 2002; Raiswell and Canfield, 2012). Hydrothermal systems represent another potentially important flux of Fe to the ocean (Tagliabue et al., 2010; Conway and John, 2014; Lough et al., 2019), and this flux is likely to be elevated when ocean chemistry is pervasively anoxic and relatively low in SO_4^{2-} (Kump and Seyfried, 2005). Our simulations therefore also include a hydrothermal flux of Fe^{2+} , broadly comparable to a plausible input flux to Proterozoic oceans ($15.1 \times 10^{12} \text{ mol yr}^{-1}$; Thompson et al., 2019). This Fe^{2+} flux in our model setup is equally distributed along a 'hydrothermal ridge' located in the middle of the ocean (a straight line from $k=9, l=3$ to $k=9, l=15$; Fig. 2a). In order to balance the S:Fe flux ratio at 2:1 (see above), we must also specify a uniform surface flux of SO_4^{2-} of $30.2 \times 10^{12} \text{ mol yr}^{-1}$. This is an extremely high S flux, and is unlikely to be realistic on long timescales. However, it allows us to quickly diagnose spatial patterns in reducing Fe-S cycling at steady state, without *a priori* introducing a bias towards ferruginous or euxinic redox states. We initialise the model with a semi-arbitrary marine phosphate inventory that is 50% of today in order to reduce marine primary productivity (as productivity in the Precambrian ocean was likely lower than today; Crockford et al., 2018; Ozaki et al., 2019), and run the ocean circulation to steady state for 20,000 years. From that spin-up, we then run each individual experiment for 40,000 years and present model output from the 40,000th year of integration. It is important to emphasise that the idealised configuration we implement here is not meant to represent any specific period or event in Earth's history, but is rather meant to serve as a broadly plausible and computationally efficient set of boundary conditions for testing the extended model code.

The sea surface temperature and ocean circulation generated by our configuration of cGENIE are shown in Fig. 5. Even though we have a symmetric continent, the overturning patterns are skewed to the southern hemisphere, with a strong anti-clockwise circulation at around -60°N (Fig. 5c). In contrast, the barotropic streamfunction shows a large degree of symmetry (Fig. 5d). Since the aim of this manuscript is to describe the newly developed Fe-S chemistry, and not ocean circulation in real or fake worlds, we will not further discuss the emerging patterns of ocean circulation.

4.2 Spatial output

In the following section, we briefly discuss the spatial model output and modelled redox cycling for our baseline configuration. Precipitation fluxes of oxidised Fe^{3+} are specified according to the scavenging scheme (Eq. 11). We focus here on: (i) three depth slices of particle sinking fluxes of POC , $FeOOH$, FeS_2 , and $Fe_3Si_2O_5(OH)_4$ (Fig. 6; $FeCO_3$ fluxes were negligible and are not shown); (ii) the average and three vertical longitudinal slices of O_2 , dissolved Fe_{total}^{3+} , Fe_{total}^{2+} and dissolved ΣH_2S (Fig. 7); and (iii) an overview of globally averaged reaction rates and iron mineral burial fluxes (Table 5).




Figure 5. (a) Topography, (b) sea surface temperature and (c,d) ocean circulation patterns for all model runs. (b) Overturning streamfunction. (c) Barotropic streamfunction. Positive values indicate clockwise circulation.

4.2.1 Bulk fluxes and concentrations

The *POC* flux decreases with depth in the model from a maximum of $\sim 10 \text{ mmol C m}^{-2} \text{ d}^{-1}$ immediately below the surface layer, to near-zero values at the seafloor (Fig. 6a-c). The spatial pattern of solid *FeOOH* flux (following the scavenging scheme described earlier – see section 3.1.1) matches the *POC* flux pattern immediately below the surface layer, with higher values at the equator and poles (light brown shading in Fig. 6d). Most of this *FeOOH* is reduced in anoxic subsurface layers and the flux declines with depth. However, at the poles where deep convection allows for greater oxygen (and Fe^{3+}) penetration into the ocean interior (Fig. 7a-h), sinking particles have more time to continue to scavenge and accumulate Fe^{3+} and the flux increases with depth (Fig. 6e,f). The maximum *FeOOH* flux at the seafloor reaches $\sim 300 \mu\text{mol m}^{-2} \text{ d}^{-1}$ (Fig. 6f), comparable to that observed in typical modern ocean margin sediments (see e.g.; van de Velde and Meysman, 2016). Fluxes of *FeS₂* reach their maximum near the seafloor (where a source of deep Fe^{2+} is supplied via the imposed hydrothermal flux), with maximum fluxes similar to those of *FeOOH* (Fig. 6g-i). The spatial pattern of $\text{Fe}_3\text{Si}_2\text{O}_5(\text{OH})_4$ formation is very similar to *FeS₂*, but the fluxes are several orders of magnitude lower (Fig. 6j-l). Fluxes of *FeCO₃* where near-zero everywhere (data not shown), consistent with recent work suggesting that water column precipitation of *FeCO₃* is difficult to achieve, even in iron-dominated oceans (Jiang and Tosca, 2019; Tosca et al., 2019).

We additionally find that even in an idealised ocean with a simple symmetrical continental configuration, complex spatial patterns emerge in the Fe-S redox chemistry (Fig. 7). For instance, the poles are more well-ventilated, allowing oxygenated




Figure 6. Model output of (a)-(c) Particulate organic carbon flux (J_{POC}), (d)-(f) solid iron oxide flux (J_{FeOOH}), (g)-(i) pyrite flux (J_{FeS_2}) and (j)-(l) greenalite flux ($J_{Fe_3Si_2O_5(OH)_4}$) for an ocean with 50 % of the modern phosphate inventory. J_{POC} is given in $mmol\ m^{-2}\ d^{-1}$, J_{FeOOH} and J_{FeS_2} are given in $\mu mol\ m^{-2}\ d^{-1}$ and $J_{Fe_3Si_2O_5(OH)_4}$ is given in $nmol\ m^{-2}\ d^{-1}$. The siderite flux (J_{FeCO_3}) was near-zero everywhere and is not shown here.

conditions and persistence of Fe^{3+} to a few kilometres depth in our benchmark simulation (Fig. 7a-h). The concentrations of Fe^{3+} are much higher than the nano - picomolar concentrations we observe in the ocean today (Tagliabue et al., 2016) – a consequence of much higher rates of iron delivery to the surface ocean (through upwelling of reduced Fe^{2+}), which allows Fe^{3+} to accumulate to higher concentrations before it is eventually scavenged. Our model predicts concentrations in the hundreds of nanomolar range (up to micromolar at the oxic/anoxic interface), which compare well to oxic water layers overlying anoxic deep water (Taillefert and Gaillard, 2002; Crowe et al., 2008). Note that this Fe^{3+} is likely in colloidal or nanoparticulate form and not truly dissolved. At eastward latitudes, deep convection at the poles is less intense, and upwelling on the eastward edge of the ocean leads to higher export production, which subsequently leads to build-up of reduced Fe^{2+} and ΣH_2S (Fig. 7l,p). Dissolved Fe_{total}^{2+} reaches higher concentrations in the deeper ocean, largely as a result of deep hydrothermal inputs (Fig. 7i-l), whereas the highest ΣH_2S concentrations are spatially constrained to areas of more intense POC degradation along the equator, just below the oxic zone (Fig. 7m-p).

Because pervasive anoxia is not present in modern open ocean environments (see above), evaluation of the realism of our globally integrated reaction rates must rely on comparison to modern process analogues for ancient oceans (e.g., anoxic lake and restricted marine systems). Though we consider this a valid approach, it must be borne in mind that the transport




Figure 7. Model output of (a)-(d) dissolved oxygen (O_2), (e)-(h) dissolved ferric iron (Fe^{3+}), (i)-(l) total dissolved ferrous iron (Fe^{2+}) and (m)-(p) total dissolved sulphide (ΣH_2S) for an ocean with 50% of the modern phosphate inventory. All concentrations are in μM .

processes in particular are very different in stratified lacustrine and marine systems, and there are reasons to assume that the biogeochemical dynamics of these systems will not strictly map onto pervasively anoxic open ocean environments. However, though the absolute values are not directly comparable, we can qualitatively compare our results with rates derived from anoxic lake systems. For instance, our default model suggests that DIR is a negligible contributor to POC mineralisation (Table 5), consistent with previous observations from ferruginous Lake Matano and the euxinic Black Sea (Konovalov et al., 2006; Crowe et al., 2011). However, Crowe et al. (2011) found that rates of DIR were roughly an order magnitude lower than those of methanogenesis, whereas we find that DIR is 6-7 orders of magnitude less important than all other mineralisation pathways (Table 5). A possible reason for this discrepancy is the importance of sediment recycling in the natural lake system, which our model does not currently represent. Due to the shallower water column (e.g., $\sim 300m$ in Lake Matano versus $5000m$ in our idealized ocean), there is a much stronger coupling between sedimentary processes (i.e., the recycling of Fe as a benthic flux) and water column processes. Indeed, when the total rate of DIR is corrected for iron recycling occurring in sediments within Lake Matano the importance of DIR decreases by several orders of magnitude (Crowe et al., 2008).

Our model predicts a globally integrated FIO rate of $\sim 0.30 \times 10^{-6} mol Fe kg^{-1} yr^{-1}$ (Table 5), lower than rates estimated for anoxic lakes ($12 - 51 \times 10^{-6} mol Fe kg^{-1} yr^{-1}$; Crowe et al., 2008; Walter et al., 2014). However, rates measured near the oxycline ($6 - 10 \times 10^{-6} mol Fe kg^{-1} yr^{-1}$) are of the same order of magnitude as rates measured in anoxic lake systems. Interestingly, more than half of the ΣH_2S is re-oxidised via SMI (Table 5), which indicates that Fe is able to act as a relatively



Table 5. Globally integrated reaction rates and burial fluxes for our benchmark simulation.

Reaction number	Reaction name	Units	Rate
R1	Aerobic respiration	$mol\ POC\ kg^{-1}\ yr^{-1}$	0.80×10^{-7}
R2	Dissimilatory iron reduction	$mol\ POC\ kg^{-1}\ yr^{-1}$	0.48×10^{-13}
R3	Dissimilatory sulphate reduction	$mol\ POC\ kg^{-1}\ yr^{-1}$	0.12×10^{-6}
R4	Methanogenesis	$mol\ POC\ kg^{-1}\ yr^{-1}$	0.17×10^{-6}
R5	Ferrous iron oxidation	$mol\ Fe\ kg^{-1}\ yr^{-1}$	0.30×10^{-6}
R6	Canonical sulphide oxidation	$mol\ S\ kg^{-1}\ yr^{-1}$	0.67×10^{-7}
R7a	Sulphide-mediated iron reduction (dissolved)	$mol\ S\ kg^{-1}\ yr^{-1}$	0.37×10^{-7}
R7b	Sulphide-mediated iron reduction (solid)	$mol\ S\ kg^{-1}\ yr^{-1}$	0.83×10^{-7}
R8	Iron oxide precipitation	$mol\ FeOOH\ kg^{-1}\ yr^{-1}$	0.67×10^{-6}
R9	Pyrite precipitation	$mol\ FeS_2\ kg^{-1}\ yr^{-1}$	0.43×10^{-8}
R10	Siderite precipitation	$mol\ FeCO_3\ kg^{-1}\ yr^{-1}$	0.79×10^{-63}
R11	Greenalite precipitation	$mol\ Fe_3Si_2O_5(OH)_4\ kg^{-1}\ yr^{-1}$	0.13×10^{-13}
-	Iron oxide burial	$mol\ FeOOH\ m^{-2}\ yr^{-1}$	0.12×10^{-1}
-	Pyrite burial	$mol\ FeS_2\ m^{-2}\ yr^{-1}$	0.22×10^{-1}
-	Siderite burial	$mol\ FeCO_3\ m^{-2}\ yr^{-1}$	0.41×10^{-56}
-	Greenalite burial	$mol\ Fe_3Si_2O_5(OH)_4\ m^{-2}\ yr^{-1}$	0.48×10^{-7}

efficient intermediate between ΣH_2S oxidation and O_2 reduction. This also occurs in the Black Sea, where metal oxides are responsible for $\sim 60\%$ of the sulphide re-oxidation (Konovalov et al., 2006).

4.2.2 Isotope patterns

Figure 8a-e shows the modelled stable Fe isotope patterns for all key Fe-bearing dissolved and solid-phase species (with the exception of $FeCO_3$, which is a negligible component in our benchmark simulation). In our model simulations, all dissolved Fe that enters the ocean is assigned an isotope signature of 0.0 ‰. This allows us to observe the isotope fractionation of all Fe phases relative to the Fe that entered the ocean (Fig. 9a). The dissolved iron phases show similar isotopic signatures ($\sim 1.1\%$; Fig. 8a,b; Fig. 9a), which can be explained by the large amount of isotopically light FeS_2 burial (Fig. 9a), which drives $\delta^{56}Fe - Fe^{2+}$ to heavier values. The isotope signature of buried oxidised iron ($FeOOH$) is around 1.8 ‰ heavier than the Fe that entered the ocean, whereas the major burial fraction of reduced iron (FeS_2) has an isotope signature that is $\sim 1\%$ lighter, which reflects their relative importance as an Fe burial phase (Fig. 8c,e; Fig. 9a). These isotope values are broadly comparable to phase-specific stable Fe isotope observations from ancient sedimentary rocks (Heard and Dauphas, 2020). Similarly, Fig. 8f-g shows the modelled stable S isotope patterns for all key S-bearing dissolved and solid-phase species. All dissolved S that enters the ocean is assigned an isotope signature of 0.0 ‰. Sulphate is isotopically enriched relative to its input value (~ 3.3




Figure 8. (a-e) Modelled stable Fe isotope signatures of key dissolved and solid-phase Fe species. Shown are zonally averaged values for (a) dissolved ferric Fe ($\delta^{56}Fe - Fe^{3+}$) and (b) dissolved ferrous iron ($\delta^{56}Fe - Fe^{2+}$), and the isotope compositions at the seafloor of (c) iron oxides ($\delta^{56}Fe - FeOOH$), (d) greenalite ($\delta^{56}Fe - Fe_3Si_2O_5(OH)_4$) and (e) pyrite ($\delta^{56}Fe - FeS_2$) for an ocean with 50 % of the modern phosphate inventory. All values are in ‰, relative to IRMM-14. (f-g) Modeled stable S isotope signatures of key dissolved and solid-phase S species. Shown are zonally averaged values for (f) sulphate ($\delta^{34}S - SO_4^{2-}$) and (g) dissolved free sulphide ($\delta^{34}S - H_2S$), and (h) the isotope composition at the seafloor of pyrite ($\delta^{34}S - FeS_2$). All values are in ‰, relative to VCDT.

‰; Fig. 8f; Fig. 9b), and this difference compares well to the geological record (SO_4^{2-} is ~ 5 ‰ heavier than its input value; Canfield and Farquhar, 2009). Additionally, free sulphide is isotopically lighter than sulphate (~ -5 ‰ on a global scale; up to ~ -15 ‰ locally), while buried FeS_2 expresses an isotope signature of ~ -3.25 ‰ (Fig. 9b). Our baseline simulation thus suggests that the expressed isotope fractionation between SO_4^{2-} and FeS_2 is only around 6.5 ‰, which is roughly consistent with what is observed in the geological record (Canfield and Farquhar, 2009). Therefore, we conclude that our model has strong potential for tracking iron and sulphur isotope signatures for comparison with Earth's rock record.

4.3 Sensitivity analysis

We evaluate model output sensitivity to four key parameters of our modelled iron-sulphur cycle ($K_{0,FeOOH}$, $K_{i,FeOOH}$, $k_{SMI,s}$, k_{PyP}) using the Elementary Effect Test (EET; Morris, 1991). These parameters were chosen because they are either unconstrained by laboratory experiments ($K_{0,FeOOH}$, $K_{i,FeOOH}$; Section 3.1.2), represent a complex mixture of different iron minerals with different reactivities ($k_{SMI,s}$; Poulton et al., 2004a) or are expected to have a strong and potentially difficult to forecast influence on other reactions (k_{PyP} ; van de Velde et al., in review). Other parameters introduced in the model are either relatively well-constrained by laboratory studies and calibrated on field data (k_{CSO} , k_{FIO} ; Millero et al., 1987a, b; Ridgwell




Figure 9. Isotope mass balance for the (a) ^{56}Fe and (b) ^{34}S systems from our baseline simulation. Purple arrows are influxes, black arrows outfluxes. Bubbles represent the inventory of the whole ocean. Note that in our simulation, the S system is not in steady-state (because FeS_2 burial is too low).

et al., 2007), have been calibrated extensively in previous work (k_{scav} ; Tagliabue et al., 2016) or are likely to be of secondary importance to the iron-sulphur cycle (k_{AFC} , b_{AFC} , $k_{greenalite}$, $b_{greenalite}$).

The EET method estimates global sensitivity by calculating the mean of r finite differences (‘Elementary Effects’) (Pianosi et al., 2016),

$$S_i = \frac{1}{r} \sum_{j=1}^r EE^j$$

$$= \frac{1}{r} \sum_{j=1}^r \frac{f(\bar{x}_1^j, \dots, \bar{x}_i^j + \Delta_i^j, \dots, \bar{x}_M^j) - f(\bar{x}_1^j, \dots, \bar{x}_i^j, \dots, \bar{x}_M^j)}{\Delta_i^j} c_i \quad (38)$$

where \bar{x}_i^j represents the j^{th} value of the i^{th} parameter, Δ_i^j the variation on the i^{th} parameter, $f()$ is the model output for a given set of parameters and c_i is a scaling factor. A higher mean value S_i indicates that a given model output is more sensitive to variations in parameter i . The standard deviation can also be calculated, with a high standard deviation indicating that a parameter interacts with others because its sensitivity changes across the variability space (inlay in Fig. 10; Pianosi et al., 2016).

We use the EET method, as implemented within the Sensitivity Analysis For Everyone (SAFE) toolbox (Pianosi et al., 2015), to investigate the four chosen model parameters across the ranges specified in Table 6. We vary the limitation and inhibition constants $K_{0,FeOOH}$ and $K_{i,FeOOH}$ over 6 orders of magnitude around the default/baseline value due to the high uncertainty associated with these parameters. The lower-bound of the reaction constant of solid iron oxide with sulphide ($k_{SMI,s}$) is defined as the reactivity of hematite ($5.34 \cdot 10^{-3} M^{-0.5} h^{-1}$; Poulton et al., 2004a), and the upper bound is taken to be 5 orders of magnitude higher ($5.34 \cdot 10^2 M^{-0.5} h^{-1}$, which is comparable to freshly precipitated hydrous ferric oxide).



Table 6. Sensitivity range and baseline values of each parameter tested.

Constant	Symbol	Unit	Baseline	Minimum	Maximum
Limitation constant DIR	$K_{0,FeOOH}$	M	1.0×10^{-2}	1.0×10^{-5}	1.0×10^1
Inhibition constant DIR	$K_{i,FeOOH}$	M	1.0×10^{-2}	1.0×10^{-5}	1.0×10^1
Sulphide-mediated iron reduction (solid)	$k_{SMI,s}$	$M^{-0.5}h^{-1}$	1.98×10^0	5.34×10^{-3}	5.34×10^2
Pyrite precipitation	k_{PyP}	$M^{-1}h^{-1}$	0.3708×10^0	0.3708×10^{-3}	0.3708×10^2

We test the pyrite precipitation constant (k_{PyP}) across a range between $0.3708 \times 10^{-3} M^{-1}h^{-1}$ and $0.3708 \times 10^2 M^{-1}h^{-1}$, which corresponds to the range of kinetic constants commonly used in diagenetic models (see e.g.; Van Cappellen and Wang, 1996; Meysman et al., 2003; Dale et al., 2009; van de Velde et al., 2020). Our sensitivity ensemble consists of one hundred individual model experiments, using Latin-Hypercube sampling approach (using the SAFE toolbox; Pianosi et al., 2015) to select random starting points x^j ($j = 1, \dots, r$) and parameter variations Δ_i . For more information on the sampling strategy we refer the interested reader to Campolongo et al. (2011).


The EET analysis suggests that changes in $K_{0,FeOOH}$ can significantly impact DIR, whereas the inhibition constant $K_{i,FeOOH}$ has a relatively minor impact on model output (Fig. 10). This is expected, as both parameters act only on the mineralisation pathways (Eq. 10). It is also consistent with some of the literature on anoxic systems, which suggests that in many cases the majority of iron reduction in the water column is coupled to sulphide (Mikucki et al., 2009). Even though both $K_{0,FeOOH}$ and $K_{i,FeOOH}$ are the least constrained by experimental results (see section 3.1.2), the EET analysis indicates that they are relatively unimportant for the overall model output, despite their impact on DIR. In contrast, k_{PyP} and $k_{SMI,s}$ both exerted more notable impact on model output across a range of diagnostics (Fig. 10). In particular, k_{PyP} had a significant impact across all model output diagnostics analysed here both in terms of model sensitivity and interactivity with other parameters (Fig. 10). Because pyrite precipitation controls the inventories of both dissolved Fe^{2+} and dissolved ΣH_2S , reducing or increasing the kinetic precipitation parameter will affect the ambient concentrations, re-oxidation pathways, and eventual mineral burial for all phases across the Fe-S system.

Unfortunately, an Elementary Effect Test gives no quantitative metric to evaluate the magnitude with which a parameter affects overall model outcome. Therefore, to illustrate the quantitative impact of the possible parameter choices, we ran a separate set of experiments where we changed a parameter from its baseline value to its lower and upper bound, whilst keeping the other 3 parameters at their baseline values (Table 6).


Figure 11 reveals that AR, DSR, mean surface O_2 concentrations and CSO are relatively insensitive to parameter choices (Fig. 11a,c,d,i). Dissimilatory iron reduction is only important when $K_{0,FeOOH}$ is set at its lowest value, and even then it is three orders of magnitude smaller than AR or SR (Fig. 11a-c). Consistent with the EET analysis (Fig. 10), $K_{0,FeOOH}$ and $K_{i,FeOOH}$ have no influence on the model output, aside from the magnitude of DIR, which in itself is of less importance as mineralisation pathway (Fig. 11b and Section 4.2.1). In contrast, changes in the pyrite precipitation constant k_{PyP} impact




Mineralisation pathways




Mean ocean concentrations



Reoxidation pathways



Mineral burial



■ $K_{SMI,s}$ ● K_{PyP} ▲ $K_{i,FeOOH}$ ◆ $K_{0,FeOOH}$

Figure 10. Sensitivity analysis of the four key-parameters of the iron-sulphur cycle, for a range of different outputs. Inset shows how to read the graph; points that plot more to the right indicate that the specific output is more sensitive to changes in parameter values, points that plot higher indicate that the parameter is more interactive with other parameters. Data processing was done with the SAFER toolbox of Pianosi et al. (2015).

several model outputs. When pyrite precipitation rates are elevated, Fe^{2+} and ΣH_2S are removed from local seawater more rapidly, which results in:

1. a decreased build-up of Fe^{2+} and ΣH_2S in the ocean interior (Fig. 11f,g), leading to




Figure 11. Global model output for the three different values of the four key parameters ($K_{0,FeOOH}$, $K_{i,FeOOH}$, k_{PyP} and $k_{SMI,s}$). Values for low, baseline and high values are given in Table 6. Panels (a)-(c) and (h)-(n) are in $mol\ yr^{-1}$, panels (d)-(g) are in $mol\ kg^{-1}$.

2. a decrease in both aerobic and anaerobic re-oxidation pathways (FIO, SMI_d and SMI_s ; Fig. 11h-k), which then results in
3. more O_2 available for aerobic respiration at the expense of less thermodynamically favourable electron acceptors (i.e. AR increases and DSR decreases; Fig. 11a,c) and
- 5 4. a decrease in Fe^{3+} concentrations in the surface ocean (Fig. 11e), and
5. less burial of $FeOOH$, increasing the burial of reduced iron minerals (Fig. 11l-n)

The effect on greenaltite burial is non-linear (Fig. 11o), which indicates that at higher values of k_{PyP} , pyrite precipitation is competing with greenaltite precipitation. Overall, our sensitivity analysis suggests that k_{PyP} is an important parameter for the model output, and should be chosen with care. Fortunately, pyrite precipitation has been well studied in laboratory experiments

10 (Rickard, 1997, 2006), with the result that our baseline value for this parameter is relatively well constrained.



The fourth parameter, which also influences the model output, is the reactivity parameter of solid iron oxides ($k_{SMI,s}$). Here, the parameter choice is more complex. Laboratory experiments have shown that different iron oxide minerals exhibit a wide reactivity range (spanning several orders of magnitude) (Canfield, 1992; Poulton et al., 2004a). Therefore, we explore the sensitivity of this parameter in more detail using a range of measured reactivity constants by Poulton et al. (2004a), whilst
5 keeping all other parameters at their baseline values (Fig. 12). Increasing the reactivity of particulate $FeOOH$

1. increases the anaerobic re-oxidation reaction of sulphide with $FeOOH$ (SMI_s) at the expense of the aerobic re-oxidation reaction (CSO) (Fig. 12i,j), which then leads to
2. an increase in the Fe_{total}^{2+} inventory and a decrease in the ΣH_2S inventory (Fig. 12f,g), more Fe_{total}^{2+} leads to
3. more FIO, and thus a higher surface Fe^{3+} concentration, and more re-oxidation of ΣH_2S with dissolved Fe^{3+} (SMI_d ;
10 Fig. 12e,h,k)
4. Because of the higher reactivity of the $FeOOH$ particles, less $FeOOH$ is buried, and more reduced Fe -minerals are buried (Fig. 12l-o)

Although it is clear that changing $k_{SMI,s}$ impacts model output, the overall magnitude of the effect is moderate when compared to changing k_{PyP} . Nevertheless, the choice of $k_{SMI,s}$ is critical. We choose a baseline reactivity value (Table 3)
15 comparable to lepidocrocite for several reasons. Firstly, we assume all Fe^{3+} that is not scavenged represents a 'colloidal' pool, with a reactivity of similar to that of hydrous ferric oxide. When Fe^{3+} becomes scavenged (and is thus in solid state), it has likely undergone some ageing, and it will be less reactive than hydrous ferric oxide. Secondly, any $FeOOH$ that does not react in the water column will end up in the sediment, and will, at least in part, be recycled back to the water column (even under oxic conditions; Dale et al., 2015). Our model currently lacks a sedimentary iron cycle (see section 5), and would thus
20 tend to underestimate the overall importance of the iron cycle were we to select a reactivity constant that is too low. Finally, field evidence suggests that $FeOOH$ that is freshly precipitated is highly reactive (Picard et al., 2015; Beam et al., 2018) and thus iron precipitating from the surface ocean is expected to react on relatively short timescales. Any $FeOOH$ minerals that would resist reduction passing through a sulphidic water column are likely unreactive, and are thus presumably inert on early diagenetic timescales. Indeed, iron oxide minerals in sediments underlying euxinic water columns tend to show no depth trend,
25 indicating that very little iron reduction is occurring at depth in such systems (see e.g., Xiong et al., 2019). Taken together, these observations support the presumption that once Fe^{3+} is scavenged, its reactivity is less than hydrous ferric oxide but higher than that of goethite, motivating our default $k_{SMI,s}$ value. Nevertheless, the value used for this parameter should be considered carefully depending on the model application and assumed boundary conditions.

5 Outlook and conclusions

30 The principal aim of this paper is to provide a detailed description of our extension of the cGENIE biogeochemistry module to include coupled, anoxic Fe and S biogeochemical cycles. Because direct tuning with actual measured concentrations and




Figure 12. Global model output for a range of $k_{SMI,s}$ values (units are $M^{-1/2} h^{-1}$). Lines indicate experimental reactivity parameters as presented in (Poulton et al., 2004a). Panels (a)-(c) and (h)-(n) are in $mol yr^{-1}$, panels (d)-(g) are in $mol kg^{-1}$. Note that the x-axis is logarithmic.

rates is not possible, we have relied heavily on kinetic constants and solubility values extracted from laboratory incubations. While care should be exercised in the application of these kinetic constants to reactions under *in-situ* conditions, our sensitivity analysis indicates that our key model results are robust across a wide range of possible parameter values. In addition, our proposed baseline parametrisation yields reaction rates, concentrations, burial fluxes, and stable Fe isotope compositions that

5 broadly compare well to both field measurements of process analogues for ancient ocean systems (i.e., anoxic lakes) and observations from the geologic record. Therefore, we believe that our model description of the anoxic Fe-S cycle is a valuable tool and an important step forward in simulating ocean redox landscapes during periods of Earth's history in which the ocean interior was pervasively anoxic. However, below we highlight some important limitations to our current model architecture, and also give some examples of possible future developments.

10 Most notably, our model is currently unable to resolve any sedimentary processes that would contribute to the global iron cycle (see e.g., Dale et al., 2015). In particular, the model does not include a representation of benthic iron reduction and



recycling back into the water column. Building on the improvements in the biogeochemistry of cGENIE described here, in the future we plan to extend the organic matter enabled sediment component of cGENIE (OMEN-SED; Hülse et al., 2018) to include an explicit representation of the benthic iron cycle. We anticipate that this will both improve the realism of tracer fields within the ocean interior and will make comparisons between predicted sedimentary signals and observations from Earth's sedimentary rock record more accurate and robust. Second, there are likely to be important mechanistic links between the biogeochemistry of Fe and S within the ocean and the local and global recycling and bioavailability of key nutrient species for the biosphere (Bjerrum and Canfield, 2002; Laakso and Schrag, 2014; Jones et al., 2015; Reinhard et al., 2017). Future work will thus also focus on explicitly coupling the anoxic Fe and S biogeochemistry to the phosphorus (P) and nitrogen (N) cycles, and in particular the scavenging and remobilisation of P under different redox states and the impact of dissolved Fe availability on nitrogen fixation. The modularity of cGENIE also allows the substitution of an explicit plankton ecological model ('ECOGEM') for the default biological export scheme (Ward et al., 2018), enabling the exploration of feedbacks between marine ecosystems, nutrient availability and ocean redox conditions (Reinhard et al., 2020b). Lastly, future work will seek to include other redox-sensitive proxies and bioessential elements, such as molybdenum, uranium or vanadium (Tribovillard, 2006) within the model code, which will further extend the applicability of our model and help to validate it against observations from modern anoxic systems and the geologic record.

Code availability. The specific version used of the cGENIE.muffin model used in this paper is tagged as release v0.9.16, and has been assigned a DOI:10.5281/zenodo.4033262 The code is hosted on GitHub and can be obtained by cloning:

```
https://github.com/derpycode/cgenie.muffin
```

changing the directory to `cgenie.muffin` and then checking out the specific release:

```
20 $ git checkout v0.9.16
```

Configuration files for the specific experiments presented in the paper can be found in the directory:

```
genie-userconfigs/MS/vandeveldeetal.GMD.2020
```

Details of the experiments, plus the command line needed to run each one, are given in the `readme.txt` file in that directory. All other configuration files and boundary conditions are provided as part of the release. A manual, detailing code installation, basic model configuration, plus an extensive series of tutorials covering various aspects of muffin capability, experimental design, and results output and processing, is provided on GitHub. The latex source of the manual, along with a pre-built PDF format version, can be obtained by cloning:

```
25 https://github.com/derpycode/muffindoc
```

Author contributions. SJV and AR developed the model with input from all authors. CR provided geological context. SJV and DH ran the sensitivity tests and analysed the output. SJV wrote the paper with input from all authors.

30 *Competing interests.* The authors declare that they have no conflict of interest.



Acknowledgements. SVDV was supported by a post-doctoral Fellowship of the Belgian American Educational Foundation (2018-2019) and a NASA Postdoctoral Program fellowship at the University of California, Riverside, administered by Universities Space Research Association under contract with NASA (2019-2020). DH is supported by the Simons Foundation (Postdoctoral Fellowship in Marine Microbial Ecology, Award 653829). CR gratefully acknowledges funding from the NASA Exobiology Program and the NASA Nexus for Exoplanet System Science (NExSS).



References

- Adloff, M., Ridgwell, A., Monteiro, F. M., Parkinson, I. J., Dickson, A., Pogge von Strandmann, P. A. E., Fantle, M. S. and Greene, S. E.: Inclusion of a suite of weathering tracers in the cGENIE Earth System Model - muffin release v.0.9.10, *Geoscientific Model Development Discussions*, 1-48, <https://doi.org/10.5194/gmd-2020-233>, 2020.
- 5 Alcott, L. J., Mills, B. J. W. and Poulton, S. W.: Stepwise Earth oxygenation is an inherent property of global biogeochemical cycling, *Science*, 366, 1333-1337, <https://doi.org/10.1126/science.aax6459>, 2019.
- Balistrieri, L. S., Murray, J. W., and Paul, B.: The cycling of iron and manganese in the water column of Lake Sammamish, Washington, *Limnol. Oceanogr.*, 37, 510-528, 1992.
- Balzano, S., Statham, P. J., Pancost, R. D. and Lloyd, J.R.: Role of microbial populations in the release of reduced iron to the water column
10 from marine aggregates, *Aquat. Microb. Ecol.*, 54, 291-303, <https://doi.org/10.1111/10.3354/ame01278>, 2009.
- Beam, J. P., Scott, J. J., McAllister, S. M., Chan, C. S., McManus, J., Meysman, F. J. R. and Emerson, D.: Biological rejuvenation of iron oxides in bioturbated marine sediments, *The ISME Journal*, 12, 1389–1394, <https://doi.org/10.1038/s41396-017-0032-6>, 2018.
- Beard, B. L., Johnson, C. M., Cox, L., Sun, H., Nealson, K. H. and Aguilar, C.: Iron isotope biosignatures, *Science*, 285, 1889-1892, 1999.
- Beard, B. L., Johnson, C. M., Skulan, J. L., Nealson, K. H., Cox, L. and Sun, H.: Application of Fe isotopes to tracing the geochemical and
15 biological cycling of Fe, *Chem. Geol.*, 195, 87-117, 2003.
- Bekker, A., Slack, J. F., Planavsky, N., Krapez, B., Hofmann, A., Konhauser, K. O. and Rouxel, O. J.: Iron Formation: The sedimentary product of a complex interplay among mantle, tectonic, oceanic, and biospheric processes, *Economic Geology*, 105, 467-508, 2010.
- Berg, J. S., Michellod, D., Pjevac, P., Martinez-Perez, C., Buckner, C. R. T., Hach, P. F., Schubert, C. J., Milucka, J. and Kuypers, M. M. M.:
20 Intensive cryptic microbial iron cycling in the low iron water column of the meromictic Lake Cadagno, *Environmental Microbiology*, 18, 5288-5302, <https://doi.org/10.1111/1462-2920.13587>, 2016.
- Berner, R. A.: Phosphate removal from sea water by adsorption on volcanogenic ferric oxides, *Earth Planet. Sci. Lett.*, 18, 77–86, 1973.
- Berner, R. A.: Biogeochemical cycles of carbon and sulfur and their effect on atmospheric oxygen over phanerozoic time. *Paleogeography, Paleoclimatology, Paleoecol.*, 75, 97–122, 1989.
- Berner, R. A.: A model for calcium, magnesium and sulfate in seawater over Phanerozoic time, *American Journal of Science*, 304, 438-453,
25 <https://doi.org/10.2475/ajs.304.5.438>, 2004.
- Bjerrum C. J. and Canfield D. E.: Ocean productivity before about 1.9 Gyr ago limited by phosphorus adsorption onto iron oxides, *Nature*, 417, 159–162, <https://doi.org/10.2475/10.1038/417159a>, 2002.
- Böttcher, M. E., Smock, A. M. and Cypionka, H.: Sulfur isotope fractionation during experimental precipitation of iron(II) and manganese(II) sulfide at room temperature, *Chem. Geol.*, 146, 127–134, 1998.
- 30 Brocks J. J., Jarrett A. J. M., Sirantoine E., Hallmann C., Hoshino Y. and Liyanage T.: The rise of algae in Cryogenian oceans and the emergence of animals, *Nature*, 548, 578–581, <http://dx.doi.org/10.1038/nature23457>, 2017.
- Bullen, T. D., White, A. F., Childs, C. W., Vivit, D. V. and Schulz, M. S.: Demonstration of significant abiotic iron isotope fractionation in nature, *Geology*, 29(8), 699-702, 2001.
- Campolongo, F., Saltelli, A. and Cariboni, J.: From screening to quantitative sensitivity analysis. A unified approach, *Computer Physics Communications*, 182, 978–988, <http://dx.doi.org/10.1016/j.cpc.2010.12.039>, 2011.
- 35 Canfield, D.E., Raiswell, R. and Bottrell, S.: The reactivity of sedimentary iron minerals toward sulfide, *American Journal of Science*, 292, 659-683, 1992.



- Canfield, D. E. and Teske, A.: Late proterozoic rise in atmospheric oxygen concentration inferred from phylogenetic and sulphur-isotope studies, *Nature*, 382, 127–132, <http://dx.doi.org/10.1038/382127a0>, 1996.
- Canfield, D.E.: A new model for Proterozoic ocean chemistry, *Nature*, 396, 450-453, 1998.
- Canfield, D.E., Rosing, M. T. and Bjerrum, C.: Early anaerobic metabolisms, *Phil. Trans. R. Soc. B*, 361, 1819-1836, 2006.
- 5 Canfield, D.E., Poulton, S. W., Knoll, A. H., Narbonne, G. M., Ross, G., Goldberg, T. and Strauss, H.: Ferruginous Conditions Dominated Later Neoproterozoic Deep-Water Chemistry, *Science*, 321, 949-952, 2008.
- Canfield D. E. and Farquhar J.: Animal evolution, bioturbation, and the sulfate concentration of the oceans, *Proc. Natl. Acad. Sci.*, 106, 8123–8127, 2009.
- Cao, L., Eby, M., Ridgwell, A., Caldeira, K., Archer, D., Ishida, A., Joos, F., Matsumoto, K., Mikolajewicz, U., Mouchet, A., Orr, J. C., Plattner, G.-K., Schlitzer, R., Tokos, K., Totterdell, I., Tschumi, T., Yamanaka, Y., and Yool, A.: The role of ocean transport in the uptake of anthropogenic CO₂, *Biogeosciences*, 6, 375–390, <https://doi.org/10.5194/bg-6-375-2009>, 2009.
- Cloud, P.: A working model of the primitive Earth, *Am. J. Sci.*, 272, 537–548, <https://doi.org/10.2475/ajs.272.6.537>, 1972.
- Colbourn, G., Ridgwell, A. and Lenton, T. M. The Rock Geochemical Model (RokGeM) v0.9, *Geoscientific Model Development*, 6(5), 1543-1573, <https://doi.org/10.5194/gmd-6-1543-2013>, 2013.
- 15 Conway, T. M. and John, S. G. Quantification of dissolved iron sources to the North Atlantic Ocean, *Nature*, 511(7508), 212–215, <https://doi.org/10.1038/nature13482>, 2014.
- Cosmidis, J., Benzerara, K., Morin, G., Busigny, V., Lebeau, O., Jézéquel, D., Noël, V., Dublet, G. and Othmane, G.: Biomineralization of iron-phosphates in the water column of Lake Pavin (Massif Central, France), *Geochim. Cosmochim. Acta*, 126, 78-96, <http://dx.doi.org/10.1016/j.gca.2013.10.037>, 2014.
- 20 Crichton, K. A., Wilson, J. D., Ridgwell, A. and Pearson, P. N.: Calibration of key temperature-dependent ocean microbial processes in the cGENIE.muffin Earth system model, *Geoscientific Model Development Discussions*, 1-26, <http://dx.doi.org/10.5194/gmd-2019-344>, 2020.
- Crockford, P. W., Hayles, J. A., Bao, H., Planavsky, N. J., Bekker, A., Fralick, P. W., Halverson, G. P., Bui, T. H., Peng, Y. and Wing, B. A.: Triple oxygen isotope evidence for limited mid-Proterozoic primary productivity, *Nature*, 559, 613–616, <https://doi.org/10.1038/s41586-018-0349-y>, 2018.
- 25 Crosby, S. A., Glasson, D. R., Cuttler, A. H., Butler, I., Turner, D. R., Whitfield, M. and Millward, G. E.: Surface areas and porosities of Fe(III)- and Fe(II)-derived oxyhydroxides, *Environ. Sci. Technol.*, 17, 709-713, 1983.
- Crosby, S. A., Roden, E. E., Johnson, C. M. and Beard, B. L.: The mechanisms of iron isotope fractionation produced during dissimilatory Fe(III) reduction by *Shewanella putrefaciens* and *Geobacter sulfurreducens*, *Geobiology*, 5, 169-189, 2007.
- 30 Crowe, S. A., O'Neill, A. H., Katsev, S., Hehanussa, P., Haffner, G. D., Sundby, B., Mucci, A. and Fowle, D. A.: The biogeochemistry of tropical lakes: A case study from Lake Matano, Indonesia, *Limnol. Oceanogr.*, 53, 319-331, 2008.
- Crowe, S. A., , Katsev, S., Leslie, K., Sturm, A., Magen, C., Nomosatryo, S., Pack, M. A., Kessler, J. D., Reeburgh, W. S., Roberts, J. A., González, L., Douglas Haffner, G., Mucci, A., Sundby, B. and Fowle, D. A.: The methane cycle in ferruginous Lake Matano, *Geobiology*, 9, 61-78, <http://dx.doi.org/10.1111/j.1472-4669.2010.00257.x>, 2011.
- 35 Crowe, S. A., Jones, C., Canfield, D. E., Paris, G., Adkins, J. F., Sessions, A. L., Katsev, S., Kim, S.-T., Zerkle, A. L., Nomosatryo, S., Fowle, D. A. and Farquhar, J.: Sulfate was a trace constituent of Archean seawater, *Science*, 346(6210), 735–739, <https://doi.org/10.1126/science.1258966>, 2014.



- Dale, A. W., Brüchert, V., Alperin, M. and Regnier, P.: An integrated sulfur isotope model for Namibian shelf sediments, *Geochim. Cosmochim. Acta*, 73, 1924–1944, <http://dx.doi.org/10.1016/j.gca.2008.12.015>, 2009.
- Dale, A. W., Nickelsen, L., Scholz, F., Hensen, C., Oschlies, A. and Wallmann, K.: A revised global estimate of dissolved iron fluxes from marine sediments. *Global Biogeochem. Cycles*, 29, 1–17 (2015).
- 5 Dehairs, F., Baeyens, W. and Goeyens, L.: Accumulation of Suspended Barite at Mesopelagic Depths and Export Production in the Southern Ocean, *Science*, 20, 1332–1335, <http://dx.doi.org/10.1126/science.258.5086.1332>, 1990.
- Detmers, J., Bruchert, V., Habicht, K. S., and Kuever, J.: Diversity of sulfur isotope fractionations by sulfate-reducing prokaryotes, *Appl. Environ. Microbiol.*, 67(2), 888–894, 2001.
- Diaz, R. J., and Rosenberg, R.: Spreading dead zones and consequences for marine ecosystems, *Science* 321, 926–929, 2008.
- 10 Doney, S. C., Lindsay, K., Fung, I., and John, J.: Natural variability in a stable, 1000-yr global coupled climate-carbon cycle simulation, *Journal of Climate*, 19, 3033–3054, 2006.
- Edwards, N. R. and Shepherd, J. G.: Bifurcations of the thermohaline circulation in a simplified three-dimensional model of the world ocean and the effects of interbasin connectivity. *Clim. Dyn.* 19, 31–42, 2002.
- Edwards, N., and Marsh, R.: Uncertainties due to transport-parameter sensitivity in an efficient 3-D ocean-climate model, *Clim. Dynam.*, 24, 415–433, 2005.
- 15 Fakhraee, M., Hancisse, O., Canfield, D. E., Crowe, S. A. and Katsev, S.: Proterozoic seawater sulfate scarcity and the evolution of ocean–atmosphere chemistry, *Nat. Geosci.*, 12, 375–380, <http://dx.doi.org/10.1038/s41561-019-0351-5>, 2019.
- Ferreira, D., Marshall, J. and Campin, J.: Localization of Deep Water Formation: Role of Atmospheric Moisture Transport and Geometrical Constraints on Ocean Circulation. *J. Climate*, 23, 1456–1476, <https://doi.org/10.1175/2009JCLI3197.1>, 2010
- 20 Froelich, P. N., Klinkhammer, G. P., Bender, M.L., Luedtke, N. A., Heath, G. R., Cullen, D. and Dauphin, P.: Early oxidation of organic matter in pelagic sediments of the eastern equatorial Atlantic: suboxic diagenesis. *Geochim. Cosmochim. Acta*, 43, 1075–1090, 1979.
- Fry, B., Gest, H. and Hayes, J. M.: 34S/32S fractionation in sulfur cycles catalyzed by anaerobic bacteria. *Appl. Environ. Microbiol.*, 54(1), 250–256, 1988.
- Gomes, M. L. and Johnston, D. T.: Oxygen and sulfur isotopes in sulfate in modern euxinic systems with implications for evaluating the extent of euxinia in ancient oceans, *Geochim. Cosmochim. Acta*, 205, 331–359, 2017.
- 25 Gorby, Y. A., Yanina, S., McLean, J. S., Rosso, K. M., Moyles, D., Dohnalkova, A., Beveridge, T. J., Chang, I. S., Kim, B. H., Kim, K. S., Culley, D. E., Reed, S. B., Romine, M. F., Saffarini, D. A., Hill, E. A., Shi, L., Elias, D. A., Kennedy, D. W., Pinchuk, G., Watanabe, K., Ishii, S., Logan, B., Neelson, K. H. and Fredrickson, J. K.: Electrically conductive bacterial nanowires produced by *Shewanella oneidensis* strain MR-1 and other microorganisms, *PNAS*, 103, 11358–11363, <https://doi.org/10.1073/pnas.0604517103>, 2006.
- 30 Grotzinger J. P. and Kasting J. F.: New Constraints on Precambrian Ocean composition, *J. Geol.*, 101, 235–243, 1993.
- Guilbaud, R., Butler, I. B. and Ellam, R. M.: Abiotic pyrite formation produces a large Fe isotope fractionation, *Science*, 332, 1548–1551, 2011.
- Guilbaud, R., Poulton, S. W., Butterfield, N. J., Zhu M. and Shields-Zhou, G. A.: A global transition to ferruginous conditions in the early Neoproterozoic oceans, *Nat. Geosc.*, 8, 466–470, <https://doi.org/10.1038/NGEO2434>, 2015.
- 35 Guilbaud, R., Poulton, S. W., Thompson, J., Husband, K. F., Zhu, M., Zhou, Y., Shields, G. A. and Lenton, T. M.: Phosphorus-limited conditions in the early Neoproterozoic ocean maintained low levels of atmospheric oxygen, *Nat. Geosc.*, 13, 296–301, 2020.
- Gustafsson, J.P.: Visual MINTeq, version 3.1. KTH Royal Institute of Technology, Stockholm, Sweden, 2019 available at: <https://vminteq.lwr.kth.se/>



- Hargreaves, J. C., Annan, J. D., Edwards, N. R., and Marsh, R.: An efficient climate forecasting method using an intermediate complexity Earth System Model and the ensemble Kalman filter, *Clim. Dynam.*, 23, 745–760, <https://doi.org/10.1007/s00382-004-0471-4>, 2004.
- Heard, A. W. and Dauphas, N.: Constraints on the coevolution of oxic and sulfidic ocean iron sinks from Archean–Paleoproterozoic iron isotope records. *Geology*, 48, 358–362, 2020.
- 5 Henson, S. A., Sanders, R., Madsen, E., Morris, P. J., Le Moigne, F. and Quartly, G. D.: A reduced estimate of the strength of the ocean’s biological carbon pump, *Geophys. Res. Lett.*, 38, L04606, <https://doi.org/10.1029/2011GL046735>, 2011.
- Holland, H. D.: *The Chemical Evolution of the Atmosphere and Oceans*, (Princeton Univ Press, Princeton), p 582, 1984.
- Hülse, D., Arndt, S., Wilson, J. D., Munhoven, G. and Ridgwell, A.: Understanding the causes and consequences of past marine carbon cycling variability through models, *Earth-Science Rev.*, 171, 349–382, <https://doi.org/10.1016/j.earscirev.2017.06.004>, 2017.
- 10 Hülse, D., Arndt, S., Daines, S., Regnier, P. and Ridgwell, A.: OMEN-SED 1.0: a novel, numerically efficient organic matter sediment diagenesis module for coupling to Earth system models, *Geosci. Model Dev.*, 11, 2649–2689, <https://doi.org/10.5194/gmd-11-2649-2018>, 2018.
- Hülse, D., Arndt, S. and Ridgwell, A.: Mitigation of extreme Ocean Anoxic Event conditions by organic matter sulfurization, *Paleoceanography and Paleoclimatology*, 34(4), 476–489, <https://doi.org/10.1029/2018PA003470>, 2019.
- 15 Jiang, C. Z. and Tosca, N. J.: Fe(II)-carbonate precipitation kinetics and the chemistry of anoxic ferruginous seawater, *Earth and Planetary Science Letters*, 506, 231–242, <https://doi.org/10.1016/j.epsl.2018.11.010>, 2019.
- Jimenez-Lopez, C., and Romanek, C. S.: Precipitation kinetics and carbon isotope partitioning of inorganic siderite at 25°C and 1 atm, *Geochim. Cosmochim. Acta*, 68(3), 557–571, 2004.
- Johnson, K.S.: Carbon dioxide hydration and dehydration kinetics in seawater, *Limnology and Oceanography*, 27, 849–855, <https://doi.org/10.4319/lo.1982.27.5.0849>, 1982.
- 20 Johnson, C. M., Roden, E. E., Welch, S. A. and Beard, B. L.: Experimental Constraints on Fe isotope fractionation during magnetite and Fe carbonate formation coupled to dissimilatory hydrous ferric oxide reduction, *Geochim. Cosmochim. Acta*, 69(4), 963–993, 2004.
- Jones C., Nomosatryo S., Crowe S. A., Bjerrum C. J. and Canfield D. E.: Iron oxides, divalent cations, silica, and the early earth phosphorus crisis, *Geology*, 43, 135–138, <https://doi.org/10.1038/10.1130/G36044.1>, 2015.
- 25 Kaplan, I. R. and Rittenberg, S. C.: Microbiological fractionation of sulphur isotopes, *J. Gen. Microbiol.*, 34, 195–212, 1964.
- Keeling, R. F., Körtzinger, A. and Gruber, N.: Ocean Deoxygenation in a Warming World. *Ann. Rev. Mar. Sci.*, 2, 199–229, <https://doi.org/10.1146/annurev.marine.010908.163855>, 2010.
- Kharecha, P., Kasting, J. and Siefert, J.: A coupled atmosphere-ecosystem model of the early Archean Earth. *Geobiology*, 2, 53–76, <https://doi.org/10.1111/j.1472-4669.2005.00049.x>, 2005.
- 30 Konhauser, K. O., Planavsky, N. J., Hardisty, D. S., Robbins, L. J., Warchola, T. J., Haugaard, R., Lalonde, S. V., Partin, C. A., Onk, P. B. H., Tsikos, H., Lyons, T. W., Bekker, A. and Johnson, C. M.: Iron formations: A global record of Neoproterozoic to Palaeoproterozoic environmental history, *Earth-Science Reviews*, 172, 140–177, <http://dx.doi.org/10.1016/j.earscirev.2017.06.012>, 2017.
- Konovalov, S.K., Murray, J.W., Luther, G.W. and Tebo, B.M.: Processes controlling the redox budget for the oxic/anoxic water column of the Black Sea, *Deep-Sea Research II*, 53, 1817–1841, <http://dx.doi.org/10.1016/j.dsr2.2006.03.013>, 2006.
- 35 Kump, L. R. and Seyfried, W. E.: Hydrothermal Fe fluxes during the Precambrian: Effect of low oceanic sulfate concentrations and low hydrostatic pressure on the composition of black smokers, *Earth and Planetary Science Letters*, 235(3–4), 654–662. <https://doi.org/10.1016/j.epsl.2005.04.040>, 2005.



- Laakso, T. A. and Schrag, D. P.: Regulation of atmospheric oxygen during the Proterozoic, *Earth Planet. Sci. Lett.*, 388, 81–91, <https://doi.org/10.1016/j.epsl.2013.11.049>, 2014.
- Laakso, T. A. and Schrag, D. P.: A small marine biosphere in the Proterozoic, *Geobiology*, 17, 161–171, <https://doi.org/10.1111/gbi.12323>, 2019.
- 5 LaRowe, D. E. and Van Cappellen, P.: Degradation of natural organic matter: A thermodynamic analysis, *Geochimica et Cosmochimica Acta*, 75(8), 2030–2042, <https://doi.org/10.1016/j.gca.2011.01.020>, 2011.
- Lenton T. M., Daines S. J. and Mills B. J. W.: COPSE reloaded: An improved model of biogeochemical cycling over Phanerozoic time, *Earth-Science Rev.*, 178, 1–28, <https://doi.org/10.1016/J.EARSCIREV.2017.12.004>, 2018.
- Liu, X. and Millero, F. J.: The solubility of iron in seawater, *Marine Chemistry*, 77, 43–54, 2002.
- 10 Lough, A. J. M., Connelly, D. P., Homoky, W. B., Hawkes, J. A., Chavagnac, V., Castillo, A., Kazemian, M., Nakamura, K. I., Araki, T., Kaulich, B. and Mills, R. A.: Diffuse hydrothermal venting: A hidden source of iron to the oceans, *Frontiers in Marine Science*, 6, 1–14. <https://doi.org/10.3389/fmars.2019.00329>, 2019.
- Lovley, D.: Dissimilatory Fe(III) and Mn(IV) Reduction, *Microbiological Reviews*, 55, 259–287, 1991
- Luther III, G. W., Rickard, D. T., Theberge, S. and Olroyd, A.: Determination of metal (bi)sulfide stability constants of Mn^{2+} , Fe^{2+} , Co^{2+} , Ni^{2+} , Cu^{2+} , and Zn^{2+} by voltammetric methods, *Environ. Sci. Technol.*, 30, 671–679, 1996.
- 15 Lyons, T. W., Reinhard, C. T. and Planavsky, N. J.: The rise of oxygen in Earth’s early ocean and atmosphere, *Nature*, 506, 307–315, 2014.
- Marsh, R., Müller, S. A., Yool, A., and Edwards, N. R.: Incorporation of the C-GOLDSTEIN efficient climate model into the GENIE framework: “eb_go_gs” configurations of GENIE, *Geosci. Model Dev.*, 4, 957–992, <https://doi.org/10.5194/gmd-4-957-2011>, 2011.
- Martin, J., Knauer, G. A., Karl, D., and Broenkow, W.: VERTEX: carbon cycling in the northeast Pacific, *Deep. Res.*, 34, 267–285, <https://doi.org/10.7591/9781501723667>, 1987.
- 20 Marion, G.M., Millero, F.J., Camões, M.F., Spitzer, P., Feistel, R. and Chen, C.T.A.: pH of seawater, *Marine Chemistry*, 126, 89–96, <https://doi.org/10.1016/j.marchem.2011.04.002>, 2011.
- Meyer, K. M., Ridgwell, A. and Payne, J. L.: The influence of the biological pump on ocean chemistry: Implications for long-term trends in marine redox chemistry, the global carbon cycle, and marine animal ecosystems, *Geobiology*, 14(3), 207–219, <https://doi.org/10.1111/gbi.12176>, 2016.
- 25 Meysman, F. J. R., Middelburg, J. J., Herman, P. M. J. and Heip, C. H. R.: Reactive transport in surface sediments. II. Media: an object-oriented problem-solving environment for early diagenesis, *Comput. Geosci.*, 29, 301–318, [https://doi.org/10.1016/S0098-3004\(03\)00007-4](https://doi.org/10.1016/S0098-3004(03)00007-4), 2003.
- Michard, G., Viollier, E., Jézéquel, D. and Sarazin, G.: Geochemical study of a crater lake: Pavin Lake, France - Identification, location and quantification of the chemical reactions in the lake, *Chem. Geol.*, 115, 103–115, 1994.
- 30 Mikucki, J. A., Pearson, A., Johnston, D. T., Turchyn, A. V., Farquhar, J., Schrag, D. P., Anbar, A. A., Prisco, J. C. and Lee, P. A.: A Contemporary Microbially Maintained Subglacial Ferrous “Ocean”, *Science*, 324, 397–400, 2009.
- Millero, F. J., Hubinger, S., Fernandez, M. and Garnett, S.: Oxidation of H_2S in seawater as a function of temperature, pH, and ionic strength, *Environ. Sci. Technol.*, 21, 439–443, <https://doi.org/10.1021/es00159a003>, 1987.
- 35 Millero, F. J., Sotolongo, S. and Izaguirre, M.: The oxidation kinetics of Fe(II) in seawater, *Geochim. Cosmochim. Acta*, 51, 793–207, [https://doi.org/10.1016/0016-7037\(87\)90093-7](https://doi.org/10.1016/0016-7037(87)90093-7), 1987.
- Millero, F. J., Wensheng, Y. and Aicher, J.: The speciation of Fe(II) and Fe(III) in natural waters, *Mar. Chem.*, 50, 21–39, 1995.



- Monteiro, F. M., Pancost, R. D., Ridgwell, A., and Donnadieu, Y.: Nutrients as the dominant control on the spread of anoxia and euxinia across the Cenomanian-Turonian oceanic anoxic event (OAE2): Model-data comparison, *Paleoceanography*, 27, PA4209, <https://doi.org/10.1029/2012PA002351>, 2012.
- Morris, M.: Factorial sampling plans for preliminary computational experiments., *Technometrics*, 33, 161-174, 1991.
- 5 Nisbet, E. G. and Sleep, N. H.: The habitat and nature of early life, *Nature*, 409, 1083–1091, <https://doi.org/10.1038/35059210>, 2001.
- Olson S. L., Kump L. R. and Kasting J. F.: Quantifying the areal extent and dissolved oxygen concentrations of Archean oxygen oases, *Chem. Geol.*, 362, 35–43, <https://doi.org/10.1016/j.chemgeo.2013.08.012>, 2013.
- Olson, S. L., Reinhard, C. T. and Lyons, T. W.: Limited role for methane in the mid-Proterozoic greenhouse, *PNAS*, 113, 11447-11452, 2016.
- Ozaki K., Tajima S. and Tajika E.: Conditions required for oceanic anoxia/euxinia: Constraints from a one-dimensional ocean biogeochemical
10 cycle model, *Earth Planet. Sci. Lett.* 304, 270–279, <https://doi.org/10.1016/J.EPSL.2011.02.011>, 2011.
- Ozaki, K., Tajika, E., Hong, P. K., Nakagawa, Y. and Reinhard, C. T.: Effects of primitive photosynthesis on Earth's early climate system, *Nature Geoscience*, 11(1), 55–59, <https://doi.org/10.1038/s41561-017-0031-2>, 2018.
- Ozaki K., Reinhard C. T. and Tajika E.: A sluggish mid-Proterozoic biosphere and its effect on Earth's redox balance, *Geobiology*, 17, 3–11, <http://doi.wiley.com/10.1111/gbi.12317>, 2019.
- 15 Parekh, P., Follows, M. J. and Boyle, E.: Modeling the global ocean iron cycle, *Global Biogeochemical Cycles*, 18, GB1002, <http://dx.doi.org/10.1029/2003GB002061>, 2004.
- Pellerin, A., Anderson-Trocmé, L., Whyte, L. G., Zane, G. M., Wall, J. D. and Wing, B. A.: Sulfur Isotope Fractionation during the Evolutionary Adaptation of a Sulfate-Reducing Bacterium, *Appl. Environ. Microbiol.*, 81, 2676-2689, <http://dx.doi.org/10.1128/AEM.03476-14>, 2015.
- 20 Pellerin, A., Wenk, C. B., Halevy, I. and Wing, B. A.: Sulfur Isotope Fractionation by Sulfate-Reducing Microbes Can Reflect Past Physiology, *Environ. Sci. Technol.*, 52, 4013-4022, <http://dx.doi.org/10.1021/acs.est.7b05119>, 2018.
- Pellerin, A., Antler, G., Holm, S. A., Findlay, A. J., Crockford, P. W., Turchyn, A. V., Jørgensen, B. B. and Finster, K.: Large sulfur isotope fractionation by bacterial sulfide oxidation, *Science Advances*, 5, eaaw1480, <https://doi.org/10.1126/sciadv.aaw1480>, 2019.
- Pianosi, F., Sarrazin, F. and Wagener, T.: A Matlab toolbox for Global Sensitivity Analysis, *Environmental Modelling & Software*, 70, 80-85,
25 2015.
- Pianosi, F., Beven, K., Freer, J., Hall, J. W., Rougier, J., Stephenson, D. B. and Wagener, T.: Sensitivity analysis of environmental models: A systematic review with practical workflow, *Environmental Modelling & Software*, 79, 214-232, 2016.
- Picard, A., Kappler, A., Schmid, G., Quaroni, L. and Obst, M.: Experimental diagenesis of organo-mineral structures formed by microaerophilic Fe(II)-oxidizing bacteria, *Nature Communications*, 6, 6277, <http://dx.doi.org/10.1038/ncomms7277>, 2015.
- 30 Planavsky, N. J., McGoldrick, P., Scott, C. T., Li, C., Reinhard, C. T., Kelly, A. E., Chu, X., Bekker, A., Love, G. D. and Lyons, T. W.: Widespread iron-rich conditions in the mid-Proterozoic ocean, *Nature*, 477, 448–451, <https://doi.org/10.1038/nature10327>, 2011.
- Poser, A., Vogt, C., Knöller, K., Ahlheim, J., Weiss, H., Kleinstüber, S. and Richnow, H.-H.: Stable Sulfur and Oxygen Isotope Fractionation of Anoxic Sulfide Oxidation by Two Different Enzymatic Pathway, *Environ. Sci. Technol.*, 48, 9094-9102, <https://doi.org/10.1021/es404808r>, 2014.
- 35 Postma, D. and Jakobsen, R.: Redox zonation: equilibrium constraints on the Fe(III)/SO₄-reduction interface, *Geochim. Cosmochim. Acta*, 60, 3169–3175, [https://doi.org/10.1016/0016-7037\(96\)00156-1](https://doi.org/10.1016/0016-7037(96)00156-1), 1996.
- Poulton, S. W. and Raiswell, R.: The low-temperature geochemical cycle of iron: from continental fluxes to marine sediment deposition, *American Journal of Science*, 302, 774–805, 2002.



- Poulton, S. W., Krom, M. D. and Raiswell, R.: A revised scheme for the reactivity of iron (oxyhydr)oxide minerals towards dissolved sulfide, *Geochim. Cosmochim. Acta*, 68(18), 3703–3715, 2004.
- Poulton, S. W. and Canfield, D. E.: Development of a sequential extraction procedure for iron: Implications for iron partitioning in continentally derived particulates, *Chem. Geol.*, 214, 209–221, <https://doi.org/10.1016/j.chemgeo.2004.09.003>, 2005.
- 5 Poulton, S. W., Fralick, P. W. and Canfield, D. E.: Spatial variability in oceanic redox structure 1.8 billion years ago, *Nat. Geosci.*, 3, 486–490, 2010.
- Poulton, S. W. and Canfield, D. E.: Ferruginous Conditions: A Dominant Feature of the Ocean through Earth’s History, *Elements*, 7, 107–112, <https://doi.org/10.2113/gselements.7.2.107>, 2011.
- Price, F. T. and Shieh, Y. N.: Fractionation of sulfur isotopes during laboratory synthesis of pyrite at low temperatures, *Chem. Geol.*, 27, 245–253, 1979.
- 10 Raiswell, R. and Canfield, D. E.: The iron biogeochemical cycle past and present, *Geochemical Perspectives*, 1(1), 2012.
- Rasmussen, B., Krapež, B., Muhling, J.R. and Suvorova, A.: Precipitation of iron silicate nanoparticles in early Precambrian oceans marks Earth’s first iron age, *Geology*, 43, 303–306, <https://doi.org/10.1130/G36309.1>, 2015.
- Reinhard C. T., Planavsky N. J., Robbins L. J., Partin C. A., Gill B. C., Lalonde S. V., Bekker A., Konhauser K. O. and Lyons T. W.: Proterozoic ocean redox and biogeochemical stasis, *Proc. Natl. Acad. Sci. U.S.A.*, 110, 5357–5362, <https://doi.org/10.1038/10.1073/pnas.1208622110>, 2013.
- 15 Reinhard C. T., Planavsky N. J., Olson S. L., Lyons T. W. and Erwin D. H.: Earth’s oxygen cycle and the evolution of animal life., *Proc. Natl. Acad. Sci. U.S.A.*, 113, 8933–8, <https://doi.org/10.1038/10.1073/pnas.1521544113>, 2016.
- Reinhard C. T., Planavsky N. J., Gill B. C., Ozaki K., Robbins L. J., Lyons T. W., Fischer W. W., Wang C., Cole D. B. and Konhauser K. O.: Evolution of the global phosphorus cycle, *Nature*, 541, 386–389, <https://doi.org/10.1038/nature20772>, 2017.
- Reinhard, C. T., Olson S. L., Pälke, C., Kirtland Turner, S., Kanzaki, Y. and Ridgwell, A.: Oceanic and atmospheric methane cycling in the cGENIE Earth System model, *Geosci. Model Dev. Disc.*, 1–45, <https://doi.org/10.5194/gmd-2020-32>, 2020.
- Reinhard, C.T., Planavsky, N.J., Ward, B.A., Love, G.D., Le Hir, G., Ridgwell, A.: The impact of marine nutrient abundance on early eukaryotic ecosystems, *Geobiology*, 18, 139–151, <https://doi.org/10.1111/gbi.12384>, 2020.
- 25 Rickard, D.: Kinetics of pyrite formation by the H_2S oxidation of iron(II) monosulfide in aqueous solutions between 25 and 125°C: The rate equation, *Geochim. Cosmochim. Acta*, 61, 115–134, 1997.
- Rickard, D.: The solubility of FeS, *Geochim. Cosmochim. Acta*, 70, 5779–5789, 2006.
- Ridgwell, A., Hargreaves, J. C., Edwards, N. R., Annan, J. D., Lenton, T. M., Marsh, R., Yool, A. and Watson, A.: marine geochemical data assimilation in an efficient Earth System Model of global biogeochemical cycling, *Biogeosciences*, 4, 87–104, 2007.
- 30 Ridgwell, A. and Hargreaves, J. C.: Regulation of atmospheric CO₂ by deep-sea sediments in an Earth system model, *Global Biogeochem. Cycles*, 21, GB2008, doi:10.1029/2006GB002764, 2007.
- Ridgwell, A.: Evolution of the ocean’s “biological pump”, *Proceedings of the National Academy of Sciences*, 108(40), 16485–16486, doi:10.1073/pnas.1112236108, 2011.
- Ridgwell, A. and Death, R.: Iron limitation in an efficient model of global carbon cycling and climate, in preparation.
- 35 Rolison, J. M., Stirling, C. H., Middag, R., Gault-Ringold, M., George, E., Rijkenberg, M. J. A.: Iron isotope fractionation during pyrite formation in a sulfidic Precambrian ocean analogue, *Earth and Planetary Science Letters*, 488, 1–13, <https://doi.org/10.1016/j.epsl.2018.02.006>, 2018.



- Rouxel, O. J., Bekker, A. and Edwards, K. J.: Iron isotope constraints on the Archean and Paleoproterozoic ocean redox state. *Science*, 307, 1088–1091, 2005.
- Sim, M. S., Bosak, T. and Ono, S.: Large sulfur isotope fractionation does not require disproportionation, *Science*, 333, 74–77, 2011.
- Sharp, J. H., Benner, R., Bennett, L., Carlson, C. A., Fitzwater, S. E., Pletzer, E. T. and Tupas, L., M.: Analyses of dissolved organic carbon
5 in seawater: the JGOFS EqPac methods comparison, *Mar. Chem.*, 48, 91–108, 1995.
- Soetaert, K., Herman, P. M. J. and Middelburg, J. J.: A model of early diagenetic processes from the shelf To abyssal depths, *Geochim. Cosmochim. Acta*, 60, 1019–1040, 1996.
- Sperling, E. A., Wolock, C. J., Morgan, A. S., Gill, B. C., Kunzmann, M., Halverson, G. P., Macdonald, F. A., Knoll, A. H. and
10 Johnston, D. T.: Statistical analysis of iron geochemical data suggests limited late Proterozoic oxygenation, *Nature*, 523, 451–454,
<https://doi.org/10.1038/nature14589>, 2015.
- Steeffel, C.I. and Van Cappellen, P.: A new kinetic approach to modeling water–rock interaction: the role of nucleation, precursors, and
Ostwald ripening, *Geochim. Cosmochim. Acta*, 54, 2657–2677, [https://doi.org/10.1016/0016-7037\(90\)90003-4](https://doi.org/10.1016/0016-7037(90)90003-4), 1990.
- Suits, N. S. and Wilkin, R. T.: Pyrite formation in the water column and sediments of a meromictic lake, *Geology*, 26, 1099–1102, 1998.
- Tagliabue, A., Bopp, L., Dutay, J. C., Bowie, A. R., Chever, F., Jean-Baptiste, P., Bucciarelli, E., Lannuzel, D., Remenyi, T., Sarthou, G.,
15 Aumont, O., Gehlen, M. and Jeandel, C.: Hydrothermal contribution to the oceanic dissolved iron inventory, *Nature Geoscience*, 3(4),
252–256, <https://doi.org/10.1038/ngeo818>, 2010.
- Tagliabue, A., Aumont, O., DeAth, R., Dunne, J. P., Dutkiewicz, S., Galbraith, E., Misumi, K., Moore, J. K., Ridgwell, A., Sherman, E.,
Stock, C., Vichi, M., Völker, C., and Yool, A.: How well do global ocean biogeochemistry models simulate dissolved iron distributions?,
Global Biogeochem. Cy., 30, 149–174, <https://doi.org/10.1002/2015GB005289>, 2016.
- 20 Taillefert, M. and Gaillard, J.-F.: Reactive transport modeling of trace elements in the water column of a stratified lake: iron cycling and
metal scavenging, *Journal of Hydrology*, 256, 16–34, 2002.
- Thamdrup, B.: Chapter 2: Bacterial Manganese and Iron Reduction in Aquatic Sediments, In: *Advances in Microbial Ecology*, Schink, B.
ed., Luwer Academic/Plenum Publishers, 2000.
- Thompson, K. J., Kenward, P. A., Bauer, K. W., Warchola, T., Gauger, T., Martinez, R., Sinister, R. L., Michiels, C. C., Llíros, Reinhard, C.
25 T., Kappler, A., Konhauser, K. O. and Crowe, S. A.: Photoferrotrophy, deposition of banded iron formations, and methane production in
Archean oceans, *Science Advances*, 5, eaav2869, <https://doi.org/10.1126/sciadv.aav2869>, 2019.
- Tosca, N. J., Guggenheim, S. and Pufahl, P. K.: An authigenic origin for Precambrian greenalite: Implications for iron formation and the
chemistry of ancient seawater, *Geological Society of America Bulletin*, <https://doi.org/10.1130/B31339.1>, 2015.
- Tosca, N. J., Jiang, C. Z., Rasmussen, B. and Muhling, J.: Products of the iron cycle on the early Earth, *Free Radical Biology and Medicine*,
30 140, 138–153, <https://doi.org/10.1016/j.freeradbiomed.2019.05.005>, 2019.
- Tribovillard, N., Algeo, T. J., Lyons, T. and Riboulleau, A.: Trace metals as paleoredox and paleoproductivity proxies: An update, *Chem.
Geol.*, 232, 12–32, <https://doi.org/10.1016/j.chemgeo.2006.02.012>, 2006.
- Tucker, M. E.: Chapter 7: Sedimentary iron deposits, In: *Sedimentary petrology: an introduction to the origin of sedimentary rocks*, Blackwell
Science, 2001
- 35 Van Cappellen, P. and Ingall, E. D.: Redox stabilization of the Atmosphere and Oceans by Phosphorus-Limited Marine Productivity, *Science*,
271, 493–496, <https://doi.org/10.1126/science.271.5248.493>, 1996.
- Van Cappellen, P. and Wang, Y. F.: Cycling of iron and manganese in surface sediments - A general theory for the coupled transport and
reaction of carbon, oxygen, nitrogen, sulfur, iron, and manganese, *Am. J. Sci.*, 296, 197–243, 1996.



- van de Velde, S. and Meysman, F. J. R.: The Influence of Bioturbation on Iron and Sulphur Cycling in Marine Sediments: A Model Analysis, *Aquat. Geochem.*, 22, 469–504, 2016.
- van de Velde, S., Mills, B. J. W., Meysman, F. J. R., Lenton, T. M. and Poulton, S. W.: Early Palaeozoic ocean anoxia and global warming driven by the evolution of shallow burrowing, *Nat. Comm.*, 9, 2554, <https://doi.org/10.1038/s41467-018-04973-4>, 2018.
- 5 van de Velde, S. J., Hidalgo-Martinez, S., Callebaut, I., Antler, G., James, R. K., Leermakers, M. and Meysman, F. J. R.: Burrowing fauna mediate alternative stable states in the redox cycling of salt marsh sediments. *Geochim. Cosmochim. Acta*, 276, 31–49, 2020
- van de Velde, S., Reinhard, C., Ridgwell, A. and Meysman, F. J. R.: Bistability in the redox chemistry of sediments and oceans, in review at *Proceedings of the National Academy of Science*.
- Vervoort, P., Kirtland Tuner, S., Rochholz, F., and Ridgwell, A.: Earth System Model Analysis of how Astronomical Forcing is Imprinted
10 onto the Marine Geological Record, in review at *Paleoceanography and Paleoclimatology*.
- Wallmann K., Flögel S., Scholz F., Dale A. W., Kemena T. P., Steinig S. and Kuhnt W.: Periodic changes in the Cretaceous ocean and climate caused by marine redox see-saw, *Nat. Geosci.*, 12, 456–461, <https://doi.org/10.1038/s41561-019-0359-x>, 2019.
- Walter, X. A., Picazo, A., Miracle, M. R., Vicente, E., Camacho, A., Aragno, M. and Zopfi, J.: Phototrophic Fe(II)-oxidation in the chemocline of a ferruginous meromictic lake, *Front. Microbiol.*, 5, 713, 2014.
- 15 Ward, B.A., Wilson, J.D., Death, R., Monteiro, F.M., Yool, A. and Ridgwell, A.: EcoGENIE 1.0: plankton ecology in the cGENIE Earth system model. *Geosci. Model Dev.*, 11, 4241–4267, <https://doi.org/10.5194/gmd-11-4241-2018>, 2018.
- Wiesli, R. A., Beard, B. L. and Johnson, C. M.: Experimental determination of Fe isotope fractionation between aqueous Fe(II), siderite and "green rust" in abiotic systems, *Chem. Geol.*, 211, 343–362, 2004.
- Wing, B. A. and Halevy, I.: Intracellular metabolite levels shape sulfur isotope fractionation during microbial sulfate respiration, *Proceedings
20 of the National Academy of Sciences*, 111(51), 18116–18125, <https://doi.org/10.1073/pnas.1407502111>, 2014.
- Xiong, Y., Guilbaud, R., Peacock, C. L., Cox, R. P., Canfield, D. E., Krom, M. D. and Poulton, S. W.: Phosphorus cycling in Lake Cadagno, Switzerland: A low sulfate euxinic ocean analogue, *Geochimica et Cosmochimica Acta*, 251, 116–135, <https://doi.org/10.1016/j.gca.2019.02.011>, 2019.
- Zegeye, A., Bonneville, S., Benning, L. G., Sturm, A., Fowle, D. A., Jones, C., Canfield, D. E., Ruby, C., MacLean, L. C., Nomosatryo, S.,
25 Crowe, S. and Poulton, S. W.: Green rust formation controls nutrient availability in a ferruginous water column, *Geology*, 40, 599–602, <https://doi.org/10.1130/G32959>, 2012.
- Zhang, J. Z. and Millero, F. J.: The products from the oxidation of H_2S in seawater, *Geochimica et Cosmochimica Acta*, 57, 1705–1718, 1993.