

BARRA v1.0: Kilometre-scale downscaling of an Australian regional atmospheric reanalysis over four midlatitude domains

Chun-Hsu Su¹, Nathan Eizenberg², Dörte Jakob¹, Paul Fox-Hughes³, Peter Steinle¹, Christopher J. White^{4,5}, Charmaine Franklin¹

5 ¹Bureau of Meteorology, Docklands, Victoria 3008, Australia

²Department of Earth Sciences, The University of Melbourne, Parkville, Victoria 3010, Australia

³Bureau of Meteorology, Hobart, Tasmania 7000, Australia

⁴Department of Civil and Environmental Engineering, University of Strathclyde, Glasgow, Scotland, UK

⁵School of Engineering, University of Tasmania, Hobart, Australia

10

Correspondence to: Chun-Hsu Su (chunhsu.su@bom.gov.au)

Abstract. Regional reanalyses provide a dynamically consistent recreation of past weather observations at scales useful for local-scale environmental applications. The development of convection-permitting models (CPMs) in numerical weather prediction has facilitated the creation of kilometre-scale (1-4 km) regional reanalysis and climate projections. The Bureau of

15 Meteorology Atmospheric high-resolution Regional Reanalysis for Australia (BARRA) also aims to realise the benefits of these high-resolution models over Australian sub-regions for applications such as fire danger research, by nesting them in BARRA's 12 km regional reanalysis (BARRA-R). Four mid-latitude sub-regions are centred on Perth in Western Australia, Adelaide in South Australia, Sydney in New South Wales (NSW), and Tasmania. The resulting 29-year 1.5 km downscaled reanalyses (BARRA-C) are assessed for their added skill over BARRA-R and global reanalyses for near-surface parameters

20 (temperature, wind and precipitation) at observation locations and against independent 5 km gridded analyses. BARRA-C demonstrates better agreement with point observations for temperature and wind, particularly in topographically complex regions and coastal regions. BARRA-C also improves upon BARRA-R in terms of intensity and timing of precipitation during the thunderstorm seasons in NSW, and spatial patterns of sub-daily rain fields during storm events. BARRA-C reflects known issues of CPMs: overestimation of heavy rain rates and rain cells, and underestimation of light rain occurrence. However, as a hindcast-only system, BARRA-C largely inherits the domain-averaged biases and temporal variations of biases pattern from BARRA-R but simulates different climatological extremes for temperature and precipitation. An Added value analysis of temperature and precipitation extremes shows that BARRA-C provides additional skill over BARRA-R when compared to gridded observations. The for the spatial patterns of BARRA-C the warm temperature extremes and wet precipitation extremes are more highly correlated with observations. and bias in

25 BARRA-C adds value in representation of the spatial pattern of cold extremes over coastal regions- but remains biased in terms of magnitude, but simulates different climatological extremes for temperature and precipitation. FurtherIn particular, BARRA-C reflects known issues of CPMs: overestimation of heavy rain rates and rain cells, and underestimation of light rain occurrence.

1 Introduction

35 At horizontal kilometre-scales (1-4 km), convection-permitting models (CPMs) have provided a step-change in weather forecasting capabilities, particularly for forecasting rainfall and cloud cover (e.g., Lopez et al., 2009; Mailhot et al., 2010; Brousseau et al., 2016; Clark et al., 2016) and over local regions with complex terrain or land-sea boundaries (Calmet et al., 2018). Similarly, CPMs have provided new insights in regional climate projections (e.g., Argüeso et al., 2014; Prein et al., 2015; Kendon et al., 2017; 2019) beyond current global models. For instance, regional CPMs have suggested that future 40 increases in short-duration precipitation extremes are larger than what can be expected from increases in atmospheric moisture alone (Kendon et al., 2021 and references therein). Major efforts are underway toward refining the horizontal resolution of global climate models to kilometre-scale (Schär et al., 2020). Extreme weather events such as thunderstorms, damaging winds, and hailstorms, are better represented in higher resolution models (Walsh et al., 2016). Current gGeneral practice is that grid spacings less than about four kilometres are required to explicitly model small convective cloud 45 processes, replacing parameterizations of moist convection. This avoids several issues seen in parameterized convection schemes used in models with a grid spacing greater than 10 km (Lean et al., 2008) and the “grey zone” issues in mesoscale (4-10 km) scale models (Gerard et al., 2009). A common assumption of traditional convective parameterizations is that cloud fields adjust so much more rapidly than the processes forcing it that this adjustment can be modelled as instantaneous. Such schemes thus have no “memory” of the meteorological flow, leading to unrealistic model behaviours. Models with 50 parameterized convection exhibit including premature convective initiation, misrepresented diurnal cycle of precipitation, over-estimation of drizzle occurrence, under-estimation of extreme rainfall (Lean et al., 2008; Clark et al., 2016), fewer identifiable mesoscale convective systems with less structure (Done et al., 2004), and rainfall coastal locking where precipitation generated over the sea does not penetrate inland (Bureau of Meteorology, 2018). When the parameterization scheme is used at a finer resolution than 10 km, it also tends to produce intermittent on-off behaviour of deep convection 55 (Gerard et al., 2009).

By contrast, CPMs can represent deep convection and mesoscale convective organization explicitly on the model grid. Explicit representation-modelling of convection improves the physical nature of better captures precipitation persisting across orographic or land-sea boundaries by the advection of clouds/precipitation. Better representation of topography in 60 CPMs also leads to improved wind circulation patterns and resulting vertical velocities (e.g., Fosser et al., 2015). Improved modelling of the interactions between storm cells and their organisations should improve the estimation of damaging winds. Many studies have found a better diurnal cycle of tropical convection over land, cloud vertical structure, and coupling between moisture convection and convergence in CPMs (Stein et al., 2015; Leutwyler et al., 2017). A finer grid resolution can improve the flow and wind simulation over the recirculation zone behind the escarpment of a hill and higher vertical grid resolution improves simulation on the lee side of hills (Ma and Liu, 2017).

65 These benefits from using CPMs are however yet to be fully realized in many atmospheric reanalyses. Atmospheric reanalyses combine prior knowledge of physical processes captured in the models with observations from a diverse range of

instruments to form spatially complete representations of the historical atmospheric conditions. They are therefore invaluable for revisiting applications concerned with the local weather processes, climate signals or events that were not fully observed, for applications such as climate monitoring and change assessments (Kendon et al., 2017; 2019), renewable energy 70 assessment (e.g., Frank et al., 2020), and hazard management (e.g., Vitolo et al., 2019). Global-scale reanalyses have advanced in quality and quantity during the past three decades with improvements to models, data assimilation methods, the number of observations and ensemble methods (Kalnay et al., 1996; Ebital et al., 2011; Gelaro et al., 2017; Dee et al., 2011), and also with increasing spatial resolutions. The latest addition, ERA5 (Hersbach et al., 2020), has a horizontal spacing of 31 km. Users of reanalyses have called for development towards finer spatial and temporal scales, i.e., below 10 km horizontal 75 spacing and sub-daily time intervals (Gregow et al., 2016). Such scales are needed in localized climate monitoring where local-scale mechanisms influenced by complex topography, coastlines and convective processes are responsible for local climate features and feedbacks.

Departing markedly from the global reanalyses are the regional reanalyses that use limited area models at higher horizontal resolutions over sub-regions, e.g., such as North America (Mesinger et al., 2006), the Arctic polar region 80 (Bromwich et al., 2016), Europe (Borsche et al., 2015 and references therein), India (Mahmood et al., 2018) and Australia (Su et al., 2019). These reanalyses use grid lengths in the order of 10 km to improve the representation of sub-daily variability and near-surface weather. These are generally produced with global atmosphere model configurations that include convection parameterizations (e.g., Su et al., 2019). Recently, Wahl et al. (2017) overcame this with a 7-year 2 km reanalysis 85 over Germany; with the assimilation of conventional observations and radar-derived rain rates and, showed demonstrating improved spatiotemporal variability and intensity frequency of precipitation. Such a direction in the development of the reanalyses, combined with the higher resolution regional projections, can offer a more accurate picture of changes in regional meteorology and extreme weather in the changing climate.

Dynamical downscaling is frequently used to estimate the dynamic variables at scales finer than those of coarser-resolution climate or weather models. This approach is undertaken at the Bureau of Meteorology (Bureau) in Australia to 90 produce kilometre-scale weather forecasts and/or ensemble forecasts over major cities and a 1.5 km forecast-only model has been used since 2017 for added value over the Bureau's lower resolution global system. This goal is also pursued in the Bureau of Meteorology Atmospheric high-resolution Regional Reanalysis for Australia (BARRA; Jakob et al., 2017) project. Within this context, this paper is a companion paper to Su et al. (2019) where an Australian regional 12 km reanalysis system (BARRA-R) was presented. Here we describe dynamical downscaling of BARRA-R using the UK Met Office 95 (UKMO) Unified Model (UM) at a 1.5 km horizontal grid length over four mid-latitude sub-regions of Australia (Figure 1) over 29 years from January 1990 to February 2019. These regions are chosen in partnership with state fire and emergency management agencies, because of the important advantages that dynamically downscaled reanalyses can provide for local-scale planning and management to reduce future risks due to extreme weather events such as bushfires. The four downscaling models, collectively referred to as BARRA-C, yield gridded products that include a variety of 10 minute to

100 hourly surface parameters describing both weather and land-surface conditions as well as ~~and~~ hourly upper-air parameters covering the troposphere and stratosphere with a 40 km model top on 70 model levels and 37 pressure levels.

This paper describes the model and the experimental design in Sect. 2, and Sect. 3 provides the first assessment of the downscaled reanalysis with focus on screen-level temperature, 10 m wind and precipitation. Comparisons with ~~the~~ BARRA-R and global reanalyses are also made to illustrate the added value of BARRA-C. Our findings are further 105 discussed in Sect. 4, with an overall summary in Sect. 5.

2 BARRA-C

The development of BARRA is based on the Bureau's operational deterministic NWP forecasting over the Australian region using the Australian Community Climate and Earth-System Simulator systems ACCESS-R and ACCESS-C (Puri et al., 2013). The operational version at the time (Australian Parallel Suite 2) of ACCESS-R is the national 12 km 6-hourly

110 analysis/assimilation and 3-day forecasting system (Bureau of Meteorology, 2016). ~~It~~-ACCESS-R has provided the initial and boundary conditions to initialize and constrain ACCESS-C over 6 smaller domains centred at the Australian cities ~~up~~ until 2020 (Bureau of Meteorology, 2018). The APS2 ACCESS-C dynamically downscale ACCESS-R to provide 6 hourly, 1.5-day forecasts at 1.5 km horizontal resolution. The relation between BARRA-R and BARRA-C mirrors this system but is implemented with shorter forecast (or hindcast) range, and a newer version of the meteorological forecast model and science 115 configuration (Section 2.1). In particular, BARRA-R is nested in ERA-Interim reanalysis (Dee et al., 2011) and includes four assimilation and hindcast cycles per day (Su et al., 2019), ~~and~~ BARRA-C-is a hindcast-only system which inherits the analysis from BARRA-R as initial conditions. While BARRA-C refers to the collection of the four sub-domain models, we use BARRA-AD, BARRA-PH, BARRA-SY and BARRA-TA to denote individual domains centred at Adelaide (South Australia, AD), Perth (Western Australia, PH), Sydney (New South Wales, SY), and Tasmania (TA) (Figure 1).

120 The PH and AD domains are similar in terms of climate, with having arid deserts north of their domains, and temperate dry hot or warm summers near coasts, and arid steppe climate in-between (Peel et al., 2007). SY has a temperate climate with warm to hot summers and lacks a dry season, while TA differs with a cooler summer. Cool-season (C3) perennial grass (C3) is the dominant vegetation over the southwestern region of PH and near-coast region of AD, and broadleaf trees are widespread in the SY and TA domains (Figure S1 in the Supplement). There are several large ephemeral

125 salt lakes (e.g., Lake Torrens, Lake Gairdner) in the AD domain, and these are modelled as land points with bare soil. Of the four domains, ~~It~~ is only SY ~~that~~ has a distinct thunderstorm season which occurs during November-March. Thunderstorms are far less frequent in the other three domains due to lower incidence of warm, humid air masses, favourable for storm development and also prevalent stable conditions during the potentially favourable warmer months owing to subtropical high-pressure belt over or near these areas (Kuleshov et al., 2002). In contrast to PH and AD, ~~t~~ The SY and TA domains are 130 topographically complex, with ~~t~~ The Great Dividing Range extending north to south through the SY domain and the TA domain features low mountains and a landscape of plateaus, and low mountain ranges in the TA domain.

2.1 Forecast model

The UM (Davies et al., 2005; version 10.6) is the grid-point atmospheric model used in BARRA and ACCESS. It uses a non-hydrostatic, fully compressible, deep atmosphere formulation and its dynamical core (Even Newer Dynamics for 135 General atmospheric modelling of the environment, ENDGame) solves the equations of motion using mass-conserving, semi-implicit, semi-Lagrangian (SL), time integration methods (Wood et al., 2014). The prognostic variables are three-dimensional wind components, virtual dry potential temperature and Exner pressure, dry density, and mixing ratios of moist quantities. These variables are discretized horizontally onto a regular longitude-latitude grid with Arakawa-C staggering (Arakawa and Lamb, 1977); and vertically with the Charney-Phillips staggered grid (Charney and Phillips, 1953). THere the 140 BARRA-C model has a horizontal spacing of $0.0135^\circ \times 0.0135^\circ$ (about 1.5 km at the equator) and its vertical levels follow the modelled orography at the surface and relax to surfaces of uniform radial height after 62 model levels (~ 17 km above ground) in the upper atmosphere with the model top height of 40 km. At this resolution, the model is run with an integration time step of 60 seconds.

The science configuration of the model in BARRA-C is based on the UK Met Office operational suite OS36, while 145 BARRA-R is based on Global Atmosphere (GA6) configuration of Walters et al. (2017). While the OS36 model configurations preceded the release of the first UM Regional Atmosphere and Land (RAL1) configuration of Bush et al. (2020), BARRA-C implements some of the improvements from RAL1. Table 1 summarizes the differences between BARRA-C, BARRA-R, and RAL1. The physical parameterization schemes common to BARRA-C and BARRA-R include a variant of Wilson and Ballard (1999) for mixed-phase cloud microphysics, the large-scale cloud scheme of Smith (1990), 150 and the radiation scheme of Edwards and Slingo (1996), all of which have been improved since publication. The UM BARRA-R uses a convection parameterization scheme based on Gregory and Rowntree (1990); which is not used in BARRA-CC. At With the a grid length of 1.5 km, the horizontal grid length approaches the depth of the boundary layer (Hanley et al., 2015) and as such it is no longer appropriate to use the 1D boundary layer parameterization that restricts mixing to the vertical. BARRA-C therefore uses a blended boundary layer parameterization (Boutle et al. 2014) where the 155 scheme transitions from the 1D vertical turbulence scheme of Lock et al. (2000) to a 3D subgrid turbulence scheme based on Smagorinsky (1963) as a function of the grid length to the turbulent length scale. The mixing length of 300 m, which can be tuned to control the smoothness of the fields and the number of small cells, is taken as 300 m from which is used in the operational systems.

The cloud scheme uses a profile of critical relative humidity values (RHcrit), above which a grid box contains some 160 cloud if the relative humidity is exceeded. Based on the assumption that there should be less subgrid variability in humidity in for smaller grid boxes, BARRA-C uses higher RHcrit values than BARRA-R in the lowest few layers, decreasing smoothly above to 0.8.

Without the convection parameterization scheme, BARRA-C relies on the model dynamics to represent convective motions. While convection remains unresolved in 1.5 km models, removal of the cumulus parameterization has shown to

165 result in more realistic behaviour (Clark et al., 2016). In particular, the model can explicitly capture processes with convective-like characteristics, which can subsequently drive scales that the model can properly resolve. BARRA-C also reduces the appearance of unrealistically strong vertical velocities and “grid-point storms” seen in BARRA-R due to the inability of convective parameterization to stabilize the air column (Su et al., 2019). Nevertheless, convection can remain under-resolved, leading to cases of too-early small, shallow showers or no rainfall at all. The mid-latitude version of RAL1
170 therefore includes stochastic perturbations of temperature and moisture and relative weak turbulent mixing, to encourage the model fields to be less uniform and help convection to initiate. It is of note that the stochastic perturbations of moisture are absent in BARRA-C does not use stochastic perturbations for moisture, and thus may still suffer from convection the initiation issues.

Another distinguishing feature of BARRA-C is the handling of mass conservation during the advection of moisture
175 prognostic variables. This is one of the key science developments in RAL1. BARRA-C and RAL1 use the zero-lateral flux scheme of Zerroukat and Shipway (2017) for moisture conservation at the model’s lateral boundaries, avoiding spurious extreme precipitation caused by the SL treatment of moisture variables near partially-resolved convection.

BARRA-C is missing some of the configuration improvements introduced in RAL1 because production runs had already commenced. BARRA-C however does not include a set of changes to the representation of the land surface and the
180 canopy radiation model in RAL1, which have shown to improve the issue of the damped diurnal cycle issue in near-surface temperatures. BARRA-C also does not benefit from the improved treatment of gaseous absorption in both long- and short-wave regimes in GA7 and RAL1, which improves interaction with band-by-band aerosol and cloud forcing.

BARRA uses the land surface scheme of Best et al. (2011), implemented in the Joint UK Land Environment and Simulator (JULES). It describes a 3 m four-layer soil column, with sub-surface temperature updated using a heat diffusion
185 equation and, with the vertical moisture flux is estimated using the Richard’s equation and Darcy’s law. The soil hydraulics is are computed using the van Genuchten equation. It uses a nine-tile approach to represent subgrid-scale heterogeneity in land cover, with the surface of each land point subdivided into five vegetation types (broadleaf trees, needle-leaved trees, temperate cool-season (C3) grass, tropical warm-season (C4) grass, and shrubs) and four non-vegetated surface types (urban, inland water, bare soil, and land ice). In particular, the urban surfaces are represented only by a single urban tile, where and such that street canyons and roofs are not distinguished.

The characteristics of the lower boundary, climatological fields, and natural and anthropogenic emissions are specified using static ancillary fields. These are created as per Bush et al. (2020; Table A1), with the exceptions of ancillaries for the land-sea mask, canopy tree heights, and land usage. The land-sea mask is created from the 1 km resolution International Geosphere–Biosphere Programme (IGBP) land cover data (Loveland et al., 2000) for SY and TA, and from
195 Shuttle Radar Topography Mission (SRTM) orography data for AD and PH. Land cover data based on Climate Change Initiative (CCI, Hartley et al., 2017) is not adopted here as its mapping to the nine land surface tiles over the Australian region remains untested. The canopy tree heights are derived from satellite light detection and ranging (lidar; Simard et al., 2011; Dharssi et al., 2015). The land usage ancillary, created from IGBP, is modified for AD and PH to match the water

fractions in the Water Observations from Space (WOfS, Mueller et al., 2016). Aerosol absorption and scattering in the
200 radiation scheme assume climatological aerosol properties. A climatological ozone field is also prescribed.

2.2 Initial and boundary conditions

The BARRA-C model hindcast is re-initialized with 6-hourly initial conditions at the synoptic hours $t_0 = 00:00, 06:00, 12:00$,
and 18:00 UTC created by downscaling from BARRA-R analyses (Figure S2 in the Supplement). These fields are taken
205 from the centre of BARRA-R's 6-hour analysis windows. A two-component reconfiguration approach is used~~taken~~, in which
BARRA-R winds, moisture and temperature are downscaled separately with different resolution topography sets, to remove
210 model instability ~~ever due to high topography~~~~large horizontal topography gradients~~. BARRA-C is further constrained by
BARRA-R at the lateral boundaries without nudging, based on the prescription described in Bush et al. (2020) and a
boundary rim width of 0.34° . The boundary conditions force the development of the larger-scale features within the
BARRA-C domains. These setups follow the Bureau's NWP system~~s~~ and ensure that the benefits of the BARRA-R analysis
215 ~~is~~are inherited by BARRA-C, where ~~BARRA-C~~the nested model is treated as a physically consistent interpolator of
~~BARRA-R~~the driving model.

The JULES soil moisture and temperature are prescribed by BARRA-R. Consistent with BARRA-R, daily sea
surface temperature and sea ice $0.05 \times 0.05^\circ$ analysis from reprocessed (1985–2007; Roberts-Jones et al., 2012) and near-
real-time Operational Sea Surface Temperature and Ice Analysis (OSTIA; Donlon et al., 2012) are used as lower boundaries
215 over the water after being interpolated to the BARRA-C ~~UM~~ grid. The ~~NRT~~near-real-time data are used from January 2007.

Each hindcast in BARRA-C is a 9-hour simulation ~~but only 6-hours are used~~, ~~where~~The model data during the
first 3-hours ~~period~~ is discarded as the fine detail is only partially established from the coarse-resolution initial conditions
220 due to model spin-up ~~from the coarse-resolution initial conditions~~. Therefore, the hindcast fields between t_0+4h and t_0+9h
form the BARRA-C data sets. Such a hindcast length is considered short, but is chosen to meet computational constraints
~~with~~when regular reinitialization is needed for running the model for such an extended period. One clear limitation of our
setup is that model spin up artefacts are expected to be still present, particularly for convective clouds and rain.

3 Assessment

Our assessment focuses on near-surface variables and precipitation as the aim of BARRA-C is to capture small-scale local
225 weather phenomena ~~that which are is~~ most apparent near the surface. BARRA-C hindcasts are evaluated against point-scale
station observations for screen-level temperature, 10 m wind speed (Sec. 3.1) and precipitation (3.3). They are also compared
with gridded daily analyses of these observations for temperature (Sec. 3.2 and 3.6) and precipitation (Sec. 3.4 and 3.6).
Added skills in BARRA-C ~~are is~~ illustrated by comparing ~~these variables~~ against BARRA-R~~_~~and ERA-Interim hindcasts,
~~and against ERA5 hindcasts for precipitation~~and ERA5 hourly analyses ~~(ERA5 hindcasts only for precipitation)~~for the
~~other variables~~. To increase the diversity of models used in our inter-comparison, we also include the Modern-Era

230 Retrospective analysis for Research and Applications-2 (MERRA2, Gelaro et al., 2017) hindcasts. A scale-selective evaluation of extreme storms is conducted in Sec. 3.5 using radar observations available over the SY domain. Finally, an added value (AV) method is used to quantify improvements between BARRA-C and BARRA-R in the representation of extreme daily maximum and minimum temperature and daily rainfall from gridded observations. Readers are referred to Sec. A of the Supplement for details of the various reference data sets considered in our assessment.

235 **3.1 Point evaluation of screen temperature, 10 m wind speed, surface pressure**

The t_0+6h model hindcasts of screen-level temperature, 10 m wind speed, and surface pressure are evaluated against land station observations during the 2010-2012 period, following the approach of Su et al. (2019). These observations have no direct relation to BARRA-C, since there is no analysis in BARRA-C and they are not used in the associated BARRA-R cycle t_0 . These fields are interpolated from the model levels using surface similarity theory (Walters et al., 2017). Our benchmarks 240 include BARRA-R and ERA-Interim t_0+6h hindcasts, the MERRA2 hourly time-averaged hindcast fields, and the ERA5 hourly analysis. The models are interpolated to be coincident with the observed locations and times. As the observations are irregularly distributed in time, all observations within a t_0+5h to t_0+7h time window for $t_0 = 00$ and 12 UTC are considered. Root mean square difference (RMSD), Pearson's linear correlation, additive bias, and variance bias are calculated at each 245 station between observed (d_o) and model (d_m) data. Additive bias is defined as, with $Bias = E(d_m) - E(d_o)$, where $E(\cdot)$ is the expectation operator, and the variance bias as $Mbias = \frac{var(d_m)}{var(d_o)} - 1$, to capture differences in the dispersion, where $E(\cdot)$ is the expectation operator and so as to capture differences in the dispersion, where $var(\cdot)$ computes the variance in time. This assessment does not serve to inform the true quality of the various reanalyses at their native resolutions. rather, it indicates whether the models contain finer-scale information captured by point measurements. Based on Di Luca et al., 250 (2016), we distinguish three distinct regions with characteristics of complex topography (stations with an elevation higher than 500 m – *topo*), land-sea contrasts (stations that are within 1.5° of the coast – *coast*), or a relatively smooth terrain (stations far from the coast – *flat*) (Figure S3 in the Supplement).

The comparisons of scores across all BARRA-C domains are shown in Figure 2. For temperature, the BARRA (i.e., BARRA-R and BARRA-C) and ERA5 show better agreement with the station data than the other coarser reanalyses for most metrics. For instance, BARRA-C shows lower RMSD than ERA-Interim at 80% of stations. BARRA shows greater contrast 255 from the global reanalyses than between them. ERA5 shows warm (additive) bias, while the BARRA appears cooler. ERA-Interim and ERA5 generally show less variability in temperature than observations ($Mbias < 0$) while the other models tend to have more similar temperature variability with observations. This is related to the cold bias in ERA during high temperature (shown in the next section). On average, BARRA scores lower RMSD than ERA5 at elevated stations (e.g., Snowy Mountains in SY) and smaller $Mbias$ at near-coast stations. Similarly In general, BARRA-C shows more visible 260 improvements to BARRA-R at stations near coasts or over complex topography in terms of RMSD, correlation and $Mbias$ (Figure S3 in the Supplement). Consequently, BARRA-TA scores higher than BARRA-R on average. However, BARRA-C

shows higher RMSD in the flat regions than in the other regions, unlike the other reanalyses. The degradation is small, within 0.6 K in terms of RMSD, and for AD, this is related to over-dispersion (MBias > 1).

For 10 m wind speed, BARRA-C, BARRA-R and ERA5 similarly exhibit lower RMSD and higher correlation with the station data than the other global reanalyses, and the differences between these three models are not pronounced. BARRA's largest enhancement to ERA-Interim is found at elevated stations and near coasts, benefitting Tasmania ~~specifically~~particularly. Contrasting BARRA-R, BARRA-C tends to show lower RMSD at these stations (Figure S3, Supplement), and where we observe higher RMSD in BARRA-C, the difference is within 1 m/s. The wind estimated by all the models tends to be under-dispersed (MBias < 1), relating to positive (negative) bias during light (strong) wind conditions. Such a model under-dispersion is more striking in the TA and SY domains than in the other domains, and also over coastal regions.

~~For the~~For surface pressure, the higher resolution models, including ERA5, show markedly lower RMSD near coasts. There is very good agreement between ERA5 and the observations. ~~Some improvements to BARRA-R from BARRA-C shows some improvements over BARRA-R in~~ are mainly in correlation and Mbias, and over coastal regions and mountains.

3.2 Comparison with gridded analysis of daily maximum and minimum screen temperature

The reanalyses are compared against a gridded daily $0.05^\circ \times 0.05^\circ$ analysis of observed maximum and minimum screen temperature from the Australian Water Availability Project (AWAP; Jones et al., 2009) in Figure 3. BARRA outperforms the driving model ERA-Interim in reducing the cold (warm) bias during summer DJF (winter JJA), particularly over the SY and TA domains. ~~BARRA avoids the bias in ERA over the salt lakes in SA, by modelling them with land characteristics based on IGBP.~~ BARRA-C shows smaller extent of summer cold bias in daily maximum temperature over the Great Dividing Range than both BARRA-R and ERA5, but shares similar bias with BARRA-R elsewhere. BARRA and the global reanalyses also exhibit a considerable warm bias in the northwest of the AD domain—the Nullarbor Plain, ~~but this is likely an artefact of the AWAP station density and discussed later. In this region, AWAP analysis is generally poorer (Sec. A of the Supplement) due to very few observing stations (less than 2 per deg²).~~ The differences between BARRA and ERA reanalyses can also be related to the differences in their land cover classification based on IGBP and CCI.

The warm bias in daily minimum temperature in winter is also similar between BARRA-C and BARRA-R. BARRA-C has largely inherited the biases from BARRA-R, but with small local-scale differences. Despite such similarities in summer bias, ~~when comparing the number of hot days exceeding 35 °C (308.15 K) in Figure 3(c),~~ there are more hot days (i.e., days exceeding 35 °C (308.15 K)) in Figure 3(c) in BARRA-C than in BARRA-R over inland Australia. By contrast, the summer cold temperature bias in both ERA reanalyses is also reflected by fewer hot days, and vice versa for MERRA2. Further analysis of the temperature extremes is considered in Sec. 3.6.

Figure 4 and Figure 5 examine the inter-seasonal and inter-annual variations in temperature bias with respect to AWAP for daily maximum and minimum temperature respectively. They are similar between BARRA-C and BARRA-R,

Commented [CHS1]: Moved to discussion section.

Commented [CHS2]: Moved to discussion section.

Formatted: Indent: First line: 1.27 cm

295 with BARRA-C showing slightly wider inter-seasonal variability. The inter-seasonal range of bias in BARRA is around 2 K, which is similar to ERA-Interim and MERRA2 in most domains but is larger than ERA5 with the exception for TA. For AD and PH, the daily maximum temperature is positively biased during summer months (DJF) and is negatively biased during winter (JJA). The negative bias in daily maximum temperature is smallest during summer for SY and TA, and is largest during winter for SY. For daily minimum temperature, these are reversed; e.g., the associated positive bias peaks during 300 winter for AD, PH and SY, and the negative bias is maximum during summer for AD and PH.

~~There is both *i*The inter-annual variability and a trend of the temperature biases ~~do~~ exist~~ in BARRA. For daily maximum temperature bias, there is a cooling trend in AD and PH, and a warming trend in TA. These trends can also be seen in ERA5 and MERRA2. For daily minimum temperature bias, trends in BARRA are less apparent than in ERA5 and MERRA2. Here We also observe in the TA domain that BARRA shows a small warming trend with respect to AWAP.

305 This analysis of variability of bias is ~~also~~ repeated for the standard deviation of the modelled temperature and AWAP in Figure S4 and S5 of the Supplement. BARRA-C shows a slightly wider dispersion of daily maximum temperatures than AWAP (by 0.4 K) and BARRA-R (by 0.1 K), with the exception for the TA domain. For BARRA-TA, the standard deviation of BARRA is similar to AWAP and is higher than the global reanalyses. For daily minimum temperature, both BARRA are similar and ~~they are~~ generally under-dispersed by 0.3 K compared to AWAP.

Formatted: Indent: First line: 1.27 cm

310 3.3 Comparison with raingauges over Sydney

Hourly modelled precipitation from BARRA and ERA5 are compared against observations from 27 raingauges within 1° radius around Sydney during the warmer months (NDJF) in 2008-2013 in Figure 6. During these months, convection processes dominate and can produce a distinct diurnal distribution in thunderstorm activity, ~~with~~ The h ~~the~~ greatest frequency of severe thunderstorms ~~occurring occurs~~ in November and December (Griffiths, et al., 1993). ~~BARRA-R and ERA5 and for a lesser extent BARRA-R, both underestimate the frequency of heavy rain rate > 8 mm/h, with a lesser extent for BARRA-R.~~ By contrast, BARRA-C underestimates the frequency of light rain rate and overestimates heavy rates. BARRA and ERA5 also distribute rainfall differently over a day. BARRA-C shows a bimodal distribution similar to the observations, albeit showing too much rain leading up to the 06 UTC peak and too little rain during the daily minimum around 18 UTC. ~~The more pronounced diurnal cycle in precipitation is consistent with the over and under estimation of different rain rates.~~ 320 BARRA-R shows less diurnal variation in rainfall with too much rain distributed during 00-06 UTC, whereas ERA5 shows a pronounced early timing bias.

Commented [NE3]: Moved to discussion

3.4 Comparison with daily rainfall analysis

Figure 7(a) compares the modelled precipitation against daily raingauge analysis from AWAP, including MERRA2's hourly time-averaged precipitation (PRECTOTCORR) product. BARRA-C shows a wet bias over the Great Dividing Range and the southeast area of the AD domain, but improves the dry bias in BARRA-R and ERA reanalyses over the eastern and western seaboards ~~and, and~~ the Fleurieu and Yorke Peninsulas of South Australia. BARRA-C also shows dry biases on the western

borders of the AD and SY domains possibly due to inconsistencies with the zero-lateral moisture mass flux on the boundary conditions (Sect. 2.1). A striking difference between BARRA and the global reanalyses is over western Tasmania where the latter displays a dry bias.

330 ~~Next-in~~ In Figure 7(b), BARRA-R, ERA5 and ERA-Interim show too few heavy rain days (> 10 mm/day) over the coastlines, SA peninsula, and western Tasmania. BARRA-C improves on this, but generally simulates more heavy rain days than other reanalyses, and too few moderate-light rain days (< 10 mm/day, not shown) in all domains. BARRA-R and MERRA2 generally show too many light rain days, and the ERA reanalyses show too many light rain days in SY and eastern Tasmania, and too few in AD, PH, and western Tasmania.

335 The inter-seasonal and inter-annual variations in precipitation bias with respect to AWAP are plotted in Figure 8. As with temperature (Figure 4), they are similar between the BARRA-R and BARRA-C, although the latter shows a larger range in all the BARRA-C domains ~~but except~~ TA. In particular, the wet bias is generally observed during the wet season (JJA for AD, DJF for PH), wetter months (JJA for TA) or thunderstorm season (DJF for SY), and the dry bias generally occurs during the dry season or drier months, (i.e., e.g., SON for AD, PH and TA). This is consistent with the tendency of

340 BARRA-C to overestimate heavy rain rates and underestimate light rain occurrence. Some of the inter-annual variations in the bias are clearly common amongst BARRA and the global reanalyses; examples e.g., are in AD and PH domains during the Millennium drought (1996–2009) – where the various models are drier share a dry bias during the Millennium drought (1996–2009). BARRA ~~can~~ also shows different trends to the global reanalyses. For instance, There is a wetting trend post-2009 for BARRA in AD, but this is opposite for the other models. In SY, BARRA also displays a wetting trend, while ERA trends drier.

345 It should however be noted that, as is often found for gridded interpolated data, AWAP tends to underestimate the intensity of extreme rainfall events, and overestimate the frequency and intensity of low rainfall events (King et al., 2012). The errors are larger at high elevations (SY and TA) where gauges are fewer, and when there is frozen precipitation, and/or topography is exposed to prevailing winds (Chubb et al., 2016).

350 3.5 Storms over Sydney

The point gauge-based assessment in Sect. 3.3 is harsher to higher resolution models than coarser models, due to the compound error of space and time near-misses which increases as the grid cells shrink. Therefore, we compare the simulated rain-~~fields~~fall from BARRA-SY with the Bureau's radar nowcasting rainfall product (Rainfields2; Seed et al., 2007) using and use the fractions skill scores (FSS) to allow assessment at different spatial scales, following the approach described in Roberts and Lean (2007), Jermy and Renshaw (2016) and Acharya et al. (2020). The FSS provides an evaluation of the rainfall skill as a function of spatial resolution. The radar product, blended with gauge observations using conditional merging (Sinclair and Pegram, 2005), is available from 2014-onwards on a mosaic grid consisting of the domains of multiple radars. Following Acharya et al. (2020), the largest 36 storm events during 2014–2016 are selected based on domain-averaged daily precipitation.

360 FSS is categorised as a 'neighbourhood verification' metric (Ebert, 2009) in which fractional coverages of grid cells
close to observation are valued equally. The FSS tallies the relative number of 'hits' between the model and the observation
at different spatial scales and different rain thresholds. An FSS of 1 represents a perfect forecast where the number of cells
with precipitation above a threshold within a neighbourhood is identical between the model and observation grids for all
possible neighbourhoods. Here, BARRA hourly rain rates are regridded to the radar grid of 1.5 km ~~is~~ and the accumulated
365 rain amounts over moving 6-hour windows are analysed. From the 36 multi-day storm event set, 1323 different 6-hour
events are produced using a moving window. FSS is computed for each 6-hour event for each model and then the scores are
aggregated to give an average for all events. Given that inherent bias between the observation and the models exists due to
~~differences in~~ their representativity, ~~differences~~ and also to focus on the spatial accuracy of the models, we use percentile-
based thresholds computed across all the storm events. This ensures that the model and observed rain fields have an identical
370 fraction of rain events for each threshold value (see explained further in Sec. E of the Supplement). Figure 9 illustrates the
striking differences between ~~the~~ BARRA-R and BARRA-SY for five events in 2014. BARRA-SY can show more realistic
organisation in the 1.5 km model owing to the explicit modelling of convection and can produce higher rainfall intensity.
The event on 7 December 2014 in Figure 9(v) illustrates a summer storm case where BARRA-R shows rainfall
375 accumulations which lack the spatial pattern common ~~with to~~ convective organisation ~~which and are~~ evident in BARRA-SY
and in observations. BARRA-R also shows excessive grid-point precipitation over the mountains, which are absent in ~~both~~
observations and BARRA-SY. At the same time, BARRA-SY can show too many cells (Figure 9(ii)), which can produce
streaks of light rainfall (Figure 9(iv)).

380 The FSS results in Figure 10 show that BARRA-SY is more skilful over all scales than BARRA-R for all threshold
levels. $FSS_{uniform}$ is the FSS of a forecast field with a uniform fractional coverage equal to the fraction of points observed
with any rain (>0.2mm/hr). Scores greater than $FSS_{uniform}$ is considered skilful. For the lowest threshold (56%, i.e. 4 mm in
the observed radar values), the uniform score ($FSS_{uniform}$) is reached at scales of 0.3° (BARRA-SY), and 0.65° (BARRA-R).
At the highest threshold (99.9%, 64 mm), the uniform score is reached at scales of 2.4° and 3.35°, respectively. The contrast
385 between the two BARRA FSS~~s~~ is therefore greater at the higher precipitation thresholds. FSS_{s} for higher rainfall thresholds
~~is are~~ also generally lower as the area of rain being sampled becomes more localized and is more challenging to be
reproduced ~~correctly~~ in the models.

3.6 Added value analysis for temperature and rainfall extremes

390 We apply an approach similar to Di Luca et al. (2015) to quantify the added value (AV) in the representation of
climatological extremes from BARRA-C by comparing ~~the its skill between the BARRA-C and to the skill in~~ BARRA-R.
The warm extremes of daily maximum temperature, the cold extremes of daily minimum temperature and the wet extremes
of daily precipitation are assessed against AWAP, ~~noting that the true AV from BARRA-C at its native resolution is not~~
~~fully determined here.~~ The statistics for extremes (X) are given by the percentiles of the daily temperature and precipitation
values over the 29-year time period. We use $AV_d =$

Commented [CHS4]: Already noted in the Discussion.

$[d(X_{BARRA-R}, X_{AWAP}) - d(X_{BARRA-C}, X_{AWAP})]/[d(X_{BARRA-R}, X_{AWAP}) + d(X_{BARRA-C}, X_{AWAP})]$ of Di Luca et al. (2016) where d defines a distance metric between the model-derived and AWAP-derived statistics computed across the grid cells.

395 To capture both the total errors and spatial patterns of the statistics, we let $d \equiv MSE(A, B) = E[(A - B)^2]$ to define the mean squared error and also use $d \equiv Corr(A, B) = 1 - R(A, B)$ with R as Pearson's correlation. Larger positive AV values suggest smaller errors in BARRA-C than in BARRA-R and thus substantial added value by the downscaling of BARRA-R.

Figure 11 plots AV score for different BARRA-C domains, showing that AV is not gained consistently across the percentiles, variables and domains. For warm extremes of daily maximum temperature, BARRA-C shows positive AV_{MSE} over BARRA-R in the TA and AD domains. Low or negative AV_{MSE} for AD, PH and SY (inland region) is mainly due to its the warm and wet bias in BARRA-C seen, also seen in Figure 3(c) and 6(a,b). With The positive AV_{Corr} indicates that, BARRA-C captures the spatial patterns of the warm extremes across the domains, particularly over the coastal and high topography regions (see also Figure S6 of the Supplement).

405 For cold extremes in Figure 11(b), BARRA-C still shows positive AV_{MSE} over all domains except but the SY domain, due to AV over the coastal regions. This AV is mostly contributed by coastal regions as seen in Figure S6. Negative AV_{MSE} in SY is related to warmer cold extremes, particularly over the Great Dividing Range. The pPositive AV_{Corr} is seen in TA but not in the other domains. However, although it should be noted that the BARRAs are generally strongly correlated with AWAP with R mostly between 0.7 to 0.9.

410 AV from BARRA-C for wet extremes of precipitation relates more to the spatial patterns of the extremes (Figure 11(c)). Given the tendency of BARRA-C to overestimate heavy rainfall, the wet bias relative to AWAP, particularly over the PH domains (Figure 7(b)), is responsible for the low AV_{MSE}. There is negative AV_{MSE} for all domains except TA which remains near zero, highlighting the BARRA-C rainfall bias. On the other hand the AV_{Corr} is positive for all domains except AD for the highest three rainfall percentiles, which indicates better spatial correlation of rainfall than BARRA-R. For the SY domain, positive AV_{corr} for precipitation agrees with the above FSS analysis, which somewhat avoids the issue of bias through percentile-based thresholding. Assessing AV for wet extremes may also be problematic with AWAP. As an interpolated dataset, AWAP tends to underestimate the intensity of extreme heavy rainfall observed at stations and the issue is more pronounced at locations with sparse observational sampling or high topography, particularly in SY and TA (Chubb et al. 2016; King et al., 2012).

Formatted: Indent: First line: 1.27 cm

Commented [CHS5]: Moved to the Discussion section.

Commented [CHS6]: Moved to Discussion section.

Commented [CHS7]: Moved to discussion section.

4 Discussion

420 The BARRA-C 1.5 km models are strongly forced by BARRA-R with both initial conditions every 6 hours, and hourly boundary conditions. BARRA-C has therefore inherited much of the same quality of the BARRA-R, however, it The dynamical downscaling of BARRA's 12 km reanalysis, BARRA-R, with the BARRA-C 1.5 km models has been shown to does provide additional information about local near-surface meteorological conditions. BARRA-C provides better representative point-scale estimates of screen temperature, 10 m wind speed and surface pressure at some areas with

425 complex topography or near coastlines, and mainly inherits the skills of BARRA-R over other areas. The degradation from BARRA-R is slight, within (RMSD) 0.6 K for temperature and 1 m/s for wind speed.

BARRA-C also shows a ~~2–10~~ m wind speed bias that is positive (negative) ~~bias~~ during light (strong) wind conditions, similar to ~~the bias in~~ BARRA-R. Many factors such as boundary layer mixing, form drag for subgrid orography and surface properties can influence wind estimation over land. The representation of the stable boundary layer remains challenging due 430 to the multiplicity of physical processes ~~and their complex interactions, i.e., (including~~ turbulence, radiation, land surface coupling and heterogeneity ~~and,~~ turbulent orographic form drag) involved and their complex interactions, such that ~~models~~ Models typically suffer biases in 2 m temperature and wind speed under such conditions (Steeneveld, 2014 and references therein).

BARRA-C also inherits the domain-averaged biases in daily maximum and minimum temperature from BARRA-R. It 435 reduces some bias over the Great Dividing Range but simulates more hot days ~~than seen in observations~~—particularly over inland Australia. ~~However, in some inland regions the AWAP analyses are poorer quality due to low observing station density. For example, in the northwest of the AD domain - the Nullarbor Plain - both BARRA and the global models show a large warm bias in daily maximum temperature, however, the station density used in AWAP is less than 2 per deg² (Sec. A of the Supplement).~~

Commented [CHS8]: Moved here from the Results section.

440 The ~~daily temperature~~ bias varies ~~differently in time~~—between the four domains, with AD and PH showing a change of sign in bias between summer and winter months, while SY and TA show persisting negative (positive) bias for daily maximum (minimum) temperatures. Such similarities between the domains may be related to their similarities in terms of climate and land cover. Bush et al. (2020) discussed that changes in RAL1 for land surface representations (Table 1, Sec. 2.1) are important to improve the diurnal biases in pre-RAL1 configurations. These could benefit the biases seen over 445 vegetated areas, particularly for daily minimum temperature in SY and TA.

Commented [CHS9]: Moved here from the Results section.

450 The ~~reduced~~ dry bias of higher rain rates seen in the coarser scale models during the thunderstorm seasons in ~~the SY domain~~ is alleviated by BARRA-C. The underestimation of the peak rain rates in BARRA-R and ERA5 was expected from the lack of convection organisation due to the use of a cumulus parameterization, whereas BARRA-C evidently shows more realistic organization ~~and does not underestimate peak rain rates~~. However, the latter also exhibits too much heavy rain and not enough light rain ~~which is~~, likely due to the ~~still~~ under-resolved convection and the model's inability to resolve detrainment from convective updrafts. This is consistent with the findings reported in other studies. For example, Lean et al. 455 (2008) and Hanley et al. (2016) found that 1 km grid length UM simulations tend to produce cells that were too intense, too far apart and with not enough light rain. The latter also noted insufficient small storms in both shower cases and large storm cases, and too many large cells in shower cases.

460 The short hindcast length in BARRA-C (Sect. 2.2) poses a further limitation. The rainfall excess could result from model spin-up, as extra energy (i.e. CAPE) builds up during the early timesteps when there is insufficient convection, which is then finally released in the form of convective precipitation in later timesteps (Lean et al., 2008). Champion and Hodges (2014) have also noted that modelled precipitation intensities are most accurate when the model is initialised 12 hours before the rain maxima. The moisture conserving zero-lateral mass flux boundary conditions in BARRA-C exacerbate this issue. Moisture variables are not advected across boundaries and instead allowed to develop via physical processes in the model. These processes take some time to spin-up in each hindcast leading to near-boundary downstream moisture bias, e.g., 465 for example, the western boundary of the annual rainfall maps of AD and SY domains (Figure 7(a)). These issues of precipitation with short hindcasts can be improved with an assimilation system that will allow high resolution features to propagate from one hindcast cycle to the next (Dixon et al., 2008). In spite of these limitations, we find that BARRA-C provides a more representative rainfall climatology for heavy rain days near the coastal or and mountainous regions, and as well as sub-daily rain spatial patterns.

470 BARRA-C simulates peaks in the diurnal distribution of precipitation better than BARRA-R and ERA5. However, we also find that precipitation may be initiated too early and grow too rapidly. Consequently, BARRA-C under-represents off-peak rainrates resulting in an over-pronounced diurnal cycle, seen in Figure 4(b) for BARRA-SY in summer. This is contrary to the expectation for all models to initiate too late since subgrid-scale initial plumes cannot be represented. The early initiation bias in BARRA-R is due to the CAPE-based trigger mechanism of the convection scheme (Lean et al., 2008).

475 In the case of the kilometre-scale UM, the reasons are likely several. Hanley et al. (2015) partly attributed timing bias in convection initiation, which is too early in shower cases and too late in the larger storm cases, to unresolved convection at the kilometre-scale grid length. Other reasons may be that stochastic perturbations (Sect. 2.1) or model responses to the pre-convective profile are too strong, or that the profile has inadequate convective inhibition (CIN). The various aspects (intensity, size and timing) of simulated cells have shown to improve with adjustments to the mixing length used in the 480 subgrid turbulence scheme, but not all aspects improve simultaneously (Hanley et al., 2015).

There are trends and/or inter-annual variability of bias in BARRA against analyses of temperature and precipitation observations, and some of these trends are also apparent in the global reanalyses. BARRA-C largely mirrors the has similar bias variability in to BARRA-R, and its magnitude is of the order is similar or less than the global reanalyses. Spurious trends or artificial shifts in reanalyses could result from abrupt changes to the amount of satellite data assimilated, e.g., at the start 485 and end of satellite missions and or the various observational data archives. In BARRA-R, corrections were also made to the observation screening and thinning rules mid-production (Su et al., 2019). It is however However, it is outside of the scope of this work to assess the impacts of various observational changes.

BARRA-C shows better agreement with the pattern and the relative distribution of radar-derived rainfall during storms over Sydney. This improvement is, owing due to the use of explicit convection (Sect 2.1) and a higher resolution model and 490 this is consistent with earlier studies with UM (e.g., Lean et al., 2008). Comparisons of FSS from the same events including ERA5 show that its lower resolution leads to larger representation errors and lower FSS than BARRA-R despite both

parametrising convection (Figure S6, the Supplement). While BARRA-C still shows considerable bias compared to both rain gauges and radar observations, **BARRA-Cit** adds value to BARRA-R and ERA by providing more realistic and accurate spatial representations of rainfall during storms at various spatial scales and percentile thresholds.

495 The AV analysis of temperature and precipitation extremes shows that BARRA-C provides **only** some value over BARRA-R in various aspects including the spatial patterns of the warm temperature extremes and wet precipitation extremes and bias in cold extremes over coastal regions. -Low AV can be related to temperature and precipitation biases, which differ between the regions. **For example, the BARRA-C wet bias relative to AWAP, particularly over the PH domain (Figure 7(b)), is responsible for the low AV_{MSE} for rainfall. The positive AV_{corr} for precipitation in BARRA-SY agrees with the above FSS analysis, which somewhat avoids the wet bias issue through percentile-based thresholding.**

500 Assessing AV for wet extremes may also be problematic with AWAP. As an interpolated dataset, AWAP tends to underestimate the intensity of extreme heavy rainfall observed at stations and the issue is more pronounced at locations with sparse observational sampling or high topography, particularly in SY and TA (Chubb et al. 2016; King et al., 2012).

505 -While this analysis suggests that limited value is added by the downscaling of BARRA-R for these extremes, the true AV of BARRA-C at its native resolution is not assessed here given the limited resolution of AWAP and can be explored further with scale-dependent AV analysis of Di Luca et al. (2016). -Determining AV at the kilometre scale is also expected to be challenging as more accurate and representative observational data sets are needed. ▲

Commented [CHS10]: Moved here from the Results section.

Formatted: Font: (Asian) Times New Roman

Formatted: Font: (Asian) Liberation Serif

510 5 Conclusions

The recent development of CPMs in NWP has facilitated the creation of kilometre-scale regional reanalysis and climate projections. BARRA is the first regional reanalysis that focuses on the Australasian region, **which, It** has been developed with significant co-investment from state-level emergency service agencies across Australia. BARRA-C is the critical component of the project that provides these agencies with the means for developing a deeper understanding of past extreme 515 weather at local scales, especially in areas that were not adequately served by observation networks (e.g., Figure S3, the Supplement). The four mid-latitude domains of BARRA-C are designed to address these needs; and BARRA-R is needed to establish a driving model for BARRA-C **that is of higher resolution than ERA-Interim**, and **to** utilise more of the Australian local observations (Su et al., 2019). Completed in June 2019, the 29-year BARRA-R reanalysis (1990 to February 2019) and its downscaled counterparts BARRA-C₁ form a collection of high-resolution gridded meteorological datasets with 12 and 1.5 520 km horizontal grid lengths and 10 minutes to hourly time resolution, produced using systems closely related to the Bureau's present (as of October 2020) regional NWP systems. The hybrid model-level and pressure-level gridded data from BARRA-C are also available to drive/force sub-**kilometrekm** weather or non-weather models.

This paper describes the experimental configuration of BARRA-C and provides a preliminary assessment to illustrate its skills over BARRA-R and the global reanalyses at their subgrid scales. As expected from a hindcast-only system, it inherits the domain-averaged biases from BARRA-R. On the other hand, our added value analysis ~~however~~ shows that BARRA-C simulated different climatological extremes for temperature. Altogether, there exists added skill at the local-scale for temperature and wind, particularly in topographically complex regions in SY and TA, and coastal regions in all domains. As expected, the contrasts in skills and biases are most apparent between BARRA and the coarser-scale reanalyses (ERA-Interim, MERRA2). ~~The~~ BARRA-R and BARRA-C produce more distinctive precipitation estimates for intensity, 525 sub-daily timing and hourly spatial patterns that are characteristics of their physical schemes. BARRA-C also provides different spatial distribution of precipitation over complex terrains and more skilful representations of sub-daily rainfall fields. The latter suggests that BARRA-C is more suited for studies of extreme rainfall events, albeit ~~with still has a~~ high rainfall bias. The high rainfall bias ~~therefore also~~ manifests ~~as biases~~ in the climatological extremes of precipitation. These 530 findings highlight ~~that~~ improvements are still needed in ~~our for future~~ kilometre-scale downscaled reanalysis, ~~such as e.g.,~~ ~~adding kilometre-scale through~~ data assimilation and ~~further~~ model development. At this stage, BARRA-R and BARRA-C can be used conjunctively to improve individual estimates of temperature and precipitation. Some of their biases, including 535 for 10 m wind, could also be addressed via post-processing using multi-variate regression models or quantile matching methods such as those of Glahn and Lowry (1972), and Cattoën et al. (2020). Users of BARRA are strongly encouraged to undertake a local evaluation to ascertain the skills of BARRA-C for their regions and parameters of interest.

540 BARRA lays some of the important groundwork for future reanalysis-related activities and developing national climate risk services at the Bureau. Some of the issues identified in this work are being actively researched by collaborating national meteorological centres and academic institutions, within the “Regional Atmosphere” configuration development framework (Bush et al., 2020). Future reanalyses will also benefit from the recent advances in Bureau’s NWP, whereby an assimilation system (Rennie et al., 2020) and ensemble are introduced in its upcoming kilometre-scale models, to allow 545 propagation of high-resolution information between hindcast cycles and estimation of uncertainties.

Code availability

All code, including the UM (version 10.6) and JULES (version 4.7), used to produced BARRA-C is version-controlled under the Met Office Science Repository Service. The UM is available for use under license, <http://www.metoffice.gov.uk/research/modelling-systems/unified-model>. JULES is available under licence free of charge, 550 http://jules-lsm.github.io/access_req/JULES_access.html. The infrastructure for building and running UM-JULES simulations uses the Rose suite engine (<https://metomi.github.io/rose/doc/html/index.html>) and scheduling using the CycL work flow engine (<https://cylc.github.io/>, Oliver et al., 2019). Both Rose and CycL are available under Version 3 of the GNU General Public License. The BARRA-C Rose/CycL suite, with an identifier u-ak499, is version-controlled under the Met Office Science Repository Service and contains the UM-JULES science namelist and simulation configurations. Output from

555 the model simulations was converted from UM fieldsfile format to netCDF4 format using Iris (<https://scitools-iris.readthedocs.io/en/stable/><https://scitools.org.uk/iris/docs/latest/>).

Data availability

The BARRA datasets for the period of January 1990 to February 2019 are available for academic use. Readers are referred
560 to <http://www.bom.gov.au/research/projects/reanalysis> (last access: 31 August 2020; Bureau of Meteorology, 2020) for information on available parameters, access and licensing. The BARRA-R datasets used to initialise and constrain BARRA-C at the boundaries and the BARRA-C ancillary files can be requested by contacting the authors directly and are subject to the same licensing conditions.

Author contributions

565 PS, DJ, PFH and CJW conceived and/or designed BARRA. NE, CHS and PS developed the BARRA-C system with inputs from CF. NE performed the production, and CHS and NE the evaluation. CHS and NE prepared the paper with contributions from all co-authors.

Acknowledgments

Funding for this work was provided by emergency service agencies (New South Wales Rural Fire Service, Western Australia
570 Department of Fire and Emergency Services, South Australia Country Fire Service, South Australia Department of Environment, Water and National Resources) and research institutions (Antarctic Climate and Ecosystems Cooperative Research Centre (ACE CRC) and the University of Tasmania). Funding from Tasmania is supported by the Tasmanian Government and the Australian Government, provided under the Tasmanian Bushfire Mitigation Grants Program.

BARRA-C is set up with assistance from the UKMO colleagues (Stuart Webster) and many colleagues at the
575 Bureau of Meteorology (Greg Kociuba, Gary Dietachmayer, Hongyan Zhu, Yimin Ma, Ilia Bermous, Robin Bowen), the Commonwealth Scientific and Industrial Research Organisation (CSIRO; Martin Dix), and National Computational Infrastructure (NCI; Dale Roberts, Grant Ward). The FSS analysis with the Rainfields2 product is undertaken with assistance from Susan Rennie, Kevin Cheong, and Alan Seed (Bureau of Meteorology), and Suwash Acharya (University of Melbourne). We also thank Mitchell Black and Vinodkumar for their feedback on drafts of the paper. The BARRA project
580 was undertaken with the assistance of resources and services from NCI, which is supported by the Australian Government. This study uses the ERA-Interim and ERA5 data provided through the ARC Centre of Excellence for Climate System Science (Paola Petrelli) at NCI.

ERA-Interim can be retrieved from the ECMWF at <https://www.ecmwf.int/en/forecasts/datasets/archive-datasets/reanalysis-datasets/era-interim>. ERA5 can be retrieved from Copernicus Climate Data Store, at

585 <https://cds.climate.copernicus.eu/>. The AWAP data can be requested from <http://www.bom.gov.au/climate> (last access: 31 August 2020). The Rainfields2 radar product is retrieved from the Rainfields Archiving System provided by the Bureau of Meteorology.

References

Acharya, S. C., Nathan, R., Wang, Q. J., Su, C.-H., and Eizenberg, N.: Ability of an Australian reanalysis dataset to 590 characterise sub-daily precipitation, *Hydrol. Earth Syst. Sci.*, 24, 2951–2962, doi:10.5194/hess-24-2951-2020, 2020.

Arakawa, A. and Lamb, V. R.: Computational design of the basic dynamical processes of the UCLA general circulation model, *Methods of Comp. Phys.: Adv. Res. Appl.*, 17, 173–265, doi:10.1016/B978-0-12-460817-7.50009-4, 1977.

Argüeso, D., Evans, J.P., Fita, L., and Bormann, K. J.: Temperature response to future urbanization and climate change. *Clim Dyn* 42, 2183–2199. doi:10.1007/s00382-013-1789-6, 2014.

595 Bermejo, R., and Staniforth, A.: The conversion of semi-Lagrangian advection schemes to quasi-monotone schemes. *Mon. Wea. Rev.*, 120, 2622–2632, doi: 10.1175/1520-0493(1992)120<2622:TCOSLA>2.0.CO;2, 1992.

Best, M. J., Pryor, M., Clark, D. B., Rooney, G. G., Essery, R. L. H., Ménard, C. B., Edwards, J. M., Hendry, M. A., Porson, A., Gedney, N., Mercado, L. M., Sitch, S., Blyth, E., Boucher, O., Cox, P. M., Grimmond, C. S. B., and Harding, R. J.: The Joint UK Land Environment Simulator (JULES), model description – Part 1: Energy and water fluxes, *Geosci. Model Dev.*, 600 4, 677–699, doi:10.5194/gmd-4-677-2011, 2011.

Borsche, M., Kaiser-Weiss, A. K., Unden, P., and Kaspar, F.: Methodologies to characterize uncertainties in regional reanalyses, *Adv. Sci. Res.*, 12, 207–218, doi:10.5194/asr-12-207-2015, 2015.

Boutle, I. A., Abel, S. J., Hill, P. G., and Morcrette, C. J.: Spatial variability of liquid cloud and rain: observations and microphysical effects, *Q. J. R. Meteorol. Soc.*, 140, 583–594, doi:doi.org/10.1002/qj.2140, 2014a.

605 Boutle, I. A., Eyre, J. E. J., and Lock, A. P.: Seamless stratocumulus simulation across the turbulent gray zone, *Mon. Weather Rev.*, 142, 1655–1668, doi:10.1175/MWR-D-13-00229.1, 2014b.

Bromwich, D. H., A. B. Wilson, L. Bai, G. W. K. Moore, and P. Bauer: A comparison of the regional Arctic System Reanalysis and the global ERA-Interim Reanalysis for the Arctic. *Q. J. R. Meteorol. Soc.*, 142, 644-658, doi:10.1002/qj.2527, 2016.

610 Brousseau, P., Seity, Y., Ricard, D., and Léger, J.: Improvement of the forecast of convective activity from the AROME-France system. *Q.J.R. Meteorol. Soc.*, 142: 2231-2243. doi:10.1002/qj.2822, 2016.

Bureau of Meteorology: APS2 upgrade of the ACCESS-R numerical weather prediction system, NOC Operations Bulletin Number 107, <http://www.bom.gov.au/australia/charts/bulletins/apob107-external.pdf> (last access: 31 August 2020), 2016.

Bureau of Meteorology: APS2 upgrade of the ACCESS-C numerical weather prediction system, NOC Operations Bulletin 615 Number 114, http://www.bom.gov.au/australia/charts/bulletins/BNOC_Operations_Bulletin_114.pdf (last access: 31 August 2020), 2018.

Bureau of Meteorology: Atmospheric high-resolution regional reanalysis for Australia, available at: <http://www.bom.gov.au/research/projects/reanalysis>, last access: 1 May 2020.

Bush, M., Allen, T., Bain, C., Boulte, I., Edwards, J., Finnenkoetter, A., Franklin, C., Hanley, K., Lean, H., Lock, A., 620 Manners, J., Mittermaier, M., Morcrette, C., North, R., Petch, J., Short, C., Vosper, S., Walters, D., Webster, S., Weeks, M., Wilkinson, J., Wood, N., and Zerroukat, M.: The first Met Office Unified Model–JULES Regional Atmosphere and Land configuration, RAL1, Geosci. Model Dev., 13, 1999–2029, <https://doi.org/10.5194/gmd-13-1999-2020>, 2020.

Calmet, I., Mestayer, P. G., van Eijk, A. M. J., and Herlédant, O.: A coastal day summer breeze study, Part 2: High-resolution numerical simulation of sea-breeze local influences. Boundary-Layer Meteorol. 167, 27–51, doi:10.1007/s10546-625 017-0319-1, 2018.

Cattoën, C., D. E. Robertson, J. C. Bennett, Q. J. Wang, and T. K. Carey-Smith: Calibrating Hourly Precipitation Forecasts with Daily Observations. J. Hydrometeorol., 21, 1655–1673, doi:10.1175/JHM-D-19-0246.1, 2020.

Champion, A. J., and Hodges, K.: Importance of resolution and model configuration when downscaling extreme precipitation, Tellus A, 66, 23993. doi: doi:10.3402/tellusa.v66.23993, 2014.

630 Charney, J. G. and Phillips, N. A.: Numerical integration of the quasi-geostrophic equations for barotropic and simple baroclinic flows, J. Meteorol., 10, 71–99, doi:10.1175/1520-0469(1953)010<0071:NIOTQG>2.0.CO;2, 1953.

Chubb, T., Manton, M., Siems, S., & Peace, A. D.: Evaluation of the AWAP daily precipitation spatial analysis with an independent gauge network in the Snowy Mountains. Journal of Southern Hemisphere Earth Systems Science, 66(1), 55–67, doi:10.22499/3.6601.006, 2016.

635 Clark, P., Roberts, N., Lean, H., Ballard, S.P. and Charlton-Perez, C.: Convection-permitting models: a step-change in rainfall forecasting. Met. Apps, 23, 165–181, doi:10.1002/met.1538, 2016.

Davies, T., Cullen, M. J. P., Malcolm, A. J., Mawson, M. H., Staniforth, A., White, A. A., and Wood, N.: A new dynamical core for the Met Office's global and regional modelling of the atmosphere, Q. J. Roy. Meteorol. Soc., 131, 1759–1782, doi:10.1256/qj.04.101, 2005.

640 Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Holm, E. V., Isaksen, L., Kallberg, P., Kohler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J. J., Park, B. K., Peubey, C., de Rosnay, P., Tavolato, C., Thepaut, J. N.,

and Vitart, F.: The Era-Interim reanalysis: Configuration and performance of the data assimilation system, *Q. J. Roy. Meteorol. Soc.*, 137, 553–597, doi:10.1002/qj.828, 2011.

645 Dharssi, I., Steinle, P., and Fernon, J.: Improved numerical weather predictions by using optimised urban model parameter values and satellite derived tree heights, in: MODSIM2015, 21st International Congress on Modelling and Simulation, edited by: Weber, T., McPhee, M. J., and Anderssen, R. S., Modelling and Simulation Society of Australia and New Zealand, December 2015, 1161–1167, ISBN: 978-0-9872143-5-5, available at: <https://www.mssanz.org.au/modsim2015/M4/dharssi.pdf> (last access: 31 August 2020), 2015.

650 Di Luca, A., de Elía, R., and Laprise, R.: Challenges in the quest for added value of regional climate dynamical downscaling, *Curr. Clim. Change Rep.*, 1(1), 10–21, doi:10.1007/s40641-015-0003-9, 2015.

Di Luca, A., Argüeso, D., Evans, J. P., de Elía, R., and Laprise, R.: Quantifying the overall added value of dynamical downscaling and the contribution from different spatial scales, *J. Geophys. Res. Atmos.*, 121, 1575– 1590, doi:10.1002/2015JD024009, 2016.

655 Dixon, M., Z. Li, H. Lean, N. Roberts, and S. Ballard: Impact of data assimilation on forecasting convection over the United Kingdom using a high resolution version of the Met Office Unified Model. *Mon. Wea. Rev.*, 137, 1562–1584, doi:10.1175/2008MWR2561.1, 2009.

Done, J., Davis, C.A. and Weisman, M.: The next generation of NWP: explicit forecasts of convection using the weather research and forecasting (WRF) model. *Atmos. Sci. Lett.*, 5, 110–117. doi:10.1002/asl.72, 2004.

660 Donlon, C. J., Martin, M., Stark, J. D., Roberts-Jones, J., Fiedler, E., and Wimmer, W.: The Operational Sea Surface Temperature and Sea Ice analysis (OSTIA) system, *Remote Sens. Environ.*, 116, 140–158, doi:10.1016/j.rse.2010.10.017, 2012.

Ebert, E. E.: Neighborhood verification: A strategy for rewarding close forecasts. *Weather and Forecasting*, 24(6), 1498–1510. doi:10.1175/2009WAF222251.1, 2009.

665 Ebita, A., Kobayashi, S., Ota, Y., Moriya, M., Kumabe, R., Onogi, K., Harada, Y., Yasui, S., Miyaoka, K., Takahashi, K., Kamahori, H., Kobayashi, C., Endo, H., Soma, M., Oikawa, Y., and Ishimizu, T.: The Japanese 55-year reanalysis JRA-55: An interim report, *SOLA*, 7, 149–152, doi:10.2151/sola.2011-038, 2011.

Edwards, J. M. and Slingo, A.: Studies with a flexible new radiation code. I: Choosing a configuration for a largescale model, *Q. J. Roy. Meteorol. Soc.*, 122, 689–719, doi:10.1002/qj.49712253107, 1996.

670 Fosser, G., Khodayar, S., and Berg, P: Benefit of convection permitting climate model simulations in the representation of convective precipitation. *Clim. Dyn.* 44, 45–60, doi:10.1007/s00382-014-2242-1, 2015.

Frank, C. W., Pospichal, B., Wahl, S., Keller, J. D., Hence, A., and Crewell, S.: The added value of high resolution regional reanalyses for wind power applications, *Renewable Energy*, 148, 1094-1109, doi: 10.1016/j.renene.2019.09.138, 2020.

675 Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., da Silva, A. M., Gu, W., Kim, G., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S. D., Sienkiewicz, M., and Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2), *J. Climate*, 30, 5419–5454, doi:10.1175/JCLI-D-16-0758.1, 2017.

680 Gerard, L., J. Piriou, R. Brožková, J. Geleyn, and D. Banciu, 2009: Cloud and precipitation parameterization in a meso-gamma-scale operational weather prediction model. *Mon. Wea. Rev.*, 137, 3960–3977, doi:10.1175/2009MWR2750.1, 2009.

Glahn, H. R. and Lowry, D. A.: The use of model output statistics (MOS) in objective weather forecasting, *J. Appl. Meteor.*, 11, 1203–1211, doi:10.1175/1520-0450(1972)011<1203:TUOMOS>2.0.CO;2, 1972.

685 Gregory, D. and Rowntree, P. R.: A mass flux convection scheme with representation of cloud ensemble characteristics and stability-dependent closure, *Mon. Weather Rev.*, 118, 1483–1506, [https://doi.org/10.1175/1520-0493\(1990\)118<1483:AMFCSW>2.0.CO;2](https://doi.org/10.1175/1520-0493(1990)118<1483:AMFCSW>2.0.CO;2), 1990.

Gregow, H., K. Jylhä, H.M. Mäkelä, J. Aalto, T. Manninen, P. Karlsson, A.K. Kaiser-Weiss, F. Kaspar, P. Poli, D.G. Tan, A. Obregon, and Z. Su.: Worldwide survey of awareness and needs concerning reanalyses and respondents views on climate services. *Bull. Amer. Meteor. Soc.*, 97, 1461–1473, doi:10.1175/BAMS-D-14-00271.1, 2016.

690 Griffiths, D. J., Colquhoun, J. R., Batt, K. L., and Casinader, T. R.: Severe thunderstorms in New South Wales: Climatology and means of assessing the impact of climate change. *Climatic Change* 25, 369–388. doi:10.1007/BF01098382, 1993.

Halliwell, C.: Subgrid turbulence scheme, *Unified Model Documentation Paper 28*. Met Office: Exeter, UK, 2007.

Hanley, K.E., Plant, R.S., Stein, T.H.M., Hogan, R.J., Nicol, J.C., Lean, H.W., Halliwell, C. and Clark, P.A.: Mixing-length controls on high-resolution simulations of convective storms. *Q. J. R. Meteorol. Soc.*, 141: 272-284. doi:10.1002/qj.2356, 695 2015.

Hartley, A., MacBean, N., Georgievski, G., and Bontemps, S.: Uncertainty in plant functional type distributions and its impact on land surface models, *Remote Sens. Environ.*, 203, 71–89, doi:10.1016/j.rse.2017.07.037, 2017.

Hersbach, H., Bell, B., Berrisford, P., et al.: The ERA5 global reanalysis. *Q. J. R. Meteorol. Soc.*, 146: 1999– 2049, doi:10.1002/qj.3803, 2020.

700 Jakob, D., Su, C.-H., Eizenberg, N., Kociuba, G., Steinle, P., Fox-Hughes, P., and Bettio, L.: An atmospheric high-resolution regional reanalysis for Australia, *B. Aus. Meteorol. Oceanog. Soc.*, 30, 16–23, 2017.

Jermey, P. M. and Renshaw, R. J.: Precipitation representation over a two-year period in regional reanalysis. *Q. J. R. Meteorol. Soc.*, 142: 1300–1310. doi:10.1002/qj.2733, 2016.

Jones, D. A., Wang, W., and Fawcett, R.: High-quality spatial climate data-sets for Australia, *Aust. Meteorol. Oceanogr. J.*, 705 58, 233–248, 2009.

Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Leetmaa, A., Reynolds, R., Jenne, R., and Joseph, D.: The NCEP/NCAR 40-Year Reanalysis Project, *B. Am. Meteorol. Soc.*, 77, 437–472, doi:10.1175/1520-0477(1996)077%3C0437:TNYRP%3E2.0.CO;2, 1996.

710 Kendon, E. J., Ban, N., Roberts, N. M., Fowler, H. J., Roberts, M. J., Chan, S. C., Evans, J. P., Fosser, G., and Wilkinson, J. M.: Do convection-permitting regional climate models improve projections of future precipitation change? *Bull. Amer. Meteorol. Soc.*, 98, 79–93, doi:10.1175/BAMS-D-15-0004.1, 2017.

Kendon, E.J., Stratton, R.A., Tucker, S., Marsham, J. H., Berthou, S., Rowell, D. P., and Senior, C. A.: Enhanced future changes in wet and dry extremes over Africa at convection-permitting scale. *Nat Commun.* 10, 1794, doi:10.1038/s41467-715 019-09776-9, 2019.

Kendon, E. J., Prein, A. F., and Senior, C. A.: Challenges and outlook for convective-permitting climate modelling. *Phil. Trans. R. Soc. A.* 379, 20190547, doi:10.1098/rsta.2019.0547, 2021.

King, A.D., Alexander, L.V. and Donat, M.G.: The efficacy of using gridded data to examine extreme rainfall characteristics: a case study for Australia. *Int. J. Climatol.*, 33: 2376–2387, doi:10.1002/joc.3588, 2012.

720 Kuleshov, Y., de Hoedt, G., Wright, W., and Brewster, A.: Thunderstorm distribution and frequency in Australia, *Aust. Met. Mag.*, 51, 145–154, 2002.

Louis, J.-F.: A parametric model of vertical eddy fluxes in the atmosphere. *Boundary-Layer Meteorol.*, 17, 187–202, 1979.

Lean, H. W., Clark, P. A., Dixon, M., Roberts, N. M., Fitch, A., Forbes, R., Halliwell, C.: Characteristics of high-resolution versions of the Met Office Unified Model for forecasting convection over the United Kingdom. *Mon. Weather Rev.* 136: 725 3408–3424, doi: 10.1175/2008MWR2332.1, 2008.

Leutwyler, D., Lüthi, D., Ban, N., Fuhrer, O., and Schär, C.: Evaluation of the convection-resolving climate modeling approach on continental scales, *J. Geophys. Res. Atmos.*, 122, 5237–5258, doi:10.1002/2016JD026013, 2017.

Lock, A. P., Brown, A. R., Bush, M. R., Martin, G. M., and Smith, R. N. B.: A new boundary layer mixing scheme. Part I: Scheme description and single-column model tests, *Mon. Weather Rev.*, 128, 3187–3199, doi:10.1175/1520-730 0493(2000)128<3187:ANBLMS>2.0.CO;2, 2000.

Lock, A., Edwards, J., and Boutle, I.: The parametrization of boundary layer processes, Unified Model Documentation Paper 024, vn10.6, 2016.

Lopez, M. A., D. L. Hartmann, P. N. Blossey, R. Wood, C. S. Bretherton, and T. L. Kubar: A test of the simulation of tropical convective cloudiness by a cloud-resolving model. *J. Climate*, 22, 2834–2849, doi:10.1175/2008JCLI2272.1, 2009.

735 Louis, J.-F.: A parametric model of vertical eddy fluxes in the atmosphere, *Bound.-Layer Meteor.*, 17, 187–202, doi:10.1007/BF00117978, 1979.

Loveland, T. R., Reed, B. C., Brown, J. F., Ohlen, D. O., Zhu, Z., Yang, L., and Merchant, J. W.: Development of a global land cover characteristics database and IGBP DISCover from 1 km AVHRR data, *Int. J. Remote Sens.*, 21, 1303–1330, doi:10.1080/014311600210191, 2000.

740 Ma, Y., and Liu, H.: Large-eddy simulations of atmospheric flows over complex terrain using the immersed-boundary method in the Weather Research and Forecasting model. *Boundary-Layer Meteorol.* 165, 421–445, doi:10.1007/s10546-017-0283-9, 2017.

Mahmood, S., Davie, J., Jermy, P., Renshaw, R., George, J. P., Rajagopal, E. N., and Rani, S. I.: Indian monsoon data assimilation and analysis regional reanalysis: Configuration and performance. *Atmos. Sci. Lett.* 19:e808.

745 doi:10.1002/asl.808, 2018.

Mailhot, J., Bélair, S., Charron, M., Doyle, C., Joe, P., Abrahamowicz, M., Bernier, N. B., Denis, B., Erfani, A., Frenette, R., Giguère, A., Issac, G. A., McLennan, N., McTaggart-Cowan, R., Milbrandt, J., and Tong, L.: Environment Canada's experimental numerical weather prediction systems for the Vancouver 2010 Winter Olympic and Paralympic Games. *Bull. Amer. Meteor. Soc.*, 91, 1073–1086, doi:10.1175/2010BAMS2913.1, 2010.

750 Mesinger, F., DiMego, G., Kalnay, E., Mitchell, K., Shafran, P. C., Ebisuzaki, W., Jović, D., Woollen, J., Rogers, E., Berbery, E. H., Ek, M. B., Fan, Y., Grumbine, R., Higgins, W., Li, H., Lin, Y., Manikin, G., Parrish, D., and Shi, W.: North American Regional Reanalysis. *B. Am. Meteorol. Soc.*, 87, 343–360, doi:10.1175/BAMS-87-3-343, 2006.

Mueller, N., Lewis, A., Roberts, D., Ring, S., Melrose, R., Sixsmith, J., Lymburner, L., McIntyre, A., Tan, P., Curnow, S., and Ip, A.: Water observations from space: Mapping surface water from 25 years of Landsat imagery across Australia, *Rem.*

755 *Sens. Env.*, 174, 341–352, doi: 10.1016/j.rse.2015.11.003, 2016.

Oliver, H., Shin, M., Matthews, D., Sanders, O., Bartholomew, S., Clark, A., Fitzpatrick, B., van Haren, R., Hut, R., and Drost, N.: Workflow automation for cycling systems, *Comp. in Sci. & Eng.*, 21(4), 7-21, doi:10.1109/MCSE.2019.2906593, 2019.

Peel, M. C., Finlayson, B. L., and McMahon, T. A.: Updated world map of the Köppen-Geiger climate classification, 760 *Hydrol. Earth Syst. Sci.*, 11, 1633–1644, doi:10.5194/hess-11-1633-2007, 2007.

Prein, A. F., Langhans, W., Fosser, G., Ferrone, A., Ban, N., Goergen, K., Keller, M., Tölle, M., Gutjahr, O., Feser, F., Brisson, E., Kollet, S., Schmidli, J., Van Lipzig, N. P. M., and Leung, R.: A review on regional convection-permitting climate modeling: Demonstrations, prospects, and challenges, *Rev. Geophys.*, 53, 323–361. doi:10.1002/2014RG000475, 2015.

Puri, K., Dietachmayer, G., Steinle, P., Dix, M., Rikus, L., Logan, L., Naughton, M., Tingwell, C., Xiao, Y., Barras, V.,
765 Bermous, I., Bowen, R., Deschamps, L., Franklin, C., Fraser, J., Glowacki, T., Harris, B., Lee, J., Le, T., Roff, G., Sulaiman, A., Sims, H., Sun, X., Sun, Z., Zhu, H., Chattopadhyay, M. and Engel, C.: Implementation of the initial ACCESS numerical weather prediction system, *Aust. Meteorol. Oceanogr. J.*, 63, 265–284, 2013.

Rennie, S., L. Rikus, N. Eizenberg, P. Steinle, and M. Krysta: Impact of Doppler radar wind observations on Australian high-resolution numerical weather prediction. *Wea. Forecasting*, 35, 309–324, doi:10.1175/WAF-D-19-0100.1, 2020.

770 Roberts-Jones, J., Fiedler, E. K., and Martin, M. J.: Daily, global, high-resolution SST and sea ice reanalysis for 1985–2007 Using the OSTIA system, *J. Climate*, 25, 6215–6232, doi:10.1175/JCLI-D-11-00648.1, 2012.

Roberts, N. M., and Lean, H. W.: Scale-selective verification of rainfall accumulations from high-resolution forecasts of convective events, *Mon. Weath. Rev.*, 136, 78–97, doi:10.1175/2007MWR2123.1, 2007.

Schär, C., Fuhrer, O., Arteaga, A., Ban, N., Charpiloz, C., Di Girolamo, S., Hentgen, L., Hoefler, T., Lapillonne, X.,
775 Leutwyler, D., Osterried, K., Panosetti, D., Rüdisühli, S., Schlemmer, L., Schulthess, T. C., Sprenger, M., Ubbiali, S., and Wernli, H.: Kilometer-scale climate models: Prospects and challenges. *Bull. Amer. Meteor. Soc.*, 101, E567–E587, doi:10.1175/BAMS-D-18-0167.1, 2020.

Seed, A., Duthie, E., and Chumchean, S.: Rainfields: The Australian Bureau of Meteorology system for quantitative precipitation estimation. *Proceedings of the 33rd AMS Conference on Radar Meteorology*, Cairns, Australia, 2007.

780 Simard, M., Pinto, N., Fisher, J. B., and Baccini, A.: Mapping forest canopy height globally with spaceborne lidar, *J. Geophys. Res.-Biogeosci.*, 116, G04021, <https://doi.org/10.1029/2011JG001708>, 2011.

Sinclair, S. and Pegram, G.: Combining radar and rain gauge rainfall estimates using conditional merging, *Atmos. Sci. Lett.*, 6, 19–22, doi:10.1002/asl.85, 2005.

785 Smagorinsky, J.: General circulation experiments with the primitive equations. I: The basic experiment. *Mon. Weather Rev.* 91: 99–164, 1963.

Smith, R. N. B.: A scheme for predicting layer cloud and their water content in a general circulation model, *Q. J. R. Meteorol. Soc.*, 116, 435–460, doi:10.1002/qj.49711649210, 1990.

Steenneveld, G.-J.: Current challenges in understanding and forecasting stable boundary layers over land and ice, *Front. Environ. Sci.*, 2, doi: 10.3389/fenvs.2014.00041, 2014.

790 Stein, T. H. M., R. J. Hogan, P. A. Clark, C. E. Halliwell, K. E. Hanley, H. W. Lean, J. C. Nicol, and R. S. Plant: The DYMECS Project: A statistical approach for the evaluation of convective storms in high-resolution NWP models. *Bull. Amer. Meteor. Soc.*, 96, 939–951, doi:10.1175/BAMS-D-13-00279.1, 2015.

795 Su, C.-H., Eizenberg, N., Steinle, P., Jakob, D., Fox-Hughes, P., White, C. J., Rennie, S., Franklin, C., Dharssi, I., and Zhu, H.: BARRA v1.0: the Bureau of Meteorology Atmospheric high-resolution Regional Reanalysis for Australia, *Geosci. Model Dev.*, 12, 2049–2068, doi:10.5194/gmd-12-2049-2019, 2019.

800 Vitolo, C., Napoli, C. D., Giuseppe, F. D., Cloke, H. L., and Pappenberger, F.: Mapping combined wildfire and heat stress hazards to improve evidence-based decision making, *Environment International*, 127, 21–34, doi:10.1016/j.envint.2019.03.008, 2019.

Wahl, S., Bollmeyer, C., Crewell, S., Figura, C., Friederichs, P., Hense, A., Keller, J. D., and Ohlwein, C.: A novel 800 convective-scale regional reanalysis COSMO-REA2: Improving the representation of precipitation, *Meteorologische Zeitschrift*, 26(4), 345–361, doi:10.1127/metz/2017/0824, 2017.

Walsh, K. J. E., White, C. J., McInnes, K. L., Holmes, J., Schuster, S., Richter, H., Evans, J. P., Di Luca, A. and Warren, R.A., Natural hazards in Australia: storms, wind and hail, *Climatic Change*, 139, 55–67, doi:10.1007/s10584-016-1737-7, 2016.

805 Walters, D., Boutle, I., Brooks, M., Melvin, T., Stratton, R., Vosper, S., Wells, H., Williams, K., Wood, N., Allen, T., Bushell, A., Copsey, D., Earnshaw, P., Edwards, J., Gross, M., Hardiman, S., Harris, C., Heming, J., Klingaman, N., Levine, R., Manners, J., Martin, G., Milton, S., Mittermaier, M., Morcrette, C., Riddick, T., Roberts, M., Sanchez, C., Selwood, P., Stirling, A., Smith, C., Suri, D., Tennant, W., Vidale, P. L., Wilkinson, J., Willett, M., Woolnough, S., and Xavier, P.: The Met Office Unified Model Global Atmosphere 6.0/6.1 and JULES Global Land 6.0/6.1 configurations, *Geosci. Model Dev.*, 810 10, 1487–1520, <https://doi.org/10.5194/gmd-10-1487-2017>, 2017.

Walters, D., Baran, A. J., Boutle, I., Brooks, M., Earnshaw, P., Edwards, J., Furtado, K., Hill, P., Lock, A., Manners, J., Morcrette, C., Mulcahy, J., Sanchez, C., Smith, C., Stratton, R., Tennant, W., Tomassini, L., Van Weverberg, K., Vosper, S., Willett, M., Browne, J., Bushell, A., Carslaw, K., Dalvi, M., Essery, R., Gedney, N., Hardiman, S., Johnson, B., Johnson, C., Jones, A., Jones, C., Mann, G., Milton, S., Rumbold, H., Sellar, A., Ujiie, M., Whitall, M., Williams, K., and Zerroukat, M.: 815 The Met Office Unified Model Global Atmosphere 7.0/7.1 and JULES Global Land 7.0 configurations, *Geosci. Model Dev.*, 12, 1909–1963, <https://doi.org/10.5194/gmd-12-1909-2019>, 2019.

Wilkinson, J.M. and Jorge Bornemann, F.: A lightning forecast for the London 2012 Olympics opening ceremony. *Weather*, 69, 16–19. doi:10.1002/wea.2176, 2014.

820 Wilson, D. R. and Ballard, S. P.: A microphysically based precipitation scheme for the UK Meteorological Office Unified Model, *Q. J. Roy. Meteorol. Soc.*, 125, 1607–1636, doi:10.1002/qj.49712555707, 1999.

Wood, N., Staniforth, A., White, A., Allen, T., Diamantakis, M., Gross, M., Melvin, T., Smith, C., Vosper, S., Zerroukat, M., and Thuburn, J.: An inherently mass-conserving semi-implicit semi-Lagrangian discretization of the deep-atmosphere global nonhydrostatic equations, *Q. J. Roy. Meteorol. Soc.*, 140, 1505–1520, doi:10.1002/qj.2235, 2014.

Zerroukat, M. and Shipway, B. J.: ZLF (Zero Lateral Flux): a simple mass conservation method for semi-Lagrangian-based limited-area models. *Q.J.R. Meteorol. Soc.*, 143, 2578-2584, doi:10.1002/qj.3108, 2017.




Figure 1: Domains of BARRA-C, (left to right) BARRA-PH (over Perth), BARRA-AD (Adelaide), BARRA-TA (Tasmania), and BARRA-SY (Sydney), showing the modelled orography. Red circles indicate the locations of the state capital cities.




Figure 2: Box plots showing the distribution of evaluation scores of various models for (a) screen-level temperature, (b) 10 m wind speed, and (c) surface pressure across the four BARRA-C domains. Three regions are analysed separately: coastal ('coast'), complex topography ('topo'), and flat, and the models distinguished by colours. The scores are calculated on model hindcasts valid between 05-07 UTC, and 17-19 UTC against observations during 2010–2012.




Figure 3: Mean difference in (a) summer (DJF) daily maximum temperature, (b) winter (JJA) daily minimum temperature and (c) number of days with temperature exceeding 35 °C, in various models during 1990-2018, with respect to AWAP. The models are regressed onto the AWAP grid using the nearest-neighbour interpolation.





Figure 4: Timeseries of monthly mean difference in daily maximum temperature averaged over various BARRA-C domains, with respect to AWAP. The timeseries are shaded around their individual 1990-2018 means.



855 Figure 5: As with Figure 4, but for daily minimum temperature.




Figure 6: Distribution of (a) hourly rain rate (mm/h) and (b) rain over 24 hours in UTC over Sydney during November to February of 2006-2018.




Figure 7: Mean difference in (a) annual precipitation and (b) annual count of wet days with depth ≥ 10 mm. The models are regressed onto the AWAP grid using the nearest-neighbour interpolation.





Figure 8: Mean difference in seasonal precipitation totals over various BARRA-C domains, with respect to AWAP. Black curves are shaded around the 1990-2018 means. Note that the y-axes in (a)-(d) are different.



870 Figure 9: Simulated 6-hour rainfall accumulation [mm] in BARRA-SY and BARRA-R, compared with rainfall derived from the
 871 composite radar network in-the-round the Sydney area for five events.




Figure 10: Aggregated FSS across 1323 6-hour storm events as a function of neighbourhood distance (degrees) for 6-hour rainfall above three percentile thresholds (distinguished by colours, percentile values, and observed amount in mm). The solid curves indicate the score for BARRA-SY, dotted curves for BARRA-R, and the dashed horizontal lines the uniform score ($FSS_{uniform}$) for each threshold as specified by Roberts and Lean (2007).




Figure 11: Added value (AV) analysis of the (a) warm extreme of daily maximum temperature, (b) cold extreme of daily minimum temperature, and (c) wet extreme of daily precipitation, performed for different all four BARRA-C domains.

880

885

Table 1: An overview of major differences between BARRA-C, BARRA-R and the mid-latitude version of RAL1 (RAL1-M) models. The configurations for BARRA-R are described in Su et al. (2019) and Walters et al. (2017), and those for RAL1-M in Bush et al. (2020).

Aspects	BARRA-R	BARRA-C	RAL1-M
Nesting setup	Nested in 6-hourly ERA-Interim boundary conditions	Nested in hourly BARRA-R boundary conditions	NA
Horizontal grid length in radial resolution	0.11°	0.0135°	0.0135 to 0.04°
Vertical model level set	70 levels, with 50 levels below 18 km, and 20 levels above this, fixed model lid of 80 km above the sea level.	70 levels, with 61 levels below 18 km, 9 levels above this, fixed model lid of 40 km above sea level	
Model timestep	300 seconds	60 seconds	60-100 seconds, depending on the model resolution
UM model version	10.2	10.6	≥ 10.6

← Formatted Table

JULES model version	3.0	4.7	≥ 4.8
Data assimilation	6-hourly 4D variational analysis	None	NA
Moisture variable SL advection schemes	Quasi-monotone (Bermejo and Staniforth, 1992)		Posteriori monotonicity filter (PMF)
Convective parameterization scheme	Mass-flux convection scheme of Gregory and Rowntree (1990)		None
Gaseous absorption (radiation) scheme	GA6 (Walters et al., 2017)		GA7 (Walters et al., 2019)
Include spectral land-surface albedo	No		Yes
Canopy radiation back-scatter scheme	Isotropic		Anisotropic
Cloud microphysics scheme	Single moment scheme based on Wilson and Ballard (1999)	Wilson and Ballard (1999), with prognostic graupel (Wilkinson and Bornemann, 2014) and improved warm rain scheme (Boutle et al., 2014a)	
Boundary layer scheme	1D vertical turbulent mixing scheme of Lock et al. (2000)	Blended boundary layer parameterization (Boutle et al., 2014b)	
Land surface and hydrology	GA6 (Walters et al., 2017), PDM subgrid-scale heterogeneity, JULES urban parameters are optimized for Australia (Dharsri et al., 2015)		
BL stochastic perturbations	None	Perturbation to temperature	Perturbation to temperature and moisture
BL stability functions	For stable BL, the “sharp” function of Lock et al. (2016) is used over the sea, and over land is a blended combination of the Louis (1979) and the “sharpest” function, for heights below 200 m. The convective BL stability functions are based on UKMO Large-Eddy model simulations.		
Critical relative humidity profile	0.92 in the lowest layer, with a gradual decrease to 0.8 at model level 17 (~2100 km).		