Grid-independent High Resolution Dust Emissions (v1.0) for Chemical Transport Models: Application to GEOS-Chem (version-12.5.0)

Jun Meng^{1,2}, Randall V. Martin^{2,1,3}, Paul Ginoux⁴, Melanie Hammer^{2,1}, Melissa P. Sulprizio⁵,

5 David A. Ridley⁶, Aaron van Donkelaar^{1,2}

¹Department of Physics and Atmospheric Science, Dalhousie University, Halifax, Nova Scotia, B3H 4R2, Canada ² Department of Energy, Environmental & Chemical Engineering, Washington University in St.

Louis, St. Louis, Missouri 63130, United States

10 ³Smithsonian Astrophysical Observatory, Harvard-Smithsonian Center for Astrophysics, Cambridge, MA 02138, USA

⁴NOAA Geophysical Fluid Dynamics Laboratory, Princeton, New Jersey 08540, USA

⁵School of Engineering and Applied Science, Harvard University, Cambridge, MA 02138, USA

⁶California Environmental Protection Agency, Sacramento, CA 95814, USA

15

Correspondence to: Jun Meng (jun.meng@ucla.edudal.ca)

Abstract. The nonlinear dependence of the dust saltation process on wind speed poses a

- 20 challenge for models of varying resolutions. This challenge is of particular relevance for the next generation of chemical transport models with nimble capability for multiple resolutions. We develop and apply a method to harmonize dust emissions across simulations of different resolutions by generating offline grid independent dust emissions driven by native high resolution meteorological fields. We implement into the GEOS-Chem chemical transport model
- 25 a high resolution dust source function to generate updated offline dust emissions. <u>These</u> <u>updated offline dust emissions based on high resolution meteorological fields strengthen dust</u> <u>emissions over relatively weak dust source regions</u><u>These updated offline dust emissions based</u> <u>on high resolution meteorological fields can better resolve weak dust source regions</u>, such as in southern South America, southern Africa and the southwestern United States. Identification of
- 30 an appropriate dust emission strength is facilitated by the resolution independence of offline emissions. We find that the performance of simulated aerosol optical depth (AOD) versus measurements from the AERONET network and satellite remote sensing improves significantly when using the updated offline dust emissions with the total global annual dust emission strength of 2,000 Tg yr⁻¹ rather than the standard online emissions in GEOS-Chem. The offline
- 35 high resolution dust emissions are easily implemented in chemical transport models. The source code isand_available online through GitHub: https://github.com/Jun-Meng/geos-chem/tree/v11-01-Patches-UniCF-vegetation. The global offline high_-resolution dust emission inventory are publicly is freely available.e (see Code and Data Availability section).

1 Introduction

Mineral dust, as one of the most important natural aerosols in the atmosphere, has significant impacts on weather and climate by absorbing and scattering solar radiation (Bergin et al., 2017;

- 45 Kosmopoulos et al., 2017), on atmospheric chemistry by providing surfaces for heterogeneous reaction of trace gases (Chen et al., 2011; Tang et al., 2017), on the biosphere by fertilizing the tropical forest (Bristow et al., 2010; Yu et al., 2015), and on human health by increasing surface fine particulate matter (PM_{2.5}) concentrations (De Longueville et al., 2010; Fairlie et al., 2007; Zhang et al., 2013). Dust emissions are primarily controlled by surface wind speed to the third
- or fourth power, vegetation cover and soil water content. The principal mechanism for natural dust emissions is saltation bombardment (Gillette and Passi, 1988; Shao et al., 1993), in which sand-sized particles creep forward and initiate the suspension of smaller dust particles when the surface wind exceeds a threshold. The nonlinear dependence of dust emissions on meteorology ity of this process introduces an artificial dependence of simulations upon model
 resolution (Ridley et al., 2013). For example, dust emissions in most numerical models are parameterized with an empirical method (e.g. Ginoux et al., 2001; Zender et al., 2003), which

requires a critical wind threshold to emit dust particles. Smoothing meteorological fields to

coarse resolution can lead to wind speeds falling below the emission threshold in regions that

do emit dust. Methods are needed to address the artificial dependence of simulations upon

60 model resolution that arises from nonlinearity in dust emissions.

Addressing this nonlinearity is especially important for the next generation of chemistry transport models that is emerging with nimble capability for a variety of resolutions at the

global scale. For example, the high performance version of GEOS-Chem (GCHP) (Eastham et al., 2018) currently offers simulation resolutions that vary by over a factor of 100 from C24 (~

- 4°x45°) to C360 (~0.25°x0.25°), with progress toward even finer resolution and toward a variable stretched grid capability_(Bindle et al., 2020). Resolution-dependent mineral dust emissions would vary by a factor of 3 from C360 to C24, and inhibit interpretation (Ridley et al., 2013). Such large resolution-dependent biases would undermine applications of CTMs to assess dust effects, and would lead to large within-simulation inconsistency for stretched grid
- 70 <u>simulations that can span the entire resolution range simultaneously.</u> Grid-independent high resolution dust emissions offer a potential solution to this <u>concernissue</u>.

An important capability in global dust evaluation is ground-based and satellite remote sensing. The Aerosol Robotic Network (AERONET), a global ground-based remote sensing aerosol monitoring network of Sun photometers (Holben et al., 1998), has been widely used to

- 75 evaluate dust simulations. Satellite remote sensing provides additional crucial information across arid regions where in-situ observations are sparse (Hsu et al., 2013). Satellite aerosol retrievals have been used extensively in previous studies to either evaluate the dust simulation (Ridley et al., 2012, 2016) or constrain the dust emission budget (Zender et al., 2004). Satellite aerosol products have been used to identify dust sources worldwide (Ginoux et al., 2012;
- 80 Schepanski et al., 2012; Yu et al., 2018), especially for small-scale sources (Gillette, 1999).

The objective of this study is to develop a method to mitigate the <u>large</u> inconsistency of total dust emissions across different resolutions of simulations by generating and archiving_an offline dust emissions using native high resolution meteorological fields. We apply this method to the GEOS-Chem chemical transport model. As part of this effort, we implement an updated

85 high resolution satellite-identified dust source function into the dust mobilization module of GEOS-Chem to better represent the spatial structure of dust sources. We apply this new capability to assess the source strength that best represents observations.

2 Materials and Methods

90 2.1 Description of Observations

We use both ground-based and satellite observations to evaluate our GEOS-Chem simulations. AERONET is a global ground-based remote sensing aerosol monitoring network of sun photometers with direct sun measurements every 15 minutes (Holben et al., 1998). We use Level 2.0 Version 3 data that has improved cloud screening algorithms (Giles et al., 2019).

95 Aerosol optical depth (AOD) at 550 nm is interpolated based on the local <u>Aangstrom exponent</u> at the 440 nm and 670 nm channels.

Twin Moderate-Resolution Imaging Spectroradiometer (MODIS) instruments aboard both on the Terra and Aqua NASA satellite platforms and provide near daily measurements globally. We use the AOD at 550 nm retrieved from Collection 6.1 (C6) of MODIS product (Sayer et al., 2014). We use AOD from the Deep Blue (DB) retrieval algorithm (Hsu et al., 2013; Sayer et al., 2014) designed for bright surfaces, and the Multi-Angle Implementation of Atmospheric Correction (MAIAC) algorithm (Lyapustin et al., 2018), which provides global AOD retrieved from MODIS C6 radiances at a resolution of 1 km. The MAIAC AOD used in this study is interpolated to the AOD value at 550 nm.

 105
 We use ground-based surface fine dust concentration measurements over the US from

 the Interagency Monitoring of Protected Visual Environments (IMPROVE,

http://views.cira.colostate.edu/fed/DataWizard/) network. The IMPROVE network provides 24hr average fine dust concentrations data every third day over the national parks in the United States. We also include a climatology of dust surface concentrations measurements over 1981-

- 110 2000 from independent dust measurement sites over the globe (Kok et al. 2020). We use those sites (12 in total) (Figure S1) that either in the dust belt across Northern Hemisphere or sites relatively close to the weak emission regions in the Southern Hemisphere to evaluate our dust simulation. (Kok et al., 2020)
- We compare the simulated fine-AOD and dust concentrations with measurements using reduced major axis linear regression. We report root mean square error (E), correlation (R) and slope (M).

2.2 Dust mobilization module

We use the dust <u>endetrainment and deposition (DEAD) scheme (Zender et al., 2003) in the</u>
 GEOS-Chem model to calculate dust emission<u>s</u>. The saltation process is dependent on the critical threshold wind speed, which is determined by surface roughness, soil type and soil moisture. We use a fixed soil clay fraction of 0.2 as suggested in Zender et al. (2003). Dust aerosol is transported in four size bins (0.1-1.0, 1.0-1.8, 1.8-3.0, and 3.0-6.0 µm radius). Detailed description of the dust emission parameterization is in Sect. S1 of the supplemental material.

The fractional area of land with erodible dust is represented by a source function. The dust source function used in the dust emission module plays an important role in determining the spatial distribution of dust emissions. The standard GEOS-Chem model (version 12.5.0) uses

a source function at 2° x 2.5° resolution from Ginoux et al. (2001) as implemented by Fairlie et

- 130 al. (2007). We implement an updated high resolution version of the dust source function in this study at 0.25° x 0.25° resolution <u>(Sect. S2)</u>. Figure S24 shows a map of the original and updated version of the dust source function. The updated source function exhibits more spatially resolved information due to its finer spatial resolution resulting in a higher fraction of erodible dust over in the eastern Arabian Peninsula, the Bodélé depression, and the central Asian
- deserts. The dust module dynamically applies this source function, together with information
 on soil moisture, vegetation, and land use to calculate hourly emissions using the <u>Harmonized</u>
 <u>Emissions Component (-HEMCO)</u> module described below.

2.3 Offline dust emissions at the native meteorological resolution

- HEMCO (Keller et al., 2014) is a stand-alone software module for computing emissions in global atmospheric models. We run the HEMCO standalone version using native meteorological resolution (0.25° x 0.3125°) data for wind speed, soil moisture, vegetation, and land use to archive the offline dust emissions at the same resolution as the meteorological data. The computational time required for calculating offline dust emission fluxes at 0.25° x 0.3125°
 resolution is around 6 hours for one-year of offline dust emissions on a compute node with 32 cores on 2 Intel CPUs at 2.1 GHz. (Graham CC Doc, 2021) In this study, we generate two offline dust emission datasets at 0.25° x 0.3125° resolution. One, referred to as the default offline dust emissions, uses the existing dust source function in the GEOS-Chem dust module; the other, referred to as the updated offline dust emissions, uses the updated dust source function
- 150 implemented here. Both datasets are at the hourly resolution of the parent meteorological

fields. The archived native resolution offline dust emissions can be <u>conservatively regridded to</u> <u>coarser resolution for consistent input to</u> <u>-an input emission inventory for</u> chemical transport models <u>with scalable dust source strengthsat multiple resolutions</u>. We use the GEOS-Chem model to evaluate the dust simulations and the emission strength.

155

2.4 GEOS-Chem chemical transport model and simulation configurations

GEOS-Chem (The International GEOS-Chem User Community, 2019) is a three-dimensional chemical transport model driven by assimilated meteorological data from the Goddard Earth Observation System (GEOS) of the NASA Global Modelling and Assimilation Office (GMAO). The

- GEOS-Chem aerosol simulation includes the sulfate-nitrate-ammonium (SNA) aerosol system
 (Fountoukis and Nenes, 2007; Park et al., 2004), carbonaceous aerosol (Hammer et al., 2016;
 Park et al., 2003; Wang et al., 2014), secondary organic aerosols (Marais et al., 2016; Pye et al., 2010), sea salt (Jaeglé et al., 2011) and mineral dust (Fairlie et al., 2007) with updates to aerosol size distribution (Ridley et al., 2012; Zhang et al., 2013). Aerosol optical properties are based on
- the Global Aerosol Data Set (GADS) as implemented by Martin et al. (2003) for externally mixed aerosols as a function of local relative humidity with updates based on measurements (Drury et al., 2010; Latimer and Martin, 2019). Wet deposition of dust, including the processes of scavenging from convections and large scale precipitations, is presented by the scheme developed byfollows Liu et al. (2001). Dry deposition of dust includes the effects of
- 170 gravitational settling (Seinfeld and Pandis, 1998) and turbulent resistance to the surface, (Zhang et al., 2001), which are represented with deposition velocities in the parameterization, implemented into GEOS-Chem by Fairlie et al. (2007).

The original GEOS-Chem simulation used online dust emissions by coupling the dust mobilization module online. We develop the capability to use offline dust emissions based on

175 the archived fields described in Sect. 2.3. We conduct global simulations with GEOS-Chem (version 12.5.0) at a horizontal resolution of 2° by 2.5° for the year 2016. Simulations using the online and offline dust emissions are conducted to evaluate the offline dust emissions. We conduct two simulations using online dust emissions with different dust source functions. The first is with the original version of dust source function, hereafter noted as the original online 180 dust simulation. The other one is with the updated version of source function, in which the updated fine resolution source function is interpolated to 2° by 2.5° resolution. The annual total emissions for the online dust emissions are at the original value of 909 Tg yr⁻¹. We conduct another two-four sets of simulations using offline dust emissions. The first uses the default offline dust emissions with the annual total dust emission of 909 Tg yr⁻¹. The remaining second 185 usesuse the updated offline dust emissions with the annual total dust emission scaled to 1,500, 2,000 and 2,500 Tg yr⁻¹, which areis in the range of the current dust emission estimates of over 426 — 2,726 Tg yr⁻¹ (Huneeus et al., 2011). We focus on the simulation with 2,000 Tg yr⁻¹ which and better represents observations as will be shown below.

190 **3 Results and Discussion**

3.1 Spatial and seasonal variation of the offline dust emissions

Figure 1 shows the spatial distribution of the annual and seasonal dust emission flux rate for the updated offline dust emissions. The annual dust emission flux rate is high over major deserts, such as the northwestern Sahara, the Bodélé Depression in northern Chad, eastern

- 195 Arabian Peninsula and central Asian Taklimakan and Gobi deserts. There are also hotspots of dust emission flux rate over relatively smaller deserts, such as the Mojave Desert of the southwestern United States, the Atacama desert of southern South America, the Kalahari desert on the west coast of southern Africa and the deserts in central Australia. Those features reflect the fine resolution of the updated dust source function and of the offline dust emissions.
- 200 Seasonally, the dust emission flux rate resembles the annual distribution, however, with a lower dust emission flux rate over the Bodélé Depression in northern Chad in summer and higher dust emission flux rate over the Middle East and central Asian deserts in spring and summer.



Figure <u>1</u>4. Annual and seasonal mean dust emission flux rate for the offline high resolution dust emissions with updated dust source function and updated annual total dust emission of 2,000 Tg.





Figure 22. Annual mean dust emission flux rate for 2016. (a), The original online dust emissions with original dust source function and annual total dust emissions of 909 Tg. (b), Online dust emissions with updated dust source function. (c), Difference of flux rate between online dust emissions using original and updated dust source functions. (d), Offline dust emissions with updated dust source function. (e), Offline dust emissions with updated dust emissions with updated dust source functions. (d), Offline dust emissions with updated dust source function. (e)
 Offline dust emissions with updated dust source function and updated annual total dust emissions of 2,000 Tg. (f), Difference of flux rate between offline and online dust emissions. The online dust emissions are in 2° x 2.5° resolution. The offline dust emissions shown in (b), (d), (f) are regridded from 0.25° x 0.3125° resolution to -2° x 2.5° for comparison with online dust emissions.

Figure 2 shows the spatial distribution of the annual dust emission flux rate for the

online and offline dust emissions with the original and updated dust source functions with

220 original and updated global total dust source strengths. All simulations exhibit high dust emission flux rates over major desert regions, such as the Sahara, Middle East and Central Asian deserts, with local enhancements over the western Sahara and northern Chad. The simulation with the updated source function exhibits stronger emissions in the Sahara and Persian Gulf regions (Fig. 2c). The difference between the online and offline dust emissions, shown in Fig. 2f,

225 <u>can be considered the error in the online approach arising from coarse resolution</u>

meteorological fields. Tindicates that the offline dust emissions based on native resolution meteorological fields have lower dust emission flux rates over northwest Africa, but higher dust emission flux rates over the Middle East and Central Asia. Higher annual dust emission flux rates over the southwestern United States, southern South America, the west coast of southern Africa and central Australia in the offline dust emissions reflect that the native resolution offline dust emissions are strengthened over relatively weaker dust emission regions. Generally, coastal and minor desert regions emit more dust when calculating emissions at the native meteorological resolution.

Figures S<u>3</u>2–S<u>6</u>5 show the seasonal variations of dust emission flux rates for online and
offline emissions. The offline dust emissions have lower emission flux rates than the online dust
emissions during spring (March, April and May) (MAM) and winter (December, January and
February) (DJF) over the Sahara Desert. The offline dust emission flux rate is higher than the
online dust emission flux rate over the Middle East and Central Asian deserts during spring and
summer (June, July and August) (JJA). Emission flux rates are low over Central Asian deserts
during winter. The strengthening of offline dust emissions over weaker dust emitting regions
persists throughout all seasons.

3.2 The performance of AOD simulations over desert regions

Figure 3 shows simulated AOD using the original online and updated offline dust emissions. Difference maps of simulated AOD between online and offline dust emissions are shown in Figure S7. We select for evaluation the AERONET sites where the ratio of simulated dust optical depth (DOD) to simulated total AOD exceeds 0.5 in the simulation using the updated offline dust emissions with annual dust strength of 2,000 Tg. Annually, the simulated DOD has the highest value over the Bodélé Depression. This feature persists in all seasons except summer

250 when DOD has the highest values over the western Sahara and eastern Arabian Peninsula. The scatter plots show that annually the simulated AOD from both simulations are highly correlated with AERONET measurements across the dust regions (R = 0.86-0.88). The simulation with updated offline dust emissions has an improved slope and smaller root mean square error than the simulation using the original online dust emissions. AOD from the simulation with updated offline dust emissions is also more consistent with the measurements in different seasons,

especially in the spring (MAM) and fall (SON) with slopes close to unity and R exceeding 0.9.





Figure <u>3</u>. Annual and seasonal mean simulated dust optical depth (DOD) fraction (left column) and aerosol optical depth (AOD) (middle column) from GEOS-Chem simulations for 2016, and AERONET measured AOD at sites where the ratio of simulated DOD and AOD exceeds 0.5, which are shown as filled circles in the middle column. Boxes in the left top panel outline the three major deserts examined in Figure 4. The right column shows the corresponding scatter plot with root mean square error (E),
 correlation coefficient(R) and slope (M) calculated with reduced major axis linear regression. N is the number of valid ground-based monitoring records. The results for the simulation using the original dust emissions are shown in blueack; the results for the simulation using updated dust emissions with dust strength of 2,000 Tg yr⁻¹-are shown in red. The best fit lines are dashed. The 1:1 line is solid.

270 We further evaluate the performance of simulated AOD over major desert regions using the MODIS Deep Blue (DB) and MAIAC AOD products. Figure 4 shows annual and seasonal scatter plots comparing GEOS-Chem simulated AOD using original online dust emissions and updated offline dust emissions against retrieved AOD from MODIS DB and MAIAC satellite products over the three major desert regions outlined in Fig. 3. Figure S&6 shows the annual
and seasonal AOD distribution from MODIS DB and MAIAC. Annually, the simulation using
updated offline dust emissions exhibits greater consistency with satellite AOD than does the
simulation using original online dust emissions across all three desert regions. The simulation
using updated offline dust emission performs better across all three desert regions and in all
four seasons except for the Sahara in summer, during which AOD is overestimated. Both
simulations underestimate AOD over central Asian deserts during winter when dust emissions
are low and other sources may be more important. Overall, the simulation using original online
dust emissions underestimates AOD over all three major desert regions, especially over the
Middle East and central Asian deserts. The simulation using updated offline dust emissions
exhibits greater consistency with satellite observations with higher slopes and correlations.

3.3 Evaluation of the simulations against surface dust concentration measurements
 We also evaluate our simulations using different dust emissions against measurements of
 surface dust concentrations. Figure 5 shows the comparison of modeled fine dust surface
 concentration against the fine dust concentration observation from the IMPROVE network. The
 simulations using the updated offline dust emissions can better represent the observed surface
 fine dust concentration measurements than the simulation using the original online dust
 emissions with higher correlations and slopes across all seasons. Annually, the correlation
 between the simulation and observation increases from 0.39 to 0.68, and the slope increases
 from 0.31 to 0.71 when using the updated offline dust emissions with annual dust strength of
 909 Tg compared to the simulation using the original online dust emissions. Scaling the annual

	dust strength to 2,000 Tg/yr marginally improves the performance of the model simulation of
	fine dust concentrations in all seasons except winter, during which the surface fine dust
	concentrations are overestimated. Given the specificity and density of the dust measurements,
	and the disconnect of North American dust emissions from the global source, we conduct an
300	additional sensitivity simulation with North American dust emissions reduced by 30%. The right
	column shows that the annual slope in the resultant simulation versus observations improves
	to 1.07, minor improvements to annual and seasonal correlations. Future efforts should focus
	on better representing the seasonal variation of dust emissions.
	Figure 6 shows the comparison of seasonal averaged modeled and measured surface
305	dust concentrations from 12 independent sites across the globe. The simulation using the
	updated offline dust emissions with dust strength of 2,000 Tg yr ¹ is more consistent with the
	observations at almost all sites. The remaining bias at sites distant from source regions, for
	example sites in the Southern Hemisphere and East Asia, likely reflects remaining uncertainty in
	representing dust deposition. Further research is needed to address remaining knowledge gaps,
310	such as better representing the dust size distribution and deposition during transport.



3.3 Discussion of the dust source strength

- One of the advantages of the offline dust emissions is that the dust source strengths are scalable. We have found that the simulation with global total annual dust emission scaled to 2,000 Tg better represents observations than does the default simulation with global total annual dust emission of 909 Tg. We also evaluate simulations with global total annual dust emission scaled to 1,500 Tg and 2,500 Tg. Figure S7 indicates that the simulation with global total annual dust emission scaled to 2,000 Tg is more consistent with satellite observations over
- 320 the Sahara and Middle East. Although the central Asian deserts and regions with AERONET observations (Fig. S8) are better represented by the simulation with global total annual dust emission scaled to 2,500 Tg, since the Sahara has the highest dust emissions (Huneeus et al., 2011), and AOD over the Sahara is most likely dominated by dust, we scale global total annual dust emissions to best match this source region. Additional development and evaluation should
- 325 be conducted to further narrow the uncertainty of dust emissions, especially at the regional scale.



corresponding marker signs and colors. The 1:1 line solid black line.

335

330

Figure 4. Scatter plots and statistics of comparing GEOS-Chem simulated AOD with satellite AOD over desert regions annually (the first column) and seasonally (the right four columns). The results for Sahara, Middle East and Central Asia deserts are shown in the top, middle and bottom rows respectively. The results for the simulation using the original dust emissions are shown in blueack; the results for the simulation using updated dust emissions with dust strength of 2,000 Tg yr⁻¹ are shown in red. Open circles Dots represent the comparison with MODIS Deep Blue AOD; the plus signs represent the comparison with MAIAC AOD. Correlation coefficient(R), root mean square error (E), and Slope (M) are reported, in which R1, E1 and M1 show the results of the comparison with MODIS Deep Blue AOD; R2, E2 and M2 show the results of the comparison with MAIAC AOD. The best fit lines are dashed lines with



with different dust emissions for 2016, and IMPROVE fine dust measurements, which are shown as filled circles. Root mean square error (E), correlation coefficient(R) and slope (M) calculated with reduced major axis linear regression are reported. The results for the simulation using the original dust emissions are shown in blue (left column); the results for the simulation using updated dust emissions with dust strength of 909 Tg yr⁻¹ are shown in red (second column); the results for the simulation using updated dust emissions updated dust emissions with dust strength of 2,000 Tg yr⁻¹ are shown in magenta (third column); the right column is the sensitivity simulation with North America dust emission reduced by 30%.



Figure 6. Comparison of modeled and measured seasonal averaged surface dust concentrations at 12 independent globally distributed sites for the years 1981-2000. Nine sites are in the dust belt across Northern hemisphere. The remaining 3 sites are relatively close to the weak dust emission regions in Southern Hemisphere. The results for the simulation using the original dust emissions are shown in blue; the results for the simulation using updated dust emissions with dust strength of 909 Tg yr⁻¹ are shown in red; the results for the simulation using updated dust emissions with dust strength of 2,000 Tg yr⁻¹ are shown in magenta. The measurements are in black.

3.4 Discussion of the dust source strength

360	One of the advantages of the offline dust emissions is that the same dust source strength can
	be readily applied to all model resolutions, facilitating evaluation of dust source strength
	independent of resolutions. We have found that the simulation with global total annual dust
	emission scaled to 2,000 Tg better represents observations than does the default simulation
	with global total annual dust emissions of 909 Tg. We also evaluate simulations with global total
365	annual dust emission scaled to 1,500 Tg and 2,500 Tg. Figure S9 indicates that the simulation
	with global total annual dust emission scaled to 2,000 Tg is more consistent with satellite
	observations over the Sahara and Middle East. Although the central Asian deserts and regions
	with AERONET observations (Fig. S10) are better represented by the simulation with global total
	annual dust emission scaled to 2,500 Tg, since the Sahara has the highest dust emissions
370	(Huneeus et al., 2011), and AOD over the Sahara is most likely dominated by dust, we scale
	global total annual dust emissions to best match this source region. Additional development
	and evaluation should be conducted to further narrow the uncertainty of dust emissions,
	especially at the regional scale.

375

3.54-Advantages of high resolution offline dust emissions for model development

Uncertainty remains in the estimated global annual total dust emissions. Direct dust emission flux observations are few. Current atmospheric and chemical transport models apply a global scale factor to optimize with a specific set of ground observations. Because of the non-linear

380 dependence on resolution of the dust emissions, the source strength has historically depended

upon model resolution, which inhibits general evaluation. The native resolution offline dust emissions facilitate consistent evaluation and application across all model resolutions. <u>Such</u> <u>consistency is particularly important for stretched-grid simulations with the capability for</u> <u>factors of over 100 variation in resolution within a single simulation (Bindle et al., 2020).</u>

385

4 Summary and Conclusions

The nonlinear dependence of dust emission parameterizations upon model resolution poses a challenge for the next generation of chemical transport models with nimble capability for multiple resolutions. In this paper we have developed and tested a The method explored here 390 to calculate offline dust emissions at the native meteorological resolution to promotes consistency of dust emissions across different model resolutions. We take advantage of the capability of HEMCO standalone module to calculate dust emission offline at native meteorological resolution using the DEAD dust emission scheme combined with an updated high resolution dust source function. We evaluate the performance of the simulation with 395 native resolution offline dust emissions and an updated dust source function with source strength of 2,000 Tg yr⁻¹. We find better agreement with measurements, including satellite and AERONET AOD, and surface dust concentrations. The offline fine resolution dust emissions strengthen the dust emissions over smaller better resolve smaller desert regions. The independence of source strength from simulation resolution facilitates evaluation with 400 observations. Sensitivity simulations with an annual global source strength of either 1,500 or 2,500 Tg generally degraded the performance. A sensitivity simulation with North American emissions reduced by 30% improved the annual mean slope versus observations. Futureurther

work should continue to develop and evaluate the representation of dust <u>deposition and</u> regional seasonal variation<u>emissions</u>.

405 **5 Code and Data Availability**

The source code for generating the offline dust emissions is available on GitHub (<u>https://github.com/Jun-Meng/geos-chem/tree/v11-01-Patches-UniCF-vegetation</u>) and Zenodo repository (<u>https://doi.org/10.5281/zenodo.4062003</u>) (Meng et al., 2020b). The instruction of how to generate the emission files is in the README.md file in the GitHub repository

- (https://github.com/Jun-Meng/geos-chem/tree/v11-01-Patches-UniCF-vegetation). The global high resolution (0.25°x0.3125°) dust emission inventory is available on Zenodo
 (https://doi.org/10.5281/zenodo.4060248) (Meng et al., 2020a), containing netCDF format files of global gridded hourly mineral dust emission flux rate. Currently, the dataset (version1.0) is available for the year 2016. The dataset for other years since 2014 will be available in future
- 415 versions.

The base GEOS-Chem source code in version 12.5.0 is available on Github
(<u>https://github.com/geoschem/geos-chem/tree/12.5.0</u>) and Zenodo repository
(<u>https://zenodo.org/record/3403111#.X7PKv5NKiF0,%202019</u>). The GEOS-Chem simulation

420 output data and AOD observations used to evaluate the model performance, including MODIS Deep Blue, MODIS MAIAC and AERONET AOD, can be accessed via this Zenodo repository (<u>https://doi.org/10.5281/zenodo.4312944</u>) (Meng et al., 2020c).

Information about the Supplement

425 The supplement related to this article describes the details of the dust emission scheme used in this project, <u>the updated high resolution dust source function</u>, as well as additional figures described in the main text.

Author contributions

430 RVM and JM conceived the project. JM developed the dust emission dataset using data and algorithms from DAR, PG, MH, AvD, and MPS. JM prepared the manuscript with contributions from all coauthors. All authors helped revise the manuscript.

Competing interests

435 The authors declare that they have no conflict of interest.

Acknowledgement

This work was supported by the Natural Science and Engineering Research Council of Canada. Jun Meng was partially supported by a Nova Scotia Research and Innovation Graduate
 Scholarship. Martin acknowledges partial support from NASA AIST-18-0011. We are grateful to Compute Canada and Research Infrastructure Services in Washington University in St. Louis for computing resources. The meteorological data (GEOS-FP) used in this study have been provided by the Global Modeling and Assimilation Office (GMAO) at NASA Goddard Space Flight Center. We thank Jasper Kok and Longlei Li for providing the compilation of independent surface dust concentrations measurements. We thank the four anonymous reviewers for their constructive comments and suggestions. All figures are produced with the MATLAB -2019a-software.

References

Graham - CC Doc: https://docs.computecanada.ca/wiki/Graham, last access: 6

450 March 2021.

465

470

Bergin, M. H., Ghoroi, C., Dixit, D., Schauer, J. J., and Shindell, D. T.: Large Reductions in Solar Energy Production Due to Dust and Particulate Air Pollution, Environ. Sci. Technol. Lett., 4, 339– 344, https://doi.org/10.1021/acs.estlett.7b00197, 2017.

Bindle, L., Martin, R. V., Cooper, M. J., Lundgren, E. W., Eastham, S. D., Auer, B. M., Clune, T. L.,
Weng, H., Lin, J., Murray, L. T., Meng, J., Keller, C. A., Pawson, S., and Jacob, D. J.: Grid-Stretching Capability for the GEOS-Chem 13.0.0 Atmospheric Chemistry Model, 1–21, https://doi.org/10.5194/gmd-2020-398, 2020.

Bristow, C. S., Hudson-Edwards, K. A., and Chappell, A.: Fertilizing the Amazon and equatorial Atlantic with West African dust, 37, https://doi.org/10.1029/2010GL043486, 2010.

460 Chen, H., Navea, J. G., Young, M. A., and Grassian, V. H.: Heterogeneous Photochemistry of Trace Atmospheric Gases with Components of Mineral Dust Aerosol, J. Phys. Chem. A, 115, 490–499, https://doi.org/10.1021/jp110164j, 2011.

De Longueville, F., Hountondji, Y.-C., Henry, S., and Ozer, P.: What do we know about effects of desert dust on air quality and human health in West Africa compared to other regions?, Sci. Total Environ., 409, 1–8, https://doi.org/10.1016/j.scitotenv.2010.09.025, 2010.

Drury, E., Jacob, D. J., Spurr, R. J. D., Wang, J., Shinozuka, Y., Anderson, B. E., Clarke, A. D., Dibb, J., McNaughton, C., and Weber, R.: Synthesis of satellite (MODIS), aircraft (ICARTT), and surface (IMPROVE, EPA-AQS, AERONET) aerosol observations over eastern North America to improve MODIS aerosol retrievals and constrain surface aerosol concentrations and sources, 115, https://doi.org/10.1029/2009JD012629, 2010.

Eastham, S. D., Long, M. S., Keller, C. A., Lundgren, E., Yantosca, R. M., Zhuang, J., Li, C., Lee, C. J., Yannetti, M., Auer, B. M., Clune, T. L., Kouatchou, J., Putman, W. M., Thompson, M. A., Trayanov, A. L., Molod, A. M., Martin, R. V., and Jacob, D. J.: GEOS-Chem High Performance (GCHP v11-02c): a next-generation implementation of the GEOS-Chem chemical transport

475 model for massively parallel applications, 11, 2941–2953, https://doi.org/10.5194/gmd-11-2941-2018, 2018.

Fairlie, T. D., Jacob, D. J., and Park, R. J.: The impact of transpacific transport of mineral dust in the United States, Atmospheric Environment, 41, 1251–1266, https://doi.org/10.1016/j.atmosenv.2006.09.048, 2007.

480 Fountoukis, C. and Nenes, A.: ISORROPIA II: A computationally efficient thermodynamic equilibrium model for K⁺-Ca²⁺-Mg²⁺-NH₄⁺-Na⁺-SO₄²⁻-NO₃⁻-Cl⁻-H₂O aerosols, 7, 4639–4659, https://doi.org/10.5194/acp-7-4639-2007, 2007.

Giles, D. M., Sinyuk, A., Sorokin, M. G., Schafer, J. S., Smirnov, A., Slutsker, I., Eck, T. F., Holben,
B. N., Lewis, J. R., Campbell, J. R., Welton, E. J., Korkin, S. V., and Lyapustin, A. I.: Advancements
in the Aerosol Robotic Network (AERONET) Version 3 database – automated near-real-time
quality control algorithm with improved cloud screening for Sun photometer aerosol optical
depth (AOD) measurements, 12, 169–209, https://doi.org/10.5194/amt-12-169-2019, 2019.

Gillette, D. A.: A qualitative geophysical explanation for hot spot dust emitting source regions, Contrib. atmos. phys, 72, 67–77, 1999.

490 Gillette, D. A. and Passi, R.: Modeling dust emission caused by wind erosion, 93, 14233–14242, https://doi.org/10.1029/JD093iD11p14233, 1988.

Ginoux, P., Chin, M., Tegen, I., Prospero, J. M., Holben, B., Dubovik, O., and Lin, S.-J.: Sources and distributions of dust aerosols simulated with the GOCART model, 106, 20255–20273, https://doi.org/10.1029/2000JD000053, 2001.

495 Ginoux, P., Prospero, J. M., Gill, T. E., Hsu, N. C., and Zhao, M.: Global-scale attribution of anthropogenic and natural dust sources and their emission rates based on MODIS Deep Blue aerosol products, 50, https://doi.org/10.1029/2012RG000388, 2012.

Hammer, M. S., Martin, R. V., van Donkelaar, A., Buchard, V., Torres, O., Ridley, D. A., and Spurr,R. J. D.: Interpreting the ultraviolet aerosol index observed with the OMI satellite instrument to understand absorption by organic aerosols: implications for atmospheric oxidation and direct

radiative effects, 16, 2507–2523, https://doi.org/10.5194/acp-16-2507-2016, 2016.

500

Holben, B. N., Eck, T. F., Slutsker, I., Tanré, D., Buis, J. P., Setzer, A., Vermote, E., Reagan, J. A.,
Kaufman, Y. J., Nakajima, T., Lavenu, F., Jankowiak, I., and Smirnov, A.: AERONET—A Federated
Instrument Network and Data Archive for Aerosol Characterization, Remote Sensing of
Environment, 66, 1–16, https://doi.org/10.1016/S0034-4257(98)00031-5, 1998.

Hsu, N. C., Jeong, M.-J., Bettenhausen, C., Sayer, A. M., Hansell, R., Seftor, C. S., Huang, J., and Tsay, S.-C.: Enhanced Deep Blue aerosol retrieval algorithm: The second generation, 118, 9296–9315, https://doi.org/10.1002/jgrd.50712, 2013.

Huneeus, N., Schulz, M., Balkanski, Y., Griesfeller, J., Prospero, J., Kinne, S., Bauer, S., Boucher,
O., Chin, M., Dentener, F., Diehl, T., Easter, R., Fillmore, D., Ghan, S., Ginoux, P., Grini, A.,
Horowitz, L., Koch, D., Krol, M. C., Landing, W., Liu, X., Mahowald, N., Miller, R., Morcrette, J.-J.,
Myhre, G., Penner, J., Perlwitz, J., Stier, P., Takemura, T., and Zender, C. S.: Global dust model
intercomparison in AeroCom phase I, 11, 7781–7816, https://doi.org/10.5194/acp-11-7781-2011, 2011.

515 Jaeglé, L., Quinn, P. K., Bates, T. S., Alexander, B., and Lin, J.-T.: Global distribution of sea salt aerosols: new constraints from in situ and remote sensing observations, 11, 3137–3157, https://doi.org/10.5194/acp-11-3137-2011, 2011.

520

525

530

550

Keller, C. A., Long, M. S., Yantosca, R. M., Da Silva, A. M., Pawson, S., and Jacob, D. J.: HEMCO v1.0: a versatile, ESMF-compliant component for calculating emissions in atmospheric models, 7, 1409–1417, https://doi.org/10.5194/gmd-7-1409-2014, 2014.

Kok, J. F., Adebiyi, A. A., Albani, S., Balkanski, Y., Checa-Garcia, R., Chin, M., Colarco, P. R., Hamilton, D. S., Huang, Y., Ito, A., Klose, M., Leung, D. M., Li, L., Mahowald, N. M., Miller, R. L., Obiso, V., Pérez García-Pando, C., Rocha-Lima, A., Wan, J. S., and Whicker, C. A.: Improved representation of the global dust cycle using observational constraints on dust properties and abundance, 1–45, https://doi.org/10.5194/acp-2020-1131, 2020.

Kosmopoulos, P. G., Kazadzis, S., Taylor, M., Athanasopoulou, E., Speyer, O., Raptis, P. I., Marinou, E., Proestakis, E., Solomos, S., Gerasopoulos, E., Amiridis, V., Bais, A., and Kontoes, C.: Dust impact on surface solar irradiance assessed with model simulations, satellite observations and ground-based measurements, 10, 2435–2453, https://doi.org/10.5194/amt-10-2435-2017, 2017.

Latimer, R. N. C. and Martin, R. V.: Interpretation of measured aerosol mass scattering efficiency over North America using a chemical transport model, 19, 2635–2653, https://doi.org/10.5194/acp-19-2635-2019, 2019.

Liu, H., J. Jacob, D., Bey, I., and Yantosca, R.: Constraints from 210Pb and 7Be on wet deposition
 and transport in a global three-dimensional chemical tracer model driven by assimilated
 meteorological fields, 106, https://doi.org/10.1029/2000JD900839, 2001.

Lyapustin, A., Wang, Y., Korkin, S., and Huang, D.: MODIS Collection 6 MAIAC algorithm, 11, 5741–5765, https://doi.org/10.5194/amt-11-5741-2018, 2018.

Marais, E. A., Jacob, D. J., Jimenez, J. L., Campuzano-Jost, P., Day, D. A., Hu, W., Krechmer, J.,
Zhu, L., Kim, P. S., Miller, C. C., Fisher, J. A., Travis, K., Yu, K., Hanisco, T. F., Wolfe, G. M.,
Arkinson, H. L., Pye, H. O. T., Froyd, K. D., Liao, J., and McNeill, V. F.: Aqueous-phase mechanism for secondary organic aerosol formation from isoprene: Application to the southeast United States and co-benefit of SO₂ emission controls, 16, 1603–1618, https://doi.org/10.5194/acp-16-1603-2016, 2016.

545 Martin, R. V., Jacob, D. J., Yantosca, R. M., Chin, M., and Ginoux, P.: Global and regional decreases in tropospheric oxidants from photochemical effects of aerosols, 108, https://doi.org/10.1029/2002JD002622, 2003.

Meng, J., Martin, R. V., Ginoux, P., Ridley, D. A., and Sulprizio, M. P.: Global High Resolution Dust Emission Inventory for Chemical Transport Models (Version 2020_v1.0), https://doi.org/10.5281/zenodo.4060248, 2020a.

Meng, J., Martin, R. V., and Ridley, D. A.: Offline_Dust_Emissions_SourceCode_2020_v1.0, https://doi.org/10.5281/zenodo.4062003, 2020b.

Park, R. J., Jacob, D. J., Chin, M., and Martin, R. V.: Sources of carbonaceous aerosols over the United States and implications for natural visibility, 108, 4355, https://doi.org/10.1029/2002JD003190, 2003.

Park, R. J., Jacob, D. J., Field, B. D., Yantosca, R. M., and Chin, M.: Natural and transboundary pollution influences on sulfate-nitrate-ammonium aerosols in the United States: Implications for policy, 109, https://doi.org/10.1029/2003JD004473, 2004.

Pye, H. O. T., Chan, A. W. H., Barkley, M. P., and Seinfeld, J. H.: Global modeling of organic
aerosol: The importance of reactive nitrogen (NOx and NO₃)., 10, 11261–11276, https://doi.org/10.5194/acp-10-11261-2010, 2010.

555

570

Ridley, D. A., Heald, C. L., and Ford, B.: North African dust export and deposition: A satellite and model perspective, Journal of Geophysical Research: Atmospheres, 117, https://doi.org/10.1029/2011JD016794, 2012.

565 Ridley, D. A., Heald, C. L., Pierce, J. R., and Evans, M. J.: Toward resolution-independent dust emissions in global models: Impacts on the seasonal and spatial distribution of dust, 40, 2873– 2877, https://doi.org/10.1002/grl.50409, 2013.

Ridley, D. A., Heald, C. L., Kok, J. F., and Zhao, C.: An observationally constrained estimate of global dust aerosol optical depth, 16, 15097–15117, https://doi.org/10.5194/acp-16-15097-2016, 2016.

Sayer, A. M., Munchak, L. A., Hsu, N. C., Levy, R. C., Bettenhausen, C., and Jeong, M.-J.: MODIS Collection 6 aerosol products: Comparison between Aqua's e-Deep Blue, Dark Target, and "merged" data sets, and usage recommendations, 119, 13,965-13,989, https://doi.org/10.1002/2014JD022453, 2014.

575 Schepanski, K., Tegen, I., and Macke, A.: Comparison of satellite based observations of Saharan dust source areas, Remote Sensing of Environment, 123, 90–97, https://doi.org/10.1016/j.rse.2012.03.019, 2012.

Shao, Y., Raupach, M. R., and Findlater, P. A.: Effect of saltation bombardment on the entrainment of dust by wind, 98, 12719–12726, https://doi.org/10.1029/93JD00396, 1993.

580 Tang, M., Huang, X., Lu, K., Ge, M., Li, Y., Cheng, P., Zhu, T., Ding, A., Zhang, Y., Gligorovski, S., Song, W., Ding, X., Bi, X., and Wang, X.: Heterogeneous reactions of mineral dust aerosol: implications for tropospheric oxidation capacity, 17, 11727–11777, https://doi.org/10.5194/acp-17-11727-2017, 2017.

The International GEOS-Chem User Community: geoschem/geos-chem: GEOS-Chem 12.5.0 (Version 12.5.0), Zenodo, https://doi.org/10.5281/zenodo.3403111, 2019.

Wang, Q., Jacob, D. J., Spackman, J. R., Perring, A. E., Schwarz, J. P., Moteki, N., Marais, E. A., Ge, C., Wang, J., and Barrett, S. R. H.: Global budget and radiative forcing of black carbon aerosol: Constraints from pole-to-pole (HIPPO) observations across the Pacific, 119, 195–206, https://doi.org/10.1002/2013JD020824, 2014.

590 Yu, H., Chin, M., Yuan, T., Bian, H., Remer, L. A., Prospero, J. M., Omar, A., Winker, D., Yang, Y., Zhang, Y., Zhang, Z., and Zhao, C.: The fertilizing role of African dust in the Amazon rainforest: A first multiyear assessment based on data from Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observations, 42, 1984–1991, https://doi.org/10.1002/2015GL063040, 2015.

Yu, Y., Kalashnikova, O. V., Garay, M. J., Lee, H., and Notaro, M.: Identification and
 Characterization of Dust Source Regions Across North Africa and the Middle East Using MISR
 Satellite Observations, 45, 6690–6701, https://doi.org/10.1029/2018GL078324, 2018.

Zender, C. S., Bian, H., and Newman, D.: Mineral Dust Entrainment and Deposition (DEAD) model: Description and 1990s dust climatology, 108, https://doi.org/10.1029/2002JD002775, 2003.

600 Zender, C. S., Miller, R. L. R. L., and Tegen, I.: Quantifying mineral dust mass budgets:Terminology, constraints, and current estimates, 85, 509–512, https://doi.org/10.1029/2004EO480002, 2004.

605

Zhang, L., Kok, J. F., Henze, D. K., Li, Q., and Zhao, C.: Improving simulations of fine dust surface concentrations over the western United States by optimizing the particle size distribution, 40, 3270–3275, https://doi.org/10.1002/grl.50591, 2013.