dh2loop 1.0: an open-source Python library for automated processing and classification of geological logs

Ranee Joshi^{1,2}, Kavitha Madaiah^{1,2}, Mark Jessell^{1,2}, Mark Lindsay^{1,2} and Guillaume Pirot^{1,2}

¹Centre of Exploration Targeting, School of Earth Sciences, University of Western Australia, 35 Stirling Highway, Crawley 6009 Western Australia

²Mineral Exploration Cooperative Research Centre (MinEx CRC), School of Earth Sciences, University of Western Australia,
 35 Stirling Highway, Crawley 6009 Western Australia

Correspondence to: Ranee Joshi (ranee.joshi@research.uwa.edu.au)

- 10 Abstract. A huge amount of legacy drilling data is available in geological survey but cannot be used directly as it is compiled and recorded in an unstructured textual form and using different formats depending on the database structure, company, logging geologist, investigation method, investigated materials and/or drilling campaign. It is subjective and plagued with uncertainty as it is likely to have been conducted by tens to hundreds of geologists, all of whom would have their own personal biases. *dh2loop* (https://github.com/Loop3D/dh2loop) is an open-source python library for extracting and standardizing geologic drill
- 15 hole data and export it into readily importable interval tables (collar, survey, lithology). In this contribution, we extract, process and classify lithological logs from the Geological Survey of Western Australia Mineral Exploration Reports Database in the Yalgoo-Singleton Greenstone Belt (YSGB) region. For this study case, the extraction rate for collar, survey and lithology data is respectively 93%, 86% and 34%. It also addresses the subjective nature and variability of nomenclature of lithological descriptions within and across different drilling campaigns by using thesauri and fuzzy string matching. 86% of the extracted
- 20 lithology data is successfully matched to lithologies in the thesauri. Since this process can be tedious, we attempted to test the string matching with the comments, which resulted to a matching rate of 16% (7,870 successfully matched records out of 47,823 records). The standardized lithological data is then classified into multi-level groupings that can be used to systematically upscale and downscale drill hole data inputs for multiscale 3D geological modelling. *dh2loop* formats legacy data bridging the gap between utilization and maximization of legacy drill hole data and drill hole analysis functionalities
- 25 available in existing python libraries (lasio, welly, striplog).

1 Introduction

Drilling is a process of penetrating through the ground that is capable of extracting information about rocks from various depths below the surface. This is useful for establishing the geology beneath the surface. Drill core or cuttings can be collected

- 30 thus providing samples for description, interpretation and analysis. The location of where drilling starts is referred to as the collar. As the drilling progresses, survey orientation measurements are taken to be able to convert the specific depths to exact coordinate locations of the drill core being retrieved. In a hard rock setting, geological drill core logging is the process whereby the recovered drill core sample is systematically studied to determine the lithology, mineralisation, structures, and alteration zones of a potential mineral deposit. It is usually performed by geologists who classify a rock unit into a code, based on one
- 35 or multiple properties such as rock type, alteration intensity and mineralisation content. Exploration and mining companies rely on the diverse geoscientific information obtained by drill core logging techniques to target and to build models for prospectivity mapping or mine planning. This work focuses on lithological logs which is the component of a geological log that refers to the geological information on the dominant rock type in a specific downhole interval. Inevitably, lithological drill core logging is subjective and plagued with uncertainty as all logging geologists have their own personal biases (Lark et al.,
- 40 2014). The information and level of detail contained in logs is highly dependent on the purpose of the study, this already makes geological logging subjective. This subjectivity is also influenced by the lack of a standards between projects and/or companies combined with the personal biases of the logging geologist. Furthermore, it can be difficult to recognize lithology with confidence and to establish subtle variations or boundaries in apparently homogeneous sequences.
- 45 With the advent of the digital age, semi-automated drill core logging techniques such as X-Ray Diffraction (XRD), X-Ray Fluorescence spectrometry (XRF) and Hyperspectral (HS) imaging have provided higher detail of data collection and other properties such as conductivity, volumetric magnetic susceptibility, density using gamma-ray attenuation, and chemical elements during logging (Zhou et al., 2003; Rothwell and Rack, 2006; Ross et al., 2013). This has prompted a shift towards using numerical data rather than depending on traditional geological drill core logging procedures (Culshaw, 2005). Multiple
- 50 methods have been recently applied to geological drill core logging such as wavelet transform analysis or data mosaic (Arabjamaloei et al., 2011; Hill et al., 2020; Le Vaillant et al., 2017; Hill et al., 2015), artificial neural network model (Lindsay, 2019; Zhou et al., 2019; Emelyanova et al., 2017) and inversion (Zhu et al., 2019). Relying solely on these semi-automatic methods comes with drawbacks as it excludes some of the subjective interpretations that cannot be replaced. The semiautomatic methods also are poor at describing textural characteristics (foliation, banding, grain size variation). Furthermore, a
- 55 rich amount of legacy data is collected in the traditional drill core logging method and disregarding this information limits the dataset.

Legacy data are information collected, compiled and/or stored in the past into many different old or obsolete formats or systems, such as handwritten records, aperture cards, floppy disks, microfiche, transparencies, magnetic tapes and/or newspaper clippings making it difficult to access and/or process (Smith et al., 2015). Legacy digital data also suffer from lack of standardisation and inconsistency. In geoscience, these are currently scattered amongst unpublished company reports, departmental reports, publications, petrographic reports, printed plans and maps, aerial photographs, field notebooks, sample ticket books, drill core samples, tenement information and geospatial data providing a major impediment to their efficient use. This includes geological drill core logs that are the outcome of most expensive part of most mineral exploration campaigns:

65 drilling. This is valuable information source and key assets that can be used to add value to geoscientific data for research and exploration; design mapping programs and research questions of interest; more efficiently target remapping and sustainable new discoveries; and provide customers with all existing information at the start of the remapping program. It should not be abandoned for it may have lower intrinsic quality than observations made with more modern equipment, its recovery and translation to a digital format is too tedious. Griffin (2015) argues that there is no distinction in principle between legacy data

- 70 and 'new' data, as all of it is data. The intention of recovering legacy data is to a) upcycle information with integration into modern datasets, b) use salvaged data for new scientific applications and c) allow reuse of that information into utility downstream applications (Vearncombe et al., 2017). Furthermore, extracting information from legacy datasets is valuable and relatively low-risk as geoscientific insight is added to a project for little or no cost compared to those of drilling (Vearncombe et al., 2016).
- 75

80

The primary challenge in dealing with geological legacy datasets is that a large amount of important data, information and knowledge are recorded in an unstructured textural form, such as host rock, alteration types, geological setting, ore-controlled factors, geochemical and geophysical anomaly patterns, and location (Wang and Ma, 2019). To acknowledge the ambiguity in the context of "unstructured textual form", we define it in this paper as, "descriptive text that lacks a pre-defined format and/or metadata thus cannot be readily indexed and mapped into standard database fields". The geological drill core logging forms and formats also vary depending on the company, logging geologist, investigation method, investigated materials and/or

- drilling campaign. Natural language processing (NLP) also known as computational linguistics has been used for information extraction, text classification and automatic text summarization (Otter et al., 2020). NLP applications on legacy data have been demonstrated in the fields of taxonomy (Rivera-Quiroz and Miller, 2019), biomedicine (Liu et al., 2011) and legal services
- 85 (Jallan et al., 2019). Qiu et al. (2020) proposed an ontology-based methodology to support automated classification of geological reports using word embeddings, geoscience dictionary matching and bidirectional long short-term memory model (Dic-Att-BiLSTM) that assists in identifying the difference in relevance from a report. Padarian and Fuentes (2019) also introduced the use of domain-specific word embeddings (GeoVec) which is used to automate and reduce subjectivity of geological mapping of drill hole descriptions (Fuentes et al., 2020).
- 90

110

Similarity matching has many applications in natural language processing as it is one of the best techniques for improving retrieval effectiveness (Park et al., 2005). The use of text similarity is beneficial for text categorization (Liu and Guo, 2005) and text summarization (Erkan and Radev, 2004; Lin and Hovy, 2003). It has been used to extract lithostratigraphic markers from drill lithology logs (Schetselaar and Lemieux, 2012). Fuzzy string matching, also known as approximate string matching, is the process of finding strings that approximately match a given pattern (Cohen, 2011; Gonzalez et al., 2017). It has been

- 95 is the process of finding strings that approximately match a given pattern (Cohen, 2011; Gonzalez et al., 2017). It has been used in language syntax checker, spell-checking, DNA analysis and detection, spam detection, sport and concert event ticket search (Higgins and Mehta, 2018), text re-use detection (Recasens et al., 2013) and clinical trials (Kumari et al., 2020).
- Most of the available python libraries available have been built to process extracted and standardized drill hole data. The most common of these are: *lasio* (https://lasio.readthedocs.io/en/latest/) which deals with reading and writing Log ASCII Standard (LAS) files, a drill hole format commonly used in the oil and gas industry, *welly* (https://github.com/agile-geoscience/welly) which deals with loading, processing, and analysis of drill holes and *striplog* (https://github.com/agile-geoscience/striplog) which digitizes, visualizes and archives stratigraphic and lithological data. *Striplog* (Hall and Keppie, 2016) also parses natural language 'descriptions', converting them into structured data via an arbitrary lexicon which allows further querying and analysis
 - 105 on drill hole data. The main limitation of these existing libraries, with respect to legacy data in the mining sector is that they assume that the data is already standardized and pre-processed.

dh2loop provides the functionality to extract and standardize geologic drill hole data and export it into readily importable interval tables (collar, survey, lithology). It addresses the subjective nature and variability of nomenclature of lithological descriptions within and across different drilling campaigns by integrating published dictionaries, glossaries and/or thesauri

that are built to improve resolution of poorly defined or highly subjective use of terminology and idiosyncratic logging

methods. It is however important to highlight that verifying the accuracy and/or correctness of the geological logs being standardized is outside the scope of this tool, thus we assume logging has been conducted to the best of the geologist's ability.

115 Furthermore, it classifies lithological data into multi-level groupings that can be used to systematically upscale and downscale drill hole data inputs in multiscale 3D geological model. It also provides drill hole desurveying (computes the geometry of a drillhole in three-dimensional space) and log correlation functions so that the results can be plotted in 3D and analysed against each other. It also links the gap between utilization and maximization of legacy drill hole data and the drill hole analysis functionalities available in existing python libraries.

120 2 dh2loop Drillhole Data Extraction

2.1 Conventions and Terminologies

This paper involves multiple python libraries, database tables and fields. For clarity, the following conventions are used for this paper (Appendix A1):

- 1. Python libraries are written in italics: *dh2loop*
- 2. Python functions are written in italics followed by an open and close parenthesis: token_set_ratio()
- 3. Database tables are written in Lucida Console Italics: *dhgeology*
- 4. Database table fields are written in Lucida Console: CollarID
- 5. Workflows are written in Century Gothic Bold: Lithology Code workflow

2.2 Dependencies

125

135

140

- 130 dh2loop stands for drill hole data extracted into a 3D modelling input format, compatible with/for the Loop platform (Ailleres et al., 2019). It is a drill hole processing tool that integrates published dictionaries, glossaries and/or thesauri to and improve standardize highly subjective use of terminology and idiosyncratic logging methods and classify lithological logs. It primarily depends on a number of external open-source libraries (Appendix A2):
 - 1. fuzzywuzzy (https://github.com/seatgeek/fuzzywuzzy) which uses fuzzy logic for string matching (Cohen, 2011)
 - 2. *pandas* (<u>https://pandas.pydata.org/</u>) for data analysis and manipulation (McKinney, 2011)
 - psycopg2 (<u>https://pypi.org/project/psycopg2/</u>), a PostgreSQL database adapter for python (Gregorio and Varrazzo, 2018)
 - 4. numpy (<u>https://github.com/numpy/numpy</u>)
 - nltk (<u>https://github.com/nltk/nltk</u>), the Natural Language Toolkit is a suite of open source Python modules, data sets, and tutorials supporting research and development in Natural Language Processing (Loper and Bird, 2002).
 - 6. *pyproj* (<u>https://github.com/pyproj4/pyproj</u>), python interface to PROJ (cartographic projections and coordinate transformations library)

Code describing basic drill hole operations, such as desurveying (process of translating collar (location) and survey data (azimuth, inclination, length) of drill holes into XYZ coordinates in order to define its 3D geometry of the non-vertical

145 borehole), is heavily inspired from *pyGSLIB* drill hole module (Martínez-Vargas, 2016). The *pyGSLIB* drillhole module is rewritten into python to make it more compact with less dependencies and tailor it to the data extraction output.

2.3 Data Source

The Geological Survey of Western Australia Mineral Exploration Reports Database contains open-file reports submitted as a compliance to the Sunset Clause, Regulation 96(4) of the Western Australia legislation Mining Regulations 1981. These reports

150 contain valuable exploration information in hardcopy (1957-2000), hardcopy and digital format (2000-2007) and digital format

(2000-present) (Riganti et al., 2015). The minimum contents of a drilling report comprise a collar file which describe the geographic coordinates of the collar location (Fig. 1). Additional files may be included, such as a survey file describing the depth, azimuth and inclination measurements for the drilling path; assays; downhole geology and property surveys (e.g. downhole geochemistry, petrophysics) may also be available depending on the company's submission (Riganti et al., 2015).

- 155 The data in the drilling reports are extracted with spatial attribution and imported to a custom-designed relational database (also called the Mineral Drillhole Database) curated by the GSWA that allows easy retrieval and spatial querying. For simplicity, we will refer to this database as the WAMEX database in this text.
- The WAMEX database contains more than 50 years' worth of mineral exploration drill hole data with more than 2.05 million drill holes, imported from over 1,514 companies. Each drill hole is identified by its surface coordinates and its unique ID (CollarID) in the *collar* table (Fig. 2). The drill hole 3D geometry is described in the survey tables (*dhsurvey*, *dhsurveyattr*). The lithology along the drill hole is described as a function of depth in the lithology tables (*dhgeology* and *dhgeologyattr*). However, it is important to emphasize that the drill hole data is of variable quality and reliability and that no validation has been done. The necessary amendments and reformatting enabling to extract and utilize data from the WAMEX database are part of the functionalities provided by *dh2loop*.



170

Figure 1. Simplified example of a drill hole (1.A) and its corresponding interval tables collar (1.B), survey (1.C) and lithology (1.D). The black circle denotes the collar location of the drill hole which is obtained from a collar table (1.B). The purple line represents
the first downhole interval taking its deviation data from the survey table (1.C) and the lithology information from the lithology table (1.D). The same applies for the second interval (orange line) and third interval (blue line). The orange line follows the same trajectory as the first interval as it uses the same entry in the survey table (1.C). The blue line has no lithology data as this information is not present in the lithology table (1.D). The MaxDepth denotes the total drill length (1.B).



175

Figure 2. Simplified WAMEX database schema showing the one-to-many relationship between the collar table and the collarattr table (red solid line). collarattr stores other attributes that describe each unique drill hole, such as maximum depth and elevation. The figure also shows the relationship between the collar table and the other interval tables such as dhsurvey, dhsurveyattr, dhqeology, dhqeologyattr. The deviation of the drill hole is stored in a table, dhsurvey, 180 with a primary key (DHSurveyID) that refers to each unique depth of a drill hole. This primary key has a many-to-one relationship with collar, as there are multiple depth measurements for each drill hole. Furthermore, *dhsurvey* also has a one-to-many relationship with table *dhsurveyattr*, which stores additional attribute information regarding survey, such as azimuth and inclination readings. The example shows the relationship between tables for the first (red dashed line) and second interval (red dashed-dot line). Each drill hole in the WAMEX database is identified by its geographic coordinates and a unique ID (CollarID) 185 in the collar table. The drill hole 3D geometry is described in the survey tables (dhsurvey, dhsurveyattr). This similar relationship is maintained with interval tables, except that the primary key (e.g.DHGeologyID) is used to refer a unique downhole interval rather than a depth measurement. For lithological information, we refer to tables: dhgeology and dhgeologyattr. *dhqeologyattr* which contain information such as rock names and free text descriptions while *dhqeology* provides information to which hole and interval depth that data refers to. This information can be joined and extracted through SQL (Structured Query 190 Language) queries.

2.4 Thesauri

Since most exploration companies have their own nomenclature and systems, which could also change between drilling campaigns, it is necessary to build thesauri: dictionaries that list equivalent and related nomenclature (or synonyms) for different attribute names and values. Synonyms include terminologies that share a similar intent, for example, RL (relative

- 195 level) terms, whether elevation or relative level, as long as the words are recording a vertical height. These thesauri are stored as additional tables in the database. For example, if we are interested in the major lithology in a specific interval, this information can be tabulated as "Major Rock Type", "Lithology_A" or "Main_Geology_Unit" depending on the drill core logging system used. The resulting thesauri considers change in cases, abbreviations, addition of characters, typographical errors and a combination of these. Although listing out these terms is manual and tedious, it only needs to be done once and
- 200 can be re-used and forms the basis for future text matching and as a training set to automate finding similar terms. This is preferred over selection based on regular expressions as when parsing these terms, there are complex patterns in the terms used and the inconsistencies in the way they are written that can be understood by a person with a geological background but not by a simple regular expression. The complexity of the regular expression required to catch all the terms of interest means an optimal expression is difficult, if not impossible, to define, and also tends to be computationally burdensome. *dh2loop* provides
- 205 several thesauri that can easily be updated (if needed) for the following attributes (

A1 Conventions and Terminologies):

- 1. Drill Hole Collar Elevation Thesaurus: 360 synonyms such as "elevation" and "relative level"
- 2. Drill Hole Maximum Depth Thesaurus: 160 synonyms such as "end of hole", "final depth" and "total depth"
- 3. Drill Hole Survey Azimuth Thesaurus: 142 synonyms
 - 4. Drill Hole Survey Inclination Thesaurus: 8 synonyms such as "dip"
 - Drill Hole Lithology Thesaurus: 688 synonyms such as "geology", "Lithology_A", "Major_Geology_Unit" and "Major_Rock_Type"
 - 6. Drill Hole Comments Thesaurus: 434 synonyms such as "description"
- 215 The thesauri created specifically for further processing lithology and comments information are:
 - 7. Drill Hole Lithology Codes Thesaurus (discussed further in Sect. 2.4.1)
 - 8. Clean-up Dictionary (discussed further in Sect. 2.4.2)
 - 9. Lithology Hierarchical Thesaurus (discussed further in Sect. 2.4.3)

In order to extract the other attributes we envisage developing other thesauri, following the same workflow.

220

210

2.4.1 Drill Hole Lithology Codes Thesaurus

This is a thesaurus compiling the equivalent lithology for a given lithological code based on the reports submitted to GSWA. This thesaurus is identified by a company id and report number. The current thesaurus covers 41 out of the 168 companies in the study area with a total of 352 entries (Appendix A1). It is important to note that the Company_LithoCode varies

225 depending on the CompanyID. For example, "Company 551" refers to "Saprolite" as "CS" while Company "2551" uses CS to refer to "Cambrian Sediment". It is also common to for companies to use the different Company_LithoCode to refer to the same lithology. For example, a company may use "AMPH" to refer to "Amphibolite" while another company may use "MAA". Basically, there is a many-to-many relationship between Company_LithoCode and Company_Litho.

230 2.4.2 Clean-up Dictionary

The Clean-up Dictionary is a list of words and non-alphabetic characters that are used as descriptions in the geological logging syntax. This dictionary is used to remove these terms from the Company_Litho and/or Comments free text descriptions prior to the fuzzy string matching. The dictionary is composed of 1662 records, most of which are compiled from abbreviations in field and mine geological mapping (Chace, 1956) and the CGI-IUGS geoscience vocabularies accessible at http://geosciml.org/resource/def/voc/ (Simons et al., 2006; Richard et al., 2007; Raymond et al., 2012). 353 of these records

235 http://geosciml.org/resource/def/voc/ (Simons et al., 2006; Richard et al., 2007; Raymond et al., 2012). 353 of these records are original to *dh2loop* and are added to accommodate the geological logging syntax in Western Australia. The dictionary includes terms that describe age, location, structural forms, textures, amount/distribution, minerals, colors, symbols and common phrases.

240 2.4.3 Lithology Hierarchical Thesaurus

The Lithology Hierarchical Thesaurus is a list of 757 rock names (Detailed_Lithology), their synonyms and a twogrouping (Lithology_Subgroup and Lithology_Group) (Fig 3). level upscale Each row in Detailed_Lithology refers to a rock name. Each rock name row lists the standardized terminology first, followed by its synonyms. The two corresponding columns for this row indicated the two-level upscale grouping. 169 of these rock names 245 CGI-IUGS are compiled from the Simple Lithology vocabulary available at: http://resource.geosciml.org/classifier/cgi/lithology (Simons et al., 2006; Richard et al., 2007; Raymond et al., 2012). The synonyms are obtained from mindat.org (Ralph and Chau, 2014; Ralph, 2004). The hierarchical classification is inherited from both mindat.org (Ralph and Chau, 2014; Ralph, 2004) and the British Geological Survey (BGS) Classification Scheme (Gillespie and Styles, 1999; Robertson, 1999; Hallsworth and Knox, 1999; McMillan and Powell, 1999; Rosenbaum et al.,

250 2003). It is important to use multiple libraries to be able to build an exhaustive thesauri as some libraries are limited by the nomenclature, level of interest and presence of the lithology or rock group in a geographic area. For example, the BGS classification did not have a comprehensive regolith dictionary. Thus, regolith has been classified using the regolith glossary (Eggleton, 2001).



- 255 Figure 3. Lithology Hierarchical Thesaurus showing the 7 major Lithology_Groups: Igneous rocks (pink), Sedimentary rocks (light brown), Metamorphic rocks (green), Surficial Rocks (light yellow), Texture and Structure (blue), Mineralisation (purple) and Unclassified Rocks (dark yellow) and their corresponding Lithology_Subgroups. Many of the Lithology_Subgroups listed have parent-child relationships e.g. 'mafic_fine_grained_crystalline' is a child of 'mafic'. Parents in parent-child relationships are included in their children as catch-all groups to capture free text descriptions that do not include details that would be captured by 260 only using the child terms alone. Igneous rocks Lithology_Subgroups are further classified to 12 lithology subgroups, considering grain size, composition and a combination of both. Sedimentary rocks are subclassified to 16 Lithology_Subgroups based on genetic source and composition (carbonate, clastic, evaporate, hybrid, hydroxide, ironstone, non-clastic siliceous, organicrich, phosphate, siliceous, siliciclastic, volcaniclastic, glacigenic). Metamorphic rocks are subdivided into Lithology_Subgroups based on the degree and type of metamorphism (metasomatic, contact, low-grade, schist, gneiss, high-grade, granofels, greenschist, 265 metacarbonate). Surficial rocks are subdivided into 13 Lithology_Subgroups based on the depositional environment and composition. The residual deposit Lithology_Subgroup includes the regolith detailed lithologies. Mineralisation is considered as a separate classification to be able to classify ore zones. Structure and texture addresses situations that structures are logged as
- lithologies in geological logging. Structure and Texture is divided into five Lithology_Subgroups: fault rock, breaks, contact, fillings and sedimentary structures. The final classification is a catch-all for unclassified rocks. The matching is done at the Detailed_Lithology level, thus not causing confusion in the Lithology_Subgroup and Lithology_Group level. Volcaniclastics are present in both lithological groups as although volcaniclastics are volcanic in origin and are categorized as igneous rocks, ambiguous lithologies such as "metavolcaniclastic sandstone" is more sedimentary than igneous.

2.5 Data Extraction

- 275 Currently, the *dh2loop* library extracts collar, survey and lithology information. It uses a configuration file that allows the user to define the inputs, which are:
 - 1. Region of interest (in WGS 1984 lat/long); and/or
 - 2. List of drill hole ID codes codes, if known.
 - 3. If reprojection is desired, the EPSG code of the projected coordinate system (e.g. EPSG:28350 for MGA Zone 50;

280 http://epsg.io)

- 4. The connection credentials to the local copy of the WAMEX database
- 5. Input and output file directories/location

2.5.1 Collar Extraction

- 285 With the minimum input of a region of interest, the *dh2loop* library exports a Comma-Separated Values file (CSV) listing the drill holes in the area with the following information (Fig. 4):
 - 1. **CollarID**: The **CollarID** for a drill hole is identical in all tables in order for data to be associated with that drill hole.
 - 2. HoleID: This is the drill hole name, as the company would internally identify the drill hole.
 - 3. Longitude and Latitude: Both values are expressed in WGS 1984 lat/long (EPSG:4326).
 - 4. Relative level (RL): We use RL here to refer to elevations of survey points with reference to the mean sea level. This definition of RL is equivalent to the elevation values used in DEMs. This value is extracted by using the Drill Hole Collar Elevation Thesaurus to filter the values referring to relative level (Fig. 4b). More than one value can be fetched due to duplicate company submissions or multiple elevation measurements, in which case the code retains the value with most decimal places assuming higher precision corresponds to better accuracy. If no elevation values are fetched from the database the entire record is skipped. Non-numeric values are also ignored.
 - 5. Maximum depth (MaxDepth): This value is extracted by using the Drill Hole Maximum Depth Thesaurus (Fig. 4c). Due to duplicate company submissions, there can be more than one value fetched. Since there is no submission date information, the code takes the value with largest value assuming it is the latest submission.
 - 6. Calculated X, Y values of projected coordinates: These values are commonly calculated and used to be able to plot the drill hole in a metric system to be able to accurate display and measure distance within and between drill holes. The projection system used in the calculation is based on the input specified in the configuration file.

WAMEX DATABASE							THE	SAURI				
co]]ar					co]]arat	tr					Elevation	Maximum Depth
CollarID	HoleID	Longitude	Latitude		CollarI	D Attr	ibute Column	Attri	bute Valı	le	Synonyms	Synonyms
001	DDH_001	117.30	-29.30		001	elevati	on	450			elevation	end of hole
002	DDH_002	117.00	-29.00	1	002	relativ	e level	500			relative level	final depth
					001	total d	epth	120				total depth
					002	end of	hole	150				TOTAL_DEPTH
			ď								C	
								OL	JTPUT			
				Co	ollar Export							
				Co	ollarID	HoleID	Longitude	Latitud	de RL	MaxDept	th X	Y
				• 00	01	DDH_001	117.30	-29.30	450	120	529135.28	6758738.88
				00)2	DDH_002	117.00	-29.00	500	150	500000.00	6792014.38

Figure 4. Collar extraction workflow showing the CollarID, HoleID, Longitude and Latitude information is fetched from 305 the *collar* table (a, red), the corresponding RL and MaxDepth values are fetched from the *collaratt* table using the Drill Hole Collar Elevation Thesaurus (b, blue) and Drill Hole Maximum Depth Thesaurus (c, orange).

290

295

2.5.2 Survey Extraction

With the same inputs defined in the configuration file, the *dh2loop* library outputs a survey CSV file containing the following information: CollarID, Depth, Azimuth, Inclination and Calculated X, Y, Z values (Fig. 5). The workflow

- 310 accommodates for underground holes drilled upwards as long as the metadata and data appropriately describe them as such. For all properties, all non-numeric values are ignored. For Depth, negative values are replaced by their absolute value. This assumption is made as some drill holes have negative depth information and it is technically not possible to have a negative length. This is done by some companies to denote that the depth measure is going upwards (usually for underground probing drill holes) rather than downhole. For Azimuth, the code fetches values between 0-360 degrees, thus ignoring values greater
- 315 than 360. Values between -360 to 0 are assumed to be counter-clockwise from the north. If there is no survey information for a drill hole present in collar, the azimuth value is set to 0. The X, Y, Z, values are calculated using the minimum curvature basing the code off the *pyGSLIB* drill hole module.



320 Figure 5. Survey extraction workflow showing the DHSurveyID, CollarID and Depth information is fetched from the *dhsurvey* table (a, red), the corresponding Azimuth and Inclination values are fetched from the *dhsurveyattr* table using the Drill Hole Survey Azimuth Thesaurus (b, blue) and Drill Hole Survey Inclination Thesaurus (c, orange).

2.5.3 Lithology Extraction

The lithology extraction is divided into two workflows: Lithology Code Workflow and Comments Workflow. Both

- 325 workflows output a lithology CSV file containing the following information (Fig. 6):
 - 1. **CompanyID**: The primary key to link the lithology code to the Drill Hole Lithology Codes Thesaurus and decode the lithologies.
 - 2. CollarID: The primary key to link the lithology information to the collar file.
 - 3. FromDepth and ToDepth: If the ToDepth is null, we assume ToDepth to be equal to FromDepth + 0.01. If the FromDepth is larger than ToDepth, the FromDepth and ToDepth values are switched.
 - 4. Detailed_Lithology: This value is the lithology matched through fuzzy string matching. The string that serves as input to the fuzzy string matching may either be the Company_Litho (decoded lithology from Company_LithoCode) or from the Comments (free text descriptions).
 - 4.1. Decoding Lithological Codes
 - 4.1.1. **Company_LithoCode**: This fetches the lithology codes that are typically three-letter codes using the Drill Hole Lithology Thesaurus.
 - 4.1.2. Company_Litho: The Company_Litho is fetched by matching the CompanyID and Company_LithoCode to the Drill Hole Lithology Codes Thesaurus.

335

4.2. Comments: This fetches the free text descriptions using the Drill Hole Comments Thesaurus.

- 5. Lithology_Subgroup and Lithology_Group: Upscales the lithological information to more generic rock groups. For example, Detailed_Lithology: "basalt" is upscaled to Lithology_Subgroup:"mafic_fine-grained crystalline" and further upscaled to Lithology_Group:"igneous rock".
 - 6. Calculated X, Y, Z for the start, mid and endpoint also using the minimum curvature algorithm. The desurveying code is heavily based on the *pyGSLIB* drill hole module.



350 Figure 6. Lithology extraction is done through the Lithology Code workflow and Comments workflow. The values are fetched from the *dhgeology* and *dhgeologyattr* table (green) using either the Drill Hole Lithology Thesaurus (blue) and Drill Hole Lithology Codes Thesaurus (light blue) or the Drill Hole Comments Thesaurus (blue). The string fetched is then cleaned prior to the fuzzy string matching using the Clean-up Dictionary (dark yellow). The result is then matched against the Detailed_Lithology level of the Lithological Hierarchical Thesaurus. If there is a match with a score greater or equal to 80, the match is taken and matched with the rest of the columns in the Lithology Hierarchical thesaurus. If not, it is labelled as unclassified rock.

Once the Company_Litho (decoded lithology from Company_LithoCode or from the Comments (free text descriptions) have been extracted from the database, the lithology strings are pre-processed such that:

a) The strings are converted to lowercase form.

360

365

- b) The string inside parenthesis, brackets and braces are removed, as these are found to reduce the accuracy of the matching.
- c) The string preceded by key phrases such as "with", "possibly", "similar to" are removed.
- d) If any of the words listed in the Clean-up Dictionary are present in the string, these words are removed.
- e) Lemmatization, the removal of the inflections at the end of the words in order the "lemma" or root of the words, is applied to all nouns (Müller et al., 2015).
- f) All words with non-alphabetic characters and tokens with less than three characters are removed. This include twoletter words such as "to", "in", "at".
 - g) Stopwords, a set of words frequently used in language which are irrelevant for text mining purposes (Wilbur and Sirotkin, 1992), are removed. Examples on stopwords are: as the, is, at, which, and on.
- 370 This is followed by fuzzy string matching, a technique that finds the string that matches a pattern approximately. Fuzzy string matching is typically divided into two sub-problems: 1) finding approximate substring matches inside a given string, and 2) finding dictionary strings that match the pattern approximately. Fuzzy string matching uses the Levenshtein Distance to calculate the differences between sequences and patterns (Okuda et al., 1976; Cohen, 2011). The Levenshtein distance measures the minimum number of single-character edits (insertion, deletion, substitution) necessary to convert a given string
- into an exact match with the dictionary string (Levenshtein, 1965).

We utilize *fuzzywuzzy* (<u>https://github.com/seatgeek/fuzzywuzzy</u>) for this. *fuzzywuzzy* provides two methods to calculate a similarity score between two strings: *ratio()* or *partial_ratio()*. It also provides two functions to pre-process the strings: *token_sort()* and *token_set()*. In this work, we used the *token_set_ratio()* scorer to do fuzzy string matching to classify the company. Little or Company. Little or Company.

- 380 **Company_Litho** or **Comments** entries into one of the Lithology Hierarchical Thesaurus entries (Table 1). *token_set()* preprocesses the strings by: 1) splitting the string on white-spaces (tokenization), 2) turning to lowercase and 3) removing punctuations, non-alpha non-numeric characters and unicode symbols. It tokenizes both strings (given string and dictionary string), splits the tokens into: intersection and remainder, then sort and compare the strings. The sorted intersection component refers to the similar tokens between the two strings. Since the sorted intersection component (similar tokens between two
- 385 strings) of token_set(), will result in an exact match, the score will tend to increase when: 1) the sorted intersection makes up a larger percentage of the full string, and 2) the remainder component are more similar. The ratio() method then computes the standard Levenshtein distance between two strings. token_set_ratio() is found to be effective in addressing harmless misspelling and duplicated words but sensitive enough to calculate lower scores for longer strings (3-10 word labels), inconsistent word order and missing or extra words. partial ratio() which takes the "best partial" of two strings or the best
- 390 matching on the shorter substring is not preferred as it does not address the difference and order in substring construction. token_sort() is not preferred as it alphabetically sorts the tokens that ignores word order and does not weight intersection tokens which does not address the behavior of the strings in the logs.

- 395 Table 1. Examples of fuzzy string matching output using different combinations of the *fuzzywuzzy* functions. The table demonstrates the corresponding effect of these functions to the given string. token set ratio () which is used by dh2loop, works best on geological free text descriptions as it weights the intersection tokens, honors substring construction and word order and ignores misspelling, extra and duplicated words (in bold). partial ratio () ignores substring construction and order and is more sensitive to misspellings. token_sort_ratio () also ignores substring order and does not recognize duplicate and extra words. The ticks and crosses indicated 400 beside the score indicates the preferred (ticks) result between the methods clustered together.

<i>fuzzywuzzy</i> Function	Given String	Dictionary String	Score		Remarks	
ratio ()	diorite	granodiorite rock	58	\checkmark	partial_ratio () ignores	
partial_ratio ()	diorite	granodiorite rock	100	×	substring construction	
ratio ()	granodoirit rcok granodiorite rock 85 v		\checkmark	ratio () mitigates		
partial_ratio ()	granodoirit rcok	granodiorite rock	81	x	misspelling	
ratio ()	rock felsic granodiorite	granodiorite rock	59	\checkmark	partial_ratio () ignores	
partial_ratio ()	rock felsic granodiorite	granodiorite rock	83	x	substring order	
token_set_ratio ()	rock felsic granodiorite	granodiorite rock	83	\checkmark	token_sort_ratio () ignores	
token_sort_ratio ()	rock felsic granodiorite	granodiorite rock	100	x	substring order	
token_set_ratio ()	intermediate granodiorite rock	granodiorite rock	100	\checkmark	token_set_ratio () weights	
token_sort_ratio ()	intermediate granodiorite rock	granodiorite rock	72	x	intersection tokens	
token_set_ratio ()	gray granodiorite granodiorite	granodiorite rock	83	\checkmark	token_set_ratio () ignores	
token_sort_ratio ()	gray granodiorite granodiorite	granodiorite rock	64	x	extra and duplicate words	
token_set_ratio ()	gray granodiorite granodiorite rckso	granodiorite rock	83	\checkmark	<pre>token_set_ratio () weights intersection tokens,</pre>	
partial_token_set_ratio ()	gray granodiorite granodiorite rckso	granodiorite rock	100	×	addresses substring construction and word order, ignores misspelling, extra and duplicate words	

dh2loop calculates the token set ratio() between the Company_Litho or Comments (given string) and the entries in the Lithology Hierarchical Thesaurus (dictionary string). The tendency is to enumerate the descriptors before the rock name. For

- 405 example, if the lithology in the logged interval is "basalt", the free text description could be something like "Dark gray to dark reddish brown, with olivine phenocrysts, largely altered andesitic basalt". After processing the string, it will be left with "andesitic basalt". To avoid, misclassifying the rock to "andesite", a bonus score is also added to add weight to the last word (in this case, "basalt"). Furthermore, the reader may worry that "basaltic andesite" will be simplified and classified into "andesite". Since "basaltic andesite" is an established volcanic rock name, it will remain as "basaltic andesite". For the pair
- 410 between Company_Litho or Comments and the entries in the Lithology Hierarchical Thesaurus with the highest score, the first synonym is stored as Detailed_Lithology. If the score is less than 80, it is classified as "unclassified rock". The cut-off value is user-defined and can be chosen based on the performance of the matching on the subset of the desired region. If the performance is significantly lower, this indicates that the thesauri used in *dh2loop* may not be suitable to your area. The user may opt to update these thesauri to suit their needs. Once matched on Detailed_Lithology, the corresponding
- Lithology_Subgroup and Lithology_Group classifications are also fetched. 415

2.6 Fuzzy String Matching Assessment

The objective is to compare the Detailed_Lithology classification results obtained from two independent workflows: 1) Lithology Code Workflow and 2) Comments Workflow. Using the Company_LithoCode, Company_Litho,

420 Lithology Code Workflow: Detailed_Lithology and Comments Workflow: Detailed_Lithology from the dataset for the fuzzy string matching assessment, we can assess if matches using the Comments workflow alone can sufficiently decode lithology.

To be able to assess the matching we take a look at the type of matches between Lithology Code Workflow:

- 425 Detailed_Lithology and Comments Workflow: Detailed_Lithology. First, we define a match as retrieving an answer from the fuzzy string matching with a score greater than 80. It is important to note here that it only suggests that it succeeded to find an answer above the score threshold but not necessarily mean that it is the correct answer. To further describe the quality of a match, we modified for this purpose the following terminologies from the Simple Knowledge Organization System (Miles and Bechhofer, 2009):
- 430 a) Exact Match suggests that both Lithology Code workflow and Comments workflow resulted in the same classification at all 3 levels. The match at the Detailed_Lithology level has an exact match, thus resulting to an exact match on the other two levels.
 - b) Close Match suggests that the results at the Detailed_Lithology level are related rocks and belong to the same Lithology_Subgroup. This is usually caused by differing use of lithological nomenclature.
- c) Related Match suggests that the results at the Detailed_Lithology level are related rocks and belong to the same Lithology_Group.
 - d) Broad Match refers to the Detailed_Lithology from Lithology Code workflow matches to a Lithology_Subgroup in the Comments workflow.
 - e) Narrow Match is the logical equivalent of a Broad Match. In this case, the Comments workflow resulted in a Detailed_Lithology level while the Lithology Code workflow resulted in a Lithology_Subgroup level.
 - f) Broader Match is similar to a broad match except that the Detailed_Lithology from Lithology Code workflow matches to a Lithology_Group instead of a Lithology_Subgroup in the Comments workflow.
- 445 g) Narrower Match is the logical equivalent of Broader Match. The Comments workflow results to a Detailed_Lithology while the Lithology Code workflow results to a Lithology_Group level.
 - h) **Failed Match** suggests all levels of both workflows do not match. This is usually attributed to contrasting information from both fields or the algorithm fails. This category is an addition to the SKOS reference.
- 450 For better understanding of these relationships, examples are shown in Table 2 and Fig 7.

440

Table 2. Fuzzy string matching terminology used to describe the quality of matches based on the Simple Knowledge Organization System (SKOS) (Miles and Bechhofer, 2009). The values being compared are the Detailed_Lithology level for both Lithology Code workflow and Comments workflow. The level at which the records are considered to match are in bold. A Match retrieves an answer from the fuzzy string matching with a score greater than 80. An Exact Match suggests that both workflows resulted in the same classification at all 3 levels. A Close Match suggests that the results at the Detailed Lithology level are related rocks.

455

460

the same classification at all 3 levels. A Close Match suggests that the results at the Detailed_Lithology level are related rocks and belong to the same Lithology_Subgroup. A Related Match suggests that the results at the Detailed_Lithology level are related rocks and belong to the same Lithology_Group. A Broad Match refers to the Detailed_Lithology from Lithology Code workflow matches to a Lithology_Subgroup in the Comments workflow. Narrow Match is the logical equivalent of a Broad Match. Broader Match is similar to a broad match except that the Detailed_Lithology from Lithology Code workflow matches to a Lithology_Group instead of a Lithology_Subgroup in the Comments workflow. Narrower Match is the logical equivalent of Broader Match. A Failed Match suggests all levels of both workflows do not match.

Lithology Code Workflow: Detailed_ Lithology	Comments Workflow: Detailed_ Lithology	Lithology Code Workflow: Lithology_ Subgroup	Comments Workflow: Lithology_ Subgroup	Lithology Code Workflow: Lithology_ Group	Comments Workflow: Lithology_ Group	Type of Match
basalt	basalt					Exact Match
basalt	basaltoid	mafic fine grained crystalline	mafic fine grained crystalline			Close Match
basalt	gabbro	mafic fine grained crystalline	mafic coarse grained crystalline	igneous	igneous	Related Match
basalt	mafic fine grained crystalline	mafic fine grained crystalline	mafic fine grained crystalline			Broad Match
mafic fine grained crystalline	basalt	mafic fine grained crystalline	mafic fine grained crystalline			Narrow Match
basalt	mafic	mafic fine grained crystalline	mafic	igneous	igneous	Broader Match
mafic	basalt	mafic	mafic fine grained crystalline	igneous	igneous	Narrower Match
basalt	sandstone	mafic fine grained crystalline	clastic	igneous	sedimentary	Failed Match



Figure 7. SKOS graph showing the semantic, associative and hierarchical relationship in the Lithology Hierarchical Thesaurus. In this example, terms "basalt" and "alkali basalt" are judged to be sufficiently the same to assert an Exact Match relationship (in green). "basic volcanic rock" however is considered a Close Match (in cyan) and "gabbro" a Related Match (in blue). "mafic fine grained crystalline" and "mafic coarse grained crystalline" are broader concepts, thus considered a Broad Match (in orange) to "basalt" and "gabbro" respectively. Broader Match (in brown) are similar to Broad Matches but are used to refer a wider semantic difference between the two concepts. Narrow Matches (in light purple) and Narrower Matches (in dark purple) are the logical equivalent of Broad Match and Broader Match. Failed Matches is used to describe unrelated matches.

- 475 The matching results can be visualized as confusion matrices, which are typically used in machine learning to compare the performance of an algorithm versus a known result. In this case, we are comparing the performance of the string matching using the Comments workflow against the results from the Lithology Code workflow. Each row of the matrix represents the matched lithology from the Comments workflow while each column represents the matched lithology from the Lithology Code workflow. The diagonal elements represent the count for which the Comments workflow class is equal
- 480 to the Lithology Code workflow. The off-diagonal elements are those that are misclassified by the Comments workflow. The higher the diagonal values of the confusion matrix the better, indicating many correct matches. The confusion matrices show normalisation by class support size. This kind of normalisation addresses the class imbalance and allows better visual interpretation of which class is being misclassified. The color of the cell represents the normalised count of the records to address the uneven distribution of records across different classes. Relying on one metric to assess the matching can be
- 485 misleading, therefore, we would like to use four metrics: accuracy, precision, recall and F1 score. It is worth mentioning that a small support influences the precision and/or recall. However, this is the nature of using real-world geological logs as more detail is given to particular lithologies or areas depending on the interest of the study.

3 Case Study: Yalgoo-Singleton Greenstone Belt

490 3.1 Study Area

In this paper, we demonstrate the application of *dh2loop* to data from the Yalgoo-Singleton greenstone belt (YSGB) (Fig. 8), a geologically complex, largely heterogeneous and highly mineralized arcuate granite-greenstone terrane, in the western Youanmi Terrane, Yilgarn Craton in Western Australia (Anand and Butt, 2010). The YSGB has good range of different lithologies in the area. Igneous rocks occur as extensive granitoid intrusions emplaced between 2700 and 2630 Ma (Myers,

- 495 1993), as well as ultramafic to mafic volcanic rocks formed as extensive submarine lavas and local eruptive centres of felsic and mafic volcanic rocks. Some layered gabbroic sills intruding the greenstone are also observed. Sedimentary rocks formed in broad basins during tectonic and volcanic quiescence consist of mostly banded iron formation (BIF) and felsic volcaniclastic rocks. The greenstone belt is metamorphosed to greenschist facies (Barley et al., 2008). The area is also covered by deeply weathered regolith which conceals mineral deposits hosted by the underlying bedrock. Regolith contains signatures of
- 500 mineralisation that are distal signatures of possible economically significant deposits (Cockbain, 2002). Furthermore, the YSGB is a major target for exploration as it has considerable resources of gold, nickel, bauxite, as well as lesser amounts of a wide range of other commodities (Cockbain, 2002). It hosts multiple mineral deposits ranging from volcanogenic massive sulphide (Golden Grove, Gossan Hill), orogenic gold (Mt. Magnet), banded iron formations (Mount Gibson, Karara, Extension Hill). The geological and structural complexity, including its relevance to mineral exploration makes the YSGB a reasonable
- 505 and sensible area to test the *dh2loop* thesauri, matching and upscaling.



Figure 8. The map shows the Yalgoo-Singleton greenstone belt highlighting the different mines and prospects in the area. The inset map shows the heterogeneous distribution and drill hole density from the legacy data available from the WAMEX database.

510 3.2 Data Extraction Results

3.2.1 Collar

530

Extraction of the collar data for YSGB resulted in a collar file with 68,729 drill holes (Fig. 9). This information is extracted from the *collar* table with 73,881 drill holes with 769,981 rows of information from *collarattr*. It includes the location of the collar both in geographic and projected coordinated systems, relative level (RL) and maximum depth (MaxDepth). A

- 515 total of 136,100 records for RL are retrieved from the database, 1,526 of which are disregarded: 846 records for having an RL value greater than 10,000 meters and 680 non-numeric records. These discarded values are retrieved from the attribute column "RL_Local". In spite of it being an isolated issue for "RL_Local", the attribute column is retained as it is retrieved sensible values for other companies. The discarded values are limited to data from two companies (4085, 4670) for RL attribute columns "TD" and "DEPTH". A total of 58,706 records for MaxDepth are retrieved from the database: 58,642 of which are extracted
- 520 as is, while 64 entries are disregarded for having a value of -999. The discarded values come from 8 companies. Null values are disregarded and absent RL or MaxDepth values. The "clean" collar export file contains at least either a value for RL or MaxDepth. The reasoning behind keeping records with at least one of the two field is there are other ways to extract for RL or MaxDepth from the database. RL values can be extracted from digital terrain models and MaxDepth values can be taken for the largest ToDepth values from the other tables.



Figure 3. Extraction of the collar, survey and lithology data for the YSGB. The collar extraction resulted in a collar file with 68,729 drill holes from the *co77ar* table with 73,881 drill holes with 769,981 rows of information from *co77arattr*. A total of 136,100 records for RL are retrieved from the database, 1,526 of which are disregarded: 846 records for having an RL value greater than 10,000 meters and 680 non-numeric records. A total of 58,706 records for MaxDepth are retrieved from the database: 58,642 of

- which are extracted as is, while 64 entries are disregarded for having a value of -999. The "clean" collar export file contains at least either a value for RL or MaxDepth. Survey extraction in YSGB resulted in 126,669 survey depth information across 45,708 drill holes. The *dhsurvey* table contained 146,713 survey depth intervals (from 45,708 drill holes) with corresponding 850,507 entries of supplementary survey information in *dhsurveyattr*. 77 Azimuth values greater than 360 and 152 values are non-numeric
- 535 values. Lithology extraction is divided into two workflows. For the Lithology Code workflow, the extraction starts with filtering the *dhgeology* and *dhgeologyattr* table by the location extents and the Drill Hole Lithology Thesaurus. The *dhgeology* table contained 47,062 drill holes across 115 companies with 797,975 lithology depth intervals with corresponding 820,612 entries of lithology information in *dhgeologyattr*. These records are matched with the entries from the Lithology Code thesaurus resulting to 273,684 matched records. The FromDepth and ToDepth for these records are then validated. 74 records had equal FromDepth
- 540 and ToDepth values. 654 had values for FromDepth but null values for ToDepth. For both cases, ToDepth is calculated as FromDepth+0.01. The Lithology Code workflow resulted to 273,684 intervals across 12,793 drill holes wherein 235,606 records

are successfully matched in the fuzzy string matching. The Comments workflow extracts the records from the *dhgeology* and *dhgeologyattr* table as well, but this time using the Drill Hole Comments Thesaurus (262,567 records across 22,766 drill holes with free text descriptions). 47,823 records are present in both workflows, 7,870 records of which are successfully matched. The 3,074 unique entries from this is used as the dataset for the fuzzy string matching assessment.

3.2.2 Survey

545

For the survey extraction, the *dhsurvey* table contained 146,713 survey depth intervals (from 45,708 drill holes) with corresponding 850,507 entries of supplementary survey information in *dhsurveyattr* (Fig. 9). Survey extraction in YSGB resulted in 126,669 survey depth information across 45,708 drill holes with azimuth (-52.5 to 359) and inclination

- 550 measurements (0-90) for each depth interval. A total of 517,592 records for Azimuth are retrieved from the database. 77 Azimuth values greater than 360 are retrieved and thus disregarded. 152 values are non-numeric values and are also disregarded. These discarded values involved 228 holes across 10 companies. A value of 0 is assigned to missing Azimuth values. A total of 118,223 records for Inclination are fetched from the database, 118,138 of which are extracted as is, while 95 entries are disregarded for having a value greater than 90. A values of -90 is assigned as the default for
- 555 Inclination. The discarded values correspond to 94 drill holes across 5 companies.

3.2.3 Lithology: Lithology Code Workflow and Comments Workflow

Lithology extraction is divided into two workflows. For the **Lithology Code workflow**, the extraction starts with filtering the *dhgeology* and *dhgeologyattr* table by the location extents and the Drill Hole Lithology Thesaurus. The *dhgeology* table contained 47,062 drill holes across 115 companies with 797,975 lithology depth intervals with corresponding 820,612 entries of lithology information in *dhgeologyattr*. These records are matched with the entries from the Drill Hole Lithology Codes Thesaurus resulting to 273,684 matched records. The FromDepth and ToDepth for these records are then validated. 74 records had equal FromDepth and ToDepth values. 654 had values for FromDepth but null values for ToDepth. For both cases, ToDepth is calculated as FromDepth+0.01. The cut-off value of 80 is used for the string matching based on the performance of the matching on a subset of 1,548 unique lithology codes from the Golden

Grove area (Fig. 10). The **Lithology Code workflow** resulted to 273,684 intervals across 12,793 drill holes wherein 235,606 records are successfully matched in the fuzzy string matching. The remaining 546, 819 entries did not obtain a match with a score greater than 80. An example of unmatched entries is provided in Table 2.

Cut-off Score Determination



Figure 10. The user-defined cut-off score of 80 is chosen based on the results of the testing different cut-offs on a smaller dataset within the YSGB area. As seen in this figure, the number of exact matches plateau at a score of 80. This relationship may vary depending on the datasets available in the area. Thus, this cut-off value is user-defined and is best to test the matching performance on a subset in the user's area. The number of exact matches plateau at a score of 80 giving a realistic cut-off and avoids an overly stringent threshold of 100, where useful matches may not be captured.

The **Comments workflow** extracts the records from the *dhgeology* and *dhgeologyattr* table as well, but this time using the Drill Hole Comments Thesaurus. For YSGB, the database has 262,567 records across 22,766 drill holes with free text descriptions. 47,823 records are present in both workflow. Since the free text descriptions are extracted here to compare their results from fuzzy string matching, only 7,870 records that also matched (both have a score greater than 80) in the **Lithology Code workflow** are retained.

585 3.3 Fuzzy String Matching Results

We present results from the data extraction using both workflows: Lithology Code and Comments. The dataset for the fuzzy string matching assessment consists only of the unique records matched on both Lithology Code workflow and Comments workflow (3,074 records). It is visually checked from the records that the Lithology Code workflow: Detailed_Lithology results are sound classifications of the Company_Litho. This is done to make sure that these results could be considered as the "true value" in the fuzzy string matching assessment. The overlaps between these two workflows suggest that the user may need to make choices to identify which is better suited for matching in their area of interest. To better understand the difference between these results, we looked at the matching overlaps between the two workflows (3,074 entries). These matching overlaps are used to compare and describe the fuzzy string matching using the decoding the Company_LithoCode and using Comments.

595

590

We also take a look at the unique combinations of Company_LithoCode, Company_Litho, Lithology Code workflow: Detailed_Lithology and Comments workflow: Detailed_Lithology (53 unique records from the 3,074 records). 34 out of the 53 unique entries (64%) result to matches between the Lithology Code Workflow: Detailed_Lithology and Comments Workflow: Detailed_Lithology. 26 of which are Exact Matches, 19

- 600 unique entries are Close Matches and 26% percent are Failed Matches. The Failed Matches are due to unrelated descriptions in the Comments field which is used to obtain the results in **Comments Workflow**: Detailed_Lithology. An example of this is the interval is logged as "ironstone" (Company_Litho) but Comments contains "mafic schist". Another less common reason is the Company_LithoCode is repeated in the Comments. An example of this is would be an interval logged as "colluvium" and the Comments as "COL". The **Comments workflow** will result to "coal" instead.
- 605 **Exact Matches**: Of the total matched entries, 944 are Exact Matches (31%) (Table 2). The Exact Matches are ideal outcomes as both workflows resulted in exactly the same answers.

Close Matches: The Close Matches are common for coarse-grained igneous rocks, clastic sedimentary rocks, surficial residual rocks and filling structures. The coarse-grained igneous rocks such as gabbro, gabbroid and dolerites are used interchangeably in both fields. Comments can contain terminologies such as "gabbroic", "granophyric gabbro to dolerite", "intrusive granitoid to gabbro" resulting to close matches. Similar cases are observed between granodiorite and granite and between peridotite and coarse-grained ultramafic rocks. For clastic sedimentary rocks, the Close Matches are a result of gradation of grain size in the Comments. For example, an interval logged as mudstone (Company_Litho) is then described in Comments as "mudstone to sandstone" or "intercalated with siltstone". Comments entries like this will result in "sandstone" and "siltstone", "

- 615 respectively. Both clastic sedimentary rocks but not an Exact Match to mudstone. Metasediments and quartz veins occur together and what is described last dictates the Detailed_Lithology classification. Surficial rocks such as soil, duricrust, colluvium, laterite, calcrete, ferricrete and cover are used loosely or occur together resulting to multiple combination of these Close Matches.
- 620 Related Matches: 60 entries (3%) resulted in related matches. For igneous rocks, this result is observed when Comments use rock type descriptors such as "komatitic", "basaltic" and "doleritic". An example would be an interval logged as dolerite and is then described in Comments as "dolertic basalt". This would result in dolerite in the Lithology Code workflow and "basalt" in the Comments workflow. Both lithologies are igneous, however have different composition and textural implications. For sedimentary rocks, Lithology Code workflow results to sedimentary rocks classified based on grain size
- 625 as they have been logged ("gravel", "mud"). The comments contains compositional descriptions such as "with silcrete" or "minor chert". In this case, the Comments workflow will result in "silcrete" and "chert". Both workflows will result in sedimentary rocks, but the Lithology Code workflow will result in "clastic" rocks while the Comments workflow will classify these to "siliceous" at the Lithology_Subgroup level. The related matches for structures occur across coincident lithologies such as "mylonite", "vein", "fault" and "breccia" which could either be "fillings" or "fault_rock" at the 630 Lithology_Sugbroup.

Broad and Narrow Matches: No broad matches are noted and only one narrow match is obtained (Table 3). The interval is logged as "ironstone" with "BIF" in **comments**, "ironstone" being a more general description for "banded iron formation".

635 Broader and Narrower Matches: More common cases are Broader and Narrower Matches indicate that there is a bigger relationship gap between the data in Company_Litho and Comments. Broad matches are a result of low detail free text descriptions in Comments. For example, an interval logged as "gabbro" is described as "medium-grained mafic", "massive mafic", "rich mafic". The inverse is noted for narrower matches, the interval is logged as "sediment" but in Comments the interval is described as "siliceous sediments".

Failed Matches: 1,694 entries resulted in Failed Matches (55%). Failed Matches occur when Company_Litho and Comments contain different information. This could be because the Company_Litho contains the main lithology while Comments contains all other lithologies intercalated in the interval. Another reason is the Company_Litho is relogged based on adjacent intervals without amending Comments. "Mudstone" had failed matches with a wide range of lithologies, such as: "amphibolite", "dolerite", "saprolite", "duricrust", "laterite", "banded iron formation", "chert", "phyllite", "schist", "vein". The same is observed for igneous rocks such as: "coarse-grained-ultramafic-rock". For "chert", the failed matches are within a range of sedimentary rocks: "alluvium" and "mud", "amphibolite" and "massive sulphide", "carbonate", "vein", "pegmatite".

650 Table 3. Distribution of matches across the Fuzzy String Matching Dataset. A total of 45% of the unique records are matched reasonably, 31% of which are Exact Matches, 6% Close Matches, 3% Related Matches, 3% Broader Matches and 3% Narrower Matches.

Type of Match	Number of Entries	Percent
Exact Match	944	31%
Close Match	197	6%
Related Match	60	3%
Broad Match	0	0%
Narrow Match	1	0%
Broader Match	84	3%
Narrower Match	95	3%
Failed Match	1694	55%
TOTAL	3,074	100%

The matching results are visualized as confusion matrices, comparing the performance of the string matching using the 655 **Comments workflow** against the results from the **Lithology Code workflow**. From the 3,074 unique records, we use a total of 1,200 samples for the confusion matrices. The reason for this difference is the limitation of building a confusion matrix wherein both workflows look at the same classes, and ensuring that both workflows produce a match.

3.3.1 Structure and Texture

- 660 While geological structures are not lithologies, they are sometimes described in lithological logs (Fig 11). Structures common in the YSGB area are faults and veins. Figure 11 shows the confusion matrix for the structures and textures. The vertical axis represents the matches from the Lithology Code workflow while the horizontal axis for the results from the Comments workflow. We consider a dataset of 52 unique records where we are trying to assess if the Comments workflow results to the same classification as the Lithology Code workflow. Figure 11 shows that there are 6 records classified as "fault" and 46 records as "vein". When looking at the classification of "faults" we can say that there are 2 records that are true positives.
- 46 records are true negative pairs, as in this 2x2 matrix, if it is not a "fault", it is a "vein". True negatives together with true positives are the Exact Matches and suggests that the **Comments workflow** identified it correctly. To have a better look at the parts that are not classified correctly we look at the false positives and false negatives. False positives represent the number of records classified as "fault" but based on the **Lithology Code workflow** are not. In this case, there are no false positive
- values. False negatives represent number of records classified as "vein" but are actually "faults" based on the LithologyCode workflow.

A total of 48 Exact Matches are noted, 46 records of which are "veins" and 2 records are "faults". This can be surmised by looking into the diagonal cells. The rest of the "veins" (4 records) are Related Matches as "faults". They are considered Related

675

Matches as faults and veins tend to coexist in nature. In addition, faults often occur as fault zones, with infill clay or silica vein sulphides which are described in **Comments** that then obscures the classification. These structure-related lithological descriptions can be used as proxies in further geological studies.



Figure 4. Confusion matrix for structure and texture comparing the fuzzy string matching results from the Lithology Code workflow (vertical axis) and Comments workflow (horizontal axis). The heatmap shows the values normalised to the support size to address the imbalance between classes. The values shown in the cells indicate the number of samples classified for the class. Empty cells indicate zero samples. The Structures and Texture Lithology_Group had an accuracy of 92.3% across 52 samples, 46 for veins and 6 for faults.

3.3.2 Igneous Rocks

- The confusion matrix for igneous rocks considers a dataset of 218 unique records (Fig 12). Dealing with a larger matrix is not as straight-forward as the previous matrix. When looking at the classification of a single lithology, the true positives are where both axes refer to the same class. For example, for "basalt" there are 15 records of true positives which correspond to the Exact Matches. The false positives are the sum of all the other entries along the corresponding vertical axis and the false negatives are the sum of all the entries along the corresponding horizontal axis. The sum of all the other cells represent the true negatives.
- 690 For "basalt", there are 15 true positives, 13, false positives, 15 false negatives and 175 true negatives. This results to 54% classification precision for "basalt".

This statistic is helpful in quantifying the performance of the classification. However, what it does not capture is the semantic and hierarchical relationship of the false negative pairs. As shown in Figure 12, 3 records are classified as "komatiite" and 12

records are classified as "mafic". The "komatiite" matches are a result of when Comments describe the basalts as "komatiitic 695 basalts". This can be considered as a Related Match. The 12 records which are classified as "mafic: are considered "Broader Match". For the false positive values, the "mafic" records are Narrower Matches while the "dolerite" is a Related Match. These quantitative assessment of the matches show us that although the matching is not perfect, the context of the misclassification is not severe.

700

"Dolerite" is the most common igneous rock matched. This could be attributed to the sampling bias towards dolerite as it is often targeted by drilling as they are used as targeting criteria for gold mineralisation (Groves et al., 2000). Given that dolerites can be described by their mafic component or be confused as gabbro when weathered, the descriptions contain strings "mafic" and "gabbro" which explain Close and Broader Matches. Gabbros are also common in the YSGB. Some of the "gabbros" are 705 classified as "mafic" in the Comments Detailed_Lithology. This is another example of a Broader Match. However, it is important to note that although it is not an Exact Match, a Broader Match can be useful in geological studies relating to rock composition as gabbros are members of mafic rocks. 40% of the igneous rock that are mismatched at the

Detailed_Lithology level are Broader Matches (matches correctly at Lithology_Group).



Lithology Code Workflow

Accuracy=0.596

	precision	recall	f1-score	support
basalt	0.54	0.50	0.52	30
coarse-grained- ultramafic-rock	1.00	0.20	0.33	10
dolerite	0.86	0.65	0.74	65
felsic	0.50	1.00	0.67	1
gabbro	0.68	0.62	0.65	45
komatiite	0.20	1.00	0.33	1
mafic	0.45	0.62	0.52	65
peridotite	0.33	1.00	0.50	1
macro avg	0.57	0.70	0.53	218
weighted avg	0.66	0.60	0.60	218

710 Figure 5. Confusion matrix for igneous rocks comparing the fuzzy string matching results from the Lithology_Code workflow (vertical axis) and Comments workflow (horizontal axis). The heatmap shows the values normalised to the support size to address the imbalance between classes. The values shown in the cells indicate the number of samples classified for the class. Empty cells indicate zero samples. The accuracy is 59.6%, with a weighted average precision of 66% and recall of 60%. These results are taken from a subset of 218 samples across 8 classes. "Coarse-grained-ultramafic-rock" has a precision of 1 that implies there are no False 715 Positives.

3.3.3 Sedimentary Rocks

The largest Lithology_Group of the lithological entries relates to sedimentary rocks (800 entries) (Fig 13). 457 of the 800 entries are true positive classification of mudstones. Mudstones are common as shale beds. Mudstones resulted in Related Matches with "chert" and "ironstone". The misclassification occurs when the logs describe intervals wherein the mudstone

720 occurs together and is intercalated with these lithologies. A few mudstones (17) are matched as sandstone due to textural and grain-size descriptors (Close Match). 48% of the cherts are resulted in Exact Matches. 39 records of cherts resulted in Failed Matches as their Detailed_Lithogy level matched with "banded iron formation", it occurs when intercalated together such as "cherts with BIF" or as include string descriptors such as "BIF-fy".



Accuracv=0.739

	precision	recall	f1-score	support
banded_iron_formation	0.03	0.67	0.05	3
chert	0.65	0.48	0.55	156
conglomerate	0.33	1.00	0.50	1
ironstone	0.41	0.49	0.45	47
mudstone	0.92	0.83	0.88	548
sandstone	0.63	0.73	0.67	44
silcrete	0.25	1.00	0.40	1
macro avg	0.46	0.74	0.50	800
weighted avg	0.82	0.74	0.77	800

725 Figure 6. Confusion matrix for sedimentary rocks comparing the fuzzy string matching results from the Lithology Code workflow (vertical axis) and Comments workflow (horizontal axis). The heatmap shows the values normalised to the support size to address the imbalance between classes. The values shown in the cells indicate the number of samples classified for the class. Empty cells indicate zero samples. The accuracy is 73.9%, with a weighted average precision of 82% and recall of 74%. These results are taken from a subset of 800 samples across 7 classes.

3.3.4 Metamorphic Rocks

Out of a total of 61 metamorphic rock entries, 60 are matched correctly (Fig 14). Most of these are "schists" as the YSGB area is rich in talc-carbonate schists. The Company_Litho entry "amphibolite mica schist" which is matched as "amphibolite" matches as "schist" in the Comments workflow. This is considered a Related Match.

735



740

Figure 7. Confusion matrix for metamorphic rocks comparing the fuzzy string matching results from the Lithology Code workflow (vertical axis) and Comments workflow (horizontal axis). The heatmap shows the values normalised to the support size to address the imbalance between classes. The values shown in the cells indicate the number of samples classified for the class. Empty cells indicate zero samples. The accuracy is 98.4%, with a weighted average precision of 98% and recall of 98%. These results are taken from a subset of 61 samples across 4 classes.

3.3.5 Surficial Rocks

Fuzzy string matching accuracy of surficial rocks scored a 45% on a total of 69 entries (Fig 15). Saprolites are matched as saprolite (Exact Match), rock (Failed Match) and saprock (Close Match). In instances where saprock is inputted as "sap rock",

- 745 it results to a failed match as "rock". "Soil" is commonly used in logs to refer to the first intercept of highly weathered, clayrich and unidentifiable intercept. "Soil" is classified with the highest variability of terms: "soil" (Exact Match), "rock" (Failed Match), "duricrust" (Close Match), "colluvium" (Related Match) and "calcrete" (Close Match). "Laterite" is matched to "colluvium" (Related Match), "duricrust" (Close Match) and "lag" (Close Match). "Lag" generally matches with "colluvium: (Related Match). However, when described in Comments, it can be associated with its protolith which results into a Failed
- 750 Match as "rock".



Figure 8. Confusion matrix for surficial comparing the fuzzy string matching results from the Lithology Code workflow (vertical axis) and Comments workflow (horizontal axis). The heatmap shows the values normalised to the support size to address the imbalance between classes. The values shown in the cells indicate the number of samples classified for the class. Empty cells indicate zero samples. The accuracy is 44.9%, with a weighted average precision of 57% and recall of 45%. These results are taken from a subset of 69 samples across 9 classes.

4 Discussion

4.1 dh2loop Functions and Notebooks

The *dh2loop* library supports a workflow that extracts, processes and classifies lithological logs (Appendix A4). This library is built to extract drill hole logs from the WAMEX database. The assumptions made in the entire workflow attempts to replicate the thought process of a geologist performing the data extraction, data quality checks and lithological log classification manually. However, it can be adapted for other geological relational databases or from other table formats. An example using comma separated values tables (CSVs) is shown in the notebook: Exporting and Text Parsing of drill hole Data Demo.

- 765 In addition to the data extraction, downhole desurveying and lithological matching functions discussed, *dh2loop* also provides functionalities and a notebook demonstrating harmonization of drill hole data. This is useful for combining and correlating drill hole exports of different properties such as lithology, assays and alteration. It is also possible to export this information in Visualization Toolkit format (.VTK). It also provides a notebook that demonstrates the application of *lasio* and *striplog* on *dh2loop* interval table exports. WAMEX reports can also be interactively downloaded through a notebook provided in the
- 770 package.

4.2 Thesauri

dh2loop provides the user with 9 thesauri that deal with the extraction of collar, survey and lithology interval tables. For extraction of other properties, such as downhole alteration, geochemistry, mineralogy and structures, at least one thesaurus is needed for each attribute we would like to export. These thesauri are built manually by inspecting all the terminologies

- 775 available in the database. Although, creating them can be tedious, updating an existing thesaurus is as simple as adding and/or removing a word to the list. There are many other properties available in the database that could be exploited using the existing methodology, thus there is an incentive in finding a way to improve the methodology of building these thesauri. Analysis on the syntax of the existing thesauri may help in automating creation of other thesauri.
- 780 The Hierarchical Lithology thesaurus puts equal weight on each of the entries in the thesaurus. Knowing the geology in a user's area, the matching can be improved by adding more weight to prevalent lithologies through adding a bonus score.

4.3 Data Extraction

dh2loop supports data extraction of collar, survey and lithology interval tables. The main consideration in the data extraction
 is that the data retrieved is complete, relevant and useful. We would rather throw erroneous or questionable data out and have the rest with a high level of confidence, than the other way around. 93% of the available collar data in the area is extracted successfully. This can be improved by implementing alternative ways for retrieving RL and MaxDepth values. For example, if no RL values are fetched from the database, it could be fetched from open-source digital terrain models (DTM) and/or SRTM (Shuttle Radar Topography Mission). As for missing MaxDepth values, the maximum ToDepth values in the survey and/or

790 interval tables could be used.

The survey extraction rate of 86% is fairly good. *dh2loop* ensures that the Azimuth and Inclination values are sensible measurements before including them into the extracted output file. An improvement that could be implemented is to run an assessment on the deflection angles for each drill hole and flag intervals with unrealistic deflection angles.

795

The lithology extraction using the **Lithology Code workflow** shows that the bottle neck to its extraction rate is the extensiveness of the Drill Hole Lithology Codes Thesaurus. Since the thesaurus did not have the information for all companies

in the area, only 34% of the available information is retrieved. The extraction results for the **Comments workflow** cannot be compared with the **Lithology Code workflow** as only the intersection of both workflows is considered in this study.

800

4.4 Assessment of String Matching Results

The number of successful matches are dependent on the selected cut-off score. The selection of a cut-off score is a balance between the number of matched records and the exact match percentage. In this case study, we selected a cut-off score of 80 since this is where the number of exact matches plateaus (Fig. 10). A lower cut-off score could be used, depending on the

805 familiarity to the data and/or purpose of drillhole processing. For our case, we wanted to be as conservative as possible without being too stringent (cut-off score 100).

The string matching results highlights that geological drill core logging is prone to human error and bias, and result to incorrect logs. Sometimes even if the data is available and correct, it is not in format that can be directly extracted. For example, **Comments** are filled with a string description such as "same as above" and "-do-". Currently, for this case, *dh2loop* returns without a match, as replacing "same as above" requires building a dictionary for all possible permutations to refer to this. This is not included in the scope of this work. In the future, we could be able to search through the previous entries to retrieve the correct lithology. Furthermore, the code does not handle and check for inconsistencies in the logs. It only addresses the inconsistencies in nomenclature and not the logging itself. The string matching misclassification results illustrate that importance in the consistency and level of detail being put into logging and identifies differences in convention or uncoordinated logging among geologists. *dh2loop* provides a notebook that demonstrates using *striplog* to improve the consistency of the logs through data pruning and annealing. In the future, the geochemical compositions can be used to counter check and lithology assigned to the interval.

- 820 Comparing the string matching between the Lithology Code workflow and Comments workflow, the Lithology code workflow results to a higher matching rate, 86% of the extracted data is successfully matched. Comparing this subset to the Comments workflow, the matching rate is much lower at 16%. This shows that the Lithology Code workflow, while potentially tedious, results into a higher percentage of successful matches. However, if we are considering a regional study involving multiple companies and drilling campaigns, building thesauri can be time-consuming depending on the size of the
- 825 region being studied, number of attributes of interest, number of companies and drilling campaigns. This could range from a couple of hours to months. It can also be tedious as it involves inputting errors and inconsistencies as well as exhausting all permutations for decision-tree based logging systems. The thesauri provided by *dh2loop* could serve as a starting point to automate this process using recent advances in NLP and machine learning.
- 830 String matching using **Comments** provides a quicker way to standardize and classify rocks. The comprehensive Clean-up Dictionary allows assists in improving the matching accuracy. Given the context that we are dealing with legacy data, an extraction rate of 16% Although it is a low extraction rate, there is value in being able to obtain 7,870 records more than what is previously deemed "unusable". With minimal effort, we obtain additional geological data wherein, although of a smaller percentage (31% of Exact Matches) but with reasonably high confidence in its quality. It is important to note that most of the
- time Failed Matches are not a result of the limitations of the algorithm but of the legacy geological logs itself. Inconsistent logs (Company_Litho data is different from Comments) usually occur when:
 - 1. The logs are post-processed and correlated with the rest of the hole or neighbouring drill holes and changes are made to the Company_Litho but none on the Comments field.

- 840
- The Comments would have more level of detail than the Company_Litho. In this case, we may get a lithology at Lithology_Subgroup from the Lithology Code workflow and a Detailed_Lithology from the Comments workflow.
- 3. The Company_Litho would have more level of detail than the Comments
- 4. **Comments** contains the description of the whole intercept, which could include a contact of two lithologies or intercalating lithologies.

845

850

855

860

From the results of the confusion matrix (Sect. 3.4), some rock groups are more sensitive to these inconsistencies than others. There is higher confidence in the classification of structures and textures and metamorphic rocks in the study area dataset, not necessarily in others. There could be metamorphic-dominated terranes where the subordinate igneous rocks will be classified with higher confidence. The user should be more careful when dealing with sedimentary and surficial rocks. They are more difficult to classify as the way they are described are highly variable between different geologists. For structure-related lithological descriptions the small number of misclassifications occur where faults, veins and fillings coexist. For metamorphic rocks, entries like "mica amphibolite schist" can cause Broader Matches with the confusion of whether to classify it as "amphibolite" or "schist". "Schist" is a textural term of medium grade metamorphic rock with a medium to coarse-grained foliation defined by micas while "amphibolite" is a compositional term representing a granular metamorphic rock which mainly consists of hornblende and plagioclase. One should be wary about these possibilities as they may impact the interpretation of the geology in the area. For sedimentary rocks, descriptions of intercalated lithologies or presence of major and minor lithology can result to Failed Matches. The lack of a standard syntax as to how free text descriptions are recorded impacts the classification. This procedure provides a basis for creating a pre-standard. Not so much providing a guide of practice but highlighting what should not be done and what practices create ambiguity. Standardization will definitely reduce subjectivity and is for the geological surveys to decide and implement. It is also important to note that a "standard" would be tricky to achieve as the information and level of detail contained in logs is highly dependent on the purpose of the study. Igneous rocks perform fairly well, most of what is not captured as Exact Matches are captured at least as Broader Matches. These are usually related to either an inconsistent level of detail between the fields or rock types used as descriptors

865

("komatiitic", "andesitic", basaltic").

Low matching accuracy in surficial rocks can be attributed to the lack of universally agreed terminology for: deeply weathered regolith; poorly-defined and misapplied surficial rock nomenclature; wide range and variation of materials within the regolith and; difficulty in bulk mineral identification from macroscopic samples. Furthermore, since the degree of weathering of minerals generally increases from the bottom to the top of in-situ weathering profiles, the intermixing of strongly weathered

- and less weathered grains may cause confusion (Cockbain, 2002). Ubiquitous, highly variable and less interesting lithologies also cause mismatches. An example of this is "soil". Soils are technically are not rocks but is commonly used in logs to refer to the first intercept of the regolith or to describe highly weathered, clay-rich and unidentifiable intercept. Soils vary in character from thin, coarse-grained, poorly differentiated lithosols to thick, well-differentiated silt and clay-rich soils. Soils are classified with the highest variability of terms: "soil", "rock", "duricrust", "colluvium" and "calcrete". There are also certain
- 875 lithologies with ambiguous nomenclature conventions, like "laterite", "duricrust", "lag". Some geologists use laterite to refer to the whole lateritic profile (ferruginous zone, mottled zone, and saprolite) while others to refer to the ferruginous zone (Eggleton, 2001). Ironcrust, duricrust, lateritic gravels and lag are commonly used interchangeably. Duricrust and ironcrust are terms to describe ferruginous indurated accumulations at or just below the surface. The difference in usage of the term laterite and the interchangeability of duricrust and lag explains the misclassification of "laterite" to "colluvium", "duricrust"
- and "lag". Another example is "saprolite" and "saprock". They are ambiguous terminologies as they both represent the lower horizons of lateritic weathering profiles, with saprolites having more than 20% of weatherable minerals altered and saprock

having less than 20% of the weatherable minerals being altered (Eggleton, 2001). This arbitrary limit makes the terminology used in the logs easily interchangeable, thus affecting the Detailed_Lithology matching.

885 Ideally, a combination of the **Lithology Code workflow** and **Comments workflow** should result in a more robust classification. This will also allow the user to have a better look at the result of both workflows and decide what is appropriate for one's purpose.

4.5 Value of the Lithological Information Extracted for Multiscale Analyses

- 890 The *dh2loop* lithology export provides a standardized lithological log across different drilling campaigns. This information can readily imported into 3D visualization and modelling software. This allows for drill hole data to be incorporated into 3D modelling, providing better subsurface constraints, especially at a regional scale. It also allows the user to decide on the lithological resolution necessary for their purpose. It provides a three-level hierarchical scheme: Detailed_Lithology, Lithology_Subgroup and Lithology_Group that can be used as an input to multiscale geological modelling.
- 895 *dh2loop* can be improved by correlating the these lithologies to their corresponding stratigraphic formations. Having the spatial extents of the different geological formations and their lithological assemblages (GSWA Explanatory Notes System) as well stratigraphic drill holes, it may be possible to infer the corresponding stratigraphic formation.

5 Conclusions

The *dh2loop* library is an open-source library that extracts geological information from a legacy drill hole database. This workflow has the following advantages:

- 1. Maximizes the amount of legacy geoscientific data available for analysis and modelling.
- 2. Provides better subsurface characterization and critical inputs to 3D geological modelling
- 3. Standardizes geological logs across different drilling campaigns, a necessary but typically time-consuming and errorprone activity
- 905
- 4. Provides a set of complementary thesauri that are easily updated and are individually useful references
 - 5. Implements a hierarchical classification scheme that can be used as an input to multiscale geological modelling
 - 6. Classification results can also be used as a tool to improve future geological logging works by revealing common errors and sources of inconsistencies

Code and Data Availability

910 *dh2loop* is a free, open-source python library licensed under the MIT License. It is hosted on the GitHub repository <u>https://github.com/Loop3D/dh2loop</u> and can be cited as http://doi.org/10.5281/zenodo.4043568.

Author Contribution

M. Jessell contributed the original idea, which ed further developed by R. Joshi. K. Madaiah developed the code. M. Jessell, M. Lindsay, G. Pirot provided guidance and direction in the research. R. Joshi prepared the manuscript with contributions

915 from all co-authors. Lastly, M. Jessell supervised the entire process.

Acknowledgements

The research was carried out while the first author was supported in receipt of Scholarship for International Research Fees (Australian Government Research Training Program Scholarship) and Automated 3D Geology Modelling PhD Scholarship (University Postgraduate Award) at the University of Western Australia. The work has been supported by the Mineral Exploration Cooperative Research Centre whose activities are funded by the Australian Government's Cooperative Research Centre Program. This is MinEx CRC Document 2020/***. This work was also done with the Loop Consortium

920

Centre Program. This is MinEx CRC Document 2020/***. This work was also done with the Loop Consortium (http://loop3d.org) as part of an international effort to found a new open-source platform to build the next generation of 3D geological modelling tools. Mark Lindsay is funded by ARC Discovery DE190100431. We would also like to acknowledge Tim Ivanic for his inputs on the geology of the Yalgoo-Singleton greenstone belt.

925 Appendix A: *dh2loop* package information

A1 Conventions and Terminologies

Convention	Usage in the paper	Description/Repository
Python libraries are	dh2loop	
written in italics	fuzzywuzzy	Python package for fuzzy logic for string matching
	nandas	Python package for data analysis and manipulation
	punuus	(McKinney 2011)(McKinney 2011)(McKinney
		2011)(McKinney, 2011)
	psvcopg2	Python package for PostgreSQL database adapter for
		python
	питру	питру
	nltk	Python package for Natural Language Toolkit
	pyproj	Python package for cartographic projections and
		coordinate transformations library
Python functions are	ratio ()	<i>fuzzywuzzy</i> functions
written in italics followed	partial_ratio ()	-
by an open and close	token_set_ratio ()	4
parentnesis	token_sort_ratio ()	4
D.4.1	partial_token_set_ratio ()	
Database tables are	collarattr	It contains main collar information
Italias	dhsurvey	It contains collar additional information
Italies	dhsurvevattr	It contains main survey information
	dhaeology	It contains survey additional information
	dhaeology	It contains geology information
Database table fields are	CollarID	It is the primary key from the collar table. It is the
written in Lucida Console		Unique ID field that identifies drill hole. It is used to
written in Euclua Console		associate data in different tables with a single drill hole
	HoleID	This is the drill hole name as the company would
		internally identify the drill hole.
	Longitude	The geographical longitude coordinate locating the collar
		of the drill hole.
	Latitude	The geographical latitude coordinate locating the collar of
		the drill hole.
	CompanyID	Unique ID field that identifies the company used
	DHSurveyID	Unique ID field that identified unique drill hole and depth
	Donth	location
	Depth	It refers to the downhole depth where the survey
	DHGeologyTD	Linique ID field that identified unique drill hole and denth
		interval
	FromDepth	The start/from and end/to downhole depth values (meters)
	ToDepth	The end/to downhole depth values (meters)
Output fields are written	KL	Relative Level refers to the Z coordinate of the collar
in Lucida Console	MayDonth	location (meters).
	махдертп	This refers to the maximum downhole length (meters)
		drilled for a drill hole, commonly referred as the end-of-
	X	noie. It is the coloulated Northing (maters)
	× v	It is the calculated Northing (meters)
	7	It is the calculated 7 position (meters)
	Azimuth	It is the trend direction indicated by an angle between 0
		360 degrees from the north going clockwise
	Inclination	It is the plunge angle of the drill hole relative to horizontal
		indicated by an angle between -90 to 90. It is measured
		from the horizontal plane, thus a positive value indicates
		an upward-directed drill hole and a negative value
		indicates a drill hole directed downwards.

	Company_LithoCode	This fetches the lithology codes that are typically three-
		letter codes using the Drill Hole Lithology Thesaurus.
	Company_Litho	This value is fetched by matching the CompanyID and
		Company_LithoCode to the Drill Hole Lithology
		Codes Thesaurus.
	Comments	It is the free text descriptions from <i>dhgeologyattr</i>
	Detailed_Lithology	This value is the lowest level lithology matched through
		fuzzy string matching.
	Lithology_Subgroup	This value is the subgroup level lithology matched
		through fuzzy string matching.
	Lithology_Group	This value is the highest/group level lithology matched
		through fuzzy string matching.
Workflows are written in	Lithology Code workflow	Workflow to decode Company_LithoCode
Century Gothic Bold	Comments workflow	Workflow to decode Comments
Thesurus	Drill Hole Collar Elevation Thesaurus	https://github.com/Loop3D/dh2loop/blob/master/thesauri
(https://github.com/Loop		/thesaurus_collar_elevation.csv
3D/dh2loop/blob/master/	Drill Hole Maximum Depth Thesaurus	https://github.com/Loop3D/dh2loop/blob/master/thesauri
<u>thesauri/)</u>		/thesaurus_collar_maxdepth.csv
	Drill Hole Survey Azimuth Thesaurus	https://github.com/Loop3D/dh2loop/blob/master/thesauri
		/thesaurus_survey_azimuth.csv
	Drill Hole Survey Inclination Thesaurus	https://github.com/Loop3D/dh2loop/blob/master/thesauri
		/thesaurus_survey_inclination.csv
	Drill Hole Lithology Thesaurus	https://github.com/Loop3D/dh2loop/blob/master/thesauri
		/thesaurus_geology_lithology.csv
	Drill Hole Comments Thesaurus	https://github.com/Loop3D/dh2loop/blob/master/thesauri
		/thesaurus_geology_comment.csv
	Drill Hole lithology Codes Thesaurus	https://github.com/Loop3D/dh2loop/blob/master/Thesaur
		i/thesaurus_geology_lithology_code.csv
	Clean-up Dictionary	https://github.com/Loop3D/dh2loop/blob/master/thesauri
		/thesaurus_cleanup.csv
	Lithology Hierarchical Thesaurus	https://github.com/Loop3D/dh2loop/blob/master/thesauri
		/thesaurus_geology_hierarchical.csv

A2 Installation and Dependencies

Installing *dh2loop* can be done by cloning the GitHub repository with \$ git clone https://github.com/Loop3D/dh2loop.git and

930 then manually installing it by running the python setup script in the repository: \$ python setup.py install

It primarily depends on a number of external open-source libraries:

- 1. fuzzywuzzy (https://github.com/seatgeek/fuzzywuzzy) which uses fuzzy logic for string matching (Cohen, 2011)
- 2. pandas (https://pandas.pydata.org/) for data analysis and manipulation (McKinney, 2011)
- psycopg2 (<u>https://pypi.org/project/psycopg2/</u>), a PostgreSQL database adapter for python (Gregorio and Varrazzo, 2018)
 - 4. numpy (<u>https://github.com/numpy/numpy</u>)
 - 5. *nltk* (<u>https://github.com/nltk/nltk</u>), the Natural Language Toolkit is a suite of open source Python modules, data sets, and tutorials supporting research and development in Natural Language Processing (Loper and Bird, 2002).
- 940 6. *pyproj* (<u>https://github.com/pyproj4/pyproj</u>), python interface to PROJ (cartographic projections and coordinate transformations library)

A3 Documentation

935

dh2loop's documentation provides a general overview over the library and multiple in-depth tutorials. The tutorials are provided as Jupyter Notebooks, which will provide the convenient combination of documentation and executable script blocks in one document. The notebooks are part of the repository and located in the notebooks folder. See http://jupyter.org/ for more information on installing and running Jupyter Notebooks.

A4 Jupyter notebooks

- 950 Jupyter notebooks are provided as part of the online documentation. These notebooks can be executed in a local python environment (if the required dependencies are correctly installed). In addition, static versions of the notebooks can currently be inspected directly on the *github* repository web page or through the use of *nbviewer*.
 - WAMEX Interactive report downloads
 (https://github.com/Loop3D/dh2loop/blob/master/notebooks/0_WAMEX_Downloads_Interactive.ipynb)
- 2. Exporting and text parsing of drill hole data from PostgreSQL database (https://github.com/Loop3D/dh2loop/blob/master/notebooks/1_Exporting_and_Text_Parsing_of_Drillhole_Data_Fr om_PostgreSQL.ipynb)
 - Exporting and Text Parsing of drill hole Data Demo
 (https://github.com/Loop3D/dh2loop/blob/master/notebooks/2_Exporting_and_Text_Parsing_of_Drillhole_Data_D
- 960 <u>emo.ipynb</u>)
 - Harmonizing drill hole data (<u>https://github.com/Loop3D/dh2loop/blob/master/notebooks/3_Harmonizing_Drillhole_Data.ipynb</u>)

Ailleres, L., Jessell, M., de Kemp, E., Caumon, G., Wellmann, F., Grose, L., Armit, R., Lindsay, M., Giraud, J., Brodaric, B., Harrison, M., and Courrioux, G.: Loop - Enabling 3D stochastic geological modelling, ASEG Extended Abstracts, 2019, 1-3, 10.1080/22020586.2019.12072955, 2019.

Anand, R. R., and Butt, C. R. M.: A guide for mineral exploration through the regolith in the Yilgarn Craton, Western Australia, 970 Australian Journal of Earth Sciences, 57, 1015-1114, Pii 929860728

10.1080/08120099.2010.522823, 2010.
Arabjamaloei, R., Edalatkha, S., Jamshidi, E., Nabaei, M., Beidokhti, M., and Azad, M.: Exact Lithologic Boundary Detection Based on Wavelet Transform Analysis and Real-Time Investigation of Facies Discontinuities Using Drilling Data, Petroleum Science and Technology, 29, 569-578, Pii 933125287
975 10 1080/10916460903419206 2011

10.1080/10916460903419206, 2011. Barley, M. E., Brown, S. J. A., Krapez, B., and Kositcin, N.: Physical volcanology and geochemistry of a Late Archaean volcanic arc: Kurnalpi and Gindalbie Terranes, Eastern Goldfields Superterrane, Western Australia, Precambrian Research, 161, 53-76, 10.1016/j.precamres.2007.06.019, 2008.

Chace, F. M.: Abbreviations in field and mine geological mapping, Economic Geology, 51, 712-723, 1956.

980 Cockbain, A. E.: Regolith geology of the Yilgarn Craton - Introduction, Australian Journal of Earth Sciences, 49, 1-1, DOI 10.1046/j.1440-0952.2002.00913.x, 2002.

Cohen, A.: FuzzyWuzzy: Fuzzy string matching in python, ChairNerd Blog, 22, 2011.

Culshaw, M. G.: From concept towards reality: developing the attributed 3D geological model of the shallow subsurface, Quarterly Journal of Engineering Geology and Hydrogeology, 38, 231-+, Doi 10.1144/1470-9236/04-072, 2005.

985 Eggleton, R. A.: The regolith glossary, Cooperative Centre for Landscape Evolution and Mineral Exploration, National Capital Printing: Canberra, 2001.

Emelyanova, I., Pervukhina, M., Clennell, M., and Dyt, C.: Unsupervised identification of electrofacies employing machine learning, 2017.

Erkan, G., and Radev, D. R.: LexRank: Graph-based lexical centrality as salience in text summarization, Journal of Artificial Intelligence Research, 22, 457-479, DOI 10.1613/jair.1523, 2004.

Fuentes, I., Padarian, J., Iwanaga, T., and Vervoort, R. W.: 3D lithological mapping of borehole descriptions using word embeddings, Computers & Geosciences, 141, ARTN 104516 10.1016/j.cageo.2020.104516, 2020.

Gillespie, M., and Styles, M.: BGS rock classification scheme, Volume 1. Classification of igneous rocks, 1999.

- 995 Griffin, R. E.: When are Old Data New Data?, GeoResJ, 6, 92-97, 10.1016/j.grj.2015.02.004, 2015. Groves, D. I., Goldfarb, R. J., Knox-Robinson, C. M., Ojala, J., Gardoll, S., Yun, G. Y., and Holyland, P.: Late-kinematic timing of orogenic gold deposits and significance for computer-based exploration techniques with emphasis on the Yilgarn Block, Western Australia, Ore Geology Reviews, 17, 1-38, Doi 10.1016/S0169-1368(00)00002-0, 2000. Hall, M., and Keppie, F.: Striplog: new open source software for handling and analysing discontinuous and qualitative data,
- 1000 2016.

Hallsworth, C. R., and Knox, R.: BGS rock classification scheme. Volume 3, classification of sediments and sedimentary rocks, 1999.

Higgins, R. F., and Mehta, S.: SeatGeek, 2018.

Hill, E. J., Robertson, J., and Uvarova, Y.: Multiscale hierarchical domaining and compression of drill hole data, Computers & Geosciences, 79, 47-57, 10.1016/j.cageo.2015.03.005, 2015.

Hill, E. J., Pearce, M. A., and Stromberg, J. M.: Improving Automated Geological Logging of Drill Holes by Incorporating Multiscale Spatial Methods, Mathematical Geosciences, 1-33, 10.1007/s11004-020-09859-0, 2020.

Jallan, Y., Brogan, E., Ashuri, B., and Clevenger, C. M.: Application of Natural Language Processing and Text Mining to Identify Patterns in Construction-Defect Litigation Cases, Journal of Legal Affairs and Dispute Resolution in Engineering and Construction, 11, Unsp 04519024

- 10.1061/(Asce)La.1943-4170.0000308, 2019.
 Kumari, S., Mohan, A., and Saberwal, G.: Hidden duplicates: 10s or 100s of Indian trials, registered with ClinicalTrials.gov, have not been registered in India, as required by law, PLoS One, 15, e0234925, 10.1371/journal.pone.0234925, 2020.
 Lark, R. M., Thorpe, S., Kessler, H., and Mathers, S. J.: Interpretative modelling of a geological cross section from boreholes:
- sources of uncertainty and their quantification, Solid Earth, 5, 1189-1203, 10.5194/se-5-1189-2014, 2014.
 Le Vaillant, M., Hill, J., and Barnes, S. J.: Simplifying drill-hole domains for 3D geochemical modelling: An example from the Kevitsa Ni-Cu-(PGE) deposit, Ore Geology Reviews, 90, 388-398, 10.1016/j.oregeorev.2017.05.020, 2017.
 Lin, C.-Y., and Hovy, E.: Automatic evaluation of summaries using n-gram co-occurrence statistics, Proceedings of the 2003 Human Language Technology Conference of the North American Chapter of the Association for Computational Linguistics,
- Human Language Technology Conference of the North American Chapter of the Association for Computational Linguistics,
 2003, 150-157,
 Lindsay, M.: Geological Interpretation of Geophysics Support from Machine Learning of Drillcore, 2019.
 Liu, K., Hogan, W. R., and Crowley, R. S.: Natural Language Processing methods and systems for biomedical ontology learning, J Biomed Inform, 44, 163-179, 10.1016/j.jbi.2010.07.006, 2011.
 Liu, T. and Guo, L: Text similarity computing based on standard deviation. International Conference on Intelligent
 - Liu, T., and Guo, J.: Text similarity computing based on standard deviation, International Conference on Intelligent Computing, 2005, 456-464,
 - Loper, E., and Bird, S.: NLTK: the natural language toolkit, arXiv preprint cs/0205028, 2002.

McKinney, W.: pandas: a foundational Python library for data analysis and statistics, Python for High Performance and Scientific Computing, 14, 2011.

McMillan, A., and Powell, J.: British Geological Survey Rock Classification Scheme: The Classification of Artificial (man made) Ground and Natural Superficial Deposits: Applications to Geological Maps and Datasets in the UK, British Geolgoical Survey Research Report RR/99/4, 1999.
 Wiles A. and Bachhofm S.: SKOS simple knowledge experimentation system references. W2C recommendation, 18, W2C, 2000.

Miles, A., and Bechhofer, S.: SKOS simple knowledge organization system reference, W3C recommendation, 18, W3C, 2009.
 Müller, T., Cotterell, R., Fraser, A., and Schütze, H.: Joint lemmatization and morphological tagging with lemming,
 Proceedings of the 2015 Conference on Empirical Methods in Natural Language Processing, 2015, 2268-2274,

Myers, J.: Precambrian Tectonic History of the West Australian Craton and Adjacent Orogens, Annu Rev Earth Pl Sc, 21, 453-485, 10.1146/annurev.ea.21.050193.002321, 1993.
 Okuda, T., Tanaka, E., and Kasai, T.: A method for the correction of garbled words based on the Levenshtein metric, IEEE

Transactions on Computers, 100, 172-178, 1976.

Otter, D. W., Medina, J. R., and Kalita, J. K.: A Survey of the Usages of Deep Learning for Natural Language Processing, 1040 IEEE Trans Neural Netw Learn Syst, 10.1109/TNNLS.2020.2979670, 2020.

Padarian, J., and Fuentes, I.: Word embeddings for application in geosciences: development, evaluation and examples of soil-related concepts, 10.5194/soil-2018-44, 2019.

Park, S. H., Ryu, K. H., and Gilbert, D.: Fast similarity search for protein 3D structures using topological pattern matching based on spatial relations, Int J Neural Syst, 15, 287-296, 10.1142/S0129065705000244, 2005.

- Qiu, Q., Xie, Z., Wu, L., and Tao, L.: Dictionary-Based Automated Information Extraction From Geological Documents Using a Deep Learning Algorithm, Earth and Space Science, 7, 10.1029/2019ea000993, 2020.
 Ralph, J.: Mindat. org-the mineral database, Mindat, Surrey, England, 2004.
 Raymond, O., Duclaux, G., Boisvert, E., Cipolloni, C., Cox, S., Laxton, J., Letourneau, F., Richard, S., Ritchie, A., and Sen, M.: GeoSciML v3. 0-a significant upgrade of the CGI-IUGS geoscience data model, EGUGA, 2711, 2012.
- 1050 Recasens, M., Danescu-Niculescu-Mizil, C., and Jurafsky, D.: Linguistic models for analyzing and detecting biased language, Proceedings of the 51st Annual Meeting of the Association for Computational Linguistics (Volume 1: Long Papers), 2013, 1650-1659,

Richard, S. M., Boisvert, E., Brodaric, B., Cox, S., and Duffy, T.: GeoSciML-a GML application for geoscience information interchange, Philadelphia Annual Meeting, 2007, 47-59,

- 1055 Riganti, A., Farrell, T. R., Ellis, M. J., Irimies, F., Strickland, C. D., Martin, S. K., and Wallace, D. J.: 125 years of legacy data at the Geological Survey of Western Australia: Capture and delivery, GeoResJ, 6, 175-194, 10.1016/j.grj.2015.02.015, 2015. Rivera-Quiroz, F. A., and Miller, J.: Extracting Data from Legacy Taxonomic Literature: Applications for planning field work, Biodiversity Information Science and Standards, 3, 10.3897/biss.3.37082, 2019. Robertson, S.: BGS rock classification scheme. Volume 2, Classification of metamorphic rocks, 1999.
- Rosenbaum, M. S., McMillan, A. A., Powell, J. H., Cooper, A. H., Culshaw, M. G., and Northmore, K. J.: Classification of artificial (man-made) ground, Engineering Geology, 69, 399-409, 10.1016/S0013-7952(02)00282-X, 2003.
 Ross, P. S., Bourke, A., and Fresia, B.: A multi-sensor logger for rock cores: Methodology and preliminary results from the Matagami mining camp, Canada, Ore Geology Reviews, 53, 93-111, 10.1016/j.oregeorev.2013.01.002, 2013.
 Rothwell, R. G., and Rack, F. R.: New techniques in sediment core analysis: an introduction, New Techniques in Sediment
- 1065 Core Analysis, 267, 1-29, Doi 10.1144/Gsl.Sp.2006.267.01.01, 2006. Schetselaar, E. M., and Lemieux, D.: A drill hole query algorithm for extracting lithostratigraphic contacts in support of 3D geologic modelling in crystalline basement, Computers & Geosciences, 44, 146-155, 10.1016/j.cageo.2011.10.015, 2012. Simons, B., Boisvert, E., Brodaric, B., Cox, S., Duffy, T. R., Johnson, B. R., Laxton, J. L., and Richard, S.: GeoSciML: enabling the exchange of geological map data, ASEG Extended Abstracts, 2006, 1-4, 2006.
- 1070 Smith, M. J., Keesstra, S., and Rose, J.: Use of legacy data in geomorphological research, GeoResJ, 6, 74-80, 10.1016/j.grj.2015.02.008, 2015.
 Vearncombe, J., Conner, G., and Bright, S.: Value from legacy data, T I Min Metall B, 125, 231-246, 10.1080/03717453.2016.1190442, 2016.

Vearncombe, J., Riganti, A., Isles, D., and Bright, S.: Data upcycling, Ore Geology Reviews, 89, 887-893, 1075 10.1016/j.oregeorev.2017.07.009, 2017.

Wang, C., and Ma, X.: Text Mining to Facilitate Domain Knowledge Discovery, in: Text Mining-Analysis, Programming and Application, IntechOpen, 2019.

Wilbur, W. J., and Sirotkin, K.: The Automatic Identification of Stop Words, Journal of Information Science, 18, 45-55, Doi 10.1177/016555159201800106, 1992.

 Zhou, C. Y., Ouyang, J. W., Ming, W. H., Zhang, G. H., Du, Z. C., and Liu, Z.: A Stratigraphic Prediction Method Based on Machine Learning, Appl Sci-Basel, 9, ARTN 3553
 10.3390/app9173553, 2019.
 Zhou, O. Liu, H. H. Bodyarsson, G. S. and Oldenburg, C. M.: Flow and transport in unsaturated fractured rock: effects of

Zhou, Q., Liu, H. H., Bodvarsson, G. S., and Oldenburg, C. M.: Flow and transport in unsaturated fractured rock: effects of multiscale heterogeneity of hydrogeologic properties, J Contam Hydrol, 60, 1-30, 10.1016/s0169-7722(02)00080-3, 2003.

1085 Zhu, G., Gao, M., Kong, F., and Li, K.: Application of Logging While Drilling Tool in Formation Boundary Detection and Geo-steering, Sensors (Basel), 19, 10.3390/s19122754, 2019.