

Investigating the importance of sub-grid particle formation in point source plumes over eastern China using IAP-AACM v1.0 with a sub-grid parameterization

Ying Wei^{1,2}, Xueshun Chen^{2,3,*}, Huansheng Chen², Yele Sun^{2,3,5}, Wenyi Yang^{2,3},
5 Huiyun Du², Qizhong Wu⁴, Dan Chen¹, Xiujuan Zhao¹, Jie Li^{2,3}, Zifa Wang^{2,3,5}

¹ Institute of Urban Meteorology, China Meteorology Administration, Beijing, 100089, China

² The State Key Laboratory of Atmospheric Boundary Layer Physics and Atmospheric Chemistry, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing
10 100029, China

³ Center for Excellence in Regional Atmospheric Environment, Institute of Urban Environment, Chinese Academy of Sciences, Xiamen 361021, China

⁴ College of Global Change and Earth System Science, Beijing Normal University, Beijing 100875

15 ⁵ University of Chinese Academy of Sciences, Beijing 100049, China

Correspondence: Xueshun Chen (chenxsh@mail.iap.ac.cn)

Abstract:

20 The influence of sub-grid particle formation (SGPF) in point source plumes on aerosol particles over eastern China was firstly illustrated by implementing a SGPF scheme into a global-regional nested chemical transport model with aerosol microphysics module. The key parameter in the scheme was optimized based on the observations in eastern China. With the parameterization of SGPF, the spatial
25 heterogeneity and diurnal variation of particle formation processes in sub-grid scale were well resolved. The SGPF scheme can significantly improve the model performance in simulating aerosol components and new particle formation processes at typical sites influenced by point sources. The comparison with observations at Beijing, Wuhan, and Nanjing showed that the normal mean bias (NMB) of sulfate and
30 ammonium could be reduced by 23%-27% and 12%-14%, respectively. When wind

fields were well reproduced, the correlation of sulfate between simulation and observation can be increased by 0.13 in Nanjing. Considering the diurnal cycle of new particle formation, the SGPF scheme can greatly reduce the overestimation of particle number concentration in nucleation and Aitken mode at night caused by fixed-fraction parameterization of SGPF. In the regional scale, downwind areas of point source got an increase of sulfate concentration by 25%-50%. The results of this study indicate the significant effects of SGPF on aerosol particles over areas with the point source and necessity of reasonable representation of SGPF processes in chemical transport models.

Key words: IAP-AACM, sub-grid particle formation, secondary inorganic aerosol, particle number concentration, China

1. Introduction

Air pollution caused by high concentrations of aerosol particles has become a ~~major~~primary environmental problem in metropolis and ~~caused-drew~~ widespread public ~~concern~~ attention in China (Zhang and Cao, 2015; Sun and Chen, 2017). Atmospheric aerosol particles **not only** affect atmospheric visibility and public health, **but also** has significant climatic and ecological effects (Zhang et al., 2010; Boucher et al., 2013; Powell et al., 2015; Yang et al., 2019, 2020), which are closely related to their size distribution and chemical composition (Spracklen et al., 2005; Dusek et al., 2006). As one of the main inorganic aerosol components, sulfate directly changes the energy budget of the earth-atmosphere system by scattering solar radiation, and indirectly affects the climate **by** acting as cloud condensation nuclei (CCN) (IPCC, 2013). Sulfuric acid (H_2SO_4) is the core material in the nucleation and growth of particles. The hygroscopicity of sulfate triggers heterogeneous reactions of gas precursors by changing the aerosol water content (Zhuang et al., 2014). Sulfate can also enhance ~~the~~ extinction and contribute to haze pollution by mixing with other components (Zhu et al., 2010; Zanatta et al., 2018), affecting the formation of secondary inorganic aerosol (Adams et al., 1999) and secondary organic aerosol (SOA) (He et al., 2018). Coal consumption contributed about 80% to sulfur dioxide (SO_2)

emission in China over the last decade, with 70%–90% being from power plants and industrial emissions (Wang et al., 2014; Ma et al., 2017; Zheng et al., 2018). Although emissions in China have declined since 2007 and ~~now~~ India has replaced ~~ds~~ China as the world's largest SO₂ emitter (Li et al., 2017), SO₂ from energy and industrial sectors is still the major source of sulfate in China.

For point sources, the oxidation from SO₂ to sulfate is a typical sub-grid phenomenon with a ~~more~~ rapid conversion rate in the plume. The concentration of nitrogen oxides (NO_x) and volatile organic compounds (VOCs) can influence ~~the~~ atmospheric oxidation through gas-phase chemical reactions and thus the OH concentration and the oxidation of SO₂. In the sulfur-rich plume, a swiftly gas-to-particle process occurs through the reaction $2\text{OH} + \text{SO}_2 \rightarrow \text{H}_2\text{SO}_4$ (Kulmala and Kerminen, 2008), due to the higher concentration of ~~nitrogen oxides (NO_x)~~ and ~~volatile organic compounds (VOCs)~~ in the plume than in the ambient atmosphere, and also due to the inhomogeneous meteorological condition (i.e., temperature, relative humidity (RH)) and aerosol concentrations within and external to the plume (Yu, 2010; Lonsdale et al., 2012; Stevens et al., 2012). H₂SO₄ can either condense onto pre-existing particles or nucleate to form new particles, which are treated as primary sulfate, or so-called sub-grid sulfate (SG-ASO₄). A significant increase in both mass and number concentrations of particles has been observed downwind of coal-fired power plants (Richards et al., 1981; Gillani et al., 1998; Brock et al., 2002). Yu (2010) showed that particle formation in the plume depends on hydroxyl radicals (OH) concentration, ~~which varies diurnally~~. The OH concentration varies diurnally and thus ~~the~~ fraction converted to sulfate ~~thus~~ has a clear diurnal variation, leading to spatial-temporal heterogeneity in local particle mass concentration and particle number size distribution (PNSD).

The ~~modeling-simulated~~ distributions of both particle number and component mass are sensitive to the fraction of SO₂ oxidized to sulfate in the plume. Currently, for SG-ASO₄, an average proportion of 0%–5% of total SO₂ emissions is often taken as H₂SO₄ emitted to every grid in chemical transport models (CTMs) (Textor et al., 2006). Furthermore, a ~~fracit~~ion of 0%–15% of H₂SO₄ is taken as the newly formed

90 particles ~~through nucleation~~ in aerosol microphysical models (Luo and Yu, 2011; Chen et al., 2018). The assumption of an averaged fraction of oxidation and nucleation in the grid neglects the diurnal cycle of sub-grid particle formation (SGPF). ~~and furthermore, the scheme~~ cannot capture the spatial-temporal variation of particle formation in the plume, nor does it account for the effect of H₂SO₄ condensing onto
95 pre-existing particles, which may have a significant impact on PNSD. A simulation in North China employed 26.5% of primary fine particulate matter as SG-ASO₄ indicated a monthly averaged contribution to sulfate of ~10%–20% (Zhang et al., 2012). The assumed proportion of SG-ASO₄ caused an uncertainty in global CCN concentration of up to 40%, and over 100% in polluted regions (Spracklen et al.,
100 2005). As SGPF may occur on a spatial scale of 10s km⁻¹, this leads to large uncertainties in predicting the spatio-temporal variation of particle number and mass concentration during dilution in the plume when it is calculated as a grid-averaged concentration, as in most CTMs (Spracklen et al., 2005, 2008).

To solve the uncertainties caused by grid-averaged fraction of SO₂-to-H₂SO₄, a
105 plume-in-grid (PinG) model ~~coupled coupling the smoke plume model~~ into the Eulerian Model has been developed in air quality models such as the Community Multiscale Air Quality model, the Comprehensive Air quality Model with extensions and the Weather Research and Forecasting/Chemistry (Gillani et al., 1999; Karamchandani et al., 2002, 2010). However, the implementation of PinG greatly
110 increases the computational burden due to the large amount of fine grids resolved. Due to this reason, PinG is not suitable for complicated aerosol models involving microphysical processes. Stevens et al. (2013) developed a computationally efficient sub-grid parameterized scheme, the ‘Predicting Particle Production in Power-Plant Plumes’ (P6) scheme, based on physicochemical processes of particle formation. The
115 scheme was incorporated in a global CTM with aerosol microphysics module and the sub-grid effect on particle number concentration was evaluated against observations over North America and Europe. However, urban sites were excluded from the evaluation and the sub-grid impact on aerosol components was not described. Up to now, ~~There are, in fact,~~ few modeling studies involving sub-grid particle

characteristics in high polluted regions such as China. The severe air pollution in China means that atmospheric chemical characteristics ~~there-in China~~ are quite different to those in other countries. Higher atmospheric oxidability and particle growth rates have been reported in recently researches (Wang et al., 2017; Tan et al., 2019; Yang et al., 2020; Liu et al., 2021). ~~This means~~ Therefore, OH concentration (key parameter of the oxidation process) parameterized by NO_x concentration in the P6 scheme is not suitable for the atmospheric condition in China, and the characteristics of SGPF in plumes should be different here.

In this study, we coupled the P6 scheme to a global-regional nested atmospheric chemistry model with an aerosol microphysics module to better describe the process of SGPF in plumes. Moreover, the localized SGPF scheme (refers to SGPF scheme) was ~~developed carried-out~~ based on the observed high level of radicals caused by the polluted background in central-eastern China. With the updated model, the improvements in simulating aerosol composition and PNSD were evaluated by comparing with abundant observations in eastern China. The original model and its updating are described in Sections 2.1–2.3. Simulation experiments and observations are introduced in Section 2.4 and 2.5, respectively. Meteorological fields are verified in Section 3.1. The evaluation of the updated model against observations and the model improvements ~~to the updated model against observations~~ are described ~~described~~ in Section 3.2–3.4. The influence of SGPF on regional scale is analyzed in Section 3.5.

2. Methods

2.1 Description of IAP-AACM

The Aerosol and Atmospheric Chemistry Model of Institute of Atmospheric Physics (IAP-AACM) is a multi-scale nested three-dimensional chemistry transport model coupled to the Earth System Model of the Chinese Academy of Sciences (CAS-ESM) (Wei et al., 2019; Zhang et al., 2020). The IAP-AACM was developed on the basis of the Nested Grid Air Quality Prediction Model System (NAQPMS) (Wang et al. 2006b) and the Global Nested Grid Air Quality Prediction Model System

(GNAQPMS) (Chen et al., 2015). NAQPMS/GNAQPMS are widely used in the simulation of dust (Li et al., 2012; Wei et al., 2019), ozone (Wang et al., 2006a; Li et al., 2007), deposition (Ge et al., 2014), air pollution control policy (Wu et al., 2011; Li et al., 2016; Wei et al., 2017) and global transportation of mercury (Chen et al., 2015). In the IAP-AACM, dimethyl_sulfide, sea salt and dust emissions are calculated online. The dust scheme originates from the wind erosion model developed by Wang et al. (2000) and improved by Luo et al. (2006).

The gas phase chemistry is calculated with the Carbon-Bond Mechanism Z model (CBM-Z; Zaveri and Peters, 1999). The calculation of photolysis rate ~~is calculated~~ mainly ~~considering~~ considered altitude, latitude, longitude and the effects of clouds. The rates of photolysis reactions depend on the spectral actinic flux and the spectral actinic flux depends on the absorption and scattering of incident solar radiation by gaseous molecules, clouds and aerosols. The photolytic rate constants typically increase with height due to the reduction in the total integrated optical depth (OD) with lower pressure, less aerosols and clouds (Seinfeld and Pandis, 2012; Williams et al., 2012). The gas-phase chemical mechanism has important impacts on NO_x and ultimately the OH concentration calculated in the SGPF scheme. Different mechanism may have different impacts on the parameterizedation ~~edation~~ result. Zhang et al. (2012) compared simulations conducted in summer with three different gas-phase mechanisms (i.e., CBM-Z, CB05 and SAPRC-99) in WRF-Chem and found that simulations with all three gas-phase mechanisms well reproduced the surface concentrations of O₃, CO, NO₂, and PM_{2.5}. Prediction discrepancies caused by different mechanisms were of mass concentrations of O₃ (up to 5 ppb), PM_{2.5} (up to 0.5 μg m⁻³), secondary inorganic PM_{2.5} species (up to 1.1 μg m⁻³). Overall, the simulation discrepancy between model with the CBM-Z mechanism and other widely used new mechanisms should be reasonably acceptable.

The aqueous chemistry and wet deposition scavenging is simulated with the Regional Acid Deposition Model chemical mechanism (Stockwell et al., 1990). The heterogeneous chemistry uses the scheme described by Li et al. (2012). For aerosol microphysical processes, the IAP-AACM ~~in this study~~ describes the size distribution

of aerosol particles using the Advanced Particle Microphysics (APM; Yu and Luo, 2009) module, as reported in previous studies (Chen et al., 2014; Chen et al., 2018). The APM in IAP-AACM uses 40 sectional bins to represent secondary particles formed from nucleation and subsequent growth with dry diameters of 0.0012–12 μm . Black carbon (BC) and organic carbon (OC) particles are represented by 28 bins. Sea salt and dust particles are represented by 20 bins and 4 bins, respectively. The APM assumes these particles are the cores of particles and they are coated with secondary species. Both the cores and coating species are tracked in the model. Semi-volatile aerosol species, including nitrate, ammonium and SOA, are simulated by the bulk method. Only their total mass concentrations are tracked and the concentrations apportioned to particles in different sizes are assumed to be proportional to the mass concentration of associated sulfate. Nitrate and ammonium are simulated by ISORROPIA version 1.7 (Nenes et al., 1998, 1999). SOA concentration is calculated with the scheme described by Strader et al. (1999). A comprehensive evaluation of the simulation of IAP-AACM from global to regional was shown in Wei et al. (2019).

2.2 Implementing sub-grid scheme into IAP-AACM

In the original version of the IAP-AACM, H_2SO_4 was emitted directly into grids ~~in-by using~~ a fixed proportion with 5% being emitted into nucleation mode and 95% condensing onto the existing accumulation mode particles in the APM module (Yu and Luo, 2009). We updated the IAP-AACM+APM ~~with-by coupling~~ the P6 sub-grid particle parameterization scheme (Stevens and Pierce, 2013) to resolve the dynamic variation of SGPF ~~in the global-nested model~~.

The P6 scheme ~~includes-simulates~~ the rapid conversion from SO_2 to sulfate within the plume, considering both computational efficiency and physical basis. The training data for constructing the P6 scheme are based on results of the large-eddy simulation/cloud-resolving model incorporated with a microphysics module, System for Atmospheric Modeling with Two Moment Aerosol Sectional (SAM-TOMAS) (Adams and Seinfeld, 2002; Kairoutdinov and Randall, 2003). The model results used to construct the P6 scheme have been tested against aircraft observations (Lonsdale et

al., 2012; Stevens et al., 2012). ~~For M~~more information on the model, ~~can~~ refer to Stevens et al. (2012).

210 The sub-grid scheme resolves SGPF into two key processes, namely oxidation and nucleation, ~~involving with involved~~ parameters from the meteorological field, emission source and environmental background as inputs. Accordingly, the oxidation of SO₂ emitted from a point source is constructed with meteorological conditions (i.e., wind speed (v_g), boundary layer height (BLH), downward shortwave radiative flux (DSWRF)), emissions of SO₂ and NO_x (NO_xemis) from the source, mean background
215 concentrations of SO₂ and NO_x (bgNO_x), and the distance from the source (d). Nucleation of H₂SO₄ in the plume is constructed with factors mentioned above and the mean background condensation sink. A scaling factor is used to allow the equations to fit the data when calculating the effective concentrations of NO_x and SO₂ within the
220 plume, based on the reality that the emitted fluxes and the resulting- of NO_x- and SO₂ concentrations in the plume are ~~very-much~~ different ~~from~~ the grid-mean values. A detailed calculation of the parameterization scheme is provided by Stevens et al. (2013).

Oxidation and nucleation process in the P6 scheme were integrated into the chemical
225 reaction and aerosol microphysics modules, respectively. The key parameter predicted during the oxidation process is the oxidation fraction of the emitted SO₂ (f_{ox}). The P6 scheme is combined with the gas-phase chemistry module to describe the dynamic variation of H₂SO₄ production within and outside the plume, based on the variation of meteorological conditions and environmental backgrounds. Emissions separated by
230 sector is used in IAP-AACM to analyze the impact of different emission sources. We used real-time online calculation of f_{ox} to repartition ~~the total SO₂ sub-grid H₂SO₄ production and SO₂~~ from energy and industrial sector sources to sub-grid H₂SO₄ production and SO₂.

For the nucleation process, the H₂SO₄ produced in oxidation process is distributed
235 to nucleate to new particles and condense onto pre-existing particles by the key parameter of new particle formation fraction (f_{new}). In the APM, the condensation of H₂SO₄ onto pre-existing particles (sulfate, BC, OC, dust and sea salt) are proportional

to the ~~corresponding ratio of~~ condensation sink ratio of the corresponding particles. Furthermore, the H₂SO₄ nucleating to newly formed particles is partitioned into 40
240 bins of secondary particles in the APM with a lognormal distribution as follows:

$$n_N(D_p) = \frac{dN}{dD_p} = \frac{N}{(2\pi)^{1/2} D_p \ln \sigma_g} \exp\left(-\frac{(\ln D_p - \ln \bar{D}_m)^2}{2 \ln^2 \sigma_g}\right) \quad (1)$$

where D_p and N represent the particle size and total number of particles, respectively; σ_g is the geometric standard deviation of the aerosol size distribution, ~~of~~ 1.4 in this study; and D_m is the number-median diameter [μm] calculated by

$$245 \quad D_m = \left(\frac{M_m}{\rho} \frac{6}{\pi}\right)^{\frac{1}{3}} \exp(-1.5 \ln^2 \sigma_g) \quad (2)$$

where M_m is the ~~the~~ mean mass of per newly formed particles [kg] predicted by the P6 scheme; ρ is the density of dry aerosol (1.7 g cm⁻³ in our model). After this repartitioning, the tracers of secondary particles are updated through moving mass concentrations of sulfate across bins in the APM module.

250 2.3 Optimization of the key parameter in the sub-grid scheme

Since OH is very important for the diurnal cycle of SG-ASO₄ conversion in the plume, the determination of OH concentration is crucial to the sub-grid scheme. In parameterizingation ~~of~~ the oxidation process, the key step is calculating f_{ox} in the plume, which depends on the rate constant, k , time elapsed, t , and the effective OH
255 concentration in the plume, OH_{eff}, [molecules cm⁻³]. OH_{eff} is calculated by:

$$\text{OH}_{\text{eff}} = 0.82 \cdot 10^{P1 \cdot \log(P2)/6.8} \quad (3)$$

where P1 and P2 are the function of the effective NO_x concentration in the plume (NO_{x,eff}, [ppb]) and the DSWRF, respectively, constructed from excessive training data ~~that~~ generated from SAM-TOMAS. P1 and P2 are dimensionless and calculated

260 as:

$$P1 = -0.014x^6 + 0.0027x^5 + 0.1713x^4 - 0.0466x^3 - 0.7893x^2 - 0.1739x + 6.9414 \quad (4)$$

$$P2 = (-1345y^3 + 4002y^2 - 471.8y + 42.72) \times 10^4 \quad (5)$$

$$y = \frac{DSWRF}{S_0 \cdot T} \quad (6)$$

where S_0 is the solar constant at the top of the atmosphere, 1370 W m^{-2} , and T is an assumed transmittance of the clear atmosphere, 0.76. The relationship between DSWRF and OH concentration in SGPF scheme was displayed in Fig. S1. The simulated OH concentration is under $1 \times 10^6 \text{ cm}^{-3}$ when DSWRF varies between 0-200 W m^{-2} . Thus, the fraction of total SO_2 emitted converted into particles in the plume much lower in cloudy days.

In the original P6 scheme, x was calculated from $\text{NO}_{x,\text{eff}}$ by:

$$x = \log([\text{NO}_x, \text{eff}]) - 0.195 \quad (7)$$

where $\text{NO}_{x,\text{eff}}$ is related to $\text{NO}_{x,\text{emis}}$ [kg N s^{-1}], bgNO_x [ppb], BLH [m], v_g [m s^{-1}] and d [m] as mentioned in Section 2.2. for the low-VOC case for the isoprene mixing ratio of <0.15 ppb, or:

$$x = \log([\text{NO}_x, \text{eff}] \times 0.6) - 0.195 \quad (8)$$

in the high-VOC case with the isoprene mixing ratio of 1.5 ppb. The effect of VOC within the plume is not explicitly taken into account in the parameterization of OH_{eff} . Stevens et al. (2012) discussed the potential effect of low and high VOC concentrations on OH production and related it to the concentration of NO_x that corresponds to the peak concentration of OH. The large amount of oxidation of isoprene has material effect on ~~peroxy~~ radicals and tends to shift the peak of OH production to a higher level. It is reasonable to expect that the true peak OH concentrations should be shifted to a higher NO_x than suggested by the P6 scheme. The upper limit of the background concentration of NO_x in the P6 scheme is ~ 8 ppb (Stevens and Pierce, 2013).

The NO_x concentration in eastern China is commonly at a high level of >20 ppb, so there will be an underestimation of OH production in polluted regions if the original P6 scheme is applied directly. We therefore adjusted the OH concentration parameter in the scheme to take into account the high oxidizability in the polluted atmosphere in eastern China. Although the OH_{eff} calculated in the SGPF scheme is independent of the OH concentration calculated by CBM-Z and only used to calculate the SO_2 oxidation fraction to sulfate in the plume, the grid-averaged OH concentration

calculated by CBM-Z has large impact on NO_x and VOCs and ultimately changes OH_{eff}. The simulation of NO₂ and OH in the base model has been validated in Wei et al. (2019). Overall, the model well reproduced the seasonal variation of NO₂ and captured the daily variations with R of 0.49-0.7 in most cities in eastern China. The global distribution of OH concentration in the IAP-AACM is similar with other CTMs. In addition, the simulations of the IAP-AACM with observations took at rural sites located in the North China Plain near Beijing (Tan et al., 2017) and the Pearl River Delta near Guangzhou City (Lu et al., 2012) in summer were compared in Fig. S2. Although the observation in Guangzhou was took in 2006, the characteristics of atmospheric oxidation under the regional atmospheric background can still be roughly reflected. As shown in Fig. S2, the model reproduced the diurnal characteristics of OH concentration well from south to north of China in summertime.

Field observations in North China indicate a considerable underestimation of OH levels in simulations for metropolitan areas. For warm seasons, Lu et al. (2013) reported daytime peak OH concentrations of $(4-17) \times 10^6 \text{ cm}^{-3}$ at a suburban site in Beijing in summer 2006. Tan et al. (2017) observed a daily OH maxima of $(5-15) \times 10^6 \text{ cm}^{-3}$ at a rural site in the North China Plain in summer 2014. In cold seasons, the radical concentrations are expected to be much lower than in summertime due to limited photochemistry, but high concentrations are still found in recently-measured records. Tan et al. (2018) observed the average daytime maximum OH concentrations of $2.5 \times 10^6 \text{ cm}^{-3}$ at another suburban site in Beijing in January 2016. Slater et al. (2020) found daily maximum OH concentrations of $(1-8) \times 10^6 \text{ cm}^{-3}$ with averaged daytime peak of $2.7 \times 10^6 \text{ cm}^{-3}$ over the whole observation period.

In view of these observations, we changed the coefficient of $[\text{NO}_{x,\text{eff}}]$ to 0.3 in the scheme as shown in Eq. (7) to avoid an underestimation of SGPF, on the basis of observations of OH and NO_x in eastern China:

$$x = \log([\text{NO}_{x,\text{eff}}] \times 0.3) - 0.195 \quad (9)$$

The NO_x-OH concentration parameterization in both the original scheme and the localized scheme at the site of IAP in Beijing is illustrated in Fig. 1. Before

localization, the OH is depleted more when the NO_x is high. The NO_x–OH curve without localization fits to the atmospheric condition in Europe and America. In the updated parameterization, the OH concentration decreases as the NO_x concentration increases when the NO_x concentration is higher than ~5ppb. The OH concentration falls to $\sim 2 \times 10^6 \text{ cm}^{-3}$ when NO_x concentration is higher than 30 ppb. The maximum of annual averaged OH concentration is still $\sim 7 \times 10^6 \text{ cm}^{-3}$, but the value of OH concentration corresponding to an NO_x concentration of 15–30 ppb is revised from $(0.5\text{--}1.5) \times 10^6 \text{ cm}^{-3}$ to $2\text{--}4 \times 10^6 \text{ cm}^{-3}$. Obviously, there are uncertainties ~~to~~ in adjusting the variation curve of OH concentration with respect to the NO_x concentration in the plume based on the surface observations. The variation curve can be further updated when observations in the plume are available.




Fig. 1 the localized (blue line) and original (yellow line) parameterization of the NO_x–OH concentration curve in the P6 scheme. The input parameters used to calculate the curve were annually averaged.

2.4 Simulation design

In this study, a nested domain over China with $0.33^\circ \times 0.33^\circ$ resolution was implemented to analyze the impact of SGPF on aerosol pollution in China, with the first domain covering the globe at $1^\circ \times 1^\circ$ resolution. Vertically, the model uses 20 layers, from the bottom layer centered at 50m to the top layer at ~20km, with 10

layers below 3 km. Meteorology was ~~driven-produced~~ by the global version of Weather Research and Forecasting (GWRf) version 3.7 (Zhang et al., 2012). The WRF was driven by the National Centers for Environmental Prediction Final Analysis (FNL) datasets with the calculation nudged to FNL data. The input frequencies are 3
 345 hourly in the global domain and 1 hourly in the nested domain. The top boundary conditions for ozone, NO_x and carbon ~~monoxide~~ ~~monoixde~~ were prescribed by the Model for Ozone and Related Chemical Tracers version4 (Emmons et al., 2010). For model performance evaluation and analysis of aerosol components, several cases
 350 corresponding to observation periods in 2014 (shown in Section 2.5) were simulated with one month ~~asfor~~ spin-up time. One simulation with the SGPF scheme (SG) and a control experiment with $f_{ox}=0\%$ (F0) were conducted in the simulating period to evaluate the sub-grid particles' impact on aerosol mass concentration simulation. The fraction of $f_{ox}=0\%$ represents the simulation without sub-grid particles. In addition, we
 355 conducted two simulations that both considered sub-grid particles to explore the impact of SGPF scheme on the model performance in PNSD at a typical urban site for winter 2016 (described in Section 2.5), ~~but in which~~ one is with the SGPF scheme and the other is not. The simulation without the SGPF scheme employed the $f_{ox}=2.5\%$ (F2.5) which refers to the Aerosol Comparisons between Observations and Models
 360 ~~AeroCom~~-recommendation by Dentener et al. (2006). The simulation~~ong~~ of F2.5 was also implemented in the comparison of the diurnal characteristics of SGPF in January and July 2014 in Section 3.2.

Table 1. Experiments configuration.

Experiment	Description
SG	Simulation with localized sub-grid particle formation scheme
F0	Simulation with 0% sub-grid particles
F2.5	Simulation with 2.5% sub-grid particles

A global emission dataset of source categories (with 29 species and 14 sectors)
 365 was applied with anthropogenic emissions from Hemispheric Transport of Air Pollution version 2. Detailed information for the emissions is available from Wei et al. (2019). The SO₂ ~~oxidized~~ ~~oxidated~~ to SG-ASO₄ was from ~~en~~energy and industry sectors (shown in Fig. 2) , and was emitted into the first five and three layers of the

model, respectively. The emissions in China were scaled to the level of 2014 based on
 370 emission trends reported by Zheng et al. (2018). The emissions of China for the
 simulation of 2016 were updated to the Multi-resolution Emission Inventory for
 China-2016 published by Tsinghua University (Zhang et al., 2019).




Fig. 2 The nested domain with annual mean SO₂ emission flux (unit : $\mu\text{g S m}^{-2}\text{s}^{-1}$)
 375 from energy and industry sectors in 2014. The black circles are locations of the
 observation sites.

2.5 Observation data

The observations of aerosol components and PNSD were obtained from the
 Atmospheric Pollution and Human Health in a Chinese Megacity (APHH-Beijing)
 380 campaign conducted at an urban site (the meteorological tower of IAP) in central
 Beijing during November–December 2016. Particle size ranges of 3–25, 25–100, and
 100–1000 nm were applied for the nucleation mode, Aitken mode and accumulation
 mode, respectively. In addition, we collected the mass concentrations of
 sulfate-nitrate-ammonium (SNA) at urban sites in the central of Nanjing and Wuhan
 385 (see in Fig. 2) to evaluate model performance in simulating aerosol components. The
 Nanjing and Wuhan sites were respectively located on the east and west banks of
 different reaches of the Yangtze River, with several power plants being located to their
 northeast. Locations and observation periods are given in Table 2. Observations to test
 the meteorological fields were collected from the National Climate Data Center at

sites given in Table 2. In this study, the observation periods were classed as warm (summer and autumn) and cold (winter) seasons to investigate the SGPF under different meteorological conditions. Noting that October-November in Beijing and May in Nanjing were categorized as cold and warm seasons, respectively.

Table 2. Information of observations for aerosol components and particle number concentrations.

Site name	Longitude (°)	Latitude (°)	Observation period	Observed species
Aerosol observation				
Beijing	116.37	39.97	2014. 6.3-2014.7.8	Mass concentration of SNA
			2014. 10.15-2014.11.6	
			2016.11.21-2016.12.13	PNSD and mass concentration of SNA, BC and organic matter (OM)
Nanjing	118.75	32.06	2014.5.1-2014.5.31	Mass concentration of SNA
			2014.1.1-2014.1.31	
Wuhan	114.28	30.62	2014. 10.1-2014.10.21	Mass concentration of SNA
Meteorological observation				
Beijing	116.47	39.80	June and October 2014	Temperature at 2m, relative humidity at 2m and wind at 10m
Nanjing	118.90	31.93	January and May 2014	
Wuhan	114.05	30.60	October 2014	

3 Results

3.1 Evaluation of the simulated meteorological fields

The SGPF is closely related to meteorological conditions, especially solar radiation and wind. Stronger radiation leads to more rapid gas-phase reactions, which means more SGPF. Wind speed and direction control plume diffusion and particle transportation in the grids. Furthermore, the concentrations of gases and aerosols in the air are related to temperature, RH, boundary layer height and other meteorological factors. We therefore compared the simulation of several meteorological factors with ground observations in corresponding cities during the same period as shown in Fig. 3. The correlation coefficient (R) of meteorological factors are given in Table 3. The simulation of temperature and RH agreed well with observations, with correlation coefficients (R) of >0.8. Regarding wind field, the model displays more uncertainties, with R values of 0.30–0.73 for wind speed and

<0.30 for wind direction in Beijing and Wuhan. The worse performance in Beijing warm season than cool season should be related to the poor simulation of wind field in warm season. Particularly, there is an obvious discrepancy between modeled and observed variations in Wuhan, with a relatively lower R value of 0.34. This may lead to a poor performance in modeling of aerosol distribution.




Fig. 3 Comparison between observed and simulated meteorological factors (i.e., temperature, relative humidity and wind speed) in: (a1) Beijing; (a2) Nanjing; (a3)

Wuhan in warm seasons and (b1) Beijing; (b2) Nanjing in cold seasons of 2014. The red dots and blue lines represent hourly observations and simulations, respectively. Monthly mean values of the observation and simulation are also given on subplots in blue texts, abbreviated to MO and MM, respectively.

420

Table 3 Summary of correlation coefficients for hourly meteorological factors in different cities. RH, U and V represent relative humidity, U wind and V wind, respectively.

Meteorological factors	Warm seasons			Cold seasons	
	Beijing	Nanjing	Wuhan	Beijing	Nanjing
Tempreature Temperature	0.84	0.94	0.85	0.88	0.88
RH	0.78	0.82	0.77	0.76	0.82
U	0.06	0.65	-0.18	0.60	0.48
V	0.19	0.48	0.22	0.52	0.30
Wind speed	0.30	0.73	0.34	0.62	0.58

3.2 Diurnal variation of sub-grid particle formation

The diurnal variation of oxidation rate and mass concentration of SG-ASO₄ ~~is-are~~ shown in Fig. 4. ~~with In summer,~~ a clear diurnal pattern for f_{ox} at different latitudes ~~is displayed,(Fig. 4a, e);~~ with initial upward then downward trends from 08:00 to 20:00 (Beijing Time) ~~in summer,~~ ~~with~~ The f_{ox} varies ~~respectively~~ from ~0% to over 10% and from 0% to 14% in Beijing (Fig. 4a) and Wuhan (Fig. 4c), respectively, due to the variation of solar radiation. At night, f_{ox} remained at almost 0% due to the much lower OH concentration. In winter, f_{ox} increased to ~5% during daytime and fell to ~0% at night in Wuhan, with the time for f_{ox} being >1% narrowing to 10:00–18:00. For the same period, the f_{ox} in Wuhan is about 1.5 times that of Beijing, ~~and~~ ~~W~~within a city, the maximum f_{ox} in summer is ~3 times that in winter. Particularly, ~~The-the~~ maximum f_{ox} in summer is about 6 times and 4 times that of the simulation of F2.5 for Wuhan and Beijing, respectively. Overall, there is an evident discrepancy for f_{ox} in the spatial-temporal distribution. The rate of sulfate formation at noon is significantly higher than at other time, leading to more sulfur deposition (Xu et al., 2014).

The diurnal variation in the mass concentration of SG-ASO₄ associated with different f_{ox} and emission flux is shown in Fig. 4b ~~and~~ d. The modeling results of SG-ASO₄ in Beijing ranged between 0–2 and 0–1 $\mu\text{g m}^{-3}$ in summer and winter, respectively. In Wuhan, the maximum value exceeded 10 $\mu\text{g m}^{-3}$ and ~4 $\mu\text{g m}^{-3}$ in summer and winter, respectively. This indicates that SFPF is substantially important

in areas with large point sources. On the other hand, simulation with fixed f_{ox} (2.5%) maintained a constant SG-ASO₄ concentration of about 0.5–0.9 $\mu\text{g m}^{-3}$ and 2 $\mu\text{g m}^{-3}$ in Beijing and Wuhan, respectively. It is worth noting that, in Beijing, the model results of SG-ASO₄ with fixed f_{ox} is slightly higher in winter than summer, due to the higher emission flux of SO₂ in heating season. However, the simulation with the SGPF scheme suggests the opposite due to higher SG-ASO₄ conversion in summer, demonstrating the superiority of the scheme in resolving temporal heterogeneity of SGPF.




Fig. 4 Monthly averaged diurnal variation of the oxidation rate (f_{ox} , a, c) and mass concentration ($\mu\text{g m}^{-3}$) of sub-grid sulfate (b, d) in January (blue line) and July (red line) from the IAP-AACM. The orange line represents constant f_{ox} of 2.5%. The solid line and dash-dotted line represent simulation with the sub-grid scheme (SG) and constant f_{ox} of 2.5% (F2.5), respectively. The top row and bottom row are for Beijing and Wuhan, respectively.

3.3 Improvement in aerosol components simulation

–The testing of the simulated time-series of SNA mass concentration against site observations in warm seasons is illustrated in Fig. 5, which displays the IAP-AACM simulation with and without the SGPF scheme. The statistics for different model

results are given in Table 4. The model reproduced SNA concentration and their temporal evolution well, with R values generally over 0.5. Overall, the model ~~somewhat under-predicted secondary inorganic aerosols~~ the concentration of SNA in different degrees. The simulation in Beijing matched observations most closely, likely due to the more accurate emission inventory available for that area.

The SG simulation ~~indicates can increase~~ sulfate increases of $1\text{--}5\text{ }\mu\text{g m}^{-3}$ in warm seasons (Fig. 5). Compared with the F0 simulation, the SG simulation leads to an obvious improving on model bias (shown in Table 4), on the background of underestimation of ~~secondary inorganic aerosols~~ SNA in the model. The normalized model bias (NMB) in Nanjing and Wuhan were narrowed from -41% to -18% and from -60% to -33% respectively. The overestimation of the Beijing NMB increased by 20% (it was overestimated by 6% in F0). The simulation of SG significantly improved the correlation of sulfate in Nanjing with the R increasing by 0.13. The SG correlation is similar to that of F0 in Beijing, but lower in Wuhan where the R value decreased by 0.15, with this being related to some extent to the poor wind simulation in these areas (as shown in Table 3). The SGPF scheme also has an obvious impact on the simulation of ammonium, with the NMB ~~narrowing-reduced~~ by 14% and R ~~increas~~ed by 0.08 in Nanjing and the NMB ~~decreas~~ed by 12% in Wuhan. The correlation of nitrate in SG is similar to that in the simulation of F0, with the model bias increasing by 1–9%. Overall, the IAP-AACM exhibits good performance ~~with~~ in SNA simulation, and the SGPF scheme improves the simulation on sulfate and ammonium.

The SGPF scheme significantly narrowed the gap between model and observation in Wuhan, with the NMB decreasing from -0.6 to -0.33 due to the high emission rate of the point source. We found that the sulfate concentration simulated by the IAP-AACM with the SGPF scheme increased significantly during daytime, especially on October 1 and 15. The concentration of sulfate was up to $20\text{ }\mu\text{g m}^{-3}$ higher on those days than the model results without SG- ASO_4 . This could be related to the influence of the local wind direction on the plume spreading and accumulation of pollutants. We compared the average diurnal variation of simulated SNA

concentrations in Wuhan with observations in Fig. 6. The simulated diurnal profiles of both SG and F0 reproduced the variation of sulfate proportions in SNA (ASO₄/SNA) as a unimodal. Simulation with the SGPF scheme provides diurnal SNA mass concentration profiles more similar to observations than that of F0. The averaged SNA concentration was increased by ~5 $\mu\text{g m}^{-3}$ at night and 10 $\mu\text{g m}^{-3}$ in the afternoon. The simulated ASO₄/SNA ratio of SG increased by 5%–10% or more at daytime and about 5% at night than F0. The simulated ASO₄/SNA ratio ~~dropped~~ dropped—much more dramatically than observed one at noon due to over-decomposition of nitric acid under high-temperature conditions.

The simulation of SNA in cold seasons is shown in Fig. 7. In general, the mass concentration of sulfate was obviously underestimated, with the SG simulation showing weaker improvement in the negative bias than in warm seasons. As shown in Table 4, the NMB of sulfate decreased by 0.16 and 0.05 in Beijing and Nanjing, respectively. The correlation increased slightly (by 0.02–0.04) due to the favorable performance of wind simulation (Table 3). The SGPF scheme slightly improved the simulation of ammonium but had ~~ambiguous-neglectable~~ ambiguous-neglectable effect on the simulation of nitrate. The NMB of ammonium was reduced by 0.06 in Beijing but not changed in Nanjing as the SGPF is weakest in January. For nitrate, the simulation of SG gave little changing in correlation but increased the NMB by 0.05 in Nanjing. The evaluation of SNA simulations in SG and F0 indicates that the SGPF scheme improves model performance for sulfate and ammonium, especially in warm seasons. In particular, the consideration of SG-ASO₄ has a ~~great-considerable~~ great-considerable influence on the spatial-temporal distribution of SNA near sulfur-rich stacks on condition that the simulation of wind field is reliable.

As displayed in Fig. 8, sulfate accounted for over 40% of SNA concentration in warm seasons and about 20%–30% in cold seasons in 2014. In Nanjing and Wuhan in warm seasons, ~~simulated sulfate from the modeling results~~ simulated sulfate without SGPF made up only ~30% of SNA. The simulation of SG raised the simulated proportion of sulfate by ~10% in warm seasons and <5% in cold seasons. We concluded that the model coupled with the SGPF scheme delivers ~~superior-performance~~ superior-performance remarkable

advantages in determining the concentration of sulfate and its proportion in SNA.




Fig. 5 Comparison of hourly simulated aerosol components against site observations in Beijing, Nanjing and Wuhan in warm seasons: (a1–a3) sulfate, (b1–b3) nitrate, (c1–c3) ammonium. The blue and orange line represent simulations with SG scheme and with constant f_{ox} of 0% (F0), respectively. The red dots represent observations.

Table 4 Summary of statistics for hourly mass concentration of aerosol components in different seasons. MO, MM, NMB, and R represent mean value of the observations, mean value of the model, normalized mean bias, and correlation coefficients respectively. ASO4, ANO3 and ANH4 represent sulfate, nitrate and ammonium, respectively.

Period	Site name	Species	MO (μgm^{-3})	SG			F0		
				MM (μgm^{-3})	NMB	R	MM (μgm^{-3})	NMB	R
Warm seasons	Beijing	ASO4	10.30	12.99	0.26	0.59	10.95	0.06	0.60
		ANO3	6.84	5.94	-0.13	0.46	6.03	-0.12	0.47
		ANH4	5.41	5.72	0.06	0.56	5.01	-0.07	0.55
	Nanjing	ASO4	10.72	8.84	-0.18	0.61	6.33	-0.41	0.48
		ANO3	8.58	7.00	-0.18	0.48	7.47	-0.13	0.50
		ANH4	5.36	3.43	-0.36	0.60	2.69	-0.50	0.52
	Wuhan	ASO4	21.53	14.34	-0.33	0.31	8.68	-0.60	0.46

Cold seasons	Beijing	ANO3	14.52	13.05	-0.10	0.53	14.37	-0.01	0.54
		ANH4	11.60	8.84	-0.24	0.48	7.47	-0.36	0.50
		ASO4	10.53	9.43	-0.10	0.57	7.79	-0.26	0.53
		ANO3	20.60	18.18	-0.12	0.52	17.89	-0.13	0.54
	Nanjing	ANH4	8.77	7.49	-0.15	0.58	6.91	-0.21	0.58
		ASO4	23.65	9.09	-0.62	0.65	7.91	-0.67	0.63
		ANO3	29.41	18.02	-0.39	0.62	19.28	-0.34	0.62
		ANH4	19.82	7.03	-0.65	0.66	6.91	-0.65	0.63




Fig. 6 Averaged diurnal variation for the observed and simulated total SNA concentrations (solid lines), observed ASO4/SNA ratio (black dot line) and the percentage of sulfate, nitrate and ammonium in total SNA concentrations (color bars, i.e., observations, simulations of SG and F0) in Wuhan, October 1-21, 2014.




Fig. 7 The same as Fig. 5, but for Beijing and Nanjing in cold seasons.




Fig. 8 Mean percentage of the simulated (SG and F0) and observed sulfate, nitrate and ammonium (i.e., red, blue and orange color bars) and averaged SNA concentration (black dots) in Beijing, Nanjing and Wuhan in (a) warm seasons and (b) cold seasons.

3.4 Improvement in PNSD simulation

Besides the mass concentration of component, particle number concentration is an essential parameter of aerosol particles in evaluating their climatic and environmental effects. Most aerosol models use constant fractions for f_{ox} and f_{new} to describe the SGPF, causing large uncertainties in simulating particle formation processes and particle number concentration. To investigate the influence of SGPF scheme on particle number concentration, we ~~undertook~~performed two experiments using IAP-AACM+APM with a fixed scheme ($f_{ox}=2.5\%$ and $f_{new} = 5\%$) and the SGPF scheme. Modeling results were ~~tested~~evaluated against observations at the site in Beijing during the APHH-Beijing, winter 2016. ~~The evaluation of SNA against~~

~~observations is shown in Fig. S3.~~ The model reproduced the mass concentration of SNA well (Fig. S3), with R ranging from 0.59 to 0.66 and NMB ranging from -0.68 to 0.01. The PNSDs of the two experiments ~~and the observation are~~ shown in Fig. 9a and Fig. 9b. ~~The observed PNSD during the same period is shown in Fig. 9e.~~ The model with constant SG-ASO₄ formation over-predicted the number of particles in Aitken mode at night (Fig. 9a). In fact, nucleation is negligible at night due to very low OH concentration in the plume. Compared with observations, the sub-grid scheme significantly optimized the overestimation of particles number concentration in small size bins (Fig. 9b). Specifically, considering the diurnal variation of nucleation process reduced the positive bias by more than 3 times in nucleation mode and by about a half in Aitken mode (see in Table 5). This indicates that IAP-AACM coupled with the SGPF scheme reasonably captured the process of new particle formation in clean periods (e.g. November 27–28, December 5–6) and growth in haze periods (e.g. November 28–30). Here we also display the simulations of full spectrum of PNSD in F2.5 and SG in Fig. S4. Particles in the simulation with F2.5 (Fig. S4a) were produced continuously within the particle size range of 3–10nm, due to constant nucleation of SG-ASO₄. To avoid this unrealistic particle formation, the fraction of SG-ASO₄ is always set as 0% in aerosol models. However, it will result in a considerable ~~ey~~ underestimation of particle number concentration in in nucleation mode and Aitken mode. Taking into account the influence of solar radiation on SGPF, the model with the SGPF scheme reproduces the diurnal cycle of particle number in small-sized bins and provides more reasonable estimation of the particle number concentration originated from new particle formation~~numbers concentration of CCN.~~




Fig. 9 Particle number size distribution of simulations from IAP-AACM with (a) F2.5 and $f_{new} = 5\%$, (b) SG and (c) observations during the APHH-Beijing in 2016.

Table 5 Mean number concentrations of the observations and simulations and the NMB for different modes at the site in Beijing during the APHH-Beijing in 2016.

Experiments	Nucleation		Aitken		Accumulation	
	Con(cm^{-3})	NMB(-)	Con(cm^{-3})	NMB (-)	Con(cm^{-3})	NMB (-)
Observation	1312		10223		3754	
SG	2208	0.68	15183	0.49	7022	0.87
F2.5	4412	2.36	18507	0.81	7160	0.91

3.5– Regional impacts of sub-grid particle formation

~~The significant impact of SGPF on sulfate concentration near site is shown in Fig. 5.~~ The distribution of sub-grid particles in ~~local source~~ areas is strongly influenced strongly by ~~the wind-driven field due to~~ dilution ~~of high and transport of~~ plumes. To explore the spatial inhomogeneity in regions around point sources, the regional impact of SGPF during typical periods is illustrated in Fig. 10. In Nanjing and Wuhan, the area represented by the observation site is located downwind of the point source when the easterly wind prevails. For Nanjing, there is ~~then~~ an increase in sulfate mass concentration by 25%–50% around the observation site. ~~in~~ For Wuhan, the increase can be reaching-reaching over 50% in the downwind area. This should be ascribed to the high capacity of local power plants with large SO₂ emission of over 2 μg m⁻² s⁻¹ in suburban Wuhan, indicating the significant impact of point source ~~with high emission rate~~ on both the spatial scale and material concentration of the aerosol distribution. On the other hand, the regional influence in Beijing was relatively small (Fig. 10a, d), due to the upwind location and much lower emission rate of plants. Comparison of the three scenarios indicates that, although sub-grid particles from elevated sources are discharged ~~to~~ at high altitudes, they can still contribute significantly to ~~local~~ ground layer pollution under unfavorable wind condition.

The influence of the SGPF diurnal profile (Section 3.2) on regional scales was investigated at two particular times (i.e., 14:00 and 02:00 Beijing time). To exclude the influence of boundary layer height on aerosol concentration, we used sulfate/BC ratios to normalize the comparison between day and night. As shown in Fig. 11, ~~in~~ over industrial areas in eastern China there is a larger increase in sulfate concentration due to SGPF in daytime than at night. The increase in large areas of Shanxi Province indicates the important role of coal burning as the main energy source. ~~In~~ Over central China, there are several maxima which indicates the local influence of SGPF. ~~Note that the highest increase occurred in northwest China because the concentration of BC is almost zero in such clean areas.~~ Overall, the temporal variation of SGPF affect the sulfate concentration on a large-scale in eastern China, especially in areas with high-emission sources, leading to heterogeneous spatial distribution.




Fig. 10 Daily averaged differences with ~~overlaid~~ overlaid wind field (left column, $\mu\text{g m}^{-3}$) and relative change (right column, %) of sulfate concentrations on regional scale between SG and F0 at surface, around: (a, d) Beijing in June 22, (b, e) Nanjing in May 16 and (c, f) Wuhan in October 15. The difference was calculated as the simulation of SG minus F0. Black dots and blue triangles represent locations of the ground sites and plants, respectively.




Fig. 11 Difference of sulfate/BC at July 1st (a) 14:00 and (b) 02:00 Beijing time between SG and F0 at surface. The difference is calculated as the simulation of SG minus F0.

4. Conclusions and discussions

In this study, the P6 sub-grid parameterization scheme was coupled ~~with~~into the global nested aerosol model IAP-AACM to resolve the SGPF characteristics in both oxidation and nucleation processes over eastern China. Furthermore, the key parameter of the scheme, effective OH concentration in the plume, was modified to fit the local chemical background based on ~~the basis of~~ extensive field observations in eastern China, ~~on account of~~according to ~~the its~~ dependence ~~of radicals~~ on precursor gases. With the updated model integrated with the localized new scheme, the spatio-temporal inhomogeneity of SGPF was well parameterized in the grids. The significance of sub-grid particles in simulating the distribution of secondary inorganic aerosols was demonstrated in comparisons against site observations for both aerosol mass concentration and PNSD in different seasons.

The major findings were as follows. (1) The SGPF scheme provides a more reasonable description of oxidation and nucleation processes for SGPF, including the diurnal variation of f_{ox} and f_{new} . The spatio-temporal inhomogeneity of sub-grid particles distribution is well characterized. The spatial and temporal discrepancy is obvious, as for the same area, the f_{ox} in summer can be three times that of winter. (2) simulations with the SGPF scheme better reproduce nanoparticle formation than the

fixed oxidation fraction as the physicochemical parameterization nature and diurnal cycle of the nucleation in plumes are realistically represented~~profiles more accurately describe the nucleation source at night~~. The overestimation of particle number in small size was reduced by over 3 times in nucleation mode and roughly one half in Aitken mode. (3) The SGPF gave more significant improvement ~~in model performance~~ in warm seasons as it reduced the NMB of sulfate and ammonium by 12%–27% at most sites, while increasing that of nitrate by 1%–9%. In cold seasons, the influence of SGPF was limited, with the NMB for sulfate decreasing by only 5%–16%. ~~In cold seasons, the influence of SGPF is limited with the NMB for sulfate only reduced by 5%–16%.~~ The simulating improvement on correlation will ~~experience get~~ a significant increase by 0.13 if the ~~model performance of~~ wind fields are nicewell reproduced. (4) The SGPF scheme can improve model performance in determining the concentration of sulfate and its proportion in SNA, increasing the proportion of sulfate by ~10% in warm seasons and <5% in cold seasons. Specifically, in Wuhan, the averaged SNA concentration increased by ~5 $\mu\text{g m}^{-3}$ at night and 10 $\mu\text{g m}^{-3}$ in the afternoon, ~~which has implicatingons thefor obvious impact of SGPF on~~ sulfate simulating ~~overin~~ areas ~~near-with~~ stacks. (5) The SGPF has a significant impact ~~in localover~~ areas near point sources, with the sulfate concentration increasing by 25%–50% or even >50% under downwind conditions. This indicates that the impact of SGPF should be taken into account in studies of air pollution and aerosol formation in extensive industrial cities, not only in China, but also in other developing countries facing PM_{2.5} and ozone pollution like India (Chen et al., 2020).

As the IAP-AACM was driven by the global WRF off-line, the aerosol feedback was not taken into account in this work. On the other hand, ~~the~~ SGPF scheme didn't ~~contain-incorporate~~ the variation in OD associated with droplets in the plume. Without the aerosol-radiation interaction (ARI), the impact on increasing atmospheric stability by cooling the surface but heating the air aloft was excluded. ~~The impact of aerosols serving as cloud condensation nuclei on optical properties and lifetime of clouds and precipitation was also not included.~~ Moreover, the scattering and absorption of ultraviolet radiation by aerosols influence the photolysis rates and reduce the

675 formation of O₃ and other oxidants (He and Carmichael, 1999). In high polluted area,
ignoring the ~~aerosol-photolysis interaction (API)impact of OD~~ will cause a certain
degree of overestimation of atmospheric oxidation capacity. Li et al. (2018)
incorporated recently reported heterogeneous chemical mechanisms into the regional
version of IAP-AACM (~~NAQPMS~~) and found that perturbations in photolysis
680 frequencies reduced O₃ concentrations by 1–5 ppb in winter and 1–3 ppb in summer.
Considering both the ARI and ~~the aerosol-photolysis interaction (API)~~, Wu et al.
(2020) conducted model experiments in North China Plain and found that ARI
contributed to a 7.8% increase in near-surface PM_{2.5}, while API suppressed secondary
aerosol formation to a 3% decrease of PM_{2.5}. Therefore, the overestimation of
685 atmospheric oxidation capacity caused by ignoring the OD impact should be under 5%
in eastern China. To better understand the impact and uncertainty of sub-grid particle,
the impact of OD should be included in the model in our future work.

Code and data availability

All of the observation data needed to evaluate the conclusion of this paper are
690 provided in the main text. The source codes of IAP-AACM with SGPF scheme are
available online via ZENODO (<https://doi.org/10.5281/zenodo.4383361>; Wei et al.,
2020). Please contact Ying Wei (weiyang@mail.iap.ac.cn) to obtain the model data for
IAP-AACM.

Author contribution

695 YW developed the model, did the simulation, and wrote the paper. XC developed the
model and designed the study. HC developed the model and prepared the gridded
emission data. YS provided the observation data. WY, HD, QW, DC, XZ, JL and ZW
modified the paper. YW and XC prepared the manuscript with contributions from all the
co-authors.

Acknowledgments:

700 We sincerely thank Prof. Fangqun Yu at State University of New York at Albany
for providing the code of original APM box model. We also thank the Jiangsu
Environmental Monitoring Center and Hubei Environmental Monitoring Center for
their supports with aerosol composition data of Nanjing and Wuhan respectively. This

705 research is supported by the National Key R&D Program of China (Grant NO. 2020YFA0607801), the National Natural Science Foundation of China (Grant NO. 42007199; 41907200) and the National Key Scientific and Technological Infrastructure project “Earth System Science Numerical Simulator Facility”.

References

- Adams, P. J., Seinfeld, J. H., and Koch, D. M.: Global concentrations of tropospheric sulfate, nitrate, and ammonium aerosol simulated in a general circulation model, *Journal of Geophysical Research-Atmospheres*, 104, 13791-13823, 1999.
- Boucher, O., D. Randall, P. Artaxo, C. Bretherton, et al.: Clouds and Aerosols. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
- Brock, C. A., Washenfelder, R. A., Trainer, M., Ryerson, T. B., Wilson, J. C., Reeves, J. M., Huey, L. G., Holloway, J. S., Parrish, D. D., Hubler, G., and Fehsenfeld, F. C.: Particle growth in the plumes of coal-fired power plants, *Journal of Geophysical Research-Atmospheres*, 107, 2002.
- Chen, H. S., Wang, Z. F., Li, J., Tang, X., Ge, B. Z., Wu, X. L., Wild, O., and Carmichael, G. R.: GNAQPMS-Hg v1.0, a global nested atmospheric mercury transport model: model description, evaluation and application to trans-boundary transport of Chinese anthropogenic emissions, *Geoscientific Model Development*, 8, 2857-2876, 2015.
- Chen, X. S., Wang, Z. F., Li, J., et al.: Simulation of particle number size distribution in Beijing in winter using NAQPMS+APM model, *Climatic and Environmental Research (in Chinese)*, 20 (6): 611–619, doi:10.3878/j.issn.1006-9585.2015.15095, 2015.
- Chen, X., Wang, Z., Li, J., Yang, W., Chen, H., Wang, Z., Hao, J., Ge, B., Wang, D., and Huang, H.: Simulation on different response characteristics of aerosol particle number concentration and mass concentration to emission changes over mainland China, *Science of the Total Environment*, 643, 692-703, 2018.
- Chen, X., Wang, Z., Li, J., and Yu, F.: Development of a Regional Chemical Transport Model with Size-Resolved Aerosol Microphysics and Its Application on Aerosol Number Concentration Simulation over China, *Sola*, 10, 83-87, 2014.
- Chen, X., Yu, F., Yang, W., Sun, Y., Chen, H., Du, W., Zhao, J., Wei, Y., Wei, L., Du, H., Wang, Z., Wu, Q., Li, J., An, J., and Wang, Z.: Global-regional nested simulation of particle number concentration by combing microphysical processes with an evolving organic aerosol module, *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2020-759>, in review, 2020.
- Chen, Y., Wild, O., Ryan, E., Sahu, S.K., Beig, G.J.A.C., Physics, 2020. Mitigation of PM2.5 and ozone pollution in Delhi: a sensitivity study during the pre-monsoon period. 20, 499-514.
- Dentener, F., Kinne, S., Bond, T., Boucher, O., Cofala, J., Generoso, S., Ginoux, P., Gong, S., Hoelzemann, J. J., Ito, A., Marelli, L., Penner, J. E., Putaud, J. P., Textor, C., Schulz, M., van der Werf, G. R., and Wilson, J.: Emissions of primary aerosol and precursor gases in the years 2000 and 1750 prescribed data-sets for AeroCom, *Atmospheric Chemistry and Physics*, 6, 4321-4344, 2006.
- Dusek, U., Frank, G. P., Hildebrandt, L., Curtius, J., Schneider, J., Walter, S., Chand, D., Drewnick, F., Hings, S., Jung, D., Borrmann, S., and Andreae, M. O.: Size matters more than chemistry for cloud-nucleating ability of aerosol particles, *Science*, 312, 1375-1378, 2006.
- Emmons, L. K., Walters, S., Hess, P. G., J.-F. L., Pfister, G. G., Fillmore, D., Granier, C., Guenther, A., Kinnison, D., and Laepple, T. J. G. M. D.: Description and evaluation of the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4), 3, 43-67, 2010.

- Ge, B. Z., Wang, Z. F., Xu, X. B., Wu, J. B., Yu, X. L., and Li, J.: Wet deposition of acidifying substances in different regions of China and the rest of East Asia: Modeling with updated NAQPMS, *Environmental Pollution*, 187, 10-21, 2014.
- 755 Gillani N V, Godowitch J M.: Plume-in-Grid Treatment of Major Point Source Emissions. In Science Algorithms of the EPA Models-3 Community Multiscale Air Quality (CMAQ) Modeling System. U.S. Environmental Protection Agency: Washington, DC, USA, 1999; Chapter 9.76.
- 760 Gillani, N. V., Meagher, J. F., Valente, R. J., Imhoff, R. E., Tanner, R. L., and Luria, M.: Relative production of ozone and nitrates in urban and rural power plant plumes 1. Composite results based on data from 10 field measurement days, *Journal of Geophysical Research-Atmospheres*, 103, 22593-22615, 1998.
- 765 He, Q.-F., Ding, X., Fu, X.-X., Zhang, Y.-Q., Wang, J.-Q., Liu, Y.-X., Tang, M.-J., Wang, X.-M., and Rudich, Y.: Secondary Organic Aerosol Formation From Isoprene Epoxides in the Pearl River Delta, South China: IEPOX- and HMML-Derived Tracers, *Journal of Geophysical Research-Atmospheres*, 123, 6999-7012, 2018.
- He, S., Carmichael, G.R.: Sensitivity of photolysis rates and ozone production in the troposphere to aerosol properties. *J. Geophys. Res.* 104 (26) (307,23324), doi:10.1029/2004JD005359, 1999.
- 770 Hu, M., Shang, D., Guo, S., and Wu, Z.: Mechanism of New Particle Formation and Growth as well as Environmental Effects under Complex Air Pollution in China, *Acta Chimica Sinica*, 74, 385-391, 2016.
- IPCC, 2013: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1535 pp., doi:10.1017/CBO9781107415324.
- 775 Karamchandani, P., Seigneur, C., Vijayaraghavan, K., and Wu, S. Y.: Development and application of a state-of-the-science plume-in-grid model, *Journal of Geophysical Research-Atmospheres*, 107, 2002.
- 780 Karamchandani, P., Vijayaraghavan, K., Zhang, Y., Chen, S-Y.: Development of a Plume-in-Grid Version of Global-through-Urban WRF/Chem. In Proceedings of the 9th Annual CMAS Conference. Chapel Hill, NC, USA, October 2010.
- Khairoutdinov, M. F. and Randall, D. A.: Cloud resolving modeling of the ARM summer 1997 IOP: Model formulation, results, uncertainties, and sensitivities, *Journal of the Atmospheric Sciences*, 60, 607-625, 2003.
- 785 Kulmala, M. and Kerminen, V.-M.: On the formation and growth of atmospheric nanoparticles, *Atmospheric Research*, 90, 132-150, 2008.
- Kulmala, M., Kontkanen, J., Junninen, H., Lehtipalo, K., Manninen, H. E., Nieminen, T., Petaja, T., Sipila, M., Schobesberger, S., Rantala, P., Franchin, A., Jokinen, T., Jarvinen, E., Aijala, M., Kangasluoma, J., Hakala, J., Aalto, P. P., Paasonen, P., Mikkila, J., Vanhanen, J., Aalto, J., Hakola, H., Makkonen, U., Ruuskanen, T., Mauldin, R. L., III, Duplissy, J., Vehkamaki, H., Back, J., Kortelainen, A., Riipinen, I., Kurten, T., Johnston, M. V., Smith, J. N., Ehn, M., Mentel, T. F., Lehtinen, K. E. J., Laaksonen, A., Kerminen, V.-M., and Worsnop, D. R.: Direct Observations of Atmospheric Aerosol Nucleation, *Science*, 339, 943-946, 2013.
- 795 Li, C., McLinden, C., Fioletov, V., Krotkov, N., Carn, S., Joiner, J., Streets, D., He, H., Ren, X.,

Li, Z., Dickerson, R.R., 2017. India Is Overtaking China as the World's Largest Emitter of Anthropogenic Sulfur Dioxide. *Scientific Reports* 7.

Li, J., Wang, Z., Akimoto, H., Gao, C., Pochanart, P., and Wang, X.: Modeling study of ozone seasonal cycle in lower troposphere over east Asia, *Journal of Geophysical Research-Atmospheres*, 112, 2007.

Li, J., Wang, Z., Zhuang, G., Luo, G., Sun, Y., Wang, Q. J. A. C., and Physics: Mixing of Asian mineral dust with anthropogenic pollutants over East Asia: a model case study of a super-duststorm in March 2010, 12, 7591-7607, 2012.

Li, J., Yang, W., Wang, Z., Chen, H., Hu, B., Li, J., Sun, Y., Fu, P., and Zhang, Y.: Modeling study of surface ozone source-receptor relationships in East Asia, *Atmospheric Research*, 167, 77-88, 2016.

Li, J., Chen, X., Wang, Z., Du, H., Yang, W., Sun, Y., Hu, B., Li, J., Wang, W., Wang, T., Fu, P., and Huang, H.: Radiative and heterogeneous chemical effects of aerosols on ozone and inorganic aerosols over East Asia, *Science of the Total Environment*, 622, 1327-1342, 2018.

Liu, J., Liu, Z., Ma, Z., Yang, S., Yao, D., Zhao, S., Hu, B., Tang, G., Sun, J., Cheng, M., Xu, Z., and Wang, Y.: Detailed budget analysis of HONO in Beijing, China: Implication on atmosphere oxidation capacity in polluted megacity, *Atmospheric Environment*, 244, 117957, 2021.

Lonsdale, C. R., Stevens, R. G., Brock, C. A., Makar, P. A., Knipping, E. M., and Pierce, J. R.: The effect of coal-fired power-plant SO₂ and NO_x control technologies on aerosol nucleation in the source plumes, *Atmospheric Chemistry and Physics*, 12, 11519-11531, 2012.

Lu, K. D., Rohrer, F., Holland, F., Fuchs, H., Bohn, B., Brauers, T., Chang, C. C., Haeseler, R., Hu, M., Kita, K., Kondo, Y., Li, X., Lou, S. R., Nehr, S., Shao, M., Zeng, L. M., Wahner, A., Zhang, Y. H., and Hofzumahaus, A.: Observation and modelling of OH and HO₂ concentrations in the Pearl River Delta 2006: a missing OH source in a VOC rich atmosphere, *Atmospheric Chemistry and Physics*, 12, 1541-1569, 2012.

Luo, G. and Wang, Z. F.: A Global Environmental Atmospheric Transport Model (GEATM) : Model Description and Validation, *Chinese Journal of Atmospheric Sciences*, 30, 504-518, 2006.

Luo, G. and Yu, F.: Sensitivity of global cloud condensation nuclei concentrations to primary sulfate emission parameterizations, *Atmospheric Chemistry and Physics*, 11, 1949-1959, 2011.

Ma, Q., Cai, S., Wang, S., Zhao, B., Martin, R. V., Brauer, M., Cohen, A., Jiang, J., Zhou, W., Hao, J., Frostad, J., Forouzanfar, M. H., and Burnett, R. T.: Impacts of coal burning on ambient PM_{2.5} pollution in China, *Atmospheric Chemistry and Physics*, 17, 4477-4491, 2017.

Nenes, A., Pandis, S. N., and Pilinis, C.: ISORROPIA: A new thermodynamic equilibrium model for multiphase multicomponent inorganic aerosols, *Aquatic Geochemistry*, 4, 123-152, 1998.

Nenes, A., Pandis, S. N., and Pilinis, C.: Continued development and testing of a new thermodynamic aerosol module for urban and regional air quality models, *Atmospheric Environment*, 33, 1553-1560, 1999.

Peng, J. F., Hu, M., Wang, Z. B., Huang, X. F., Kumar, P., Wu, Z. J., Guo, S., Yue, D. L., Shang, D. J., Zheng, Z., and He, L. Y.: Submicron aerosols at thirteen diversified sites in China: size distribution, new particle formation and corresponding contribution to cloud condensation

- nuclei production, *Atmospheric Chemistry and Physics*, 14, 10249-10265, 2014.
- Powell, H., Krall, J. R., Wang, Y., Bell, M. L., and Peng, R. D.: Ambient Coarse Particulate Matter and Hospital Admissions in the Medicare Cohort Air Pollution Study, 1999-2010, *Environmental Health Perspectives*, 123, 1152-1158, 2015.
- 845 Powell, H., Krall, J. R., Wang, Y., Bell, M. L., and Peng, R. D.: Ambient Coarse Particulate Matter and Hospital Admissions in the Medicare Cohort Air Pollution Study, 1999-2010, *Environmental Health Perspectives*, 123, 1152-1158, 2015.
- Riccobono, F., Schobesberger, S., Scott, C. E., Dommen, J., Ortega, I. K., Rondo, L., Almeida, J., Amorim, A., Bianchi, F., Breitenlechner, M., David, A., Downard, A., Dunne, E. M., Duplissy, J., Ehrhart, S., Flagan, R. C., Franchin, A., Hansel, A., Junninen, H., Kajos, M., 850 Keskinen, H., Kupc, A., Kuerten, A., Kvashin, A. N., Laaksonen, A., Lehtipalo, K., Makhmutov, V., Mathot, S., Nieminen, T., Onnela, A., Petaja, T., Praplan, A. P., Santos, F. D., Schallhart, S., Seinfeld, J. H., Sipila, M., Spracklen, D. V., Stozhkov, Y., Stratmann, F., Tome, A., Tsagkogeorgas, G., Vaattovaara, P., Viisanen, Y., Vrtala, A., Wagner, P. E., Weingartner, E., Wex, H., Wimmer, D., Carslaw, K. S., Curtius, J., Donahue, N. M., Kirkby, 855 J., Kulmala, M., Worsnop, D. R., and Baltensperger, U.: Oxidation Products of Biogenic Emissions Contribute to Nucleation of Atmospheric Particles, *Science*, 344, 717-721, 2014.
- Richards, L. W., Anderson, J. A., Blumenthal, D. L., Brandt, A. A., McDonald, J. A., Waters, N., Macias, E. S., and Bhardwaja, P. S.: THE CHEMISTRY, AEROSOL PHYSICS, AND 860 OPTICAL-PROPERTIES OF A WESTERN COAL-FIRED POWER-PLANT PLUME, *Atmospheric Environment*, 15, 2111-2134, 1981.
- Seinfeld, J.H., Pandis, S.N.: *Atmospheric Chemistry and Physics: from Air Pollution to Climate Change*, 2nd ed. John Wiley & Sons, Hoboken, New Jersey, 2012.
- Slater, E. J., Whalley, L. K., Robert, Woodward-Massey, Ye, C., Lee, J. D., Squires, F., et al.: 865 Elevated levels of OH observed in haze events during wintertime in central Beijing. *Atmos. Chem. Phys.*, 20, 14847–14871, <https://doi.org/10.5194/acp-20-14847-2020>, 2020.
- Spracklen, D. V., Carslaw, K. S., Kulmala, M., Kerminen, V. M., Mann, G. W., and Sihto, S. L.: The contribution of boundary layer nucleation events to total particle concentrations on regional and global scales, *Atmospheric Chemistry and Physics*, 6, 5631-5648, 2006.
- 870 Spracklen, D. V., Carslaw, K. S., Kulmala, M., Kerminen, V.-M., Sihto, S.-L., Riipinen, I., Merikanto, J., Mann, G. W., Chipperfield, M. P., Wiedensohler, A., Birmili, W., and Lihavainen, H.: Contribution of particle formation to global cloud condensation nuclei concentrations, *Geophysical Research Letters*, 35, 2008.
- Spracklen, D. V., Pringle, K. J., Carslaw, K. S., Chipperfield, M. P., and Mann, G. W.: A global 875 off-line model of size-resolved aerosol microphysics: I. Model development and prediction of aerosol properties, *Atmospheric Chemistry and Physics*, 5, 2227-2252, 2005.
- Stevens, R. G. and Pierce, J. R.: The contribution of plume-scale nucleation to global and regional aerosol and CCN concentrations: evaluation and sensitivity to emissions changes, *Atmospheric Chemistry and Physics*, 14, 13661-13679, 2014.
- 880 Stevens, R. G. and Pierce, J. R.: A parameterization of sub-grid particle formation in sulfur-rich plumes for global-and regional-scale models, *Atmospheric Chemistry and Physics*, 13, 12117-12133, 2013.
- Stevens, R. G., Pierce, J. R., Brock, C. A., Reed, M. K., Crawford, J. H., Holloway, J. S., Ryerson, T. B., Huey, L. G., and Nowak, J. B.: Nucleation and growth of sulfate aerosol in coal-fired

- 885 power plant plumes: sensitivity to background aerosol and meteorology, *Atmospheric Chemistry and Physics*, 12, 189-206, 2012.
- Stockwell, W. R., Kirchner, F., Kuhn, M., and Seefeld, S.: A new mechanism for regional atmospheric chemistry modeling, *Journal of Geophysical Research-Atmospheres*, 102, 25847-25879, 1997.
- 890 Strader, R., Lurmann, F., and Pandis, S. N.: Evaluation of secondary organic aerosol formation in winter, *Atmospheric Environment*, 33, 4849-4863, 1999.
- Sun, K. and Chen, X.: Spatio-temporal distribution of localized aerosol loading in China: A satellite view, *Atmospheric Environment*, 163, 35-43, 2017.
- 895 Tan, Z., Fuchs, H., Lu, K., Hofzumahaus, A., Bohn, B., Broch, S., Dong, H., Gomm, S., Haeseler, R., He, L., Holland, F., Li, X., Liu, Y., Lu, S., Rohrer, F., Shao, M., Wang, B., Wang, M., Wu, Y., Zeng, L., Zhang, Y., Wahner, A., and Zhang, Y.: Radical chemistry at a rural site (Wangdu) in the North China Plain: observation and model calculations of OH, HO₂ and RO₂ radicals, *Atmospheric Chemistry and Physics*, 17, 663-690, 2017.
- 900 Tan, Z., Lu, K., Dong, H., Hu, M., Li, X., Liu, Y., Lu, S., Shao, M., Su, R., Wang, H., Wu, Y., Wahner, A., and Zhang, Y.: Explicit diagnosis of the local ozone production rate and the ozone-NO_x-VOC sensitivities, *Science Bulletin*, 63, 1067-1076, 2018.
- Tan, Z., Lu, K., Jiang, M., Su, R., Wang, H., Lou, S., Fu, Q., Zhai, C., Tan, Q., Yue, D., Chen, D., Wang, Z., Xie, S., Zeng, L., and Zhang, Y.: Daytime atmospheric oxidation capacity in four Chinese megacities during the photochemically polluted season: a case study based on box model simulation, *Atmospheric Chemistry and Physics*, 19, 3493-3513, 2019.
- 905 Textor, C., Schulz, M., Guibert, S., Kinne, S., Balkanski, Y., Bauer, S., Berntsen, T., Berglen, T., Boucher, O., Chin, M., Dentener, F., Diehl, T., Easter, R., Feichter, H., Fillmore, D., Ghan, S., Ginoux, P., Gong, S., Kristjansson, J. E., Krol, M., Lauer, A., Lamarque, J. F., Liu, X., Montanaro, V., Myhre, G., Penner, J., Pitari, G., Reddy, S., Seland, O., Stier, P., Takemura, T., and Tie, X.: Analysis and quantification of the diversities of aerosol life cycles within AeroCom, *Atmospheric Chemistry and Physics*, 6, 1777-1813, 2006.
- 910 Wang, H., Lu, K., Chen, X., Zhu, Q., Chen, Q., Guo, S., Jiang, M., Li, X., Shang, D., Tan, Z., Wu, Y., Wu, Z., Zou, Q., Zheng, Y., Zeng, L., Zhu, T., Hu, M., and Zhang, Y.: High N₂O₅ Concentrations Observed in Urban Beijing: Implications of a Large Nitrate Formation Pathway, *Environmental Science & Technology Letters*, 4, 416-420, 2017.
- 915 Wang, S. X., Zhao, B., Cai, S. Y., Klimont, Z., Nielsen, C. P., Morikawa, T., Woo, J. H., Kim, Y., Fu, X., Xu, J. Y., Hao, J. M., and He, K. B.: Emission trends and mitigation options for air pollutants in East Asia, *Atmospheric Chemistry and Physics*, 14, 6571-6603, 2014.
- 920 Wang, Z., Li, J., Wang, X., Pochanart, P., and Akimoto, H.: Modeling of regional high ozone episode observed at two mountain sites (Mt. Tai and Huang) in East China, *Journal of Atmospheric Chemistry*, 55, 253-272, 2006a.
- Wang, Z. F., Xie, F. Y., Wang X. Q., An J. L., Zhu, J.: Development and Application of Nested Air Quality Prediction Modeling System, *Chinese Journal of Atmospheric Sciences*, 30(5), 778-790, 2006b. (in Chinese).
- 925 Wang, Z. B., Hu, M., Mogensen, D., Yue, D. L., Zheng, J., Zhang, R. Y., Liu, Y., Yuan, B., Li, X., Shao, M., Zhou, L., Wu, Z. J., Wiedensohler, A., and Boy, M.: The simulations of sulfuric acid concentration and new particle formation in an urban atmosphere in China, *Atmospheric Chemistry and Physics*, 13, 11157-11167, 2013.

- Wang, Z. F., Ueda, H., and Huang, M. Y.: A deflation module for use in modeling long-range
930 transport of yellow sand over East Asia, *Journal of Geophysical Research-Atmospheres*, 105,
26947-26959, 2000.
- Wei, Y., Chen H. S., Liu H. et al.: Multi-scale Simulation of the Influence of Spring Dust on
PM2.5 Concentration in Central Shaanxi Area, China. *Chinese Journal of Atmospheric
Sciences (in Chinese)*, 44(1): 76-92. doi:10.3878/j.issn.1006-9895.1902.18224, 2020.
- 935 Wei, Y., J., Li, Z. F., Wang, H. S., Chen, Q. Z., Wu, & J. J., Li, et al. : Trends of surface PM2.5
over Beijing-Tianjin-Hebei in 2013-2015 and their causes: emission controls vs.
meteorological conditions, *Atmospheric and Oceanic Science Letters*, 10, 276-283,
<https://doi.org/10.1080/16742834.2017.1315631>, 2017.
- Wei, Y., Chen, X., Chen, H., Li, J., Wang, Z., Yang, W., Ge, B., Du, H., Hao, J., Wang, W., Li, J.,
940 Sun, Y., and Huang, H.: IAP-AACM v1.0: a global to regional evaluation of the atmospheric
chemistry model in CAS-ESM, *Atmospheric Chemistry and Physics*, 19, 8269-8296, 2019.
- Wei, Y., Chen, X., Chen, H., Yang, W., Wu Q., Li, J., Wang, Z. IAP-AACM+sub-grid particle
formation (SGPF) scheme v1.0, <https://doi.org/10.5281/zenodo.4383361>, 2020.
- Williams, J. E., Strunk, A., Huijnen, V., and van Weele, M.: The application of the Modified Band
945 Approach for the calculation of on-line photodissociation rate constants in TM5: implications
for oxidative capacity, *Geoscientific Model Development*, 5, 15-35, 2012.
- Wu, J., Bei, N., Hu, B., Liu, S., Wang, Y., Shen, Z., Li, X., Liu, L., Wang, R., Liu, Z., Cao, J., Tie,
X., Molina, L. T., and Li, G.: Aerosol-photolysis interaction reduces particulate matter during
wintertime haze events, *Proceedings of the National Academy of Sciences of the United
950 States of America*, 117, 9755-9761, 2020.
- Wu, Q. Z., Wang, Z. F., Gbaguidi, A., Gao, C., Li, L. N., and Wang, W.: A numerical study of
contributions to air pollution in Beijing during CAREBeijing-2006, *Atmospheric Chemistry
and Physics*, 11, 5997-6011, 2011.
- Xu, W. Y., Zhao, C. S., Ran, L., Lin, W. L., Yan, P., and Xu, X. B.: SO2 noontime-peak
955 phenomenon in the North China Plain, *Atmospheric Chemistry and Physics*, 14, 7757-7768,
2014.
- Yang, Y., Ren, L., Li, H., Wang, H., Wang, P., Chen, L., Yue, X., Liao, H., 2020. Fast Climate
Responses to Aerosol Emission Reductions During the COVID-19 Pandemic. *Geophysical
Research Letters* 47.
- 960 Yang, Y., Smith, S.J., Wang, H., Mills, C.M., Rasch, P.J., 2019. Variability, timescales, and
nonlinearity in climate responses to black carbon emissions. *Atmospheric Chemistry and
Physics* 19, 2405-2420.
- Yang, Y., Wang, Y., Zhou, P., Yao, D., Ji, D., Sun, J., Wang, Y., Zhao, S., Huang, W., Yang, S.,
Chen, D., Gao, W., Liu, Z., Hu, B., Zhang, R., Zeng, L., Ge, M., Petaja, T., Kerminen, V.-M.,
965 Kulmala, M., and Wang, Y.: Atmospheric reactivity and oxidation capacity during summer at
a suburban site between Beijing and Tianjin, *Atmospheric Chemistry and Physics*, 20,
8181-8200, 2020.
- Yu, F.: Diurnal and Seasonal Variations of Ultrafine Particle Formation in Anthropogenic SO2
Plumes, *Environmental Science & Technology*, 44, 2011-2015, 2010.
- 970 Yu, F. and Luo, G.: Simulation of particle size distribution with a global aerosol model:
contribution of nucleation to aerosol and CCN number concentrations, *Atmospheric
Chemistry and Physics*, 9, 7691-7710, 2009.

- Yue, D. L., Hu, M., Zhang, R. Y., Wu, Z. J., Su, H., Wang, Z. B., Peng, J. F., He, L. Y., Huang, X. F., Gong, Y. G., and Wiedensohler, A.: Potential contribution of new particle formation to cloud condensation nuclei in Beijing, *Atmospheric Environment*, 45, 6070-6077, 2011.
- 975 Zanatta, M., Laj, P., Gysel, M., Baltensperger, U., Vratolis, S., Eleftheriadis, K., Kondo, Y., Dubuisson, P., Winiarek, V., Kazadzis, S., Tunved, P., and Jacobi, H.-W.: Effects of mixing state on optical and radiative properties of black carbon in the European Arctic, *Atmospheric Chemistry and Physics*, 18, 14037-14057, 2018.
- 980 Zaveri, R. A. and Peters, L. K.: A new lumped structure photochemical mechanism for large-scale applications, *Journal of Geophysical Research-Atmospheres*, 104, 30387-30415, 1999.
- Zhang, H., Li, J., Ying, Q., Yu, J. Z., Wu, D., Cheng, Y., He, K., and Jiang, J.: Source apportionment of PM_{2.5} nitrate and sulfate in China using a source-oriented chemical transport model, *Atmospheric Environment*, 62, 228-242, 2012.
- 985 Zhang, H., Zhang, M., Jin, J., Fei, K., Ji, D., Wu, C., Zhu, J., He, J., Chai, Z., Xie, J., Dong, X., Zhang, D., Bi, X., Cao, H., Chen, H., Chen, K., Chen, X., Gao, X., Hao, H., Jiang, J., Kong, X., Li, S., Li, Y., Lin, P., Lin, Z., Liu, H., Liu, X., Shi, Y., Song, M., Wang, H., Wang, T., Wang, X., Wang, Z., Wei, Y., Wu, B., Xie, Z., Xu, Y., Yu, Y., Yuan, L., Zeng, Q., Zeng, X., Zhao, S., Zhou, G., and Zhu, J.: CAS-ESM 2: Description and Climate Simulation
- 990 Performance of the Chinese Academy of Sciences (CAS) Earth System Model (ESM) Version 2, n/a, e2020MS002210.
- Zhang, H., Zhang, M., Jin, J., Fei, K., Ji, D., Wu, C., Zhu, J., He, J., Chai, Z., Xie, J., Dong, X., Zhang, D., Bi, X., Cao, H., Chen, H., Chen, K., Chen, X., Gao, X., Hao, H., Jiang, J., Kong, X., Li, S., Li, Y., Lin, P., Lin, Z., Liu, H., Liu, X., Shi, Y., Song, M., Wang, H., Wang, T.,
- 995 Wang, X., Wang, Z., Wei, Y., Wu, B., Xie, Z., Xu, Y., Yu, Y., Yuan, L., Zeng, Q., Zeng, X., Zhao, S., Zhou, G., and Zhu, J.: CAS-ESM 2: Description and Climate Simulation
- Performance of the Chinese Academy of Sciences (CAS) Earth System Model (ESM) Version 2, n/a, e2020MS002210.
- Zhang, Q., Zheng, Y., Tong, D., Shao, M., Wang, S., Zhang, Y., Xu, X., Wang, J., He, H., Liu, W.,
- 1000 Ding, Y., Lei, Y., Li, J., Wang, Z., Zhang, X., Wang, Y., Cheng, J., Liu, Y., Shi, Q., Yan, L., Geng, G., Hong, C., Li, M., Liu, F., Zheng, B., Cao, J., Ding, A., Gao, J., Fu, Q., Huo, J., Liu, B., Liu, Z., Yang, F., He, K., and Hao, J.: Drivers of improved PM_{2.5} air quality in China from 2013 to 2017, *Proceedings of the National Academy of Sciences of the United States of America*, 116, 24463-24469, 2019.
- 1005 Zhang, Q. H., Zhang, J. P., and Xue, H. W.: The challenge of improving visibility in Beijing, *Atmospheric Chemistry and Physics*, 10, 7821-7827, 2010.
- Zhang, Y., Chen, Y., Sarwar, G., and Schere, K.: Impact of gas-phase mechanisms on Weather Research Forecasting Model with Chemistry (WRF/Chem) predictions: Mechanism implementation and comparative evaluation, *Journal of Geophysical Research-Atmospheres*,
- 1010 117, 2012.
- Zhang, Y., Hemperly, J., Meskhidze, N., and Skamarock, W. C.: The Global Weather Research and Forecasting (GWRF) Model: Model Evaluation, Sensitivity Study, and Future Year Simulation, *Atmospheric and Climate Sciences*, 02, 231-253, 10.4236/acs.2012.23024, 2012.
- Zhang, Y.-L. and Cao, F.: Fine particulate matter (PM_{2.5}) in China at a city level, *Scientific Reports*, 5, 2015.
- 1015 Zheng, B., Tong, D., Li, M., Liu, F., Hong, C., Geng, G., Li, H., Li, X., Peng, L., Qi, J., Yan, L.,

Zhang, Y., Zhao, H., Zheng, Y., He, K., and Zhang, Q.: Trends in China's anthropogenic emissions since 2010 as the consequence of clean air actions, *Atmospheric Chemistry and Physics*, 18, 14095-14111, 2018.

1020 Zhu, T., Shang, J., Zhao D. F.: The roles of heterogeneous chemical processes in the formation of an air pollution complex and gray haze. *SCIENTIA SINICA Chimica*, 40, 1731-1740, <https://doi.org/10.1360/zb2010-40-12-1731>, 2010.

1025 Zhuang, X., Wang, Y., He, H., Liu, J., Wang, X., Zhu, T., Ge, M., Zhou, J., Tang, G., and Ma, J.: Haze insights and mitigation in China: An overview, *Journal of Environmental Sciences*, 26, 2-12, 2014.