### CoupModel (v6.0): an ecosystem model for coupled phosphorus, nitrogen, and carbon dynamics – evaluated against empirical data from a climatic and fertility gradient in Sweden

5

Hongxing He<sup>1</sup>, Per-Erik Jansson<sup>2</sup>, Annemieke I. Gärdenäs<sup>1</sup>

<sup>1</sup>Department of Biological and Environmental Sciences, University of Gothenburg, PO Box 460, Gothenburg 40530, Sweden

<sup>2</sup>Department of Land and Water Resources Engineering, Royal Institute of Technology

10 (KTH), 100 44 Stockholm, Sweden Correspondence to: Hongxing He (<u>hongxing-he@hotmail.com</u>) and Annemieke Gärdenäs (<u>annemieke.gardenas@bioenv.gu.se</u>)

#### 15 Abstract

This study presents the integration of the phosphorus (P) cycle into CoupModel (v6.0, referred to as Coup-CNP). The extended Coup-CNP, which explicitly considers the symbiosis between soil microbes and plant roots, enables simulations of coupled carbon (C), nitrogen (N) and P dynamics for terrestrial ecosystems. The model was evaluated against 20 observed forest growth and measured leaf C/P, C/N and N/P ratios in four managed forest regions in Sweden. The four regions form a climatic and fertility gradient from 64°N (northern Sweden) to 56°N (southern Sweden), with mean annual temperature varying from 0.7-7.1 °C and soil C/N and C/P ratios varying between 19.8-31.5 and 425-633, respectively. The growth of the southern forests was found to be P-limited, with harvested biomass representing the largest P losses over the studied rotation period. The simulated P budgets 25 revealed that southern forests are losing P, while northern forests have balanced P budgets. Symbiotic fungi accounted for half of total plant P uptake across all four regions, which highlights the importance of fungal-tree interactions in Swedish forests. The results of a sensitivity analysis demonstrated that optimal forest growth occurs at a soil N/P ratio between

30 15-20. A soil N/P ratio above 15-20 will result in decreased soil C sequestration and P leaching, along with a significant increase in N leaching. Simulations from the new Coup-

CNP model provide strong evidence that P fluxes need to be further considered in studies of how ecosystems and C turnover react to climate change. We conclude that biogeochemical models should include the P cycle to reliably assess how both climate change and N deposition will affect C sequestration and nutrient leaching over long time periods.

#### **1** Introduction

35

Phosphorus (P) is an essential element for photosynthetic plants in terrestrial ecosystems, with the P cycle coupled to Carbon (C) and Nitrogen (N) fluxes through processes such as
decomposition of soil organic matter and nutrient uptake (Lang et al., 2016; Vitousek et al., 2010). A steep increase in the anthropogenic release of C and N to the atmosphere relative to P release has altered plant and soil nutrient stoichiometry, leading to new forcing conditions (Elser et al., 2007; Penuelas et al., 2013). For instance, numerous monitoring studies have revealed increasing N/P ratios in plants and soils, especially in forests from North America

- 45 (Crowley et al., 2012; Gress et al., 2007; Tessier and Raynal, 2003) and central and northern Europe (Braun et al., 2010; Jonard et al., 2015; Talkner et al., 2015). Such trends are generally assumed to indicate that these ecosystems are shifting from being N-limited to either co-limited by both N and P or P-limited (Elser et al., 2007; Saito et al., 2008; Vitousek et al., 2010; Du et al., 2020). Human activities are expected to continue increasing
- 50 atmospheric N deposition; as such, P availability and P cycle dynamics will become progressively more important in regulating the biogeochemistry of terrestrial ecosystems and amplifying feedbacks relevant to climate change, e.g. limiting the growth response of plants to increased temperature (Deng et al., 2017; Fleischer et al., 2019; Goll et al., 2017). Nevertheless, the P cycle is seldom incorporated into ecosystem model structures.
- 55 Incorporating the P cycle is essential to improving how global models can assess climate-C cycling interactions (Reed et al., 2015). Most of the process-based models that can simulate P cycling were specifically developed for agricultural systems and focus on the soil processes, e.g., EPIC (Jones et al., 1984, Gassman et al., 2005), ANIMO (Groenendijk et al., 2005), and GLEAMS (Knisel and Turtola, 2000). A few catchment-scale models focus on surface water
- 60 quality, e.g. SWAT (Arnold et al., 2012), HYPE (Arheimer et al., 2012), and INCA-P (Jackson-Blake et al., 2016), aim to simulate how crop management influences P leaching and thus, consider processes such as nutrient retention, leaching, and transport. The C response to P limitation has recently been studied through several empirical and field studies (Van Sundert et al., 2019; Du et al., 2020). For example, Van Sundert et al. (2019) showed
- 65 that the productivity of European beech (Fagus sylvestris) forests is negatively related to soil

organic carbon concentrations and mineral C/P ratios. Several global vegetation models have included the P cycle to study how it affects the C cycle (Goll et al., 2012, 2017; Wang et al., 2010; Yang et al., 2014; Zhu et al., 2016; Thum et al., 2019). These P enabled models differ in how they describe soil P dynamics, i.e., implicitly or explicitly through symbiotic 70 relationships with mycorrhiza and other soil microbes, plant P use and acquisition strategies, ultimately leading to considerable uncertainty in the C response (Fleischer et al., 2019; Medlyn et al., 2016; Reed et al., 2015). Medlyn et al. (2016) applied six global vegetation models - including two coupled Carbon-Nitrogen-Phosphorus (CNP) models (CABLE and CLM4.0-CNP) - to study how the C cycle of the Eucalyptus-Free-Air CO<sub>2</sub> Enrichment experiment responds to elevated CO<sub>2</sub> (eCO<sub>2</sub>) levels. The results demonstrated notable 75 variations, ranging from 0.5 to 25%, in predicted net primary productivity. The CNP models that explicitly considered the P dependency of C assimilation predicted the lowest eCO<sub>2</sub> response. Yu et al. (2018) included the P cycle in the ForSAFE field-scale biogeochemical model to study the P budget of a southern Swedish Spruce forest site. They concluded that 80 internal turnover from mineralization of soil organic matter affects the P supply more than weathering. Fleischer et al. (2019) demonstrated that four CNP models, when applied to the Amazon forest, provide up to 50% lower estimates of the eCO<sub>2</sub>-induced biomass increment than the 10 coupled C-N models. They suggested that the inclusion of flexible tissue stoichiometry and enhanced plant P acquisition could improve the ability of terrestrial

85 ecosystem models to simulate C-P cycle coupling. Most terrestrial plants live in symbiosis with mycorrhizal fungi to increase the uptake capacity of P, among other nutrients (Smith and Read, 2008). Several studies have shown that the depletion zone around plant roots, which is caused by plant uptake and the immobile nature of mineral P, increases when a plant interacts with mycorrhizal fungi (Bolan, 1991;

- 90 Schnepf and Roose, 2006; Smith, 2003). Global meta-analysis studies have highlighted that the symbiosis between plants and soil mycorrhizal fungi strongly influences plant P availability and, subsequently, affects plant growth (Terrer et al., 2016; 2019). Previous research has shown that mycorrhizal fungi can receive between 1 and 25% of plant photosynthates and constitute as much as 70% of the total soil microbial biomass; thus, it is
- 95 clear that this symbiont has a major impact on soil C sequestration (Averill et al., 2014; Clemmensen et al., 2013; Staddon et al., 2003). Even though there is a well-established link between mycorrhizal fungi and plant P nutrition (Bucher, 2007; Read and Perez-Moreno, 2003; Rosling et al., 2016), this factor is seldom included in ecosystem models (Smith and Read, 2008). To the best of our knowledge, only Orwin et al. (2011) have presented an

- 100 ecosystem model that consider C, N and P together with symbiotic fungi. They found that considering organic nutrient uptake by symbiotic fungi in an ecosystem model can significantly increase soil C storage, with this effect more pronounced under nutrient-limited conditions. In this model, organic nutrient uptake reflects a pathway through which plants can utilize organic nutrients by biochemical mineralization, either in symbiosis with mycorrhizal
- 105 fungi or via root exudates (e.g. Schachtman et al., 1998, Gärdenäs et al. 2011, Richardson et al. 2009). However, plant growth is static in the model presented by Orwin et al. (2011); as such, plant-soil or plant-environment interactions are largely ignored. Our model (Eckersten and Beier, 1998; He et al., 2018) also includes a short-cut for nutrient uptake that relies on rhizosphere processes. The assumption is that nutrients released by biochemical
- 110 mineralization are instantly taken up by symbiotic microbes and/or the plants, thereby bypassing the soil matrix solution. He et al. (2018) integrated the MYCOFON model (Meyer et al., 2009) into CoupModel v5 to ensure that the symbiosis between plant roots and mycorrhiza would be sufficiently considered, and compared the results with a previous implicit representation of N uptake in forest ecosystems with limited N availability.
- 115 CoupModel v5 assumes that carbohydrates provided by plants are the primary driver of mycorrhizal responses to N availability, and that fungal uptake of N will influence host plant photosynthesis. We argued that terrestrial ecosystem models which explicitly consider mycorrhizal interactions should also take into account P cycling due to the significant role of symbiont mycorrhiza for P uptake in P-limited environments. For this reason, we developed a new version of CoupModel that includes the P cycle.
- The overall objective of this study was to improve the current understanding of C, N and P cycle interactions in forest ecosystems by presenting a new scheme for modeling P dynamics. More specifically, the study had the following aims: 1) to present the new CoupModel v6.0, which explicitly includes the P cycle and interactions between the N and P cycles; 2) to
- 125 estimate the regional C, N, and P budgets of Swedish forests along a climatic and fertility gradient; and 3) to demonstrate how soil N and P availability influence growth, soil C and nutrient leaching. Hence, we present a new version of CoupModel (v6.0), hereafter referred to as Coup-CNP, which explicitly simulates the P cycle. The key features of the new Coup-CNP model are: 1) coupled C, N, and P dynamics; 2) explicit representations of symbiosis
- 130 between plant roots and mycorrhiza, along with implicit routes through which non-symbiotic microbes contribute to N and P uptake from the soil; 3) flexible CNP stoichiometry for plant components, soil organic matter, and symbiotic microbes; 4) dynamic nutrient demand and uptake, as well as photosynthesis and growth rates, all of which are regulated by N and P

availability; and 5) simultaneous uptake of nutrients to roots or symbiotic mycorrhizae from
both inorganic and organic pools. The Coup-CNP model was evaluated using four forest
regions situated along a climatic and fertility gradient in Sweden that has earlier been
considered by He et al. (2018) and Svensson et al. (2008).

#### 2 Description of model structure and phosphorus model

#### 140 **2.1 Brief description of CoupModel (v5)**

The CoupModel platform (coupled heat and mass transfer model for soil-plant-atmosphere systems) is a process-based model designed to simulate water and heat fluxes, along with C and N cycles, in terrestrial ecosystems (Jansson, 2012). The main model structure is a one-dimensional, vertical model, with one or two layers of vegetation (for example a tree and

- 145 field layer as in this application) on a multi-layered soil profile. The core of the model consists of five sets of coupled partial differential equations that cover water, heat, C, N, and P cycles (the latter is only included in v6.0). They are numerically solved using an explicit forward difference model scheme (Euler integration, for more details see pp 400-401 in Jansson and Karlberg (2011)), which means that the current size of a state variable is
- 150 calculated based on fluxes to and from the state variable during the previous time step. In this application, we used a daily time step for each equation, although a smaller time step was applied for the water and heat calculations during specific events representing peaks in water and/or heat flow, e.g., snow melting, to ensure numerical stability and accuracy. The model is driven by climatic data precipitation, air temperature, relative humidity, wind speed, and
- 155 global radiation, i.e. sum of direct and diffuse shortwave incoming radiation- and can simulate ecosystem dynamics with daily resolution. Vegetation is described using the "multiple-big leaves" concept, i.e. two vegetation layers (trees and understory plants) are simulated taking into account mutual competition for light interception, water uptake and soil N (Jansson and Karlberg, 2011). The model and technical description (Jansson and Karlberg,
- 160 2011) is freely available at <u>www.coupmodel.com</u>. Furthermore, Jansson (2012) has previously presented CoupModel use, calibration and validation, while He et al. (2018) introduced an explicit plant-mycorrhizal representation (CoupModel v5).

#### 2.2 Phosphorus cycle representation in CoupModel (v6.0)

165 Coup-CNP was extended with P cycle representation to enable simulations of coupled C, N and P dynamics for terrestrial ecosystems while explicitly considering the symbiosis between soil microbes and plant roots. Coup-CNP has P state variables and fluxes representing different plant parts, symbiotic microbes, soil organic P forms ( $P_o$ , P that is bound to organic C in the soil) and soil inorganic P forms ( $P_i$ ) (Fig. 1). To clarify the coupling between C, N

- and P cycles, the C and N state variables and major N and N+P fluxes are given in Fig. 1.
   P within plants is partitioned into grain, leaf, stem, coarse root, fine root, along with P in symbiotic microbes, which is analogous to how C and N are represented in CoupModel v5.0 (Fig. 1). In this paper, we use mycorrhizal fungi as the main representation of plant-microbe symbiosis; the same concept is also applicable for other symbiosis microbes. Soil organic P is
- divided into three state variables litter (*P<sub>Litter</sub>*), humus (*P<sub>Humus</sub>*), and dissolved organic (*P<sub>DOP</sub>*)
   in every soil layer, which is analogous to how C and N are represented in CoupModel v2.0 (Fig. 1). Non-symbiosis microbes is implicitly included in litter. Soil inorganic P is represented by both new and renewed state variables. A new state variable is soil solid inorganic P, *P<sub>solid</sub>*, which is a lumped pool containing primary and secondary mineral compounds containing P such as apatite (and occluded P) (Smeck, 1985; Wang et al., 2007).
- $P_{ilab}$  is the sum of phosphate ions absorbed by soil and those in soil solution, which follows how the mineral pool is represented in the salt tracer module of CoupModel (Gärdenäs et al., 2006). Instantaneous equilibrium between adsorbed and soil solution P is assumed. Plants and microbes take up phosphate ions from the  $P_{ilab}$  pool.  $P_{isol}$ , which is an intrinsic part of  $P_{ilab}$ ,
- 185 can be compared with the sum of the N state variables NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> (Fig. 1). We developed the P model in a way that 1) focuses on the P processes that are most relevant to biogeochemical cycling, e.g., dynamic plant growth and P leaching, and 2) follows the conceptual structure of CoupModel as closely as possible. The P processes that are relevant to biogeochemical cycling are described in more detail below. Appendix A further describes
- 190 processes that are analogous to those of the N cycle, e.g., atmospheric deposition, fertilization (A.1), mineralization-immobilization (A.2), plant growth and uptake (A.3), litterfall (A.4), leaching and surface runoff (A.5), and removal of plant harvest (A.6). For simplicity, the equations are given in a form that reflects one time step and one of the layers of the entire soil profile. This paper has been formatted in a way that conforms to the CoupModel
- 195 nomenclature; more specifically, a capitalized P refers to state variables while a lowercase p refers to parameters that are related to P processes.

#### 2.2.1 Weathering

During weathering soil solid inorganic P (Psolid) is transformed into labile P (Pilab) (Fig. 1; eq.

200 1). The weathering rate depends on soil pH and temperature (Guidry and Machenzie, 2000), and is calculated as,

$$P_{solid \to ilab} = k_w \times f(T_s) \times f(pH) \times P_{solid}$$
<sup>(1)</sup>

Where  $P_{solid \rightarrow ilab}$  is the flux rate of weathering (g P m<sup>-2</sup> day<sup>-1</sup>),  $k_w$  is a first-order integrated weathering rate coefficient (day<sup>-1</sup>) which depends on lithology, rates of physical erosion and soil properties (Table 3). The erosion affect the weathering rate by reducing the pool size of  $P_{solid}$  (eq. A.14).  $f(T_s)$  and f(pH) are response functions of soil temperature and pH, respectively, while  $P_{solid}$  is the size of the  $P_{solid}$  pool (g P m<sup>-2</sup>), determined by,

$$P_{solid} = \delta_P \times \rho_{bulk} \times \Delta z_{laver} \times 10^6 \tag{2}$$

of each soil layer (g cm<sup>-3</sup>), and  $\Delta z_{layer}$  is the thickness of the simulated soil layer (m).

Where  $\delta_P$  is the prescribed  $P_{solid}$  content for each soil layer (g P g dry soil<sup>-1</sup>), with reported ranges from  $1 \times 10^{-4}$  to  $1.5 \times 10^{-3}$  g P g soil<sup>-1</sup> (Yang et al., 2014),  $\rho_{bulk}$  is the dry bulk density

The temperature effect can be expressed as an Arrhenius function (3), where  $E_{a,wea}$  is the activation energy parameter (J mol<sup>-1</sup>) for minerals (i.e., apatite; available from empirical studies), *R* is the gas constant (J K<sup>-1</sup> mol<sup>-1</sup>), *T<sub>s</sub>* is the simulated soil temperature in °C, *T<sub>s,0</sub>* is a parameter (°C) which normalizes the function  $f(T_s)=1$ , and  $T_{abszero}$  is -273.15 °C.

$$f(T) = e^{\left(-\frac{E_{a,wea}}{R} \times \left(\frac{1}{T_s + T_{abszero}} - \frac{1}{T_{s,0} + T_{abszero}}\right)\right)}$$
(3)

Alternatively, the existing Ratkowsky function, O'Neill function or  $Q_{10}$  method can be used to determine the temperature response in CoupModel.

The effect of soil pH on weathering can be calculated as,

220 
$$f(pH) = 10^{n_H \times |pH_{opt} - pH|}$$
 (4)

where  $n_H$  is a parameter that describes the sensitivity of the soil to pH changes when it deviates from  $pH_{opt}$ , an optimal pH value for weathering (Table 3).

#### 2.2.2 Inorganic soluble phosphorus dynamics

When  $P_i$  is added to a soil ecosystem, the soluble ( $P_{isol}$ ) and adsorbed P pools reach equilibrium in less than one hour (Cole et al., 1977; Olander and Vitousek, 2005). As a daily time step is applied to this model, we assume that  $P_{isol}$  and the adsorbed part of  $P_{ilab}$  are

205

210

always in equilibrium (eq. 5). The modified Langmuir isotherm (Barrow, 1979) was used to model the fast and reversible sorption process within  $P_{ilab}$ .

$$230 \qquad P_{ilab,con} = p_{\max,ads} \times \frac{P_{isol}}{c_{50,ads} + P_{isol}}$$
(5)

Where  $P_{ilab,con}$  is the concentration of labile pool (g P g soil<sup>-1</sup>),  $p_{max,ads}$  is the maximum sorption capacity of the labile pool (g P g soil<sup>-1</sup>), and  $c_{50,ads}$  is an empirical parameter corresponding to 50% of P saturation (g P m<sup>-2</sup>) (Table 3). It should be noted that  $P_{ilab,con}$  can also be calculated using equation (2) as  $P_{ilab,con} = P_{ilab} / (\rho_{bulk} \times \Delta z_{layer} \times 10^6)$ .

235

#### 2.2.3 Soil inorganic phosphorus dynamics and nutrient short-cut uptake

Atmospheric P deposition is assumed to directly flow into the labile inorganic P pool ( $P_{ilab}$ ) in the uppermost soil layer (eq. A.1 in Appendix A). If mineral  $P_i$  fertilizer is applied at the soil surface, the  $P_i$  first enters an undissolved fertilizer pool, after which  $P_i$  from this pool gradually dissolves into the labile P pool following a decay-type function (eq. A.1). P can also be added as an external organic substrate (faeces or manure). In this case, P moves to the surface faeces ( $P_{ofae}$ ), litter ( $P_{Litter}$ ), and labile ( $P_{ilab}$ ) P pools according to the composition of the manure.  $P_i$  within  $P_{isol}$  and dissolved organic P ( $P_{DOP}$ ) can be transported by water flows between layers or from a layer to a drainage outlet (eq. A. 12-13). The soil surface layer may also lose solid inorganic P ( $P_{solid}$ ) by erosion, which is driven by surface runoff (eq. A.14).

- also lose solid inorganic P (*P<sub>solid</sub>*) by erosion, which is driven by surface runoff (eq. A.14).
  P mineralization is conceptually divided into biological and biochemical mineralization (eq. A.2-A.6) following McGill and Cole (1981). Biological mineralization, which is regulated by temperature and moisture, represents the microbe-mediated oxidation of organic matter, during which nutrients (P and N) are immobilized by non-symbiotic microbes or transferred
- 250 from litter to humus (Fig. 1; eq. A.2). Biochemical mineralization, on the other hand, describes the release of  $P_i$  through extracellular enzymatic action (e.g., phosphatases from root exudates), which is driven by plant demand for nutrients (Richardson and Simpson, 2011). In Coup-CNP, biochemical mineralization is conceptually included in the short-cut uptake of nutrients (called organic uptake in earlier CoupModel publications) and assumed to
- 255 be driven by the unfulfilled plant P demand after  $P_{ilab}$  root uptake (eq. A.8) but regulated by the availability in other P pools (i.e. short-cut uptake coefficients in eq. A.4). The assumption is that under P-limited conditions, plant roots and symbiotic fungi bypass  $P_{ilab}$ , and obtain mineralized  $P_i$  directly from organic  $P_{Litter}$  and  $P_{Humus}$  (Fig. 1; eq. A.4).

#### 260 **2.2.4 Plant growth under phosphorus and nitrogen limitation**

Plant photosynthesis is modeled by a "light use efficiency" approach (Monteith, 1965, eq. 6)). We adopted Liebig's law of minimum to simulate how plants respond to multiple nutrient stress (Liebig, 1840). This approach assumes that the nutrient (N, P) which has the smallest supply relative to the corresponding plant demand will limit growth (eq. 7).

265 
$$C_{a \to plant} = \varepsilon_L \times f(T_{leaf}) \times f(nutrient) \times f(\frac{E_{ta}}{E_{tp}}) \times R_s$$
 (6)

$$f(nutrient) = min\left(f(C / N_{leaf}); f(C / P_{leaf})\right)$$
(7)

Where  $C_{a \rightarrow plant}$  is the plant carbon assimilation rate (g C m<sup>-2</sup> day<sup>-1</sup>),  $\varepsilon_L$  is the coefficient for radiation use efficiency (g C J<sup>-1</sup>),  $f(T_{leaf})$ , f(nutrient) and  $f(E_{ta}/E_{tp})$  are response functions of leaf temperature, leaf nutrient status (N<sub>leaf</sub>, P<sub>leaf</sub>) in proportion to its C content, and water,

- 270 respectively, and  $R_s$  represents radiation absorbed by the canopy (J m<sup>-2</sup> day<sup>-1</sup>). Details concerning  $f(T_{leaf})$ ,  $f(E_{ta}/E_{tp})$ , as well as growth and maintenance respiration, can be found in Jansson and Karlberg (2011). Plant demand for nutrients was estimated through defined optimum ratios (equ A.9). The nutrient response function f(nutrient), which includes P, is described below.
- As is the case with N, the photosynthetic process responds to the leaf C/P ratio, a dynamic which has been modeled by Ingestad and Ågren (1992). Hence, photosynthesis is not limited by P below an optimum C/P ratio ( $p_{CP,opt}$ ), while between  $p_{CP,opt}$  and  $p_{CP,th}$  the response function decreases linearly until it reaches zero,

$$f(C / P_{leaf}) = \begin{cases} 1 & C / P_{leaf} < p_{CP,opt} \\ 1 + (\frac{C / P_{leaf} - p_{CP,opt}}{p_{CP,opt} - p_{CP,th}}) & p_{CP,th} \le C / P_{leaf} \ge p_{CP,opt} \\ 0 & C / P_{leaf} > p_{CP,th} \end{cases}$$
(8)

280 Where C/P<sub>leaf</sub> is the actual leaf C/P ratio and  $p_{CP,opt}$  and  $p_{CP,th}$  are parameters that vary between plant species (Table 3). The leaf C/P ratio is calculated at each time step using the leaf state variables C and P.

#### 2.2.5 Symbiotic mycorrhizal fungal growth and phosphorus dynamics

The following section describes the fungal processes that are specific to P. Plant C allocation to mycorrhizal fungi is influenced by soil  $P_i$  concentrations. We thus introduce a response function  $f_{a \rightarrow fungi}(P_i)$  to account for reductions in plant C allocation to mycorrhizal fungi when soil  $P_i$  concentrations are high, which is analogous to the N response function in He et al. (2018),

290 
$$f_{a \to fungi}(P_i) = e^{(-p_{avail} \times P_{isol}^2)^3}$$
 (9)

Where  $P_{isol}$  is the total soluble  $P_i$  in the soil (g P m<sup>-2</sup>) (eq. 5), and  $p_{avail}$  is a reduction parameter (m<sup>4</sup> g<sup>-2</sup> P) (Table 3). According to Bahr et al. (2015), mycorrhizal fungal biomass decreases when either N or P is added to the system, while the addition of both N and P leads to the most significant decrease. These multiple responses were integrated into the model so that potential fungal growth would decline as a result of either increasing soil N or P.

$$C_{a \to fungi,\max} = C_{a \to root} \times p_{fmax} \times (f_{a \to fungi}(P_i) \times f_{a \to fungi}(N))$$
(10)

Where  $C_{a \rightarrow fungi,max}$  is the defined maximum C flow that plants allocate to fungi (g C m<sup>-2</sup> day<sup>-</sup> <sup>1</sup>),  $C_{a \rightarrow root}$  is the total C allocated to both root and mycorrhiza (g C m<sup>-2</sup> day<sup>-1</sup>) (eq. A.10),  $p_{fmax}$ is a parameter that defines the maximum C fraction allocated to mycorrhiza from the total

300 root and mycorrhiza C pool (Table 3), and  $f_{a \rightarrow fungi}(P_i)$  and  $f_{a \rightarrow fungi}(N)$  are response functions that describe how soil N and P availability regulates maximum mycorrhizal fungal growth (eq. 9).

The actual growth of mycorrhizal fungi,  $C_{a \rightarrow fungi}$  (g C m<sup>-2</sup> day<sup>-1</sup>), is limited by the defined maximum growth,  $C_{a \rightarrow fungi, max}$  (eq. 10) calculated as,

305 
$$C_{a \to fungi} = \min\left\{ \left[ \left( (C_{root} \times p_{fopt}) - C_{fungi} \right) \times \min(f(N_{supply}); f(P_{supply})) \right]; C_{a \to fungi, max} \right\}$$
(11)

Where  $C_{root}$  is the total root C content (g C m<sup>-2</sup>),  $p_{fopt}$  is the defined optimum ratio parameter between fungal and root C content (Table 3), C<sub>fungi</sub> is the total C content of fungi (g C m<sup>-2</sup>), and  $f(N_{supply})$  and  $f(P_{supply})$  are response functions of fungal growth to the amount of N and P (both uptake from inorganic pools and short-cut uptake from organic pools) transferred from 310 fungi to plants (eq. 12). In this way, mycorrhizal fungal growth is also influenced by how efficiently the fungi transfer nutrients to the host plant (eq. 11). The model follows the assumption that plants provide fungi with C as long as their investment is outweighed by the benefits (i.e., acquired N or P) (Nasholm et al., 2013; Nehls, 2008). We further assume the C investment will be limited by the minimum nutrient supply efficiency provided by fungi.

 $f(P_{supply})$  is calculated as, 315

$$f(P_{supply}) = \begin{cases} 1 & P_{fungi \rightarrow plant} \\ \hline P_{fungi \rightarrow plant} + P_{ilab \rightarrow root} \\ \hline P_{fungi \rightarrow plant} + P_{ilab \rightarrow root} \\ \hline P_{fungi \rightarrow plant, th} > P_{fungi \rightarrow plant} \end{cases}$$
(12)

$$P_{fungi \to plant,th} = p_{fth} \times (P_{fungi \to plant} + P_{ilab \to root})$$
(13)

Where  $P_{fungi \rightarrow plant,th}$  is the defined threshold rate of fungal P supply (g P m<sup>-2</sup> day<sup>-1</sup>), below which plant C investment is limited,  $p_{fth}$  is a threshold fraction determined by fungal and plant species (Table 3).  $P_{fungi \rightarrow plant}$  is the actual mycorrhizal fungal P supply to the plant (g P m<sup>-2</sup>

320

species (Table 3).  $P_{fungi \rightarrow plant}$  is the actual mycorrhizal fungal P supply to the plant (g l day<sup>-1</sup>) (eq. 16),  $P_{ilab \rightarrow root}$  describes plant uptake by roots (g P m<sup>-2</sup> day<sup>-1</sup>) (eq. A.8). P in the fungal biomass,  $P_{fungi}$  (g P m<sup>-2</sup>), is calculated as,

$$P_{fungi} = P_{soil \to fungi} - P_{fungi \to litter} - P_{fungi \to plant}$$
(14)

Where fungal P litter production ( $P_{fungi \rightarrow litter}$ , g P m<sup>-2</sup> day<sup>-1</sup>) is estimated from a first-order rate 325 equation,

$$P_{fungi \to litter} = P_{fungi} \times p_{lrate} \times (1 - p_{fret})$$
(15)

Where  $P_{fungi}$  stands for fungal P content (g P m<sup>-2</sup>),  $p_{lrate}$  is the litterfall rate parameter (day<sup>-1</sup>) (Table 3), and  $p_{fret}$  is a parameter describing the fraction of P retained in fungal tissue during senescence (Table 3).

330 P transfer from mycorrhizal fungi to plants, P<sub>fungi→plant</sub> (g P m<sup>-2</sup> day<sup>-1</sup>), is driven by plant P demand (eq. A.9) after root uptake (eq. A.8), but regulated by P availability to fungi,

$$P_{fungi \to plant} = \begin{cases} P_{Demand} - P_{ilab \to root} & P_{Demand} - P_{ilab \to root} \leq P_{fungiavail} \\ P_{fungiavail} & P_{Demand} - P_{ilab \to root} > P_{fungiavail} \end{cases}$$
(16)

Where  $P_{fungiavail}$  is the P that can be acquired by fungi and transferred to the plant (g P m<sup>-2</sup>), calculated as,

335 
$$P_{fungiavail} = P_{fungi} - \frac{C_{fungi}}{p_{cp fungi \max}}$$
(17)

Where  $P_{fungi}$  is fungal P content (g P m-2),  $p_{cpfungimax}$  is a parameter describing the predefined maximum C/P ratio of fungal tissue (Table 3). This is based on the assumption that mycorrhizal fungi will only supply the plant with P as long as fungal C demand is fulfilled (Nehls, 2008).

340

#### 2.2.6 Phosphorus uptake by mycorrhizal fungi

The total and partial uptake of P by mycorrhizal fungi is calculated in a way that is analogous to how He et al. (2018) calculated N uptake by mycorrhizal fungi,

$$P_{soil \to fungi} = P_{ilab \to fungi} + P_{Litter \to fungi} + P_{Humus \to fungi}$$
(18)

- 345 The mycorrhiza is further distinguished into the mycelia, responsible for N and P uptake (both in inorganic forms and nutrient short-cut from organic pools), and the fungal mantle, which covers the fine-root tips (He et al., 2018).  $P_{ilab}$  uptake is first limited by the potential uptake rate  $P_{ilabpot \rightarrow fungi}$  (g P m<sup>-2</sup> day<sup>-1</sup>), which is determined by the biomass of fungal mycelia,  $P_{ilabpot \rightarrow fungi} = p_{i,rate} \times C_{fungi} \times p_{fmyc}$  (19)
- 350 Where  $P_{ilabpot \rightarrow fungi}$  describes the potential fungal  $P_i$  uptake rate (g P m<sup>-2</sup> day<sup>-1</sup>), and  $p_{i,rate}$  is a parameter which describes the potential mycorrhizal fungal uptake rate of  $P_i$  per unit  $C_{fungi}$  (g P g C<sup>-1</sup> day<sup>-1</sup>) (Table 3), and  $p_{fmyc}$  is the fraction of fungal mycelia in total fungal biomass (Table 3).

The actual fungal uptake of  $P_{ilab}$ ,  $P_{ilab \rightarrow fungi}$  (g P m<sup>-2</sup> day<sup>-1</sup>), is calculated based on the potential uptake rate (eq. 19), which is further regulated by soil  $P_{ilab}$  availability,

$$P_{ilab \to fungi} = \begin{cases} P_{ilabpot \to fungi} \times f(P_{fungidef}) & P_{ilabpot \to fungi} \le P_{ilab} \times f(P_{fungiavail}) \\ P_{ilab} \times f(P_{fungiavail}) & P_{ilabpot \to fungi} > P_{ilab} \times f(P_{fungiavail}) \end{cases}$$
(20)

Where  $f(P_{fungiavail})$  is an availability function determining the fraction of  $P_{ilab}$  that fungi can directly obtain (eq. 21), and  $f(P_{fungidef})$  is the function determining the deficiency fraction that fungi can possibly uptake, which is determined by the fungal C/P ratio (eq. 22),

$$360 f(P_{fungiavail}) = p_{iavail} \times upt_{f,enh} (21)$$

Where  $p_{iavail}$  defines the fraction of  $P_{ilab}$  that can be directly obtained by roots (Table 3, also see eq. A.8),  $upt_{f,enh}$  is an enhanced uptake coefficient that accounts for the fact that fungal mycelia have higher uptake efficiency than roots (He et al., 2018).

The function of uptake deficiency fraction,  $f(P_{fungidef})$ , scales the unfulfilled capacity of fungi for P uptake, and is calculated as

$$f(P_{fungidef}) = 1 - \frac{p_{cpfungimax}}{C_{fungi} / P_{fungi}}$$
(22)

Where  $p_{cpfungimin}$  is the defined minimum fungal C/P ratio parameter (Table 3).

In our model, we assume that  $P_i$  derived from the enzymatic hydrolysis of organic  $P_o$  is directly taken up by fungi (termed nutrient short-cut uptake in this study). Similar to

355

365

....

370  $P_{ilab \rightarrow fungi}$  (eq. 20), fungal uptake of  $P_{Litter}$  is first limited by the potential uptake rate  $P_{Litterpot \rightarrow fungi}$  (g P m<sup>-2</sup> day<sup>-1</sup>), which is determined by the biomass of fungal mycelia.

$$P_{Litterpot \to fungi} = p_{Litter,rate} \times C_{fungi} \times p_{fmyc}$$
<sup>(23)</sup>

Where  $p_{Litter,rate}$  is a parameter that describes the potential rate at which fungal mycelia acquire P from soil litter (g P g C<sup>-1</sup> day<sup>-1</sup>) (Table 3). The actual uptake from  $P_{Litter}$  to fungi,  $P_{Litter \rightarrow fungi}$  (g P m<sup>-2</sup> day<sup>-1</sup>), is calculated by,

$$P_{Litter \to fungi} = \begin{cases} P_{Litterpot \to fungi} \times f(P_{fungidef}) \times frac_{P,lit} & P_{Litterpot \to fungi} < p_{Litterf} \times P_{Litter} \\ p_{Litterf} \times P_{Litter} \times frac_{P,lit} & P_{Litterpot \to fungi} \ge p_{Litterf} \times P_{Litter} \end{cases}$$
(24)

Where  $p_{Litterf}$  is the nutrient short-cut uptake parameter that describes the uptake rate of soil litter  $P_{Litter}$  that can be hydrolyzed and directly acquired by fungi (day<sup>-1</sup>) (Table 3), *frac*<sub>P,lit</sub> is introduced to ensure that fungal nutrient short-cut uptake is less than the missing plant demand after  $P_{ilab}$  uptake, as well as to avoid uptake from only one organic pool, calculated



385

as,

375

$$P_{of,max} = P_{Litter} \times p_{Litterf} + P_{Humus} \times p_{Humusf}$$
  
$$frac_{P,lit} = \min\left\{\frac{P_{Demand} - P_{ilab \to plant}}{P_{of,max}}; \frac{P_{Litter} \times p_{Litterf}}{P_{of,max}}\right\}$$
(25)

Where  $p_{Humusf}$  is the fungal nutrient short-cut uptake parameter that describes the uptake rate of soil humus  $P_{Humus}$  that can be hydrolyzed and directly acquired by fungi (day<sup>-1</sup>). The same approach can be used to quantify fungal P uptake from the humus pool by replacing terms that include the litter P pool with the humus P pool in eqs. (23), (24) and (25).

The fungal mantle prevents contact between roots and the soil, and thereby limits the rate at which roots can directly acquire nutrients from the soil. The plant root  $P_i$  uptake response to P availability and the fungal mantle is calculated as,

$$390 f(P_{iavail}) = p_{iavail} \times e^{(-fm \times m)} (26)$$

Where  $p_{iavail}$  is a parameter that describes the maximum fraction of  $P_{ilab}$  that is available for uptake by plant roots, (eq. A.8) (i.e., not covered by the fungal mantle), *fm* is an uptake reduction parameter that describes cover by the fungal mantle, and *m* is the mycorrhization degree, see He et al. (2018).

395

#### **3** Description of the region used for simulation and model setup

#### 3.1 Description of the region

The Coup-CNP model was tested on four managed forest regions - Västerbotten 64°N, Dalarna 61°N, Jönköping 57°N, and Skåne 56°N - situated along a climatic, N and P

400

deposition, and fertility gradient across Sweden. These are the same four regions that were investigated by Svensson et al. (2008) and He et al. (2018). An overview of the climatic, geological, plant and soil characteristics of the four regions is provided in Table 1. In general, the four regions represent a North-South transect characterized by increasing mean air temperature (from 0.7 to 7.1 °C), precipitation (613-838 mm), and atmospheric N deposition

- $(1.5 12.5 \text{ kg N ha}^{-1} \text{ year}^{-1})$ . The measured annual P deposition ranges from 0.06 to 0.28 kg P 405 ha<sup>-1</sup>, with the lowest and highest deposition rates observed in the 61°N and 57°N regions, respectively. To ensure comparability, all sites selected in the four regions are characterized by podzol soil (Jahn et al., 2006) and dominated by Scots pine (Pinus sylvestris) and/or Norway spruce (Picea abies) (Table 1). Soil fertility, indicated by C-to-nutrient ratios,
- 410 exhibited an increasing trend from North to South; however, the highest soil organic C/P ratio (thus the poorest P content) was measured in the 61°N region (Table 1). Soil mineral P content varied with geology (Table 1). The aqua regia extraction method was used to determine total soil mineral P content from regional till samples collected by the Geological Survey of Sweden (SGU) (Andersson et al., 2014). Samples were taken from the C-horizon at
- 415 a depth of approximately 0.8 m, where the till is generally not disturbed by weathering. In general, Swedish till soils belong to the youngest and least weathered soils in Europe. High total mineral P contents can be found in southern (i.e. 57 and 56°N) and northern parts of the country (i.e. 64°N), which include apatite-iron ore districts (Table 1). Total mineral P content in central Sweden (e.g., 61°N) is much lower than in other parts due to the occurrence of 420 marine and postglacial clays that cover, for example, the Mälaren region.

#### 3.2 Datasets for model evaluation

Literature data concerning tree biomass, leaf nutrient content, water flow and P leaching were compiled from sites representing coniferous forests on Podzol soil within the major moisture 425 classes (mesic and moist), according to the Swedish National Forest Soil Inventory (NFSI) (Olsson et al. 2009, Stendahl et al., 2010). The corresponding forest biomass data were based on measured standing stock volumes of different age classes presented in the Swedish Forest Inventory (SFI) (SLU, 2003); for more details, see Svensson et al. (2008).

The leaf nutrient data used in the evaluation were based on measurements from forest 430 monitoring sites of the Swedish Forest Agency (Wijk, 1997; Akselsson et al. 2015) that represented the studied regions (some forest sites are also part of the ICP FOREST LEVEL II monitoring program, <u>www.icp-forests.org</u>). The data representing the northern 64°N region include two Scots pine stand sites, Gransjö (N 64°30', E17°24') and Brattfors (N64°29', E18°28'). The 61°N region was represented by two sites with Scots pine stands - Kansbo

- 435 (N61°7', E 14°21') and Furudalsbruk (N61°12', E15°11'). The 57°N region was represented by the Fagerhult (N57°30', E15°20') site, which is dominated by Norway spruce, and the Gynge Scots pine stand (N57°52', E14°44'). The data representing the 56°N region comprised a Scots pine stand in Bjärsgård (N56°10', E13°8'), a Norway spruce stand in Västra Torup (N56°8', E13°30') and a European Beech stand in Kampholma (N56°6', E13°30').
- 440 To compare the model outputs with empirical measures of P leaching, PO<sub>4</sub> and total P data in stream water were obtained from the open database of environmental monitoring data (MVM, <u>https://miljodata.slu.se/mvm/</u>). Thus, the observations of P leaching also contain P leaching from upstream sources. DOP was not measured for the regions; instead, DOP was calculated as the difference between the measured total P and PO<sub>4</sub>. This means that the
- 445 "measured DOP" may contain both our simulated DOP fractions and particular phosphorus. We used measured water outflow rates from the regional outlet from the Swedish Meteorological and Hydrological Institute (SMHI, <u>https://vattenwebb.smhi.se/station/)</u> to convert the concentrations into fluxes.

#### 450 **3.3 Model design and setup**

- The results were based on simulated forest development with daily resolution over a rotation period from a stand age of 10 years to 10 years after final harvesting. The 10 years after final harvesting, as per recommendations from Gärdenäs et al. (2003), were included to cover potential nutrient leaching during the regeneration phase. The trees across all of the investigated regions were assumed to have been planted in 1961; thus, the period from 1961 to 1970 was used as a spin-up period. The harvesting intensities and rotation lengths were specified for each region following recommendations from SLU (2012). The simulated rotational period was 120, 110, 90 and 70 years from the northern to southern regions, respectively. Two thinnings were conducted in the two northern regions while three thinnings
- 460 were conducted in the two southern regions (Table 1). Following general forest management guidelines, it was assumed that 20% of the stems are removed while 5% is transformed into litter during thinning (Swedish Forest Agency, 2005). For leaves and roots, it was assumed that 25% is transformed into litter. For all of the regions, one clearance during which 60% of the stand is removed was applied at the end of the spin-up period, i.e., when the stand is
- 465 10-years-old. During final felling, 5% of trees remain intact, and it was assumed that 90% of

the stems are harvested - with 5% becoming litter - and all of the leaves and roots become litter.

#### 3.4 Model forcing, initial and boundary conditions

- 470 Historical weather data were derived from the nearby SMHI weather station data through spatial interpolation for each region. Projections of future weather data were generated by the climate change and environmental objective (CLEO) project, using ECHAM5 projections and bias correction of regional climatic data (Personal communication with Thomas Bosshard, SMHI). Concerning P deposition, the P deposition rate from each region (Table 1)
  475 was kept constant over the simulation period, as was also the case for N deposition.
- As was performed by Svensson et al. (2008), an 11.3-meter deep soil profile containing 20 layers was simulated for all four regions. An assumed constant heat flow was used to define the lower boundary condition for heat and no water flow was assumed at the bottom soil layer. Part of the model setup and initial conditions, e.g., soil physical properties, drainage,
- 480 initial soil C content and C/N ratio, followed what was reported by Svensson et al. (2008), who also relied on NSFI data. He et al. (2018) additionally described explicit mycorrhizal fungi settings. The following section will only describe the initial conditions for the newly developed P model.
- The two vegetation layers were initialized as bare ground with small amounts of C, N and P from the seedling phase to the start of vegetation growth. The initial solid inorganic P content, soil organic matter content, and soil stoichiometry conditions are reported in Table 1. The initial soil organic P pool (Table 1) was partitioned between the soil litter (5%) and humus pools (95%), which is analogous to N partition in Svensson et al. (2008). The total amount of soil organic P decreased exponentially with depth (Fransson and Bergkvist, 2000).
- 490 Litter and humus were assumed to be distributed down to depths of 0.5 m and 1 m, respectively. The initial labile P<sub>i</sub> concentrations were set according to previous data from similar Swedish forest sites (Kronnäs et al., 2019; Fransson and Bergkvist, 2000). Soil pH was set according to the NSFI data and kept constant over the simulation period (Table 1). The initial value of soil organic P for the soil profile was estimated based on measurements of
- 495 soil organic matter N/P ratios from the same forest monitoring sites of the Swedish Forest Agency (Wijk, 1995; Akselsson et al., 2015) at which leaf nutrient content had been sampled. However, only the organic N/P ratio at the O horizon was measured at most sites (Table 1). Thus, in our calculations of the total stock of soil organic P, we assumed that the mean N/P ratio measured for the O horizon also extends to the other horizons in the default model run.

500 Model uncertainties associated with this assumption were assessed by including various soil N/P ratios (10-25) in the sensitivity analysis (e.g. Fig. 5).

#### 3.5 Sensitivity analysis

The C and N parameters for these regions have been calibrated in previous versions of
CoupModel (Svensson et al., 2008), while the parameters describing fungal processes, humus decomposition rate and short-cut N uptake rate from humus pool were calibrated by He et al. (2018) (Table 2). The surface cover parameters and litterfall rates of understory vegetation were modified from Svensson et al. (2008) to achieve more realistic understory dynamics for the analyzed regions (Table 3). Most of the default values of the newly introduced P parameters were derived from literature (Table 3). For instance, the optimal leaf C/P ratios for forest growth and the C/P ratios of individual plant components were obtained from empirical measurements from Swedish forests (e.g. Thelin et al., (1998; 2002)). The weathering and surface runoff parameters were defined according to data from laboratory experiments (Guidry and Machenzie, 2000).

- 515 We conducted a global sensitivity analysis of various parameterization schemes (n=34) for the new Coup-CNP model using a Monte-Carlo based sensitivity analysis method to assess the stability and robustness of the model with respect to its parameter values. The sampled parameters and their ranges (Table S.1), model design and global sensitivity results (Tables S.2, S.3, S.4) are reported in detail in the Supplement. Based on these simulations and
- 520 parameter sensitivity rankings, we selected the three parameters which had the strongest effect on the model outcome to serve as the basis for a new set of model runs – which underlie the sensitivity analysis results discussed in this paper. These three parameters are initial soil humus P, the short-cut N uptake rate and the short-cut P uptake rate (Tables S.2, S.3, Table 2), all of which strongly regulate soil N and P availability. The sensitivity of plant
- growth, soil C, and leaching losses in response to soil N and P availability was then assessed by varying the soil N/P ratio from 10 to 25 for the study regions (see Table 2 and Figure 5). These ranges were set according to previously published Swedish forest soil data (Lagerström et al., 2009; Giesler et al., 2002; Kronnäs et al., 2019) and additional soil humus P data from the Swedish Forestry Agency inventory (Table 1). The ranges of the short-cut uptake
  coefficients for N and P were based on regional minima and maxima for N and P short-cut

#### 4 Results

uptake rates (Table 2).

#### 4.1 Model assessment

- 535 The new Coup-CNP model was able to reproduce the observed development of forest tree biomass (SLU, 2003) over the rotation period well (Fig. 2). It should be noted that the dips in the simulated biomass are related to the timing of forestry operations in the model that are not represented in the empirical measurements. The regional biomass data show an increasing trend from North to South, which the model captured clearly (Fig. 2). However, when the
- 540 predictions were compared with observed plant biomass prior to final harvesting, the model showed a slight underestimation (12%) for the northern 61°N region and slight overestimations for the other regions (7%, 13% and 1% for the 64, 57 and 56 °N regions, respectively).

The simulated leaf C/P ratios agree fairly well with the available Swedish Forest Agency data (Wijk, 1997; Akselsson et al. 2015), despite general overestimation of 10%, 32%, 30% and 21% from North to South, respectively. The average measured leaf C/P ratio across the four regions was 396 (standard deviation 48), 398 (59), 355 (45) and 396 (72) from North to South, respectively. The model found that the 56 and 61 °N regions have higher C/P ratios than the other regions, which can also be seen from the observational data (Fig. 2). The average measured leaf C/N ratios across the four regions were 44 (4), 41 (3), 36 (5) and 31

- (7) from North to South, respectively. As such, the model provided accurate simulations of leaf C/N ratios, and captured a decreasing leaf C/N trend from North to South. The exception was a slight leaf C/N overestimation for the 57 °N region (Fig. 2). Regarding leaf N/P ratios, the average values across the regions from North to South were 9.1 (1), 9.6 (1.3), 9.9 (1.4)
- and 13.4 (3.8), respectively. As such, the Coup-CNP model was also able to accurately reproduce the measured leaf N/P ratios, revealing that leaf N/P increases as the latitude decreases (Fig. 2). In terms of the climatic variables, the radiation absorbed by the tree canopy increased from North to South, while the temperature and water limitation of Gross Primary Production (GPP) declined from North to South (Table 4). Forest growth in the 64
- <sup>560</sup> °N and 57 °N regions were primarily limited by N, while forest growth in the 61 °N and 56 °N regions was predominantly limited by P (Table 4). The limiting effect of P availability could be seen in the relatively high predicted N/P ratios, as the 56 °N region and, to a lesser extent, the 61 °N region showed high N/P ratios (Fig. 2).
- Total annual plant N and P uptake rates in the northernmost region were modeled to be 3.7 g
  565 N m<sup>-2</sup> year<sup>-1</sup> and 0.4 g P m<sup>-2</sup> year<sup>-1</sup>. The southernmost region demonstrated N and P uptake rates that were three and two times higher, respectively, than what was modeled for the northernmost region (Table 4). Total N uptake via short-cut uptake from the organic N pools

decreased from North to South (Table 4). The modeling results also indicated that short-cut uptake of P is necessary to satisfy the demands of the plant. However, the fraction of total P

- 570 uptake from the short-cut pathway was found to be neither associated with latitude nor the C/N ratio. Instead, it is regulated by soil C/P ratio and geology (Tables 1 and 5). The contribution of fungi to total N litter production decreased from North to South, but this was not the case for P, as fungi contributed a stable amount to the P litter pool across all four regions (Table 4).
- 575 The simulated annual soil C sequestration rates were 2, -2, 9, and 15 g C m<sup>-2</sup> year<sup>-1</sup> from North to South, respectively (Figs. 3a, 4a, Table 5). Thus, the soil C stock was generally in a steady-state over the forest rotation period, with slightly higher C sequestration rates predicted for the southern regions (Figs. 3a, 4a). The soil C/N ratios of all of the regions were in a steady-state over the forest rotation period. In contrast, the C/P ratios and N/P ratios

580 showed a slightly increasing trend over the rotation period, with the exception of the soil N/P ratio in the 64 °N region (Fig. 3b, c, d). The modeled P leaching generally reflected the observational data; however, the mean estimated concentrations were often lower than the measurements available for each region (MVM, https://miljodata.slu.se/mvm/, Table 6). It should be noted that the observed stream P

- 585 concentrations include P from the entire watershed, whereas our model only includes upstream P sources. The data show that P losses through leaching were small compared to the internal fluxes, i.e., they account for approximately one-third of the annual deposition input, while DOP losses were more dominant in the northern systems (Table 6). However, the simulated proportion of DOP in total losses through leaching was much lower than what had
- 590 been measured, and the decreasing trend from North to South identified in the simulations was not supported by the observational data (Table 6).

#### 4.2 Modeled forest C, N and P budgets

Regarding C assimilation, average plant growth over the rotation period in the southernmost region was predicted to be three times higher than that of the northernmost region (Fig. 4a). As most of the forest productivity was transformed into harvested products, the change in plant C was small as the simulation started when the plants were 10 years old and ended 10 years after they had been harvested. Regarding the N budget, the northernmost ecosystem showed a slight loss while the southern ecosystems showed N gains. The N sequestration for rates generally increased towards the southern latitudes (Fig. 4b). The P budget showed an opposite pattern, as the northernmost ecosystem was in balance while the other three ecosystems showed P losses, with total losses increasing from North to South (Fig. 4c).

Most of the C captured from the atmosphere was stored in the harvested plants (Fig. 4a). Our model predicted small losses of DOC through leaching, and the forest soil in all of the regions was found to be in a quasi-steady-state with generally low sequestration. An exception was the region with the lowest P-availability (61 °N), which showed soil C losses (Tables 1, 5).

Our results identified atmospheric deposition as the main N input. When accounting only for harvested N, 60%, 53%, 35% and 36% of the deposited N was removed from the 64, 61, 57

- and 56 °N regions, respectively (Fig. 4b). The N accumulated in standing plants and harvested plants accounted for 104%, 80%, 54% and 55% of the annual N deposition in the 64, 61, 57 and 56 °N regions, respectively. The model results indicate that the soils of the two northern regions will lose N while soils of the two southern regions will accumulate N. Annual average losses through leaching were predicted to increase from North to South, and
- ranged from 0.09, 0.19, 0.27, 0.47 g N m<sup>-2</sup> from North to South, which corresponds to 60%, 45%, 21%, 41% of the annual N deposition, respectively (Fig. 4b).
  The simulated annual P weathering fluxes ranged from 0.009 to 0.025 g P m<sup>-2</sup> year<sup>-1</sup>, and showed similar magnitudes as the deposition inputs (Fig. 4c). The most significant source of
- P losses over the rotation period was harvesting, which removed 89%, 255%, 108% and 167% of the deposited P from the 64, 61, 57 and 56 °N regions, respectively (Fig. 4c). When the P that accumulated in standing and harvested plants is considered together, this accounts for 85%, 147%, 90% and 114% of the total P input through deposition and weathering for the 64, 61, 57 and 56 °N regions, respectively. Thus, the simulation revealed that soils from all four studied regions are slightly losing P, with annual losses ranging from 0.01 to 0.03 g P m<sup>-</sup>
   <sup>2</sup> from North to South (Fig. 4c).
  - 4.3 Impacts of forest growth, soil C and leaching on soil N and P levels

Forest growth, measured through harvested biomass, increased as the soil N/P ratios increased from 10 to 15, but decreased once an optimum soil N/P ratio of around 15-20 was reached. This trend was noted for three studied regions (Fig. 5a), becoming less pronounced from North to South and was ultimately undetectable for the northernmost region, where GPP was strongly limited by radiation (average absorbed radiation 3.89×10<sup>6</sup> J m<sup>-2</sup> day<sup>-1</sup> at 64 °N and 6.57×10<sup>6</sup> J m<sup>-2</sup> day<sup>-1</sup> at 56 °N, Table 4). The 64 °N region had the lowest air temperature

and amount of precipitation, which also contributed to GPP limitation (temperature/water limitation of GPP 0.47/0.45 at 64 °N and 0.67/0.65 at 56 °N, Table 4). Soil C sequestration across the 56-61 °N latitudes was found to be highly sensitive to soil N/P ratios, with the model predicting that soil C sequestration would consistently decrease as the soil N/P ratio increases (Fig. 5b). In addition, total P losses through leaching generally decreased as soil N/P ratios increased; an exception was the 57 °N region, where P losses through leaching also

- 640 increased for soil N/P ratios above 15-18, a range which coincides with maximum plant harvest (Fig 5a). In contrast, total N losses through leaching were found to be positively correlated with the soil N/P ratio above 15, with this relationship more pronounced for the southern regions (Fig. 5c, d). Thus, the sensitivity analysis results indicate that strong C-N-P interactions are also prevalent in forest ecosystems (Fig. 5). Forest soils with soil N/P ratios
- 645 exceeding 15-18 were predicted to exhibit slower forest growth rates, lower soil C sequestration (potentially even losses), and high N leaching risk.

#### **5** Discussion

#### 5.1 Modeled P budgets and comparison with other modeling studies

- It is important to compare our modeled P fluxes with previously reported values. In a study that compared results from the PROFILE model with empirical data, Akselsson et al. (2008) estimated the average weathering rate in Swedish forests (down to a depth of 0.5 m) to be 0.009 g P m<sup>-2</sup> year<sup>-1</sup>, ranging from 0.001 to 0.024 g m<sup>-2</sup> year<sup>-1</sup> (5% and 95% percentiles, respectively). This can be compared with our simulated P weathering rate (down to a depth of
- 655 1 m soil depth) range of 0.009 to 0.025 g P m<sup>-2</sup> year<sup>-1</sup>. Both our estimations and those by Akselsson et al. (2008) were far lower than the 0.071 g P m<sup>-2</sup> year<sup>-1</sup> (0.5 m depth) reported by Yu et al. (2018) for a spruce forest on Podzol soil in southern Sweden. It is important to mention that Yu et al. (2018) suggested that the weathering rate they provided was an overestimation.
- The modeled total plant P uptake rates in this study ranged from 0.4 to 1 g P m<sup>-2</sup> year<sup>-1</sup> (Table 4), which is slightly higher than the  $0.5 \pm 0.4$  to 0.96 g P m<sup>-2</sup> year<sup>-1</sup> reported by Johnson et al. (2003) and Yanai (1992) for temperate forests, and the 0.5 g P m<sup>-2</sup> year<sup>-1</sup> reported for a southern Swedish forest by Yu et al. (2018). One explanation for this discrepancy could be that Coup-CNP explicitly considers mycorrhizal processes related to P uptake, e.g., the presented estimates revealed that mycorrhizal fungi accounted for more than
- half of total plant P uptake (Table 4). This highlights that mycorrhizal fungi are crucial to

plant P acquisition in forest ecosystems. The estimated P uptake by fungi was - to a large extent - proportional to the rates estimated for N (Table 4). He et al. (2018) compared explicit and implicit models and found that CoupModel v5.0 predictions of plant N uptake were

- 670 higher when mycorrhizal fungi were explicitly included in the model. Furthermore, it is important to note that previous accounts of empirical data (Johnson et al., 2003; Yanai, 1992), as well as the ForSAFE model (Yu et al., 2018), did not account for P uptake by understory vegetation. In this study, understory vegetation was estimated to contribute to c.a. one-third of total P uptake in northern regions and one-sixth of total P uptake in southern
- 675 regions (data not shown).

Akselsson et al. (2008) reported that, in Swedish forests, whole-tree harvesting causes average P removal of 0.054 g P m<sup>-2</sup> year<sup>-1</sup> with a range from 0.016 to 0.13 g P m<sup>-2</sup> year<sup>-1</sup>. This agrees well with our modeled range (0.012 to 0.038 g P m<sup>-2</sup> year<sup>-1</sup>) as well as the value reported by Yu et al. (2018), 0.037 g P m<sup>-2</sup> year<sup>-1</sup>.

- The P balances estimated for the ecosystem in this study ranged from 0 to -0.02 g P m<sup>-2</sup> year<sup>-1</sup> (with the negative value representing P losses). This agrees with what has been reported by Akselsson et al. (2008), i.e., an average P balance of -0.029, ranging from 0.008 g P m<sup>-2</sup> year<sup>-1</sup> in northern Sweden to -0.1 g P m<sup>-2</sup> year<sup>-1</sup> in southern Sweden. The modeling by Yu et al. (2018) yielded P accumulation of 0.004 g P m<sup>-2</sup> year<sup>-1</sup> over a 300-year period in south
- 685 Sweden. This predicted gain in P over the simulation period, which was very low, could have been due to relatively high P inputs via weathering. Our modeled regional P budget implies that clear-felling will result in a negative P balance for most Swedish forests even when P uptake by mycorrhizal fungi in nutrient-poor forests is accounted for. The exception to this prediction is the northernmost region.

690

#### 5.2 Implications of P availability on forest C and N dynamics

Our results demonstrate that Swedish forests are increasingly P-limited with decreasing latitude, a trend that was especially noticeable at southern latitudes (Table 4). N limitation was more severe than P limitation at the 64 and 57 °N regions. Furthermore, the canopy of

695 forests in the northernmost region intercepted far less radiation than the canopies at lower latitudes. This may partly mask the response to nutrient limitation, and cause forests in these regions to appear less sensitive to nutrient limitation. This was supported by the observed leaf N/P ratios (average values between ca. 9-14), which are recognized to reflect the state of nutrient limitation in forest trees (e.g. Jonard et al. 2015). In Swedish spruce forests, leaf N/P

signal P limitation (Rosengren-Brinck and Nihlgård, 1995; Yu et al., 2018). Linder (1995) has previously reported an optimal N/P ratio of 10 for spruce forests in northern Sweden. Similar optimal N/P ratio for pine forest was also reported (Ingestad, 1979; Tarvainen et al., 2016). Our leaf N/P ratio estimates were within these ranges, with the exception of the 705 southernmost region (Fig. 1). The ratio of total plant P uptake to total N uptake in the southernmost region was much lower than what was measured for the other regions (Table 4), which further suggests P limitation in the southernmost region. The 61 °N region, which was characterized by the lowest P inputs among the studied regions due to geology and deposition (Table 1) (Fig. 4), was also shown to be P-limited (Table 4). This low P input also 710 explains why this region showed the highest simulated fraction of short-cut P uptake from soil organic P (Table 4). Our modeling suggests that northern regions, which have traditionally been conceived as N-limited (Högberg et al., 2017), may experience P limitation or co-limitation by N and P. For instance, the 57 °N region showed N limitation as the average response of N on GPP, 0.30, is lower than the average response of P on GPP, 0.34, 715 over the rotation period (Table 4). However, our model results for the same region also yielded average values for the response of P on GPP that were lower than the average values for the response of N on GPP during initial stand development. This suggests that colimitation could still occur, e.g., at some points during the first 10 years of the simulation, and

- indicates that nutrient limitations may shift during forest development stages. For example,
  Tarvainen et al. (2016) reported a decrease in needle P following N fertilization in a Scots pine forest in northern Sweden. Several groundwater discharge areas were also shown to be P-limited (Giesler et al., 2002). Sundqvist et al. (2014) and Vincent et al. (2014) reported that alpine ecosystems in northern Sweden may also be P-limited.
- The removal of harvest residues from final fellings for use in biofuel production is common, and expected to increase in southern and central regions of Sweden (Cintas et al., 2017, Ortiz et al., 2014; Stendahl et al., 2010). Our modeling indicates that clear-cutting or final-felling will significantly impact the forest P balance and soil C sequestration (Figs. 4c, 5b). Furthermore, it is important to note that this practice was found to affect P availability more than N availability, especially in southern Sweden (Figs. 4b, 4c). Simulations with earlier
- 730 versions of CoupModel have also revealed N depletion for final-felling/clear-cutting scenarios in northern Sweden, but reported N gains for southern Sweden (He et al., 2018; Svensson et al., 2008; Gärdenäs et al., 2003).

The soil C sequestration simulated by the Coup-CNP model is generally comparable with what has been reported in previous studies (Table 5). Plants in the northern regions need to

- 735 acquire nutrients to meet demands for growth, but the Coup-CNP model showed that plants acquire a smaller fraction of total nutrients than what was previously estimated (Coup-CN model; see Table 8 in Svensson et al. (2008)). Our results further suggest that P regulates SOC, as an increasing soil N/P ratio will decrease soil C sequestration rates (Fig. 5b).
- Our global parameter sensitivity analysis showed that initial soil organic P content and the short-cut uptake coefficients for N and P have a larger influence on model outputs than parameters related to weathering rates (see the Supplement). This again confirms that the internal cycling of P is more important in regulating ecosystem C, N and P dynamics than current weathering inputs in Swedish forests, a finding that is similar to what has been reported by Yu et al. (2018). The sensitivity analysis results revealed that the optimum soil
- N/P ratio for forest production on podzol soils across the 61-56 °N regions is between 15-20 (Fig. 5). Manzoni et al. (2010) reviewed the forest litter decomposition process and found that litter C-to-nutrient ratios decreased towards a C/N ratio of 20 and C/P ratio of 350 (thus an N/P ratio of 17.5) as decomposition proceeded. A synthesis of long-term decomposition studies covering northern forests also showed that the N/P ratio of both fine litter and woody
- 750 residues converges to c.a. 20 (Laiho and Prescott, 2004). The optimum range identified by the Coup-CNP model is thus similar to these observed convergence ratios, which generally represent the shift from immobilization during the initial decomposition phase to net mobilization (Penuelas et al., 2013; Güsewell, 2004; Cleveland and Liptzin, 2007). Lagerström et al. (2009) measured soil and microbial nutrient contents in 30 diversified forest
- 755 islands in northern Sweden that vary considerably in terms of fertility. Surprisingly, they found that microbial biomass N/P ratios remained unchanged across the gradient, suggesting that nutrient availability is mainly determined by soil organic N/P ratios. The identified bell shape response of plant growth to the soil N/P ratio thus highlights the importance of nutrient stoichiometry. This implies that forests with N/P below the optimal range can benefit from N
- fertilization, which will stimulate forest growth and reduce the P leaching risk. In contrast, P fertilization in forests with N/P above the optimal range will stimulate forest growth, promote soil C sequestration and reduce N leaching (Figure 5). A synthesis of long-term water quality measurements from forest streams in the geochemical monitoring network (GEOMON) found total N fluxes to be tightly linked to DON/TP ratios (Oulehle et al., 2017). As such,
   total N leaching increased with the DON/TP ratio, a finding which agrees with the results
- obtained in this modeling study. The presented modeling predictions thus corroborate that decreased P availability can profoundly affect the N cycle and catchment retention.

770

To summarize, the presented model (CoupModel v6.0) demonstrated that integrating the P cycle into ecosystem models can significantly impact estimations of forest C and N dynamics. This is an important finding in the context of climate change and forest management, as researchers need to have tools that will reliably model the C-N-P dynamics in an ecosystem. Climate change research strives to maximize C accumulation in terrestrial ecosystems, but this may currently be limited by P availability, which will be further jeopardized by the removal of forest residues for bioenergy production. The presented results 775 show that forest growth in southern regions, which are characterized by high N deposition and already show limitation by P, will be most affected (Fig. 4c, Table 4) (Akselsson et al., 2008; Yu et al., 2018; Almeida et al., 2018).

#### **6** Conclusions

- 780 This paper describes the most recent version (6.0) of CoupModel, which explicitly considers the phosphorus cycle and mycorrhizal interactions. The simulations of the C, N, and P budgets for four forest regions were complete and accurate based on comparisons using empirical forest biomass, leaf nutrient ratio, and P leaching data. The development and evaluation of this new model demonstrates that P availability needs to be considered when 785 studying how climate change will influence C turnover and ecosystem responses. Simulations
- that do not factor in P availability may overlook important feedback mechanisms and overestimate the C sink. Thus, the detailed description of all the Coup-CNP components and their interactions - including the water, heat, C, N, and P cycles - are highly relevant to future studies.
- 790 Our model results showed that N was the most limiting nutrient for the 64 °N and 57 °N regions, while P was the most limiting nutrient for the 61 °N and 56 °N regions (Table 4). The N limitation at the 64 and 57 °N regions was more severe than the P limitation at the 61 °N and 56 °N regions. Furthermore, the northernmost region was characterized by lower temperature, precipitation, and radiation intercepted by the canopy relative to other regions,
- 795 all of which may have skewed the estimated sensitivity to nutrient limitation. During the simulated rotation period, southern forests showed P losses, mainly through harvesting and changes in soil storage, while northern forests were close to a steady-state in P availability. Mycorrhizal fungi accounted for half of total plant P uptake in all of the regions, which highlights the crucial role of the mycorrhiza in Swedish forests. A sensitivity analysis 800 determined that a soil N/P ratio of 15 to 20 is optimal for forest growth. Furthermore, soil

N/P ratios above 15-20 decreased soil C sequestration and total P leaching, and significantly increased N leaching. The largest P outflow over the rotation period was found to be removal via final-felling.

We conclude that the potential P-limitation of terrestrial ecosystems highlights the need for 805 biogeochemical models to comprehensively consider the P cycle. The inclusion of the P cycle will enable a model to account for various feedback mechanisms that have a significant impact on C sequestration and N leaching under climate change and/or elevated N deposition.

#### 7 Code and data availability

- 810 The model and extensive documentation, including tutorial exercises, are freely available from the CoupModel home page: http://www.coupmodel.com/ (CoupModel, 2019). CoupModel is written in the C++ programming language and runs with a GUI under the Windows systems, but can also be run on other platforms without GUI. Version 6.0, from 03 July 2019, was used for the presented simulations. This version is archived on Zenodo
- 815 (https://zenodo.org/record/3547628#.Xn3Bc0F7IEZ), as are the simulation files including the model and calibration set-up, parameterization settings, and corresponding input and validation files.

## Appendix A: Equations and parameterization regarding the phosphorus processes that are analogous to nitrogen processes

The following section provides the equations for P processes that are analogous to those describing N processes, as well as discusses parameterization aspects. The inclusion of the N cycle in CoupModel was previously described by Gärdenäs et al. (2003), Jansson and Karlberg (2011), and He et al. (2018).

825

#### A1 Deposition and fertilization

830

835

Atmospheric deposition,  $P_{dep \rightarrow ilab}$  is treated as a model input using the parameter  $p_{dep}$ . In contrast to N deposition, only dry P deposition is considered since wet deposition is generally neglectable. Fertilization  $P_{fert \rightarrow ilab}$  is also treated as a model input and calculated as,

$$P_{fert \to ilab} = p_{kfert} P_{fert} \tag{A.1}$$

Where  $P_{fert \rightarrow ilab}$  is the rate of fertilizer P addition (g P m<sup>-2</sup> day<sup>-1</sup>) and  $p_{kfert}$  is the specific dissolution rate of commercial fertilizer (day<sup>-1</sup>). The value of  $p_{kfert}$  depends on fertilizer type and moisture conditions, e.g., in our model, a value of 0.15 corresponds to a half-time of 5 days, and that 90% of the fertilizer is dissolved into the  $P_{ilab}$  pool within 15 days. If manure

fertilizer is used, the organic  $P_o$  in the manure is added into a separate organic pool  $P_{ofae}$ , termed faeces. Fecal processes are similar to those of soil litter, described below.

According to a global compilation of published data, the average annual global P deposition 840 is 0.027 g P m<sup>-2</sup> year<sup>-1</sup> (0.033 for Europe), which equals to 0.000074 g P m<sup>-2</sup> day<sup>-1</sup> (Tipping et al., 2014; Schlesinger, 1997).

#### A2 Mineralization-Immobilization & decomposition

The  $P_{ilab}$  pool is also controlled by biological demand and turnover (Olander and Vitousek, 2005). The P flux of biological mineralization-immobilization is calculated precisely as for N, in that C fluxes from litter (or faeces) to humus - or from humus to atmosphere - are driven by the microbial need for energy. The non-symbiotic microbes are implicitly simulated using a fixed microbe C/P ratio parameter. The C/P ratio for microbes ( $cp_m$ ) can vary widely, ranging from approximately 25–400 (see review by Manzoni et al. 2010).

$$C_{DecomL} = k_{l} \times f(T) \times f(\theta) \times C_{Litter}$$

$$P_{Litter \to ilab} = C_{DecomL} \left( \frac{1}{C_{litter} / P_{Litter}} - \frac{f_{e,l}}{cp_{m}} \right)$$

$$P_{Litter \to Humus} = \frac{C_{Litter \to humus}}{cp_{m}}$$
(A.2)

Where k<sub>l</sub> is the decomposition coefficient for soil litter (day<sup>-1</sup>), C<sub>litter</sub> is the size of the litter pool (g C m<sup>-2</sup>), while f(T) and f(θ) are common temperature and water content response functions for decomposition; for more details see Jansson and Karlberg (2011). Humus decomposition is calculated by changing the pool size and decomposition coefficient in the previous equation into terms that describe humus. P<sub>Litter→ilab</sub> is the mineralization flux from the soil litter pool to the P<sub>ilab</sub> pool (g P m<sup>-2</sup> day<sup>-1</sup>). P<sub>Litter→Humus</sub> is the humufication flux rate. C<sub>DecomL</sub> is the C decomposition flux of soil litter (g C m<sup>-2</sup> day<sup>-1</sup>), whereas C<sub>litter</sub>/P<sub>litter</sub> and cpm represent the C-to-P ratio in the litter pool and microbes, respectively. f<sub>e,l</sub> is a microbial efficiency parameter which represents the fraction of mineralized C that remains in the soil.
Corresponding fluxes are calculated by changing the efficiency parameter to f<sub>e,f</sub> or f<sub>e,h</sub>, along with changing the litter C/P ratio to a fecal C/P ratio or humus C/P ratio, to provide the corresponding flows from the faecal pool, P<sub>ofae→ilab</sub>, or the humus pool, P<sub>Humus→ilab</sub>, respectively. A negative value means that net immobilization takes place.

Total biological mineralization is calculated as,

865 
$$P_{biomin} = P_{Litter \to ilab} + P_{Humus \to ilab} + P_{ofae \to ilab}$$
(A.3)

The biochemical mineralization process includes the release of root exudates, e.g., efflux of protons and organic anions, phosphatase and cellulolytic enzymes required for the hydrolysis or mineralization of  $P_o$  (Richardson and Simpson, 2011; Bünemann, 2015; Hinsinger, 2001). This additional mineralization process is driven by plant demand for P (Richardson et al., 2009). Bünemann (2008) reviewed the existing enzyme addition experiments and showed, for example, that the phosphatase enzyme has low substrate specificity and that up to 60% of total organic  $P_o$  in soil can be hydrolyzed and mineralized. We, therefore, assume that biochemical mineralization can occur from both the soil litter and humus pools. The flux rate is calculated as a first-order function regulated by pool size and uptake rate. Furthermore, it is

875 assumed that the flux rate will not exceed the remaining plant demand after root  $P_i$  uptake (eq. A.8). The following equation (A. 4) is used when symbiotic microbes are implicitly simulated.

$$P_{Litter \rightarrow plant} = frac_{P,litter} \times o_{uptPlitter} \times P_{Litter}$$

$$P_{Humus \rightarrow plant} = frac_{P,humus} \times o_{uptPhumus} \times P_{Humus}$$

$$P_{bioche,max} = P_{Litter} \times o_{uptPlitter} + P_{Humus} \times o_{uptPhumus}$$

$$frac_{P,litter} = \min\left\{\frac{P_{Demand} - P_{ilab \rightarrow plant}}{P_{bioche,max}}; \frac{P_{Litter} \times o_{uptPlitter}}{P_{bioche,max}}\right\}$$

$$frac_{P,humus} = \min\left\{\frac{P_{Demand} - P_{ilab \rightarrow plant}}{P_{bioche,max}}; \frac{P_{Humus} \times o_{uptPhumus}}{P_{bioche,max}}\right\}$$

880

885

Where  $P_{Litter \rightarrow plant}$  and  $P_{Humus \rightarrow plant}$  represent the biochemical mineralization fluxes from the litter and humus pools (g P m<sup>-2</sup> day<sup>-1</sup>), respectively, assuming immediate uptake by the plant roots after mineralization.  $o_{uptPlitter}$  and  $o_{uptPhumus}$  are coefficient parameters that define the maximum plant uptake rates from the soil litter and humus pools, respectively.  $P_{Litter}$  and  $P_{Humus}$  are the P pool sizes (g P m<sup>-2</sup>), frac<sub>P,litter</sub> and frac<sub>P,humus</sub> are introduced to ensure that biochemical mineralization remains less than the missing plant demand after  $P_{ilab}$  uptake, as well as to ensure proportional uptake from  $P_{Litter}$  and  $P_{Humus}$ . In this modeling framework, the inorganic  $P_i$  released by enzymatic activity is acquired directly by plants rather than entering the  $P_i$  pool.

Total biochemical mineralization is calculated as,

$$P_{biochem} = P_{Litter \to plant} + P_{Humus \to plant}$$
(A.5)

890 The total mineralization-immobilization flux is calculated as,

$$P_{totmin} = P_{biochem} + P_{biomin} \tag{A.6}$$

As is the case with DOC/DON, in Coup-CNP, organic P dissolution is described as a microbial decomposition process. The redistribution is done following that of water flow, as the DOM is assumed to have full mobility with water. The formation of DOM is from litter and humus (equ A.7). The dissolved organic matter can be fixed by humus via adsorption and precipitation, among other processes. A fixation coefficient,  $d_{DOD}$ , which varies between layers, was introduced (Kalbitz et al., 2000; Kaiser and Kalbitz, 2012). Parameterization from Svensson et al. (2008) were used in this study. The equation for DOP is similar to that for DOC, and is calculated as,

900 
$$P_{Litter \to DOP} = d_{DO,l} \times f(T) \times f(\theta) \times P_{Litter}$$

$$P_{Humus \to DOP} = f(T) \times f(\theta) \times (d_{DO,h} \times P_{Humus} - d_{DOD}(z) \times P_{DOP})$$
(A.7)

Where  $d_{DO,l}$  and  $d_{DO,h}$  are the dissolution rate coefficients (day<sup>-1</sup>) for the litter and humus, respectively, f(T) and  $f(\theta)$  are common response functions for soil temperature and water content, and identical to those used for the decomposition process (eq. A.2).

#### 905 A3 Plant growth and P uptake

Plants can acquire  $P_i$  through both the roots and mycorrhizal fungi; for this reason, both of these processes were simulated. We assume that  $P_i$  uptake by the roots is driven by net photosynthesis and determined by plant demand, yet constrained by the  $P_{ilab}$  pool size.

$$P_{ilab \to root} = min \ (p_{iavail} \times P_{ilab}; P_{demand}) \tag{A.8}$$

910 Where P<sub>demand</sub> is the plant P demand, based on the C/P ratios of various plant compartments (*iplant* includes leaf, stem, fine roots and coarse roots),

$$P_{demand} = \sum_{iplant} \frac{C_{a \to iplant} - C_{iplant \to a}}{cp_{iplant \min}}$$
(A.9)

Where  $C_{a \rightarrow iplant}$  is the photosynthetic assimilation for each compartment  $_{iplant}$  (g C m<sup>-2</sup> day<sup>-1</sup>),  $C_{iplant \rightarrow a}$  is the respiration of each compartment, and  $c_{piplantmin}$  is the defined minimum C/P

- 915 ratio for each plant compartment. Empirical measurements show that the C/P ratio of leaves generally varies between 200-600, while the stem and roots require C/P ratios between 1000-3000 and 500-1500, respectively (Bell et al., 2014; Tang et al., 2018). The model provides flexible stoichiometry because the C/P ratios for distinct compartments are calculated for each time step.
- 920 In addition, increasing soil P abundance, particularly when P fertilizer is added, is known to decrease belowground C allocation (Ericsson, 1995). We assume that an increasing C/P ratio (i.e. decreasing P content) in the leaf,  $C/P_l$ , will increase belowground allocation (e.g., *frac(root)*).

$$\begin{aligned} & frac_{a \to root}(C / P_l) = r_{cpc1} + r_{cpc2} \times C / P_l \\ & frac(root) = frac_{a \to root}(C / P_l) \times frac_{a \to root}(C / N_l) \\ & C_{a \to root} = C_{a \to plant} \times frac(root) \end{aligned}$$
(A.10)

925 Where  $r_{cpc1}$  and  $r_{cpc2}$  are the plant allocation pattern parameters, determined by plant species and a similar equation as what was used to calculate  $frac_{a\to root}(C/N_l)$  (He et al. 2018). CoupModel can additionally account for the effects of water stress on plant allocation. In this study, C allocation to roots is assumed to be constrained by both N and P contents in the leaves, i.e.,  $frac_{a\to root}(C/N_l)$  and  $frac_{a\to root}(C/P_l)$ .

930

#### A4 Plant litterfall

Plant litterfall P fluxes are proportional to the corresponding C fluxes, and determined by the C/P ratio of each compartment *iplant* (*iplant*=leaf, steam, grain, fine roots, and coarse roots), calculated as,

935 
$$P_{plant \to soil} = \sum_{iplant} \frac{c_{liplant} \times C_{iplant} \times (1 - m_{retain})}{C / P_{iplant}}$$
(A.11)

Where  $c_{liplant}$  is the litterfall rate (day<sup>-1</sup>) for plant compartment  $_{iplant}$ ,  $C_{iplant}$  is the C stock in that compartment (g C m<sup>-2</sup>), and  $c_{iplantret}$  is a parameter defined as the fraction of C that was retained before litterfall. Total litterfall also includes inputs from mycorrhizal fungi. The litterfall flux is directly added to the surface soil litter pool, or to the layer in which it formed when it was produced by roots and fungi. The average C/P ratio of fresh litter varies widely, e.g., between 100 - 4100 (Manzoni et al. 2010). The retention of nutrients prior to leaf senescence is one of the main factors that affect the C/P ratio of fresh litter. During litterfall seasons, plants can reallocate P and N from leaves to an internal, mobile storage form to prepare for rapid growth in the spring. This mechanism is known to increase the efficient use of nutrients (e.g. Aerts, 1996) (also see  $m_{retain}$  in Table S.1 in the Supplement).

945

940

#### A5 Leaching and surface runoff

The losses of soluble  $P_{isol,loss}$  (g P m<sup>-2</sup> day<sup>-1</sup>) are modeled through the transport of water,

$$P_{isol,loss} = \sum_{jlayer} P_{isoldrainage,j} + P_{isolpercolation}$$

$$P_{isoldrainage,jlayer} = \frac{P_{isol,j}}{\theta_{,j} \times \Delta z_{,j}} \times q_{drainage,j}$$

$$P_{isolpercolation} = \frac{P_{isol}}{\theta \times \Delta z} \times q_{percolation}$$
(A.12)

950 Where  $q_{drainage}$  is the water flow (mm day<sup>-1</sup>) due to drainage, and  $q_{percolation}$  is the deep percolation flow (mm day<sup>-1</sup>),  $\theta_{ij}$  is the water content (volume %) of soil layer *j*, and  $\Delta z_{ij}$  is the layer thickness (m) of soil layer *j*. The vertical  $P_i$  flow between layers is calculated through a similar equation.

DOP losses from the system is calculated as,

955 
$$P_{DOP,loss} = \frac{P_{DOP}}{\theta \times \Delta z} \times \left( \sum_{j \mid ayer} q_{drainage,j} + q_{percolation} \right)$$
(A.13)

In addition, we also accounted for particulate phosphorus (*PP*) losses, e.g., due to soil erosion, subsidence and lateral losses of secondary minerals and occluded P due to surface runoff. We assume the *PP* loss is simply proportional to the water flow. When surface runoff

occurs, for example, during snow melting, the loss is assumed to occur only for the first soil

960 layer (soil surface),

970

$$P_{solid,loss} = q_{surfacerunoff} \times k_{scale}$$

$$k_{scale} = \min(1, \frac{q_{surfacerunoff}}{q_{thr}}) \times (P_{\Delta,i} - P_{base,i}) + P_{base,i}$$
(A.14)

Where  $q_{surfacerunoff}$  is the surface runoff flow (mm day<sup>-1</sup>). An empirical scale factor  $k_{scale}$  is introduced to account for the concentration of erodible  $P_{solid}$  over the flow rate of surface runoff.  $P_{\Delta}$  (mg P l<sup>-1</sup>),  $P_{base}$  (mg P l<sup>-1</sup>) and  $q_{thr}$  (mm day<sup>-1</sup>) are empirical coefficients.

965 Therefore, the total P losses are calculated as,

$$P_{totloss} = P_{isol,loss} + P_{DOP,loss} + P_{solid,loss}$$
(A.15)

#### A6 P removal during plant harvest

The removal of P during harvesting was calculated in a similar way as C losses through harvesting, and depends on the C/P ratio of the specific plant compartment.

# Appendix B: Simulated annual mean P, N and C budgets, generated by varying three regional key parameters, including soil N/P ratio and the short-cut uptake rates for N and P

975	Table B1 Simulated annual P budget, including the associated uncertainty range (mean $\pm$ SD,
	$g P m^{-2} year^{-1}$ )

P budget	64°N	61°N	57°N	56°N
Weathering	0.014 (0.0002)	0.0094 (0.0002)	0.024 (0.0002)	0.025 (0.001)
Deposition	0.013	0.0065	0.028	0.023
Leaching	0.0025 (0.0003)	0.0015 (0.0003)	0.004 (0.0004)	0.006 (0.0009)
Harvest export	0.01 (0.002)	0.018 (0.003)	0.03 (0.004)	0.045 (0.01)
Change in plant	0.0125 (0.003)	0.007 (0.002)	0.018 (0.006)	0.017 (0.006)
Change in soil	-0.012 (0.005)	-0.02 (0.002)	-0.024 (0.005)	-0.045 (0.01)
Change in ecosystem	0.0005 (0.0004)	-0.013 (0.003)	-0.006 (0.003)	-0.028 (0.007)

N budget	64°N	61°N	57°N	56°N
Deposition	0.15	0.35	0.78	1.26
Leaching	0.10 (0.004)	0.15 (0.02)	0.16 (0.05)	0.45 (0.15)
Harvest export	0.08 (0.01)	0.19 (0.02)	0.27 (0.04)	0.5 (0.19)
Change in plant	0.09 (0.009)	0.10 (0.008)	0.15 (0.03)	0.17 (0.01)
Change in soil	-0.12 (0.03)	-0.09 (0.02)	0.20 (0.05)	0.14 (0.03)
Change in ecosystem	-0.03 (0.005)	0.01 (0.008)	0.35 (0.03)	0.31 (0.09)

Table B2 Simulated annual N budget, including the associated uncertainty range (mean $\pm$  SD, g N m<sup>-2</sup> year<sup>-1</sup>)

980

Table B3 Simulated annual C budget, including the associated uncertainty range (mean $\pm$  SD, g C m<sup>-2</sup> year<sup>-1</sup>)

C budget	64°N	61°N	57°N	56°N
Net ecosystem productivity	63 (7)	90 (10)	170 (19)	237 (21)
Leaching	0.8 (0.07)	0.6 (0.06)	0.4 (0.1)	0.3 (0.05)
Harvest export	50 (6)	81 (9)	145 (17)	201 (19)
Change in plant	10 (2)	12 (1)	18 (2)	20.7 (3)
Change in soil	2 (1)	-3.6 (6)	6.5 (4)	15 (7)
Change in ecosystem	12.2 (1)	8.4 (1)	24.5 (3)	35.7 (5)

Appendix C: Sensitivity of annual harvested biomass to changes in the short-cut uptake rates for N and P



990 Fig. C1 Simulated annual mean harvested biomass amounts for the four studied regions at different short-cut uptake rates for N and P. The bar indicates the standard deviation for each data point (Table 2)

Author contributions. AG formulated the project, HH conducted the literature review and developed the phosphorus model concepts with AG and PEJ, HH and PEJ implemented the

995 phosphorus cycle code into CoupModel. HH performed the simulations and analysis. HH drafted the manuscript, to which AG and PEJ contributed.

Competing interests. The authors declare that they have no conflicts of interest.

- 1000 Acknowledgements. The authors express their gratitude to all persons who provided data and background information. Gunilla Pihl-Karlsson and Per-Erik Karlsson at IVL Göteborg provided regional P deposition data for the studied regions, whereas the Swedish Forest Agency (Olle Kellner) provided data on needle P and N contents, soil P, and regional P deposition across the studied regions. Magnus Svensson provided the regional simulation files with CoupModel v4.0, while Thomas Bosshard at SMHI provided the regional climate
- files. The Department of Aquatic Sciences and Assessment (SLU) contributed P concentrations from open water data, while the Swedish Geological Survey (SGU) provided regional mineral P content data. The Swedish Forest Soil Inventory provided soil C/N data. Cecilia Akeselson collected data for the SWETHRO project.
- 1010 The Department of Biological and Environmental Sciences, University of Gothenburg and the strategic research area, along with ModElling the Regional and Global Earth system (MERGE), funded the study. MERGE provided SP project funding for "incorporating phosphorus cycle into ecosystem models". This research was also funded by the Formasbacked strong research environment IMPRESS. Further, we acknowledge the comments from
- 1015 participants of the N-P interaction workshops for the new model concepts, which were funded by MERGE and BECC (Biodiversity and Ecosystem services in a Changing Climate).



1020 Fig. 1 Conceptual figure of the simplified Coup-CNP and its relationship to the N cycle. The green pools represent plant-symbiotic microbes (e.g., mycorrhiza fungi), while brown pools represent soil organic matter, grayish-yellow pools represent water solutions and blue pools represent soil inorganic P. Within the pools, Croot stands for coarse roots and Froot stands for fine roots.

Regional characteristics	Västerbotten	Dalarna	Jönköping	Skåne
Latitude	64°N	61°N	57°N	56°N
Mean annual air temperature (°C) <sup>a</sup>	0.7	3.3	5.2	7.1
Mean annual precipitation (mm) <sup>a</sup>	613	630	712	838
Annual N deposition (kg N ha <sup>-1</sup> ) <sup>b</sup>	1.5	3.5	7.5	12.5
Annual P deposition (kg P ha <sup>-1</sup> ) <sup>b</sup>	0.13	0.06	0.28	0.23
Studied soil type	Podzol	Podzol	Podzol	Podzol
Quaternary deposit, SGU <sup>c</sup>	Glacial till	Glacial till	Glacial till	Glacial till
Bedrock geology, SGU <sup>c</sup>	Gneiss	Sandstone, Rhyolite	Gneiss	Gneiss
Solid inorganic P content of till (mg kg <sup>-1</sup> ) <sup>d</sup>	881	428	859	773
Major tree species: Pine/Spruce/broadleaved trees (%) <sup>f</sup>	45/37/16	49/40/9	31/54/13	12/46/41
Rotation period (years)	120	110	90	70
Years in which thinning is performed $(1^{\text{st}}/2^{\text{nd}}/3^{\text{rd}} \text{ thinning})^{\text{h}}$	50/100	40/90	25/40/70	25/40/55
Measured plant biomass for the 100- year age class $(g C m^{-2})^{f}$	5371	7815	10443	11501
Soil organic matter C/N (-) <sup>e</sup>	31.5	29.1	27.2	19.8
C/N humus <sup>b</sup>	43	40	31	25
Soil organic matter C/P (-) <sup>b</sup>	494	633	425	425
C/P humus <sup>b</sup>	325	400	410	550
Soil organic matter N/P (-)	15.7	21.8	15.6	21.5
Initial Soil C (g C m <sup>-2</sup> ) <sup>f</sup>	7006	8567	9995	10666
Litter C (g C m <sup>-2</sup> ) <sup>f</sup>	350	428	500	533
Humus C (g C m <sup>-2</sup> ) <sup>f</sup>	6655	8139	9495	10133
Initial Soil N (g N m <sup>-2</sup> )	223	295	367	539
Litter N (g N m <sup>-2</sup> ) <sup>f</sup>	11	15	18	27
Humus N (g N m <sup>-2</sup> ) <sup>f</sup>	212	280	349	512
Initial Soil P (g P m <sup>-2</sup> )	14.2	13.5	23.5	25.1
Litter P (g P m <sup>-2</sup> ) <sup>g</sup>	0.7	0.7	1.2	1.3
Humus P (g P m <sup>-2</sup> ) <sup>g</sup>	13.5	12.8	22.3	23.8
Soil pH <sup>e</sup>	5.1	5.1	5.1	4.9

Table 1 Overview of climatic, geological, plant and soil characteristics of the four forest regions

<sup>a</sup> 30-year (1961 to 1991) annual average from regional SMHI stations

<sup>b</sup> N and P deposition data were obtained from the SWETHTRO project, while the soil organic
 C/P ratio was estimated with available Swedish Forest Agency data, data of C/N and C/P

humus was from the additional survey of Swedish Forest Agency.

<sup>c</sup> Geological Survey of Sweden (SGU), <u>https://apps.sgu.se/kartvisare/</u>

<sup>d</sup> according to Geochemical Atlas of Sweden. 2014, measured till samples from the C-

1035 horizon, c.a. 0.8 m below the soil surface

<sup>e</sup> calculated based on Swedish Forest Soil Inventory data (SFSI. <u>https://www.slu.se/en/Collaborative-Centres-and-Projects/Swedish-Forest-Soil-Inventory/</u>)

<sup>f</sup> Svensson et al. (2008)

<sup>g</sup> assumption that 5% of the total organic pool is litter and 95% is humus, as reported for N in Svensson et al. (2008)

<sup>h</sup> Skogsdata (2012)

Region	Humus	Short-cut P uptake rate from	Short-cut N uptake rate
	decomposition rate,	humus pool (day <sup>-1</sup> ) <sup>b</sup>	from humus pool (day <sup>-1</sup> ) <sup>a</sup>
	$k_h (\mathrm{day}^{-1})^{\mathrm{a}}$		
Västerbotten 64°N	0.00048	1.5×10 <sup>-5</sup>	1.5×10 <sup>-5</sup>
Dalarnas 61°N	0.00042	2.75×10 <sup>-5</sup>	1.2×10 <sup>-5</sup>
Jönköpings 57°N	0.0004	1.0×10 <sup>-5</sup>	1.0×10 <sup>-5</sup>
Skåne 56°N	0.00038	1.5×10 <sup>-5</sup>	0.5×10 <sup>-5</sup>

Table 2 Parameters with specific values for different regions

<sup>a</sup>: from He et al. (2018)

<sup>&</sup>lt;sup>b</sup>: a high short-cut P uptake rate was assumed for regions with high soil organic matter C/P ratios

Table 3 Parameters used for the default model run for the P processes with common values across all four studied regions. Note that the same parameter values were applied for tree and understory layers unless otherwise specified.

Symbol	Parameter	Equation	Value	Unit	Reference
$k_w$	Integrated weathering rate	(1)	8×10 <sup>-7</sup>	day-1	Guidry and
n <sub>H</sub>	Weathering pH response coefficient	(4)	0.27	-	Machenzie, (2000);
pH <sub>opt</sub>	Weathering pH response base coefficient	(4)	7	-	Sverdrup and Warfvinge, (1993)
p <sub>max,ads</sub>	Langmuir max sorption capacity	(5)	0.0002	g P g soil <sup>-1</sup>	Adjusted from
C50,ads	Langmuir half saturation coefficient	(5)	5×10 <sup>-5</sup>	g P m <sup>-2</sup>	Wang et al. (2007)
$p_{cp,opt}$	C/P optimal (leaf)	(8)	250	gC gP <sup>-1</sup>	Thelin et al. (1998; 2002)
$p_{cp,th}$	C/P threshold (leaf)	(8)	600	gC gP <sup>-1</sup>	
$p_{avail}$	Coefficient describing reduced C allocation under P availability	(9)	0.0009	-	Assumed
$p_{fopt}$	The optimum ratio between C allocation between fungi and root	(11)	0.22	-	He et al. (2018); Orwin et al.
k <sub>rm</sub>	Fungal respiration coefficient		0.01	day <sup>-1</sup>	(2011)
p <sub>lrate</sub>	Fungal litterfall rate	(15)	0.0045	day <sup>-1</sup>	
n <sub>avail</sub>	Coefficient describing reduced C allocation under N availability		0.00039	-	
Pi,rate	Potential unit fungal mycelia uptake rate for PO <sub>4</sub>	(19)	0.0001	g P g C <sup>-1</sup> m <sup>-2</sup> day <sup>-1</sup>	Smith and Read (2008)
n <sub>NH4rate</sub> / n <sub>NO3rate</sub>	Potential unit fungal mycelia uptake rate for NH <sub>4</sub> /NO <sub>3</sub>		0.0004	$g N g C^{-1} m^{-2}$ $day^{-1}$	He et al. (2018)
N <sub>Litter,rate</sub> / N <sub>Humus,rate</sub>	Potential unit fungal mycelia uptake rate for organic N		0.00002	$g N g C^{-1} m^{-2}$ day <sup>-1</sup>	
$p_{cpfungimax}$	Fungal maximum C/P	(17)	200	gC gP <sup>-1</sup>	Wallander et al.
<b>P</b> iavail	Maximum PO4 uptake fraction for roots	(21)	0.008	-	(2003); Zhang and Elser,
p <sub>cpfungimin</sub>	Fungal minimum C/P	(22)	100	-	(2017)
PLitter,rate / PHumus,rate	Potential unit fungal mycelia uptake rate for organic P	(23)	0.00002	$\frac{g P g C^{-1} m^{-2}}{day^{-1}}$	Assumed to be the same as N
Soil organ	nic P processes	•	•	• •	•
cp <sub>m</sub>	C/P of non-symbiotic microbes	(A.3)	350	gC gP <sup>-1</sup>	Manzoni et al. (2010)
Uptake de	emand of P				
CPleaf, min	Minimum C/P (leaf)	(A.9)	220	-	Bell et al. (2014): Tang et

					al. (2018)
<b>cp</b> <sub>stem, min</sub> / <b>cp</b> croot min	Minimum C/P for stem and coarse roots	(A.9)	4000/ 800	-	
cp <sub>root, min</sub>	Minimum C/P ratio (fine roots)	(A.9)	400	-	
Plant Litte	erfall processes		•		
	Leaf litterfall rate for understory		0.0015	day <sup>-1</sup>	Calibrated
Plant surf	face cover		•	•	
	Maximum canopy cover, forest		0.8	$m^2 m^{-2}$	Assumed
	Maximum canopy cover, understory		1	$m^2 m^{-2}$	Assumed
Erosion	· ·		•		
pbase	P concentration scaling coefficient for surface erosion 1	(A.14)	2.7×10 <sup>-6</sup>	mg l <sup>-1</sup>	Assumed
$p_{\Delta}$	P concentration scaling coefficient for surface erosion 2	(A.14)	7×10 <sup>-6</sup>	mg l <sup>-1</sup>	
$q_{\it thr}$	Critical surface flow rate for erosion	(A.14)	10	mm day <sup>-1</sup>	



Fig. 2 Simulated (lines) and measured (symbols) plant biomass and leaf C/P, C/N and N/P ratios over the rotation period across the four regions. The x-axis denotes the stand age in years. The right axis of charts showing leaf C/P and C/N ratios shows the minimum (f(nutrient)=0) and optimum (f(nutrient)=1) values in terms of nutrient response to gross primary production, respectively. Biomass data and leaf nutrient data were from SFI (SLU, 2003) and the Swedish Forest Agency.

Table 4 Summary of the plant-fungal internal C, N and P variables (shown as average values
over the rotations period) of the simulated forest ecosystems. Bold values indicate a limiting
factor for GPP, according to Liebig's law of the minimum. The scale describing the response
on GPP includes temperature, water, N and P ranges from 0 (no assimilation) to 1 (optimal
growth conditions).

Variable	Unit	64°N	61 °N	57 °N	56 °N
Net primary production, tree layer	g C m <sup>-2</sup> year <sup>-1</sup>	205	302	486	600
Radiation adsorbed, tree layer	$\times 10^{6} \text{ J m}^{-2} \text{ day}^{-1}$	3.89	5.35	6.50	6.57
Temperature response on GPP, tree layer	-	0.47	0.52	0.63	0.67
Water response on GPP, tree layer	-	0.45	0.50	0.63	0.65
N response on GPP, tree layer	-	0.22	0.45	0.30	0.80
P response on GPP, tree layer	-	0.56	0.23	0.34	0.33
Total plant uptake, N	g N m <sup>-2</sup> year <sup>-1</sup>	3.67	5.76	9.00	13.8
Total plant uptake, P	g P m <sup>-2</sup> year <sup>-1</sup>	0.42	0.49	0.87	1.08
Short-cut N uptake fraction (of total)	-	0.34	0.21	0.17	0.05
Short-cut P uptake fraction (of total)	-	0.14	0.23	0.10	0.14
Fungal N uptake fraction (of total)	-	0.68	0.69	0.66	0.65
Fungal P uptake fraction (of total)	-	0.56	0.57	0.56	0.56
Fungal N transfer to plant (of total)	-	0.31	0.31	0.33	0.34
Fungal P transfer to plant (of total)	-	0.44	0.43	0.43	0.43
Total plant litter, N	g N m <sup>-2</sup> year <sup>-1</sup>	3.50	5.47	8.61	13.2
Total plant litter, P	g P m <sup>-2</sup> year <sup>-1</sup>	0.40	0.47	0.82	1.02
Fungi N litter (of total litter)	-	0.38	0.40	0.35	0.33
Fungi P litter (of total litter)	-	0.13	0.15	0.14	0.13

1065 Table 5 Simulated annual average soil C changes (g C m<sup>-2</sup> year<sup>-1</sup>, positive mean sequestration, negative mean losses), including comparison with previously reported results. Values in parentheses indicate uncertainties due to certain model parameters.

Study for comparison	Approach	64°N	61 °N	57 °N	56 °N
Svensson et al. (2008)	Coup-CN implicit mycorrhizal interactions	-5	-2	9	23
$\mathbf{H}_{2} \neq \mathbf{a}^{1}  (2018)$	Coup-CN implicit mycorrhizal interactions	-6 (10)	-5 (11)	3 (13)	13 (13)
ne et al. (2018)	Coup-CN explicit mycorrhizal interactions	-8 (11)	-9 (12)	-5 (15)	-1 (19)
This study	Coup-CNP explicit mycorrhizal interactions	2	-2	9	15



Fig 3. Simulated a) soil C stocks, b) soil C/P, c) soil C/N and d) soil N/P ratios over the rotation period (i.e., from the relative age of 10 years to 10 years after the final harvest). The rotation period increases from southern to northern Sweden and the small peaks in soil C were related to forest operations, which were more frequent in southern Sweden. A clearance at year 10 was conducted across all four sites. Thinnings varied from four in southern regions to two in northern regions.

Table 6 Simulated and measured annual P losses through leaching. Measured TP is larger than the simulated DOP and PO4 fraction due to the presence of particulate phosphorus and because the measured value contains P leaching from upstream sources.

P leaching	64°N	61 °N	57 °N	56 °N
Annual regional total P leaching, measured (kg P ha <sup>-1</sup> )	0.04	0.02	0.09	0.08
Annual regional total P leaching, simulated (kg P ha <sup>-1</sup> )	0.03	0.01	0.05	0.07
Average TP concentration, measured $(mg l^{-1})$	0.0067	0.0066	0.03	0.02
Average PO <sub>4</sub> +DOP concentration, simulated (mg l <sup>-1</sup> )	0.0056	0.002	0.003	0.006
Fraction of dissolved organic P in total leached P, measured	63%	64%	83%	61%
Fraction of dissolved organic P in total leached P, simulated	56%	74%	15%	12%



Fig. 4 Simulated mean annual fluxes in a) C, b) N and c) P in the four regions. The numbers above the stacks indicate the annual mean change in the ecosystem. The simulation period
starts from year 10 and ends 10 years after final felling. Plant growth in a) represents net primary production.



Fig. 5 Simulated annual mean a) harvested biomass, b) soil C change (positive values denote sequestration, negative values denote losses), c) total N leaching and d) total P leaching in response to changes in soil organic N/P ratio across the four regions. The bars demonstrate the standard deviation of a mean value based on changes in short-cut uptake rates of N and P (see Table 2).

#### References

Aerts, R.: Nutrient resorption from senescing leaves of perennials: are there general patterns?, Journal of Ecology, 84(4), 597-608, 1996.

Akselsson, C., Pihl Karlsson, G., Karlsson, P.-E., and Ahlstrand, J.: Miljöövervakning på

1105 Obsytorna 1984-2013, Beskrivning, resultat, utvärdering och framtid, Skogsstyrelsen Rapport 2015:1, 2015.

Akselsson, C., Westling, O., Alveteg, M., Thelin, G., Fransson, A. M., and Hellsten, S.: The influence of N load and harvest intensity on the risk of P limitation in Swedish forest soils, Sci Total Environ, 404, 284-289, 10.1016/j.scitotenv.2007.11.017, 2008.

- 1110 Almeida, J. P., Rosenstock, N. P., Forsmark, B., Bergh, J., and Wallander, H.: Ectomycorrhizal community composition and function in a spruce forest transitioning between nitrogen and phosphorus limitation, Fungal Ecol, 10.1016/j.funeco.2018.05.008, 2018.
- Andersson, M., Carlsson, M., Ladenberger, A., Morris, G., Sadeghl, M., and Uhlbäck, J.:
  Geokemisk atlas över sverige [Geochemical atlas of Sweden], Sveriges Geologiska Undersökning, Uppsala, Sweden, 2014.
  Arheimer, B., Dahné, J., Donnelly, C., Lindström, G., and Strömqvist, J.: Water and nutrient

Arheimer, B., Dahné, J., Donnelly, C., Lindström, G., and Strömqvist, J.: Water and nutrient simulations using the HYPE model for Sweden vs. the Baltic Sea basin – influence of inputdata quality and scale, Hydrol Res, 43, 315-329, 10.2166/nh.2012.010, 2012.

1120 Arnold, J. G., Daniel, N. M., Gassman, P. W., Abbaspour, K. C., and White, M. J.: SWAT: Model use, calibration, and validation, Transactions of the American Society of Agricultural and Biological Engineers, 55, 1491-1508, 2012. Averill, C., Turner, B. L., and Finzi, A. C.: Mycorrhiza-mediated competition between plants

Averill, C., Turner, B. L., and Finzi, A. C.: Mycorrhiza-mediated competition between plants and decomposers drives soil carbon storage, Nature, 505, 543-545, 10.1038/nature12901, 2014.

Bahr, A., Ellström, M., Bergh, J., and Wallander, H.: Nitrogen leaching and ectomycorrhizal nitrogen retention capacity in a Norway spruce forest fertilized with nitrogen and phosphorus, Plant and Soil, 390, 323-335, 10.1007/s11104-015-2408-6, 2015.

Barrow, N. J.: The description of desorption of phosphate from soil, J Soil Sci, 30, 259-270, 1130 1979.

Baum, C., and Makeschin, F.: Effects of nitrogen and phosphorus fertilization on mycorrhizal formation of two poplar clones *(Populus trichocarpa and P. tremula x tremuloides)*, J Plant Nutr Soil Sci, 163, 491-497, 2000.

Bell, C., Carrillo, Y., Boot, C. M., Rocca, J. D., Pendall, E., and Wallenstein, M. D.:
Rhizosphere stoichiometry: are C : N : P ratios of plants, soils, and enzymes conserved at the plant species-level?, New Phytol, 201, 505-517, 10.1111/nph.12531, 2014.
Bolan, N. S.: A critical review on the role of mycorrhizal fungi in the uptake of phosphorus

by plants, Plant and Soil, 134, 189-207, 1991.
Braun, S., Thomas, V. F., Quiring, R., and Fluckiger, W.: Does nitrogen deposition increase
forest production? The role of phosphorus, Environ Pollut, 158, 2043-2052, 10.1016/j.envpol.2009.11.030, 2010.

Bucher, M.: Functional biology of plant phosphate uptake at root and mycorrhiza interfaces, New Phytol, 173, 11-26, 10.1111/j.1469-8137.2006.01935.x, 2007.

Bünemann, E. K.: Enzyme additions as a tool to assess the potential bioavailability of organically bound nutrients, Soil Biol and Biochem, 40, 2116-2129, 10.1016/j.soilbio.2008.03.001, 2008.
Bünemann, E. K.: Assessment of gross and net mineralization rates of soil organic

phosphorus-A review, Soil Biol Biochem, 89, 82-98, 10.1016/j.soilbio.2015.06.026, 2015.

Cintas O., Berndes, G., Hansson, J., Poudel, B. C., Bergh, J., Börjesson, P., Egnell, G., 1150 Lundmark, T., and Nordin, A.: The potential role of forest management in Swedish scenarios towards climate neutrality by mid century, Forest Ecol Manag, 383, 73-84. doi: 10.1016/j.foreco.2016.07.015, 2017.

Clemmensen, K. E., Bahr, A., Ovaskainen, O., Dahlberg, A., Ekblad, A., Wallander, H., Stenlid, J., Finlay, R. D., Wardle, D. A., and Lindahl, B. D.: Roots and associated fungi drive long-term carbon sequestration in boreal forest, Science, 339, 1615-1618,

- 1155 long-term carbon sequestration in boreal forest, Science, 339, 1615-1618, 10.1126/science.1231923, 2013.
  Cleveland, C. C., and Liptzin, D.: C:N:P stoichiometry in soil: is there a "Redfield ratio" for the microbial biomass?, Biogeochemistry, 85, 235-252, 10.1007/s10533-007-9132-0, 2007. Cole, C. V., Innis, G. S., and Stewart, J. W. B.: Simulation of Phosphorus cycling in semiarid
- grasslands, Ecology, 58, 2-15, 1977.
  Crowley, K. F., McNeil, B. E., Lovett, G. M., Canham, C. D., Driscoll, C. T., Rustad, L. E., Denny, E., Hallett, R. A., Arthur, M. A., Boggs, J. L., Goodale, C. L., Kahl, J. S., McNulty, S. G., Ollinger, S. V., Pardo, L. H., Schaberg, P. G., Stoddard, J. L., Weand, M. P., and Weathers, K. C.: Do nutrient limitation patterns shift from Nitrogen toward Phosphorus with
- increasing Nitrogen deposition across the northeastern United States?, Ecosystems, 15, 940-957, 10.1007/s10021-012-9550-2, 2012.
  Deng, Q., Hui, D., Dennis, S., and Reddy, K. C.: Responses of terrestrial ecosystem phosphorus cycling to nitrogen addition: A meta-analysis, Global Ecol Biogeogr, 26, 713-728, 10.1111/geb.12576, 2017.
- Du, E., Terrer, C., Pellegrini, A.F.A. Ahlström, A., van Lissa, C.J., Zhao, X., Xia N., Wu, X. 1170 and Jackson, R.B.: Global patterns of terrestrial nitrogen and phosphorus limitation. Nature Geosci, 13, 221–226. https://doi.org/10.1038/s41561-019-0530-4, 2020. Eckersten, H., and Beier, C.: Comparison of N and C dynamics in two Norway spruce stands oriented model, Eniron using process simulation Pollut. 102. 395-401. а https://doi.org/10.1016/S0269-7491(98)80059-6, 1998.
- https://doi.org/10.1016/S0269-7491(98)80059-6, 1998.
   Ekblad, A., Wallander, H., Carlsson, R., and Huss-Danell, K.: Fungal biomass in roots and extramatrical mycelium in relation to macronutrients and plant biomass of ectomycorrhizal *Pinus Sylvestris* and *Alnus incana*, New Phytol, 131, 443-451, 1995.
   Elser, J. J., Bracken, M. E., Cleland, E. E., Gruner, D. S., Harpole, W. S., Hillebrand, H.,
- Ngai, J. T., Seabloom, E. W., Shurin, J. B., and Smith, J. E.: Global analysis of nitrogen and phosphorus limitation of primary producers in freshwater, marine and terrestrial ecosystems, Ecol Lett, 10, 1135-1142, 10.1111/j.1461-0248.2007.01113.x, 2007.
   Ericsson, T.: Growth and shoot: root ratio of seedlings in relation to nutrient availability, Plant Soil, 168, 205-214, 1995.
- 1185 Fleischer, K., Rammig, A., De Kauwe, M. G., Walker, A. P., Domingues, T. F., Fuchslueger, L., Garcia, S., Goll, D. S., Grandis, A., Jiang, M., Haverd, V., Hofhansl, F., Holm, J. A., Kruijt, B., Leung, F., Medlyn, B. E., Mercado, L. M., Norby, R. J., Pak, B., von Randow, C., Quesada, C. A., Schaap, K. J., Valverde-Barrantes, O. J., Wang, Y.-P., Yang, X., Zaehle, S., Zhu, Q., and Lapola, D. M.: Amazon forest response to CO2 fertilization dependent on plant
- phosphorus acquisition, Nature Geosci, 12(9), 736-741, 10.1038/s41561-019-0404-9, 2019.
   Fransson, A.-M., and Bergkvist, B.: Phosphorus fertilisation causes durable enhancement of phosphorus concentrations in forest soil, Forest Ecol Manag, 130, 69-76, 10.1016/S0378-1127(99)00184-X, 2000.

Gassman, P.W., Williams, J.R., Benson, V.W., César lzaurralde, R., Hauck, L.M., Jones,
C.A., Atwood, J.D., Kiniry, J.D., and Flowers, J.D.: Historical development and applications of the EPIC and APEX models, CARD working paper 05-WP 397. Center for Agricultural and Rural Development, Iowa State University, Ames, IA, DOI: <u>10.13031/2013.17074</u>, 2005. Giesler, R., Petersson, T., and Högberg, P.: Phosphorus limitation in boreal forests: effects of

aluminum and iron accumulation in the humus layer, Ecosystems, 5, 300-314, 1200 10.1007/s10021-001-0073-5, 2002.

Goll, D. S., Brovkin, V., Parida, B. R., Reick, C. H., Kattge, J., Reich, P. B., van Bodegom, P. M., and Niinemets, Ü.: Nutrient limitation reduces land carbon uptake in simulations with a model of combined carbon, nitrogen and phosphorus cycling, Biogeosciences, 9, 3547-3569, 10.5194/bg-9-3547-2012, 2012.

- 1205 Goll, D. S., Vuichard, N., Maignan, F., Jornet-Puig, A., Sardans, J., Violette, A., Peng, S., Sun, Y., Kvakic, M., Guimberteau, M., Guenet, B., Zaehle, S., Penuelas, J., Janssens, I., and Ciais, P.: A representation of the phosphorus cycle for ORCHIDEE (revision 4520), Geosci Model Dev, 10, 3745-3770, 10.5194/gmd-10-3745-2017, 2017.
- Gower, S. T., and Vitousek, P. M.: Effects of nutrient amendments on fine root biomass in a primary successional forest in Hawai'i, Oecologia, 81, 566-568, 1989.
- Gress, E. S., Nichols, T. D., Northcraft, C. C., and Peterjohn, W. T.: Nutrient limitation in soils exhibiting differing nitrogenavailabilities: what lies beyond nitrogen saturation?, Ecology, 88, 119-130, 10.1890/0012-9658(2007)88[119:NLISED]2.0.CO;2, 2007.
- Groenendijk, P., Renaud, L.V., and Roelsma, J.: Prediction of nitrogen and phosphorus leaching to groundwater and surface waters; process descriptions of the Animo4.0 model. Wageningen, Alterra–Report 983, 114 pp., 2005.

Guidry, M. W., and Machenzie, F. T.: Apatite weathering and the phanerozoic phosphorus cycle, Geology, 28, 631-634, 2000.

Güsewell, S.: N : P ratios in terrestrial plants: variation and functional significance, New Phytol, 164, 243-266, 10.1111/j.1469-8137.2004.01192.x, 2004.

- Gärdenäs, A., Eckersten, H., and Lillemägi, M.: Modeling long-term effects of N fertilization and N deposition on the N balances of forest stands in Sweden, Swedish University of Agricultural Sciences, 1651-7210, 34, 2003.
- Gärdenäs, A. Jansson, P.-E. and Karlberg, L.: A model of accumulation of radionuclides in
  biosphere originating from groundwater contamination. SKB report. R-06-47, SKB, Solna,
  Sweden, 2006.

Gärdenäs, A., Ågren, G., Bird, J., Clarholm, M., Hallin, S., Ineson, P., Kätterer, T., Knicker, H., Nilsson, I., Näsholm, T., Ogle, S., Paustian, K., Persson, T., and Stendahl, J.: Knowledge gaps in soil carbon and nitrogen interactions – From molecular to global scale, Soil Biology and Biochemistry, 43 (4), 702-717, 10.1016/j.soilbio.2010.04.006, 2011.

- He, H., Meyer, A., Jansson, P.-E., Svensson, M., Rütting, T., and Klemedtsson, L.: Simulating ectomycorrhiza in boreal forests: implementing ectomycorrhizal fungi model MYCOFON in CoupModel (v5), Geosci Model Dev, 11, 725-751, 10.5194/gmd-11-725-2018, 2018.
- Hinsinger, P.: Bioavailability of soil inorganic P in the rhizosphere as affected by root-induced chemical changes: a review, Plant Soil, 237, 173-195, 2001.
  Högberg, P., Näsholm, T., Franklin, O., and Högberg, M. N.: Tamm Review: On the nature of the nitrogen limitation to plant growth in Fennoscandian boreal forests, Forest Ecol Manag, 403, 161-185, 10.1016/j.foreco.2017.04.045, 2017.
- Ingestad, T., and Ågren, G. I.: Theories and methods on plant nutrient and growth, Physiol Plantarum, 84, 177-184, 1992.
  Ingestad, T.: Mineral nutrient requirements of *Pinus silvestris* and *Picea abies* Seedlings, , Physiologia Plantarum, 45, 373-380, 1979.
- 1245 Jackson-Blake, L. A., Wade, A. J., Futter, M. N., Butterfield, D., Couture, R. M., Cox, B. A., Crossman, J., Ekholm, P., Halliday, S. J., Jin, L., Lawrence, D. S. L., Lepistö, A., Lin, Y., Rankinen, K., and Whitehead, P. G.: The INtegrated CAtchment model of phosphorus dynamics (INCA-P): Description and demonstration of new model structure and equations, Environ mental Modelling & Software, 83, 356-386, 10.1016/j.envsoft.2016.05.022, 2016.

Jahn, R., Blume, H.-P., Asio, V.-B., Spaargaren, O., and Schad, P.: Guidelines for soil description, 4<sup>th</sup> ed. Food and Agriculture Organization of the United Nations, Rome, Italy, 2006.
 Jansson, P. F. and Karlberg, J. : User manual of Coupled heat and mass transfer model for

Jansson, P.-E., and Karlberg, L.: User manual of Coupled heat and mass transfer model for soil-plant-atmosphere systems, Royal Institute of Technology, Department of Land and Water Resources, Stockholm, Sweden, 2011.

Jansson, P. E.: CoupModel: model use, calibration, and validation, T ASABE, 4, 1335-1344, 2012.

Johnson, A. H., Frizano, J., and Vann, D. R.: Biogeochemical implications of labile phosphorus in forest soils determined by the Hedley fractionation procedure, Oecologia, 135, 487-499, 10.1007/s00442-002-1164-5, 2003.

- Jonard, M., Furst, A., Verstraeten, A., Thimonier, A., Timmermann, V., Potocic, N., Waldner, P., Benham, S., Hansen, K., Merila, P., Ponette, Q., de la Cruz, A. C., Roskams, P., Nicolas, M., Croise, L., Ingerslev, M., Matteucci, G., Decinti, B., Bascietto, M., and Rautio, P.: Tree mineral nutrition is deteriorating in Europe, Glob Chang Biol, 21, 418-430, 10.1111/gcb.12657, 2015.
- Jones, C. A., Cole, C. V., Sharpley, A. N., and Williams, J. R.: A simplified soil and plant phosphorus model: I. Documentation, Soil Sci Soc Am J, 48, 800-805, 10.2136/sssaj1984.03615995004800040020x, 1984.

Kaiser, K., and Kalbitz, K.: Cycling downwards – dissolved organic matter in soils, Soil Biol Biochem, 52, 29-32, 10.1016/j.soilbio.2012.04.002, 2012.

Kalbitz, K., Solinger, S., Park, J.-H., Michalzik, B., and Matzner, E.: Controls on the dynamics of dissolved organic matter in soils: a review, Soil Sci, 165, 277-304, 2000. Kleja, D. B., Svensson, M., Majdi, H., Jansson, P.-E., Langvall, O., Bergkvist, B., Johansson, M.-B., Weslien, P., Truusb, L., Lindroth, A., and Ågren, G. I.: Pools and fluxes of carbon in

three Norway spruce ecosystems along a climatic gradient in Sweden, Biogeochemistry, 89, 7-25, 10.1007/s10533-007-9136-9, 2008.
Knisel, W. G., and Turtola, E.: Gleams model application on a heavy clay soil in Finland, Agr Water Manage, 43, 285-309, 10.1016/S0378-3774(99)00067-0, 2000.

Kronnäs, V., Akselsson, C., and Belyazid, S.: Dynamic modelling of weathering rates – the benefit over steady-state modelling, Soil, 5, 33-47, 10.5194/soil-5-33-2019, 2019.

Lagerström, A., Esberg, C., Wardle, D. A., and Giesler, R.: Soil phosphorus and microbial response to a long-term wildfire chronosequence in northern Sweden, Biogeochemistry, 95, 199-213, 10.1007/s10533-009-9331-y, 2009.

Laiho, R., and Prescott, C. E.: Decay and nutrient dynamics of coarse woody debris in northern coniferous forests: a synthesis, Can J Forest Res, 34, 763-777, https://doi.org/10.1139/x03-241, 2004.

Lang, F., Bauhus, J., Frossard, E., George, E., Kaiser, K., Kaupenjohann, M., Krüger, J., Matzner, E., Polle, A., Prietzel, J., Rennenberg, H., and Wellbrock, N.: Phosphorus in forest ecosystems: New insights from an ecosystem nutrition perspective, J Plant Nutr Soil Sci, 179, 129-135, 10.1002/jpln.201500541, 2016.

1290 129-135, 10.1002/jpln.201500541, 2016.
Liebig, J.: Die Chemie in ihrer Andwendung auf Agrikultur und Physiologie, Vieweg und Söhne, Braunschweig, 1840.
Linder, S.: Foliar analysis for detecting and correcting nutrient imbalances in Norway spruce, Ecol Bull, 44, 178-190, <u>https://www.jstor.org/stable/20113161</u>, 1995.

Manzoni, S., Trofymow, J. A., Jackson, R. B., and Porporato, A.: Stoichiometric controls on carbon, nitrogen, and phosphorus dynamics in decomposing litter, Ecol Monogr, 80, 89-106, <a href="https://doi.org/10.1890/09-0179.1">https://doi.org/10.1890/09-0179.1</a>, 2010.

McGill, W. B., and Cole, C. V.: Comparative aspects of cycling of organic C, N, S and P through soil organic matter, Geoderma, 26, 267-286, 1981.

1255

- Medlyn, B. E., De Kauwe, M. G., Zaehle, S., Walker, A. P., Duursma, R. A., Luus, K., Mishurov, M., Pak, B., Smith, B., Wang, Y. P., Yang, X., Crous, K. Y., Drake, J. E., Gimeno, T. E., Macdonald, C. A., Norby, R. J., Power, S. A., Tjoelker, M. G., and Ellsworth, D. S.: Using models to guide field experiments: a priori predictions for the CO2 response of a nutrient- and water-limited native Eucalypt woodland, Glob Chang Biol, 22, 2834-2851, 10.1111/gcb.13268, 2016.
- 1305 10.1111/gcb.13268, 2016.
   Meyer, A., Grote, R., Polle, A., and Butterbach-Bahl, K.: Simulating mycorrhiza contribution to forest C- and N cycling-the MYCOFON model, Plant and Soil, 327, 493-517, 10.1007/s11104-009-0017-y, 2009.

Monteith, J. L.: Evaporation and environment, Symp Soc Exp Biol, 16, 205-234, 1965.

1310 Nasholm, T., Hogberg, P., Franklin, O., Metcalfe, D., Keel, S. G., Campbell, C., Hurry, V., Linder, S., and Hogberg, M. N.: Are ectomycorrhizal fungi alleviating or aggravating nitrogen limitation of tree growth in boreal forests?, New Phytol, 198, 214-221, 10.1111/nph.12139, 2013.

Nehls, U.: Mastering ectomycorrhizal symbiosis: the impact of carbohydrates, J Exp Bot, 59, 1315 1097-1108, 10.1093/jxb/erm334, 2008.

Nieminen, T., and Helmisaari, H.-S.: Nutrient retranslocation in the foliage of *Pinus sylvestris L*. growing along a heavy metal pollution gradient, Tree Physiol, 16 (10): 825-831, <u>10.1093/treephys/16.10.825</u>, 1996.

Nylund, J.-E., and Wallander, H.: Ergosterol Analysis as a means of quantifying mycorrhizal biomass, Methods Microbiol, 24, 77-88, 1992.

- Olander, L., and Vitousek, P.: Short-term controls over inorganic phosphorus during soil and ecosystem development, Soil Biol and Biochem, 37, 651-659, 10.1016/j.soilbio.2004.08.022, 2005.
- Olsson, M. T., Erlandsson, M., Lundin, L., Nilsson, T., Nilsson, Å., and Stendahl, J.: Organic carbon stocks in Swedish Podzol soils in relation to soil hydrology and other site characteristics, Silva Fennica, 43, 209-222, 2009.

Ortiz, C. A., Lundblad, M., Lundström, A., and Stendahl, J.: The effect of increased extraction of forest harvest residues on soil organic carbon accumulation in Sweden, Biomass Bioenergy, 70, 230-238, 10.1016/j.biombioe.2014.08.030, 2014.

- Orwin, K. H., Kirschbaum, M. U., St John, M. G., and Dickie, I. A.: Organic nutrient uptake by mycorrhizal fungi enhances ecosystem carbon storage: a model-based assessment, Ecol Lett, 14, 493-502, 10.1111/j.1461-0248.2011.01611.x, 2011.
  Oulehle, F., Chuman, T., Hruška, J., Krám, P., McDowell, W. H., Myška, O., Navrátil, T.,
- and Tesař, M.: Recovery from acidification alters concentrations and fluxes of solutes from
  Czech catchments, Biogeochemistry, 132, 251-272, 10.1007/s10533-017-0298-9, 2017.
- Penuelas, J., Poulter, B., Sardans, J., Ciais, P., van der Velde, M., Bopp, L., Boucher, O., Godderis, Y., Hinsinger, P., Llusia, J., Nardin, E., Vicca, S., Obersteiner, M., and Janssens, I. A.: Human-induced nitrogen-phosphorus imbalances alter natural and managed ecosystems across the globe, Nat Commun, 4, 2934, 10.1038/ncomms3934, 2013.
- 1340 Read, D. J., and Perez-Moreno, J.: Mycorrhizas and nutrient cycling in ecosystems a journey towards relevance?, New Phytol, 157, 475-492, 10.1046/j.1469-8137.2003.00704.x, 2003.

Reed, S. C., Yang, X., and Thornton, P. E.: Incorporating phosphorus cycling into global modeling efforts: a worthwhile, tractable endeavor, New Phytol, 208, 324-329, 10.1111/nph.13521, 2015.

Richardson, A. E., Barea, J.-M., McNeill, A. M., and Prigent-Combaret, C.: Acquisition of phosphorus and nitrogen in the rhizosphere and plant growth promotion by microorganisms, Plant Soil, 321, 305-339, 10.1007/s11104-009-9895-2, 2009.

1345

Richardson, A. E., and Simpson, R. J.: Soil microorganisms mediating phosphorus 1350 availability microbial phosphorus, Plant Physiol, 989-996, update on 156. 10.1104/pp.111.175448, 2011. Rosengren-Brinck, U., and Nihlgård, B.: Nutritional status in needles of Norway spruce in nutrient relation water and supply, Ecol Bull. 44. 168-177, to

https://www.jstor.org/stable/20113160, 1995.

- 1355 Rosling, A., Midgley, M. G., Cheeke, T., Urbina, H., Fransson, P., and Phillips, R. P.: Phosphorus cycling in deciduous forest soil differs between stands dominated by ecto- and arbuscular mycorrhizal trees, New Phytol, 209, 1184-1195, 10.1111/nph.13720, 2016. Saito, M. A., Goepfert, T. J., and Ritt, J. T.: Some thoughts on the concept of colimitation: three definitions and the importance of bioavailability, Limnol Oceanogr, 53, 276-290, 2008.
- Schachtman, D. P., Reid, R. J., and Ayling, S. M.: Phosphorus uptake by plants: from soil to cell, Plant Physiol, 116, 447-453, 1998.
  Schlesinger, W. H.: Biogeochemistry. An analysis of global change, 2nd ed., Academic Press, Cambridge, MA, 588 pp., 1997.
- Schnepf, A., and Roose, T.: Modelling the contribution of arbuscular mycorrhizal fungi to
  plant phosphate uptake, New Phytol, 171, 669-682, 10.1111/j.1469-8137.2006.01771.x,
  SLU: Skogsdata: Aktuella uppgifter om de svenska skogarna från Riksskogstaxeringen, SLU,

Umeå, Sweden, 2003, available at https://www.slu.se/globalassets/ew/org/centrb/rt/dokument/skogsdata/skogsdata2003\_webb.p df (last accessed 2020-02-10)

- SLU: Skogsdata: Aktuella uppgifter om de svenska skogarna från Riksskogstaxeringen, SLU, Umeå, Sweden, 2012, available at https://pub.epsilon.slu.se/9266/1/SkogsData2012\_webb.pdf (last accessed 2020-02-10). Smeck, N. E.: Phosphorus dynamics in soils and landscapes, Geoderma, 36, 185-199, 1985.
- 1375 Smith, S. E.: Mycorrhizal fungi can dominate phosphate supply to plants irrespective of growth responses, Plant Physiol, 133, 16-20, 10.1104/pp.103.024380, 2003. Smith, S. E., and Read, D. J.: Mycorrhizal symbiosis, 3rd ed., Academic Press, Cambridge, MA, 2008.
- Staddon, P. L., Thompson, K., Jakobsen, I., Grime, J. P., Askew, A. P., and Fitter, A. H.:
  Mycorrhizal fungal abundance is affected by long-term climatic manipulations in the field, Glob Change Biol, 9, 186-194, 10.1046/j.1365-2486.2003.00593.x, 2003.
  Stendahl, J., Johansson, M.-B., Eriksson, E., Nilsson, Å., and Langvall, O.: Soil organic carbon in Swedish spruce and pine forests- differences in stock levels and regional patterns, Silva Fennica, 44, 5-21, 2010.
- Sundqvist, M. K., Liu, Z., Giesler, R., and Wardle, D. A.: Plant and microbial responses to nitrogen and phosphorus addition across an elevational gradient in subarctic tundra, Ecology, 95, 1819-1835, <u>https://doi.org/10.1890/13-0869.1</u>, 2014. Sverdrup, H., and Warfvinge, P., Calculating field weathering rates using a mechanistic

geochemical model PROFILE, Appl Geochem, 8, 273-283, 10.1016/0883-2927(93)90042-F, 1390 1993.

Swedish Forest Agency: Grundbok for skogsbrukare. Swedish Forest Agency, Jönköping, Sweden, 190 pp., 2005.

Svensson, M., Jansson, P.-E., and Berggren Kleja, D.: Modelling soil C sequestration in spruce forest ecosystems along a Swedish transect based on current conditions, Biogeochemistry, 89, 95-119, 10.1007/s10533-007-9134-y, 2008.

Biogeochemistry, 89, 95-119, 10.1007/s10533-007-9134-y, 2008.
Talkner, U., Meiwes, K. J., Potočić, N., Seletković, I., Cools, N., De Vos, B., and Rautio, P.: Phosphorus nutrition of beech (Fagus sylvatica L.) is decreasing in Europe, Ann For Sci, 72, 919-928, 10.1007/s13595-015-0459-8, 2015.

Tang, Z., Xu, W., Zhou, G., Bai, Y., Li, J., Tang, X., Chen, D., Liu, Q., Ma, W., Xiong, G.,

- He, H., He, N., Guo, Y., Guo, Q., Zhu, J., Han, W., Hu, H., Fang, J., and Xie, Z.: Patterns of plant carbon, nitrogen, and phosphorus concentration in relation to productivity in China's terrestrial ecosystems, Proc Natl Acad Sci, 201700295, 10.1073/pnas.1700295114 %J, 2018. Tarvainen, L., Lutz, M., Rantfors, M., Nasholm, T., and Wallin, G.: Increased needle nitrogen contents did not improve shoot photosynthetic performance of mature nitrogen-poor
- Scots pine trees, Front Plant Sci, 7, 1051, 10.3389/fpls.2016.01051, 2016.
   Terrer, C., Vicca, S., Hungate, B. A., Phillips, R. P., and Prentice, I. C.: Mycorrhizal association as a primary control of the CO2 fertilization effect, Science, 353, 72-74, 10.1126/science.aaf4610, 2016.
  - Terrer, C., Jackson, R. B., Prentice, I. C., Keenan, T. F., Kaiser, C., Vicca, S., Fisher, J. B.,
- Reich, P. B., Stocker, B. D., Hungate, B. A., Peñuelas, J., McCallum, I., Soudzilovskaia, N. A., Cernusak, L. A., Talhelm, A. F., Van Sundert, K., Piao, S., Newton, P. C. D., Hovenden, M. J., Blumenthal, D. M., Liu, Y. Y., Müller, C., Winter, K., Field, C. B., Viechtbauer, W., Van Lissa, C. J., Hoosbeek, M. R., Watanabe, M., Koike, T., Leshyk, V. O., Polley, H. W., and Franklin, O.: Nitrogen and phosphorus constrain the CO2 fertilization of global plant biomass, Nat Clim Change, 10.1038/s41558-019-0545-2, 2019.
- Tessier, J. T., and Raynal, D. J.: Use of nitrogen to phosphorus ratios in plant tissue as an indicator of nutrient limitation and nitrogen saturation, J Appl Ecol, 40, 523-534, 10.1046/j.1365-2664.2003.00820.x, 2003.
- Thelin, G., Rosengren-Brinck, U., Nihlgård, B., and Barkman, A.: Trends in needle and soil
  chemistry of Norway spruce and Scots pine stands in South Sweden 1985-1994, Environ
  Poll, 99, 149-158, 10.1016/S0269-7491(97)00192-9, 1998.
  - Thelin, G., Rosengren, U., Callesen, I., and Ingerslev, M.: The nutrient status of Norway spruce in pure and in mixed-species stands, For Ecol and Manage, 160, 115-125, 10.1016/S0378-1127(01)00464-9, 2002.
- 1425 Thum, T., Caldararu, S., Engel, J., Kern, M., Pallandt, M., Schnur, R., Yu, L., and Zaehle, S.: A new model of the coupled carbon, nitrogen, and phosphorus cycles in the terrestrial biosphere (QUINCY v1.0; revision 1996), Geosci Model Dev, 12, 4781–4802, https://doi.org/10.5194/gmd-12-4781-2019, 2019.
- Tipping, E., Benham, S., Boyle, J. F., Crow, P., Davies, J., Fischer, U., Guyatt, H., Helliwell,
  R., Jackson-Blake, L., Lawlor, A. J., Monteith, D. T., Rowe, E. C., and Toberman, H.: Atmospheric deposition of phosphorus to land and freshwater, Environ Sci Process Impacts, 16, 1608-1617, 10.1039/c3em00641g, 2014.

Treseder, K. K.: A meta-analysis of mycorrhizal responses to nitrogen, phosphorus, and atmospheric CO2 in field studies, New Phytol, 164, 347-355, 10.1111/j.1469-8137.2004.01159.x, 2004.

- Van Sundert, K., Radujkovic, D., Cools, N., De Vos, B., Etzold, S., Fernandez-Martinez, M., Janssens, I., Merila, P., Penuelas, J., Sardans, J., Stendahl, J., Terrer, C., and Vicca, S.: Towards comparable assessment of the soil nutrient status across scales review and development of nutrient metrics, Glob Chang Biol, 10.1111/gcb.14802, 2019.
- 1440 Wallander, H., Mahmood, S., Hagerberg, D., Johansson, L., and Pallon, J., Elemental composition of ectomycorrhizal mycelia identified by PCR-RFLP analysis and grown in contact with apatite or wood ash in forest soil, FEMS Microbiol Ecol, 44, 57-65, <u>10.1111/j.1574-6941.2003.tb01090.x</u>, 2003.
- Wang, Y. P., Houlton, B. Z., and Field, C. B.: A model of biogeochemical cycles of carbon,
  nitrogen, and phosphorus including symbiotic nitrogen fixation and phosphatase production,
  Glob Biogeochem Cycles, 21, 10.1029/2006gb002797, 2007.

Wang, Y. P., Law, R. M., and Pak, B.: A global model of carbon, nitrogen and phosphorus cycles for the terrestrial biosphere, Biogeosciences, 7, 2261-2282, 10.5194/bg-7-2261-2010, 2010.

- Wijk, S.: Skogsvårdsorganisationens skogliga observationsytor. Anvisningar för analys av markprover, version 1995-11-15. Skogsstyrelsen, Jönköping, Sweden, 1995.
   Wijk, S.: Skogsvårdsorganisationens skogliga observationsytor. Manual – anvisningar för urval av träd för barrprovtagning. Version 1997-08-14. Skogsstyrelsen, Jönköping, Sweden, 1997.
- 1455 Vincent, A. G., Sundqvist, M. K., Wardle, D. A., and Giesler, R.: Bioavailable soil phosphorus decreases with increasing elevation in a subarctic tundra landscape, PLoS One, 9, e92942, 10.1371/journal.pone.0092942, 2014.

Vitousek, P. M., Porder, S., Houlton, B. Z., and Chadwick, O. A.: Terrestrial phosphorus limitation: mechanisms, implications, and nitrogen-phosphorus interactions, Ecol Appl, 20, 5-1460 15, 2010.

- Yanai, R. D.: Phosphorus budget of a 70-year-old northern hardwood forest, Biogeochemistry, 17, 1-22, <u>https://doi.org/10.1007/BF00002757</u>, 1992.
  Yang, X., Thornton, P. E., Ricciuto, D. M., and Post, W. M.: The role of phosphorus dynamics in tropical forests – a modeling study using CLM-CNP, Biogeosciences, 11, 1667-
- 1465 1681, 10.5194/bg-11-1667-2014, 2014.
  Yu, L., Zanchi, G., Akselsson, C., Wallander, H., and Belyazid, S.: Modeling the forest phosphorus nutrition in a southwestern Swedish forest site, Ecol Model, 369, 88-100, 10.1016/j.ecolmodel.2017.12.018, 2018.
- Zhang, J., and Elser, J. J.: Carbon:Nitrogen:Phosphorus stoichiometry in fungi: a metaanalysis, Front Microbiol, 8:1281, <u>10.3389/fmicb.2017.01281</u>, 2017.
- Zhu, Q., Riley, W. J., Tang, J., and Koven, C. D., Multiple soil nutrient competition between plants, microbes, and mineral surfaces: model development, parameterization, and example applications in several tropical forests, Biogeosciences, 13(1), 341-363, http://dx.doi.org/10.5194/bg-13-341-2016, 2016.

1475