

## **Response to Reviewer #1**

### **General comment:**

*“There is a plethora of observational evidences supporting that diffuse d light favors higher plant productivity thanks to radiation being more evenly distributed and accessible across the shaded part of the canopy. Yet, few Land Surface Models ( have a representation of light transmission within the canopy that explicitly accounts for the quality of light (direct vs diffuse) and its effect of primary productivity. Zhang et al. implemented such a capability in the ORCHIDEE LSM (ORCHIDEE\_DF) and this paper provides the details of their modelling framework as well as a solid evaluation of ORCHIDEE\_DF performance against a large set of ground site measurements (FLUXNET).*

*This paper reads very easily as it is well written and well structured.*

*The first part of the paper describes the model framework and the parameterisations. It is generally well written, although some symbols equations could be improved, and a couple of sections swapped together (see general comments).*

*The second part of the paper presents a very good evaluation of ORCHIDEE\_DF The added value of introducing a representation of diffuse light fraction is convincingly exposed and rigorous efforts were made to disentangle this from other cofounding effects (e.g. VDP, Temperature). Despite the calibration of model parameter being sub-optimal for this new configuration of ORCHIDEE, the analysis and supporting plots are very useful and effectively achieve to highlight where the model performs well, and which future development efforts should be prioritized. This is a useful evaluation effort which will also benefit the wider LSM community beyond the ORCHIDEE user base. The effort that the authors went through in evaluating ORCHIDEE\_DF against a large ensemble of observations goes much further than previous attempts published in the literature and is greatly appreciated.*

*Adding a representation of diffuse light fraction in the canopy can only be useful if the boundary conditions that is the fraction of diffuse radiation hitting the top of the canopy is known. This information is usually lacking from the dataset that are used to drive LSMs. Technically this is a problem that is external to land surface modelling, but it is*

*great to see that Zhang et al. provide a practical framework to retrieve that missing information and could offer some insight to the terrestrial carbon cycle community for a developing a harmonized framework in future LSMs inter-comparisons.*

*The topic covered in this paper is absolutely relevant to GMD and I therefore strongly support its publication after addressing those minor very few points.”*

**[Response]** We thank the reviewer for the careful review and helpful comments and suggestions, which helped us to significantly improve our manuscript. We have addressed all the suggestions and comments in our revision. Please find below the reviewer’s comments, followed by our responses and relevant changes in the manuscript. We hope that the revised version addresses all the issues raised by the reviewer.

**Comments:**

*“1. I believe it will be improved at production stage, but some equations are not easy to read in current form. Use of upper script and lower script could help bringing better separation between the terms in the equations (e.g.  $Km_C$  instead of  $KmC$ ,  $C_c$  instead of  $Cc$ , etc).”*

**[Response]** We thank the reviewer for this suggestion, the notations have been improved throughout the manuscript. ( $A_c$  to  $A_c$ ,  $A_j$  to  $A_j$ ,  $R_d$  to  $R_d$ ,  $C_c$  to  $C_c$ ,  $KmC$  to  $Km_C$ ,  $KmO$  to  $Km_O$ )

*“2. Would it make more sense to introduce section 2.1.3 (Light transmission in ORCHIDEE\_DF) before section 2.1.2 (Light partitioning in ORCHIDEE\_DF) so it follows naturally section 2.1.1 (Light transmission in ORCHIDEE\_trunk) especially given that the calculation of the fraction of diffuse light hitting the canopy top could eventually be treated by the radiative transfer of the driving atmospheric model as it is done in an Earth System Framework (e.g. Yue et al., 2017; Malavelle et al. 2019) making a specific parameterization for this not necessary in ORCHIDEE.”*

**[Response]** We agree with the reviewer that it is more reasonable to put the two light

transmission sections together. Sections 2.1.2 and 2.1.3 have been swapped in the updated manuscript.

*“3. For the evaluation framework described at P 11 L 324 to 326 - Getting the same level of PPFD that way may involves comparing GPPs at different time during the day which might not capture vegetation in similar physiological states. Wouldn't it be easier to simply normalize the cloudy and sunny GPPs by their respective PPFDs rather than removing a part of the dataset (likely the midday data for the sunny GPP when insolation is maximal and light saturation of the sunlit leaves possible)?”*

**[Response]** Thanks for this question. We considered carefully and tried to use the proposed normalization method (Fig R1). The results are similar to what we found controlling PPFD level in the manuscript. However, we did not use it in the manuscript due to some concerns. It is known that the light response curve is not linear. Therefore, the LUE (GPP/PPFD) should depend on PPFD level. Due to the nature of atmospheric




Figure R1. Site distribution of the observed and modeled LUE difference between cloudy and sunny conditions at midday during the peak growing season (monthly GPP>90% of the mean monthly GPP maximum).

light transmission, the cloudy PPFD should be smaller than the sunny PPFD for a given solar zenith angle. If the PPFD level is not controlled, it would become difficult to explain whether the difference in LUE is due to diffuse radiation fraction or to the PPFD level. Therefore, in the manuscript, we compared the GPP and LUE with PPFD controlled at different times of the day (Fig. 7), which, we think, has ensured the vegetation to have similar physiological states in each period.

*“4. P 13 L 387 389 It is interesting to note that both ORCHIDEE trunk and DF underestimate the dGPP and the dLUE around midday. Could it be related to the relative high proportion of sunlit leaves which is primarily a function of the solar zenith angle in the DF configuration? Segregating the dataset in to latitudes may help to appreciate if this behaviour occurs more in the tropic or the mid latitude sites”*

**[Response]** Thanks for this point. We had made an extra analysis to investigate the latitude dependence of the ratio between modeled dGPP vs observed dGPP at midday (Fig. R2, two outlier sites not shown on the plot). There are no sites having similar midday PPFD level under sunny and cloudy conditions in low latitudes. According to the remaining data, positive relationship between the dGPP ratios and latitudes is not significant for both trunk and DF simulations. From our perspective, the underestimation in midday dGPP could be a result of parameterizations of processes other than diffuse radiation in ORCHIDEE because both the trunk and DF configurations have this problem. This will be added to the manuscript (Line 450-451). “The underestimation of midday  $\Delta$ GPP could be a result of error in current ORCHIDEE parameterizations”. With better parametrization and/or calibration done in the future,

this midday underestimation could be corrected.

**Minor comment:**


*P02 L050 VPD acronym has not been defined yet.*

**[Response]** It is now defined.

*SC2: P02 L 55 “large scale aerosol changes”. [optional] You could add “and long term changes in cloudiness”.*

**[Response]** It is added accordingly to the manuscript.

*P03 L 075 How come? Is it because of the large reduction in radiation under cloudy*



*sky that tends to outweigh beneficial the effect of increased diffuse light?*

**[Response]** The reduction in radiation under cloudy sky can change the radiation budget at land surface and cause a cooling effect. This effect may decrease the VPD and mitigate its stress on stomatal conductance and finally affect GPP. The cooling itself can also influence directly photosynthesis rates in the model. Therefore, in the manuscript we wrote: “The covariance of these environmental factors may also cause the GPP to increase under cloudier conditions, although not being a direct effect of diffuse light”.

*P03 L 076 My bad, explanations for my comment above are provided in the following sentences I would remove the word “Finally”, which creates confusion during the transition between the two sentences.*

**[Response]** The word “Finally” has been changed to “Lastly” to avoid confusion.

*P03 L 077 Williams et al. 2016 (year not matching the reference at the end i.e. 2014).*

**[Response]** The year has been correctly accordingly.

*P03 L 087 Le Quere et al. 2018 missing from the reference list.*

**[Response]** The reference has been added to the list.

*P05 L 143 You can maybe point the reader towards Fig 3 as well. This schematic is useful for visualizing what eq. 4 calculates. I initially misunderstood what the cumulative LAI represente. It only represents the cumulative LAI above the current layer but does not include the current layer (if I got it right).*

**[Response]** Fig 3 is cited here. And yes the cumulative LAI above the current layer but does not include the current layer.

*P05 L 152 Shouldn't it be  $dI_i / dLAI_{-c_i}$  instead of  $dI/dLAI_{-c}$  in eq. 6 What does the vertical bar | symbol represents? Is it a derivative at fixed  $LAI_{-c_i}$ ?*

**[Response]** Here  $dI/dLAI_{-c}$  indicates the derivative of light with respect to cumulative LAI from the top of the canopy. Since this equation is continuous and for all canopy position, no subscript  $i$  is added here. To calculate  $I_{abs_i}$  which is the absorption at layer  $i$ , the derivative is calculated at layer  $i$ , noted  $|LAI_{-c_i}$ . This calculation is based on the assumption that all canopy layers are thin enough to neglect the difference in light absorption within each canopy layer (explained after Eq. 6).

P06 L 170 *Either explicitly provide the relationships or give a reference where those are documented.*

**[Response]** The reference has been added to the manuscript.

P06 L 175 *“forcing datasets”: Do you mean dataset used to drive LSMs?*

**[Response]** Yes, the manuscript has been clarified to use “datasets to drive LSMs”.

P 06 L 179  $Fdf_{PAR}$  *Should it be rewritten  $fPAR_{df}$  to be consistent with the notation in other equations?*

P 07 L 216 *Same as above,  $fPPFD_{df}$  instead of  $Fdf_{PPF}$*

**[Response]** In the manuscript, we use  $Fdf_{PAR}$  or  $Fdf_{PPFD}$  to distinguish the fraction of diffuse light from the radiation variables in  $W\ m^{-2}$  or in  $\mu mol\ m^{-2}\ s^{-1}$  using subscript “ $_{df}$ ” for diffuse light (see Table 1).

P08-09 eq. 26, 28, 30. *Should it be  $LAI_{-c_i}$  instead of  $LAI_{-a_i}$  in the exponentials?*

**[Response]** Thanks for finding this error, the equations have been corrected.

P08 L 253 *Ref to Hikosaka et al. (missing from the reference list)*

**[Response]** The reference has been added to the list.

P09 L 2 68 27 0 *Same as eq. 6. The notation for the derivative is not clear to me. Could you explain?*

**[Response]** Please see the response to the above comment.

P09 L 289 *Change to “from 252 sites in total”.*

**[Response]** The manuscript has been changed accordingly.

P10 L 292 *Good job getting the references for all the sites!*

P10 L 292 *“annual climate” sounds weird. Could be rephrased by saying, “(climatological) annual mean temperature span the range xx to yy while (climatological) annual mean precipitation ... span the range” Same for Fig S2 legend.*

**[Response]** It has been rephrased as: “The annual mean temperature of the sites spans from -9 to 27°C, while the annual precipitation spans from 67 to over 3000 mm yr<sup>-1</sup>” in the text. Fig S2 legend has been changed accordingly.

P13 L 411 *This (fig 9 & 10) is an extremely useful way of presenting the sensitivity of the two models*

P14 L 422 *Could that result be related to similarities in parameter traits and optimum*

points (e.g.  $V_{cmax}$ ) between PFTs used to represent temperate and tropical biomes?


**[Response]** The ORCHIDEE model calculates the optimum points for  $V_{cmax}$  according to the growth temperature the vegetation is adapting to during the season. The range of the acclimation spans from 11 to 35°C in the current ORCHIDEE model. Therefore, the model should be capable of distinguishing temperate and tropical biomes. However, considering the limiting observation data for calibration (P16 L 499), it is possible that current parameters are not good enough to represent sufficiently well temperature acclimation.

*Figure 2 The subtle light gradient makes it hard to appreciate the density of points. Could you maybe add a Probability Density Function along the x (respectively y) axis to represent the distribution of modelled (respectively observed) fraction of PPFD.*

**[Response]** Figure 2 has been improved accordingly (also Fig R3).

*Figure 6 “is controlled the same” feels a bit clunky Could be rephrased by just saying that the sunny and cloudy days are sampled at equal light levels*

**[Response]** The caption has been modified accordingly.



## **Response to Reviewer #2**

### **General comment:**

*“This manuscript led by Zhang presented a study on improving the ORCHIDEE land surface model with specific consideration of the impacts of diffuse light fraction on vegetation photosynthesis, a well recognized phenomenon but poorly represented in the existing version of ORCHIDEE model. The new model, named after ORCHIDEE\_DF, has included a scheme for partitioning light into direct and diffuse components, and separated the existing multi-layer canopy into sunlit and shaded leaves with a two-stream radiative transfer model following Spitters 1986. Then the authors used global fluxnet observations to evaluate the new model and found that the new model better simulates GPP under different illumination conditions. Examinations on the effects of diffuse light on GPP and light use efficiency and the interactions between diffuse light and other environmental factors such as temperature and vapor pressure deficit were conducted. The new model is suggested to have great potential in investigating aerosol effect on global biogeochemical cycles.*

*Overall the manuscript is very well organized and written, and easy to read. The description of the model development is clear, and the evaluation strategy is comprehensive and convincing. The analyses sections provide insightful understanding of the interactions of diffuse light and environmental factors. I don't really have much to add, but here I provide some minor suggestions and hope they can help further improve the quality of the manuscript.”*

**[Response]** We thank the reviewer for the review and helpful comments and suggestions, which helped us to further improve our manuscript. We have addressed all the suggestions and comments in our revision. Please find below the reviewer's comments, followed by our responses and relevant changes in the manuscript. We hope that the revised version addresses all the issues and satisfies the reviewer.

### **Comments:**

*1. Line 42-43: "However, this effect remains poorly represented in current land*

*surface models". This is not accurate, at least CLM (Oleson et al., 2013), JULES (Mercado et al., 2009), CoLM (Dai et al 2004), iTem (Chen et al., 2014), and YIBs (Strada et al., 2016) have included processes that account for the diffuse light effect.*

*Oleson, K., Lawrence, D. M., Bonan, G. B., Drewniak, B., Huang, M., Koven, C. D., ... Yang, Z. -L. (2013). Technical description of version 4.5 of the Community Land Model (CLM) (No. NCAR/TN-503+STR). doi:10.5065/D6RR1W7M Mercado LM, Bellouin N, Sitch S, et al. Impact of changes in diffuse radiation on the global land carbon sink. Nature. 2009;458(7241):1014-1017. doi:10.1038/nature07949 Dai, Y., R. E. Dickinson, and Y. Wang, 2004: A Two-Big-Leaf Model for Canopy Temperature, Photosynthesis, and Stomatal Conductance. J. Climate, 17, 2281–2299 Min Chen & Qianlai Zhuang (2014) Evaluating aerosol direct radiative effects on global terrestrial ecosystem carbon dynamics from 2003 to 2010, Tellus B: Chemical and Physical Meteorology, 66:1, DOI: 10.3402/tellusb.v66.21808 Strada, S. and Unger, N.: Potential sensitivity of photosynthesis and isoprene emission to direct radiative effects of atmospheric aerosol pollution, Atmos. Chem. Phys., 16, 4213–4234, <https://doi.org/10.5194/acp-16-4213-2016>, 2016.*

*The first three have been introduced in the paragraph of Line 88-100, but latter two were directly applied for examining aerosol impacts and should be discussed as well.*

**[Response]** Thanks for the suggestion. We have added these studies to our updated manuscript: Line 42-43: “this effect remains poorly represented or evaluated in current land surface models.” Line 94-95: “This two-big-leaf scheme was further used in iTem LSM (Chen and Zhuang, 2014) and got partly inherited in later CLM models (Oleson et al., 2013).” Line 102-105: “Apart from JULES, the Yale Interactive terrestrial Biosphere model (YIBs) also included a two-stream multilayer canopy light transmission scheme, but few efforts have been made to evaluate the ability of YIBs model to capture the observed diffuse light fertilization effect, especially at sub-daily time scales (Yue and Unger, 2015).”

*2. I would suggest the authors provide a table of acronyms in Section 2.1.2 and 2.1.3 as an appendix so that the readers are easier to follow the equations.*

**[Response]** An appendix section of acronym list has been added to the manuscript.

After Line 594:

**“Appendix A**

List of acronyms:

Fdf: Fraction of diffuse radiation  
GPP: Gross Primary Production  
LAI: Leaf Area Index  
LSM: Land Surface Model  
LUE: Light Use Efficiency  
NIR: Near-Infrared Radiation  
PAR: Photosynthetically Active Radiation  
PFT: Plant Functional Type  
PPFD: Photosynthetic Photon Flux Density  
SW: downward Shortwave Radiation at the top of canopy  
TOA: Top of Atmosphere  
TOC: Top of Canopy  
VPD: Vapor Pressure Deficit

\* The variable names in Section 2 are listed in Table 1”

*3. Section 4.2 discussed factors affecting response of GPP to diffuse light and the authors suggested that the lower temperature and VPD may be the main cause of the higher midday GPP under cloudier conditions. Does ORCHIDEE simulate leaf temperature at different canopy layers? If not, it is not very convincing to me, as the short-term air temperature and VPD variations are mainly determined by the meteorological system, rather than the radiation regime.*

**[Response]** Thanks for raising this concern. In current ORCHIDEE trunk and DF model, the air temperature is taken directly as leaf temperature and does not vary

within the canopy. We agree that the short-term air temperature and VPD variations are mainly determined by the meteorological system. The explanation of leaf temperature is added to the manuscript: (Line 179-181) “Because in current ORCHIDEE, there is only one energy budget per grid cell, from which we cannot determine the leaf temperature, the air temperature is used to represent the leaf temperature in current model.”

We compared the observed Tair and VPD under cloudy and sunny conditions at midday time and found that the cloudy midday Tair and VPD is lower than the sunny ones (Fig. R4). Therefore, the lower midday temperature and VPD could be the main cause of the detected midday  $\Delta$ GPP in the manuscript. This lower cloudy midday Tair and VPD at site level might be because the time scales of weather systems which cause overcast conditions are often long enough to affect Tair. As a result, dynamics in canopy leaf temperature are not necessary to explain the simulated effect shown in the manuscript in line with the FLUXNET observations.

*4. Section 4.3. I think another important limitation of the developed ORCHIDEE\_DF model for examining aerosol impacts is that it does not consider the impacts of the changing radiation regime on leaf temperature. This might be a second-order effect, but could be potentially important as shown in Chen and Zhuang, 2014 Tellus B.*

**[Response]** Thanks for pointing out this limitation. Indeed, there remains no representation of the impacts of the changing radiation regime on leaf temperature in the current model, which may be potentially important. We have added some discussion of this point in Line 567-573 “Besides the possible bias in parameters, both ORCHIDEE trunk and ORCHIDEE\_DF lack a representation of the response of leaf temperature to radiation. Instead, the air temperature is used directly to represent the leaf temperature throughout the canopy for simulating gas exchange processes in current model. As shown by Chen and Zhuang (2014), the changes of radiation regime due to aerosols can significantly affect leaf temperature, which could potentially affect

GPP. For now, ORCHIDEE\_DF remains not capable of dealing with this response of leaf temperature. Further developments are needed for disentangling the role of leaf temperature and diffuse light on GPP”. This will be a future direction of our model development work.




Figure R4. Site distribution of (a) the Tair difference (b) the VPD difference between cloudy (diffuse light fraction >0.8) and sunny (diffuse light fraction <0.4) conditions between 11:00 and 13:00 during peak growing season.

# Modeling the impacts of diffuse light fraction on photosynthesis in ORCHIDEE (v5453) land surface model

Yuan ZHANG<sup>1, 2</sup>, Ana BASTOS<sup>3, \*</sup>, Fabienne MAIGNAN<sup>2, \*</sup>, Daniel GOLL<sup>4</sup>, Olivier BOUCHER<sup>5</sup>,  
Laurent LI<sup>1</sup>, Alessandro CESCATTI<sup>6</sup>, Nicolas VUICHARD<sup>2</sup>, Xiuzhi CHEN<sup>7</sup>, Christof AMMANN<sup>8</sup>,  
5 Altaf ARAIN<sup>9</sup>, T. Andrew BLACK<sup>10</sup>, Bogdan CHOJNICKI<sup>11</sup>, Tomomichi KATO<sup>12,13</sup>, Ivan  
MAMMARELLA<sup>14</sup>, Leonardo MONTAGNANI<sup>15,16</sup>, Olivier ROUPSARD<sup>17,18,19</sup>, Maria J SANZ<sup>20,21</sup>,  
Lukas SIEBICKE<sup>22</sup>, Marek URBANIAK<sup>11</sup>, Francesco Primo VACCARI<sup>23</sup>, Georg WOHLFAHRT<sup>24</sup>, Will  
WOODGATE<sup>25,26</sup>, Philippe CIAIS<sup>2</sup>

<sup>1</sup>Laboratoire de Météorologie Dynamique, IPSL, Sorbonne Université/CNRS, Paris, France

10 <sup>2</sup>Laboratoire des Sciences du Climat et de l'Environnement (LSCE), IPSL, CEA/CNRS/UVSQ, Gif sur Yvette, France

<sup>3</sup>Department of Geography, Ludwig-Maximilian University of Munich, Munich, Germany

<sup>4</sup>Department of Geography, University of Augsburg, Augsburg, Germany

<sup>5</sup>Institut Pierre-Simon Laplace, CNRS/Sorbonne Université Paris, France

<sup>6</sup>Institute for Environment and Sustainability, Joint Research Centre, European Commission, Ispra, Italy

15 <sup>7</sup>Guangdong Province Key Laboratory for Climate Change and Natural Disaster Studies, School of Atmospheric Sciences, Sun  
Yat-sen University, Guangzhou 510275, China

<sup>8</sup>Climate and Agriculture Group, Agroscope, Zürich, 8046, Switzerland

<sup>9</sup>School of Geography and Earth Sciences and McMaster Centre for Climate Change, McMaster University, Hamilton, ON,  
Canada

20 <sup>10</sup>Faculty of Land and Food Systems, University of British Columbia, Vancouver, BC, Canada

<sup>11</sup>Poznan University of Life Sciences. Piatkowska 94, 60-649 Poznan, Poland

<sup>12</sup>Research Faculty of Agriculture, Hokkaido University, Sapporo 060-8589, Japan

<sup>13</sup>Global Institution for Collaborative Research and Education, Hokkaido University, Sapporo 060-8589, Japan

<sup>14</sup>Institute for Atmospheric and Earth System Research/Physics, Faculty of Sciences, University of Helsinki, Finland

25 <sup>15</sup>Autonomous Province of Bolzano, Forest Services, Via Brennero 6, Bolzano 39100, Italy

<sup>16</sup>Faculty of Science and Technology, Free University of Bolzano, Piazza Università 5, Bolzano 39100, Italy

<sup>17</sup>CIRAD, UMR Eco&Sols, BP1386, CP18524, Dakar, Senegal

<sup>18</sup>Eco&Sols, Univ Montpellier, CIRAD, INRAE, IRD, Montpellier SupAgro, Montpellier, France

<sup>19</sup>LMI IESOL, Centre IRD-ISRA de Bel Air, BP1386, CP18524 Dakar, Senegal

30 <sup>20</sup>Basque Centre for Climate Change, Sede Building 1, Scientific Campus of the University of the Basque Country, 48940,  
Leioa, Spain

<sup>21</sup>Ikerbasque, Basque Science Foundation, 48013 Bilbao, Spain

<sup>22</sup>University of Göttingen, Bioclimatology, Büsgenweg 2, 37077 Göttingen, Germany

<sup>23</sup>Institute of BioEconomy, National Research Council, 50145 Firenze, Italy

35 <sup>24</sup>Department of Ecology, University of Innsbruck, Innsbruck, Austria

<sup>25</sup>CSIRO Land & Water, Canberra, ACT, Australia

<sup>26</sup>School of Earth and Environmental Sciences, University of Queensland, St Lucia, 4067, Qld, Australia

*\*Contribute equally*

Correspondence to: Yuan ZHANG (yuan.zhang@lmd.jussieu.fr)

**Abstract.** Aerosol and cloud-induced changes in diffuse light have important impacts on the global land carbon cycle by changing light distribution and photosynthesis in vegetation canopies. However, this effect remains poorly represented or evaluated in current land surface models. Here we add a light partitioning module and a new canopy light transmission module to the ORCHIDEE land surface model (trunk version, v5453) and use the revised model, ORCHIDEE\_DF, to estimate the fraction of diffuse light and its effect on gross primary production (GPP) in a multi-layer canopy. We evaluate the new parameterizations using flux observations from 159 eddy covariance sites over the globe. Our results show that compared to the original model, ORCHIDEE\_DF improves the GPP simulation under sunny conditions and captures the observed higher photosynthesis under cloudier conditions in most plant functional types (PFTs). Our results also indicate that the larger GPP under cloudy conditions compared to sunny conditions is mainly driven by increased diffuse light in the morning and in the afternoon, and by decreased vapor pressure deficit (VPD) and air temperature at midday. The observations show strongest positive effects of diffuse light on photosynthesis are found in the range 5-20 °C and VPD<1 kPa. This effect is found to decrease when VPD becomes too large, or temperature falls outside that range likely because of increasing stomatal resistance to leaf CO<sub>2</sub> uptake. ORCHIDEE\_DF underestimates the diffuse light effect at low temperature in all PFTs and overestimates this effect at high temperature and high VPD in grasslands and croplands. The new model has the potential to better investigate the impact of large-scale aerosol changes and long term changes in cloudiness on the terrestrial carbon budget, both in the historical period and in the context of future air quality policies and/or climate engineering.

## 1 Introduction

Process-based Land Surface Models (LSMs), which simulate the water and energy balance, and biogeochemical processes on land, have been widely used to attribute past changes in carbon (C) fluxes (Piao et al., 2009; Sitch et al., 2013) and to project the future land C budget (Ciais et al., 2013). Despite being useful and widely applied tools, large uncertainties are a limitation of LSMs (Sitch et al., 2008). One of the sources of the uncertainties is the omission or oversimplification of important processes that affect primary production. For instance, the impacts of light quality on photosynthesis is not currently represented in most LSMs, limiting the possibility to predict the variability of the carbon budget driven by changes in the atmospheric aerosol load which may be triggered by volcanic eruptions or variation in air pollution levels.

It has been found by in situ observations that under the same light level, the increase of diffuse light fraction can enhance light use efficiency and ultimately photosynthesis, or gross primary production (GPP) (Gu et al., 2003; Niyogi et al., 2004; Misson et al., 2005; Alton, 2007a; Knohl and Baldocchi, 2008; Mercado et al. 2009; Oliphant et al., 2011; Kanniah et al., 2013; Williams et al., 2014; Cheng et al., 2015; Wang et al., 2018). Several mechanisms explaining this GPP enhancement have been proposed and tested. First, the more isotropic nature of diffuse light means that it penetrates deeper into the canopy to become available for photosynthesis of the lower canopy leaves, which would otherwise be shaded and light limited (Roderick et al., 2001; Urban et al, 2012). Second, the multi-directionality of diffuse light produces a more homogeneous distribution of radiation between sunlit and shaded leaves, enhancing the photosynthesis of upper canopy shaded leaves and limiting the waste

of energy in light-saturated sunlit leaves (Li et al., 2014; Williams et al., 2014). Third, higher diffuse light fraction is often accompanied with less stressing temperature and vapor pressure deficit (VPD) for photosynthesis. The covariance of these environmental factors may also cause the GPP to increase under cloudier conditions, although not being a direct effect of diffuse light (Gu et al., 2002; Cheng et al., 2015; Li et al., 2014). ~~Finally~~Lastly, plant LAI (leaf area index, the area of leaves per unit land area) maximum may get acclimated to the cloudier seasons, which also contributes to higher GPP (Williams et al., 2016).

Currently, most process-based LSMs simulate leaf photosynthesis using equations and parameterizations derived from Farquhar et al. (1980) with different formulations of stomatal conductance, usually with stomatal closure under high VPD or low relative humidity (Ball et al., 1987; Yin et al., 2009; Medlyn et al., 2011). These parameterizations calculate photosynthesis per unit LAI considering the stress from temperature, VPD and soil water, and then integrate it over the entire canopy volume. Therefore, the effects of temperature and VPD change under cloudier conditions have been usually implicitly considered in current LSMs (e.g. Zhang et al., 2019). However, for the sake of simplicity and computational efficiency and for the lack of diffuse light fraction data, most global LSMs assumed a single extinction coefficient for both direct and diffuse light (Sellers et al., 1997; Sitch et al., 2008). These LSMs are therefore incapable to investigate the effect of diffuse light fraction changes on photosynthesis. This limit of LSMs is thought to cause considerable underestimation of land C sink after the eruption of Mount Pinatubo (le Quere et al., 2018).

To better simulate the diffuse light impacts, several earlier works have developed photosynthesis models that considers different light transmission of diffuse and direct radiation (Spitters, 1986; Leuning et al., 1995; de Pury and Farquhar, 1997). Based on these models, a few studies have tried to address the influence of light quality on GPP in LSMs. Dai et al. (2004) introduced a two-big-leaf canopy model to simulate the effects of diffuse and direct radiation in the Common Land Model (CLM 2L). ~~However, this model~~This two-big-leaf scheme was further used in iTem LSM (Chen and Zhuang, 2014) and got partly inherited in later CLM models (Oleson et al., 2013). However, this light transmission scheme assumes a single-layer canopy and can therefore not simulate the vertical profile of leaf traits. A multilayer canopy model is more suitable to represent the vertical heterogeneity of leaf traits and radiation transfer (Alton et al., 2007b; Bonan et al., 2012). Differentiating sunlit and shaded leaves in a multilayer canopy LSM was firstly considered in the Joint UK Land Environment Simulator (JULES) LSM (Alton et al., 2007a; Mercado et al., 2009). Using this version of JULES, Mercado et al. (2009) investigated the diffuse light effect and suggested that diffuse light fraction change enhanced the global land C sink during the 1960-1999 period by about a quarter. However, Mercado et al. (2009)'s model was only tested at two forest sites which cannot represent well global terrestrial ecosystems. Thus, there remains need to obtain well-evaluated LSMs that distinguish diffuse and direct light to test the results of Mercado et al. (2009), and to further investigate the diffuse radiation effect of aerosols. Apart from JULES, the Yale Interactive terrestrial Biosphere model (YIBs) also included a two-stream multilayer canopy light transmission scheme, but few efforts have been made to evaluate the ability of YIBs model to capture the observed diffuse light fertilization effect, especially at sub-daily time scales (Yue and Unger, 2015).

Here we introduce a modified version of the LSM ORCHIDEE (Organizing Carbon and Hydrology In Dynamic Ecosystems, Krinner et al., 2005), referred to as ORCHIDEE\_DF, which uses a semi-empirical method to calculate the fraction of diffuse light (Weiss and Norman, 1985), and a process-based multilayer canopy light transmission model to simulate the effects of diffuse light fraction on photosynthesis (Spitters, 1986). We evaluated the GPP simulated by ORCHIDEE\_DF and the same version of the ORCHIDEE code without diffuse light (trunk version, v5453) using observations collected from 159 eddy covariance flux sites over 11 plant functional types (PFT) (Baldocchi et al., 2001). Using both model simulations and observations at the flux sites, we also investigated the interactions between diffuse light fraction and biotic and abiotic factors on GPP, with the objective of understanding when and how much does light quality affect photosynthesis. Because diffuse light is expected to enhance photosynthesis of shaded leaves in deep canopy, we would also test whether the enhancement of GPP due to diffuse radiation is larger in canopies with larger LAI and whether environmental factors such as temperature or VPD affect this enhancement from diffuse light.

## 2 Data and Methods

### 2.1 Model description

#### 2.1.1 Canopy light transmission and photosynthesis in the ORCHIDEE trunk

The ORCHIDEE\_DF model is based on ORCHIDEE trunk version 5453 (updated in September 2018). A general description of the physical processes related to energy and water balance, vegetation dynamics and biogeochemical processes in ORCHIDEE can be found in Krinner et al. (2005). The ORCHIDEE trunk version 5453 brings a number of improvements, and photosynthesis parameters were recently re-calibrated against FLUXNET data (Baldocchi et al., 2001) and atmospheric CO<sub>2</sub> observations for the IPSL Earth System Model (IPSL-CM6) and the CMIP6 simulations.

The leaf-scale photosynthesis calculation in the ORCHIDEE trunk is based on the scheme of Yin and Struik (2009). This scheme is an adaptation of the biophysical model of Farquhar et al. (1980) with a specific parameterization of stomatal conductance. The Farquhar et al. model calculates assimilation ( $A$ ) as the minimum of the Rubisco-limited rate of CO<sub>2</sub> assimilation ( $A_c$ ) and the electron transport-limited rate of CO<sub>2</sub> assimilation ( $A_j$ ):

$$A = \min\{A_c, A_j\} \quad (1)$$

Here  $A_c$  is mainly affected by the maximum carboxylation capacity of Rubisco ( $V_{cmax}$ ), which is temperature dependent (Yin and Struik., 2009), and the CO<sub>2</sub> concentration at the carboxylation site ( $C_c$ ):

$$A_c = \frac{(C_c - \Gamma^*) V_{cmax}}{C_c + K_m C (1 + O/K_m O)} - R_d \quad A_c = \frac{(C_c - \Gamma^*) V_{cmax}}{C_c + K_m C (1 + O/K_m O)} - R_d \quad (2)$$

where  $\Gamma^*$  is the CO<sub>2</sub> compensation point in the absence of dark respiration ( $R_d$ ).  $K_m C$  and  $K_m O$  are the Michaelis-Menten constants for CO<sub>2</sub> and O<sub>2</sub>,  $O$  is the O<sub>2</sub> concentration at the carboxylation site.

带格式的: 下标

带格式的: 下标

带格式的: 下标

带格式的: 下标

带格式的: 下标

带格式的: 下标

带格式的: 下标

$A_i$  is calculated as a function of  $C_c$  and electron transport rate ( $J$ ):

$$A_i = \frac{J(C_c - \Gamma^*)}{4.5C_c + 10.5\Gamma^*} - R_d \quad A_i = \frac{J(C_c - \Gamma^*)}{4.5C_c + 10.5\Gamma^*} - R_d \quad (3)$$

Here  $J$  is determined by a temperature-dependent maximum electron transport rate ( $J_{max}$ ) and the photosynthetic photons absorbed by leaves, calculated following Yin and Struik (2009). Due to the attenuation of photosynthetically active radiation (PAR) with depth in the canopy,  $J$  also varies vertically. In addition, to account for the distribution of light and maximize the assimilation, plants tend to allocate nitrogen unevenly in the canopy profile (Niinemets et al., 1997; Meir et al., 2002), resulting in a vertical gradient in enzyme concentration and consequently in  $V_{cmax}$  and  $J_{max}$ . The vertical heterogeneity of canopy photosynthetic properties highlights the need to represent the canopy in a multilayer way.

In order to simulate the vertical transmission and absorption of light within the canopy, ORCHIDEE trunk uses a multilayer canopy with a big leaf approximation in each layer. The canopy is geometrically divided into up to a maximum number of 20 layers depending on the leaf area index (LAI). The discretization is represented in Fig. 1a and the LAI at the interface of the layers are given by:

$$LAI_{-c_i} = 12 \times \frac{e^{0.15 \times (i-1)} - 1}{e^{0.15 \times 20} - 1} \quad (4)$$

where  $LAI_{-c_i}$  is the cumulative LAI above layer  $i$ , ( $1 \leq i \leq 20$ ) and the layers are numbered from top to bottom: (Fig. 3). It should be noted that 20 layers are only for canopies with total LAI larger than 12. The number of layers decreases with total LAI. For instance, if the LAI is 2, only the first 10 layers are used to calculate the light distribution and photosynthesis (Fig. 1a).

Light transmission in the multilayer canopy is calculated using the Beer-Lambert law (Monsi and Saeki, 1953) without distinguishing direct and diffuse light. The downward shortwave radiation arriving at the top of canopy (TOC) layer  $i$  ( $I_i$ ) is:

$$I_i = I_0 e^{-k \times LAI_{-c_i}} \quad (5)$$

where  $k$  is the light extinction coefficient, taken equal to 0.5.  $I_0$  is the TOC downward shortwave radiation ( $W m^{-2}$ ).

Because the radiation attenuation between one layer and the one just below is assumed to be due to leaf absorption, the absorbed radiation per leaf area at the top of layer  $i$  ( $Iabs_i$ ) can be estimated as in Saeki (1960):

$$Iabs_i = \frac{-dI}{dLAI_{-c}} | LAI_{-c_i} = k I_0 e^{-k LAI_{-c_i}} \quad (6)$$

Here we assume that all canopy layers are thin enough to neglect the difference in light absorption within each canopy layer. i.e. the absorbed radiation does not attenuate within each canopy layer and  $Iabs_i$  is used for all leaves in layer  $i$ .

It should be noted that the radiation considered to calculate the  $J$  term in Eq. (3) is not shortwave radiation in  $W.m^{-2}$  but photosynthetic photon flux density (PPFD) in  $\mu mol m^{-2} s^{-1}$ . A translation from  $Iabs_i$  to the absorbed PPFD per leaf area in canopy layer  $i$  ( $PPFDabs_i$ ) is thus needed. Currently, there is no standard definition of the wavelength range for shortwave radiation (e.g. Howell et al, 1982; Zhang et al., 2004; Chen et al., 2012). In ORCHIDEE trunk, shortwave radiation in  $W m^{-2}$

带格式的: 下标

带格式的: 下标

带格式的: 字体: 倾斜

is multiplied by a factor of 0.5 to calculate photosynthetically active radiation (PAR) in  $\text{W m}^{-2}$ , and then a quanta-to-energy ratio of  $4.6 \text{ mmol J}^{-1}$  is used to convert PAR into PPFD in  $\mu\text{mol m}^{-2} \text{ s}^{-1}$ .

ORCHIDEE accounts for a vertical gradient in enzyme concentration in canopy.  $V_{\text{cmax}}$  and  $J_{\text{max}}$  are assumed in the model to be linearly related to photosynthetically active leaf nitrogen concentration (per leaf area) (Kattge et al. 2007). Meir et al. (2002) found a decreasing leaf nitrogen concentration, as well as  $V_{\text{cmax}}$  and  $J_{\text{max}}$  with increasing canopy depth in different ecosystems, suggesting an acclimation of plants to maximize photosynthesis in a canopy with unevenly distributed radiation. ORCHIDEE trunk lacks an explicit model of dynamic N allocation to leaves in the canopy, instead, it uses an empirical relationship to represent the impact of leaf nitrogen concentration on  $V_{\text{cmax}}$  and  $J_{\text{max}}$  using the vertical profile of radiation:

$$V_{\text{cmax}_i} = V_{\text{cmax}_0} \left(1 - 0.7 \times \left(1 - \frac{I_i}{I_0}\right)\right) \quad (7)$$

$$J_{\text{max}_i} = J_{\text{max}_0} \left(1 - 0.7 \times \left(1 - \frac{I_i}{I_0}\right)\right) \quad (8)$$

It should be noted that in ORCHIDEE trunk, the leaf-scale assimilation variables (e.g.  $V_{\text{cmax}}$ ) are also affected by the instantaneous ~~air~~leaf temperature and the temperature of the last month which plants have adapted to (Kattge and Knorr, 2007).

Because in current ORCHIDEE, there is only one energy budget per grid cell, from which we cannot determine the leaf temperature, the air temperature is used to represent the leaf temperature in current model. The calculation of  $C_a$  depends on VPD and also on whether the vegetation follows the C3 or C4 photosynthesis pathway (Yin and Struik, 2009). For simplicity, the near surface air temperature and humidity are used for the calculation of assimilation in all canopy layers. Furthermore, there are 13 PFTs in ORCHIDEE (Table S1) and  $V_{\text{cmax}}$  and  $J_{\text{max}}$  are PFT-dependent.

带格式的: 下标

## 2.1.2 Light partitioning in ORCHIDEE\_DF

The lack of light quality (diffuse light fraction) information in most forcing datasets is one of the main difficulties when simulating the diffuse light effect. Here we partition the half hourly downward PAR, which can be derived from the shortwave radiation, into diffuse and direct components following the Weiss and Norman (1985) empirical equations. Compared with another empirical method (Spitters et al., 1986), we found that this method reproduces better the observed diffuse light fraction at the flux sites used in this study (Fig. 2, Fig. S1). The diffuse PAR fraction ( $Fdf_{\text{PAR}}$ ) above the canopy is estimated as:

$$Fdf_{\text{PAR}} = 1 - \frac{PAR_{\text{pot,direct}}}{PAR_{\text{pot}}} \left(1 - \left(\frac{a-R}{b}\right)^{\frac{1}{n}}\right) \quad (9)$$

where  $PAR_{\text{pot}}$  and  $PAR_{\text{pot,direct}}$  are the potential total and direct PAR, i.e. the total and direct PAR which would arrive at land surface under clear sky conditions.  $a$  and  $b$  are parameters, which take values of 0.9 and 0.7, and  $R$  is the ratio of observed to potential total downward shortwave radiation ( $SW_{\text{obs}}$  and  $SW_{\text{pot}}$ ) reaching the top of the canopy:

$$R = \frac{SW_{\text{obs}}}{SW_{\text{pot}}} \quad (10)$$

The potential downward shortwave radiation consists of potential downward PAR (visible, range 0.4–0.7 μm) and potential downward near-infrared radiation (NIR, range 0.7–5 μm). Also the potential PAR and NIR are the sum of direct ( $PAR_{p,dir}$ ,  $NIR_{p,dir}$ ) and diffuse ( $PAR_{p,dif}$ ,  $NIR_{p,dif}$ ) components, given by:

$$SW_p = PAR_p + NIR_p = PAR_{p,dir} + PAR_{p,dif} + NIR_{p,dir} + NIR_{p,dif} \quad (11)$$

A simple atmospheric light transfer model modified from Weiss and Norman (1985) is used to estimate potential radiation. The potential direct PAR,  $PAR_{p,dir}$ , is calculated as:

$$PAR_{p,dir} = PAR_{TOA} e^{-0.185(p/p_0)^m \cos \theta} \quad (12)$$

where  $PAR_{TOA}$  is the PAR at top of atmosphere (TOA),  $p$  and  $p_0$  indicate the local and standard sea level air pressure,  $m$  is the optical air mass, calculated using the solar zenith angle  $\theta$ :

$$m = (\cos \theta)^{-\frac{1}{2}} \quad (13)$$

The potential diffuse TOC PAR,  $PAR_{p,dif}$ , is assessed as:

$$PAR_{p,dif} = 0.4(PAR_{TOA} \cos \theta - PAR_{p,dir}) \quad (14)$$

which expresses that 40% of the PAR flux that is extinguished in the atmosphere through scattering and absorption is available as diffuse PAR at the surface. Similarly, the potential direct and diffuse NIR at the top of the canopy ( $NIR_{p,dir}$  and  $NIR_{p,dif}$  respectively), can be estimated as:

$$NIR_{p,dir} = (NIR_{TOA} e^{-0.06(p/p_0)^m} - \omega) \cos \theta \quad (15)$$

$$NIR_{p,dif} = 0.6(NIR_{TOA} \cos \theta - NIR_{p,dir} - \omega \cos \theta) \quad (16)$$

where  $\omega$  is a flux term accounting for atmospheric water vapor absorption, calculated as a function of the solar constant ( $SC$ , in  $Wm^{-2}$ ) and  $m$ :

$$\omega = SC \times 10^{(-1.195 + 0.1159 \log_{10} m - 0.0215 (\log_{10} m)^2)} \quad (17)$$

Using the results from Eqs. (12, 14, 15 and 16), we are able to calculate the  $SW_p$  to obtain the value of  $R$  in Eq. (10).

It should be noted that the quanta-to-energy ratio (in  $mmol J^{-1}$ ) is different under different sky conditions, because atmospheric scattering varies spectrally with the air mass and the cloud amount (Dye, 2004). For this consideration, the calculation of PPFD from PAR in ORCHIDEE\_DF uses the observation-oriented empirical equations from Dye (2004):

$$\beta_e = 4.576 - 0.03314 Fdf_{PAR} \quad (18)$$

$$\beta_{dif} = \frac{4.5886 Fdf_{PAR}}{0.010773 + Fdf_{PAR}} \quad (19)$$

where the  $\beta_e$  is the quanta-to-energy ratio for the total PAR ( $PAR_e$ ) at the top of the canopy, while  $\beta_{dif}$  is for its diffuse component ( $PAR_{dif}$ ):

$$PPFD_e = \beta_e PAR_e \quad (20)$$

$$PPFD_{dif} = \beta_{dif} PAR_{dif} \quad (21)$$

The diffuse PPFD fraction ( $Fdf_{PPFD}$ ) can thus be calculated as:

$$Fdf_{PPFD} = \frac{PPFD_{dif}}{PPFD_e} = \frac{\beta_{dif}}{\beta_e} Fdf_{PAR} \quad (22)$$

### 2.1.3 Canopy light transmission in ORCHIDEE\_DF

In ORCHIDEE\_DF, we use the same stratification of canopy as in the trunk version (Eq. (4)). But for the light transmission, we use a two-stream radiative transfer model [with direct and diffuse radiation treated separately](#) following Spitters (1986). For convenience, we use radiation and  $I$  in this section to refer to the PPFD derived from the light partitioning step- [\(see section 2.1.3\)](#).

An assumption of the model is that leaves are bi-Lambertian surfaces for radiation, i.e. the reflection and transmission are isotropic. This reflection and transmission are together referred to as leaf scattering. This assumption implies that once direct radiation encounters a leaf, it gets either absorbed or scattered as diffuse light. While for diffuse radiation, the scattered light remains diffuse. The scattering coefficient,  $\sigma$ , is assumed equal to 0.2 following Spitters (1986), meaning 20% of the light encountering a leaf is scattered (80% is absorbed).

Based on this assumption, the radiation penetrating the canopy can be divided into three components (Fig. 3): the direct light which has not been intercepted by leaves ( $I_{dr,dr}$ ), the diffuse light generated by leaf scattering of intercepted direct light ( $I_{dr,df}$ ), and the diffuse light in the canopy provided by the TOC diffuse radiation ( $I_{df}$ ). It should be noted that the diffuse light generated by multiple times of scattering of the direct light is grouped into  $I_{dr,df}$ , while those from the scattering of TOC diffuse radiation belong to  $I_{df}$  (Fig. 3). The sum of  $I_{dr,dr}$  and  $I_{dr,df}$  hereafter noted as  $I_{dr}$  represents the total radiation in each canopy layer derived from the TOC direct radiation, hereafter  $I_{dr,0}$ .

If we also consider direct radiation as parallel beams, only the first leaves on the way of direct light can absorb  $I_{dr,dr}$ . These leaves are referred to as sunlit leaves. The fraction of sunlit leaves in each canopy layer can be calculated by applying Beer-Lambert law using an extinction coefficient for opaque, non-reflective “black” leaves (Fig. 1b):

$$LAI f_{sun,i} = e^{-k_b LAI_{-c,i}} \quad (239)$$

here  $LAI f_{sun,i}$  is the fraction of sunlit LAI in canopy layer  $i$ .  $LAI_{-c,i}$  is the cumulative LAI in Eq. (4).  $k_b$  is the extinction coefficient if the leaves are assumed “black”. A function of  $\theta$ , leaf angle distribution index ( $LA$ ) and leaf clumping index ( $LC$ ) is used to represent the geometry between the direct radiation and leaves:

$$k_b = \frac{LA \cdot LC}{\cos \theta} \quad (2410)$$

For spherically distributed leaves,  $LA$  equals 0.5 (Goudriaan, 1977; Bodin and Franklin, 2012).  $LC$  is defined as in Myneni et al. (1989) and Baldocchi and Wilson (2001), varying between 0 and 1. Here we use the value 0.85 instead of 0.84 as recommended by an observationally-based study (Baldocchi and Wilson, 2001).

The leaves which cannot absorb  $I_{dr,dr}$  are referred to as shaded leaves. The fraction of shaded LAI in canopy layer  $i$  ( $LAI f_{shd,i}$ ) is thus the complement of  $LAI f_{sun,i}$ :

$$LAI f_{shd,i} = 1 - LAI f_{sun,i} \quad (2511)$$

Because  $I_{dr,dr}$  is assumed not to be transmitted as direct radiation through leaves,  $I_{dr,dr,i}$ , which represents  $I_{dr,dr}$  at layer  $i$  can be calculated similarly as in Eq. (239) using the downward direct radiation at the top of the canopy ( $I_{dr,0}$ ):

$$I_{dr,dr,i} = I_{dr,0} e^{-k_b LAI - \rho_i} \quad (26)$$

The transmission of  $I_{dr,df}$  is difficult to estimate directly. Here we calculate it as the difference between  $I_{dr}$  and  $I_{dr,dr}$  in each layer:

$$I_{dr,df,i} = I_{dr,i} - I_{dr,dr,i} \quad (27)$$

where  $I_{dr,df,i}$  and  $I_{dr,i}$  represent net (downward minus upward)  $I_{dr,df}$  and net  $I_{dr}$  at layer  $i$ , respectively.

The calculation of  $I_{dr,i}$  is based on Goudriaan (1982) and Hikosaka et al. (2016) under the assumptions that there is no difference in optical traits between leaves from different canopy layers and that the canopy is deep enough to neglect the reflection of the soil:

$$I_{dr,i} = (1 - \rho) I_{dr,0} e^{-\sqrt{1-\sigma} k_b LAI - \rho_i} \quad (28)$$

where  $\rho$  indicates the canopy reflection coefficient (i.e., the ratio between the TOC downward and upward radiation), calculated as:

$$\rho = \left( \frac{1 - \sqrt{1 - \sigma}}{1 + \sqrt{1 - \sigma}} \right) \left( \frac{2}{1 + 1.6 \cos \theta} \right) \quad (29)$$

In contrast to the direct light transmission, the diffuse light will not change its directional characteristics when scattered by leaves. Similar to Eq. (5), net  $I_{df}$  at canopy layer  $i$  ( $I_{df,i}$ ) can be estimated using TOC downward diffuse radiation ( $I_{df,0}$ ) in a Beer-Lambert equation:

$$I_{df,i} = (1 - \rho) I_{df,0} e^{-k_d LAI - \rho_i} \quad (30)$$

where  $k_d$  is the light extinction coefficient for diffuse light, calculated following Spitters (1986) as:

$$k_d = 0.8 \sqrt{1 - \sigma} \quad (31)$$

Similar to Eq. (6), the flux of light that is absorbed per canopy leaf area in layer  $i$  from  $I_{df}$  ( $I_{abs,df,i}$ ),  $I_{dr}$  ( $I_{abs,dr,i}$ ), and  $I_{dr,dr}$  ( $I_{abs,dr,dr,i}$ ) can be written respectively as:

$$I_{abs,df,i} = \frac{-dI_{df}}{dLAI_c} | LAI_c = k_d I_{df,i} \quad (32)$$

$$I_{abs,dr,i} = \frac{-dI_{dr}}{dLAI_c} | LAI_c = \sqrt{1 - \sigma} k_b I_{dr,i} \quad (33)$$

$$I_{abs,dr,dr,i} = \frac{-dI_{dr,dr}}{dLAI_c} | LAI_c = k_b I_{dr,dr,i} \quad (34)$$

The  $I_{dr,df}$  absorbed per canopy leaf area by layer  $i$  ( $I_{abs,dr,df,i}$ ) is:

$$I_{abs,dr,df,i} = I_{abs,dr,i} - I_{abs,dr,dr,i} \quad (35)$$

It should be noted that all leaves can absorb diffuse radiation. Therefore Eq. (32) and Eq. (35) also represent the absorption

of  $I_{df}$  and  $I_{dr,df}$  at the leaf scale. However,  $I_{dr,dr}$  is only absorbed by sunlit leaves, thus the absorption of  $I_{dr,dr}$  per sunlit leaf

area does not equal to  $Iabs_{dr,dr,i}$ , which is the average at canopy scale. Instead, because  $I_{dr,dr}$  does not change its intensity, the absorption of  $I_{dr,dr}$  per sunlit leaf area can be written as:

$$Iabs_{dr,dr,i,sun} = (1 - \sigma)k_b I_{dr,0} \quad (3622)$$

We have assumed that shaded leaves can only absorb diffuse light. Then, the radiation absorbed (per leaf area) by shaded leaves layer  $i$  ( $I_{shd,i,abs}$ ) is:

$$Iabs_{shd,i} = Iabs_{df,i} + Iabs_{dr,df,i} \quad (3723)$$

The sunlit leaves also absorb the direct light besides diffuse light. The radiation received by sunlit leaves can thus be calculated as:

$$Iabs_{sun,i} = Iabs_{shd,i} + Iabs_{dr,dr,i,sun} \quad (3824)$$

Apart from light transmission, all other parameters (e.g.  $Vcmax$ ,  $Jmax$ ) in ORCHIDEE\_DF are kept the same as in ORCHIDEE trunk.

### 2.1.3 Light partitioning in ORCHIDEE\_DF

The lack of light quality (diffuse light fraction) information in most datasets to drive LSMs is one of the main difficulties when simulating the diffuse light effect. This field can be calculated in atmospheric light transmission models when aerosol and cloud information is available (Yue et al., 2017; Malavelle et al., 2019). However, the aerosol and cloud information is not always available. Here we use the empirical equations following Weiss and Norman (1985) to partition the half-hourly downward PAR, which can be derived from the shortwave radiation, into diffuse and direct components. Compared with another empirical method (Spitters et al., 1986), we found that this method reproduces better the observed diffuse light fraction at the flux sites used in this study (Fig. 2, Fig. S1). The diffuse PAR fraction ( $Fdf_{PAR}$ ) above the canopy is estimated as:

$$Fdf_{PAR} = 1 - \frac{PAR_{p,dr}}{PAR_p} \left(1 - \left(\frac{a-R}{b}\right)^{\frac{2}{3}}\right) \quad (25)$$

where  $PAR_p$  and  $PAR_{p,dr}$  are the potential total and direct PAR, i.e. the total and direct PAR which would arrive at land surface under clear sky conditions.  $a$  and  $b$  are parameters, which take values of 0.9 and 0.7, and  $R$  is the ratio of observed to potential total downward shortwave radiation ( $SW_{obs}$  and  $SW_p$ ) reaching the top of the canopy:

$$R = \frac{SW_{obs}}{SW_p} \quad (26)$$

The potential downward shortwave radiation consists of potential downward PAR (visible, range 0.4-0.7 $\mu$ m) and potential downward near-infrared radiation (NIR, range 0.7-5 $\mu$ m). Also the potential PAR and NIR are the sum of direct ( $PAR_{p,dr}$ ,  $NIR_{p,dr}$ ) and diffuse ( $PAR_{p,df}$ ,  $NIR_{p,df}$ ) components, given by:

$$SW_p = PAR_p + NIR_p = PAR_{p,dr} + PAR_{p,df} + NIR_{p,dr} + NIR_{p,df} \quad (27)$$

A simple atmospheric light transfer model modified from Weiss and Norman (1985) is used to estimate potential radiation.

The potential direct PAR,  $PAR_{p,dr}$  is calculated as:

$$PAR_{p,dr} = PAR_{TOA} e^{-0.185(p/p_0)^m} \cos \theta \quad (28)$$

where  $PAR_{TOA}$  is the PAR at top of atmosphere (TOA),  $p$  and  $p_0$  indicate the local and standard sea level air pressure,  $m$  is the optical air mass, calculated using the solar zenith angle  $\theta$ :

$$m = (\cos \theta)^{-1} \quad (29)$$

325 The potential diffuse TOC PAR,  $PAR_{p,df}$  is assessed as:

$$PAR_{p,df} = 0.4(PAR_{TOA} \cos \theta - PAR_{p,dr}) \quad (30)$$

which expresses that 40% of the PAR flux that is extinguished in the atmosphere through scattering and absorption is available as diffuse PAR at the surface. Similarly, the potential direct and diffuse NIR at the top of the canopy ( $NIR_{p,dr}$  and  $NIR_{p,df}$  respectively), can be estimated as:

330  $NIR_{p,dr} = (NIR_{TOA} e^{-0.06(p/p_0)m} - \omega) \cos \theta \quad (31)$

$$NIR_{p,df} = 0.6(NIR_{TOA} \cos \theta - NIR_{p,dr} - \omega \cos \theta) \quad (32)$$

where  $\omega$  is a flux term accounting for atmospheric water vapor absorption, calculated as a function of the solar constant ( $SC$ , in  $Wm^{-2}$ ) and  $m$ :

$$\omega = SC \times 10^{(-1.195 + 0.4459 \log_{10} m - 0.0345 (\log_{10} m)^2)} \quad (33)$$

335 Using the results from Eqs. (28, 30, 31 and 32), we are able to calculate the  $SW_p$  to obtain the value of  $R$  in Eq. (26).

It should be noted that the quanta-to-energy ratio (in  $mmol J^{-1}$ ) is different under different sky conditions, because atmospheric scattering varies spectrally with the air mass and the cloud amount (Dye, 2004). For this consideration, the calculation of PPFD from PAR in ORCHIDEE\_DF uses the observation-oriented empirical equations from Dye (2004):

$$\beta_t = 4.576 - 0.03314 Fdf_{PAR} \quad (34)$$

340  $\beta_{df} = \frac{4.5886 Fdf_{PAR}}{0.010773 + Fdf_{PAR}} \quad (35)$

where the  $\beta_t$  is the quanta-to-energy ratio for the total PAR ( $PAR_t$ ) at the top of the canopy, while  $\beta_{df}$  is for its diffuse component ( $PAR_{df}$ ):

$$PPFD_t = \beta_t PAR_t \quad (36)$$

$$PPFD_{df} = \beta_{df} PAR_{df} \quad (37)$$

345 The diffuse PPFD fraction ( $Fdf_{PPFD}$ ) can thus be calculated as:

$$Fdf_{PPFD} = \frac{PPFD_{df}}{PPFD_t} = \frac{\beta_{df}}{\beta_t} Fdf_{PAR} \quad (38)$$

## 2.2 Flux data and site level simulations

To evaluate ORCHIDEE\_DF, we collected flux site measurements from the La Thuile dataset, which includes 965 site-year observations from ~~in total~~ 252 sites in total (<https://fluxnet.fluxdata.org/data/la-thuile-dataset/>). Because our ORCHIDEE

350 simulations assume that the ecosystems are in equilibrium and do not experience disturbances (e.g., logging, fire), we selected flux sites without strong disturbances during the last 10 years. For sites that also provided growing season LAI information, we also removed forests site with  $LAI < 2$ , which may be considered as sparse forests with understory vegetation. In the end,

observations of 655 site-years from 159 sites were retained (Table S2). The annual ~~climate~~mean temperature of the sites spans from -9 °C to 27°C~~-in temperature, and, while the annual precipitation spans~~ from 67 mm yr<sup>-1</sup> to over 3000 mm yr<sup>-1</sup> ~~in precipitation~~ (Fig. S2), which is representative to most of the climate conditions over the globe. The dataset provides in situ meteorology, net ecosystem exchange (NEE), gross primary productivity (GPP), and data quality information at 30-min time steps. The GPP provided by this dataset is partitioned from NEE and gap filled using the method of Reichstein et al. (2005). Specifically, 64 of the 159 sites provided measurements of both total and diffuse PPFD, which allows us to evaluate the light partitioning parametrization (Eqs. (9-2025-38)). The gaps and missing variables in meteorology are filled using the approach from Vuichard and Papale (2015) to meet the model input requirements.

Because ORCHIDEE has different photosynthesis parameters for different PFTs, we classified the vegetation at each site into the 13 ORCHIDEE PFTs (Table S1) according to the IGBP land cover types specified on the website of FluxNet ([www.fluxdata.org](http://www.fluxdata.org)). If the IGBP land cover type is not specified or may match more than one ORCHIDEE PFTs (e.g. shrublands, savannas and wetlands), the PFT is determined according to the dominant plant species described in related references. Specifically, the mixed forests (MF) type exists in the IGBP classification but not in the ORCHIDEE PFTs. Because MF sites are mostly located in temperate regions, we assume that they are composed of 50% temperate broadleaf deciduous forests and 50% temperate evergreen needle-leaf forests. Detailed information of flux sites is found in Table S2.

To evaluate the model, spinup simulations of 30 years are firstly conducted on ORCHIDEE\_DF at each site to equilibrate the leaf area index with site conditions. Then the simulations with 30 min output are conducted with ORCHIDEE trunk and ORCHIDEE\_DF, using the full span of the Fluxnet la Thiule series respectively at each site. It should be noted that we use the same spinup for ORCHIDEE trunk and ORCHIDEE\_DF to ensure the same initial states of the two simulations. A test has shown that different spinup simulations do not affect the simulation of GPP in the following years (not shown).

### 2.3 Analyses

When evaluating the modeled GPP response to diffuse light we have not used all the 30-min data points due to several concerns. First, all night time data points were excluded from the analyses given that GPP is zero at night. Second, all data points flagged with poor quality in the FLUXNET archive have been removed. Third, ORCHIDEE might not be perfect in capturing the seasonality of leaf flushing and shedding. In order to minimize the uncertainty from phenology, we used only data from the growing season at each site, which is defined as months when:

$$GPP_m > GPP_{m,min} + (GPP_{m,max} - GPP_{m,min}) / 4 \quad (39)$$

where  $GPP_m$  is the observed monthly GPP,  $GPP_{m,min}$  and  $GPP_{m,max}$  are the observed minimum and maximum monthly GPP at the corresponding sites.

To assess the effect of variable diffuse light fraction on both GPP and light use efficiency (LUE, the ratio of GPP to incoming shortwave radiation), we look at the difference in GPP and LUE during sunny and cloudy conditions. We define sunny and

带格式的: 字体: 倾斜

带格式的: 字体: 倾斜

带格式的: 字体: 倾斜

cloudy conditions as those when the fraction of diffuse PPFD at the top of the canopy ( $Fdf_{PPFD}$ ) is smaller than 0.4 and greater than 0.8, respectively, and calculate the average sunny and cloudy GPP and LUE at each site. To ensure that the comparison between sunny and cloudy conditions are at the same PPFD level, the sunny time steps with PPFD larger than the maximum PPFD under cloudy conditions are removed from the average, and vice versa. In addition, to make sure that the difference in GPP between sunny and cloudy is not an artifact of different LAI, sites with average ~~modelled~~modeled LAI under cloudy and sunny conditions differing by more than 0.3 are excluded from this analysis.

### 3 Results

#### 3.1 Diffuse light modeling

Fig. 2 shows the relationship between 30-min modeled and measured  $Fdf_{PPFD}$  at flux sites (64 sites). The data points are generally distributed along the 1:1 line, indicating an unbiased estimation of our diffuse light model. In total, our simple model explains over 51% of the variance in observed diffuse PPFD fraction. Although this model is imperfect, we currently have no better way to reproduce the diffuse PPFD at the flux site scale.

#### 3.2 General model performance

The performance of both ORCHIDEE trunk and ORCHIDEE\_DF for 30-min GPP from each PFT (all sites) is presented in Fig. 4. Generally, ORCHIDEE trunk underestimated the standard deviation (STD) of GPP at 30-min time-step compared with observations, and across all PFTs except Boreal evergreen needleleaf forests and C4 Croplands (Fig. 4a). The correlation coefficients between ORCHIDEE trunk GPP and observations are generally between 0.5 and 0.7 among PFTs (Fig. 4b). In tropical broadleaf forests, this correlation coefficient is about 0.2, which is much smaller than in other PFTs and likely due to the limited seasonality of primary production in the tropics. The GPP simulated by ORCHIDEE\_DF shows comparable performance with ORCHIDEE trunk, but with slightly smaller STD (Fig. 4a). Similar evaluations on the GPP from the two models are performed under cloudy and sunny conditions respectively (Fig. 4c-f). Under cloudy conditions, ORCHIDEE trunk and ORCHIDEE\_DF both underestimated GPP STD. The correlation coefficients to observations are generally between 0.5 and 0.8 (Fig. 4d). Compared with ORCHIDEE trunk, ORCHIDEE\_DF shows slightly worse correlation coefficients but improves STD for most of the PFTs except Tropical broad-leaved evergreen forests (TrEBF) and Temperate needleleaf evergreen forests (TeENF) (Fig. 4c). Compared with cloudy conditions, the GPP simulated by the two models under sunny conditions show weaker correlation to observations. The correlation coefficients generally vary between 0.3 and 0.6 among PFTs. However, it should be noted that ORCHIDEE\_DF better reproduced GPP variation under sunny conditions compared with ORCHIDEE trunk in most PFTs except TeDBF and C4Cro (Fig. 4f). The GPP STD derived from ORCHIDEE trunk simulations under sunny conditions show larger variability among PFTs than under cloudy conditions. While for ORCHIDEE\_DF, the GPP STD under sunny and cloudy conditions show similar bias compared with observations (Fig. 4e).

### 3.3 Effects of diffuse light on GPP and LUE

Because the modification of ORCHIDEE\_DF was limited to light transmission, the pertinent process-oriented evaluation of the two models should focus on their ability to capture the observed GPP differences between cloudy and sunny conditions (hereafter  $\Delta\text{GPP}$ ), rather than on correlations or RMSE with observations, that may result from different structural and parametric errors of the model, not related to diffuse light.

Figure 5 shows the observed and modeled GPP under sunny and cloudy conditions at different PPFD levels at flux sites with relatively long time series of observations from each PFT. For all the sites selected, the observed GPP under cloudy conditions is larger than under sunny conditions. However, the GPP simulated by ORCHIDEE trunk shows no or small difference between cloudy and sunny conditions at most sites. In contrast, ORCHIDEE\_DF reproduces this GPP difference in most PFTs except TrDBF, BoDBF and C4Gra. However, there is only one TrDBF site and very few C4Gra sites in our dataset. Furthermore, at most C4Gra sites, we are not able to find PPFD levels where sunny and cloudy conditions co-exist. Therefore, we are not able to make further evaluation of cloudy-minus-sunny GPP differences for TrDBF and C4Gra. At three of the four BoDBF sites, the modeled GPP difference under cloudy and sunny conditions is relatively small (not shown). This might be because the model overestimated the deleterious effect of low temperature on photosynthesis at the BoDBF sites (mean annual temperature  $< 3^{\circ}\text{C}$ ). In total, observations from about 70% of the sites show remarkable higher GPP under cloudy than sunny conditions. This percentage is only 30% in ORCHIDEE trunk simulations but 60% in ORCHIDEE\_DF simulations.

To summarize the site level results, we investigated the distribution of GPP difference between cloudy and sunny conditions (here after refer to as  $\Delta\text{GPP}$ ) (Fig. 6a). Observations and ORCHIDEE\_DF show a positive bias in  $\Delta\text{GPP}$ , with  $\Delta\text{GPP}$  values between  $0.3 \times 10^{-4} \text{ gC m}^{-2} \text{ s}^{-1}$  at most sites. However, for ORCHIDEE trunk,  $\Delta\text{GPP}$  is near zero at most sites. This result confirms that ORCHIDEE\_DF performs much better than ORCHIDEE trunk in simulating differences in GPP under different light conditions.

It should be noted that  $\Delta\text{GPP}$  can be affected by PPFD. At sites where sunny and cloudy conditions only coexist at a relatively low PPFD level, the  $\Delta\text{GPP}$  should be also small. To remove the effect of PPFD level on  $\Delta\text{GPP}$ , we analyzed the difference in LUE, i.e.  $\Delta\text{LUE}$ , between the two conditions (Fig. 6b). Compared with  $\Delta\text{GPP}$ , positive  $\Delta\text{LUE}$  values are more evenly distributed around  $0.15 \times 10^{-8} \text{ gC } \mu\text{mol}^{-1} \text{ photon}$  for observation and ORCHIDEE\_DF simulation. For ORCHIDEE trunk, the  $\Delta\text{LUE}$  has the range of  $0.8 \times 10^{-8} \text{ gC } \mu\text{mol}^{-1} \text{ photon}$ , with the upper range smaller than in the observations and ORCHIDEE\_DF. We further refined this analysis to investigate if the effects of diffuse light differ at different times of the day (Fig. 7). Results for three different periods in the day show that in the morning and afternoon, cloudy conditions result in higher GPP of  $0.5 \times 10^{-4} \text{ gC m}^{-2} \text{ s}^{-1}$  than sunny conditions at most sites, which is generally captured by ORCHIDEE\_DF but missed by ORCHIDEE trunk in the morning (Fig. 6a, c). At midday, due to the dependence of  $F_{\text{df}}$  on PPFD (Eqs. (925 and 4026)), we fail at many sites to find PPFD levels where sunny and cloudy conditions coexist. Nevertheless, the result generally indicates larger mid-day  $\Delta\text{GPP}$  than those in the morning and afternoon; although the modeled  $\Delta\text{GPP}$  is slightly smaller than the observation. It should be noted that this large difference is captured by both ORCHIDEE\_DF and ORCHIDEE trunk (Fig. 7b).

Because direct and diffuse light are not distinguished in ORCHIDEE trunk, this midday  $\Delta$ GPP should be mainly contributed by environmental factors other than diffuse light fraction. The underestimation of midday  $\Delta$ GPP could be a result of error in current ORCHIDEE parameterizations. The  $\Delta$ LUE derived by ORCHIDEE\_DF also shows a largely similar distribution as in observations, but ORCHIDEE trunk underestimates the morning and afternoon  $\Delta$ LUE (Fig. 7d-f).

### 3.4 Interactions between diffuse light and environmental factors

As implied by Fig. 7, the diffuse light fraction is not the only factor causing  $\Delta$ GPP. Other possible factors include temperature and VPD (Gu et al., 2002; Cheng et al., 2015; Li et al., 2014). Here, we thus investigate the diffuse light effect along temperature and VPD gradients in Fig. 8. To remove the effect of PPFD level, we only show  $\Delta$ LUE.

$\Delta$ LUE shows a unimodal curve along the temperature gradient for observation and the two models (Fig. 8a). At low temperature, both models indicate a very low  $\Delta$ LUE of  $1 \text{ gC } \mu\text{mol}^{-1} \text{ photon}$ , which is about 1/3 of the  $\Delta$ LUE derived from observations. With increasing temperature, the observed  $\Delta$ LUE shows a maximum at 10-20 °C, with a magnitude of  $\sim 8 \times 10^{-8} \text{ gC } \mu\text{mol}^{-1} \text{ photon}$  and declines slightly at higher temperatures. The peak of  $\Delta$ LUE simulated by ORCHIDEE\_DF has a magnitude comparable to that of observations, but at higher temperature (20-25 °C) than for observations. The  $\Delta$ LUE simulated by ORCHIDEE trunk is much smaller, with a peak of  $\sim 4 \times 10^{-8} \text{ gC } \mu\text{mol}^{-1} \text{ photon}$  at 10-15 °C.

The effect of VPD on  $\Delta$ LUE is shown in Fig. 8b. For observations and both model simulations, a monotonic decreasing trend of  $\Delta$ LUE along the VPD gradient is found. The  $\Delta$ LUE from observations and ORCHIDEE\_DF show a comparable magnitude, from  $8 \times 10^{-8} \text{ gC } \mu\text{mol}^{-1} \text{ photon}$  at VPD < 0.5 kPa to  $5 \times 10^{-8} \text{ gC } \mu\text{mol}^{-1} \text{ photon}$  at 2-4 kPa VPD level. The  $\Delta$ LUE simulated by ORCHIDEE trunk is smaller than observations.

Apart from environmental factors, the effects of diffuse light may also differ among PFTs because different PFTs have different canopy structures and photosynthetic parameters (e.g. Vcmax). Here we analyzed the  $\Delta$ LUE in forests and short vegetation (grasslands and croplands) separately (Fig. 8c-f). In forests (Fig. 8c, d), ORCHIDEE\_DF underestimates  $\Delta$ LUE at temperatures lower than 20 °C. It also largely captures the observed  $\Delta$ LUE trend with VPD, while ORCHIDEE trunk underestimates  $\Delta$ LUE at all cases. Compared with forests, in short vegetation (Fig. 8e, f), observations show a stronger decline of  $\Delta$ LUE at high temperatures (>25 °C) and high VPD conditions (>0.5 kPa). However, for ORCHIDEE\_DF, the short vegetation  $\Delta$ LUE remains as high as for forests.

Fig. 9 shows the distribution of  $\Delta$ LUE in the Temperature-VPD dimensions. Observations indicate that the largest  $\Delta$ LUE is reached under conditions when temperature is in the range 5-20 °C and VPD < 1 kPa (Fig. 9a). This temperature is thought more favorable for photosynthesis as it is generally consistent with the photosynthesis optimum temperature detected by Huang et al. (2019) in latitudes where most of the sites are located. Under these conditions, the  $\Delta$ LUE is usually over  $7 \times 10^{-8} \text{ gC } \mu\text{mol}^{-1} \text{ photon}$ . When the temperature is lower than 5 °C or higher than 20 °C, or VPD becomes larger than 1 kPa,  $\Delta$ LUE tends to decline. Compared with observations, the  $\Delta$ LUE simulated by ORCHIDEE\_DF shows a similar decreasing trend with VPD at all temperature levels (Fig. 9c), however, no obvious decline of  $\Delta$ LUE is found at high temperatures. The  $\Delta$ LUE simulated by ORCHIDEE trunk is much smaller compared with observations (Fig. 9b).

The  $\Delta$ LUE from forests and short vegetation are shown separately in Fig. 10. Based on site level observations (Fig. 10a), both vegetation types show a larger  $\Delta$ LUE at lower VPD between 5-20 °C. In forests, there is also large  $\Delta$ LUE at high temperature conditions, which mainly occurs in tropical forests (Fig. S3). Nevertheless, ORCHIDEE\_DF still overestimates the  $\Delta$ LUE at high temperatures (Fig. 10e), which is mainly due to the overestimation of  $\Delta$ LUE at high temperatures for temperate forests (Fig. S3).

Compared with forests, the short vegetation shows a much stronger decline of  $\Delta$ LUE at higher VPD level (Fig. 10b), however, it is not well captured by ORCHIDEE\_DF (Fig. 10f). In most cases, ORCHIDEE trunk tends to strongly underestimate  $\Delta$ LUE unless the observed  $\Delta$ LUE is small or negative (e.g. VPD > 2kPa for short vegetation).

#### 4 Discussion

##### 4.1 Improvement of ORCHIDEE\_DF

The role of diffuse light on photosynthesis has been found and modeled in different vegetation types (Gu et al., 2003; Niyogi et al., 2004; Misson et al., 2005; Alton et al., 2007b; Knohl and Baldocchi, 2008; Mercado et al. 2009; Oliphant et al., 2011; Kanniah et al., 2013; Williams et al., 2014; Cheng et al., 2015; Wang et al., 2018). However, very few studies have attempted to account for the diffuse light effect in a global land surface model, and fewer studies have used large FLUXNET datasets for evaluation. Here, by using flux observations from 159 sites over the globe, we show that by separating the direct and diffuse light, ORCHIDEE\_DF improves the simulation of GPP under sunny conditions and, more importantly, reproduced the observed impacts of diffuse light on GPP and LUE for most of the PFTs (Figs. 4-6). Under cloudy conditions, ORCHIDEE\_DF seems to perform slightly worse than ORCHIDEE trunk (Fig. 4). However, it should be noted that ORCHIDEE\_DF has not been recalibrated and all parameters are those from ORCHIDEE trunk despite the substantial changes in the code with respect to light partitioning and canopy light transmission. Furthermore, the GPP simulated by ORCHIDEE trunk shows different GPP STD biases under sunny and cloudy conditions, while ORCHIDEE\_DF gives a more systematically underestimated GPP STD, which should be more easily corrected in a future calibration. The site level comparison (Fig. 5) also explains how ORCHIDEE\_DF reproduces the GPP increase compared to ORCHIDEE trunk. At most sites, the GPP simulated by the two models show similar magnitude under cloudy conditions. While under sunny conditions, the GPP simulated by ORCHIDEE\_DF is significantly smaller. This is because in the one-stream canopy light transmission model in ORCHIDEE trunk, all light is considered as diffuse light and evenly distributed in each leaf layer. This simplified approach to the modelling of light distribution leads to larger GPP under sunny conditions because the effect of light saturation on sunlit leaves is ignored. Since ORCHIDEE trunk was calibrated using both sunny and cloudy data, but ORCHIDEE\_DF corrected the overestimation under sunny conditions, ORCHIDEE\_DF may give an overall underestimation using current parameters.

#### 4.2 Factors affecting the response of GPP to diffuse light

Although diffuse light can increase photosynthesis of shaded leaves, the GPP increase under cloudy conditions is not contributed only by this effect. A recent field study suggested that photosynthesis from part of the canopy (especially sunlit leaves) benefits from the lower VPD rather than the higher diffuse light fraction under cloudier conditions (Wang et al., 2018).

Our results show that during the morning and the afternoon, higher diffuse PAR fraction is the main factor causing larger GPP under cloudy conditions compared with sunny conditions, as only ORCHIDEE\_DF reproduced the observed positive  $\Delta$ GPP during the two periods (Fig. 7). While at midday, the larger GPP under cloudy conditions should be mainly due to lower T or VPD other than to diffuse light because ORCHIDEE trunk, which does not simulate the diffuse light effect, also reproduces this effect (Fig. 7). A similar effect is also reported by Cheng et al. (2015), who found that in croplands the midday GPP increase under cloudier conditions is mainly caused by lower temperature and lower VPD rather than by diffuse light. Photosynthesis is often considered as limited by either carboxylation or electron transportation (Farquhar et al., 1980). It is when the shaded leaf photosynthesis is limited by light that diffuse light can increase GPP. At midday, large VPD may cause stomatal closure, leading to a carboxylation-limited photosynthesis. Our results imply that it might be important to consider the diurnal cycle of environmental factors to better understand the effect of diffuse light.

It should be noted that the covariation of environmental factors with more diffuse light under cloudier conditions does not always benefit photosynthesis. For instance, if the vegetation is cold stressed under cooler conditions, the decrease of temperature under cloudier condition may strengthen this stress and offset the effect of diffuse light. Our analyses indicate that under most stressed conditions, the effect of diffuse light on photosynthesis is weakened (Figs. 9, 10).

Another important factor is the light itself. When there is no light saturation of shaded leaves, under the same diffuse light fraction, stronger light levels are likely to benefit the shaded leaves more, resulting in higher  $\Delta$ GPP (Fig. 5,  $\Delta$ GPP tends to be larger at higher PPFD level at most sites). Nevertheless, apart from GPP, in this study we also investigated LUE (the photosynthesis per unit PPFD), which has removed this effect.

Besides environmental factors, canopy structure is also very important. Theoretically, thicker canopies with large LAI tend to be more sensitive to diffuse light because a larger fraction of leaves are light limited due to shading (Fig. 1). As expected, ORCHIDEE\_DF has shown an increasing  $\Delta$ LUE with LAI (Fig. 11). However, the analyses based on LAI observations suggested a very weak positive effect of LAI on  $\Delta$ LUE (Fig. 11). This insensitive response of  $\Delta$ LUE to LAI detected here should be treated with caution because the LAI observations are not well defined (maximum or average) and remain very limited in the current FLUXNET dataset (less than 10 in each LAI interval). Using more detailed LAI and CO<sub>2</sub> flux observations, Wohlfahrt et al. (2008) has clearly exhibited the influence of LAI on diffuse light-induced photosynthesis changes at a grassland.

### 4.3 Uncertainty and Limitations

Many empirical methods have been proposed to partition solar radiation into diffuse and direct light (e.g. Spitters et al., 1986; Weiss and Norman, 1985; Erbs et al., 1982). However, biases remain in the predicted diffuse light fraction under all aerosol and cloud conditions, which inevitably introduce some uncertainties to our analyses. Nevertheless, such methods are currently the most feasible approach at flux site level. More continuous measurements of direct and diffuse surface radiation at more sites are desirable.

Another source of uncertainty is from the light transmission model. In ORCHIDEE\_DF, we used a two-stream radiative transfer approximation. In this model, the canopy trait parameters such as leaf scattering, leaf orientation and leaf clumping factors are assumed the same for all PFTs, however real canopies are very diverse (Smith et al., 2004). In situ observations are required to obtain better parameters. Furthermore, the validity of the light transmission model in ORCHIDEE\_DF depends on the several assumptions described in the model description section. These assumptions are not always valid. For example, because direct solar beams are not exact parallels, leaves in canopies are not always sunlit or shaded, they may also fall in penumbra regions, (i.e. regions where only part of the incoming direct solar beams are blocked, Smith et al., 1989; Cescatti and Niinemets, 2005). These more complex processes should be considered in future model development. Nevertheless, our simplified light transmission already succeeds in reproducing the observed diffuse light impact.

There are other sources of uncertainties in complex land surface models. Although ORCHIDEE\_DF reproduces the magnitude of the diffuse light effects, it fails to reproduce the response of  $\Delta$ LUE to temperature. For all PFTs, ORCHIDEE\_DF underestimates the  $\Delta$ LUE at low temperatures, and overestimates  $\Delta$ LUE at high temperatures (Fig. 8). The low temperature underestimation is also found in ORCHIDEE\_trunk, indicating that the models may have underestimated the tolerance of plants to low temperatures. While at high temperatures, ORCHIDEE\_DF tends to underestimate the impact of heat stress. This bias might be due to the parameterization of temperature acclimation which is based on observations mainly from a narrow temperature range (11-29 °C) (Kattge et al. 2007). For short vegetation, the introduction of diffuse light into the model results in an increase of  $\Delta$ LUE at high temperatures and high VPD (Figs. 8, 10), indicating the vegetation simulated by ORCHIDEE\_trunk remains light limited under such conditions. However, the strong decreasing trend of observed  $\Delta$ LUE along temperature and VPD gradients indicates heat and VPD stress. This implies that parameters in current ORCHIDEE version may have underestimated the response of grassland and cropland photosynthesis to heat and VPD stress.

Besides the possible bias in parameters, both ORCHIDEE\_trunk and ORCHIDEE\_DF lack a representation of the response of leaf temperature to radiation. Instead, the air temperature is used directly to represent the leaf temperature throughout the canopy for simulating gas exchange processes in current model. As shown by Chen and Zhuang (2014), the changes of radiation regime due to aerosols can significantly affect leaf temperature, which could potentially affect GPP. For now, ORCHIDEE\_DF remains not capable of dealing with this response of leaf temperature. Further developments are needed for disentangling the role of leaf temperature and diffuse light on GPP.

585

**5 Conclusion**

In this study, we added to the ORCHIDEE trunk a module to partition the downward surface solar radiation into diffuse and direct components, and a new canopy radiative transfer model, which separates the existing multilayer canopy into sunlit and shaded leaves. The resulting new land surface model, ORCHIDEE\_DF, is evaluated using the La Thuile flux dataset over 159 sites over 11 PFTs. Compared with ORCHIDEE trunk, ORCHIDEE\_DF improves the GPP simulation under sunny conditions. This improvement successfully reproduces the observed enhancement of GPP under cloudier conditions at most of the sites. Using observed and modeled GPP, we found an increase of GPP under cloudier conditions at all times of the day; however, the mechanisms causing this effect are different at midday from morning and afternoon. During morning and afternoon, the increase in GPP is mainly caused by increased diffuse light fraction, while at midday, the GPP increase is mainly due to weaker stress from temperature and VPD.

585 Observations indicate that under cloudy and sunny conditions for the same light level, the maximum LUE difference can be over  $7 \times 10^{-8}$  gC  $\mu\text{mol}^{-1}$  photon. The maximum LUE is found at temperature and VPD conditions more favorable for photosynthesis (5-20 °C for temperature and < 1 kPa for VPD). With increasing VPD, or under lower or higher temperatures, the LUE may decrease. Compared with observations, ORCHIDEE\_DF underestimates the diffuse light effect at low temperature and overestimates it at high temperatures, possibly due to imperfect temperature acclimation parameterization in the current ORCHIDEE model. In grasslands and croplands, ORCHIDEE\_DF overestimates the diffuse light effect on LUE, which might be due to an overestimation of their tolerance to dry conditions.

590 As ORCHIDEE\_DF is a land surface model which is able to capture the effect of diffuse light for a large number of sites over the globe, we are confident that, with this improved model framework and proper calibration, we can investigate the effect of aerosols on global biogeochemical cycles, and assess the impact of aerosol emission policies and aerosol related climate engineering on such cycles.

**Appendix A**

595 List of acronyms:

Fdf: Fraction of diffuse radiation

GPP: Gross Primary Production

LAI: Leaf Area Index

LSM: Land Surface Model

600 LUE: Light Use Efficiency

NIR: Near-Infrared Radiation

PAR: Photosynthetically Active Radiation

PFT: Plant Functional Type

PPFD: Photosynthetic Photon Flux Density

605 SW: downward Shortwave Radiation at the top of canopy  
TOA: Top of Atmosphere  
TOC: Top of Canopy  
VPD: Vapor Pressure Deficit  
\* The variable names in Section 2 are listed in Table 1

610 **Code and data availability**

The code of the ORCHIDEE\_DF is available at [https://forge.ipsl.jussieu.fr/orchidee/wiki/GroupActivities/CodeAvailabilityPublication/ORCHIDEE\\_DFv1.0\\_site](https://forge.ipsl.jussieu.fr/orchidee/wiki/GroupActivities/CodeAvailabilityPublication/ORCHIDEE_DFv1.0_site)[doi.org/10.14768/20200407003.1](https://doi.org/10.14768/20200407003.1). Flux data (La Thuile) used in this study is available at <https://fluxnet.fluxdata.org/data/la-thuile-dataset/>.

615 **Acknowledgement**

The authors acknowledge support from European Research Council Synergy project SyG-2013-610028 IMBALANCE-P and the ANR CLAND Convergence Institute- [\(16-CONV-0003\)](#). The authors are very grateful to the FLUXNET communities for their efforts at making sites and collecting data, and specially to flux site PIs who are not in the author list but have given constructive suggestions on this manuscript. The authors also acknowledge Dr. Yves Balkanski and Dr. Nicolas Viovy for  
620 their suggestions during this work.

**Author contributions.**

PC, OB and LL designed the project. YZ developed the model code with help from AB, FM, DG and AC. NV provided the code for data gap filling. YZ prepared the paper with contributions from all the co-authors.

**References**

625 Alton, P., North, P., and Los, S.: The impact of diffuse sunlight on canopy light-use efficiency, gross photosynthetic product and net ecosystem exchange in three forest biomes, *Global Change Biol.*, 13(4), 776-787, 2007a  
Alton, P., Ellis, R., Los, S., and North, P.: Improved global simulations of gross primary product based on a separate and explicit treatment of diffuse and direct sunlight, *J. Geophys. Res.-Atmos.*, 112, D07203, doi:10.1029/2006JD008022.1, 2007b  
630 Baldocchi, D. D., and Wilson, K. B.: Modeling CO<sub>2</sub> and water vapor exchange of a temperate broadleaved forest across hourly to decadal time scales, *Ecol. Model.*, 142(1-2), 155-184, 2001

Bodin, P., and Franklin, O.: Efficient modeling of sun/shade canopy radiation dynamics explicitly accounting for scattering, *Geosci. Model Dev.*, 5(2), 535-541, 2012

Bonan, G. B., Oleson, K. W., Fisher, R. A., Lasslop, G., and Reichstein, M.: Reconciling leaf physiological traits and canopy flux data: Use of the TRY and FLUXNET databases in the Community Land Model version 4, *J. Geophys. Res.-Biogeo.*, 117, G02026, doi:10.1029/2011JG001913.1, 2012

Chen, L., Yan, G., Wang, T., Ren, H., Calb ́ J., Zhao, J., and McKenzie, R.: Estimation of surface shortwave radiation components under all sky conditions: Modeling and sensitivity analysis, *Remote Sens. Environ.*, 123, 457-469, 2012

Chen, M., and Zhuang, Q.: [Evaluating aerosol direct radiative effects on global terrestrial ecosystem carbon dynamics from 2003 to 2010, \*Tellus B: Chemical and Physical Meteorology\*, 66, 21808, 10.3402/tellusb.v66.21808, 2014.](#)

Cheng, S. J., Bohrer, G., Steiner, A. L., Hollinger, D. Y., Suyker, A., Phillips, R. P., and Nadelhoffer, K. J.: Variations in the influence of diffuse light on gross primary productivity in temperate ecosystems, *Agr. Forest Meteorol.*, 201, 98-110, 2015

Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R., Galloway, J., Heimann, M., Jones, C., Le Qu ́ C., Myneni, R.B., Piao, S., and Thornton, P.: Carbon and other biogeochemical cycles, in *Climate change 2013: the physical science basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 465-570, 2013

Dai, Y., Dickinson, R. E., and Wang, Y.-P.: A two-big-leaf model for canopy temperature, photosynthesis, and stomatal conductance, *J. Climate*, 17(12), 2281-2299, 2004

de Pury, D., and Farquhar, G.: Simple scaling of photosynthesis from leaves to canopies without the errors of big-leaf models, *Plant Cell Environ.*, 20, 537– 557, 1997

Dye, D. G.: Spectral composition and quanta-to-energy ratio of diffuse photosynthetically active radiation under diverse cloud conditions, *J. Geophys. Res.-Atmos.*, 109, D10203, doi:10.1029/2003JD004251, 2004

Erbs, D., Klein, S., and Duffie, J.: Estimation of the diffuse radiation fraction for hourly, daily and monthly-average global radiation, *Sol. Energy*, 28(4), 293-302, 1982

Farquhar, G. D., von Caemmerer, S. v., and Berry, J.: A biochemical model of photosynthetic CO<sub>2</sub> assimilation in leaves of C3 species, *Planta*, 149(1), 78-90, 1980

Goudriaan, J.: *Crop micrometeorology: a simulation study*, Pudoc, Wageningen, the Netherlands 1977

Goudriaan, J.: Potential production processes, in *Simulation of plant growth and crop production*, Pudoc, Wageningen, the Netherlands, pp. 98-113, 1982

Gu, L., Baldocchi, D., Verma, S. B., Black, T., Vesala, T., Falge, E. M., and Dowty, P. R.: Advantages of diffuse radiation for terrestrial ecosystem productivity, *J. Geophys. Res.-Atmos.*, 107(D6), ACL 2-1-ACL 2-23, 2002

Gu, L., Baldocchi, D. D., Wofsy, S. C., Munger, J. W., Michalsky, J. J., Urbanski, S. P., and Boden, T. A.: Response of a deciduous forest to the Mount Pinatubo eruption: Enhanced photosynthesis, *Science*, 299(5615), 2035-2038, 2003

Hikosaka, K., Niinemets, U., and Anten, N. P.: Canopy photosynthesis: from basics to applications, Springer, 2016. Doi: [10.1007/978-94-017-7291-4](https://doi.org/10.1007/978-94-017-7291-4)

Howell, T., Meek, D., and Hatfield, J.: Relationship of photosynthetically active radiation to shortwave radiation in the San Joaquin Valley, Agr. Meteorol., 28(2), 157-175, 1983

670 Huang, M., Piao, S., Ciais, P., Peñuelas, J., Wang, X., Keenan, T. F., Peng, S., Berry, J. A., Wang, K., Mao, J., Alkama, R., Cescatti, A., Cuntz, M., De Deurwaerder, H., Gao, M., He, Y., Liu, Y., Luo, Y., Myneni, R. B., Niu, S., Shi, X., Yuan, W., Verbeeck, H., Wang, T., Wu, J., and Janssens, I. A.: Air temperature optima of vegetation productivity across global biomes, Nature Ecology & Evolution, 3, 772-779, [10.1038/s41559-019-0838-x](https://doi.org/10.1038/s41559-019-0838-x), 2019.

675 Kanniah, K. D., Beringer, J., and Hutley, L.: Exploring the link between clouds, radiation, and canopy productivity of tropical savannas, Agr. Forest Meteorol., 182, 304-313, 2013.

Kattge, J., and Knorr, W.: Temperature acclimation in a biochemical model of photosynthesis: a reanalysis of data from 36 species, Plant Cell Environ., 30, 1176-1190, 2007.

Knohl, A., and Baldocchi, D. D.: Effects of diffuse radiation on canopy gas exchange processes in a forest ecosystem, J. Geophys. Res.-Biogeo., 113, G02023, doi:10.1029/2007JG000663, 2008

680 Krinner, G., Viovy, N., de Noblet-Ducoudré N., Ogée, J., Polcher, J., Friedlingstein, P., Ciais, P., Sitch, S., and Prentice, I. C.: A dynamic global vegetation model for studies of the coupled atmosphere-biosphere system, Global Biogeochem. Cy., 19, 2005.

685 Leuning, R., Kelliher, F. M., de Pury, D. G. G., and Schulze, E.-D.: Leaf nitrogen, photosynthesis, conductance and transpiration: scaling from leaves to canopies, Plant Cell Environ., 18, 1183-1200, [10.1111/j.1365-3040.1995.tb00628.x](https://doi.org/10.1111/j.1365-3040.1995.tb00628.x), 1995.

Li, T., Heuvelink, E., Dueck, T., Janse, J., Gort, G., and Marcelis, L.: Enhancement of crop photosynthesis by diffuse light: quantifying the contributing factors, Ann. Bot.-London, 114(1), 145-156, 2014

690 Le Quéré, C., Andrew, R. M., Friedlingstein, P., Sitch, S., Hauck, J., Pongratz, J., Pickers, P. A., Korsbakken, J. I., Peters, G. P., Canadell, J. G., Arneth, A., Arora, V. K., Barbero, L., Bastos, A., Bopp, L., Chevallier, F., Chini, L. P., Ciais, P., Doney, S. C., Gkritzalis, T., Goll, D. S., Harris, I., Haverd, V., Hoffman, F. M., Hoppema, M., Houghton, R. A., Hurtt, G., Ilyina, T., Jain, A. K., Johannessen, T., Jones, C. D., Kato, E., Keeling, R. F., Goldewijk, K. K., Landschützer, P., Lefèvre, N., Lienert, S., Liu, Z., Lombardozzi, D., Metz, N., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S., Neill, C., Olsen, A., Ono, T., Patra, P., Peregon, A., Peters, W., Peylin, P., Pfeil, B., Pierrot, D., Poulter, B., Rehder, G., Resplandy, L., Robertson, E., Rocher, M., Rödenbeck, C., Schuster, U., Schwinger, J., Séférian, R., Skjelvan, I., Steinhoff, T., Sutton, A., Tans, P. P., Tian, H., Tilbrook, B., Tubiello, F. N., van der Laan-Luijkx, I. T., van der Werf, G. R., Viovy, N., Walker, A. P., Wiltshire, A. J., Wright, R., Zaehle, S., and Zheng, B.: Global Carbon Budget 2018, Earth Syst. Sci. Data, 10, 2141–2194, <https://doi.org/10.5194/essd-10-2141-2018>, 2018.

- Meir, P., Kruijt, B., Broadmeadow, M., Barbosa, E., Kull, O., Carswell, F., Nobre, A., and Jarvis, P. G.: Acclimation of photosynthetic capacity to irradiance in tree canopies in relation to leaf nitrogen concentration and leaf mass per unit area, *Plant Cell Environ.*, 25, 343-357, 10.1046/j.0016-8025.2001.00811.x, 2002.
- 700 Mercado, L. M., Bellouin, N., Sitch, S., Boucher, O., Huntingford, C., Wild, M., and Cox, P. M.: Impact of changes in diffuse radiation on the global land carbon sink, *Nature*, 458(7241), 1014-1017, 2009
- Misson, L., Lunden, M., McKay, M., and Goldstein, A. H.: Atmospheric aerosol light scattering and surface wetness influence the diurnal pattern of net ecosystem exchange in a semi-arid ponderosa pine plantation, *Agr. Forest Meteorol.*, 129(1-2), 69-83, 2005
- 705 Monsi, M., and Saeki, T.: On the factor light in plant communities and its importance for matter production, *Ann. Bot.-London*, 95(3), 549-567, 2005
- Myneni, R. B., Ross, J., and Asrar, G.: A review on the theory of photon transport in leaf canopies, *Agr. Forest Meteorol.*, 45(1-2), 1-153, 1989
- 710 Niinemets, U., Kull, O., and Tenhunen, J. D.: An analysis of light effects on foliar morphology, physiology, and light interception in temperate deciduous woody species of contrasting shade tolerance, *Tree Physiol.*, 18(10), 681-696, 1998
- Niyogi, D., Chang, H.-I., Saxena, V. K., Holt, T., Alapaty, K., Booker, F., Chen, F., Davis, K. J., Holben, B., Matsui, T., Meyers, T., Oechel, W. C., Pielke Sr., R. A., Wells, R., Wilson, K., and Xue, Y.: Direct observations of the effects of aerosol loading on net ecosystem CO<sub>2</sub> exchanges over different landscapes, *Geophys. Res. Lett.*, 31, 10.1029/2004gl020915, 2004.
- 715 [Oleson, K., Lawrence, D., Bonan, G., Drewniak, B., Huang, M., Koven, C., Levis, S., Li, F., Riley, W., Subin, Z., Swenson, S., Thornton, P., Bozbiyik, A., Fisher, R., Heald, C., Kluzek, E., Lamarque, J., Lawrence, P., Leung, L., Lipscomb, W., Muszala, S., Ricciuto, D., Sacks, W., Tang, J., and Yang, Z.: Technical Description of version 4.5 of the Community Land Model \(CLM\), Technical Note #NCAR/TN-503+STR, 2013.](#)
- 720 Oliphant, A., Dragoni, D., Deng, B., Grimmond, C., Schmid, H.-P., and Scott, S.: The role of sky conditions on gross primary production in a mixed deciduous forest, *Agr. Forest Meteorol.*, 151(7), 781-791, 2011
- Piao, S., Ciais, P., Friedlingstein, P., de Noblet-Ducoudré, N., Cadule, P., Viovy, N., and Wang, T.: Spatiotemporal patterns of terrestrial carbon cycle during the 20th century, *Global Biogeochem. Cy.*, 23, GB4026, doi:10.1029/2008GB003339, 2009
- 725 Reichstein, M., Falge, E., Baldocchi, D., Papale, D., Aubinet, M., Berbigier, P., Bernhofer, C., Buchmann, N., Gilmanov, T., Granier, A., Grünwald, T., Havránková K., Ilvesniemi, H., Janous, D., Knohl, A., Laurila, T., Lohila, A., Loustau, D., Matteucci, G., Meyers, T., Miglietta, F., Ourcival, J.-M., Pumpanen, J., Rambal, S., Rotenberg, E., Sanz, M., Tenhunen, J., Seufert, G., Vaccari, F., Vesala, T., Yakir, D., and Valentini, R.: On the separation of net ecosystem exchange into assimilation and ecosystem respiration: review and improved algorithm, *Global Change Biol.*, 11, 1424-1439,
- 730 10.1111/j.1365-2486.2005.001002.x, 2005.

- Ripullone, F., Grassi, G., Lauteri, M., and Borghetti, M.: Photosynthesis–nitrogen relationships: interpretation of different patterns between *Pseudotsuga menziesii* and *Populus × euroamericana* in a mini-stand experiment, *Tree Physiol.*, 23(2), 137-144, 2003
- Roderick, M. L., Farquhar, G. D., Berry, S. L., and Noble, I. R.: On the direct effect of clouds and atmospheric particles on the productivity and structure of vegetation, *Oecologia*, 129(1), 21-30, 2001
- 735 Saeki, T.: Interrelationships between leaf amount, light distribution and total photosynthesis in a plant community, *Bot. Mag. Tokyo*, 73(860), 55-63, 1960
- Sellers, P. J., Dickinson, R. E., Randall, D. A., Betts, A. K., Hall, F. G., Berry, J. A., Collatz, G. J., Denning, A. S., Mooney, H. A., Nobre, C. A., Sato, N., Field, C. B., and Henderson-Sellers, A.: Modeling the Exchanges of Energy, Water, and Carbon Between Continents and the Atmosphere, *Science*, 275, 502-509, 10.1126/science.275.5299.502, 1997.
- 740 Sitch, S., Huntingford, C., Gedney, N., Levy, P. E., Lomas, M., Piao, S. L., Betts, R., Ciais, P., Cox, P., Friedlingstein, P., Jones, C. D., Prentice, I. C., and Woodward, F. I.: Evaluation of the terrestrial carbon cycle, future plant geography and climate-carbon cycle feedbacks using five Dynamic Global Vegetation Models (DGVMs), *Global Change Biol.*, 14, 2015-2039, 10.1111/j.1365-2486.2008.01626.x, 2008.
- 745 Smith, W.K., Knapp, A.K., Reiners, W.A.: Penumbral effects on sunlight penetration in plant communities. *Ecology* 70,1603–1609, 1989
- Smith, W. K., Vogelmann, T. C., and Critchley, C.: *Photosynthetic adaptation: chloroplast to landscape*. Springer, New York, NY, USA, 2010
- Spitters, C.: Separating the diffuse and direct component of global radiation and its implications for modeling canopy photosynthesis Part II. Calculation of canopy photosynthesis, *Agr. Forest Meteorol.*, 38(1-3), 231-242, 1986
- 750 Spitters, C., Toussaint, H., and Goudriaan, J.: Separating the diffuse and direct component of global radiation and its implications for modeling canopy photosynthesis Part I. Components of incoming radiation, *Agr. Forest Meteorol.*, 38(1-3), 217-229, 1986
- Urban, O., Klem, K., Ač, A., Havráňková, K., Holišová, P., Navrátil, M., Zitová, M., Kozlová, K., Pokorný, R., Šprtová, M., Tomášková, I., Špunda, V., and Grace, J.: Impact of clear and cloudy sky conditions on the vertical distribution of photosynthetic CO<sub>2</sub> uptake within a spruce canopy, *Funct. Ecol.*, 26, 46-55, 10.1111/j.1365-2435.2011.01934.x, 2012.
- Vuichard, N., and Papale, D.: Filling the gaps in meteorological continuous data measured at FLUXNET sites with ERA-Interim reanalysis, *Earth Syst. Sci. Data*, 7(2), 157-171, 2015
- Wang, X., Wu, J., Chen, M., Xu, X., Wang, Z., Wang, B., Wang, C., Piao, S., Lin, W., Miao, G., Deng, M., Qiao, C., Wang, J., Xu, S., and Liu, L.: Field evidences for the positive effects of aerosols on tree growth, *Global Change Biol.*, 24, 4983-4992, 10.1111/gcb.14339, 2018.
- 760 Weiss, A., and Norman, J.: Partitioning solar radiation into direct and diffuse, visible and near-infrared components, *Agr. Forest Meteorol.*, 34(2-3), 205-213, 1985

765

Williams, M., Rastetter, E. B., Van der Pol, L., and Shaver, G. R.: Arctic canopy photosynthetic efficiency enhanced under diffuse light, linked to a reduction in the fraction of the canopy in deep shade, *New Phytol.*, 202(4), 1267-1276, 2014

Wohlfahrt, G., Hammerle, A., Haslwanter, A., Bahn, M., Tappeiner, U., and Cernusca, A.: Disentangling leaf area and environmental effects on the response of the net ecosystem CO<sub>2</sub> exchange to diffuse radiation. *Geophys. Res. Lett.*, 35, L16805, 2008

Yin, X., and Struik, P.: C3 and C4 photosynthesis models: an overview from the perspective of crop modelling, *NJAS-Wagen. J. Life Sc.*, 57(1), 27-38, 2009


770

[Yue, X. and Unger, N.: The Yale Interactive terrestrial Biosphere model version 1.0: description, evaluation and implementation into NASA GISS ModelE2, \*Geosci. Model Dev.\*, 8, 2399–2417, <https://doi.org/10.5194/gmd-8-2399-2015>, 2015.](#)




Zhang, Y., Rossow, W. B., Lacis, A. A., Oinas, V., and Mishchenko, M. I.: Calculation of radiative fluxes from the surface to top of atmosphere based on ISCCP and other global data sets: Refinements of the radiative transfer model and the input data, *J. Geophys. Res.-Atmos.*, 109, D19105, doi:10.1029/2003JD004457, 2004

775


Zhang, Y., Goll, D., Bastos, A., Balkanski, Y., Boucher, O., Cescatti, A., Collier, M., Gasser, T., Ghattas, J., Li, L., Piao, S., Viovy, N., Zhu, D., and Ciais, P.: Increased Global Land Carbon Sink Due to Aerosol-Induced Cooling, *Global Biogeochem. Cy.*, 33, 439-457, 10.1029/2018gb006051, 2019.



**Figure 1: The distribution of light and leaves in canopy. (a) light distribution in ORCHIDEE trunk. (b) distribution of sunlit and shaded leaves in canopy in ORCHIDEE\_DF. (c) light absorbed by shaded leaves in each canopy layer under different solar zenith angle (SZA) and fraction of diffuse light (Fdf) in ORCHIDEE\_DF. (d) Same as (c) but for sunlit leaves.  $I$ , downward PPFD at the top of the canopy;  $I_{abs}$ , PPFD absorption per leaf area in ORCHIDEE trunk;  $I_{shd,abs}$ , PPFD absorption per leaf area in shaded leaves;  $I_{sun,abs}$ , PPFD absorption per leaf area in sunlit leaves.**



790 Figure 2: Modeled and observed diffuse PPFD fraction. The (a) Scatter plot with the dark area indicates high data density, while light area indicates low data density, (b) Density distribution of the observed diffuse PPFD fraction, (c) Density distribution of the modeled diffuse PPFD fraction.



**Figure 3: The diagram of the canopy light transmission in ORCHIDEE\_DF.**  $I_{dr,0}$ , downward direct PPFD at the top of the canopy;  $I_{df,0}$ , downward diffuse PPFD at the top of the canopy;  $LAI_{c,i}$ , cumulative LAI above canopy layer  $i$ ;  $I_{dr,dr,i}$ , downward direct PPFD at the top of canopy layer  $i$ ;  $I_{dr,df,i}$ , net diffuse PPFD derived from the scattering of  $I_{dr,0}$  at the top of canopy layer  $i$ , equals to the difference of its downward ( $I_{dr,df,down,i}$ ) and upward ( $I_{dr,df,up,i}$ ) components;  $I_{df,i}$ , net diffuse PPFD derived from  $I_{df,0}$  at the top of canopy layer  $i$ , equals to the difference of its downward ( $I_{df,down,i}$ ) and upward ( $I_{df,up,i}$ ) components.




Figure 4: Performance of ORCHIDEE trunk and ORCHIDEE\_DF at different PFTs. (a) the Taylor plot of GPP, all valid 30min observations are used as reference, the filled circles indicate ORCHIDEE trunk, opened circles indicate ORCHIDEE\_DF. (b) comparison of the correlation coefficients between the two models against observations. (c) and (e) same as (a) but for cloudy (diffuse light fraction >0.8) and sunny (diffuse light fraction <0.4) conditions only. (d) and (f) same as (b) but for cloudy and sunny conditions only.




Figure 5: Observed GPP and GPP modeled by ORCHIDEE\_trunk and ORCHIDEE\_DF under cloudy (diffuse light fraction >0.8) and sunny (diffuse light fraction <0.4) conditions at selected sites (with relatively long time series) from each PFT.




Figure 6: Site distribution of (a) the GPP difference between cloudy (diffuse light fraction >0.8) and sunny (diffuse light fraction <0.4) conditions. (b) same as (a) but for LUE. ~~It should be noted that the light level is controlled the same for~~The sunny and cloudy conditions ~~time steps are sampled at each site~~equal light levels.







Figure 7: Same as Figure 6 but differentiated for three times of the day.



**Figure 8:** The dependence of LUE difference between cloudy and sunny conditions on climate factors. In observation (blue), ORCHIDEE trunk (red) and ORCHIDEE DF (green), the average and error bars indicate statistics of site level means (a) dependence of LUE difference on temperature, (b) dependence of LUE difference on VPD. (c) and (e) the same as (a) but for only forest sites and short vegetation (grasslands and croplands) sites. (d) and (f) the same as (b) but for forest sites and short vegetation sites.



825 **Figure 9: The distribution of LUE difference between cloudy and sunny conditions (ΔLUE) in temperature-VPD field. The upper numbers in each grid indicate the average of site level ΔLUE, while numbers in brackets indicate the number of sites with valid data. (a) the ΔLUE based on observations, (b) the ΔLUE based on ORCHIDEE trunk, (c) the ΔLUE based on ORCHIDEE\_DF**



830 Figure 10: Same as Figure 9 but for forests (a, c, e) and for short vegetation (b, d, f).




Figure 11: Same as Figure 8a, but for LAI.

835 **Table 1.** Variables in this study

Variable	Definition	Unit
$A$	Net photosynthesis rate	$\mu\text{molCO}_2 \text{ m}^{-2} \text{ s}^{-1}$
$A_e$	Rubisco activity limited net photosynthesis rate	$\mu\text{molCO}_2 \text{ m}^{-2} \text{ s}^{-1}$
$A_t$	Electron transport limited net photosynthesis rate	$\mu\text{molCO}_2 \text{ m}^{-2} \text{ s}^{-1}$
$C_e$	Chloroplast $\text{CO}_2$ partial pressure	$\mu\text{bar}$
$Fdf_{PAR}$	The fraction of diffuse PAR in total PAR	-
$Fdf_{PPFD}$	The fraction of diffuse PPFD in total PPFD	-
$i$	Leaf layer in canopy, for the top layer, $i=1$	-
$I_0$	Downward shortwave radiation at the top of the canopy	$\text{W m}^{-2}$
$I_{abs,df,i}$	Average absorption of $I_{df}$ per unit leaf area in canopy layer $i$	$\mu\text{mol m}^{-2}\text{s}^{-1}$
$I_{abs,dr,df,i}$	Average absorption of $I_{dr,df}$ per unit leaf area in canopy layer $i$	$\mu\text{mol m}^{-2}\text{s}^{-1}$
$I_{abs,dr,dr,i}$	Average absorption of $I_{dr,dr}$ per unit leaf area in canopy layer $i$	$\mu\text{mol m}^{-2}\text{s}^{-1}$
$I_{abs,dr,dr,i,sun}$	Absorption of $I_{dr,dr}$ per sunlit unit leaf area in canopy layer $i$	$\mu\text{mol m}^{-2}\text{s}^{-1}$
$I_{abs,dr,i}$	Average absorption of $I_{dr}$ per unit leaf area in canopy layer $i$	$\mu\text{mol m}^{-2}\text{s}^{-1}$
$I_{abs,i}$	Average radiation absorption per unit leaf area in canopy layer $i$	$\text{W m}^{-2}$
$I_{abs,shd,i}$	PPFD absorbed by shaded leaves per unit leaf area in canopy layer $i$	$\mu\text{mol m}^{-2}\text{s}^{-1}$
$I_{abs,sun,i}$	PPFD absorbed by sunlit leaves per unit leaf area in canopy layer $i$	$\mu\text{mol m}^{-2}\text{s}^{-1}$
$I_{df,0}$	Diffuse downward <u>PPFDphotosynthetic photon flux density</u> at the top of the canopy	$\mu\text{mol m}^{-2}\text{s}^{-1}$
$I_{df,i}$	Net <u>PPFDphotosynthetic photon flux density</u> derived from $I_{df,0}$ at the top of canopy layer $i$	$\mu\text{mol m}^{-2}\text{s}^{-1}$
$I_{dr,0}$	Direct downward <u>PPFDphotosynthetic photon flux density</u> at the top of the canopy	$\mu\text{mol m}^{-2}\text{s}^{-1}$
$I_{dr,df,i}$	Net diffuse <u>PPFDphotosynthetic photon flux density</u> derived from the scattering of $I_{dr,0}$ at the top of canopy layer $i$	$\mu\text{mol m}^{-2}\text{s}^{-1}$
$I_{dr,dr,i}$	Downward direct <u>PPFDphotosynthetic photon flux density</u> at the top of canopy layer $i$	$\mu\text{mol m}^{-2}\text{s}^{-1}$
$I_{dr,i}$	Net PPFD derived from $I_{dr,0}$ at the top of canopy layer $i$ , the sum of $I_{dr,dr,i}$ and $I_{dr,df,i}$	$\mu\text{mol m}^{-2}\text{s}^{-1}$
$I_i$	Downward shortwave radiation arriving at canopy layer $i$	$\text{W m}^{-2}$
$J$	Rate of electron transport	$\mu\text{mol e}^{-} \text{ m}^{-2} \text{ s}^{-1}$
$J_{max}$	Maximum value of $J$ under saturated light, depending on temperature	$\mu\text{mol e}^{-} \text{ m}^{-2} \text{ s}^{-1}$

带格式的: 下标

带格式的: 下标

带格式的: 下标

<i>Jmax<sub>0</sub></i>	<i>Jmax</i> at the top of the canopy	μmol e <sup>-</sup> m <sup>-2</sup> s <sup>-1</sup>
<i>Jmax<sub>i</sub></i>	<i>Jmax</i> at the canopy layer <i>i</i>	μmol e <sup>-</sup> m <sup>-2</sup> s <sup>-1</sup>
<i>k</i>	Light extinction coefficient in ORCHIDEE trunk	-
<i>k<sub>b</sub></i>	Light extinction coefficient when leaves are assumed black	-
<i>k<sub>d</sub></i>	Light extinction coefficient for diffuse PPFD	-
<i>Km<sub>C</sub></i>	Michaelis constants for CO <sub>2</sub> , depending on temperature	μbar
<i>Km<sub>O</sub></i>	Michaelis constants for O <sub>2</sub> , depending on temperature	μbar
<i>LAI<sub>c<sub>i</sub></sub></i>	Cumulative LAI above canopy layer <i>i</i>	m <sup>2</sup> m <sup>-2</sup>
<i>LAI<sub>fshd,i</sub></i>	Fraction of shaded leaf area in total leaf area in canopy layer <i>i</i>	-
<i>LAI<sub>fsum,i</sub></i>	Fraction of sunlit leaf area in total leaf area in canopy layer <i>i</i>	-
<i>m</i>	Optical air mass	-
<i>NIR<sub>p</sub></i>	Potential total downward near infrared radiation at the top of the canopy	W m <sup>-2</sup>
<i>NIR<sub>p,df</sub></i>	Potential diffuse downward near infrared radiation at the top of the canopy	W m <sup>-2</sup>
<i>NIR<sub>p,dr</sub></i>	Potential direct downward near infrared radiation at the top of the canopy	W m <sup>-2</sup>
<i>NIR<sub>TOA</sub></i>	Downward near infrared radiation at the top of the atmosphere	W m <sup>-2</sup>
<i>O</i>	Chloroplast O <sub>2</sub> partial pressure	μbar
<i>p</i>	Air pressure near surface	Pa
<i>p0</i>	Standard sea level air pressure	Pa
<i>PAR<sub>p</sub></i>	Potential total downward photosynthetically active radiation at the top of the canopy	W m <sup>-2</sup>
<i>PAR<sub>p,df</sub></i>	Potential diffuse downward photosynthetically active radiation at the top of the canopy	W m <sup>-2</sup>
<i>PAR<sub>p,dr</sub></i>	Potential direct downward photosynthetically active radiation at the top of the canopy	W m <sup>-2</sup>
<i>PAR<sub>TOA</sub></i>	Downward photosynthetically active radiation at the top of the atmosphere	W m <sup>-2</sup>
<i>PPFDabs<sub>i</sub></i>	Average photosynthetic photon flux density absorption per unit leaf area in canopy layer <i>i</i>	μmol m <sup>-2</sup> s <sup>-1</sup>
<i>PPFD<sub>df, I<sub>df,0</sub></sub></i>	Diffuse downward photosynthetic photon flux density above canopy	μmol m <sup>-2</sup> s <sup>-1</sup>
<i>PPFD<sub>t</sub></i>	Total downward photosynthetic photon flux density above canopy	μmol m <sup>-2</sup> s <sup>-1</sup>
<i>R</i>	Ratio of actual to potential downward shortwave radiation at the top of the canopy	-
<i>R<sub>d</sub></i>	Dark respiration	gC m <sup>-2</sup> s <sup>-1</sup>

带格式的：下标

带格式的：下标

带格式的：字体: 倾斜

带格式的：字体: 倾斜

带格式的：字体: 倾斜

带格式的：字体: 倾斜

带格式的：字体: 倾斜

带格式的：字体: 倾斜

带格式的：字体: 倾斜

带格式的：字体: 倾斜, 下标

带格式的：字体: 倾斜

$SW_{obs}$	Actual (observed) downward shortwave radiation at the top of the canopy	W m <sup>-2</sup>
$SW_p$	Potential (under clearsky conditions without clouds and aerosols) downward shortwave radiation at the top of the canopy	W m <sup>-2</sup>
$V_{cmax}$	Maximum rate of Rubisco activity-limited carboxylation, depending on temperature	μmolCO <sub>2</sub> m <sup>-2</sup> s <sup>-1</sup>
$V_{cmax0}$	Vcmax at the top of the canopy	μmolCO <sub>2</sub> m <sup>-2</sup> s <sup>-1</sup>
$V_{cmaxi}$	Vcmax at the canopy layer i	μmolCO <sub>2</sub> m <sup>-2</sup> s <sup>-1</sup>
$\beta_{df}$	Quanta-to-energy ratio for diffuse PAR	-
$\beta_t$	Quanta-to-energy ratio for total PAR	-
$\Gamma^*$	CO <sub>2</sub> compensation point in the absence of Rd	μbar
$\theta$	Solar zenith angle	degree
$\rho$	The reflection coefficient of the canopy, i.e. the ratio between the downward and upward radiation at the top of the canopy	-
$\rho_w$	Term accounting for atmospheric water vapor absorption	W m <sup>-2</sup>

带格式的: 字体: 倾斜

带格式的: 字体: 倾斜

带格式的: 字体: 倾斜

带格式的: 字体: 倾斜

带格式的: 字体: 倾斜

带格式的: 字体: 倾斜

带格式的: 字体: 倾斜

带格式的: 字体: 倾斜

带格式的: 字体: 倾斜

带格式的: 字体: 倾斜

带格式的: 字体: 倾斜