
NEMO-Bohai 1.0: a high-resolution ocean and sea ice modelling system for the Bohai Sea, China

Yu Yan^{1,2,3}, Wei Gu², Andrea M. U. Gierisch⁴, Yingjun Xu², Petteri Uotila³

¹School of Ocean Sciences, China University of Geosciences, Beijing 100083, China

5 ²Academy of Disaster Reduction and Emergency Management, Faculty of Geographical Science, Beijing Normal University, Beijing 100875, China

³Institute for Atmospheric and Earth System Research (INAR), Faculty of Science, University of Helsinki, 00014 Helsinki, Finland

⁴Danish Meteorological Institute, DK-2100 Copenhagen, Denmark

10 *Correspondence to:* Yu Yan (yuyan@cugb.edu.cn) and Petteri Uotila (petteri.uotila@helsinki.fi)

Abstract. Severe ice ~~condition~~conditions in the Bohai Sea could cause serious harm to maritime traffic, offshore oil exploitation, aquaculture, and other economic activities in the surrounding regions. In addition to providing sea ice forecasts for disaster prevention and risk mitigation, sea ice numerical models could help explain the sea ice variability within the context of climate change in marine ecosystems, such as ~~that of~~ spotted seals, which are the only ice-dependent ~~sea~~ animal that breeds in Chinese waters. Here, we developed NEMO-Bohai, an ocean-ice coupled model based on the Nucleus for European Modelling of the Ocean (NEMO) model version 4.0 and Sea Ice modelling Integrated Initiative (SI³) (NEMO4.0-SI³) for the Bohai Sea. This study will present the scientific design and technical choices of the parameterizations for the NEMO-Bohai model. The model was calibrated and

15 evaluated with in-situ and satellite observations of the ocean and sea ice. The model simulations agree with the observations with respect to sea surface height (SSH), temperature (SST), ~~and salinity (SSS)~~, currents, and temperature and salinity stratification. The seasonal variation of the sea ice area is well simulated by the model compared to the satellite remote sensing data for the period of 1996-2017, ~~and there are similar overall statistics in~~. Overall agreement is found for the occurrence dates of the annual

20 maximum sea ice area. The simulated sea ice thickness and volume are in general agreement with the observations with slight over-estimations. NEMO-Bohai ~~is able to~~can simulate seasonal sea ice evolution and long-term interannual variations. Hence, Nemo-Bohai is ~~intended to be a~~ useful/valuable tool for long-term ocean and ice simulations as well as ~~the ocean and~~ climate change studies.

30 **1. Introduction**

The Bohai Sea is the southernmost seasonal frozen sea in the Northern Hemisphere (Yan et al., 2017). The formation of sea ice in the Bohai Sea mainly depends on the geographical environment and hydrometeorological characteristics (Ding, 1999). Specifically, the Bohai Sea is located in the continental shelf area, and the average water depth is only 18 m (Su and Wang, 2012), which indicates low oceanic heat content in winter. Following a northern continental climate, the Bohai Sea is affected by the cold Siberian air every winter, which causes the sea surface temperature of the Bohai Sea to be significantly lower than that at the same latitude (Zhang et al., 2016; Donlon et al., 2012). In addition, the sea surface salinity of the Bohai Sea is about 30 PSU, which is the lowest in the entire coastal waters in China (Yan et al., 2020). It means that the Bohai seawater freezes before reaching the maximum density. Therefore, it even more easily convects and loses heat just before freezing. The sea ice season is generally from December to March, ~~and the~~. The sea ice in the Bohai Sea also exhibits significant interannual variability, which could cover half of the sea area during severe winters (Su and Wang, 2012).-

With four of China's top ten ports (Tianjin, Tangshan, Dalian, Yingkou), the Bohai Sea is an important economic zone in China (Fu et al., 2017). Under severe sea ice conditions, such as during the 45 2009/2010 winter, 296 ports in this region were frozen. In particular, the 42 km artificial waterway of Huanghua Port was covered by sea ice; thus, the maritime traffic was severely restricted, and 7157 fishing vessels were damaged (Gu et al., 2013). ~~Two hundred seven thousand~~ About 2×10^5 hectares offshore aquatic farms were covered by sea ice, and the aquaculture industry losses reached ~6 billion yuan (~878 million U.S. dollars) (Zhang et al., 2013). In addition, ice floes drift on the Bohai Sea surface under 50 the drive of winds, tides, and ocean currents. When the ice floes meet offshore structures (oil and gas platforms, navigation lights, and lighthouses) or ships, convergent ice motion seriously threatens marine transportation and oil and gas exploration (Leppäranta, 2011; Li et al., 2011; Yan et al., 2019). Thus, sea ice monitoring and forecasting is essential for assessing sea-ice-hazard risk and preventing ice disasters.

55 In addition, spotted seals (*Phoca largha*), ~~which are distributed in the Northwest Pacific Ocean where Liaodong Bay of the Bohai Sea represents the southernmost geographic breeding concentration of this species, are listed as~~ are listed as the least concern species by the International Union for the Conservation of Nature Red List of Threatened Species and as a Grade II state protection species by

China's Wild Animal Protection Law (1988-), ~~which are distributed in the Northwest Pacific Ocean with~~
60 ~~the Liaodong Bay of the Bohai Sea as the southernmost geographic breeding site.~~ At the end of October
each year, the seals traveled from the Yellow Sea to Liaodong Bay. However, the population of the spotted
seal has experienced several drastic declines from ~2300 to ~1000 individuals over the past three decades
due to hunting and environmental pollution (Yan et al., 2018; Yang et al., 2017). They den in coastal
areas and on offshore drifting ice. When the sea ice melts, they gradually leave Liaodong Bay and embark
65 on a long journey back to the Pacific Ocean. Therefore, ~~the realization of it is crucial to study~~ long-term
changes ~~in of~~ Bohai Sea ice ~~is crucial to learn how sea ice changes impact to study its impacts on~~ spotted
seals.

Most research ~~have~~has focused on the polar region, however, sea ice in mid-high latitude is also
sensitive to climate change (Bai et al., 2011; Gong et al., 2007; Yan et al., 2017). Knowledge of regional
70 sea ice variability is ~~important~~vital to ~~study~~studying how climate change will affect regional basins. The
traditional view on the Bohai Sea ice is concentrated on extracting sea ice data from satellite imagery
(Karvonen et al., 2017; Shi and Wang, 2012; Su et al., 2019). As the use of remote sensing data to retrieve
sea ice data has limitations in providing a ~~consistent~~ continuous record for a long time series, numerical
~~modeling~~modelling is an effective tool to understand the sea ice ~~continuous process~~development from
75 freezing to thawing (Hordoier et al., 2019; Pemberton et al., 2017).

Sea ice models can reasonably simulate sea ice conditions, which can ~~largely~~primarily supplement
remote Earth observations. To our knowledge, there are only very few studies on developing regional
coupled ocean-ice models for the Bohai Sea, and most of them have focused on short-duration
simulations, such as one-week ~~and~~or one-year case studies. Wang et al. (1984) reported the first sea ice
80 dynamic-thermodynamic model for the Bohai Sea, which simulated the sea ice freezing-melting cycle at
a resolution of 20 km×20 km driven by monthly climatic forcing. Wu (1991) and Wu et al. (1997) tried
to simulate sea ice evolution based on a dynamic-thermodynamic model with consideration of the ice
thickness distribution function consisting of three types (level ice, drift ice, and open water). Bai and Wu
(1998) combined the sea ice dynamic-thermodynamic model, atmospheric boundary layer model, and
85 tide model at a resolution of 8.64 km×11.11 km. ~~The model was initialized with sea ice data from NOAA
and driven by weather forecast data from the Japan Meteorological Agency for each 4 day period
between 1990 and 1996.~~ Liu et al. (2003) applied a high-resolution atmosphere model (T213,

~~0.5625°×0.5625°~~ to reproduce sea ice conditions in 2002 by a sea ice model initialized by high-resolution satellite data from HY-1 and MODIS (~~250 m~~). Su et al. (~~2004~~) simulated the winters of 1998 and 2000 with the coupled Princeton Ocean Model (POM) and viscous-plastic dynamic-thermodynamic sea ice model, and simulated the winters of 1998 and 2000. Tang et al. (2010) simulate the sea ice freezing-melting process in 2003 with the coupled POM with a sea ice model considering the thickness distribution function, which was driven by MM5 atmospheric forcing provided by the NMEFC, to simulate the sea ice freezing-melting process in 2003. Zhang and Zhang (2013) used the finite-volume, primitive equation community ocean model to simulate sea ice in 2003. Li et al. (2021) simulated sea ice variations in the severe winter of 2009–2010 in the Bohai Sea based on version 3.6 of the Nucleus for European Modelling of the Ocean (NEMO). To the best of our knowledge, there is has been no sea long-term simulation study on Bohai Sea ice prediction system for the Bohai Sea for a long enough time to reproduce so far aiming to understand the interannual or interdecadal variability.

The Nucleus for European Modelling of the Ocean (NEMO) is a state-of-the-art numerical modeling framework designed for research activities and forecasting services in the ocean and climate sciences (Madec et al., 2016). The NEMO code, including its sea ice component, enables the investigation of ocean-ice dynamics/thermodynamics and their interactions with the atmosphere. NEMO offers a wide range of applications from short-term forecasts and climate projections (Drouard and Cassou, 2019; Obermann-Hellhund et al., 2018; Uotila et al., 2017; Voltaire et al., 2013) to process studies (Courtois et al., 2017; Declerck et al., 2016; Feucher et al., 2019). While global models offer a poor representation of regional ocean processes, regional models have been developed based on the framework of NEMO in recent years for various seas, e.g., models for the Atlantic marginal basins (Graham et al., 2018; O'Dea et al., 2017), the Baltic and North seas (Hordoier et al., 2019; Pemberton et al., 2017), the Northwestern Mediterranean Sea (Declerck et al., 2016), the South Indian Ocean (Schwarzkopf et al., 2019), the South China Sea (Thompson et al., 2019), and the Black Sea (Gunduz et al., 2020). However, a NEMO-based regional model for the Bohai Sea based on NEMO has not yet been developed, attempted for long-term climate studies until now.

Clearly, a coupled ocean-ice model Long-term sea ice simulations could help improve our understanding of thermodynamic and dynamic sea ice processes in the Bohai Sea, which is crucial for sea ice disaster prevention, spotted seal habitat studies, and regional climate change studies. The aim of

the paper [is aims](#) to report the development of NEMO-Bohai and assess its performance. The paper is organized as follows: Sec. 2 introduces the model and observation dataset. Model comparison and validation [is are](#) carried out in Sec. 3. The analysis of sea ice variation based on the 22-year hindcast simulations of NEMO-Bohai is presented in Sec. 4. Discussion and a ~~short~~ summary are provided in Sec. 5.

2. Model setup: NEMO-Bohai

NEMO-Bohai is localized to the Bohai Sea based on the NEMO ocean engine (Madec et al., 2016). ~~Here, we~~We apply the NEMO 4.0. ~~b~~ [beta](#) and Sea Ice Integrated Initiative (SI³) model in a regional configuration covering the Bohai Sea. The regional configuration, NEMO-Bohai, is nested into the global configuration (~~ORCA025~~) using one-way lateral boundary conditions with the flow relaxation scheme algorithm (Engedahl, 1995). ~~The~~The sea surface heights (SSHs), temperatures, salinity levels, and velocities are relaxed to outer model values over a 1-point zone at the unstructured open boundary of the model domain. ~~In addition, NEMO 4.0 and the use of the XIOS server is a key element in a more user-friendly version of NEMO-Bohai.~~

2.1 Global ocean-ice model ORCA025

A global 1/4° ocean-ice model (ORCA025) maintained by the DRAKKAR group is ~~presented. The fundamental conservation laws of mass, momentum and energy in NEMO are resolved in a curvilinear coordinate system with discretization on a staggered grid (Arakawa C grid). The horizontal grid has a 0.25° resolution at the equator, leading to 1442×1207 grid points horizontally, and there are 75 vertical levels in the model configuration with a grid spacing between 1–200 m from the top to the bottom.~~ performed. A detailed description of ORCA025 is provided by Bernard et al. (2006). In our simulation, forcing fields are provided from the Japanese 55-year Reanalysis (Harada et al., 2016; Kobayashi et al., 2015), covering the 55 years from 1958 with 0.5°×0.5° spatial resolution. ~~The variables include of wind speed (u/v component), 2-m air temperature, surface downwelling shortwave/longwave radiation, near surface specific humidity, rainfall flux, snowfall flux, and sea level pressure at 3-hourly intervals. The generic bulk formula (Large and Yeager, 2009) is used for the surface boundary condition (SBC) and the frequency of the SBC and sea ice model calls every time step (ea. every 15 min).~~ A monthly

climatological runoff flux with Antarctic ice shelves and iceberg meltwater flux (Dai et al., 2009; Depoorter et al., 2013) is used. ~~Penetrative solar radiation is applied through the RGB light penetration formulation to split visible light to Red Green Blue wavebands.~~ To limit the atmospheric forcing biases from propagating into the upper ocean and avoid salinity and temperature drift, which impacts the overturning circulation, the sea surface temperature (SST) and salinity (SSS) are restored monthly based on the World Ocean Atlas-2013 (Locarnini et al., 2013; Zweng et al., 2013).

~~The split explicit free surface formulation is implemented, and no-slip lateral boundary conditions are chosen. The linear top/bottom friction parameterization, which assumes that the top/bottom friction is proportional to the interior velocity, is applied in the simulation. The Laplacian operator is used in the iso-level direction in the lateral diffusion scheme for tracers, and the eddy coefficient is set to a constant. Tracer advection is achieved using a flux-corrected transport (FCT) scheme (Zalesak, 1979). The turbulent kinetic energy (TKE) closure scheme (Blanke and Delecluse, 1993) is used for vertical physics setting. The simulation is spun up in January 1987 with~~ The global simulation is spun up in January 1987 ~~with the~~ rest state and ended in 2017. The time step is set to 900 s. The ORCA025 configuration is performed using 240 CPU cores on a Cray XC40 system on the Sisu supercomputer and requires ~6000 CPU hours per simulated year. The model output frequency is set to every ~~five~~ days ~~for the period~~ from 1987 to 1994 for the test run and increases to every day for the period of 1995–2017. ~~We generate approximately 2 TB~~ The size of the output files per simulated year ~~for the period of 1995–2017. For the regional model, is approximately 2 TB. The open boundary for NEMO-Bohai, the interface between open boundaries~~ is forced by barotropic and baroclinic modes. The sea surface heights (SSHs), temperatures, salinity levels, and velocities are relaxed to outer-model values over a 1-point zone at the unstructured open boundary of the model domain. ~~In addition, the use of the XIOS server makes NEMO-Bohai more user-friendly.~~

2.21 NEMO-Bohai: The ocean model

The NEMO-Bohai domain consists of one central zone and three bays (see Fig. 1). The area covers 117°–122° E and 37°–41° N. The horizontal resolution is 0.025° in orthogonal curvilinear coordinates, which is equivalent to a resolution of approximately 2 km ($204 \times 244 = 49\,776$ grid points). The bathymetry is interpolating into the target grids according to the ETOPO1 1 arc-minute dataset. We further manually remove the isolated land cells with a number less than 4. The vertical discretization

scheme adopts the z^* formulation. There are 10 vertical levels, and the top layer is set to 1 m. The partial step is applied to the bottom topography. The time step of the ocean model is set to 90 s, while it calls the sea ice model every three time steps. The split-explicit time stepping method (`ln_dynspg_ts = .true.`) is used to compute barotropic/baroclinic mode and their interactions.

175





Figure 1: The location and bathymetry of the Bohai Sea. The black dots denote the coastal tide gauge stations. The red dots and black triangles indicate the station locations with sea surface temperature observations. The red dots and black triangles indicate the station locations with sea surface salinity observations, respectively; black squares denote the oil platforms where the sea ice thickness observations were conducted. The black triangles indicate Line AB

180

185 represents the profile section along which temperature and salinity were presented (figures 6 and 7, respectively), while line CD indicates the sea-surface salinity-observation-stations. Bohai Strait. The straight line in the east denotes an open boundary for the regional ocean model.

The initial conditions of the numerical simulation in the Bohai Sea for July 1, 1995, including SST and SSS, are obtained by interpolating the temperature and salinity fields from the global ocean simulations with the ORCA025 configuration, as shown in Fig. 2. The regional model is forced with lateral ocean boundary conditions, tidal forcing, atmospheric forcing, and river runoff during the study period. Two kinds of boundary ~~condition~~conditions are used for the setting of lateral open ocean boundary. Flow relaxation scheme is applied to baroclinic velocities and tracers (Engedahl, 1995), while Flather boundary condition (Flather, 1976) is used for barotropic dynamics, such as SSH and barotropic velocities ($u2d$, $v2d$). NEMO-Bohai has one open boundary located 100 km away from the Bohai Strait, and we set the relaxation zone width to 1 ($nn_rimwidth = 1$). The lateral open ocean boundary conditions for 1995 to 2017 are also taken from a series of global ocean simulations with the ORCA025 hindcast. The Oregon State University Tidal Inversion Model is used for the barotropic mode (Egbert and Erofeeva, 2002), ~~which includes~~including 11 tidal harmonics (M2, S2, N2, K2, K1, O1, P1, Q1, M4, MS4, MN4). We choose the TPXO9.1 global tide model, which is the latest version with a $0.25^\circ \times 0.25^\circ$ resolution.



200 **Figure 2: Examples of (a) initial sea surface temperature (SST) field and (b) initial sea surface salinity (SSS) field interpolated from the global ORCA025 model simulation on July 1, 1995.**

The NEMO-Bohai atmospheric forcing is derived from the ERA5 dataset, ~~which is~~ the ECMWF's latest reanalysis product covering the period from 1979 to the present ~~and, which~~ has replaced the widely

205 used ERA-Interim dataset. The data cover the Earth with a horizontal spatial resolution of 30 km and represent the atmosphere using 137 levels from the surface up to a height equaling 0.01 hPa. The forcing files consist of 1 hourly instantaneous ~~fieldsfield~~ of the 2 m air temperature, 10 m wind speed, downward short/long wave radiation, sea-level pressure, specific humidity, and precipitation. The equivalent inverse barometer SSH is calculated from the atmospheric pressure (`ln_apr_dyn = .true.`) with the
 210 referenced pressure of $101,000 \text{ N m}^{-2}$. In addition, the river runoff, which provides a significant volume of freshwater to the Bohai Sea and influences stratification, circulation, and primary productivity, is considered in the model. The daily runoff volume data for the rivers flowing into the Bohai Sea, such as the Yellow River, Liao River, and Hai River, are acquired from JRA-55 from 1995 to 2018. The runoff forcing is considered as ~~fresh-water~~freshwater with a salinity of $0 \text{ g}\cdot\text{kg}^{-1}\text{PSU}$ and ~~the~~ same temperature
 215 as the ocean surface.

The bottom roughness impacts the dynamics of the tide, ocean circulation, and storm surges in the Bohai Sea. A constant bottom roughness (`rn_z0`) is used to calculate the drag coefficient for all bottom grids. To better reflect changes in the tide and circulation, we tune the bottom roughness accordingly. ~~With a reference to~~Regarding the configuration in the Gulf of Finland (`rn_z0 = 5.0\times 10^{-3} \text{ m}`)
 220 (Westerlund, 2019), we set `rn_z0` to 5.0×10^{-1} , 5.0×10^{-2} , 5.0×10^{-3} , 5.0×10^{-4} , 5.0×10^{-5} , and 5.0×10^{-8} while keeping all the other variables consistent. Results show that reducing the bottom friction can increase the phase difference and amplitude of the tide. In NEMO-Bohai, the smaller the roughness value is, the more accurate the tide is. However, it also needs to be combined with the actual situation at the bottom of the Bohai Sea. With reference to (Zhang and Zhang, ~~2013~~) (2013), the bottom roughness had a
 225 magnitude of $1\times 10^{-3} \text{ m}$. Therefore, the bottom roughness in NEMO-Bohai is determined to be $5.0 \times 10^{-3} \text{ m}$. ~~For the~~The bottom drag coefficient, ~~we set~~ `(rn_Cd0)` is set to the default value after a set of experiments ~~similar to the abovementioned~~. The turbulent kinetic energy (TKE) closure scheme is used for vertical mixing (Blanke and Delecluse, 1993) with the default values of vertical eddy viscosity and diffusivity coefficients, and the ~~FCT~~flux-corrected transport scheme (Zalesak, 1979) is responsible for
 230 the tracer advection. Some keycritical parameters of the ocean part of NEMO-Bohai are listed in Table 1.

Table 1: Key physical parameters in the ocean namelist (`namelist_oce`) in NEMO-Bohai.

Parameter	Namelist	Setting	Parameter	Namelist	Setting
-----------	----------	---------	-----------	----------	---------

Time step	rn_rdt	90 s	Activate tides	ln_tide	true
Frequency of surface module call	nn_fsbc	3	Hydrostatic pressure gradient option	ln_hpg_sco	true
Relaxation zone width	nn_rimwidth	1	Ocean equation of seawater	ln_eos80	true
Bottom roughness	rn_z0	5.0×10^{-3} m	Penetrative solar radiation	ln_qsr_rgb	true
Top drag coefficient	rn_Cd0	1.0×10^{-3}	Advection scheme for tracer	ln_traadv_fct	true
Bottom drag coefficient	rn_Cd0	1.0×10^{-3}	Standard operator of lateral diffusion scheme for tracers	ln_traldf_iso	true
Lateral momentum boundary condition	rn_shlat	0	Laplacian operator of lateral diffusion on momentum	ln_dynldf_lap	true
Vertical eddy viscosity	rn_avm0	$1.2 \times 10^{-4} \text{ m}^2 \cdot \text{s}^{-1}$	Surface pressure gradient	ln_dynspg_ts	true
Vertical eddy diffusivity	rn_avt0	$1.2 \times 10^{-5} \text{ m}^2 \cdot \text{s}^{-1}$	Vertical physics	ln_zdftke	true
Ocean initialization	ln_tsd_init	true	Top/bottom drag coefficient	ln_loglayer	true

2.32 NEMO-Bohai: The sea ice model

SI³ is recommended by the Sea Ice Working Group to reduce duplication and better use development resources (Aksenov et al., 2019). SI³ merges the capabilities of three sea ice models previously used in NEMO (CICE, GELATO, and LIM). SI³ is a horizontal dynamic and vertical thermodynamic sea ice model. The thermodynamics and dynamics are separated and rely upon different frameworks and sets of hypotheses: thermodynamics ~~uses~~ the ice thickness distribution and the mushy-layer frameworks, whereas dynamics ~~assume~~ continuum mechanics, e.g., Leppäranta (2011). Thermodynamics and dynamics interact in two ways: first, advection impacts the ice state variables; second, the horizontal momentum equation depends on, among other things, the ice state. The modified elastic-viscous-plastic rheology is used for sea ice dynamics (Bouillon et al., 2013; Kimmritz et al., 2017). Bohai Sea ice is relatively thin and has low salinity, ~~which is~~ different ~~with~~ polar ice (Gu et al., 2013). Therefore, the parameters in NEMO-Bohai ~~is~~ required to be modified accordingly.

In NEMO-Bohai, we selected and adjusted a series of sea ice model parameters. We increase the number of ice categories (jpl) and the number of ice layers (nlay_i) since Massonnet et al. (2019) suggested that increasing the resolution of the ice thickness distribution results in small but non-negligible variations in the ice extent and volume. The ice thickness is discretized into ten categories (0.0, 0.0476, 0.0976, 0.150, 0.205, 0.263, 0.324, 0.388, 0.455, 0.526, 0.601 m), and the thermodynamic calculations used five ice layers (nlay_i = 5). The minimum ice thickness used in remapping is set at the minimum value (rn_himin = 0.01 m) ~~so that we can better~~ capture new thin ice ~~formation. Thickness formations better. The thickness~~ of new ice formed in open waters (rn_hinew) is set to 0.02

m as the value must be larger than `rn_hnewice`. According to the test run of NEMO-Bohai, the sea ice thickness is generally overestimated. Compared to the referenced value of 2.0 m, we reset the average
255 ice thickness (`rn_himean`) as 0.20 m according to the observations (Li et al., 2004; Yuan et al., 2012).

The ice initialization is activated (`ln_iceini = .true.`), and the initial ice salinity (`rn_smi_ini_n`) is set to ~~7 $\text{g}\cdot\text{kg}^{-1}$~~ PSU, and the initial real snow thickness (`rn_hts_ini_n`) is set to 0.1 m according to in-situ observations (Bai and Wu, 1998; Gu et al., 2014). For the snow and ice albedos, we set them to lower values compared to the referenced ones based on the in-situ observations
260 (Zheng et al., 2017). The ice strength parameter (`rn_pstar`) is defined as $2.75 \times 10^4 \text{ N m}^{-2}$ based on previous in-situ studies (Li et al., 2011). A ridging scheme is considered in SI³, ~~thus he~~. Thus, the parameter `rn_hstar`, which adjusts the maximum thickness of ridged ice, is reduced from 100.0 m to 10.0 m following the NEMO-NORDIC 1.0 configuration (Pemberton et al., 2017) to reflect the general thinner Bohai Sea ice. In addition, the free-slip lateral boundary condition is chosen for sea ice dynamics
265 (`rn_ishlat = 0.`), which is synchronized with the ocean model, and a landfast parameterization is set (`ln_landfast_L16 = .true.`) because there exists landfast ice in the eastern Liaodong Bay.

3. Model comparison and validation

270 In this section, we analyze how well NEMO-Bohai reproduces ocean properties (SSH, SST, SSS, temperature and SSSsalinity stratification, currents and volume exchanges with the Yellow Sea) and sea ice properties (area, thickness, and volume). To evaluate the model's performance, we compare our model results to in-multiple sources in-situ and satellite observations from 1996 to 2017.

3.1 Observational data

275 To evaluate the SSH, the modeled tide amplitude and phase were compared to the tide tables with hourly intervals of Yantai (37.550° N, 121.383° E), Dalian (38.867° N, 121.683° E), and Jingtanggang (39.217° N, 119.017° E) from National Marine Data and Information Service during the period of October 1 to 31, 2012. In addition, we use in-situ-observed in-situ data from the National Marine Environmental Monitoring Center. The sampling station is at Bayuquan (39.804° N, 121.456° E), and the sampling period is from January 1 to 31, 2013. For the SST evaluation, we use data from the National
280 Marine Science Data Center. The coastal stations are located at Laohutan in Dalian (38.90° N, 121.70° E) and Zhifudao in Yantai (37.60° N, 121.40° E). The observed records cover the periods of 2000–2001, 2010 (July–December), and 2011–2015. For the sea surface salinity, the in-situ-observed in-situ data for eight ocean stations (Wentuozi, Huludao, Zhimaowan, Qinhuangdao, Tanggu, Longkou, Penglai, Beihuangcheng) in the Bohai Sea isare from Yuan et al. (2015). Specifically, the observation period in
285 Huludao, Zhimaowan, Qinhuangdao, Tanggu, Longkou, and Beihuangcheng is from 1960 to 2013, while the period in Wentuozi and Penglai is 1987-2001 and 2012-2013, respectively. The climatological vertical profiles of temperature and salinity are from the marine atlas of the Bohai Sea based on the observed data from the 1950s to 1990s (Chen, 1992).

290 The sea ice model is evaluated byagainst a series of observational datasets. A satellite-derived dataset eoveredcovering the winters from 1988 to 2017 was retrieved. Sea ice area is-extractingwas extracted from two datasets; the first based on the zonal threshold method for Advanced Very High Resolution Radiometer (AVHRR) during 1988-2000, while; and the second based on an object-based feature extraction method is-applied for Moderate Resolution Imaging Spectroradiometer (MODIS) from 2001 to 2017, and the. A detailed description can be seen fromfound in Yan et al. (2017). The shapefiles
295 derived by the abovementioned methods are further modified by visual interpretation with-the aim-to

build a more accurate sea ice area dataset. For the sea ice thickness evaluation, we use the in-situ observations from four offshore oil platforms (see Fig. 4) on January 2, 6, 9, 16, 26, February 2, 9, 12, 16 during 2012–2013 (Zeng et al., 2016; Karvonen et al., 2017). The data were collected on January 2, 6, 9, 16, 26, February 2, 9, 12, 16.


300 3.2 Ocean

3.2.1 Sea surface height

Reliable precision of the tidal model is pre-required for subsequent simulations in NEMO-Bohai. The tidal model is validated by in-situ hourly tidal observations at Bayuquan and tide table data at Yantai, Dalian, and Jingtanggang. Fig. 3 provides a robust description of the comparison of SSH time evolution at hourly intervals at the four stations. The tidal range in the Bohai Strait (Yantai and Dalian) is larger than that at other tide gauging stations. It is clear that the modeled water elevation at Bayuquan and Jingtanggang stations agrees less with the simulated results. As presented in tide table data than at the other two stations, as shown in Table 2. Nevertheless, the Fig. 3; model reproduced well the semidiurnal M2 tidal cycle with exact phase and amplitude at all four stations. The with a mean correlation of 0.92, mean absolute error of 0.17 m, and root-mean-square error of 0.22 m. The model reproduces SSH standard deviation well, and the difference is within 6 cm in Yantai, Dalian, and Jingtanggang (Table 2); the largest deviation at Bayuquan is approximately 11 cm. Simulated SSH depends on the model's bathymetry, open boundary forcing, and freshwater flux (Kärnä et al., 2021). The errors are partly due to the deviation of positions between the tidal gauges and the model grid points, as the model has a horizontal grid size of ~2 km. Also, errors can be partly due to the inaccurate bottom topography, especially in the shallow water depth zone. As shown in Fig. 6, there is a clear topographic error between the model and the atlas for the four stations are 0.14, 0.18, 0.17 and 0.20 m, respectively. The profile section AB. In general, the reasonable accuracy of the hydrodynamic model can partially assure the rather realistic simulations of the sea ice process of growth and melting, as described in the following section 3.3.

Table 2: Sea surface height representation, in terms of standard deviation (SD, meters), correlation, mean absolute error (MAE, meters), and root-mean-square error (RMSE, meters), made by NEMO-Bohai for four Bohai Sea stations.

<u>Station</u>	<u>SD Observation</u> <u>(meters)</u>	<u>SD NEMO-Bohai</u> <u>(meters)</u>	<u>Correlation</u>	<u>MAE</u> <u>(meters)</u>	<u>RMSE</u> <u>(meters)</u>
<u>Yantai</u>	<u>0.65</u>	<u>0.59</u>	<u>0.96</u>	<u>0.14</u>	<u>0.18</u>
<u>Dalian</u>	<u>0.81</u>	<u>0.85</u>	<u>0.96</u>	<u>0.18</u>	<u>0.23</u>
<u>Jingtanggang</u>	<u>0.36</u>	<u>0.37</u>	<u>0.84</u>	<u>0.17</u>	<u>0.21</u>
<u>Bayuquan</u>	<u>0.61</u>	<u>0.50</u>	<u>0.92</u>	<u>0.20</u>	<u>0.25</u>



325 **Figure 3: Comparison of sea surface height (SSH) between the observations and NEMO-Bohai simulations at (a) Yantai, (b) Dalian, (c) Jingtanggang, and (d) Bayuquan stations in the Bohai Sea.**

3.2.2 Sea surface temperature and salinity

As shown in Fig. 4, the modeled SST followed well the seasonal cycle, and the mean absolute error (MAE) was typically less than 1°C at Laohutan and Zhifudao stations during the observed period. The modeled variations of SST are smoother than the observations, observed, and they particularly failed fail to capture the high fluctuations of daily variations in summer/autumn at both stations. It is also noticeable that the simulated SST at Zhifudao are is generally colder in winter and warmer in summer than the observations.

330




Figure 4: The simulated sea surface temperature (SST) compared with in-situ observations at Laohutan, Dalian (a-h) and Zhifudao, and Yantai (i-p) for the period of 2000-2001 and 2010-2015, respectively.

Table 3 displays the comparisons of multi-year average sea surface salinities at ~~8~~eight ocean stations in the Bohai Sea between simulations from NEMO-Bohai and observations reported by Yuan et al. (2015). Generally speaking, NEMO-Bohai is able to capture the main variations of the sea surface salinity in the Bohai Sea. Values at six ocean stations agree with observations with the relative deviation less than 5%, while the modeled values at Huludao and Tanggu are less salty than observed. This demonstrates that the model faces ~~larger~~more significant challenges in the low salinity areas. ~~As we know~~Primarily, the freshwater ~~inputs from rivers would lead~~river runoff leads to lower salinity in the coastal ~~areas~~regions. In Nemo-Bohai, the ~~model, the river~~river runoff is based on climatological estimates without interannual variability. The ~~river~~ salinity ~~in NEMO-Bohai is nevertheless~~is assumed to be $0 \text{ g}\cdot\text{kg}^{-1}$, ~~which~~PSU, and ~~the river temperature is set to the same value as the SST at the closest grid point~~. ~~All these reasons~~ might cause underestimations. In addition, the biases of river runoff and shallow water depth (generally $< 3 \text{ m}$) also need to be taken into considerations.

Table 3: Comparison of multi-year average sea surface salinity between the observations and simulations.

Station	longitude (°)	Latitude (°)	Observations (g·kg ⁻¹ PSU)	Simulations* (g·kg ⁻¹ PSU)	Relative deviation
Wentuozi	121.41	39.72	30.5	30.1	-1.3%
Huludao	120.81	40.56	29.6	23.2	-21.6%
Zhimaowan	119.92	40.02	30.8	31.0	+0.6%
Qinhuangdao	119.62	39.92	30.4	31.0	+0.6%
Tanggu	117.78	38.98	29.3	25.4	-13.3%
Longkou	120.32	37.65	28.2	28.0	-0.7%
Penglai	120.73	37.82	30.2	30.4	+0.6%
Beihuangcheng	120.92	38.40	31.5	32.8	+4.1%

*The observed sea surface salinity data is referenced from Yuan et al. (2015).

3.2.3 Sea current

355 The circulation in the Bohai Sea is mainly barotropic and results from the combined effects of tides and winds (Li et al., 2015). The simulated monthly mean current velocities at the surface and 16 m depth in February and August are shown in Fig. 5. The monthly mean current velocities are calculated based on hourly model output during August 2012 and February 2013. The figure shows that both the sea surface and 16 m depth current are usually less than 0.4 m s⁻¹. Due to the blocking effects of the bays,
360 currents are weak at the head of the three bays, which is consistent with the observations by Chen et al. (1992). The surface currents in the Bohai Sea also show apparent seasonal variations. In August, relatively strong currents flow from the southeastern part of Liaodong Bay to the southern Bohai Strait with a solid outflow to the Yellow Sea. Specifically, the current field distribution at Bohai Strait shows that the inflow is mainly in the narrow channel in its northern part, which agrees with the observations of Wan et al. (2015). Also, Lin et al. (2011) suggested that persistent winds drive a cyclonic coastal current into the northern Yellow Sea, and one branch of the current enters the Bohai Sea at the northern Bohai Strait, which transports warm and saline water from the Yellow Sea.

365
370 The monthly mean water volume exchange at the Bohai Strait (see Fig.1) based on hourly model simulations during August 2012 and February 2013 are also calculated to evaluate the model's performance. The Bohai Sea water exchange with the Yellow Sea is weak due to its half-closed shape and a relatively independent circulation system. The model results show that the inflow from the Yellow

Sea to the Bohai Sea in August reaches $6.9 \times 10^4 \text{ m}^3 \text{ s}^{-1}$, almost double than that in February ($3.5 \times 10^4 \text{ m}^3 \text{ s}^{-1}$), which lies in the range of 5×10^3 to $8 \times 10^4 \text{ m}^3 \text{ s}^{-1}$ indicated by Bian et al. (2016). Our results show that the outflow exists in both months (August: $8.3 \times 10^4 \text{ m}^3 \text{ s}^{-1}$; February: $6.5 \times 10^4 \text{ m}^3 \text{ s}^{-1}$), with a larger amount than the inflow. The net flow appears to be outflow both in winter and summer, which is consistent with other model simulation results (Lin et al., 2002; Ji et al., 2019).




Figure 5: Simulated monthly mean current velocities at surface and 16 m depth in August 2012 and February 2013. The monthly mean current velocities are calculated based on the outputs with hourly intervals.

3.2.4 Vertical profile

NEMO-Bohai and observed water temperature and salinity profiles along the transect AB (see Fig.1) are shown in Fig. 6 and Fig. 7, respectively. Observations are from the atlas by Chen (1992), which is based on data from the 1950s to 1990s. The temporally closest 5-year period from 1995 to 2000 of NEMO-Bohai simulations was selected for model-observation comparisons. Common features are found both in the model and observations. The Bohai Sea waters are vertically well-mixed in autumn and winter, and they have a remarkable homogeneous vertical distribution for both temperature and salinity. In spring and summer, thermal stratification occurs with a significant cold-water core at depth, eventually eroded in autumn. As apparent in Fig. 6 and Fig. 7, the stratification in shallow coastal waters is generally homogeneous. Similar features were reported by Wang et al. (2008), who analyzed the seasonal

variations of the vertical profiles in the Bohai Sea.

The model results, however, show some discrepancies compared to the atlas. Although the model reproduces the summer saline stratification, it is weaker than in the atlas. Nonetheless, Li et al. (2015) reported that the summer salinity stratification in the Bohai Sea is possibly weaker than in the atlas, with an observed top-to-bottom salinity difference of 0.6 PSU. The modeled salinity distribution along with transect AB during summer is possibly affected by the high vertical diffusivity. In the north part of the transect, which corresponds to the northern Liaodong Bay, a negative salinity bias is visible compared to the atlas. In addition to the reasons mentioned in section 3.2.2, inaccuracies in the ETOPO1 bathymetry, especially in the low water depth region seen from Fig. 6 and Fig. 7, may also cause these underestimations.





Figure 6: Comparison of vertical profiles of water temperature (°C) along with transect AB (locations shown in figure 1) between NEMO-Bohai (a-d) and the atlas (Chen, 1992) (e-h) in February, May, August, and November.




405 Figure 7: Similar to figure 6 but for salinity (PSU).

3.3 Sea ice

Sea ice is the focus of our study and is, in this section, directly validated against observations with respect to the area, thickness, volume, and their variations.

3.3.1 Sea ice area

410 To evaluate the performance of NEMO-Bohai in simulating the characteristics of sea ice area, a series of variables were validated, including the daily sea ice area (DSIA), annual maximum sea ice area (AMSIA), and spatial patterns. ~~Figure 5~~Fig. 8 shows that the model-estimated DSIA agrees well with the satellite-derived observations ($r = 0.85$, $p < 0.01$). In addition to the good agreement of inter-annual variability, the occurrence dates of the AMSIA are quite close to the observed dates from 1996 to 2017, as listed in Table 4. The percentage of the occurrence dates of the AMSIA less than 3 days accounted for 72.7%, and the mean absolute error is 3.5 days. In 2013, the ~~NEMO-Bohai-based~~ maximum sea ice area was simulated ~~as occurring~~ on February 9, 2013, which is only ~~one~~ day later than reported (North China Sea Branch of State Oceanic Administration, 2013). ~~The~~However, the date retrieved from MODIS was ~~however~~ determined on January 17 ~~due to missing as the~~ data on February 8, 2013 ~~as was missing~~ due to severe cloud ~~covered~~cover (Yan et al., 2017). The largest difference happened in the winter of 2010 with 18 days earlier than the observations. Specifically, DSIA exhibited a ~~bi-modal~~bimodal distribution with two remarkable peaks in ~~the~~ winter of 2010. Satellite showed that the AMSIA happened on February 12, followed by the second maximum on January 26, while NEMO-Bohai simulated AMSIA was on January 26, followed by ~~the~~ second maximum on February 12. It is worth noting that ~~the~~ satellite shows the largest DSIA happened in ~~the~~ winter of 2010, while NEMO-Bohai simulated DSIA reached the maximum in 2011. On the contrary, the modeled occurrence date of the AMSIA was ~~six~~ days later than the observed date (February 7) in 2008.



430 | **Figure 58:** Comparison of daily sea ice area (DSIA) between NEMO-Bohai simulations (black line) and satellite-derived data (blue circles) from 1996 to 2017.

Table 4: Comparison of the occurrence dates of the annual maximum sea ice area between the observations and model simulations.

Year	The occurrence dates of the annual maximum sea ice area			Year	The occurrence dates of the annual maximum sea ice area		
	Observed	NEMO-Bohai	Error		Observed	NEMO-Bohai	Error
1996	2.10	2.10	0	2007	2.1	2.2	+1
1997	2.3	2.3	0	2008	2.7	2.13	+6
1998	1.28	1.25	-3	2009	1.26	1.26	0
1999	2.3	2.4	+1	2010	2.13	1.26	-18
2000	2.8	2.3	-5	2011	1.30	1.31	+1
2001	2.7	2.7	0	2012	2.2	2.9	-7
2002	1.25	1.24	-1	2013	2.8	2.9	+1
2003	2.2	1.30	-3	2014	2.12	2.11	-1
2004	2.5	2.7	+2	2015	2.4	2.9	+5
2005	2.19	2.21	+2	2016	2.2	2.3	-1
2006	2.8	2.9	+1	2017	2.12	2.11	-1

435

440

The comparison of the spatial distribution of sea ice from the NEMO-Bohai simulation and remote sensing inversion during the freeze, severe freeze, and thaw periods in light ice year (2007), normal ice year (2009), and heavy ice year (2010) is shown in Fig. 69. The simulated spatial distributions reflect general characteristics of sea ice evolution in the Bohai Sea except for the bias at the ice edge zones. Sea ice is mainly located in Liaodong Bay in light and normal ice years with extension to Bohai Bay and Laizhou Bay in heavy ice years. It is worthy to mention that NEMO-Bohai has well reproduced the development cycle of the sea ice but with a relatively slow thawing process.




Figure 69: Comparison of the spatial distribution of sea ice from NEMO-Bohai simulation and remote sensing inversion in freeze, severe freeze, and thaw periods for light (a-f), normal (g-l), and heavy (m-r) ice years.

3.3.2 Sea ice thickness and volume

445

450

Fig. 710 shows that the modeled sea ice thickness based on NEMO-Bohai mostly lies in the range of in-situ observations with a slight overestimation. The mean relative bias of sea ice thickness between the simulations and observations is 4.6 cm ($n = 37$). Other studies also revealed that sea ice thickness is overestimated by NEMO in regional studies, such as in the Baltic Sea (Tedesco et al., 2017) and the Canadian Arctic Archipelago (Hu et al., 2018). When comparing ice thickness, the discrepancy of spatial scales makes it difficult to directly compare the modeled average of grids (~2 km) to the exact location in-situ measurement as sea ice thickness varies greatly in the Bohai Sea, especially in the floating ice zone.





Figure 710: The comparison of sea ice thickness between the simulated and observation data in the Bohai Sea. The red vertical bars represent the range of the observations, while the blue triangles denote the simulations based on NEMO-Bohai.

Sea ice volume is defined as the total ice over the whole Bohai Sea, which is calculated through sea ice concentration times multiplied by ice thickness in all grids. As demonstrated in Fig. 811, modeled daily sea ice volume is in reasonable agreement with satellite-derived data between 1996 and 2017. The modeled sea ice volume is higher than the observed, with the mean relative bias of $15.1 \times 10^8 \text{ m}^3$ (RMSE= $20.3 \times 10^8 \text{ m}^3$). The correlation coefficient between simulations and satellite-derived data is quite relatively high ($r=0.80$, $p<0.01$, $n=432$), indicating reasonable agreement at daily sea ice volume variations. In 2010, NEMO-Bohai simulations reproduced the range of $0.1 \sim 73.0 \times 10^8 \text{ m}^3$, which agrees with the range of $4.4 \sim 63.0 \times 10^8 \text{ m}^3$ reported by Liu et al. (2013) based on AVHRR satellite images. However, it is noticeable that modeled there is a lag in modelling sea ice volume change is slightly later than the observed by satellite formation during the early freezing period, which is similar compared to the later phase indicated by the aforementioned comparison of sea ice area satellite observations.



470 **Figure 811: Comparison of daily sea ice volume between satellite-derived data (circles) and NEMO-Bohai simulations (grey bars) from 1996 to 2017.**


Results of model comparison with in-situ data and satellite-derived data confirm the robustness of the developed model, which allows us to use it in a more detailed evaluation of the spatial and temporal changes of Bohai Sea ice and study the continuous processes of freezing and melting at daily scales.

4. Initial results and applications

475 4.1 Ocean

4.1.1 Sea surface temperature

480 Fig. 912 shows the monthly seasonal ehangescycle of monthly-average SST in the Bohai Sea. There is an obvious ‘warm tongue’ starting from the Bohai Strait to the central Bohai Sea in winter (December, January, and February) due to the warm current from the Yellow Sea, which. It turns into a ‘cold tongue’ in summer due to faster temperature increases in shallow coastal shallow-waters as also-the influence of the warm current from the Yellow Sea also weakens. During January and February, the SST is below 0 °C in all three bays. There are large areas with SSTs below -1 °C in Liaodong Bay, which is the coldest sea in China during the winter (Ju and Xiong, 2013). This extremely cold water provides an ideal platform for sea ice formation.




485

Figure 912: Spatial distributions of multi-year monthly average sea surface temperature in the Bohai Sea for the period of 1996-2017.

4.1.2 Sea surface salinity

490

As shown in Fig. 1013, except for estuaries, the salinity isolines of the Bohai Sea are roughly parallel to the coast as the salinity in the coastal water is generally low due to river runoff. The average salinity of the whole Bohai Sea is low in summer (lowest in August at $29.5 \text{ g}\cdot\text{kg}^{-1}\text{PSU}$) and high in winter (highest in February at $30.0 \text{ g}\cdot\text{kg}^{-1}\text{PSU}$). The Bohai Strait in the east, ~~that~~which connects the Bohai Sea with the Yellow Sea, exhibits the highest salinity in the entire Bohai Sea, with an average of about $32.8 \text{ g}\cdot\text{kg}^{-1}\text{PSU}$.




495

Figure 1013: Spatial distributions of multi-year monthly average sea surface salinity (PSU) in the Bohai Sea during 1996-2017.

4.2 Sea ice

4.2.1 Temporal variation of Bohai sea ice

500 The variation of annual average and maximum sea ice area in the Bohai Sea based on NEMO-Bohai from 1996 to 2017 are shown in Fig. 14a and Fig. 14b, respectively. Annual average sea ice area (AASIA) is averaged by the DSIA during the ice period for each year. The sea ice area exhibits obvious interannual and decadal variability in the study period. The mean AASIA during the period 1996-2017 is $1.07 \times 10^4 \text{ km}^2$, accounting for 14.3% of the total sea area. The 2010/11 and 2011/12 winters exhibited high AASIA values of $1.71 \times 10^4 \text{ km}^2$ and $1.73 \times 10^4 \text{ km}^2$, which cover 22.8% and 23.1%, respectively. The minimum AASIA reaches $0.54 \times 10^4 \text{ km}^2$ (7.2% coverage) in 2007 winter, only accounting for half of the multi-year average AASIA. Analogously to AASIA, AMSIA is the maximum of the DSIA during the period for each year. The mean AMSIA during 1996-2017 is $2.07 \times 10^4 \text{ km}^2$ (27.6% coverage), and the AMSIA values in 2001/02, 2010/11, 2011/12 and 2016/17 winters are particularly high, larger than $3.00 \times 10^4 \text{ km}^2$ (40% coverage). No significant trend in AASIA was observed, while AMSIA showed a non-significant increasing trend with $0.1 \times 10^4 \text{ km}^2 \text{ decade}^{-1}$ ($r=0.08$, $p=0.72$) from 1996 to 2017. In addition, a strong positive correlation between AASIA and AMSIA ($r=0.94$, $p<0.01$) was found.




515 **Figure 14:** The annual average sea ice area (a), the annual maximum sea ice area (b), the simulated length of ice period (c), and the annual average sea ice volume (d) in the Bohai Sea from 1996 to 2017. The blue dots

present the annual average values, and the error bars stand for one standard deviation, and the dotted lines represent 22 a-trends obtained through the linear fitting.

520 The length of the ice period is defined as days with a sea ice area greater than 100 km² from December to March. It varies from 60 to 110 days during the period 1996-2017, as shown in Fig. 14c. The mean of the length of the ice period is 87 days, with the standard deviation of 12 days, and there is a slight increasing but statistically insignificant trend ($r=0.12$, $p=0.59$). The annual average sea ice volume (AASIV, defined as the average of daily sea ice volume in the ice period for each year) from 1996 to 2017 is 2.24 billion m³. As shown in Fig. 14d, no significant trends can be found for AASIV 525 during the study period. The highest value of AASIV appeared in 2010, which was 3.76 billion m³, and the lowest value appeared in 2007 with 0.7 billion m³, accounting for only 18.6% of the ice volume of the highest year. The strong interannual variability of sea ice volume is determined by the strong interannual variability of sea ice area ($r=0.96$, $p<0.01$). Interestingly, it can be noticed that the ice period is not highly correlated with sea ice area or thickness, implying complicated processes of sea ice forming, 530 growth, and thaw in the Bohai Sea.

The climatological seasonal cycles of the sea ice area and volume in the Bohai Sea averaged over 1996-2017 both show unimodal variations (Fig. 1215). Ice usually starts to form in mid-December and reaches the maximum in early February. Then it starts to melt until the Bohai Sea becomes totally ice-free by mid-to-late March. The climatological mean of the length of the ice period is about three and a 535 half months, and the freezing period (~8 weeks) is longer than the thawing period (~7 weeks). The asymmetrical process is related to changing rates of air temperature in freezing and thawing periods (Yan et al., 2020). Similarly, the seasonal variation of sea ice volume is an asymmetrical process. It is worth mentioning that the standard deviations of sea ice area and volume during the freezing period are quite pretty large, showing that the Bohai Sea ice has a large inter-annual variability.



540

Figure 4.2.15: Seasonal cycles of daily sea ice area (a) and sea ice volume (b) in the Bohai Sea. The lines show the daily averages, and the shading denotes one standard deviation. The seasonal cycles are calculated for the periods 1996–2017.

4.2.2 Spatial variation of Bohai sea ice

545

This section will mainly discuss the spatial features of Bohai Sea ice. As shown in Fig. 4.3.16, there are strong seasonal and spatial differences in sea ice concentrations. To be specific, seawater freezes first in Liaodong Bay, next in Bohai Bay, followed by Laizhou Bay, showing significant variation with latitude. On the contrary, the melting process happens exactly at the reverse order. Similar to the aforementioned temporal changes in the last section, the sea ice concentration reaches the peak in February. Seawater in Liaodong Bay freezes most severely, and the high concentration zone moves from the north in January to the east in February, where it survives the longest until March.

550




Figure 1316: Climatological monthly mean sea ice concentration in the Bohai Sea during 1996–2017, which is calculated as the average of all daily sea ice concentrations for each month during the ice period.

555 As shown from Fig. 1417, sea ice thickness shows quite similar seasonal and spatial features with sea ice concentration. The monthly mean Bohai Sea ice thickness simulated by NEMO-Bohai usually reaches its maximum in February, with a monthly mean thickness of 16.9 cm, following the second-- highest monthly mean thickness of 15.8 cm in January. Sea ice is thicker in on the east coast of Liaodong Bay than that in on the west coast in January and February. The maximum sea ice thickness appears near 560 the Bayuquan on the east coast of Liaodong Bay, where the thickness reaches up to ~60.0 cm in February.




Figure 1417: Similar to figure 1316 but for sea ice thickness.

According to the guideline for risk assessment and zoning of sea ice disaster issued by the State Oceanic Administration in 2016, a high-risk level is reached when sea ice thickness becomes greater than 25 cm, while the risk level is low when sea ice thickness is lower than 10 cm. When sea ice thickness is between 10 cm and 25 cm, the risk level is moderate. Accordingly, in this study, the thresholds for moderate-risk and high-risk level levels of sea ice disaster are set at 10 cm and 25 cm, respectively. In Fig. 1518, sea ice risk maps are calculated based on daily sea ice thickness from 1996 to 2017, and they clearly show that the high-risk area is mainly located at Liaodong Bay, with the highest risk area in the eastern Liaodong Bay. Thus, the seas around Yingkou, Bayuquan, and Wafangdian bear high exposure to sea ice disasters.




Figure 1518: Frequency distribution of daily sea ice thickness larger than 10 cm (a) and 25 cm (b) in the Bohai Sea during 1996–2017.

575 As the drift ice is a major component in Bohai Sea ice, studying its motion characteristics is also quite important. As shown in Fig. 1619, the direction is mainly northeast-southwesterly, and the drift exhibits a high spatial variability. The zone with the highest sea ice velocity ($\sim 0.15 \text{ m s}^{-1}$ and mainly directed to southwest) is visible at the southeast edge of Liaodong Bay, and is next to the high concentration and thickness area. In February, the velocity in the northern Liaodong Bay is significantly lower than that in the southern part near the ice edge.

580





Figure 1619: Similar to figure 1316 but for sea ice velocity.

4.2.3 Correlations between sea ice and climate factors

585 The synoptic forcing may play an essential role in the changes of Bohai Sea ice which is primarily
 influenced by the Eurasian continental climate. To explore the potential regional climate drivers on the
 evolution of sea ice, correlations between sea ice and various indices based on detrended air temperature,
 air pressure, wind speed, and precipitation were examined. These indices were calculated based on daily
 meteorological measurements during the winter (DJFM) from 1996 to 2017 at 12 weather stations
 (Dalian, Wafangdian, Xiongyue, Yingkou, Jinzhou, Xingcheng, Suizhong, Qinhuangdao, Laoting,
 590 Tanggu, Huanghua, Dongying) surrounding the Bohai Sea. These measurements were obtained from the
 China Meteorological Data Service Center (<http://data.cma.cn/en>). As shown in Fig. 20a, a strong
 negative correlation ($r = -0.78$, $p < 0.01$) between the detrended time series of AASIA and the mean
 winter air temperature (MWAT) is observed. Similarly, the detrended AASIV exhibits even a stronger
 negative correlation ($r = -0.87$, $p < 0.01$, Fig. 20e) with MWAT, suggesting that MWAT is an essential
 regional climate factor for sea ice changes. The detrended Bohai Sea ice area exhibits a weak but
 insignificant positive correlation ($r = 0.28$, $p = 0.21$, Fig. 20b) with the mean winter relative humidity
 (MWRH). The correlation between the detrended AASIV and the detrended MWRH is also found to be
 weak and insignificant ($r = 0.36$, $p = 0.10$, Fig. 20f). In addition, the correlations between the detrended
 AASIA and the detrended time series of mean winter wind speed (MWWS, $r = 0.13$, $p = 0.56$, Fig. 20c)
 and mean winter air pressure (MWAP, $r = 0.04$, $p = 0.85$, Fig. 20d) were found to be very weak and
 insignificant. Similar very weak and insignificant correlations were also found between AASIV and
 600 MWWS/MWAP (see Fig. 20g, 20h). The above correlation analysis indicates that the local temperature
 is the controlling factor for the sea ice evolution in the Bohai Sea.




605

Figure 20: Pearson correlations between annual average sea ice area (AASIA) (a-d), annual average sea ice volume (AASIV) (e-h) and mean winter air temperature (MWAT), mean winter relative humidity (MWRH), mean winter wind speed (MWWS), mean winter air pressure (MWAP) through detrended analysis for the period 1996–2017. The climate parameters are calculated based on the measured daily meteorological data at 12 meteorological stations surrounding the Bohai Sea during the winter (DJFM).

610

Fig. 21 illustrates the spatial correlations between daily sea ice concentration with vertically integrated ocean heat/salt content from the surface to the bottom of the mixed layer during the ice period for each grid point. Ocean heat content strongly correlates with sea ice concentration in Liaodong Bay and moderately correlates near the coastal areas. The spatial pattern corresponds quite nicely with climatological monthly mean sea ice concentration (shown in Fig. 16), with the highest values appearing at the east of Liaodong Bay. On the other hand, a very weak positive or negative correlation between daily sea ice concentration and integrated ocean salt content was found in most sea areas, indicating a more complicated relationship throughout the whole Bohai Sea. Therefore, the interannual variability of sea ice is more dominated by ocean heat content than salt.

615



620

Figure 21: Spatial correlation between daily sea ice concentration in the Bohai Sea and (a) vertically integrated ocean heat content, and (b) vertically integrated ocean salt content from the surface to the bottom of the mixed layer during the ice period (DJFM).

5. Conclusion and perspectives for the modelling platform

625 In this study, we provided a detailed description of NEMO-Bohai, a newly developed ~~set-up~~ setup of an ocean-ice model for the Bohai Sea. ~~Primary~~The primary intent of our study was to test how NEMO-Bohai represents the ocean characteristics ~~mainly in terms of~~and sea ice properties.

Comparisons with observational data confirm that NEMO-Bohai is able to reproduce reasonably well the ocean ~~properties, including ocean~~ surface information, ~~including (e.g.,~~ SSH, SST, ~~SSS), currents,~~ and ~~SSS-temperature and salinity stratification~~. However, the ongoing development of ~~the~~ NEMO ocean engine, soon providing an updated version with the inclusion of wetting and drying processes, could improve the model performance in terms of sea surface height in shallow areas (Hordoir et al., 2019), especially for the Bohai Sea with ~~an~~ average depth of only 18 m. In addition, a higher-precision bathymetry could ~~possibly~~ improve the SSH performance. In this study, the monthly climatology of coastal runoff flux from Dai et al. (2009) ~~was~~ recorded only for 1948–2004, ~~which is~~ different from the study period. ~~Moreover~~More importantly, the river runoff is based on climatological estimates without ~~interannual variability~~. In addition, we assume that the river salinity is set to 0 ~~g~~-kg⁻¹PSU and the runoff temperature is ~~the~~ same as the ocean surface. ~~A major~~Major future development would be ~~the use of~~using gauge records with observations of ~~salinity and~~ temperature ~~and salinity~~ assimilated into the NEMO-Bohai model, which would reduce uncertainties in temperature and salinity. Furthermore, a possible implementation of multiple embedded ~~method~~methods (Hvatov et al., 2019; Schwarzkopf et al., 2019), such as Global Ocean-West Pacific Ocean-East China Sea-Bohai Sea rather than current Global Ocean-Bohai Sea direct nesting, should be investigated in the future. Moreover, interactive feedback from ~~the~~ Bohai Sea to ~~the~~ global ~~Ocean~~ocean could also be considered in a two-way coupling method compared to current one-way nesting.

645 For the sea ice component, NEMO-Bohai reproduced satisfactorily the seasonal and interannual variabilities of ~~sea~~ ice area compared to the satellite remote sensing data for the period 1996-2017. The modeled dates of ~~the~~ annual maximum sea ice area were ~~on average~~ 0.9 days earlier than the observed ones. Spatially, the simulation results realistically reflect the main characteristics of Bohai Sea ice evolution ~~as~~ compared to the satellite data. Therefore, NEMO-Bohai can reliably be used to provide ice information during the dates without satellite-derived data. Nevertheless, it is worth mentioning that the simulated sea ice area tends to be ~~a~~ somewhat overestimated, which is also reported for other seas in

earlier NEMO-related publications (Blockley et al., 2014; Massonnet et al., 2011; Rjazin et al., 2019). ~~In~~
~~spring, the~~ NEMO-Bohai melting process is delayed, particularly in the land-fast ice zone in the Eastern
655 Liaodong Bay. It is likely that ~~too~~-thick ice melts slower away than thinner observed ice. The applied
~~parametrization~~parameterization of the surface albedo may cause a slow melting process, which ~~is~~does
not ~~taking~~take realistically into account the relevant physical processes, such as surface melt ponds, as
the surface albedo continues to decrease until sea ice is completely disappeared (Mortin et al., ~~2016~~). ~~It~~
~~is also possible related to the model unrealism of solar radiation penetration into ice, which strongly~~
660 ~~affects the surface energy balance (Light et al., 2008), thus~~2016). Thus, regional atmospheric forcing
data with higher accuracy and resolution can be used for further development. In addition, the space
discrepancy between modeled sea ice thickness and extremely limited in-situ observations makes their
comparison difficult and introduces significant uncertainties.

In conclusion, NEMO-Bohai ~~is able to~~can simulate ocean and sea ice properties with
665 ~~good~~reasonable skill in a broad spatiotemporal context, especially in terms of seasonal evolution and
long-term interannual variations of sea ice. This finding implies that NEMO-Bohai well complements
the discontinuous satellite data in sea ice hazard risk analysis. Therefore, NEMO-Bohai is a
usefulvaluable tool, ~~not only~~ for long-term ocean and ice simulations, ~~but also for the ocean~~ and climate
change studies.

670 **Code and data availability**

The NEMO-Bohai is built upon the standard NEMO code (~~release~~NEMO 4.0.6 beta, revision 10226).
The reference code is available at the NEMO website (<http://www.nemo-ocean.eu/>) through the
following link: <http://forge.ipsl.jussieu.fr/nemo/svn/NEMO/trunk> (last access: ~~20~~ May 20, 2021). The
parameters and configuration files running the NEMO-Bohai, such as namelists, bathymetry, boundary
675 coordinates, and definition files, forcing data etc., together with the scripts to run the model, are
available at <https://doi.org/10.5281/zenodo.4892454>. The forcing data for the year 2010 are included in
the Zenodo archive.

Author contributions

YY, WG, and PU designed the study. YY and AG ran the experiments. YY and YX analyzed the model

680 and observational data. YY and PU wrote the manuscript. All authors provided scientific input.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

This study was financially supported by the National Natural Science Foundation of China (Nos. 41977406, 41571510), the Academy of Finland (grant 322432), the China Scholarship Council (No. 201806040130), [111 project \(No. B20011\)](#), and the National Key Research and Development Program of China (No. 2017YFA0604903). This work was granted access to the HPC resources of CSC - IT Center for Science. We gratefully thank Juha Lento, Robinson Hordoier, Yongmei Gong ~~and~~, Yu Zhang, [Xiaoqiao Wang](#) for sharing the scripts for input data processing.

690 References

- Aksenov, Y., Blockley, E., Chevallier, M., Feltham, D., Fichefet, T., Garric, G., Holland, P., Iovino, D., Madec, G., Massonnet, F., Ridley, J., Rousset, C., Salas, D., Schroeder, D., Tietsche, S., and Vancoppenolle, M.: Sea Ice modelling Integrated Initiative (SI³) – The NEMO sea ice engine, Scientific Notes of Climate Modelling Center, ISSN 1288-1619, Institut Pierre-Simon Laplace (IPSL), doi:10.5281/zenodo.1471689, 2019.
- 695 Bai, S. and Wu, H.: Numerical sea ice forecast for the Bohai Sea, Acta Meteorologica Sinica (in Chinese), 56, 139-153, 1998.
- Bai, X., Wang, J., Liu, Q., Wang, D., and Liu, Y.: Severe ice conditions in the Bohai Sea, China, and mild ice conditions in the great lakes during the 2009/10 winter: Links to El Nino and a strong negative arctic oscillation, J Appl Meteorol Clim, 50, 1922-1935, 2011.
- 700 [Bernard, B., Madec, G., Penduff, T., Molines, J. M., Treguier, A. M., Le Sommer, J., Beckmann, A., Biastoch, A., Böning, C., Dengg, J., Derval, C., Durand, E., Gulev, S., Remy, E., Talandier, C., Theetten, S., Maltrud, M., McClean, J., and De Cuevas, B.: Impact of partial steps and momentum advection schemes in a global ocean circulation model at eddy-permitting resolution, Ocean Dynam, 56\(5\), 543-567, 2006.](#)
- 705 [Bian, C., Jiang, W., Pohlmann, T., and Sündermann, J.: Hydrography-physical description of the Bohai Sea, J Coast Res, 74, 1-12, 2016.](#)
- Blanke, B. and Delecluse, P.: Variability of the tropical Atlantic Ocean simulated by a general circulation model with two different mixed-layer physics, [Journal of Physical Oceanography J Phys Oceanogr](#), 23, 1363-1388, 1993.
- 710 Blockley, E. W., Martin, M. J., McLaren, A. J., Ryan, A. G., Waters, J., Lea, D. J., Mirouze, I., Peterson, K. A., Sellar, A., and Storkey, D.: Recent development of the Met Office operational ocean

- forecasting system: an overview and assessment of the new Global FOAM forecasts, [Geoscientific Model Development](#), 7, 2613-2638, 2014.
- 715 Bouillon, S., Fichet, T., Legat, V., and Madec, G.: The elastic–viscous–plastic method revisited, [Ocean Modelling](#), 71, 2-12, 2013.
- [Chen, D. X.: Marine atlas of Bohai Sea, Yellow Sea and East China Sea: Hydrology, Ocean Press, Beijing, China, 1992.](#)
- 720 Courtois, P., Hu, X., Pennelly, C., Spence, P., and Myers, P. G.: Mixed layer depth calculation in deep convection regions in ocean numerical models, [Ocean Modelling](#), 120, 60-78, 2017.
- Dai, A., Qian, T., Trenberth, K., and Milliman, J.: Changes in continental freshwater discharge from 1948 to 2004, [J Climate](#), 22, 2773-2792, 2009.
- 725 Declerck, A., Ourmières, Y., and Molcard, A.: Assessment of the coastal dynamics in a nested zoom and feedback on the boundary current: the North-Western Mediterranean Sea case, [Ocean Dynamics](#), 66, 1529-1542, 2016.
- Depoorter, M. A., Bamber, J. L., Griggs, J. A., Lenaerts, J. T., Ligtner, S. R., van den Broeke, M. R., and Moholdt, G.: Calving fluxes and basal melt rates of Antarctic ice shelves, [Nature](#), 502, 89-92, 2013.
- [Ding, D.W.: Introduction to sea ice engineering \(in Chinese\), Ocean Press, Beijing, 1999.](#)
- 730 [Donlon, C. J., Martin, M., Stark, J., Roberts-Jones, J., Fiedler, E., and Wimmer, W.: The operational sea surface temperature and sea ice analysis \(OSTIA\) system, \[Remote Sens Environ.\]\(#\), 116: 140-158, 2012.](#)
- Drouard, M. and Cassou, C.: A modeling-and process-oriented study to investigate the projected change of ENSO-forced wintertime teleconnectivity in a warmer world, [J Climate](#), 32, 8047-8068, 2019.
- 735 Egbert, G. D. and Erofeeva, S. Y.: Efficient inverse modeling of barotropic ocean tides, [Journal of Atmospheric and Oceanic Technology](#), 19, 183-204, 2002.
- Engedahl, H.: Use of the flow relaxation scheme in a three-dimensional baroclinic ocean model with realistic topography, [Tellus A](#), 47, 365-382, 1995.
- 740 Feucher, C., Garcia-Quintana, Y., Yashayaev, I., Hu, X., and Myers, P. G.: Labrador sea water formation rate and its impact on the local meridional overturning circulation, [Journal of Geophysical Research: J Geophys Res-Oceans](#), 124, 5654-5670, 2019.
- Flather, R. A.: A tidal model of the north-west European continental shelf, [Memoires Societe Royale des Sciences de Liege](#), 10, 141-164, 1976.
- 745 Fu, M., Liu, H., Jin, X., and He, K.: National-to port-level inventories of shipping emissions in China, [Environ Res Lett](#), 12, 114024, 2017.
- Gong, D., Kim, S., and Ho, C.: Arctic oscillation and ice severity in the Bohai Sea, East Asia, [Int J Climatol](#), 27, 1287-1302, 2007.
- Graham, J. A., O'Dea, E., Holt, J., Polton, J., Hewitt, H. T., Furner, R., Guihou, K., Brereton, A., Arnold, A., and Wakelin, S.: AMM15: a new high-resolution NEMO configuration for operational simulation of the European north-west shelf, [Geoscientific Model Development](#), 11, 681-696, 2018.
- 750 Gu, W., Liu, C., Yuan, S., Li, N., Chao, J., Li, L., and Xu, Y.: Spatial distribution characteristics of sea-ice-hazard risk in Bohai, China, [Ann Glaciol](#), 54, 73-79, 2013.
- 755 Gu, W., Shi, P., Chen, W., Xie, F., Xu, Y., and Yuan, S.: Sea ice reserves estimates and quality evaluation, Science Press, Beijing, 2014.
- Gunduz, M., Özsoy, E., and Hordoir, R.: A model of Black Sea circulation with strait exchange (2008–

- 2018), [Geoscientific Model Development](#), 13, 121-138, 2020.
- 760 Harada, Y., Kamahori, H., Kobayashi, C., Endo, H., Kobayashi, S., Ota, Y., Onoda, H., Onogi, K., Miyaoka, K., and Takahashi, K.: The JRA-55 reanalysis: Representation of atmospheric circulation and climate variability, [Journal of the Meteorological Society of Japan. Ser. III Meteorol Soc Jpn](#), 94, 269-302, 2016.
- 765 Hordoir, R., Axell, L., Höglund, A., Dieterich, C., Fransner, F., Gröger, M., Liu, Y., Pemberton, P., Schimanke, S., Andersson, H., Ljungemyr, P., Nygren, P., Falahat, S., Nord, A., Jönsson, A., Lake, I., Döös, K., Hieronymus, M., Dietze, H., Löptien, U., Kuznetsov, I., Westerlund, A., Tuomi, L., and Haapala, J.: Nemo-Nordic 1.0: a NEMO-based ocean model for the Baltic and North seas – research and operational applications, [Geosci. Model Dev.](#), 12, 363-386, 2019.
- 770 Hvatov, A., Nikitin, N. O., Kalyuzhnaya, A. V., and Kosukhin, S. S.: Adaptation of NEMO-LIM3 model for multigrid high resolution Arctic simulation, [Ocean Modelling](#), 141, 101427, 2019.
- [Ji, C. Z., Li, K., Yu, B., Dong, L., and Liu, Q. R.: The multi-time scale variations of water exchange across the Bohai Strait, \[Oceanol Limnol Sin \\(in Chinese\\)\]\(#\), 50, 24-30, 2019.](#)
- Ju, X. and Xiong, X.: Distributions and seasonal changes of water temperature in the Bohai Sea, Yellow Sea and East China Sea, [Advances in Marine Science \(in Chinese\)](#), 31, 55-68, 2013.
- 775 [Kärnä, T., Ljungemyr, P., Falahat, S., Ringgaard, I., Axell, L., Korabel, V., Murawski, J., Maljutenko, I., Lindenthal, A., Jandt-Scheelke, S., Verjovkina, S., Lorkowski, I., Lagema, P., She, J., Tuomi, L., Nord, A., and Huess, V.: Nemo-Nordic 2.0: operational marine forecast model for the Baltic Sea, \[Geosci Model Dev\]\(#\), 14\(9\), 5731-5749, 2021.](#)
- Karvonen, J., Shi, L., Cheng, B., Simila, M., Makynen, M., and Vihma, T.: Bohai sea ice parameter estimation based on thermodynamic ice model and earth observation data, [Remote Sens-Basel](#), 9, 234, 2017.
- 780 Kimritz, M., Losch, M., and Danilov, S.: A comparison of viscous-plastic sea ice solvers with and without replacement pressure, [Ocean Modelling](#), 115, 59-69, 2017.
- Kobayashi, S., Ota, Y., Harada, Y., Ebata, A., Moriya, M., Onoda, H., Onogi, K., Kamahori, H., Kobayashi, C., Endo, H., Miyaoka, K., and Takahashi, K.: The JRA-55 reanalysis: General specifications and basic characteristics, [Journal of the Meteorological Society of Japan. Ser. III Meteorol Soc Jpn](#), 93, 5-48, 2015.
- 785 Large, W. G. and Yeager, S. G.: The global climatology of an interannually varying air-sea flux data set, [Climate Dynamics](#), 33, 341-364, 2009.
- Leppäranta, M.: The drift of sea ice, Springer Science & Business Media, 2011.
- 790 [Li, R., Lu, Y., Hu, X., Guo, D., Zhao, P., Wang, N., Lee, K., and Zhang, B.: Space-time variations of sea ice in Bohai Sea in the winter of 2009–2010 simulated with a coupled ocean and ice model, \[J Oceanogr\]\(#\), 77\(2\), 243-258, 2021.](#)
- [Li, Y., Wolanski, E., and Zhang, H.: What processes control the net currents through shallow straits? A review with application to the Bohai Strait, China, \[Estuar Coast Shelf S\]\(#\), 158, 1-11, 2015.](#)
- 795 [Li, Z., Lu, P., and Sodhi, D. S.: Ice engineering sub-areas in Bohai from ice physical and mechanical parameters, \[Advances in Water Science \\(in Chinese\\)\]\(#\), 15, 598-602, 2004.](#)
- Li, Z., Zhang, L., Lu, P., Leppäranta, M., and Li, G.: Experimental study on the effect of porosity on the uniaxial compressive strength of sea ice in Bohai Sea, [ScienceSci China Technol Sci](#), 54, 2429-2436, 2011.
- 800 [Light, B., Grenfell, T., Lin, X., Wu, D., Bao, X., and Perovich, D. K.: Transmission and absorption flux variation of solar radiation by Arctic sea ice during the](#)

- ~~melt-season~~Bohai Strait, *Journal of Geophysical Research: Ocean University of Qingdao (in Chinese)*, 32(3), 355-360, 2002.
- Lin, X., Yang, J., Guo, J., Zhang, Z., Yin, Y., Song, X., and Zhang, X.: An asymmetric upwind flow, *Yellow Sea warm current: 1. New observations in the western Yellow Sea*, *J Geophys Res-Oceans*, 113, C03023, 2008116(C4), C04026, 2011.
- 805 Liu, C., Gu, W., Chao, J., Li, L., Yuan, S., and Xu, Y.: Spatio-temporal characteristics of the sea-ice volume of the Bohai Sea, China, in winter 2009/10, *Ann Glaciol*, 54, 97-104, 2013.
- Liu, Q., Liu, Y., Bai, S., Zhang, Q., Li, B., and Li, C.: Numerical sea ice forecast for the Bohai Sea in the winter of 2002-2003, *Marine Forecasts (in Chinese)*, 20, 60-67, 2003.
- 810 Locarnini, R. A., Mishonov, A. V., Antonov, J. I., Boyer, T. P., Garcia, H. E., Baranova, O. K., Zweng, M. M., Paver, C. R., Reagan, J. R., Johnson, D. R., Hamilton, M., and Seidov, D.: World ocean atlas 2013, volume 1: Temperature, NOAA Atlas NESDIS 73, 40 pp., 2013.
- Madec, G., and the NEMO team: NEMO ocean engine. Note du Pôle de modélisation, Institut Pierre-Simon Laplace No 27, France, ISSN No No 1288-1619, 2016.
- 815 Massonnet, F., Barthélemy, A., Worou, K., Fichet, T., Vancoppenolle, M., Rousset, C., and Moreno-Chamarro, E.: On the discretization of the ice thickness distribution in the NEMO3.6-LIM3 global ocean-sea ice model, *Geoscientific Model Development Dev*, 12, 3745-3758, 2019.
- Massonnet, F., Fichet, T., Goosse, H., Vancoppenolle, M., Mathiot, P., and König Beatty, C.: On the influence of model physics on simulations of Arctic and Antarctic sea ice, *The Cryosphere*, 5, 687-699, 2011.
- 820 Mortin, J., Svensson, G., Graversen, R. G., Kapsch, M.-L., Stroeve, J. C., and Boisvert, L. N.: Melt onset over Arctic sea ice controlled by atmospheric moisture transport, *Geophysical Research Letters*, 43, 6636-6642, 2016.
- O'Dea, E., Furner, R., Wakelin, S., Siddorn, J., While, J., Sykes, P., King, R., Holt, J., and Hewitt, H.: The CO5 configuration of the 7 km Atlantic Margin Model: large-scale biases and sensitivity to forcing, physics options and vertical resolution, *Geoscientific Model Development Dev*, 10, 2947, 2017.
- 825 Obermann-Hellhund, A., Conte, D., Somot, S., Torma, C. Z., and Ahrens, B.: Mistral and Tramontane wind systems in climate simulations from 1950 to 2100, *Climate dynamics*, 50, 693-703, 2018.
- 830 Pemberton, P., Löptien, U., Hordoir, R., Höglund, A., Schimanke, S., Axell, L., and Haapala, J.: Sea-ice evaluation of NEMO-Nordic 1.0: a NEMO-LIM3.6-based ocean-sea-ice model setup for the North Sea and Baltic Sea, *Geoscientific Model Development Dev*, 10, 3105-3123, 2017.
- Rjazin, J., Hordoir, R., and Parn, O.: Evaluation of the NEMO-Nordic model by comparing the sea-ice concentration values in the Baltic Sea, *Journal of Ocean Technology*, 14, 183, 2019.
- 835 Schwarzkopf, F. U., Biastoch, A., Böning, C. W., Chanut, J., Durgadoo, J. V., Getzlaff, K., Harlaß, J., Rieck, J. K., Roth, C., Scheinert, M. M., and Schubert, R.: The INALT family – a set of high-resolution nests for the Agulhas Current system within global NEMO ocean/sea-ice configurations, *Geoscientific Model Development Dev*, 12, 3329-3355, 2019.
- 840 Shi, W. and Wang, M.: Sea ice properties in the Bohai Sea measured by MODIS-Aqua: 2. Study of sea ice seasonal and interannual variability, *Journal of Marine Systems*, 95, 41-49, 2012.
- Su, H., Ji, B., and Wang, Y.: Sea ice extent detection in the Bohai Sea using Sentinel-3 OLCI data, *Remote Sens-Basel*, 11, 2436, 2019.
- Su, H. and Wang, Y.: Using MODIS data to estimate sea ice thickness in the Bohai Sea (China) in the

- 2009-2010 winter, *J Geophys Res-Oceans*, 117, C10018, 2012.
- 845 Su, J., Wu, H., Zhang, Y., Liu, Q., and Bai, S.: A coupled ice-ocean model for the Bohai Sea: I. Study on model and parameter, *Acta Oceanol Sin*, 23, 597-608, 2004.
- Tang, M., Liu, Q., Liu, Y., and Bai, S.: The numerical simulation of sea ice in the Bohai Sea, *Marine Forecasts (in Chinese)*, 27, 48-52, 2010.
- 850 Tedesco, L., Miettunen, E., An, B. W., Haapala, J., Kaartokallio, H., and Miller, L. A.: Long-term mesoscale variability of modelled sea-ice primary production in the northern Baltic Sea, *Elementa: Science of the Anthropocene-Sci Anthropol*, 5, 29, 2017.
- Thompson, B., Sanchez, C., Sun, X., Song, G., Liu, J., Huang, X., and Tkalic, P.: A high-resolution atmosphere-ocean coupled model for the western Maritime Continent: development and preliminary assessment, *Climate DynamicsClim Dynam*, 52, 3951-3981, 2019.
- 855 Uotila, P., Iovino, D., Vancoppenolle, M., Lensu, M., and Rousset, C.: Comparing sea ice, hydrography and circulation between NEMO3.6 LIM3 and LIM2, *Geosci- Model Dev*, 10, 1009-1031, 2017.
- Voldoire, A., Sanchez-Gomez, E., y Méliá, D. S., Decharme, B., Cassou, C., Sénési, S., Valcke, S., Beau, I., Alias, A., and Chevallier, M.: The CNRM-CM5. 1 global climate model: description and basic evaluation, *Climate dynamicsClim Dynam*, 40, 2091-2121, 2013.
- 860 [Wan, K., Bao, X., Wang, Y., Wan, X., Li, H., and Liu, K.: Barotropic current fluctuations coupled with sea level drawdown in Yellow and Bohai Seas, *Chin J Oceanol Limn*, 33\(1\), 272-281, 2015.](#)
- [Wang, Q., Guo, X., and Takeoka, H.: Seasonal variations of the Yellow River plume in the Bohai Sea: A model study, *J Geophys Res*, 113, C08046, 2008.](#)
- 865 Wang, R., Liu, X., and Zhang, L.: Numerical tests of sea ice, *Acta Oceanologica Sinica (in Chinese)*, 6, 572-582, 1984.
- Westerlund, A.: A Gulf of Finland (GoF) configuration for the NEMO ocean model, available at: <https://github.com/fmidev/nemo-gof/>, last access: 15 December 2019. ~~2019.~~
- Wu, H.: Mathematical representations of sea ice dynamic-thermodynamic processes, *Oceanologia et*
- 870 *Limnologia Sinica (in Chinese)*, 22, 221-228, 1991.
- Wu, H., Bai, S., Zhang, Z., and Li, G.: Numerical simulation for dynamical processes of sea ice, *Acta Oceanol Sin*, 16, 303-325, 1997.
- Yan, H., Wang, N., Wu, N., and Lin, W.: Abundance, habitat conditions, and conservation of the largha seal (*Phoca largha*) during the past half century in the Bohai Sea, China, *MammalMamm Study*, 43, 1-10, 2018.
- 875 Yan, Y., Gu, W., Xu, Y., and Li, Q.: The in situ observation of modelled sea ice drift characteristics in the Bohai Sea, *Acta Oceanol Sin*, 38, 17-25, 2019.
- Yan, Y., Shao, D., Gu, W., Liu, C., Li, Q., Chao, J., Tao, J., and Xu, Y.: Multidecadal anomalies of Bohai Sea ice cover and potential climate driving factors during 1988-2015, *Environ Res Lett*, 12, 094014, 2017.
- 880 Yan, Y., Uotila, P., Huang, K., and Gu, W.: Variability of sea ice area in the Bohai Sea from 1958 to 2015, *Science of TheSci Total EnvironmentEnviron*, 709, 136164, 2020.
- Yang, L., Xu, X., Zhang, P., Han, J., Li, B., and Berggren, P.: Classification of underwater vocalizations of wild spotted seals (*Phoca largha*) in Liaodong Bay, China, *The Journal of the Acoustical Society of AmericaJ Acoust Soc Am*, 141, 2256-2262, 2017.
- 885 Yuan, B., Huang, R., Jiao, Y., Guo, D., Shang, J., and Zhang, H.: Analysis of sea surface temperature and salinity based on the observed buoys data in the Bohai Sea, *Marine Forecasts (in Chinese)*, 32, 44-50, 2015.

-
- 890 Yuan, S., Gu, W., Xu, Y., Wang, P., Huang, S., Le, Z., and Cong, J.: The estimate of sea ice resources quantity in the Bohai Sea based on NOAA/AVHRR data, *Acta Oceanol Sin*, 31, 33-40, 2012.
- Zalesak, S. T.: Fully multidimensional flux-corrected transport algorithms for fluids, ~~*Journal of Computational Physics*~~*J Comput Phys*, 31, 335-362, 1979.
- Zeng, T., Shi, L., Marko, M., Cheng, B., Zou, J., and Zhang, Z.: Sea ice thickness analyses for the Bohai Sea using MODIS thermal infrared imagery, *Acta Oceanol Sin*, 35, 96-104, 2016.
- 895 Zhang, ~~Q. and Zhang, N.N.~~, Wu, Y., and Zhang, Q.: Forecasting the evolution of the sea ice in the Liaodong Bay using meteorological data, *Cold Reg Sci Technol*, 125, 21-30, 2016.
- Zhang, Q. and Zhang, N.: A three-dimensional numerical model of sea ice evolution in the Bohai Sea, *Journal of Tianjin University (Science and Technology)* (in Chinese), 46, 333-341, 2013.
- 900 Zhang, X., Zhang, Z., Xu, Z., Li, G., Sun, Q., and Hou, X.: Sea ice disasters and their impacts since 2000 in Laizhou Bay of Bohai Sea, China, *Nat Hazards*, 65, 27-40, 2013.
- Zheng, J., Ke, C., and Shao, Z.: Winter sea ice albedo variations in the Bohai Sea of China, *Acta Oceanol Sin*, 36, 56-63, 2017.
- Zweng, M. M., Reagan, J. R., Antonov, J. I., Locarnini, R. A., Mishonov, A. V., Boyer, T. P., Garcia, H. E., Baranova, O. K., Johnson, D. R., and Seidov, D.: *World ocean atlas 2013, volume 2: Salinity*, NOAA Atlas NESDIS 74, 39 pp., 2013.
- 905