

Added value of the EURO-CORDEX high-resolution downscaling over the Iberian Peninsula revisited. Part I: Precipitation

João António Martins Careto¹, Pedro Miguel Matos Soares¹, Rita Margarida Cardoso¹, Sixto Herrera² and Jose Manuel Gutiérrez³

5 ¹Instituto Dom Luiz, Faculdade de Ciências, Universidade de Lisboa, 1749-016, Portugal

²Meteorology Group. Dept. of Applied Mathematics and Computer Science. Universidad de Cantabria. Santander, Spain.

³Meteorology Group. Instituto de Física de Cantabria, CSIC-University of Cantabria, Santander, Spain.

Corresponding author João A. M. Careto (jacareto@fc.ul.pt)

10 *Faculdade de Ciências da Universidade de Lisboa, Campo Grande, Ed. C8 (3.01)*

1749-016 Lisboa, Portugal

15

20

25

30 **Abstract.** Over the years higher resolution Regional Climate Model simulations have emerged owing to the large increase in computational resources. The 12 km resolution from the Coordinated Regional Climate Downscaling Experiment for the European domain (EURO-CORDEX) is a reference, which includes a larger multi-model ensemble at a continental scale while spanning at least a 130-year period. These simulations are computationally demanding but not always revealing added value. In this study, a recently developed regular gridded dataset and a new metric for added value quantification, the Distribution Added Value (DAV), are used to assess the precipitation of all available EURO-CORDEX Hindcast (1989-2008) and Historical (1971-2005) simulations. 35 This approach enables a direct comparison between the higher resolution regional model runs against their forcing global model or ERA-Interim reanalysis, with respect to their probability density functions. This assessment is performed for the Iberian Peninsula. Overall, important gains are found for most cases, particularly in precipitation extremes. Most Hindcast models reveal gains above 15%, namely for wintertime, while for precipitation extremes values above 20% are reached for the summer and autumn. As for the Historical models, although most pairs display gains, regional models forced by 2 GCMs reveal losses, sometimes around -5% or lower, for the entire year. However, the spatialization of the DAV is clear in terms of added value for 40 precipitation, particularly in precipitation extremes with gains well above 100%.

1 Introduction

From the last decades of the 20th century up to today, climate change due to anthropogenic gas emissions has become a major concern for mankind. General Circulation Models (GCMs) are the primary tool used by the IPCC (Intergovernmental Panel on 45 Climate Change) to assess past, present, and future climate conditions. Overall, GCMs can capture the large-scale circulations of the atmosphere and the ocean, together with their centennial to decadal variability and synoptic weather (Meehl et al., 2007; Randall et al., 2007; Stocker et al., 2014). However, the GCMs coarse resolution does not allow for a good representation of orography, land-ocean-atmosphere interactions, or sub-grid processes (Randall et al., 2007; Rummukainen, 2010; Soares et al., 2012). Instead, these local processes rely often on parametrisations resulting in a poor description of processes such as convection and 50 thermal circulations (Prein et al., 2015). Therefore, to bridge the gap between large to local-scale climate, downscaling techniques were developed. These include statistical downscaling (Wilby et al., 1998; Khan et al., 2006) and dynamical downscaling (Giorgi and Bates, 1989; McGregor, 1997; Christensen et al., 2007; Rummukainen, 2010), in which the latter makes use of Regional Climate Models (RCMs). RCMs are run over a geographical domain (continental, national or regional) driven by a GCM, including reanalysis, by means of the boundary conditions. RCMs are an important tool for the representation of regional to local climates, 55 since they are run at much higher resolutions, nowadays between tenths of kilometres to the convection permitting scales, and therefore can capture physically consistent regional to local processes and circulations (Giorgi and Mearns, 1991; 1999; Leung et al., 2003; Laprise, 2008; Heikkilä et al., 2010; Soares et al., 2012a; 2012b; Cardoso et al., 2013; Rios-Entenza et al., 2014; Soares et al., 2014). The gains for an individual variable or process of the higher-resolution RCM simulations against lower-resolutions, given by the GCM or reanalysis driving the RCM, relative to observation, are commonly known as added value (Di Luca et al., 60 2012; 2013; Prein et al., 2013a; 2016; Torma et al., 2015; Rummukainen, 2016; Soares and Cardoso, 2018; Cardoso and Soares, 2021; Careto et al 2021).

In recent years, the increase of computational resources has allowed researchers to run simulations at larger domains and resolutions (Prein et al., 2015; Soares et al., 2017; Jacob et al., 2020; Coppola et al., 2020). These often encompass an entire continent, spanning larger periods from a few decades to over a century. For instance, from the 20th to the end of the 21st century or from the 50 km 65 horizontal resolution from the PRUDENCE project (Christensen and Christensen, 2007) or the 25 km from the ENSEMBLES (van der Linden and Mitchell, 2009), down to the 12 km resolution from the World Research Climate Program Coordinated Regional Downscaling Experiment (WRCP-CORDEX, Jacob et al., 2014; 2020) for the European domain (EURO-CORDEX, hereafter). Moreover, other examples are the ideal case studies, employing simulations at kilometre-scale (Hohenegger et al., 2009; Kendon et al., 2012; 2014; Prein et al., 2013b; Ban et al., 2014; Froidevaux et al., 2014; Fosser et al., 2017; Imamovic et al., 2017; Leutwyler 70 et al., 2017; Liu et al., 2017; Kirshbaum et al., 2018; Fumière et al., 2020; Berthou et al., 2020), or the convection permitting

simulations from the WRCP-CORDEX Flagship Pilot Studies focused over the Alps (Coppola et al., 2020; Ban et al., 2021; Pichelli et al., 2021).

The evaluation and added value of higher resolution simulations constitute an important step to gauge their quality and usefulness. Soares and Cardoso (2018) proposed a new metric to quantify the added value of higher resolutions with respect to their forcing
75 or lower resolution counterpart simulations. This metric is based on the ability of models in representing the observed probability density functions (PDF). It relies on a distribution added value (DAV) which can be applied to either the full PDF or to PDF sections, thus enabling an easy evaluation of extremes or any section of the PDF.

In the past, the hindcast simulations from the EURO-CORDEX were extensively evaluated, revealing gains for the main meteorological variables (Kotlarski et al., 2014; Casanueva et al., 2016a; 2016b; Prein et al., 2016; Soares and Cardoso, 2018;
80 Herrera et al., 2020; Cardoso and Soares, 2021; Careto et al., 2021). Kotlarski et al (2014) assessed temperature and precipitation at monthly and seasonal timescales for the hindcast simulations, reporting slight improvements from EURO-CORDEX relative to the ENSEMBLES (Van der Linden and Mitchell, 2009). Overall, the models showed to be able to capture the space-time variability of the European climate. However, when considering averages over large subdomains and at the seasonal timescale, the higher resolution simulations did not reveal noticeable improvements. Prein et al (2016) also assessed precipitation for both resolutions
85 of the hindcast EURO-CORDEX (50 km and 12 km) and found improvements, mostly in regions characterised by complex terrain and in summertime precipitation, due to the better resolved convective features. More recently, Herrera et al (2020) performed an assessment for precipitation and temperature for an ensemble of 8 Hindcast EURO-CORDEX RCMs over the Iberian Peninsula. The authors report a good spatial agreement between models and observations, namely for temperature. On the other hand, this agreement decreases when extremes are considered. Nevertheless, the authors also report a larger uncertainty related to
90 observations for precipitation relative to temperature.

The first to quantify the added value of the EURO-CORDEX hindcast runs were Soares and Cardoso (2018), evaluating 5 RCMs for precipitation at both resolutions (50 km and 12 km) considering their probability density functions with the station-based dataset ECAD (Klein Tank et al 2002, Klok & Klein Tank 2009) as observational benchmark. This study reported relevant added value of the RCMs against the driving ERA-Interim reanalysis (Dee et al., 2011). Nonetheless, when comparing both resolutions, the
95 improvements are not as significant, with the exception for extreme precipitation. More recently, other studies such as Cardoso and Soares (2021) and Careto et al (2021), used a similar DAVs technique to assess the added value for other variables, simulations, and domains.

The precipitation historical period EURO-CORDEX simulations were also assessed for specific regions (Torma et al., 2015; Soares et al., 2017; Ciarlo et al., 2020). For instance, Torma et al (2015) evaluated precipitation over an alpine area, whereas Soares et al
100 (2017) assessed the same variable but for Portugal. Both studies describe the ability of the higher resolution runs in simulating the mean spatial and temporal patterns of precipitation, as well as their distributions. More recently, Ciarlo et al (2020) assessed the added value of all available EURO-CORDEX and CORDEX-CORE (Gutowski et al., 2016) simulations for precipitation, also considering a probability density function metric. The authors found added value, particularly at the tail of the distributions, however, they also report a significant uncertainty linked to the observational datasets in the results.

In this study, the DAV metric is used to assess the added value of precipitation for all available 12 km resolution simulations from the EURO-CORDEX Hindcast (1989-2008) and Historical (1971-2005) set. The added value is then computed comparing the RCMs precipitation results versus their corresponding driver GCM or ERA-Interim reanalysis, where the recently developed Iberia Gridded Dataset (IGD, Herrera et al., 2019) is considered as baseline. The IGD is a high-resolution dataset, with 0.1° resolution, and is based on a large number of weather stations covering the entire Iberian Peninsula. Thus, a new and unprecedented assessment

110 of the added value in the high-resolution EURO-CORDEX regional simulations is performed for this domain, with observations
at a similar resolution scale. The next section introduces the data and a description of the methods considered. The results and
discussion are presented in the following section. Finally, the main conclusions are drawn in the last section.

2 Data and Methods

2.1 Iberian Gridded Dataset

115 A recently developed dataset, the Iberian Gridded Dataset at 0.1° resolution (IGD, Herrera et al 2019) is used as a baseline for the
added value assessment. This dataset was built by considering an unprecedented number of weather stations – 3486 (275) for
precipitation (temperatures) over the entire Iberian Peninsula for daily precipitation, maximum, minimum and mean temperatures,
spanning over 45 years, from 1971 until the end of 2015. The authors performed a comparison with E-OBS v17 and v17e,
confirming the ability of this new dataset in reproducing the mean and extreme precipitation and also the temperature regimes.
120 Both datasets are comparable, yet statistically different. Since a large number of stations were considered, particularly for
precipitation, IGD should reproduce more realistically the climate of the Iberian Peninsula.

2.2 EURO-CORDEX

The aim of CORDEX is to develop a coordinated ensemble of high-resolution Regional Climate projections to provide detailed
climate data for all land regions of the world, at user-relevant scales, and support climate change impact and adaptation research
125 (Giorgi et al., 2009; Gutowski et al., 2016). All model data is available at the Earth System Grid Federation portal (Williams et al.,
2011). The EURO-CORDEX (Jacob et al., 2014; 2020) is a branch from the international CORDEX initiative and consists of a
multi-model ensemble of simulations at 50 km, 25 km, or 12 km resolutions for a European domain. These simulations consist in
Hindcast for the 1989-2008 period, forced by the ERA-Interim Reanalysis (Dee et al., 2011), and Historical/Scenario simulations
driven by the Intergovernmental Panel on Climate Change Coupled Model Intercomparison Project – Phase 5 (IPCC-CMIP5)
130 GCMs covering the 1971-2100 period. All simulations are available at the Earth System Grid Federation portal (Williams et al.,
2011; <https://esgf.llnl.gov/>).

The information regarding the simulations used are summarised in Table S1 for Hindcast and Table S2 for the Historical. For all
models, the added value is computed for the common Iberian Peninsula domain shown in Fig. 1, where prior to all computations,
all RCM model data was first conservatively interpolated (Schulzweida et al., 2009) into this observational domain, while the
135 observations were interpolated into each low-resolution grid. Thus, the evaluation of the EURO-CORDEX regional models is
performed at the 0.1° regular grid, while at the same time the GCMs or ERA-Interim (0.75°) are evaluated at their native resolutions
(see table S2 for each GCM resolution).

2.3 Distribution Added Value

The Distribution Added Value (DAV) is a metric put forward by Soares and Cardoso (2018) which allows assessing in a direct
140 way the gains or losses of using higher against lower resolution models relying on their Probability Density Functions (PDF), by
having an observational dataset as reference. DAV uses the PDF skill score proposed by Perkins et al (2007) to measure the
similarity between two different PDFs. In order to compute this metric, first, a PDF must be built from the data. In this work, two
slightly different methods are considered for building the PDFs to assess the daily precipitation from the EURO-CORDEX models.




Figure 1. Orography from the Iberian Gridded Dataset for the Iberian Peninsula at 0.1° horizontal resolution

145

In the first method the precipitation values are accumulated within each bin, thus returning a precipitation intensity distribution. While the second one considers the number of events that fall into each bin, thus returning a precipitation frequency distribution. Then, a normalisation is carried out by dividing each bin by the sum of all data considered as input for the PDF (Gutowski et al., 2007; Boberg et al., 2009; 2010). With this normalisation, one can more accurately compare the results between seasons or regions (Soares and Cardoso, 2018), but also, changes in PDF are identified more straightforwardly (Gutowski et al., 2007). Each bin has a width of 1 mmday^{-1} to avoid excessively fine and potential noisy steps in both methodologies, thus satisfying the criteria proposed by Wilks, (1995). All DAVs are computed by only considering the wet days, i.e., days with precipitation equal or above to 1 mm, as models tend to overestimate the days with very small precipitation amounts (Boberg et al., 2009; 2010; Soares and Cardoso, 2018). For either methodology, the score is given by the sum of the minimum value obtained at each bin between the models PDF and the observational PDF:

155

$$S = \sum_1^n \min(Z^m, Z^{obs}) \quad (1)$$

Where n is the number of bins for the PDFs, m denotes the high or the low-resolution simulation and obs is the observational PDF. For precipitation, the limits are bounded between 1 and 300 mm, roughly corresponding to the maximum precipitation rate in IGD. Subsequently, the DAV metric is then computed as follows:

$$160 \quad DAV = 100 * \frac{S_{hr} - S_{lr}}{S_{lr}} \quad (2)$$

with the subscript *hr* depicting the high resolution and *lr* the low resolution. The DAVs return the fraction or percentage of gains or losses of value by downscaling the low-resolution runs. With the normalisation of the PDFs, the contribution from each bin to the overall score of a particular model is more relevant for the lower bins, decreasing when approaching the tails of the distribution. If for a specific bin there is no model or observation data, then the contribution of that bin would be 0. By definition, the maximum value for S is 1, where if a specific model overestimates the observable PDF in one section, then it will inevitably underestimate in another section. Both these scenarios lower the score of individual models. DAVs is a versatile metric with the advantage of being able to be computed for PDF sections, which is useful for the extremes added value characterisation. In this study, the added value assessment is performed by considering not only the whole PDF but also for an extreme precipitation PDF section, where only values above the observational 95th percentile are accounted for. Since the resolution difference between observations and the high-resolution models is approximately 0.01°, this threshold is computed from the observations at the original resolution, while for the low-resolution driving models, the percentile is obtained from the interpolated observations.

For the DAVs assessment, firstly, a regional approach is considered by pooling together all data from the Iberia Peninsula, thus computing the added value for the entire domain. Secondly, a spatial approach is performed, where all data within each grid cell from the low-resolution simulation are pooled together, returning a DAV's spatial view, instead. Therefore, the Iberian overall value does not represent a mean from the spatial DAVs. Although the results should be similar, one must consider that different behaviour is expected, and care must be taken when comparing the results.

3 Results and Discussion

3.1 Hindcast (1989-2008)

This section presents the results for the EURO-CORDEX Hindcast (1989-2008) simulations, by applying the DAVs metric to precipitation and precipitation extremes. All results have the IGD as a reference. The precipitation PDFs are shown in the supplementary material. Figure S1 root to the results displayed in Fig. 2. Two different approaches are performed, one following a precipitation intensity PDF (left panels in Fig S1) and a precipitation frequency (right panels in Fig S1). Overall, the high-resolution RCM simulations capture better the observable PDFs, contrary to their lower-resolution counterparts. This behaviour suggests an expected and overall added value of the high-resolution runs relative to the coarser resolution over the Iberian Peninsula domain, for both the annual and seasonal timescales. Contrasts are visible for both methodologies, where for precipitation intensity, the differences between the low- and high-resolution PDFs are more perceptible, particularly at bins below the percentile thresholds. Thus, one can anticipate a generalised larger added value. On the other end, for precipitation frequency, and for the lower bins, the pdfs show a closer representation, almost overlapping, resulting in lower DAVs. Moreover, low-resolution runs tend to overestimate considerably the lower rainfall bins, and in consequence of the normalisation, the higher bins, roughly above the 15 *mm day*⁻¹ are underestimated. The same occurs for the high-resolution runs, however at a lower degree, hence reproducing more reliably the observable PDF.

In Fig. 2, the DAVs for the entire Iberian Peninsula are shown, revealing important gains of the EURO-CORDEX Hindcast high-resolution simulations in comparison to the driving ERA-Interim. Fig. 2a displays the DAVs for precipitation intensity considering the whole PDF with larger gains at the annual scale for 11 models, which have a DAV equal or above to 10%. From these CCLM, ETHZ, CNRM63, and SMHI models stand out surpassing 18%. CNRM53, ICTP, and IPSL RCMs show lower gains, ranging




Figure 2. Yearly and seasonal distribution added values (DAV) of the Iberian Peninsula, between the RCMs and the ERA-Interim reanalysis for the 1989-2008 period, taken from the Hindcast EURO-CORDEX simulations, with the IGD regular dataset as a reference for (a) daily precipitation intensity, considering the whole PDF shown in the left panels of Figure S1, (b) daily precipitation frequency considering the whole PDF shown in the right panels of Figure S1, (c) daily precipitation intensity extremes, only considering the values above the observational 95th percentile shown in Figure S1 left side and (d) daily precipitation frequency extremes, only considering the values above the observational 95th percentile shown in the right side of Figure S1. All RCM data were previously interpolated to 0.1° regular resolution from the observations, while the observations were interpolated into the ERA-Interim resolution

from ~5 to 10%. In winter, spring, and autumn the models roughly reproduce similar DAV values seen at the annual scale, however this is not the case for summer. Of all seasons, summer has the lowest performances, particularly CNRM53, DHMZ, and IPSL showing a detrimental effect ranging from -0.2 % to -6%. In fact, summer is the season where models display more difficulty in capturing the precipitation features, since it is the driest season for the entire Iberian Peninsula and precipitation is mostly associated to water recycling through soil moisture atmosphere feedbacks (Rios-Entenza et al., 2014)., In addition to the added importance of lower precipitation rates which models overestimate (Boberg et al. 2009; 2010; Soares and Cardoso, 2018), the representation of soil moisture in any model is still very challenging thus the weaker performance of the RCMs is not surprising. In fact, the summer PDF for the precipitation intensity (Fig. S1), in comparison with the other seasons, reveal a higher overestimation for the lowest bins and an underestimation in the tails, thus reducing the downscaling added value. Additionally, ERA-interim assimilates soil moisture and temperature, near surface temperature and humidity thus constraining the local land-atmosphere feedbacks and improving its added value.

While Fig. 2a shows the added value for the precipitation intensities, Fig. 2b considers the precipitation frequencies. The overall DAVs are lower, yet the models reveal similar differences, with the same models showing maximum DAVs in Figure 2a also present in Figure 2b. In this case, 4 models reveal some detrimental effects associated with the downscaling of ERA-Interim at the annual scale. For instance, the underperformance of the downscaling ICTP is highlighted with the negative values at the yearly


scale derived from the poorer performance for winter and autumn. Similar to the precipitation intensity, summer has the lowest
220 DAV values, but in this case 11 out of 13 models reveal losses, in particular DHMZ, KNMI, and MOHC with negative DAVs
superior to -7%. In fact, Herrera et al (2020) reported a bad performance for the DHMZ RCM for all the metrics. In the opposite
sense for most RCMs, winter and to some extent spring and autumn reveal gains, particularly winter with DAVs above 7% for 4
RCMs. In Herrera et al (2020), all regional models assessed against the same IGD still reveal strong biases for precipitations.
225 Nevertheless, the gains found here for either precipitation intensities (Fig. 2a) and frequencies (Fig. 2b) still reveal improvements
in comparison to the driving ERA-Interim reanalysis. Moreover, in Soares and Cardoso (2018) the Iberia displayed larger added
value in comparison to the other regions analysed.

The next panel shows the DAV metric but applied only for values above the observational 95th percentile, where the normalisation
is carried out considering just the bins above this threshold, thus only inspecting the added value related to the extreme precipitation
tail. Previous studies such as Soares and Cardoso (2018) and Ciarlo et al (2020) reported more relevant gains when looking into
230 extremes, with a few exceptions. Instead, results from Fig. 2c reveal lower DAVs compared to the whole PDF case (Fig. 2a), yet
noteworthy. The same was reported by Soares and Cardoso (2018) for the Iberia Peninsula, despite the low station density
considered, the DAVs reveal smaller values for the extremes and higher for the PDF as a whole. At the annual scale, 4 RCMs
reveal similar performances in comparison to the driving simulation, while the other RCMs have added value above ~10%. From
all seasons, winter reveals 4 RCMs with slight detrimental effects, contrasting with the results obtained from the other seasons and
235 against Fig. 2a. The downscaling shows a better performance, i.e., a larger added value, for the summer and autumn season for
more than half of the RCMs, of which they display gains equal or superior to 15%. The gains for the summer season where most
precipitation is convective (Azorin-Molina et al., 2014) are relevant, as low-resolution models have trouble in capturing these
highly spatial and temporal heterogeneities, due to shortcomings associated with the parametrisation of convection (Prein et al.,
2015). Spring also revealed high added value for almost all RCMs, ranging from 7.4% to 16.6%. The exception is for the DMI
240 model which has limited to no added value throughout the year, apart from autumn. In fact, this RCM is the only one showing a
similar performance to the driving simulation for 3 seasons. In Herrera et al (2020) most models overestimate the 50-year return
period for precipitation extremes for a large part of the domain, which is in line with the results shown in Fig. S1. However, the
low-resolution models cannot reproduce such high precipitation rates, which in the end results into added value, despite the
overestimation.

245 For the precipitation frequency extremes (Fig. 2d), the DAV values are almost always slightly lower in comparison to precipitation
intensity (Fig. 2c), but larger than Fig. 2b. The similarity across both methodologies at the yearly and seasonal scale is clear, where
summer and autumn reveal larger gains, while for spring and winter most RCMs display gains close to 10%. The exceptions are
DMI, GERICS, MPI, and MOHC, namely for winter and autumn with more limited DAVs.

Figure S2 from the supplemental material displays a slightly different approach than in Fig. 2. In this case, all model data was
250 previously interpolated to the 0.1° resolution from the observations. A conservative remapping of precipitation was considered
(Schuzweida et al., 2009), resulting in a smoothing of the precipitation field, which do not significantly impact the PDFs from
ERA-Interim. On the other hand, by upscaling the IGD, part of the original variability is conserved, thus changing the intensity or
frequency within each bin. These differences result in larger gains for Fig. S2 in comparison to Fig. 2. Nevertheless, the overall
inter-model variability is close between these two metrics.

255 Figure 3 displays the same metric used in Fig. 2, but from a spatial overview. Here a different approach is implemented, where the
percentiles and PDFs are computed by pooling together all information only within each grid cell from the low-resolution driving
model. Fig. 3a displays a spatialisation of the added value for the full precipitation intensity PDF, revealing a notable added value



260 **Figure 3. Yearly and seasonal spatial distribution added values (DAV) of the Iberian Peninsula, between the RCMs and the ERA-Interim**
reanalysis for the 1989-2008 period, taken from the Hindcast EURO-CORDEX simulations, with the IGD regular dataset as a reference,
for (a) daily precipitation intensity, considering the whole PDF, (b) daily precipitation frequency considering the whole PDF, (c) daily
precipitation intensity extremes, only considering the values above the observational 95th percentile from each ERA-Interim grid point
and (d) daily precipitation frequency extremes, only considering the values above the observational 95th percentile from each ERA-
265 **Interim grid point. All RCM data were previously interpolated to 0.1° regular resolution from the observations, while the observations**
were interpolated into the ERA-Interim resolution.

for most of the domain, as expected from the results obtained from Fig. 2a. In general, the gains are larger for the coastal areas and
the Pyrenees throughout the year, due to a better representation by the higher-resolutions of the land-sea and topographically
induced circulations. The more expressive gains for coastal areas, particularly for the Mediterranean, were also observed in Careto
270 et al (2021) for maximum and minimum temperatures. Similar to Fig. 2a, most models for winter, spring, and autumn mostly
display locations with large added value. For summer, the gains are more focused in the southern part of the peninsula, albeit most
models still reveal noteworthy percentages for the central and northern locations. RCMs such as CNRM53, ICTP, and to some
extent IPSL reveal sites in the Iberian interior with low or even negative added values. Those have a clear impact on the results
shown in Fig. 2a. Moreover, DHMZ which was revealed to have the minimum value in the summer regional overview, displays
275 some points in Fig. 3a with having a slightly detrimental effect, ~-10%.

In contrast with Fig. 3a and following the results from relative precipitation frequency in Fig. 2b, Fig. 3b shows an overall smaller
added value, following a very similar inter-model difference. While for precipitation intensity, values easily surpassed 50%, namely
near the coast, here the gains are more limited, going up to 30% in some coastal sites. Moreover, for the same models which had
a lower performance in the regional overview, particularly over the summer season, locations emerge having DAVs of ~-10% to
280 -15% in comparison with the driving simulation. Nevertheless, for the same season, all models reveal large added value for the
southern Peninsula, mirroring Fig. 3a. However, these positive values are not enough to reverse the losses found for the entire
domain in Fig. 2b. The regional overview of the DAVs is not a mean from the spatialisations and thus, care must be taken when
comparing both figures. For instance, the ICTP model in Fig. 2b had the worst overall performance with negative values for all
time scales, while in Fig. 3b the picture is different. ICTP only shows negative percentages at the annual and for the winter and
285 autumn seasons. On the other hand, DHMZ which had the minimum DAV in Fig. 2b, still reveals locations with large gains.

The picture for the extreme precipitation intensity and frequencies is completely different (Figs. 3c and 3d). Here the gains of the
high-resolution relative to the ERA-Interim reanalysis are evident when moving to more local scales. Moreover, the contrast of
values with the extremes shown in Fig. 2, highlights the difference in both methodologies. For the spatialisation, each point of the
low resolution only considers the amount of information available within, whereas, in Fig. 2, all data is considered. In the former,
290 there are fewer values above the observational percentiles and even less for the seasons, contrasting with the high-resolution. This
fact highlights the difficulty in the representation of extremes by the lower resolutions models, resulting in added value. Moreover,
in Fig. 2, precipitation intensity showed higher DAVs, while for Figs. 3c and 3d, the results are similar between themselves, with
points revealing gains above 250%. Still, at the annual scale, in comparison to the individual seasons, it is possible to see lower
percentages over the centre of Iberia, highlighting the improvements near the coast.

295 **3.2 Historical (1971-2005)**


This section displays the same metric, but applied to the Historical simulation, covering the 1971-2005 period. For this case, the
same RCM could be forced by different GCMs, however, the results do not necessarily have to agree. In fact, following the values

from Fig. 4, the different performances are more closely related to the different GCMs themselves than across the same high-resolution models. This enforces a weak or even no relationship between a single RCM forced by different GCMs. Moreover, any comparison with the previous Hindcast (Figs. 2 and 3) is hindered, not only owing to these differences, but also due to different time periods. Nevertheless, the range of DAVs for the Historical is considerably higher. Figs. S3 and S4 display the PDFs of models and observations. In the first all model data were previously interpolated to the 0.1° resolution of the IGD observations, as for the last, the PDFs from the GCMs are kept at their original resolutions, while the observations were interpolated into each GCM grid. Thus, the PDFs for the RCMs in Fig. S3, together with the PDFs in Fig. S4 root to the DAVs shown in Fig. 4. Following the results from the Hindcast simulations, there is a stronger agreement between all PDFs for precipitation frequency (right side of Figs. S3 and S4) in comparison to precipitation intensity (left side of Figs. S3 and S4), namely for the lower bins, anticipating a more difficulty in obtaining added value for the first.


The precipitation intensity with respect to the whole PDF is shown in Fig. 4a. For most cases, there exists some or even a larger added value. From those, models forced by the NCC GCM are highlighted with having the highest added values for all time scales, namely at the annual, with values above the 30%. For this GCM group, only the winter season displays slightly lower gains for some RCMs, yet still noteworthy. IPSL-LR-GERICS also display large gains at the annual scale, reaching 28%, contrasting with the slightly lower results for the individual seasons. The other GCM groups do not reveal such a relevant added value, namely models forced by CNRM, ICHEC1, ICHEC2, IPSL-MR and NOAA with gains ranging between 5% to 20%. Despite the overall gains, some models driven by CNRM for winter, spring, and summer reveal an absence of added value. Moreover, all ICHEC1 driven RCMs for winter, ICHEC1-DMI for spring, and some RCMs driven by ICHEC2, particularly for summer, reveal small and sometimes slightly negative percentages bounded between -5.7% to 0.4%. While the previous pairs still reveal some gains, MPI1 mostly show neutral percentages, oscillating between -6.9 % and 8 % indicating a closer performance between low and high resolutions in comparison with the cases before mentioned. As for MPI2 and MPI3, the RCMs driven by these two GCMs also reveal neutral percentages for almost all time-scales, with values up to 9.5 % for MPI2-ETHZ for autumn. Following the previous results, the MOHC driven RCMs also show lower gains, but in this case also some detrimental effects in 7 RCMs at the annual scale, thus having the worst results amongst all GCM-RCM pairs. From these same RCMs, at least one season also displays losses up to -13.8%. Nevertheless, MOHC-CCLM and MOHC-ETHZ reveal values close to 0 % throughout the year.

The different behaviour across the downscaling of each GCM group may be related not only to their resolution but also to the performance and quality of the GCMs itself, mainly within the lateral boundary forcing zone and also on the storm track positioning. For instance, Brands et al (2013) describes the MPI and MOHC as the best performing GCMs over these regions. This implies a more correct representation of the circulation in comparison with other GCMs. In fact, Zappa et al (2013) described the MOHC GCM by having a correct placement of storm tracks. However, a good performing GCM in terms of circulation, from which the RCMs inherit, does not translate into added value for the downscaling products. In fact, the Perkins skill score of these lower resolution models is higher, with 0.95 for MOHC for instance. This high score value implies an added difficulty for the downscaling RCMs in revealing added value. On the other hand, Jury et al. (2015) refers to the IPSL_MR as having a poor performance for upper-air variables over the same forcing region. Moreover, table 6 from McSweeney et al. (2015) resumes the overall performance for the individual GCMs, with CNRM, NOAA, MOHC and MPI as having a good performance; IPSL_MR and NCC with an intermediate performance; ICHEC and IPSL_LR with a poorer performance. In fact, the RCMs driven by either the MOHC or MPI GCMs have a difficulty in obtaining added value, whereas the models driven by NCC or IPSL_LR clearly display added value. The resolution of the GCMs can also play a major role in the added value of precipitation. For instance, although NOAA GCM reveals a good performance (McSweeney et al., 2015), at the same time it has one of the lowest resolutions,


(a)



(b)



(c)



(d)




Figure 4. Yearly and seasonal distribution added values (DAV) of the Iberian Peninsula, between the RCMs and the CMIP5 GCMs for the 1989-2008 period, taken from the Historical EURO-CORDEX simulations, with the IGD regular dataset as a reference for (a) daily precipitation intensity, considering the whole PDF shown in the left panels of Figure S3, (b) daily precipitation frequency considering the whole PDF shown in the right panels of Figure S3, (c) daily precipitation intensity extremes, only considering the values above the observational 95th percentile shown in Figure S3 left side and (d) daily precipitation frequency extremes, only considering the values above the observational 95th percentile shown in the right side of Figure S1. All RCM data were previously interpolated to 0.1° regular resolution from the observations, while the observations were interpolated into each CMIP5 GCM resolution. The thick blue lines separate the RCMs driven by different GCM.

which may be a possible reasoning behind the gains found in Fig. 4a for the NOAA-GERICS pair. An important note is the fact that the precipitation variable is not directly linked between the GCMs and RCMs, as only the circulation information is passed for the dynamical downscaling models.

350 The next panel displays the precipitation frequency relatively to the whole PDF (Fig. 4b). As with the Hindcast simulation (Fig. 2), the frequency reveals limited gains, but with similar inter-model differences, correlating well with the precipitation intensity approach. Thus, the overall results for precipitation frequency are lower and closer to 0%. In other words, the negative values are not as expressive, where for instance the losses for the MOHC driven RCMs are slightly less notable in comparison to Fig. 4a. At the other end of the spectrum, models forced by the NCC GCM still reveal some of the largest added value, albeit lower when compared to precipitation intensity. The exception is for winter where 6 RCMs display minor and slightly negative DAVs.

355 Following this reasoning, the other GCM-RCM pairs reveal a similar inter-model variability across Figs. 4a and 4b. The main reason for the similar behaviour, yet lower results, come from the closer PDFs seen for precipitation frequency in comparison to precipitation intensity (Figs. S3 and S4) and higher Perkins skill score obtained for the lower resolutions.

Figure 4c displays the results for the precipitation intensity extremes by considering only the values above the observationa 195th percentile. Comparing with the Hindcast results from Fig. 2, some GCM-RCM pairs do display larger added values, while at the same time showing a higher DAVs variability. Mirroring Fig. 4a, the NCC forced RCMs reveals a very noticeable added value at the annual scale, derived from the strong signal found seasonally, particularly for winter. At the same time, 4 RCMs from this group are highlighted for having gains above 30% for almost all seasons. The other GCM-RCM pairs do not reveal such expressive added value. Nevertheless, 2 RCMs driven by ICHEC1, 6 by ICHEC2, and all models forced by the IPSL GCMs, display at least one season with percentages above 20%. From these, 3 GCM-RCM pairs reveal gains superior to 30% for a single season. On the contrary, the DMI RCM forced by both ICHEC reveals lower gains, namely for spring and summer. Moreover, the IPSL-MR RCMs display small percentages for spring and even losses for winter, which overshadows the gains found in summer and autumn. The CNRM driven RCMs reveals 5 models with similar percentages found in Fig. 4a. However, CNRM-DMI and CNRM-GERICS display losses at the annual scale derived from a noticeable detrimental effect for summer and winter respectively. The losses found in extremes hint towards a lower accuracy for the RCMs in representing the higher bins. On the other hand, the models forced by

360 MPI1 show an overall intensification of either gains or losses found in Fig. 4a, meaning that the gains for the whole PDF are possibly derived from the gains obtained for the extremes. The same occurs for the NOAA-GERICS pair. As for models driven by MPI2 or MPI3, while in the whole PDF the DAVs for the high resolution are more neutral, for Fig.4c the values are more expressive surpassing 20% in two situations. Finally, 6 RCMs driven by MOHC reveal added value for the second half of the year, but with more modest values for winter and spring. From all pairs, MOHC-DMI reveals the lowest DAV, particularly for the winter season,

375 having repercussions at the annual scale.

The last panel from Fig. 4 shows the precipitation frequency. In this case, the similarity across precipitation intensity and precipitation frequency are evident as there is a good agreement between Figs. 4c and 4d, although as expected from before, with more limited DAVs, where the negative values are not as negative, e.g., winter season in MOHC-DMI, and the positive values are not as expressive, e.g., annual and winter season for models driven by the NCC GCM.

380 Similar to the Hindcast simulations, a second metric was implemented, where all data were interpolated to the 0.1° resolution from the observations (Fig. S5). The results here reveal an overall higher added value due to the stronger effect in upscaling the observations in comparison to the downscaling of the low-resolution. The gains are even more evident for both precipitation intensity and frequency extremes, with percentages well above 100% for models forced by the NCC GCM at the annual and winter season. Nevertheless, the DAVs in Fig. S5 correlate well with the results obtained in Fig. 4.


385 The next set of figures shows a spatialisation of the DAV metric, where individual PDFs and percentiles thresholds were considered
for each low-resolution grid point. As with the Hindcast in Fig.3, the spatialisation of the technique allows the emergence of points
with either high positive added value or losses, which would be masked otherwise. Figure 5 displays the results for precipitation
intensity, and overall, important gains are found for all models, even for those which underperformed at the Iberia Peninsula scale
in Fig. 4a. Nevertheless, it is possible to verify higher added value in coastal areas relative to inland points. This situation also
390 occurred for the Hindcast simulations and is owed mainly to a better representation of the land-sea circulations. Moreover, RCMs
forced by IPSL-LR, NCC, and NOAA, reveal most grid points with large positive gains, corroborating the results from Fig. 4a. On
the contrary, the GCMs RCMs pairs that displayed gains not as relevant as those listed before all display points with limited values
and sometimes small losses for sites in the interior, thus lowering the joint performance. Nevertheless, these pairs reveal substantial
added value, in particular on the Mediterranean coast. Similar to the previous cases, models forced by all three MPI GCM versions
395 reveal a similar behaviour, with higher gains in coastal areas and lower values in the interior. However, these results contrast with
the DAVs found in Fig. 4a, namely for models forced by MPI2, i.e., when assessing the precipitation at a more local scale, the
gains become even more evident. Lastly, models forced by the MOHC GCM still reveal relevance for the most part, namely for
summer in coastal areas, contrasting with the lower values in the other seasons. MOHC-ICTP had the overall worst performance,
yet some added value is still shown for points located on the Atlantic coast for winter, spring, and autumn.

400 Fig. 6 displays the same results, but for precipitation frequency instead. As expected from the previous cases, the overall gains are
more limited for both the positive and negative percentages. Still, a good agreement with Figs. 4b and 5 are found. From all GCM-
RCM pairs, 36 models reveal a better performance for most points within the domain for the summer season. Although a good
performance is revealed for most locations for the ICHEC driven GCMs in Figure 5, a widespread small DAVs occurs for
precipitation frequency. IPSL driven RCMs also show relevant gains, at the annual and seasonal scales, namely over Portugal and
405 on the Atlantic coast. As for the models forced by the NCC GCM, precipitation frequency revealed smaller DAVs in comparison
to Fig. 5. Only a single point located over Gibraltar displays a very large added value, above 30% for all seasons but summer.
The next figures show the results for precipitation intensity extremes (Fig. 7) and precipitation frequency extremes (Fig. 8). In
comparison to the Hindcast simulations, the results here reveal more spatial variability. Nevertheless, the added value is more
noteworthy for the DAVs spatialisation, rather than for the whole domain (Fig. 4), particularly for precipitation extremes due to
410 the fact this variable is highly localised. In contrast between Figs. 5 and 6, where some differences arose, the precipitation intensity
and frequency extremes (Figs. 7 and 8) reveal a closer behaviour. Similar to the previous cases, since the data is further split up
when assessing the seasons, the DAVs at the annual scale are usually slightly lower. In fact, all models in Figs. 7 and 8 clearly
reveal this nature. Moreover, from all model pairs, only the MOHC, NCC, and NOAA groups display smaller DAVs for points in
the interior, while the other pairs mostly show large gains throughout the entire domain.

415 **4 Summary and conclusions**

In this study, the performance of RCMs from the Hindcast (1989-2008) and Historical (1971-2005) simulations is assessed
relatively to their PDFs, by using a distribution added value metric proposed by Soares and Cardoso (2018). This assessment has
the IGD regular gridded dataset observations as a reference, covering the entire Iberian Peninsula. To this end, all RCM information
was first interpolated to the 0.1° resolution from the observations, while the low-resolution is assessed at their native resolution.
420 Two slightly different approaches were considered here, one following a precipitation intensity PDF and the other a precipitation
frequency. Between both, the results reveal very similar inter-model differences, however, a stronger signal is found for
precipitation intensity. Nevertheless, all RCMs reveal a notable added value, particularly in the representation of extremes, where

Precipitation Intensity 0-100



425 **Figure 5. Yearly and seasonal distribution added values (DAV) of the Iberian Peninsula for the Historical (1971-2005) EURO-CORDEX RCMs, with the IGD as a reference for the daily precipitation intensity, considering the whole PDF. All RCM data were previously interpolated to 0.1° regular resolution from the observations, while the observations were interpolated into each CMI5 GCM resolution.**

the global models have more difficulty in describing the higher precipitation rates. This result is expected and shows the importance of considering regional models with higher resolution. However, in some isolated cases, the RCMs instead display neutral or even a slight deterioration effect. On the other hand, by considering a more local scale in the spatialisation of the DAVs, in particular for extreme precipitation, very large added values within the entire Iberia Peninsula are revealed. These gains are more relevant for coastal sites possibly owed to the better representation of the land-sea boundary.

430 Previous works warned about the uncertainty owed to interpolation procedures (Ciarlo et al., 2020). In a way interpolating the GCMs to higher resolutions could generate unrealistic values, whereas upscaling the high resolution degrades the spatial information, affecting primarily the tail end of the distributions (Torma et al., 2015; Prein et al., 2016). To gauge these differences a second methodology was investigated, where all data were interpolated to the 0.1° resolution from the observations. In this case, 435 the overall DAVs revealed higher added value. These results hint towards a stronger effect in the upscaling of the high-resolution

Precipitation Frequency 0-100





Figure 6. Yearly and seasonal distribution added values (DAV) of the Iberian Peninsula for the Historical (1971-2005) EURO-CORDEX RCMs, with the IGD as a reference for the daily precipitation frequency, considering the whole PDF. All RCM data were previously interpolated to 0.1° regular resolution from the observations, while the observations were interpolated into each CMI5 GCM resolution.

PDF, against the effect of the generation of spurious values when interpolating lower-resolution datasets, not considering local
 440 feedback systems and the effect that higher resolution topography has on precipitation. Nonetheless, since unrealistic values may be created, the uncertainty associated to this second approach is higher.

While the DAVs metric allows for quantification of the gains or losses by the downscaling of the global models, no relationship is found when the same RCM is forced by multiple GCMs. More importantly a strong connection is observed for high-resolution models driven by the same GCM. The performance of the GCM along the regions of lateral forcing for the EURO-CORDEX plays
 445 an important role in the ability of the RCMs to downscale precipitation. This study clearly shows that the gains obtained from the use of higher resolution RCMs are paramount, not only owed to finer details in the representation of variables by itself, but also due to the increased description of orography, and land-ocean-atmosphere feedbacks, which all have important impacts on precipitation. The added value associated to the higher resolution gives credence to the growing effort to perform increasingly

Precipitation Intensity 95-100



450 **Figure 7. Yearly and seasonal distribution added values (DAV) of the Iberian Peninsula for the Historical (1971-2005) EURO-CORDEX RCMs, with the IGD as a reference for the daily precipitation intensity, only considering the values above the observational 95th percentile. All RCM data were previously interpolated to 0.1° regular resolution from the observations, while the observations were interpolated into each CMI5 GCM resolution.**

455 higher resolution simulations up to convection permitting scales. However, the inter-model variability supports the need for a coordinated ensemble of simulations similar to the one of the CORDEX Flagship Pilot Study on “Convective phenomena at high resolution over Europe and the Mediterranean”. Increasing resolution implies higher computational costs; thus, in the last years the CORDEX community identified as a major challenge the objective quantification of RCM added value in respect to the GCM forcing. Added value assessments will allow the detection of future model development needs.

460

Precipitation Frequency 95-100




Figure 8. Yearly and seasonal distribution added values (DAV) of the Iberian Peninsula for the Historical (1971-2005) EURO-CORDEX RCMs, with the IGD as a reference for the daily precipitation frequency, only considering the values above the observational 95th percentile. All RCM data were previously interpolated to 0.1° regular resolution from the observations, while the observations were interpolated into each CMI5 GCM resolution.

465

Data availability

All model and observational datasets are publicly available. The regional and global model data is available through the Earth System Grid Federation portal (Williams et al., 2011; <https://esgf.llnl.gov/>). The ERA-Interim reanalysis is available at the ECMWF portal (<https://www.ecmwf.int/>). The Iberia01 dataset is publicly available through the DIGITAL.CSIC open science service (Herrera et al., 2019a, <https://doi.org/10.20350/digitalCSIC/8641>).

470

Author contribution

J Careto computed all results and developed the manuscript with contributions from all co-authors. PMM Soares and RM Cardoso developed the metric in which this paper is based on. S Herrera, J Gutiérrez, PMM Soares and R M Cardoso previously developed the observation based Iberian Gridded Dataset.

475 Competing interests

The authors declare that they have no conflict of interest.

Acknowledgments

The authors would like to thank all the individual participating institutes, listed in Tables S1 and S2, and the Earth System Grid Federation infrastructure for providing all the model data used in this study. The authors also acknowledge the Iberian Gridded
480 dataset (IGD) <http://hdl.handle.net/10261/183071>.

Financial support

J Careto is supported by the Portuguese Foundation for Science and Technology (FCT) with the Doctoral Grant SFRH/BD/139227/2018 financed by national funds from the MCTES, within the Faculty of Sciences, University of Lisbon. PMM Soares would like to acknowledge the financial support of FCT through project UIDB/50019/2020 – IDL and EEA-Financial
485 Mechanism 2014-2021 and the Portuguese Environment Agency through Pre-defined Project-2 National Roadmap for Adaptation XXI (PDP-2). RM Cardoso are supported by the FCT under the project LEADING (PTDC/CTA-MET/28914/2017). This work was also supported by project FCT UIDB/50019/2020 - Instituto Dom Luiz (IDL).

References

- Azorin-Molina, C., Tijm, S., Ebert, E.E., Vicente-Serrano, S.M. and Estrela, M.J.: Sea breeze thunderstorms in the eastern Iberian
490 Peninsula. Neighborhood verification of HIRLAM and HARMONIE precipitation forecasts. *Atmospheric research*, 139, 101-115, <https://doi.org/10.1016/j.atmosres.2014.01.010>, 2014.
- Ban, N., Schmidli, J., Schär, C.: Evaluation of the convection-resolving regional climate modeling approach in decade-long simulations. *Journal of Geophysical Research: Atmospheres*, 119, 7889–7907, <https://doi.org/10.1002/2014JD021478>, 2014.
- Ban, N., Caillaud, C., Coppola, E., Pichelli, E., Sobolowski, S., Adinolfi, M., Ahrens, B., Alias, A., Anders, I., Bastin, S. and
495 Belušić, D.: The first multi-model ensemble of regional climate simulations at kilometer-scale resolution, Part I: Evaluation of precipitation. *Climate Dynamics*, 1-28, <https://doi.org/10.1007/s00382-021-05708-w>, 2021.
- Berthou, S., Kendon, E.J., Chan, S.C., Ban, N., Leutwyler, D., Schär, C., and Fosser, G.: Pan-European climate at convection-permitting scale: a model intercomparison study. *Climate Dynamics*, 55, 35-59, <https://doi.org/10.1007/s00382-018-4114-6>, 2020.
- Boberg, F., Berg, P., Thejll, P., Gutowski, W.J. and Christensen, J.H.: Improved confidence in climate change projections of
500 precipitation evaluated using daily statistics from the PRUDENCE ensemble. *Climate dynamics*, 32, 1097-1106, <https://doi.org/10.1007/s00382-008-0446-y>, 2009.

- Boberg, F., Berg, P., Thejll, P., Gutowski, W.J. and Christensen, J.H.: Improved confidence in climate change projections of precipitation further evaluated using daily statistics from ENSEMBLES models, *Climate dynamics*, 35, 1509-1520. <https://doi.org/10.1007/s00382-009-0683-8>, 2010.
- 505 Brands, S., Herrera, S., Fernández, J. and Gutiérrez, J.M., How well do CMIP5 Earth System Models simulate present climate conditions in Europe and Africa?. *Climate dynamics*, 41, 803-817, <https://doi.org/10.1007/s00382-013-1742-8>, 2013.
- Cardoso, R.M. and Soares, P.M.: Is there added value in the EURO-CORDEX hindcast temperature simulations? Assessing the added value using climate distributions in Europe, Submitted to *Climate Dynamics*, <https://doi.org/10.1002/joc.7472>, 2021.
- Cardoso, R.M., Soares, P.M.M., Miranda, P.M.A. and Belo-Pereira, M.: WRF high resolution simulation of Iberian mean and extreme precipitation climate. *International Journal of Climatology*, 33, 2591-2608, <https://doi.org/10.1002/joc.3616>, 2013.
- 510 Careto, J.A.M., Soares, P.M.M., Cardoso, R.M., Herrera, S. and Gutiérrez, J.M.. Added value of the EURO-CORDEX high-resolution downscaling over the Iberian Peninsula revisited. Part II: Max and Min Temperature. *Geoscientific Model Development Discussions*, pp.1-25, <https://doi.org/10.5194/gmd-2021-208>, 2021.
- Casanueva, A., Herrera, S., Fernández, J. and Gutiérrez, J.M.: Towards a fair comparison of statistical and dynamical downscaling in the framework of the EURO-CORDEX initiative. *Climatic Change*, 137, 411-426, <https://doi.org/10.1007/s10584-016-1683-4>, 2016a.
- Casanueva, A., Kotlarski, S., Herrera, S., Fernández, J., Gutiérrez, J.M., Boberg, F., Colette, A., Christensen, O.B., Goergen, K., Jacob, D. and Keuler, K.: Daily precipitation statistics in a EURO-CORDEX RCM ensemble: added value of raw and bias-corrected high-resolution simulations. *Climate dynamics*, 47, 719-737, <https://doi.org/10.1007/s00382-015-2865-x>, 2016b.
- 520 Christensen, J.H. and Christensen, O.B.: A summary of the PRUDENCE model projections of changes in European climate by the end of this century. *Climatic change*, 81, 7-30, <https://doi.org/10.1007/s10584-006-9210-7>, 2007.
- Christensen, O.B., Drews, M., Christensen, J.H., Dethloff, K., Ketelsen, K., Hebestadt, I. and Rinke, A.: The HIRHAM regional climate model. Version 5 (beta), 2007, <https://www.dmi.dk/fileadmin/Rapporter/TR/tr06-17.pdf>, last access: 29 April 2021.
- Christensen, J. H., Hewitson, B., Busuioc, A., Chen, A., Gao, X., Held, I., Jones, R., Kolli, R. K., Kwon, W.-T., Laprise, R., 525 Magaña Rueda, V., Mearns, L., Menéndez, C. G., Räisänen, J., Rinke, A., Sarr, A., Whetton, P.: Regional climate projections. In *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., Miller, H. L., Cambridge University Press: Cambridge, UK and New York, NY, TRN: GB07CC205, 2007.
- Ciarlo, J.M., Coppola, E., Fantini, A., Giorgi, F., Gao, X., Tong, Y., Glazer, R.H., Alavez, J.A.T., Sines, T., Pichelli, E. and 530 Raffaele, F.: A new spatially distributed added value index for regional climate models: the EURO-CORDEX and the CORDEX-CORE highest resolution ensembles. *Climate Dynamics*, 1-22, <https://doi.org/10.1007/s00382-020-05400-5>, 2020.
- Colin, J., Déqué, M., Radu, R. and Somot, S.: Sensitivity study of heavy precipitation in Limited Area Model climate simulations: influence of the size of the domain and the use of the spectral nudging technique. *Tellus A: Dynamic Meteorology and Oceanography*, 62, 591-604, <https://doi.org/10.1111/j.1600-0870.2010.00467.x>, 2010.
- 535 Coppola, E., Sobolowski, S., Pichelli, E., Raffaele, F., Ahrens, B., Anders, I., Ban, N., Bastin, S., Belda, M., Belusic, D. and Caldas-Alvarez, A.: A first-of-its-kind multi-model convection permitting ensemble for investigating convective phenomena over Europe and the Mediterranean. *Climate Dynamics*, 55, 3-34, <https://doi.org/10.1007/s00382-018-4521-8>, 2020.
- Daniel, M., Lemonsu, A., Déqué, M., Somot, S., Alias, A. and Masson, V.: Benefits of explicit urban parameterization in regional climate modeling to study climate and city interactions. *Climate Dynamics*, 52, 2745-2764, [https://doi.org/10.1007/s00382-018-](https://doi.org/10.1007/s00382-018-4289-x)
- 540 [4289-x](https://doi.org/10.1007/s00382-018-4289-x), 2019.

- Dee, D.P., Uppala, S.M., Simmons, A.J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M.A., Balsamo, G., Bauer, D.P. and Bechtold, P.: The ERA-Interim reanalysis: Configuration and performance of the data assimilation system. *Quarterly Journal of the royal meteorological society*, 137, 553-597, <https://doi.org/10.1002/qj.828>, 2011.
- Di Luca, A., de Elía, R., and Laprise, R.: Potential for added value in precipitation simulated by high-resolution nested regional climate models and observations. *Climate dynamics*, 38, 1229-1247, <https://doi.org/10.1007/s00382-011-1068-3>, 2012.
- Di Luca, A., de Elía, R., and Laprise, R.: Potential for small scale added value of RCM's downscaled climate change signal. *Climate Dynamics*, 40, 601-618, <https://doi.org/10.1007/s00382-012-1415-z>, 2013.
- Fosser, G., Khodayar, S., and Berg, P.: Climate change in the next 30 years: What can a convection-permitting model tell us that we did not already know?. *Climate Dynamics*, 48, 1987-2003, <https://doi.org/10.1007/s00382-016-3186-4>, 2017.
- 545 Froidevaux, P., Schlemmer, L., Schmidli, J., Langhans, W. and Schär, C.: Influence of the background wind on the local soil moisture-precipitation feedback. *Journal of the atmospheric sciences*, 71, 782-799, <https://doi.org/10.1175/JAS-D-13-0180.1>, 2014.
- Fumière, Q., Déqué, M., Nuissier, O., Somot, S., Alias, A., Caillaud, C., Laurantin, O. and Seity, Y.: Extreme rainfall in Mediterranean France during the fall: added value of the CNRM-AROME Convection-Permitting Regional Climate Model. *Climate Dynamics*, 55, 77-91, <https://doi.org/10.1007/s00382-019-04898-8>, 2020.
- 555 Giorgi, F. and Bates, G.T.: The climatological skill of a regional model over complex terrain. *Monthly Weather Review*, 117, 2325-2347, [https://doi.org/10.1175/1520-0493\(1989\)117<2325:TCSOAR>2.0.CO;2](https://doi.org/10.1175/1520-0493(1989)117<2325:TCSOAR>2.0.CO;2), 1989.
- Giorgi, F. and Mearns, L.O.: Approaches to the simulation of regional climate change: a review. *Reviews of geophysics*, 29, 191-216, <https://doi.org/10.1029/90RG02636>, 1991.
- 560 Giorgi, F. and Mearns, L.O.: Introduction to special section: Regional climate modeling revisited. *Journal of Geophysical Research: Atmospheres*, 104, 6335-6352, <https://doi.org/10.1029/98JD02072>, 1999.
- Giorgi, F., Jones, C. and Asrar, G.R.: Addressing climate information needs at the regional level: the CORDEX framework. *World Meteorological Organization (WMO) Bulletin*, 58, 175-183, 2009.
- Giorgi, F., Coppola, E., Solmon, F., Mariotti, L., Sylla, M.B., Bi, X., Elguindi, N., Diro, G.T., Nair, V., Giuliani, G. and 565 Turuncoglu, U.U.: RegCM4: model description and preliminary tests over multiple CORDEX domains. *Climate Research*, 52, 7-29, <https://doi.org/10.3354/cr01018>, 2012.
- Gutowski Jr, W.J., Takle, E.S., Kozak, K.A., Patton, J.C., Arritt, R.W. and Christensen, J.H.: A possible constraint on regional precipitation intensity changes under global warming. *Journal of Hydrometeorology*, 8, 1382-1396, <https://doi.org/10.1175/2007JHM817.1>, 2007.
- 570 Gutowski Jr, W. J., Giorgi, F., Timbal, B., Frigon, A., Jacob, D., Kang, H.-S., Raghavan, K., Lee, B., Lennard, C., Nikulin, G., O'Rourke, E., Rixen, M., Solman, S., Stephenson, T., and Tangang, F.: WCRP COordinated Regional Downscaling EXperiment (CORDEX): a diagnostic MIP for CMIP6, *Geosci. Model Dev.*, 9, 4087-4095, <https://doi.org/10.5194/gmd-9-4087-2016>, 2016.
- Heikkilä, U., Sandvik, A. and Sorteberg, A.: Dynamical downscaling of ERA-40 in complex terrain using the WRF regional climate model. *Climate dynamics*, 37, 1551-1564, <https://doi.org/10.1007/s00382-010-0928-6>, 2011.
- 575 Herrmann, M., Somot, S., Calmanti, S., Dubois, C., and Sevault, F.: Representation of spatial and temporal variability of daily wind speed and of intense wind events over the Mediterranean Sea using dynamical downscaling: impact of the regional climate model configuration, *Nat. Hazards Earth Syst. Sci.*, 11, 1983-2001, <https://doi.org/10.5194/nhess-11-1983-2011>, 2011

- Herrera, S., Cardoso, R.M., Soares, P.M., Espírito-Santo, F., Viterbo, P. and Gutiérrez, J.M.: Iberia01: a new gridded dataset of daily precipitation and temperatures over Iberia. *Earth System Science Data*, 11, 1947-1956, <https://doi.org/10.5194/essd-11-1947-2019>, 2019.
- Herrera, S., Soares, P.M., Cardoso, R.M. and Gutiérrez, J.M.: Evaluation of the EURO-CORDEX Regional Climate Models Over the Iberian Peninsula: Observational Uncertainty Analysis. *Journal of Geophysical Research: Atmospheres*, 125, p.e2020JD032880, <https://doi.org/10.1029/2020JD032880>, 2020.
- Hohenegger, C., Brockhaus, P., Bretherton, C.S. and Schär, C.: The soil moisture–precipitation feedback in simulations with explicit and parameterized convection. *Journal of Climate*, 22, 5003-5020, <https://doi.org/10.1175/2009JCLI2604.1>, 2009.
- Imamovic, A., Schlemmer, L. and Schär, C.: Collective impacts of orography and soil moisture on the soil moisture-precipitation feedback. *Geophysical Research Letters*, 44, 11-682, <https://doi.org/10.1002/2017GL075657>, 2017.
- Jacob, D., Elizalde, A., Haensler, A., Hagemann, S., Kumar, P., Podzun, R., Rechid, D., Remedio, A.R., Saeed, F., Sieck, K. and Teichmann, C.: Assessing the transferability of the regional climate model REMO to different coordinated regional climate downscaling experiment (CORDEX) regions. *Atmosphere*, 3, 181-199, <https://doi.org/10.3390/atmos3010181>, 2012.
- Jacob, D., Petersen, J., Eggert, B., Alias, A., Christensen, O.B., Bouwer, L.M., Braun, A., Colette, A., Déqué, M., Georgievski, G. and Georgopoulou, E.: EURO-CORDEX: new high-resolution climate change projections for European impact research. *Regional environmental change*, 14, 563-578, <https://doi.org/10.1007/s10113-013-0499-2>, 2014.
- Jacob, D., Teichmann, C., Sobolowski, S., Katragkou, E., Anders, I., Belda, M., Benestad, R., Boberg, F., Buonomo, E., Cardoso, R.M., Casanueva, A., Christensen, O.B., Christensen, J.H., Coppola, E., De Cruz, L., Davin, E.L., Dobler, A., Domínguez, M., Fealy, R., Fernandez, J., Gaertner, M.A., García-Díez, M., Giorgi, F., Gobiet, A., Goergen, K., Gómez-Navarro, J.J., Alemán, J.J.G., Gutiérrez, C., Gutiérrez, J.M., Güttler, I., Haensler, A., Halenka, T., Jerez, S., Jiménez-Guerrero, P., Jones, R.G., Keuler, K., Kjellström, E., Knist, S., Kotlarski, S., Maraun, D., van Meijgaard, E., Mercogliano, P., Montávez, J.P., Navarra, A., Nikulin, G., Noblet-Ducoudré, N., Panitz, H.J., Pfeifer, S., Piazza, M., Pichelli, E., Pietikäinen, J.P., Prein, A.F., Preuschmann, S., Rechid, D., Rockel, B., Romera, R., Sánchez, E., Sieck, K., Soares, P.M.M., Somot, S., Srnec, L., Sørland, S.L., Termonia, P., Truhetz, H., Vautard, R., Warrach-Sagi, K. and Wulfmeyer, V.: Regional climate downscaling over Europe: perspectives from the EURO-CORDEX community. *Reg Environ Change* 20, 51. <https://doi.org/10.1007/s10113-020-01606-9>, 2020.
- Jury, M.W., Prein, A.F., Truhetz, H. and Gobiet, A.: Evaluation of CMIP5 models in the context of dynamical downscaling over Europe. *Journal of Climate*, 28, 5575-5582, <https://doi.org/10.1175/JCLI-D-14-00430.1>, 2015
- Kendon, E.J., Roberts, N.M., Senior, C.A. and Roberts, M.J.: Realism of rainfall in a very high-resolution regional climate model. *Journal of Climate*, 25, 5791-5806. <https://doi.org/10.1175/JCLI-D-11-00562.1>, 2012.
- Kendon, E.J., Roberts, N.M., Fowler, H.J., Roberts, M.J., Chan, S.C. and Senior, C.A.: Heavier summer downpours with climate change revealed by weather forecast resolution model. *Nature Climate Change*, 4, 570-576, <https://doi.org/10.1038/nclimate2258>, 2014.
- Keuler, K., Radtke, K., Kotlarski, S., and Lüthi, D.: Regional climate change over Europe in COSMO-CLM: Influence of emission scenario and driving global model, *Meteorologische Zeitschrift*, 25, 121–136, <https://doi.org/10.1127/metz/2016/0662>, 2016.
- Khan, M.S., Coulibaly, P. and Dibike, Y.: Uncertainty analysis of statistical downscaling methods. *Journal of Hydrology*, 319, 357-382, <https://doi.org/10.1016/j.jhydrol.2005.06.035>, 2006.
- Kirshbaum, D.J., Adler, B., Kalthoff, N., Barthlott, C. and Serafin, S.: Moist orographic convection: Physical mechanisms and links to surface-exchange processes. *Atmosphere*, 9, 80, <https://doi.org/10.3390/atmos9030080>, 2018.

- Klein Tank, A.M.G., Wijngaard, J.B., Können, G.P., Böhm, R., Demarée, G., Gocheva, A., Mileta, M., Pashiardis, S., Hejkrlik, L., Kern-Hansen, C. and Heino, R.: Daily dataset of 20th-century surface air temperature and precipitation series for the European Climate Assessment. *International Journal of Climatology: A Journal of the Royal Meteorological Society*, 22, 1441-1453, <https://doi.org/10.1002/joc.773>, 2002.
- 620 Klok, E.J. and Klein Tank, A.M.G.: Updated and extended European dataset of daily climate observations. *International Journal of Climatology: A Journal of the Royal Meteorological Society*, 29, 1182-1191, <https://doi.org/10.1002/joc.1779>, 2009.
- Kotlarski, S., Keuler, K., Christensen, O. B., Colette, A., Déqué, M., Gobiet, A., Goergen, K., Jacob, D., Lüthi, D., van Meijgaard, E., Nikulin, G., Schär, C., Teichmann, C., Vautard, R., Warrach-Sagi, K., and Wulfmeyer, V.: Regional climate modeling on European scales: a joint standard evaluation of the EURO-CORDEX RCM ensemble, *Geosci. Model Dev.*, 7, 1297–1333, 625 <https://doi.org/10.5194/gmd-7-1297-2014>, 2014.
- Laprise, R.: Regional climate modelling. *Journal of Computational Physics*, 227, 3641-3666, <https://doi.org/10.1016/j.jcp.2006.10.024>, 2008.
- Leung, L.R., Mearns, L.O., Giorgi, F. and Wilby, R.L.: Regional climate research: Needs and opportunities in *Bulletin of the American Meteorological Society*, 84, 89-95, Retrieved April 29, 2021, from <http://www.jstor.org/stable/26215433>, 2003.
- 630 van der Linden, P. and Mitchell, J.E.: ENSEMBLES: Climate Change and its Impacts: Summary of research and results from the ENSEMBLES project. Met Office Hadley Centre, FitzRoy Road, Exeter EX1 3PB, UK, 160, 2009.
- Leutwyler, D., Lüthi, D., Ban, N., Fuhrer, O. and Schär, C.: Evaluation of the convection-resolving climate modeling approach on continental scales. *Journal of Geophysical Research: Atmospheres*, 122, 5237-5258. <https://doi.org/10.1002/2016JD026013>, 2017.
- Liu, C., Ikeda, K., Rasmussen, R., Barlage, M., Newman, A.J., Prein, A.F., Chen, F., Chen, L., Clark, M., Dai, A. and Dudhia, J.: 635 Continental-scale convection-permitting modeling of the current and future climate of North America. *Climate Dynamics*, 49, 71-95, <https://doi.org/10.1007/s00382-016-3327-9>, 2017.
- McGregor, J.L.: Regional climate modelling. *Meteorology and Atmospheric Physics*, 63, 105-117, <https://doi.org/10.1007/BF01025367>, 1997.
- McSweeney, C.F., Jones, R.G., Lee, R.W. and Rowell, D.P., Selecting CMIP5 GCMs for downscaling over multiple regions. 640 *Climate Dynamics*, 44, 3237-3260, <https://doi.org/10.1007/s00382-014-2418-8>, 2015.
- Meehl, G.A., Stocker, T.F., Collins, W.D., Friedlingstein, P., Gaye, A.T., Gregory, J.M., Kitoh, A., Knutti, R., Murphy, J.M., Noda, A. and Raper, S.C.; Global climate projections In *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., Miller, H. L, Cambridge University Press: Cambridge, 645 UK and New York, NY, TRN: GB07CC205, 2007.
- Nabat, P., Somot, S., Cassou, C., Mallet, M., Michou, M., Bouniol, D., Decharme, B., Drugé, T., Roehrig, R., and Saint-Martin, D.: Modulation of radiative aerosols effects by atmospheric circulation over the Euro-Mediterranean region, *Atmos. Chem. Phys.*, 20, 8315–8349, <https://doi.org/10.5194/acp-20-8315-2020>, 2020.
- Perkins, S.E., Pitman, A.J., Holbrook, N.J. and McAneney, J.: Evaluation of the AR4 climate models' simulated daily maximum temperature, minimum temperature, and precipitation over Australia using probability density functions. *Journal of climate*, 20, 4356-4376, <https://doi.org/10.1175/JCLI4253.1>, 2007.
- Pichelli, E., Coppola, E., Sobolowski, S., Ban, N., Giorgi, F., Stocchi, P., Alias, A., Belušić, D., Berthou, S., Caillaud, C. and Cardoso, R.M.: The first multi-model ensemble of regional climate simulations at kilometer-scale resolution part 2: historical and future simulations of precipitation. *Climate Dynamics*, 1-22, <https://doi.org/10.1007/s00382-021-05657-4>, 2021.

- 655 Pothapakula, P. K., Primo, C., Sørland, S., and Ahrens, B.: The synergistic impact of ENSO and IOD on the Indian Summer Monsoon Rainfall in observations and climate simulations - an information theory perspective, *Earth Syst. Dynam*, 11, 903–923, <https://doi.org/10.5194/esd11-903-2020>, 2020.
- Prein, A.F., Gobiet, A., Suklitsch, M., Truhetz, H., Awan, N.K., Keuler, K. and Georgievski, G.: Added value of convection permitting seasonal simulations. *Climate Dynamics*, 41, 2655–2677, <https://doi.org/10.1007/s00382-013-1744-6>, 2013a.
- 660 Prein, A.F., Holland, G.J., Rasmussen, R.M., Done, J., Ikeda, K., Clark, M.P., and Liu, C.H.: Importance of regional climate model grid spacing for the simulation of heavy precipitation in the Colorado headwaters. *Journal of climate*, 26(13), 4848–4857, <https://doi.org/10.1175/JCLI-D-12-00727.1>, 2013b
- Prein, A.F., Langhans, W., Fosse, G., Ferrone, A., Ban, N., Goergen, K., Keller, M., Tölle, M., Gutjahr, O., Feser, F. and Brisson, E.: A review on regional convection-permitting climate modeling: Demonstrations, prospects, and challenges. *Reviews of geophysics*, 53, 323–361 <https://doi.org/10.1002/2014RG000475>, 2015.
- 665 Prein, A.F., Gobiet, A., Truhetz, H., Keuler, K., Goergen, K., Teichmann, C., Maule, C.F., Van Meijgaard, E., Déqué, M., Nikulin, G. and Vautard, R.: Precipitation in the EURO-CORDEX 0.11° and 0.44° simulations: high resolution, high benefits?, *Climate Dynamics*, 46, 383, <https://doi.org/10.1007/s00382-015-2589-y>, 2016.
- Randall, D.A., Wood, R.A., Bony, S., Colman, R., Fife, J., Kattsov, V., Pitman, A., Shukla, J., Srinivasan, J. and Stouffer, R.J., 2007. *Climate models and their evaluation*. In *Climate change 2007: The physical science basis*. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., Miller, H. L, Cambridge University Press: Cambridge, UK and New York, NY, TRN: GB07CC205, 2007.
- 670 Remedio, A.R., Teichmann, C., Bunttemeyer, L., Sieck, K., Weber, T., Rechid, D., Hoffmann, P., Nam, C., Kotova, L. and Jacob, D.: Evaluation of new CORDEX simulations using an updated Köppen–Trewartha climate classification. *Atmosphere*, 10, 726, <https://doi.org/10.3390/atmos10110726>, 2019.
- Rios-Entenza, A., Soares, P.M.M., Trigo, R.M., Cardoso, R.M. and Miguez-Macho, G.: Precipitation recycling in the Iberian Peninsula: spatial patterns and temporal variability. *Journal of Geophysical Research: Atmospheres*, 119, 5895–5912, <https://doi.org/10.1002/2013JD021274>, 2014.
- 680 Rummukainen, M.: State-of-the-art with regional climate models. *Wiley Interdisciplinary Reviews: Climate Change*, 1, 82–96, <https://doi.org/10.1002/wcc.8>, 2010.
- Rummukainen, M.: Added value in regional climate modeling. *Wiley Interdisciplinary Reviews: Climate Change*, 7, 145–159, <https://doi.org/10.1002/wcc.378>, 2016.
- Samuelsson, P., Jones, C.G., Willén, U., Ullerstig, A., Gollvik, S., Hansson, U.L.F., Jansson, E., Kjellström, M., C., Nikulin, G. and Wyser, K.: The Rossby Centre Regional Climate model RCA3: model description and performance. *Tellus A: Dynamic Meteorology and Oceanography*, 63, 4–23, <https://doi.org/10.1111/j.1600-0870.2010.00478.x>, 2011.
- Schulzweida, U.: *Climate Data Operators, User’s Guide*. Version 1.1.9, Max-Planck Institute for Meteorology, Hamburg Germany, <https://code.mpimet.mpg.de/projects/cdo/embedded/cdo.pdf>, last access 29 April 2021.
- Soares, P.M. and Cardoso, R.M.: A simple method to assess the added value using high-resolution climate distributions: application to the EURO-CORDEX daily precipitation. *International Journal of Climatology*, 38, 1484–1498, <https://doi.org/10.1002/joc.5261>, 2018.
- 690

- Soares, P.M., Cardoso, R.M., Miranda, P.M., de Medeiros, J., Belo-Pereira, M. and Espirito-Santo, F: WRF high resolution dynamical downscaling of ERA-Interim for Portugal. *Climate dynamics*, 39, 2497-2522, <https://doi.org/10.1007/s00382-012-1315-2>, 2012a.
- 695 Soares, P.M., Cardoso, R.M., Miranda, P.M., Viterbo, P. and Belo-Pereira, M: Assessment of the ENSEMBLES regional climate models in the representation of precipitation variability and extremes over Portugal. *Journal of Geophysical Research: Atmospheres*, 117, D07114, <https://doi.org/10.1029/2011JD016768>, 2012b.
- Soares, P.M., Cardoso, R.M., Semedo, Á., Chinita, M.J. and Ranjha, R.: Climatology of the Iberia coastal low-level wind jet: weather research forecasting model high-resolution results. *Tellus A: Dynamic Meteorology and Oceanography*, 66, 22377, <https://doi.org/10.3402/tellusa.v66.22377>, 2014.
- 700 Soares, P.M., Cardoso, R.M., Lima, D.C. and Miranda, P.M.: Future precipitation in Portugal: high-resolution projections using WRF model and EURO-CORDEX multi-model ensembles. *Climate Dynamics*, 49, 2503-2530. <https://doi.org/10.1007/s00382-016-3455-2>, 2017.
- Stocker, T.F., Qin, D., Plattner, G.K., Tignor, M.M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V. and Midgley, P.M. (Eds.): *Climate Change 2013: The physical science basis. contribution of working group I to the fifth assessment report of IPCC the intergovernmental panel on climate change*, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/CBO9781107415324>, 2014
- 705 Tinker, J., Lowe, J., Holt, J., Pardaens, A. and Wiltshire, A.: Validation of an ensemble modelling system for climate projections for the northwest European shelf seas. *Progress in Oceanography*, 138, 211-237, <https://doi.org/10.1016/j.pocean.2015.07.002>, 2015.
- 710 Torma, C., Giorgi, F. and Coppola, E.: Added value of regional climate modeling over areas characterized by complex terrain—Precipitation over the Alps. *Journal of Geophysical Research: Atmospheres*, 120, 3957-3972, <https://doi.org/10.1002/2014JD022781>, 2015.
- Vautard, R., Gobiet, A., Jacob, D., Belda, M., Colette, A., Déqué, M., Fernández, J., García-Díez, M., Goergen, K., Güttler, I. and Halenka, T.: The simulation of European heat waves from an ensemble of regional climate models within the EURO-CORDEX project. *Climate Dynamics*, 4, 2555-2575, <https://doi.org/10.1007/s00382-013-1714-z>, 2013.
- 715 Vautard, R., Kadyrov, N., Iles, C., Boberg, F., Buonomo, E., Buelow, K., Coppola, E., Corre, L., van Meijgaard, E., Nogherotto, R., Sandstad, M., Schwingshagl, C., Somot, S., Aalbers, E. E., Christensen, O., Ciarlo, J., Demory, M.-E., Giorgi, F., Jacob, D., Jones, R. G., Keuler, K., Kjellström, E., Lenderink, G., Levvasseur, G., Nikulin, G., Sillmann, J., Solidoro, C., Sørland, S., Steger, C., Teichmann, C., Warrach-Sagi, K., and Wulfmeyer, V.: Evaluation of the large EURO-CORDEX regional climate model ensemble, *Journal of Geophysical Research: Atmospheres*, <https://doi.org/10.1029/2019JD032344>, 2020.
- van Meijgaard, E., Van Ulft, L.H., Van de Berg, W.J., Bosveld, F.C., Van den Hurk, B.J.J.M., Lenderink, G. and Siebesma, A.P., 2008. The KNMI regional atmospheric climate model RACMO, version 2.1 (p. 43). De Bilt, Netherlands: KNMI, <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.475.9494&rep=rep1&type=pdf>, last access: 29 April 2021.
- 725 Wilby, R.L., Wigley, T.M.L., Conway, D., Jones, P.D., Hewitson, B.C., Main, J. and Wilks, D.S.: Statistical downscaling of general circulation model output: A comparison of methods. *Water resources research*, 34, 2995-3008, <https://doi.org/10.1029/98WR02577>, 1998.
- Wilks D.S.: *Statistical Methods in the Atmospheric Sciences*. Academic Press: Oxford, UK, 467, 1995.
- 730 Williams, D.N., Taylor, K.E., Cinquini, L., Evans, B., Kawamiya, M., Lautenschlager, M., Lawrence, B., Middleton, D. and Contributors, E.S.G.F., 2011. The Earth System Grid Federation: Software framework supporting CMIP5 data analysis and dissemination. *CIIVAR Exchanges*, 56, 40-42, http://centaur.reading.ac.uk/25732/1/WilEA11_CE.pdf, last access: 29 April 2021.