

RADiv1: a non-steady-state early diagenetic model for ocean sediments in Julia and MATLAB/GNU Octave

Olivier Sulpis^{1,2}, Matthew P. Humphreys³, Monica M. Wilhelmus^{4,5}, Dustin Carroll^{5,6}, William M. Berelson⁷, Dimitris Menemenlis⁵, Jack J. Middelburg¹, Jess F. Adkins⁸

- 5 ¹Department of Earth Sciences, Utrecht University, Utrecht, The Netherlands
²Department of Earth and Planetary Sciences, McGill University, Montreal, Canada
³Department of Ocean Systems (OCS), NIOZ Royal Netherlands Institute for Sea Research, Texel, The Netherlands
⁴Center for Fluid Mechanics, Brown University, Providence, USA
⁵Jet Propulsion Laboratory, California Institute of Technology, Pasadena, USA
10 ⁶Moss Landing Marine Laboratories, San José State University, Moss Landing, USA
⁷Department of Earth Sciences, University of Southern California, Los Angeles, USA
⁸Geological and Planetary Sciences, California Institute of Technology, Pasadena, USA

Correspondence to: Olivier Sulpis (o.j.t.sulpis@uu.nl)

Abstract. We introduce a time-dependent, one-dimensional model of early diagenesis that we term RADI, an acronym
15 accounting for the main processes included in the model: chemical Reactions, Advection, molecular and bio-Diffusion, and
bio-Irrigation. RADI is targeted for study of deep-sea sediments, in particular those containing calcium carbonates (CaCO₃).
RADI combines CaCO₃ dissolution driven by organic matter degradation with a diffusive boundary layer and integrates state-
of-the-art parameterizations of CaCO₃ dissolution kinetics in seawater, thus serving as a link between mechanistic surface-
reaction modelling and global-scale biogeochemical models. RADI also includes CaCO₃ precipitation, providing a continuum
20 between CaCO₃ dissolution and precipitation. RADI integrates components rather than individual chemical species for
accessibility and is straightforward to compare against measurements. RADI is the first diagenetic model implemented in Julia,
a high-performance programming language that is free and open source, and it is also available in MATLAB/GNU Octave.
Here, we first describe the scientific background behind RADI and its implementations. Then, we evaluate its performance in
three selected locations and explore other potential applications, such as the influence of tides and seasonality on early
25 diagenesis in the deep ocean. RADI is a powerful tool to study the time-transient and steady-state response of the sedimentary
system to environmental perturbation, such as deep-sea mining, deoxygenation or acidification events.

1 Introduction

The seafloor, which covers ~70% of the surface of the planet and modulates the transfer of materials and energy from the biosphere to the geosphere, remains for the vast majority unexplored. Today, this rich, poorly understood ecosystem is threatened locally by deep-sea mining activities (e.g. ploughing of the seabed), because it contains abundant valuable minerals and metals essential for the energy transition (Thompson et al., 2018). The deep ocean is also being perturbed globally by climate change, including seawater acidification caused by the uptake of ~10 billion tons of anthropogenic carbon dioxide (CO₂) into the ocean each year (Perez et al., 2018; Gruber et al., 2019), roughly a quarter of our total annual emissions (Friedlingstein et al., 2020). In this context, it is important to improve our understanding of the seafloor's response to environmental change.

Accumulation of sinking biogenic aggregates and lithogenic particles at the seafloor provides reactive material that regulates the chemical composition of sediment porewaters. Whereas biogenic particles typically sink through the water column at rates from a few meters to hundreds of meters per day (Riley et al., 2012), the same particles accumulate in sediments much more slowly, typically a few centimeters per thousand years (Jahnke, 1996). The residence time of solid particles in the top centimeter of sediments is therefore very long (a few hundred or thousand years) compared to their residence time in the water column (a few weeks). Additionally, while solutes are dispersed by advection in the water column, molecular diffusion dominates in porewaters, which is slower. The long residence time of reactive solid material in surface sediments, coupled with the slow diffusive transport of dissolved species, can lead to large gradients in chemical composition between sediment porewaters and the overlying seawater, inducing solute fluxes between the two (Hammond et al., 1996). Thus, the top few millimeters of the seafloor play a significant role in many major marine biogeochemical cycles.

The overall rate of biogeochemical reactions is determined by the slowest, 'rate-limiting' step, which can be (i) transport to or from the reaction site or (ii) the reaction kinetics of the particle at the mineral-water interface. At the seafloor, the rate-limiting step for many biogeochemical reactions is solute transport via molecular diffusion through the sediment porewaters or through the diffusive boundary layer (DBL). The DBL is a thin film of water extending up to a few millimeters above the sediment-water interface in which molecular diffusion is the dominant mode of solute transport. The presence of a DBL above the sediment-water interface (Fig. 1) has been reported by several investigators (Morse, 1974; Archer et al., 1989b; Gundersen and Jørgensen, 1990; Santschi et al., 1991; Glud et al., 1994) and its thickness depends on the composition and roughness of the substrate, as well as on the flow speed of the overlying seawater (Chriss and Caldwell, 1982; Dade, 1993; Røy et al., 2002; Han et al., 2018). Diffusive fluxes of solutes across the sediment-water interface are driven by concentration gradients between the overlying seawater and the sediment column being considered. If most of the concentration gradient for a given solute occurs within the porewaters, rather than within the DBL, then the diffusive flux of this solute is termed 'internal' or 'sediment-side controlled' (Boudreau and Guinasso, 1982). Conversely, if the majority of the concentration gradient for a given solute is within the DBL, the chemical flux across the sediment-water interface is termed 'external' or 'water-side transport-controlled'. In practice, the chemical exchange of most solutes is controlled by a combination of both regimes termed

65 ‘mixed-control’, such as dissolved oxygen (Jørgensen and Revsbech, 1985; Hondzo, 1998), radon (Homoky et al., 2016; Cook et al., 2018), and the products of calcium carbonate dissolution (Sulpis et al., 2018; Boudreau et al., 2020), which have concentration gradients on both sides of the sediment-water interface. Despite the importance of the DBL in controlling diffusive fluxes across the sediment-water interface, DBLs are not explicitly included in most models that simulate early diagenesis in the deep ocean.

70 Multiple numerical models simulating early diagenesis have previously been published (Burdige and Gieskes, 1983; Rabouille and Gaillard, 1991; Boudreau, 1996b; Van Cappellen and Wang, 1996; Soetaert et al., 1996b; Archer et al., 2002; Munhoven, 2007; Couture et al., 2010; Yakushev et al., 2017; Hülse et al., 2018; Munhoven, 2021), each with its own assumptions and best area of application (Paraska et al., 2014). For instance, most existing models are limited to a steady state and are thus unable to predict the transient sediment response to time-dependent phenomena such as tides, seasonal change,
75 ocean deoxygenation, or acidification. Moreover, most of these models do not take the presence of a DBL into account, even though diffusion through the DBL may control the overall rate of many biogeochemical reactions. Finally, as the landscape of computing software and programming languages evolves and improves computing efficiency and code accessibility, it is important to leverage emerging developments to implement new biogeochemical models. Here, we describe a new sediment porewater model built upon earlier work termed RADI, an acronym accounting for the main processes included in the model
80 that control the vertical distribution of solutes and solids: chemical Reactions, Advection, molecular and bio-Diffusion, and bio-Irrigation. The novelty of RADI is that it combines organic matter degradation-driven CaCO_3 dissolution (Archer et al., 2002) with a diffusive boundary layer (Boudreau, 1996b) and integrates the state-of-the art parameterization of CaCO_3 dissolution kinetics in seawater (Dong et al., 2019; Naviaux et al., 2019a). RADI thus links mechanistic surface-reaction modelling to global-scale biogeochemical models (Carroll et al., 2020). By integrating components (e.g., total alkalinity) rather
85 than individual chemical species (e.g., carbonate and bicarbonate ions), RADI is easy to compare to observations. RADI is implemented in two popular scientific programming languages: Julia and MATLAB/GNU Octave. To our knowledge, this is the first diagenetic model implemented in Julia (<https://julialang.org>), a high-level, high-performance, and cross-platform programming language that is free and open source (Bezanson et al., 2017). Here, we first describe the scientific background behind RADI and its implementations. Then, we evaluate its performance in three selected locations and explore other potential
90 applications, such as the influence of tides and seasonality on early diagenesis in the deep ocean.

2 Model description

In the following section, we describe how reactions, advection, diffusion, and irrigation are implemented in RADIv1. Model variables are *italicized* and their name as coded in the model are shown in `monospaced` font. Tables 1 and 2 include an inventory of model variables and parameters and a list of nomenclature for chemical species, respectively.

95

Table 1. Nomenclature of model parameters and variables

Variable	Model notation	Description	Equation #
General			
Z	z_max	Total height of the sediment column	
dz	z_res	Depth resolution	
z	depths	Array of modelled depths within the sediment	
T	stoptime	Total simulation time	
dt	interval	Time steps	
t	timesteps	Array of modelled timepoints	
ϕ_z	phi	Porewater porosity	3
β	phiBeta	Porosity attenuation coefficient	3
$\phi_{s,z}$	phiS	Solid volume fraction	4
θ^2	tort2	Squared tortuosity	24
F_v	Fvar	Solid deposition flux	
v_w	var_w	Bottom waters solute concentration	
δ	dbl	Diffusive boundary layer thickness	
T_w	T	Temperature	
Advection			
u	u	Porewater burial velocity	17
w	w	Solid burial velocity	15, 16, 18
$Pe_{h,z}$	Peh	One-half of the cell Peclet number	22
σ_z	sigma	Number from Fiadero and Veronis (1977)	21
Reactions			
c/p	RC	“Redfield” ratio for carbon	
n/p	RN	“Redfield” ratio for nitrogen	
p/p	RP	“Redfield” ratio for phosphorus	
K_v	Kvar	Half-saturation constant for a given electron acceptor	
K_v'	Kvari	Inhibition constant for a given electron acceptor	
$k_{reaction}$	kvar	Rate constant for a given chemical reaction	
$f_{v,z}$	fvar	Fractions of organic matter degraded by a given oxidant	7, 8
$\eta_{diss. ca.}$	order_diss_ca	Reaction order for calcite dissolution	12
$\eta_{diss. ar.}$	order_diss_ar	Reaction order for aragonite dissolution	13
$\eta_{prec. ca.}$	order_prec_ca	Reaction order for calcite precipitation	14
Ω_{ca}	OmegaCa	Seawater saturation state with respect to calcite	12, 14
Ω_{ar}	OmegaAr	Seawater saturation state with respect to aragonite	13
Diffusion			
$d_z(v)$	D_var_tort2	Effective molecular diffusion coefficient	23, 27
$d_z^0(v)$	D_var	“Free-solution” molecular diffusion coefficient	23
b_z	D_bio	Bioturbation coefficient	25, 26
λ_b	lambda_b	Characteristic bioturbation depth	26
Irrigation			
α_z	alpha	Irrigation coefficient	30, 31
λ_i	lambda_i	Characteristic depth for irrigation	31

2.1 Model structure and fundamental equation

130 RADI uses the same set of reactive-transport partial differential equations as implemented in CANDI (Boudreau, 1996b), i.e., for each solute component,

$$\frac{\partial v}{\partial t} = \frac{1}{\varphi} \frac{\partial}{\partial z} \left(\varphi d \frac{\partial v}{\partial z} - \varphi uv \right) + \alpha(v_w - v) + \Sigma R \quad (1),$$

and for each solid component,

$$\frac{\partial v}{\partial t} = \frac{1}{\varphi_s} \frac{\partial}{\partial z} \left(\varphi_s b \frac{\partial v}{\partial z} - \varphi_s wv \right) + \Sigma R \quad (2),$$

135 where v is the concentration of a given component, t is time, φ is sediment porosity and φ_s is the solid-volume fraction, d is the effective molecular diffusion coefficient and b is the bioturbation coefficient, z is depth, u is the porewater burial velocity and w is the solid burial velocity, α is the irrigation coefficient, v_w is the concentration of a solute in the bottom waters and ΣR is the net production rate from all biogeochemical reactions for a given component. Each of these terms will be described in detail later in the *Model description* section. These partial differential equations are solved numerically using the method of

140 lines described in Boudreau (1996b). Instead of searching for steady-state solutions directly, RADI computes the concentrations of a set of solids and solutes at each depth and time steps following a time vector set by the user. The user determines the simulation time depending on the objectives, e.g., multimillennial to predict a steady state, or a few days to study the response of the sedimentary system to high frequency cyclic phenomena such as tides. For initial conditions, the user can choose between predefined uniform values (e.g., set all concentrations to zero) or a set of saved concentrations (e.g., from

145 a previous simulation that has reached steady state). T is the total simulation time, dt is the temporal resolution, i.e., the interval between each timestep, and t refers to the array of modelled timepoints. All time units are in years (a). The interface between the surface sediment and overlying seawater, conventionally set at a sediment depth $z = 0$, represents the top layer of RADI's vertical axis (Fig. 1). The bottom layer of the model is at a sediment depth Z . Between these limits, n layers are present, each being separated by a constant vertical gap dz . Depth units are in meters. The values assigned to dz and dt depend on the nature

150 of the problem and on the kinetics of the chemical reactions. In the present study, all cases use $dz = 2$ mm and $dt = 1/128000$ a, i.e., ~4 minutes. If a lower dz is used, dt needs to be lowered as well to preserve numerical stability. In general, the ratio dz/dt should be kept below a value of 256 m/a. If dz is divided by two, dt needs to be divided by two as well, and the speed at which RADI runs will be reduced by a factor of four.

155

160

Table 2. Nomenclature of modelled chemical species. All variables are concentrations, expressed in mol per m³ of solid for solid species and mol per m³ of water for solute species.

Variable v	Model notation	Description
[O ₂]	dO2	Dissolved oxygen
[TAlk]	dalk	Total alkalinity
[ΣCO ₂]	dtCO2	Dissolved inorganic carbon
[Ca ²⁺]	dCa	Dissolved calcium
[ΣNO ₃]	dtNO3	Dissolved inorganic nitrogen
[ΣSO ₄]	dtSO4	Dissolved inorganic sulfate
[ΣPO ₄]	dtPO4	Dissolved inorganic phosphorus
[ΣNH ₄]	dtNH4	Dissolved inorganic nitrogen
[ΣH ₂ S]	dtH2S	Dissolved inorganic sulfide
[Fe ²⁺]	dFe	Dissolved iron
[Mn ²⁺]	dMn	Dissolved manganese
[POC _{refractory}]	proc	Refractory particulate organic carbon
[POC _{slow}]	psoc	Slow-decay particulate organic carbon
[POC _{fast}]	pfoc	Fast-decay particulate organic carbon
[<i>Calcite</i>]	pcalcite	Calcite
[<i>Aragonite</i>]	paragonite	Aragonite
[MnO ₂]	pMnO2	Manganese (IV) oxide
[Fe(OH) ₃]	pFeOH3	Iron (III) hydroxide
[<i>Clay</i>]	pclay	Clay ¹

¹We consider all clay minerals to be montmorillonite (Al₂H₂O₁₂Si₄, molar mass = 360.31 g mol⁻¹)

RADI operates on a static, user-defined porosity profile. Sediment porosity, φ_z in Fig. 1, refers to the porewater volume fraction in the sediment (dimensionless) and typically decreases exponentially with sediment depth due to steady-state compaction. The sediment porosity profile is parametrized following Boudreau (1996b) as:

$$\varphi_z = \varphi_\infty + (\varphi_0 - \varphi_\infty)e^{-\beta z} \quad (3),$$

where φ_∞ is the porosity at great depth, φ_0 is the porosity at the sediment-water interface, and β is an attenuation coefficient expressed in m⁻¹. A typical deep-sea sediment porosity profile is shown in Fig. 1. Here the measured porosity profile at station 7 of cruise NBP98-2 (Sayles et al., 2001) is fit using $\varphi_\infty = 0.87$, $\varphi_0 = 0.915$, and $\beta = 33$ m⁻¹. The solid volume fraction (φ_s , dimensionless) is defined as:

$$\varphi_{s,z} = 1 - \varphi_z \quad (4),$$

and increases with sediment depth (as compaction forces squeeze porewaters out).




Figure 1. Schematic of RADI's vertical structure alongside steady-state depth profiles of porosity ϕ_z , see Eqs. (3,4), porewater (u , light blue solid line) and solid (w , brown solid line) burial velocities at in situ conditions taken at station 7 of Sayles et al. (2001). Burial velocity varies with depth due to porosity, as described in *Advection* Section 2.3. The open circles in the porosity profile are porosity measurements from Sayles et al. (2001).

195

Within this grid and for each time step, RADI computes the concentrations of 11 solute variables (Talk , ΣCO_2 , O_2 , Ca^{2+} , ΣNO_3 , ΣSO_4 , ΣPO_4 , ΣNH_4 , $\Sigma\text{H}_2\text{S}$, Fe^{2+} , and Mn^{2+}) and 8 solid variables (*Calcite*, *Aragonite*, $\text{Fe}(\text{OH})_3$, MnO_2 , *Clay*, and three kinds of particulate organic carbon collectively termed POC). Note that clay is simply modelled as a non-reactive solid, that is included because the clay accumulation flux to the sediment-water interface participates to the calculation of the solid burial velocity, see *Advection* section 2.3. Concentration units are in mol per m^3 of water for solutes and in mol per m^3 of solid for solid species, For each modelled solute or solid concentration v at time t and sediment-depth z :

$$v_{(t+d\delta),z} = v_{t,z} + [R(v_{t,z}) + A(v_{t,z}) + D(v_{t,z}) + I(v_{t,z})] \cdot dt \quad (5),$$

where $R(v_{t,z})$ quantifies the rate of change of $v_{t,z}$ due to chemical reactions, $A(v_{t,z})$ quantifies the rate of change of $v_{t,z}$ due to advection, $D(v_{t,z})$ quantifies the rate of change of $v_{t,z}$ due to molecular and bio-diffusion, and $I(v_{t,z})$ quantifies the rate of change of $v_{t,z}$ due to bio-irrigation. In general, only the subscript zs are explicitly written out in this document, for variables and parameters that vary with depth. The ts are implicit but excluded for clarity.

205

Table 3. Diagenetic reactions, reaction rates, and reactions contributions to porewater Talk and ΣCO_2 .

Reaction	Rate [mM a^{-1}]	ΔTalk	$\Delta\Sigma\text{CO}_2$
Organic matter degradation			
$(\text{CH}_2\text{O})(\text{NH}_3)_{\frac{m}{c}}(\text{H}_3\text{PO}_4)_{\frac{p}{c}} + \text{O}_2 \leftrightarrow \text{CO}_2 + \frac{m}{c}\text{NH}_3 + \frac{p}{c}\text{H}_3\text{PO}_4 + \text{H}_2\text{O}$	$(k_{\text{POC}_{fast}}[\text{POC}_{fast}] + k_{\text{POC}_{slow}}[\text{POC}_{slow}])f_{\text{O}_2}$	$+n/c - p/c$	+1
$(\text{CH}_2\text{O})(\text{NH}_3)_{\frac{m}{c}}(\text{H}_3\text{PO}_4)_{\frac{p}{c}} + 0.8\text{NO}_3^- \leftrightarrow 0.2\text{CO}_2 + 0.4\text{N}_2 + 0.8\text{HCO}_3^- + \frac{m}{c}\text{NH}_3 + \frac{p}{c}\text{H}_3\text{PO}_4 + 0.6\text{H}_2\text{O}$	$(k_{\text{POC}_{fast}}[\text{POC}_{fast}] + k_{\text{POC}_{slow}}[\text{POC}_{slow}])f_{\Sigma\text{NO}_3}$	$+0.8 + n/c - p/c$	+1
$(\text{CH}_2\text{O})(\text{NH}_3)_{\frac{m}{c}}(\text{H}_3\text{PO}_4)_{\frac{p}{c}} + 2\text{MnO}_2 + 3\text{CO}_2 + \text{H}_2\text{O} \leftrightarrow 4\text{HCO}_3^- + 2\text{Mn}^{2+} + \frac{m}{c}\text{NH}_3 + \frac{p}{c}\text{H}_3\text{PO}_4$	$(k_{\text{POC}_{fast}}[\text{POC}_{fast}] + k_{\text{POC}_{slow}}[\text{POC}_{slow}])f_{\text{MnO}_2}$	$+4 + n/c - p/c$	+1
$(\text{CH}_2\text{O})(\text{NH}_3)_{\frac{m}{c}}(\text{H}_3\text{PO}_4)_{\frac{p}{c}} + 4\text{Fe}(\text{OH})_3 + 7\text{CO}_2 \leftrightarrow 8\text{HCO}_3^- + 4\text{Fe}^{2+} + \frac{m}{c}\text{NH}_3 + \frac{p}{c}\text{H}_3\text{PO}_4 + 3\text{H}_2\text{O}$	$(k_{\text{POC}_{fast}}[\text{POC}_{fast}] + k_{\text{POC}_{slow}}[\text{POC}_{slow}])f_{\text{Fe}(\text{OH})_3}$	$+8 + n/c - p/c$	+1
$(\text{CH}_2\text{O})(\text{NH}_3)_{\frac{m}{c}}(\text{H}_3\text{PO}_4)_{\frac{p}{c}} + 0.5\text{SO}_4 \leftrightarrow \text{HCO}_3^- + 0.5\text{H}_2\text{S} + \frac{m}{c}\text{NH}_3 + \frac{p}{c}\text{H}_3\text{PO}_4$	$(k_{\text{POC}_{fast}}[\text{POC}_{fast}] + k_{\text{POC}_{slow}}[\text{POC}_{slow}])f_{\Sigma\text{SO}_4}$	$+1 + n/c - p/c$	+1
$(\text{CH}_2\text{O})(\text{NH}_3)_{\frac{m}{c}}(\text{H}_3\text{PO}_4)_{\frac{p}{c}} \leftrightarrow 0.5\text{CO}_2 + 0.5\text{CH}_4 + \frac{m}{c}\text{NH}_3 + \frac{p}{c}\text{H}_3\text{PO}_4$	$(k_{\text{POC}_{fast}}[\text{POC}_{fast}] + k_{\text{POC}_{slow}}[\text{POC}_{slow}])f_{\text{CH}_4}$	$+n/c - p/c$	+0.5
Redox reactions			
$\text{Fe}^{2+} + 0.25\text{O}_2 + 2\text{HCO}_3^- + 0.5\text{H}_2\text{O} \leftrightarrow \text{Fe}(\text{OH})_3 + 2\text{CO}_2$	$k_{\text{Fe ox}}[\text{Fe}^{2+}][\text{O}_2]$	-2	0
$\text{Mn}^{2+} + 0.5\text{O}_2 + 2\text{HCO}_3^- \leftrightarrow \text{MnO}_2 + 2\text{CO}_2 + \text{H}_2\text{O}$	$k_{\text{Mn ox}}[\text{Mn}^{2+}][\text{O}_2]$	-2	0
$\text{H}_2\text{S} + 2\text{O}_2 + 2\text{HCO}_3^- \leftrightarrow \text{SO}_4^{2-} + 2\text{CO}_2 + 2\text{H}_2\text{O}$	$k_{\text{S ox}}[\Sigma\text{H}_2\text{S}][\text{O}_2]$	-2	0
$\text{NH}_3 + 2\text{O}_2 + \text{HCO}_3^- \leftrightarrow \text{NO}_3^- + \text{CO}_2 + 2\text{H}_2\text{O}$	$k_{\text{NH ox}}[\Sigma\text{NH}_4][\text{O}_2]$	-2	0
CaCO₃ dissolution and precipitation			
$\text{CaCO}_3 \leftrightarrow \text{Ca}^{2+} + \text{CO}_3^{2-}$	$[\text{Calcite}] \cdot k_{\text{diss. ca.}} \cdot (1 - \rho_{ca})^{\eta_{\text{diss. ca.}}}$	+2	+1
$\text{CaCO}_3 \leftrightarrow \text{Ca}^{2+} + \text{CO}_3^{2-}$	$[\text{Aragonite}] \cdot k_{\text{diss. ar.}} \cdot (1 - \rho_{ar})^{\eta_{\text{diss. ar.}}}$	+2	+1
$\text{Ca}^{2+} + \text{CO}_3^{2-} \leftrightarrow \text{CaCO}_3$	$k_{\text{prec. ca.}} \cdot (\rho_{ca} - 1)^{\eta_{\text{prec. ca.}}}$	-2	-1

2.2 Reactions

In RADI, biogeochemical reactions operate on solutes and solids throughout the entire sediment column, including the very top and bottom layers. $R(v_z)$ is the net rate at which v_z is being consumed (negative R) or produced (positive R) by these reactions. Biogeochemical reactions in RADI (Table 3) are grouped into three categories: (i) organic matter degradation, (ii) oxidation of reduced metabolites (organic matter degradation byproducts), and (iii) dissolution or precipitation of calcium carbonate minerals. RADI has been designed for early diagenesis in deep-sea sediments, so formation and re-oxidation of metal sulfide minerals are not considered.

2.2.1 Organic matter degradation

Organic carbon deposited on the seafloor originates mainly from primary production in the upper ocean or on land and, to a lesser extent, from the ocean interior via chemoautotrophy. Despite the differences in origin, detrital organic matter found in marine sediments typically has the same composition: ~60% proteins, ~20% lipids, ~20% carbohydrates, and a fraction of other compounds (Hedges et al., 2002; Burdige, 2007; Middelburg, 2019). Here, the stoichiometry of organic matter is represented by the coefficients c (for carbon), n (for nitrogen), and p (for phosphorus). By default, the $c:p$ ratio is set to 106:1 and the $n:p$ ratio set to 16:1, following the Redfield ratio that describes the average composition of phytoplankton biomass (Redfield, 1958), but these values can easily be adjusted. In RADI, c/p is denoted RC , n/p is denoted RN , and p/p is denoted RP , which is unity. Organic matter is also simplified here as an elementary carbohydrate (CH_2O). In reality, loss of H and O during biosynthesis of proteins, lipids, and polysaccharides occurs (Anderson, 1995; Hedges et al., 2002; Middelburg, 2019), which results in an effective molar ratio of O_2 consumed to C degraded of ~1.2 during aerobic respiration (Anderson and Sarmiento, 1994) instead of 1 as assumed here (Table 3).

Observations show that some organic compounds are preferentially degraded and become selectively depleted (Cowie and Hedges, 1994; Lee et al., 2000). As a result, the bulk reactivity of organic matter decreases with increasing age (Middelburg, 1989). Degradation of organic matter deposited at the seafloor typically follows a sequential utilization of available oxidants, O_2 , NO_3^- , MnO_2 , $Fe(OH)_3$, and SO_4^{2-} , followed by methanogenesis (Froelich et al., 1979; Berner, 1980; Arndt et al., 2013). All organic matter degradation reactions implemented in RADI are shown in Table 3.

To account for the decrease in organic matter degradation rate with sediment depth, we separate organic matter into fractions of different reactivity, and we assign a rate constant to each of the degradable fractions. Following Jørgensen (1978), Westrich and Berner (1984), and Soetaert et al. (1996b), three different classes of organic matter are considered: refractory, slow-decay, and fast-decay. The refractory organic matter class is not reactive during the timescales considered here. The fast- and slow-decay organic matter fractions each have a depth-dependent, oxidant-independent reactivity. The overall organic matter degradation rate decreases with depth because the quantity of organic matter and the relative proportions of fast and slow-decay materials decline with depth. Organic matter is degraded following the sequential utilization of available oxidants. The oxidant limitation is represented by a Michaelis-Menten-type (also termed ‘Monod’) function, in which each oxidant has

an associated half-saturation constant (K_{oxidant} in mol m⁻³) that symbolizes the oxidant concentration at which the process proceeds at half its maximal speed (Soetaert et al., 1996b). The presence of some oxidants may also inhibit other metabolic pathways; this is represented by an inhibition constant (K'_{oxidant} in mol m⁻³) that is specific to each oxidant. These limiting and inhibitory functions have been widely used (Boudreau, 1996b; Van Cappellen and Wang, 1996; Soetaert et al., 1996b; Couture et al., 2010) and they allow a single equation to be used for each component across the entire model depth range, and also permit some overlap between the different pathways, as observed in nature (Froelich et al., 1979). In RADI, the overall degradation of fast- or slow-decay organic carbon occurs at a rate:

$$R_{\text{POC}_{\text{fast or slow},z}} = f_{\text{oxidant},z} \cdot k_{\text{POC}_{\text{fast or slow}}} \cdot [\text{POC}_{\text{fast or slow}}]_z \quad (6)$$

where k_{POC} is the rate constant for the degradation of a given type of organic carbon (fast- or slow-decay) expressed in a⁻¹, $[\text{POC}_{\text{fast or slow}}]$ is the concentration of organic carbon (fast- or slow-decay) in sediments, and f_{Ox} is the sum of the fractions of organic carbon degraded by each oxidant (dimensionless, always equal to one), given by:

$$f_{\text{oxidant},z} = f_{\text{O}_2,z} + f_{\Sigma\text{NO}_3,z} + f_{\text{MnO}_2,z} + f_{\text{Fe(OH)}_3,z} + f_{\Sigma\text{SO}_4,z} + f_{\text{CH}_4,z}, \quad (7)$$

where

$$f_{\text{O}_2,z} = \frac{[\text{O}_2]_z}{K_{\text{O}_2} + [\text{O}_2]_z} \quad (8a),$$

$$f_{\Sigma\text{NO}_3,z} = \frac{[\Sigma\text{NO}_3]_z}{K_{\Sigma\text{NO}_3} + [\Sigma\text{NO}_3]_z} \frac{K_{\text{O}_2}'}{K_{\text{O}_2}' + [\text{O}_2]_z} \quad (8b),$$

$$f_{\text{MnO}_2,z} = \frac{[\text{MnO}_2]_z}{K_{\text{MnO}_2} + [\text{MnO}_2]_z} \frac{K_{\Sigma\text{NO}_3}'}{K_{\Sigma\text{NO}_3}' + [\Sigma\text{NO}_3]_z} \frac{K_{\text{O}_2}'}{K_{\text{O}_2}' + [\text{O}_2]_z} \quad (8c),$$

$$f_{\text{Fe(OH)}_3,z} = \frac{[\text{Fe(OH)}_3]_z}{K_{\text{Fe(OH)}_3} + [\text{Fe(OH)}_3]_z} \frac{K_{\text{MnO}_2}'}{K_{\text{MnO}_2}' + [\text{MnO}_2]_z} \frac{K_{\Sigma\text{NO}_3}'}{K_{\Sigma\text{NO}_3}' + [\Sigma\text{NO}_3]_z} \frac{K_{\text{O}_2}'}{K_{\text{O}_2}' + [\text{O}_2]_z} \quad (8d),$$

$$f_{\Sigma\text{SO}_4,z} = \frac{[\Sigma\text{SO}_4]_z}{K_{\Sigma\text{SO}_4} + [\Sigma\text{SO}_4]_z} \frac{K_{\text{Fe(OH)}_3}'}{K_{\text{Fe(OH)}_3}' + [\text{Fe(OH)}_3]_z} \frac{K_{\text{MnO}_2}'}{K_{\text{MnO}_2}' + [\text{MnO}_2]_z} \frac{K_{\Sigma\text{NO}_3}'}{K_{\Sigma\text{NO}_3}' + [\Sigma\text{NO}_3]_z} \frac{K_{\text{O}_2}'}{K_{\text{O}_2}' + [\text{O}_2]_z} \quad (8e),$$

$$f_{\text{CH}_4,z} = \frac{K_{\Sigma\text{SO}_4}'}{K_{\Sigma\text{SO}_4}' + [\Sigma\text{SO}_4]_z} \frac{K_{\text{Fe(OH)}_3}'}{K_{\text{Fe(OH)}_3}' + [\text{Fe(OH)}_3]_z} \frac{K_{\text{MnO}_2}'}{K_{\text{MnO}_2}' + [\text{MnO}_2]_z} \frac{K_{\Sigma\text{NO}_3}'}{K_{\Sigma\text{NO}_3}' + [\Sigma\text{NO}_3]_z} \frac{K_{\text{O}_2}'}{K_{\text{O}_2}' + [\text{O}_2]_z} \quad (8f).$$

Half-saturation and inhibition constants for each oxidant used in RADI are given in Table 4. The degradation rate constant of organic carbon, $k_{\text{POC}_{\text{fast or slow}}}$, is computed as a function of the flux of organic carbon reaching the seafloor and is sediment-depth dependent (Archer et al., 2002):

$$k_{\text{POC}_{\text{fast}}} = (1.5 \times 10^{-1}) (F_{\text{POC}} \cdot 10^2)^{0.85} \quad (9a),$$

$$k_{\text{POC}_{\text{slow}}} = (1.3 \times 10^{-4}) (F_{\text{POC}} \cdot 10^2)^{0.85} \quad (9b),$$

where F_{POC} is the total flux of organic carbon reaching the seafloor (i.e., fast, slow, and refractory), in mol m⁻² a⁻¹. The numbers 1.3×10^{-4} and 1.5×10^{-1} have been tuned to best fit observations of both a Southern Ocean and North Atlantic Station, see the *Model evaluation* section 3.

2.2.2 Oxidation of organic-matter degradation by-products

Organic matter degradation reactions primarily change oxidants (e.g., O₂, NO₃⁻, MnO₂, Fe(OH)₃, SO₄²⁻) into their reduced forms (e.g., H₂O, N₂, Mn²⁺, Fe²⁺, H₂S; Table 3). If oxygen is introduced into the system, or if the reduced metabolites diffuse upwards in oxic porewaters, then these reduced byproducts are converted back into their oxidized form and the energy contained in them becomes available to the microbial community, though these energetics are not considered in RADI.

Here, four redox reactions involving organic-matter degradation byproducts are implemented (Table 3): oxidation of each of Fe²⁺, Mn²⁺, ΣH₂S and ΣNH₃. These four reactions consume porewater total alkalinity (TAlk) but do not alter porewater ΣCO₂ (Table 3), thus locally acidifying porewaters. Here, we use the TAlk definition of Dickson (1981), in which TAlk is defined as “the number of moles of hydrogen ion equivalent to the excess of proton acceptors (bases formed from weak acids with a dissociation constant $K \leq 10^{-4.5}$ and zero ionic strength) over proton donors (acids with $K > 10^{-4.5}$) in one kilogram of sample”. This scheme should be sufficient for all open-ocean applications but may not be suitable for coastal and anoxic environments with extensive metal-sulfide mineral turnover, which require a more complete set of redox reactions such as that from the CANDI model of Boudreau (1996b). Additional components and reactions can easily be implemented in future versions (see *Future Developments* section 5). The rate constants for these four redox reactions are taken from Boudreau (1996b) and reported in Table 4.

Table 4. Suggested values for model parameters.

Parameter	Model notation	Value	Unit	Source
K_{O_2} / K_{O_2}'	KdO2/KdO2i	3 / 10	μM	Soetaert et al. (1996)
$K_{\Sigma NO_3} / K_{\Sigma NO_3}'$	KdtNO3/KdtNO3i	30 / 5	μM	Soetaert et al. (1996)
$K_{MnO_2} = K_{MnO_2}'$	KpMnO2/KpMnO2i	42.4	mM	Van Cappellen and Wang (1996) ¹
$K_{Fe(OH)_3} = K_{Fe(OH)_3}'$	KpFeOH3/KpFeOH3i	265	mM	Van Cappellen and Wang (1996) ¹
$K_{\Sigma SO_4} = K_{\Sigma SO_4}'$	KdtSO4/KdtSO4i	1.6	mM	Van Cappellen and Wang (1996) ¹
$k_{Fe\ ox}$	kFeox	10 ⁶	mM ⁻¹ a ⁻¹	Boudreau (1996b) ²
$k_{Mn\ ox}$	kMnox	10 ⁶	mM ⁻¹ a ⁻¹	Boudreau (1996b) ²
$k_{S\ ox}$	kSox	3 x 10 ⁵	mM ⁻¹ a ⁻¹	Boudreau (1996b) ²
$k_{NH\ ox}$	kNHox	10 ⁴	mM ⁻¹ a ⁻¹	Boudreau (1996b) ²
β	phiBeta	33	m ⁻¹	Tuned
λ_b	lambda_b	0.08	m	Archer et al. (2002)
λ_i	lambda_i	0.08	m	Archer et al. (2002)

¹Assuming a solid density of 2.65 g cm⁻³; ²Values for the “deep-sea”

2.2.3 CaCO₃ dissolution and precipitation

310 RADI includes two CaCO₃ polymorphs: low-Mg calcite and aragonite, but more could be added in future versions, e.g., high-Mg calcite and/or vaterite. Calcite and aragonite both have different dissolution kinetics, in which their dissolution rates increase as the undersaturation state of seawater with respect to calcite ($1 - \Omega_{ca,z}$) or aragonite ($1 - \Omega_{ar,z}$) increases (Keir, 1980; Walter and Morse, 1985; Sulpis et al., 2017; Dong et al., 2019; Naviaux et al., 2019b). Here, Ω_z is the sediment-depth-dependent saturation state of seawater with respect to calcite or aragonite, computed as $[Ca^{2+}]_z \cdot [CO_3^{2-}]_z / K_{sp}^*$, where K_{sp}^* is the stoichiometric solubility constant of calcite or aragonite at in situ temperature, pressure, and salinity, as given in Mucci (1983) and Millero (1995). At each time step, Ω_z is computed using porewater $[Ca^{2+}]_z$ and $[CO_3^{2-}]_z$ from the previous time step, the latter being calculated as a function of TAlk and the proton concentration $[H^+]$. At each model time step, the total hydrogen ion concentration $[H^+]$ is computed from TAlk and $\sum CO_2$ using a single Newton-Raphson iteration from the previous time step (Humphreys et al., 2021):

$$320 [H^+]_t = [H^+]_{t-1} - \frac{[TAlk]([H^+]_{t-1}, [\sum CO_2]) - [TAlk]}{d[TAlk]([H^+]_{t-1}, [\sum CO_2]) / d[H^+]_{t-1}} \quad (10),$$

where $[H^+]_t$ is the new $[H^+]$ value and $[H^+]_{t-1}$ is the $[H^+]$ from the previous time step. TAlk($[H^+]_{t-1}$, $\sum CO_2$) is the total alkalinity computed from user-specified total dissolved silicate, $[\sum PO_4]$ and total borate calculated from salinity (Uppström, 1974), plus equilibrium constants for silicic acid (Sillén et al., 1964) and phosphoric acid (Yao and Millero, 1995). Its derivative is computed following the approach of CO2SYS, see Humphreys et al. (2021). The carbonate ion concentration is then computed as:

$$325 [CO_3^{2-}] = \frac{[\sum CO_2] \times K_1^* \times K_2^*}{K_1^* \times K_2^* + K_1^* \times [H^+]_t + [H^+]_t^2} \quad (11),$$

where K_1^* and K_2^* are the first and second dissociation constants for carbonic acid, respectively, taken from Lueker et al. (2000).

The dissolution rates (R_{diss} , in mol m⁻³ a⁻¹) of calcite (solid blue line in Fig. 2a) and of aragonite (solid red line in Fig. 2a) as a function of $(1 - \Omega_{ca})$ are empirically defined as:


$$330 R_{diss. ca., z} = [Calcite] \cdot k_{diss. ca.} \cdot (1 - \Omega_{ca})^{\eta_{diss. ca.}} \quad (12),$$

$$R_{diss. ar., z} = [Aragonite] \cdot k_{diss. ar.} \cdot (1 - \Omega_{ar})^{\eta_{diss. ar.}} \quad (13).$$

In these expressions, the dissolution rate constant (k_{diss} , in a⁻¹) and the reaction order (η_{diss} , unitless) are mineral-specific. The dissolution rate constants implicitly account for each mineral's specific surface area. Similar formulations have previously been implemented to describe calcite dissolution rates (e.g. Archer et al. 2002) but, in most cases, with a high reaction order and with a tuned rate constant independent of solution chemistry (Fig. 2). Such discretizations are convenient but lack a mechanistic description of the controls on calcite dissolution in seawater (Adkins et al., 2021).

The latest advances using isotope-labelling approaches to study carbonate dissolution kinetics show abrupt changes in dissolution mechanism depending on solution saturation state with either calcite or aragonite (Subhas et al., 2017; Dong et al., 2019; Naviaux et al., 2019a; Naviaux et al., 2019b). Close to equilibrium, dissolution occurs primarily at sites on the crystal

surfaces that are most exposed to the solution, e.g., steps and kinks. Further away from equilibrium, dissolution etch pits are activated at surface sites associated with defects and impurity atoms. Far away from equilibrium, there is enough free energy for dissolution etch pits to occur anywhere on the mineral surface, without the aid of crystal defects (Adkins et al., 2021). However, at temperatures most relevant to the deep oceans, ~ 5 °C or less, the defect-assisted dissolution mechanism is skipped (Naviaux et al., 2019b) and only the step-edge retreat (close to equilibrium) and homogeneous etch-pit formation (far away from equilibrium) dissolution regimes remain (Naviaux et al., 2019b) (Fig. 2). For both aragonite and calcite, while homogeneous etch-pit formation is indeed associated with a high-order dependency on the solution saturation state, step-edge retreat dissolution rates dominating near equilibrium show very little dependence on seawater saturation (Dong et al., 2019; Naviaux et al., 2019a). This could have significant consequences for the predicted carbonate dissolution rate near equilibrium: saturation-state independent step-edge retreat dissolution will always be predicted to be faster close to equilibrium than dissolution associated with a high reaction order because a high reaction order forces the dissolution rate to converge to zero as the solution gets closer to equilibrium (Fig. 2).



355 Figure 2. (a) Dissolution rate of calcite as computed using Eq. (12) and $[Calcite] = 10^4 \text{ mol m}^{-3}$ (solid blue line) and dissolution rate
of aragonite as computed using Eq. (13) and $[Aragonite] = 10^4 \text{ mol m}^{-3}$ (solid red line). Note that for each dissolution rate profile,
two different rate constants (k_{diss}) and reaction orders (η_{diss}) are used, depending on the seawater saturation state; each accounting
for a separate dissolution mechanism, i.e., step-edge retreat or homogeneous edge pit formation. The dashed black line stands for a
“traditional” dissolution rate profile obtained using $[Calcite] = 10^4 \text{ mol m}^{-3}$, a single dissolution rate constant for the entire $(1 - \Omega_{ca})$
360 range $k_{diss} = 100 \text{ \% day}^{-1}$, and a reaction order η_{diss} of 4.5. (b) Precipitation rate of calcite as computed from Eq. (14). Note that
dissolution rates are normalized here per total solid sediment volume, not per CaCO₃ surface area as in traditional kinetics
studies.

Naviaux et al. (2019a) derived reaction orders for two separate regions of the $(1 - \Omega_{ca})$ spectrum: the Ω_{ca} threshold
365 value dividing these two regions was $\Omega_{ca, critical} \approx 0.8$. Here, based on the results of Naviaux et al. (2019a), we set $\eta_{diss, ca} = 0.11$
for $0.828 < \Omega_{ca} < 1$ and $\eta_{diss, ca} = 4.7$ for $\Omega_{ca} \leq 0.828$. The $\Omega_{ca, critical}$ value used here is slightly higher than the ~ 0.8 value given
in Naviaux et al. (2019a) in order to have a smooth transition between defect-assisted and homogeneous dissolution. For
aragonite, based on the results of Dong et al. (2019), we set $\eta_{diss, ar} = 0.13$ for $0.835 < \Omega_{ar} < 1$ and $\eta_{diss, ar} = 1.46$ for $\Omega_{ar} \leq 0.835$.

The rate constants are tuned to best fit the observations in the two stations presented in the *Model Evaluation* sections 3. We use $k_{diss. ca.} = 6.3 \times 10^{-3} \text{ a}^{-1}$ for $0.828 < \Omega_{ca} < 1$, $k_{diss. ca.} = 20 \text{ a}^{-1}$ for $\Omega_{ca} \leq 0.828$, $k_{diss. ar.} = 3.8 \times 10^{-3} \text{ a}^{-1}$ for $0.835 < \Omega_{ar} < 1$, and $k_{diss. ar.} = 4.2 \times 10^{-2} \text{ a}^{-1}$ for $\Omega_{ar} \leq 0.835$. Both calcite and aragonite dissolution rate constants are lower than the values reported in the original publications. We suspect that i) the reactive surface area of grains in sediments is much smaller than their specific surface area measured using adsorption isotherms via the BET method and ii) unaccounted dissolution inhibitors are present in sediments, such as dissolved organic carbon (Naviaux et al., 2019a). A comparison of the steady-state $[\text{CO}_3^{2-}]$ and *Calcite* porewater profiles predicted by RADI using the tuned rate constants implemented in RADIv1 and the original rate constants is shown in Fig. S1.

Calcite precipitation is also included in the model and its rate (solid blue line in Fig. 2b) is parameterized with the following function:

$$R_{prec. ca., z} = k_{prec. ca.} \cdot (\Omega_{ca} - 1)^{\eta_{prec. ca.}} \quad (14),$$

where $k_{prec. ca.}$ is the precipitation rate constant set to $0.4 \text{ mol m}^{-3} \text{ a}^{-1}$ and η is equal to 1.76. The precipitation reaction order is taken from Zuddas and Mucci (1998), corrected for a seawater-like ionic strength of 0.7 mol kg^{-1} . The precipitation / dissolution rate continuum implemented in RADI (see Fig. 2) is very different from what a classic model with only calcite dissolution following high-reaction order kinetics would display. For comparison, the dissolution rate of calcite using a dissolution rate constant k_{diss} of 100 % per day and a reaction order η of 4.5, as implemented in most diagenetic models, including Archer (1991), is shown in Fig. 2a. Mechanistic interpretations of the “kinks” in the dissolution rate profiles and of a non-zero dissolution rate near equilibrium still require more research but the implications of these features for our understanding of marine CaCO_3 cycles can be explored with the present model.

2.3 Advection

The solid burial velocity at the sediment-water interface, w_0 in m a^{-1} , is given by:

$$w_0 = \frac{\sum F_v \cdot M_v}{\rho_v} / \varphi_{s,0} \quad (15),$$

where F_v is the flux of a solid species at the sediment-water interface ($\text{mol m}^{-2} \text{ a}^{-1}$), M_v is the molar mass of that solid (g mol^{-1}), and ρ_v is its solid density (g m^{-3}). The solid and porewater burial velocity at greater depth are assumed to be equal, and are computed as:

$$w_\infty = u_\infty = w_0 \varphi_{s,0} / \varphi_{s,\infty} \quad (16).$$

Thus, the porewater burial velocity, u , at all depths is:

$$u_z = u_\infty \varphi_\infty / \varphi_z \quad (17)$$

and the solid burial velocity, w :

$$w_z = w_\infty \varphi_{s,\infty} / \varphi_{s,z} \quad (18).$$

Depth profiles of u and w are shown in Fig. 1, computed from the solid fluxes at station 7 of cruise NBP98-2 (Sayles et al., 2001), seen in *Model Evaluation* Section 3.2. In Fig. 1, the sharp porosity decline in the top centimeters of the sediments causes

the solid fraction at ~5-cm depth to be roughly 50% higher than just below the interface. This leads to a solid burial velocity decrease of about the same magnitude (Fig. 1).

Advection is implemented following Boudreau (1996b), where the advection rate (A_z , in $\text{mol m}^{-3} \text{a}^{-1}$) for solutes is given by:

$$405 \quad A_z(v) = - \left(u_z - \frac{d_z(v)}{\varphi_z} \cdot \frac{d\varphi_z}{dz} - d^\circ(v) \cdot \frac{d\left(\frac{1}{\theta_z^2}\right)}{dz} \right) \cdot \frac{v_{(z+dz)} - v_{(z-dz)}}{2dz} \quad (19),$$

where d_z is the effective diffusion coefficient for a given solute at a given depth (in $\text{m}^2 \text{a}^{-1}$), d° is the “free-solution” molecular diffusion coefficient for a given solute (in $\text{m}^2 \text{a}^{-1}$), and θ_z is the depth-dependent tortuosity (unitless) defined in the *Diffusion section 2.4*. For solids, a more sophisticated weighted-difference scheme is employed (Fiadeiro and Veronis, 1977; Boudreau, 1996b):

$$410 \quad A_z(v) = - \left(w_z - \frac{db_z}{dz} - \frac{b_z}{\varphi_{s,z}} \cdot \frac{d\varphi_{s,z}}{dz} \right) \cdot \frac{(1-\sigma_z)v_{(z+dz)} + 2\sigma_z v_z - (1+\sigma_z)v_{(z-dz)}}{2dz} \quad (20),$$

where b_z is the depth-dependent bioturbation coefficient ($\text{m}^2 \text{a}^{-1}$) and

$$\sigma_z(v) = 1/\tanh(Pe_{h,z}) - 1/Pe_{h,z} \quad (21),$$

where

$$Pe_{h,z} = w_z \cdot dz/2b_z \quad (22).$$

415 The parameter Pe_h is half of the ‘cell Peclet number’, which expresses the influence of advection relative to bioturbation across a distance separating two points of the grid, centered at the depth z . If bioturbation dominates ($Pe_h \ll 1$), e.g., near the sediment-water interface, σ_z tends toward zero and a centered-difference discretization is implemented. If advection dominates ($Pe_h \gg 1$), e.g., deeper in sediments, σ_z tends toward unity and backward-difference discretization prevails, see Eq. (20). This differencing scheme, originally developed by Fiadeiro and Veronis (1977), maintains stability in the entire sediment column
420 (Boudreau, 1996b).

2.4 Diffusion

The diffusion flux of any species depends on its effective diffusion coefficient, $d_z(v)$, which varies with depth within the sediment.

425 For each solute, free-solution diffusion coefficients, denoted $d_z^\circ(v)$, were computed at in-situ temperatures (Li and Gregory, 1974; Boudreau, 1997; Schulz, 2006). For solute variables representing several individual species (e.g., ΣPO_4 , ΣCO_2), the diffusion coefficient of the dominant species was considered. Given the high proportion of HCO_3^- relative to CO_3^{2-} and $\text{CO}_2(\text{aq})$ in seawater and porewaters (see Fig. S2), the diffusion coefficient of HCO_3^- was adopted for both TALK and ΣCO_2 . However, this approach may not be suited for sedimentary environments in which pH is lower than 7, because a greater proportion of dissolved inorganic species would then be under the form of carbonic acid, or $\text{CO}_2(\text{aq})$, which has a higher

430 diffusion coefficient than HCO_3^- (Fig. S2). Free-solution diffusion coefficients, their temperature dependencies and their sources are reported in Table 5. The diffusion of solutes in the porewaters is slower than in an equivalent volume of water as a result of the physical barriers caused by the presence of solid grains in a sediment. To correct for this effect, we follow Boudreau (1996b) and compute the effective diffusion coefficient for a given solute as:

$$d_z(v) = d^\circ(v)/(\theta_z^2) \quad (23),$$

435 where so-called tortuosity (θ) is defined as (Boudreau, 1996a):

$$\theta_z = \sqrt{1 - 2 \ln(\phi_z)} \quad (24).$$

Table 5. Temperature-dependent molecular diffusion coefficients ($\text{m}^2 \text{a}^{-1}$).

Diffusion coefficient	Value	Source
$d_z^\circ(\text{TAlk})$	$0.015179 + 0.000795 \times T_w$	(Boudreau, 1997; Schulz, 2006) ¹
$d_z^\circ(\Sigma\text{CO}_2)$	$0.015179 + 0.000795 \times T_w$	(Boudreau, 1997; Schulz, 2006) ¹
$d_z^\circ(\text{Ca}^{2+})$	$0.011771 + 0.000529 \times T_w$	(Li and Gregory, 1974)
$d_z^\circ(\text{O}_2)$	$0.031558 + 0.001428 \times T_w$	(Boudreau, 1997; Schulz, 2006)
$d_z^\circ(\Sigma\text{NO}_3)$	$0.030863 + 0.001153 \times T_w$	(Li and Gregory, 1974) ²
$d_z^\circ(\Sigma\text{SO}_4)$	$0.015779 + 0.000712 \times T_w$	(Li and Gregory, 1974) ³
$d_z^\circ(\Sigma\text{PO}_4)$	$0.009783 + 0.000513 \times T_w$	(Boudreau, 1997; Schulz, 2006) ⁴
$d_z^\circ(\Sigma\text{NH}_4)$	$0.030926 + 0.001225 \times T_w$	(Li and Gregory, 1974) ⁵
$d_z^\circ(\Sigma\text{H}_2\text{S})$	$0.028938 + 0.001314 \times T_w$	(Boudreau, 1997; Schulz, 2006)
$d_z^\circ(\text{Fe}^{2+})$	$0.001076 + 0.000466 \times T_w$	(Li and Gregory, 1974)
$d_z^\circ(\text{Mn}^{2+})$	$0.009625 + 0.000481 \times T_w$	(Li and Gregory, 1974)

¹value for HCO_3^- ion, ²value for NO_3^- ion, ³value for SO_4^{2-} ion, ⁴value for HPO_4^{2-} ion, ⁵value for NH_4^+ ion

For each solid, effective diffusion occurs through the mixing action of burrowing microorganisms, quantified using a bioturbation coefficient that decreases with depth. Archer et al. (2002) used a dataset including 53 sediment sites ranging in depth from 47 to 5668 m to derive an optimal bioturbation rate profile, in which the rate of bioturbation increases with increasing flux of total organic carbon reaching the seafloor (F_{POC}) and attenuates in low-oxygen conditions. This pattern was also observed by Smith et al. (1997) and Smith and Rabouille (2002). As in Archer et al. (2002), we couple both bioturbation and irrigation to the incoming carbon deposition flux (Fig. 3) rather than water depth or sediment accumulation rate (Boudreau, 1994; Middelburg et al., 1997; Soetaert et al., 1996c), although all these quantities are related to each other. From an ecological perspective, more carbon to the seafloor represents more food available to benthic communities, hence more biological transport. Linking bioturbation activity to carbon deposition flux also allows for a direct coupling with Earth system models

simulating carbon sinking fluxes in the ocean. Following Archer et al. (2002), we express the surficial bioturbation mixing rate (b_0 , in $\text{m}^2 \text{a}^{-1}$) as:

$$b_0 = (2.32 \cdot 10^{-6})(F_{\text{POC}} \cdot 10^2)^{0.85} \quad (25),$$

465 where F_{POC} is expressed in $\text{mol m}^{-2} \text{a}^{-1}$. The bioturbation mixing rate at all depths (b_z , in $\text{m}^2 \text{a}^{-1}$) is:

$$b_z = b_0 e^{-\left(\frac{z}{\lambda_b}\right)^2} \frac{[\text{O}_2]_w}{[\text{O}_2]_w + 0.02} \quad (26),$$

where the characteristic depth $\lambda_b = 8 \text{ cm}$, following Archer et al. (2002), and $[\text{O}_2]_w$ is the oxygen concentration in the bottom waters. This depth-dependent bioturbation mixing rate is common to all solids and its depth distribution is shown in Fig. 3 as a function of in situ $[\text{O}_2]_w$ and F_{POC} . The effective diffusion coefficient for solids is then set as:

470 $d_z(v) = b_z \quad (27).$

(Bio)diffusion is implemented in RADI following the centered difference discretization scheme from Boudreau (1996b). At sediment depth z , where $0 < z < Z$, for both solutes and solids:

$$D_z(v) = d_z(v) \cdot (v_{(z-dz)} - 2v_z + v_{(z+dz)}) / (dz)^2 \quad (28),$$

where $d_z(v)$ is the relevant effective diffusion coefficient.

475




Figure 3. Bioturbation mixing rate b_z and irrigation coefficients α_z as a function of sediment depth z , organic carbon deposition flux F_{POC} , and dissolved oxygen concentration in the bottom waters $[\text{O}_2]_w$.

The mixing of solutes caused by burrow flushing or ventilation occurs through an ensemble of processes collectively termed *irrigation*. Macroscopic burrows are often present in the seafloor sediment, with a complex three-dimensional structure and filled with oxygenated water that is ventilated for aerobic respiration. In a one-dimensional framework, this causes apparent internal sources or sinks of porewater solutes at particular depths (Boudreau, 1984; Emerson et al., 1984; Aller, 2001).

485 Mathematically, this is parameterized as a non-local exchange function, i.e., a first-order kinetic reaction:

$$I_{t,z}(v) = \alpha_z(v_w - v_z) \quad (29),$$

where α_z is an irrigation coefficient common to all solutes expressed in a^{-1} . Following Archer et al. (2002), who used a dataset of 53 sediment sites comprised of microelectrode oxygen profiles and chamber oxygen fluxes across the sediment-water interface to derive an irrigation-rate profile, we express the surficial irrigation coefficient as a function of the organic carbon

490 deposition flux and the oxygen concentration of the overlaying waters:

$$\alpha_0 = 11 \left(\frac{\tan^{-1} \left(\frac{5F_{\text{POC}} \cdot 10^2 - 400}{400} \right)}{\pi} + 0.5 \right) - 0.9 + \frac{20[\text{O}_2]_w}{[\text{O}_2]_w + 0.01} \cdot \frac{F_{\text{POC}} \cdot 10^2}{F_{\text{POC}} \cdot 10^2 + 30} \cdot e^{\frac{-[\text{O}_2]_w}{0.01}} \quad (30)$$

and the irrigation coefficient at all depths as:

$$\alpha_z = \alpha_0 e^{-(z/\lambda_i)^2} \quad (31),$$

where the characteristic depth λ_i is 5 cm (Archer et al., 2002). The depth-distribution of the irrigation coefficient is shown in

495 Fig. 3 as a function of in-situ $[\text{O}_2]_w$ and F_{POC} .

2.6 Boundary conditions

Modeling of advection and diffusion processes requires appropriate boundary conditions in the layers above and below ($z - dz$ and $z + dz$, respectively). Effective values of each variable immediately adjacent to the modelled depth domain are calculated following Boudreau (1996b) and used to compute the effects of advection and diffusion in the top and bottom

500 layers using the same equations as within the sediment itself.

At the sediment-water interface, RADI enables prescribed solid fluxes and a diffusive boundary layer control for solutes. Following Boudreau (1996b), we calculate advection and diffusion at $z = 0$ for solutes and solids as:

$$v_{(-dz)} = v_{dz} + \frac{2\theta^2 dz}{\delta} (v_w - v_0) \quad (32)$$

and

505
$$v_{(-dz)} = v_{dz} + \frac{2dz}{b_0} \left(\frac{F_v}{\varphi_{s,oz}} - w_0 v_0 \right) \quad (33),$$

respectively. Here, θ is the tortuosity, δ is the boundary layer thickness (expressed in m, see Fig. 1), and v_w is the solute concentration above the diffusive boundary layer, i.e., in the bottom waters. At the sediment depth $z = Z$, $v_{(Z+dz)}$ falls outside

the depth range of the model. The bottom boundary condition demands that concentration gradient disappear, which can be translated by the following for both solutes and solids:

$$510 \quad v_{(z+dz)} = v_{(z-dz)} \quad (34).$$

This ‘no-flux’ bottom boundary condition should be appropriate here because we set Z so that all action occurs at shallower depth. However, if anaerobic methane oxidation or subsurface weathering are included in future versions, a ‘constant’ flux boundary conditions might need to be included.

2.7 Julia and MATLAB/GNU Octave implementations

515 We have implemented RADI both in Julia (Humphreys and Sulpis, 2021) and in MATLAB/GNU Octave (Sulpis et al., 2021). Both implementations use similar nomenclature and provide identical results. Documentation for both is available from <https://radi-model.github.io>. The Julia implementation is available from <https://github.com/RADI-model/Radi.jl> and the MATLAB/GNU Octave implementation is available from <https://github.com/RADI-model/Radi.m>.

520 Julia (<https://julialang.org>) is a high-level, high-performance, and cross-platform programming language that is free and open source (Bezanson et al., 2017). Its high-performance stems primarily from just-in-time (JIT) compilation of code before execution, which has been built-in since its origin. RADI uses Julia’s multiple-dispatch paradigm, a core feature of the language, which improves the readability of the code and reduces the scope for errors. Specifically, each modelled component of the sediment column is either a porewater solute or a solid. These components are initialized in the model as variables either of `Solute` or `Solid` type. Advection and diffusion are governed by different equations for porewater solutes than for solids
525 but the same top-level functions (`advect!` and `diffuse!`) can be used within RADI to calculate the effects of these processes for both component types; the multiple-dispatch paradigm ensures that the correct equations are automatically used on the basis of the type of the input variable. While the model has been designed to solve a single profile at a time, Julia’s support for parallelized computation (across multiple processors) would also support efficient computations across a series or grid of vertical profiles.

530 As of version R2015b, MATLAB also features JIT compilation with a corresponding execution speed-up. However, MATLAB is an expensive, proprietary software, which limits how widely it can be used. The MATLAB implementation also runs in GNU Octave (<https://www.gnu.org/software/octave/>), which is a free and open-source clone of MATLAB. However, GNU Octave executes more slowly than MATLAB for a variety of reasons, including a lack of JIT compilation.

535 For a model that necessarily includes long simulations with relatively short time-steps, computational speed is an important consideration. Our testing indicates that the Julia implementation runs ~3 times faster than the MATLAB (R2020a) implementation and ~70 times faster than the GNU-Octave implementation.

540 Simultaneously developing the model in two languages allowed us to quickly identify and remedy bugs and typographical errors in both implementations. Each was coded independently, with equations and parameterizations written-out from the original sources, thus avoiding code copy-and-paste errors. Frequent comparisons were made throughout this process to ensure that the results were consistent. For a typographical error to survive to the final models would therefore

require an identical mistake to have been made independently in both implementations. The risk of such errors is thus substantially reduced by our dual-language approach. Where errors were identified, in some cases they were subtle enough that they may otherwise not have been noticed, while still causing meaningful errors in final model results.

3 Model evaluation

545 To evaluate the performance of RADI, we used in-situ data obtained at three different locations and compared our predictions to the measured porewater and sediment solid phase composition profiles. We used these comparisons to tune the CaCO_3 dissolution and POC degradation rate constants; all other parameters were assigned a priori using values from the literature. Thus, we did not aim to reproduce observations as well as possible by tuning a wide selection of parameters. Instead, we evaluated whether a generic approach using measured deposition fluxes and bottom-water conditions could explain
550 observations while tuning only the inorganic and organic reactivity constants.

3.1 North-western Atlantic Ocean

First, RADI was compared to the porewater and sediment composition measurements of station #9 described in Hales et al. (1994), located in the North-western Atlantic Ocean ($24.33^\circ\text{N} / 70.35^\circ\text{W}$) at a 5210-m depth. The bottom-water TAlk and ΣCO_2 were $2342 \mu\text{mol kg}^{-1}$ and $2186 \mu\text{mol kg}^{-1}$, respectively, bottom-water in situ temperature was 2.2°C , salinity was
555 34.9 , and oxygen concentration was $266.6 \mu\text{mol kg}^{-1}$ (Hales et al., 1994). The computed bottom-water saturation state with respect to calcite was 0.88 . The only CaCO_3 polymorph reaching the seafloor was assumed to be calcite. The calcite flux to the seafloor was set to $0.20 \text{ mol m}^{-2} \text{ a}^{-1}$ ($20.02 \text{ g m}^{-2} \text{ a}^{-1}$) and the POC flux to $0.18 \text{ mol m}^{-2} \text{ a}^{-1}$, which correspond to the low end of fluxes measured by sediment traps on the continental slope (Hales et al., 1994). The clay flux was set to a value of $26 \text{ g m}^{-2} \text{ a}^{-1}$ to fit the calcite sediment surface concentration measured by Hales et al. (1994). The porosity at the sediment-water
560 interface was set to that measured by Sayles et al. (2001) in the Southern Pacific Ocean station, see Fig. 1. Following the diffusive boundary layer distribution from Sulpis et al. (2018), δ at the station location was set to $938 \mu\text{m}$. This value represents an annual-mean estimate derived using a number of assumptions, e.g., considering the sediment-water interface to be a horizontal surface and neglecting sediment roughness. A complete description of the environmental parameters for this North-Atlantic station, along with their sources, is available in Table S1.

565 RADI was run using the environmental conditions described above and the steady-state concentration profiles of O_2 , ΣNO_3 , calcite, and POC were compared with observations. Complete methods for solute and solids measurements are described in Hales et al. (1994). Briefly, porewater O_2 concentration was measured both in-situ using microelectrodes and on-board (along with ΣNO_3) from the retrieved box core (Hales et al., 1994). The steady-state calcite, TAlk and ΣCO_2 profiles were compared to those obtained from a RADI simulation with “traditional”, 4.5-order calcite dissolution kinetics, see Fig. 2,
570 all other variables being unchanged.




Figure 4. Comparison of RADI with measurements from the North-western Atlantic station #9 described in (Hales et al., 1994). The lower panels represent (left) the computed CO_3^{2-} concentrations in porewaters (solid black line) and at equilibrium with respect to calcite (dashed blue line), and (center) the fractions of organic matter degraded by a given oxidant.

575

RADI predicts a porewater O_2 concentration decreasing from the bottom-water value to zero at ~20 cm-depth (Fig. 4). In the top 2 cm, the RADI porewater O_2 predictions near the surface are in good agreement with the in-situ microelectrode measurements. The RADI-predicted $[\text{O}_2]$ is lower than that measured on-board, but $[\Sigma\text{NO}_3]$ is well reproduced by RADI. RADI predicts that organic matter respiration is mainly aerobic (see Table 3a) until about 20 cm-depth. Between 20 and 35-cm depth, ΣNO_3 is the preferred oxidant for organic matter degradation (see Table 3b), which leads to a decrease in porewater

580

[ΣNO_3] in both RADI predictions and observations, followed by ΣSO_4 deeper than 35-cm depth. The calcite profile is relatively well reproduced by RADI in the top 20 cm but the measured calcite concentration drop below 20-cm depth is not well reproduced. When “traditional” 4.5-order calcite dissolution kinetics are implemented, calcite concentrations are underestimated by about a third in the top 20 cm. The observed lower calcite concentrations below 20-cm depth may be attributed to a lower calcite accumulation rate to the seafloor in the past, whereas the model considers accumulation of solids to be unchanged through time. Calcite concentration predicted by RADI does increase again below 25-cm depth due to porewater supersaturation (Ω_{ca} at 40 cm-depth is about 1.05), but this increase is too small to be noticed on the figure.


3.2 Southern Pacific Ocean

RADI was also compared with data collected at the station #7 mooring #3 described in Sayles et al. (2001), located in the Southern Pacific Ocean (60.15°S / 170.11°W), where the seafloor lies at the 3860-m depth. This dataset (http://usjgofs.whoi.edu/jg/dir/jgofs/southern/nbp98_2/) constrains the sedimentary system well: sediment-trap CaCO_3 and POC fluxes, CaCO_3 , and POC sediment-composition, sediment-porosity, and porewater solute depth profiles are all available from the same cruise and location.

The bottom-water chemical composition was taken from the GLODAPv2 $1 \times 1^\circ$ climatologies (Lauvset et al., 2016), linearly interpolated over depth, latitude, and longitude to match the station location and seafloor depth. The bottom-water in-situ temperature was 0.84°C, salinity was 34.696, oxygen concentration was 215.7 $\mu\text{mol kg}^{-1}$, and calculated saturation state with respect to calcite was 0.85. Solid fluxes at this station were measured by Sayles et al. (2001) using sediment traps collecting sinking particles between the months of November and December 1997. Their deepest sediment trap available was at a depth of 3257 m, i.e. 600 meters above the seafloor, which we assume to be representative of sinking fluxes to the seafloor, although the loss of material after collection usually causes sediment traps to underestimate the real sinking fluxes (Buesseler et al., 2007). The only CaCO_3 polymorph reaching the seafloor was assumed to be calcite and its flux was set to 0.25 $\text{mol m}^{-2} \text{a}^{-1}$ (25.02 $\text{g m}^{-2} \text{a}^{-1}$), rather than using the sediment-trap value, in order to fit the calcite sediment surface concentration measured by Sayles et al. (2001). This CaCO_3 flux to the seafloor is slightly higher than the measured CaCO_3 flux at 600 meters above the seafloor in mid-January 1997 (0.19 $\text{mol m}^{-2} \text{a}^{-1}$; Sayles et al. 2001). The POC flux was set to 0.14 $\text{mol m}^{-2} \text{a}^{-1}$ (4.62 $\text{g OM m}^{-2} \text{a}^{-1}$), which is also slightly higher than the measured POC flux averaged between the months of November and December 600 meters above the seafloor (0.11 $\text{mol m}^{-2} \text{a}^{-1}$). The clay flux, which we considered to be the total measured sediment flux minus the assumed POC and calcite fluxes, was 32 $\text{g m}^{-2} \text{a}^{-1}$. The porosity profile was tuned to best fit the porosity measurements at this station (Sayles et al., 2001, see Fig. 1). Finally, using the diffusive boundary layer world map computed in Sulpis et al. (2018) based on bottom current speeds at in-situ temperature and pressure measurements, the diffusive boundary layer thickness (δ) at the station location was set to 715 μm . A complete description of the environmental parameters for this station, along with their sources, is available in Table S2.

RADI was run using the environmental conditions described above to compare the steady-state concentration profiles of TALK, ΣCO_2 , O_2 , ΣNO_3 , calcite, and POC to in-situ measurements. Methods for solutes and solids concentration analyses

615 are described in Sayles et al. (2001). Briefly, TALK, ΣCO_2 , and ΣNO_3 were sampled in situ using the Woods Hole Interstitial Marine Probe (Sayles, 1979) while O_2 was sampled at a higher depth-resolution but in the ship laboratory using whole-core squeezing (Bender et al., 1987). We also compare the RADI steady-state concentration profiles with those obtained from a simulation with “traditional” 4.5-order calcite dissolution kinetics, all other variables being the same.



620 **Figure 5. Comparison of RADI with measurements from the station #7, mooring #3 MC-1 described in Sayles et al. (2001). The lower panels represent (left) the computed CO_3^{2-} concentrations in porewaters (solid black line) and at equilibrium with respect to calcite (dashed blue line), and (center) the fractions of organic matter degraded by a given oxidant.**

RADI predicts porewater O_2 concentrations that are slightly higher than observed (Fig. 5). Because RADI does not predict porewater O_2 to go to zero until about the 30-cm depth, the dominant organic matter degradation pathway switches

625 from mainly aerobic to ΣNO_3 at about the 30-cm depth. Nevertheless, the RADI-predicted ΣNO_3 profile is lower than observed values, particularly toward the bottom of the resolved depth. The TALK and ΣCO_2 porewater profiles predicted by a RADI simulation using 4.5-order calcite dissolution kinetics are very similar to those using the new calcite dissolution kinetics scheme, but the predicted calcite concentration is lower.

3.3 Central equatorial Pacific Ocean

630 As a third case study to evaluate the performance of RADI, solute fluxes through the sediment-water interface computed from model steady-state runs were compared to fluxes measured using benthic chambers. The comparison took place at station #W-2 described in Berelson et al. (1994) and Hammond et al. (1996), located in the Central equatorial Pacific Ocean ($0^\circ\text{N} / 139.9^\circ\text{W}$) at a depth of 4370 m. Bottom-water in-situ temperature was set to 1.40°C and salinity to 34.69 (Lauvset et al., 2016). Bottom-water oxygen concentration was set to $159.7 \mu\text{mol kg}^{-1}$ and the bottom-water saturation state with respect
635 to calcite computed using the carbonate system solver within RADI was 0.78. For the purposes of this evaluation, the CaCO_3 flux to the seafloor was assumed to be entirely calcite. The calcite flux was set to $0.22 \text{ mol m}^{-2} \text{ a}^{-1}$, which represents $22.02 \text{ g of calcite m}^{-2} \text{ a}^{-1}$, the POC flux was $0.20 \text{ mol m}^{-2} \text{ a}^{-1}$, that is, $6.6 \text{ g of organic matter m}^{-2} \text{ a}^{-1}$, and the clay flux was set to $2.0 \text{ g m}^{-2} \text{ a}^{-1}$. The steady-state calcite content within the top cm was 61 dry wt %, in line with CaCO_3 contents observed in this area (Archer, 1996; Hammond et al., 1996). The porosity profile was built using an attenuation coefficient $\beta = 33 \text{ m}^{-1}$, $\varphi_0 = 0.85$,
640 which is the measured surface porosity (Hammond et al., 1996), and $\varphi_\infty = 0.74$, which is the measured porosity at depth (Berelson et al., 1994), see Eq. (3). The DBL thickness, δ , was fixed to a value of 1 mm. A complete description of the environmental parameters for this station, along with their sources, is available in Table S3.

The diffusive fluxes for a given solute (J_v) between the sediment-water interface and the bottom waters occur as a response to the concentration gradient within the DBL and can be expressed by:

$$645 \quad J_v = \varphi_0 D_v \times \frac{v_0 - v_w}{\delta} \quad (35),$$

where v_0 and v_w are solute concentrations at the sediment-water interface and in bottom waters, respectively. In this definition, a positive J_v indicates a solute release from the sediment porewaters to the bottom waters while a negative J_v represents a solute flux towards the sediment.

The predicted TALK, ΣCO_2 , ΣPO_4 , and O_2 fluxes ($0.30 \text{ mol m}^{-2} \text{ a}^{-1}$, $0.32 \text{ mol m}^{-2} \text{ a}^{-1}$, $1.9 \text{ mmol m}^{-2} \text{ a}^{-1}$, and $-0.23 \text{ mol m}^{-2} \text{ a}^{-1}$, respectively) are all within the uncertainty bounds of the fluxes measured by benthic chambers at the same location
650 ($0.28 \pm 0.09 \text{ mol m}^{-2} \text{ a}^{-1}$, $0.24 \pm 0.09 \text{ mol m}^{-2} \text{ a}^{-1}$, $1.4 \pm 0.5 \text{ mmol m}^{-2} \text{ a}^{-1}$, and $-0.26 \pm 0.03 \text{ mol m}^{-2} \text{ a}^{-1}$, respectively, see Fig. 6). Nevertheless, the predicted ΣNO_3 flux ($9.0 \text{ mmol m}^{-2} \text{ a}^{-1}$) is lower than its measured value ($18 \pm 5 \text{ mmol m}^{-2} \text{ a}^{-1}$).




Figure 6. Comparison of fluxes computed from RADI with benthic-chamber measurements from the station #W-2 described in Hammond et al. (1996). Error bars are included for all measured fluxes but not always visible.

655

3.4 Discussion on model performance

These three model evaluation examples allowed us to determine a set of organic carbon degradation-rate constants, CaCO_3 dissolution-rate constants and organic-carbon flux composition (fast-decay, slow-decay or refractory) that can best reproduce sediment and porewater measurements in all stations while keeping all other model parameters to values from the literature. In each station, bottom-water composition was fixed from observations. Due to a lack of adequate data, solid

660

deposition fluxes were tuned to best fit observed CaCO_3 and POC contents in sediments, except in the North-western Atlantic station where the CaCO_3 and POC fluxes were taken from measurements and in the Southern Pacific station where the clay flux was inferred from observations. The POC composition that allows to best fit porewater and sediment data in the three stations was as follows: 70% of fast-decay POC, 27% of slow-decay POC, and 3% of refractory POC. The tuned fast- and slow-decay POC degradation rate constants are reported in *Organic matter degradation* section 2.2.1 and are of similar orders of magnitude as in most other models (Arndt et al., 2013). The tuned CaCO_3 dissolution rate constants are reported in *CaCO₃ dissolution and precipitation* section 2.2.3 and are two orders of magnitude lower than their laboratory-based values (Fig. S1), which we attribute to the presence of dissolution inhibitors (e.g., dissolved organic carbon, Naviaux et al., 2019a) or to the reactive surface area of natural grains in situ being lower than in laboratory experimental settings. Thus, with its current settings, RADI should be able to accurately predict porewater chemistry and sediment composition in deep-sea environments, provided that the POC and CaCO_3 deposition fluxes are known.

In the central equatorial Pacific, all RADI diffusive fluxes are within the uncertainty range of observations except the ΣNO_3 flux, which is underestimated by RADI. The low ΣNO_3 flux could be attributed, for example, to the presence of organic matter with a stoichiometry different than the Redfield ratio used in the current version of RADI or to errors in the nitrification parameters.

In addition, we note that the choice of calcite dissolution kinetics implemented in RADI does not seem to have a large impact on TALK and ΣCO_2 porewater profiles, but rather affect the predicted calcite concentration. RADI's step-edge retreat dissolution regime and its low reaction order induce calcite dissolution rates near equilibrium that are orders of magnitude higher than what is predicted in a high-order rate law (Fig. 2). That may cause the predicted calcite concentration to be higher when step-edge retreat dissolution is included in the kinetics than when a high-order rate law is implemented, because porewaters can get closer to equilibrium with respect to calcite more easily.

4 Potential model applications

In the following section, we continue to analyze the results obtained using the environmental conditions of the equatorial Pacific station #W-2 and present a few examples of relevant model applications. RADI can be used to study both steady state and transient conditions but in the following subsections we focus on time-dependent problems, since transient diagenetic models are underrepresented in the literature.

4.1 Seasonal variability

At the seafloor, the fluxes of sinking material regulating the chemical composition of sediment porewaters are patchy in both space and time. Seafloor microbes and macrofauna respond quickly to pulses of organic matter delivery to the seafloor (Smith et al., 1992), causing short-term variability of sediment oxygen consumption (Smith and Baldwin, 1984; Smith et al., 1994). In addition, both the POC and CaCO_3 fluxes to the deep seafloor are strongly affected by seasonal flux variability

(Billett et al., 1983; Smith and Baldwin, 1984; Lampitt, 1985; Lampitt et al., 1993; Lampitt et al., 2010). In the northeast Atlantic Ocean, at 3000-m depth, Lampitt et al. (2010) have shown that the summer POC and CaCO₃ fluxes can be ~ 10 and
695 ~ 4 times higher, respectively, than the winter-time minima. The seasonal coupling between organic matter and CaCO₃ fluxes to the seafloor and the state of upper-ocean ecosystem is the result of rapid vertical transport of these materials (Sayles et al., 1994). If the fluxes of reactive material reaching the seafloor is affected by seasons, early diagenesis could display a seasonal signal and this should be taken into account when interpreting sedimentary data (Martin and Bender, 1988; Sayles et al., 1994; Soetaert et al., 1996a). Here we use RADI to assess how the porewater chemistry and solid composition of deep-sea sediments
700 may be impacted by seasonally varying fluxes.

Seasonally time-varying solid fluxes to the seafloor (F , in mol m⁻² a⁻¹) can be represented with the following sinusoidal function:

$$F(t) = F_{average} + \Delta F \sin\left(\frac{2\pi t}{\Delta t}\right) \quad (36),$$

where t is time in years, Δt is the time period separating two maxima (here set to 1 year), and ΔF is the amplitude. We assume
705 that all CaCO₃ settling at the seafloor is calcite and set the mean F_{Calcite} to 0.22 mol m⁻² a⁻¹, its amplitude change $\Delta F_{\text{Calcite}}$ to 0.11 mol m⁻² a⁻¹, the mean F_{POC} to 0.20 mol m⁻² a⁻¹, and its amplitude change ΔF_{POC} to 0.10 mol m⁻² a⁻¹. All other parameters correspond to values from the Central equatorial Pacific stations described in the *Model evaluation* section 3.3.

Seasonal variations in the calcite and POC fluxes reaching the seafloor are visible in sediment profiles of both solids and solutes (Fig. 7). Nonetheless, the amplitude of concentration changes separating an annual minimum from an annual
710 maximum is very small, barely if (at all) detectable by observations. The annual amplitude is about 0.5 wt% for calcite, 0.07 wt% for POC, 2 μmol kg⁻¹ for [O₂], and 7 μmol kg⁻¹ for [ΣCO₂]. While concentrations at the sediment-water interface respond quickly to seasonal flux changes, there is a phase lag that increases with depth between the concentrations of solids and porewater solutes and the seasonally changing fluxes to the seafloor. Thus, it is possible that porewaters and solid particles at the top mm-thick sediment layers are never really at a steady state but always lagging behind seasonal changes, even if these
715 are minimal. This is in agreement with earlier modeling (Martin and Bender, 1988) and observational (Sayles et al., 1994) studies, indicating that biogeochemical reactions and bioturbation at the deep seafloor are too slow to show a discernible seasonal signal. However, this might not be the case for sites receiving more reactive organic matter (Soetaert et al., 1996a).




Figure 7. Response of porewater O_2 , ΣCO_2 , calcite, and POC concentrations to seasonal fluctuations in the calcite and POC fluxes reaching the seafloor

720

4.2 Tidal cycles

This section explores the applicability of RADI for studying the response of sediments to higher frequency phenomena such as tides. The DBL thickness is dependent on the overlying current speed (Levich, 1962; Santschi et al., 1983; Lorke et al., 2003): slower currents generate thicker DBLs whereas faster currents cause the DBL to thin (Larkum et al., 2003; Lorke et al., 2003; Higashino and Stefan, 2004). In the deep-sea, an important contributor of benthic current speeds are tidal forces, which makes tidal currents a potentially important contributor to biogeochemical exchanges across the sediment-water interface (Egbert and Erofeeva, 2002; Sulpis et al., 2019). If tidal current-speed fluctuations induce DBL thickness fluctuations, they may induce solute concentration fluctuations at the sediment-water interface, thus affecting early diagenesis. The strongest

730 tidal currents occur during the transition from high to low tides. For semidiurnal tides, the time period separating a low from a high tide is ~ 6 hours (Pugh, 1987). Setting the average DBL thickness to $\delta = 1$ mm and assuming that tides generate δ fluctuations with an amplitude $\Delta\delta = 0.5$ mm, the time-dependent δ can be expressed as:

$$\delta(t) = \delta_{average} + \Delta\delta \sin\left(\frac{2\pi t}{\Delta t}\right) \quad (37),$$

735 where t is time in years, and Δt is set to 1/1461 a (~ 6 hours). RADIV1 was run using the steady state solutes and solids depth profiles from the equatorial Pacific station #W-2 as initial conditions, with a DBL thickness fluctuating in response to tidal currents computed using Eq. (37).




Figure 8. Response of porewater O_2 , ΣCO_2 , calcite, and POC concentrations to fluctuations in DBL thickness (δ) driven by tidal currents.

740

While none of the solids seem to respond to tidal velocity fluctuation, due to their slow accumulation rate, solutes show a clear response (Fig. 8). At the sediment-water interface, the simulated porewater $[\Sigma\text{CO}_2]$ variation amplitude within a single tidal cycle is about $25 \mu\text{mol kg}^{-1}$ while $[\text{O}_2]$ oscillates with an amplitude of about $8 \mu\text{mol kg}^{-1}$. Deeper than a few millimeters below the sediment-water interface, the amplitude of both $[\Sigma\text{CO}_2]$ and $[\text{O}_2]$ changes become very small and tidal cycles are not measurable in the concentration profiles.


The implications are potentially important for our interpretation of porewater microprofiles. pH (Archer et al., 1989b; Cai and Reimers, 1993; Zhao and Cai, 1999; Cai et al., 2000), pCO_2 (Cai et al., 2000; Zhao and Cai, 1997), CO_3^{2-} (de Beer et al., 2008; Han et al., 2014; Cai et al., 2016), O_2 (Revsbech et al., 1980; Reimers, 1987; Archer et al., 1989a; Sosna et al., 2007), and even dissolved Fe, Mn or S(-II) (Brendel and Luther, 1995) microelectrodes have been developed during the past decades. According to the results presented here, microprofiles, which capture instantaneous snapshots of porewater chemistry, should show appreciable differences depending on when they are carried out during a tidal cycle. That organic matter degradation rates inferred from oxygen microprofiles span a wide range (Archer et al., 1989a; Arndt et al., 2013; Wenzhöfer et al., 2016) may be, among other factors, due to the dependency on tidal and other ocean-bottom current fluctuations. To adequately capture O_2 consumption rate in sediments, O_2 fluxes should be measured and integrated over a period of time longer than a tidal cycle (Berg et al., 2022).

4.3 Benthic chambers

RADI can also be used in the calibration of sensors and the optimization of sampling protocols and experimental designs. In the DBL, molecular diffusion is the dominant mode of solute transport and laboratory experiments of CaCO_3 dissolution in seawater suggest that diffusion through the DBL is the rate-limiting step for CaCO_3 dissolution at the seafloor in the absence of organic matter respiration (Sulpis et al., 2017; Boudreau et al., 2020). Nevertheless, earlier assessments of in-situ CaCO_3 dissolution at the sediment-water interface in the central equatorial Pacific indicated that DBL thickness does not impact overall dissolution rates (Berelson et al., 1994).

In their study, Berelson et al. (1994) deployed a set of free-sinking benthic chambers onto the seafloor. In each chamber, the portion of the chamber exposed above the sediment-water interface was sealed and isolated from external bottom waters, and water samples were drawn during the incubation period. Each incubation lasted between 80 and 120 hours. Chambers were stirred with a paddle at various rates to quantify the dependency of the measured diffusive fluxes across the sediment-water interface on the DBL thickness, which were calibrated via anhydrite dissolution to the 300-to-600 μm range. Seeing no influence of the stirring rate on the measured diffusive fluxes, Berelson et al. (1994) discarded the hypothesis of fast, surficial carbonate dissolution and instead argued for slow, high-reaction-order calcite dissolution kinetics at the seafloor, as subsequently implemented in most models. To better interpret the results from a benthic-chamber study such as that of Berelson et al. (1994), RADI can be used to predict the time-response of diffusive fluxes across the DBL following an instantaneous change of DBL thickness due, for instance, to a change in paddle stirring rate within a chamber.

RADI was run using the steady-state solutes and solids depth profiles from the equatorial Pacific station #W-2 as initial conditions. In the initial run, the DBL thickness was set to 1 mm. We simulated the response of this sediment to an instantaneous change in the DBL thickness, with one model run representing a situation where δ increases from 1 to 5 mm (e.g., a slow stirring rate) and one model run representing a δ drop from 1 mm to 200 μm (e.g., a fast stirring rate). Following a 5-fold increase in δ , diffusive fluxes of TALK, ΣCO_2 , and O_2 initially decrease by a factor ~ 5 but then increase back as the solute concentrations at the interface adapt to the new DBL (Fig. 9). Two hours after the δ increase, diffusive fluxes converge toward a new steady state. Following a 5-fold decrease in δ , diffusive fluxes immediately increase by the same magnitude, but go back close to their initial value within an hour as the interfacial porewater concentrations adjust to the new DBL. These results suggest that the incubation periods of the Berelson et al. (1994) benthic chamber experiments were long enough to let porewaters adjust to the changes caused by the paddle stirrers and confirm that, in an organic-matter and CaCO_3 -rich sediment such as in the Equatorial Pacific, the influence of a DBL on diffusive fluxes across the sediment-water interface should indeed be limited. Additionally, these results confirm the observation by Berelson et al. (1994) that, under CaCO_3 and POC deposition fluxes encountered in the equatorial Pacific, changing the stirring rate in a benthic chamber does not alter steady-state diffusive fluxes by much. Part of the reason may be the quick adjustment of porewater concentrations to the new diffusive boundary layer (see Fig. S3).



790 **Figure 9. Response of TALK, ΣCO_2 , and O_2 diffusive fluxes through the DBL to instantaneous changes in the DBL thickness (δ), for instance, as caused by a stirred benthic chamber. Positive values represent solute fluxes toward the bottom waters while negative values represent solute fluxes toward the sediments.**

4.4 Additional applications

795 The time-dependent problems presented above focus on relatively short timescales (from minutes to months). A non-steady-state model such as RADI can also be used to project the sediment response to perturbations over longer periods of

time. Examples include estimating the effect of negative emission technologies such as coastal enhanced weathering with olivine on early diagenesis (Meysman and Montserrat, 2017; Montserrat et al., 2017), of deep-sea mining (Haffert et al., 2020) or bottom trawling (Trimmer et al., 2005; van de Velde et al., 2018; De Borger et al., 2021), the impacts of a decadal bottom-water deoxygenation event such as in the Saint Lawrence estuary (Jutras et al., 2020), or the present anthropogenic CO₂ transient. However, the current version of RADI cannot deal with long-term major dissolution (erosion) events, because the burial velocity calculation scheme currently implemented does not account for solid mass gain or loss within the sediment. To study the response of sediments to a global-scale ocean acidification event such as the Paleocene–Eocene Thermal Maximum (Zachos et al., 2005; Cui et al., 2011), a different burial velocity calculation scheme would have to be implemented, such as that adopted by Munhoven (2021).

5 Future developments

One advantage of RADI is that it is easily tunable by the user: adding new components is straightforward as long as the chemical reactions are known. In future releases, we plan to add oxygen, carbon, and calcium isotopes as individual components in order to predict the diagenetic response of isotopic signals. Additionally, adsorption/desorption reactions on clay surfaces could be a critically important advance, especially regarding the prediction of sedimentary pH profiles (Meysman et al., 2003), as RADI currently treats clay minerals as non-reactive.

The representation of organic matter in the current version of the model is oversimplified. All reactive organic matter in RADI is associated with a ‘Redfield’ stoichiometry but marine organic matter can considerably deviate from this ideal (Martiny et al., 2013; Teng et al., 2014).

Finally, a model is only as good as its assumptions. RADI is targeted to study deep-sea, carbonate sediments. To be used in coastal environments, additional biogeochemical reactions would be necessary, particularly those involving methane and iron sulfide. Close to the shore, sediments become more permeable and the assumption of molecular diffusion as the dominant mode of solute transport in porewaters does not hold. In very shallow environments that are subject to high wave energy, pressure-induced advection in the sediment porewaters also needs to be included (Huettel et al., 2014). Moreover, coastal sediments have typically lower pH than open-ocean sediments, which may render our assumption of both ΣCO_2 and TAlk diffusing with a fixed diffusion coefficient set to that of the HCO₃⁻ ion inaccurate, see Fig. S2.

Code availability

The current versions of RADI in both Julia and MATLAB/GNU Octave are freely available from GitHub (<https://github.com/RADI-model>) under the GNU General Public License v3. The exact version of the model used to produce the results used in this paper is archived on Zenodo (RADI.jl v0.3; <https://doi.org/10.5281/zenodo.5005650>; v1 will be released

after review), along with input data and scripts to run the model for all the simulations presented in this paper. RADI users should cite both this publication and the relevant Zenodo reference (Humphreys and Sulpis, 2021; Sulpis et al., 2021).

Data availability

830 Sediment and porewater composition, porosity, and solid fluxes data for the Southern Pacific Ocean station described in Sayles et al. (2001) are available at http://usjgofs.who.edu/jg/dir/jgofs/southern/nbp98_2/. Sediment and porewater composition for the North-western Atlantic Ocean station described in Hales et al. (1994) are available at <https://doi.pangaea.de/10.1594/PANGAEA.730420>. The GLODAPv2 dataset used in this study is available at <https://www.glodap.info/>.

Authors contributions

835 **O.S.:** Conceptualization, methodology, software, validation, formal analysis, investigation, writing – original draft, visualization; **M.P.H.:** Conceptualization, methodology, software, formal analysis, investigation, writing – original draft; **M.M.W.:** Conceptualization, methodology, software, formal analysis, writing – review and editing; **D.C.:** Methodology, software, formal analysis, writing – review and editing; **W.M.B.:** Validation, writing – review and editing; **D.M.:** Writing – review and editing, supervision; **J.J.M.:** Methodology, validation, writing – review and editing, supervision; **J.F.A.:**
840 Conceptualization, methodology, validation, resources, writing – review and editing, supervision

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

845 Thanks are due to Bernard P. Boudreau whose CANDI model (Boudreau, 1996b) was a large source of inspiration during the creation of the present RADI model, and to Daniel L. Johnson for fruitful discussions. We thank David Burdige and one anonymous reviewer for their constructive feedback. We also thank Lukas van de Wiel for assistance with the Utrecht Geoscience computer cluster. O.S. and J.J.M. were supported by the Dutch Ministry of Education via the Netherlands Earth System Science Centre (NESSC). O.S. also acknowledges the Dept. of Earth and Planetary Sciences at McGill University for financial support during his residency in the graduate program and the Faculty of Science at McGill University for a graduate
850 mobility award. M.M.W, D.C., and D.M., carried out research at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with NASA, with support from the Biological Diversity, Carbon Cycle, Physical Oceanography, and Modeling, Analysis, and Prediction Programs.

References

- Adkins, J.F., Naviaux, J.D., Subhas, A.V., Dong, S. and Berelson, W.M.: The Dissolution Rate of CaCO₃ in the Ocean. Annual Review of Marine Science, 13, 57-80, <https://doi.org/10.1146/annurev-marine-041720-092514>, 2021
- Aller, R.C.: Transport and reactions in the bioirrigated zone, in: Boudreau, B.P., Jørgensen, B.B. (Eds.), The benthic boundary layer: transport processes and biogeochemistry, pp. 269-301. 2001
- Anderson, L.A.: On the hydrogen and oxygen content of marine phytoplankton. Deep Sea Research Part I, 42, 1675-1680, [https://doi.org/10.1016/0967-0637\(95\)00072-E](https://doi.org/10.1016/0967-0637(95)00072-E), 1995
- 860 Anderson, L.A., Sarmiento, J.L.: Redfield ratios of remineralization determined by nutrient data analysis. Global Biogeochemical Cycles, 8, 65-80, <https://doi.org/10.1029/93GB03318>, 1994
- Archer, D.: Modeling the calcite lysocline. Journal of Geophysical Research, 96, C9, 17037-17050, <https://doi.org/10.1029/91JC01812>, 1991
- Archer, D.E.: An atlas of the distribution of calcium carbonate in sediments of the deep sea. Global Biogeochemical Cycles, 865 10, 159-174, <https://doi.org/10.1029/95GB03016>, 1996
- Archer, D., Emerson, S. and Reimers, C.: Dissolution of calcite in deep-sea sediments: pH and O₂ microelectrode results. Geochimica et Cosmochimica Acta, 53, 2831-2845, [https://doi.org/10.1016/0016-7037\(89\)90161-0](https://doi.org/10.1016/0016-7037(89)90161-0), 1989a
- Archer, D., Emerson, S. and Smith, C.R.: Direct measurement of the diffusive sublayer at the deep sea floor using oxygen microelectrodes. Nature, 340, 623-626, <https://doi.org/10.1038/340623a0>, 1989b
- 870 Archer, D.E., Morford, J.L. and Emerson, S.R.: A model of suboxic sedimentary diagenesis suitable for automatic tuning and gridded global domains. Global Biogeochemical Cycles, 16, 171-1721, <https://doi.org/10.1029/2000GB001288>, 2002
- Arndt, S., Jørgensen, B.B., LaRowe, D.E., Middelburg, J.J., Pancost, R.D. and Regnier, P.: Quantifying the degradation of organic matter in marine sediments: A review and synthesis. Earth-Science Reviews, 123, 53-86, <https://doi.org/10.1016/j.earscirev.2013.02.008>, 2013
- 875 Bender, M., Martin, W., Hess, J., Sayles, F., Ball, L. and Lambert, C.: A whole-core squeezer for interfacial pore-water sampling. Limnology and Oceanography, 32, 1214-1225, <https://doi.org/10.4319/lo.1987.32.6.1214>, 1987
- Berelson, W.M., Hammond, D.E., McManus, J. and Kilgore, T.E.: Dissolution kinetics of calcium carbonate in equatorial Pacific sediments. Global Biogeochemical Cycles, 8, 219-235, <https://doi.org/10.1029/93GB03394>, 1994
- Berg, P., Huettel, M., Glud, R.N., Reimers, C.E. and Attard, K.M. Aquatic Eddy Covariance: The Method and Its Contributions to Defining Oxygen and Carbon Fluxes in Marine Environments 14, <https://doi.org/10.1146/annurev-marine-042121-012329>, 2022
- 880 Berner, R.A.: Early diagenesis: A theoretical approach. Princeton University Press, 256 pp. 1980
- Bezanson, J., Edelman, A., Karpinski, S. and Shah, V.B.: Julia: A Fresh Approach to Numerical Computing. SIAM Review, 59, 65-98, <https://doi.org/10.1137/141000671>, 2017
- 885 Billett, D.S.M., Lampitt, R.S., Rice, A.L. and Mantoura, R.F.C.: Seasonal sedimentation of phytoplankton to the deep-sea benthos. Nature, 302, 520-522, <https://doi.org/10.1038/302520a0>, 1983
- Boudreau, B.P.: On the equivalence of nonlocal and radial-diffusion models for porewater irrigation. Journal of Marine Research, 42, 731-735, <https://doi.org/10.1357/002224084788505924>, 1984
- Boudreau, B.P.: Is burial velocity a master parameter for bioturbation? Geochimica et Cosmochimica Acta, 58, 1243-1249, 890 [https://doi.org/10.1016/0016-7037\(94\)90378-6](https://doi.org/10.1016/0016-7037(94)90378-6), 1994
- Boudreau, B.P.: The diffusive tortuosity of fine-grained unlithified sediments. Geochimica et Cosmochimica Acta, 60, 3139-3142, [https://doi.org/10.1016/0016-7037\(96\)00158-5](https://doi.org/10.1016/0016-7037(96)00158-5), 1996a
- Boudreau, B.P.: A method-of-lines code for carbon and nutrient diagenesis in aquatic sediments. Computers & Geosciences, 22, 479-496, [https://doi.org/10.1016/0098-3004\(95\)00115-8](https://doi.org/10.1016/0098-3004(95)00115-8), 1996b

- 895 Boudreau, B.P.: Diagenetic Models and Their Implementation. Springer-Verlag, Berlin, 414 pp. 1997
- Boudreau, B.P. and Guinasso, N.L., Jr.: The influence of a diffusive boundary layer on accretion, dissolution, and diagenesis at the sea floor, in: Fanning, K.A., Manheim, F.T. (Eds.), *The Dynamic Environment of the Ocean Floor*. Lexington Books, Lexington, pp. 115-145. 1982
- Boudreau, B.P., Sulpis, O. and Mucci, A.: Control of CaCO₃ dissolution at the deep seafloor and its consequences. *Geochimica et Cosmochimica Acta*, 268, 90-106, <https://doi.org/10.1016/j.gca.2019.09.037>, 2020
- 900 Brendel, P.J. and Luther, G.W.I.: Development of a Gold Amalgam Voltammetric Microelectrode for the Determination of Dissolved Fe, Mn, O₂, and S(-II) in Porewaters of Marine and Freshwater Sediments. *Environmental Science & Technology*, 29, 751-761, <https://doi.org/10.1021/es00003a024>, 1995
- Buesseler, K.O., Antia, A.N., Chen, M., Fowler, S.W., Gardner, W.D., Gustafsson, O., Harada, K., Michaels, A.F., Loeff, M.R.v.d., Sarin, M., Steinberg, D.K. and Trull, T.: An assessment of the use of sediment traps for estimating upper ocean particle fluxes. *Journal of Marine Research*, 65, 345-416, 2007
- 905 Burdige, D.J.: Preservation of Organic Matter in Marine Sediments: Controls, Mechanisms, and an Imbalance in Sediment Organic Carbon Budgets? *Chemical Reviews*, 107, 467-485, <https://doi.org/10.1021/cr050347q>, 2007
- Burdige, D.J. and Gieskes, J.M.: A pore water/solid phase diagenetic model for manganese in marine sediments. *American Journal of Science*, 283, 29-47, <https://doi.org/10.2475/ajs.283.1.29>, 1983
- 910 Cai, W.-J., Ma, Y., Hopkinson, B.M., Grottole, A.G., Warner, M.E., Ding, Q., Hu, X., Yuan, X., Schoepf, V., Xu, H., Han, C., Melman, T.F., Hoadley, K.D., Pettay, D.T., Matsui, Y., Baumann, J.H., Levas, S., Ying, Y. and Wang, Y.: Microelectrode characterization of coral daytime interior pH and carbonate chemistry. *Nature Communications*, 7, 11144, <https://doi.org/10.1038/ncomms11144>, 2016
- 915 Cai, W.-J. and Reimers, C.E.: The development of pH and pCO₂ microelectrodes for studying the carbonate chemistry of pore waters near the sediment-water interface. *Limnology and Oceanography*, 38, 1762-1773, <https://doi.org/10.4319/lo.1993.38.8.1762>, 1993
- Cai, W.-J., Zhao, P. and Wang, Y.: pH and pCO₂ microelectrode measurements and the diffusive behavior of carbon dioxide species in coastal marine sediments. *Marine Chemistry*, 70, 133-148, [https://doi.org/10.1016/S0304-4203\(00\)00017-7](https://doi.org/10.1016/S0304-4203(00)00017-7), 2000
- 920 Van Cappellen, P. and Wang, Y.: Cycling of iron and manganese in surface sediments; a general theory for the coupled transport and reaction of carbon, oxygen, nitrogen, sulfur, iron, and manganese. *American Journal of Science* 296, 197-243, <https://doi.org/10.2475/ajs.296.3.197>, 1996
- Carroll, D., Menemenlis, D., Adkins, J.F., Bowman, K.W., Brix, H., Dutkiewicz, S., Fenty, I., Gierach, M.M., Hill, C., Jahn, O., Landschützer, P., Lauderdale, J.M., Liu, J., Manizza, M., Naviaux, J.D., Rödenbeck, C., Schimel, D.S., Van der Stocken, T. and Zhang, H.: The ECCO-Darwin Data-Assimilative Global Ocean Biogeochemistry Model: Estimates of Seasonal to Multidecadal Surface Ocean pCO₂ and Air-Sea CO₂ Flux. *Journal of Advances in Modeling Earth Systems*, 12, <https://doi.org/10.1029/2019MS001888>, 2020
- 925 Chriss, T.M. and Caldwell, D.R.: Evidence for the influence of form drag on bottom boundary layer flow. *Journal of Geophysical Research*, 87, <https://doi.org/10.1029/JC087iC06p04148>, 1982
- 930 Cook, P.G., Rodellas, V., Andrisoa, A. and Stieglitz, T.C.: Exchange across the sediment-water interface quantified from porewater radon profiles. *Journal of Hydrology*, 559, 873-883, <https://doi.org/10.1016/j.jhydrol.2018.02.070>, 2018
- Couture, R.-M., Sfafei, B., Van Cappellen, P., Tessier, A. and Gobeil, C.: Non-Steady State Modeling of Arsenic Diagenesis in Lake Sediments. *Environmental Science & Technology*, 44, 197-203, <https://doi.org/10.1021/es902077q>, 2010
- 935 Cowie, G.L. and Hedges, J.I.: Biochemical indicators of diagenetic alteration in natural organic matter mixtures. *Nature*, 369, 304-307, <https://doi.org/10.1038/369304a0>, 1994
- Cui, Y., Kump, L.R., Ridgwell, A.J., Charles, A.J., Junium, C.K., Diefendorf, A.F., Freeman, K.H., Urban, N.M. and Harding, I.C.: Slow release of fossil carbon during the Palaeocene–Eocene Thermal Maximum. *Nature Geoscience*, 4, 481-485, <https://doi.org/10.1038/ngeo1179>, 2011

- 940 Dade, W.B.: Near-bed turbulence and hydrodynamic control of diffusional mass transfer at the sea floor. *Limnology and Oceanography*, 38, 52-69, <https://doi.org/10.4319/lo.1993.38.1.0052>, 1993
- de Beer, D., Bissett, A., Wit, R.d., Jonkers, H., Köhler-Rink, S., Nam, H., Kim, B.H., Eickert, G. and Grinstain, M.: A microsensor for carbonate ions suitable for microprofiling in freshwater and saline environments. *Limnology and Oceanography: Methods*, 6, 532-541, <https://doi.org/10.4319/lom.2008.6.532>, 2008
- 945 De Borger, E., Tiano, J., Braeckman, U., Rijnsdorp, A. D., and Soetaert, K.: Impact of bottom trawling on sediment biogeochemistry: a modelling approach, *Biogeosciences*, 18, 2539–2557, <https://doi.org/10.5194/bg-18-2539-2021>, 2021
- Dickson, A.G.: An exact definition of total alkalinity and a procedure for the estimation of alkalinity and total inorganic carbon from titration data. *Deep Sea Research*, 28A, 609-623, [https://doi.org/10.1016/0198-0149\(81\)90121-7](https://doi.org/10.1016/0198-0149(81)90121-7), 1981
- Dong, S., Berelson, W.M., Rollins, N.E., Subhas, A.V., Naviaux, J.D., Celestian, A.J., Liu, X., Turaga, N., Kemnitz, N.J., Byrne, R.H. and Adkins, J.F.: Aragonite dissolution kinetics and calcite/aragonite ratios in sinking and suspended particles in the North Pacific. *Earth and Planetary Science Letters*, 515, 1-12, <https://doi.org/10.1016/j.epsl.2019.03.016>, 2019
- 950 Egbert, G.D. and Erofeeva, S.Y.: Efficient Inverse Modeling of Barotropic Ocean Tides. *Journal of Atmospheric and Oceanic Technology*, 19, 183-204, [https://doi.org/10.1175/1520-0426\(2002\)019<0183:EIMOBO>2.0.CO;2](https://doi.org/10.1175/1520-0426(2002)019<0183:EIMOBO>2.0.CO;2), 2002
- Emerson, S., Jahnke, R. and Heggie, D.: Sediment-water exchange in shallow water estuarine sediments. *Journal of Marine Research*, 42, 709-730, <https://doi.org/10.1357/002224084788505942>, 1984
- 955 Fiadeiro, M.E. and Veronis, G.: On weighted-mean schemes for the finite-difference approximation to the advection-diffusion equation. *Tellus*, 29, 512-522, <https://doi.org/10.3402/tellusa.v29i6.11385>, 1977
- Friedlingstein, P., O'Sullivan, M., Jones, M.W., Andrew, R.M., Hauck, J., Olsen, A., Peters, G.P., Peters, W., Pongratz, J., Sitch, S., Le Quéré, C., Canadell, J.G., Ciais, P., Jackson, R.B., Alin, S., Aragão, L.E.O.C., Arneeth, A., Arora, V., Bates, N.R., Becker, M., Benoit-Cattin, A., Bittig, H.C., Bopp, L., Bultan, S., Chandra, N., Chevallier, F., Chini, L.P., Evans, W., Florentie, L., Forster, P.M., Gasser, T., Gehlen, M., Gilfillan, D., Gkritzalis, T., Gregor, L., Gruber, N., Harris, I., Hartung, K., Haverd, V., Houghton, R.A., Ilyina, T., Jain, A.K., Joetzjer, E., Kadono, K., Kato, E., Kitidis, V., Korsbakken, J.I., Landschützer, P., Lefèvre, N., Lenton, A., Lienert, S., Liu, Z., Lombardozzi, D., Marland, G., Metzl, N., Munro, D.R., Nabel, J.E.M.S., Nakaoka, S.-I., Niwa, Y., O'Brien, K., Ono, T., Palmer, P.I., Pierrot, D., Poulter, B., Resplandy, L., Robertson, E., Rödenbeck, C., Schwinger, J., Séférian, R., Skjelvan, I., Smith, A.J.P., Sutton, A.J., Tanhua, T., Tans, P.P., Tian, H., Tilbrook, B., van der Werf, G., Vuichard, N., Walker, A.P., Wanninkhof, R., Watson, A.J., Willis, D., Wiltshire, A.J., Yuan, W., Yue, X. and Zaehle, S.: Global Carbon Budget 2020. *Earth System Science Data*, 12, 3269-3340, <https://doi.org/10.5194/essd-12-3269-2020>, 2020
- 965 Froelich, P.N., Klinkhammer, G.P., Bender, M.L., Luedtke, N.A., Heath, G.R., Cullen, D., Dauphin, P., Hammond, D., Hartman, B. and Maynard, V.: Early oxidation of organic matter in pelagic sediments of the eastern equatorial Atlantic: suboxic diagenesis. *Geochimica et Cosmochimica Acta*, 43, 1075-1090, [https://doi.org/10.1016/0016-7037\(79\)90095-4](https://doi.org/10.1016/0016-7037(79)90095-4), 1979
- Glud, R.N., Gundersen, J.K., Revsbech, N.P. and Jørgensen, B.B.: Effects on the benthic diffusive boundary layer imposed by microelectrodes. *Limnology and Oceanography*, 39, 462-467, <https://doi.org/10.4319/lo.1994.39.2.0462>, 1994
- 975 Gruber, N., Clement, D., Carter, B.R., Feely, R.A., Heuven, S.v., Hoppema, M., Ishii, M., Key, R.M., Kozyr, A., Lauvset, S.K., Monaco, C.L., Mathis, J.T., Murata, A., Olsen, A., Perez, F.F., Sabine, C.L., Tanhua, T. and Wanninkhof, R.: The oceanic sink for anthropogenic CO₂ from 1994 to 2007. *Science*, 363, 1193-1199, <https://doi.org/10.1126/science.aau5153>, 2019
- Gundersen, J.K. and Jørgensen, B.B.: Microstructure of diffusive boundary layers and the oxygen uptake of the sea floor. *Nature*, 345, 604-607, <https://doi.org/10.1038/345604a0>, 1990
- 980 Haffert, L., Haeckel, M., de Stigter, H. and Janssen, F.: Assessing the temporal scale of deep-sea mining impacts on sediment biogeochemistry. *Biogeosciences*, 17, 2767-2789, <https://doi.org/10.5194/bg-17-2767-2020>, 2020
- Hales, B., Emerson, S. and Archer, D.: Respiration and dissolution in the sediments of the western North Atlantic: estimates from models of in situ microelectrode measurements of porewater oxygen and pH. *Deep Sea Research Part I: Oceanographic Research Papers*, 41, 695-719, [https://doi.org/10.1016/0967-0637\(94\)90050-7](https://doi.org/10.1016/0967-0637(94)90050-7), 1994

- 985 Hammond, D.E., McManus, J., Berelson, W.M., Kilgore, T.E. and Pope, R.H.: Early diagenesis of organic material in equatorial Pacific sediments: stoichiometry and kinetics. *Deep Sea Research Part II: Topical Studies in Oceanography*, 43, 1365-1412, [https://doi.org/10.1016/0967-0645\(96\)00027-6](https://doi.org/10.1016/0967-0645(96)00027-6), 1996
- Han, C., Cai, W.-J., Wang, Y. and Ye, Y.: Calibration and evaluation of a carbonate microsensor for studies of the marine inorganic carbon system. *Journal of Oceanography*, 70, 425-433, <https://doi.org/10.1007/s10872-014-0243-7>, 2014
- 990 Han, X., Fang, H., He, G. and Reible, D.: Effects of roughness and permeability on solute transfer at the sediment water interface. *Water Res*, 129, 39-50, <https://doi.org/10.1016/j.watres.2017.10.049>, 2018
- Hedges, J.I., Baldock, J.A., Gélinas, Y., Lee, C., Peterson, M.L. and Wakeham, S.G.: The biochemical and elemental compositions of marine plankton: A NMR perspective. *Marine Chemistry*, 78, 47-63, [https://doi.org/10.1016/S0304-4203\(02\)00009-9](https://doi.org/10.1016/S0304-4203(02)00009-9), 2002
- 995 Higashino, M. and Stefan, H.G.: Diffusive boundary layer development above a sediment - water interface. *Water Environment Research*, 76, 292-300, <https://doi.org/10.2175/106143004X141870>, 2004
- Homoky, W.B., Weber, T., Berelson, W.M., Conway, T.M., Henderson, G.M., van Hulten, M., Jeandel, C., Severmann, S. and Tagliabue, A.: Quantifying trace element and isotope fluxes at the ocean-sediment boundary: a review. *Philosophical Transactions of the Royal Society*, 374, <https://doi.org/10.1098/rsta.2016.0246>, 2016
- 1000 Hondzo, M.: Dissolved oxygen transfer at the sediment-water interface in a turbulent flow. *Water Resources Research*, 34, 3525-3533, <https://doi.org/10.1029/1998WR900009>, 1998
- Huettel, M., Berg, P. and Kostka, J.E.: Benthic exchange and biogeochemical cycling in permeable sediments. *Ann Rev Mar Sci*, 6, 23-51, <https://doi.org/10.1146/annurev-marine-051413-012706>, 2014
- 1005 Hülse, D., Arndt, S., Daines, S., Regnier, P. and Ridgwell, A.: OMEN-SED 1.0: a novel, numerically efficient organic matter sediment diagenesis module for coupling to Earth system models. *Geoscientific Model Development*, 11, 2649-2689, <https://doi.org/10.5194/gmd-11-2649-2018>, 2018
- Humphreys, M.P. and Sulpis, O.: Radi.jl: the reactive-advective-diffusive-irrigative diagenetic sediment module in Julia. Zenodo. <https://doi.org/10.5281/zenodo.5005650>, 2021
- Humphreys, M.P., Lewis, E.R., Sharp, J.D. and Pierrot, D.: PyCO2SYS v1.7: marine carbonate system calculations in Python. *Geoscientific Model Development Discussions* [preprint], <https://doi.org/10.5194/gmd-2021-159>, in review, 2021.
- Jahnke, R.A.: The global ocean flux of particulate organic carbon: Areal distribution and magnitude. *Global Biogeochemical Cycles*, 10, 71-88, <https://doi.org/10.1029/95GB03525>, 1996
- Jørgensen, B.B.: A comparison of methods for the quantification of bacterial sulfate reduction in coastal marine sediments. *Geomicrobiology Journal*, 1, 29-47, <https://doi.org/10.1080/01490457809377722>, 1978
- 1015 Jørgensen, B.B. and Revsbech, N.P.: Diffusive boundary layers and the oxygen uptake of sediments and detritus. *Limnology and Oceanography*, 30, 111-122, <https://doi.org/10.4319/lo.1985.30.1.0111>, 1985
- Jutras, M., Dufour, C.O., Mucci, A., Cyr, F. and Gilbert, D.: Temporal Changes in the Causes of the Observed Oxygen Decline in the St. Lawrence Estuary. *Journal of Geophysical Research: Oceans*, 125, <https://doi.org/10.1029/2020JC016577>, 2020
- 1020 Keir, R.S.: The dissolution kinetics of biogenic calcium carbonates in seawater. *Geochimica et Cosmochimica Acta*, 44, 241-252, [https://doi.org/10.1016/0016-7037\(80\)90135-0](https://doi.org/10.1016/0016-7037(80)90135-0), 1980
- Lampitt, R.S.: Evidence for the seasonal deposition of detritus to the deep-sea floor and its subsequent resuspension. *Deep Sea Research Part A. Oceanographic Research Papers*, 32, 885-897, [https://doi.org/10.1016/0198-0149\(85\)90034-2](https://doi.org/10.1016/0198-0149(85)90034-2), 1985
- Lampitt, R.S., Hillier, W.R. and Challenor, P.G.: Seasonal and diel variation in the open ocean concentration of marine snow aggregates. *Nature* 362, 737-739, <https://doi.org/10.1038/362737a0>, 1993
- 1025 Lampitt, R.S., Salter, I., de Cuevas, B.A., Hartman, S., Larkin, K.E. and Pebody, C.A.: Long-term variability of downward particle flux in the deep northeast Atlantic: Causes and trends. *Deep Sea Research Part II: Topical Studies in Oceanography*, 57, 1346-1361, <https://doi.org/10.1016/j.dsr2.2010.01.011>, 2010

- 1030 Larkum, A.W.D., Koch, E.M.W. and Köhl, M.: Diffusive boundary layers and photosynthesis of the epilithic algal community of coral reefs. *Marine Biology*, 142, 1073-1082, <https://doi.org/10.1007/s00227-003-1022-y>, 2003
- Lauvset, S.K., Key, R.M., Olsen, A., van Heuven, S., Velo, A., Lin, X., Schirnick, C., Kozyr, A., Tanhua, T., Hoppema, M., Jutterström, S., Steinfeldt, R., Jeansson, E., Ishii, M., Perez, F.F., Suzuki, T. and Watelet, S.: A new global interior ocean mapped climatology: the 1° × 1° GLODAP version 2. *Earth System Science Data*, 8, 325-340, <https://doi.org/10.5194/essd-8-325-2016>, 2016
- 1035 Lee, C., Wakeham, S.G. and Hedges, J.I.: Composition and flux of particulate amino acids and chloropigments in equatorial Pacific seawater and sediments. *Deep Sea Research Part I. Oceanographic Research Papers*, 47, 1535-1568, [https://doi.org/10.1016/S0967-0637\(99\)00116-8](https://doi.org/10.1016/S0967-0637(99)00116-8), 2000
- Levich, V.G.: *Physicochemical Hydrodynamics*. Prentice-Hall Inc., Englewood Cliffs. 1962
- 1040 Li, Y.-H. and Gregory, S.: Diffusion of ions in sea water and in deep-sea sediments. *Geochimica et Cosmochimica Acta*, 38, 703-714, [https://doi.org/10.1016/0016-7037\(74\)90145-8](https://doi.org/10.1016/0016-7037(74)90145-8), 1974
- Lorke, A., Müller, B., Maerki, M. and Wüest, A.: Breathing sediments: The control of diffusive transport across the sediment—water interface by periodic boundary-layer turbulence. *Limnology and Oceanography*, 48, 2077-2085, <https://doi.org/10.4319/lo.2003.48.6.2077>, 2003
- 1045 Lueker, T.J., Dickson, A.G. and Keeling, C.D.: Ocean $p\text{CO}_2$ calculated from dissolved inorganic carbon, alkalinity, and equations for K_1 and K_2 : validation based on laboratory measurements of CO_2 in gas and seawater at equilibrium. *Marine Chemistry*, 70, 105-119, [https://doi.org/10.1016/S0304-4203\(00\)00022-0](https://doi.org/10.1016/S0304-4203(00)00022-0), 2000
- Martin, W.R. and Bender, M.L.: The variability of benthic fluxes and sedimentary remineralization rates in response to seasonally variable organic carbon rain rates in the deep sea; a modeling study. *American Journal of Science*, 288, 561-574, <https://doi.org/10.2475/ajs.288.6.561>, 1988
- 1050 Martiny, A.C., Pham, C.T.A., Primeau, F.W., Vrugt, J.A., Moore, J.K., Levin, S.A. and Lomas, M.W.: Strong latitudinal patterns in the elemental ratios of marine plankton and organic matter. *Nature Geoscience*, 6, 279-283, <https://doi.org/10.1038/ngeo1757>, 2013
- Meysman, F.J. and Montserrat, F.: Negative CO_2 emissions via enhanced silicate weathering in coastal environments. *Biol Lett*, 13, <https://doi.org/10.1098/rsbl.2016.0905>, 2017
- 1055 Meysman, F.J.R., Middelburg, J.J., Herman, P.M.J. and Heip, C.H.R.: Reactive transport in surface sediments. II. Media: an object-oriented problem-solving environment for early diagenesis. *Computers & Geosciences*, 29, 301-318, [https://doi.org/10.1016/S0098-3004\(03\)00007-4](https://doi.org/10.1016/S0098-3004(03)00007-4), 2003
- Middelburg, J.J.: A simple rate model for organic matter decomposition in marine sediments. *Geochimica et Cosmochimica Acta*, 53, 1577-1581, [https://doi.org/10.1016/0016-7037\(89\)90239-1](https://doi.org/10.1016/0016-7037(89)90239-1), 1989
- 1060 Middelburg, J.J.: *Marine Carbon Biogeochemistry: A Primer for Earth System Scientists*. Springer International Publishing, 118 pp. 2019
- Middelburg, J.J., Soetaert, K. and Herman, P.M.J.: Empirical relationships for use in global diagenetic models. *Deep Sea Research Part I. Oceanographic Research Papers*, 44, 327-344, [https://doi.org/10.1016/S0967-0637\(96\)00101-X](https://doi.org/10.1016/S0967-0637(96)00101-X), 1997
- 1065 Millero, F.J.: Thermodynamics of the carbon dioxide system in the oceans. *Geochimica et Cosmochimica Acta*, 59, 661-677, [https://doi.org/10.1016/0016-7037\(94\)00354-O](https://doi.org/10.1016/0016-7037(94)00354-O), 1995
- Montserrat, F., Renforth, P., Hartmann, J., Leermakers, M., Knops, P. and Meysman, F.J.: Olivine Dissolution in Seawater: Implications for CO_2 Sequestration through Enhanced Weathering in Coastal Environments. *Environ Sci Technol*, 51, 3960-3972, <https://doi.org/10.1021/acs.est.6b05942>, 2017
- Morse, J.W.: Calculation of diffusive fluxes across the sediment-water interface. *Journal of Geophysical Research*, 79, 5045-5048, <https://doi.org/10.1029/JC079i033p05045>, 1974
- 1070 Mucci, A.: The solubility of calcite and aragonite in seawater at various salinities, temperatures and one atmosphere total pressure. *American Journal of Science*, 283, 780-799, <https://doi.org/10.2475/ajs.283.7.780>, 1983

- Munhoven, G.: Glacial–interglacial rain ratio changes: Implications for atmospheric and ocean–sediment interaction. *Deep Sea Research Part II: Topical Studies in Oceanography*, 54, 722–746, <https://doi.org/10.1016/j.dsr2.2007.01.008>, 2007
- 1075 Munhoven, G.: Model of Early Diagenesis in the Upper Sediment with Adaptable complexity – MEDUSA (v. 2): a time-dependent biogeochemical sediment module for Earth system models, process analysis and teaching. *Geoscientific Model Development*, 14, 3603–3631, <https://doi.org/10.5194/gmd-14-3603-2021>, 2021
- Naviaux, J.D., Subhas, A.V., Dong, S., Rollins, N.E., Liu, X., Byrne, R.H., Berelson, W.M. and Adkins, J.F.: Calcite dissolution rates in seawater: Lab vs. in-situ measurements and inhibition by organic matter. *Marine Chemistry*, 215, <https://doi.org/10.1016/j.marchem.2019.103684>, 2019a
- 1080 Naviaux, J.D., Subhas, A.V., Rollins, N.E., Dong, S., Berelson, W.M. and Adkins, J.F.: Temperature dependence of calcite dissolution kinetics in seawater. *Geochimica et Cosmochimica Acta*, 246, 363–384, [https://doi.org/10.1016/0016-7037\(84\)90276-X](https://doi.org/10.1016/0016-7037(84)90276-X), 2019b
- Paraska, D.W., Hipsey, M.R. and Salmon, S.U.: Sediment diagenesis models: Review of approaches, challenges and opportunities. *Environmental Modelling & Software*, 61, 297–325, <https://doi.org/10.1016/j.envsoft.2014.05.011>, 2014
- 1085 Perez, F.F., Fontela, M., García-Ibáñez, M., Mercier, H., Velo, A., Lherminier, P., Zunino, P., de la Paz, M., Alonso-Pérez, F., Guallart, E.F. and Padin, X.A.: Meridional overturning circulation conveys fast acidification to the deep Atlantic Ocean. *Nature* 554, 515–518, <https://doi.org/10.1038/nature25493>, 2018
- Pugh, D.T.: Tides, surges and mean sea level. John Wiley & Sons Ltd. 472 pp. 1987
- 1090 Rabouille, C. and Gaillard, J.-F.: Towards the EDGE: Early diagenetic global explanation. A model depicting the early diagenesis of organic matter, O₂, NO₃, Mn, and PO₄. *Geochimica et Cosmochimica Acta*, 55, 2511–2525, [https://doi.org/10.1016/0016-7037\(91\)90369-G](https://doi.org/10.1016/0016-7037(91)90369-G), 1991
- Redfield, A.C.: The biological control of chemical factors in the environment. *American Scientist*, 46, 205–221, 1958
- 1095 Reimers, C.E.: An *in situ* microprofiling instrument for measuring interfacial pore water gradients: methods and oxygen profiles from the North Pacific Ocean. *Deep Sea Research Part A. Oceanographic Research Papers*, 34, 2023–2035, [https://doi.org/10.1016/0198-0149\(87\)90096-3](https://doi.org/10.1016/0198-0149(87)90096-3), 1987
- Revsbech, N.P., Jørgensen, B.B., Blackburn, T.H.: Oxygen in the Sea Bottom Measured with a Microelectrode. *Science*, 207, 1355–1356, <https://doi.org/10.1126/science.207.4437.1355>, 1980
- 1100 Riley, J.S., Sanders, R., Marsay, C., Le Moigne, F.A.C., Achterberg, E.P. and Poulton, A.J.: The relative contribution of fast and slow sinking particles to ocean carbon export. *Global Biogeochemical Cycles*, 26, <https://doi.org/10.1029/2011GB004085>, 2012
- Røy, H., Hüttel, M. and Jørgensen, B.B.: The role of small-scale sediment topography for oxygen flux across the diffusive boundary layer. *Limnology and Oceanography*, 47, 837–847, <https://doi.org/10.4319/lo.2002.47.3.0837>, 2002
- 1105 Sala, E., Mayorga, J., Bradley, D., Cabral, R.B., Atwood, T.B., Auber, A., Cheung, W., Costello, C., Ferretti, F., Friedlander, A.M., Gaines, S.D., Garilao, C., Goodell, W., Halpern, B.S., Hinson, A., Kaschner, K., Kesner-Reyes, K., Leprieur, F., McGowan, J., Morgan, L.E., Mouillot, D., Palacios-Abrantes, J., Possingham, H.P., Rechberger, K.D., Worm, B. and Lubchenco, J.: Protecting the global ocean for biodiversity, food and climate. *Nature*, 592, 397–402, <https://doi.org/10.1038/s41586-021-03371-z>, 2021
- 1110 Santschi, P.H., Anderson, R.F., Fleisher, M.Q. and Bowles, W.: Measurements of diffusive sublayer thicknesses in the ocean by alabaster dissolution, and their implications for the measurements of benthic fluxes. *Journal of Geophysical Research: Oceans*, 96, 10641–10657, <https://doi.org/10.1029/91JC00488>, 1991
- Santschi, P.H., Bower, P., Nyffeler, U.P., Azevedo, A. and Broecker, W.S.: Estimates of the resistance to chemical transport posed by the deep-sea boundary layer. *Limnology and Oceanography*, 28, 899–912, <https://doi.org/10.4319/lo.1983.28.5.0899>, 1983
- 1115 Sayles, F.L.: The composition and diagenesis of interstitial solutions—I. Fluxes across the seawater-sediment interface in the Atlantic Ocean. *Geochimica et Cosmochimica Acta*, 43, 527–454, [https://doi.org/10.1016/0016-7037\(79\)90163-7](https://doi.org/10.1016/0016-7037(79)90163-7), 1979

- Sayles, F.L., Martin, W.R., Chase, Z. and Anderson, R.F.: Benthic remineralization and burial of biogenic SiO₂, CaCO₃, organic carbon, and detrital material in the Southern Ocean along a transect at 170 West. *Deep Sea Research II*, 48, 4323-4383, [https://doi.org/10.1016/S0967-0645\(01\)00091-1](https://doi.org/10.1016/S0967-0645(01)00091-1), 2001
- 1120 Sayles, F.L., Martin, W.R. and Deuser, W.G.: Response of benthic oxygen demand to particulate organic carbon supply in the deep sea near Bermuda. *Nature*, 371, 686-689, <https://doi.org/10.1038/371686a0>, 1994
- Schulz, H.D.: Quantification of Early Diagenesis: Dissolved Constituents in Pore Water and Signals in the Solid Phase, in: Schulz, H.D., Zabel, M. (Eds.), *Marine Geochemistry*. Springer, Berlin, Heidelberg. 2006
- 1125 Sillén, L. G., Martell, A. E., and Bjerrum, J.: Stability constants of metal-ion complexes, Chemical Society, London, UK, special publication 17 edn., 1964.
- Smith, C.R., Berelson, W., Demaster, D.J., Dobbs, F.C., Hammond, D., Hoover, D.J., Pope, R.H. and Stephens, M.: Latitudinal variations in benthic processes in the abyssal equatorial Pacific: control by biogenic particle flux. *Deep Sea Research Part II: Topical Studies in Oceanography*, 44, 2295-2317, [https://doi.org/10.1016/S0967-0645\(97\)00022-2](https://doi.org/10.1016/S0967-0645(97)00022-2), 1997
- 1130 Smith, C.R. and Rabouille, C.: What controls the mixed-layer depth in deep-sea sediments? The importance of POC flux. *Limnology and Oceanography*, 47, 418-426, <https://doi.org/10.4319/lo.2002.47.2.0418>, 2002
- Smith, K.L., Baldwin, R.J. and Williams, P.M.: Reconciling particulate organic carbon flux and sediment community oxygen consumption in the deep North Pacific. *Nature*, 359, 313-316, <https://doi.org/10.1038/359313a0>, 1992
- Smith, K.L., Jr. and Baldwin, R.J.: Seasonal fluctuations in deep-sea sediment community oxygen consumption: central and eastern North Pacific. *Nature*, 307, 624-626, <https://doi.org/10.1038/307624a0>, 1984
- 1135 Smith, K.L., Kaufmann, R.S. and Baldwin, R.J.: Coupling of near-bottom pelagic and benthic processes at abyssal depths in the eastern North Pacific Ocean. *Limnology and Oceanography*, 39, 1101-1118, <https://doi.org/10.4319/lo.1994.39.5.1101>, 1994
- Soetaert, K., Herman, P.M.J. and Middelburg, J.J.: Dynamic response of deep-sea sediments to seasonal variations: A model. *Limnology and Oceanography*, 41, 1651-1668, <https://doi.org/10.4319/lo.1996.41.8.1651>, 1996a
- 1140 Soetaert, K., Herman, P.M.J. and Middelburg, J.J.: A model of early diagenetic processes from the shelf to abyssal depths. *Geochimica et Cosmochimica Acta*, 60, 1019-1040, [https://doi.org/10.1016/0016-7037\(96\)00013-0](https://doi.org/10.1016/0016-7037(96)00013-0), 1996b
- Soetaert, K., Herman, P.M.J., Middelburg, J.J., Heip, C., deStigter, H.S., Weering, T.C.E.v., Epping, E. and Helder, W.: Modeling ²¹⁰Pb-derived mixing activity in ocean margin sediments: Diffusive versus nonlocal mixing. *Journal of Marine Research*, 54, 1207-1227, <https://doi.org/10.1357/0022240963213808>, 1996c
- 1145 Sosna, M., Denuault, G., Pascal, R.W., Prien, R.D. and Mowlem, M.: Development of a reliable microelectrode dissolved oxygen sensor. *Sensors and Actuators B: Chemical*, 123, 344-351, <https://doi.org/10.1016/j.snb.2006.08.033>, 2007
- Subhas, A.V., Adkins, J.F., Rollins, N.E., Naviaux, J., Erez, J. and Berelson, W.M.: Catalysis and chemical mechanisms of calcite dissolution in seawater. *Proceedings of the National Academy of Sciences*, 114, 8175-8180, <https://doi.org/10.1073/pnas.1703604114>, 2017
- 1150 Sulpis, O., Boudreau, B.P., Mucci, A., Jenkins, C.J., Trossman, D.S., Arbic, B.K. and Key, R.M.: Current CaCO₃ dissolution at the seafloor caused by anthropogenic CO₂. *Proceedings of the National Academy of Sciences*, 115, 11700-11705, <https://doi.org/10.1073/pnas.1804250115>, 2018
- 1155 Sulpis, O., Dufour, C.O., Trossman, D.S., Fassbender, A.J., Arbic, B.K., Boudreau, B.P., Dunne, J.P. and Mucci, A.: Reduced CaCO₃ Flux to the Seafloor and Weaker Bottom Current Speeds Curtail Benthic CaCO₃ Dissolution Over the 21st Century. *Global Biogeochemical Cycles* 33, 1654-1673, <https://doi.org/10.1029/2019GB006230>, 2019
- Sulpis, O., Humphreys, M.P., Wilhelmus, M.M. and Carroll, D.: Radi.m: the reactive-advective-diffusive-irrigative diagenetic sediment module in MATLAB/ GNU Octave. Zenodo. <http://doi.org/10.5281/zenodo.4739205>. 2021
- Sulpis, O., Lix, C., Mucci, A. and Boudreau, B.P.: Calcite dissolution kinetics at the sediment-water interface in natural seawater. *Marine Chemistry*, 195, 70-83, <https://doi.org/10.1016/j.marchem.2017.06.005>, 2017

- 1160 Teng, Y.-C., Primeau, F.W., Moore, J.K., Lomas, M.W. and Martiny, A.C.: Global-scale variations of the ratios of carbon to phosphorus in exported marine organic matter. *Nature Geoscience*, 7, 895-898, <https://doi.org/10.1038/ngeo2303>, 2014
- Thompson, K.F., Miller, K.A., Currie, D., Johnston, P. and Santillo, D.: Seabed Mining and Approaches to Governance of the Deep Seabed. *Frontiers in Marine Science*, 5, <https://doi.org/10.3389/fmars.2018.00480>, 2018
- 1165 Trimmer, M., Petersen, J., Sivyer, D.B., Mills, C., Young, E., Parker, E.R.: Impact of long-term benthic trawl disturbance on sediment sorting and biogeochemistry in the southern North Sea. *Marine Ecology Progress Series* 298, 79-94, <https://doi.org/10.3354/meps298079>, 2005
- Uppström, L. R.: The boron/chlorinity ratio of deep-sea water from the Pacific Ocean, *Deep-Sea Res.*, 21, 161-162, [https://doi.org/10.1016/0011-7471\(74\)90074-6](https://doi.org/10.1016/0011-7471(74)90074-6), 1974.
- 1170 van de Velde, S., Van Lancker, V., Hidalgo-Martinez, S., Berelson, W.M. and Meysman, F.J.R.: Anthropogenic disturbance keeps the coastal seafloor biogeochemistry in a transient state. *Scientific Reports*, 8, 5582., <https://doi.org/10.1038/s41598-018-23925-y>, 2018
- Walter, L.M. and Morse, J.W.: The dissolution kinetics of shallow marine carbonates in seawater: A laboratory study. *Geochimica et Cosmochimica Acta*, 49, 1503-1513, [https://doi.org/10.1016/0016-7037\(85\)90255-8](https://doi.org/10.1016/0016-7037(85)90255-8), 1985
- 1175 Wenzhöfer, F., Oguri, K., Middelboe, M., Turnewitsch, R., Toyofuku, T., Kitazato, H. and Glud, R.N.: Benthic carbon mineralization in hadal trenches: Assessment by in situ O₂ microprofile measurements. *Deep Sea Research Part I: Oceanographic Research Papers*, 116, 276-286, <https://doi.org/10.1016/j.dsr.2016.08.013>, 2016
- Westrich, J.T. and Berner, R.A.: The role of sedimentary organic matter in bacterial sulfate reduction: The G model tested. *Limnology and Oceanography*, 29, 236-249, <https://doi.org/10.4319/lo.1984.29.2.0236>, 1984
- 1180 Yakushev, E.V., Protsenko, E.A., Bruggeman, J., Wallhead, P., Pakhomova, S.V., Yakubov, S.K., Bellerby, R.G.J. and Couture, R.-M.: Bottom RedOx Model (BROM v.1.1): a coupled benthic-pelagic model for simulation of water and sediment biogeochemistry. *Geoscientific Model Development*, 10, 453-482, <https://doi.org/10.5194/gmd-10-453-2017>, 2017
- Yao, W. and Millero, F. J.: The chemistry of the anoxic waters in the Framvaren Fjord, Norway, *Aquat. Geochem.*, 1, 53-88, <https://doi.org/10.1007/BF01025231>, 1995.
- 1185 Zachos, J.C., Röhl, U., Schellenberg, S.A., Sluijs, A., Hodell, D.A., Kelly, D.C., Thomas, E., Nicolo, M., Raffi, I., Lourens, L.J., McCarren, H. and Kroon, D.: Rapid Acidification of the Ocean During the Paleocene-Eocene Thermal Maximum. *Science*, 308, 1611-1615, <https://doi.org/10.1126/science.1109004>, 2005
- Zhao, P. and Cai, W.-J.: An Improved Potentiometric pCO₂ Microelectrode. *Anal Chem*, 69, 2052-5058, <https://doi.org/10.1021/ac970747g>, 1997
- 1190 Zhao, P. and Cai, W.-J.: pH polymeric membrane microelectrodes based on neutral carriers and their application in aquatic environments. *Analytica Chimica Acta*, 395, 285-291, [https://doi.org/10.1016/S0003-2670\(99\)00359-1](https://doi.org/10.1016/S0003-2670(99)00359-1), 1999
- Zuddas, P. and Mucci, A.: Kinetics of calcite precipitation from seawater: II. The influence of the ionic strength. *Geochimica et Cosmochimica Acta*, 62, 757-766, [https://doi.org/10.1016/S0016-7037\(98\)00026-X](https://doi.org/10.1016/S0016-7037(98)00026-X), 1998