

# Validation of turbulent heat transfer models against eddy covariance flux measurements over a seasonally ice covered lake

Joonatan Ala-Könni, Kukka-Maaria Kohonen, Matti Leppäranta, and Ivan Mammarella

Institute for Atmospheric and Earth System Research / Physics, Faculty of Science, University of Helsinki, Helsinki, Finland

**Correspondence:** Joonatan Ala-Könni (joonatan.ala-konni@helsinki.fi)

**Abstract.** In this study we analyzed turbulent heat fluxes over a seasonal ice cover on boreal lake located in southern Finland. Eddy covariance (EC) ~~measurements~~ flux measurements of sensible (H) and latent heat (LE) from four ice-on seasons between 2014 and 2019 are compared to three different bulk transfer models: one with a constant transfer coefficient, and two with stability adjusted transfer coefficients: the Lake Heat Flux Analyzer and SEA-ICE. All three models correlate to the EC results well in general, although typically underestimating the magnitude and the ~~variance~~ standard deviation of the flux in comparison to the EC observations. Differences between the models are small, with the constant transfer coefficient model performing slightly better than the stability adjusted models. Small difference in temperature and humidity between surface and air results in low correlation between models and EC. During melting periods (surface temperature  $T_0 > 0^\circ\text{C}$ ), the model performance for LE decreases when comparing to the freezing periods ( $T_0 < 0^\circ\text{C}$ ), while the opposite is true for H. At low wind speed, EC shows relatively high fluxes ( $\pm 20 \text{ W m}^{-2}$ ) for H and LE due to non-local effects that the bulk models are not able to reproduce. The complex topography of the lake surroundings create local violations of the Monin-Obukhov similarity theory, which help explain this counter-intuitive result. Finally, the uncertainty in the estimation of the surface temperature and humidity affects the bulk heat fluxes, especially when the difference between surface and air values are small.

## 1 Introduction

15 According to latest satellite based estimates, there are approximately 117 million lakes larger than  $0.002 \text{ km}^2$  globally (?). About 95 million of these are either above latitude  $60^\circ\text{N}$  or below  $56^\circ\text{S}$ . The seasonal lake ice zone extends on the northern hemisphere from  $40^\circ\text{N}$  to  $80^\circ\text{N}$  (?), so it can safely be estimated that over 80 % of all lakes on Earth receive a seasonal ice cover. A very defining property of lakes with seasonal ice cover is that they display two starkly different states of their surface during the annual cycle. As the ice cover forms, the lake water is effectively isolated from the atmosphere, and the already low amount of short wave radiation inherent for winter is almost completely attenuated in the snow and ice cover (?). Snow/ice–air boundary replaces the water–air boundary, and radical changes in albedo, emissivity, surface roughness, energy balance and gas exchange occur. Depending on the lake and the local climate, the snow–air boundary layer can be the dominating mode of exchange between the lake and the atmosphere. Thus, understanding the physics of seasonally ice covered lakes is an important, yet often an overlooked aspect, of understanding the behaviour of lakes (?).

25 It is easy to understand why ice covered lakes have been overlooked in the past, as most clearly observable activity on lakes happens during the open water season, but also the remote nature of many of the seasonally ice covered lakes has made their research difficult from a practical and technical standpoint (?). Nevertheless, while physical and biological processes in lakes slow down under the ice cover, they do not stop completely. Circulation is driven by sediment heat accumulated there during the summer, meltwater streaming from the surrounding catchment area and solar radiation, with additional mixing produced  
30 via the breaking of internal waves promoted by changes in air pressure and wind, and although primary production is minimal, other biological processes still continue (?) and affect especially the gas fluxes of the lake (?). Regardless of the season, lakes ~~have an effect in the very large scale of global climate as well as in the very local scale~~ affect the climate both at regional and global scales. In the large scale, lakes affect the global climate by acting as small net sources of carbon (CO<sub>2</sub> and CH<sub>4</sub>) into the atmosphere and also sequestering organic carbon into their sediments from internal biogeochemical processes and from  
35 the surrounding environment (?). In the small scale, due to their large heat storage, thermal inertia and evaporation, lakes can significantly affect local and regional weather patterns and microclimate, like rain and snowfall, temperature and cloudiness (???). As lakes contribute significantly to ~~their local the~~ climate, understanding their surface heat balance more precisely has ~~use in many scales~~ relevant implications, from local short term weather and ice cover forecasting (?) to long term global circulation models, where the effect of lakes has been neglected almost completely (?).

40 The yearly cycle of a lake is driven by external forcing, which follow changes in the patterns of the components of the surface energy and water balance. The surface energy balance constitutes of incoming and reflected solar radiation (also known as short wave radiation), incoming and outgoing terrestrial radiation (also called long wave radiation), turbulent heat fluxes (latent and sensible heat flux) and the precipitation heat flux (?).


Annual changes in the solar radiation drive the changes in seasons, and during the summer it dominates the energy balance. In  
45 autumn the incoming solar radiation decreases every day, and eventually enough heat will be lost through turbulent heat fluxes and outgoing terrestrial radiation to lower the temperature of the water column to the temperature of maximum density, which ~~for fresh water~~ is +4 °C for freshwater. Then, the lake mixes completely, while cooling continues. At high latitude, seasonal ice cover begins to form usually in the late autumn during clear sky and low wind conditions associated with anticyclonal weather patterns, although frazil ice can also form in the turbulent surface layer of the lake as well. During nights in calm, cloud free  
50 conditions there is significant loss of heat from the water through long wave radiation and freezing water can accumulate to the surface without being mixed with the warmer water below forming primary ice under which the more permanent congelation ice can form. Later during the winter snow can accumulate over the ice and freeze into solid, opaque snow ice. It insulates the lake more effectively from the solar radiation than the clear congelation ice. Melting begins when the radiation balance turns positive, and the surface absorbs more radiation. The length of ice season varies significantly depending on the local climate at  
55 the lake. Lakes in Southern Finland spend less than half of the year under an ice cover, but above ~~the latitude~~ 65°N lakes ~~are on average have~~ have on average a longer ice-on than ice-off season (?). Although statistics can be drawn, every winter on a lake is unique in regards to the length of the ice cover period, the layering of the ice cover, amount on snow accumulation and precipitation.

Radiative components of the energy balance are relatively simple to measure due to the passive instruments with low power consumption required to measure them, but turbulent heat transfer poses more challenges. During last four decades the eddy covariance (EC) technique has become a very popular method in many fields of environmental and geophysical sciences. It is an accurate, proven and well established method for directly measuring vertical fluxes of heat, momentum, gases and particles over a wide variety of surface types and ecosystems (?). Its strong points are the ability to collect long, continuous time series in many different environmental and meteorological conditions. Although it has been extensively used over terrestrial environments, lately EC has been applied in marine and fresh water environments as well.

As the EC setup is not suited for all applications due to the technical complexity of its installation, simpler methods to compute the turbulent heat fluxes from more basic meteorological observations have been developed, with bulk aerodynamic method and profile method being the most popular. They originated from the need to estimate turbulent heat fluxes in situations where only basic meteorological parameters were available, like with remote buoy based ~~oceanographical~~ oceanographic measurement stations with very low power available. They are also commonly used in global and regional climate models and as well as in Numerical Weather Prediction models due to their computational simplicity. For different applications (marine, land etc.), the parameterization ~~parameterization~~ of the stability, aerodynamic roughness and other parameters of the model can be adjusted accordingly. ~~for example, some models have a completely theoretical approach, while others are empirical.~~

Estimating turbulent heat fluxes by the bulk aerodynamic method is simpler than measuring them with EC, as only basic meteorological measurements are required, but it inherently contains some limitations and uncertainty. Stable boundary layer conditions and surface heterogeneity are especially troublesome, and the assumptions made in the Monin-Obukhov similarity theory (MOST) do not take into account all meteorological phenomena, like non-local effects produced by the ~~boundary between the ice cover and the surrounding forest, present over lake ice cover (??). Eddy covariance promises to issue some of these deficiencies of the bulk method. As the source area it measures is significantly larger than that of the radiation sensors required by the bulk method, it has the possibility to overcome the challenges of heterogeneous surface~~ surface heterogeneity present over small lakes surrounded by forests (??).

While few studies have reported short field campaign measurements of EC turbulent heat fluxes over seasonal ice covered lakes (??), long term turbulent heat flux measurements have not been reported so far. In this study, we present ~~an~~ a unique data set collected over a boreal lake in southern Finland over four ice-on seasons between 2014 and 2019. Previous studies of turbulent heat fluxes over lakes have been performed mostly in the open water season, like northern boreal lake in Finnish Lapland (?), boreal lake in Southern Finland (?) and the lake in question in this study, Lake Kuivajärvi (?). Ice-on lake energy balance has been studied, for example, on Lake Kilpisjärvi in NW Finnish Lapland (?) and Lake Pääjärvi in Southern Finland (??), but these experiments were done without EC equipment and estimated turbulent heat fluxes by bulk aerodynamic formulae and the profile method. EC over seasonal lake ice cover was performed over a thermokarst lake in Siberia (?), but it was also only for one winter. Thus, the data set presented here gives us a unique look into the dynamics of turbulent heat fluxes over seasonal lake ice cover as well as a possibility to validate the functionality of bulk transfer models in this environment.



**Figure 1.** Bathymetric map of Lake Kuivajärvi with the position of the EC raft shown as a white square (a). Photo (b) shows the raft as it stood in March 2021. Map adapted from ?.

## 2 Material & methods

### 2.1 Site

Lake Kuivajärvi is a dimictic, mesotrophic lake in southern Finland (lon.  $24^{\circ} 16' E$ , lat.  $61^{\circ} 50' N$ , 141 m above mean sea level). It is located in the Kokemäenjoki water system, which drains into the Baltic Sea. The lake has a strongly elongated shape, 2.6 km in the north – south direction and 200–400 m in the east–west direction (Fig. 1). Due to this shape, wind is usually channeled along the lake (?). The lake has two basins, the southern basin being the deeper one with a maximum depth of 13.2 m. The raft used for EC and most of the meteorological measurements is located approximately over this deepest point. Lake Kuivajärvi is surrounded mostly by managed Scots pine forest, which is also the home for the SMEAR II station (?).  
 100 Typical ice cover period in Lake Kuivajärvi lasts for about five months, starting in late November – early December and ending in late April – early May. A mild decreasing trend in the length of the ice-on season has been observed here since the start of observations in 1929 (?). The ice cover thickness at the start of the melting period is typically 40–50 cm.

### 2.2 Measurement of fluxes and meteorology

Fluxes of momentum, sensible ( $H$ ) and latent heat ( $LE$ ) and supporting meteorological measurements were performed on a raft anchored at the deepest point of the lake, as well as on the nearby SMEAR II station (?). The EC system consists of  
 105

**Table 1.** Ice-on periods at Lake Kuivajärvi used in this study (2014 – 2019) and corresponding number of 30 minute EC flux values used in this study.

Period	N. of 30 min fluxes (H) & coverage	N. of 30 min fluxes (LE) & coverage
9.12.2014 – 20.4.2015 (132 d)	3 662 (57.8 %)	3 264 (51.5 %)
23.1.2017 – 3.5.2017 (100 d)	2 315 (48.2 %)	1 956 (40.8 %)
3.12.2017 – 23.4.2018 (141 d)	3 632 (53.7 %)	2 905 (42.9 %)
16.11.2018 – 25.4.2019 (160 d)	2 250 (29.3 %)	1 967 (25.6 %)
Total: 533 days	11 859 (46.3 %)	10 092 (39.4 %)

Metek (Metek GmbH, Elmshorn, Germany) USA-1 3-axis anemometer providing the three component wind speed and sonic temperature and a LI-COR (LI-COR Inc., Nebraska, USA) 7200 measuring the water vapor (H<sub>2</sub>O) mixing ratio at 10 Hz frequency. A Kipp & Zonen (Kipp & Zonen, Delft, Netherlands) CNR1 net radiometer and pyranometer was used to acquire the radiation balance. This single instrument measured both directions (incoming and outgoing) of the short wave (305 – 2 800 nm) and long wave (5 000 – 50 000 nm) radiation. Relative humidity and air temperature are measured by Rotronic (Rotronic Instrument Corp., NY, USA) MP102H sensor at a height of 1.8 m.

The sonic anemometer and gas inlets were installed at a height of ~~1.7~~ 1.8 m on the western side of the raft, facing sideways of the prevalent wind directions in order to avoid the structure of the raft interfering with the wind and its measurement.

Data from four winters between 2014 and 2019 was used. Ice season was considered to begin on the day when the surface albedo had risen to 0.5, and to end on the day when it had reached values  $\alpha < 0.1$  permanently. Dates of ice-on and ice-off for these winters are presented in Table 1.

Surface temperature was derived from outgoing long wave radiation by the Stefan–Boltzmann law, and a constant emissivity of  $\epsilon = 0.997$  was assumed, which is a typical value for a snowy surface (?).

### 2.3 EC data processing

Eddy covariance raw data were processed with the EddyUH software (?), and fluxes were calculated using 30 minute averaging time as

$$H = \rho_a c_p \overline{w' T_a'} \quad (1)$$

$$LE = \rho_a L_e \overline{w' q_a'} \quad (2)$$

where  $T_a$  is air temperature,  $w$  is the vertical wind speed [ $\text{m s}^{-1}$ ],  $q_a$  is the specific humidity of air [ $\text{kg kg}^{-1}$ ],  $\rho_a$  is the density of air [ $\text{kg m}^{-3}$ ],  $L_e$  is the latent heat of ~~evaporation~~ evaporation-vaporization of water [ $\text{J kg}^{-1}$ ] and  $c_p$  is the heat capacity of air [ $\text{J kg}^{-1} \text{K}^{-1}$ ]. The prime marks the fluctuation of the corresponding value from its mean. Micrometeorological notation was used for

the sign of the flux: negative fluxes are downward and positive fluxes upwards. State of the art methodologies for the data processing typically used in land based flux ~~tower-towers~~ (?) were applied and adapted following ?. ~~Only evaporation was used instead of evaporation and sublimation in order to be able concentrate more on the errors produced by the calculation of stability.~~

In short, 2-D coordinate rotation was applied to the anemometer data in order to direct the  $u$ -component along the mean horizontal wind direction and to result in mean vertical velocity of  $\bar{w} = 0$ . Linear detrending was applied in the calculation of turbulent fluctuations. Removal of spikes was performed by setting limits for the difference between subsequent values. Half-hourly blocks of raw data were rejected if they contained over 3 000 spikes. Time lag of H<sub>2</sub>O was determined from the maximum of cross-covariance function between vertical wind velocity and H<sub>2</sub>O mixing ratio and cross-wind correction was applied to the sonic temperature data (?). High-frequency spectral corrections were done in accordance to (?). Flux quality flags were based on flux stationarity (FST), skewness (SK) and kurtosis (KU). Only flux values that had the highest quality flag "0" were used. For this quality class, the conditions were for flux stationarity  $FST \leq 0.3$ , for skewness  $-2 < SK < 2$  and for kurtosis  $1 < KU < 8$ . Wind direction was also used as a criterion for usable data, and only wind blowing along the lake ( $130^\circ < WD < 180^\circ$  and  $320^\circ < WD < 350^\circ$ ) were accepted (?). A total of 11 859 flux values were accepted for H and 10 092 for LE, resulting in a data coverage of 46.3 % and 39.4 % respectively (Tab. 1).

## 2.4 Flux footprint

A two-dimensional footprint analysis was conducted on the dataset using the model described in ?. Winter 2016 – 2017 was chosen to be a representative year and the footprint was calculated from all accepted 30-min flux values from this ice-on season. In Fig. 2 the footprint is presented for three stability classes: stable ( $zL^{-1} > 0.1$ ), neutral ( $-0.1 < zL^{-1} < 0.1$ ) and unstable ( $zL^{-1} < -0.1$ ), where  $z$  is the measurement height and  $L$  the Obukhov length.


The contour lines represent the limit of 80 % footprint area. From Fig. 2 it can be seen, that the footprint stays well within the boundaries of the lake due to the strongly channeled wind. The 80 % footprint reaches approximately 300 m from the raft in north-south direction and 100 m in east-west direction.

## 2.5 Bulk transfer models

Various forms of the bulk transfer models have been used to estimate vertical turbulent fluxes for decades. In its simplest form the sensible (latent) heat flux is written as a linear function of wind speed, temperature (humidity) difference between the surface and the air and their corresponding transfer coefficients. In order for the bulk fluxes to have the same sign as the EC data has, in this study they are calculated as

$$H_b = \rho_a c_{ap} C_H (T_0 - T_a) U \text{ and} \quad (3)$$

$$LE_b = \rho_a L_e C_E (q_0 - q_a) U \quad (4)$$



**Figure 2.** Footprint of EC measurements calculated from ice-on season 2016 – 2017 in three stability classes: stable ( $zL^{-1} > 0.1$ ), neutral ( $-0.1 \leq zL^{-1} \leq 0.1$ ) and unstable ( $zL^{-1} < -0.1$ ). The plotted lines represent 80 % footprint. Scale is in meters. Calculated by footprint script described in Kljun et al. (2015). Map: the National Land Survey of Finland, 1.4.2022.

160 where  $L_e$  is the heat of vaporization of water (see Eq. ??),  $C_H \approx C_E$  and  $C_E$  are the transfer coefficients of sensible and latent heat, respectively,  $U$  is the wind speed,  $T_a$  and  $T_0$  are the air and surface temperature respectively and  $q_a$  and  $q_0$  are the air and surface humidity. In Eq. 4-(4) surface humidity can be assumed to be at saturation due to the watery / icy surface (?).

In this study three different bulk transfer models are compared against EC measurements: one model where the transfer coefficient is kept constant, one developed specifically for open water lake environments (Lake Heat Flux Analyzer) and one  
165 developed for sea ice (SHEBA Bulk Turbulent Flux Algorithm for Sea Ice v. 2.0).

### 2.5.1 Constant transfer coefficient model

~~A The use of a constant transfer coefficient model takes in no way into account the stability of atmospheric boundary layer since the values for neutral conditions are applied to all stability conditions. By including this simplified model we can better see the effect of different ways to include the boundary layer stability into the model. Values between  $C_E = C_H \approx 1.0 - 1.5 \cdot 10^{-3}$~~   
170 neutral value is a simplified approach, which assumes negligible effect of atmospheric stability. A range of values between  $1.0 \cdot 10^{-3}$  and  $1.5 \cdot 10^{-3}$  have been reported for  $C_E$  and  $C_H$  over the years for neutral conditions at 10 m height, for example

in ?. ~~In our study a mean~~ Due to the fact that underestimation of turbulent heat fluxes is typical, a value of  $1.5 \cdot 10^{-3}$  was chosen, which corresponds to the high end of the values reported in ? was taken and the value was scaled represented in the literature. This value was then scaled from 10 m height to a height of 1.7-1.8 m, resulting in a value value of  $C_E = C_H = 1.8 \cdot 10^{-3}$  :-

175 For  $L_e$  in the constant coefficient model a temperature dependent model was applied (?):-

$$L_e = 2500.8 - 2.36T_0 + 0.0016T_0^2 - 0.00006T_0^3.$$

being chosen for our study. Fluxes are then calculated by using Eqs. (3) and (4).

## 2.5.2 Lake Heat Flux Analyzer (LHFA)

The second model used in this study is the Lake Heat Flux Analyser software, described in ?. It was originally developed in order to create a standardized way to compute turbulent flux values acquired from lake measurement networks. The calculation of roughness length of momentum is based on ?:-

$$z_0 = \left( \frac{\alpha_1 u_{*a}^2}{g} \right) + \left( \frac{\alpha_2 \nu_a}{u_{*a}} \right)$$

185 where  $g$  is gravitational acceleration  $m \cdot s^{-2}$ ,  $\alpha_1 = 0.013$  is the Charnock constant,  $\alpha_2 = 0.11$ ,  $\nu_a = 1.5 \cdot 10^{-5} m^2 \cdot s^{-1}$  is kinematic viscosity of air and  $u_{*a}$  is the friction velocity in air  $m \cdot s^{-1}$ . Roughness lengths of heat ( $z_{0T}$ ) and moisture ( $z_{0q}$ ) are estimated based on ? as :-

$$z_{0q} = z_{0T} = z_0 \exp(-b_1 Re^{0.25} - b_2),$$

where  $b_1 = 2.67$ ,  $b_2 = -2.57$  and  $Re = u_{*a} z_0 \nu_a^{-1}$  is roughness Reynolds number. Stability  $\zeta$  is calculated as a ratio of measurement height  $z$  and Obukhov length  $L$ , which is defined following (?) as-

$$L = \frac{-\rho_a u_{*a}^3 T_v}{kg \left( \frac{H_b}{c_a} + 0.61 \frac{(T_a + 273.16) L E_b}{L_e} \right)}$$

190 where  $T_v = (T_a + 273.16) \cdot (1 + 0.61 q_z)$  is the virtual temperature K, Following stability classes were assigned for the flux-gradient relations for sensible ( $\Phi_h(\zeta)$ ) and latent heat ( $\Phi_e(\zeta)$ )

$$\Phi_{e,h}(\zeta) = 5 + \zeta \text{ for } \zeta > 1 \text{ (very stable)}$$

$$\Phi_{e,h}(\zeta) = 1 + 5\zeta \text{ for } 0 \leq \zeta \leq 1 \text{ (stable)}$$



$$\Phi_{e,h}(\zeta) = (1 - 16\zeta)^{-1/2} \text{ for } -0.465 \leq \zeta \leq 0 \text{ (unstable)}$$

$$\Phi_{e,h}(\zeta) = 0.9k^{4/3}(-\zeta)^{-1/3} \text{ for } \zeta < -0.465 \text{ (very unstable)}$$

where  $k = 0.4$  is the von Kármán constant. The final transfer coefficient for latent heat flux for the measurement height  $z$  was calculated by-

$$C_{Ez} = \frac{-u_{*a}q_*}{u_z(q_0 - q_z)}$$

where  $q_*$  is the dimensionless scaling parameter for air humidity,  $u_{*a}$  is software uses an iterative approach to calculate H and LE with stability corrections based on the friction velocity of air,  $u_z$  MOST, as described in ?. As an input to calculate the turbulent fluxes the model uses air temperature, air humidity, surface temperature and wind speed. The model is developed especially for open water season, but for the sake of comparison it was decided to include this model as well. The open water optimization is apparent in the calculation of roughness length, which is done according to ? and thus includes the Charnock term as well, which in certain conditions takes into account the effect of surface waves. Roughness lengths for heat and  $q_z$  are the wind speed and air humidity at height  $z$  respectively. The transfer coefficient for sensible heat flux  $C_{Hz}$  is calculated similar to this, but replacing the scaling parameter  $q_*$  by dimensionless temperature  $T_*$  and replacing  $(q_0 - q_z)$  by  $(T_0 - T_z)$ . The software uses an iterative approach to calculate H and LE moisture are calculated according to ?.

### 2.5.3 SHEBA Bulk Turbulent Flux Algorithm for Sea Ice v. 2.0 (SEA-ICEHICE)

The third model applied in this study was developed for sea ice and is based on the data set acquired from the SHEBA (Surface Heat Budget of the Arctic Ocean) experiment in the Beaufort Sea (?). In this model, It is similar to the LHFA model in the sense that it also uses an iterative approach to the stability corrections, but in this model in very stable cases ( $\zeta > 1$ ) the flux gradient relations rely on empirical approximations from (?), for near-neutral cases no stability correction is applied and for unstable cases formulation from (?) is applied. The integral forms of stability functions and their corresponding stability classes are as follows:-

$$\Psi_{e,h}(\zeta) = \frac{b_h}{2}(1 + c_h\zeta + \zeta^2) + \left(-\frac{a_h}{B_h} + \frac{b_h c_h}{2B_h}\right) \cdot \left(\ln \frac{2\zeta + c_h - B_h}{2\zeta + c_h + B_h} - \ln \frac{c_h - B_h}{c_h + B_h}\right) \text{ for } \zeta > 1$$

$$\Psi_{e,h}(\zeta) = 0 \text{ for } 0 \leq \zeta \leq 1$$

$$\Psi_{e,h}(\zeta) = 2 \ln \left(\frac{1 - x^2}{2}\right) \text{ where } x = (1 - \gamma\zeta)^{1/4} \text{ for } \zeta < 0$$

Here,  $a_h = 5$ ,  $b_h = 5$ ,  $c_h = 3$ ,  $B_h = \sqrt{5}$  and  $\gamma = 16$ . The transfer coefficients for sensible and latent heat fluxes were calculated as-

$$225 \quad C_H = \frac{k^2}{[\ln(z/z_0) - \Psi_m(\zeta)][\ln(z/z_T)] - \Psi_h(z_{0T}/L)}$$

$$C_E = \frac{k^2}{[\ln(z/z_0) - \Psi_m(\zeta)][\ln(z/z_Q)] - \Psi_e(z_{0q}/L)}$$

The roughness lengths were calculated by an analytical method described in ?.

230 ~~The model has settings for both summer and winter sea ice, but for this study it was run in the "winter" setting with the sea ice concentration set at~~ roughness length calculation (described in ?) is optimized for ice cover, unlike the LHFA which is optimized for open water. In this study the SEA-ICE model was run at the 'winter' setting, corresponding to 100 % ice coverage.


### 3 Results

#### 3.1 Environmental drivers of diurnal, seasonal and interannual variation of turbulent heat fluxes

235 ~~Several meteorological phenomena affect and drive turbulent heat fluxes over a lake ice cover. Gradient in specific humidity or temperature between the surface and the air above it is required for vertical transport of heat or water vapor and wind or convection is needed to keep the air above the surface in movement and generating turbulence. These phenomena do not propagate themselves, but heat is required to drive the wind, bring temperature differences and provide the energy required for evaporation.~~

240 Diurnal and seasonal variation as well as the response of the turbulent fluxes to external forcing were studied by dividing the fluxes and meteorological data for all years by month and by hour of day (Fig. 3). Early in the winter (December to February) no diurnal pattern of turbulent fluxes is visible, but as the sun gets higher over the horizon in the spring and short wave radiation begins to dominate the surface energy balance the absolute flux values rise up as well and a diurnal pattern for H (Fig. 3a), LE (Fig. 3b) and wind speed (Fig. 3h) develops. Thus, the ice-on season can be divided into two phases: early winter with no  
245 diurnal pattern and late winter with a diurnal pattern with the sunlight affecting the fluxes.

The highest median value of sensible heat flux ( $-34.3 \text{ W m}^{-2}$ ) is seen in April at 16:00 UTC+2 (Fig. 3a). The flux peak is observed at the same time as the peak of temperature gradient (Fig. 3i), and lagging the maximum of net radiation by about four hours (Fig. 3g). In the darkest winter months the surface temperature of the ice cover follows the air temperature very closely, thus keeping the sensible heat flux low as well ( $0 \text{ W m}^{-2} > H > -5 \text{ W m}^{-2}$ ). More difference between the two can  
250 occur especially in the spring, when the air temperature can reach values well above freezing, but the melting ice and snow surface remains close to  $0^\circ\text{C}$ , resulting in negative flux values, i.e. heat transferred from the air into the melting ice surface. Winters at Lake Kuivajärvi (and in Southern Finland in general) are defined by cold spells separated by days where the air temperature reaches values above freezing. Peaks in sensible heat flux occur then as well, and the number of such peaks varies from year to year.



**Figure 3.** Hourly medians of key variables measured over the lake ice: (a) H and (b) LE measured by EC, all four components of the radiation balance separately ((c) through (f)), (g) net radiation, (h) wind speed, (i) temperature difference ( $T_0 - T_a$ ) and (j) specific humidity difference ( $q_0 - q_a$ ). Each curve represents a month of data separated into each hour of the day. Notice, that the turbulent fluxes have the micrometeorological notation, where negative is flux towards the surface, while net radiation and radiation flux components has the opposite notation. Net radiation plot has two vertical axes to accommodate the higher fluxes of March and April (dashed lines, scale on the right side), solid lines represent months from December to February (scale on the left). All times are in UTC+2.

255 A similar pattern is observable for the latent heat flux (Fig. 3b), but with a positive flux developing along with the intensifying radiation instead of a negative flux. During the darkest winter months there is weak median deposition or sublimation of ice or snow at all hours of the day with no clear diurnal pattern (median  $-1.5 \text{ W m}^{-2} < \text{LE} > 1.5 \text{ W m}^{-2}$ , or  $\sim \pm 0.05 \text{ mm}$  per day), which turns into daytime evaporation/sublimation in March, when daytime net radiation turns positive. This amount of evaporation or sublimation plays no significant role in the ice and snow mass balance of Lake Kuivajärvi, as the ice cover is in the order of 40 cm with centimeters of snow on top of that, and just the transport of snow by wind can be larger than this evaporation / sublimation. Peak median value,  $18.4 \text{ W m}^{-2}$ , is in afternoon at 14:00 (UTC+2), following the peak in the humidity difference (Fig. 3j), again lagging some four hours behind the peak in net radiation. Diurnal pattern in the humidity difference develops later than that of the temperature difference, and hence the diurnal pattern of LE is visible a month later than that of H.

260

265 ~~Surface energy balance (Fig. 4f) was calculated as the sum of the four components of radiation and turbulent heat fluxes. Between the months of January and February the surface energy balance had values typically ranging from  $-50 \text{ W m}^{-2}$  and  $50 \text{ W m}^{-2}$ . After March up until ice off surface energy balance varied between  $-100 \text{ W m}^{-2}$  and  $700 \text{ W m}^{-2}$ .~~

~~In the darkest winter months the surface temperature of the ice cover follows the air temperature very closely, thus keeping the sensible heat flux low as well. More difference between the two can occur especially in the spring, when the air temperature can reach values well above freezing, but the melting ice and snow surface is stuck at  $0^\circ\text{C}$ , resulting in heat transferred from the air into the melting ice surface. Winters at Lake Kuivajärvi (and in Southern Finland in general) are defined by cold spells separated by days where the air temperature reaches values above freezing. Peaks in sensible heat flux occur then as well, and the number of such peaks varies from year to year.~~

270

All timeseries used in this study are presented in Figure 4. Interannual differences in the pattern of turbulent heat fluxes occur mostly in the number of cold and warm periods and differences in their lengths. Low temperatures result in low values of LE for two reasons: one is the very dry nature of cold air and the exponential relation of dew/frost point to air temperature, and the low vertical gradient of specific humidity which follows from this. The second reason is the low wind speeds that are typically associated with the anticyclonal weather patterns that result in low air temperatures during winters. Low air temperatures result in lower values of H as well, but to a smaller degree than for LE. During winters the stability of the atmospheric boundary layer is mostly stable ( $zL^{-1} > 0$ ), as can be seen in Fig. 4e. In this data set stable conditions were measured for 73 % of the time and unstable conditions for 27 %. The change into the higher late-winter flux values ( $H \& LE > 10 \text{ W m}^{-2}$ ) quite consistently begins in mid to late March regardless of year. Peaks of  $\pm 50 \text{ W m}^{-2}$  are observed almost daily in spring, while in the early winter they are more intermittent and associated with situations where the air temperature reaches values significantly above freezing.

280

Surface energy balance (Fig. 4f) was calculated as the sum of the four components of radiation and turbulent heat fluxes. Between the months of January and February the surface energy balance had values typically ranging from  $-50 \text{ W m}^{-2}$  and  $50 \text{ W m}^{-2}$ . After March up until ice off surface energy balance varied between  $-100 \text{ W m}^{-2}$  and  $700 \text{ W m}^{-2}$ . The excess heat in the spring is used up in melting of the ice cover as well as heating the underlying water.

285


The onset of ice cover varies from year to year (range of 45 days in the four years studied here), ~~as~~ and it is controlled by the ~~temporally random nature of the~~ meteorological conditions suitable for ice formation. Ice-off is more predictable (range of 13 days), as it is driven by the incoming solar radiation which regardless of meteorological factors always begins to dominate the surface energy balance in March.

290

### 3.2 Comparison of turbulent fluxes derived by EC and bulk transfer models

For the comparison of the EC data with the models the data ~~set~~ was divided into two sets: one where the surface temperature was below freezing and one where it was above freezing. The data are compared in four ways: First, by boxplots where the data are divided, in addition to surface conditions, into each hour of the day (Fig. 5). Second, by scatter plots (Figs. 6 & 8), third by studying the correlation and centered root mean square error (CRMSE) as functions of wind speed, temperature difference and specific humidity difference (Fig. 10), and ~~finally~~ by Taylor plots (?), showcasing the differences of correlation,

295




**Figure 4.** All four winters of (a) turbulent flux data, (b) air and surface temperature, (c) air and surface humidity, (d) wind speed, (e) natural logarithm of stability and (f) surface energy balance (SEB). Each column represents one winter. All data are presented as half hourly values, except for wind, which is presented as a six hour moving mean. Notice that the surface energy balance notation is such that positive values indicate heat flux into the surface and vice versa.

CRMSE and standard deviation between the models and the EC measurements (Figs. 7 & 9). Finally, the behaviour of neutral transfer coefficients are studied in relation to wind speed and normalized transfer coefficients are examined against atmospheric stability.

### 3.2.1 Sensible heat flux (H)

EC results of sensible heat flux exhibit negative values for most of the time, i.e. heat deposited onto the surface (Figs. 5a & c). During Sensible heat flux modeling was observed to function better in melting than freezing surface conditions (Fig. 5e), the daily cycle of H is not as pronounced as it is during melting conditions, but this is mostly due to the low net radiation values of the early winter months. Frozen conditions result in worse overall agreement between the models and EC and all models




**Figure 5.** Daily variation of fluxes in EC and the three models. (a) H in melting surface conditions, (b) LE in melting surface conditions, (c) H in freezing surface conditions and (d) LE in freezing surface conditions. The line in the middle of each bar indicates median, bar edges indicate 25th and 75th percentiles and the whiskers extending from the bars indicate  $2.7\sigma$ , or 99.7 % of values are within these boundaries. All times are in UTC+2.

are quite consistently underestimating the magnitude (a & c). The models also correctly recreate the diurnal cycle of the flux. All models result in same diurnal cycle and differ rather little from each other, but with smaller values and smaller standard deviation than the EC data, but underestimation of the flux magnitude as well as standard deviation is common for all of the models studied (Table 2).

310

The scatter plots (Fig. 6) reveal how all of the models tend to underestimate the EC fluxes of sensible heat by about 15 – 45 % (Table 2). These plots also reveal the tendency of stability adjusted models to result in near-zero flux, while the EC system measures a significantly non-zero flux. Much higher variability in the model agreement is present in the negative flux values than in positive values, which indicates better agreement during unstable boundary layer conditions. Of the three models, SEA-ICE performs slightly better than others during unstable conditions (positive flux values for H), as other models show underestimation in comparison to the EC measurements (Fig. 6). Incorrect flux sign reproduction is much more prevalent in

315



**Figure 6.** Scatter plots of the 30 minute sensible heat flux values of the three included models against corresponding EC measurements. All axes have unit of  $\text{W m}^{-2}$ . Black line indicates a 1:1 fit and red line best linear fit. Linear fit parameters are listed in Table 2.

Bias in turbulent flux direction (upward or downward) is more evident during freezing surface conditions (Figs. 6a, c & e) than in during melting surface conditions (Figs. 6b, d & f).

Taylor plots (Fig. 7) of  $H$  reveal ~~that~~ that the simpler ~~constant~~  $C_H$  model has higher correlation and smaller centered root mean square error (CRMSE) than either of the stability adjusted models when compared with the EC measurements. This is true for all surface conditions, but ~~melting surface results are in~~ better correlation between models and EC data ~~than are found during~~ melting surface conditions.


### 3.2.2 Latent heat flux (LE)

All three models performed the best for LE in freezing surface conditions, with the sign, magnitude and variability all in fairly good agreement with the EC data (Fig. 5b & d) and small variability between the models (Table 3). The LE flux values are also the smallest in these conditions.

Biggest discrepancies between the models and EC data are found during nights in melting surface conditions (Fig. 5b). Here also the largest differences between the models are present. EC data indicates condensation/deposition on the surface for most of the time during nightly melting surface conditions (median of  $-3 \text{ W m}^{-2}$  and values of up to  $-15 \text{ W m}^{-2}$ ). These flux values are almost never reproduced by any of the included models, which typically show a median flux of  $1 \text{ W m}^{-2}$  and peak negative fluxes of  $-5 \text{ W m}^{-2}$ . LHFA model performs slightly better than the other two models when LE is negative, but the variability

**Table 2.** Linear fit parameters ( $ax + b$ ) and statistics of 30 minute sensible heat flux (H) values between EC and model data.

	Slope $a$	Offset $b$ [ $\text{W m}^{-2}$ ]	Standard deviation [ $\text{W m}^{-2}$ ]	CRMSE [ $\text{W m}^{-2}$ ]	Correlation
EC H, $T_0 < 0^\circ\text{C}$	-	-	12.09	-	-
EC H, $T_0 > 0^\circ\text{C}$	-	-	21.30	-	-
H constant $C_H$ , $T_0 < 0^\circ\text{C}$	0.79	-3.29	12.28	8.08	0.78
H constant $C_H$ , $T_0 > 0^\circ\text{C}$	0.86	-1.52	21.27	11.19	0.86
H LHFA $C_H$ , $T_0 < 0^\circ\text{C}$	0.56	-1.69	9.33	8.89	0.68
H LHFA $C_H$ , $T_0 > 0^\circ\text{C}$	0.59	-0.45	15.92	14.16	0.75
H SEA-ICE, $T_0 < 0^\circ\text{C}$	0.68	-2.55	11.37	8.69	0.73
H SEA-ICE, $T_0 > 0^\circ\text{C}$	0.70	-1.62	18.93	13.18	0.79




**Figure 7.** Taylor plots comparing H bulk flux models to EC measurements. Subplot (a) has all the cases where the surface was freezing ( $T_0 < 0^\circ\text{C}$ ), and in subplot (b) are the cases where the surface was melting ( $T_0 > 0^\circ\text{C}$ ). A denotes EC measurements, B is the constant  $C_H$  model, C denotes the LHFA model and D stands for the SEA-ICE model. Green lines represent isolines of root mean square error (RMSE), blue lines represent isolines of correlation and dashed black lines are the isolines of standard deviation.

and the magnitude of the fluxes are smaller than the EC measurements. During daytime the range of EC flux values is two to four times greater than the models, with the greatest standard deviation found in the SEA-ICE model and least found in LHFA. The LHFA model has consistently a lower median and smaller ~~variance~~ standard deviation than the two other models in freezing surface conditions (Figs. 5b & d). The constant  $C_E$  model and the SEA-ICE model have very similar performance.

Scatter plots (Fig. 8) reveal that similar pattern of very low bulk fluxes are given when the EC system reports significantly non-zero fluxes. Also, the flux sign is not correctly reproduced sometimes, regardless of surface temperature. The SEA-ICE model results in best linear fit (Table 3), with the static model performing almost as well and LHFA resulting in the most underestimation.






**Figure 8.** Scatter plots of the three included models against latent heat flux EC measurements. Black line indicates a 1:1 fit and red line best linear fit. All axes have unit of  $\text{W m}^{-2}$ . Linear fit parameters are listed in Table 3

**Table 3.** Linear fit parameters ( $ax + b$ ) and statistics of 30 minute latent heat flux (LE) values between EC and model data.

	Slope $a$	Offset $b$ [ $\text{W m}^{-2}$ ]	Standard deviation [ $\text{W m}^{-2}$ ]	CRMSE [ $\text{W m}^{-2}$ ]	Correlation
EC LE, $T_0 < 0^\circ\text{C}$	-	-	9.95	-	-
EC LE, $T_0 > 0^\circ\text{C}$	-	-	14.99	-	-
LE constant $C_E$ , $T_0 < 0^\circ\text{C}$	0.90	3.39	10.03	4.73	0.89
LE constant $C_E$ , $T_0 > 0^\circ\text{C}$	0.84	4.53	15.30	8.98	0.82
LE LHFA, $T_0 < 0^\circ\text{C}$	0.70	2.65	8.18	5.13	0.86
LE LHFA, $T_0 > 0^\circ\text{C}$	0.58	3.00	10.85	9.78	0.76
LE SEA-ICE, $T_0 < 0^\circ\text{C}$	0.95	2.75	10.74	5.19	0.88
LE SEA-ICE, $T_0 > 0^\circ\text{C}$	0.86	5.01	16.43	10.48	0.78

340 Taylor plots (Fig. 9) of LE show that the three models perform quite similarly in regards to correlation and root mean square error, with greatest differences visible in the standard deviation of the fluxes. By these numbers not one model is clearly better than the others, and they all share the same property of losing some of their predictive ability when the surface is melting. Constant  $C_E$  model (Fig. 7b) has a slight advantage in all statistics presented in the Taylor plot over the other two models.



**Figure 9.** Taylor plots comparing LE flux models to EC measurements. Subplot (a) has all the cases where the surface was freezing ( $T_0 < 0^\circ\text{C}$ ), and in subplot (b) are the cases where the surface was melting ( $T_0 > 0^\circ\text{C}$ ). *A* denotes EC measurements, *B* is the constant  $C_H C_E$  model, *C* denotes the LHFA model and *D* stands for the SEA-ICE model.


### 3.3 Correlation analysis

345 The behaviour of the turbulent heat fluxes was also studied by calculating their correlation and centered root mean square error (CRMSE) as a function of corresponding meteorological variables. This analysis revealed two conditions, where the models have difficulties: low correlation when the temperature/humidity difference between the air and the surface is small and low correlation and high CRMSE during low wind speed.

Figures 10a-e-a-c show the correlation and error for sensible heat flux as a function of temperature difference between surface and air ( $T_0 - T_a$ ). A clear depression in correlation can be observed at  $\pm 0.5$  K temperature difference for all models. Error for all models is at its lowest then (CRMSE  $< 5$  W m $^{-2}$ ), as sensible heat flux values themselves are small then as well.

Figures 10d-f-d-f shows the correlation and CRMSE of latent heat flux against the difference of specific humidity between surface and air. Similar to the temperature difference, low correlation is observed below  $0.2-0.2 \cdot 10^{-3}$  kg kg $^{-1}$  humidity difference. It can be seen that the correlation remains at around 0 for negative humidity differences as well, i.e. in cases where deposition of water/ice should occur over the lake ice surface. This behaviour was noted previously in Fig. 8 as the inability of the models to reproduce the scale of nighttime negative EC flux values of LE.

The second type of cases with both high error and low correlation can be found in for low wind speed cases ( $U \leq 2$  m s $^{-1}$ ) shown in Figs. 10g and 10h. Bulk models always result in low fluxes in these cases, which follows from the linear dependency of the bulk flux on the wind speed (Eqs. (3) and (4)), but EC measures sometimes relatively high fluxes of around  $\pm 20$  W m $^{-2}$  in these conditions. The expected result would have been a similar situation as was with the temperature and humidity: in low wind conditions correlation drops as the absolute value of the flux drops near zero. In these previously described cases, CRMSE would also remain small. Figure 10h shows that while correlation drops significantly in low wind conditions, but that CRMSE increases when compared to cases with higher wind speeds between 2–6 m s $^{-1}$  (Fig. 10g). This behaviour is present in all stability conditions.




**Figure 10.** The centered root mean square error (CRMSE) (top row) and correlation (middle row) of H (solid line) and LE (dashed line) for 100 equidistant bins of temperature difference, humidity difference and wind speed for each of the three models (red-blue for constant  $C_H/EC_{H/E}$ , blue-red for LHFA and black for SEA-ICE). Lowest row of figures are histograms of temperature and humidity difference and wind speed values.

### 365 3.4 Transfer coefficients

The behaviour of the transfer coefficients for heat ( $C_H$ ) and humidity ( $C_E$ ), estimated from observations and models, was studied in two ways: first by plotting the neutral values ( $-0.1 < zL^{-1} < 0.1$ ) of transfer coefficients as a function of wind speed (Fig. 11), and then by investigating how the normalized transfer coefficients depend on atmospheric stability (Fig. 12).

Neutral values of transfer coefficients, estimated from observations (EC), were characterized by similar patterns in their wind speed dependence (Fig. 11). At high wind speeds ( $U > 2 \text{ m s}^{-1}$ ),  $C_{HN}$  and  $C_{EN}$  remained at relatively constant values of  $1.9 \cdot 10^{-3}$  and  $1.3 \cdot 10^{-3}$ , respectively, which are in relatively good agreement with model values. Both transfer coefficients increased towards the lowest wind speeds. The strongest increase was found for  $C_{HN}$ , which was more than twice higher ( $4.8 \cdot 10^{-3}$ ) at the lowest wind speed bin. A similar, but less pronounced increase was observed for  $C_{EN}$ , whose value at the lowest wind speed bin was  $2.9 \cdot 10^{-3}$ . Both of the dynamic models (LHFA and SEA-ICE) fail to reproduce this increase during low wind speed, showing nearly constant values for low and high wind speed ranges (Table 4).

The dependency of transfer coefficients, normalized by their neutral values ( $-0.1 < zL^{-1} < 0.1$ ), on stability parameter  $zL^{-1}$  is presented in Fig. 12. Transfer coefficients for H, calculated from measurements, show slightly higher values than the




**Figure 11.** Dependency of neutral transfer coefficients ( $-0.1 < zL^{-1} < 0.1$ ) as a function of wind speed. Subplots a) and b) show this dependency for heat ( $C_{HN}$ ) and evaporation ( $C_{EN}$ ), as estimated from models (LHFA and SEA-ICE) and EC observations (EC), respectively. The error bars indicate the standard deviation in the bins. Subplots c) and d) show the distribution of neutral conditions as a function of wind speed.  $C_{HN}$  data was filtered for small temperature difference ( $|T_0 - T_a| < 0.5$  K) and  $C_{EN}$  data was filtered for very small and negative values of humidity difference ( $q_0 - q_a < 2 \cdot 10^{-4}$   $\text{kg kg}^{-1}$ ). The 30 min values of coefficients were also not included in case of very low wind speed ( $U < 0.5$   $\text{m s}^{-1}$ ). Bins with less than 10 values were not included. The values are for the measurement height of 1.8 m.

**Table 4.** Neutral ( $-0.1 < zL^{-1} < 0.1$ ) transfer coefficients estimated at  $z = 1.8$  m.

Model / measurement	Transfer coefficient	Transfer coefficient ( $U > 2$ $\text{m s}^{-1}$ )	Transfer coefficient ( $U < 2$ $\text{m s}^{-1}$ )
$C_{HN}$ EC	$1.9 \cdot 10^{-3}$	$1.9 \cdot 10^{-3}$	$3.0 \cdot 10^{-3}$
$C_{HN}$ LHFA	$1.4 \cdot 10^{-3}$	$1.4 \cdot 10^{-3}$	$1.4 \cdot 10^{-3}$
$C_{HN}$ SEA-ICE	$1.8 \cdot 10^{-3}$	$1.8 \cdot 10^{-3}$	$1.7 \cdot 10^{-3}$
$C_{EN}$ EC	$1.0 \cdot 10^{-3}$	$1.0 \cdot 10^{-3}$	$1.5 \cdot 10^{-3}$
$C_{EN}$ LHFA	$1.4 \cdot 10^{-3}$	$1.4 \cdot 10^{-3}$	$1.4 \cdot 10^{-3}$
$C_{EN}$ SEA-ICE	$1.8 \cdot 10^{-3}$	$1.8 \cdot 10^{-3}$	$1.8 \cdot 10^{-3}$

380

model ones, and they do not follow the expected dependency on stability, as  $C_H/C_{HN}$  estimated from EC measurements does not follow the expected decreasing trend for  $zL^{-1} > 0$ . During unstable conditions  $C_H/C_{HN}$  values acquired by EC show increasing trend, but on average have higher values than the bulk models.  $C_E$  determined from measurements shows good agreement during unstable conditions and increasingly higher values than the models towards higher stability.



**Figure 12.** Normalized and binned values of transfer coefficients as a function of stability parameter for  $C_H$  (a) and  $C_E$  (b).  $C_H$  data was filtered for small temperature difference ( $|T_0 - T_a| < 0.5$  K) and  $C_E$  data was filtered for very small and negative values of humidity difference ( $q_0 - q_a < 2 \cdot 10^{-4}$  kg kg $^{-1}$ ). The 30 min values of coefficients were also not included in case of very low wind speed ( $U < 0.5$  m s $^{-1}$ ). Bins were rejected if they had less than 10 values within them. Subplots c) and d) correspondingly present the distribution of transfer coefficient values as a function of stability parameter.

#### 4 Discussions & conclusions

Turbulent heat fluxes were studied with an EC setup for four winters over the ice cover of a boreal lake and these results were compared to three bulk aerodynamic models, one that does not take into account the atmospheric boundary-layer stability and two that do take it into account. Since EC measurement sites are much more seldom used than are regular weather stations, bulk transfer models are still going to be used in numerical weather prediction and general circulation models for the foreseeable future. Our data set spanning four ice-on seasons provided a good opportunity to verify and compare the accuracy of bulk transfer models over direct measurements of turbulent heat fluxes by an EC setup on seasonal lake ice cover.

Both, the bulk transfer models and the EC setup function the best. The best agreement between measured and modelled fluxes of H and LE was found for cases with high wind speeds and large gradients of speed and large water-air temperature and humidity, as these conditions give the best agreement between models and EC differences. Lake ice surface is a challenging environment for eddy covariance due to the relatively low amount of turbulence in the air above it. This is due to several reasons: the boundary layer is stable for most of the time during winter which leads to underdeveloped turbulence and decoupling of the flow from the surface, the surface has a very low roughness and fluxes are usually low. Despite these problems-challenges, the EC setup was able to record good quality flux values in a wide range meteorological states, and the data coverage was

sufficient. ~~An unfortunate property of the bulk flux calculation was the fact that it functioned the best when the atmospheric boundary layer was unstable, which is the minority of cases studied here.~~

The bulk aerodynamic method is technically and computationally much simpler than the eddy covariance method, but it comes with some limitations. The greatest error producing effect can be attributed to the fact that estimating the skin temperature of a snowy surface is difficult, which has been previously reported as a major issue in modeling turbulent heat fluxes over snow and ice (??). This is due to melting and refreezing and the consequent horizontally heterogeneous changes in the surface properties, like albedo, emissivity and phase (liquid or frozen). Errors in determination of the surface temperature affect both sensible and latent heat flux computation especially when the difference between air and surface is small. It results in incorrect surface humidity values, which is an exponential function of the surface temperature and thus very sensitive to errors. Also, the emissivity  $\epsilon$  of the ice/snow surface is difficult to determine accurately, and it can change with the metamorphosis of snow (?), which happens constantly over the course of the winter. Thus, the calibration and proper installation of especially the long wave radiation sensors is very important in campaigns performed over ice or snow. Due to the horizontally and vertically heterogeneous nature of the surface, point measurements performed in one location, like net radiation, are not always representative of the whole lake and there is a possibility that very biased results for the surface albedo and outgoing long wave radiation are recorded, especially during the melting period. This partially explains differences in EC and bulk flux results, as the footprint of EC measurements is at least an order of magnitude larger than the source area measured by the radiation sensors.

~~Second issue noticed with the bulk algorithms were the low correlation and high error in low wind conditions, which can be explained by non-local effects on turbulence above the lake. In ? it was shown that a sharp decrease in aerodynamic roughness, like the transition between dense forest and lake ice commonly found on boreal lakes, can cause significant fluxes on EC measurements while bulk algorithms show very low values. When the wind blows from the forest towards the lake, a significant increase in turbulent kinetic energy is observed near the center of the lake, and heat and moisture are transported from the upper boundary layer towards the surface. Bulk algorithms are not taking this phenomenon into account, and thus fail to reproduce these situations. On Lake Kuivajärvi this non-local effect on turbulent fluxes can be observed at all wind directions due to the small size and strongly elongated shape of the lake and the steep shoreline.~~

Similar behaviour of the models ~~has studied here have~~ been observed in previous studies. In ? it was noted that LHFA and SEA-ICE models tended to underestimate and result in lower standard deviation than the turbulent heat fluxes acquired by EC over a Siberian thermokarst lake, which is in line with the results of ~~this our~~ study. Correlation of these models ranged between 0.7 and 0.9 over the thermokarst lake, which are similar to our findings.

In a study conducted over landfast sea ice (?), the bulk transfer models were noticed to underestimate negative fluxes, but unlike our results, were found to overestimate positive fluxes for both H and LE. Correlation coefficients were found to be slightly larger (0.88) for LE than for H (0.82), which is similar to our results. ~~What was also reported in this article is the~~ The same study reported better accuracy of models with a constant transfer coefficient over dynamic coefficients in the winter ~~— spring transition period, although in general they observed the dynamic model was observed to perform marginally better. In our study the slight advantage of the static model was present for both fluxes and during winter and spring performing marginally better over the static one.~~ Differentiation between frozen and melting surface was not performed in the aforementioned studies,

but the results of ~~this-our~~ study indicate that modeling of H works better in melting conditions, while the opposite holds for LE.

435 ~~In conclusion, it can be said that in general bulk transfer models are able to reproduce turbulent heat fluxes measured with an EC setup over seasonal lake ice cover for most of the time with relatively little difference between the models tested. In lack of more accurate measurements of the surface temperature it might not be worth the effort to use stability-adjusted models. While they are more physically sound than the certainly incorrect assumption made in the static model, additional errors are introduced into the flux values from the stability calculations. Turbulent~~ Issues with the bulk algorithms were noticed as low correlation and high error in low wind speed conditions, which can possibly be explained by non-local effects on turbulence above the lake. In ? it was shown that a sharp decrease in aerodynamic roughness, like the transition between dense forest and  
440 ~~lake ice commonly found on boreal lakes, can cause significant fluxes on EC measurements while bulk algorithms show very low values. When the wind blows from the forest towards the lake, a significant increase in turbulent kinetic energy is observed near the center of the lake, and heat and moisture are transported from the upper boundary layer towards the surface. Bulk algorithms are not taking this local violation of MOST into account, and thus fail to reproduce these situations.~~

445 ~~Discrepancies at low wind speed can also be seen when studying the neutral transfer coefficients as a function of wind speed. Dependency of neutral transfer coefficient values on wind speed have been studied previously during open water season, most recently and applicably on lake Kasumigaura in Japan (?). In this study, observations similar to ours were made, namely that the transfer coefficients increase during low wind speed conditions ( $U < 3 \text{ m s}^{-1}$ ). This effect has been attributed to several possible mechanisms, but the study performed over lake Kasumigaura showed that capillary waves and the averaging method used for the calculation of mean wind speed only had a small effect in the increase of neutral transfer coefficients towards~~  
450 ~~lower wind speed. Larger effects were noted to be most likely caused by increase in turbulent kinetic energy over the lake in low wind speed conditions. Our data is in agreement, as in our case the coefficients increase in the same range of wind speed, and any effect of waves and underlying currents in the water can be excluded as possible reasons due to the ice cover.~~

455 ~~Neutral values in high wind speeds were all within the same range between the bulk models and EC. In high wind speed conditions ( $U > 2 \text{ m s}^{-1}$ ), neutral transfer coefficient for heat and water vapor calculated from EC observations is very close to the values estimated from models, with slightly lower values for evaporation than for heat, which indicates dissimilarity between temperature and humidity.~~

~~The fact that EC observations were only made on one level also somewhat limited our ability to perform analysis regarding, for example the roughness length over the lake ice cover or the applicability of the logarithmic profile for wind speed and scalars. With multiple levels of measurement cases where MOST is locally violated could be identified much better.~~

460 ~~Although the stability corrected bulk transfer models have a more sound physical and theoretical basis than an uncorrected static model has, it is possible to get lower agreement between EC and dynamic models than between EC and static models. This is not to say that the stability corrected models are inherently wrong, but that conditions violating MOST are not uncommon over lakes. Thus, although generally good, turbulent~~ heat flux values obtained by bulk transfer models on very small lakes surrounded by forest ~~should be interpreted with caution,~~ are generally underestimating the flux as the heterogeneity of the

465 surrounding environment ~~causes errors in the models~~ and measurement errors over the complex and dynamic ice cover causes  
incorrect output of the models in certain conditions.

*Code availability.* The Lake Heat Flux Analyzer (version 1.1.0, released May 19, 2015) is available on Zenodo:

<https://doi.org/10.5281/zenodo.5534907>

SEA-ICE (version 2.0, released November 13, 2014) is available on Zenodo:

470 <https://doi.org/10.5281/zenodo.5534911>

*Data availability.* All data used in this article can be downloaded from the University of Helsinki Aava SmartSMEAR database  
(<https://smear.avaa.csc.fi/>).

DOI identifier of the Kuivajärvi data set: doi:10.23729/9b209b52-2ea0-4d89-b059-062b734142d8

DOI identifier for the Hyytiälä forest data set: doi:10.23729/2001890a-2f0b-4e37-8c70-4d2cb5f40273

475 *Author contributions.* IM designed the study and supervised the research, JAK performed the data analysis and wrote the article, KMK  
performed the data processing, ML supported data analysis and results interpretation, All authors provide comments to the manuscript.

*Competing interests.* There are no competing interests present in this study.

*Acknowledgements.* The authors thank ICOS-Finland (3119871), ACCC Flagship (337549) and N-PERM project (341348) funded by the  
Academy of Finland, ~~the Tyumen region government in accordance with the Program of the World-Class West Siberian Inter-regional~~

480 ~~Scientific and Educational Center (National Project “Nauka”) and the European Commission H2020 Research Project RINGO (730944).~~



## References