The Effects of Ocean Surface Waves on Global Intraseasonal Prediction: Case Studies with a Coupled CFSv2.0-WW3

Ruizi Shi[1], Fanghua Xu[1*], Li Liu[1], Zheng Fan[1], Hao Yu[1], Hong Li[1,3], Xiang Li[2] and Yunfei Zhang[2]

[1] Ministry of Education Key Laboratory for Earth System Modeling, and Department of Earth System Science, Tsinghua University, Beijing, 100084, China

[2] Key Laboratory of Marine Hazards Forecasting, National Marine Environmental Forecasting Center, Ministry of Natural Resources, Beijing, 100081, China

[3] Department of Atmospheric and Oceanic Sciences and Institute of Atmospheric Sciences, Fudan University, Shanghai, 200433, China.

Correspondence to: Fanghua Xu (fxu@mail.tsinghua.edu.cn)

Abstract. This article describes the implementation of a coupling between a global forecast model (CFSv2.0) and a wave model (WW3), and investigates the effects of ocean surface waves on air-sea interface in the new framework. Several major wave-related processes, including the Langmuir mixing, Stokes-Coriolis force with entrainment, air-sea fluxes modified by Stokes drift and momentum roughness length, are evaluated in two groups of 56-day experiments, one for boreal winter and the other for boreal summer. Comparisons are made against in-situ buoys, satellite measurements and reanalysis data, to evaluate the influence of waves on intraseasonal prediction of sea surface temperature (SST), 2-m air temperature (T02), mixed layer depth (MLD), 10-m wind speed (WSP10) and significant wave height (SWH). The wave-coupled experiments show that overestimated SSTs and T02s, as well as underestimated MLDs at mid-high latitudes in summer from original CFSv2.0 are significantly improved due to enhanced vertical mixing generated by Stokes drift. For WSP10s and SWHs, the wave-related processes generally lead to reduction of biases in regions where WSP10s and SWHs are overestimated. On one hand, the decreased SSTs stabilize marine atmospheric boundary layer, weaken WSP10s and then SWHs. On the other hand, the increased roughness length due to waves leads to reduction in the originally overestimated WSP10s and SWHs. In addition, the effects of Stokes drift and current on air-sea fluxes also rectify WSP10s and SWHs. These cases are helpful for the future development of the two-way CFSv2.0-wave coupled system.

1 Introduction

Ocean surface gravity waves play an important role in modifying physical processes at the atmosphere–ocean interface, which can influence momentum, heat and freshwater fluxes across the air-sea interface (Li and Garrett 1997; Taylor and Yelland, 2001; Moon et al., 2004; Janssen 2004; Belcher et al., 2012; Moum and Smyth, 2019). For instance, ocean surface waves modify ocean surface roughness to influence the marine atmospheric boundary layer and thus change the momentum, latent heat, and sensible heat transfer (Janssen 1989, 1991; Taylor and Yelland, 2001; Moon et al., 2004; Drennan et al., 2003, 2005). The breaking waves inject turbulent kinetic energy in the upper ocean, which enhances the mixing process (Terray et al. 1996). Nonbreaking surface waves also affect mixing in the upper ocean by adding a wave-related Reynolds stress (Qiao et al., 2004; Ghantous and Babanin, 2014). The wave-related Stokes drift interacts with Coriolis force and produces the Coriolis-Stokes force (Hasselmann 1970). The shear of Stokes drift is critical for generation of Langmuir circulation, which significantly deepens the mixed layer by strong vertical mixing both at climate scales (Li and Garrett 1997; Belcher et al., 2012) and at weather scales (Kukulka et al., 2009).

Various wave-related parameterizations have been proposed and applied in modeling. The wave-related Charnock parameter (C_{ch}) defines sea surface roughness and affects wind stress estimates (Pineau-Guillou et al. 2018; Sauvage et al. 2020). There are primarily three methods for C_{ch}, assessed from the wave-induced kinematic stress (Janssen 1989, 1991), the wave age (Drennan et al., 2003, 2005; Moon et al., 2004), or the steepness (Taylor and Yelland, 2001). The former two are based on the wind-sea conditions, whereas the latter includes both swells and wind-sea waves. Modifications to these Charnock parameterizations were suggested in recent studies for the leveling off roughness under high winds (e.g. Fan et al., 2012; Bidlot et al., 2020; ECMWF, 2020; Li et al., 2021). In the oceanic boundary

56 layer, waves influence upper ocean mixing via wave dissipation and Stokes drift-induced processes. In

57 Breivik et al. (2015), the wave dissipation-related turbulent kinetic energy flux is found to yield the

58 largest sea surface temperature (SST) differences in the extratropics. The Stokes drift-induced Langmuir

59 turbulence can improve temperature simulation over most of the world oceans, particularly in the

60 Southern Ocean (Belcher et al., 2012; Li et al., 2016). Polonichko (1997), Van Roekel et al. (2012) and

61 Li et al. (2017) indicated that the Langmuir cell intensity strongly depends on the alignment of winds

62 and waves, reaching maximum when they are aligned. Li et al. (2016) found the effect of Langmuir cell

63 can be further enhanced by entrainment. In Couvelard et al. (2020), the Stokes drift–related forces can

64 also contribute to deepening of the mixed layer depth (MLD) modestly. In the First Institute of

65 Oceanography Earth System Model, Bao et al. (2019) indicated that the non-breaking wave-induced

66 mixing, Stokes drift-affected air-sea fluxes, as well as sea spray are all important for climate estimates.

67 The wave-related processes at the air-sea interface are complex and important in global coupled

68 systems (e.g., Breivik et al. 2015; Law-Chune and Aouf, 2018; Bao et al. 2019; Couvelard et al. 2020).

69 Most of the coupled models with a wave component at global scale were developed for climate research

70 (e.g., Law-Chune and Aouf, 2018; Bao et al. 2019; Couvelard et al. 2020). Exceptionally, an Integrated

71 Forecasting System (IFS) with fully coupled atmosphere, ocean and wave components, developed by

72 European Centre for Medium-Range Weather Forecasts (ECMWF) (Janssen 2004; Bidlot et al. 2019,

73 2020), has been released with great flexibility for global forecasts from medium-range weather scales to

74 seasonal scales (Breivik et al. 2015).

75 The effects of wave-related processes are worth further evaluation in different global coupled

76 modelling systems. Since it takes sufficient periods for the wave energy to develop (Janssen 2004), we

77 investigate the impact of individual wave process at intraseasonal timescale in a new global atmosphere-

78   ocean-wave system. To achieve this, we coupled the WAVEWATCH III (WW3) to the Climate Forecast

79   System model version 2.0 (CFSv2.0) and then conducted sensitivity experiments in boreal winter and

80   summer for comparison. The effects of upper ocean mixing modified by Langmuir cell, Stokes-Coriolis

81   force and entrainment, air-sea fluxes modified by surface current and Stokes drift, and momentum

82   roughness length are evaluated. The CFSv2.0 is a coupled system with main application for intraseasonal

83   and seasonal prediction (e.g. Saha et al. 2014). The National Centers for Environmental Prediction

84   (NCEP) is establishing its own atmosphere-ocean-wave system, in which the Global Forecast System

85   (GFS; the atmosphere module in CFSv2.0 system) is one-way coupled with WW3. Our work can provide

86   insights for two-way wave coupling of CFSv2.0, and is helpful for the future development of the

87   CFSv2.0-wave coupling system. Two groups of 56-day predictions were conducted for boreal winter and

88   boreal summer, respectively. The predictions were then compared with observations and reanalysis data.

89   For each group, sensitivity experiments with different wave parameterizations were carried out to

90   evaluate the effects of individual wave-related process.

91   The rest of the paper is structured as follows: methods and numerical experiments with different

92   parameterizations are described in Section 2; the observations and reanalysis data are introduced in

93   Section 3, and the results of experiments are evaluated and compared in Section 4. Finally, a summary

94   and discussion are given in Section 5.

95   **2 Methods and Experiments**

96   **2.1 Coupling WAVEWATCH III with CFSv2.0**

97   The version 5.16 of WW3 (WAVEWATCH III Development Group, 2016) developed by the National

98   Oceanic and Atmospheric Administration (NOAA)/NCEP has been incorporated into the CFSv2.0 (Saha

99 et al., 2014) as a new model component. The latitude range of WW3 is 78°S–78°N with a spatial

100 resolution of 1/3°; the frequency range is 0.04118-0.4056Hz and the total number of frequencies is 25;

101 the number of wave directions is 24 with a resolution of 15°; the maximum global time step and the

102 minimum source term time step are both 180 s.

103 The CFSv2.0 contains two components, the GFS (details are available at

104 http://www.emc.ncep.noaa.gov/GFS/doc.php) as the atmosphere component and the Modular Ocean

105 Model version 4 (MOM4; Griffies et al., 2004) as the ocean component. The MOM4 is integrated on a

106 nominal 0.5° horizontal grid with enhanced horizontal resolution to 0.25° in the tropics, and has 40

107 vertical levels; the vertical spacing is 10 m in the upper 225 m, and then increases in unequal intervals to

108 the bottom at 4478.5 m. A three-layer sea ice model is included in MOM4 (Wu et al. 2005). The GFS

109 uses a spectral triangular truncation of 382 waves (T382) in the horizontal, which is equivalent to a grid

110 resolution of nearly 35 km, and 64 sigma-pressure hybrid layers in the vertical. The time steps of both

111 MOM4 and GFS are 180 s. The ocean and atmosphere components are then coupled at the same rate. In

112 the original two-way coupled system, the GFS receives SST from MOM4 and sends fluxes of heat,

113 momentum, freshwater to MOM4 (black arrows in Fig. 1).

114 The Chinese Community Coupler version 2.0 (C-Coupler2; Liu et al., 2018) is applied to interpolate

115 and pass variables between atmosphere and wave components as well as ocean and wave components.

116 Each component receives inputs and supplies outputs on its own grids. The C-Coupler2 is a common,

117 flexible and user-friendly coupler, which contains a dynamic 3-D coupling system and enables variables

118 to remain conserved after interpolation.

119 A schematic diagram of the coupled atmosphere-ocean-wave system is shown in Fig. 1. As illustrated,

120 WW3 is two-way coupled with MOM4 and GFS, through the C-Coupler2. WW3 is forced by 10-m wind

121   from GFS (green arrows) and surface current from MOM4 (blue arrows), and then generates and evolves

122   the wave action density spectrum. Meanwhile, the surface Stokes drift velocity, the Stokes transport and

123   the turbulent Langmuir number are passed to MOM4 (red arrows; see Section 2.3) from WW3, and the

124   surface Stokes drift velocity and the Charnock parameter are passed to GFS (red arrows; see Section 2.4

125   and 2.5). The high frequency tail assumption for Stokes drift in WW3 is used with a spectral level

126   decaying as f^{-5} (frequency). Additionally, the regular ocean surface current velocities from MOM4 are

127   also passed to GFS, to calculate the relative wind velocity for the turbulent fluxes together with surface

128   Stokes drift (blue arrows; see Section 2.4).

129   Both the CFSv2.0 and WW3 use warm starts; the initial fields at 00:00 UTC of the first day in each

130   experiment for CFSv2.0 were generated by the real time operational Climate Data Assimilation System

131   (Kalnay   et   al.,   1996),   downloaded   from   the   CFSv2.0   official   website

132   (http://nomads.ncep.noaa.gov/pub/data/nccf/com/cfs/prod). To get initial conditions for WW3, a stand-

133   alone WW3 model is set up synchronously (see Section 2.2). Since the interactions between waves and

134   sea ice are complicated and beyond the scope of the study, we turn off the coupling between WW3 and

135   CFSv2.0 in areas with sea ice.

136   In addition, to properly select the coupling frequency between CFSv2.0 and WW3, the root mean

137   square errors (RMSEs) of SST, significant wave height (SWH) and 10-m wind speed (WSP10) with

138   different coupling steps for the fully coupled experiment (ALL; details in Section 2.6) are calculated and

139   compared (Table S1 of the supplementary). From Table S1, the 10_STEP_WW3 experiment has a

140   relatively short runtime and small RMSEs. Therefore, the time steps of the 10_STEP_WW3 are selected

141   to compromise computing time consumption and model RMSEs.

2.2 Initialization of WAVEWATCH III

In WW3, input of momentum and energy by wind, and dissipation for wave-ocean interaction are two important terms (combined as input-dissipation source term) in the energy balance equation (WAVEWATCH III Development Group, 2016), which include the Charnock parameter related estimation. Several different packages to calculate the input-dissipation source term (ST) are offered in the WW3 version 5.16, including ST2 (Tolman and Chalikov, 1996), ST3 (Janssen, 2004; Bidlot, 2012), ST4 (Ardhuin et al., 2010), and ST6 (Zieger et al., 2015).

The initial wave fields were generated from 10-day simulation starting from rest in a stand-alone WW3 model. To minimize the biases of initial wave fields, we tested simulations with ST2, ST3, ST4, and ST6 schemes respectively, and compared the results with Janson-3 observations. Two 10-m wind datasets, the Cross-Calibrated Multi-Platform (CCMP; Atlas et al., 2011) data and the fifth generation European Centre for Medium-Range Weather Forecasts (ECMWF) Reanalysis (ERA5; Hersbach et al., 2020) data, were used to drive the wave model respectively. Compared all results, the ST4 scheme with ERA5 wind forcing generates the minimum SWH bias (Table S2 in the supplementary), consistent with findings in Stopa et al. (2016). Thus, the ST4 scheme was chosen to calculate the input and dissipation term, and generate initial wave fields with ERA5 wind forcing for experiments listed in Table 1. The parameters used for ST4 scheme followed TEST471f from WAVEWATCH III Development Group (2016), which is the CFSR (CFS Reanalysis) tuned setup and is commonly-used at global scale.

2.3 Parameterizations of Stokes Drift-Related Ocean Mixing

The full Stokes drift profile used in MOM4 is obtained by the method of Couvelard et al. (2020), which is based on the work of Breivik et al. (2014; 2016). Breivik et al. (2016) derived the full Stokes

163   drift profile as

$$u_sB16(z) = u_s(0)[\exp(2k_pz) - \sqrt{-2\pi k_pz}\,\text{erfc}(\sqrt{-2k_pz})], \qquad (1)$$

164   The $u_s(0)$ is the surface Stokes drift velocity, $k_p = \frac{u_s(0)}{6V_s}$, V_s is the Stokes transport, and erfc is the

165   complementary error function. Eqn. 1 is depth-averaged within each vertical grid interval as

$$u_s(z) = \frac{u_s(0)}{(th)_k}[I(z_{k+1/2}, k_p) - I(z_{k-1/2}, k_p)], \qquad (2)$$

$$I(z, k_p) = \frac{1}{6k_p}[e^{2k_pz} + 4k_pz\frac{u_sB16(z)}{u_s(0)}], \qquad (3)$$

166   where *th* is the thickness of layer *k*, following Li et al., (2017), Wu et al., (2019) and Couvelard et al.,

167   (2020).

168   **2.3.1 Mixing of Langmuir Turbulence**

169   McWilliams and Sullivan (2000) modified the turbulent velocity scale *W* in K-Profile Parameterization

170   (KPP) for vertical mixing by introducing an enhancement factor ε, to account for both boundary layer

171   depth changes and nonlocal mixing by Langmuir turbulence. Based on their work, Van Roekel et al.

172   (2012) improved the enhancement factor corresponding to alignment and misalignment of winds and

173   waves. Li et al. (2016) evaluated these parameterizations in a coupled global climate model, and found

174   that the difference between parameterizations with alignment and with misalignment was not significant,

175   owing to the relatively coarse resolution which cannot accurately represent the refraction by coasts and

176   current features. We applied the parameterization from Van Roekel et al. (2012) as well. Because the

177   resolution in our model is relatively coarse too, and the angles between winds and waves are less than

178   30° in most areas (Fig. S1i&j in the supplementary), we didn't consider misalignment in the study.

179   *W* ($W=ku_*/\phi$, where u_* is the surface friction velocity, ϕ is the dimensionless flux profile, and

180   *k*=0.4 is the von Kármán constant) varies in proportion to the turbulent Langmuir number, that is,

$$W = \frac{ku_*}{\phi}\varepsilon, \tag{4}$$

$$\varepsilon = \sqrt{1 + (3.1La_t)^{-2} + (5.4La_t)^{-4}}, \tag{5}$$

181   where La_t is the turbulent Langmuir number, defined as

$$La_t = \sqrt{\frac{u_*}{|u_s(0)|}}, \tag{6}$$

182   with $u_s(0)$ is the surface Stokes drift velocity.

183   Furthermore, the enhanced W will influence the calculation of boundary layer depth. In KPP the

184   boundary layer depth is determined as the smallest depth at which the bulk Richardson number equals

185   the critical value $Ri_{cr} = 0.3$, that is,

$$Ri_b(h) = \frac{gh[\rho_r - \rho(h)]}{\rho_0[|u_r - u(h)|^2 + W^2]} = Ri_{cr}, \tag{7}$$

186   where g is acceleration of gravity, ρ is density, u is velocity, ρ_r is surface density, u_r is surface

187   velocity, ρ_0 is an average value and h is the boundary layer depth. Hence, when W is enhanced, the

188   boundary layer depth h is deepened accordingly.

189   **2.3.2 Stokes–Coriolis Force and Associated Entrainment**

190   Because the Stokes drift velocity is an increment superimposed on the original current velocity, the

191   Coriolis force and the Stoke drift together produce an additional so-called Stokes–Coriolis (SC) force

192   (Hasselmann 1970), that is,

$$SC\ Force = \overrightarrow{u_s(z)} \times f\vec{z}. \tag{8}$$

193   Here $\overrightarrow{u_s(z)}$ is the Stokes drift velocity vector, f is the Coriolis frequency, and $\vec{z}$ is the vertical unity

194   vector. For consistency, the Stokes drift velocity is also included in advection terms of tracers (e.g.

195   temperature, salinity) and convergence terms (Law-Chune and Aouf, 2018; Couvelard et al., 2020). And

196   the free surface condition for barotropic mode is correspondingly modified to

$$\frac{\partial \eta}{\partial t} = -\nabla M_{curr} - \nabla M_{st}, \tag{9}$$

197   where η is surface elevation, M_{curr} and M_{st} are the total vertical integral of regular Eulerian current

198   and Stokes drift, respectively.

199   To depict the entrainment below the ocean surface boundary layer induced by Stokes drift, Li et al.

200   (2016) suggested to add the square of surface Stokes drift velocity ($|u_s(0)|^2$) to the denominator of Eqn.

201   7, that is,

$$Ri_b(h) = \frac{gh[\rho_r - \rho(h)]}{\rho_0[|u_r - u(h)|^2 + W^2 + |u_s(0)|^2]} = Ri_{cr}. \tag{10}$$

202   The boundary layer depth h in KPP from Eqn. 10 is then enhanced due to Stokes drift velocity.

203   **2.4 Stokes Drift and Sea Surface Current on Air–Sea Fluxes**

204   At air-sea boundary layer, the momentum flux (τ), sensible heat flux (SH) and freshwater flux (E) are

205   calculated as

$$\tau = \rho_a C_d |\Delta \vec{V}| \Delta \vec{V}, \tag{11}$$

$$SH = \rho_a C_h |\Delta \vec{V}| \Delta \theta, \tag{12}$$

$$E = \rho_a C_e |\Delta \vec{V}| \Delta q, \tag{13}$$

206   where C_d, C_h, C_e are surface exchange coefficients for momentum, sensible heat and freshwater. ρ_a

207   is air density. $\Delta \theta, \Delta q$ are potential temperature and humidity differences between air and sea, and $\Delta \vec{V}$

208   is velocity of air relative to water flow.

209   In CFSv2.0, $\Delta \vec{V}$ is set to be wind speed ($\overrightarrow{U_{wind}}$). However, the effect of ocean surface current should

210   not be ignored. Luo et al. (2005) first indicated that including ocean surface current ($\overrightarrow{U_{surf}}$) improves

211   estimates of τ and subsequent ocean response. Renault et al. (2016) further indicated that the

212   improvements of τ by $\overrightarrow{U_{surf}}$ also feed back into atmosphere. At present, $\Delta \vec{V} = \overrightarrow{U_{wind}} - \overrightarrow{U_{surf}}$ is

213   widely used in coupled ocean-atmosphere models (e.g., Hersbach and Bidlot, 2008; Takatama et al.,

214   2017; Renault et al., 2021). Furthermore, Bao et al. (2019) indicated that as a part of the sea surface water

215   movement with speed magnitude comparable to surface current in mid-high latitudes, the surface Stokes

216   drift $(\overrightarrow{u_s(0)})$ should also be included, that is,

$$\Delta\vec{V} = \overrightarrow{U_{wind}} - \overrightarrow{U_{surf}} - \overrightarrow{u_s(0)}. \qquad (14)$$

217   To account for the effects of Stokes drift velocity, the Eqn. 14 was applied in the coupled experiments

218   (Table 1). To complete the coupling, the corresponding modification of the tridiagonal matrix (Lemarié

219   2015) is also conducted in CFSv2.0. Note that the direction of Stokes drift is generally consistent with

220   10-m wind (Fig. S1i&j in supplementary), but the directions of surface current and 10-m wind are usually

221   with an angle due to Coriolis effect (Fig. S1g&h). Consequently, the effects of $\overrightarrow{U_{surf}}$ and $\overrightarrow{u_s(0)}$ on

222     $\Delta\vec{V}$ depend on the angles between them and $\overrightarrow{U_{wind}}$.

223   **2.5 Parameterizations of Momentum Roughness**

224   In CFSv2.0, the fluxes of momentum, heat, and freshwater are passed from atmosphere to ocean, and

225   the estimates of them are critically important. The fluxes are in part determined by surface roughness

226   length, which can be converted to surface exchange coefficients in Eqn. 11-13.

227   **2.5.1 The Momentum Roughness Length in GFS**

228   In GFS, the momentum roughness length z_0 has two terms. The first term z_{ch} is parameterized by

229   the Charnock relationship (Charnock, 1955) representing wave-resulted sea surface roughness, and the

230   second term z_{vis} is the viscous contribution (Beljaars, 1994) for low winds and smooth surface, that is,

$$z_0 = z_{ch} + z_{vis} = \frac{C_{ch}u_*^2}{g} + \frac{0.11\nu}{u_*}. \qquad (15)$$

231   Here $C_{ch} = 0.014$ is the constant Charnock parameter, ν is the air kinematic viscosity. The relation of

232   z_0 in GFS versus 10-m wind speed is shown in Fig.2 (black line).

233   **2.5.2 The Charnock Relationship Related to Wave State**

234   When ocean surface waves are explicitly considered, the Charnock parameter C_{ch} is not a constant

235   (Janssen 1989, 1991; Taylor and Yelland, 2001; Moon et al., 2004; Drennan et al., 2003, 2005). In the

236   study, we adopted a method developed by Moon et al. (2004), which considered the surface roughness

237   leveling off under extremely high wind speed (Powell et al., 2003; Donelan et al., 2004). Based on

238   observations, Moon et al. (2004) proposed Eqn. 16 to estimate the Charnock parameter by the wave age

239    $\frac{c_{pi}}{u_*}$ (c_{pi} is the peak phase speed of the dominant wind-forced waves) with constant values of a and b

240   changing with 10-m wind speed every 5 m/s in the range of 10 m/s to 50 m/s.

$$C_{ch} = a\left(\frac{c_{pi}}{u_*}\right)^b, \qquad (16)$$

241   To obtain continuous values of a and b, we derive a new relationship (Eqn. 17) to estimate a and b from

242   10-m wind speed U_{10} by fitting the values in Table 1 of Moon et al. (2004),

$$a = \frac{1}{0.1477U_{10}{}^2 - 0.7395U_{10} - 10.9995},$$
$$ \qquad (17)$$
$$b = 1.5661E^{-5}U_{10}{}^3 - 0.002U_{10}{}^2 + 0.1017U_{10} - 1.6182.$$

243   Because the observations in Moon et al. (2004) were obtained under tropical cyclones, Eqn. 17 is used

244   for U_{10}>15 m/s, whereas the original Charnock relationship of WW3 ST4 scheme (Janssen 1989, 1991)

245   is used for $U_{10} \leq 15$ m/s. The revised parameterization is called ST4-M04. Figure S2 in supplementary

246   shows the C_{ch} distribution obtained by Eqn. 16-17. In general small wind direction variations at low

247   latitudes lead to large wave age and thus low C_{ch}. The situation is opposite at mid-high latitudes.

248   The relationships between z_0 and U_{10} in GFS, WW3 ST4 scheme (Janssen 1989, 1991) and ST4-

249 M04 scheme were compared in Fig.2. The z_0 in GFS increases relatively slowly with increasing wind

250 speed (black). The value of z_0 from ST4 scheme (purple) increases rapidly with wind speed at high

251 winds. In comparison, in ST4-M04 scheme (blue) the rapid increase of z_0 at high wind speed is

252 obviously restrained, although the mean z_0 is slightly higher than that in GFS at wind speed >10 m/s

253 due to larger C_{ch} (>0.014 in Fig. S2). Furthermore, since the Charnock number is constant in GFS, the

254 standard deviation (STD) of z_0 at a given wind speed is near zero. Since the z_0 is determined only by

255 wind-sea conditions in ST4 and ST4-M04 scheme, the STD at a given wind speed is mainly owing to

256 variations in wind fetch and development stage of sea state. The reduced STDs in ST4-M04 scheme,

257 compared to ST4, imply less sensitivity of z_0 to fetch and sea state. Note that the ST4-M04 is used in

258 GFS, while the z_0 in WW3 is still calculated by the ST4 source term to avoid affecting the balance of

259 adjusted wind input and dissipation.

260 **2.6 Set of Experiments**

261   A series of numerical experiments was conducted to evaluate the effects of aforementioned wave-

262 related processes on ocean and atmosphere in two 56-day periods, from January 3 to February 28, 2017

263 and from August 3 to September 28, 2018 for boreal winter and boreal summer, respectively.

264   The reference experiment (CTRL) is a one-way coupled experiment, in which CFSv2.0 provides 10-

265 m wind and surface current to WW3, whereas no variable is transferred from WW3 to CFSv2.0. The

266 results of CFSv2.0 in CTRL are consistent with the corresponding CFS Reanalysis data (Saha et al.,

267 2010). For each period, four sensitivity experiments were carried out (Table 1). Based on CTRL, the first

268 is VR12-AL-SC-EN experiment, in which the Langmuir mixing parameterization is applied with Stokes–

269 Coriolis force and entrainment in MOM4. The second is Z0-M04 experiment, in which the constant C_{ch}

270   in GFS is replaced by C_{ch} from WW3 ST4-M04 scheme. The effect of fluxes in GFS generated by $\Delta \vec{V}$

271   (Eqn. 14) is tested in the FLUX experiment. The last experiment is the ALL, which includes all three

272   parameterizations.

273   **3 Data**

274   Due to the availability of in situ and reanalysis data in the simulation periods, only sea surface

275   temperature (SST), ocean subsurface temperature and salinity (T/S), 2-m air temperature (T02), 10-m

276   wind speed (WSP10), and significant wave height (SWH) were used to evaluate the simulation results.

277   The daily average satellite Optimum Interpolation SST (OISST) data were obtained from NOAA, with

278     $0.25° \times 0.25°$ resolution (Reynolds et al., 2007; https://www.ncdc.noaa.gov/oisst). The global Argo

279   observational profiles of T/S (Li et al., 2019) were from China Argo Real-time Data Center

280   (www.argo.org.cn). The ERA5 datasets of T02, WSP10 and SWH with a spatial resolution of 0.5° were

281   also used (Hersbach et al., 2020; https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-

282   single-levels), which assimilated huge amounts of historical data and thus provided reliable hourly

283   estimates. Additionally, the WSP10 and SWH observations from the available National Data Buoy

284   Center (NDBC) buoy data (https://www.ndbc.noaa.gov) were applied for comparison.

285   **4 Experimental Results**

286   In this section, an evaluation of simulation results was presented. Comparisons were made between

287   model results and observations/reanalysis data. The results in the first three days were excluded in the

288   evaluation, since the initial wave influences were too weak.

4.1 Sea Surface Temperature (SST) and 2-m Air Temperature (T02)

Figure 3a shows the spatial distribution of 53-day (day 4 to day 56) averaged SST biases in CTRL in boreal winter, defined as SST in CTRL minus OISST. The global mean SST bias is approximately 0.32℃, and the average RMSE is about 1.09℃ from day 4 to day 56 in CTRL (Fig. 3a). The simulated SSTs are generally overestimated, and the large biases (>1.0℃) are mainly distributed in the Southern Ocean. In Fig. 3b, the global-averaged RMSEs of CTRL (black) increase with time in the first month and then gradually level off. Compared with CTRL, the RMSEs are reduced continuously in VR12-AL-SC-EN and ALL (yellow and red), but not in Z0-M04 and FLUX (purple and blue).

To understand the key process responsible for the bias reduction in ALL, the SST differences are compared across all four experiments (Fig.3c-f). Clearly, the difference in experiment VR12-AL-SC-EN is similar to that in ALL (Fig. 3c&3f). The spatial correlation coefficient between the SST differences with OISST of the two experiments (Fig. 3c&3f) is 0.67, significant at 99% confidence level, and the RMSEs of SST are not different significantly (red and yellow lines in Fig. 3b), indicating the Stokes drift-related parameterizations in VR12-AL-SC-EN mainly contribute to the SST positive bias reduction. This is different with Couvelard et al. (2020), where SST overestimations and MLD underestimations are reduced mainly due to the directly modified turbulence kinetic energy scheme. The global mean SST bias in ALL is 0.02 ℃ with RMSE of 1.03℃, and in most areas the SST differences compared with CTRL are significant (P ≤0.05) (dotted areas in Fig. 3f). Large SST improvements mainly appear in the Southern Ocean, with a regional RMSE decreases from 1.27 to 1.04 ℃ south of 45°S (Fig. 3f and red line in Fig. 5a). The reduction of overestimated SSTs in CTRL (red in Fig 3a) is because the Stokes drift-related parameterizations in MOM4 inject turbulent kinetic energy into the ocean, which enhance vertical mixing, and subsequently cool the surface waters (Belcher et al., 2012; Li et al. 2016). The modified

311   roughness and relative velocity in Z0-M04 and FLUX also influence upper ocean mixing (Fig. 3d&e)

312   via changing momentum flux, and lead to generally warmer SSTs (purple and blue lines in Fig. 3b&5a).

313   The effect from Stokes drift-related ocean mixing parameterizations dominates SST changes in ALL.

314   In boreal summer, the global mean SST bias in CTRL is overestimated approximately 0.29℃, and the

315   averaged RMSE from day 4 to day 56 is about 1.19℃. The overestimated SSTs (>1.0℃) mainly occur

316   in the Northern Hemisphere (Fig. 4a). The global-averaged RMSEs are also generally lower in VR12-

317   AL-SC-EN and ALL than in CTRL (Fig. 4b). The cooling effects in VR12-AL-SC-EN lead to a global

318   mean bias of 0.06℃, and the large SST improvements mainly occur north of 50°N (Fig. 4c and yellow

319   line in Fig. 5b). The changes of SST in Z0-M04 and FLUX (Fig. 4d&e; purple and blue lines in Fig.

320   4b&5b) are relatively small. The global mean bias in ALL is 0.04℃ with RMSE of 1.14℃ (Fig. 4f).

321   As aforementioned, large improvements of overestimated SST mainly occur at mid-high latitudes in

322   local summer. The time series of RMSEs and correlation coefficients of SST between model and

323   observation in the region (0-360°E, 45°-78°S in boreal winter and 0-360°E, 50°-78°N in boreal summer)

324   are shown in Fig. 5c-f. The RMSEs in CTRL (blue in Fig. 5c&d) increase in the first few weeks and then

325   gradually decrease afterward. Compared with CTRL, RMSEs in VR12-AL-SC-EN (yellow) and ALL

326   (red) are significantly (P ≤0.01) reduced by about 0.3℃. The spatial correlation coefficients decrease

327   with time but remain high (>0.90) for all experiments (Fig. 5e&f) with higher values in experiment

328   VR12-AL-SC-EN (yellow).

329   We also compared T02 from experiments with ERA5 (Fig. 6). Warm biases of T02 appear in both

330   winter and summer in CTRL (Fig. 6a&b). The changes of T02 in sensitivity experiments (Fig. 6c-j) are

331   generally consistent with the changes of SST in the same experiments (Fig.3&4). The correlation

332   coefficients between the SST and the T02 changes for the ALL experiment in boreal winter and summer

333   (Fig. 3f&6f and Fig. 4f&6j) are 0.61 and 0.53 respectively, significant at 99% confidence level. In boreal

334   winter, all wave-coupled experiments except FLUX reduce the T02 mean bias (Fig.6c-f). VR12-AL-SC-

335   EC has the largest T02 bias reduction compared with CTRL, from 0.55℃ to 0.17℃ (Fig.6c). In boreal

336   summer, both VR12-AL-SC-EC and ALL have the largest T02 bias reduction, from 0.29℃ to 0.08℃

337   (Fig.6g&j). Noticeably, the improvements in RMSEs are not large for all experiments, because the

338   improvements mainly occur in areas with overestimated temperature.

339   **4.2 Mixed Layer Depth (MLD)**

340   To further evaluate the direct effect of the wave-related processes on upper ocean, we compared the

341   MLD of all experiments with that estimated from Argo profiles in summer. The simulated T/S were

342   interpolated onto the positions of Argo profiles at the nearest time. The MLD was estimated as the depth

343   where the change of potential density reaches the value corresponding to a 0.2℃ decrease of potential

344   temperature with unchanged salinity from surface (de Boyer Montégut et al., 2004; Wang and Xu, 2018).

345   The time series of MLDs from numerical experiments and Argo south of 45°S in boreal winter (north

346   of 45°N in boreal summer) are compared in Fig. 7a (7b). The simulated MLDs are generally within the

347   STD of Argo MLDs (shading in Fig. 7). In CTRL, the mean bias (CTRL minus Argo) with STD is -

348   13.15±7.82 m (-6.75±5.29 m) in boreal winter (summer). The correlation coefficient of MLDs in CTRL

349   with Argo MLDs is 0.55 (0.68) with P ≤0.01, and the mean RMSE is 15.30 m (8.55 m) in boreal winter

350   (summer). In ALL, the mean bias (ALL minus Argo) with STD is 7.70±10.42 m (3.30±7.78 m) in boreal

351   winter (summer), and the correlation coefficient of MLDs enhances to 0.63 (0.78). The RMSE south of

352   45°S decreases from 15.30 m in CTRL to 12.96 m in ALL. The RMSE north of 45°N decreases from

353   6.71 m in CTRL to 5.55 m in ALL in the first six weeks but the value increases in the last two weeks due

354 to overestimation of MLDs. Compared with CTRL (orange in Fig. 7), VR12-AL-SC-EN (yellow) and

355 ALL (dark blue) show significantly improvements (P≤0.01) on the underestimated MLDs time series,

356 whereas the MLDs difference between CTRL and Z0-M04 (purple)/FLUX (blue) is non-significant.

357 **4.3 Wind Speed at 10 m (WSP10) and Significant Wave Height (SWH)**

358 Compared with ERA5, the WSP10s in CTRL are generally overestimated in both winter and summer

359 (Fig.8a&9a). The global averaged RMSEs of WSP10s in CTRL are 4.25 m/s (4.26 m/s) in boreal winter

360 (summer). The global averaged RMSEs of WSP10s in all experiments increase with time in the first two

361 weeks and then gradually level off with perturbations (Fig. 8b&9b). The differences of RMSEs between

362 CTRL and other experiments are tiny in the first 10 days, and afterwards the RMSEs in Z0-M04 and

363 ALL (purple and red) become clearly smaller than in CTRL over most of the time.

364 The comparisons of the simulated SWHs in CTRL with the ERA5 also show that the SWHs are

365 overestimated in both winter and summer (Fig. 10a&11a). In boreal winter, the global mean SWH bias

366 in CTRL is approximately 0.20 m with overestimates (> 0.30 m) in the Pacific, the North Atlantic and

367 the Southern Ocean (Fig.10a), and the average RMSE is about 1.29 m. In boreal summer, the global

368 mean bias in CTRL is approximately 0.17 m with 1.22 m RMSE (Fig. 11a). Similar to WSP10s, the

369 RMSEs of SWHs also increase in the first two weeks and then gradually level off with perturbations (Fig.

370 10b&11b). The RMSEs in Z0-M04 and ALL (purple and red) are smaller than in CTRL over most of the

371 time, consistent with changes of WSP10s. The correlation coefficients between changes of WSP10s and

372 changes of SWHs in ALL are 0.77 and 0.73 in boreal winter and summer respectively (Fig. 8f&10f and

373 Fig. 9f&11f), significant at 99% confidence level, indicating that the SWHs changes are closely related

374 to changes of wind speeds.

375   In VR12-AL-SC-EN, the reduction of SST warm biases affects air temperature and stabilizes marine

376   atmospheric boundary layer (Sweet et al. 1981; O'Neill et al. 2003), and subsequently reduces WSP10s

377   and SWHs with decreased global bias in boreal winter (Fig.8c&10c). In Z0-M04, the overestimated

378   WSP10s and SWHs are also reduced (Fig. 8d&10d) due to the larger z_0 with the ST4-M04 scheme at

379   wind speed > 10 m/s (Fig. 2). The increase of z_0 enhances wind stress and momentum transferred into

380   the ocean, and therefore reduces surface winds (Pineau-Guillou et al. 2018; Sauvage et al. 2020) and

381   consequently reduces SWHs. In FLUX (Fig. 8e&10e), $\overrightarrow{U_{surf}}$ and $\overrightarrow{u_s(0)}$ decrease wind stress and

382   momentum transfer when their directions are consistent with wind directions, and vice versa (Hersbach

383   and Bidlot, 2008; Renault et al., 2016). For instance, the angles between wind and current are relatively

384   small (<90°) in the northeastern Pacific, reducing the wind stress and thus enhancing WSP10s (Fig. 8e).

385   In contrast, the large angles (>90°) between the northwesterlies and the Kuroshio in the northwestern

386   Pacific enhance wind stress, and decrease WSP10s (Fig. 8e). Consequently, the improvements occur in

387   areas with misalignment of winds and currents. With all combined effects, the biases of WSP10s and

388   SWHs in ALL in most regions are decreased (Fig. 8f&10f), with the reduced global RMSEs of 4.17 m/s

389   and 1.18 m respectively. In boreal summer, the improvements of WSP10s and SWHs are relatively small

390   in terms of global averaged RMSEs, because of smaller positive biases in CTRL (Fig. 9a&11a). In ALL,

391   the global averaged bias of WSP10s (SWHs) is -0.01 m/s (0.03 m). The largest reduction primarily

392   appears in the Southern Ocean (Fig. 9f&11f) to improve the overestimated westerlies and SWHs in CTRL

393   (Fig. 9a&11a).

394   Previous studies indicated that ocean surface winds in ERA5 are underestimated in some regions

395   (Belmonte Rivas and Stoffelen 2019; Kalverla et al. 2020; Sharmar and Markina 2020). To better

396   demonstrate the effects of waves on WSP10s and SWHs, comparisons of WSP10s and SWHs with the

397   NDBC buoy data are made (Table 2 and Fig. 12). The differences between sensitivity experiments and

398   CTRL are all statistically significant at 95% confidence level. Buoys are mainly located in the

399   northeastern Pacific, the tropical Pacific and the northwestern Atlantic oceans (Fig. S3), and buoy

400   identifiers with total numbers, longitudes and latitudes are listed in Table S3. The method from Hsu et al.

401   (1994) was used to adjust wind speeds from buoy data to the reference height of 10 m.

402   Compared to the NDBC data, the WSP10s and the SWHs in CTRL are generally overestimated in both

403   winter and summer with positive mean biases (Table 2 and Fig. 12). The reduction of mean biases appears

404   in all experiments except FLUX in boreal winter. The wave-related processes are most effective in areas

405   with positive biases, consistent with previous comparisons with ERA5. In boreal winter, the angles

406   between winds and currents are small. The wind stresses are then reduced in FLUX, and the WSP10s are

407   enhanced. So the positive bias is further enhanced. The improvements in ALL are generally the largest

408   (Table 2), with the WSP10s RMSE of 1.04 m/s (1.15 m/s) and the SWHs RMSE of 0.36 m (0.24 m) in

409   boreal winter (summer). As shown in Fig. 12, with the increase of WSP10s and SWHs, the reduction of

410   overestimation in ALL compared with CTRL is more prominent.

411   **5 Summary and Discussion**

412   To investigate the individual role played by wave-related processes on atmosphere and ocean interface

413   in a coupled global atmosphere-ocean-wave modeling system on intraseasonal scale, we implemented

414   the version 5.16 of WW3 into CFSv2.0 for global oceans from 78°S-78°N, using the C-Coupler2. In this

415   coupled system, the WW3 was forced by 10-m wind and surface current generated in CFSv2.0. Stokes

416   drift-related Langmuir mixing, Stokes-Coriolis force and entrainment in ocean, air-sea fluxes modified

417   by surface current and Stokes drift, and momentum roughness length (z_0) were considered separately,

418 and the results of sensitivity experiments were compared against in-situ buoys, satellite measurements

419 and ERA5 reanalysis. The effects of waves on intraseasonal prediction were examined in two 56-day

420 cases, one for boreal winter and the other one for boreal summer.

421 The following key results were found:

422   1. Overestimated SST, T02 and underestimated MLD in mid-high latitudes in CFSv2.0 are

423   significantly improved, particularly in local summer. Because enhanced vertical mixing

424   generated by Langmuir turbulence, Stokes-Coriolis force and entrainment in VR12-AL-SC-

425   EN changes temperature structure in the upper ocean, and further affects air temperature. In

426   boreal winter, the regional RMSE of SST (T02) in the Southern Ocean decreases from 1.27

427   (1.93) in CTRL experiment to 1.04 (1.67) ℃ in ALL experiment. In boreal summer, the effect

428   is weaker because of the smaller ocean areas in mid-high latitudes of the Northern Hemisphere.

429   2. In general, all wave-related processes lead to reduction of biases for WSP10s and SWHs,

430   particularly in regions where WSP10s and SWHs are overestimated. The decreased SSTs in

431   VR12-AL-SC-EN stabilize marine atmospheric boundary layer, and lead to weakened WSP10s

432   and SWHs. The modified roughness in Z0-M04 generally enhances momentum transfer into

433   the ocean, and so decreases WSP10s and SWHs. The relative wind-wave-current speed in

434   FLUX also affects wind stress, and further influences WSP10s and SWHs. Compared with

435   NDBC buoy observations and ERA5, the ALL experiment shows significant improvements.

436 In addition to the variables aforementioned, the changes of simulated enthalpy fluxes were also

437 compared, which mainly depend on the WSP10s changes. However, the wave-related effects on enthalpy

438 fluxes are non-significant for the 2-month simulation, so the results are not shown.

439 The wave-related parameterizations used in the study mainly improve model biases at mid-high

440 latitudes, and SST biases in tropical oceans are only slightly improved (Fig. 3&4). Breivik et al. (2015)

441 improved SST as well as subsurface temperature simulations in Nucleus for European Modelling of the

442 Ocean (NEMO) with parametrizations including the wave-related Charnock parameter, modification of

443 water-side stress with wind input and wave dissipation, wave dissipation-related turbulent kinetic energy

444 flux and the Stokes-Coriolis force. Based on a global NEMO-WW3 coupled framework, Couvelard et al.

445 (2020) modified the Charnock parameter, the Stokes drift-related forces and the Langmuir cell with

446 misalignment of winds and waves, the oceanic surface momentum flux and the turbulence kinetic energy

447 to reduce SST and MLD biases. In addition, sea sprays can enhance air-sea heat fluxes in the tropics

448 (Andreas et al. 2008; Andreas et al. 2015). We will consider more processes in the future study.

449 Different parameterizations for the same wave-related process also deserve discussion. For ocean

450 surface roughness, the most classic parametrizations are those developed by Janssen (1989, 1991), Taylor

451 and Yelland (2001) and Drennan et al. (2003). The method of Taylor and Yelland (2001) requires the

452 peak wavelength for the total spectrum, whereas that of Drennan et al. (2003) only requires the peak of

453 wind-sea waves. This difference leads to the fact that the former is more suitable for a mixed sea state,

454 while the latter is more suitable for a young sea state (Drennan et al., 2005). And the effect of Janssen's

455 parameterization (1989, 1991) is similar to that of Drennan et al. (2003), since it is also based on the

456 wind-sea conditions (Shimura et al., 2017).

457 The case studies indicate that there remain significant biases in the coupled system, probably owing

458 to inaccuracy of coarse resolution, absence of a coupled wave-ice modular, and deficiency of initial fields.

459 In addition, to further improve the model and eliminate the biases, as Breivik et al. (2015) proposed,

460 extra adjusting of the individual model components in the coupled systems is also necessary. All of these

461 require further efforts to investigate efficient methods to improve fully coupled systems.

Code and data availability

The code developed for the coupled system can be found under https://doi.org/10.5281/zenodo.5811002

(Shi et al., 2021), including the coupling, preprocessing, run control and postprocessing scripts. The

initial fields for CFSv2.0 are generated by the real time operational Climate Data Assimilation System,

downloaded from the CFSv2.0 official website

(http://nomads.ncep.noaa.gov/pub/data/nccf/com/cfs/prod). The daily average satellite Optimum

Interpolation SST (OISST) data are obtained from NOAA (https://www.ncdc.noaa.gov/oisst), and the

National Data Buoy Center (NDBC) buoy data are also obtained from NOAA

(https://www.ndbc.noaa.gov). The Argo observational profiles of T/S are available at China Argo Real-

time Data Center (www.argo.org.cn). The ERA5 reanalysis are available at the Copernicus Climate

Change Service (C3S) Climate Date Store

(https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-single-levels).

Author contribution

FX and RS designed the experiments and RS carried them out. RS developed the code of coupling

parametrizations and produced the figures. ZF contributed to the installation and operation of CFSv2.0.

LL and HY contributed to the application of C-Coupler2. XL and YZ provided the original code of

CFSv2.0. RS prepared the manuscript with contributions from all co-authors. FX and HL contributed to

review and editing.

Acknowledgments

The authors would like to extend thanks to all developers of the CFSv2.0 model

(https://cfs.ncep.noaa.gov/cfsv2/downloads.html). The work was supported by National Key Research

483   and Development Program of China (no. 2016YFC1401408), and Tsinghua University Initiative

484   Scientific Research Program (2019Z07L01001). We also thank two anonymous reviewers for their

485   constructive comments.

486   **References**

487   Andreas, E. L., Mahrt, L., and Vickers, D.: An improved bulk air-sea surface flux algorithm, including

488   spray-mediated transfer, Quarterly Journal of the Royal Meteorological Society, 141, 642-654, 2015.

489   Andreas, E. L., Persson, P. O. G., and Hare, J. E.: A bulk turbulent air–sea flux algorithm for high-wind,

490   spray conditions, Journal of Physical Oceanography, 38, 1581-1596, 2008.

491   Ardhuin, F., Rogers, E., Babanin, A. V., Filipot, J., Magne, R., Roland, A., Der Westhuysen, A. V.,

492   Queffeulou, P., Lefevre, J. M., and Aouf, L.: Semiempirical Dissipation Source Functions for Ocean

493   Waves. Part I: Definition, Calibration, and Validation, Journal of Physical Oceanography, 40, 1917-1941,

494   http://dx.doi.org/10.1175/2010JPO4324.1, 2010.

495   Atlas, R., Hoffman, R. N., Ardizzone, J., Leidner, S. M., Jusem, J. C., Smith, D. K., and Gombos, D.: A

496   Cross-calibrated, Multiplatform Ocean Surface Wind Velocity Product for Meteorological and

497   Oceanographic Applications, Bulletin of the American Meteorological Society, 92, 157-174,

498   http://dx.doi.org/10.1175/2010BAMS2946.1, 2011.

499   Bao, Y., Song, Z., & Qiao, F.: FIO-ESM version 2.0: Model description and evaluation. Journal of

500   Geophysical Research: Oceans, 125, e2019JC016036. https://doi.org/10.1029/2019JC016036, 2019.

501   Belcher, S. E., Grant, A. L. M., Hanley, K., Foxkemper, B., Van Roekel, L., Sullivan, P. P., Large, W. G.,

502   Brown, A. R., Hines, A., and Calvert, D.: A global perspective on Langmuir turbulence in the ocean

503   surface boundary layer, Geophysical Research Letters, 39, http://dx.doi.org/10.1029/2012GL052932,

504   2012.

505   Beljaars, A. C. M.: The parametrization of surface fluxes in large-scale models under free convection. Q.

506   J. R. Meteorol. Soc., 121, 255{270. https://doi.org/10.1002/qj.49712152203, 1994.

507   Belmonte Rivas, M., and Stoffelen, A.: Characterizing ERA-Interim and ERA5 surface wind biases using

508   ASCAT, Ocean Science, 15, 831-852, 10.5194/os-15-831-2019, 2019.

509   Bidlot, J.-R., Prates, F., Ribas, R., et al.: Enhancing tropical cyclone wind forecasts, ECMWF newsletter,

510   164, 33-37. https://www.ecmwf.int/en/newsletter/164/meteorology/enhancing-tropical-cyclonewind-

511   forecasts, 2020.

512   Bidlot, J.-R.: Model upgrade improves ocean wave forecasts. ECMWF newsletter, 159, 10-10.

513   https://www.ecmwf.int/en/newsletter/159/news/modelupgrade-improves-ocean-wave-forecasts, 2019.

514   Bidlot, J.-R.: Present status of wave forecasting at ECMWF, Workshop on ocean waves, 25-27, 2012.

515   Breivik, Ø., Bidlot, J.-R., and Janssen, P. A.: A Stokes drift approximation based on the Phillips spectrum,

516   Ocean Modelling, 100, 49-56, 2016.

517   Breivik, Ø., Janssen, P. A., and Bidlot, J.-R.: Approximate Stokes drift profiles in deep water, Journal of

518   Physical Oceanography, 44, 2433-2445, 2014.

519   Breivik, Ø., Mogensen, K., Bidlot, J., Balmaseda, M., and Janssen, P. A. E. M.: Surface wave effects in

520   the NEMO ocean model: Forced and coupled experiments, Journal of Geophysical Research, 120, 2973-

521   2992, http://dx.doi.org/10.1002/2014JC010565, 2015.

522   Charnock, H.: Wind stress on a water surface, Quarterly Journal of the Royal Meteorological Society, 81,

523   639-640, http://dx.doi.org/10.1002/qj.49708135027, 1955.

524   Couvelard, X., Lemarié, F., Samson, G., Redelsperger, J.-L., Ardhuin, F., Benshila, R., and Madec, G.:

525   Development of a two-way-coupled ocean-wave model: assessment on a global NEMO(v3.6)-

526   WW3(v6.02) coupled configuration, Geosci. Model Dev., 13, 3067-3090, https://doi.org/10.5194/gmd-

527   13-3067-2020, 2020.

528   de Boyer Montégut, C., Madec, G., Fischer, A. S., Lazar, A., and Iudicone, D.: Mixed layer depth over

529   the global ocean: An examination of profile data and a profile‐based climatology, Journal of

530   Geophysical Research, 109, http://dx.doi.org/10.1029/2004JC002378, 2004.

531   Donelan, M. A., Haus, B. K., Reul, N., Plant, W. J., Stiassnie, M., Graber, H. C., Brown, O. B., and

532   Saltzman, E. S.: On the limiting aerodynamic roughness of the ocean in very strong winds, Geophysical

533   Research Letters, 31, 10.1029/2004gl019460, 2004.

534   Drennan, W. M., H. C. Graber, D. Hauser, and C. Quentin: On the wave age dependence of wind stress

535   over pure wind seas, J. Geophys. Res., 108(C3), 8062, doi:10.1029/2000JC000715, 2003.

536   Drennan, W. M., P. K. Taylor, and M. J. Yelland: Parameterizing the sea surface roughness, J. Phys.

537   Oceanogr., 35(5), 835-848, 2005.

538   ECMWF: Official IFS Documentation CY47R1. In chap. PART VII: ECMWF wave model. Reading,

539   UK: ECMWF, 10.21957/31drbygag, 2020.

540   Fan, Y., Lin, S., Held, I. M., Yu, Z., and Tolman, H. L.: Global Ocean Surface Wave Simulation Using a

541   Coupled Atmosphere–Wave Model, Journal of Climate, 25, 6233-6252, http://dx.doi.org/10.1175/JCLI-

542   D-11-00621.1, 2012.

543   Ghantous, M., and Babanin, A. V.: One-dimensional modelling of upper ocean mixing by turbulence due

544   to wave orbital motion, Nonlinear Processes in Geophysics, 21, 325-338, 10.5194/npg-21-325-2014,

545   2014.

546   Griffies, S. M., Harrison, M. J., Pacanowski, R. C., and Rosati, A.: A technical guide to MOM4, GFDL

547   Ocean Group Tech. Rep, 5, 342, 2004.

548   Hasselmann, K.: Wave-driven inertial oscillations, Geophysical and Astrophysical Fluid Dynamics, 1,

549   463-502, http://dx.doi.org/10.1080/03091927009365783, 1970.

550   Hersbach H. and Bidlot, J.-R.: The relevance of ocean surface current in the ECMWF analysis and

551   forecast system. Proceeding from the ECMWF Workshop on Atmosphere-Ocean Interaction, 10-12,

552   https://www.ecmwf.int/sites/default/files/elibrary/2009/9866-relevance-ocean-surface-current-ecmwf-

553   analysis-and-forecast-system.pdf, 2008.

554   Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., et al.: The ERA5 Global Reanalysis, Quarterly Journal

555   of the Royal Meteorological Society, Soc. 00: 363 (2020), https://doi.org/10.1002/qj.3803, 2020.

556   Hsu, S. A., Eric A. Meindl, and David B. Gilhousen: Determining the Power-Law Wind-Profile Exponent

557   under Near-Neutral Stability Conditions at Sea, Applied Meteorology, Vol. 33, No. 6,

558   https://doi.org/10.1175/1520-0450(1994)033<0757:DTPLWP>2.0.CO;2, 1994.

559   Janssen, P. A. E. M.: The interaction of ocean waves and wind, Cambridge University Press, pp 312,

560   2004.

561   Janssen, P. A. E. M.: The Quasi-linear theory of wind wave generation applied to wave forecasting. J.

562   Phys. Oceanogr. 21, 1631-1642. https://doi.org/10.1175/1520-

563   0485(1991)021<1631:QLTOWW>2.0.CO;2, 1991.

564   Janssen, P. A. E. M.: Wave-induced stress and the drag of air flow over sea waves. J. Phys. Oceanogr.

565   19(6), 745-754. https://doi.org/10.1175/1520-1310 0485(1989)019<0745:WISATD>2.0.CO;2, 1989.

566   Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W. D., Deaven, D. G., Gandin, L. S., Iredell, M. D., Saha,

567   S., White, G. H., and Woollen, J.: The NCEP/NCAR 40-Year Reanalysis Project, Bulletin of the

American Meteorological Society, 77, 437-471, http://dx.doi.org/10.1175/1520-0477(1996)077%3C0437:TNYRP%3E2.0.CO;2, 1996.

Kalverla, P. C., Holtslag, A. A. M., Ronda, R. J., and Steeneveld, G.-J.: Quality of wind characteristics in recent wind atlases over the North Sea, Quarterly Journal of the Royal Meteorological Society, 146, 1498-1515, 10.1002/qj.3748, 2020.

Kukulka, T., Plueddemann, A. J., Trowbridge, J. H., and Sullivan, P. P.: Significance of Langmuir circulation in upper ocean mixing: comparison of observations and simulations, Geophysical Research Letters, 36, http://dx.doi.org/10.1029/2009GL037620, 2009.

Law-Chune, S., and Aouf, L.: Wave effects in global ocean modeling: parametrizations vs. forcing from a wave model, Ocean Dynamics, 68, 1739-1758, http://dx.doi.org/10.1007/s10236-018-1220-2, 2018.

Lemarié, F.: Numerical modification of atmospheric models to include the feedback of oceanic currents on air-sea fluxes in ocean-atmosphere coupled models, INRIA Grenoble-Rhône-Alpes, Laboratoire Jean Kuntzmann, https://hal.inria.fr/hal-01184711/document, 2015.

Li, D., Staneva, J., Bidlot, J.-R., Grayek, S., Zhu, Y., and Yin, B.: Improving Regional Model Skills During Typhoon Events: A Case Study for Super Typhoon Lingling Over the Northwest Pacific Ocean, Frontiers in Marine Science, 8, 10.3389/fmars.2021.613913, 2021.

Li, M., and Garrett, C.: Mixed Layer Deepening Due to Langmuir Circulation, Journal of Physical Oceanography, 27, 121-132, http://dx.doi.org/10.1175/1520-0485(1997)027%3C0121:MLDDTL%3E2.0.CO;2, 1997.

Li, Q., Foxkemper, B., Breivik, O., and Webb, A.: Statistical models of global Langmuir mixing, Ocean Modelling, 113, 95-114, http://dx.doi.org/10.1016/j.ocemod.2017.03.016, 2017.

589   Li, Q., Webb, A., Foxkemper, B., Craig, A., Danabasoglu, G., Large, W. G., and Vertenstein, M.:

590   Langmuir mixing effects on global climate: WAVEWATCH III in CESM, Ocean Modelling, 103, 145-

591   160, http://dx.doi.org/10.1016/j.ocemod.2015.07.020, 2016.

592   Li, Z., Liu, Z., & Xing, X.: User Manual for Global Argo Observational data set (V3.0) (1997-2019),

593   China Argo Real-time Data Center, Hangzhou, 37pp, 2019.

594   Liu, L., Zhang, C., Li, R., and Wang, B.: C-Coupler2: a flexible and user-friendly community coupler for

595   model coupling and nesting, Geoscientific Model Development Discussions, 11, 1-63,

596   http://dx.doi.org/10.5194/gmd-11-3557-2018, 2018.

597   Luo, J.J., Masson, S., Roeckner, E., Madec, G., & Yamagata, T.: Reducing climatology Bias in an ocean-

598   atmosphere CGCM with improved coupling physics. Journal of Climate, 18(13), 2344-2360.

599   https://doi.org/10.1175/JCLI3404.1, 2005.

600   Mcwilliams, J. C., and Sullivan, P. P.: Vertical Mixing by Langmuir Circulations, Spill Science &

601   Technology Bulletin, 6, 225-237, http://dx.doi.org/10.1016/S1353-2561(01)00041-X, 2000.

602   Moon, I., Ginis, I., and Hara, T.: Effect of surface waves on Charnock coefficient under tropical cyclones,

603   Geophysical Research Letters, 31, http://dx.doi.org/10.1029/2004GL020988, 2004.

604   Moum J.N., and Smyth W.D.: Upper Ocean Mixing. In Cochran, J. Kirk; Bokuniewicz, J. Henry; Yager,

605   L. Patricia (Eds.) Encyclopedia of Ocean Sciences, 3rd Edition. vol. 1, pp. 71-79, Elsevier. ISBN: 978-

606   0-12-813081-0, 2019.

607   O'Neill, L. W., Chelton, D. B., and Esbensen, S. K.: Observations of SST-induced perturbations of the

608   wind stress field over the Southern Ocean on seasonal timescales, Journal of Climate, 16, 2340-2354,

609   10.1175/2780.1, 2003.

610   Pineau-Guillou L., Ardhuin,F., Bouin,M.N.,Redelsperger,J.L.,Chapron,B., Bidlot, J. R., & Quilfen, Y.:

611   Strong winds in a coupled wave-atmosphere model during a North Atlantic storm event: evaluation

612   against observations. Quarterly Journal of the Royal Meteorological Society, 144(711), 317-332, 2018.

613   Polonichko, V.: Generation of Langmuir circulation for nonaligned wind stress and the Stokes drift, J.

614   Geophys. Res., 102, 15773– 15780, https://doi.org/10.1029/97JC00460, 1997.

615   Powell, M. D., Vickery, P. J., and Reinhold, T. A.: Reduced drag coefficient for high wind speeds in

616   tropical cyclones, Nature, 422, 279-283, 10.1038/nature01481, 2003.

617   Qiao, F., Yuan, Y., Yang, Y., Zheng, Q., Xia, C., and Ma, J.: Wave‐induced mixing in the upper ocean:

618   Distribution and application to a global ocean circulation model, Geophysical Research Letters, 31,

619   http://dx.doi.org/10.1029/2004GL019824, 2004.

620   Renault, L., Arsouze, T., and Ballabrera-Poy, J.: On the Influence of the Current Feedback to the

621   Atmosphere on the Western Mediterranean Sea Dynamics, Journal of Geophysical Research-Oceans, 126,

622   10.1029/2020jc016664, 2021.

623   Renault, L., Molemaker, M. J., McWilliams, J. C., Shchepetkin, A. F., Lemarie, F., Chelton, D., Illig, S.,

624   and Hall, A.: Modulation of Wind Work by Oceanic Current Interaction with the Atmosphere, Journal of

625   Physical Oceanography, 46, 1685-1704, 10.1175/jpo-d-15-0232.1, 2016.

626   Reynolds, R. W., Smith, T. M., Liu, C., Chelton, D. B., Casey, K. S., and Schlax, M. G.: Daily High-

627   Resolution-Blended Analyses for Sea Surface Temperature, Journal of Climate, 20, 5473-5496,

628   http://dx.doi.org/10.1175/2007JCLI1824.1, 2007.

629   Saha, S., Moorthi, S., Pan, H., Wu, X., Wang, J., Nadiga, S., Tripp, P., Kistler, R., Woollen, J., and

630   Behringer, D.: The NCEP Climate Forecast System Reanalysis, Bulletin of the American Meteorological

631   Society, 91, 1015-1057, http://dx.doi.org/10.1175/2010BAMS3001.1, 2010.

632   Saha, S., Moorthi, S., Wu, X., Wang, J., Nadiga, S., Tripp, P., Behringer, D., Hou, Y., Chuang, H., and

633   Iredell, M. D.: The NCEP Climate Forecast System Version 2, Journal of Climate, 27, 2185-2208,

634   http://dx.doi.org/10.1175/JCLI-D-12-00823.1, 2014.

635   Sauvage, C., Lebeaupin Brossier, C., Bouin, M. N., & Ducrocq, V.: Characterization of the air-sea

636   exchange mechanisms during a Mediterranean heavy precipitation event using realistic sea state

637   modelling. Atmospheric Chemistry & Physics, 20(3), 2020.

638   Sharmar, V., and Markina, M.: Validation of global wind wave hindcasts using ERA5, MERRA2, ERA-

639   Interim and CFSRv2 reanalyzes, IOP Conference Series: Earth and Environmental Science, 606, 012056

640   (012059 pp.)-012056 (012059 pp.), 10.1088/1755-1315/606/1/012056, 2020.

641   Shimura, T., Mori, N., Takemi, T., et al.: Long term impacts of ocean wave-dependent roughness on

642   global   climate   systems.   Journal   of   Geophysical   Research: Oceans,   122(3),

643   https://doi.org/10.1002/2016JC012621, 2017.

644   Stopa, J. E., Ardhuin, F., Babanin, A. V., and Zieger, S.: Comparison and validation of physical wave

645   parameterizations   in   spectral   wave   models,   Ocean   Modelling,   103,   2-17,

646   http://dx.doi.org/10.1016/j.ocemod.2015.09.003, 2016.

647   Sweet, W., Fett, R., Kerling, J., and Laviolette, P.: Air–sea interaction effects in the lower troposphere

648   across the north wall of the Gulf Stream, Mon. Weather Rev., 109, 1042-1052, 10.1175/1520-

649   0493(1981)109<1042:Asieit>2.0.Co;2, 1981.

650   Takatama, K., and Schneider, N.: The Role of Back Pressure in the Atmospheric Response to Surface

651   Stress Induced by the Kuroshio, Journal of the Atmospheric Sciences, 74, 597-615, 10.1175/jas-d-16-

652   0149.1, 2017.

653 Taylor, P. K., and Yelland, M. J.: The Dependence of Sea Surface Roughness on the Height and Steepness

654 of the Waves, Journal of Physical Oceanography, 31, 572-590, http://dx.doi.org/10.1175/1520-

655 0485(2001)031%3C0572:TDOSSR%3E2.0.CO;2, 2001.

656 Terray, E. A., Donelan, M. A., Agrawal, Y. C., Drennan, W. M., Kahma, K. K., Williams, A. J., Hwang,

657 P. A., and Kitaigorodskii, S. A.: Estimates of Kinetic Energy Dissipation under Breaking Waves, Journal

658 of Physical Oceanography, 26, 792-807, http://dx.doi.org/10.1175/1520-

659 0485(1996)026%3C0792:EOKEDU%3E2.0.CO;2, 1996.

660 Tolman, H. L., and Chalikov, D. V.: Source Terms in a Third-Generation Wind Wave Model, Journal of

661 Physical Oceanography, 26, 2497-2518, http://dx.doi.org/10.1175/1520-

662 0485(1996)026%3C2497:STIATG%3E2.0.CO;2, 1996.

663 Van Roekel, L. P., Foxkemper, B., Sullivan, P. P., Hamlington, P. E., and Haney, S.: The form and

664 orientation of Langmuir cells for misaligned winds and waves, Journal of Geophysical Research, 117,

665 http://dx.doi.org/10.1029/2011JC007516, 2012.

666 Wang, L., and Xu, F.: Decadal variability and trends of oceanic barrier layers in tropical Pacific, Ocean

667 Dynamics, 68, 1155-1168, 10.1007/s10236-018-1191-3, 2018.

668 WAVEWATCH III Development Group.: User manual and system documentation of WAVEWATCH III

669 version 5.16, NOAA/NWS/NCEP/MMAB Technical Note 329, 326, 2016.

670 Wu, L., Staneva, J., Breivik, Ø., Rutgersson, A., Nurser, A. G., Clementi, E., and Madec, G.: Wave effects

671 on coastal upwelling and water level, Ocean Modelling, 140, 101405, 2019.

672 Wu, X., K. S. Moorthi, K. Okomoto, and H. L. Pan: Sea ice impacts on GFS forecasts at high latitudes,

673 Proceedings of the 85th AMS Annual Meeting, 8th Conference on Polar Meteorology and Oceanography,

674 San Diego, CA, 2005.

675   Zieger, S., Babanin, A. V., Rogers, W. E., and Young, I. R.: Observation-based source terms in the third-

676   generation wave model WAVEWATCH, Ocean Modelling, 96, 2-25,

677   http://dx.doi.org/10.1016/j.ocemod.2015.07.014, 2015.

Table 1. List of numerical experiments: setups different from CTRL are marked with bold

Experiments	Physical Process/Parameterization		
	Langmuir Cell with Stokes-Coriolis Force and Entrainment	Roughness (Charnock Parameter)	Relative Velocity in Flux
CTRL	Off	Off	Off
VR12-AL-SC-EN	**Eqn. 1-6, 8-10**	Off	Off
Z0-M04	Off	**C_{ch} from Eqn. 16, 17**	Off
FLUX	Off	Off	**$\Delta\vec{V}$ from Eqn. 14**
ALL	**Eqn. 1-6, 8-10**	**C_{ch} from Eqn. 16, 17**	**$\Delta\vec{V}$ from Eqn. 14**

Table 2. The 53-day mean bias with standard deviation (STD) and RMSE for WSP10 and SWH compared with NDBC buoy observation: the bias is calculated as simulation minus NDBC.

Boreal Winter WSP10	Bias with STD	RMSE
CTRL	0.16±1.23	1.24
VR12-AL-SC-EN	0.01±1.12	1.12
Z0-M04	-0.01±1.07	1.07
FLUX	0.39±1.20	1.26
ALL	0.07±1.04	1.04
Boreal Winter SWH	**Bias with STD**	**RMSE**
CTRL	0.21±0.38	0.44
VR12-AL-SC-EN	0.14±0.35	0.37
Z0-M04	0.10±0.30	0.32
FLUX	0.24±0.34	0.42
ALL	0.12±0.34	0.36
Boreal Summer WSP10	**Bias with STD**	**RMSE**
CTRL	0.15±1.23	1.24
VR12-AL-SC-EN	-0.03±1.22	1.22
Z0-M04	-0.04±1.21	1.21
FLUX	-0.22±1.18	1.20
ALL	-0.17±1.14	1.15
Boreal Summer SWH	**Bias with STD**	**RMSE**
CTRL	0.28±0.25	0.38
VR12-AL-SC-EN	0.19±0.24	0.30
Z0-M04	0.22±0.26	0.34
FLUX	0.14±0.25	0.29
ALL	0.12±0.21	0.24

Figure 1. A schematic diagram of the atmosphere-ocean-wave coupled modeling system. The arrows indicate the coupled variables that are passed between the model components. In the diagram, C_{ch}, La_t, $u_s(0)$, V_s, U_{10}, and U_{surf} are Charnock parameter (red arrows), turbulent Langmuir number (red arrows), surface Stokes drift velocity (red arrows), Stokes transport (red arrows), 10-m wind (green arrows) and surface current (blue arrows), respectively.

Figure 2. Relationships between momentum roughness length z_0 (m) in the coupled system and 10-m wind speed (m/s); error bars indicate twice the standard deviations for each point.

Figure 3. The 53-day average SST (℃) bias in CTRL (a; CTRL minus OISST), the time series of global-averaged RMSE (b), and the differences between VR12-AL-SC-EN (c)/Z0-M04 (d)/ FLUX (e)/ ALL (f) and CTRL in Jan-Feb, 2017 (VR12-AL-SC-EN/Z0-M04/FLUX/ALL minus CTRL). The first 3-day simulation is discarded. The dotted areas are statistically significant at 95% confidence level.

Figure 4. The same as Figure 3 but for Aug-Sep, 2018.

Figure 5. The 53-day averaged latitudinal distribution of SST root mean square errors (RMSE), time series of domain-averaged SST RMSE and correlation coefficient: **a/b** the latitudinal RMSE in boreal winter/summer compared with OISST, **c/d** the time series of domain-averaged (0-360°E, 45-78°S/50-78°N) SST RMSE in boreal winter/summer, **e/f** the time series of domain-averaged (0-360°E, 45-78°S/50-78°N) SST correlation coefficient in boreal winter/summer; differences of RMSE and correlation coefficient time series between VR12-AL-SC-EN/ALL and CTRL are statistically significant at 99% confidence level, except those in Fig. e.

Figure 6. The 53-day average T02 (°C) bias in CTRL (a&b; CTRL minus ERA5), and the differences between VR12-AL-SC-EN (c&g)/Z0-M04 (d&h)/ FLUX (e&i)/ ALL (f&j) and CTRL (VR12-AL-SC-EN/Z0-M04/FLUX/ALL minus CTRL). The first 3-day simulation is discarded. The dotted areas are statistically significant at 95% confidence level. a/c/d/e/f are for Jan-Feb, 2017, and b/g/h/i/j are for Aug-Sep, 2018.

Figure 7. The 53-day time series of domain-averaged (0-360°E, 45-78°S/N) mixed layer depth (MLD; m) in boreal winter/summer: the difference between CTRL and VR12-AL-SC-EN/ALL passes the student's t-test at 99% confidence level; the time intervals are 6 hours; shaded areas indicate twice the standard deviations for Argo.

Figure 8. The 53-day average WSP10 (m/s) bias in CTRL (a; CTRL minus ERA5), the time series of global-averaged RMSE (b), and the differences between VR12-AL-SC-EN (c)/Z0-M04 (d)/ FLUX (e)/ ALL (f) and CTRL in Jan-Feb, 2017 (VR12-AL-SC-EN/Z0-M04/FLUX/ALL minus CTRL). The first 3-day simulation is discarded. The dotted areas are statistically significant at 95% confidence level.

Figure 9. The same as Figure 8 but for Aug-Sep, 2018.

Figure 10. The 53-day average SWH (m) bias in CTRL (a; CTRL minus ERA5), the time series of global-averaged RMSE (b), and the differences between VR12-AL-SC-EN (c)/Z0-M04 (d)/ FLUX (e)/ ALL (f) and CTRL in Jan-Feb, 2017 (VR12-AL-SC-EN/Z0-M04/FLUX/ALL minus CTRL). The first 3-day simulation is discarded. The dotted areas are statistically significant at 95% confidence level.

Figure 11. The same as Figure 10 but for Aug-Sep, 2018.

Figure 12. Scatter plots of simulated WSP10/SWH (y-axis) vs buoy WSP10/SWH (x-axis): (a) the WSP10 in Jan-Feb, 2017, (b) the SWH in Jan-Feb, 2017, (c) the WSP10 in Aug-Sep, 2018, and (d) the SWH in Aug-Sep, 2018. The dotted line is y=x. The corresponding mean biases with standard deviations and RMSEs for every experiment are shown in Table 2.