

# RavenR v2.1.4: an open source R package to support flexible hydrologic modelling

Robert Chlumsky<sup>1</sup>, James R. Craig<sup>1</sup>, Simon G.M. Lin<sup>1</sup>, Sarah Grass<sup>2</sup>, Leland Scantlebury<sup>1,3</sup>, Genevieve Brown<sup>4</sup>, and Rezgar Arabzadeh<sup>1</sup>

<sup>1</sup>Department of Civil and Environmental Engineering, University of Waterloo, Waterloo, ON, Canada

<sup>2</sup>Geoprocess Research Associates, Edmonton, AB, Canada

<sup>3</sup>Hydrologic Sciences Graduate Group, University of California, Davis, Davis, CA, United States

<sup>4</sup>Northwest Hydraulic Consultants Ltd., North Vancouver, BC, Canada

**Correspondence:** Robert Chlumsky (rchlumsk@uwaterloo.ca)

**Abstract.** In recent decades, advances in the flexibility and complexity of hydrologic models hashave enhanced their utility in scientific studies and practice alike. However, the increasing complexity of these tools leads to a number of challenges, including steep learning curves for new users and in the reproducibility of modelling studies. Here, we present the RavenR package, an R package that leverages the power of scripting to both enhance the usability of the Raven hydrologic modelling

5 framework and provide complimentary analyses that are useful for modellers. The RavenR package contains functions that may be useful in each step of the model-building process, particularly for preparing input files and analyzing model outputs, and these tools may be useful even for non-Raven users. The utility of the RavenR package is demonstrated with the presentation of six use cases for a model of the Liard River basin in Canada. These use cases provide examples of visually reviewing the model configuration, preparing input files for observation and forcing data, simplifying the model discretization, performing realityre-  
10 alism checks on the model output, and evaluating the performance of the model. All of the use cases are fully reproducible, with additional reproducible examples of RavenR functions included with the package distribution itself. It is anticipated that the RavenR package will continue to evolve with the Raven project, and will provide a useful tool to new and experienced users of Raven alike.

## 1 Introduction

15 Hydrologic models are used for numerous applications, including streamflow prediction, flood forecasting, reservoir level forecasting, and in a scientific capacity to advance our understanding of hydrologic systems. Historically, most hydrologic models have been designed with a fixed model structure comprised of a predefined set of environmental processes, while the input data and model parameters may vary from watershed to watershed. While these fixed model structures (e.g., GR4J; Perrin et al., 2003) may provide sufficient performance in some catchments, they are not adequate in all catchments, environments, 20 or hydrologic applications (Hoey et al., 2014). Numerous studies have called this fixed structure paradigm into question, and have instead called for the development of flexible modelling frameworks (Leavesley et al., 2002; Clark et al., 2011; Fenicia et al., 2011), which would allow the modeller to possess more control over the model-building process. This has resulted in

the emergence of flexible modelling frameworks in the literature (e.g., Orellana et al., 2008; Clark et al., 2008; Kavetski and Fenicia, 2011; Clark et al., 2015; Knoben et al., 2019; Coxon et al., 2019; Craig et al., 2020), and recent studies have been 25 extensively supported by the use of these frameworks (Pilz et al., 2020; Remmers et al., 2020; Chadalawada et al., 2020; Knoben et al., 2020; Spieler et al., 2020; Mai et al., 2020; Chlumsky et al., 2021b).

The power contained in these flexible hydrologic models is limited in part by the modeler's ability to take advantage of it. In an ideal setup, a modeller would find converting their system conceptual model to a numerical model a seamless process; in actuality, setting up a numerical model often involves data wrangling and other tedious tasks, with decisions ranging from those 30 with relatively little impact on the final modelling results (e.g., how to combine dozens of text files) to potentially problematic and highly impactful decisions (e.g., time series interpolation or model structure adjustments). ~~Even among hydrological modelling software that have graphical user interfaces (GUIs),~~ ~~f~~Few hydrologic models offer the ability to easily deploy and compare successive model runs, resulting in a potentially large amount of time devoted to relatively trivial tasks, such as organizing model files and comparing successive model runs.

35 In order to address some of these challenges, new tools must be developed to bridge the gap between complex, customizable tools and the ability for modellers (in particular, new users) to fully understand and deploy these tools. Increasingly, freely available, open-source scripting languages, such as Python (Van Rossum and Drake, 2009) and R (R Core Team, 2021), are being employed by modellers to create, visualize, and evaluate their models (Jackson et al., 2019; Slater et al., 2019; Astagneau et al., 2021). While the languages themselves can carry significant learning curves, they unlock a wide range of time-saving 40 features due to their ability to reproducibly automate tasks in concise code that can be re-used between projects. ~~Additionally, these languages can be greatly expanded through the straightforward installation of downloadable packages that allow the performing of complex statistical analyses, obtaining and manipulating data, producing publication-ready plots, and even building interactive web visualizations, to be reduced to a handful of lines of code.~~ Many features and tools that would be inappropriate or inadvisable to build into the source code of hydrologic models (e.g., downloading and quality controlling input data) are perfectly suited to scripting languages.

45 R, in particular, has gained significant ground in hydrology, entering the toolbox of many in both consulting and academia (Anderson et al., 2018; Slater et al., 2019; Astagneau et al., 2021). This is due in part to the already robust package ecosystem awaiting hydrologists; the Comprehensive R Archive Network (CRAN), which hosts and tests R packages, allows for easy access to packages for a variety of data processing tasks such as downloading data (`tidyhydat` (Albers, 2017)), examining data (`trend` (Pohlert, 2020)), manipulating shapefiles and spatial data (`sf`, `raster` (Pebesma, 2018; Hijmans et al., 2021)), 50 evaluating model outputs (`hydroGOF` (Mauricio Zambrano-Bigiarini, 2020)), and visualizing data (`ggplot2` (Wickham, 2016)). Many hydrology-specific open-source packages have been developed in recent years, such as the `CSHShydRology` package (Shook et al., 2021), `Evapotranspiration` (Guo et al., 2020), and many other packages reviewed in the literature (Slater et al., 2019; Astagneau et al., 2021) and listed on CRAN Task View for Hydrology (<https://cran.r-project.org/web/views/Hydrology.html>). R is also being used extensively in teaching hydrology to professionals and graduate students, and packages 55 have also been used directly as educational tools, such as the `airGRteaching` package (Delaigue et al., 2020, 2018).

Reproducibility is essential to scientific advancement: to build a usable body of knowledge, we must be able to trust, rely on, learn from, and when necessary, upend the lessons of past experiments (Hutton et al., 2016). There are studies suggesting

that large proportions of scientific studies may not be reproducible (Baker, 2016; Camerer et al., 2018), including the field of computational hydrology (Hutton et al., 2016; Chawanda et al., 2020). Studies deploying proprietary or even free but not 60 open-source software, such as HEC-HMS (Hydrologic Engineering Center, 2020), pose issues in their reproducibility. While open-source modelling software provides a transparent codification of the model, the reliability of the resulting model depends heavily on the preparation of model input files and required data, which may not necessarily undergo the same level of scrutiny and transparency in a given modelling exercise. ~~The e~~Communication of modelling methodology and workflow in science has historically been primarily accomplished through ~~the inclusion of a detailed methods section within~~ research manuscripts. However, in 65 hydrology, the code and data that are needed to generate published results are often not made available, and the complexity of many hydrologic models and analyses make it infeasible for authors to include all necessary details for full reproducibility in their manuscripts (Hutton et al., 2016). Increasingly, scientists and engineers are being encouraged to use (and in some cases publish) computationally reproducible versions of their analyses (National Academies of Sciences, Engineering, and Medicine, 2019). Scripts, as a perfect record of the data manipulation, model setup, post-processing, and even figure creation steps, are 70 the ideal tools to accomplish this.

R, in particular, has gained significant ground in hydrology, entering the toolbox of many in both consulting and academia (Anderson et al., 2018; Slater et al., 2019; Astagneau et al., 2021). This is due in part to the already robust package ecosystem awaiting hydrologists; the Comprehensive R Archive Network (CRAN), which hosts and tests R packages, allows for easy access to packages for a variety of data processing tasks such as downloading data (`tidyhydat` (Albers, 2017)), examining 75 data (`trend` (Pohlert, 2020)), manipulating shapefiles and spatial data (`sf`, `raster` (Pebesma, 2018; Hijmans et al., 2021)), evaluating model outputs (`hydroGOF` (Mauricio Zambrano-Bigiarini, 2020)), and visualizing data (`ggplot2` (Wickham, 2016)). Many hydrology-specific open-source packages have been developed in recent years, such as the `CSHShydRology` package (Shook et al., 2021), `Evapotranspiration` (Guo et al., 2020), and many other packages reviewed in the literature 80 (Slater et al., 2019; Astagneau et al., 2021) and listed on CRAN Task View for Hydrology (<https://cran.r-project.org/web/views/Hydrology.html>). R is also being used extensively in teaching hydrology to professionals and graduate students, and packages have also been used directly as educational tools, such as the `airGRteaching` package (Delaigue et al., 2020, 2018).

Here, we introduce an R package with a collection of tools to aid a modeller in preparing, running, and post-processing results from custom hydrologic models developed with the hydrologic modelling framework Raven. Many of the tools are not solely Raven-specific: functions exist to plot time series, analyze yearly patterns, and compute relevant statistics. However, the 85 package importantly contains a robust suite of functions for creating, reading, and manipulating Raven model files. Specific attention has been paid to supporting the testing, comparison, and diagnosis of models built with variable model structure; many of these tools are unique. The intended purpose of the `RavenR` package is to enable modelers to simplify, automate, and document their model creation process, effortlessly facilitate model visualization and evaluation, and to expand the flexibility of the Raven hydrological modelling framework through scripting.

Section 2.1 briefly discusses Raven, and provides context for the `RavenR` package as a tool that enables improved workflows with Raven. Section 2.2 discusses `RavenR` in more detail, including a typical model-building workflow (Section 2.2.1), the installation and documentation available within `RavenR` (Section 2.2.2), and a description of the sample data sets available within the package and external to the package (Section 2.2.3).

95 **2.1 Raven hydrologic modelling framework**

Raven is an open-source software framework that can be used to build models from a selection of more than 100 available process algorithms (Craig et al., 2020). It is estimated that at least  $8 \times 10^{12}$  different hydrologic model configurations may be setup using Raven (Mai et al., 2020) and this number is continuously increasing as new options are added to the software. Raven is built for flexibility not only in process representation, but also in enabling multiple numerical schemes, discretization 100 schemes, input data types, and in providing the user control over output options. Raven is a fully object-oriented code written in C++ (Stroustrup, 2013), and is typically run from a command line. The input and output files are generally stored as text files (\*.txt or \*.csv) or in NetCDF (Network Common Data Format) format. This allows all model files to be stored as non-proprietary formats, and to be read and processed with any number of available tools for manipulating files.

The primary input files required for Raven (listed by file extension) include:

- 105 1. \*.rvi - primary input file which defines the model structure, timestep, duration, and a number of additional options
2. \*.rvp - model parameter specification and soil/vegetation/land class definitions
3. \*.rvh - model discretization, including all subbasin and hydrologic response unit (HRU) information
4. \*.rvt - time series data, including forcing and observational data; this file often points to other \*.rvt files with data sets for various stations and locations
- 110 5. \*.rvc - initial conditions for the model run

Raven provides complete control over its output generation (Craig et al., 2020), a relatively uncommon feature in hydrologic modelling software. ~~A large proportion of the computational cost of a model run is often used in the writing of output files, and thus a substantial computational burden can be alleviated if only the required outputs are written. This could be a single diagnostic metric for calibration, a single time series, or the complete mass and energy balances of the model for debugging or auditing purposes. Raven~~ Additionally, it also allows for custom outputs to 115 be generated for a given statistical, spatial and temporal specification and state variable, such as the monthly average of daily snow depth for a particular set of subbasins (Craig et al., 2020). -

This flexibility of Raven over the modelling process provides the modeller with a lot of power in configuring and running their hydrologic model, but also provides some challenges in preparing files and working with the many possible outputs.

The command-line execution of the program and the lack of a user interface can present a learning curve for new users, but 120 also enables scripting languages to easily interface with Raven, and for Raven to be deployed in high-performance computing environments.

A number of utilities exist to support the usage of Raven models, including the RavenPy (<https://github.com/CSHS-CWRA/RavenPy>) for creating, running, and post-processing Raven models within Python, and HydroGlyph (<http://raven.uwaterloo.ca/hydroglyph/>) for visualizing Raven time series output data. Hydrologic model support is also provided by many model-independent packages, such as the CSHS-hydRology package (Anderson et al., 2018). However, RavenR is the most comprehensive tool for preparing input files and performing a range of analyses with Raven output files.

## 2.2 RavenR software description

### 2.2.1 RavenR overview

The RavenR package is developed in R, and is a collection of tools to aid the modeller in preparing, running, and post-processing files associated with a hydrologic model developed using Raven (Craig et al., 2020). Unlike other software implementations, such as SuperflexPy (Dal Molin et al., 2021), in which the model code is contained within R or Python, the RavenR package is independent of the Raven model code, and operates only on related model input/output files and/or calling the compiled Raven executable. The primary benefits of this separation are 1) maintaining the computational speed of the compiled model code, and 2) allowing other users to implement their own non-R workflows in Python, shell scripts, or system-specific forecasting frameworks, such as DELFT-FEWS (Werner et al., 2013). The development of analysis and plotting tools in R benefits from a rich array of libraries and relatively fast development when compared to a compiled language such as C++, which is part of the reason for choosing to develop the external analysis tools in R.

RavenR is not intended to provide every tool needed to manipulate and analyze Raven input and output files, as the flexibility of the Raven framework would require a vast collection of highly specific scripts to accommodate the needs of all modellers. Instead, the package aims to reduce the effort required to use Raven, and allows the modeller to more effectively use the open-source scripting environment of R in their workflows. This may also reduce the learning curve of Raven that is created by its flexibility, as the package provides the means to guide new users through the generation and manipulation of common files, and reduces the burden in analyzing the model results. This can be particularly helpful for users migrating from GUI-based software such as HEC-HMS (Hydrologic Engineering Center, 2020).

The available functions within RavenR can be broadly categorized by their utility into the main categories of: 1) preparing input files, 2) reading output files, 3) running Raven, 4) tools for hydrologic analyses, and 5) support utilities (e.g. time series processing, water year analysis, etc.). The typical workflow for RavenR is closely related to the workflow required for the development and use of any hydrologic model, including one developed with Raven. This includes the collecting and processing of data for the model, determining the model structure, creating model input files in the format required by the modelling software, executing the model, and analyzing the results of the model for hydrologic consistency and performance. This can include exercises in model calibration and validation, uncertainty analysis, identifiability analysis, and project-specific simulations or adjustments to the model runs.

The typical workflow for developing a hydrologic model and examples of RavenR functions that may be used to support each step are provided in **Table 1**.

**Table 1.** Typical workflow table for building hydrologic models and connection to RavenR.

Step	Activity	Description	RavenR Functions
1	Collect/Prepare Data	Preparation and quality control of Raven input files (e.g., *.rvi files from template, *.rvt files), often from public data sources	rvn_rvi_write_template, rvn_rvt_tidyhydat, + 19 others
2	Discretize Watershed	Quality control of implemented discretization scheme and further simplification (e.g., aggregating very small or similar HRUs)	rvn_rvh_clean_hrus, rvn_subbasin_network_plot
3	Identify and Describe Processes	Model structure development and process algorithm selection	rvn_rvi_connections, rvn_rvi_process_diagrammer, rvn_rvi_process_ggplot
4	Parameterize the Model	Model parameter definition and parameter value specification	rvn_rvi_get_params
5	Execute the Model	Running the Raven (or other hydrologic) model	rvn_download & rvn_run
6	Processing Model Outputs	Reading and processing model output files for analysis	rvn_hyd_read, rvn_custom_read, + 7 others
7	Plots and Model Diagnostics	Checking model performance with a number of analyses, <a href="#">realism</a> checks, and diagnostics (often in conjunction with model calibration and validation)	rvn_annual_peak_flow, rvn_monthly_vbias, + 24 others
8	Report Results	Generating quality graphics and workflows to communicate results	Functions from step 7 + additional R libraries (e.g., ggplot2 & rmarkdown)

155     Although the model-building process is listed in **Table 1** as a series of steps, in practice it is not linear, but rather iterative and cyclic. For example, a model diagnostic (step 7) may show that inadequate model performance can be remedied by the inclusion of additional forcing data, requiring new data to be written to file (step 1). It is also recommended or common practice in modelling to begin with a simpler model and proceed to a more complex one (e.g., Fenicia et al., 2008), which may require iteration on steps 2-6 to potentially modify the structure (e.g. spatial and temporal discretization, hydrologic processes) after  
160 a basic model has been established. Model calibration would typically involve an iteration upon steps 4-6 with a calibration algorithm, and a calibration that includes model structure (e.g., Spieler et al., 2020; Chlumsky et al., 2021b) would effectively iterate upon steps 3-6. The iterative need for these model-building steps emphasizes the benefit of tools (including those in RavenR) that can reduce the overhead in simple but repetitive tasks, such as producing figures and writing data to a specific file format.

165     The functions within the RavenR package are named, where appropriate, by the three letter Raven file name or short abbreviation corresponding to the output file that they interact with, e.g., `rvn_rvi_connections` for processing the \*.rvi file structure or `rvn_res_read` for reading the output ReservoirStages.csv file. Other functions simply use illustrative names to convey their purpose (e.g. `rvn_budyko_plot`). This naming convention provides some navigability of the package functions to the new user, even before the package documentation is reviewed (see Section 2.2.2).

170     The RavenR package has a number of preferred data formats and related package dependencies. Most plots are generated using the `ggplot2` (Wickham, 2016) and related libraries from the so-called tidyverse, including `dplyr` (Wickham et al., 2021a) and `tidyrr` (Wickham, 2021) for data manipulation. This allows all plots to be exported as plot objects and further manipulated by the user as desired, and removes the need for all plot options to be wrapped into RavenR functions. Time series handling is done through the `lubridate` (Grolemund and Wickham, 2011) and `xts` (Ryan and Ulrich, 2020) packages,  
175 where the extensible time series (`xts`) format is generally expected for time series data. Finally, support for network analysis is done through the `igraph` package (Csardi and Nepusz, 2006), which primarily supports the organization of watershed discretization connections (\*.rvh file) and the network of model structure connections (\*.rvi file), including the related plot functions, e.g., `rvn_subbasin_network_plot` and `rvn_rvi_process_diagrammer`.

## 2.2.2 Installation and documentation

180     The package is developed as a free and open-source software tool, which is ideal for maintaining transparency and reproducibility in workflows related to hydrologic modelling and all steps involved. The stable package version is available for download through CRAN (currently version 2.1.4), which can be installed in R using the command `install.packages ("RavenR")`. The development version of the package is available on Github (<https://github.com/rchlumsk/RavenR>) and may be installed using the `devtools` library (Wickham et al., 2021b) as `devtools::install_github('rchlumsk/RavenR')`. Both  
185 installation commands resolve the dependencies associated with the package.

The RavenR package is fully documented and contains a description of inputs, outputs, with an usage example for each function consistent with the standards for CRAN packages. In addition to the package documentation, an introductory vignette *Introduction to RavenR*, is included with the package, which discusses getting started with the package and how it may be used

in a manner that is more useful to new users of Raven and RavenR. The introductory vignette is available with the command

190 `install.packages ("RavenR")`.

### 2.2.3 Sample data sets

In the interest of reproducible examples, the RavenR package contains a number of sample data sets and raw data files embedded within the package which are used within the function examples. Sample data is embedded directly as imported

195 data (accessible with the `data` function in R) for a number of file output types (hydrograph, watershed storage, etc.), as well as sample data for the `tidyhydat` and `weathercan` packages. The latter sample data allows the function examples to run without a dependency on the mentioned data-retrieval packages. Raw data files are also included (accessible with the `system.file` function in R), which allow for the testing of reading raw data directly. The examples where raw data files are first read into R using RavenR functions may be more helpful than examples which call sample data directly with the `data` command, since the workflow will be closer to the one applied in practice.

200 The sample Raven output files and data that is distributed with the RavenR package were generated from a model of the Nith watershed, which is located immediately west of Kitchener-Waterloo in Ontario, Canada. The Raven model of the Nith watershed can be found in full on the Raven webpage (<http://raven.uwaterloo.ca/downloads.html>) in the Tutorials 1-4 download set. Numerous additional Raven models are available from this page, including the model of the Liard River basin (Brown and Craig, 2020), which is used in the RavenR case studies in this manuscript (**Section 3**).

205 **3 Use cases of the RavenR package**

In this section, we present a number of use cases of the RavenR package. These cases are not intended to be a comprehensive review of all the applications for the RavenR package, but to provide the reader with a partial demonstration of how the package may be used in conjunction with Raven. The cases are discussed in the context of hydrologic modelling with flexible frameworks more broadly, and provide cases and checks that are likely to be useful when deploying both Raven and non-Raven

210 hydrologic models.

The use cases are presented in approximate order of the model-building process (**Table 1**), beginning with the generation of model input files and proceeding to the analysis of output files. These use cases include examples and discussion of ~~almost all~~  
~~almost of the~~ steps of in the model-building process, ~~with the exception of steps 4 and 5. Tools for these steps, such as running Raven from within R using `rvn_run` (step 4), exist within the RavenR package but are not discussed in detail in this section, with additional examples available in the use cases markdown file.~~

215 All R code and model files required to generate the results and figures in this section are provided in an open-source Zenodo repository (<https://doi.org/10.5281/zenodo.5534818><https://doi.org/10.5281/zenodo.5534817>), and utilize the version of RavenR currently available on CRAN (version 2.1.4).

### 3.1 Liard River basin

220 The use cases presented here utilize the Liard River basin model built with Raven. The Liard River basin is located in northern Canada, spanning the Yukon Territory, Northwest Territories, British Columbia and Alberta. The Liard River is the largest tributary to the Mackenzie River, with a total contributing area of 275 000 km<sup>2</sup> (Brown and Craig, 2020). The basin includes a variety of landforms, including mountainous regions and wetland-dominated regions. There are varying degrees of difficulty when trying to accurately represent these various landforms in a hydrologic model. Additional details on the Liard River basin, 225 and the corresponding hydrologic model developed for the basin with Raven, can be found in Brown and Craig (2020), which also describes the manual calibration process that was undertaken for the model.

### 3.2 Input file processing

An early step in the model-building process is the collection of data and preparation of model input files (step 1 in **Table 1**). While the supporting data analysis may not require expert knowledge of hydrology *per se*, the data preparation can require 230 a substantial amount of time and effort in the modelling process. Further, the reproducibility and merit of research may rest on the ability to access and reproduce the original and processed intermediate data, which is vastly improved by the use of a scripting environment that in effect documents the steps taken to prepare the data files (Anderson et al., 2018). As such, the use of scripting tools, such as those that will be discussed in this section, may be used to both reduce the effort required to prepare input data files and improve the reproducibility of the research or applied project.

235 This section discusses the preparation of the model structure configuration, the preparation of forcing data and observation data, and modifications to the model discretization file. Additional utilities related to input files exist within the package (such as providing parameter information), but these applications discuss a large proportion of the workflow that would be required in developing a set of Raven input files.

#### 3.2.1 Model configuration

240 One of the key characteristics of Raven is flexibility, including the ability to customize the model structure in terms of the organization of water storage units and selection of process equations (Craig et al., 2020). The hydrologic process schematic is specified through the main Raven input file, in which the number of soil layers, the list of hydrologic processes and the set of ‘to’ and ‘from’ compartments, etc., are defined.

245 ~~For new and even more experienced users of Raven, understanding the model structure and making changes within the primary input file may be a somewhat daunting task, particularly when the model must be initially developed. Fortunately, the Raven User’s Manual (Craig and the Raven Development Team, 2022) provides the templates for a number of model structures, which can be used as a starting point for constructing a customized hydrologic model. These are largely based on emulations of published hydrologic models in the literature (e.g. UBCWM, HMETS, etc.), although some are based on research models that have been developed within Raven (e.g. the Canadian Shield model). Once a base model has been selected, components of the model may be modified 250 using the many process options available within Raven which are documented in the Raven User’s Manual (Craig and the~~

Raven Development Team, 2022). The large number of process options available to the user provide no shortage of model structure tweaks to customize their model. A critical step in making these adjustments to model structure is understanding the structure and ensuring that it is consistent with the modeller's conceptual understanding of the system (step 3 in **Table 1**). While Raven itself does not currently have a user interface deployed that can visualize the model structure, functions within the RavenR package can generate a model schematic from the contents of the model input (\*.rvi) file. The ability to visualize this structure can be critical in understanding the model structure and ensuring the conceptual understanding is consistent with the implemented structure. This can also be used to check for state variables or storage units with an improper number of connections, such as a soil layer with no outflow mechanism.

255 The general workflow within RavenR to generate a model \*.rvi file and visualize the contents is as follows:

260 1. A template model structure is selected and written to file using the `rvn_rvi_write_template` function.

2. The file may be manually modified in consultation with the Raven User's Manual (Craig and the Raven Development Team, 2022)

3. The file may be read into R using the `rvn_rvi_read` function

4. The process connections from the file can be processed using the `rvn_rvi_connections` function


265 5. The process diagram can be generated either in ggplot format using the `rvn_rvi_process_ggplot` function, or as a diagrammer plot using the `rvn_rvi_process_diagrammer` function

6. (optional) the ':CreateRVPTemplate' command can be used to generate a template \*.rvp (parameter) file when Raven is executed

7. (optional) the `rvn_rvi_get_params` function may be used to obtain a data frame of parameters, ranges, and default parameter values for parameters that are included in the hydrologic model, based on the model structure configuration

270

An example of the process diagram is provided for the Liard River basin in **Figure 1**. From this figure, the directional connections between water storage compartments in the model can easily be ascertained and verified, allowing modelers building a new model and modelers inheriting a model alike to quickly understand the movement of water in their current setup. For instance, in the Liard model we can see the model has capacity for precipitation to enter specific wetland and depression compartments, snow can melt and refreeze, and fast and slow (upper and lower) reservoirs exist to represent groundwater processes as different time scales (Brown and Craig, 2020). A single-layer topsoil compartment is used to connect the surface water and subsurface domains in the model along with a vadose zone reservoir to help represent a karst structure within the model. We can see that all processes that move water to glacier are conditional based on the HRU type (glacier HRU). Ponded water is moved to depression storage under the condition of being a wetland and surface water only directly evaporates to the atmosphere if the HRU is a lake. The karst groundwater structure which was implemented in the model is only applicable to a subset of the HRUs which accounts for the conditional connections between SOIL[0], surface water, the vadose reservoir, and



**Figure 1.** The model configuration of the Liard basin, generated from the Liard model \*.rvi input file with the `rvn_rvi_process_diagrammer` function. Solid grey lines indicate connections between state variables, and dashed orange lines indicate conditional connections.

the fast and slow reservoirs. Reviewing and verifying these conditional exceptions along with the connections between other state variables can help ensure that the model is appropriately structured.

Typically, diagrams such as these are arduous to produce for highly-flexible modelling software such as Raven. Here, the 285 `function` has been automated to create publication-ready diagrams for most model setups.

### 3.2.2 Forcing data

Meteorological forcings (e.g. precipitation, temperature, wind speed) drive the hydrologic model responses. When not collected as part of a project, these data are often obtained from online, freely available public sources generally collected, processed, and maintained by local and/or public agencies. These data are likely to require some quality control before ingestion into 290 the model, such as addressing data flags, removing erroneous data, and converting units (step 1 in **Table 1**). This process can be quite tedious, especially when combining multiple data sets of various formats, time steps, and quality. The `RavenR` package offers the `rvn_rvt_write_met` function for writing forcing data directly to the Raven \*.rvt format: the function defaults are configured to accept outputs from the `weathercan` R package, which automatically downloads data for Canadian meteorological stations maintained by Environment Canada (LaZerte and Albers, 2018).

```

fort_liard <- c(60.241711, -123.467377)
stns <- weathercan::stations_search(coords=fort_liard, interval="day",
                                      dist = 500, starts_latest = 1985, ends_earliest = 2006)

weather_dl(stns$station_id[1:10], interval="day",
           start="1985-10-01", end="2005-10-01") %>%
  rvn_met_interpolate(cc=c("max_temp", "min_temp", "total_precip"),
                       key_stn_ids = stns$station_id[1:5]) %>%
  rvn_rvt_write_met()

```

**Figure 2.** Minimum code required for the use case described in **Section 3.2.2** of downloading, interpolating, and writing meteorological data into Raven \*.rvt format using the `weathercan` and `RavenR` R packages. The pipe operator (%>%) from the `dplyr` package is used for readability. Additional code comments are provided in the accompanying repository.

295 In this use case, daily meteorological data for a 20-year period is downloaded, interpolated, and written to Raven \*.rvt format. The `weathercan` R package is used to search for stations within 500 km of Fort Liard and with data records spanning from 1985 to 2005. A subset of stations meeting these criteria is downloaded for pre-processing. Missing values in the meteorological data are then interpolated using data from nearby stations, and a fix is also applied to any interpolated data where the maximum daily temperature is less than the daily minimum. The data from five of the selected stations are then written to Raven \*.rvt format. This workflow would be of substantial time and effort if performed manually or scripts for this task were adapted with each new application; in this use case, the entire workflow is performed with two main functions from the `weathercan` package and two from the `RavenR` package. The code required to accomplish this is provided in **Figure 2**.

300 The advantages of this workflow are 1) the ease of implementation, which can process any number of stations with only a few lines of R code; 2) the transparency and reproducibility of the \*.rvt file generation, which is useful for both review of the data and possible future corrections to all \*.rvt files (e.g., extending the time series to incorporate more recent data). The code may be extended to any supplied set of stations and any meteorological variable that is recognized by Raven. The function also assumes standardized Raven parameter units for all meteorological variables (see reference tables in Appendix C of the Raven User's Manual (Craig and the Raven Development Team, 2022)).

### 3.2.3 Observation data

310 Observation data, such as streamflow records, are generally not required to run hydrologic models; an exception to this may be for truncated model domains, where the model simulates a portion of the watershed and is supplemented by upstream measured flow data. However, observed time series are key to evaluating model performance (history matching) in both calibration and validation exercises and may also be used to enable data assimilation in forecasting applications.

315 Similar to the use of the `weathercan` R package for downloading Canadian meteorological data, the `tidyhydat` R package may be used to download stream gauge data from Canadian stations maintained by the Water Survey of Canada

(Albers, 2017). RavenR provides the `rvn_rvt_tidyhydat` function to process `tidyhydat` inputs directly by wrapping the `rvn_rvt_write` function, which can write any non-meteorological time series to `*.rvt` format. Possible types of time series supported by Raven `*.rvt` files include reservoir inflows, irregular observations, observation weights, temporal reservoir operation rules, etc. The entire list of available formats can be found in the Raven User's Manual (Craig and the Raven 320 Development Team, 2022).

In this use case, the `tidyhydat` package is used to prepare `*.rvt` files of observed streamflow for 9 specified stations (consistent with the stations listed in Table 2 of Brown and Craig (2020)) used in the Liard model. The daily streamflow for these stations are downloaded using `tidyhydat` from 1985 to present day, and written to `*.rvt` format using the `rvn_rvt_tidyhydat` function (a wrapper for the `rvn_rvt_write` function). Raven will automatically exclude any missing values from the calculation of diagnostics, thus missing values in observation data generally do not need to be interpolated or infilled in the same manner that meteorological forcing data needs to be processed. However, the user may still wish to be aware of and avoid large 325 gaps in observation data that may bias the calculation of diagnostic metrics (e.g., consistent winter gaps or multi-year gaps).

```

obs_stns <- read.csv("observation_stations.csv")
tidyhydat::hy_daily_flows(station_number = obs_stns$stnID,
                           start_date = "1985-01-01") %>%
  rvn_rvt_tidyhydat(subIDs=obs_stns$subID)

```


**Figure 3.** Minimum code required for the use case described in [Section 3.2.3](#) of downloading and writing observed flow data into Raven \*.rvt format using the `tidyhydat` and `RavenR` R packages. The pipe operator (`%>%`) from the `dplyr` package is used for readability. Observation station IDs and associated model subbasin IDs are provided in the ‘observation\_stations.csv’ file for brevity. Additional code comments are provided in the accompanying repository.

The same `rvn_rvt_write` function may be used to write other \*.rvt data types by adjusting the `rvt_type` parameter, which may be useful for writing the observation weights generated from the `rvn_gen_obsweights` function to exclude certain data periods from Raven diagnostics, as was done in the Liard model for winter periods with unreliable data records (Brown and Craig, 2020).

### 3.2.4 Model discretization file

The development of distributed and semi-distributed models requires the discretization of a basin into homogeneous units representing hydrologically similar areas. This is typically completed through overlaying a number of spatial data sets which have a dominant effect on the hydrological response of the basin, such as land use, elevation, or soil information (step 2 in [Table 1](#)). In overlaying the spatial data sets, a large number of small computational units, insignificant to the model function, can be created. Since the model runtime is scaled with the number of HRUs, these small areas can increase computational and calibration run times and are not necessary to simulate the dominant hydrological response of the basin. The `RavenR` package offers a way to effectively eliminate small computational units using the `rvn_rvh_cleanhrus` function. This function may merge units based on a set area threshold, and can also merge similar HRUs based on similarity in HRU properties such as landcover, slope, elevation and aspect. HRUs which are significant to the model can be locked or protected. Locked HRUs cannot be removed from the model or increase in size and protected HRUs cannot be removed but may increase in size (to maintain the total watershed area) if other HRUs are removed. This is useful in cases where a point observation is available at a given location (snow survey data) or if the HRUs are part of a significant hydrological response (glaciers).

In this use case, the reduction in the number of model HRUs is demonstrated for a subset of the initial HRUs within subbasin 3 only (initially with 172 HRUs). [In the ‘basic’ reduction of HRUs, a simple area threshold is applied. In subsequent examples](#), HRUs that are of land use type GLACIER are locked HRUs (i.e. cannot be removed or change in size), and HRUs that are either WETLAND or WATER are protected (i.e. cannot be removed but can still increase in size if other HRUs are removed). This operation is applied using area thresholds of [0.5% \(with no locked or protected HRUs\)](#), 0.5% and 2.0% of the subbasin area, resulting in [56](#), 87 and 44 HRUs, respectively. The impact of this operation on land use distribution within the subbasin is summarized in [Figure 4](#).



**Figure 4.** A barplot of total areas by land use for three sets of HRU configurations, including 1) prior to any 'cleaning', 2) following a 'basic' 0.5% area threshold merging criteria with the `rvn_rvh_cleanrhus` function ~~with no locked or protected HRUs~~, 2) following the same operation but with locked and protected HRUs specified, and 3) following a 2.0% area threshold merging criteria. ~~The GLACIER land use is locked, and the WETLAND and WATER land uses are protected~~. The example is done for a single subbasin in the watershed for demonstration purposes, and shows how the land use in the subbasin changes when the removal of subbasins below the area percentage threshold is performed, and when locked/protected HRUs are introduced, using the `rvn_rvh_cleanrhus` function.

In the figure, ~~the total area of the GLACIER type decreases in the basic application, but otherwise remains the same when it is locked~~ the total area of all GLACIER land use type HRUs remains the same, as this land use type was locked. The WATER and WETLAND land use types ~~either decrease or reduce to zero in the basic application, and otherwise increase slightly with each individual cleaning, relative to their proportion of the total area and the total area of removed HRUs based on the respective area threshold~~.

It is noted that, ~~were the WATER and WETLAND HRUs not protected (or locked), the area of these HRUs would be reduced with a 0.5% threshold and removed entirely with a 2.0% threshold~~. Literature has shown that hydrologic areas such as wetlands that are small in size can still have a disproportionately large effect on the hydrologic and biogeochemical response of the watershed (McLaughlin et al., 2014), thus retaining particular HRUs or HRU types may be critical in the cleaning of the HRUs. Finally, the plot shows how the other land use types change with these operations. The FOREST type increases in each case, suggesting that the proportion of small forested HRUs may be small, and that forested HRUs tend to be larger in size. The SHRUBLAND HRUs decrease in represented proportion in each case. This type of analysis could be repeated for other HRU properties (e.g. slope, aspect, vegetation type, etc.). This analysis should be done in conjunction with the use of the `rvn_rvh_cleanrhus` function to

ensure that the reduction in the number of HRUs does not unexpectedly alter the overall representation of HRUs within the  
365 model, and inhibit the ability of the model to capture the dominant hydrologic response.

### 3.3 Output file processing and analysis

A number of functions within RavenR are included to handle the reading of common Raven output files, such as the Hydrographs file (`rvn_hyd_read`), the WatershedStorage file (`rvn_watershed_read`), and other output files (forcings, custom output, etc.). In addition, functions to analyze the output data with typical hydrologic checks and diagnostics are  
370 included in the package. While these functions are built to work with the Raven-specific read functions they are otherwise not specific to Raven, and may be used for any hydrologic model given a means of reading time series output is provided.


This section provides use cases where the realism of the Liard model is assessed, [providing insight to the question, ‘is the model getting the right answers for the right reasons?’](#) (Kirchner, 2006; Euser et al., 2013). These checks provide the modeller with an understanding of the model dynamics and provide more confidence that the model is functioning as expected without  
375 model compensation errors (step 7 in **Table 1**). This section also provides a demonstration of tools for evaluating model performance.

#### 3.3.1 Evaluation of model realism

The flexibility of Raven in the generation of model outputs, including customized outputs that may be specified by the user, can be leveraged to undertake rigorous checks of the hydrologic model. Tools have been built into the RavenR package to  
380 capitalize on this feature, and facilitate a set of model [realism](#) checks. [Here, we define model realism as the model’s ability to replicate and be consistent with anticipated hydrologic behaviour, such as reproducing snowpack measurements, producing reasonable evapotranspiration and runoff coefficients, etc. This definition echoes the one provided in the literature by Euser et al. \(2013\).](#) These checks can be considered semi-automatic, since a script may be deployed to generate the figures but they still require interpretation by a modeller with an understanding of both the natural system and the developed model. Here, the  
385 focus is on the realism of the model to ensure that it is providing hydrologically plausible results; the actual performance of the model [with respect to streamflow](#) is discussed further in **Section 3.3.2**.

The checks that are applied to the Liard River basin in this use case include: 1) plotting the Budyko curve (Budyko, 1974) for the annual average watershed indices, 2) plotting the annual regime curve with monthly averages, 3) examining the stationarity of moisture content in soil storage layers, and 4) plotting the model performance with respect to snowpack storage as  
390 snow water equivalent (SWE). Additional checks supported (not demonstrated here) include: plotting the forcing functions to understand how the inputs may be influencing the model results (i.e. wet and dry years, erroneous temperature readings, etc.), checking the annual water balance, examining baseflow characteristics by comparing modelled and [observed](#)[estimated](#) baseflow [using](#)[from](#) baseflow separation techniques, plotting annual hydrographs in an overlay (i.e. spaghetti plot), and checking the modelled hydrographs and reservoir levels, if any reservoirs or lakes are included in the model.

395 The four plots associated with the stated checks performed in this example are provided in **Figure 5**.



**Figure 5.** Multiple diagnostic plots generated from the RavenR package that may be useful in evaluating the realism of any hydrologic model; A) budyko curve with annual average data points for the watershed, B) a series of regime curves, C) soil storage time series showing the stationarity in long-term storage within soil layers and D) plots of observed and simulated snowpack measurements at Frances River. All data are generated from the Raven model averaged at the watershed scale unless otherwise indicated (i.e. snowpack SWE).

The Budyko plot in **Figure 5A** was generated using the `rvn_budyko_plot` function. The Budyko Curve (Budyko, 1974) shows the relationship that quantifies how mean annual precipitation is partitioned into discharge or evapotranspiration (ET), where the aridity index is plotted on the x-axis and the evaporative index on the y-axis. The Budyko pattern has been observed in multiple catchments around the world (Vereecken et al., 2015). The x-axis of the Budyko curve is the aridity index which represents the ratio between potential evapotranspiration and precipitation. The y-axis is the evaporative index and represents the ratio between actual evapotranspiration and precipitation. The curve is bound by two lines, shown in Figure 4a as the solid lines: the energy limiting line, which is when the evaporative index is equal to the aridity index, and the water limiting line which occurs when the actual evapotranspiration is equal to precipitation. Certain catchment characteristics, such as significant basin storage or the presence of wetlands (which are present in the Liard model), can cause deviations from the Budyko curve. Deviation from the line may also indicate that actual evapotranspiration is underestimated, which may prompt further examination of the model.

In a traditional application of the Budyko Curve it is expected that the plotted points would fall closer to the theoretical line shown in **Figure 5A** as the dashed line. However, certain traits of a basin, such as significant basin storage, can result in deviation from this line. The interpretation of the plot in Figure 4a could also indicate that actual evapotranspiration is being underestimated in the model. Alternatively, if there is significant inter-annual storage in the basin, perhaps due to the presence of wetlands, then the plot would be in line with reality. Ultimately, it is up to the modeller to decide whether this plotted behaviour is expected or if there is a possible misrepresentation of hydrologic processes in the model.

The regime curve can be used to examine the quantities and timing of some of the key model functions. For example, the **Figure 5B** shows that the PET in the Liard model(based on the Hargreaves 1985 calculation method, see Hargreaves and Samani (1985)) peaks at the same time as the AET in June, and maintains a similar pattern over the other months of the year on average. The simulated and observed flows are close in value, and both peaking prior to the peak in precipitation. This aligns with the fact that peaks in the Liard River basin are typically freshet driven. mismatch Mismatches in elements of the regime curve, such as AET and PET or precipitation and flow that is not expected by the modeller would provide a point of investigation and validation into the model.

The soil storage information can be retrieved from Raven in either the `WatershedStorage.csv` file (generated with the `:WriteMassBalanceFile` command), or with the custom output options for specific soil layers (e.g., `:CustomOutput DAILY AVERAGE SOIL[0] ENTIRE_WATERSHED`). Plots such as **Figure 5C** may be applied to any storage compartment in the model to verify the general assumption of long-term stationarity in storage within the hydrologic model, such as lake or reservoir storage. The stationarity assumption for a continuous simulation model is that over a long duration the soil storage should reach a quasi-equilibrium, oscillating around a steady mean. Therefore, a continuous simulation model which is continuously accumulating soil moisture during the simulation period may indicate that, for example, there is insufficient evapotranspiration or baseflow, resulting in the soil storage continuously increasing to compensate for this deficiency. A similar check for reservoir or lake storage could also be warranted. In the plot provided, all four soil storage units have a pattern that generally repeats annually, indicating the storage is stationary in the long term. In addition to verifying the stationarity assumption in storage and examining model compensation effects, **Figure 5C** may be used to better understand the soil dynamics in the model, and ensure that it is consistent with the understanding of the natural system. The pattern indicates that the top soil layer (layer 1) peaks in May-June (presumably with snowmelt) and rapidly depletes, while the fourth soil layer representing deep groundwater storage receives the infiltration at a more delayed rate and has a slower release from storage.


The snow plot provides a method to evaluate the snow balance representation in the model for a particular station. The simulated snow series is produced in Raven with the custom output command (e.g., `:CustomOutput DAILY AVERAGE SNOW BY_HRU`), and is compared against the observed snow course measurements. The plot in **Figure 5D** was generated for the Frances River station. The snow plot provides a visual representation of the model's ability to represent the snow processes and compares it directly to observations. The model provides a reasonable representation of the snowpack SWE with no consistent bias in estimation. ~~The same plots may be created wherever snow measurements are available, and provide a method for directly evaluating the snow routines in the model. In RavenR, the custom output snow is read into R using the `rvn_custom_read` function, and irregular observations may be read using any native R read function (such as `read.table`) or the `rvn_rvt_read` function if they are provided as irregular observations in an `*.rvt` file. Raven may also directly calculate any model diagnostics for irregular snow observations (such as NSE) if they are provided to the model, which may be useful in building objective functions for automatic calibration of the watershed. A similar custom output request for any state variable over a specified temporal and spatial resolution may be produced by Raven at the user's request, and processed using RavenR.~~

~~Although not included in the plots above, examination of the model input forcings is often an insightful step in diagnosing potential model issues and realism. The forcing functions file that is outputted by Raven may be quickly read and visualized with RavenR (i.e. `rvn_forcings_read()` %>% `rvn_forcings_plot`) as a check. Issues such as errors in the temperature record or in the units of precipitation (mm/h instead of mm/d, for example) may be quickly determined with a visual check on the array of forcings that are both supplied to and determined by the Raven model.~~

### 3.3.2 Evaluation of model performance

The RavenR package provides a broad suite of tools for analyzing the results of any Raven hydrologic model, including many tools that can be considered model independent (step 7 in **Table 1**). For example, hydrograph plots, calculation of runoff coefficients, and flow duration curve plots are available within RavenR but may be computed for any time series of flows. The calculation of diagnostics, such as the commonly used Nash-Sutcliffe Efficiency (Nash and Sutcliffe, 1970) and Kling-Gupta Efficiency (Gupta et al., 2009) metrics, are not included in the RavenR package as they can be calculated directly within Raven, and are also available comprehensively in existing packages such as hydroGOF (Mauricio Zambrano-Bigiarini, 2020).

In this use case, a number of diagnostic plots based on simulated and observed hydrographs are presented for the Liard River basin model. These diagnostic plots are computed at the outlet of the Liard River basin (at the outlet near Water Survey of Canada station 10ED1002), and are provided in **Figure 6**. These plots are provided for a portion of the simulation period (where the plot is time-based), and in practice these plots may be applied in both calibration and validation periods for comparison.



**Figure 6.** Multiple diagnostic plots generated from the RavenR package that may be useful in evaluating model performance; A) a hydrograph plot for a subset of the simulation period, B) a scatterplot of simulated and observed annual peak flows, C) plot of timing annual peak timing errors, and D) a plot of cumulative annual flow volumes in time. In plots A and D, the observed is plotted in black and the simulated in orange.

460 In **Figure 6A**, a simple hydrograph plot for a subset of the simulation period is provided. The hydrograph shows good agreement in the magnitude and timing of summer peaks for the years shown as well as the rising limb of the hydrograph, which was the focus of the calibration in the work of Brown and Craig (2020), with a tendency to overestimate the recession from the peak in June until late December/early January. The underestimation tends to continue until the next peak. The hydrograph is shown for a subset of a few years, allowing for a more critical evaluation of the model performance, since examining rather than the entire period, as the full period can obscure the important deviations of the simulated hydrograph from observations and mask deficiencies. Examining a smaller section of the plot (e.g. with the `prd` argument in many RavenR functions), such as only a few years of simulation or a particular event of interest, allows for a more critical examination of the nature of the model errors. A subset of a hydrograph can also be viewed dynamically as a dygraph in RavenR with the `rvn_hyd_dygraph` function, which is supported by the `dygraphs` package (Vanderkam et al., 2018).

465 The peak flow scatterplot (**Figure 6B**) is a scatterplot of the simulated and observed annual peak flows, calculated based on the October 1st water year and produced using the `rvn_annual_peak` function. This figure provides a visual assessment of the performance of modelled peak flow magnitudes, including any systematic bias in over- or under-predicting peaks as a function of peak magnitude. Here, the model appears to estimate peaks with reasonable performance and without systematic bias, although which can be seen as a function of the flow magnitudes themselves (i.e. does the model simulate peak flow magnitudes well at lower and higher flow values?). A model that is able to perfectly simulate peak flow magnitudes will have all points fall on the 1:1 line included in the plot. The plot also provides insight into the nature of the peak error i.e. whether there is a systematic bias in over- or under-predicting peaks, or whether the errors are normally distributed. The plot provided shows that points are relatively close to the 1:1 line and there is no strong evidence of a systematic bias in flow magnitude prediction. However, since only 20 points are included on the plot, additional data may be required to produce conclusions that are statistically valid.

470 While (**Figure 6C**)**Figure 6B** captures the performance with respect to the magnitude of the flow peaks, the timing of peak flows is not assessed. The plot in **Figure 6D****Figure 6C** assesses the error in peak timing (rather than magnitude) with the `rvn_annual_peak_timing_error` function. A perfect model would have all points fall along the zero line, indicating that there is no error in the timing of predicted peaks. The results for the Liard simulation indicate that the model tends to predict peaks slightly later than the observed data, while some of the larger errors in timing tend to be in early peak prediction. In a forecasting framework, a data assimilation technique may reduce the timing (and magnitude) errors that are present in the continuous simulation evaluated here. However, this tendency of the model may still be useful information for forecasters. 475 The use of multiple functions in tandem within RavenR to examine both the peak magnitude and timing errors can be used to evaluate the model performance more comprehensively than a single function (see multiple RavenR functions named as `rvn_annual_*`).

480 Finally, **Figure 6D** provides a comparison of cumulative flow volumes between the simulated and observed model in time. This plot is generated by the `rvn_cum_plot_flow` function. The plot shows clearly where deviations in the overall volume arise in time. For example, the plot for the Liard model shows that the December-March winter period of each year is a time of deviation in cumulative volumes, while the freshet-driven summer peak periods tend to match volume quite well overall. This is likely a result of the calibration procedure in (Brown and Craig, 2020), where ice affected flows in the winter were not considered in the

calibration procedure due to high levels of uncertainty associated with the measurements. Additional functions that examine the relative volumes of simulated and observed flows, but aggregate them rather than examining the differences in time, are the 495 `rvn_monthly_vbias` and the `rvn_annual_volume` functions, which provide the monthly average volume differences and the annual volume differences in a scatterplot for each year, respectively. The volume is generally a useful diagnostic metric as it integrates the modelled hydrograph performance in time, and allows the modeller to identify periods of poor cumulative error or systematic errors (e.g. underestimating overall volume) that may be not clear or obvious when only examining flows.

## 4 Conclusions

500 This paper presented the `RavenR` package, an R-based set of tools that is designed to support the development, use, and analysis of hydrologic models developed using Raven but can be readily adapted for any hydrologic modelling output. `RavenR` is a free, open-source software that is intended to support the wealth of options in a flexible modelling framework while maintaining or improving the transparency and reproduciblity of the analyses undertaken by the modeller.

The tools within `RavenR` may be used in any stage of the typical modelling workflow. Although the tools are designed for 505 use with Raven, the analysis and utility functions may be useful in conjunction with any hydrologic model that has similar requirements and workflows as Raven. The `RavenR` tools provide the means for preparing Raven input files, visualizing and processing input data, executing Raven, and generating a vast array of model checks and performance-related graphics from the Raven output files. All functions in the package are supplemented by additional information and examples (consistent with CRAN requirements), and the package is further accompanied by the introductory documentation in the form of a vignette. 510 This paper illustrates how the `RavenR` functions may be used in both academic and industrial projects, including generating model input `*.rvt` files, visualizing the model structure, and exploring and assessing the hydrologic model results. This includes aiding the modeller in building an understanding and trust of the constructed hydrologic model.

A set of `RavenR` use cases are presented for the Liard River basin, for which a Raven model has previously been built and 515 thoroughly tested (Brown and Craig, 2020). The use cases present how a subset of tools may be used to generate input files for, or analyze the results of, the Raven model of the Liard river basin. The examples are bolstered by an interpretation of the functions and results, which may be useful in interpreting and building confidence in any hydrologic model. The accompanying data repository and code for this manuscript can fully recreate the figures and analyses presented in the use cases, demonstrating best practices for reproducibility in hydrologic and scientific publications.

Due to the open-source nature of the Raven project, the code is transparent and accessible to users and is being continuously 520 supplemented with new functionalities and improvements. Similarly, the `RavenR` package is open-source and is in active development. It is anticipated that the `RavenR` project will also continue to improve and expand its functionality in order to meet its goal of supporting Raven users from all backgrounds and experience levels while improving the reproducibility of their work.

525 *Code and data availability.* The RavenR package is free and open-source software, and the version of the package (v.2.1.4) used to produce  
the results of this paper is archived on Zenodo (Chlumsky et al., 2021a). All R code and data used to generate the results and figures  
presented in this manuscript is also archived on Zenodo at [doi 10.5281/zenodo.5534817](https://doi.org/10.5281/zenodo.5534817) (Chlumsky et al., 2022), and is also available on  
Github ([https://github.com/rchlumsk/RavenR\\_manuscript\\_2021](https://github.com/rchlumsk/RavenR_manuscript_2021)). The RavenR package is currently available as v2.1.4 on the Comprehensive  
R Archive Network (CRAN) (<https://cran.r-project.org/package=RavenR>), and the development version of the package is also available on  
Github (<https://github.com/rchlumsk/RavenR>). The Raven hydrologic modelling framework is open-source and may be downloaded from  
530 <http://raven.uwaterloo.ca/>.

*Author contributions.* RC initiated the concept of the RavenR package. RC and JRC contributed the bulk of the package scripts, with  
contributions and development efforts from all authors. GB and JRC provided the Liard River model files. The use cases were prepared by  
RC, LS, SL, SG, and GB. The article was prepared by RC with contributions from all authors.

*Competing interests.* The authors declare that they have no conflict of interest.

535 *Acknowledgements.* Mr. Chlumsky would like to acknowledge the financial support provided by the NSERC Canada Graduate Scholarship  
(CGSD3-558879-2021) and the Engineering Excellence Doctoral Fellowship provided at the University of Waterloo that helped make this  
work possible. The authors would like to thank all those who have contributed to the RavenR project since its inception, including Larry  
(Haobo) Liu and Joel Trubilowicz. [The authors would also like to thank Paul C. Astagneau and one anonymous reviewer for their comments  
on improving our manuscript and software.](#)

Albers, S.: tidyhydat: Extract and Tidy Canadian Hydrometric Data, *The Journal of Open Source Software*, 2, <https://doi.org/10.21105/joss.00511>, 2017.

Anderson, E., Chlumsky, R., McCaffrey, D., Trubilowicz, J., Shook, K. R., and Whitfield, P. H.: R-functions for Canadian hydrologists: a Canada-wide collaboration, *Canadian water resources journal*, 44, 108–112, 2018.

545 Astagneau, P. C., Thirel, G., Delaigue, O., Guillaume, J. H. A., Parajka, J., Brauer, C. C., Viglione, A., Buytaert, W., and Beven, K. J.: Technical note: Hydrology modelling R packages – a unified analysis of models and practicalities from a user perspective, *Hydrology and Earth System Sciences*, 25, 3937–3973, <https://doi.org/10.5194/hess-25-3937-2021>, 2021.

Baker, M.: 1,500 scientists lift the lid on reproducibility, *Nature*, 533, 452–454, <https://doi.org/10.1038/533452a>, 2016.

Brown, G. and Craig, J. R.: Structural calibration of an semi-distributed hydrological model of the Liard River basin, *Canadian Water Resources Journal / Revue canadienne des ressources hydriques*, pp. 1–17, <https://doi.org/10.1080/07011784.2020.1803143>, 2020.

550 Budyko, M. I.: Climate and life, *International Geophysics Series*, 18, 508, 1974.

Camerer, C. F., Dreber, A., Holzmeister, F., Ho, T.-H., Huber, J., Johannesson, M., Kirchler, M., Nave, G., Nosek, B. A., Pfeiffer, T., Altmejd, A., Buttrick, N., Chan, T., Chen, Y., Forsell, E., Gampa, A., Heikensten, E., Hummer, L., Imai, T., Isaksson, S., Manfredi, D., Rose, J., Wagenmakers, E.-J., and Wu, H.: Evaluating the replicability of social science experiments in *Nature* and *Science* between 2010 and 2015, *Nature Human Behaviour*, 2, 637–644, <https://doi.org/10.1038/s41562-018-0399-z>, 2018.

555 Chadalawada, J., Herath, H. M. V. V., and Babovic, V.: Hydrologically Informed Machine Learning for Rainfall-Runoff Modeling: A Genetic Programming-Based Toolkit for Automatic Model Induction, *Water Resources Research*, 56, 2020.

Chawanda, C. J., George, C., Thiery, W., van Griensven, A., Tech, J., Arnold, J., and Srinivasan, R.: User-friendly workflows for catchment modelling: Towards reproducible SWAT+ model studies, *Environmental Modelling & Software*, 134, 104812, <https://doi.org/10.1016/j.envsoft.2020.104812>, 2020.

560 Chlumsky, R., Craig, J. R., Brown, G., Scantlebury, L., Grass, S., Lin, S., and Arabzadeh, R.: rchlumsk/RavenR: v2.1.4 release, <https://doi.org/10.5281/zenodo.5525041>, 2021a.

Chlumsky, R., Mai, J., Craig, J. R., and Tolson, B. A.: Simultaneous Calibration of Hydrologic Model Structure and Parameters Using a Blended Model, *Water Resources Research*, 57, e2020WR029229, <https://doi.org/10.1029/2020WR029229>, 2021b.

565 Chlumsky, R., Craig, J. R., Brown, G., Scantlebury, L., Grass, S., Lin, S., and Arabzadeh, R.: rchlumsk/RavenR\_manuscript\_2021: Initial pre- release v0.2, <https://doi.org/10.5281/zenodo.6421692>, 2022.

Clark, M. P., Slater, A. G., Rupp, D. E., Woods, R. A., Vrugt, J. A., Gupta, H. V., Wagener, T., and Hay, L. E.: Framework for Understanding Structural Errors (FUSE): A modular framework to diagnose differences between hydrological models, *Water Resources Research*, 44, <https://doi.org/10.1029/2007WR006735>, 2008.

570 Clark, M. P., Kavetski, D., and Fenicia, F.: Pursuing the method of multiple working hypotheses for hydrological modeling, *Water Resources Research*, 47, <https://doi.org/10.1029/2010WR009827>, 2011.

Clark, M. P., Nijssen, B., Lundquist, J. D., Kavetski, D., Rupp, D. E., Woods, R. A., Freer, J. E., Gutmann, E. D., Wood, A. W., Brekke, L. D., Arnold, J. R., Gochis, D. J., and Rasmussen, R. M.: A unified approach for process-based hydrologic modeling: 1. Modeling concept, *Water Resources Research*, 51, 2498–2514, 2015.

575 Coxon, G., Freer, J., Lane, R., Dunne, T., Knoben, W. J. M., Howden, N. J. K., Quinn, N., Wagener, T., and Woods, R.: DE-  
CIPHeR v1: Dynamic fluxEs and ConnectIvity for Predictions of HydRology, *Geoscientific Model Development*, 12, 2285–2306,  
<https://doi.org/10.5194/gmd-12-2285-2019>, 2019.

Craig, J. R. and the Raven Development Team: Raven: User's and Developer's Manual v3.5, <http://raven.uwaterloo.ca/>, 2022.

Craig, J. R., Brown, G., Chlumsky, R., Jenkinson, R. W., Jost, G., Lee, K., Mai, J., Serrer, M., Sgro, N., Shafii, M., Snowdon, A. P., and  
580 Tolson, B. A.: Flexible watershed simulation with the Raven hydrological modelling framework, *Environmental Modelling and Software*,  
129, <https://doi.org/10.1016/j.envsoft.2020.104728>, 2020.

Csardi, G. and Nepusz, T.: The igraph software package for complex network research, *InterJournal, Complex Systems*, 1695, <http://igraph.org>, 2006.

Dal Molin, M., Kavetski, D., and Fenicia, F.: SuperflexPy 1.3.0: an open-source Python framework for building, testing, and improving  
585 conceptual hydrological models, *Geoscientific Model Development*, 14, 7047–7072, <https://doi.org/10.5194/gmd-14-7047-2021>, 2021.

Delaigue, O., Thirel, G., Coron, L., and Brigode, P.: airGR and airGRteaching: Two Open-Source Tools for Rainfall-Runoff Modeling and  
Teaching Hydrology, in: HIC 2018. 13th International Conference on Hydroinformatics, edited by La Loggia, G., Freni, G., Puleo, V., and  
De Marchis, M., vol. 3 of *EPiC Series in Engineering*, pp. 541–548, EasyChair, <https://doi.org/10.29007/qsqj>, 2018.

Delaigue, O., Coron, L., and Brigode, P.: airGRteaching: Teaching Hydrological Modelling with GR (Shiny Interface Included),  
590 <https://doi.org/10.15454/WOSSKT>, r package version 0.2.9.25, 2020.

Euser, T., Winsemius, H. C., Hrachowitz, M., Fenicia, F., Uhlenbrook, S., and Savenije, H. H. G.: A framework to assess the realism of  
model structures using hydrological signatures, *Hydrology and Earth System Sciences*, 17, 1893–1912, <https://doi.org/10.5194/hess-17-1893-2013>, 2013.

Fenicia, F., Savenije, H. H. G., Matgen, P., and Pfister, L.: Understanding catchment behavior through stepwise model concept improvement,  
595 *Water Resources Research*, 44, <https://doi.org/10.1029/2006WR005563>, 2008.

Fenicia, F., Kavetski, D., and Savenije, H. H. G.: Elements of a flexible approach for conceptual hydrological modeling: 1. Motivation and  
theoretical development, *Water Resources Research*, 47, <https://doi.org/10.1029/2010WR010174>, 2011.

Grolemund, G. and Wickham, H.: Dates and Times Made Easy with lubridate, *Journal of Statistical Software*, 40, 1–25, <https://www.jstatsoft.org/v40/i03/>, 2011.

600 Guo, D., Westra, S., and Peterson, T.: Evapotranspiration: Modelling Actual, Potential and Reference Crop Evapotranspiration, <https://CRAN.R-project.org/package=Evapotranspiration>, r package version 1.15, 2020.

Gupta, H. V., Kling, H., Yilmaz, K. K., and Martinez, G. F.: Decomposition of the mean squared error and NSE performance criteria: Implications for improving hydrological modelling, *Journal of Hydrology*, 377, 80–91, <https://doi.org/10.1016/j.jhydrol.2009.08.003>, 2009.

Hargreaves, G. H. and Samani, Z. A.: Reference Crop Evapotranspiration from Temperature, *Applied Engineering in Agriculture*, 1, 96–99,  
605 <https://doi.org/10.13031/2013.26773>, 1985.

Hijmans, R. J., van Etten, J., Sumner, M., Cheng, J., Baston, D., Bevan, A., Bivand, R., Busetto, L., Canty, M., Fasoli, B., Forrest, D., Ghosh,  
A., Golicher, D., Gray, J., Greenberg, J. A., Hiemstra, P., Hingee, K., for Mathematics Applied Geosciences, I., Karney, C., Mattiuzzi,  
M., Mosher, S., Naimi, B., Nowosad, J., Pebesma, E., Lamigueiro, O. P., Racine, E. B., Rowlingson, B., Shortridge, A., Venables, B., and  
Wueest, R.: raster: Geographic Data Analysis and Modeling, <https://rspatial.org/raster>, r package version 3.4-13, 2021.

610 Hoey, S. V., Seuntjens, P., van Der Kwast, J., and Nopens, I.: A qualitative model structure sensitivity analysis method to support model  
selection, *Journal of Hydrology*, 519, 3426–3435, <https://doi.org/10.1016/j.jhydrol.2014.09.052>, 2014.

Hutton, C., Wagener, T., Freer, J., Han, D., Duffy, C., and Arheimer, B.: Most computational hydrology is not reproducible, so is it really science?, *Water Resources Research*, 52, 7548–7555, <https://doi.org/10.1002/2016WR019285>, 2016.

Hydrologic Engineering Center: HEC-HMS User's Manual version 4.8., <https://www.hec.usace.army.mil/confluence/hmsdocs/hmsum/4.8>, 615 2020.

Jackson, E. K., Roberts, W., Nelsen, B., Williams, G. P., Nelson, E. J., and Ames, D. P.: Introductory overview: Error metrics for hydrologic modelling – A review of common practices and an open source library to facilitate use and adoption, *Environmental Modelling & Software*, 119, 32–48, <https://doi.org/10.1016/j.envsoft.2019.05.001>, 2019.

Kavetski, D. and Fenicia, F.: Elements of a flexible approach for conceptual hydrological modeling: 2. Application and experimental insights, 620 *Water resources research*, 47, 2011.

Kirchner, J. W.: Getting the right answers for the right reasons: Linking measurements, analyses, and models to advance the science of hydrology, *Water Resources Research*, 42, <https://doi.org/https://doi.org/10.1029/2005WR004362>, 2006.

Knoben, W. J. M., Freer, J. E., Fowler, K. J. A., Peel, M. C., and Woods, R. A.: Modular Assessment of Rainfall–Runoff Models Toolbox (MARRMoT) v1.2: an open-source, extendable framework providing implementations of 46 conceptual hydrologic models as continuous 625 state-space formulations, *Geoscientific model development*, 12, 2463–2480, 2019.

Knoben, W. J. M., Freer, J. E., Peel, M. C., Fowler, K. J. A., and Woods, R. A.: A brief analysis of conceptual model structure uncertainty using 36 models and 559 catchments, *Water Resources Research*, <https://doi.org/10.1029/2019WR025975>, 2020.

LaZerte, S. E. and Albers, S.: weathercan: Download and format weather data from Environment and Climate Change Canada, *The Journal of Open Source Software*, 3, 571, <https://joss.theoj.org/papers/10.21105/joss.00571>, 2018.

Leavesley, G. H., Markstrom, S. L., Restrepo, P. J., and Viger, R. J.: A modular approach to addressing model design, scale, and parameter estimation issues in distributed hydrological modelling, *Hydrological Processes*, 16, 173–187, 2002.

Mai, J., Craig, J. R., and Tolson, B. A.: Simultaneously determining global sensitivities of model parameters and model structure, *Hydrology and Earth System Sciences*, 24, 5835–5858, <https://doi.org/10.5194/hess-24-5835-2020>, 2020.

Mauricio Zambrano-Bigiarini: hydroGOF: Goodness-of-fit functions for comparison of simulated and observed hydrological time series, 635 <https://doi.org/10.5281/zenodo.839854>, r package version 0.4-0, 2020.

McLaughlin, D. L., Kaplan, D. A., and Cohen, M. J.: A significant nexus: Geographically isolated wetlands influence landscape hydrology, *Water Resources Research*, 50, 7153–7166, <https://doi.org/10.1002/2013WR015002>, 2014.

Nash, J. E. and Sutcliffe, J. V.: River flow forecasting through conceptual models part I — A discussion of principles, *Journal of Hydrology*, 10, 282–290, [https://doi.org/10.1016/0022-1694\(70\)90255-6](https://doi.org/10.1016/0022-1694(70)90255-6), 1970.

National Academies of Sciences, Engineering, and Medicine: Reproducibility and Replicability in Science, The National Academies Press, 640 Washington, D.C., <https://doi.org/10.17226/25303>, 2019.

Orellana, B., Pechlivanidis, I., McIntyre, N., Wheater, H., and Wagener, T.: A Toolbox for the Identification of Parsimonious Semi-Distributed Rainfall-Runoff Models: Application to the Upper Lee Catchment, in: International Congress on Environmental Modelling and Software, <https://scholarsarchive.byu.edu/cgi/viewcontent.cgi?article=2723&context=iemssconference>, 2008.

Pebesma, E.: Simple Features for R: Standardized Support for Spatial Vector Data, *The R Journal*, 10, 439–446, <https://doi.org/10.32614/RJ-2018-009>, 2018.

Perrin, C., Michel, C., and Andréassian, V.: Improvement of a parsimonious model for streamflow simulation, *Journal of Hydrology*, 279, 275–289, 2003.

650 Pilz, T., Francke, T., Baroni, G., and Bronstert, A.: How to Tailor my Process-based Hydrological Model? Dynamic Identifiability Analysis of Flexible Model Structures, Water resources research, 2020.

Pohlert, T.: Non-Parametric Trend Tests and Change-Point Detection, <https://cran.r-project.org/package=trend>, r package version 1.1.4, 2020.

R Core Team: R: A Language and Environment for Statistical Computing, R Foundation for Statistical Computing, Vienna, Austria, <https://www.R-project.org/>, 2021.

655 Remmers, J. O., Teuling, A. J., and Melsen, L. A.: Can model structure families be inferred from model output?, Environmental modelling and software : with environment data news, pp. 104 817–, 2020.

Ryan, J. A. and Ulrich, J. M.: xts: eXtensible Time Series, <https://CRAN.R-project.org/package=xts>, r package version 0.12.1, 2020.

Shook, K., Whitfield, P., Chlumsky, R., Albers, S., and Munoz, V.: Canadian Hydrological Analyses, <https://github.com/CSHS-CWRA/CSHShydRology>, r package version 1.1.2, 2021.

660 Slater, L. J., Thirel, G., Harrigan, S., Delaigue, O., Hurley, A., Khouakhi, A., Prosdocimi, I., Vitolo, C., and Smith, K.: Using R in hydrology: a review of recent developments and future directions, Hydrology and earth system sciences, 23, 2939–2963, 2019.

Spieler, D., Mai, J., Craig, J. R., Tolson, B. A., and Schütze, N.: Automatic Model Structure Identification for Conceptual Hydrologic Models, Water resources research, 56, 2020.

Stroustrup, B.: The C++ programming language, Addison-Wesley, Upper Saddle River, NJ, fourth edition. edn., 2013.

Van Rossum, G. and Drake, F. L.: Python 3 Reference Manual, CreateSpace, Scotts Valley, CA, 2009.

665 Vanderkam, D., Allaire, J., Owen, J., Gromer, D., and Thieurmel, B.: dygraphs: Interface to 'Dygraphs' Interactive Time Series Charting Library, <https://CRAN.R-project.org/package=dygraphs>, r package version 1.1.1.6, 2018.

Vereecken, H., Huisman, J. A., Hendricks Franssen, H. J., Brüggemann, N., Bogena, H. R., Kollet, S., Javaux, M., van der Kruk, J., and Vanderborght, J.: Soil hydrology: Recent methodological advances, challenges, and perspectives, Water Resources Research, 51, 2616–2633, <https://doi.org/10.1002/2014WR016852>, 2015.

670 Werner, M., Schellekens, J., Gijsbers, P., van Dijk, M., van den Akker, O., and Heynert, K.: The Delft-FEWS flow forecasting system, Environmental Modelling & Software, 40, 65–77, <https://doi.org/https://doi.org/10.1016/j.envsoft.2012.07.010>, 2013.

Wickham, H.: ggplot2: Elegant Graphics for Data Analysis, Springer-Verlag New York, <https://ggplot2.tidyverse.org>, 2016.

Wickham, H.: tidyverse: Tidy Messy Data, <https://CRAN.R-project.org/package=tidyr>, r package version 1.1.3, 2021.

675 Wickham, H., François, R., Henry, L., and Müller, K.: dplyr: A Grammar of Data Manipulation, <https://CRAN.R-project.org/package=dplyr>, r package version 1.0.5, 2021a.

Wickham, H., Hester, J., and Chang, W.: devtools: Tools to Make Developing R Packages Easier, <https://CRAN.R-project.org/package=devtools>, r package version 2.4.0, 2021b.