



Atmosphere-Ocean-Aerosol-Chemistry-Climate Model SOCOLv4.0: description and evaluation

Timofei Sukhodolov^{1,2,3}, Tatiana Egorova^{1,2}, Andrea Stenke², William T. Ball⁴, Christina Brodowsky², Gabriel Chiodo^{2,5}, Aryeh Feinberg^{2,6,7}, Marina Friedel², Arseniy Karagodin-Doyennel^{1,2}, Thomas Peter², Sandro Vattioni², and Eugene Rozanov^{1,2,3}

¹Physikalisch-Meteorologisches Observatorium Davos and World Radiation Center, Davos, Switzerland

²Institute for Atmospheric and Climate Science, ETH Zurich, Zurich, Switzerland

³St. Petersburg State University, St. Petersburg, Russia.

⁴Department of Geoscience and Remote Sensing, Faculty of Civil Engineering and Geosciences, TU Delft, Delft, the Netherlands

⁵Department of Applied Physics and Applied Mathematics, Columbia University, New York (NY), USA

⁶Institute of Biogeochemistry and Pollutant Dynamics, ETH Zurich, Zurich, Switzerland

⁷Eawag, Swiss Federal Institute of Aquatic Science and Technology, Dübendorf, Switzerland

Correspondence to: Timofei Sukhodolov (timofei.sukhodolov@pmodwrc.ch)

Abstract. This paper features the new Atmosphere-Ocean-Aerosol-Chemistry-Climate Model SOCOLv4.0 and its validation. The new model was built by interactively coupling the MPI-ESM1.2 Earth System Model (T63, L47) with the chemistry (99 species) and size-resolving (40 bins) sulfate aerosol microphysics modules from the Aerosol-Chemistry-Climate Model SOCOL-AERv2. We evaluate its performance against reanalysis products and observations of atmospheric circulation, temperature, and trace gases distribution, with a focus on stratospheric processes. We show that SOCOLv4.0 captures the low- and mid-latitude stratospheric ozone well in terms of the climatological state, variability and evolution. The model provides an accurate representation of climate change, showing a global surface warming trend consistent with observations as well as realistic cooling in the stratosphere caused by greenhouse gas emissions, although, as in previous model versions, a too fast residual circulation and exaggerated mixing in the surf zone are still present. The stratospheric sulfur budget for moderate volcanic activity is well represented by the model, albeit with slightly underestimated aerosol lifetime after major eruptions. The presence of the interactive ocean and a successful representation of recent climate and ozone layer trends make SOCOLv4.0 ideal for studies devoted to future ozone evolution and effects of greenhouse gases and ozone-destroying substances, as well as the evaluation of potential solar geoengineering measures through sulfur injections. Potential further model improvements could be to increase the vertical resolution, which is expected to allow better meridional transport in the stratosphere, as well as to update the photolysis calculation module and budget of mesospheric odd nitrogen. In summary, this paper demonstrates that SOCOLv4.0 is well suited for applications related to the stratospheric ozone and sulfate aerosol evolution, including its participation in ongoing and future model intercomparison projects.



1 Introduction

Global modelling of the atmosphere and its interaction with oceans, cryosphere, biosphere, and land surface dates back several decades (e.g., Manabe and Bryan, 1969). The numerical approximation of each of these Earth system components is a complex task by itself and therefore their development often began and continued independently from each other, with simplified descriptions of missing, but important, processes included in the form of boundary conditions. With the rapid development of computational facilities and methods, the models grew in their complexity in terms of the number of processes described and the quality of their description. Motivated primarily through the context of climate change research, scientific advances in global numerical modelling have shown that, even though stand-alone approximations are still sufficient for some specific modelling tasks, the interaction of Earth system components is required for reasonable model performance in most cases. These advances can be tracked in the history of the Climate Model Intercomparison Project requirements for models in its different phases (<https://www.wcrp-climate.org/wgcm-cmip>). Due to the importance of the dynamical links between the troposphere and the stratosphere (Kidston et al., 2015), many climate groups have also extended their state-of-the-art models vertically into the mesosphere (e.g., Manzini et al., 2006) or even the thermosphere (e.g., WACCM (Marsh et al., 2013) and HAMMONIA (Schmidt et al., 2006)). For example, variations in the stratospheric polar night jet can induce significant changes in surface weather on timescales ranging from daily to long-term climate effects (e.g., Gerber et al., 2012 and references therein). Variability of this dynamical coupling can be induced by the earth system itself, i.e., responding to ocean temperature changes and vertically propagating wave-forcing from the troposphere, or by external factors such as volcanic eruptions, variations in the solar UV irradiance, and greenhouse gas changes (Kidston et al., 2015).

Middle atmosphere studies are closely linked to the representation of atmospheric chemistry since the ozone layer primarily determines the temperature structure of the stratosphere through the absorption of solar ultraviolet (UV) irradiance. This has an influence on the general circulation of the stratosphere and subsequently also on the tropospheric climate. Stratospheric ozone itself is influenced by many factors, such as the heterogeneous chemistry intensification after volcanic eruptions (e.g., Revell et al., 2016) or the acceleration of ozone destruction cycles after energetic particle precipitation events (Rozanov et al., 2012; Mironova et al., 2015). Moreover, it is also largely affected by climate change, via radiatively-induced changes in upper stratospheric chemistry, as well as changes in the Brewer Dobson circulation (BDC, Chiodo et al., 2018). Changes in ozone can also in turn affect the BDC (e.g., Polvani et al., 2019). Therefore, changes in stratospheric ozone and dynamics feedback on each other. Ozone-circulation feedbacks have been assessed in several studies, showing their importance for stratosphere-troposphere coupling in mid-winter (Haase and Matthes, 2019; Oehrlein et al., 2020) and polar stratospheric temperature variability in springtime (Rieder et al., 2019). Long-term stratospheric ozone variations (e.g., depletion and recovery) are able to significantly affect the tropospheric climate (Previdi and Polvani, 2014; Brönnimann et al., 2017), and even the climate response to global warming might be biased if ozone feedbacks are not taken into account (Nowack et al., 2015).

The main driving issue in middle atmosphere chemistry research was the discovery of the ozone hole in the 1980s (Farman et al., 1985). The ozone layer plays an important role in shielding the biosphere from dangerous solar ultraviolet radiation and



65 the risk of related increasing cases of skin cancer and other diseases induced progress in atmospheric ozone science that led to
strong limitations on the production of halogen containing ozone depleting substances (hODS) in 1987 through the Montreal
Protocol and its Amendments. Since then, observations and models have demonstrated the positive role of these restrictive
measures (e.g. Velders et al., 2007; Egorova et al., 2013) and some signs of the ozone recovery have already been observed
(Chipperfield et al., 2017). However, the expected recovery in the lower stratosphere has been questioned, based on the updated
70 observations (Ball et al., 2018; 2019). This issue is one of many requiring further investigation and deeper understanding.
Other issues and research fields include the appearance of an unprecedentedly large ozone hole over the Northern hemisphere
in spring 2020 (Witze, 2020; Manney et al., 2020); the formation of a large and deep Antarctic ozone hole in autumn 2020
(NASA Ozone Watch, <https://ozonewatch.gsfc.nasa.gov/>); continuous unexpected CFC-11 emissions (Fleming et al., 2020);
a potential decline of the solar activity (Arsenovic et al., 2018); the potential stratospheric injection of sulfur-containing species
75 for solar geoengineering purposes (Tilmes et al., 2009; Vattioni et al., 2019); and a potential impact of increasing trends of
iodine in the stratosphere (Koenig et al., 2020). These examples underline that our understanding of atmospheric ozone
specifically, and atmospheric chemistry in general, is far from being fully resolved and inspires further model developments
and studies of the ozone layer evolution, in the present and future.

The need to represent the large number of processes involved in the state evolution of the ozone layer led to the development
80 of atmospheric chemistry models ranging from simple box models to chemistry-transport models and finally to chemistry-
climate models that include at least interactive chemistry and atmospheric dynamics, but may include ocean dynamics, aerosol
microphysics and other components (<https://www.sparc-climate.org/activities/ccm-initiative>). The chemistry climate model
SOCOL (SOlar Climate Ozone Links) was initially developed for studies related to the ozone layer (Egorova et al., 2005).
Through its versions from v1 to v3 it was used with prescribed sea surface temperature and sea ice coverage fields, advancing
85 over time in terms of model numerics, stratospheric chemistry, and transport representation. Since the publication of the base
version SOCOLv3 (Stenke et al., 2013), the atmospheric-chemistry part has undergone many further improvements, such as
an addition of the volatile organic compound (VOC) chemistry, an interactive lightning NO_x parameterization, corrections in
schemes for solar heating rates and photolysis rates, parameterization of energetic particles, and interactive deposition schemes.
The base version of Stenke et al. (2013), however, further branched into two significant sub-versions SOCOL-MPIOM with
90 interactive ocean (Muthers et al., 2014) and SOCOL-AER with interactive aerosol microphysics (Sheng et al., 2015), each of
them receiving further, independent, upgrades (Arsenovic et al., 2018; Feinberg et al., 2019), and several smaller variants,
such as an improved tropospheric ozone budget (Revell et al., 2015, Revell et al., 2018), detailed methane sources and sinks
(Feinberg et al., 2018), and atmospheric selenium cycling (Feinberg et al., 2020).

The natural next step was to combine the multiple improvements and model versions into a single fourth version of the SOCOL
95 model by coupling these updated modules onto an upgraded dynamical core, since the core itself also underwent many
improvements in recent years. As a basis for this, we used the Max Planck Institute Earth System Model (MPI-ESM1.2), so
that the chemistry (MEZON) and aerosol (AER) models are attached to the atmosphere- (ECHAM6.3) ocean- (MPIOM1.6.3)
land surface- (JSBACH3.2) ocean biogeochemistry- (HAMOCC6) model coupled through the OASIS3-MCT coupler. In this



paper we describe the new Atmosphere-Ocean-Aerosol-Chemistry-Climate SOCOLv4.0 model in detail (section 2) and
100 validate its performance against available observations and reanalysis products. The main motivation is to provide a solid
reference of model performance for future improvements and applications, including model intercomparison projects (MIPs).
The validation is split into two main parts: atmospheric dynamics (section 3.2) and atmospheric chemistry with the primary
focus on stratospheric ozone (section 3.3).

2 Model description

105 SOCOLv4.0 (SOCOLv4 hereafter) consists of the Earth System Model MPI-ESM1.2 (Mauritsen et al., 2019), the chemistry
model MEZON (Egorova et al., 2003), and the sulfate aerosol microphysical model AER (Weisenstein et al., 1997), with all
these parts being interactively coupled to each other, as schematically presented in Fig. 1. In simple terms, chemistry and
aerosol microphysics rely on atmospheric temperature, winds, and relative humidity and in turn influence the atmosphere and
ocean through the short- and long-wave radiation schemes, while aerosol microphysics depend on sulfur chemistry and provide
110 the aerosol surface area density and number density necessary for heterogeneous chemistry calculations. Transport of
individual gases and aerosols is performed by the flux-form semi-Lagrangian scheme of Lin and Rood (1996) in the dynamical
core of ECHAM6 that has remained unchanged from its predecessor ECHAM5. Transport is calculated every dynamical time
step (15 min). The dry and wet deposition of gases and aerosols is also based on the ECHAM6 parameters such as near-surface
turbulence and precipitation. In the following we discuss each of these main components separately and describe the latest
115 changes.

SOCOLv4 is based on the low resolution (LR) configuration of the MPI-ESM model. This configuration corresponds to a
spectral truncation at T63 providing an approximate horizontal grid spacing of $1.9^\circ \times 1.9^\circ$. The vertical resolution of the
atmosphere is set to 47 levels from the surface to 0.01 hPa, using a hybrid sigma-pressure coordinate system. Although other
higher horizontal and vertical resolutions of MPI-ESM are also tuned and available for use, we chose the LR configuration
120 since it is the most used and better tuned one (Mauritsen et al., 2019), and better suited for long-term climate simulations in
terms of required computational resources and storage.




Figure 1. Components and information flow in the atmosphere-ocean-aerosol-chemistry-climate-model SOCOLv4. Green boxes symbolize prescribed boundary conditions.

125 2.1 Earth system model MPI-ESM1.2

The earth system model MPI-ESM1.2 (Mauritsen et al., 2019) is a further development of its predecessor MPI-ESM (Giorgetta et al., 2013). The main components of MPI-ESM are highlighted by the dark blue boxes in Fig. 1. The ocean dynamical model, MPIOM1.6.3, transports tracers of the ocean biogeochemistry model, HAMOCC6. The atmosphere model, ECHAM6.3, is directly coupled to the land model, JSBACH3.2, through surface exchange of mass, momentum, and heat. These two major model blocks are then coupled via the OASIS3-MCT coupler (Craig et al., 2017). The coupler aggregates, interpolates, and exchanges fluxes and state variables once a day between ECHAM6-JSBACH and MPIOM-HAMOCC. Here we only describe the latest states of the MPI-ESM components that are used in SOCOLv4 and do not focus on the differences between MPI-ESM versions, as this is already discussed in greater detail by Mauritsen et al. (2019). Hereafter we refer to MPI-ESM1.2 as MPI-ESM, and ignore version numberings for other components, unless otherwise stated. In terms of differences to the earlier versions, we only focus on those between the atmospheric part of the latest version, ECHAM6, and the atmospheric part used in SOCOLv3, ECHAM5.4, as changes between versions contribute the differences between the chemical response of SOCOLv4 and all sub-versions of SOCOLv3.

2.1.1 Ocean model MPIOM

The oceanic part MPIOM is formulated on an Arakawa-C grid in the horizontal and on z-levels in the vertical direction and solves the primitive equations with the hydrostatic and Boussinesq approximations (Jungclaus et al., 2006; 2013). Subgrid-scale parameterizations include lateral mixing on isopycnals and tracer transports by unresolved eddies. Vertical mixing is represented as a combination of the Richardson number-dependent scheme and the wind-driven turbulent mixing in the mixed layer (for details see Jungclaus et al., 2013). The horizontal grid is consistent with the MPI-ESM LR configuration, which



implies a bipolar grid GR1.5 featuring one grid pole under Greenland and one under Antarctica. The resolutions are then
145 regionally enhanced in the deep water formation regions and the overflows across the Greenland-Scotland ridge so that the
grid varies between 22 and 350 km. In the vertical, 40 levels are unevenly placed in the water column, with the first 20 levels
distributed over the top 700 m. The bottom topography is represented by a partial-step formulation (Wolff et al., 1997).
The sea ice model combines the codes of MPIOM and ECHAM. In ECHAM, a simplified thermodynamic sea ice model is
incorporated to provide at each atmospheric time step a physically consistent surface temperature in ice-covered regions. This
150 part also contains a melt-pond scheme, which divides the surface of the sea ice into snow, bare ice, and melt pond with
individual albedos (Pedersen et al., 2009). The atmospheric part of the sea ice model then integrates all surface fluxes into ice
and provides this information to the oceanic part of the code, which uses it to calculate the sea ice surface energy balance and
related changes in ice thickness. Parametrizations of changes in ice concentration and ice thickness can be adjusted by two
tuning parameters, which have been used to tune the preindustrial Arctic sea ice volume of MPI-ESM to an annual average of
155 roughly 20–25 thousands of km³ (Mauritsen et al., 2019). Sea ice dynamics is calculated following a viscoplastic approach of
Hibler (1979).

2.1.2 Marine biogeochemistry model HAMMOCC

Ocean biogeochemistry in MPI-ESM is represented by the Hamburg Ocean Carbon Cycle (HAMOCC) model (Ilyina et al.,
2013; Paulsen et al., 2017). It simulates the oceanic cycles of carbon and other biogeochemical elements such as nutrients
160 (phosphate, nitrate, and iron), oxygen, silicate, phytoplankton, zooplankton, and detritus. HAMOCC includes biogeochemical
processes in the water column, the sediment, and at the air-sea interface. Biogeochemical tracers in the water column are fully
advected, mixed, and diffused by the flow field of MPIOM. In total, the model has 17 state variables calculated prognostically
in the water column and 12 state variables in the sediment. Nitrogen-fixing cyanobacteria was added to the model as an
additional prognostic phytoplankton class by Paulsen et al. (2017).

165 2.1.3 Land surface model JSBACH

The Jena Scheme for Biosphere-Atmosphere Coupling in Hamburg (JSBACH) is the land component of MPI-ESM1.2. It
provides the lower boundary conditions for the atmosphere over land and describes the dynamics of the land biogeochemistry
in interaction with global climate. JSBACH treats processes like soil hydrology (five-layer scheme of Hagemann & Stacke,
2015), soil and litter decomposition, land use change (tiling approach with 12 plant functional types and two types of bare
170 surface), fires, and a nitrogen cycle (Goll et al., 2017). Note that soil and marine chemical schemes are not yet combined with
the atmospheric chemistry scheme in the current model version.

2.1.4 ECHAM6

ECHAM6 is an atmospheric general circulation model (GCM) that describes the large-scale circulation and its coupling to
diabatic processes, both of which are ultimately driven by radiative forcing. It consists of a dry spectral-transform dynamical



175 core, a transport model, and a suite of physical parameterizations for the representation of diabatic processes. The prognostic variables are temperature, vorticity, divergence, logarithm of surface pressure, humidity as well as cloud ice and water. Tracer transport and diabatic processes (also referred to as “model physics”) are calculated on a Gaussian transform grid. The adiabatic core of ECHAM6 consists of a mixed finite-difference/spectral discretization of the primitive equation that is identical to that employed in ECHAM5 (Stevens et al., 2013). All major changes relative to ECHAM5 are therefore related to the model

180 physics. These changes include: an improved representation of radiative transfer in the shortwave (or solar) part of the spectrum; a completely new description of tropospheric aerosol; an improved representation of surface albedo, including the treatment of melt ponds on sea ice (see section 2.1.1); and an improved representation of the middle atmosphere through the gravity wave forcing. In addition, minor changes have been made in the representation of convective processes. Several coding errors in model physics were also corrected in the latest version (see details in Mauritsen et al., 2019).

185 Transport of species is performed with the flux-form semi-Lagrangian scheme of Lin and Rood (1996). Though this scheme is mass-conservative by design, its application on the sigma-pressure coordinate system can cause a violation of the mass-conservation especially in case of large spatial gradients (Jöckel et al., 2001; Stenke et al., 2013). Turbulent mixing adopts an eddy diffusivity and viscosity approach following Brinkop and Roeckner (1995). Moist convection is parameterized according to Tiedtke (1989), with extensions by Nordeng (1994) and Möbis and Stevens (2012). Stratiform clouds are computed

190 diagnostically based on a relative humidity threshold (Sundqvist et al., 1989). Gravity waves are generated from a subgrid orography scheme (Lott, 1999) and as Doppler waves following Hines (1997a, b), and they are treated according to the formulation of Palmer et al. (1986) and Miller et al. (1989). Compared to ECHAM5, the tropospheric gravity wave sources are now prescribed as a function of latitude (Schmidt et al., 2013). Both long-wave and short-wave radiative transfer calculations are now described by the k-correlated method of RRTM-G (Iacono et al., 2008). Additional extra-heating

195 parameterization in the Hartley, Huggins, and Schumann-Runge bands and the Lyman- α line is now applied for better representation of the solar cycle in the mesosphere and stratosphere (Sukhodolov et al., 2014). The optical properties for radiation are updated every 2 hours. In contrast to the base model version, which applies climatological fields for this purpose, the radiation calculation of SOCOLv4 uses the prognostic tracer concentrations of sulfate aerosol and all radiatively active species (e.g., ozone and methane) except CO₂. Cloud scattering is parameterized according to Mie theory using maximum-

200 random cloud overlap and an inhomogeneity parameter to account for three-dimensional effects. Surface albedo is parameterized according to Brovkin et al. (2013). The radiative properties of tropospheric aerosols are now represented by the MACv2-SP parameterization (Stevens et al., 2017; Fiedler et al., 2017). MACv2-SP is formulated in terms of nine spatial plumes associated with different major anthropogenic source regions. It prescribes the aerosol optical depth, asymmetry factor and single scattering albedo as a function of geographical position, height above ground level, time, and wavelength. The

205 evolution and distribution of anthropogenic aerosol is approximated with mathematical functions, while the natural aerosol is prescribed climatologically based on Kinne et al. (2013). MACv2-SP also prescribes aerosol-cloud interactions in the form of a Twomey effect (Twomey, 1977), thus allowing the increase in the cloud droplet number concentration and associated reduction in droplet size under constant liquid water path.



2.2 Atmospheric chemistry model MEZON

210 The atmospheric chemistry part of the new model is a modified version of the chemistry-transport model MEZON (Model for
Evaluation of oZONe trends; Rozanov et al., 2001; Egorova et al., 2003; Schraner et al., 2008). The last base state of this code
is described in detail by Stenke et al. (2013). It underwent major upgrades for participation in the Chemistry Climate Model
Initiative Phase 1 (CCMI) presented in Revell et al. (2015). Further upgrades were made in Revell et al. (2018) and Feinberg
et al. (2019). Below we summarize the current state of MEZON, while the history of main updates that are now included in
215 SOCOLv4 is schematically illustrated in Fig. 2. In this paper, the direct comparison with SOCOLv4 is made only for the CCMI
version (Revell et al., 2015) in sections 3.2.2 and 3.2.4, since it was the most widely used one in publications.




Figure 2. History of main updates through different versions of SOCOL, which contributed to the version described in this paper.

220 Note that there were also other minor changes, adjustments, and error corrections.

MEZON and ECHAM6 are interactively coupled by the 3-dimensional fields of temperature and wind, and by the radiative forcing induced by water vapor, ozone, methane, nitrous oxide, and chlorofluorocarbons (CFCs). The chemistry scheme is called every 2 h. The chemical solver is based on the implicit iterative Newton–Raphson scheme (Ozolin, 1992; Stott and
225 Harwood, 1993). The model includes 99 chemical species of the oxygen, hydrogen, nitrogen, carbon, chlorine, bromine, and sulfur groups, which are determined by 216 gas-phase reactions, 72 photolysis reactions, and 16 heterogeneous reactions in/on aqueous sulfuric acid aerosols as well as three types of polar stratospheric clouds (PSCs): supercooled ternary solution (STS) droplets, water ice, and nitric acid trihydrate (NAT). Chemical reaction rate coefficients and absorption cross sections of all reactions follow the recommendations from the NASA JPL data evaluation no. 18 (Burkholder et al., 2015). Photolysis rates
230 are calculated at every chemical time step using a look-up-table approach (Rozanov et al., 1999), including effects of the solar irradiance variability. Based on the stand-alone photolysis codes intercomparison study (Sukhodolov et al., 2016) and



additional sensitivity tests, the photodissociation rates of molecular oxygen and ozone ($O(^1D)$ path) have been supplemented by the correction factors of 1.2 for oxygen in the 10-1 hPa region and 1.5 for ozone below 100 hPa.

235 Dry deposition of species is based on the surface resistance approach for the estimation of dry deposition velocities proposed by Wesely (1989). This scheme takes into account actual meteorological conditions, different surface types and trace gas properties like solubility and reactivity. Further details are given by Kerkweg et al. (2006). The wet deposition scheme is based on the SCAV scheme proposed by Tost et al. (2006). Wet deposition velocities are calculated using cloud and precipitation variables from ECHAM6, including cloud cover, liquid and ice water contents, precipitation fluxes and the convective upward mass flux. Scavenging coefficients of gaseous species depend on their Henry's law constants. Upon evaporation of clouds and
240 rainwater, scavenged species are transferred back into the gas phase. An additional sink for HNO_3 , in the form of a constant removal rate of $1 \times 10^{-6} \text{ s}^{-1}$, was introduced in the upper troposphere (above 300 hPa), compensating the missing uptake of HNO_3 by ice particles (e.g., Voigt et al., 2006).

The parameterization of heterogeneous chemistry in the stratosphere is based on Carslaw et al. (1995). It coincides with HNO_3 uptake by aqueous sulfuric acid aerosols resulting in the formation of STS. The parameterization of the liquid-phase reactive
245 uptake coefficients follows Hanson and Ravishankara (1994) and Hansen et al. (1996). The PSC scheme for water ice uses a prescribed particle number density of 0.01 cm^{-3} and assumes that the cloud particles are in thermodynamic equilibrium with their gaseous environment. NAT is formed if the partial pressure of HNO_3 exceeds its saturation pressure, assuming a mean particle radius of $5 \text{ }\mu\text{m}$ for NAT. The particle number densities are limited by an upper boundary of $5 \times 10^{-4} \text{ cm}^{-3}$ to account for the fact that observed NAT clouds are often strongly supersaturated. The sedimentation of NAT and water ice is based on
250 the Stokes theory as described in Pruppacher and Klett (1997). Water ice and NAT are not explicitly transported but are evaporated back to water vapor and gaseous HNO_3 after each chemical time step, transported in the vapor phase, and then depending on the saturation conditions regenerated in the next time step with the thermodynamic approximation described above. In the troposphere, the heterogeneous hydrolysis of N_2O_5 on tropospheric aerosols contributes to the sink of odd nitrogen. Note that since the MACv2-SP parameterization that is used for radiation does not provide the aerosol mass, for the
255 tropospheric chemistry we use the tropospheric aerosol climatology that considers the aerosol properties of 11 Global Aerosol Data Set types (GADS, Koepke et al., 1997) as in the previous model version. The reaction probabilities for the different aerosol types are calculated following the parametrization by Evans and Jacob (2005).

Parameterizations of the N, NO, and OH production by galactic cosmic rays, solar protons, and energetic electrons with energies of $<300 \text{ keV}$ are introduced as 0.55, 0.7, and up to 2 molecules of N, NO, and OH per ion pair respectively (Matthes
260 et al., 2017). The contribution of thermospheric NO from downward intrusions is parameterized as a flux-form upper boundary condition (Funke et al., 2016). The lightning source of NO_x is parameterized based on the cloud top approach by Price and Rind (1992). The tropospheric budget of CO is supplemented by the Mainz Isoprene Mechanism (MIM-1, Pöschl et al., 2000) that describes the degradation of isoprene, formaldehyde and acetic acid. Contributions of all other non-methane volatile organic compounds (NMVOCs) to CO is accounted for via the addition of a certain fraction of NMVOC emissions to CO. For



265 anthropogenic, biomass burning, and biogenic NMVOC emissions the conversion factors to CO are 1.0, 0.31 and 0.83, respectively (Ehhalt et al., 2001).

Several newly discovered hODS (CFC-112, CFC-112a, CFC-113a, CFC-114a and HCFC-133a) have been added to the model chemistry scheme together with some additional chlorine containing very short-lived substances (VSLs: CHCl₃, CH₂Cl₂, C₂Cl₄, C₂HCl₃, C₂H₄Cl₂) that are not controlled by the MPA. The bromine containing VSLs forcing is now also transient
270 compared to the previous model versions and includes CH₃Br, CHBr₃ and CH₂Br₂ species. The sulfur family is represented by eight gas-phase species: OCS, CS₂, H₂S, DMS, methanesulfonic acid (MSA), SO₂, SO₃, and H₂SO₄ (Sheng et al., 2015).

2.3 Sulfate aerosol microphysics model AER

The sulfate aerosol microphysical scheme AER is based on the two-dimensional sulfate aerosol model of Weisenstein et al. (1997). It was upgraded and combined with SOCOLv3 by Sheng et al. (2015) and later further improved by Feinberg et al.
275 (2019). Sulfate aerosol particles are resolved in 40 size bins, ranging in dry radius from 0.39 nm to 3.2 μm, corresponding to a range of 2.8 - 1.6×10¹² molecules of H₂SO₄ per particle (assuming an H₂SO₄ density of 1.8 g cm⁻³). H₂SO₄ molecule number doubles between bins, while the corresponding wet sulfate aerosol radii can be much larger depending on local conditions. H₂SO₄ weight percent is calculated online based on actual temperature and relative humidity. The AER scheme includes submodules for the nucleation (Vehkamäki et al., 2002), composition (Tabazadeh et al., 1997), growth, evaporation (Ayers et al., 1980; Kulmala and Laaksonen, 1990), coagulation (Fuchs, 1964; Jacobson and Seinfeld, 2004), and sedimentation of
280 sulfate aerosol (Kasten, 1968; Walcek, 2000). In addition to gas phase H₂SO₄ production, the model calculates aqueous oxidation of S(IV) by ozone (O₃) and hydrogen peroxide (H₂O₂) (Jacob, 1986). The spatial distribution of cloud pH in the aqueous phase chemistry routine is approximated based on Tost et al. (2007). The aqueous production flux of S(VI) is added directly to the scavenged aerosol flux in cloud water. Dry and wet deposition of sulfate aerosol are calculated by the similar
285 schemes as for gas-phase species (Tost et al., 2006; Kerkweg et al., 2006). Dry and wet deposition as well as gravitational velocities of sulfate aerosol are calculated through radius-dependent parametrizations. During cloud evaporation, evaporating scavenged sulfate aerosol mass is transferred to the largest aerosol size bin. Transport of aerosols in each bin is also performed in the same way as transport of gases with a timestep of 15 min. The microphysical module is called every 2 h with 20 sub-time-steps yielding an aerosol microphysical time step of 6 min.

290 The influence of the aerosol on radiation fluxes at all wavelengths (14-band shortwave and 16-band longwave) is taken into account. Extinction coefficients, single-scattering albedo, and asymmetry factors required by the radiation codes are treated following a lookup table approach with precalculated aerosol physical properties using Mie theory for actual H₂SO₄ weight percent and temperature using refraction indices from Biermann et al. (2000). The aerosol surface area density and composition are used to calculate heterogeneous reaction rates in a chemical module.



295 2.4 Model set-up and boundary conditions

As described above, SOCOLv4 is based on the latest version of MPI-ESM (v1.2.01p6) that was also used for the Climate Model Intercomparison Project Phase 6 (CMIP6) runs. To test the new coupled model, we initialized all its parts from the MPI-ESM restart files (snapshots of the model's state at a specified time) from the end of the year 1949 and then continued the calculation until the end of 2018. The initialization of chemistry was based on SOCOLv3 runs from Revell et al. (2016).

300 In 1980 the reference run was split into an ensemble of three runs by imposing a temporary (1-month long) small perturbation in atmospheric CO₂ concentrations. For the analysis performed here, we skip the first 30 years of simulations and focus on the 1980-2018 period, which is well covered by observations for comparison with.

Model boundary conditions mostly follow the recommendations of CMIP6 provided by the input4MIPs database (<https://esgf-node.llnl.gov/search/input4mips/>). All forcings are historical before 2015 and switched to the SSP2-4.5 scenario for the last
305 four years till 2018. These includes concentrations of greenhouse gases (Meinhausen et al., 2016), surface anthropogenic and biomass-burning emissions of NO_x (as well as aircraft emissions), CO, SO₂, and NMVOCs (Hoesly et al., 2018), ionization rates by galactic cosmic rays, solar protons, and energetic electrons, solar spectral irradiance variations, and the influx of thermospheric of NO (Matthes et al., 2017). Biogenic emissions of all NMVOCs use a climatology for the year 2000 based on
310 halogenated source gases (ODSs and VSLs) we used the WMO Ozone Assessment 2018 baseline mixing ratio scenario, which is a combined atmospheric observation record up to the year 2017 (Engel et al., 2018). For some ODSs we also used the input4MIPs database. Agricultural land-use changes are based on the Land Use Harmonization project data (LUHv2h, Hurtt et al., 2011).

Continuous degassing emissions of SO₂ are prescribed according to volcano locations (Andres and Kasgnoc, 1998; Dentener et al., 2006). To represent eruptive emissions, we applied a satellite-derived dataset from Carn et al. (2016). Since the emission profiles are unknown in this database, they are emitted into the upper third of the total plume height. Air-sea DMS fluxes are calculated online through a wind-driven parametrization (Nightingale et al., 2000) and climatological gridded sea surface DMS concentrations (Lana et al., 2011). 1 Tg S yr⁻¹ of CS₂ is emitted between the latitudes of 52°S and 52°N based on Weisenstein et al. (1997). The surface mixing ratios of H₂S and OCS are prescribed as 30 pptv (Weisenstein et al., 1997) and 510 pptv
320 (Timmreck et al., 2018), respectively.

Since the model vertical resolution is insufficient for a reasonable self-generation of the QBO, we nudge it to the observed equatorial wind profiles. The nudging is applied between 20°N and 20°S from 90 hPa up to 3 hPa. Within the QBO core domain (10°N–10°S, 50–8 hPa) the relaxation time is uniformly set to 7 days; outside this region the damping depends on latitude and altitude (Giorgetta, 1996).

325 Tropospheric aerosols are represented by two separate datasets for radiation and chemistry. For radiation, to be consistent with the base model and CMIP6 recommendations we used the MACv2-SP approach (Stevens et al., 2017), which however provides



only optical parameters of aerosols. Surface area densities required for the N_2O_5 hydrolysis, are based on the tropospheric aerosol climatology of 11 aerosol types from the Global Aerosol Data Set (GADS, Koepke et al., 1997), as in SOCOLv3.

3 SOCOLv4 evaluation

330 3.1 Atmospheric dynamics

3.1.1 Reference data for evaluation

We validate simulated climate variables from SOCOLv4 against observations. For this purpose, we use reanalysis data because they incorporate various observations combined in one data set using comprehensive techniques that allow gaps in space and time to be estimated. It should be noted that the combination of different data can lead to some errors in the reanalysis data, both in trends and variability, that have a purely methodological nature, like those due to changes in the observation systems or aging of instruments as well as the details of underlying models used for assimilation. Intercomparison of different reanalysis products by Long et al. (2017) showed that reanalyses agree in the lower/middle stratosphere but deviate more from each other in the upper stratosphere and lower mesosphere because of the decreased availability of direct observations and thus the increased dependence on the assimilating model details, like the model top and vertical resolution. This should be considered when making conclusions about deviations between model results and reanalysis. To validate the thermo-dynamical structure of the modelled atmosphere, we use the recent ERA5.1 reanalysis data set (Simmons et al., 2020, Hersbach et al. 2020) from ECMWF (European Center for Medium Range Weather Forecast) and MERRA-2 (Modern Era Retrospective Analysis for Research and Applications Version 2, Gelaro et al. (2017)). For surface air temperature we additionally use the BEST data (the Berkeley Earth Surface Temperatures, Cowtan et al. (2015)), which is a merged land-ocean data set.

345 3.1.2 Temperature and winds


To assess the ability of SOCOLv4 to reproduce the climate and its response to historical forcing, we show variability in annual mean surface air temperature and its anomaly (Fig. 3a, c), as well as the El Niño Southern Oscillation (ENSO) Niño3.4 index (Fig. 3b) and the Arctic sea ice extension anomaly (Fig. 3d) for the historical period 1980 to 2018. Anomalies were calculated as a deviation from the mean over the 1980-2018 period. Figure 3 (a, c) shows the global mean 2-meter temperature and its anomaly simulated with the SOCOLv4 and various reanalysis data. The model shows a warming trend similar to observations, whereas the ensemble mean curve shows less inter-annual variability, as expected. As can be seen from the figure, the reanalysis data slightly disagree in terms of the absolute level and the model data are within this spread. Model surface air temperature is somewhat warmer than the ERA5.1 and MERRA-2 data, but colder than the BEST data.

ENSO is known as the largest mode of sea surface temperature anomalies which can influence atmospheric circulation (e.g., Bulgin et al., 2020). Figure 3b shows variability of sea surface temperature in the Niño3.4 region (5N-5S, 120W-170W). The ERA5.1 reanalysis data (red curve) fluctuate in the Niño3.4 region in the range of $\pm 3\text{K}$. Positive fluctuations are associated



with El Niño events and negative with La Niña events. Observational data show several El Niño events with positive deviations stronger than +1.5 K in temperature on the time scale 1980-2018. with the strongest being those in 1982/1983, 1997/1998, and 2015/2016. The latest El Niño event gives the largest observed positive sea surface temperature anomalies. The SOCOLv4 simulations also show variability in the Niño3.4 region which is similar to observations but differs in phase among ensemble members. If we consider the simulated warm events, we can see that the free running model cannot reproduce the timing of the events as expected due to the chaotic nature of atmosphere-ocean coupling processes that give rise to ENSO. However, it can reproduce their amplitude, which is between -2 K and +3 K, as well as their frequency of about one event per 3-4 years. Spatial pattern of ENSO in the model is reasonable, but has some biases related to the asymmetry between El Niño and La Niña events (for details see Tian et al. 2019). This drawback could be minimized by increasing coupling frequency between ocean and atmosphere from daily to hourly. However, this change also increases the computational demands to levels, which are less suitable for long-term climate calculations. Therefore, the LR version of MPI-ESM which we used for SOCOLv4 was tuned with a daily coupling (Mauritsen et al., 2019).

In general, SOCOLv4 accurately reproduces Arctic sea ice extent annual anomalies in line with observations, showing a similar decline with time because of global warming. According to Ding et al. (2019), the spread in sea-ice extent between the ensemble members can be attributed to internal variability driven by fluctuations in Arctic pressure expressed as high pressure enhancing the sea ice loss and low pressure restraining it. Based on the analysis of Fig. 3d, we conclude that the model response to historical radiative forcing is sufficient and close to the observations, which proves suitability of the model for studies of past and future climate changes.



375

Figure 3. Global and annual mean 2-meter temperature (a) and its anomalies relative to 1980 (c) in comparison with reanalysis data, ENSO Niño3.4 index (b), and Arctic sea ice extent annual anomaly (d). Black thick line represents simulated mean of 3-member SOCOLv4 ensemble, thin black lines are individual ensemble members, red line is ERA5.1 reanalysis, green line is MERRA-2 reanalysis, and blue line is BEST reanalysis.

380


The left panel of Fig. 4 shows the geographical distribution of the 2-meter air temperature climatology for 1980-2018 as simulated with SOCOLv4 and derived from the ERA5.1 reanalysis, as well as their difference. ERA5.1 data have been interpolated to the model grid. SOCOLv4 properly reproduces the observed geographical pattern of the surface air temperature with cold high-latitude areas in both hemispheres and warm tropics and subtropics. However, there are some regional discrepancies between the model and the reanalysis data. The model surface air temperature is generally slightly (~0.2 K) warmer than in ERA5.1, which is also seen from the global means (Fig. 3a), but there are also several spots with cold biases. In the tropical area, the discrepancies are within ± 1 K, except in the eastern part of South America over the Brazilian Highlands, where negative bias reaches -5 K, and over the Tibetan Plateau and the Indochinese Peninsula, with up to -3 K bias. In the northern middle latitudes, we obtain negative discrepancies of about -5 K over the Rocky Mountains and the North Atlantic Ocean. The most pronounced biases of ± 6 K are located in the Southern Hemisphere over Antarctica. Most of these biases in high-altitude regions are a common feature of all coupled models, which is usually attributed to the simplified model's

385

390



topography and uncertainties in the observational data (e.g., Guo et al., 2020). The right panel of Fig. 4 shows the 2-m surface air temperature trends over 1980-2018 over the northern hemisphere simulated by SOCOLv4 and calculated from ERA5.1 reanalysis data. The linear trend and its significance were calculated using a robust nonparametric Sen–Mann–Kenndall test applying 95% confidence interval. Simulated and observed trends over the Northern Hemisphere show a good spatial agreement including the location of the warming maximum north of Siberia. The maximum simulated trend amounts to 1.69 ± 0.16 K, which is only about 0.25 K smaller than the maximum trend of 1.93 K derived from the reanalysis. Therefore, we can conclude that the climate change forcing used in SOCOLv4 and the model response to this forcing are adequately represented.



400 Figure 4. Left panels: Global distribution of 2-m surface air temperature (K, contours are -5, -3, -1, 0, 1, 3, 5, 7, 9) averaged over the period 1980-2018 simulated by SOCOLv4 (a, ensemble mean), ERA5.1 reanalysis data (b), and their difference (c, SOCOL4-ERA5.1). Right panels: 2-m surface air temperature trend over the northern hemisphere (K/10years, contours are -0.5, -0.2, 0, 0.2, 0.5, 0.7, 1.0, 1.5) simulated (d) and observed (e) over the Northern Hemisphere. White spots show where the trend is not statistically significant at 95% level.

405




Figure 5. Zonal mean zonal wind climatology (m/s) for DJF (left panel) and JJA (right panel) simulated with SOCOLv4 (a, b), mean of ERA5.1 and MERRA-2 reanalysis (c, d) and differences between simulated and reanalysis data (e, f). Contour intervals for the difference are -30, -20, -10, -5, -2, 2, 5, 10, 20, 30 m/s. White places denote regions where the difference between model and reanalysis data is not statistically significant at 95% confidence level calculated with the Student's t-test.

410



Zonal mean climatology of zonal winds and temperatures for boreal winter and summer as simulated by SOCOLv4 are presented in comparison with observations (mean data from reanalysis ERA5.1 and MERRA-2) in Fig. 5 and 6, respectively. Even though the model has an upper limit at 0.01 hPa, we analyze the results only up to a height of 0.1 hPa, since above
415 ERA5.1 data are not reliable and MERRA-2 data are not available. In general, the model reproduces the observations well, but there are deviations which are, however, typical for most coupled models (Bock et al., 2020, Gettelman et al., 2019, Matthes et al., 2020). The tropospheric jets, their position and strength are reasonably reproduced by the model with biases of ± 2 -5 m/s. There is a tendency of the westerlies in the mid-latitudes to be insufficiently poleward, which is more pronounced in the Southern Hemisphere and represented as negative and positive deviations around 60 and 40°S, respectively. The differences
420 between the model and observations increase with altitude. In the stratosphere, the simulated summer-time easterly jets are of the similar strength as in observations but have some deviations in their shapes. The deviations between about 90 and 3 hPa in the (sub)tropics are related to the hemispherically symmetric QBO nudging approach, which is applied between 20°N to 20°S. The winter-time polar night jets show larger deviations in both hemispheres. In the Northern Hemisphere, the modelled polar vortex is significantly weaker than in observations and is extended equatorward in the lower mesosphere. The southern
425 hemispheric vortex is of the similar strength as in reanalysis mean data in the stratosphere and the main model deviations are related to its positioning. The core of the southern polar jet is shifted to 60° and located between 10 hPa and 1 hPa, while in the reanalysis data the vortex core (80 m/s isoline) extends higher and shifts towards 40°S. Therefore, the modelled southern polar vortex area is more isolated in the middle stratosphere. Oppositely, in the lower mesosphere it is extended equatorwards, similar to the northern one. The latter feature might suggest that there is some overestimation in the gravity wave forcing that
430 is dominant in this region (Plumb, 2002; Butchart, 2014).

The zonal mean temperature distribution is generally well reproduced by the model (Fig. 6), but there are also strong biases in the middle atmosphere with respect to the observations that are consistent with the biases in zonal winds. Namely, the winter-time high-latitude middle and upper stratosphere and the low mesosphere are warmer by 2-10 K suggesting weaker vortices and stronger meridional transport and mixing. In the rest of the stratosphere the difference is negative (about -2 K). This is
435 consistent with a too fast Brewer-Dobson circulation (BDC), which would cause increased adiabatic cooling in the tropics and in the summer-time high-latitudes. In the lowermost extra-tropical stratosphere, there are up to -5 K temperature deviations in both hemispheres, which leads to some downward displacement of the extratropical tropopause. The modelled tropospheric temperatures agree very well with observations and only show a warm bias over Antarctica.

It has to be noted that all of the presented model deviations in thermodynamics are long-term issues of the ECHAM-family
440 models and global models in general (Stevens et al., 2013; Mauritsen et al., 2019). In Figure A1, we present the same differences as in Fig. 5 and 6 but for the pure MPI-ESM CMIP6 run. As can be seen from there, SOCOLv4 and MPI-ESM share the same deviation patterns indicating weaker and warmer polar vortices in the stratosphere. Some differences between SOCOLv4 and MPI-ESM results are mostly related to the stratospheric ozone distribution at mid- and high-latitudes, as well as to the QBO nudging in SOCOLv4 in the tropics. Many of these problematic features were shown to be greatly improved



445 using a version of ECHAM with more detailed (95-level) vertical resolution (Schmidt et al., 2013; Stevens et al., 2013; Mauritsen et al., 2019; Matthes et al., 2020), especially the strength of the northern polar vortex and the extratropical lower-stratospheric cold biases. The increase of the vertical resolution implies, however, a strong increase in computational time, because of the column physics design. It can be partly compensated by increasing the amount of processing units, but there is also a limit in model's scalability defined by horizontal resolution.

450




Figure 6. Zonal mean temperature climatology for DJF (left panel) and JJA (right panel) simulated with SOCOLv4 (a, b), mean of ERA5.1 and MERRA-2 reanalysis (c, d) and differences between simulated and reanalysis data (e, f). Contour intervals for the differences are -30, -20, -10, -5, -2, 0, 2, 5, 10, 20, 30, 40 K. White places denote regions where the difference between model and reanalysis data is not statistically significant at 95% confidence level.



3.2 Atmospheric chemistry and transport

3.2.1 Reference data for validation

All data sets that were used for the chemistry validation are listed in Table 1. For the comparison of HCl, H₂O, and N₂O we used the Global Ozone Chemistry And Related trace gas Data records for the Stratosphere database (GOZCARDS, Froidevaux et al., 2015), which is a composite based on the data from 8 satellite instruments. The ENVISAT satellite Michelson Interferometer for Passive Atmospheric Sounding (MIPAS, Funke et al., 2014) instrument data was used for NO_y, while the SCISAT Atmospheric Chemistry Experiment Fourier Transform Spectrometer (ACE-FTS) data climatology is used for NO_y and CH₄. Besides GOZCARDS, the ozone mixing ratio is also compared to the BAYesian Integrated and Consolidated composite ozone (BASIC, Alsing and Ball, 2019), and the CMIP6 Ozone forcing dataset (Checa-Garcia, 2018). The latter is a compilation of CESM1-WACCM and CMAM modelling results and is used as a standard forcing for MPI-ESM. Ozone total column is validated against Multi Sensor Reanalysis version 2 (MSRv2, Van der A et al., 2015) and SBUVv8.6 (McPeters et al., 2013) composites. Stratospheric aerosol sulfur mass is compared to MIPAS (Günther et al., 2018) and HIRS (Baran and Foot, 1994) observations and a composite recommended by CMIP6 (GloSSACv1.1, Thomason et al., 2018).

It has to be noted that observations themselves are not perfect, especially for the shorter-lived species with pronounced diurnal cycles, low concentrations, and specific regions like high-latitudes and the upper troposphere/lower stratosphere (UTLS). Different satellite instruments vary in terms of measurement method, geographical coverage, spatial and temporal sampling and resolution, time period, and retrieval algorithm. All of this has also to be dealt with while constructing the observational composites, which imposes further potential uncertainties related to the homogenization methods applied. The SPARC Data Initiative (SPARC, 2017) overviewed the existing trace gases and aerosol measurements and provided a set of useful recommendations concerning the related uncertainties that we used in the further sections for our validation.

Table 1. Data used in this study for chemistry validation.


Data	Period	Source
O ₃ mixing ratio	1985-2012	GOZCARDS composite (Froidevaux et al., 2015)
	1985-2012	BASICv3 composite (Alsing and Ball, 2019)
	1985-2012	CMIP6 composite (Checa-Garcia, 2018)
CH ₄ , NO, NO ₂ , N ₂ O ₅ , HNO ₃ , and ClONO ₂ mixing ratio	2004-2012	ACE-FTSv3.5 (Bernath et al., 2005)
H ₂ O and HCl mixing ratio	1991-2012	GOZCARDS
N ₂ O mixing ratio	2004-2012	GOZCARDS
NO _y mixing ratio	2004-2012	ACE-FTSv3.5



	2002-2012	MIPAS (Funke et al., 2014)
H ₂ O mixing ratio	1991-2012	GOZCARDS
Liquid H ₂ SO ₄ mass	1991-1993	HIRS (Baran and Foot, 1994)
	1980-2016	CMIP6 stratospheric aerosol composite (GloSSACv1.1, Thomason et al., 2018)
	2002-2012	MIPAS (Günther et al., 2018)
O ₃ total column	1980-2018	MSRv2 composite (Van der A et al., 2015)
	1980-2018	SBUVv8.6 (McPeters et al., 2013)

3.2.2 Trace gases climatology

In this section we discuss the SOCOLv4 performance in reproducing observed distributions of various trace species. Figures 7, 10, and 11 show the data from the model and observations and the relative differences between them. Figures 8 and 12 show mean seasonal cycles of several species that are also compared to observations.




485 **Figure 7.** Annually averaged zonal mean distributions of CH₄, H₂O, HCl, and N₂O (from top to bottom). First column: SOCOLv4 data. Second column: ACE-FTS and GOZCARDS observational data. Third column: relative difference between the model and observations in %. Observational missing data mask is applied to the modelling results. Note that different species have different averaging periods. White areas are either the missing data or regions where the difference between model and observations is not statistically significant at 95% confidence level.



CH₄. Methane (CH₄) is an important source of water vapour (H₂O) in the stratosphere. On average two H₂O molecules are
490 produced per CH₄ molecule oxidized. Methane mixing ratios drop very fast with height due to increasing availability of
hydroxyl radical (OH), the excited atomic oxygen (O(¹D)), and atomic chlorine (Cl). In the lower mesosphere, photolysis by
Lyman- α radiation also becomes an important sink for CH₄. Methane is also an important atmospheric tracer for tracking the
stratospheric circulation because of its long lifetime. CH₄ mixing ratio isolines in Fig. 7 are lifted upward in the low latitudes
by the ascending branch of the BDC and are pushed downward in the middle and high latitudes by the descending branch.
495 Isolines in the middle latitudes are flattened as a result of rapid mixing by breaking planetary waves. The balance between
horizontal mixing and the diabatic circulation produces the observed tracer slope and deviations in this distribution can hint to
potential problems in transport processes (e.g., Strahan, 2015).

SOCOLv4 nicely reproduces the methane distribution in the lower stratosphere with only about $\pm 5\%$ deviations from
observations. However, above ~ 10 hPa the model deviation gradually increases with height to up to 50% overestimation in the
500 lower mesosphere. The highest overestimation close to the model top in the mesosphere is likely related to the treatment of the
Lyman- α line photolysis, while the rest is due to the problems in dynamics. Thus, a too fast vertical transport in the tropics
(Fig. 7) would mean that there is less time for CH₄ loss on the way upwards and therefore more CH₄ arrives in the mesosphere,
which can be partly enhanced by underestimation of CH₄ photolysis. Dynamical lifetime got smaller with respect to chemical
lifetime and the tracer became more vertically mixed. The increased concentration in the upper stratosphere is then also
505 distributed polewards and downwards in the middle and high latitudes. Difference in skewness of the isolines suggests an
overestimation of mid-latitude mixing in the upper stratosphere/lower mesosphere, which is consistent with analysis of winds
showing that the vortices are weaker at those altitudes (Fig. 5). Middle and lower stratospheric midlatitudes, however, show
some hemispheric asymmetry with a negative bias of 5% in the SH and a positive bias of 5-10% in the NH. Figure 8 shows
the CH₄ seasonal cycle at 20 hPa and suggests that this anomaly is likely related to the deficiencies in horizontal mixing.
510 Namely, while the model reproduces very well the tropical values (1.4 ppmv isoline), the midlatitude values (1 ppmv isoline)
are excessively shifted poleward in the NH and slightly equatorward in the SH. This is also generally consistent with Fig. 5,
which shows at these altitudes a weaker NH polar vortex and the SH polar vortex that is more confined to the pole.



515 **Figure 8.** CH₄, HCl, and NO_y zonal mean seasonal cycles at 20, 70, and 6 hPa, respectively. First column: SOCOLv4 data. Second column: ACE-FTS, GOZCARDS, and MIPAS observational data. Two consecutive cycles are shown.

520 **H₂O.** Water vapour (H₂O) provides a source for hydroxyl radicals (HO_x) in the stratosphere. Hydroxyl catalytic cycles dominate the ozone destruction above the stratopause and are also very important in the lower and middle stratosphere. The two main sources of H₂O in the stratosphere are methane oxidation and the transport from the H₂O-rich troposphere. Transport from the troposphere is important for the lower stratosphere and is highly dependent on the cold point temperature at the tropical tropopause and vertical transport in the tropics. As seen from Fig. 7, SOCOLv4 agrees very well with observations around the stratopause. In the lower stratosphere, the model is 5-10% too moist, which is due to some overestimation at the entry point. In the mesosphere, the model underestimates H₂O by up to 30% (or 1 ppmv) around the model top. This partly results from the underestimated destruction of methane (about 0.2 ppmv) but mostly comes from the overestimated water vapour photolysis, which was also previously reported for SOCOL in the stand-alone photolysis codes intercomparison study (Sukhodolov et al., 2016; Karagodin-Doyennel et al., 2021). Water vapour sink (dehydration) by PSCs in the south polar lowermost stratosphere is well captured by the model.

Water vapor is a relatively long-lived species with a highly pronounced seasonality in the lower tropical stratosphere. This seasonality is related to tropospheric wave forcing, which is strongest in northern winter and weakest in northern summer,



530 leading to enhanced upwelling and colder temperatures at the tropical tropopause during boreal winter, and vice versa. In
combination with the tropical pipe and the subtropical transport barrier, this makes H₂O useful as a tracer of transport processes
in the tropical stratosphere (Mote et al., 1998). Figure 9 shows the water vapor ‘tape recorder’ signals obtained from
GOZCARDS and SOCOLv4 to illustrate the model performance in terms of vertical transport in the tropical low and middle
stratosphere. SOCOLv4 reproduces the seasonal cycle of water vapor at the tropopause level well. However, SOCOLv4 shows
535 too fast vertical propagation of the water vapor tape recorder in the observations similar to its predecessor versions (Stenke et
al., 2013), which is a common peculiarity for chemistry-climate models (e.g., Dietmüller et al., 2018). This issue is related to
the problems in horizontal and vertical mixing and diffusion and also to representation of the residual circulation in general,
all of which are largely dependent on resolution. The overestimation of H₂O of about 0.5 ppmv throughout the middle and
lower stratosphere (see also Fig. 7) is mostly due to the higher entry value during summer but can also be partly related to the
540 overestimation of mixing with higher altitudes.




Figure 9. Annual cycle of water vapor mixing ratio (ppmv) averaged between 15°N and 15°S and over 1991-2012 for the GOZCARDS satellite composite (upper panel) and SOCOLv4 (lower panel). Two consecutive cycles are shown. The white line in both panels indicates the approximate slope of the GOZCARDS tape recorder signal.

545



HCl. HCl is a chlorine reservoir that is widely used for tracking the overall chlorine abundance in the stratosphere, which also plays an important role in polar ozone chemistry. SOCOLv4 reproduces the HCl climatology pretty well with deviations mostly on the order of ± 5 -10% throughout the stratosphere. The most pronounced deviation is the $\sim 5\%$ underestimation by the model at the stratopause and the tropical mid-stratosphere. Figure 8 illustrates the seasonal cycle at 70 hPa, demonstrating a rather good representation of the southern high-latitude variations with some underestimation of the PSC activation effects, the reasons for which are yet unclear. A limiting factor here can be the amount of available ClONO₂ (Nakajima et al., 2016), whose observations are, however, rather sparse for these conditions. Note that HCl observational uncertainties are also the largest in the SH polar vortex (SPARC, 2017). In the northern polar regions, SOCOLv4 reproduces the semi-annual variability seen in observations, but overestimates the winter-time decrease and underestimates the summer-time one, which might be related to the weaker northern vortex that leads to higher temperature and lower frequency of PSCs formation.

NO_y. Figure 10 shows the odd nitrogen (NO_y=NO+NO₂+2N₂O₅+HNO₃) climatology from the model and its difference to the ACE-FTS and MIPAS instruments data. Odd nitrogen provides another important sink of ozone in the stratosphere and also negatively contributes to and is affected by other catalytic cycles of ozone destruction by deactivating odd hydrogen, chlorine, and bromine in the reservoir species HNO₃, ClONO₂, and BrONO₂, respectively. The main source of NO_y in the middle atmosphere is oxidation and photolysis of nitrous oxide (N₂O). The lowermost stratosphere is additionally highly affected by the mixing with the troposphere and therefore the NO_y budget also depends on the free tropospheric sources of odd nitrogen such as aircraft engines and lightning (Grewe, 2009). In the mesosphere, where odd nitrogen is mostly present in the form of nitric oxide (NO), the most important sources become the in-situ N₂ ionization by solar protons and middle energy electrons (MEE) and the transport from the thermosphere, where it is produced by auroral electrons and N₂ photolysis. Compared to ACE-FTS and MIPAS, which generally agree with each other, SOCOLv4 overestimates by $\sim 10\%$ the maximum concentrations in the mid-upper stratosphere, the region where the odd nitrogen catalytic cycle of ozone destruction is the most effective. In the upper stratosphere and the mesosphere, the model also overestimates NO_y everywhere except polar mesospheric regions. The overestimation increases gradually with height, reaching 80% in the tropical mesosphere, where the absolute values, however, are rather small. As there is almost no N₂O left at these altitudes, the overestimation likely comes from the underestimated sink through the NO photolysis. Despite the strong percentage deviation in the tropical mesosphere, the model underestimates NO_y in its polar regions, though the differences are almost insignificant due to a large interannual variability. The polar bias is most likely related to an underestimated contribution of either MEE or the thermospheric NO downward flux. Mironova et al. (2019) compared CMIP6 MEE ionization effects to calculations based on polar balloon measurements and found that the related odd nitrogen signal can be up to 100% too low when CMIP6 forcing was used. High- and mid-latitude upper stratospheric NO_y concentrations are then affected by both the positive bias in the low-latitudes and the negative bias in the polar high-latitudes.





Figure 10. NO_y zonal mean annual mean climatology (ppbv) calculated by (a) the SOCOLv4 model and (b) provided by MIPAS. (c and d) - relative difference between the model and the ACE-FTS and MIPAS observations. White areas are either the missing data or regions where the difference between model and observations is not statistically significant at 95% confidence level.

NO_y observational uncertainty is the largest in the lower stratosphere ($\pm 30\%$, SPARC, 2017), however both instruments suggest that the model has a pronounced low bias there. This can be explained by the insufficient production from N_2O oxidation, which shows an underestimation in the middle and upper stratosphere (Fig. 7). This underestimation is very similar to that in methane, which suggests that both are probably related to the overestimation of the vertical transport. Another potential reason could be overestimated sinks to reservoir species, however modelled HNO_3 and ClONO_2 (Fig. A2) agree reasonably well ($\pm 10\%$) with observations in the areas of their maximum concentrations in the lower stratospheric mid-latitudes. The middle stratospheric overestimation of HNO_3 is likely related to the overestimated abundance of H_2O and hence HO_x at these altitudes and the underestimated sink through photolysis (Sukhodolov et al., 2016). The tropical UTLS region shows large underestimation of both NO_y and HNO_3 , which suggests that either the free-tropospheric sink of odd nitrogen through HNO_3 washout or the source through production by lightning activity is biased. Overall, NO_y representation in the model is defined by an underestimated sink in the upper stratosphere and the mesosphere, and the underestimated production in the lower stratosphere, which results in the slight vertical shift of the maximum NO_y concentrations upwards.

Modelled NO_y seasonal cycle at 6 hPa is presented in Fig. 8. Model shows generally higher values than MIPAS due to NO_y burden being biased high, as discussed before. However, the seasonality of the variations is reasonably well reproduced at this altitude. The high and mid-latitude seasonality is controlled by the vertical motions, which either bring the NO_y -low air from above during the wintertime or the NO_y -high air from below in the summertime. This transport is modulated by horizontal mixing, which can cause some time shifts in the appearance of minima, like seen in the NH winter modelled results, where the contribution of the vertical transport is largely diminished by the mixing with midlatitudes. In the southern vortex, this is better reproduced by SOCOLv4 for the upper stratosphere, however in the middle and lower stratosphere (not shown) there are traces of the modelled southern vortex being too isolated, which is consistent with the analysis of CH_4 and N_2O . Boreal winter



minimum in the tropics appears due to increased upwelling and, thus, transport from below, which is well caught by the model.



605 **Figure 11. Ozone zonal mean annual mean climatology (1985-2012) calculated by the SOCOLv4 model (ppmv) (a) and provided by the GOZCARDS data (b). (c and d): relative difference between the model and the observational composites BASIC and GOZCARD. White areas are either the missing data or regions where the difference between model and observations is not statistically significant at 95% confidence level.**

610 **O₃.** Figure 11 represents the ozone mixing ratio climatology calculated by the model versus two different ozone datasets. Both observational datasets (BASIC and GOZCARDS) agree very well with each other. The difference plots demonstrate that SOCOLv4 has a very good representation of ozone in the tropics and subtropics throughout the stratosphere, except some overestimation in its lowermost part. The lower stratospheric overestimation spreads further to mid-latitudes and reaches values of more than 20%. This overestimation is a common feature in many CMIP6 models (Keeble et al., 2020 - see their Fig.4).

615 There is no clear chemical evidence from our previous analysis that can explain this overestimation. The low NO_y model bias (Fig. 10) would also contribute negatively to the ozone burden at these altitudes through reduced production by oxidation of carbon monoxide, methane, and higher hydrocarbons, similarly to the troposphere (Gauss et al., 2006; Grewe, 2009). Therefore, it is most likely that the high ozone bias is related to the photolysis problems in SOCOLv4. Sukhodolov et al. (2016) analyzed the stand-alone performance of the look-up table photolysis scheme applied in SOCOL and found that the ozone

620 photolysis in both channels is underestimated especially in the lower stratosphere. Although that study also showed the underestimated molecular oxygen photolysis at the same altitudes, which could partly compensate for the ozone photolysis underestimation, it appears that the latter still dominates. Comparison to the results of other codes in Sukhodolov et al. (2016) showed that the lower stratospheric problems in photolysis are probably related to the neglected Rayleigh scattering effect and temperature dependence of absorption cross sections and quantum yields. The applied photolysis corrections, as discussed in

625 the MEZON module description, seem to improve ozone throughout the stratosphere, however a more qualitative treatment of



630 photolysis is already planned for the future. Insufficient ozone photolysis also might affect HO_x formation via $\text{O}(^1\text{D}) + \text{H}_2\text{O}$ and related catalytic ozone destruction, thus intensifying the ozone high bias. Other factors that could contribute to the lower stratospheric overestimation in ozone are the ~ 2 K underestimation in local temperatures (Fig. 5), which would slow down the ozone destroying chemistry, and overestimated tropospheric ozone. In addition, SOCOLv4 so far does not include iodine chemistry, while some recent studies suggested that it can be responsible for $\sim 30\%$ of the halogen-induced ozone loss in the lowermost stratosphere (e.g., Koenig et al., 2020).


In the upper stratosphere at mid-latitudes, SOCOLv4 underestimates ozone by up to -20% . This feature has been present in SOCOL models since the first version (Egorova et al., 2005) and is related to the transport issues, caused by the insufficient vertical resolution affecting the wave propagation and breaking in the stratosphere/mesosphere. Though SOCOLv4 now has 635 47 vertical levels compared to 39 in previous versions, all additional levels are located in the troposphere, while the number of levels in the middle atmosphere remains almost the same. As was discussed earlier, SOCOLv4 and its atmospheric base ECHAM6 show stronger residual circulation that causes stronger vertical transport in the tropics and also stronger meridional and downward transport in the winter hemisphere. The increased BDC, however, also implies a stronger wave forcing for the polar vortex making it weaker and blurring the midlatitude transport barriers (Butchart, 2014). An associated increase of 640 stirring, which dominates over the horizontal transport by the meridional circulation in these conditions (Plumb, 2007) would then lead to a decrease in the meridional gradient of tracers. As depicted Figure 5, the modelled polar vortex strength in both hemispheres is notably underestimated in the uppermost stratosphere and the lower mesosphere.

Figure 12 shows that the midlatitude ozone underestimation at 1 hPa appears in the winter months and that it happens together with an underestimation in the polar most latitudes due to the more intense downward transport from the mesosphere. These 645 results are consistent with those seen from the analysis of methane. Part of the model biases here could be related to the 30% NO_y overestimation (Fig. 10) that would intensify the ozone destruction. In the lower levels (10 hPa and 70 hPa), the model shows a generally good agreement with GOZCARDS with some deviations in timing and magnitude of extremes that can also be partly attributed to the deficiencies in transport. Thus, a too deep ozone drop at 10 hPa in the southern polar vortex is consistent with the too low methane values under these conditions and can be explained by the insufficient mixing. Seasonality 650 of the tropical values at all latitudes agrees very well with observations, showing a decrease of ozone in the lower stratosphere during boreal winter and a pronounced semi-annual oscillation in the middle and upper stratosphere.

Although the observations are very uncertain in the polar night regions, both the ozone mixing ratio (Fig. 12) and the total ozone measurements (Fig. 13) suggest that in SOCOLv4 the size and intensity of the Antarctic ozone hole is overestimated. This can be related to either a too strong vortex isolation, as discussed before, or to problems in the polar heterogeneous 655 chemistry. Since the low bias is also apparent in the higher levels, another potential reason could be the poleward dislocation of the southern night jet in the upper and middle stratosphere (Fig. 5) which would imply less upper-stratospheric ozone production inside the vortex (McConnel and Jin, 2008) and, thus, less ozone available for the downward transport. The downward transport itself might also be biased high, as illustrated by small negative anomalies of N_2O and CH_4 in Fig. 7, which would negatively contribute to ozone by bringing too much chlorine from above.



660 A comparison of SOCOLv4 ozone to the prescribed CMIP6 ozone used by MPI-ESM (Fig. A3) is useful for understanding
 the differences in temperature and winds between SOCOLv4 and MPIESM. The comparison shows that SOCOLv4 has slightly
 more ozone in the tropical and subtropical upper stratosphere, but less ozone in the winter-time high latitudes. SOCOLv4 also
 has higher ozone in the mesosphere (up to 70%) and in the troposphere (up to 20%). Upper and mid-stratospheric differences
 in ozone contribute to a slightly stronger meridional temperature gradient due to less heating in the high latitudes and more
 665 heating in the tropics. This makes the polar vortices, which are too weak in MPIESM, slightly stronger in SOCOLv4. It is
 pronounced the most for the SH polar vortex, which is weaker than in observations throughout the whole stratosphere in
 MPIESM (Fig. A3d), while in SOCOLv4 (Fig. 5f) the negative anomaly is vanished in the middle stratosphere.




670 **Figure 12.** 1, 10, and 70 hPa ozone zonal mean seasonal cycles (ppmv). First column: SOCOLv4 data. Second column: GOZCARDS
 observational data. Two consecutive cycles are shown.

3.2.3 Total ozone climatology

Total ozone column (TOC) represents an aggregate of all features and model biases discussed above, as well as their
 compensating effects. Figures 13 and 14 show that SOCOLv4 has a very good representation of tropical and subtropical TOC



675 compared to MSRV2 and SBUV data, which is consistent with the results of Fig. 11 and 12. The model accurately simulates both the seasonal cycle (Fig. 13) and the annual mean climatologies (Fig. 14). Mostly due to the corrected photolysis rates, SOCOLv4's TOC is about 20 DU lower in the tropics and subtropics than in the previous model version, which results in better agreement with observations. Compared to MSRV2 data, SOCOLv4 shows almost perfect annual mean TOC in the tropics with some regional underestimation of up to 5 DU. Mid-latitudes are also generally well captured, with a slight overestimation of TOC by only 5-10 DU in both hemispheres.



680

Figure 13. Seasonal cycle of total ozone simulated by SOCOLv4 (a) against SOCOLv3 (b), MSRV2 (c), and SBUV observations (d) averaged for the 1980-2014 period.

685 Major disagreement appears in polar latitudes under wintertime conditions. It has to be mentioned that the polar night observations are subject to substantial uncertainties. SBUV data give no information for the polar night conditions at all, while MSRV2 provides almost full spatial and temporal coverage by assimilating a large variety of data including the ground-based observations into a chemistry-transport model (Van der A et al., 2015). This model, however, exploits a simplified scheme of stratospheric chemistry by Cariolle and Teysse re (2007) that tends to overestimate TOC under the ozone hole conditions by about 20 DU. However, taking this uncertainty into account while also using the direct satellite information by SBUV and
690 GOZCARDS (Fig. 11) at the edge of the polar vortices SOCOLv4 still shows some pronounced biases. Namely, the polar



695 ozone depletion in the SH starts right after the vortex formation (Fig. 13a), while the observations and the current state of
knowledge suggests that it starts later under the spring-time conditions (Fig. 13c). This feature is fully inherited from the
previous model version (Fig. 13b) and needs further detailed investigation. Likely reasons, as discussed earlier, could be related
to the transport issues, the details of polar heterogeneous chemistry, or also to the treatment of photolysis under high solar
zenith angles. In the Northern Hemisphere, Fig. 5 and 6 suggest a clear underestimation of the NH vortex strength in the model,
which can explain the difference with respect to observations. The observations suggest some local ozone decrease in the
middle of the winter, which is then followed by a fast increase in the springtime caused by inflow from the mid-latitudes. The
vortex in the model is weaker, meaning that polar air masses are less isolated and can get enriched with ozone from mid-
latitudes throughout the wintertime, and therefore the modelled field shows a more gradual increase instead of an abrupt rise.
700 This stronger mixing of polar vortex air is also responsible for the NH results in Fig. 14, where the model correctly reproduces
the shape of the high ozone values but also shows some overestimation in the modelled annual mean total ozone.





Figure 14. Global distribution of annual mean total ozone for 1980-2018 as (a) simulated by SOCOLv4 and (b) from the MSRv2 data record in DU. (c) Difference (a)-(b) in DU with contour lines -40, -30, -20, -10, -5, 0, 5, 10, 20, 30 DU.



705 3.2.4 Ozone evolution

The upper panel of Fig. 15 shows the evolution of the total ozone column (TOC). Near global TOC was calculated for this comparison, because SBUV has no data over polar night regions and the reliability of MSRv2 there is also low. SOCOLv3 and SOCOLv4 are presented as three separate ensemble members to illustrate the uncertainty due to the internal variability. SOCOLv3 data are taken from the RefC1 runs of CCMI-1, which assumes prescribed historical sea surface temperatures and sea ice coverages. Using prescribed SSTs in v3 explains why the variability among ensemble members is smaller in v3 than in v4, where the ocean is now interactive, and the ENSO variability is stochastic rather than fixed (Fig. 3b). It is clearly seen that SOCOLv3 has a large overestimation in TOC compared to other time-series. As was discussed above, this is mostly related to the problems in the photolysis scheme that are now manually corrected. SOCOLv4 and MSRv2 agree well in terms of the mean values, while the mean TOC from SBUV is lower by about 5-7 DU with respect to MSRv2. Generally, the slight overestimation of global TOC is also seen in many CMIP6 models, and SOCOL is well within the spread across CMIP6 models (Keeble et al., 2020, see their Fig. 6a). The CMIP6 ozone composite, instead, shows slightly lower TOC than SOCOLv4 and MSRv2 in Fig. 15a.



720 **Figure 15. Upper panel: Near global (55S-55N) mean ozone evolution simulated with SOCOLv4 (black), MSR2 (red), SBUV (dark blue), SOCOLv3 (light blue), and CMIP6 (green). Lower panel: the same but normalized by the mean over the whole period.**

725 For easier comparison of the TOC variability we show normalized values in Fig. 15b. The four decades shown are characterized by contributions from anthropogenic and natural forcings with varying importance during different periods. From 1980 to the mid-90s there is a strong hODSs-induced ozone decline. This negative tendency is further enhanced by the two strong volcanic eruptions of El-Chichon in 1982 and Pinatubo in 1991, which are then followed by several years of recovery. Since the mid-90s, following a recovery after Pinatubo, ozone starts to increase further, induced by the leveling-out of the hODS emissions as a consequence of the Montreal Protocol (Egorova et al., 2013; Chipperfield et al., 2015). Since the 2000s, there were no major eruptions and the ozone evolution is mostly determined by the decreasing halogen load and the global warming effects,




730 though there are still many related uncertainties and other important factors in specific regions, especially in the lower
stratosphere (Ball et al., 2018; Stone et al., 2018). Upper stratospheric ozone was shown to have the most pronounced recovery
(Petropavlovskikh et al., 2019; Chipperfield et al., 2018) due to being the most sensitive to the chlorine catalytic cycle of ozone
destruction. Furthermore, decreasing stratospheric temperatures also led to some deceleration of ozone destruction cycles. The
contribution of these positive changes to the total column is partly supplemented by an increase of the tropospheric column
735 ozone at a rate of about 6-7%/decade resulting from the continuous increase of the surface anthropogenic emissions of ozone
precursors like CH₄, CO, NO_x, and VOC (Ball et al., 2018). The recovery in the middle and lower stratosphere in the tropical
and extratropical areas is still unclear and observations even suggest negative trends in the lowermost stratosphere, which are,
however, barely significant due to a large dynamical variability of this region (Ball et al. 2018; 2019; Petropavlovskikh et al.,
2019). There are indications that these negative trends are dynamically driven (Orbe et al, 2020) and can be related to
740 acceleration of the BDC due to rising GHGs, however models mostly fail to reproduce the observations in detail (Ball et al.,
2020).

In Figure 15b, we show that SOCOLv3 had a reasonable TOC evolution under volcanically quiescent conditions, while the
volcanic effects were strongly overestimated. Unlike SOCOLv4, which uses an interactive sulphate aerosol scheme, SOCOLv3
used prescribed fields recommended for CCM1 that seem to be biased, probably because of the uncertain observations and also
745 the assumptions made during the calculation of aerosol mass from extinction observations. The interactive aerosol scheme
clearly improves the performance of SOCOLv4 in terms of volcanic effects on ozone. Sukhodolov et al. (2018) showed though
that the observed hemispheric asymmetry of volcanic signals in ozone are still difficult to properly reproduce with a free-
running model. Overall, the general behavior of the TOC evolution is very well captured by the new model both in terms of
the mean values of the near-global field and the variability.

In Figure 16, we looked into some drivers of the total column evolution. Depletion and further recovery of ozone in the middle
750 stratosphere is well captured by the model (Fig. 16a), due to the good representation of changes in stratospheric chlorine
(Figure 16e) and the continuous cooling of the stratosphere (Fig. 16b). The lower stratospheric (60 hPa) temperature and ozone
observations show a much larger uncertainty, but still somewhat pronounced negative trends that are also well captured by the
model. A thorough analysis of lower stratospheric processes requires comprehensive statistical tools, and a set of sensitivity
runs, which is already planned for future model applications.


755



760 **Figure 16.** (a and c): Near global (55°S-55°N) annual mean ozone evolution at 60 and 10 hPa compared to the BASIC composite. (b and d): Tropical annual mean evolution of temperature at 60 and 10 hPa compared to ERA5. (e): tropical monthly mean (smoothed with the 3-month running mean) evolution of HCl at 10 hPa compared to the GOZCARDS composite. Black lines: three model ensemble members. Orange lines: observations.



3.2.5 Stratospheric sulfur



765 **Figure 17.** Global total stratospheric sulfur mass in sulfate aerosols (a) and its zonal mean values (b-d) simulated with SOCOLv4 against HIRS, CMIP6, and MIPAS. Data is plotted in a logarithmic scale so that both large and small eruptions can be distinguished.



Figure 17 shows the total stratospheric sulfur load as well as the zonal mean stratospheric column of liquid H₂SO₄ for SOCOLv4 against CMIP6, MIPAS and HIRS. The tropopause that was applied to calculate the stratospheric burden for the CMIP6 and MIPAS datasets was calculated from the ERA-interim temperature data using the WMO definition of the tropopause as the lowest level at which the temperature decreases less than 2 K/km, whereas for SOCOLv4 the modelled tropopause is used. The peaks in aerosol originate from volcanic eruptions with the largest being the El Chichón (1982) and Pinatubo (1991) eruptions in the earlier time period and Kasatochi (2008), Sarychev (2009), Nabro (2011), and Calbuco (2015) as well as some others in the more recent past.

Generally, the aerosol evolution in the stratosphere is depicted quite well by the model in terms of the total aerosol amount, lifetime and meridional transport. There are, however, a few differences that need to be addressed. Prior to and after the 1982 El Chichón eruption there are some significant discrepancies between SOCOLv4 and CMIP6. In addition, the peak of stratospheric burden caused by this eruption (see Fig. 17a) is slightly shifted in time in comparison to the SOCOLv4 simulations. In Figure 17c, the delayed peak of CMIP6 is seen mostly in the northern hemisphere and the tropical aerosol levels, where the eruption took place, are lower than in SOCOLv4. The El Chichón eruption occurred between the activities of SAGE and SAGEII. Data from this time is a composite of SAMII data in the extratropics, as well several airborne lidar missions at low latitudes, which delivered data often many months apart, leading to very sparse data coverage. The gaps were then filled by means of interpolation (Thomason et al., 2018). It is difficult to make a conclusive statement on the SOCOLv4 performance of this time due to large uncertainties in observations. The next larger event, Nevado del Ruiz in 1985, already occurred during the SAGE II era, which is deemed more reliable by Thomason et al. (2018).

The 1991 Mt. Pinatubo eruption has been discussed several times, since observations show a plateaued value for several months after the eruption instead of a more distinct peak as models like SOCOLv4 suggest (e.g., Dhomse et al., 2020). Something similar has not been observed since then. However, no other well observed eruption was the size of the 1991 Mt. Pinatubo. A source for measurement artefacts during this time is the increase in opacity of the atmosphere due to large amounts of aerosol resulting from the eruption. From the modelling side, a potential deficiency could be insufficient vertical resolution of the stratosphere leading to increased vertical diffusion and faster meridional transport of aerosols. Niemeier and Schmidt (2017) used a model with a similar dynamical core, ECHAM-HAM, with 39 and 90 vertical levels. They showed that the version with a higher vertical resolution maintains aerosols longer in the tropical reservoir, which increases their lifetime after low-latitude emission events. The post-plateau e-folding time looks very similar for CMIP6 and SOCOLv4, but in the CMIP6 data there is another small increase after the main Mt. Pinatubo plateau without there having been any other major eruptions. Again this could be due to a change in the instruments used for the construction of this dataset.

The volcanically quiescent period in the late 1990s and early 2000s and the minor volcanic activity later, which is also better covered by observations, are well reproduced by the model. MIPAS observations are added here, as they are not included in the CMIP6 composite and thus represent an independent source of information. MIPAS and CMIP6 agree with each other quite well in terms of elevated values after eruptions, while the background level is slightly lower in MIPAS. Background levels in SOCOLv4 are located between CMIP6 and MIPAS and there is some spread over the ensemble members, which is



800 caused by the variable tropopause in the model combined with vertically prescribed SO₂ emission profiles. The model also captures the zonal mean distribution of elevated aerosol values, reproducing some equatorward transport after high-latitude eruptions of Kasatochi, Sarychev, and Calbuco, as well as dominant northward transport after the equatorial eruption of Nabro. It should be noted that our modelling results here are largely defined by the emission database that we used (Carn et al., 2016). Other volcanic emission databases report quite different estimates for each volcanic event, both in terms of emitted amount of
805 SO₂ and its vertical distribution (Timmreck et al., 2018).

Overall, we can conclude that SOCOLv4 nicely reproduces background aerosol levels and minor volcanic activity compared to observations, while partly underestimating the aerosol lifetime after major eruptions of El Chichón, Nevado del Ruiz, and Pinatubo. The latter can be attributed to a number of factors. First of all, the data from earlier periods is less reliable and had a lower temporal coverage (Thomason et al., 2018), which influences the forcing data used in the model, as well as the
810 observational composites for its validation. Furthermore, the physical processes influencing aerosol concentrations after these larger eruptions differ from the smaller but better observed ones that happened more recently. Since more SO₂ is injected into the stratosphere at once in a larger event, OH radicals are depleted more quickly, which can cause a delay in sulfate aerosol formation. Nevertheless, all SO₂ is eventually (in 2-3 months time) converted to sulfate aerosol and in higher concentrations the particles will collide and grow in size. These larger aerosols are more subjected to the force of gravity and more easily
815 sediment to the troposphere (Timmreck et al., 2009), while the long-term decay of aerosols after smaller eruptions is more controlled by the BDC transport (Günther et al., 2018). Given that all major eruptions were tropical, a bias in vertical numerical diffusion due to insufficient vertical resolution could further intensify the sedimentation to the lower stratosphere and the troposphere and also reduce the confinement of aerosols in the tropical reservoir via increased leakage through the shallow branch of the BDC.

820 **4 Conclusions and outlook**

This paper presents the fourth generation of the coupled model chemistry-climate SOCOL. Unlike its predecessor chemistry-climate model SOCOLv3, SOCOLv4 includes an interactive ocean and sulfate aerosol modules. The underlying general circulation model has also been updated from ECHAM5 to ECHAM6. Our validation of the new model showed that it performs very well in terms of the mean state of most variables and the large-scale variability of the system. Namely, the warming trends
825 in the troposphere, the cooling trends in the stratosphere, and variability of the stratospheric and the total column ozone agree very well with observations. In the stratosphere, SOCOLv4 shows very good results for the low- and mid-latitude ozone and reasonably well reproduces climatologies of other trace gases. Compared to SOCOLv3, the new version of the model now performs significantly better both in terms of absolute values and variability of ozone, which is mostly due to corrections in the photolysis scheme and the inclusion of interactive stratospheric aerosol scheme.

830 However, as in the previous version, there are still some issues. Stratospheric polar night jets are weaker in the model throughout the whole stratosphere in the NH and in the upper SH stratosphere in comparison with the reanalysis data.




835 Accordingly, the advective Brewer-Dobson circulation and horizontal diffusion are also biased and too strong. All this affects
the transport and distribution of tracers, introducing biases in CH₄, N₂O, ozone, and related species. Photochemistry itself also
suffers from the outdated photolysis look-up-table (LUT) scheme, which needs to be replaced either by a LUT that also
840 considers temperature dependence or by an online radiative-transfer scheme. Furthermore, photolysis parameterizations of the
short-wave UV need to be reviewed and updated, as we identified clear biases in the mesospheric budget of odd nitrogen,
hydrogen, and oxygen. SOCOLv4 still underestimates polar winter and spring ozone in the SH, which needs further detailed
investigation. Performance of the sulfate aerosol scheme is very good for the smaller eruptions and background conditions,
but the duration of volcanic peaks after major events seems to be underestimated. An increase of the vertical resolution to 95
845 levels could be a potential source for major improvements, especially for stratospheric circulation and transport. However, it
would at least double the computational needs, which might be too expensive for long-term experiments, given the number of
tracers in the model. Marine emissions of species are currently prescribed in the model, but the interactive ocean and marine
biogeochemistry modules now allow either to directly couple some emissions or to parameterize them as a function of other
parameters. Inclusion of iodine chemistry is being currently tested and will also contribute to future updates.
845 Nevertheless, we showed that the model version documented here is already in a good condition to be used as a tool for
studying interactions between the earth system components. The presence of the interactive ocean and a successful
representation of recent climate and ozone layer trends provide a strong case for this model to be applied for studies looking
at future evolution and effects of greenhouse gases and ozone-destroying substances, as well as potential geoengineering
measures through sulfur injections. SOCOLv4 will also be used in the upcoming model intercomparison project CCMI phase
850 2.

855


860




Appendix A



865 **Figure A1: Zonal mean climatology differences between data simulated by MPI-ESM and the reanalysis data for temperature (a-b) and zonal wind (c-d). White places denote regions where the difference between model and reanalysis data is not statistically significant at 95% confidence level calculated with the Student's t-test.**



870 **Figure A2.** HNO_3 and ClONO_2 zonal mean annual mean climatologies (ppbv). First column: SOCOLv4 data. Second column: ACE-FTS and GOZCARDS observational data. Third column: relative difference between the model and observations in %. Observational missing data mask is applied to the model data. Note that different species have different averaged periods.



875 **Figure A3.** Ozone zonal and annual mean climatology (1985-2012) calculated by (a) the SOCOLv4 model (ppmv) and (b) provided by the CMIP6 modelling composite. The relative difference between SOCOLv4 and CMIP6 are shown in (c). White areas are either missing data or regions where the difference between model and observations is not statistically significant at 95% confidence level.



Code and data availability

880 The SOCOLv4.0 code can be downloaded from <https://doi.org/10.5281/zenodo.4570622> (Sukhodolov et al., 2021). In case
of problems, please contact the corresponding author. ERA5 data is available on the Copernicus Climate Change Service (C3S)
Climate Data Store: <https://cds.climate.copernicus.eu/#!/search?text=ERA5&type=dataset>. MERRA-2 data are available at
MDISC, managed by the NASA Goddard Earth Sciences (GES) Data and Information Services Center (DISC):
885 <https://disc.gsfc.nasa.gov/datasets?project=MERRA-2>. The Berkeley Earth Surface Temperatures data is available at
<http://berkeleyearth.org/data/>. GOZCARDS data can be found here: <https://gozcards.jpl.nasa.gov/info.php>. The latest BASIC
ozone composite data is available at <https://data.mendeley.com/datasets/2mgx2xzzpk/3>. The ACE-FTS data can be found in
890 <http://www.ace.uwaterloo.ca/climatology/3.5/netcdf/>. The GloSSAC data v1.1 prepared for CMIP6 is available at
ftp://iacftp.ethz.ch/pub_read/luo/CMIP6_SAD_radForcing_v4.0.0/. MSRv2 data can be downloaded from
<https://www.temis.nl/protocols/O3global.php>. SBUVv8.6 merged ozone data set can be found in https://acd-ext.gsfc.nasa.gov/Data_services/merged/instruments.html. The CMIP6 ozone composite is accessible via the input4MIPs data
server.

Author contribution

TS conducted the simulations, analysed the data, and drafted most of the paper. TE, AKD, and CB drafted parts of the paper.
TS, AS, AF, and ER led the model development work. All authors participated in the model development and discussions
about the results.

895 Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

This work was supported by the Swiss National Science Foundation (SNSF) under grants 200021_169241 (VEC) and 200020-
182239 (POLE). All calculations with the Atmosphere-Ocean-Aerosol-Chemistry-Climate Model SOCOLv4.0 were supported
900 by a grant from the Swiss National Supercomputing Centre (CSCS) under projects ID S-901 and ID S-1029. SNSF is also
acknowledged for support under the Ambizione Grant PZ00P2-180043 (GC and MF). SV is supported by the ETH Research
Grant ETH-17 19-2. Part of the model development was performed on the ETH Zürich cluster EULER. We thank the Center
for Climate Systems Modeling (C2SM) at ETH Zurich, the Max Planck Institute for Meteorology in Hamburg, Germany, and
personally Sylvaine Ferrachat (ETH), Urs Beyerle (ETH) and Sebastian Rast (MPI) for technical support related to hosting
905 and development of the model code. We also thank Bei-Ping Luo (ETH) for providing the Mie theory lookup table and Bernd
Funke (CSIC) and Michael Höpfner (KIT) for providing the MIPAS data for NO_y and sulfate aerosol mass, respectively.



References

- Alsing, J. and Ball, W.: BASIC Composite Ozone Time-Series Data, Mendeley Data, V3, <https://doi.org/10.17632/2mgx2xzzpk.3>, 2019.
- 910 Andres, R. J., and Kasgnoc, A.D.: A time-averaged inventory of subaerial volcanic sulfur emissions, *J. Geophys. Res.*, 103, 25,251–25,261, <https://doi.org/doi:10.1029/98JD02091>, 1998.
- Arsenovic, P., Rozanov, E., Anet, J., Stenke, A., Schmutz, W., and Peter, T.: Implications of potential future grand solar minimum for ozone layer and climate, *Atmos. Chem. Phys.*, 18, 3469–3483, <https://doi.org/10.5194/acp-18-3469-2018>, 2018.
- Ayers, G.P., Gillett, R.W., and Gras, J.L.: On the vapor pressure of sulfuric acid, *Geophys. Res. Lett.*, 7, 433–436, <https://doi.org/10.1029/GL007i006p00433>, 1980.
- 915 Bacmeister, J. T., Phillips, A. S., Neale, R. B., Simpson, I. R., DuVivier, A. K., Hodzic, A., and Randel, W. J.: The whole atmosphere community climate model version 6 (WACCM6), *J. Geophys. Res. Atmos.*, 124, 12,380–12,403, <https://doi.org/10.1029/2019JD030943>, 2019.
- Baran, A. J. and Foot, J. S.: New application of the operational sounder HIRS in determining a climatology of sulphuric acid aerosol from the Pinatubo eruption, *J. Geophys. Res.-Atmos.*, 99, 25673–25679, <https://doi.org/10.1029/94JD02044>, 1994.
- 920 Ball, W. T., Alsing, J., Mortlock, D. J., Staehelin, J., Haigh, J. D., Peter, T., Tummon, F., Stübi, R., Stenke, A., Anderson, J., Bourassa, A., Davis, S. M., Degenstein, D., Frith, S., Froidevaux, L., Roth, C., Sofieva, V., Wang, R., Wild, J., Yu, P., Ziemke, J. R., and Rozanov, E. V.: Evidence for a continuous decline in lower stratospheric ozone offsetting ozone layer recovery, *Atmos. Chem. Phys.*, 18, 1379–1394, <https://doi.org/10.5194/acp-18-1379-2018>, 2018.
- 925 Ball, W. T., Alsing, J., Staehelin, J., Davis, S. M., Froidevaux, L., and Peter, T.: Stratospheric ozone trends for 1985–2018: sensitivity to recent large variability, *Atmos. Chem. Phys.*, 19, 12731–12748, <https://doi.org/10.5194/acp-19-12731-2019>, 2019.
- Ball, W. T., Chiodo, G., Abalos, M., Alsing, J., and Stenke, A.: Inconsistencies between chemistry–climate models and observed lower stratospheric ozone trends since 1998, *Atmos. Chem. Phys.*, 20, 9737–9752, <https://doi.org/10.5194/acp-20-9737-2020>, 2020.
- 930 Bernath, P. F., McElroy, C. T., Abrams, M. C., Boone, C. D., Butler, M., Camy-Peyret, C., Carleer, M., Clerbaux, C., Coheur, P.-F., Colin, R., DeCola, P., DeMazière, M., Drummond, J. R., Dufour, D., Evans, W. F. J., Fast, H., Fussen, D., Gilbert, K., Jennings, D. E., Llewellyn, E. J., Lowe, R. P., Mahieu, E., McConnell, J. C., McHugh, M., McLeod, S. D., Michaud, R., Midwinter, C., Nassar, R., Nichitiu, F., Nowlan, C., Rinsland, C. P., Rochon, Y. J., Rowlands, N., Semeniuk, K., Simon, P., Skelton, R., Sloan, J. J., Soucy, M.-A., Strong, K., Tremblay, P., Turnbull, D., Walker, K. A., Walkty, I., Wardle, D. A., Wehrle, V., Zander, R., and Zou, J.: Atmospheric Chemistry Experiment (ACE): Mission overview, *Geophys. Res. Lett.*, 32, L15S01, <https://doi.org/10.1029/2005GL022386>, 2005.



- Biermann, U. M., Luo, B. P., and Peter, T.: Absorption spectra and optical constants of binary and ternary solutions of H₂SO₄, HNO₃, and H₂O in the mid infrared at atmospheric temperatures, *J. Phys. Chem. A*, 104, 783–793, 940 <https://doi.org/10.1021/jp992349i>, 2000.
- Bock, L., Lauer, A., Schlund, M., Barreiro, M., Bellouin, N., Jones, C., Meehl, G.A., Predoi, V., Roberts, M.J., and Eyring, V.: Quantifying progress across different CMIP phases with the ESMValTool, *J. Geophys. Res.: Atmospheres*, 125, e2019JD032321, <https://doi.org/10.1029/2019JD032321>, 2020.
- Brinkop, S. and Roeckner, E.: Sensitivity of a general circulation model to parameterizations of cloud–turbulence interactions 945 in the atmospheric boundary layer, *Tellus A*, 47, 197–220, <https://doi.org/10.1034/j.1600-0870.1995.t01-1-00004.x>, 1995.
- Brovkin, V., Boysen, L., Raddatz, T., Gayler, V., Loew, A. and Claussen, M.: Evaluation of vegetation cover and land-surface albedo in MPI-ESM CMIP5 simulations, *J. Adv. Model. Earth Syst.*, 5, 48–57, <https://doi.org/10.1029/2012MS000169>, 2013.
- Brönnimann, S., Jacques-Coper, M., Rozanov, E., Fischer, A.M., Morgenstern, O., Zeng, G., Akiyoshi, H., and Yamashita, Y.: Tropical circulation and precipitation response to ozone depletion and recovery, *Environ. Res. Lett.*, 6, 064011, 950 <https://doi.org/10.1088/1748-9326/aa7416>, 2017.
- Bulgin, C.E., Merchant, C.J., and Ferreira, D.: Tendencies, variability and persistence of sea surface temperature anomalies. *Sci Rep*, 10, 7986, <https://doi.org/10.1038/s41598-020-64785-9>, 2020.
- Burkholder, J. B., Sander, S. P., Abbatt, J., Barker, J. R., Huie, R. E., Kolb, C. E., Kurylo, M. J., Orkin, V. L., Wilmouth, D. M., and Wine, P. H.: Chemical kinetics and photochemical data for use in atmospheric studies, Evaluation No. 18, JPL 955 Publication 15-10, Jet Propul. Lab., Pasadena, Calif. [Available at <https://jpldataeval.jpl.nasa.gov/>], 2015.
- Butchart, N.: The Brewer-Dobson circulation, *Rev. Geophys.*, 52, 157–184, <https://doi.org/10.1002/2013RG000448>, 2014.
- Engel, A., Rigby, M., Burkholder, J. B., Fernandez, R.P., Froidevaux, L., Hall, B. D., Hossaini, R., Saito, T., Vollmer, M.K., and Yao, B.: Update on Ozone-Depleting Substances (ODSs) and other gases of interest to the Montreal Protocol, Chapter 1 960 in *Scientific Assessment of Ozone Depletion: 2018*, Global Ozone Research and Monitoring, Project–Report No. 58, World Meteorological Organization, Geneva, Switzerland, 2018.
- Cariolle, D. and Teyssède, H.: A revised linear ozone photochemistry parameterization for use in transport and general circulation models: multi-annual simulations, *Atmos. Chem. Phys.*, 7, 2183–2196, <https://doi.org/10.5194/acp-7-2183-2007>, 2007.
- Carn, S., Clarisse, L., Prata, A.: Multi-decadal satellite measurements of global volcanic degassing, *J. Volcanol. Geotherm. Res.*, 311, 99–134, <https://doi.org/10.1016/j.jvolgeores.2016.01.002>, 2016. 965
- Carslaw, K. S., Luo, B. P., and Peter, T.: An analytic-expression for the composition of aqueous HNO₃–H₂SO₄ stratospheric aerosols including gas-phase removal of HNO₃, *Geophys. Res. Lett.*, 22, 1877–1880, 1995.
- Checa-Garcia, R.: CMIP6 Ozone forcing dataset: supporting information. Zenodo. <https://doi.org/10.5281/zenodo.1135127>, 2018.
- Chiodo, G., Polvani, L.M., Marsh, D.R., Ball, W., Muthers, S., Stenke, A., Rozanov, E., and Tsigaridis, K.: The ozone response 970 to quadrupled CO₂ concentrations, *J. Climate*, 31, 3893–3907, <https://doi.org/10.1175/JCLI-D-17-0492.1>, 2018.



- Chipperfield, M. P., Dhomse, S. S., Feng, W., McKenzie, R. L., Velders, G.J.M., and Pyle, J. A.: Quantifying the ozone and ultraviolet benefits already achieved by the Montreal Protocol, *Nat. Commun.*, 6, 7233, <https://doi.org/10.1038/ncomms8233>, 2015.
- 975 Chipperfield, M., Bekki, S., Dhomse, S., Harris, N.R.P., Hassler, B., Hossaini, R., Steinbrecht, W., Thiéblemont, R., and Weber M.: Detecting recovery of the stratospheric ozone layer, *Nature*, 549, 211–218, <https://doi.org/10.1038/nature23681>, 2017.
- Chipperfield, M. P., Dhomse, S., Hossaini, R., Feng, W., Santee, M. L., Weber, M., Burrows, J. P., Wild, J. D., Loyola, D., Coldewey-Egbers, M.: On the cause of recent variations in lower stratospheric ozone. *Geophysical Research Letters*, 45, 5718–
- 980 5726, <https://doi.org/10.1029/2018GL078071>, 2018.
- Cowtan, K., Hausfather, Z., Hawkins, E., Jacobs, P., Mann, M. E., Miller, S. K., Steinman, B. A., Stolpe, M. B., and Way, R. G.: Robust comparison of climate models with observations using blended land air and ocean sea surface temperatures, *Geophys. Res. Lett.*, 42, 6526–6534, <https://doi.org/10.1002/2015GL064888>, 2015.
- Dentener, F., Kinne, S., Bond, T., Boucher, O., Cofala, J., Generoso, S., Ginoux, P., Gong, S., Hoelzemann, J. J., Ito, A.,
- 985 Marelli, L., Penner, J. E., Putaud, J.-P., Textor, C., Schulz, M., van der Werf, G. R., and Wilson, J.: Emissions of primary aerosol and precursor gases in the years 2000 and 1750 prescribed data-sets for AeroCom, *Atmos. Chem. Phys.*, 6, 4321–4344, <https://doi.org/10.5194/acp-6-4321-2006>, 2006.
- Dhomse, S. S., Mann, G. W., Antuña Marrero, J. C., Shallcross, S. E., Chipperfield, M. P., Carslaw, K. S., Marshall, L., Abraham, N. L., and Johnson, C. E.: Evaluating the simulated radiative forcings, aerosol properties, and stratospheric
- 990 warmings from the 1963 Mt Agung, 1982 El Chichon, and 1991 Mt Pinatubo volcanic aerosol clouds, *Atmos. Chem. Phys.*, 20, 13627–13654, <https://doi.org/10.5194/acp-20-13627-2020>.
- Dietmüller, S., Eichinger, R., Garny, H., Birner, T., Boenisch, H., Pitari, G., Mancini, E., Visioni, D., Stenke, A., Revell, L., Rozanov, E., Plummer, D. A., Scinocca, J., Jöckel, P., Oman, L., Deushi, M., Kiyotaka, S., Kinnison, D. E., Garcia, R., Morgenstern, O., Zeng, G., Stone, K. A., and Schofield, R.: Quantifying the effect of mixing on the mean age of air in CCMVal-
- 995 2 and CCMI-1 models, *Atmos. Chem. Phys.*, 18, 6699–6720, <https://doi.org/10.5194/acp-18-6699-2018>, 2018.
- Ding, Q., Schweiger, A., L’Heureux, M., Steig, E. J., Battisti, D. S., Johnson, N. C., Blanchard-Wrigglesworth, E., Po-Chedley, S., Zhang, Q., Harnos, K., Bushuk, M., Markle, B., and Baxter, I.: Fingerprints of internal drivers of Arctic sea ice loss in observations and model simulations, *Nature Geosci.*, 12, 28–33, <https://doi.org/10.1038/s41561-018-0256-8>, 2019.
- Egorova, T., Rozanov, E., Zubov, V., and Karol, I.: Model for investigating ozone trends (MEZON), *Izvestiya, Atmos. Ocean.*
- 1000 *Phys.*, 39, 277–292, 2003.
- Egorova, T., Rozanov, E., Zubov, V., Manzini, E., Schmutz, W., and Peter, T.: Chemistry-climate model SOCOL: a validation of the present-day climatology, *Atmos. Chem. Phys.*, 5, 1557–1576, <https://doi.org/10.5194/acp-5-1557-2005>, 2005.
- Egorova, T., Rozanov, E., Gröbner, J., Hauser, M., and Schmutz, W.: Montreal Protocol Benefits simulated with CCM SOCOL, *Atmos. Chem. Phys.*, 13, 3811–3823, <https://doi.org/10.5194/acp-13-3811-2013>, 2013.



- 1005 Ehhalt, D., Prather, M., Dentener, F., Derwent, R., Dlugokencky, E., Holland, E., Isaksen, I., Katima, J., Kirchhoff, V., Matson, P., Midgley, P., Wang, M.: Atmospheric chemistry and greenhouse gases, in: *Climate Change 2001: the Scientific Basis, Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Houghton, J.T., Ding, Y., Griggs, D.J., Noguer, M., van der Linden, P.J., Dai, X., Maskell, K., Johnson, C.A., Cambridge University Press, Cambridge, United Kingdom and New York, 239–288, 2001.
- 1010 Evans, M. J., and Jacob, D. J.: Impact of new laboratory studies of N₂O₅ hydrolysis on global model budgets of tropospheric nitrogen oxides, ozone, and OH, *Geophys. Res. Lett.*, 32, L09813, <https://doi.org/10.1029/2005GL022469>, 2005.
- Farman, J., Gardiner, B., and Shanklin, J.: Large losses of total ozone in Antarctica reveal seasonal ClO_x/NO_x interaction, *Nature*, 315, 207–210, <https://doi.org/10.1038/315207a0>, 1985.
- Feinberg A. I., Coulon A., Stenke A., Schwietzke S., and Peter T.: Isotopic source signatures: Impact of regional variability on the δ¹³C_{CH₄} trend and spatial distribution, *Atm. Env.*, 174, 99–111, <https://doi.org/10.1016/j.atmosenv.2017.11.037>, 2018.
- 1015 Feinberg, A., Sukhodolov, T., Luo, B.-P., Rozanov, E., Winkel, L. H. E., Peter, T., and Stenke, A.: Improved tropospheric and stratospheric sulfur cycle in the aerosol–chemistry–climate model SOCOL-AERv2, *Geosci. Model Dev.*, 12, 3863–3887, <https://doi.org/10.5194/gmd-12-3863-2019>, 2019.
- Feinberg, A., Maliki, M., Stenke, A., Sudret, B., Peter, T., and Winkel, L. H. E.: Mapping the drivers of uncertainty in atmospheric selenium deposition with global sensitivity analysis, *Atmos. Chem. Phys.*, 20, 1363–1390, <https://doi.org/10.5194/acp-20-1363-2020>, 2020.
- 1020 Fiedler, S., Stevens, B., and Mauritsen, T.: On the sensitivity of anthropogenic aerosol forcing to model-internal variability and parameterizing a Twomey effect, *J. Adv. Model. Earth Syst.*, 9, 1325–1341, <https://doi.org/10.1002/2017MS000932>, 2017.
- Fleming, E. L., Newman, P. A., Liang, Q., & Daniel, J. S.: The impact of continuing CFC-11 emissions on stratospheric ozone. *J. Geophys. Res.: Atmospheres*, 125, e2019JD031849, <https://doi.org/10.1029/2019JD031849>, 2020.
- 1025 Froidevaux, L., Anderson, J., Wang, H.-J., Fuller, R. A., Schwartz, M. J., Santee, M. L., Livesey, N. J., Pumphrey, H. C., Bernath, P. F., Russell, III, J. M., and McCormick, M. P.: Global O₃ Chemistry And Related trace gas Data records for the Stratosphere (GOZCARDS): methodology and sample results with a focus on HCl, H₂O, and O₃, *Atmos. Chem. Phys.*, 15, 10471–10507, [doi:10.5194/acp-15-10471-2015](https://doi.org/10.5194/acp-15-10471-2015), 2015.
- 1030 Fuchs, N. A.: The mechanics of aerosols, *Q.J.R. Meteorol. Soc.*, 91, 249–249, <https://doi.org/10.1002/qj.49709138822>, 1964.
- Funke, B., López-Puertas, M., Holt, L., Randall, C. E., Stiller, G. P., and von Clarmann, T.: Hemispheric distributions and interannual variability of NO_y produced by energetic particle precipitation in 2002–2012, *J. Geophys. Res. Atmos.*, 119, 13565–13582, <https://doi.org/10.1002/2014JD022423>, 2014.
- Funke, B., López-Puertas, M., Stiller, G. P., Versick, S., and von Clarmann, T.: A semi-empirical model for meso-spheric and stratospheric NO_y produced by energetic particle precipitation, *Atmos. Chem. Phys.*, 16, 8667–8693, <https://doi.org/10.5194/acp-16-8667-2016>, 2016.
- 1035 Gauss, M., Isaksen, I. S. A., Lee, D. S., and Søvde, O. A.: Impact of aircraft NO_x emissions on the atmosphere – tradeoffs to reduce the impact, *Atmos. Chem. Phys.*, 6, 1529–1548, <https://doi.org/10.5194/acp-6-1529-2006>, 2006.



- Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G.,
1040 Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., da Silva, A. M., Gu, W., Kim,
G.-K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S.
D., Sienkiewicz, M., and Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2
(MERRA-2), *J. Climate*, 30, 5419–5454, <https://doi.org/10.1175/JCLI-D-16-0758.1>, 2017.
- Gerber, E.P., Butler, A., Calvo, N., Charlton-Perez, A., Giorgetta, M., Manzini, E., Perlwitz, J., Polvani, L.M., Sassi, F., Scaife,
1045 A.A., Shaw, T.A., Son, S.-W., and Watanabe, Sh.: Assessing and understanding the impact of stratospheric dynamics and
variability on the earth system, *Bull. Amer. Meteor. Soc.*, 93, 845-859, <https://doi.org/10.1175/BAMS-D-11-00145.1>, 2012.
- Gottelman, A., Mills, M. J., Kinnison, D. E., Garcia, R. R., Smith, A. K., Marsh, D. R., Tilmes, S., Vitt, F., Bardeen, C. G.,
McInerny, J., Liu, H.-L., Solomon, S. C., Polvani, L. M., Emmons, L. K., Lamarque, J.-F., Richter, J. H., Glanville, A. S.,
Giorgetta, M. A.: Der Einfluss der quasi-zweijährigen Oszillation: Modellrechnungen mit ECHAM4, Max-Planck-Institut für
1050 Meteorologie, Hamburg, Examensarbeit Nr. 40, MPI-Report 218, 1996.
- Giorgetta, M. A. , Jungclaus, J. , Reick, C. H. , Legutke, S. , Bader, J. , Böttinger, M. , Brovkin, V. , Crueger, T. , Esch, M. ,
Fieg, K. , Glushak, K. , Gayler, V. , Haak, H. , Hollweg, H.-D. , Ilyina, T. , Kinne, S. , Kornblueh, L. , Matei, D. , Mauritsen,
T. , Mikolajewicz, U. M. W. , Notz, D. , Pithan, F. , Raddatz, T. , Rast, S. , Redler, R. , Roeckner, E. , Schmidt, H. , Schnur,
R. , Segschneider, J. , Six, Katharina, D. , Stockhause, M. , Timmreck, C. , Wegner, J. , Widmann, H. , Wieners, K.-H. ,
1055 Claussen, M. , Marotzke, J. , and Stevens, B. : Climate and carbon cycle changes demo 1850 to 2100 in MPI-ESM simulations
for the Coupled Model Intercomparison Project phase 5, *J. Adv. Mod. Earth Sys.*, 5, 572–597,
<https://doi.org/10.1002/jame.20038>, 2013.
- Goll, D. S., Winkler, A. J., Raddatz, T., Dong, N., Prentice, I. C., Ciais, P., and Brovkin, V.: Carbon–nitrogen interactions in
idealized simulations with JSBACH (version 3.10), *Geosci. Model Dev.*, 10, 2009–2030, [https://doi.org/10.5194/gmd-10-](https://doi.org/10.5194/gmd-10-2009-2017)
1060 [2009-2017](https://doi.org/10.5194/gmd-10-2009-2017), 2017.
- Craig, A., Valcke, S., and Coquart, L.: Development and performance of a new version of the OASIS coupler, OASIS3-
MCT_3.0, *Geosci. Model Dev.*, 10, 3297–3308, <https://doi.org/10.5194/gmd-10-3297-2017>, 2017.
- Grewe, V.: Impact of lightning on air chemistry and climate, in: *Lightning: Principles, Instruments and Applications (Review
of Modern Lightning Research)*, edited by: Betz, H. D., Schumann, U., and Laroche, P., Springer Netherlands , 537-549,
1065 <https://doi.org/10.1007/978-1-4020-9079-0>, 2009.
- Günther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene
emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmos. Chem. Phys.*, 6, 3181–3210,
<https://doi.org/10.5194/acp-6-3181-2006>, 2006.
- Günther, A., Höpfner, M., Sinnhuber, B.-M., Griessbach, S., Deshler, T., von Clarmann, T., and Stiller, G.: MIPAS
1070 observations of volcanic sulfate aerosol and sulfur dioxide in the stratosphere, *Atmos. Chem. Phys.*, 18, 1217–1239,
<https://doi.org/10.5194/acp-18-1217-2018>, 2018.



- Guo, Y., Yu, Y., Lin, P., Liu, H., He, B., Bao, O., Zhao, S., and Wang, X.: Overview of the CMIP6 Historical Experiment Datasets with the Climate System Model CAS FGOALS-f3-L, *Adv. Atmos. Sci.*, 37, 1057–1066, <https://doi.org/10.1007/s00376-020-2004-4>, 2020.
- 1075 Haase, S. and Matthes, K.: The importance of interactive chemistry for stratosphere–troposphere coupling, *Atmos. Chem. Phys.*, 19, 3417–3432, <https://doi.org/10.5194/acp-19-3417-2019>, 2019.
- Hagemann, S., and Stacke, T.: Impact of the soil hydrology scheme on simulated soil moisture memory, *Clim. Dyn.*, 44, 1731–1750, <https://doi.org/10.1007/s00382-014-2221-6>, 2015.
- Hansen, J., Ruedy, M. Sato, and R. Reynolds: Global surface air temperature in 1995: Return to pre-Pinatubo level, *Geophys. Res. Lett.*, 23, 1665–1668, <https://doi.org/10.1029/96GL01040>, 1996.
- 1080 Hanson, D. R. and Ravishankara, A. R.: Reactive Uptake of ClONO₂ onto Sulfuric Acid Due to Reaction with HCl and H₂O, *J. Phys. Chem.*, 98, 5728–5735, <https://doi.org/10.1021/j100073a026>, 1994.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., 1085 Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, Ph., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I. Vamborg, F., Villaume, S., Thépaut, J.-N.: The ERA5 global reanalysis. *Q. J. R. Meteor. Soc.*, 146, 1999–2029, <https://doi.org/10.1002/qj.3803>, 2020.
- Hibler, W.D.: A Dynamic Thermodynamic Sea Ice Model, *J. Phys. Oceanogr.*, 9, 815–846, <https://doi.org/10.1175/1520-1090> 0485(1979)009, 1979.
- Hines, C. O.: Doppler-spread parameterization of gravity-wave momentum deposition in the middle atmosphere, 1, Basic formulation, *J. Atmos. Solar Terr. Phys.*, 59, 371–386, [https://doi.org/10.1016/S1364-6826\(96\)00079-X](https://doi.org/10.1016/S1364-6826(96)00079-X), 1997a.
- Hines, C. O.: Doppler-spread parameterization of gravity-wave momentum deposition in the middle atmosphere, 2, Broad and quasi monochromatic spectra, and implementation, *J. Atmos. Solar Terr. Phys.*, 59, 387–400, [https://doi.org/10.1016/S1364-6826\(96\)00080-6](https://doi.org/10.1016/S1364-6826(96)00080-6), 1997b.
- 1095 Hoesly, R. M., Smith, S. J., Feng, L., Klimont, Z., Janssens-Maenhout, G., Pitkanen, T., Seibert, J. J., Vu, L., Andres, R. J., Bolt, R. M., Bond, T. C., Dawidowski, L., Kholod, N., Kurokawa, J.-I., Li, M., Liu, L., Lu, Z., Moura, M. C. P., O'Rourke, P. R., and Zhang, Q.: Historical (1750–2014) anthropogenic emissions of reactive gases and aerosols from the Community Emissions Data System (CEDS), *Geosci. Model Dev.*, 11, 369–408, <https://doi.org/10.5194/gmd-11-369-2018>, 2018.
- 1100 Hurtt, G. C., Chini, L. P., Frohling, S., Betts, R. A., Feddema, J., Fischer, G., Fisk, J. P., Hibbard, K., Houghton, R. A., Janetos, A., Jones, C. D., Kindermann, G., Kinoshita, T., Klein Goldewijk, K., Riahi, K., Shevliakova, E., Smith, S., Stehfest, E., Thomson, A., Thornton, P., van Vuuren, D. P., and Wang, Y. P.: Harmonization of land-use scenarios for the period 1500–2100: 600 years of global gridded annual land-use transitions, wood harvest, and resulting secondary lands, *Climatic Change*, 109, 117–161, <https://doi.org/10.1007/s10584-011-0153-2>, 2011.



- 1105 Iacono, M. J., Delamere, J. S., Mlawer, E. J., Shephard, M. W., Clough, S. A., and Collins, W. D.: Radiative forcing by long-lived greenhouse gases: Calculations with the AER radiative transfer models, *J. Geophys. Res.*, 113, D13103, <https://doi.org/10.1029/2008JD009944>, 2008.
- Ilyina, T., Wolf-Gladrow, D., Munhoven, G., and Heinze, C.: Assessing the potential of calcium-based artificial ocean alkalization to mitigate rising atmospheric CO₂ and ocean acidification, *Geophys. Res. Lett.*, 40, 5909–5914, <https://doi.org/10.1002/2013GL057981>, 2013.
- 1110 Jacob, D. J.: Chemistry of OH in remote clouds and its role in the production of formic acid and peroxymonosulfate, *J. Geophys. Res.*, 91, 9807–9826, <https://doi.org/10.1029/JD091iD09p09807>, 1986.
- Jacobson, M.Z. and Seinfeld, J.H.: Evolution of nanoparticle size and mixing state near the point of emission, *Atmos. Environ.*, 38, 1839–1850, <https://doi.org/10.1016/j.atmosenv.2004.01.014>, 2004.
- 1115 Jöckel, P., von Kuhlmann, R., Lawrence, M.G., Steil, B., Brenninkmeijer, C.A.M., Crutzen, P.J., Rasch, P.J. and Eaton, B.: On a fundamental problem in implementing flux-form advection schemes for tracer transport in 3-dimensional general circulation and chemistry transport models, *Q. J. R. Meteorol. Soc.*, 127, 1035–1052, <https://doi.org/10.1002/qj.49712757318>, 2001.
- Jungclaus, J. H., Keenlyside, N., Botzet, M., Haak, H., Luo, J.-J., Latif, M., Marotzke, J., Mikolajewicz, U., and Roeckner, E.: Ocean circulation and tropical variability in the coupled model ECHAM5/MPI-OM, *J. Climate*, 19, 3952–3972, <https://doi.org/10.1175/JCLI3827.1>, 2006.
- 1120 Jungclaus, J. H., Fischer, N., Haak, H., Lohmann, K., Marotzke, J., Matei, D., Mikolajewicz, U., Notz, D., and von Storch, J. S.: Characteristics of the ocean simulations in MPIOM, the ocean component of the MPI-Earth system model, *J. Adv. Model. Earth Syst.*, 5, 422–446, doi:10.1002/jame.20023, 2013.
- 1125 Karagodin-Doyennel, A., Rozanov, E., Kuchar, A., Ball, W., Arsenovic, P., Remsberg, E., Jöckel, P., Kunze, M., Plummer, D. A., Stenke, A., Marsh, D., Kinnison, D., and Peter, T.: The response of mesospheric H₂O and CO to solar irradiance variability in models and observations, *Atmos. Chem. Phys.*, 21, 201–216, <https://doi.org/10.5194/acp-21-201-2021>, 2021.
- Kasten, F.: Falling speed of aerosol particles, *J. App. Met. Clim.*, 7, 944–947, [https://doi.org/10.1175/1520-0450\(1968\)007](https://doi.org/10.1175/1520-0450(1968)007), 1968.
- 1130 Keeble, J., Hassler, B., Banerjee, A., Checa-Garcia, R., Chiodo, G., Davis, S., Eyring, V., Griffiths, P. T., Morgenstern, O., Nowack, P., Zeng, G., Zhang, J., Bodeker, G., Cugnet, D., Danabasoglu, G., Deushi, M., Horowitz, L. W., Li, L., Michou, M., Mills, M. J., Nabat, P., Park, S., and Wu, T.: Evaluating stratospheric ozone and water vapor changes in CMIP6 models from 1850–2100, *Atmos. Chem. Phys. Discuss.* [preprint], <https://doi.org/10.5194/acp-2019-1202>, in review, 2020.
- 1135 Kerkweg, A., Buchholz, J., Ganzeveld, L., Pozzer, A., Tost, H., and Jöckel, P.: Technical Note: An implementation of the dry removal processes DRY DEPosition and SEDimentation in the Modular Earth Submodel System (MESSy), *Atmos. Chem. Phys.*, 6, 4617–4632, <https://doi.org/10.5194/acp-6-4617-2006>, 2006.



- Kidston, J., Scaife, A., Hardiman, S., Mitchel, D.M., Butchart, N., Baldwin, M.P., and Gray, L.J.: Stratospheric influence on tropospheric jet streams, storm tracks and surface weather, *Nature Geosci.*, 8, 433–440, <https://doi.org/10.1038/ngeo2424>, 2015.
- 1140 Kinne, S., O'Donnel, D., Stier, P., Kloster, S., Zhang, K., Schmidt, H., Rast, S., Giorgetta, M., Eck, T. F., and Stevens, B.: MAC-v1: A new global aerosol climatology for climate studies, *J. Adv. Model. Earth Syst.*, 5, 704–740, <https://doi.org/10.1002/jame.20035>, 2013.
- Koenig, Th.K., Baidara, S., Campuzano-Josta, P., Cuevas, C.A., Dixa, B., Fernandez, R.P., Guoa, H., Halle, S.R., Kinnison, D., Naulta, B.A., Ullmanne, K., Jimenez, L.L., Saiz-Lopez, A. and Volkamer, R.: Quantitative detection of iodine in the stratosphere, *Proc. Natl. Acad. Sci. USA*, 117, 1860–1866, <https://doi.org/10.1073/pnas.1916828117>, 2020.
- 1145 Koepke P., Hess, M., Schult, I., and Shettle, E.P.: "Global aerosol dataset", Report N 243, Max-Planck-Institut für Meteorologie, Hamburg, 44 pp., 1997.
- Kulmala, M. and Laaksonen, A.: Binary nucleation of water–sulfuric acid system: Comparison of classical theories with different H₂SO₄ saturation vapor pressures, *J. Chem. Phys.*, 93, 696–701, <https://doi.org/10.1063/1.459519>, 1990.
- 1150 Lana, A., Bell, T. G., Simó, R., Vallina, S. M., Ballabrera-Poy, J., Kettle, A. J., Dachs, J., Bopp, L., Saltzman, E. S., Stefels, J., Johnson, J. E., and Liss, P. S.: An updated climatology of surface dimethylsulfide concentrations and emission fluxes in the global ocean, *Global Biogeochem. Cycles*, 25, GB1004, <https://doi.org/10.1029/2010GB003850>, 2011.
- Lin, S. J., and Rood, R. B.: Multidimensional flux-form semiLagrangian transport schemes, *Month. Weather Rev.*, 124, 2046–2070, [https://doi.org/10.1175/1520-0493\(1996\)124](https://doi.org/10.1175/1520-0493(1996)124), 1996.
- 1155 Long, C. S., Fujiwara, M., Davis, S., Mitchell, D. M., and Wright, C. J.: Climatology and interannual variability of dynamic variables in multiple reanalyses evaluated by the SPARC Reanalysis Intercomparison Project (S-RIP), *Atmos. Chem. Phys.*, 17, 14593–14629, <https://doi.org/10.5194/acp-17-14593-2017>, 2017.
- Lott, F.: Alleviation of stationary biases in a GCM through a mountain drag parameterization scheme and a simple representation of mountain lift forces, *Mon. Weather Rev.*, 127, 788–801, [https://doi.org/10.1175/1520-0493\(1999\)127](https://doi.org/10.1175/1520-0493(1999)127), 1999.
- 1160 Manabe S. and Bryan, K.: Climate calculations with a combined ocean–atmosphere model, *J. Atmos. Sci.*, 26, 786–789, [https://doi.org/10.1175/1520-0469\(1969\)026](https://doi.org/10.1175/1520-0469(1969)026), 1969.
- Manney, G. L., Livesey, N. J., Santee, M. L., Froidevaux, L., Lambert, A., Lawrence, Z. D., Millán, L.F., Neu, J.L., Read, W.G., Schwartz, M.J., and Fuller, R.A.: Record-low Arctic stratospheric ozone in 2020: MLS observations of chemical processes and comparisons with previous extreme winters, *Geophys. Res. Lett.*, 47, e2020GL089063, <https://doi.org/10.1029/2020GL089063>, 2020.
- 1165 Manzini, E., Giorgetta, M.A., Esch, M., Kornbluh, L., and Roeckner, E.: The influence of the sea surface temperatures on the northern winter stratosphere: Ensemble Simulations with the MAECHAM5 model, *J. Climate*, 19, 3863–3881, <https://doi.org/10.1175/JCLI3826.1>, 2006.
- Marsh, D. R., Janches, D., Feng, W., & Plane, J. M. C.: A global model of meteoric sodium, *J. Geophys. Res. Atmos.*, 118, 11442–11452, <https://doi.org/10.1002/jgrd.50870>, 2013.
- 1170



- Matthes, K., Funke, B., Andersson, M. E., Barnard, L., Beer, J., Charbonneau, P., Clilverd, M. A., Dudok de Wit, T., Haberreiter, M., Hendry, A., Jackman, C. H., Kretzschmar, M., Kruschke, T., Kunze, M., Langematz, U., Marsh, D. R., Maycock, A. C., Misios, S., Rodger, C. J., Scaife, A. A., Seppälä, A., Shangguan, M., Sinnhuber, M., Tourpali, K., Usoskin, I., van de Kamp, M., Verronen, P. T., and Versick, S.: Solar forcing for CMIP6 (v3.2), *Geosci. Model Dev.*, 10, 2247–2302, 1175 <https://doi.org/10.5194/gmd-10-2247-2017>, 2017.
- Matthes, K., Biastoch, A., Wahl, S., Harlaß, J., Martin, T., Brücher, T., Drews, A., Ehlert, D., Getzlaff, K., Krüger, F., Rath, W., Scheinert, M., Schwarzkopf, F. U., Bayr, T., Schmidt, H., and Park, W.: The Flexible Ocean and Climate Infrastructure version 1 (FOCI1): mean state and variability, *Geosci. Model Dev.*, 13, 2533–2568, <https://doi.org/10.5194/gmd-13-2533-2020>, 2020.
- 1180 Mauritsen, T., Bader, J., Becker, T., Behrens, J., Bittner, M., Brokopf, R., Brovkin, V., Claussen, M., Crueger, T., Esch, M., Fast, I., Fiedler, S., Fläschner, D., Gayler, V., Giorgetta, M., Goll, D. S., Haak, H., Hagemann, S., Hedemann, C., Hohenegger, C., Ilyina, T., Jahns, Th., Jimenez-de-la-Cuesta, D., Jungclaus, J., Kleinen, Th., Kloster, S., Kracher, D., Kinne, S., Kleberg, D., Lasslop, G., Kornbluh, L., Marotzke, J., Matei, D., Meraner, K., Mikolajewicz, U., Modali, K., Möbis, B., Müller, W. A., Nabel, J. E. M. S., Nam, C. C. W., Notz, D., Nyawira, S.-S., Paulsen, H., Peters, K., Pincus, R., Pohlmann, H., Pongratz, J., 1185 Popp, M., Raddatz, Th. J., Rast, S., Redler, R., Reick, Ch. H., Rohrschneider, T., Schemann, V., Schmidt, H., Schnur, R., Schulzweida, U., Six, K. D., Stein, L., Stemmler, I., Stevens, B., von Storch, J.-S., Tian, F., Voigt, A., Vrese, Ph., Wieners, K.-H., Wilkenskjaeld, S., Winkler, A., Roeckner, E.: Developments in the MPI-M Earth System Model version 1.2 (MPI-ESM1.2) and its response to increasing CO₂, *J. Adv. Model. Earth Syst.*, 11, 998–1038, <https://doi.org/10.1029/2018MS001400>, 2019.
- 1190 McConnell, J. C. and Jin, J. J.: Stratospheric ozone chemistry, *Atmosphere-Ocean*, 46, 69–92, <https://doi.org/10.3137/ao.460104>, 2008.
- McPeters, R. D., Bhartia, P. K., Haffner, D., Labow, G. J., and Flynn, L.: The version 8.6 SBUV ozone data record: An overview, *J. Geophys. Res. Atmos.*, 118, 8032–8039, <https://doi.org/10.1002/jgrd.50597>, 2013.
- Meinshausen, N., Hauser, A., Mooij, J. M., Peters, J., Versteeg, Ph., and Bühlmann, P.: Methods for causal inference from 1195 gene perturbation experiments and validation, *Proc Natl Acad Sci USA*, 27, 7361–7368, <https://doi.org/10.1073/pnas.1510493113>, 2016.
- Meinshausen, M., Vogel, E., Nauels, A., Lorbacher, K., Meinshausen, N., Etheridge, D. M., Fraser, P. J., Montzka, S. A., Rayner, P. J., Trudinger, C. M., Krummel, P. B., Beyerle, U., Canadell, J. G., Daniel, J. S., Enting, I. G., Law, R. M., Lunder, C. R., O'Doherty, S., Prinn, R. G., Reimann, S., Rubino, M., Velders, G. J. M., Vollmer, M. K., Wang, R. H. J., and Weiss, 1200 R.: Historical greenhouse gas concentrations for climate modelling (CMIP6), *Geosci. Model Dev.*, 10, 2057–2116, <https://doi.org/10.5194/gmd-10-2057-2017>.
- Miller, M.J., Palmer, T.N., and Swinbank, R.: Parametrization and influence of subgridscale orography in general circulation and numerical weather prediction models, *Meteorol. Atmos. Phys.*, 40, 84–109, <https://doi.org/10.1007/BF01027469>, 1989.



- Mironova, I. A., Aplin, K. L., Arnold, F., Bazilevskaya, G. A., Harrison, R. G., Giles, R., Krivolutsky, A. A., Nicoll, K.A.,
1205 Rozanov, E.V., Turunen, E., and Usoskin, I. G.: Energetic Particle Influence on the Earth's Atmosphere, *Space Sci. Rev.*, 194,
1–96, <https://doi.org/10.1007/s11214-015-0185-4>, 2015.
- Mote, P. W., Dunkerton, T. J., McIntyre, M. E., Ray, E. A., Haynes, P. H., and Russell, J. M.: Vertical velocity, vertical
diffusion, and dilution by midlatitude air in the tropical lower stratosphere, *J. Geophys. Res.*, 103, 8651– 8666,
<https://doi.org/10.1029/98JD00203>, 1998.
- 1210 Möbis, B., and Stevens, B.: Factors controlling the position of the Intertropical Convergence Zone on an aquaplanet, *J. Adv.
Model. Earth Syst.*, 4, M00A04, <https://doi.org/10.1029/2012MS000199>, 2012.
- Muthers, S., Anet, J. G., Stenke, A., Raible, C. C., Rozanov, E., Brönnimann, S., Peter, T., Arfeuille, F. X., Shapiro, A. I.,
Beer, J., Steinhilber, F., Brugnara, Y., and Schmutz, W.: The coupled atmosphere–chemistry–ocean model SOCOL-MPIOM,
Geosci. Model Dev., 7, 2157–2179, <https://doi.org/10.5194/gmd-7-2157-2014>, 2014.
- 1215 Nakajima, H., Wohltmann, I., Wegner, T., Takeda, M., Pitts, M. C., Poole, L. R., Lehmann, R., Santee, M. L., and Rex, M.:
Polar stratospheric cloud evolution and chlorine activation measured by CALIPSO and MLS, and modeled by ATLAS, *Atmos.
Chem. Phys.*, 16, 3311–3325, <https://doi.org/10.5194/acp-16-3311-2016>, 2016.
- Niemeier, U. and Schmidt, H.: Changing transport processes in the stratosphere by radiative heating of sulfate aerosols, *Atmos.
Chem. Phys.*, 17, 14871–14886, <https://doi.org/10.5194/acp-17-14871-2017>, 2017.
- 1220 Nightingale, P. D., Malin, G., Law, C. S., Watson, A. J., Liss, P. S., Liddicoat, M. I., Boutin, J., and Upstill-Goddard, R. C.:
In situ evaluation of air-sea gas exchange parameterizations using novel conservative and volatile tracers, *Global Biogeochem.
Cycles*, 1, 373– 387, <https://doi.org/10.1029/1999GB900091>, 2000.
- Nordeng, T. E.: Extended versions of the convective parameterization scheme at ECMWF and their impact on the mean and
transient activity of the model in the tropics, ECMWF Research Department Tech. Memo., 41 pp., 1994.
- 1225 Nowack, P.J., Abraham, N.L., Maycock, A.C., Braesicke, P., Gregory, J.M., Joshi, M.M., Osprey, A., and Pyle, J.A.: A large
ozone-circulation feedback and its implications for global warming assessments, *Nat. Clim. Chang.*, 5, 41-45,
<https://doi.org/10.1038/nclimate2451>, 2015.
- Oehrlein, J., Chiodo, G., and Polvani, L.M.: The effect of interactive ozone chemistry on weak and strong stratospheric polar
vortex events, *Atmos. Chem. Phys.*, <https://doi.org/10.5194/acp-20-10531-2020>, 2020.
- 1230 Orbe, C., Wargan, K., Pawson, S., & Oman, L. D.: Mechanisms linked to recent ozone decreases in the Northern Hemisphere
lower stratosphere. *Journal of Geophysical Research: Atmospheres*, 125, e2019JD031631,
<https://doi.org/10.1029/2019JD031631>, 2020.
- Ozolin, Y.: Modelling of diurnal variations of gas species in the atmosphere and diurnal averaging in photochemical models,
Izv. Akad. Nauk. Phys. Atmos. Ocean., 28, 135–143, 1992.
- 1235 Palmer, T.N., Shutts, G.J. and Swinbank, R.: Alleviation of a systematic westerly bias in general circulation and numerical
weather prediction models through an orographic gravity wave drag parametrization, *Q. J. R. Meteorol. Soc.*, 112, 1001-1039,
<https://doi.org/10.1002/qj.49711247406>, 1986.



- Paulsen, J., Sekelja, M., Oldenburg, A.R., Barateau, A., Briand, N., Delbarre, E., Shah, A., Sørensen, A. L., Vigouroux, C., Buendia, B., and Collas, Ph.: Chrom3D: three-dimensional genome modeling from Hi-C and nuclear lamin-genome contacts, *Genome Biol.*, 18, 1-15, <https://doi.org/10.1186/s13059-016-1146-2>, 2017.
- Pedersen, C. A., Roeckner, E., Lüthje, M., and Winther, J.-G.: A new sea ice albedo scheme including melt ponds for ECHAM5 general circulation model, *J. Geophys. Res.*, 114, D08101, <https://doi.org/10.1029/2008JD010440>, 2009.
- Petropavlovskikh, I., Godin-Beekmann, S., Hubert, D., Damadeo, R., Hassler, B., and Sofieva, V.: SPARC/IO3C/GAW report on Long-term Ozone Trends and Uncertainties in the Stratosphere, SPARC/IO3C/GAW, SPARC Report No. 9, WCRP-17/2018, GAW Report No. 241, <https://doi.org/10.17874/f899e57a20b>, 2019.
- Plumb, R. A.: Stratospheric transport, *J. Meteorol. Soc. Jpn.*, 80, 793–809, <https://doi.org/10.2151/jmsj.80.793>, 2002.
- Plumb, R. A.: Tracer interrelationships in the stratosphere, *Rev. Geophys.*, 45, RG4005, <https://doi.org/10.1029/2005RG000179>, 2007.
- Polvani, L. M., Wang, L., Abalos, M., Butchart, N., Chipperfield, M. P., Dameris, M., Deushi, M., Dhomse, S. S., Jöckel, P., Kinnison, D., Michou, M., Morgenstern, O., Oman, L. D., Plummer, D. A., and Stone, K. A.: Large impacts, past and future, of ozone-depleting substances on Brewer-Dobson circulation trends: A multimodel assessment. *J. Geophys. Res. Atmos.*, 124, 6669–6680, <https://doi.org/10.1029/2018JD029516>, 2019.
- Pöschl, U., von Kuhlmann, R., Poisson, N., and Crutzen, P. J.: Development and Intercomparison of Condensed Isoprene Oxidation Mechanisms for Global Atmospheric Modeling, *J. Atmos. Chem.*, 37, 29–52, <https://doi.org/10.1023/A:1006391009798>, 2000.
- Previdi, M. and Polvani, L.M.: Climate system response to stratospheric ozone depletion and recovery, *Q. J. Roy. Met. Society*, 140, 2401–2419, <https://doi.org/10.1002/qj.2330>, 2014.
- Price, C. and Rind, D.: A simple lightning parameterization for calculating global lightning distributions, *J. Geophys. Res.*, 97, 9919–9933, <https://doi.org/10.1029/92JD00719>, 1992.
- Pruppacher, H. R., and Klett, J. D.: *Microphysics of Clouds and Precipitation*, 2nd Edition, Kluwer Academic, Dordrecht, 954 p., 1997.
- Revell, L. E., Tummon, F., Stenke, A., Sukhodolov, T., Coulon, A., Rozanov, E., Garny, H., Grewe, V., and Peter, T.: Drivers of the tropospheric ozone budget throughout the 21st century under the medium-high climate scenario RCP 6.0, *Atmos. Chem. Phys.*, 15, 5887–5902, <https://doi.org/10.5194/acp-15-5887-2015>, 2015.
- Revell, L. E., Stenke, A., Rozanov, E., Ball, W., Lossow, S., Peter, T.: The role of methane in projections of 21st century stratospheric water vapour, *Atmos. Chem. Phys.*, 16, 13067–13080, <https://doi.org/10.5194/acp-16-13067-2016>, 2016.
- Revell, L. E., Stenke, A., Tummon, F., Feinberg, A., Rozanov, E., Peter, T., Abraham, N. L., Akiyoshi, H., Archibald, A. T., Butchart, N., Deushi, M., Jöckel, P., Kinnison, D., Michou, M., Morgenstern, O., O'Connor, F. M., Oman, L. D., Pitari, G., Plummer, D. A., Schofield, R., Stone, K., Tilmes, S., Visionsi, D., Yamashita, Y., and Zeng, G.: Tropospheric ozone in CCM1



- models and Gaussian process emulation to understand biases in the SOCOLv3 chemistry–climate model, *Atmos. Chem. Phys.*, 18, 16155–16172, <https://doi.org/10.5194/acp-18-16155-2018>, 2018.
- Rieder, H., Chiodo, G., Fritzer, J., Wienerroither, C., and Polyani, L.: Is interactive ozone chemistry important to represent polar cap stratospheric temperature variability in Earth-System Models?, *Environ. Res. Lett.*, 14, 044026, <https://doi.org/10.1088/1748-9326/ab07ff>, 2019.
- 1275
- Rozanov, E. V., Zubov, V. A., Schlesinger, M. E., Yang, F., and Andronova, N. G.: The UIUC three-dimensional stratospheric chemical transport model: Description and evaluation of the simulated source gases and ozone, *J. Geophys. Res.*, 104, 11755–11781, <https://doi:10.1029/1999JD900138>, 1999.
- Rozanov, E. V., Schlesinger, M. E., and Zubov, V. A.: The University of Illinois, Urbana-Champaign three-dimensional stratosphere-troposphere general circulation model with interactive ozone photochemistry: Fifteen-year control run climatology, *J. Geophys. Res.*, 106, 27233–27254, <https://doi.org/10.1029/2000JD000058>, 2001.
- 1280
- Rozanov, E., Calisto, M., Egorova, T., Peter, T., and Schmutz, W.: Influence of the Precipitating Energetic Particles on Atmospheric Chemistry and Climate, *Surv. Geophys.*, 33, 483–501, <https://doi.org/10.1007/s10712-012-9192-0>, 2012.
- Schmidt, T., Alexander, P., and de la Torre, A.: Stratospheric gravity wave momentum flux from radio occultations, *J. Geophys. Res. Atmos.*, 121, 4443–4467, <https://doi.org/10.1002/2015JD024135>, 2013.
- 1285
- Schmidt, H., Brasseur, G. P., Charron, M., Manzini, E., Giorgetta, M. A., Diehl, T., Fomichev, V.I., Kinnison, D., Marsh, D., and Walters, S.: The HAMMONIA Chemistry Climate Model: Sensitivity of the Mesopause Region to the 11-Year Solar Cycle and CO₂ Doubling. *J. Clim.*, 19, 3903–3931, <https://doi.org/10.1175/JCLI3829.1>, 2006.
- Schraner, M., Rozanov, E., Schnadt Poberaj, C., Kenzelmann, P., Fischer, A. M., Zubov, V., Luo, B. P., Hoyle, C. R., Egorova, T., Fueglistaler, S., Brönnimann, S., Schmutz, W., and Peter, T.: Technical Note: Chemistry-climate model SOCOL: version 1290 2.0 with improved transport and chemistry/microphysics schemes, *Atmos. Chem. Phys.*, 8, 5957–5974, <https://doi.org/10.5194/acp-8-5957-2008>, 2008.
- Sheng, J.-X., Weisenstein, D. K., Luo, B.-P., Rozanov, E., Stenke, A., Anet, J., Bingemer, H., and Peter, T.: Global atmospheric sulfur budget under volcanically quiescent conditions: Aerosol-chemistry-climate model predictions and validation, *J. Geophys. Res. Atmos.*, 120, 256–276, <https://doi.org/10.1002/2014JD021985>, 2015.
- 1295
- Simmons, A.J., Soci, C., Nicolas, J., Bell, B., Berrisford, P., Dragani, R., Flemming, J., Haimberger, L., Healey, S.B., Hersbach, H., Horányi, A., Inness, A., Muñoz-Sabater, J., Radu, R. and Schep-ers, D.: Global stratospheric temperature bias and other stratospheric aspects of ERA5 and ERA5.1, Technical Memorandum 859, ECMWF, Reading, UK, <https://doi.org/10.21957/rcxqfmg0>, 2020.
- 1300
- SPARC: The SPARC Data Initiative: Assessment of stratospheric trace gas and aerosol climatologies from satellite limb sounders. By M. I. Hegglin and S. Tegtmeier (eds.), SPARC Report No. 8, WCRP-5/2017, available at www.sparc-climate.org/publications/sparc-reports/sparc-report-no8, 2017.



- Stenke, A., Schraner, M., Rozanov, E., Egorova, T., Luo, B., and Peter, T.: The SOCOL version 3.0 chemistry–climate model: description, evaluation, and implications from an advanced transport algorithm, *Geosci. Model Dev.*, 6, 1407–1427, 1305 <https://doi.org/10.5194/gmd-6-1407-2013>, 2013.
- Stevens, B., Giorgetta, M., Esch, M., Mauritsen, T., Crueger, T., Rast, S., Salzmann, M., Schmidt, H., Bader, J., Block, K., Brokopf, R., Fast, I., Kinne, S., Kornbluh, L., Lohmann, U., Pincus, R., Reichler, Th., and Roeckner, E.: Atmospheric component of the MPI-M Earth System Model: ECHAM6, *J. Adv. Model. Earth Syst.*, 5, 1942–2466, <https://doi.org/10.1002/jame.20015>, 2013.
- 1310 Stevens, B., Fiedler, S., Kinne, S., Peters, K., Rast, S., Müsse, J., Smith, S. J., and Mauritsen, T.: MACv2-SP: a parameterization of anthropogenic aerosol optical properties and an associated Twomey effect for use in CMIP6, *Geosci. Model Dev.*, 10, 433–452, <https://doi.org/10.5194/gmd-10-433-2017>, 2017.
- Strahan, S.E. MIDDLE ATMOSPHERE | Transport Circulation, in *Encyclopedia of Atmospheric Sciences (Second Edition)*, Editors: North, G. R., Pyle, J., and Zhang, F., 41–49, 2015.
- 1315 Stone, K. A., Solomon, S., and Kinnison, D. E.: On the identification of ozone recovery, *Geophys. Res. Lett.*, 45, 5158–5165, <https://doi.org/10.1029/2018GL077955>, 2018.
- Stott, P. A. and Harwood, R. S.: An implicit time-stepping scheme for chemical species in a global atmospheric circulation model, *Ann. Geophys.*, 11, 377–388, 1993.
- Sukhodolov T. et al.: Atmosphere-Ocean-Aerosol-Chemistry-Climate Model SOCOLv4.0 code (Version 1.0). Zenodo. 1320 <http://doi.org/10.5281/zenodo.4570622>, 2021.
- Sukhodolov, T., Rozanov, E., Shapiro, A. I., Anet, J., Cagnazzo, C., Peter, T., and Schmutz, W.: Evaluation of the ECHAM family radiation codes performance in the representation of the solar signal, *Geosci. Model Dev.*, 7, 2859–2866, <https://doi.org/10.5194/gmd-7-2859-2014>, 2014.
- Sukhodolov, T., Rozanov, E., Ball, W.T., Bais, A., Tourpali, K., Shapiro, A.I., Telford, P., Smyshlyaev, S., Fomin, B., Sander, 1325 R., Bossay, B., Bekki, S., Marchand, M., Chipperfield, M.P., Dhomse, S., Haigh, J.D., Peter, Th., and Schmutz, W.: Evaluation of simulated photolysis rates and their response to solar irradiance variability, *J. Geophys. Res. Atmos.*, 121, 6066–6084, <https://doi.org/10.1002/2015JD024277>, 2016.
- Sukhodolov, T., Sheng, J.-X., Feinberg, A., Luo, B.-P., Peter, T., Revell, L., Stenke, A., Weisenstein, D. K., and Rozanov, E.: Stratospheric aerosol evolution after Pinatubo simulated with a coupled size-resolved aerosol–chemistry–climate model, 1330 SOCOL-AERv1.0, *Geosci. Model Dev.*, 11, 2633–2647, <https://doi.org/10.5194/gmd-11-2633-2018>, 2018.
- Sundqvist, H., Berge, E., and Kristjánsson, J. E.: Condensation and cloud parameterization studies with a mesoscale numerical weather prediction model, *Mon. Weather Rev.*, 117, 1641–1657, [https://doi.org/10.1175/1520-0493\(1989\)117<1641:CACPSW>2.0.CO;2](https://doi.org/10.1175/1520-0493(1989)117<1641:CACPSW>2.0.CO;2), 1989.
- Tabazadeh, A., Toon, O. B., Clegg, S. L., and Hamill, P.: A new parameterization of H₂SO₄/H₂O aerosol composition: 1335 Atmospheric implications, *Geophys. Res. Lett.*, 24, 1931–1934, <https://doi.org/10.1029/97GL01879>, 1997.



- Tian, F., Storch, J.S.v. & Hertwig, E.: Impact of SST diurnal cycle on ENSO asymmetry. *Clim Dyn* 52, 2399–2411. <https://doi.org/10.1007/s00382-018-4271-7>, 2019.
- Tiedtke, M. : A Comprehensive Mass Flux Scheme for Cumulus Parameterization in Large Scale Models, *Mon. Wea. Rev.*, 117, 1779 -1800, [https://doi.org/10.1175/1520-0493\(1989\)117<1779:ACMFSF>2.0.CO;2](https://doi.org/10.1175/1520-0493(1989)117<1779:ACMFSF>2.0.CO;2), 1989.
- 1340 Tilmes, S., Garcia, R. R., Kinnison, D. E., Gettelman, A., and Rasch, P. J.: Impact of geoengineered aerosols on the troposphere and stratosphere, *J. Geophys. Res.*, 114, D12305, <https://doi.org/10.1029/2008JD011420>, 2009.
- Timmreck, C., Lorenz, S. J., Crowley, T. J., Kinne, S., Raddatz, T. J., Thomas, M. A., and Jungclaus, J. H.: Limited temperature response to the very large AD 1258 volcanic eruption, *Geophys. Res. Lett.*, 36, L21708, <https://doi.org/10.1029/2009GL040083>, 2009.
- 1345 Timmreck, C., Mann, G. W., Aquila, V., Hommel, R., Lee, L. A., Schmidt, A., Brühl, C., Carn, S., Chin, M., Dhomse, S. S., Diehl, T., English, J. M., Mills, M. J., Neely, R., Sheng, J., Toohey, M., and Weisenstein, D.: The Interactive Stratospheric Aerosol Model Intercomparison Project (ISA-MIP): motivation and experimental design, *Geosci. Model Dev.*, 11, 2581–2608, <https://doi.org/10.5194/gmd-11-2581-2018>, 2018.
- Thomason, L. W., Ernest, N., Millán, L., Rieger, L., Bourassa, A., Vernier, J.-P., Manney, G., Luo, B., Arfeuille, F., and Peter, T.: A global space-based stratospheric aerosol climatology: 1979–2016, *Earth Syst. Sci. Data*, 10, 469–492, <https://doi.org/10.5194/essd-10-469-2018>, 2018.
- 1350 Tost, H., Jöckel, P., and Lelieveld, J.: Influence of different convection parameterisations in a GCM, *Atmos. Chem. Phys.*, 6, 5475–5493, <https://doi.org/10.5194/acp-6-5475-2006>, 2006.
- Tost, H., Jöckel, P., and Lelieveld, J.: Lightning and convection parameterisations – uncertainties in global modelling, *Atmos. Chem. Phys.*, 7, 4553–4568, <https://doi.org/10.5194/acp-7-4553-2007>, 2007.
- 1355 Twomey, S.: The influence of pollution on the short wave albedo of clouds, *J. Atmos. Sci.*, 34, 1149–1152, [https://doi.org/10.1175/1520-0469\(1977\)034](https://doi.org/10.1175/1520-0469(1977)034), 1977.
- Van der A, R. J., Allaart, M. A. F., and Eskes, H. J. Extended and refined multi sensor reanalysis of total ozone for the period 1970-2012, *Atmos. Meas. Tech.*, 8, 3021-3035, <https://doi.org/10.5194/amt-8-3021-2015>, 2015.
- 1360 Vattioni, S., D. Weisenstein, D. Keith, A. Feinberg, T. Peter, A. Stenke, Exploring accumulation-mode H₂SO₄ versus SO₂ stratospheric sulfate geoengineering in a sectional aerosol-chemistry-climate model, *Atmos. Chem. Phys.*, 19, Iss. 7, DOI: 10.5194/acp-19-4877-2019, 2019.
- Vehkamäki, H., Kulmala, M., Napari, I., Lehtinen, K. E. J., Timmreck, C., Noppel, M., and Laaksonen, A.: An improved parameterization for sulfuric acid–water nucleation rates for tropospheric and stratospheric conditions, *J. Geophys. Res.*, 107, 4622, <https://doi.org/10.1029/2002JD002184>, 2002.
- 1365 Velders, G. J. M., Andersen, S. O., Daniel, J. S., Fahey, D. W., and McFarland, M.: The importance of the Montreal Protocol in protecting climate, *Proc. Natl. Acad. Sci. USA*, 104, 4814-4819, <https://doi.org/10.1073/pnas.0610328104>, 2007.
- Voigt, C., Schlager, H., Ziereis, H., Kärcher, B., Luo, B. P., Schiller, C., Krämer, M., Popp, P. J., Irie, H., and Kondo, Y.: Nitric acid in cirrus clouds, *Geophys. Res. Lett.*, 33, L05803, <https://doi.org/10.1029/2005GL025159>, 2006.



- 1370 Weisenstein, D. K., Yue, G. K., Ko, M. K. W., Sze, N.-D., Rodriguez, J. M., and Scott, C. J.: A two-dimensional model of sulfur species and aerosols, *J. Geophys. Res.*, 102, 13019–13035, <https://doi.org/10.1029/97JD00901>, 1997.
- Wesely, M.L.: Parameterization of surface resistances to gaseous dry deposition in regional-scale numerical models, *Atm. Environ.* (1967), 23, 1293-1304, [https://doi.org/10.1016/0004-6981\(89\)90153-4](https://doi.org/10.1016/0004-6981(89)90153-4), 1989.
- Witze A.: Rare ozone hole opens over Arctic - it's big, *Nature*, 580, 18-19, <https://doi.org/10.1038/d41586-020-00904-w>,
1375 2020.
- Walcek, C. J.: Minor flux adjustment near mixing ratio extremes for simplified yet highly accurate monotonic calculation of tracer advection, *J. Geophys. Res.*, 105, 9335– 9348, <https://doi.org/10.1029/1999JD901142>, 2000.
- Wolff, J.-O., Maier-Reimer, E., and Legutke, S.: The Hamburg Ocean Primitive Equation Model. World Data Center for Climate (WDCC) at DKRZ, https://doi.org/10.2312/WDCC/DKRZ_Report_No13, 1997.