

Earth System Model Aerosol-Cloud Diagnostics Package (ESMAC Diags) Version 1: Assessing E3SM Aerosol Predictions Using Aircraft, Ship, and Surface Measurements

Shuaiqi Tang¹, Jerome D. Fast¹, Kai Zhang¹, Joseph C. Hardin², Adam C. Varble¹, John E. Shilling¹, Fan Mei¹, Maria A. Zawadowicz³ and Po-Lun Ma¹

¹Pacific Northwest National Laboratory, Richland, WA, USA

²ClimateAI Inc, San Francisco, CA, USA

³Brookhaven National Laboratory, Upton, NY, USA

Correspondence to: Shuaiqi Tang (shuaiqi.tang@pnnl.gov)

Abstract. An Earth System Model (ESM) aerosol-cloud diagnostics package is developed to facilitate the routine evaluation of aerosols, clouds and aerosol-cloud interactions simulated by the Department of Energy's (DOE) Energy Exascale Earth System Model (E3SM). The first version focuses on comparing simulated aerosol properties with aircraft, ship, and surface measurements, most of them are measured in-situ. The diagnostics currently covers six field campaigns in four geographical regions: Eastern North Atlantic (ENA), Central U.S. (CUS), Northeastern Pacific (NEP) and Southern Ocean (SO). These regions produce frequent liquid or mixed-phase clouds with extensive measurements available from the Atmospheric Radiation Measurement (ARM) program and other agencies. Various types of diagnostics and metrics are performed for aerosol number, size distribution, chemical composition, CCN concentration and various meteorological quantities to assess how well E3SM represents observed aerosol properties across spatial scales. Overall, E3SM qualitatively reproduces the observed aerosol number concentration, size distribution and chemical composition reasonably well, but overestimates Aitken mode and underestimates accumulation mode aerosols over the CUS and ENA regions, suggesting that processes related to particle growth or coagulation might be too weak in the model. The current version of E3SM struggles to reproduce new particle formation events frequently observed over both the CUS and ENA regions, indicating missing processes in current parameterizations. The diagnostics package is coded and organized in a way that can be extended to other field campaign datasets and adapted to higher-resolution model simulations.

1. Introduction

Aerosol number, mass, size, composition, and mixing state affect how aerosol populations scatter and absorb solar radiation and influence cloud albedo, amount, lifetime, and precipitation (Twomey, 1977; Albrecht, 1989) by acting as cloud condensation nuclei (CCN) (e.g., Petters and Kreidenweis, 2007). However, there are still knowledge and measurement gaps on the physical and chemical mechanisms regulating the sources, sinks, gas-to-particle partitioning (e.g., secondary formation processes), and spatiotemporal distribution of aerosol populations. Consequently, the representation of the aerosol lifecycle and the interaction of aerosol populations with clouds and radiation in Earth system models (ESMs) still suffer from large uncertainties (Seinfeld et al., 2016; Carslaw et al., 2018), which impacts the ability of ESMs to predict the evolution of the climate system (IPCC, 2013).

To facilitate model evaluation and document the performance of parameterizations in ESMs, many modeling centers have developed standardized diagnostics packages. Some examples focus on meteorological metrics include the U.S. National Center of Atmospheric Research (NCAR) Atmospheric Model Working Group (AMWG) diagnostics package (AMWG, 2021), the U.S. Department of Energy (DOE) Energy Exascale Earth System Model (E3SM, Golaz et al., 2019) diagnostics (E3SM, 2021), the European Union (EU) Earth System Model Evaluation Tool (ESMValTool, Eyring et al., 2016), and the Program for Climate Model Diagnosis and Intercomparison (PCMDI) Metric Package (PMP, Gleckler et al., 2016). Some recent efforts focus on process-oriented diagnostics (POD) that are designed to provide insights into parameterization developments to address long-standing model biases. Maloney et al. (2019) summarizes the activities by the U.S. National Oceanic and Atmospheric Administration (NOAA) Modeling, Analysis, Prediction, and Projections program (MAPP) Model Diagnostics Task Force (MDTF) to apply community-developed PODs to climate and weather prediction models. Zhang et al. (2020) developed a diagnostics package that utilizes statistics derived from long-term ground-based measurements from the DOE Atmospheric Radiation Measurement (ARM) User Facility for climate model evaluation. Aerosol properties, however, are not included in these diagnostics packages.

The international collaborative AeroCom project (Myhre et al., 2013; Schulz et al., 2006) focuses on evaluation of aerosol predictions using available measurements and includes intercomparisons among global models to assess uncertainties in seasonal and regional variations in aerosol properties and their potential impact on climate. Their diagnostics heavily rely on satellite remote sensing products (e.g., aerosol optical depth) which have global coverage but poor spatial and temporal resolution that hinders a process-level understanding of the sources of model uncertainty. More recently, the Global Aerosol Synthesis and Science Project (GASSP, Reddington et al., 2017; Watson-Parris et al., 2019) has developed a global database of aerosol observations from fixed surface sites as well as ship and aircraft

platforms from 86 field campaigns between 1990 and 2015 that can be used for model evaluation. Recent field campaigns after year 2015 are not included in this effort.

Many aerosol properties are difficult to measure directly. Remote sensing instruments (e.g., ground and satellite radiometers) that only measure radiative properties of column-integrated aerosols, such as optical depth, are frequently used to evaluate model predictions. Instruments such as ground lidars (e.g., Campbell et al., 2002) or lidars onboard aircraft (e.g., Müller et al., 2014) and satellite (e.g., CALIPSO, Winker et al., 2009) platforms can provide vertical profiles of aerosol extinction, backscatter, and/or depolarization, but they do not directly measure aerosol number, size and composition. Therefore, the quantities measured by remote sensing instruments cannot be used alone to assess model predictions of aerosol-radiation-cloud-precipitation interactions. Surface monitoring sites provide long-term in situ aerosol property measurements but are limited to land locations with far fewer operational sites compared to those dedicated to routine meteorological sampling. Ship and aircraft platforms are commonly deployed during field campaigns to obtain in situ and remote sensing aerosol property measurements in remote or poorly sampled locations such as over the ocean and within the free troposphere, which are highly valuable when studying spatial variations of aerosols. Aircraft platforms also provide a means to obtain coincident measurements of aerosol and cloud properties needed to understand their interactions. Although in-situ ship and airborne aerosol measurements are usually limited to specific locations for short time periods, the increasing number of completed field campaigns conducted over a range of atmospheric conditions provides an opportunity to use them for model evaluation.

As noted by Reddington et al. (2017), the considerable investment in collecting field campaign measurements of aerosol properties is underexploited by the climate modeling community. This can be largely attributed to datasets located in disparate repositories and the lack of a standardized file format that requires excessive time and effort spent on manipulating the datasets to facilitate comparisons between observed and simulated values, especially for those unfamiliar with measurement techniques, assumptions, and uncertainties. With many field campaigns conducted since 2015 being available but rarely used for model evaluation, this study describes the first version of the ESM Aerosol-Cloud Diagnostics (ESMAC Diags) package to facilitate the evaluation of ESM-predicted aerosols, utilizing recent measurements from aircraft, ship and surface platforms collected by the U.S. DOE ARM and National Science Foundation (NSF) NCAR user facilities, most of which are in-situ measurements. The overall structure of ESMAC Diags is designed similar to the Aerosol Modeling Testbed for the Weather Research and Forecasting (WRF) model described in Fast et al. (2011), except that ESMAC Diags uses Python to interface the measurements with ESM output and does not preprocess the observational dataset into a common format. The diagnostics package is firstly designed with and applied to E3SM Atmosphere

Model version 1 (EAMv1, Rasch et al., 2019). EAMv1 uses an improved modal aerosol treatment implemented based on the 4-mode version of the modal aerosol module (MAM4, Liu et al., 2016), such as improved treatment of H₂SO₄ vapor for new particle formation (NPF), improved secondary organic aerosol (SOA) treatment, new marine organic aerosol (MOA) species, improvements to aerosol convective transport, wet removal, resuspension from evaporation and aerosol-affected cloud microphysical processes (Wang et al., 2020).

2. Introduction of ESMAC Diags

The workflow of ESMAC Diags v1 is illustrated in Figure 1. Most field campaign datasets are directly read by the diagnostics package. In some field campaigns, more than one instrument is used to measure aerosol size distribution over different size ranges. We therefore merge these datasets to create a more complete description of the size distribution. These data are introduced in Section 2.1. Model outputs are extracted at the ground sites and along the flight tracks or ship tracks. The simulation and preprocessing details are provided in Section 2.2. ESMAC Diags reads in these field campaign and model data with quality controls and generates a set of diagnostics and metrics listed in Section 2.3. The diagnostics package is designed to be flexible so that additional measurements and functionality can be included in the future. Figure 2 depicts the directory structure to illustrate the organization of the datasets and code. Most of the datasets used in ESMAC Diags are in a standardized netCDF format (NETCDF, 2021); however, some ARM aircraft measurements use different ASCII formats. Currently, the diagnostic package reads observational data directly from their original format. In the long term, we may standardize the observational data format in a similar manner as was done in GASSP project (Reddington et al., 2017).





Figure 1: Workflow of ESMAC Diags. Data preprocessing and input are indicated by blue; diagnostics and plotting are indicated by orange.



124

125 **Figure 2: Structure of ESMAC Diags.** The “scripts” directory contains executable scripts and user-
 126 specified settings. The “src” directory contains all source code including code used to preprocess
 127 model output, read files, merge measurements from different instruments, compute observed
 128 versus simulated statistical relationships, and plot results. All observational and model data in the
 129 “data” directory are organized by field campaign. The diagnostic plots and statistics are put in the
 130 “figures” directory, also organized by field campaign. The “testcase” directory includes a small
 131 amount of input and verify data to test if the package is installed properly. The “webpage”
 132 directory provides an interface to view diagnostics figures. Boxes in blue describe the functions of
 133 the directory. Asterisks represent boxes that follow the same format as those shown in parallel.

134 2.1 Field observations and merged aerosol size distribution

135 We initially focus on four geographical regions where liquid clouds occur frequently and extensive
 136 measurements are available from ARM and other agencies: Eastern North Atlantic (ENA), Northeastern
 137 Pacific (NEP), Central U.S. (CUS, where the ARM Southern Great Plains, SGP, site is located), and
 138 Southern Ocean (SO). Aerosol properties also vary among these regions. Six field campaigns from these
 139 four testbeds are selected in the version 1 of ESMAC Diags (Table 1). HI-SCALE and ACE-ENA are

based on long-term ARM ground sites with aircraft field campaigns sampling below, within, and above convective and marine boundary layer clouds, respectively, within a few hundred kilometers around the sites. CSET and MAGIC are field campaigns with aircraft and ship platforms, respectively, sampling transects between California and Hawaii characterized by a transition between stratocumulus and trade cumulus dominated regions. SOCRATES and MARCUS are field campaigns with aircraft and ship platforms, respectively, based out of Hobart, Australia. Aircraft transects during SOCRATES extended south to around 60°S, while ship transects during MARCUS extended southwest from Hobart to Antarctica. The aircraft (black) and ship (red) tracks for these field campaigns are shown in Figure 3.

Table 1. Descriptions of the field campaigns used in this study. Numbers after aircraft or ship represent number of flights or ship trips in each field campaign or IOP.

Campaign*	Period	Platform	Typical Conditions	Reference
HI-SCALE	IOP1: 24 Apr – 21 May 2016 IOP2: 28 Aug – 24 Sep 2016	Ground, aircraft (IOP1: 17, IOP2: 21)	Continental cumulus with high aerosol loading	(Fast et al., 2019)
ACE-ENA	IOP1: 21 Jun – 20 Jul 2017 IOP2: 15 Jan – 18 Feb 2018	Ground, aircraft (IOP1: 20, IOP2: 19)	Marine stratocumulus with low aerosol loading	(Wang et al., 2021)
MAGIC	Oct 2012 – Sep 2013	Ship (18)	Marine stratocumulus to cumulus transition with low aerosol loading	(Lewis and Teixeira, 2015; Zhou et al., 2015)
CSET	1 Jul – 15 Aug 2015	Aircraft (16)	Same as above	(Albrecht et al., 2019)
MARCUS	Oct 2017 – Apr 2018	Ship (4)	Marine liquid and mixed phase clouds with low aerosol loading	(McFarquhar et al., 2021)
SOCRATES	15 Jan – 24 Feb, 2018	Aircraft (14)	Same as above	(McFarquhar et al., 2021)

* Full names of the listed field campaigns:

HI-SCALE: Holistic Interactions of Shallow Clouds, Aerosols and Land Ecosystems

ACE-ENA: Aerosol and Cloud Experiments in the Eastern North Atlantic

MAGIC: Marine ARM GCSS Pacific Cross-section Intercomparison (GPCI) Investigation of Clouds

CSET: Cloud System Evolution in the Trades

MARCUS: Measurements of Aerosols, Radiation and Clouds over the Southern Ocean

SOCRATES: Southern Ocean Cloud Radiation and Aerosol Transport Experimental Study




Figure 3. Aircraft (black) and ship (red) tracks for the six field campaigns. Overlaid is aerosol optical depth at 550nm averaged from 2014 to 2018 simulated in EAMv1.

The instruments and measurements used in ESMAC Diags version 1 are listed in Table 2. All in-situ measurements are converted to under ambient temperature and pressure. Note that some instruments are only available for certain field campaigns or failed operationally during certain periods, so that model evaluation is limited by the availability of data collected in each field campaign. ARM data usually include quality flags indicating bad or indeterminate data. These flagged data are filtered out, except surface condensation particle counter (CPC) measurements for HI-SCALE, that data flagged as greater than maximum value (8000 cm^{-3}) are retained since aerosol loading can be higher than that during NPF events. This exception ensures a reasonable diurnal cycle shown in Section 3.3. For some data that do not have a quality flag, a simple minimum and maximum threshold is applied (e.g., 500 cm^{-3} maximum threshold is used for each UHSAS bin from the NCAR research flight measurements).

Table 2. List of instruments and measurements used in ESMAC Diags v1.

Instrument	Platform	Measurements	Available campaigns	DOIs or References
Surface meteorological station (MET)	Ground, ship	Temperature, relative humidity, wind, pressure	HI-SCALE, ACE-ENA, MAGIC, MARCUS	HI-SCALE, ACE-ENA: (Kyrouac and Shi, 2018) MAGIC: (ARM, 2014) MARCUS: 10.5439/1593144
Scanning mobility particle sizer (SMPS)	Ground	Aerosol size distribution (20-700 nm)	HI-SCALE	(Howie and Kuang, 2016)

Nano scanning mobility particle sizer (nanoSMPS)	Ground	Aerosol size distribution (2-150 nm)	HI-SCALE	(Koontz and Kuang, 2016)
Ultra-High Sensitivity Aerosol Spectrometer (UHSAS)	Ground, aircraft, ship	Aerosol size distribution (60 – 1000 nm), number concentration	HI-SCALE, ACE-ENA, MAGIC, MARCUS, CSET, SOCRATES	HI-SCALE, MAGIC, MARCUS: (Koontz and Uin, 2018) ACE-ENA: (Uin et al., 2018) CSET: 10.5065/D65Q4T96 SOCRATES: 10.5065/D6M32TM9
Condensation particle counter (CPC)	Ground, aircraft, ship	Aerosol number concentration (> 10 nm)	HI-SCALE, ACE-ENA, MAGIC, MARCUS	HI-SCALE (ground): (Kuang et al., 2016) ACE-ENA (ground), MAGIC: (Kuang et al., 2018a) MARCUS: (Kuang et al., 2018b) HI-SCALE (aircraft): (ARM, 2016b) ACE-ENA (aircraft): (Mei, 2018)
Condensation particle counter – ultrafine (CPCU)	Ground, aircraft	Aerosol number concentration (> 3 nm)	HI-SCALE, ACE-ENA	HI-SCALE (ground): 10.5439/1046186 HI-SCALE (aircraft): (ARM, 2016b) ACE-ENA (aircraft): 10.5439/1440985
Condensation nuclei counter (CNC)	Aircraft	Aerosol number concentration (11-3000 nm)	CSET, SOCRATES	CSET: 10.5065/D65Q4T96 SOCRATES: 10.5065/D6M32TM9
Cloud condensation nuclei (CCN) counter	Ground, aircraft, ship	CCN number concentration (0.1% to 0.5% supersaturation* depending on the platform)	HI-SCALE, ACE-ENA, MAGIC, MARCUS, SOCRATES	HI-SCALE (ground), ACE-ENA (ground), MARCUS: 10.5439/1342133 MAGIC: 10.5439/1227964 SOCRATES: 10.5065/D6Z036XB HI-SCALE (aircraft): (ARM, 2016a)
Aerosol chemical speciation monitor (ACSM)	Ground	Aerosol composition	HI-SCALE, ACE-ENA	10.5439/1762267
Microwave radiometer (MWR)	Ground, ship	Liquid water path, precipitable water vapor	MAGIC, MARCUS	10.5439/1027369
Counterflow virtual impactor (CVI)	Aircraft	Separates large droplets or ice crystals	HI-SCALE, ACE-ENA, SOCRATES	HI-SCALE: (ARM, 2016a) ACE-ENA: 10.5439/1406248 SOCRATES: 10.5065/D6M32TM9
Fast integrated mobility spectrometer (FIMS)	Aircraft	Aerosol size distribution (10 – 425 nm)	HI-SCALE, ACE-ENA	HI-SCALE: (ARM, 2017) ACE-ENA: (ARM, 2020)
Passive cavity aerosol spectrometer (PCA SP)	Aircraft	Aerosol size distribution (120 – 3000 nm)	HI-SCALE, ACE-ENA, CSET	HI-SCALE: (ARM, 2016a) ACE-ENA: (ARM, 2018) CSET: 10.5065/D65Q4T96

Optical particle counter (OPC)	Aircraft	Aerosol size distribution (390 – 15960 nm)	ACE-ENA	(ARM, 2018)
Interagency working group for airborne data and telemetry systems (IWG)	Aircraft	Navigation information and atmospheric state parameters	HI-SCALE, ACE-ENA	HI-SCALE: (ARM, 2017) ACE-ENA: (ARM, 2018)
High-resolution time-of-flight aerosol mass spectrometer (AMS)	Aircraft	Aerosol composition	HI-SCALE, ACE-ENA	HI-SCALE: (ARM, 2017) ACE-ENA: 10.5439/1468474
Water content measuring system (WCM)	Aircraft	Cloud liquid and total water content	HI-SCALE, ACE-ENA	HI-SCALE: (ARM, 2016a) ACE-ENA: 10.5439/1465759
Doppler lidar (DL)	Ground	Boundary layer height	HI-SCALE	10.5439/1726254
Reprocessed CN and CCN data to remove ship exhaust influence	Ship	CN, CCN number concentration	MARCUS	10.25919/ezp0-em87

* For measured supersaturations (SS) that vary over time, a $\pm 0.05\%$ window is applied (e.g., 0.5% SS includes samples with SS between 0.45% and 0.55%).

For some field campaigns (HI-SCALE and ACE-ENA), there are several instruments (e.g., FIMS, PCASP, OPC for aircraft; SMPS and nanoSMPS for ground) measuring aerosol size distribution over different size ranges. These datasets are merged to create a more complete size distribution. In ESMAC Diags v1, aerosol “size” refers to mobility and optical dry diameter of particles. The aerosol concentrations in the “overlapping” bins measured by multiple instruments are weighted by the uncertainty of each instrument based on the knowledge of the ARM instrument mentors. An example of the merged aerosol size distribution and individual measurements for one flight in ACE-ENA is shown in Figure 4. Ranging from 10^1 to 10^4 nm, the merged aerosol size distribution data account for ultrafine, Aitken, and accumulation modes.

Although these measurements are considered as “truth” when evaluating ESMs, we note that they are subject to limitations and uncertainties due to theoretical/methodological formulations, sampling representativeness, instrumental accuracy and precision, imperfect calibration, random errors, etc. In addition, sampling volumes differ between observations and model output and are not reconcilable. It is difficult to quantify every aspect of observational uncertainty within the context of interpreting comparisons with model output, but we try to discuss some of them in this study to the best of our

knowledge. Percentiles (either 25% - 75% or 5% - 95%) are used in some analyses of this study to approximate data variability that is likely to be much higher than measurement uncertainty.




Figure 4. An example of a mean aerosol number distribution merged from FIMS, PCASP and OPC instruments for ACE-ENA aircraft measurements on 29 June 2017.

2.2 Preprocessing of model output

In this study, we run EAMv1 from 2012 to 2018, covering all six field campaign periods introduced previously, with enough time for model spin-up. The model is configured to follow the Atmospheric Model Intercomparison Project (AMIP) protocol (Gates et al., 1999) with real-world forcings (e.g., greenhouse gases, sea surface temperature, aerosol emissions, etc.). For each simulation year, we use the year 2014 emission data from CMIP6, since the emission data does not cover years after 2014. The simulated horizontal winds are nudged towards the Modern-Era Retrospective analysis for Research and Applications, Version 2 (MERRA-2, Gelaro et al., 2017) with a relaxation time scale of 6 hours. Previous studies (Sun et al., 2019; Zhang et al., 2014) showed that with such nudging configuration the large-scale circulation is well constrained in the nudged simulation, especially for the mid- and high- latitude regions. The simulation uses a horizontal grid spacing of $\sim 1^\circ$ (NE30, the number of elements along a cube face of the E3SM High-Order Methods Modeling Environment, HOMME, dynamics core) with a 30-minute timestep. We saved hourly output to compare with field campaign measurements. The diagnostics package post-processes 3-D model variables associated with aerosol concentration, size, composition, optical properties, precursor concentration, CCN concentration, and atmospheric state variables. The size of output data is reduced by saving 3-D variables only over the field campaign regions. The model

configuration and execution scripts are uploaded as electronic supplement to this paper. Users can apply it in their own E3SM simulations (or output similar variables if running other models) to use this package.

We extracted model output along the aircraft (ship) tracks using an “aircraft simulator” (Fast et al., 2011) strategy to facilitate comparisons of observations and model predictions. At each aircraft (ship) measurement time, we find the nearest model grid cell, output time slice, and vertical level of the aircraft altitude (or the lowest level for ship) to obtain the appropriate model values. Since there are both spatial and temporal mismatch existing between model output and field measurements, the evaluation focuses on overall statistics. We also calculate the aerosol size distribution from 1 nm to 3000 nm at 1 nm increments from the individual size distribution modes in MAM4 to facilitate comparisons with observed aerosol number distribution that has different size ranges for different instruments. All these variables are saved in separate directories according to the specific aircraft (ship) tracks, as indicated in Figure 2.

2.3 List of diagnostics and metrics

Currently, ESMAC Diags produces the following diagnostics and metrics:

- Mean value, bias, root mean square error (RMSE) and correlation of aerosol number concentration.
- Timeseries of aerosol variables (aerosol number concentration, aerosol number size distribution, chemical composition, CCN number concentration) for each field campaign or intensive observational period (IOP) at the surface or along each flight (ship) track.
- Diurnal cycle of aerosol variables at the surface.
- Mean aerosol number size distribution for each field campaign or IOP.
- Percentiles of aerosol variables by height for each field campaign or IOP.
- Percentiles of aerosol variables by latitude for each field campaign or IOP.
- Pie/bar charts of observed and predicted aerosol composition averaged over each field campaign or IOP.
- Vertical profile of cloud fraction and LWC composite of aircraft measurements for each field campaign or IOP.
- Timeseries of atmospheric state variables.
- Aircraft and ship track maps.

In the next section we will demonstrate these diagnostics and metrics by providing several examples.

3. Examples

Aerosol number concentration, size distribution, and chemical composition (that controls hygroscopicity) are key quantities that impact aerosol-cloud interactions, such as the activation of cloud droplets. Errors in model predictions of these aerosol properties contribute to uncertainties in aerosol direct and indirect radiative forcing. These aerosol properties vary dramatically depending on location, altitude, season, and meteorological conditions due to variability in emissions, formation mechanisms, and removal processes in the atmosphere. This section shows some examples to illustrate the usage of this diagnostics package on evaluating global models.

3.1 Aerosol size distributions and number concentrations

Aerosol properties are highly dependent on location and season. Figure 5 shows the mean aerosol size distribution for each of the four testbed regions. For HI-SCALE and ACE-ENA, the two IOPs operated in different seasons are shown separately. Table 3 shows the mean aerosol number concentration from these field campaigns, for two particle size ranges: >10 nm and >100 nm. The inter-quartile range (25% and 75% percentiles) are also shown to illustrate the variability in space and time. Among the four testbed regions, the CUS region has the largest aerosol number concentrations since the other field campaigns are primarily over open ocean. Overall, EAMv1 overestimates Aitken mode (10 – 70 nm) aerosols and underestimates accumulation mode (70 – 400 nm) aerosols for the CUS and ENA regions, suggesting that processes related to particle growth or coagulation might be too weak in the model. Over the NEP region, EAMv1 overestimates aerosol number for particle sizes >100 nm and >10 nm (Table 3), both at the surface and aloft. Over the SO region, which is considered a pristine region with low aerosol concentration, observations show a significant number of particles <200 nm in both aircraft and ship measurements (Figure 5). The mean aerosol number concentration over SO region is comparable or even greater than the other ocean testbeds (Table 3). In contrast, EAMv1 simulates a clean environment with the lowest aerosol number concentrations among the four regions. These types of comparisons demonstrate the need for additional analyses to understand why SO has similar aerosol number with other ocean regions and why EAMv1 cannot simulate this feature. The observed 75% percentiles are sometimes smaller than the mean value (Table 3), indicating skewed aerosol size distribution with long tail in large aerosol size. EAMv1 usually produces smaller inter-quartile range than the observations, likely because the current model resolution is too coarse to capture the observed spatial variability in aerosol properties.




Figure 5: Mean aerosol number distribution averaged for each field campaign or IOP. Shadings denote the range between 10% and 90% percentiles.

Table 3: Mean aerosol number concentration and inter-quartile range (25% and 75% percentiles, small numbers in parenthesis) for two size ranges averaged for each field campaign (or each IOP for HI-SCALE and ACE-ENA). Aircraft measurements 30 minutes after take-off and before landing are excluded to remove possible contamination from the airport.

Unit: #/cm ³			>10 nm		>100 nm	
			CPC	E3SMv1	UHSAS/PCASP*	E3SMv1
CUS	Surface (HI-SCALE)	IOP1	4095 (2198, 4943)	4566 (2865, 5984)	675.1 (393.2, 929.5)	321.3 (229.7, 400.8)
		IOP2	N/A	N/A	N/A	N/A
	Aircraft (HI-SCALE)	IOP1	4206 (1132, 5013)	3872 (2803, 4946)	465.7 (112.6, 616.1)	159.6 (112.2, 200.5)
		IOP2	4121 (1610, 3829)	2514 (1332, 3584)	789.1 (444.4, 1088.0)	383.6 (280.7, 483.8)
ENA	Surface	IOP1	610 (343, 711)	1723 (600, 1650)	206.1 (134.5, 267.1)	209.8 (155.3, 255.5)

	(ACE-ENA)	IOP2	458 (239, 505)	843 (320, 1152)	59.6 (25.0, 71.9)	61.9 (53.6, 71.9)
	Aircraft (ACE-ENA)	IOP1	576 (264, 677)	919 (562, 917)	135.6 (65.3, 185.1)	199.9 (146.6, 266.3)
		IOP2	356 (132, 383)	521 (279, 627)	72.8 (22.2, 72.8)	50.3 (41.6, 62.3)
NEP	Ship (MAGIC)		417 (117, 285)	1271 (356, 1652)	113.6 (47.0, 139.9)	143.0 (93.7, 148.5)
	Aircraft (CSET)		408 (155, 386)	607 (353, 675)	81.5 (17.0, 73.4)	134.5 (81.2, 151.3)
SO	Ship (MARCUS)		354 (244, 415)	303 (164, 326)	68.5 (36.8, 94.0)	54.5 (32.5, 71.7)
	Aircraft (SOCRATES)		988 (327, 991)	237 (169, 270)	56.2 (14.1, 50.4)	32.3 (13.2, 42.2)

* PCASP is available only on aircraft for HI-SCALE and ACE-ENA. UHSAS is available only in surface measurements for HI-SCALE and ACE-ENA, and in other field campaigns.

Both observed and simulated aerosol size distribution and number concentration show large variability during these field campaigns. Over the period of a few weeks or longer, aerosol number can vary by an order of magnitude between the 10% and 90% percentiles, especially for small particles (Figure 5). Figure 6 shows mean aerosol size distributions for two flight days during HI-SCALE: one with a large number of small (<70 nm) particles (14 May) and the other (3 September) with fewer small particles but more accumulation mode (70 – 300 nm) particles. On both days, EAMv1 reproduced the observed planetary boundary layer (PBL) height (PBLH) reasonably well with sufficient samples below and above PBL. On 14 May, EAMv1 reproduces the observed aerosol size distribution reasonably well both within the PBL and in the lower free atmosphere. However, on 3 September EAMv1 produces too many aerosols in the Aitken mode and too few accumulation mode aerosols in the PBL. In the free atmosphere, EAMv1 reproduces the lower concentration of Aitken mode aerosols but still underestimates the accumulation mode. Such contrasting cases will be useful to help diagnose the specific processes contributing to model uncertainties in future analyses. This large day-to-day variability also indicates that long-term measurements are needed to avoid sampling bias in building robust statistics in aerosol properties. The next version of ESMAC Diags will be extended to include the available long-term ARM measurements at SGP, ENA and other sites outside of the field campaign time periods.




Figure 6: (Top) Mean aerosol number distribution for two flights during HI-SCALE: (left) 14 May 2016 and (right) 3 September 2016, for data above (dashed line) and below (solid line) observed PBLH. If there is cloud observed within a 1-hour window of the sample point, the above-PBL sample needs to be above cloud top and the below-PBL sample needs to be below cloud base for the sample point to be chosen. Shadings represent the data range between 10% and 90% percentiles. Relatively large particles with no shading indicate more than 90% of samples with zero values. (bottom) Timeseries of observed (black) and simulated (red) PBLH overlaid with flight height (blue) during the two flight periods. The observed PBLH is derived from Doppler lidar measurements.

3.2 Vertical profiles of aerosol properties

A research aircraft is the primary platform to provide information on the vertical variations of key aerosol properties that cannot be obtained accurately by remote sensing instrumentation. In this section we show an example of evaluating vertical profiles of aerosol properties using aircraft measurements as well as illustrate the capability of evaluating multiple model simulations with ESMAC Diags. In addition to the standard EAMv1 simulation described in the previous section, we performed an EAMv1 simulation using the regionally refined mesh (RRM) (Tang et al., 2019). The model is configured to run with the horizontal grid spacing of $\sim 0.25^\circ$ over the continental U.S. and $\sim 1^\circ$ elsewhere. The two model configurations are identical, except for the higher spatial resolution (including primary aerosol emissions) in the RRM over

the continental U.S. All aircraft measurements with a cloud detected simultaneously (cloud flag = 1) were excluded.

Figure 7 shows vertical percentiles of aerosol number concentration, composition and CCN number concentration among all the HI-SCALE aircraft flights. Note that aircraft rarely flew above 3 km during HI-SCALE so the sample size above that altitude is much smaller. The observed aerosol concentrations of number and chemical composition decrease with height since the major sources of aerosols (anthropogenic, biogenic, and biomass burning) (Liu et al., 2021) are from precursors emitted near the surface and chemical formation within the PBL. EAMv1 generally simulates less variability than observations except for sulfate. Overall, EAMv1 reproduces the observed mean aerosol number concentration for aerosol size > 10 nm but underestimates the number of larger particles > 100 nm during HI-SCALE (Table 3). The model also overestimates sulfate and underestimates organic matter concentrations when compared to aircraft AMS measurements. Its underestimation of CCN number concentration is consistent with underestimation of aerosol number concentration for diameter > 100 nm but contrary with overestimation of sulfate. A similar relationship is seen for ACE-ENA, to be described later in this section.




Figure 7: Vertical profiles of (from left to right): aerosol number concentration, mass concentration of sulfate, mass concentration of total organic matter and CCN number concentration under the supersaturation in the parentheses for HI-SCALE (top) IOP1 and (bottom) IOP2. The percentile box represents 25% and 75% percentiles, and the bar represents 5% and 95% percentiles.

The differences in sulfate and organic matter aloft is consistent with longer term surface measurement differences shown in Figure 8, suggesting this is a model bias. Note that near-surface measurements by aircraft are not always consistent with ground measurements (e.g., total organic matter in IOP1), which reflects the large spatial variability in aerosol properties associated with the aircraft flight paths up to a few hundred kilometers around the ARM site. The greater fraction of sulfate in EAMv1 suggests that the simulated aerosol hygroscopicity is likely higher than observed. Currently only these two species are available in both EAMv1 and AMS/ACSM observations for comparison purpose. Zaveri et al. (2021)

recently added chemistry associated with NO_3 formation in MAM4, which is expected to be implemented in a future version of EAM.




Figure 8: Bar plots of the surface average aerosol composition during HI-SCALE IOP1 (top) and IOP2 (bottom). Observations are obtained from an ACSM. Dust and black carbon (BC) are not measured in the observation. NO_3 and NH_4 are not predicted in EAMv1 and RRM.

Ongoing developments in E3SM will soon permit regional-refined meshes with grid spacings as small as ~ 3 km as well as global convection-permitting simulations ($\Delta x \sim 3$ km); therefore, this diagnostics package is designed to be flexible in scale to take advantage of higher-resolution ESM simulations that are more compatible with high-resolution in-situ aerosol observations. This study demonstrates this ability by using a 0.25° RRM simulation. Overall, the RRM analyzed here has similar biases as EAMv1, with differences that vary seasonally. The inter-quartile range in Figure 7 show that the variability of organic aerosols and CCN from the EAMv1 and RRM simulations are similar. However, the variability of sulfate in RRM is larger than EAMv1 and observations during the spring IOP (IOP1). During the summer IOP (IOP2), the variabilities of sulfate in EAMv1, RRM, and observations are similar, and the sulfate concentrations from RRM are closer to observed than EAMv1. Individual timeseries from the RRM simulation are still too smooth to capture the fine scale variability of aerosols in observations (not shown).

We expect E3SM to capture more fine scale variabilities related to urban and point sources of aerosols and their precursors when the simulation grid spacing is further reduced to ~ 3 km. A sensitivity study will be conducted when this high-resolution version of E3SM simulation becomes available.

Figure 9 shows the vertical variation in percentiles of aerosol properties for ACE-ENA. The observed aerosol number concentrations, composition masses, and CCN number concentrations are much smaller than those for HI-SCALE, representing a cleaner ocean environment. EAMv1 produces larger mean values than the observations for all these quantities. The overall variabilities in predicted aerosol number and concentrations of sulfate and organic matter are also greater than observed. Note that the observed variabilities for HI-SCALE are much larger than for ACE-ENA, indicating that EAMv1 has smaller location variation on aerosol variabilities. The observed total organic concentration shows a peak aloft between 1.6 and 2.2 km, corresponding to the level of CCN number concentration peak. This implies a major source of aerosols or precursors is free tropospheric transport (Zawadowicz et al., 2021). This peak of total organic concentration aloft is also captured by the model.

The bar plots in Figure 10 of aerosol composition at the surface during ACE-ENA from the ACSM instrument and EAMv1 illustrate a similar bias in sulfate and organic mass as aloft. While the surface sulfate measurements are like those from the aircraft at the lowest altitudes, the observed surface organic matter is much higher than aloft, particularly during IOP2. The differences in these measurements may be due to local effects or possible contamination from aircraft since the surface station is located near an airport on an island.




Figure 9: Same as Figure 7 but for ACE-ENA.




Figure 10: Same as Figure 8 but for ACE-ENA.

3.3 New particle formation events

Aerosol number concentrations and size distributions are highly impacted by NPF events (Kulmala et al., 2004), which further influence CCN concentration (e.g., Kuang et al., 2009; Pierce and Adams, 2009) and ultimately cloud properties. NPF and subsequent particle growth are frequently observed in the CUS region (Hodshire et al., 2016). As described by Fast et al. (2019) and shown in Figure 11a, several NPF events were observed during the HI-SCALE spring IOP (IOP1). Large concentrations of aerosols smaller than 10 nm were observed, with the size growing larger over the next few hours. The average diurnal variation in aerosol number distribution in Figure 12a shows that NPF events usually occur during the morning between 12 and 15 UTC (6 – 9 am local time), followed by particle growth during the rest of the morning and afternoon. This variation is also seen in the diurnally averaged CPC measurements of aerosol diameters > 3 nm and > 10 nm (Figure 12c) but diurnal changes in CCN number concentrations (Figure 12d) are more modest.




Figure 11: Time series of (a) observed and (b) simulated surface aerosol number distribution during HI-SCALE IOP1. The observed aerosol number distribution is from merged nanoSMPS and SMPS. Model data is cut off at 500 nm to compare with observation.




Figure 12: Average diurnal cycle of surface (a) observed aerosol number distribution, (b) simulated aerosol number distribution, (c) aerosol number concentration for diameters > 10 nm and > 3 nm, and (d) CCN number concentration for supersaturations of 0.1% and 0.5% for HI-SCALE IOP1.

Various NPF pathways associated with different chemical species have been proposed and implemented in models. Two NPF pathways are considered in MAM4 in EAMv1: a binary nucleation pathway and a PBL cluster nucleation pathway. However, the current simulation does not reproduce the observed large day-to-day variability of small particle concentrations due to NPF. Instead, the model produces high aerosol concentrations between 10 and 100 nm almost all the time. It also fails to reproduce

the large diurnal variability of aerosol and CCN number concentration with a peak seen in the morning near 15 UTC (9 am local time), 7 hours earlier than the observed 22 UTC (4 pm local time) afternoon peak. Its overestimation of aerosol number concentration for particle diameter >10 nm and underestimation of CCN number concentration is consistent with that shown in Figure 5. Several efforts are underway to improve the simulation of NPF by adding a nucleation mode in MAM4 to explicitly resolve ultrafine particles and implementing new chemical pathways to simulate NPF following Zhao et al. (2020). ESMAC Diags is being used to evaluate these new model developments.

Using aircraft measurements from ACE-ENA, Zheng et al. (2021) recently found evidence of NPF events occurring in the upper part of marine boundary layer between broken clouds following the passage of a cold front. 16 February 2018 is identified as a typical NPF day in Zheng et al. (2021). The vertical profiles of aerosol number and CCN concentrations measured by aircraft on 16 February 2018 are shown in Figure 13. The NPF event and particle growth happened in the upper boundary layer is shown by the large mean and variance of aerosol number concentration just below the base of the marine boundary layer clouds. EAMv1 could not simulate NPF events in the upper marine boundary layer on this day and other days during ACE-ENA, likely due to the lack of NPF mechanisms related to dimethyl sulfide (DMS) oxidation, and/or missing part in parameterizations to deal with the processes related to broken marine boundary layer clouds and sub-grid circulation. Similarly, the sharp increase of CCN number just above the level of marine boundary layer clouds is not simulated.




Figure 13: Vertical profiles of aerosol number concentration for diameters >10 nm and >3 nm, CCN number concentration, and cloud frequency measured by the 16 February 2018 flight in ACE-ENA. The percentile box represents 25% and 75% percentiles, and the bar represents 5% and 95% percentiles.

3.4 Latitudinal dependence of aerosols and clouds

Unlike some field campaigns (i.e., HI-SCALE and ACE-ENA) where aircraft missions were conducted over a relatively localized region with limited spatial variability of the meteorological conditions, ship and/or aircraft measurements over the NEP and SO testbed regions span regions > 1500 km (i.e., from California to Hawaii and from Tasmania to the far Southern Ocean, respectively). As shown in Figure 3, there are large spatial gradients in EAMv1 simulated aerosol optical depth along these ship/aircraft tracks. In ESMAC Diags version 1, we include composite plots of aerosol and cloud properties binned by latitude to assess model representation of synoptic-scale variations.




Figure 14: Percentiles of (a) air temperature, (b) grid-mean liquid water path (LWP), (c) aerosol number concentration for diameter >10 nm, and (d) aerosol number concentration for diameter >100 nm for all ship tracks in MAGIC binned by 1° latitude bins. The percentile box

represents 25% and 75% percentiles, and the bar represents 5% and 95% percentiles. The observed aerosol number concentrations for diameters >10 nm and >100 nm are obtained from CPC and UHSAS, respectively.




Figure 15: Percentiles of (a) cloud fraction, (b) aerosol number concentration for diameter >10 nm, and (c) aerosol number concentration for diameter >100 nm for all aircraft measurements between 0-3 km in CSET binned by 1° latitude bins. The percentile box represents 25% and 75% percentiles, and the bar represents 5% and 95% percentiles. The observed aerosol number concentrations for diameters >10 nm and >100 nm are obtained from CNC and UHSAS, respectively.


The research ship (aircraft) from the MAGIC (CSET) field campaign in the NEP testbed travelled between California and Hawaii, where there is frequently a transition between marine stratocumulus clouds near California and broken trade cumulus clouds near Hawaii (e.g., Teixeira et al., 2011). Although ESMAC Diags v1 focuses primarily on aerosols, we show some basic meteorological and cloud

fields here since they are important to illustrate the transition of cloud regimes along the ship (aircraft) tracks. Additional cloud properties derived from surface and satellite measurements are not included in the current analysis, but are being implemented in ESMAC Diags v2. Some of the meteorological, cloud, and aerosol properties along the ship (aircraft) tracks binned by latitude are shown in Figure 14 (Figure 15). Note that cloud fraction in Figure 15 is calculated as cloud frequency in aircraft observation and from grid-mean cloud fraction in model along the flight track. This is different from the classic definition of cloud fraction usually used for satellite measurements or models and is subject to aircraft sampling strategy. As the surface temperature increases from California to Hawaii (Figure 14a), the cloud fraction (Figure 15a) shows an decreasing trend southwestward, indicating the transition from stratocumulus to cumulus clouds. However, ship-measured LWP (Figure 14b) has no trend related to latitude, possibly because cumulus clouds at lower latitudes have smaller cloud fraction but larger LWP when clouds exist. EAMv1 shows decreasing trends of both cloud fraction and LWP from high to low latitudes along these tracks. It generally underestimates LWP and overestimates cloud fraction to the north of 30° N. For aerosol number concentrations, EAMv1 produces too many aerosols compared to measurements both at the surface (ship) and aloft (aircraft), consistent with the aerosol size distribution in Figure 5 and total number concentration in Table 3. However, EAMv1 does reproduce the increase trend in accumulation mode aerosol concentration approaching the California coast.

Similar latitudinal gradients of aerosol and CCN number concentrations along ship tracks from MARCUS and aircraft tracks from SOCRATES are shown in Figures 16 and 17, respectively. Over the SO region, NPF frequently occurs during austral summer when ample biogenic precursor gases (e.g., DMS) are released and rise into the free troposphere (McFarquhar et al., 2021; McCoy et al., 2021). Large values of ship-measured aerosol and CCN number concentration are observed near Antarctica corresponding to the coastal biological emissions of aerosol precursors, and also occur to the north of 45°S, indicating impacts from continental and anthropogenic sources. This is consistent with other studies (Sanchez et al., 2021; Humphries et al., 2021). EAMv1 underestimates aerosol and CCN number concentration near Antarctica. This bias, which may be related to too strong wet scavenging or insufficient NPF and growth, is commonly seen in many other ESMs (e.g., McCoy et al., 2020; McCoy et al., 2021). Aircraft flight paths during SOCRATES (Figure 17) do not extent as far south as the ship measurements (Figure 16). The observed aerosol properties have little latitudinal variation in general. EAMv1 underestimates aerosol number concentration for size > 10nm and CCN number concentration with SS=0.5%, but the predictions are closer to observed for aerosol size > 100 nm and CCN with SS=0.1% (Figure 17), consistent with the mean aerosol size distribution in Figure 5. This indicates that the model performs better in simulating accumulation mode than Aitken mode particles over SO. These

499 model aerosol biases are highly relevant when considering their interaction with clouds and radiations,
500 which will be included in version 2 of ESMAC Diags.

501



502

503 **Figure 16: Percentiles of (a) air temperature, (b) aerosol number concentration for diameter >10**
504 **nm, (c) aerosol number concentration for diameter >100 nm, (d) CCN number concentration for**

supersaturation $SS=0.1\%$, and (e) CCN number concentration for supersaturation $SS=0.5\%$ for all ship tracks in MARCUS binned by 1° latitude bins.




Figure 17: Percentiles of (a) aerosol number concentration for diameter $>10\text{ nm}$, (b) aerosol number concentration for diameter $>100\text{ nm}$, (c) CCN number concentration for supersaturation $SS=0.1\%$, and (d) CCN number concentration for supersaturation $SS=0.5\%$ for all aircraft measurements between 0-3 km in SOCRATES binned by 1° latitude bins.

4. Summary

A Python-based ESM aerosol-cloud diagnostics (ESMAC Diags) package is developed to quantify the performance of the DOE's E3SM atmospheric model using ARM and NCAR field campaign measurements. The first version of this diagnostics package focuses on aerosol properties. The measurements include aerosol number, size distribution, chemical composition, and CCN collected from surface, aircraft, and ship platforms needed to assess how well the aerosol lifecycle is represented across spatial and temporal scales which will subsequently impact uncertainties in aerosol radiative forcing estimates. Currently, the diagnostics cover the field campaigns of ACE-ENA, HI-SCALE, MAGIC/CSET, and MARCUS/SOCRATES over Northeastern Atlantic, Continental U.S., Northeastern Pacific, and Southern Ocean, respectively. The code structure is designed to be flexible and modular for future extension to other field campaigns or additional datasets. Since there is no one instrument that can measure the entire aerosol size distribution, we have constructed merged aerosol size distributions from two or more ARM instruments to better assess predicted size distributions. An "aircraft simulator" is used to extract aerosol and meteorological model variables along flight paths that vary in space and time. Similarly, the aircraft simulator is applied to ship tracks in which the altitude remains fixed at sea level.

The version 1 of ESMAC Diags package provides various types of diagnostics and metrics, including timeseries, diurnal cycles, mean aerosol size distribution, pie charts for aerosol composition, percentiles by height, percentiles by latitude, mean statistics of aerosol number concentration, and more. This allows quantification of model performance predicting aerosol number, size, composition, vertical distribution, spatial distribution (along ship tracks or aircraft tracks) and new particle formation events. A full set of diagnostics plots and metrics for simulations used in this paper are available at https://portal.nersc.gov/project/m3525/sqtang/ESMAC_Diags_v1/forGMD/webpage/ and are archived as an electronic supplement of this paper. This paper shows some examples to demonstrate the capability of ESMAC Diags to evaluate EAMv1 simulated aerosol properties. The diagnostics package also allows multiple simulations in one plot to compare different models or model versions. It can also be applied to evaluate other ESMs with necessary modifications to fit different model output formats.

Because in-situ aerosol measurements are usually collected at high temporal frequency (typically 1 second to a minute) over fine spatial volumes, there is a spatiotemporal scale mismatch with the standard climate model resolution (usually 1-degree grid spacing with hourly output). This is a limitation that cannot be completely overcome and must be accepted to perform model-observation comparisons necessary for identifying shortcomings in model representation of aerosol, cloud, and aerosol-cloud interaction processes that are the primary source for uncertainties in prediction of future climate. As new versions of E3SM become available that has grid spacings as small as a few kilometers via regional-refined and convection-permitting global domains (e.g., Caldwell et al., 2021), spatiotemporal

variabilities of aerosols at finer scales should be captured and be more compatible with fine resolution observations such that resolution impacts on statistical differences can be quantified. The diagnostics package will be applied to diagnose high resolution model output when the data are available.

While the current version focuses on aerosol properties, a version 2 of ESMAC Diags is being developed to include more diagnostics and metrics for cloud, precipitation, and radiation properties to facilitate the evaluation of aerosol-cloud interactions. These include inversion strength, above cloud relative humidity, cloud-surface coupling, cloud fraction, depth, LWP, optical depth, effective radius, droplet number concentration, adiabaticity, and albedo, precipitation rate, and more. Long-term surface-based and satellite retrievals will also be used to provide better statistics in model evaluation and to address limitations related to data coverage and uncertainty. Analyses are being designed to quantify relationships between these variables and relate them to effective radiative forcing, which will be used to assess and improve model parameterizations. In the future, this diagnostics package may also be extended to include other field campaigns that provide valuable data on aerosol properties and cloud-aerosol interactions, such as the ARM Layered Atlantic Smoke Interactions with Clouds (LASIC, Zuidema et al., 2018), NASA ObseRvations of Aerosols above CLouds and their intEractionS (ORACLES, Redemann et al., 2021), or NASA Atmospheric Tomography Mission (ATom, Brock et al., 2019) campaigns. As an open-source package, ESMAC Diags can also be applied by any user to other ESMs with small modifications on model preprocessing.

While there are other efforts to develop model diagnostics packages, this diagnostics package provides a unique capability for detailed evaluation of aerosol properties that are tightly connected with parameterized processes. Together with other commonly used diagnostics packages such as the ARM diagnostics package (Zhang et al., 2020), the DOE E3SM diagnostics package, and the PCMDI's metrics package (Gleckler et al., 2016), we expect to better understand the strengths and weaknesses of E3SM or other ESMs and provide insights into model deficiencies to guide future model development. This includes studies that develop a better understanding of how various processes contribute to uncertainties in aerosol number and composition predictions and subsequent representation of CCN and aerosol radiative forcing estimates.

Code availability:

The current version of ESMAC Diags is publicly available through GitHub (https://github.com/eagles-project/ESMAC_diags) under the new BSD license. The exact version (1.0.0-beta.2) of the code used to produce the results used in this paper is archived on Zenodo (<https://doi.org/10.5281/zenodo.6371596>). The model simulation used in this paper is version 1.0 of E3SM (<https://doi.org/10.11578/E3SM/dc.20180418.36>). The model configuration and execution scripts are uploaded as electronic supplement to this paper.

Data availability:

Measurements from the HI-SCALE, ACE-ENA, MAGIC, and MARCUS campaigns as well as the SGP and ENA sites are supported by the DOE Atmospheric Radiation Measurement (ARM) user facility and available at <https://adc.arm.gov/discovery/>. Measurements from the CSET and SOCRATES campaigns are supported by National Science Foundation (NSF) and obtained from NCAR Earth Observing Laboratory at https://data.eol.ucar.edu/master_lists/generated/cset/ and https://data.eol.ucar.edu/master_lists/generated/socrates/, respectively. DOI numbers or references of individual instruments are given in Table 2. All the above observational data and preprocessed model data used to produce the results used in this paper is archived on Zenodo (<https://doi.org/10.5281/zenodo.6369120>).

Author contribution:

ST, JDF and PM designed the diagnostics package; ST wrote the code and performed the analysis; JES, FM and MAZ processed the field campaign data; KZ contributed to the model simulation; JCH and ACV contributed to the package design and setup; ST wrote the original manuscript; all authors reviewed and edited the manuscript.

Competing interests:

Po-Lun Ma is a Topical Editor of Geoscientific Model Development. Other authors declare that they have no conflict of interest.

Acknowledgements:

This study was supported by the Enabling Aerosol-cloud interactions at GLObal convection-permitting scales (EAGLES) project (74358), funded by the U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research, Earth System Model Development (ESMD) program area. We thank the numerous instrument mentors for providing the data. This research used resources of the National Energy Research Scientific Computing Center (NERSC), a U.S. Department of Energy Office of Science User Facility operated under Contract No. DE-AC02-05CH11231. Pacific Northwest National Laboratory (PNNL) is operated for DOE by Battelle Memorial Institute under contract DE-AC05-76RL01830.

References:

- Albrecht, B., Ghate, V., Mohrmann, J., Wood, R., Zuidema, P., Bretherton, C., Schwartz, C., Eloranta, E., Glienke, S., Donaher, S., Sarkar, M., McGibbon, J., Nugent, A. D., Shaw, R. A., Fugal, J., Minnis, P., Paliknoda, R., Lussier, L., Jensen, J., Vivekanandan, J., Ellis, S., Tsai, P., Rilling, R., Haggerty, J., Campos, T., Stell, M., Reeves, M., Beaton, S., Allison, J., Stossmeister, G., Hall, S., and Schmidt, S.: Cloud System Evolution in the Trades (CSET): Following the Evolution of Boundary Layer Cloud Systems with the NSF–NCAR GV, *Bull. Amer. Meteor. Soc.*, 100, 93–121, <https://doi.org/10.1175/bams-d-17-0180.1>, 2019.
- Albrecht, B. A.: Aerosols, Cloud Microphysics, and Fractional Cloudiness, *Science*, 245, 1227–1230, <https://doi.org/10.1126/science.245.4923.1227>, 1989.
- AMWG Diagnostic Package: https://www.cesm.ucar.edu/working_groups/Atmosphere/amwg-diagnostics-package/, last access: 2 November 2021. 2021.
- ARM Intensive Operational Period (IOP) Data Browser: <https://iop.archive.arm.gov/arm-iop/2012/mag/magic/reynolds-marmet/>, last access: 2 November 2021. 2014.
- ARM Intensive Operational Period (IOP) Data Browser: <https://iop.archive.arm.gov/arm-iop/2016/sgp/hiscale/matthews-wcm>, last access: 2 November 2021. 2016a.
- ARM Intensive Operational Period (IOP) Data Browser: <https://iop.archive.arm.gov/arm-iop/2016/sgp/hiscale/mei-cpc>, last access: 16 September 2021. 2016b.
- ARM Intensive Operational Period (IOP) Data Browser: <https://iop.archive.arm.gov/arm-iop/2016/sgp/hiscale/wang-fims>, last access: 2 November 2021. 2017.
- ARM Intensive Operational Period (IOP) Data Browser: <https://iop.archive.arm.gov/arm-iop/2017/ena/aceena/tomlinson-pcas>, last access: 2 November 2021. 2018.
- ARM Intensive Operational Period (IOP) Data Browser: <https://iop.archive.arm.gov/arm-iop/2017/ena/aceena/wang-fims/>, last access: 2 November 2021. 2020.
- Brock, C. A., Williamson, C., Kupc, A., Froyd, K. D., Erdesz, F., Wagner, N., Richardson, M., Schwarz, J. P., Gao, R. S., Katich, J. M., Campuzano-Jost, P., Nault, B. A., Schroder, J. C., Jimenez, J. L., Weinzierl, B., Dollner, M., Bui, T., and Murphy, D. M.: Aerosol size distributions during the Atmospheric Tomography Mission (ATom): methods, uncertainties, and data products, *Atmos. Meas. Tech.*, 12, 3081–3099, <https://doi.org/10.5194/amt-12-3081-2019>, 2019.
- Caldwell, P. M., Terai, C. R., Hillman, B., Keen, N. D., Bogenschutz, P., Lin, W., Beydoun, H., Taylor, M., Bertagna, L., Bradley, A. M., Clevenger, T. C., Donahue, A. S., Eldred, C., Foucar, J., Golaz, J.-C., Guba, O., Jacob, R., Johnson, J., Krishna, J., Liu, W., Pressel, K., Salinger, A. G., Singh, B., Steyer, A., Ullrich, P., Wu, D., Yuan, X., Shpund, J., Ma, H.-Y., and Zender, C. S.: Convection-Permitting Simulations With the E3SM Global Atmosphere Model, *Journal of Advances in Modeling Earth Systems*, 13, e2021MS002544, <https://doi.org/https://doi.org/10.1029/2021MS002544>, 2021.
- Campbell, J. R., Hlavka, D. L., Welton, E. J., Flynn, C. J., Turner, D. D., Spinhirne, J. D., Scott, V. S., III, and Hwang, I. H.: Full-Time, Eye-Safe Cloud and Aerosol Lidar Observation at Atmospheric Radiation Measurement Program Sites: Instruments and Data Processing, *J. Atmos. Ocean. Technol.*, 19, 431–442, [https://doi.org/10.1175/1520-0426\(2002\)019<0431:FTESCA>2.0.CO;2](https://doi.org/10.1175/1520-0426(2002)019<0431:FTESCA>2.0.CO;2), 2002.

648 Carslaw, K. S., Lee, L. A., Regayre, L. A., and Johnson, J. S.: Climate Models Are Uncertain, but We Can
649 Do Something About It, *Eos*, 99, <https://doi.org/10.1029/2018EO093757>, 2018.

650 E3SM Diagnostics: <https://e3sm.org/resources/tools/diagnostic-tools/e3sm-diagnostics/>, last access: 2
651 November 2021. 2021.

652 Eyring, V., Righi, M., Lauer, A., Evaldsson, M., Wenzel, S., Jones, C., Anav, A., Andrews, O., Cionni, I.,
653 Davin, E. L., Deser, C., Ehbrecht, C., Friedlingstein, P., Gleckler, P., Gottschaldt, K. D., Hagemann, S.,
654 Juckes, M., Kindermann, S., Krasting, J., Kunert, D., Levine, R., Loew, A., Mäkelä, J., Martin, G., Mason,
655 E., Phillips, A. S., Read, S., Rio, C., Roehrig, R., Senftleben, D., Sterl, A., van Ulft, L. H., Walton, J., Wang,
656 S., and Williams, K. D.: ESMValTool (v1.0) – a community diagnostic and performance metrics tool for
657 routine evaluation of Earth system models in CMIP, *Geosci. Model Dev.*, 9, 1747-1802,
658 <https://doi.org/10.5194/gmd-9-1747-2016>, 2016.

659 Fast, J. D., Gustafson, W. I., Chapman, E. G., Easter, R. C., Rishel, J. P., Zaveri, R. A., Grell, G. A., and
660 Barth, M. C.: The Aerosol Modeling Testbed: A Community Tool to Objectively Evaluate Aerosol Process
661 Modules, *Bull. Amer. Meteor. Soc.*, 92, 343-360, <https://doi.org/10.1175/2010bams2868.1>, 2011.

662 Fast, J. D., Berg, L. K., Alexander, L., Bell, D., D'Ambro, E., Hubbe, J., Kuang, C., Liu, J., Long, C.,
663 Matthews, A., Mei, F., Newsom, R., Pekour, M., Pinterich, T., Schmid, B., Schobesberger, S., Shilling, J.,
664 Smith, J. N., Springston, S., Suski, K., Thornton, J. A., Tomlinson, J., Wang, J., Xiao, H., and Zelenyuk,
665 A.: Overview of the HI-SCALE Field Campaign: A New Perspective on Shallow Convective Clouds, *Bull.*
666 *Amer. Meteor. Soc.*, 100, 821-840, <https://doi.org/10.1175/bams-d-18-0030.1>, 2019.

667 Gates, W. L., Boyle, J. S., Covey, C., Dease, C. G., Doutriaux, C. M., Drach, R. S., Fiorino, M., Gleckler,
668 P. J., Hnilo, J. J., Marlais, S. M., Phillips, T. J., Potter, G. L., Santer, B. D., Sperber, K. R., Taylor, K. E.,
669 and Williams, D. N.: An Overview of the Results of the Atmospheric Model Intercomparison Project
670 (AMIP I), *Bull. Amer. Meteor. Soc.*, 80, 29-56, [https://doi.org/10.1175/1520-0477\(1999\)080<0029:Aootro>2.0.Co;2](https://doi.org/10.1175/1520-0477(1999)080<0029:Aootro>2.0.Co;2), 1999.

672 Gelaro, R., McCarty, W., Suarez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C., Darmenov, A.,
673 Bosilovich, M. G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V.,
674 Conaty, A., da Silva, A., Gu, W., Kim, G. K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka,
675 G., Pawson, S., Putman, W., Rienecker, M., Schubert, S. D., Sienkiewicz, M., and Zhao, B.: The Modern-
676 Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2), *J. Climate*, 30, 5419-
677 5454, <https://doi.org/10.1175/JCLI-D-16-0758.1>, 2017.

678 Gleckler, P. J., Doutriaux, C., Durack, P. J., Taylor, K. E., Zhang, Y., Williams, D. N., Mason, E., and
679 Servonnat, J.: A more powerful reality test for climate models, *Eos, Trans. Amer. Geophys. Union*, 97,
680 <https://doi.org/10.1029/2016EO051663>, 2016.

681 Golaz, J.-C., Caldwell, P. M., Van Roekel, L. P., Petersen, M. R., Tang, Q., Wolfe, J. D., Abeshu, G.,
682 Anantharaj, V., Asay-Davis, X. S., Bader, D. C., Baldwin, S. A., Bisht, G., Bogenschutz, P. A., Branstetter,
683 M., Brunke, M. A., Brus, S. R., Burrows, S. M., Cameron-Smith, P. J., Donahue, A. S., Deakin, M., Easter,
684 R. C., Evans, K. J., Feng, Y., Flanner, M., Foucar, J. G., Fyke, J. G., Griffin, B. M., Hannay, C., Harrop, B.
685 E., Hoffman, M. J., Hunke, E. C., Jacob, R. L., Jacobsen, D. W., Jeffery, N., Jones, P. W., Keen, N. D.,
686 Klein, S. A., Larson, V. E., Leung, L. R., Li, H.-Y., Lin, W., Lipscomb, W. H., Ma, P.-L., Mahajan, S.,
687 Maltrud, M. E., Mametjanov, A., McClean, J. L., McCoy, R. B., Neale, R. B., Price, S. F., Qian, Y., Rasch,
688 P. J., Reeves Eyre, J. E. J., Riley, W. J., Ringler, T. D., Roberts, A. F., Roesler, E. L., Salinger, A. G.,
689 Shaheen, Z., Shi, X., Singh, B., Tang, J., Taylor, M. A., Thornton, P. E., Turner, A. K., Veneziani, M.,
690 Wan, H., Wang, H., Wang, S., Williams, D. N., Wolfram, P. J., Worley, P. H., Xie, S., Yang, Y., Yoon, J.-

691 H., Zelinka, M. D., Zender, C. S., Zeng, X., Zhang, C., Zhang, K., Zhang, Y., Zheng, X., Zhou, T., and
 692 Zhu, Q.: The DOE E3SM Coupled Model Version 1: Overview and Evaluation at Standard Resolution, J.
 693 Adv. Model. Earth Syst., 11, 2089-2129, <https://doi.org/10.1029/2018ms001603>, 2019.

694 Hodshire, A. L., Lawler, M. J., Zhao, J., Ortega, J., Jen, C., Yli-Juuti, T., Brewer, J. F., Kodros, J. K.,
 695 Barsanti, K. C., Hanson, D. R., McMurry, P. H., Smith, J. N., and Pierce, J. R.: Multiple new-particle
 696 growth pathways observed at the US DOE Southern Great Plains field site, Atmos. Chem. Phys., 16, 9321-
 697 9348, <https://doi.org/10.5194/acp-16-9321-2016>, 2016.

698 Howie, J. and Kuang, C.: Scanning Mobility Particle Sizer (AOSSMPS) [dataset],
 699 <https://doi.org/10.5439/1476898>, Accessed 14 January 2020.

700 Humphries, R. S., Keywood, M. D., Gribben, S., McRobert, I. M., Ward, J. P., Selleck, P., Taylor, S.,
 701 Harnwell, J., Flynn, C., Kulkarni, G. R., Mace, G. G., Protat, A., Alexander, S. P., and McFarquhar, G.:
 702 Southern Ocean latitudinal gradients of cloud condensation nuclei, Atmos. Chem. Phys., 21, 12757-12782,
 703 <https://doi.org/10.5194/acp-21-12757-2021>, 2021.

704 IPCC, Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y.,
 705 Bex, V., and Midgley, P. M. (Eds.): Climate Change 2013: The Physical Science Basis. Contribution of
 706 Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change,
 707 Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1535 pp.,
 708 <https://doi.org/10.1017/CBO9781107415324>, 2013.

709 Koontz, A. and Kuang, C.: Scanning mobility particle sizer (AOSNANOSMPS) [dataset],
 710 <https://doi.org/10.5439/1242975>, Accessed 14 January 2020.

711 Koontz, A. and Uin, J.: Ultra-High Sensitivity Aerosol Spectrometer (AOSUHSAS) [dataset],
 712 <https://doi.org/10.5439/1333828>, Accessed 3 February 2021.

713 Kuang, C., McMurry, P. H., and McCormick, A. V.: Determination of cloud condensation nuclei production
 714 from measured new particle formation events, Geophys. Res. Lett., 36,
 715 <https://doi.org/10.1029/2009GL037584>, 2009.

716 Kuang, C., Salwen, C., and Singh, A.: Condensation Particle Counter (AOSPCF) [dataset],
 717 <https://doi.org/10.5439/1046184>, Accessed 24 November 2020.

718 Kuang, C., Andrews, E., Salwen, C., and Singh, A.: Condensation Particle Counter (AOSPC) [dataset],
 719 <https://doi.org/10.5439/1025152>, Accessed 16 September 2020.

720 Kuang, C., Salwen, C., Boyer, M., and Singh, A.: Condensation Particle Counter (AOSPCF1M) [dataset],
 721 <https://doi.org/10.5439/1418260>, Accessed 3 February 2021.

722 Kulmala, M., Vehkamäki, H., Petäjä, T., Dal Maso, M., Lauri, A., Kerminen, V. M., Birmili, W., and
 723 McMurry, P. H.: Formation and growth rates of ultrafine atmospheric particles: a review of observations,
 724 J. Aerosol Sci., 35, 143-176, <https://doi.org/10.1016/j.jaerosci.2003.10.003>, 2004.

725 Kyröuac, J. and Shi, Y.: Surface Meteorological Instrumentation (MET) [dataset],
 726 <https://doi.org/10.5439/1786358>, Accessed 11 November 2020.

727 Lewis, E. R. and Teixeira, J.: Dispelling clouds of uncertainty, Eos, Trans. Amer. Geophys. Union, 96,
 728 <https://doi.org/10.1029/2015eo031303>, 2015.

729 Liu, J., Alexander, L., Fast, J. D., Lindenmaier, R., and Shilling, J. E.: Aerosol characteristics at the
 730 Southern Great Plains site during the HI-SCALE campaign, *Atmos. Chem. Phys.*, 21, 5101-5116,
 731 <https://doi.org/10.5194/acp-21-5101-2021>, 2021.

732 Liu, X., Ma, P. L., Wang, H., Tilmes, S., Singh, B., Easter, R. C., Ghan, S. J., and Rasch, P. J.: Description
 733 and evaluation of a new four-mode version of the Modal Aerosol Module (MAM4) within version 5.3 of
 734 the Community Atmosphere Model, *Geosci. Model Dev.*, 9, 505-522, [https://doi.org/10.5194/gmd-9-505-](https://doi.org/10.5194/gmd-9-505-2016)
 735 [2016](https://doi.org/10.5194/gmd-9-505-2016), 2016.

736 Maloney, E. D., Gettelman, A., Ming, Y., Neelin, J. D., Barrie, D., Mariotti, A., Chen, C. C., Coleman, D.
 737 R. B., Kuo, Y.-H., Singh, B., Annamalai, H., Berg, A., Booth, J. F., Camargo, S. J., Dai, A., Gonzalez, A.,
 738 Hafner, J., Jiang, X., Jing, X., Kim, D., Kumar, A., Moon, Y., Naud, C. M., Sobel, A. H., Suzuki, K., Wang,
 739 F., Wang, J., Wing, A. A., Xu, X., and Zhao, M.: Process-Oriented Evaluation of Climate and Weather
 740 Forecasting Models, *Bull. Amer. Meteor. Soc.*, 100, 1665-1686, <https://doi.org/10.1175/bams-d-18-0042.1>,
 741 2019.

742 McCoy, I. L., Bretherton, C. S., Wood, R., Twohy, C. H., Gettelman, A., Bardeen, C. G., and Toohey, D.
 743 W.: Influences of Recent Particle Formation on Southern Ocean Aerosol Variability and Low Cloud
 744 Properties, *J. Geophys. Res. Atmos.*, 126, <https://doi.org/10.1029/2020jd033529>, 2021.

745 McCoy, I. L., McCoy, D. T., Wood, R., Regayre, L., Watson-Parris, D., Grosvenor, D. P., Mulcahy, J. P.,
 746 Hu, Y., Bender, F. A.-M., Field, P. R., Carslaw, K. S., and Gordon, H.: The hemispheric contrast in cloud
 747 microphysical properties constrains aerosol forcing, *Proceedings of the National Academy of Sciences*,
 748 117, 18998-19006, <https://doi.org/doi:10.1073/pnas.1922502117>, 2020.

749 McFarquhar, G. M., Bretherton, C. S., Marchand, R., Protat, A., DeMott, P. J., Alexander, S. P., Roberts,
 750 G. C., Twohy, C. H., Toohey, D., Siems, S., Huang, Y., Wood, R., Rauber, R. M., Lasher-Trapp, S., Jensen,
 751 J., Stith, J. L., Mace, J., Um, J., Järvinen, E., Schnaiter, M., Gettelman, A., Sanchez, K. J., McCluskey, C.
 752 S., Russell, L. M., McCoy, I. L., Atlas, R. L., Bardeen, C. G., Moore, K. A., Hill, T. C. J., Humphries, R.
 753 S., Keywood, M. D., Ristovski, Z., Cravigan, L., Schofield, R., Fairall, C., Mallet, M. D., Kreidenweis, S.
 754 M., Rainwater, B., D'Alessandro, J., Wang, Y., Wu, W., Saliba, G., Levin, E. J. T., Ding, S., Lang, F.,
 755 Truong, S. C. H., Wolff, C., Haggerty, J., Harvey, M. J., Klekociuk, A. R., and McDonald, A.: Observations
 756 of Clouds, Aerosols, Precipitation, and Surface Radiation over the Southern Ocean: An Overview of
 757 CAPRICORN, MARCUS, MICRE, and SOCRATES, *Bull. Amer. Meteor. Soc.*, 102, E894-E928,
 758 <https://doi.org/10.1175/bams-d-20-0132.1>, 2021.

759 Mei, F.: Condensation particle counter for ACEENA [dataset], <https://doi.org/10.5439/1440985>, Accessed
 760 5 December 2020.

761 Müller, D., Hostetler, C. A., Ferrare, R. A., Burton, S. P., Chemyakin, E., Kolgotin, A., Hair, J. W., Cook,
 762 A. L., Harper, D. B., Rogers, R. R., Hare, R. W., Cleckner, C. S., Obland, M. D., Tomlinson, J., Berg, L.
 763 K., and Schmid, B.: Airborne Multiwavelength High Spectral Resolution Lidar (HSRL-2) observations
 764 during TCAP 2012: vertical profiles of optical and microphysical properties of a smoke/urban haze plume
 765 over the northeastern coast of the US, *Atmos. Meas. Tech.*, 7, 3487-3496, [https://doi.org/10.5194/amt-7-](https://doi.org/10.5194/amt-7-3487-2014)
 766 [3487-2014](https://doi.org/10.5194/amt-7-3487-2014), 2014.

767 Myhre, G., Samset, B. H., Schulz, M., Balkanski, Y., Bauer, S., Bernsten, T. K., Bian, H., Bellouin, N.,
 768 Chin, M., Diehl, T., Easter, R. C., Feichter, J., Ghan, S. J., Hauglustaine, D., Iversen, T., Kinne, S.,
 769 Kirkevåg, A., Lamarque, J. F., Lin, G., Liu, X., Lund, M. T., Luo, G., Ma, X., van Noije, T., Penner, J. E.,
 770 Rasch, P. J., Ruiz, A., Seland, Ø., Skeie, R. B., Stier, P., Takemura, T., Tsigaridis, K., Wang, P., Wang, Z.,
 771 Xu, L., Yu, H., Yu, F., Yoon, J. H., Zhang, K., Zhang, H., and Zhou, C.: Radiative forcing of the direct

772 aerosol effect from AeroCom Phase II simulations, *Atmos. Chem. Phys.*, 13, 1853-1877,
773 <https://doi.org/10.5194/acp-13-1853-2013>, 2013.

774 NETCDF: Introduction and Overview: <https://www.unidata.ucar.edu/software/netcdf/docs/index.html>, last
775 access: 2 November 2021. 2021.

776 Petters, M. D. and Kreidenweis, S. M.: A single parameter representation of hygroscopic growth and cloud
777 condensation nucleus activity, *Atmos. Chem. Phys.*, 7, 1961-1971, [https://doi.org/10.5194/acp-7-1961-](https://doi.org/10.5194/acp-7-1961-2007)
778 [2007](https://doi.org/10.5194/acp-7-1961-2007), 2007.

779 Pierce, J. R. and Adams, P. J.: Uncertainty in global CCN concentrations from uncertain aerosol nucleation
780 and primary emission rates, *Atmos. Chem. Phys.*, 9, 1339-1356, <https://doi.org/10.5194/acp-9-1339-2009>,
781 2009.

782 Rasch, P. J., Xie, S., Ma, P.-L., Lin, W., Wang, H., Tang, Q., Burrows, S. M., Caldwell, P., Zhang, K.,
783 Easter, R. C., Cameron-Smith, P., Singh, B., Wan, H., Golaz, J.-C., Harrop, B. E., Roesler, E., Bacmeister,
784 J., Larson, V. E., Evans, K. J., Qian, Y., Taylor, M., Leung, L. R., Zhang, Y., Brent, L., Branstetter, M.,
785 Hannay, C., Mahajan, S., Mamatjanov, A., Neale, R., Richter, J. H., Yoon, J.-H., Zender, C. S., Bader, D.,
786 Flanner, M., Foucar, J. G., Jacob, R., Keen, N., Klein, S. A., Liu, X., Salinger, A. G., Shrivastava, M., and
787 Yang, Y.: An Overview of the Atmospheric Component of the Energy Exascale Earth System Model, *J.*
788 *Adv. Model. Earth Syst.*, 11, 2377-2411, <https://doi.org/10.1029/2019ms001629>, 2019.

789 Reddington, C. L., Carslaw, K. S., Stier, P., Schutgens, N., Coe, H., Liu, D., Allan, J., Browse, J., Pringle,
790 K. J., Lee, L. A., Yoshioka, M., Johnson, J. S., Regayre, L. A., Spracklen, D. V., Mann, G. W., Clarke, A.,
791 Hermann, M., Henning, S., Wex, H., Kristensen, T. B., Leaitch, W. R., Pöschl, U., Rose, D., Andreae, M.
792 O., Schmale, J., Kondo, Y., Oshima, N., Schwarz, J. P., Nenes, A., Anderson, B., Roberts, G. C., Snider, J.
793 R., Leck, C., Quinn, P. K., Chi, X., Ding, A., Jimenez, J. L., and Zhang, Q.: The Global Aerosol Synthesis
794 and Science Project (GASSP): Measurements and Modeling to Reduce Uncertainty, *Bull. Amer. Meteor.*
795 *Soc.*, 98, 1857-1877, <https://doi.org/10.1175/bams-d-15-00317.1>, 2017.

796 Redemann, J., Wood, R., Zuidema, P., Doherty, S. J., Luna, B., LeBlanc, S. E., Diamond, M. S., Shinozuka,
797 Y., Chang, I. Y., Ueyama, R., Pfister, L., Ryoo, J. M., Dobracki, A. N., da Silva, A. M., Longo, K. M.,
798 Kacenelenbogen, M. S., Flynn, C. J., Pistone, K., Knox, N. M., Piketh, S. J., Haywood, J. M., Formenti, P.,
799 Mallet, M., Stier, P., Ackerman, A. S., Bauer, S. E., Fridlind, A. M., Carmichael, G. R., Saide, P. E., Ferrada,
800 G. A., Howell, S. G., Freitag, S., Cairns, B., Holben, B. N., Knobelspiesse, K. D., Tanelli, S., L'Ecuyer, T.
801 S., Dzambo, A. M., Sy, O. O., McFarquhar, G. M., Poellot, M. R., Gupta, S., O'Brien, J. R., Nenes, A.,
802 Kacarab, M., Wong, J. P. S., Small-Griswold, J. D., Thornhill, K. L., Noone, D., Podolske, J. R., Schmidt,
803 K. S., Pilewskie, P., Chen, H., Cochrane, S. P., Sedlacek, A. J., Lang, T. J., Stith, E., Segal-Rozenhaimer,
804 M., Ferrare, R. A., Burton, S. P., Hostetler, C. A., Diner, D. J., Seidel, F. C., Platnick, S. E., Myers, J. S.,
805 Meyer, K. G., Spangenberg, D. A., Maring, H., and Gao, L.: An overview of the ORACLES (ObseRvations
806 of Aerosols above CLouds and their interActionS) project: aerosol–cloud–radiation interactions in the
807 southeast Atlantic basin, *Atmos. Chem. Phys.*, 21, 1507-1563, <https://doi.org/10.5194/acp-21-1507-2021>,
808 2021.

809 Sanchez, K. J., Roberts, G. C., Saliba, G., Russell, L. M., Twohy, C., Reeves, J. M., Humphries, R. S.,
810 Keywood, M. D., Ward, J. P., and McRobert, I. M.: Measurement report: Cloud processes and the transport
811 of biological emissions affect southern ocean particle and cloud condensation nuclei concentrations, *Atmos.*
812 *Chem. Phys.*, 21, 3427-3446, <https://doi.org/10.5194/acp-21-3427-2021>, 2021.

813 Schulz, M., Textor, C., Kinne, S., Balkanski, Y., Bauer, S., Bernsten, T., Berglen, T., Boucher, O.,
814 Dentener, F., Guibert, S., Isaksen, I. S. A., Iversen, T., Koch, D., Kirkevåg, A., Liu, X., Montanaro, V.,

815 Myhre, G., Penner, J. E., Pitari, G., Reddy, S., Seland, Ø., Stier, P., and Takemura, T.: Radiative forcing by
816 aerosols as derived from the AeroCom present-day and pre-industrial simulations, *Atmos. Chem. Phys.*, 6,
817 5225-5246, <https://doi.org/10.5194/acp-6-5225-2006>, 2006.

818 Seinfeld, J. H., Bretherton, C., Carslaw, K. S., Coe, H., DeMott, P. J., Dunlea, E. J., Feingold, G., Ghan, S.,
819 Guenther, A. B., Kahn, R., Kraucunas, I., Kreidenweis, S. M., Molina, M. J., Nenes, A., Penner, J. E.,
820 Prather, K. A., Ramanathan, V., Ramaswamy, V., Rasch, P. J., Ravishankara, A. R., Rosenfeld, D.,
821 Stephens, G., and Wood, R.: Improving our fundamental understanding of the role of aerosol–cloud
822 interactions in the climate system, *Proc. Natl. Acad. Sci. U.S.A.*, 113, 5781-5790,
823 <https://doi.org/10.1073/pnas.1514043113>, 2016.

824 Sun, J., Zhang, K., Wan, H., Ma, P.-L., Tang, Q., and Zhang, S.: Impact of Nudging Strategy on the Climate
825 Representativeness and Hindcast Skill of Constrained EAMv1 Simulations, *J. Adv. Model. Earth Syst.*, 11,
826 3911-3933, <https://doi.org/10.1029/2019MS001831>, 2019.

827 Tang, Q., Klein, S. A., Xie, S., Lin, W., Golaz, J. C., Roesler, E. L., Taylor, M. A., Rasch, P. J., Bader, D.
828 C., Berg, L. K., Caldwell, P., Giangrande, S. E., Neale, R. B., Qian, Y., Riihimäki, L. D., Zender, C. S.,
829 Zhang, Y., and Zheng, X.: Regionally refined test bed in E3SM atmosphere model version 1 (EAMv1) and
830 applications for high-resolution modeling, *Geosci. Model Dev.*, 12, 2679-2706,
831 <https://doi.org/10.5194/gmd-12-2679-2019>, 2019.

832 Teixeira, J., Cardoso, S., Bonazzola, M., Cole, J., DelGenio, A., DeMott, C., Franklin, C., Hannay, C.,
833 Jakob, C., Jiao, Y., Karlsson, J., Kitagawa, H., Köhler, M., Kuwano-Yoshida, A., LeDrian, C., Li, J., Lock,
834 A., Miller, M. J., Marquet, P., Martins, J., Mechoso, C. R., Meijgaard, E. v., Meinke, I., Miranda, P. M. A.,
835 Mironov, D., Neggers, R., Pan, H. L., Randall, D. A., Rasch, P. J., Rockel, B., Rossow, W. B., Ritter, B.,
836 Siebesma, A. P., Soares, P. M. M., Turk, F. J., Vaillancourt, P. A., Von Engel, A., and Zhao, M.: Tropical
837 and Subtropical Cloud Transitions in Weather and Climate Prediction Models: The GCSS/WGNE Pacific
838 Cross-Section Intercomparison (GPCI), *J. Climate*, 24, 5223-5256, <https://doi.org/10.1175/2011jcli3672.1>,
839 2011.

840 Twomey, S.: The Influence of Pollution on the Shortwave Albedo of Clouds, *J. Atmos. Sci.*, 34, 1149-1152,
841 [https://doi.org/10.1175/1520-0469\(1977\)034<1149:TIOPOT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1977)034<1149:TIOPOT>2.0.CO;2), 1977.

842 Uin, J., Senum, G., Koontz, A., and Flynn, C.: Ultra-High Sensitivity Aerosol Spectrometer (AOSUHSAS)
843 [dataset], <https://doi.org/10.5439/1409033>, Accessed 24 November 2020.

844 Wang, H., Easter, R. C., Zhang, R., Ma, P.-L., Singh, B., Zhang, K., Ganguly, D., Rasch, P. J., Burrows, S.
845 M., Ghan, S. J., Lou, S., Qian, Y., Yang, Y., Feng, Y., Flanner, M., Leung, R. L., Liu, X., Shrivastava, M.,
846 Sun, J., Tang, Q., Xie, S., and Yoon, J.-H.: Aerosols in the E3SM Version 1: New Developments and Their
847 Impacts on Radiative Forcing, *J. Adv. Model. Earth Syst.*, 12, e2019MS001851,
848 <https://doi.org/10.1029/2019ms001851>, 2020.

849 Wang, J., Wood, R., Jensen, M. P., Chiu, J. C., Liu, Y., Lamer, K., Desai, N., Giangrande, S. E., Knopf, D.
850 A., Kollias, P., Laskin, A., Liu, X., Lu, C., Mechem, D., Mei, F., Starzec, M., Tomlinson, J., Wang, Y.,
851 Yum, S. S., Zheng, G., Aiken, A. C., Azevedo, E. B., Blanchard, Y., China, S., Dong, X., Gallo, F., Gao,
852 S., Ghate, V. P., Glienke, S., Goldberger, L., Hardin, J. C., Kuang, C., Luke, E. P., Matthews, A. A., Miller,
853 M. A., Moffet, R., Pekour, M., Schmid, B., Sedlacek, A. J., Shaw, R. A., Shilling, J. E., Sullivan, A., Suski,
854 K., Veghte, D. P., Weber, R., Wyant, M., Yeom, J., Zawadowicz, M., and Zhang, Z.: Aerosol and Cloud
855 Experiments in the Eastern North Atlantic (ACE-ENA), *Bull. Amer. Meteor. Soc.*, 1-51,
856 <https://doi.org/10.1175/bams-d-19-0220.1>, 2021.

857 Watson-Parris, D., Schutgens, N., Reddington, C., Pringle, K. J., Liu, D., Allan, J. D., Coe, H., Carslaw, K.
858 S., and Stier, P.: In situ constraints on the vertical distribution of global aerosol, *Atmos. Chem. Phys.*, 19,
859 11765-11790, <https://doi.org/10.5194/acp-19-11765-2019>, 2019.

860 Winker, D. M., Vaughan, M. A., Omar, A., Hu, Y., Powell, K. A., Liu, Z., Hunt, W. H., and Young, S. A.:
861 Overview of the CALIPSO Mission and CALIOP Data Processing Algorithms, *J. Atmos. Ocean. Technol.*,
862 26, 2310-2323, <https://doi.org/10.1175/2009jtecha1281.1>, 2009.

863 Zaveri, R. A., Easter, R. C., Singh, B., Wang, H., Lu, Z., Tilmes, S., Emmons, L. K., Vitt, F., Zhang, R.,
864 Liu, X., Ghan, S. J., and Rasch, P. J.: Development and Evaluation of Chemistry-Aerosol-Climate Model
865 CAM5-Chem-MAM7-MOSAIC: Global Atmospheric Distribution and Radiative Effects of Nitrate
866 Aerosol, *J. Adv. Model. Earth Syst.*, 13, e2020MS002346, <https://doi.org/10.1029/2020MS002346>, 2021.

867 Zawadowicz, M. A., Suski, K., Liu, J., Pekour, M., Fast, J., Mei, F., Sedlacek, A. J., Springston, S., Wang,
868 Y., Zaveri, R. A., Wood, R., Wang, J., and Shilling, J. E.: Aircraft measurements of aerosol and trace gas
869 chemistry in the eastern North Atlantic, *Atmos. Chem. Phys.*, 21, 7983-8002, [https://doi.org/10.5194/acp-](https://doi.org/10.5194/acp-21-7983-2021)
870 [21-7983-2021](https://doi.org/10.5194/acp-21-7983-2021), 2021.

871 Zhang, C., Xie, S., Tao, C., Tang, S., Emmenegger, T., Neelin, J. D., Schiro, K. A., Lin, W., and Shaheen,
872 Z.: The ARM Data-Oriented Metrics and Diagnostics Package for Climate Models: A New Tool for
873 Evaluating Climate Models with Field Data, *Bull. Amer. Meteor. Soc.*, 101, E1619-E1627,
874 <https://doi.org/10.1175/bams-d-19-0282.1>, 2020.

875 Zhang, K., Wan, H., Liu, X., Ghan, S. J., Kooperman, G. J., Ma, P. L., Rasch, P. J., Neubauer, D., and
876 Lohmann, U.: Technical Note: On the use of nudging for aerosol-climate model intercomparison studies,
877 *Atmos. Chem. Phys.*, 14, 8631-8645, <https://doi.org/10.5194/acp-14-8631-2014>, 2014.

878 Zhao, B., Shrivastava, M., Donahue, N. M., Gordon, H., Schervish, M., Shilling, J. E., Zaveri, R. A., Wang,
879 J., Andreae, M. O., Zhao, C., Gaudet, B., Liu, Y., Fan, J., and Fast, J. D.: High concentration of ultrafine
880 particles in the Amazon free troposphere produced by organic new particle formation, *Proc. Natl. Acad.*
881 *Sci. U. S. A.*, 117, 25344-25351, <https://doi.org/10.1073/pnas.2006716117>, 2020.

882 Zheng, G., Wang, Y., Wood, R., Jensen, M. P., Kuang, C., McCoy, I. L., Matthews, A., Mei, F., Tomlinson,
883 J. M., Shilling, J. E., Zawadowicz, M. A., Crosbie, E., Moore, R., Ziemba, L., Andreae, M. O., and Wang,
884 J.: New particle formation in the remote marine boundary layer, *Nat. Commun.*, 12, 527,
885 <https://doi.org/10.1038/s41467-020-20773-1>, 2021.

886 Zhou, X., Kollias, P., and Lewis, E. R.: Clouds, Precipitation, and Marine Boundary Layer Structure during
887 the MAGIC Field Campaign, *J. Climate*, 28, 2420-2442, <https://doi.org/10.1175/jcli-d-14-00320.1>, 2015.

888 Zuidema, P., Alvarado, M., Chiu, C., de Szoeko, S., Fairall, C., Feingold, G., Freedman, A., Ghan, S.,
889 Haywood, J., Kollias, P., Lewis, E., McFarquhar, G., McComiskey, A., Mechem, D., Onasch, T.,
890 Redemann, J., Romps, D., Turner, D., Wang, H., Wood, R., Yuter, S., and Zhu, P.: Layered Atlantic Smoke
891 Interactions with Clouds (LASIC) Field Campaign Report, U.S. Department of Energy, Technical Report,
892 DOE/SC-ARM-18-018, 2018.

893