# Simulation, Precursor Analysis and Targeted Observation Sensitive Area Identification for Two Types of ENSO using ENSO-MC v1.0

Bin Mu<sup>1,\*</sup>, Yuehan Cui<sup>1,\*</sup>, Shijin Yuan<sup>1</sup>, and Bo Qin<sup>1</sup>

<sup>1</sup>School of Software Engineering, Tongji University, Shanghai, China

\*These authors contributed equally to this work.

Correspondence: Shijin Yuan (e-mail: yuanshijin2003@163.com)

**Abstract.** The global impact of an El Niño-Southern Oscillation (ENSO) event can differ greatly depending on whether it is an Eastern-Pacific-type (EP-type) event or a Central-Pacific-type (CP-type) event. Reliable predictions of the two types of ENSO are therefore of critical importance. Here we construct a deep neural network with multichannel structure for ENSO (named ENSO-MC) to simulate the spatial evolution of sea surface temperature (SST) anomalies for the two types of events.

- 5 We select SST, heat content, and wind stress (i.e., three key ingredients of Bjerknes feedback) to represent coupled oceanatmosphere dynamics that underpins ENSO, achieving skillful forecasts for the spatial patterns of SST anomalies out to one year ahead. Furthermore, it is of great significance to analyze the precursors of EP-type or CP-type events and identify targeted observation sensitive area for the understanding and prediction of ENSO. Precursors analysis is to determine what type of initial perturbations will develop into EP-type or CP-type events. Sensitive area identification is to determine the regions where
- 10 initial states tend to have greatest impacts on evolution of ENSO. We use saliency map method to investigate the subsurface precursors and identify the sensitive areas of ENSO. The results show that there are pronounced signals in the equatorial subsurface before EP events, while the precursory signals of CP events are located in the North Pacific. It indicates that the subtropical precursors seem to favor the generation of the CP-type El Niño and the EP-type El Niño is more related to the tropical thermocline dynamics. And the saliency maps show that the sensitive areas of the surface and the subsurface are
- 15 located in the equatorial central Pacific and the equatorial western Pacific, respectively. The sensitivity experiments imply that additional observations in the identified sensitive areas can improve forecasting skills. Our results of precursors and sensitive areas are consistent with the previous theories of ENSO, demonstrating the potential usage and advantages of the ENSO-MC model in improving the simulation, understanding and observations of two ENSO types.

# 1 Introduction

20 El Niño-Southern Oscillation (ENSO) is an irregular climate signal with a period of 2-7 years in the tropical Pacific Ocean and often grows up to be exceptionally strong under unstable air-sea interactions (Bjerknes, 1969; Philander, 1983), causing large global climatic anomalies and hence affecting many regions even far from the tropical area (Yu et al., 2012). Studies suggested that each ENSO event may differ in spatial structure, temporal evolution, amplitude and trigger (Capotondi et al., 2015; Timmermann et al., 2018). One view is that there may be two different types of ENSO, referred to as the Eastern-

- 25 Pacific-type (EP-type) event and the Central-Pacific-type (CP-type) event (Yu and Kao, 2007; Kao and Yu, 2009). And the differences in the details of sea surface temperature (SST) anomaly patterns between EP and CP events will lead to different remote teleconnection patterns and effects on the global climate (An et al., 2007; Ashok et al., 2007; Timmermann et al., 2018). In recent decades, with the increased occurrence of CP El Niño relative to EP El Niño, the predictability of two ENSO types has attracted widespread attentions (Lee and McPhaden, 2010). Tao et al. (2020) used the nonlinear forcing singular
- 30 vector (NFSV)-tendency assimilation approach to improve ENSO model and showed the ability of recognizing the types of El Niño at least six months in advance in predictions (Lingjiang and Wansuo, 2019). Tian and Duan (2016) demonstrated that the spring predictability barrier is weaker in CP-El Niño than in EP-El Niño when model error effects can be negligible. Improved forecasting and understanding of the two types of ENSO are therefore of great importance.
- Most studies on the simulations of two types of ENSO are based on the climate numerical models. Kug et al. (2010) used Geophysical Fluid Dynamics Laboratory Coupled Model (GFDL) to simulate the CP-type El Niño, which shows distinct spatial characteristics and dynamic processes from the EP-type El Niño. More comprehensively, Kug et al. used the climate models from the Coupled Model Intercomparison Project phase-3 (CMIP3) (Ham and Kug, 2012) and phase-5 (Kug et al., 2012) to validate the fidelity in simulating the two types of events. The results showed that a few models can simulate the two types of El Niño, and most of models tend to simulate a single type. Duan et al. (2014) proposed an optimal forcing vector (OFV) approach
- 40 to optimize the Zebiak–Cane model and reproduced several observed EP and CP events, and revealed the dominant roles of zonal advection process in the development of CP-El Niño; then Duan et al. (2017) first demonstrated that the diversity of El Niño is closely related to changes in the nonlinear characteristics of the tropical Pacific. Accurate simulations and predictions of two types of ENSO are still of a great challenge, owing to the inherent uncertainty and diversity of ENSO (Chen and Cane, 2008; Trenberth and Stepaniak, 2001; Capotondi et al., 2015).
- In the past two years, deep learning methods have paved a new and profound way to making accurate ENSO forecasts for long lead times (Huang et al., 2019; Ham et al., 2019). For example, Ham et al. (2019) used convolutional neural network (CNN) together with transfer learning method to produce higher skills in predicting the Niño 3.4 index than current dynamical and statistical models at lead times of up to one and a half years. Yan et al. (2020) used the temporal convolutional network and empirical mode decomposition to predict each subcomponents of Niño3.4 index that were then reconstructed to improve
- 50 the forecasting skills of total values. In addition to the Nino index forecasting that most models are currently focused on, deep neural networks also show great potential for a wide range of application for the pattern predictions (Mu et al., 2019, 2021). Here we develop a spatiotemporal model of multichannel structure for ENSO (named ENSO-MC) to simulate the spatial diversity and evolution of SST anomalies patterns in the equatorial Pacific. The multichannel structure containing the complex ocean-atmosphere interactions is built to achieve skillful predictions of two types of ENSO one year in advance.
- <sup>55</sup> In addition to developing forecast models, understandings and observations of ENSO are also of great significance for prediction improvement, which are two basic issues in the predictability of ENSO. In order to better understand the mechanism of ENSO occurrence, one approach is to explore the precursor of ENSO, which is the initial perturbation distribution that is most likely to develop into a CP event or an EP event. Duan et al. (2004) is one of the earliest papers that explored the precursory disturbance of ENSO events (Duan et al., 2013). These precursors help us understand the dynamic process of ENSO and provide

60 the potential to predict ENSO events and their types. In terms of observations, owing to the limited sampling frequency in time and sampling density in space of the current observation systems, especially ocean observation systems, intensive observations are usually prioritized in sensitive areas. Such a strategy is called targeted observation method. The key issue in targeted observation is the identification of the sensitive areas. The initial conditions in these sensitive areas may be more important than those in other regions when predicting ENSO (Mu et al., 2015). Due to the high cost of observation over the ocean, it is a

65

cost-effective method which can help reduce initial errors, thereby reducing prediction errors and improving prediction skills. Precursor investigation and sensitive area identification based on numerical models and optimal perturbations, such as the linear singular vector (LSV) approach (Moore and Kleeman, 1996), the linear inverse modeling (LIM) approach (Newman et al., 2011; Vimont et al., 2014) and the conditional nonlinear optimal perturbation (CNOP) technique (Mu et al., 2003), have been applied extensively and produced meaningful results. For example, Capotondi and Sardeshmukh (2015) suggested that

- 70 the initial subsurface conditions play an important distinguishing role in the generation of different ENSO types. And recent research has also recognized the critical role of some climate patterns outside the tropical Pacific (Vimont et al., 2003; Ham et al., 2013; Chikamoto et al., 2015) that precede ENSO. Based on the method of finding the optimal initial perturbation, several studies have linked the precursor analysis with the targeted observation of ENSO events (Mu et al., 2014; Hu and Duan, 2016). Hu and Duan (2016) identified the western equatorial Pacific of the subsurface and the eastern equatorial Pacific of
- 75 the surface as sensitive areas using CNOP method based on the Community Earth System Model (CESM). And it showed that eliminating the initial errors in these sensitive areas can greatly improve ENSO prediction. Despite numerous attempts for precursory signals investigation and sensitive areas identification of ENSO, the results may vary owing to the complexity of the models. The research on improved observation, understanding and forecasting of ENSO has therefore always received critical attention.
- In this paper, based on the model ENSO-MC we proposed above that can simulate spatial patterns of SST anomalies, we further analyze the subsurface precursors of different types of events and identify the sensitive areas with the help of saliency map interpretability method (Simonyan et al., 2013; Zeiler and Fergus, 2014). The obtained saliency map answers the question "which input pixels should be changed to yield a maximal increase in the considered output value with minimal change?" (Ebert-Uphoff and Hilburn, 2020). It indicates the sensitivity of the predicted results to the perturbations in each region of
- 85 the input field and can be used to discover the initial perturbation distribution that would develop into an ENSO event, which reveals the signals prior to the events captured by the ENSO-MC. Besides, the sensitive areas are the regions in which the small perturbations would have the greatest influence on the forecasts. Therefore, saliency map method can also help identify the sensitive areas. Since the original saliency map method is prone to noise, the smoothGrad method (Smilkov et al., 2017) is used here to help sharpen the saliency maps. Sensitivity experiments are then performed to verify the effectiveness of the
- 90 identified sensitive area. The results are consistent with the previous understanding of ENSO mechanism. Our results suggest that the ENSO-MC model may provide a powerful and promising method for simulating the seasonal-to-interannual variations of ENSO and analyzing the inherent predictability of ENSO.

The remainder of the paper is structured as follows. Section 2 describes the ENSO-MC model for simulating two types of ENSO, including the multichannel spatiotemporal prediction neural network, detailed descriptions of the predictor selection

95 and combined loss function. And we discuss the effect of each component on the model performance in Section 3. Section 4 provides the assessment of the ENSO pattern simulation performance based on the ENSO-MC model. Then we analyze the precursors in heat content for two types El Niño events and La Niña events in Section 5, and identify the sensitive areas of targeted observation for ENSO in Section 6. Summary and discussion are presented in Section 7.

### 2 ENSO-MC: Simulation model for two types of ENSO

# 100 2.1 Multichannel spatiotemporal structure

105

Here we develop a spatiotemporal model of multichannel structure named ENSO-MC to generate SST pattern sequence for ENSO forecasts. As shown in Fig. 1, the ENSO-MC is constructed based on the encoder-decoder architecture (Sutskever et al., 2014), whose encoder extracts the feature representations associated with ENSO over the past period and decoder generates the sea surface temperature pattern in the future. Due to the diversity of ENSO in amplitude, spatial pattern and temporal evolution, several convolutional long short-term memory (ConvLSTM) layers (Xingjian et al., 2015) form the skeleton in the encoder-decoder architecture to learn its multiple spatial and temporal representations. The encoder is the first half of the architecture (Fig. 1). A ConvLSTM layer with kernel size 3×3 followed by a 3D max-pooling layer constitutes an encoding module. The max-pooling layer downsamples the input along the spatial dimensions to extract multi-scale spatial connections. We use three encoding modules to construct the encoder, which is the network depth that perform best in ENSO forecasting problem here.

- 110 To balance model performance and computational cost, the output channels of ConvLSTM in the three modules are 8, 16, 32 in order. After three encoding modules, we use a convolution layer with kernel size 5×5 and stride 5×5, and the number of output channels is 64. The dimension of feature map output by each layer is shown in Fig. 1, and the final feature dimension of the encoder is 2×4×64. The structure of the decoder is symmetrical with the encoder. After a transposed convolution layer, there are three decoding modules. Each module consists of an upsampling layer with size 2 followed by a ConvLSTM layer
- 115 to restore the original resolution of SST pattern, where the kernel size of the network and the number of output channels are the same as those in the encoder. And the final layer in the ENSO-MC model is an additional 3×3 ConvLSTM that generates a single feature map representing the SST pattern sequence predicted by the model.

In order to preserve oceanic processes information for a long time for ENSO forecast, we add skip-layer connection and states connection between the encoder and the decoder on different spatial scales. For each ConvLSTM layer in the encoder,

120 the feature maps of all time steps are fused into one feature map and the weight of each time step is automatically determined through the attention mechanism. These fused feature maps are attached to the corresponding ConvLSTM layers in the decoder to achieve skip-layer connection. And in the states connection, the hidden states output by the ConvLSTM layers in the encoder are reserved for the corresponding layer when the decoder is initialized.

In addition, ENSO prediction requires a multiple-step forecasting strategy to achieve long-term prediction. There are two main strategies, direct multi-step forecast and one-step ahead forecast. As shown in Fig. 2, the inputs for direct multi-step forecast are fixed observations  $X_{T-m}, \ldots, X_{T-1}$ . To achieve multi-step prediction, one of the methods is to build a separate model  $G_{T+x}$  for each prediction time step T + x. In the case of predicting SST for the next *n* months, *n* models  $G_{T+1}, \ldots, G_{T+n}$ 



Figure 1. The encoder-decoder architecture of ENSO-MC for SST pattern sequence prediction. The encoder contains three ConvLSTM layers and each layer is followed by a pooling layer, and the last layer is a convolutional layer, which allows extracting spatial and temporal features. And the decoder comprises one deconvolutional layer, three ConvLSTM layers and three upsampling layer that restores the features to the same size as the initial spatial dimension (80×160). The model uses skip-connection with attention mechanism and state-connection between the encoder and the decoder to improve forecasting skills. The input variables are the SST, heat content, zonal wind and meridional wind for  $T_{in}$  consecutive months over  $40^{\circ}N - 40^{\circ}S$ ,  $120^{\circ}E - 80^{\circ}W$  (80×160,  $1^{\circ} \times 1^{\circ}$  resolution) and each type of variable is input to the model as a channel. The output of the model is SST pattern sequence for the next  $T_{out}$  months.

need to be constructed. This is also the strategy used in Ham's paper (Ham et al., 2019), which produced skillful prediction results one and a half years in advance. Another approach is to build a model that can forecast the entire prediction sequence
130 Y<sub>T+1</sub>,...,Y<sub>T+n</sub> in a one-shot manner to achieve direct multi-step forecasting, which has the advantage of significantly re-

- ducing computational and maintenance costs. While one-step ahead forecast strategy refers to the multiple use of a one-step model, in which the prediction of the previous time step is used as the input for the prediction of the next time step, i.e., recursive multi-step forecast. In general, one-step ahead forecasting models are more stable and easier to train (Shi and Yeung, 2018). However, because predictions are used instead of observations, the one-step ahead strategy allows prediction errors to
- 135 accumulate, resulting in a rapid decline in performance as the prediction time increases, while direct multi-step forecasting has more accurate results in long-term prediction (Taieb et al., 2012). In this paper, direct multi-step forecast strategy and one-step



Figure 2. Two main strategies of multi-step forecasting for ENSO prediction, direct multi-step forecast strategy and one-step ahead forecast strategy. And direct multi-step forecast strategy has two methods, the second one is used in this paper.

forecast strategy are both used for prediction and comparison. Considering the computational cost, we use the second direct forecast method instead of the first method of constructing several individual models.

# 2.2 Selection of physical variables

- 140 Using deep neural networks for ENSO simulation is essentially a data-driven method (Reichstein et al., 2019) that is good at mining complicated relations hidden in multidimensional observations of the climate system. Therefore, in addition to building a suitable network to well fit the data, it is also important to choose appropriate predictors that well represent ENSO physical processes to train the model (Reichstein et al., 2019).
- First, SST is selected as one of the predictors since it is a source of ENSO predictability and a direct reflection of the occurrence of ENSO event. As the slow-evolving thermal anomaly in the subsurface ocean that provides a key long-lasting memory for ENSO prediction (e.g., Zhang and Levitus, 1997; Tang et al., 2018), we then choose heat content (vertically averaged oceanic temperature in the upper 300 m) as the second attribute , which is closely related to the recharge-discharge oscillator point of view (Jin, 1997a, b; Meinen and McPhaden, 2000; McPhaden, 2003). Third, the westerly wind burst is believed to be an important trigger for El Niño events (Gebbie et al., 2007; Hu et al., 2014; Menkes et al., 2014; Chen et al.,
- 150 2015; Fedorov et al., 2015) and the atmospheric noise from the wind can be a limit for ENSO predictability (Latif et al., 1988; Moore and Kleeman, 1999). We therefore select SST, heat content, and wind stress, in accordance with Bjerknes feedback, to simulate the air-sea interactions responsible for ENSO.

We utilize Simple Ocean Data Assimilation (SODA) reanalysis data set consisting of sea surface temperature, heat content and wind stress gridded variables from 1871 to 2008 to train the ENSO-MC model, with the resolution  $1^{\circ} \times 1^{\circ}$ . The domain over

 $40^{\circ}N - 40^{\circ}S$ ,  $120^{\circ}E - 80^{\circ}W$  is utilized. To avoid possible overfitting, the data set is divided into training set and validation set according to the ratio of 4 to 1. Then we test the performance of model using NCEP Global Ocean Data Assimilation System (GODAS) and ERA-Interim data (2010-2019), where remove the data that is already in the training set (1981-2008), and there is a one-year gap between training set and test set to reduce the possible influence of oceanic memory.

## 2.3 Loss function

170

160 In our experiments, we combine the losses based on Mean Squared Error (MSE), Structural Similarity Index (SSIM) (Wang et al., 2002) and Gradient Difference Loss (GDL) (Mathieu et al., 2015):

$$\mathcal{L} = \lambda_{mse} \mathcal{L}_{mse}(Y, \hat{Y}) + \lambda_{ssim} \mathcal{L}_{ssim}(Y, \hat{Y}) + \lambda_{gdl} \mathcal{L}_{gdl}(Y, \hat{Y}).$$
(1)

 $Y(\hat{Y})$  denotes the observed (predicted) SST pattern sequence.  $\lambda$  represents the weight of each loss. Specifically, MSE measures the discrepancy of each pixel in the sea surface temperature field:

165 
$$\mathcal{L}_{mse}(Y, \hat{Y}) = \frac{1}{T} \sum_{t=1}^{T} ||Y_t - \hat{Y}_t||^2,$$
 (2)

where T represents the length of the prediction sequence. In addition to quantifying difference in each corresponding pixel value between the observations and predictions, we introduce a loss based on SSIM to measure the global structural differences. SSIM is widely used as a metric to measure the similarity of two images by extracting structural information. It takes into account three features: luminance (l), contrast (c) and structure (s), and its metric formula is the product of these three elements.

$$SSIM(x,y) = \left(\frac{2\mu_x\mu_y + C_1}{\mu_x^2 + \mu_y^2 + C_1}\right)_l \cdot \left(\frac{2\sigma_x\sigma_y + C_2}{\sigma_x^2 + \sigma_y^2 + C_2}\right)_c \cdot \left(\frac{\sigma_{xy} + C_3}{\sigma_x\sigma_y + C_3}\right)_s = \frac{(2\mu_x\mu_y + C_1)(2\sigma_{xy} + C_2)}{(\mu_x^2 + \mu_y^2 + C_1)(\sigma_x^2 + \sigma_y^2 + C_2)},\tag{3}$$

where μ is the mean value of a field (luminance), σ is the standard deviation (contrast) and σ<sub>xy</sub> is the covariance of the two fields (structure). C<sub>1</sub>, C<sub>2</sub>, C<sub>3</sub> are constants used to maintain the calculations stable. ENSO is associated with the interannual variations of SST anomalies in the tropical Pacific. And the Nino3.4 index is one of the most commonly used ENSO indicators, which
is the average SST anomalies in the equatorial Pacific sub-region where the maximum variance of SST is located. Therefore, SSIM metric can help evaluate important signals embedded in the SST patterns for ENSO prediction (Mo et al., 2014). The range of SSIM is from 0 to 1, and when two fields are the same, the value of SSIM is 1. Therefore, we construct the SSIM-based loss function as

$$\mathcal{L}_{ssim}(Y, \hat{Y}) = \frac{1 - \frac{1}{T} (\sum_{t=1}^{T} SSIM(Y_t, \hat{Y}_t))}{2}.$$
(4)

- 180 We also consider gradient information in the loss functions. The SST gradient represents the difference in the sea temperature across the adjacent area. Previous studies have shown that MSE loss function tends to average the values of all points in the whole prediction field to minimize the MSE error, while considering the gradient difference value can alleviate this problem(Oprea et al., 2020). Besides, the SST gradient also plays a role in the atmospheric circulation. The region with a large SST gradient will generate stronger winds, which in turn promote the further increase of the SST gradient (Bjerknes, 1969).
  185 As ENSO approaches maturity, the SST gradient increases gradually. Therefore, we use the GDL to measure the gradient
- difference of the surface sea temperature field:

$$\mathcal{L}_{gdl}(Y,\hat{Y}) = \frac{1}{T} \sum_{t=1}^{T} \sum_{i,j} \left| |Y_{i,j}^t - Y_{i-1,j}^t| - |\hat{Y}_{i,j}^t - \hat{Y}_{i-1,j}^t| \right|^2 + \left| |Y_{i,j-1}^t - Y_{i,j}^t| - |\hat{Y}_{i,j-1}^t - \hat{Y}_{i,j}^t| \right|^2,$$
(5)

Sequence length	3	6	9	12	15
	Corr/RMSE	Corr/RMSE	Corr/RMSE	Corr/RMSE	Corr/RMSE
3-month	0.78 / 0.67	0.59 / 1.02	0.58 / 1.24	0.81 / 0.54	0.68 / 1.10
6-month	0.52 / 0.96	0.46 / 1.15	0.47 / 1.16	0.64 / 0.81	0.48 / 1.72
12-month	0.29 / 1.00	0.36 / 1.39	0.28 / 1.14	0.53 / 0.84	0.31 / 2.06
18-month	0.17 / 1.06	0.25 / 1.72	0.15 / 1.17	0.44 / 0.91	0.30/2.15

Table 1. Correlation skill (Corr) and Root Mean Square Error (RMSE) of lead times with different input sequence lengths

where i, j denote the pixel position on the sea surface temperature field. Here we only consider the gradient difference in neighboring regions. In future studies, we will consider the gradient difference at a larger spatial scale according to the characteristics of ENSO, such as the difference in SST between the Western Pacific and the Central Pacific during ENSO (Zinke et al., 2021).

# **3** Model Performance evaluation

190

# 3.1 Influence of the input sequence length

Appropriate input sequence length is critical for ENSO prediction of the multichannel model. We use data from the past 3, 6,
9, 12 and 15 months as inputs to predict the development of ENSO in the next 18 months to examine the effects of different input sequence lengths on ENSO predictions. Table 1 shows the comparison of correlation skill and RMSE of lead times. For the correlation coefficient, the higher the value is, the higher the forecasting skill is. And the smaller RMSE represents the higher skill. The results show that the ENSO-MC model performs best with data from the past 12 months as input. This may be because the variables we select have long-term memory for the development of ENSO, such as oceanic heat content. A longer input sequence contains more information that is helpful to ENSO prediction, but also contains more noise that interferes with the prediction, so the improvement of forecasting skill is not positively correlated with the increase of input sequence length.

But for different models and forecasting horizons, the most appropriate input sequence lengths may not be the same.

# 3.2 Physical variable sensitivity for the multichannel structure

- With the physical variables we selected in the Section 2, we construct multichannel input that takes into account the complicated spatiotemporal interactions in the ocean and atmosphere underpinning the onset and development of ENSO events. For the grid observations of monthly SST, heat content, zonal wind and meridional wind, we treat each type of the input variable as a channel in the first ConvLSTM layer and thus there are four channels. The number of channels is the depth of the matrices involved in the convolutions, so that the cross-correlation and transmission between these ocean-atmosphere data can be calculated in the convolution operation. We design an ablation experiment to examine the contribution of predictors and the effectiveness
- of the multichannel structure. In addition to SST, the most important predictor, we remove heat content (t300) and wind from



Figure 3. The correlation skill of Nino 3.4 index of ENSO-MC model with different predictors.



**Figure 4.** (i) The detailed structure of the skip-layer connection and attention mechanism between encoder and decoder at the  $n^{th}$  layer in ENSO-MC. (ii) The correlation skill of Nino 3.4 index of the forecast lead month in models with different structures: (a) ENSO-MC model of skip-layer connection with attention mechanism and states connection, (b) ENSO-MC model without attention mechanism, (c) ENSO-MC model without states connection, (d) ENSO-MC model without skip-layer connection.

the inputs respectively to detect their effects on the correlation skill of the Nino3.4 index. As shown in Fig. 3, the model using the three key ingredients of Bjerknes feedback (SST, heat content, wind) as input produces more favorable forecast skill than the ones that remove one of them, which indicates that the multichannel structure can help to learn the ocean-atmosphere coupling between several important predictors. For the models with two predictors, the model containing wind predictor shows slightly higher forecasting skill in the first few months, while the one containing heat content predictor performs more stable skill at lead times of more than eight months. It suggests that the memory of subsurface heat plays an important role in ENSO prediction on seasonal to interannual time scales, which is consistent with previous research.

#### **3.3** Effectiveness of the model components

215

The ENSO-MC model learns the feature of ENSO at different spatial scales with the convolution and max-pooling layers in the encoder, and gradually restores the spatial dimensionality of the original SST field in the decoder. With symmetrical structure design of the encoder and decoder as shown in Fig. 1, skip-layer connection is used to transfer features form the encoder to the decoder to recover spatial information lost during downsampling (yellow line in Fig. 1). Rather than transferring the original features of all time steps obtained from the encoder, we design an attention mechanism to enable the skip-layer to automatically learn the attention weights  $\beta_1, \beta_2, \ldots, \beta_t$  on the temporal sequence because these air-sea features may have different effects on ENSO development at different time scales. As shown in Fig. 4(i), the encoder obtains the features  $f_n \in \mathbb{R}^{T_{in} \times h_n \times w_n \times c_n}$  after

225 max-pooling and convolution calculation at the  $n^{th}$  layer. Using a two-layer densely-connected neural network, we obtain the attention weight  $\beta \in \mathbb{R}^{T_{in}}$  of each time step's features according to Eq. (6), where  $f_n' \in \mathbb{R}^{T_{in} \times (h_n \times w_n \times c_n)}$  are reshaped from  $f_n$ :

$$\beta = softmax(\mathbf{W}_{\beta\alpha}tanh(\mathbf{W}_{\alpha f}f_n^{\top} + \mathbf{b}_{\alpha f}) + \mathbf{b}_{\beta\alpha}), \tag{6}$$

where  $\mathbf{W}_{\alpha f}$ ,  $\mathbf{W}_{\beta \alpha}$  are weight matrices created by the layer, and  $\mathbf{b}_{\alpha f}$ ,  $\mathbf{b}_{\beta \alpha}$  are the bias vectors.  $\beta$  represents the contribution 230 of each time step to prediction. According to Eq. (7), the feature maps of each time step are multiplied by the corresponding weights, and the fused maps  $\tilde{f}_n \in \mathbb{R}^{h_n \times w_n \times c_n}$  are obtained by adding them along the time dimension.

$$\tilde{f}_n = \sum_{T_{in}} (\beta \circ f_n),\tag{7}$$

235

where  $f_n$  are the feature maps to be transmitted in the skip-layer connection, which are connected to the features of the corresponding layer in the decoder. Besides, we also add states connection between the encoder and the decoder (grey line in Fig. 1), where the hidden states output by the ConvLSTM layers in the encoder are reserved for the corresponding layer when the decoder is initialized. With the methods of skip-layer connection and states connection, the model can make full use of the information extracted from the encoder before ENSO events, which help stabilize training and convergence.

We remove the attention mechanism (model b), states connection structure (model c) and skip-layer connection structure (model d) respectively from the constructed ENSO-MC model (model a) to analyze their effects on model performance. As 240 shown in Fig. 4(ii), the two connection structures, especially the skip-layer connection structure, have a great influence on the prediction results. In model b, we use average weights to replace the attention mechanism. Compared with model a, selfattention mechanism can play a greater role in long-term ENSO prediction.

#### 3.4 Effects of different loss functions

- In order to balance the optimization speed of each loss in the training process, we set  $\lambda_{mse}=7$ ,  $\lambda_{ssim}=9$  and  $\lambda_{qdl}=1$ . The 245 effectiveness of combined loss function is validated. As shown in Fig. 5(a), although SSIM and GDL do not significantly improve the model performance when combined with MSE alone, the combination of MSE, SSIM and GDL loss functions achieve the best performance on the correlation skill. Besides, since GDL loss function tends to retain extreme values and MSE loss function tends to smooth all values, the presence of GDL inhibits the decrease of MSE, so the MSE errors of the models
- with GDL loss function are higher than the ones without GDL (Fig. 5(b)). And comparing the results of correlation skill and 250 RMSE in Fig. 5(a) and (b), low RMSE values do not represent high correlation skills. Therefore, it is necessary to explore loss functions suitable for ENSO prediction other than MSE to balance the training of the model.





Figure 5. The performances of the ENSO-MC with different loss functions.

# 4 Analysis of ENSO forecast skill

#### 4.1 Simulation of two types of ENSO

- 255 The simulation ability of ENSO-MC for different types of events is evaluated in this section. We first select the typical EP El Niño, CP El Niño and La Niña events in recent years to validate the forecast skills on individual events in detail. The forecasts of spatial patterns and Niño 3.4 index time series are compared with observations. Besides, the correlation skills for all targeted seasons of the model are further validated.
- We validate the performance of ENSO-MC for simulating the latest extreme El Niño event in 2015/16. It can be classified
  as an EP-type event, and some studies suggested that the 2015/16 El Niño appears a mixed EP and CP patterns (Santoso et al., 2017). Figure 6(a) compares the predicted spatial evolution of SST anomalies for 2015/16 event (the first row) ahead of one year and the corresponding observed patterns (the second row). The four months shown in the Fig. 6(a) (June, September, December and April) represent the main states involved in the phases of ENSO evolution. The temperature anomalies emerge in the eastern equatorial Pacific (June. 2015), which are then amplified and spread to the central equatorial Pacific (Sept. 2015). When the event reaches the mature stage (Dec. 2015), the center of the anomalies tends to move toward the central equatorial Pacific, eventually decaying in spring 2016 (April. 2016). The prediction of SST anomalies development in the equatorial Pacific during the mature and decay stages. And in the South Pacific, there is more cooling in the model than observed. According to the predicted SST anomalies patterns, the time series of Nino3.4 index is further calculated (Figs.
- 270 6(b), (c)). The amplitude and temporal evolution of the 2015/16 El Niño for the one-year-lead forecast of ENSO-MC are almost consistent with observations, although with weaker amplitude compared to observed Nino3.4 index. And the forecasts initiated during the May-June-July (MJJ) season (Fig. 6(c)) agree a bit more favorably with the observations than initiated during February-March-April (FMA) 2015 (Fig. 6(b)).



**Figure 6.** The SSTA patterns and index prediction results for 2015/16 El Niño using ENSO-MC. (a). Spatial development of SST anomalies predicted (the first row) ahead of one year compared with the real observations (the second row) for the onset, growth, mature and decay phase. (b). Nino3.4 index time series forecast initiated in the FMA season using ENSO-MC (red) and the observed Nino3.4 index (black). (c). Nino3.4 index time series forecast initiated in the MJJ season using ENSO-MC (red) and the observed Nino3.4 index (black).

By contrast, the prediction results of 2009/2010 event, which is known as a CP El Niño, are shown in Fig. 7. The spatial
patterns of SST anomalies predicted (the first row) and observed (the second row) are shown in Fig. 7(a). The observations show a pronounced central Pacific warming. Specifically, the SST anomalies appear (June. 2009), grow (Dec. 2009) and reach the maturity (Jan. 2010) in the central Pacific, whose meridional shift on the equator is less obvious. The model captures most of these features, although the anomalies distributed over the central equatorial Pacific are smaller in amplitude and scope than observed during the growth and mature phase, and the CP characteristics of ENSO are not as pronounced as the actual ones.
And there is also some evident bias in the midlatitudes. For the Nino3.4 index results (Figs 7(b), (c)), the ENSO-MC model exhibits the similar trend but weaker amplitude compared with the observed values. Especially when initiated in FMA 2009 (Fig 7(b)), the model tends to underestimate the strength during the growth phase.

Given the different mechanisms and periods of warm El Niño and cold La Niña events, we also evaluate the simulation ability of the model for a La Nina event. Figure 8 shows the prediction results for 1988/1989 strong La Niña. The simulated evolution of SST anomalies spatial structure along the equator are in reasonable agreement with observations (Fig. 8(a)). The



**Figure 7.** The SSTA patterns and index prediction results for 2009/10 El Niño using ENSO-MC. (a). Spatial development of SST anomalies predicted (the first row) ahead of one year compared with the real observations (the second row) for the onset, growth, mature and decay phase. (b). Nino3.4 index time series forecast initiated in the FMA season using ENSO-MC (red) and the observed Nino3.4 index (black). (c). Nino3.4 index time series forecast initiated in the MJJ season using ENSO-MC (red) and the observed Nino3.4 index (black).

Table 2. Correlation skill (Corr) and Root Mean Square Error (RMSE) of lead times with different input sequence lengths

Lead time	3-month	6-month	9-month	12-month
EP (Niño3)	1.18	1.23	1.04	1.07
CP (Niño4)	0.45	0.99	0.91	0.93

cold temperature anomalies occur in the eastern Pacific, then spread to the central Pacific and reach maturity. However, the predicted cold anomalies are weaker in amplitude, and do not extend as broad area as observed. It is evident in the Nino3.4 index time series results, where the amplitudes of the predictions are weaker than the observed values regardless of whether the initialization is performed before the event (Fig. 8(b)) or in the early stage of its development (Fig. 8(c)).

290

In addition to the above three typical events in recent years, the prediction results of other events that occurred between 1984 and 2019 are also detected. For each event, we compare the spatial development of predicted and observed SST anomalies in the equatorial Pacific from the onset to the maturity stage. Fig. 9 shows the simulation results of the ENSO-MC model for



**Figure 8.** The SSTA patterns and index prediction results for 1988/89 La Niña using ENSO-MC. (a). Spatial development of SST anomalies predicted (the first row) ahead of one year compared with the real observations (the second row) for the transition, onset, growth, mature phase. (b). Nino3.4 index time series forecast initiated in the DJF season using ENSO-MC (red) and the observed Nino3.4 index (black). (c). Nino3.4 index time series forecast initiated in the FMA season using ENSO-MC (red) and the observed Nino3.4 index (black).

three EP El Niño events of 1991/1992, 1997/1998 and 2006/2007, with observations in the first row and predictions in the

295

second row for each group. The results show that the model can simulate the occurrence and development of SSTA for each event. However, for some events with less significant EP type characteristics (for example, that of 1991/1992), the SSTA center of predictions is closer to the central Pacific than observed. In addition, for some super strong events (for example, that of 1997/1998) and weak events (for example, that of 2006-2007), the amplitude of the predicted results at mature phase may be lower or higher than the observed.

300

The prediction results for three CP El Niño events in 1994/1995, 2002/2003 and 2018/2019 are displayed in Fig. 10. For the events of 1994/1995 and 2018/2019, the model can simulate the process that the SST anomalies in the northeast Pacific propagate to the southwest and finally contribute to the occurrence of CP events. The amplitude and center location of the predicted anomalies are also in agreement with the observations. However, the meridional distribution of predicted SSTA is not as broad as observed in the mature stage. The observed SSTA extends eastward to  $80^{\circ}W$ , while the predicted value extends roughly between  $100^{\circ}W$  and  $120^{\circ}W$ .



**Figure 9.** SSTAs of three EP El Niño events in (a) 1991/1992, (b) 1997/1998 and (c) 2006/2007 from the onset to the maturity stage, with observations in the first row and predictions in the second row for each event. The mature phases here are the months when the El Niño events peak. And "0" and "1" next to the calendar month denote the year when the El Niño event occurred and the following year, respectively.

305

And Fig. 11 shows the predictions of three La Niña events in 1984/1985, 1998/1999 and 2000/2001. These three events occurred under different conditions. The events of 1984/1985 and 1998/1999 were preceded by strong El Niño events, and the 1998/1999 event occurring more rapidly. The 2000/2001 La Niña was another weaker event after the previous La Niña event



Figure 10. As in Fig. 9, but for the three CP El Niño events in (a) 1994/1995, (b) 2002/2003 and (c) 2018/2019.

ended. Compared with observations, the model can simulate the occurrence, development and phase transition or persistence of La Niña events.

310

In addition to comparing the detailed spatial distribution of SSTA, the related indices and metrics are calculated to further evaluate the simulation performance of the ENSO-MC model. The Niño 3 index (average SST anomalies over  $5^{\circ}N - 5^{\circ}S$ ,  $150^{\circ}W - 90^{\circ}W$ ) and the Niño 4 index (average SST anomalies over  $5^{\circ}N - 5^{\circ}S$ ,  $160^{\circ}E - 150^{\circ}W$ ) are commonly used to define two types of El Niño events. Events with Niño 4 index greater than Niño 3 are regarded as CP El Niños, and events with



Figure 11. As in Fig. 9, but for the three CP El Niño events in (a) 1994/1995, (b) 2002/2003 and (c) 2018/2019.

315

Niño 3 index greater than Niño 4 are classified as EP El Niños. Figure 12a, b shows the distribution for Niño 3 and Niño 4
indexes calculated from the model's one-year-lead predictions of the peak periods for all EP events (Fig. 12a) and CP events (Fig. 12b) from 1984 to 2019. The results show that the model can correctly classify five EP events (1987/1988, 1991/1992, 1997/1998, 2006/2007, 2015/2016) and three CP events (1994/1995, 2002/2003, 2018/2019) in the past 30 years, but misjudge the event of 2009/2010 as EP type and no El Niño event occurred in 2004 (Niño 3=0, Niño 4=0). The CP event of 2004/2005 is much weaker than other CP ones, making it more difficult for the model to capture its development. We also make statistics on



Figure 12. Scatterplots in Nino3-Nino4 index plane of 12-month-lead predictions for all (a) EP El Niño events and (b) CP El Niño events during peak phase from 1984 to 2019. (c) Root Mean Square Error (RMSE) of the Nino3.4, Nino3 and Nino4 indexes between the forecast results of the ENSO-MC model and observations during validation period.

the RMSE between the predictions and official records of Nino3/4 index for all EP/CP El Niño events at mature phase (Table 2) and for the whole validation period (Fig. 12c), and find that although the model has a higher classification accuracy for EP events, the index error of predictions for EP events is larger than that for CP. It may be because most of the strong El Niños are EP-type events, and the prediction skills of the model for such extreme events need to be improved. The SSTA distribution in Fig. 9 also shows that for some EP events, there is a difference in amplitude between predictions and observations for the maturity stage of the event, while that of the CP events is consistent with the observations (Fig. 10).

#### 4.2 Correlation skills for different calendar months

The model also shows a reasonable forecast of ENSO ahead of one year for all targeted seasons (Fig. 13). Figure 13(a) shows the correlation skill of Nino3.4 index forecasts for the GODAS validation data from 1982 to 2019, which are initiated in each calendar month and predicted for a lead of up to 18 months. The correlation skill in the model is above 0.5 for a lead of 11 months. And Figure 13(c) shows the results for the last decade whose validating period is from 2010 to 2019. The results in Fig. 13(c) perform only slightly skillful than those in Fig. 13(a), indicating the robustness of the model in terms of validation time. We also compare the two forecast strategies in deep learning, that is, the one-step ahead forecast strategy and the multi-step forecast strategy. Figures 13(a) and 13(c) are the results of multi-step prediction, while Figures 13(b) (1982-2019) and 5(d) (2010-2019) are the results of one-step prediction. It shows that regardless of the season from which the forecast is started, numbers in Fig. 13. We also calculate the overall decline of forecasting skills in each target month compared with the previous month and the results are presented in Fig. 14. The prediction skills of the model decline most from April to May, regardless of whether the multi-step or one-step ahead forecast strategy is used. Besides, Fig. 14 shows that the performance of the model is slightly improved in winter, which leads to the improvement of skills after 12 months in Fig. 13. Since the seasonal variation

of SST anomaly variance is weaker in spring, it is difficult for the model to capture useful information, which leads to the spring predictability barrier (SPB), while the strong signals of ENSO during winter are more easily learned by the model. In



**Figure 13.** The correlation skills of the Nino3.4 index forecasts started from each calendar month in ENSO-MC using multi-step forecast strategy (a) and one-step ahead forecast strategy (b) for the GODAS data from 1982 to 2019. (c) is the same as (a), except for the GODAS data from 2010 to 2019. (b) is the same as (d), except for the GODAS data from 2010 to 2019. Hatches represent the forecasts with correlation skill exceeding 0.5, and the black numbers mean the target forecast months.

addition, the one-step ahead strategy has a larger decline after the boreal spring (Fig. 14), and the subsequent forecasts are more susceptible to the SPB due to its cumulative error (Figs. 13(b), (d)), while the method of the multi-step strategy can reduce the influence (Figs. 13(a), (c)). We can conclude that the ENSO-MC model using multi-step forecast strategy is less affected by spring predictability barriers.

It should be noted that the correlation coefficient skills of Nino3.4 index obtained by simulating the SST anomalies spatial distribution are not as high as that obtained by using deep neural network to predict the Nino3.4 index directly, such as the results of Ham et al. (2019). This is because the former has a higher prediction target dimension, and the cumulative error of each point of the spatial field is larger than the error of direct prediction of a single index. This is also one of the key issues to be solved in the future development of the ENSO-MC model.

350

345

19



Figure 14. The decline of forecasting skills for ENSO in each target month using multi-step forecast strategy and one-step ahead forecast strategy.

#### 5 Precursor analysis of two types of ENSO

Based on the ENSO-MC model that successfully simulates different types of ENSO events, we can further explore the ENSO dynamics learned by the ENSO-MC model and observe the signals before the onset of events. And since the ENSO-MC model using the multi-step forecast strategy achieves better performance than using one-step strategy, here we calculate the saliency maps based on the multi-step forecasting model for precursor analysis and sensitive area identification. Considering the important role of subsurface thermal memories in ENSO prediction, the precursory characteristics in heat content of different types of events are discussed here. We select five EP El Niño events (1987/88, 1991/92, 1997/98, 2006/07 and 2015/16), three CP El Niño events (1994/95, 2002/03 and 2009/10) and three La Niña events (1988/89, 2007/08 and 2010/11) occurred in the past 30 years, and calculate the precursor maps of heat content anomaly in the year prior to each event.

360

355

Specifically, the precursor maps of each event are obtained by computing the gradient of the regressed output with respect to the input with saliency map method. The maps tell us how the output value will change when the pixel at this position in the input image changes slightly, that is, the sensitivity of the predicted results to the perturbations in each region. In this way, we can get the initial perturbation distribution that would develop into an ENSO event. The gradient calculation is equivalent to the process of seeking the gradient of the errors with respect to the weights during training a machine learning model, which represents the contribution of each weight to the total loss. We calculate the deviations between the output  $\hat{Y}$  and the real 365 situation Y, fix the weights of the ENSO-MC model and perform back propagation for each pixel (x, y) in the input H at each moment  $\tau$ .

 $loss = \mathcal{L}(Y, \hat{Y})$ (8)

$$370 \quad M_{\tau} = \frac{\partial loss}{\partial H_{x,y}^{\tau}} \tag{9}$$

Then we obtain a series of heat maps  $M: M_1, \ldots, M_t$  over the heat content predictors  $H_1, \ldots, H_t$ . Each  $M_\tau$  indicates the perturbation sensitivity distribution in heat content  $H_{\tau}$  for  $\tau$ -lead time. For each event, twelve heat maps  $M_1, \ldots, M_{12}$  are obtained, which describe the precursor development in the year preceding the event. We add up the maps of  $\tau$ -lead time of each event for one type to obtain the composite evolution maps  $M_1^{comp}, \ldots, M_{12}^{comp}$  for each type. For example, the composite precursor map of  $\tau$ -month lead for EP-type El Niño is obtained by adding up the precursor maps of  $\tau$ -lead time for all five 375 EP events. The composite maps of CP-type El Niño and La Niña are obtained in the same way. The precursor maps from 12-month lead to 1-month lead of EP-type El Niño, CP-type El Niño and La Niña are shown in the Fig. 15, and we present results every few months for each type to see more clearly how precursors change over time. Since the composite maps are the sum of saliency maps of each event, here we focus on the distribution of perturbation without considering the intensity, and the saliency values in Fig. 15 are the standardized results of the scale between 0 and 1. For EP El Niño (Fig. 15(a)), the subsurface 380 temperature component presents large anomalies in the equatorial Pacific, especially in the central and western Pacific, the subtropical northeast Pacific and the subtropical South Pacific. With the occurrence of El Niño, the anomalies weaken in the equatorial region and slightly intensify in the subtropical area. And the large anomalies of the precursory perturbation for CP El Niño (Fig. 15(b)) are concentrated in the subtropical northeast Pacific. As El Niño approaches, the disturbance tends to spread to the southeast. Regarding the subsurface disturbance for La Niña (Fig. 15(c)), their anomalies are concentrated in equatorial regions and propagates from the western Pacific to the eastern.

385

On the whole, the subsurface signals distributed in Fig. 15(a) are more intense and more extensive than those in Fig. 15(b), indicating that the occurrence of the EP-type El Niño is more related to the subsurface dynamics, while the CP events may be more affected by the atmospheric convection. Specifically, compared with Fig. 15(b) (CP-type El Niño), Fig. 15(a) (EP-type El

- 390 Niño) shows a more pronounced signal, especially in the equatorial Pacific. It may be related to the stronger zonal tilt change of the equatorial thermocline and larger eastward movement of convection in tropical Pacific before the EP-type events. For example, as shown in Fig. 16, a series of westerly wind events along the equatorial Pacific led to an abrupt relaxation and reversal of trade winds in the western and central equatorial Pacific in early 1997. The westerly wind anomalies generated downwelling Kelvin waves, which propagated eastward and deepened the thermocline in the eastern Pacific in late 1997. The
- 395 depressed thermocline limited the upwelling of subsurface cold water, prompting the development of warm surface temperatures. Meanwhile, westward-propagating Rossby waves shallowed the thermocline in the western Pacific. These processes led to significant changes in the equatorial thermocline (Fig. 15(a)), a flattening of the thermocline and a decrease in the zonal SST gradient along the equator. The reduction of the SST gradient in turn further weakened the trade winds, leading to the rapid development of the 1997/1998 El Niño. La Niña events usually occur in the second year after a warm event. As shown in
- 400 Fig. 15(c), there are precursor signals produced by wind forcing propagating eastward from the western tropical Pacific in the subsurface from 12-month lead to the occurrence. Combined with the mechanism of the La Niña event, the signal would shoal



Figure 15. The composite evolution maps of initial perturbations in heat content before (a) EP-type El Niños, (b) CP-type El Niños and (c) La Niñas from 12-month lead to 1-month lead.

the thermocline in the eastern Pacific and enhance the upwelling of cold subsurface waters, thereby ending the El Niño event and triggering a subsequent cold event.

- While the equatorial subsurface signal is weak in Fig. 15(b), but there is an obvious signal in the North Pacific. The results
  are consistent with the previous studies that the negative phase of the North Pacific Oscillation promotes the development of SST anomalies in the central Pacific (Yu and Kim, 2011). Besides, there are robust signals over the northeastern Pacific in both types of El Niño (Figs. 15(a), (b)). The distribution is similar to the spatial structure of the Pacific meridional mode (PMM). PMM is forced by mid-latitude atmospheric variability in the Northern Hemisphere and evolves equatorward subsequently, which can affect ENSO. As shown in Fig. 17, one year before the 1994/1995 CP El Niño, there were warm subtropical SST
  anomalies extending southwest from Baja California. The SST anomalies weakened the trade winds and reduced the surface evaporation over the region via Wind-Evaporation-SST (WES) feedback. The reduction in evaporation allowed warm waters to expand further southwestward, enhancing the PMM and eventually reaching the equator, which weakened equatorial trade
- winds and triggered an El Niño event in late 1994. PMM not only appeared before CP El Niño, for example, the emergence of PMM in late 2014 contributed to the development of 2015/2016 El Niño (Fig. 17). It indicates that signals outside the tropics
- 415 play an important role in the prediction of El Niño and PMM can be regarded as a precursor to El Niño.



**Figure 16.** Longitude-time diagram of monthly surface zonal wind anomalies (left), SST anomalies (middle), and heat content (t300) anomalies (right) across the equatorial Pacific  $(2^{\circ}N - 2^{\circ}S, 120^{\circ}E - 80^{\circ}W)$  from September 1996 to April 1998. Data are based on NCEP Global Ocean Data Assimilation System and ERA-5.



Figure 17. The SST and 10-m wind-vector anomalies for the different seasons before 1994/1995 CP Niño and 2015/2016 EP Niño.



Figure 18. The composite saliency maps of surface (a) and subsurface (b) over  $40^{\circ}N - 40^{\circ}S$ ,  $120^{\circ}E - 80^{\circ}W$ . The higher the saliency value, the more sensitive the area. Since the saliency map results obtained from the surface layer and the subsurface layer may affect each other, we first select six common areas with large values as candidates to perform sensitivity experiments. The six areas are the central equatorial Pacific (0), the western equatorial Pacific (1), the northern Central Pacific (2), the northeastern Pacific (3), the southwestern Pacific (4) and the southern Central Pacific (5).

### 6 Targeted observation sensitive area identification of ENSO

A saliency map shows which input pixels produce the largest increase in output values with minimal change. The idea is similar to the targeted observation strategy, that is, prioritizing the deployment of observations in the sensitive areas where small perturbations tend to have the greatest impacts on the forecasts. The saliency map method is therefore appropriate 420 to identify the sensitive areas of targeted observation for ENSO. The areas with large values in the saliency map indicate that improving the accuracy of observations in these sensitive areas is the most efficient way to correct the output. Here we explore the sensitive areas of the surface and the subsurface layer respectively, so in addition to heat content, the saliency maps of SST are also calculated. Since the sensitive area is a common attribute, it should be universal to all ENSO events and different types of ENSO are not considered here. We select eight El Niño events (1987/88, 1991/92, 1994/95, 1997/98, 2002/03, 2006/07, 2009/10 and 2015/16) and three La Niña events (1988/89, 2007/08 and 2010/11) that occurred in the past 425 30 years, and calculate the saliency maps of SST and heat content for each event according to the method described in Sect. 4. Then all the saliency maps of SST are added up to obtain the composite saliency map of the surface (Fig. 18(a)), and that of the subsurface (Fig. 18(b)) is obtained in the same way. The saliency values in the figure are the standardized results of the scale between 0 and 1. The anomalies of surface disturbance are mainly distributed in the north central Pacific, the central Pacific and the western Pacific. The anomalies in the north Pacific are more intense than those in the south Pacific. For the subsurface 430 temperature precursory, the perturbation possesses a wide range of anomalies in the equatorial Pacific and the North Pacific, and the anomaly values are more intense than that of surface disturbance. Since ENSO-MC regards multi-variable fields as multiple channels, the saliency regions obtained from surface layer and subsurface layer may affect each other. As shown in Fig. 18, the large-value region of surface (a) overlaps with that of subsurface (b). Therefore, we first artificially define six 435 common regions with large values (black boxes in the Fig. 18), and perform sensitive experiments on the surface and subsurface respectively. Since the error is random in real observations, we use random perturbations for sensitivity experiments. For each



**Figure 19.** Sensitive areas identification results for ENSO with the saliency map method. (b) compares the sensitivity of the six candidate areas (0-5) for surface (red) and subsurface (blue). The first two most sensitive areas are selected as the sensitive areas of targeted observation for ENSO, that is, the area\_0 and area\_2 for surface and the area\_1 and area\_4 for subsurface. Specifically, the hatching areas in (a) (surface) and (c) (subsurface). In order to evaluate the effectiveness of the above identified sensitive areas, (d) compares the benefit for removing the random perturbations in the sensitive areas (blue) and removing the random perturbations outside the sensitive areas (orange, that is, retaining the random perturbations in the sensitive areas) for the eight El Niño events and three La Niña events occurred in the past 30 years.

of the 11 selected ENSO events, 30 sets of random perturbations are superimposed to the original input field. The experiment that superimposes whole-field perturbations is called "all\_rand", and the experiment that removes perturbations in the target area based on the whole-field perturbations is called "remove\_rand". For the six regions shown in Fig. 18, the sensitivity of each region is measured according to Eq. (10), that is, the reduction of prediction error caused by removing random perturbations in each region.

440

$$Sensitivity = \frac{\sum_{i} [\mathcal{L}_{RMSE}(Y_{i}, \hat{Y}_{i}^{all}) - \mathcal{L}_{RMSE}(Y_{i}, \hat{Y}_{i}^{remove})]}{\sum_{i} \mathcal{L}_{RMSE}(Y_{i}, \hat{Y}_{i}^{all})}.$$
(10)

For each event i,  $\hat{Y}_i^{all}$  represents the prediction results of experiment "all\_rand",  $\hat{Y}_i^{remove}$  represents the prediction results of experiment "remove\_rand", and  $Y_i$  represents the real observation. The results are shown in Fig. 19(b). The surface areas with the highest sensitivity for ENSO are No.0 and No.2 areas, that is, the central equatorial Pacific and the northern Central Pacific shown in Figure 19a, while the subsurface areas with the highest sensitivity are No.1 and No.4 areas, namely the western equatorial Pacific and the southwestern Pacific shown in the Fig. 19(c).

Furthermore, we perform two sets of experiments to measure the benefit of effective observations in the identified sensitive areas for improving forecast results. The first set calculates the reduction of the prediction error after removing the random

- perturbations in the identified sensitive areas ("remove"), and the second calculates the reduction after removing the pertur-450 bations outside the sensitive areas, that is, there are perturbations in the hatching areas in Figs. 19(a), (c) ("retain"). For each ENSO event, we superimpose 30 groups of random perturbations on the original input field, select the random perturbations k whose errors are larger than the mean error of all groups, and then conduct experiment "remove" and experiment "retain" respectively. Then we calculate the benefit  $B_{remove}$  and  $B_{retain}$ , where  $V_{in}$  is the volume of the identified sensitive areas,  $V_{out}$ is the volume outside the sensitive areas.
- 455

$$B_{remove} = \frac{\sum_{k} [\mathcal{L}_{RMSE}(Y_k, \hat{Y}_k^{all}) - \mathcal{L}_{RMSE}(Y_k, \hat{Y}_k^{remove})]}{V_{in} \sum_{k} \mathcal{L}_{RMSE}(Y_k, \hat{Y}_k^{all})}$$
(11)

$$B_{retain} = \frac{\sum_{k} [\mathcal{L}_{RMSE}(Y_k, \hat{Y}_k^{all}) - \mathcal{L}_{RMSE}(Y_k, \hat{Y}_k^{retain})]}{V_{out} \sum_{k} \mathcal{L}_{RMSE}(Y_k, \hat{Y}_k^{all})}$$
(12)

- $B_{remove}$  represents the degree of reduction of prediction errors after implementing target observation in per volume of 460 identified sensitive areas shown in Figs. 19(a), (c), and  $B_{retain}$  represents the reduction after implementing target observation in per volume of non-sensitive areas (here is the outside areas of hatching regions in Figs. 19(a), (c)). After removing the perturbations inside or outside the sensitive areas, the prediction errors are both reduced. However, due to the small size of sensitive areas, the benefit obtained by removing random perturbations in the sensitive areas is relatively high (Fig. 19(d)), except for the super strong El Niño event in 1997. Therefore, it is reasonable to give priority to effective observations in these identified sensitive areas, which are located in the central equatorial Pacific and the northern Central Pacific surface region, 465 and western equatorial Pacific and the southwestern Pacific subsurface region. The results for the equatorial region support the conclusions of Kumar et al. (2014) and Duan and Hu (2016) in previous studies. Kumar et al. suggested that the observations
- in the central Pacific are more crucial than those in the eastern Pacific because of their role in preserving the memory of ENSO evolution. Duan and Hu emphasized the importance of subsurface signals in the western Pacific for ENSO predictions, which 470 can influence the surface through equatorial waves and thermodynamic effects. However, due to the complexity of different
- models has a great impact on the identification of sensitive areas, there is no consistent conclusion about the sensitive areas of ENSO at present. For example, based on the outputs of CMIP5 model, Zhang et al. (2015) identified the central-eastern equatorial surface region and eastern subsurface region as the sensitive areas. Therefore, determining the most appropriate regions for target observation remains to be a long-standing challenge. Nevertheless, such studies based on interpretability can
- improve our understanding of how the ENSO-MC model works in ENSO prediction. The results of sensitive area identification 475 support the theoretical understanding that oceanic thermal anomaly in the central and western Pacific provides a key long-term memory for SST predictions. In addition, the results show that processes outside the tropical Pacific also have an impact on ENSO prediction, such as surface temperature variations in the northern Central Pacific and subsurface thermal changes in the southwestern Pacific.

#### 480 7 Conclusions

With the successful application of deep learning algorithms in ENSO forecasts, this paper attempts to expand the application scope of deep neural networks from prediction to a broader field, including the pattern simulation, understanding and observation of ENSO. For reliable forecasts of the two types of ENSO, a multichannel data-driven model ENSO-MC is proposed to simulate the diversity of spatial patterns during ENSO events. Based on the ENSO-MC model, we then provide a new promising approach to investigate the early signals of different types of ENSO events and identify the sensitive areas with the help of

485

saliency map interpretability method.

Specifically, the model ENSO-MC driven with oceanic and atmospheric predictors is proposed to simulate the oceanatmosphere coupling process and predict the changes in spatial distribution of sea surface temperature anomalies. The simulation results show that the model can predict the development of SST anomalies in the equatorial Pacific during the onset, growth, maturity and decay of the El Niño and La Niña events one year in advance. In particular, we simulate the changes in the SST anomalies field of typical EP-type El Niños and CP-type El Niños at lead time beyond one year. With the SST pattern forecasts, the all-season correlation skill of the Nino3.4 index in the ENSO-MC model is also evaluated, which is above 0.5 for a lead of 11 months. The precursor maps reveal the different pronounced characteristics of the subsurface signals before EP-type El Niño, CP-type El Niño and La Niña events. The results indicate that the EP-type El Niño is more related to the

- 495 tropical thermocline dynamics, and the subtropical precursors seem to favor the generation of the CP-type El Niño. Both types of events have pronounced precursory signals in the northeastern Pacific whose distribution is similar to PMM. Before the La Niña events, there is an obvious subsurface signal propagating eastward from the equatorial western Pacific, which would shoal the thermocline in the eastern Pacific and trigger a cold event. In addition, we present an attempt of the saliency method based on the ENSO-MC model for sensitive area identification of ENSO. The identification results show that the surface sensitive
- 500 areas are located in the central equatorial Pacific and the northern Central Pacific, and the subsurface sensitive areas are concentrated in the western equatorial Pacific and the southwestern Pacific. Additional observations in these areas are expected to better predict an event in the future. It indicates that in equatorial regions, the central surface area and the western subsurface area play an important role in the occurrence of future ENSO events, which are essential for preserving the memory of ENSO evolution. Besides, the processes in the extratropical Pacific also contribute to ENSO prediction, such as changes in the surface

505 layer of the northern Central Pacific and the subsurface layer of the southwestern Pacific.

Since the cumulative error of the Nino3.4 index calculated by predicting SST anomalies patterns is larger, the correlation skill is not as high as that obtained by predicting the index directly. Further research should be undertaken to explore how to ensure high correlation skills of Nino3.4 index while correctly simulating the spatial distribution of SST anomalous. In addition to the existing components in the ENSO-MC model, how effectively use our existing domain knowledge, such as conservation

510 of mass, conservation of salinity and other physical laws, to build physics-informed ENSO-MC model that may help reduce uncertainty and increase the credibility of predictions. For the precursor investigation, this paper focuses on verifying the known ENSO mechanisms, and the unknown inherent characteristics exploration will be considered in the future. Combining structural causal model would help to extract the unknown causality relationship among factors and phenomena in ENSO complex interaction on this matter. This would help further explore the precursors of ENSO and improve our understanding of

515 its predictability.

520

*Code and data availability.* The source code of the ENSO-MC 1.0 is available at https://doi.org/10.5281/zenodo.5725987. Thanks to Computational and Information Systems Laboratory (CISL) of NCAR, IRI/LDEO Climate Data Library, NOAA/OAR/ESRL PSL and ECMWF for providing the historical reanalysis data (https://rda.ucar.edu/) (Rayner et al., 2003); (https://iridl.ldeo.columbia.edu/) (Giese and Ray, 2011); (https://psl.noaa.gov/data/gridded/) (Reynolds et al., 2002; Behringer and Xue, 2004); (https://apps.ecmwf.int/datasets/) (Dee et al., 2011)

*Author contributions*. All authors designed the experiments and carried them out. Yuehan Cui developed the model code and performed the experiments. Yuehan Cui and Shijin Yuan prepare the manuscript with contributions from all co-authors.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. This study is supported in part by the key project fund of Shanghai 2020 "science and technology innovation action
 plan" for social development under Grant 20dz1200702, in part by the National Key Research and Development Program of China under Grant 2020YFA0608002, in part by the Fundamental Research Funds for the Central Universities under Grant 13502150039/003, and in part by the National Natural Science Foundation of China under Grant 42075141.

# References

An, S.-I., Kug, J.-S., Timmermann, A., Kang, I.-S., and Timm, O.: The influence of ENSO on the generation of decadal variability in the

- 530 North Pacific, Journal of Climate, 20, 667–680, 2007.
  - Ashok, K., Behera, S. K., Rao, S. A., Weng, H., and Yamagata, T.: El Niño Modoki and its possible teleconnection, Journal of Geophysical Research: Oceans, 112, 2007.
    - Behringer, D. and Xue, Y.: Evaluation of the global ocean data assimilation system at NCEP: The Pacific Ocean, in: Proc. Eighth Symp. on Integrated Observing and Assimilation Systems for Atmosphere, Oceans, and Land Surface, 2004.
- Bjerknes, J.: Atmospheric teleconnections from the equatorial Pacific, Monthly weather review, 97, 163–172, 1969.
   Capotondi, A. and Sardeshmukh, P. D.: Optimal precursors of different types of ENSO events, Geophysical Research Letters, 42, 9952–9960, 2015.
  - Capotondi, A., Wittenberg, A. T., Newman, M., Di Lorenzo, E., Yu, J.-Y., Braconnot, P., Cole, J., Dewitte, B., Giese, B., Guilyardi, E., et al.: Understanding ENSO diversity, Bulletin of the American Meteorological Society, 96, 921–938, 2015.
- 540 Chen, D. and Cane, M. A.: El Niño prediction and predictability, Journal of Computational Physics, 227, 3625–3640, 2008. Chen, D., Lian, T., Fu, C., Cane, M. A., Tang, Y., Murtugudde, R., Song, X., Wu, Q., and Zhou, L.: Strong influence of westerly wind bursts on El Niño diversity, Nature Geoscience, 8, 339–345, 2015.
  - Chikamoto, Y., Timmermann, A., Luo, J.-J., Mochizuki, T., Kimoto, M., Watanabe, M., Ishii, M., Xie, S.-P., and Jin, F.-F.: Skilful multi-year predictions of tropical trans-basin climate variability, Nature communications, 6, 1–7, 2015.
- 545 Dee, D. P., Uppala, S. M., Simmons, A., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M., Balsamo, G., Bauer, d. P., et al.: The ERA-Interim reanalysis: Configuration and performance of the data assimilation system, Quarterly Journal of the royal meteorological society, 137, 553–597, 2011.
  - Duan, W. and Hu, J.: The initial errors that induce a significant "spring predictability barrier" for El Niño events and their implications for target observation: results from an earth system model, Climate Dynamics, 46, 3599–3615, 2016.
- 550 Duan, W., Mu, M., and Wang, B.: Conditional nonlinear optimal perturbations as the optimal precursors for El Nino–Southern Oscillation events, Journal of Geophysical Research: Atmospheres, 109, 2004.
  - Duan, W., Yu, Y., Xu, H., and Zhao, P.: Behaviors of nonlinearities modulating the El Niño events induced by optimal precursory disturbances, Climate Dynamics, 40, 1399–1413, 2013.
  - Duan, W., Tian, B., and Xu, H.: Simulations of two types of El Niño events by an optimal forcing vector approach, Climate dynamics, 43,
- 555 1677–1692, 2014.
  - Duan, W., Huang, C., and Xu, H.: Nonlinearity modulating intensities and spatial structures of central Pacific and eastern Pacific El Niño events, Advances in Atmospheric Sciences, 34, 737–756, 2017.
  - Ebert-Uphoff, I. and Hilburn, K.: Evaluation, tuning, and interpretation of neural networks for working with images in meteorological applications, Bulletin of the American Meteorological Society, 101, E2149–E2170, 2020.
- 560 Fedorov, A. V., Hu, S., Lengaigne, M., and Guilyardi, E.: The impact of westerly wind bursts and ocean initial state on the development, and diversity of El Niño events, Climate Dynamics, 44, 1381–1401, 2015.
  - Gebbie, G., Eisenman, I., Wittenberg, A., and Tziperman, E.: Modulation of westerly wind bursts by sea surface temperature: A semistochastic feedback for ENSO, Journal of the Atmospheric Sciences, 64, 3281–3295, 2007.

Giese, B. S. and Ray, S.: El Niño variability in simple ocean data assimilation (SODA), 1871–2008, Journal of Geophysical Research:

565 Oceans, 116, 2011.

- Ham, Y.-G. and Kug, J.-S.: How well do current climate models simulate two types of El Niño?, Climate dynamics, 39, 383–398, 2012.
- Ham, Y.-G., Kug, J.-S., Park, J.-Y., and Jin, F.-F.: Sea surface temperature in the north tropical Atlantic as a trigger for El Niño/Southern Oscillation events, Nature Geoscience, 6, 112–116, 2013.
- Ham, Y.-G., Kim, J.-H., and Luo, J.-J.: Deep learning for multi-year ENSO forecasts, Nature, 573, 568–572, 2019.
- 570 Hu, J. and Duan, W.: Relationship between optimal precursory disturbances and optimally growing initial errors associated with ENSO events: Implications to target observations for ENSO prediction, Journal of Geophysical Research: Oceans, 121, 2901–2917, 2016.
  - Hu, S., Fedorov, A. V., Lengaigne, M., and Guilyardi, E.: The impact of westerly wind bursts on the diversity and predictability of El Niño events: An ocean energetics perspective, Geophysical research letters, 41, 4654–4663, 2014.

Huang, A., Vega-Westhoff, B., and Sriver, R. L.: Analyzing El Niño–Southern Oscillation Predictability Using Long-Short-Term-Memory
 Models, Earth and Space Science, 6, 212–221, 2019.

Jin, F.-F.: An equatorial ocean recharge paradigm for ENSO. Part I: Conceptual model, Journal of the atmospheric sciences, 54, 811–829, 1997a.

Jin, F.-F.: An equatorial ocean recharge paradigm for ENSO. Part II: A stripped-down coupled model, Journal of Atmospheric Sciences, 54, 830–847, 1997b.

580 Kao, H.-Y. and Yu, J.-Y.: Contrasting eastern-Pacific and central-Pacific types of ENSO, Journal of Climate, 22, 615–632, 2009. Kug, J.-S., Choi, J., An, S.-I., Jin, F.-F., and Wittenberg, A. T.: Warm pool and cold tongue El Niño events as simulated by the GFDL 2.1 coupled GCM, Journal of Climate, 23, 1226–1239, 2010.

Kug, J.-S., Ham, Y.-G., Lee, J.-Y., and Jin, F.-F.: Improved simulation of two types of El Niño in CMIP5 models, Environmental Research Letters, 7, 034 002, 2012.

585 Kumar, A., Wang, H., Xue, Y., and Wang, W.: How much of monthly subsurface temperature variability in the equatorial Pacific can be recovered by the specification of sea surface temperatures?, Journal of climate, 27, 1559–1577, 2014.

Latif, M., Biercamp, J., and Von Storch, H.: The response of a coupled ocean-atmosphere general circulation model to wind bursts, Journal of the atmospheric sciences, 45, 964–979, 1988.

Lee, T. and McPhaden, M. J.: Increasing intensity of El Niño in the central-equatorial Pacific, Geophysical Research Letters, 37, 2010.

590 Lingjiang, T. and Wansuo, D.: Using a nonlinear forcing singular vector approach to reduce model error effects in ENSO forecasting, Weather and Forecasting, 34, 1321–1342, 2019.

Mathieu, M., Couprie, C., and LeCun, Y.: Deep multi-scale video prediction beyond mean square error, arXiv preprint arXiv:1511.05440, 2015.

McPhaden, M. J.: Tropical Pacific Ocean heat content variations and ENSO persistence barriers, Geophysical research letters, 30, 2003.

- 595 Meinen, C. S. and McPhaden, M. J.: Observations of warm water volume changes in the equatorial Pacific and their relationship to El Niño and La Niña, Journal of Climate, 13, 3551–3559, 2000.
  - Menkes, C. E., Lengaigne, M., Vialard, J., Puy, M., Marchesiello, P., Cravatte, S., and Cambon, G.: About the role of westerly wind events in the possible development of an El Niño in 2014, Geophysical research letters, 41, 6476–6483, 2014.

Mo, R., Ye, C., and Whitfield, P. H.: Application potential of four nontraditional similarity metrics in hydrometeorology, Journal of Hydrom eteorology, 15, 1862–1880, 2014.

30

Moore, A. M. and Kleeman, R.: The dynamics of error growth and predictability in a coupled model of ENSO, Quarterly Journal of the Royal Meteorological Society, 122, 1405-1446, 1996.

Moore, A. M. and Kleeman, R.: Stochastic forcing of ENSO by the intraseasonal oscillation, Journal of Climate, 12, 1199–1220, 1999.

Mu, B., Peng, C., Yuan, S., and Chen, L.: ENSO forecasting over multiple time horizons using ConvLSTM network and rolling mechanism, 605 in: 2019 International Joint Conference on Neural Networks (IJCNN), pp. 1–8, IEEE, 2019.

- Mu, B., Qin, B., and Yuan, S.: ENSO-ASC 1.0. 0: ENSO Deep Learning Forecast Model with a Multivariate Air-Sea Coupler, Geoscientific Model Development Discussions, pp. 1-33, 2021.
  - Mu, M., Duan, W., and Wang, B.: Conditional nonlinear optimal perturbation and its applications, Nonlinear Processes in Geophysics, 10, 493-501, 2003.
- 610 Mu, M., Yu, Y., Xu, H., and Gong, T.: Similarities between optimal precursors for ENSO events and optimally growing initial errors in El Niño predictions, Theoretical and applied climatology, 115, 461–469, 2014.
  - Mu, M., Duan, W., Chen, D., and Yu, W.: Target observations for improving initialization of high-impact ocean-atmospheric environmental events forecasting, National Science Review, 2, 226-236, 2015.

Newman, M., Shin, S.-I., and Alexander, M. A.: Natural variation in ENSO flavors, Geophysical Research Letters, 38, 2011.

615 Oprea, S., Martinez-Gonzalez, P., Garcia-Garcia, A., Castro-Vargas, J. A., Orts-Escolano, S., Garcia-Rodriguez, J., and Argyros, A.: A review on deep learning techniques for video prediction, IEEE Transactions on Pattern Analysis and Machine Intelligence, 2020. Philander, S. G. H.: El Nino southern oscillation phenomena, Nature, 302, 295–301, 1983.

Rayner, N., Parker, D. E., Horton, E., Folland, C. K., Alexander, L. V., Rowell, D., Kent, E. C., and Kaplan, A.: Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century, Journal of Geophysical Research: Atmospheres, 108, 2003.

- 620
  - Reichstein, M., Camps-Valls, G., Stevens, B., Jung, M., Denzler, J., Carvalhais, N., et al.: Deep learning and process understanding for data-driven Earth system science, Nature, 566, 195-204, 2019.
    - Reynolds, R. W., Rayner, N. A., Smith, T. M., Stokes, D. C., and Wang, W.: An improved in situ and satellite SST analysis for climate, Journal of climate, 15, 1609-1625, 2002.
- 625 Santoso, A., Mcphaden, M. J., and Cai, W.: The defining characteristics of ENSO extremes and the strong 2015/2016 El Niño, Reviews of Geophysics, 55, 1079–1129, 2017.
  - Shi, X. and Yeung, D.-Y.: Machine learning for spatiotemporal sequence forecasting: A survey, arXiv preprint arXiv:1808.06865, 2018.
  - Simonyan, K., Vedaldi, A., and Zisserman, A.: Deep inside convolutional networks: Visualising image classification models and saliency maps, arXiv preprint arXiv:1312.6034, 2013.
- 630 Smilkov, D., Thorat, N., Kim, B., Viégas, F., and Wattenberg, M.: Smoothgrad: removing noise by adding noise, arXiv preprint arXiv:1706.03825, 2017.
  - Sutskever, I., Vinyals, O., and Le, Q. V.: Sequence to sequence learning with neural networks, in: Advances in neural information processing systems, pp. 3104-3112, 2014.
  - Taieb, S. B., Hyndman, R. J., et al.: Recursive and direct multi-step forecasting: the best of both worlds, vol. 19, Citeseer, 2012.
- 635 Tang, Y., Zhang, R.-H., Liu, T., Duan, W., Yang, D., Zheng, F., Ren, H., Lian, T., Gao, C., Chen, D., et al.: Progress in ENSO prediction and predictability study, National Science Review, 5, 826-839, 2018.
  - Tao, L., Duan, W., and Vannitsem, S.: Improving forecasts of El Niño diversity: a nonlinear forcing singular vector approach, Climate Dynamics, 55, 739-754, 2020.

Tian, B. and Duan, W.: Comparison of the initial errors most likely to cause a spring predictability barrier for two types of El Niño events,

Climate Dynamics, 47, 779–792, 2016.
 Timmermann, A., An, S.-I., Kug, J.-S., Jin, F.-F., Cai, W., Capotondi, A., Cobb, K. M., Lengaigne, M., McPhaden, M. J., Stuecker, M. F., et al.: El Niño–southern oscillation complexity, Nature, 559, 535–545, 2018.

Trenberth, K. E. and Stepaniak, D. P.: Indices of el Niño evolution, Journal of climate, 14, 1697–1701, 2001.

Vimont, D. J., Battisti, D. S., and Hirst, A. C.: The seasonal footprinting mechanism in the CSIRO general circulation models, Journal of
 Climate, 16, 2653–2667, 2003.

- 650 Xingjian, S., Chen, Z., Wang, H., Yeung, D.-Y., Wong, W.-K., and Woo, W.-c.: Convolutional LSTM network: A machine learning approach for precipitation nowcasting, in: Advances in neural information processing systems, pp. 802–810, 2015.
  - Yan, J., Mu, L., Wang, L., Ranjan, R., and Zomaya, A. Y.: Temporal convolutional networks for the advance prediction of ENSO, Scientific reports, 10, 1–15, 2020.

Yu, J.-Y. and Kao, H.-Y.: Decadal changes of ENSO persistence barrier in SST and ocean heat content indices: 1958–2001, Journal of

- 655 Geophysical Research: Atmospheres, 112, 2007.
  - Yu, J.-Y. and Kim, S. T.: Relationships between extratropical sea level pressure variations and the central Pacific and eastern Pacific types of ENSO, Journal of Climate, 24, 708–720, 2011.
  - Yu, J.-Y., Zou, Y., Kim, S. T., and Lee, T.: The changing impact of El Niño on US winter temperatures, Geophysical Research Letters, 39, 2012.
- 660 Zeiler, M. D. and Fergus, R.: Visualizing and understanding convolutional networks, in: European conference on computer vision, pp. 818– 833, Springer, 2014.
  - Zhang, J., Duan, W., and Zhi, X.: Using CMIP5 model outputs to investigate the initial errors that cause the "spring predictability barrier" for El Niño events, Science China Earth Sciences, 58, 685–696, 2015.

Zhang, R.-H. and Levitus, S.: Interannual variability of the coupled tropical Pacific ocean-atmosphere system associated with the El Niño-

- 665 Southern Oscillation, Journal of climate, 10, 1312–1330, 1997.
- Zinke, J., Browning, S., Hoell, A., and Goodwin, I.: The West Pacific Gradient tracks ENSO and zonal Pacific sea surface temperature gradient during the last Millennium, Scientific reports, 11, 1–16, 2021.

Vimont, D. J., Alexander, M. A., and Newman, M.: Optimal growth of central and east Pacific ENSO events, Geophysical Research Letters, 41, 4027–4034, 2014.

Wang, Z., Bovik, A. C., and Lu, L.: Why is image quality assessment so difficult?, in: 2002 IEEE International conference on acoustics, speech, and signal processing, vol. 4, pp. IV–3313, IEEE, 2002.