

An investigation into the processes controlling the global distribution of dissolved ^{231}Pa and ^{230}Th in the ocean and the sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios by using an ocean general circulation model COCO ver4.0

5

Yusuke Sasaki¹, Hidetaka Kobayashi¹, and Akira Oka¹

¹ Atmosphere and Ocean Research Institute, The University of Tokyo, Kashiwa, Japan

Correspondence to: Akira Oka (akira@aori.u-tokyo.ac.jp)

Abstract. Sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios provide clues to estimate the strength of past ocean circulation. For its estimation, understanding the processes controlling the distributions of ^{231}Pa and ^{230}Th in the ocean is important. However, simulations of dissolved and particulate ^{231}Pa and ^{230}Th in the modern ocean, recently obtained from the GEOTRACES project, remain challenging. Here we show an improved model simulation of ^{231}Pa and ^{230}Th in the global ocean by introducing bottom scavenging and the dependence of scavenging efficiency on particle concentration with water-column reversible scavenging. As demonstrated in a previous study, the incorporation of bottom scavenging improves the simulated distribution of dissolved ^{231}Pa and ^{230}Th in the deep ocean, which has been overestimated in models not considering the bottom scavenging. We further demonstrate that introducing the dependence of scavenging efficiency on particle concentration results in a high concentration of dissolved ^{230}Th in the Southern Ocean as observed in the GEOTRACES data. Our best simulation can well reproduce not only the oceanic distribution of ^{231}Pa and ^{230}Th but also the sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios. Sensitivity analysis reveals that oceanic advection of ^{231}Pa primarily determines sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios. On the other hand, ^{230}Th advection and bottom scavenging have an opposite effect to ^{231}Pa advection on the sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios, reducing their latitudinal contrast. Our best simulation shows the realistic residence times of ^{231}Pa and ^{230}Th , but simulation without bottom scavenging and dependence of scavenging efficiency on particle concentration significantly overestimates the residence times for both ^{231}Pa and ^{230}Th in spite of similar distribution of sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios to our best simulation.

1 Introduction

The $^{231}\text{Pa}/^{230}\text{Th}$ ratios in marine sediments are used for estimating past ocean circulation strength (Yu et al., 1996; McManus et al., 2004; Gherardi et al., 2009; Böhm et al., 2015; Waelbroeck et al. 2018; Sufke et al., 2020). Alpha decay of ^{235}U and ^{234}U produces ^{231}Pa (half-life of ~ 32.5 kyr) and ^{230}Th (half-life of ~ 75.2 kyr), respectively, at an approximately constant $^{231}\text{Pa}/^{230}\text{Th}$ ratio of 0.093 in the ocean (Henderson and Anderson, 2003). ^{231}Pa and ^{230}Th are absorbed onto and

desorbed from the surfaces of sinking particles (reversible scavenging; Bacon and Anderson, 1982) and eventually removed from the water column into marine sediments. Differential scavenging efficiencies of ^{231}Pa and ^{230}Th result in differences in their residence times in the ocean; the residence times of ^{231}Pa and ^{230}Th were estimated to be 111 and 26 years in Yu et al. (1996), and 130 and 20 years in Henderson and Anderson (2003). The shorter residence time of ^{230}Th indicates that ^{230}Th generated from ^{234}U is removed relatively quickly to marine sediments. On the other hand, the longer residence time of ^{231}Pa indicates that ^{231}Pa produced from ^{235}U is transported for a longer period by ocean advection and mixing. Therefore, the deviation of the sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios from the constant production ratio of 0.093 has been used as a proxy for ocean circulation (Yu et al., 1996). For example, the sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios from the Bermuda Rise were closer to 0.093 at the Last Glacial Maximum (LGM) than today, which suggests that the Atlantic meridional overturning circulation (AMOC) was weaker at the LGM (McManus et al., 2004; Böhm et al., 2015). Some modeling studies using a two-dimensional (2D) ocean model (Lippold et al., 2012) and a three-dimensional (3D) ocean model (Gu et al. 2020) attempted to simulate the compiled paleo data of sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ in the Atlantic during the LGM.

To use the sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios as a proxy for ocean circulation in a more quantitative manner, one needs to take into account the different scavenging efficiencies of different marine particle types (e.g., particulate organic carbon, calcite, and opal) as well as the distribution of these particles (Chase et al., 2002; Edwards et al., 2005). Sinking particles effectively scavenge ^{231}Pa and ^{230}Th in regions with high particle concentrations. In general, ^{231}Pa has a longer residence time than ^{230}Th , because sinking particles scavenge ^{230}Th more strongly. However, as for opal particles, Chase et al. (2002) argue that opal scavenges ^{231}Pa more effectively than ^{230}Th . This report is consistent with observational studies that find high $^{231}\text{Pa}/^{230}\text{Th}$ ratios in the Southern Ocean, where opal sinking flux is high (Chase et al., 2003).

Authors of previous modeling studies have tried to simulate the global distributions of ^{231}Pa and ^{230}Th by 2D ocean models (Marchal et al., 2000; Luo et al., 2010; Gu and Liu, 2017; Rempfer et al., 2017; van Hulst et al., 2018; Missiaen et al., 2020a) or 3D ocean models of LSG-OGCM (Henderson et al., 1999), Bern 3D (Siddall et al., 2005; ; Rempfer et al., 2017;), NEMO (Dutay et al., 2009; van Hulst et al., 2018), CESM (Gu and Liu, 2017) and iLOVECLIM (Missiaen et al., 2020a). There are also modeling studies that discuss the relationship between the strength of the AMOC and changes in sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios (Siddall et al., 2007; Gu and Liu, 2017; Missiaen et al., 2020a; 2020b). Siddall et al. (2005) pioneered the 3D simulation of both ^{231}Pa and ^{230}Th by incorporating reversible scavenging. Their control simulation appropriately reproduced the observed distribution of sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios; it showed high sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios in regions where the sinking opal particle flux is high. In their control simulation, the concentrations of dissolved ^{231}Pa and ^{230}Th increased linearly with depth; this pattern agreed broadly with observed features. However, simulated dissolved ^{231}Pa and ^{230}Th were both higher than observations in the deep ocean. In addition to reversible scavenging by sinking ocean particles, several studies (e.g., Anderson et al., 1983; Roy-Barman, 2009; Okubo et al., 2012) have pointed out the importance of additional scavenging at the seafloor (bottom scavenging) and the continental boundaries (boundary scavenging). The bottom scavenging has not been explicitly included in global 3D ocean models except for Rempfer et al. (2017) which used a simplified 3D ocean model of intermediate complexity similar to that used by Siddall et al. (2005) and reproduced the distributions of dissolved ^{231}Pa and

²³⁰Th more realistically by introducing the bottom scavenging. On the other hand, Henderson et al. (1999) reproduced the distribution of dissolved ²³⁰Th in their ocean general circulation model (OGCM) simulation by changing the efficiency of reversible scavenging depending on particle concentration; this effect has not been focused by recent modeling studies. Recently, the GEOTRACES project has led to a dramatic increase in the number of observations of dissolved and particulate ²³¹Pa and ²³⁰Th (Schlitzer et al., 2018). The GEOTRACES database provides an opportunity to test models describing the cycling of these two radioisotopes in the global ocean. This study reports the results of our OGCM simulations, which reproduce dissolved ²³¹Pa and ²³⁰Th more realistically than previous simulations by introducing bottom scavenging and the dependence of scavenging efficiency on particle concentration. Furthermore, we quantitatively discuss the processes that control the global distribution of sedimentary ²³¹Pa/²³⁰Th ratios; by performing a series of sensitivity simulations, we clarify how the individual processes (i.e., water-column reversible scavenging, ocean transport, and bottom scavenging) affect the global distribution of dissolved ²³¹Pa and ²³⁰Th and sedimentary ²³¹Pa/²³⁰Th ratios.

2 Materials and Methods

2.1 Ocean general circulation model

The OGCM used in this study is COCO version 4.0 (Hasumi, 2006), the ocean component of the coupled ocean-atmosphere general circulation model MIROC version 3.2 (K-1 Model Developers, 2004). The COCO is also used as the ocean part of MIROC earth system model (Hajima et al. 2020; Ohgaito et al., 2021). The model domain is global, with about one-degree horizontal resolution and 43 vertical layers. The vertical resolution varies from 5 (top) to 250 m (bottom). Surface boundary conditions are given from monthly averages of zonal and meridional components of wind stress, air temperature, specific humidity, net shortwave radiation, downward longwave radiation, freshwater flux, and wind speed. These boundary conditions are taken from the output of a pre-industrial simulation with MIROC (Kobayashi et al., 2015; Oka et al., 2012). To calculate ²³¹Pa and ²³⁰Th, we perform offline tracer simulation using physical fields obtained in advance by COCO (Oka et al., 2008, 2009). The tracer model is integrated for 3,000 years and tracer fields reach a steady state where change in ocean tracer inventory almost vanishes (less than 10⁻⁵% per 100 years). We analyze the average of the last 100 years of the integration.

The physical fields used in this study is based on MIROC climate model simulations, and its reproducibility has been discussed and confirmed in a variety of literature (e.g., K-1 Model Developers, 2004; Gregory et al., 2005; Oka et al., 2006; Stouffer et al., 2006). We also note that the physical fields used here are the same as the pre-industrial (PI) simulation reported in Kobayashi et al. (2015) and Kobayashi and Oka (2018). For reference, the Atlantic meridional overturning circulation (AMOC) simulated by the COCO used in this study is shown in Fig. S11.

2.2 Particle fields

Following Siddall et al. (2005), the distribution of biogenic particles (organic carbon, calcite, and opal) is used to evaluate the scavenging of both ²³¹Pa and ²³⁰Th. We define the concentration M of each particle type [g m⁻³] as $M = F/w_s$.

where F is the particle flux [$\text{g m}^{-2} \text{yr}^{-1}$] and w_s is the constant settling velocity [m yr^{-1}]. The vertical particle flux is calculated using the export flux from the euphotic zone and an assumed vertical profile of each particle type. The detailed procedure is explained below.

First, the particulate organic carbon (POC) export flux from the euphotic zone is calculated by multiplying the distribution of primary production derived from satellite observations (Behrenfeld and Falkowski, 1997) by the export ratio (Dunne et al., 2005). From POC export flux and $M = F/w_s$, the concentration of POC at the base of the euphotic zone, $M_{\text{POC}}(z_0)$, where z_0 is the depth of the bottom of the euphotic zone, is obtained. After obtaining $M_{\text{POC}}(z_0)$, the POC concentration in the water column is expressed (Marchal et al., 1998) as

$$M_{\text{POC}} = M_{\text{POC}}(z_0) \left(\frac{z}{z_0}\right)^{-\varepsilon}, \quad (1)$$

where ε is a remineralization exponent for POC.

Next, the calcite and opal export fluxes from the euphotic zone are calculated by multiplying the POC export flux by their rain ratios, which are estimated following formulations of Siddall et al. (2005) and Maier-Reimer (1993); please refer to Eq. (2)–(5) of Siddall et al. (2005) for detail. The calcite particle concentration is calculated by assuming an exponentially decreasing vertical profile (Henderson et al., 1999; Marchal et al., 2000; Siddall et al., 2005). Thus, we have

$$M_{\text{CaCO}_3} = M_{\text{CaCO}_3}(z_0) \exp\left(\frac{z_0 - z}{z_p}\right), \quad (2)$$

where z_p is the calcite penetration depth. While the opal concentration is expressed as an exponentially decreasing vertical profile in some previous studies (e.g., Henderson et al., 1999), we consider opal dissolution to be dependent on temperature, following Siddall et al. (2005), as

$$M_{\text{opal}} = M_{\text{opal}}(z_0) \exp\left[\frac{D_{\text{opal}}(z_0 - z)}{w_s}\right], \quad (3a)$$

$$D_{\text{opal}} = B(T - T_0), \quad (3b)$$

where D_{opal} [yr^{-1}] is the opal dissolution rate, T_0 is the minimum temperature [$^{\circ}\text{C}$] of seawater in the model, and B is a dissolution constant [$^{\circ}\text{C}^{-1} \text{yr}^{-1}$]. Table 1 lists the parameter values used in this study. Figure S10 shows particle fluxes in the surface ocean.

2.3 Reversible scavenging model

We use a tracer model of ^{231}Pa and ^{230}Th based on Siddall et al. (2005). The dissolved concentration (A_d) and particle concentration (A_p) of ^{231}Pa and ^{230}Th are calculated from the following equations:

$$\frac{\partial A_{\text{total}}^i}{\partial t} = \beta^i - \lambda^i A_{\text{total}}^i - w_s \frac{\partial A_p^i}{\partial z} + \text{Transport}, \quad (4a)$$

$$A_{\text{total}}^i = A_p^i + A_d^i. \quad (4b)$$

In Eq. (4a), the first term on the right-hand side (β^i) represents production from uranium (^{231}Pa from ^{235}U ; ^{230}Th from ^{234}U), the second term represents radioactive decay, the third term represents the effect of vertical particle settling, and the fourth

term represents ocean transport by advection, diffusion, and convection. The superscript i represents the isotope type (^{231}Pa , ^{230}Th).

By following a reversible scavenging model (Bacon and Anderson, 1982), the relationship between the radionuclide concentration in the dissolved phase (A_d^i) and particulate phase (A_p^i) is represented by the partition coefficient K_j^i as

$$A_p^i = \sum_j A_{j,p}^i, \quad (5a)$$

$$K_j^i = \frac{A_{j,p}^i}{A_d^i \cdot C_j}, \quad (5b)$$

where subscript j represents the particle type (organic carbon, calcite, opal) and C_j is the dimensionless ratio of particle concentration to the density of seawater. The formulation of the reversible scavenging was also described in Oka et al., (2009, 2021) and readers can obtain its detailed description therein. The partition coefficient depends on the type of particles (Siddall et al., 2005). The partition coefficients of ^{231}Pa and ^{230}Th for each type of particle have been estimated in previous studies (Luo and Ku, 1999; Chase et al., 2002). Chase et al. (2002) show that opal scavenges ^{231}Pa more efficiently than ^{230}Th , whereas calcite scavenges ^{230}Th more efficiently than ^{231}Pa . Here we use partition coefficients following Chase and Anderson (2004), as in other previous modeling studies (Dutay et al., 2009; Gu and Liu, 2017; Siddall et al., 2005; Table 2).

2.4 One-dimensional reversible scavenging model

In addition to the three-dimensional tracer model based on the OGCM, we use a simple, vertical, one-dimensional model to analyze simulation results in Section 4. In the one-dimensional model, we assume a steady state and ignore the effect of ocean transport in Eq. (4a). Furthermore, we do not take the radioactive decay term into account because it is much smaller than the production term. Under these assumptions, Eq. (4a) becomes

$$\beta^i - w_s \frac{\partial A_p^i}{\partial z} = 0. \quad (6)$$

In this one-dimensional model, production by uranium radioactive decay (the first term on the left side of Eq. (6)) is balanced by vertical transport through particle settling (the second term on the left side of Eq. (6)). If we assume that A_p^i is zero at the sea surface ($z = 0$), then Eq. (6) can be solved, leading to

$$A_p^i = \frac{\beta^i}{w_s} \cdot z. \quad (7)$$

From Eq. (5), we have

$$\begin{aligned} A_p^i &= \sum_j A_{j,p}^i = (K_{\text{CaCO}_3}^i \cdot C_{\text{CaCO}_3} + K_{\text{opal}}^i \cdot C_{\text{opal}} + K_{\text{POC}}^i \cdot C_{\text{POC}}) \cdot A_d^i \\ &= \sum_j (K_j^i \cdot C_j) \cdot A_d^i. \end{aligned} \quad (8)$$

The dissolved concentration can be obtained from Eq. (7) and (8):

$$A_d^i = \frac{1}{\sum_j (K_j^i \cdot C_j)} A_p^i$$

$$= \frac{1}{\sum_j (K_j^i \cdot c_j)} w_s \cdot z. \quad (9)$$

Equation (9) shows that the vertical profile of A_d^i is determined only by the particle settling speed, the partition coefficients, and the concentrations of each particle. By comparing results from the one-dimensional model and the three-dimensional tracer model, we can isolate the influence of ocean transport (i.e., advection, diffusion, and convection) on the simulated distributions of dissolved ^{231}Pa and ^{230}Th (see Section 4; Table 3).

2.5 Experimental design

This study conducts a series of OGCM experiments. First, we perform an experiment named Siddall_EXP using the same parameters and formulations as in Siddall et al. (2005).

Second, we perform an experiment named BTM_EXP, in which we additionally take bottom scavenging into account. In BTM_EXP, we assume a globally uniform concentration of lithogenic particles in the deepest layer of the OGCM, following Rempfer et al. (2017). The intensity of the bottom scavenging depends on two parameters: the partition coefficient for lithogenic particles (K_{bottom}) and the concentration of lithogenic particles (C_{bottom}). The value of C_{bottom} is taken from Rempfer et al. (2017): $C_{\text{bottom}} = 6.0 \times 10^{-8} \text{ g cm}^{-3}$. This value is within the range of values from 4.0×10^{-8} to $1.65 \times 10^{-6} \text{ g cm}^{-3}$ observed in the benthic nepheloid layers (50-130 m above the bottom) in the North Atlantic as reported by Lam et al. (2015). As for K_{bottom} , because our formulation of the reversible scavenging is not the same as Rempfer et al. (2017), we needed to find its appropriate parameter value. For this purpose, we perform a number of simulations with different bottom scavenging intensities by changing the value of K_{bottom} .

Third, we perform a sensitivity experiment named KREF_EXP concerned with the reference partition coefficient (K_{ref}). In KREF_EXP, in addition to varying the partition coefficient for lithogenic particles (K_{bottom}), we also vary the reference partition coefficients (K_{ref}) from the values assumed in Siddall_EXP and BTM_EXP.

Finally, we perform an experiment named PCE_EXP, in which we incorporate the dependence of scavenging efficiency on particle concentration. In PCE_EXP, K_{ref} is not assumed to be constant but varies according to the following formulation of Henderson et al. (1999):

$$K_{\text{ref}} = \left(\frac{C_{\text{total}}}{C_{\text{ref}}} \right)^{-0.47} \times 10^7, \quad (10)$$

where C_{total} [g cm^{-3}] is the total concentration of all sinking particles ($C_{\text{total}} = C_{\text{CaCO}_3} + C_{\text{opal}} + C_{\text{POC}}$) and C_{ref} [g cm^{-3}] is the reference concentration. Note that the value of C_{total} is differently specified on each grid whereas C_{ref} is given as a globally uniform value. Due to the dependence of K_{ref} on C_{total} , the scavenging efficiency becomes lower under higher particle concentrations and higher under lower particle concentrations. We conduct several simulations by varying C_{ref} between 10^{-9} and $10^{-6} \text{ g cm}^{-3}$ (smaller C_{ref} value leads to stronger scavenging). Although the observed decrease of the partition coefficient with increased bulk particle concentration is not entirely understood (Pavia et al., 2018), we will show that this

particle concentration effect becomes essential for controlling dissolved ^{230}Th in some ocean regions.

3 Results

3.1 Dissolved ^{231}Pa and ^{230}Th along the Atlantic meridional transects

185 First, we discuss the results of Siddall_EXP, focusing on the meridional distribution of ^{231}Pa and ^{230}Th in the Atlantic Ocean. Figure 1 shows the dissolved concentrations of ^{231}Pa and ^{230}Th simulated in Siddall_EXP along the Atlantic 30°W transect, together with GEOTRACES data (see Fig. S1 for the location of observations referenced in this study). We confirm that the distributions of dissolved ^{231}Pa and ^{230}Th in Siddall_EXP are approximately the same as those reported in Siddall et al. (2005; their Fig. 2). Because ^{231}Pa and ^{230}Th exchange reversibly with sinking particles and are transported to the deep
190 ocean, the dissolved ^{231}Pa and ^{230}Th concentrations increase with depth, both in the model simulation and in observations. However, as in Siddall et al. (2005), the model simulation overestimates dissolved ^{231}Pa and ^{230}Th concentrations at depths greater than 2,000 m and 1,000 m, respectively. For quantitative analysis, we perform a linear regression analysis between the simulation results and observed data from the GEOTRACES GA02 transect; we calculate the root mean square deviation (RMSD), the correlation coefficient (R), and the slope of the linear regression (s) of modeled concentration versus measured
195 concentration, as summarized in Table S1. The linear regression line slope indicates the model's ability to reproduce the observed distribution; it approaches 1.0 when the model simulation realistically reproduces the target distribution (Dutay et al., 2009; Gu and Liu, 2017). For Siddall_EXP, the slope of linear regression line is significantly larger than 1.0 for both ^{231}Pa (s=1.88, R=0.72 and RMSD=0.15) and ^{230}Th (s=4.44, R=0.89 and RMSD=1.31; Table S1). This overestimation in the deep ocean is also found in other previous model simulations (e.g., Dutay et al., 2009; Gu and Liu, 2017; van Hulst et al., 2018).

200 Next, to reduce the overestimation of the simulated concentrations in the deep ocean, we additionally incorporate bottom scavenging in benthic nepheloid layers (BTM_EXP). The dissolved ^{231}Pa and ^{230}Th distributions are shown in Fig. 2 and 3, respectively. As expected, the incorporation of bottom scavenging helps reduce ^{231}Pa and ^{230}Th concentrations in the deep ocean, improving the model's agreement with the data. As for the distribution of dissolved ^{231}Pa , the model results come relatively close to the GEOTRACES data if $K_{\text{bottom}}^{\text{Pa}}$ is set to 5.0×10^5 (s=1.04, R=0.90 and RMSD=0.05; see CTRL_EXP
205 in Table S1; Fig. 2c and 2d). This result confirmed the importance of the bottom scavenging, which were already reported from a previous global 3D model (Rempfer et al., 2017) and a regional eddy-permitting model (Lerner et al., 2020). On the other hand, it is difficult to reproduce the observed distribution of dissolved ^{230}Th in BTM_EXP. With $K_{\text{bottom}}^{\text{Th}} = 1.0 \times 10^6$, the concentrations of ^{230}Th in bottom waters come close to observed values (Fig. 3c and 3d), but the concentrations in the deep ocean (from 2000 m to 5000 m) remain overestimated. In the case of larger $K_{\text{bottom}}^{\text{Th}}$, the simulated ^{230}Th concentrations
210 approach observed values in the deep ocean but are significantly lower than measurements in bottom waters (e.g., $K_{\text{bottom}}^{\text{Th}} = 1.0 \times 10^7$ in Fig. 3g and 3h). These results indicate that modification of Siddall_EXP by considering bottom scavenging alone is not sufficient for accurately simulating ^{230}Th distribution in our model. As shown in Rempfer et al. (2017) and Lerner et al.

(2020), the appropriate selection of scavenging parameter coefficients is required for more realistic simulation. Because our reversible scavenging model (which is the same as Siddall et al., 2005; section 2.3) is not the same as Rempfer et al. (2017) and Lerner et al. (2020), we need to discuss the validity of a scavenging parameter coefficient in our model (i.e., K_{ref}). In the following experiments (i.e., KREF_EXP and PCE_EXP), we discuss about more appropriate treatment about K_{ref} by focusing solely on ^{230}Th .

To reproduce the distribution of ^{230}Th more realistically, we change the value of the reference partition coefficient (K_{ref}^{Th}) in addition to K_{bottom}^{Th} (KREF_EXP). Figure 4 summarizes the results of KREF_EXP and shows the simulated vertical distributions of dissolved ^{230}Th for various values of K_{ref}^{Th} and K_{bottom}^{Th} (see Fig. 4g). Note that, for example, the simulation R2_B5 means that K_{ref}^{Th} is set to 2.0×10^7 and K_{bottom}^{Th} to 5.0×10^5 . In the cases where K_{bottom}^{Th} is set to 5.0×10^5 (namely R2_B5, R4_B5, and R6_B5), the ^{230}Th concentrations systematically change depending on K_{ref}^{Th} ; as the reversible scavenging on sinking particles becomes stronger (i.e., for larger K_{ref}^{Th}), the concentrations of dissolved ^{230}Th become smaller throughout the water column (Fig. 4c, 4e, and 4f). As discussed for BTM_EXP, it is also confirmed that the stronger bottom scavenging (i.e., larger K_{bottom}^{Th}), the lower the concentrations near the sea bottom (e.g., see R2_B5, R2_B10, and R2_B20). For some combinations of water-column scavenging and bottom scavenging, simulations (e.g., R6_B5, R4_B5, R4_B10) reasonably reproduce the observed profile of dissolved ^{230}Th concentration. Among our KREF_EXP simulations, the R6_B5 simulation (Fig. 4f) shows the slope of the linear regression line nearest to 1.0 ($s=0.88$, $R=0.81$, and $\text{RMSD}=0.20$; Table S1) where K_{ref}^{Th} is higher ($K_{ref}^{Th} = 6.0 \times 10^7$) than for Siddall_EXP and BTM_EXP ($K_{ref}^{Th} = 1.0 \times 10^7$). In the R6_B5 simulation (Fig. 4f), the vertical profile of dissolved ^{230}Th is significantly improved from that of Siddall_EXP (Fig. 1d) and BTM_EXP (Fig. 3). We confirmed that the R6_B5 simulation captures the observed features of the Atlantic transects of the GEOTRACES data (Fig. 5a). However, the R6_B5 simulation still underestimates the concentrations of dissolved ^{230}Th from the surface to intermediate depths (see Fig. 4f). Also, the high concentrations of dissolved ^{230}Th observed in the Southern Ocean in GEOTRACES data are not well reproduced (Fig. 5a). To address this issue, we performed additional simulations by slightly changing the values of K_{ref}^{Th} and K_{bottom}^{Th} from the R6_B5 simulation (not shown), but found that it is difficult to remove the preceding deficiencies by merely changing the values of K_{ref}^{Th} and K_{bottom}^{Th} in KREF_EXP.

Finally, we discuss PCE_EXP, in which the dependence of scavenging efficiency on particle concentration is taken into account, according to Eq. (10). We conduct several simulations by varying the value of the reference concentration (C_{ref}) between 10^{-9} and 10^{-6} g cm^{-3} . Among these results, we here discuss the case with $C_{ref} = 10^{-7}$ g cm^{-3} , which shows the best agreement with observations. Compared to the case in which the dependence of scavenging efficiency on particle concentration is not considered (i.e., R6_B5 simulation of KREF_EXP), PCE_EXP is expected to show smaller (larger) K_{ref}^{Th} for the higher (lower) concentration of sinking particles. In Fig. 5, we compare the simulated dissolved ^{230}Th distribution obtained from PCE_EXP and R6_B5 simulation of KREF_EXP. Owing to the dependence of scavenging efficiency on particle concentration, PCE_EXP reproduces the vertical distribution of dissolved ^{230}Th slightly better than KREF_EXP (Fig. 5d). The regression

245 analysis also confirms that the agreement with the GEOTRACES data becomes improved in PCE_EXP ($r_s=0.98$ and $R=0.84$;
CTL_EXP in Table S1). It is worthy to note that the distribution in the Southern Ocean is significantly improved in PCE_EXP
(Fig. 5b) compared to KREF_EXP (Fig. 5a) as a result of the non-uniform distribution of the reference partition coefficient
 K_{ref}^{Th} (Fig. 5c). In the Southern Ocean, where particle concentration is relatively higher than in other regions (Honjo et al.,
2008), the value of K_{ref}^{Th} in PCE_EXP is lower than that in the R6_B5 simulation of KREF_EXP ($K_{ref}^{Th} = 6 \times 10^7$; i.e.,
250 $\log_{10} K_{ref}^{Th} \sim 7.8$) (Fig. 5c). Therefore, the concentration of dissolved ^{230}Th in PCE_EXP becomes high compared to the
KREF_EXP, which leads to more realistic distribution of dissolved ^{230}Th in the Southern Ocean. Hereafter, our best simulation
(i.e., $K_{bottom}^{Pa} = 5.0 \times 10^5$ case of BTM_EXP for ^{231}Pa and PCE_EXP for ^{230}Th) is called CTRL_EXP (see Table 2 for
parameter values of CTRL_EXP).

3.3 Particulate ^{231}Pa and ^{230}Th

255 By conducting a series of experiments described above, this study successfully reproduces the observed distributions
of dissolved ^{231}Pa and ^{230}Th , shown again in Fig. 6a and 6b, respectively. In addition to dissolved ^{231}Pa and ^{230}Th , particulate
 ^{231}Pa and ^{230}Th simulated in CTRL_EXP are compared with the reported observations (Fig. 6c and 6d). In addition to the
GEOTRACES dataset, we use several reported observations here (i.e., data referenced in Siddall et al., 2005, Marchal et al.,
2007, and Lerner et al., 2020; namely, from Colley et al., 1995; Moran et al., 1997; Moran et al., 2001; Rutgers van der Loeff
260 and Berger, 1993; Vogler et al., 1998; Walter et al., 1997; Cochran et al., 1987; Moran et al., 2002; Guo et al., 1995). The
model captures the observed tendency that the concentration becomes higher in the high-latitude Southern Ocean, as reported
in previous studies (e.g., see Fig. 2 in Siddall et al. 2005). The ratio of ^{231}Pa to ^{230}Th in the particulate phase in the water column
shows low concentrations in the deep ocean at low latitudes and high in the intermediate ocean at mid-latitudes and the
Southern Ocean (Fig. 6f). This feature is consistent with observational findings and recent modeling studies (e.g., Fig. 2 in Gu
265 and Liu 2017; Fig. 3 in Rempfer et al., 2017). Although the number of available observations is limited for the particulate
phase, it is confirmed that our simulation reasonably reproduces observed distributions for both dissolved and particulate
phases.

3.4 Sedimentary $^{231}Pa/^{230}Th$ ratios

Our CTRL_EXP also well reproduces the global distribution of sedimentary $^{231}Pa/^{230}Th$ ratios (Fig. 6e) compared
270 with the reported observations (Mangianini & Sonntag, 1977; Muller & Mangini, 1980; Anderson et al., 1983; Shimmield et
al., 1986; Schmitz et al., 1986; Yang et al., 1986; Shimmield & Price, 1988; Yong Lao et al., 1992; François et al., 1993; Frank
et al., 1994; Frank, 1996; Bradtmiller et al., 2014, Luo et al., 2010, and their supplemental data). Sedimentary $^{231}Pa/^{230}Th$ ratios
are high along the margin of the North Pacific and the North Atlantic, as well as in the Southern Ocean, where particle
concentrations are high. On the other hand, sedimentary $^{231}Pa/^{230}Th$ ratios are low in the low-latitude regions, including
275 subtropical gyres, where particle concentrations are low. These simulated features are consistent with observations (circles in

Fig. 6e). Previous modeling studies reported the similar distribution of sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios (e.g., Fig. 2 in Siddall et al., 2005; Fig. 11 in Dutay et al., 2009; Fig. 4 in Gu and Liu, 2017; Fig. 10 in Hulten et al., 2018; Fig. 4 in Messiaen et al., 2020) and our Siddall_EXP also reasonably reproduced the global distribution of sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios (Fig. S4a). However, as shown above, the distributions of dissolved ^{231}Pa and ^{230}Th in the ocean are significantly different between CTRL_EXP and Siddall_EXP. Thus, each experiment implies a different set of processes controlling the distribution of sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios. We will discuss this point later in the next section.

4 Discussion

4.1 Comparison with previous modeling studies

We demonstrated that our CTRL_EXP can reproduce more realistic distribution of dissolved ^{231}Pa and ^{230}Th along the Atlantic meridional transects than Siddall_EXP by considering the bottom scavenging and the dependence of scavenging efficiency on particle concentration. Here, we compared our results with previous modeling studies which showed their model results along with Atlantic meridional transects (GEOTRACES GA02 section).

As far as we know, Rempfer et al. (2017) was the only 3D global ocean model which introduces the bottom scavenging, and in our study, we introduced the bottom scavenging into the global OGCM for the first time. Models without the bottom scavenging tend to overestimate the dissolved ^{231}Pa and ^{230}Th in the deep ocean as in our Siddall_EXP. For example, in Gu and Liu (2017) in which ^{231}Pa and ^{230}Th tracer are introduced into CESM1.3, their simulated ^{231}Pa and ^{230}Th concentrations are significantly overestimated in the deep ocean along the GEOTRACES GA02 section (their Fig. 2). In Dutay et al. (2009) in which ^{231}Pa and ^{230}Th tracer are introduced into NEMO-PISCES, influences of particle size and type on ^{231}Pa and ^{230}Th are discussed by performing several sensitivity simulations but all of their simulations overestimate ^{231}Pa and ^{230}Th concentrations in the deep Atlantic Ocean (their Figs 4 and 5, respectively). In van Hulten et al. (2018) which was the updated ^{231}Pa and ^{230}Th simulation with NEMO-PISCES, the model still overestimates ^{231}Pa and ^{230}Th concentrations in the deep Atlantic Ocean (their Fig. 12) because particles in the nepheloid layers (i.e., bottom scavenging) are not included in their model.

Although the incorporation of bottom scavenging is important for controlling the scavenging efficiency, it is worthy to note that bottom scavenging is not the sole process that controls the scavenging efficiency. Therefore, for example, the model which specified the relatively stronger affinity to the particle can lead to smaller tracer concentration even if the model does not include the bottom scavenging. In fact, in Messiaen et al. (2020), their simulated ^{231}Pa and ^{230}Th are underestimated in both the upper and deep oceans (their Supplementary Fig. 3) even if the bottom scavenging was not included in their model. This is because their specified scavenging parameters are relatively stronger than the other models (see their Table 2) as a result of their parameter tuning without the bottom scavenging.

In Re3d_Bt_Bd simulation reported in Rempfer et al. (2017) where the bottom scavenging process is considered, the above-mentioned overestimation in the deep ocean was relaxed and their simulated distribution appears similar to our CTRL_EXP. Their study is the first 3D model demonstration about the importance of the bottom scavenging process, which

is confirmed again in our study (e.g., from comparison with Siddall_EXP and KREF_EXP). However, their model still tends to somewhat overestimate the dissolved ^{231}Pa compared with GEOTRACES GA02 data (their Fig.3). Because the formulation of the reversible scavenging and their model parameters are not the same as our CTRL_EXP, we expect that different choice of model parameter values leads to such differences; more specifically, our choice of $K_{\text{ref}}^{\text{Pa}}$ is based on Chase et al. (2002) and Siddall et al. (2005) whereas the scavenging efficiency parameters in Rempfer et al. (2017) are similar to those in Luo et al. (2010) and Marshall et al. (2002). In addition, as for ^{230}Th , the high concentration in the Southern Ocean is not reproduced in their model, whereas this is reproduced in our CTRL_EXP by considering the dependence of scavenging efficiency on particle concentration. Although the dependence of scavenging efficiency on particle concentration was already introduced in Henderson et al. (1995), our study demonstrates its importance for reproducing high concentration in the Southern Ocean reported in GEOTRACES GA02 data for the first time.

4.2 Reproducibility along GEOTRACES GA03 and GP16 transects

So far, we have compared our model results with observations by focusing on the Atlantic meridional GEOTRACES transects (i.e., GA02 and GIPY05). Here, we will compare our CTRL_EXP with other available GEOTRACES transects: GA03 in the subtropical North Atlantic (Hayes et al., 2015) and GP16 in the South Pacific (Pavia et al., 2018).

Figure S5 shows the results of CTRL_EXP along with the GEOTRACES GA03 data. For dissolved ^{231}Pa , the model shows a high concentration around a depth of about 3000 m and higher concentrations on the eastern/southern side of the basin as in observations (Fig. S5a). This feature was also well reproduced in Re3d_Bt_Bd simulation of Rempfer et al. (2017) as shown in their Figure 2, but not in other previous models (e.g., Fig. 8 in van Hulst et al., 2018; Fig. 3 in Gu and Li, 2017). This confirms that the consideration of the bottom scavenging is helpful for improving the model result along the GEOTRACES GA03 section. For dissolved ^{230}Th , features similar to ^{231}Pa are also found in both the model and observations although the model appears to underestimate north-south or western-eastern differences (Fig. S5b). For particulate ^{231}Pa and ^{230}Th , the model tends to simulate high concentration near the sea bottom and the continental margins where the particle concentration becomes high, but such features are not necessarily clear in the GEOTRACES data (Fig. S5c and S5d). Our model may not sufficiently reproduce the bottom and boundary scavenging associated with terrestrial particles in this region. More sophisticated treatment of bottom and boundary scavenging might be required for addressing these issues.

Figure S6 shows the results of CTRL_EXP along with the GEOTRACES GP16 data. As with the other section data, CTRL_EXP approximately reproduces the distribution of ^{231}Pa and ^{230}Th . In this transect, the observational data shows a clear signal associated with hydrothermal vents: low concentrations of dissolved ^{231}Pa and ^{230}Th and high concentrations of particulate ^{231}Pa and ^{230}Th , which are not simulated in our model. Processes related to the hydrothermal vents are not explicitly incorporated in the present ^{231}Pa and ^{230}Th model simulations; its detailed treatment is beyond our scope but appears necessary for more realistic simulations.

340 4.3 Processes controlling sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios

In this subsection, we discuss about the processes controlling the global distribution of sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios. For this purpose, we decompose the processes controlling sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios simulated in our best simulation CTRL_EXP into three parts: water-column reversible scavenging, three-dimensional ocean transport, and bottom scavenging. To evaluate how these three processes affect the distribution of $^{231}\text{Pa}/^{230}\text{Th}$ ratios, we conduct two additional experiments (see Table 3). The first experiment is 3D_EXP, which is the same as CTRL_EXP except that bottom scavenging is not taken into account (i.e., we set $K_{\text{bottom}}^{\text{Pa}} = K_{\text{bottom}}^{\text{Th}} = 0$ in 3D_EXP). The second is 1D_EXP, which is the one-dimensional reversible scavenging model experiment described in Section 2.4. The tracer distribution in 1D_EXP is determined solely by the one-dimensional vertical process of reversible scavenging; the strength of scavenging changes spatially through changes in the partition coefficient (K_j^i of Eq. (9) in Section 2.4) that depends on the specified three-dimensional particle concentration (C_j of Eq. (9)). By using results of CTRL_EXP, 3D_EXP, and 1D_EXP, we can extract the influence of three processes: the influence of the one-dimensional vertical reversible scavenging is revealed by 1D_EXP, the influence of bottom scavenging is revealed by the difference between CTRL_EXP and 3D_EXP, and the influence of ocean transport is revealed by the difference between 3D_EXP and 1D_EXP. When we focus on sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios, each process described above can be further examined for ^{231}Pa and ^{230}Th individually. For example, the difference in $^{231}\text{Pa}/^{230}\text{Th}$ ratios between CTRL_EXP and 3D_EXP represents the influence of bottom scavenging of both ^{231}Pa and ^{230}Th , whereas the influence of bottom scavenging of ^{231}Pa alone can also be evaluated from CTRL_EXP and 3D_EXP (i.e., $^{231}\text{Pa}(\text{CTRL})/^{230}\text{Th}(3\text{D})$ minus $^{231}\text{Pa}(3\text{D})/^{230}\text{Th}(3\text{D})$).

In 1D_EXP, the particulate concentration is obtained from Eq. (7); the particulate concentration increases linearly with depth (Fig. S2c and S2d). The dissolved concentration is calculated from Eq. (9), suggesting that the concentration becomes higher for a lower partition coefficient (K_j^i in Eq. (9)) and for a lower particle concentration (C_j in Eq. (9)). Mainly due to the dependency on C_j , the dissolved concentration becomes higher (lower) in the area with lower (higher) particle concentration in 1D_EXP. As a result, the dissolved concentration becomes very high in the deeper ocean, where the particle concentration becomes lower for both ^{231}Pa and ^{230}Th (Fig. S2a and S2b). It is interesting to point out that the spatial pattern of dissolved ^{231}Pa and ^{230}Th (Fig. S2a and S2b) is similar to that of K_{ref} in PCE_EXP (Fig. 5c) because both are affected by the amount of particle concentration. More importantly, we emphasize here that the sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios in 1D_EXP become uniform everywhere (0.093; Fig. S2e) because, as confirmed from Eq. (7), the ratio of particulate ^{231}Pa to particulate ^{230}Th amounts everywhere to $\beta^{\text{Pa}}\beta^{\text{Th}} = 0.093$, regardless of geographic location (Fig. S2f).

In 3D_EXP, three-dimensional ocean transport operates, in addition to water-column scavenging considered in 1D_EXP (Fig. S3). As described above, the influence of ocean transport can be evaluated from the difference between 3D_EXP and 1D_EXP (Fig. 7). On the other hand, the influence of bottom scavenging can be obtained from the difference between CTRL_EXP and 3D_EXP (Fig. 8). Note again that since the sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios in 1D_EXP are globally uniform ($^{231}\text{Pa}/^{230}\text{Th} = 0.093$), their spatial distribution is controlled not by the one-dimensional vertical process but by the ocean

transport. Figures 7e and 7f demonstrate that the ocean transport effect captures the overall features of CTRL_EXP (Figs. 6e and 6f). On the other hand, bottom scavenging tends to cancel the effects of ocean transport and weaken the spatial contrast of $^{231}\text{Pa}/^{230}\text{Th}$ ratios simulated in CTRL_EXP (Figs. 8e and 8f).

To evaluate the above processes controlling the sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios in more detail, we further decompose the ocean transport contribution into those from ^{231}Pa and ^{230}Th , separately (Fig. 9a for ^{231}Pa and 9b for ^{230}Th). Similarly, we further decompose the contribution of bottom scavenging into those for ^{231}Pa and ^{230}Th (Fig. 9c and 9d, respectively). In Fig. 9a, we demonstrate that ocean transport solely from ^{231}Pa (i.e., $^{231}\text{Pa}(3\text{D})/^{230}\text{Th}(1\text{D})$) can reproduce the overall distribution of the sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios in CTRL_EXP (Fig. 6e). This result confirms that ocean transport of ^{231}Pa primarily controls the distribution of sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios, consistent with previous studies (Yu et al., 1996; Marchal et al., 2000). These previous studies suggest that the distribution of ^{231}Pa mainly determines the global distribution of sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios because the residence time of ^{231}Pa is longer than that of ^{230}Th .

Here, we further discuss how the ocean transport of ^{231}Pa controls the distribution of sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios. Since changes in sedimentary ^{231}Pa correspond to particulate ^{231}Pa changes in the bottom ocean, we focus the ocean transport effect on particulate ^{231}Pa (Fig. 7c). Consistent with Fig. 9a, Fig. 7c indicates that ocean transport acts to decrease (increase) particulate ^{231}Pa in lower (higher) latitudes. We also found that particulate ^{231}Pa changes (Fig. 7c) are similar to those in dissolved ^{231}Pa (Fig. 7a). Because most of ^{231}Pa are in the dissolved phase, the advection of particulate ^{231}Pa itself is very small compared with that of dissolved ^{231}Pa and ocean transport takes place mainly in the form of dissolved ^{231}Pa . Therefore, it is interpreted that ocean transport first controls the dissolved ^{231}Pa , and then the corresponding changes in particulate ^{231}Pa take place so that the relationship between dissolved and particulate ^{231}Pa (i.e., Eq. (5b)) is satisfied. In other words, the changes in particulate ^{231}Pa take place as a result of changes in dissolved ^{231}Pa . Therefore, we need to focus on the processes that control the dissolved ^{231}Pa changes (Fig. 7a). As previously mentioned, in the case of no ocean transport (i.e., 1D_EXP), the dissolved ^{231}Pa concentration near the seabed in lower latitudes becomes very high (Fig. S2a). Ocean transport, which includes both advection and diffusion, reduces high concentrations of dissolved ^{231}Pa in low latitude oceans by transporting dissolved ^{231}Pa from lower latitudes to higher latitudes. As a result of the change in the dissolved ^{231}Pa (Fig. 7a), the changes in particulate ^{231}Pa (Fig. 7c) also take place by satisfying Eq. (5b); this leads to lower sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios in lower latitudes and higher ratios in higher latitudes (Figs. 7e and 7f).

Contrary to ^{231}Pa , the influences of ^{230}Th transport on sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios have been usually regarded as small because ^{230}Th is generally assumed to be scavenged very quickly everywhere. However, our results demonstrate that ocean transport of ^{230}Th also affects the distribution of sediment $^{231}\text{Pa}/^{230}\text{Th}$ to some extent. As a matter of course, ^{230}Th ocean transport acts in the opposite direction of ^{231}Pa ocean transport, reducing the spatial contrast in sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios (Fig. 9b). However, an exception is found in the Southern Ocean, where the ^{230}Th ocean transport contributes to higher sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios, in the same way as the ^{231}Pa ocean transport. Because opal scavenges ^{231}Pa more effectively than ^{230}Th (Chase et al., 2002), ^{231}Pa transported toward the Southern Ocean is expected to be immediately removed there due to the high opal flux. Therefore, previous studies concluded that ocean transport of ^{231}Pa explains high sedimentary $^{231}\text{Pa}/^{230}\text{Th}$

ratios in the Southern Ocean. On the other hand, in addition to ocean transport of ^{231}Pa , our results suggest that ocean transport of ^{230}Th also contributes to the high $^{231}\text{Pa}/^{230}\text{Th}$ ratios in the Southern Ocean. This result implies that scavenging of ^{230}Th is not so efficient in the Southern Ocean as previously expected due to the dependence of scavenging efficiency on particle concentration. This interpretation is consistent with the high concentration of dissolved ^{230}Th in the Southern Ocean (Fig. 6b).

Bottom scavenging promotes the removal of both ^{231}Pa and ^{230}Th near the seafloor and tends to cancel the influence of ocean transport. Namely, the bottom scavenging of ^{231}Pa reduces the contrast among sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios (Fig. 9c), whereas the bottom scavenging of ^{230}Th increases this contrast (Fig. 9d). Because the influences of bottom scavenging of ^{231}Pa tends to be stronger than that of ^{230}Th , bottom scavenging overall results in reducing the contrast of $^{231}\text{Pa}/^{230}\text{Th}$ ratios (Figs. 8e and 8f). Precisely speaking, the actual processes of the bottom scavenging effect on the sedimentary ^{231}Pa and ^{230}Th appear somewhat complicated compared with those of the ocean transport effect. The effect of the bottom scavenging is two-fold. First, extra particles in the bottom ocean lead to an increase of sedimentary ^{231}Pa and ^{230}Th (e.g., positive values near the bottom in low latitudes in Fig. 8c). Second, the bottom scavenging removes ^{231}Pa and ^{230}Th from the ocean, which reduces the concentration of dissolved ^{231}Pa and ^{230}Th in the ocean interior (Fig. 8a and 8b). The changes in dissolved-phase concentration then lead to changes in particulate-phase concentration in a way such that the Eq. (5b) is satisfied. The former leads to higher sedimentary ^{231}Pa and ^{230}Th , whereas the latter leads to lower sedimentary ^{231}Pa and ^{230}Th . Our results indicate that the former process becomes more important than the latter in the low latitudes, and the sedimentary ^{231}Pa increases there. In contrast, the latter dominates in the high latitudes, and the sedimentary ^{231}Pa decreases there by the bottom scavenging effect. The effect of bottom scavenging on ^{230}Th is also basically similar to ^{231}Pa .

4.4 Residence time of ^{231}Pa and ^{230}Th

Additional insights on the simulated distribution of $^{231}\text{Pa}/^{230}\text{Th}$ ratios can be obtained from a comparison of CTRL_EXP with Siddall_EXP which reproduces sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios (Fig. S4a) as realistically as does CTRL_EXP (Fig. 6e). In this subsection, we discuss this point by focusing on the difference in the residence time of ^{231}Pa and ^{230}Th between CTRL_EXP and Siddall_EXP. Assuming the mass balance of ^{231}Pa and ^{230}Th are in a steady state, we calculate the residence time of ^{231}Pa and ^{230}Th from the following formulas:

$$\tau^i = \int A_{\text{total}}^i dv / F_{\text{in}}^i, \quad (11a)$$

$$F_{\text{in}}^i = \int \beta^i dv, \quad (11b)$$

In Eq. (11a) and (11b), the integral domain is global and the parameters are described in Table 1. The residence times of ^{231}Pa and ^{230}Th are calculated to be 103 and 21 years, respectively, in CTRL_EXP, whereas they are 211 and 89 years, respectively, in Siddall_EXP (Table S2). By incorporating bottom scavenging and modifying the partition coefficient of ^{230}Th , the modeled residence time in CTRL_EXP comes close to the previous estimate based on data in Yu et al. (1996): 111 years for ^{231}Pa and 26 years for ^{230}Th . Because the reference partition coefficients for ^{231}Pa of Siddall_EXP and that of CTRL_EXP are the same

value (i.e., $K_{\text{ref}}^{\text{Pa}} = 1.0 \times 10^7$), the influence of ocean transport on ^{231}Pa is identical in both experiments (Fig. 9a). Therefore, the difference in the ^{231}Pa distribution between the model experiments must come from the bottom scavenging, which is included in CTRL_EXP but not in Siddall_EXP. The bottom scavenging reduces the residence time of ^{231}Pa in CTRL_EXP (103 years) compared to Siddall_EXP (211 years). The difference in the ^{230}Th distribution between CTRL_EXP and Siddall_EXP mainly comes from the difference in reference partition coefficients ($K_{\text{ref}}^{\text{Th}}$). The reference partition coefficient $K_{\text{ref}}^{\text{Th}}$ of CTRL_EXP, which depends on particle concentration, is larger than that of Siddall_EXP ($K_{\text{ref}}^{\text{Th}} = 6.0 \times 10^7$) in most of the ocean. Therefore, the contribution from the ocean transport of ^{230}Th becomes larger in Siddall_EXP (Fig. S4b) than in CTRL_EXP (Fig. 6b). Together with additional contribution from the bottom scavenging effect on ^{230}Th (Fig. 9d), the residence time of ^{230}Th in CTRL_EXP (21 years) is shorter than that in Siddall_EXP (89 years). Since the residence time is overestimated for both ^{231}Pa and ^{230}Th in Siddall_EXP compared to CTRL_EXP, the distribution of sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios in Siddall_EXP ends up similar to that in CTRL_EXP. The residence time in CTRL_EXP is similar to the residence time of their control simulation in Gu and Liu (2017), which does not include the bottom scavenging process. Therefore, total scavenging efficiency in the ocean is more important than the introduction of bottom scavenging to reproduce residence time. Our CTRL_EXP reproduces the distribution of sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios more realistically because our model can more realistically reproduce the dissolved ^{231}Pa and ^{230}Th than in previous studies. Besides, the residence time of ^{231}Pa and ^{230}Th is also consistent with the observational estimate in Yu et al. (1996).

This study newly introduces a $^{231}\text{Pa}/^{230}\text{Th}$ model to the existing global three-dimensional OGCM. Based on the reversible scavenging model, this study well reproduces the distribution of dissolved concentration of ^{231}Pa and ^{230}Th by considering the bottom scavenging and the dependence of the scavenging efficiency on particle concentration. The importance of bottom scavenging on the dissolved concentration of ^{231}Pa and ^{230}Th is already discussed in previous studies (Rempfer et al., 2017; Lerner et al., 2020). Therefore, our result should be viewed as a confirmation of these previous results in this meaning. However, it is emphasized that this study provides a new estimate of this contribution to the distribution of sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios compared to other processes such as advection and water-column scavenging. Rempfer et al. (2017) evaluated the performance of their ^{231}Pa and ^{230}Th simulations based on the root mean squared deviation normalized by the standard deviation of observations. In our control experiment (CTRL_EXP), the RMSD between the available GEOTRACES data is 0.57 for dissolved ^{231}Pa and 0.51 for dissolved ^{230}Th . These values lie in the range of values for the “standard” and “optimal” experiments of Rempfer et al. (2017), the latter of which considers both bottom scavenging and boundary scavenging (see Fig. 5 in Rempfer et al., 2017). Lerner et al. (2020) use a regional eddy-permitting ocean circulation model and focus on the western North Atlantic. They also point out that removal in the nepheloid layer significantly impacts the basin-scale distribution of dissolved and particulate phases of ^{231}Pa and ^{230}Th . In line with these previous studies, our result confirmed the importance of the boundary scavenging. Recently, Gardner et al. (2018) reported data on the distribution of particles in benthic nepheloid layers. If such datasets become available for specifying the global

470 distribution of particles in nepheloid layers, the effect of bottom scavenging can be introduced more realistically. It is also expected that additional consideration about boundary scavenging helps to improve our model simulation.

In addition to the bottom scavenging, our study highlights the importance of the dependence of scavenging efficiency on particle concentration. Although the decrease of the partition coefficient with increased bulk particle concentration has been reported from observations, the dependence of scavenging efficiency on particle concentration considered in PCE_EXP is not entirely understood (Honeyman et al. 1988; Henderson et al. 1999; Hayes et al. 2015). Recently, the particle concentration effect on ^{231}Pa and ^{230}Th partition coefficients in the open ocean along the GEOTRACES GA03 transect has been reported (Hayes et al., 2015; Lerner et al., 2017). Their study suggests that the dependency in the open ocean may deviate from Eq. (10). In discussing the factors responsible for the particle concentration effect, Pavia et al. (2018) point out the possibility that the particle concentration effect is an artifact caused by filtration. Further research is needed to elucidate the mechanisms that control the particle concentration effect.

As another remaining problem, as pointed out in previous studies (Rempfer et al., 2017; Lerner et al., 2020), it is not easy to reproduce the distribution of particle phase of these two radioisotopes than the dissolved phase. Part of the error comes from the oceanic flow fields simulated in the ocean model. It is also related to the particle fluxes that we give as an empirical distribution based on satellite observations. A $^{231}\text{Pa}/^{230}\text{Th}$ modeling study using an ecosystem model that considers six different particles well reproduces the distribution of ^{231}Pa and ^{230}Th with a simple reversible scavenging model (van Hulst et al., 2018). For dissolved ^{230}Th , the correlation coefficient between their model and observations is 0.80 for the GEOTRACES GA02 transect and 0.78 for the GA03 transect, comparable to our CTRL_EXP of 0.84 and 0.70, respectively. Furthermore, By examining the response of ^{231}Pa and ^{230}Th to freshwater forcing into the North Atlantic, Missiaen et al. (2020b) show that changes in biogenic particle fluxes may have caused 30% of the changes in the sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios during the Heinrich stadial 1. Also, in Gu & Liu, (2017), the particle change due to freshwater and its impact on sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios is examined. Therefore, the role of particle fields on the distribution of ^{231}Pa and ^{230}Th , which was not directly investigated in this study, needs to be further discussed in a future study.

4.5 Remaining issues

Although our model was able to generally reproduce the basin-scale distributions of ^{231}Pa and ^{230}Th , there are still some mismatches between the model results and observations. For dissolved ^{231}Pa , introducing bottom scavenging helped to reproduce the concentrations seen in the data at depths below 3000m (Fig. 2). However, the model tends to simulate lower concentration than the observations below 3000m in Figs. 2c and 2d, which needs to be improved. The improvement was not possible simply by reducing the bottom scavenging (i.e., specifying the smaller $K_{\text{bottom}}^{\text{Pa}}$ than Figs. 2c and 2d), therefore more fundamental improvement appears to be required. One possibility is that our treatment of the nepheloid layer (i.e., the thickness of the ocean deepest layer) may be too simple and needs to be modified so that the thickness of the nepheloid layer is more realistically specified. The introduction of more realistic bottom scavenging and the consideration of the effects of particles from the continental shelf and hydrothermal vents may also help to improve the model-data agreement for both ^{231}Pa and ^{230}Th .

In this study, particle fields were not calculated in the model but specified as boundary conditions in our approach where the specified distribution of biological particles is taken from satellite-based estimation. Since the bias of the particle field affects the distribution of ^{231}Pa and ^{230}Th , our approach has advantages over the other studies where the particles are explicitly simulated in the model. However, satellite-based estimation referenced here may also contain some errors and understanding about the influence of the particle field on sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios, which was not seriously discussed in this study, is also important as the previous studies pointed out (e.g. Missiaen et al. 2020b; Dutay et al. 2009; van Hulten et al., 2018).

To reconstruct past sedimentation flux, ^{230}Th normalization was used. Recently, the influence of lithogenic and authigenic ^{230}Th on ^{230}Th in sediments was evaluated (Missiaen et al., 2018; Costa et al., 2020). This is not a direct topic of our study, but we need to care about such processes which also affect the ratio of ^{231}Pa and ^{230}Th obtained from marine sediments. As more observational data and their modelling become available, we expect to make further progress in quantitative understanding of the processes governing ^{231}Pa and ^{230}Th in the ocean.

515 5 Summary and concluding remarks

In this study, we performed OGCM experiments that incorporated the bottom scavenging and the dependence of scavenging efficiency on particle concentration together with the water-column reversible scavenging. We quantitatively evaluated the processes that determine the global distribution of sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios, which is used as a proxy for the strength of paleo-ocean circulation.

520 First, we performed an OGCM experiment using the same model settings and parameters as Siddall et al. (2005), which only introduced reversible scavenging (Siddall_EXP). In Siddall_EXP, the simulated concentrations of ^{231}Pa and ^{230}Th increase with depth, consistent with data; however, this experiment significantly overestimated the concentrations observed in the deep ocean. By incorporating bottom scavenging in nepheloid layers following Rempfer et al. (2017) (BTM_EXP), we reduce this overestimation and successfully reproduced the vertical profile of dissolved ^{231}Pa . However, this experiment had difficulty in reproducing the observed vertical profile of dissolved ^{230}Th . Therefore, we modified the parameters associated with the strength of water-column scavenging (i.e., K_{ref} : the reference partition coefficient for sinking particles) with the consideration of the bottom scavenging (KREF_EXP). When we increased the reference partition coefficient of ^{230}Th ($K_{\text{ref}}^{\text{Th}} = 6.0 \times 10^7$) from that used in the Siddall_EXP with the consideration of bottom scavenging ($K_{\text{bottom}}^{\text{Th}} = 1.0 \times 10^7$), dissolved ^{230}Th was found to be more realistically simulated, but significant underestimation in the Southern Ocean remained. We found 530 that the underestimation in the Southern Ocean can be improved by introducing dependence of K_{ref} on particle concentration which was used in Henderson et al. (1999) (PCE_EXP). Although most of previous ^{231}Pa and ^{230}Th model results showed significant overestimation in the deep ocean (e.g. Siddall et al. 2005; Dutay et al. 2009; Gu and Liu 2017; van Haurten et al. 2018), our best OGCM simulation considering the reversible scavenging, bottom scavenging, and the dependence of

scavenging efficiency on particle concentration (CTRL_EXP) can reproduce the distributions of dissolved ^{231}Pa and ^{230}Th consistently with GEOTRACES data, together with the realistic distribution of sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios.

We also made quantitative assessment about the processes that determine the global distribution of sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios by decomposing the processes affecting the sediment $^{231}\text{Pa}/^{230}\text{Th}$ ratios into three parts: water-column scavenging, ocean transport (advection, diffusion, and convection), and bottom scavenging. We found that the global sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios in our best model (CTRL_EXP) are primarily determined by ocean transport of ^{231}Pa , as in previous models. Contrary to ^{231}Pa , ocean transport of ^{230}Th tends to reduce the spatial contrast of sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios. However, we found that this is not the case for the Southern Ocean; ^{230}Th advection increases the sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios in the Southern Ocean and strengthens the observed high $^{231}\text{Pa}/^{230}\text{Th}$ ratios there. This means that not only ^{231}Pa advection but also ^{230}Th advection contributes to the high $^{231}\text{Pa}/^{230}\text{Th}$ ratios in the Southern Ocean. This result implies that scavenging of ^{230}Th is not much efficient in the Southern Ocean as conventionally thought when we consider the dependence of scavenging efficiency on particle concentration. We also show that bottom scavenging works opposite to ocean transport and decreases the spatial contrast of $^{231}\text{Pa}/^{230}\text{Th}$ ratios; bottom scavenging promotes the removal of ^{231}Pa near the sea bottom more efficiently than that of ^{230}Th , and the total effect of bottom scavenging reduces spatial contrasts of the $^{231}\text{Pa}/^{230}\text{Th}$ ratios. Our best simulation shows the realistic residence times of ^{231}Pa and ^{230}Th , but simulation without bottom scavenging and dependence of scavenging efficiency on particle concentration significantly overestimates the residence times for both ^{231}Pa and ^{230}Th in spite of similar distribution of sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios to our best simulation.

The model developed in this study is useful not only for simulating $^{231}\text{Pa}/^{230}\text{Th}$ ratios in the present-day ocean but also in different climates such as glacial periods. Our OGCM experiments using the present-day physical fields can clarify the processes governing the global distribution of sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios. A similar analysis using the physical ocean fields during glacial periods may help climate scientists to understand the mechanisms for glacial changes in the sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratio observed in sediment cores. Although simulated sedimentary Pa/Th under glacial times are also discussed in a 2-D model (Lippold et al., 2012) and recently in a 3-D model (Gu et al., 2020), there is insufficient discussion of the mechanism of change in the three-dimensional distribution. Simulation of $^{231}\text{Pa}/^{230}\text{Th}$ ratios under glacial climates (e.g., Oka et al., 2011; Kobayashi and Oka, 2018) is an exciting avenue of future study.

Code and data availability

The $^{231}\text{Pa}/^{230}\text{Th}$ model code and data used to produce the results in this study are available at the repository website Zenodo: <https://doi.org/10.5281/zenodo.4600287> (Sasaki et al., 2021a) and <https://doi.org/10.5281/zenodo.4655882> (Sasaki et al., 2021b), respectively. COCO is an ocean component of MIROC and the code of COCO4 is included as a part of MIROC-ES2L. The source code of MIROC-ES2L can be obtained from <https://doi.org/10.5281/zenodo.3893386> (Ohgaito et al., 2020).

565 Supplement

The supplement related to this article is available online at: <https://doi.org/xxxx>.

Author contributions

All the authors contributed to the interpretation of the simulation results. Y.S. performed the numerical simulations. A.O. designed and supervised the study. Y.S. and H.K. analyzed the results. Y.S. wrote the first draft and the final draft was prepared with the inputs from all the coauthors.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgments

The authors acknowledge many constructive comments from reviewers, which significantly improves the manuscript. This work was supported by JSPS KAKENHI Grant Number JP19H01963.

References

- Anderson, R. F., Bacon, M. P., and Brewer, P. G.: Removal of ^{230}Th and ^{231}Pa from the open ocean, *Earth Planet. Sci. Lett.*, 62(1), 7–23, [https://doi.org/10.1016/0012-821X\(83\)90067-5](https://doi.org/10.1016/0012-821X(83)90067-5), 1983.
- Bacon, M. P., and Anderson, R. F.: Distribution of thorium isotopes between dissolved and particulate forms in the deep sea, *J. Geophys. Res.*, 87(C3), 2045, <https://doi.org/10.1029/JC087iC03p02045>, 1982.
- Behrenfeld, M. J., and Falkowski, P. G.: Photosynthetic rates derived from satellite-based chlorophyll concentration, *Limnol. Oceanogr.*, 42(1), 1–20, <https://doi.org/10.4319/lo.1997.42.1.0001>, 1997.
- Böhm, E., Lippold, J., Gutjahr, M., Frank, M., Blaser, P., Antz, B., Fohlmeister, J., Frank, N., Anderson, M. B., and Deininger, M.: Strong and deep Atlantic meridional overturning circulation during the last glacial cycle, *Nature*, 517(7532), 73–76, <https://doi.org/10.1038/nature14059>, 2015.
- Bradtmiller, L. I., McManus, J. F., and Robinson, L. F.: $^{231}\text{Pa}/^{230}\text{Th}$ evidence for a weakened but persistent Atlantic meridional overturning circulation during Heinrich Stadial 1, *Nat. Commun.*, 5(1), 5817, <https://doi.org/10.1038/ncomms6817>, 2014.
- Chase, Z., and Anderson, R. F.: Comment on “On the importance of opal, carbonate, and lithogenic clays in scavenging and fractionating ^{230}Th , ^{231}Pa and ^{10}Be in the ocean” by S. Luo and T.-L. Ku., *Earth Planet. Sci. Lett.*, 220(1–2), 213–222, [https://doi.org/10.1016/S0012-821X\(04\)00028-7](https://doi.org/10.1016/S0012-821X(04)00028-7), 2004.

- Chase, Z., Anderson, R. F., Fleisher, M. Q., and Kubik, P. W.: The influence of particle composition and particle flux on scavenging of Th, Pa and Be in the ocean, *Earth Planet. Sci. Lett.*, 204(1–2), 215–229, [https://doi.org/10.1016/S0012-821X\(02\)00984-6](https://doi.org/10.1016/S0012-821X(02)00984-6), 2002.
- 595 Chase, Z., Anderson, R. F., Fleisher, M. Q., and Kubik, P. W.: Scavenging of ²³⁰Th, ²³¹Pa and ¹⁰Be in the Southern Ocean (SW Pacific sector): the importance of particle flux, particle composition and advection, *Deep-Sea Res. II*, 50(3–4), 739–768, [https://doi.org/10.1016/S0967-0645\(02\)00593-3](https://doi.org/10.1016/S0967-0645(02)00593-3), 2003.
- Cochran, J. K., Livingston, H. D., Hirschberg, D. J., and Surprenant, L. D.: Natural and anthropogenic radionuclide distributions in the northwest Atlantic Ocean, *Earth Planet. Sci. Lett.*, 84(2–3), 135–152, [https://doi.org/10.1016/0012-821X\(87\)90081-1](https://doi.org/10.1016/0012-821X(87)90081-1), 1987.
- 600 Colley, S., Thomson, J., and Newton, P. P.: Detailed ²³⁰Th, ²³²Th and ²¹⁰Pb fluxes recorded by the 1989/90 BOFS sediment trap time-series at 48°N, 20°W, *Deep-Sea Res. I*, 42(6), 833–848, [https://doi.org/10.1016/0967-0637\(95\)00033-3](https://doi.org/10.1016/0967-0637(95)00033-3), 1995.
- Deng F., Henderson, G. M., Castrillejo, M., Perez, F. F., and Steinfeldt, R.: Evolution of ²³¹Pa and ²³⁰Th in overflow waters of the North Atlantic, *Biogeosciences*, 15, 7299–7313, <https://doi.org/10.5194/bg-15-7299-2018>, 2018.
- 605 Dunne, J. P., Armstrong, R. A., Gnanadesikan, A., and Sarmiento, J. L.: Empirical and mechanistic models for the particle export ratio, *Glob. Biogeochem. Cycles*, 19(4), <https://doi.org/10.1029/2004GB002390>, 2005.
- Dutay, J.-C., Lacan, F., Roy-Barman, M., and Bopp, L.: Influence of particle size and type on ²³¹Pa and ²³⁰Th simulation with a global coupled biogeochemical-ocean general circulation model: A first approach, *Geochem. Geophys. Geosyst.*, 10, Q01011, <https://doi.org/10.1029/2008GC002291>, 2009.
- 610 Edwards, N. R., Stocker, T. F., Joos, F., Henderson, G. M., Frank, M., Siddall, M., and Müller, S. A.: ²³¹Pa/²³⁰Th fractionation by ocean transport, biogenic particle flux and particle type, *Earth Planet. Sci. Lett.*, 237(1–2), 135–155, <https://doi.org/10.1016/j.epsl.2005.05.031>, 2005.
- Franois, R., Bacon, M. P., Altabet, M. A., and Labeyrie, L. D.: Glacial/interglacial changes in sediment rain rate in the SW Indian Sector of subantarctic Waters as recorded by ²³⁰Th, ²³¹Pa, U, and $\delta^{15}\text{N}$, *Paleoceanography*, 8(5), 611–629, <https://doi.org/10.1029/93PA00784>, 1993.
- 615 Frank, M., Eckhardt, J.-D., Eisenhauer, A., Kubik, P. W., Dittrich-Hannen, B., Segl, M., and Mangini, A.: Beryllium 10, thorium 230, and protactinium 231 in Galapagos microplate sediments: Implications of hydrothermal activity and paleoproductivity changes during the last 100,000 years, *Paleoceanography*, 9(4), 559–578, <https://doi.org/10.1029/94PA01132>, 1994.
- 620 Gardner, W. D., Mishonov, A. V., and Richardson, M. J.: Decadal Comparisons of Particulate Matter in Repeat Transects in the Atlantic, Pacific, and Indian Ocean Basins, *Geophys. Res. Lett.*, 45(1), 277–286, <https://doi.org/10.1002/2017GL076571>, 2018.
- Gherardi, J.-M., Labeyrie, L., Nave, S., Franois, R., McManus, J. F., and Cortijo, E.: Glacial-interglacial circulation changes inferred from ²³¹Pa/²³⁰Th sedimentary record in the North Atlantic region, *Paleoceanography*, 24(2), PA2204, <https://doi.org/10.1029/2008PA001696>, 2009.
- 625

- Gregory, J. M., Dixon, K. W., Stouffer, R. J., Weaver, A. J., Driesschaert, E., Eby, M., Fichefet, T., Hasumi, H., Hu, A., Jungclaus, J. H., Kamenkovich, I. V., Levermann, A., Montoya, M., Murakami, S., Nawrath, S. Oka, A., Sokolov, A. P., Thorpe, R. B.: A model intercomparison of changes in the Atlantic thermohaline circulation in response to increasing atmospheric CO₂ concentration, *Geophys. Res. Lett.*, 32(12), 1–5, <https://doi.org/10.1029/2005GL023209>, 2005.
- 630 Gu, S., and Liu, Z.: 231Pa and 230Th in the ocean model of the Community Earth System Model (CESM1.3), *Geosci. Model Dev.*, 10, 4723–4742, <https://doi.org/10.5194/gmd-10-4723-2017>, 2017.
- Gu, S., Liu, Z., Oppo, D. W. Lynch-Stieglitz, J., Jahn, A., Zhang, J., Wu, L.: Assessing the potential capability of reconstructing glacial Atlantic water masses and AMOC using multiple proxies in CESM, *Earth Planet. Sci. Lett.*, 541, 116294, <https://doi.org/10.1016/j.epsl.2020.116294>, 2020.
- 635 Guo, L., Santschi, P. H., Baskaran, M., and Zindler, A.: Distribution of dissolved and particulate 230Th and 232Th in seawater from the Gulf of Mexico and off Cape Hatteras as measured by SIMS, *Earth Planet. Sci. Lett.*, 133(1–2), 117–128, [https://doi.org/10.1016/0012-821X\(95\)00063-I](https://doi.org/10.1016/0012-821X(95)00063-I), 1995.
- Hajima, T., Watanabe, M., Yamamoto, A., Tatebe, H., Noguchi, M. A., Abe, M., Ohgaito, R., Ito, A., Yamazaki, D., Okajima, H., Ito, A., Takata, K., Ogochi, K., Watanabe, S., and Kawamiya, M.: Development of the MIROC-ES2L Earth system model and the evaluation of biogeochemical processes and feedbacks, *Geosci. Model Dev.*, 13, 2197–2244, <https://doi.org/10.5194/gmd-13-2197-2020>, 2020.
- 640
- Hasumi, H.: CCSR Ocean Component Model (COCO) version 4.0. *CCSR Rep. 25*, 103 pp., Center for Climate System Research, Univ. of Tokyo, 2006.
- Hayes, C. T., Anderson, R. F., Jaccard, S. L., François, R., Fleisher, M. Q., Soon, M., Gersonde, R.: A new perspective on boundary scavenging in the North Pacific Ocean, *Earth Planet. Sci. Lett.*, 369–370, 86–97, <https://doi.org/10.1016/j.epsl.2013.03.008>, 2013.
- 645
- Hayes, C. T., Anderson, R. F., Fleisher, M. Q., Huang, K.-F., Robinson, L. F., Lu, Y., Cheng, H., Lawrence Edwards, R., and Bradley Moran, S.: 230Th and 231Pa on GEOTRACES GA03, the U.S. GEOTRACES North Atlantic transect, and implications for modern and paleoceanographic chemical fluxes, *Deep-Sea Res. II.*, 116, 29–41, <https://doi.org/10.1016/j.dsr2.2014.07.007>, 2015.
- 650
- Henderson, G. M., and Anderson, R. F.: The U-series Toolbox for Paleoceanography. *Reviews in Mineralogy and Geochemistry*, 52(1), 493–531, <https://doi.org/10.2113/0520493>, 2003.
- Henderson, Gideon M., Heinze, C., Anderson, R. F., and Winguth, A. M. E.: Global distribution of the 230Th flux to ocean sediments constrained by GCM modeling, *Deep-Sea Res. I.*, 46(11), 1861–1893, [https://doi.org/10.1016/S0967-0637\(99\)00030-8](https://doi.org/10.1016/S0967-0637(99)00030-8), 1999.
- 655
- Honeyman, B. D., Balistreri, L. S., and Murray, J. W.: Oceanic trace metal scavenging: the importance of particle concentration, *Deep-Sea Res. A*, 35(2), 227–246, [https://doi.org/10.1016/0198-0149\(88\)90038-6](https://doi.org/10.1016/0198-0149(88)90038-6), 1988.

- Honjo, S., Manganini, S. J., Krishfield, R. A., and François, R.: Particulate organic carbon fluxes to the ocean interior and factors controlling the biological pump: A synthesis of global sediment trap programs since 1983, *Prog. Oceanogr.*, 660 76(3), 217–285, <https://doi.org/10.1016/j.pocean.2007.11.003>, 2008.
- K-1 Model Developers: K-1 coupled GCM (MIROC) description. In H. Hasumi, & S. Emori (Eds.), *K-1 Tech. Rep. I*, 34 pp., Center for Climate System Research, Univ. of Tokyo, 2004.
- Kobayashi, H., Abe-Ouchi, A., and Oka, A.: Role of Southern Ocean stratification in glacial atmospheric CO₂ reduction evaluated by a three-dimensional ocean general circulation model, *Paleoceanography*, 30(9), 1202–1216, 665 <https://doi.org/10.1002/2015PA002786>, 2015.
- Kobayashi, H., and Oka, A.: Response of atmospheric pCO₂ to glacial changes in the Southern Ocean amplified by carbonate compensation, *Paleoceanogr. Paleoclimatol.*, 33(11), 1206–1229, <https://doi.org/10.1029/2018PA003360>, 2018.
- Lam, P. J., Ohnemus, D. C., and Auro, M. E.: Size-fractionated major particle composition and concentrations from the US 670 GEOTRACES North Atlantic Zonal Transect, *Deep-Sea Res. II: Topical Studies in Oceanography*, 116, 303–320, <https://doi.org/10.1016/J.DSR2.2014.11.020>, 2015.
- Lerner P., Marchal O., Lam P., Buesseler K., Charette M.: Kinetics of thorium and particle cycling along the U.S. GEOTRACES North Atlantic transect, *Deep-Sea Research I*, 125, 106–128, <https://doi.org/10.1016/j.dsr.2017.05.003>, 2017.
- 675 Lerner, P., Marchal, O., Lam, P. J., Gardner, W., Richardson, M. J., and Mishonov, A.: A model study of the relative influences of scavenging and circulation on ²³⁰Th and ²³¹Pa in the western North Atlantic, *Deep-Sea Res. I*, 155, 103159, <https://doi.org/10.1016/j.dsr.2019.103159>, 2020.
- Lippold, J., Luo, Y., Francois, R., Allen, S. E., Gherardi, J., Pichat, S., Hickey, B., and Schulz, H.: Strength and geometry of the glacial Atlantic Meridional Overturning Circulation, *Nat. Geosci.*, 5, 813–816, <https://doi.org/10.1038/NGEO1608>, 2012.
- 680 Luo, S., and Ku, T. L.: Oceanic ²³¹Pa/²³⁰Th ratio influenced by particle composition and remineralization, *Earth Planet. Sci. Lett.*, 167(3–4), 183–195, [https://doi.org/10.1016/S0012-821X\(99\)00035-7](https://doi.org/10.1016/S0012-821X(99)00035-7), 1999.
- Luo, Y., François, R., and Allen, S. E.: Sediment ²³¹Pa/²³⁰Th as a recorder of the rate of the Atlantic meridional overturning circulation: Insights from a 2-D model, *Ocean Sci.*, 6(1), 381–400, <https://doi.org/10.5194/os-6-381-2010>, 2010.
- Maier-Reimer, E.: Geochemical cycles in an ocean general circulation model. Preindustrial tracer distributions, *Glob. 685 Biogeochem. Cycles*, 7(3), 645–677, <https://doi.org/10.1029/93GB01355>, 1993.
- Mangini, A., and Sonntag, C.: ²³¹Pa dating of deep-sea cores via ²²⁷Th counting, *Earth Planet. Sci. Lett.*, 37(2), 251–256, [https://doi.org/10.1016/0012-821X\(77\)90170-4](https://doi.org/10.1016/0012-821X(77)90170-4), 1977.
- Marchal, O., François, R., Stocker, T. F., and Joos, F.: Ocean thermohaline circulation and sedimentary ²³¹Pa/²³⁰Th ratio, *Paleoceanography*, 15(6), 625–641, <https://doi.org/10.1029/2000PA000496>, 2000.
- 690 Marchal, O., Stocker, T. F., and Joos, F.: A latitude-depth, circulation-biogeochemical ocean model for paleoclimate studies. Development and sensitivities, *Tellus B*, 50(3), 290–316, <https://doi.org/10.3402/tellusb.v50i3.16130>, 1998.

- McManus, J. F., François, R., Gherardl, J. M., Kelgwin, L., and Drown-Leger, S.: Collapse and rapid resumption of Atlantic meridional circulation linked to deglacial climate changes, *Nature*, 428(6985), 834–837, <https://doi.org/10.1038/nature02494>, 2004.
- 695 Missiaen, L., Bouttes, N., Roche, D. M., Dutay, J.-C., Quiquet, A., Waelbroeck, C., Pichat, S., Peterschmitt, J.-Y.: Carbon isotopes and Pa/Th response to forced circulation changes: a model perspective, *Clim. Past*, 16(3), 867–883, <https://doi.org/10.5194/cp-16-867-2020>, 2020a
- Missiaen, L., Menviel, L. C., Meissner, K. J., Roche, D. M., Dutay, J.-C., Bouttes, N., Lhardy, F., Quiquet, A., Pichat, S., Waelbroeck, C.: Modelling the impact of biogenic particle flux intensity and composition on sedimentary Pa/Th, *Quaternary Sci. Rev.*, 240, 106394, <https://doi.org/10.1016/j.quascirev.2020.106394>, 2020b.
- 700 Moran, S. B., Charette, M. A., Hoff, J. A., Edwards, R. L., and Landing, W. M.: Distribution of ²³⁰Th in the Labrador Sea and its relation to ventilation, *Earth Planet. Sci. Lett.*, 150(1–2), 151–160, [https://doi.org/10.1016/s0012-821x\(97\)00081-2](https://doi.org/10.1016/s0012-821x(97)00081-2), 1997.
- Moran, S. B., Shen, C. C., Edmonds, H. N., Weinstein, S. E., Smith, J. N., and Edwards, R. L.: Dissolved and particulate ²³¹Pu and ²³⁰Th in the Atlantic Ocean: Constraints on intermediate/deep water age, boundary scavenging, and ²³¹Pu/²³⁰Th fractionation, *Earth Planet. Sci. Lett.*, 203(3–4), 999–1014, [https://doi.org/10.1016/S0012-821X\(02\)00928-7](https://doi.org/10.1016/S0012-821X(02)00928-7), 2002.
- 705 Moran, S. B., Shen, C.-C., Weinstein, S. E., Hettlinger, L. H., Hoff, J. H., Edmonds, H. N., and Edwards, R. L.: Constraints on deep water age and particle flux in the equatorial and South Atlantic Ocean based on seawater ²³¹Pu and ²³⁰Th data, *Geophys. Res. Lett.*, 28(18), 3437–3440, <https://doi.org/10.1029/2001GL013339>, 2001.
- 710 Müller, P. J., and Mangini, A.: Organic carbon decomposition rates in sediments of the Pacific manganese nodule belt dated by ²³⁰Th and ²³¹Pu, *Earth Planet. Sci. Lett.*, 51(1), 94–114, [https://doi.org/10.1016/0012-821X\(80\)90259-9](https://doi.org/10.1016/0012-821X(80)90259-9), 1980.
- Oka, A., Hasumi, H., and Abe-Ouchi, A.: The thermal threshold of the Atlantic meridional overturning circulation and its control by wind stress forcing during glacial climate, *Geophys. Res. Lett.*, 39(9), <https://doi.org/10.1029/2012GL051421>, 2012.
- 715 Oka, A., Hasumi, H., Obata, H., Gamo, T., and Yamanaka, Y.: Study on vertical profiles of rare earth elements by using an ocean general circulation model, *Glob. Biogeochem. Cycles*, 23(4), GB4025, <https://doi.org/10.1029/2008GB003353>, 2009.
- Oka, A., Hasumi, H., Okada, N., Sakamoto, T. T., and Suzuki, T.: Deep convection seesaw controlled by freshwater transport through the Denmark Strait, *Ocean Model.*, 15(3–4), 157–176, <https://doi.org/10.1016/j.ocemod.2006.08.004>, 2006.
- 720 Oka, A., Kato S., and Hasumi H.: Evaluating effect of ballast mineral on deep-ocean nutrient concentration by using an ocean general circulation model, *Glob. Biogeochem. Cycles*, 22(3), <https://doi.org/10.1029/2007GB003067>, 2008.
- Oka, A., Tazoe H., and Obata, H.: Simulation of global distribution of rare earth elements in the ocean using an ocean general circulation model, *Journal of Oceanography*, <https://doi.org/10.1007/s10872-021-00600-x>, 2021.

- Okubo, A., Obata, H., Gamo, T., and Yamada, M.: ^{230}Th and ^{232}Th distributions in mid-latitudes of the North Pacific Ocean: Effect of bottom scavenging, *Earth Planet. Sci. Lett.*, 339–340, 139–150, <https://doi.org/10.1016/J.EPSL.2012.05.012>, 2012.
- Ohgaito, R., Yamamoto, A., Hajima, T., O'ishi, R., Abe, M., Tatebe, H., Abe-Ouchi, A., and Kawamiya, M.: Core code of MIROC-ES2L, Geoscientific Model Development, Zenodo, <https://doi.org/10.5281/zenodo.3893386>, 2020.
- Ohgaito, R., Yamamoto, A., Hajima, T., O'ishi, R., Abe, M., Tatebe, H., Abe-Ouchi, A., and Kawamiya, M.: PMIP4 experiments using MIROC-ES2L Earth system model, *Geosci. Model Dev.*, 14, 1195–1217, <https://doi.org/10.5194/gmd-14-1195-2021>, 2021.
- Pavia, F., Anderson, R., Vivancos, S., Fleisher, M., Lam, P., Lu, Y., Cheng, H., Zhang, P., Lawrence Edwards, R.: Intense hydrothermal scavenging of ^{230}Th and ^{231}Pa in the deep Southeast Pacific, *Mar. Chem.*, 201, 212–228, <https://doi.org/10.1016/J.MARCHEM.2017.08.003>, 2018.
- 735 Rempfer, J., Stocker, T. F., Joos, F., Lippold, J., and Jaccard, S. L.: New insights into cycling of ^{231}Pa and ^{230}Th in the Atlantic Ocean, *Earth Planet. Sci. Lett.*, 468, 27–37, <https://doi.org/10.1016/j.epsl.2017.03.027>, 2017.
- Roy-Barman, M.: Modeling the effect of boundary scavenging on Thorium and Protactinium profiles in the ocean. *Biogeosciences*, 6(12), 3091–3107, <https://doi.org/10.5194/bg-6-3091-2009>, 2009.
- Rutgers van der Loeff, M. M., and Berger, G. W.: Scavenging of ^{230}Th and ^{231}Pa near the antarctic polar front in the South Atlantic, *Deep-Sea Res. I*, 40(2), 339–357, [https://doi.org/10.1016/0967-0637\(93\)90007-P](https://doi.org/10.1016/0967-0637(93)90007-P), 1993.
- 740 Sasaki, Y., Kobayashi, H., A. Oka : Data for the figures of Sasaki et al. entitled "An investigation into the processes controlling the global distribution of dissolved ^{231}Pa and ^{230}Th in the ocean and the sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios by using an ocean general circulation model COCO ver4.0.", <https://doi.org/10.5281/zenodo.4600287>, 2021a.
- Sasaki, Y., Kobayashi, H., A. Oka : Code of Sasaki et al. entitled "An investigation into the processes controlling the global distribution of dissolved ^{231}Pa and ^{230}Th in the ocean and the sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios by using an ocean general circulation model COCO ver4.0.", <https://doi.org/10.5281/zenodo.4600287>, 2021b.
- 745 Schlitzer, R., Anderson, R. F., Dodas, E. M., Lohan, M., Geibert, W., Tagliabue, A., Bowie, A., Jeandel, C., Maldonado, M. T., Landing, W. M., Cockwell, D. et al.: The GEOTRACES Intermediate Data Product 2017, *Chem. Geol.*, 493, 210–223, <https://doi.org/10.1016/j.chemgeo.2018.05.040>, 2018.
- 750 Schmitz, W., Mangini, A., Stoffers, P., Glasby, G. P., and Plüger, W. L.: Sediment accumulation rates in the southwestern Pacific Basin and Aitutaki Passage, *Mar. Geol.*, 73(1–2), 181–190, [https://doi.org/10.1016/0025-3227\(86\)90118-0](https://doi.org/10.1016/0025-3227(86)90118-0), 1986.
- Shimmield, G. B., Murray, J. W., Thomson, J., Bacon, M. P., Anderson, R. F., and Price, N. B.: The distribution and behaviour of ^{230}Th and ^{231}Pa at an ocean margin, Baja California, Mexico, *Geochim. Cosmochim. Acta*, 50(11), 2499–2507, [https://doi.org/10.1016/0016-7037\(86\)90032-3](https://doi.org/10.1016/0016-7037(86)90032-3), 1986.
- 755 Shimmield, G. B., and Price, N. B.: The scavenging of U, ^{230}Th and ^{231}Pa during pulsed hydrothermal activity at 20°S, East Pacific Rise, *Geochim. Cosmochim. Acta*, 52(3), 669–677, [https://doi.org/10.1016/0016-7037\(88\)90329-8](https://doi.org/10.1016/0016-7037(88)90329-8), 1988.

- Siddall, M., Henderson, G. M., Edwards, N. R., Frank, M., Mu, S. A., Stocker, T. F., and Joos, F.: $^{231}\text{Pa}/^{230}\text{Th}$ fractionation by ocean transport, biogenic particle flux and particle type, *Earth Planet. Sci. Lett.*, *237*, 135–155, <https://doi.org/10.1016/j.epsl.2005.05.031>, 2005.
- 760 Stouffer, R. J., Yin, J., Gregory, J. M., Dixon, K. W., Spelman, M. J., Hurlin, W., Weaver, A. J., Eby, M., Flato, G. M., Hasumi, H., Hu, A., Jungclaus, J. H., Kamenkovich, I. V., Levermann, A., Montoya, M., Murakami, S., Nawrath, S., Oka, A., Peltier, W. R., Robitaille, D. Y., Sokolov, A., Vettoretti, G., Weber, S. L.: Investigating the Causes of the Response of the Thermohaline Circulation to Past and Future Climate Changes, *J. Climate*, *19*(8), 1365–1387, <https://doi.org/10.1175/JCLI3689.1>, 2006.
- 765 Süfke, F., Schulz, H., Scheen, J., Szidat, S., Regelous, M., Blaser, P., Pöppelmeier, F., Goepfert, T. J., Stocker, T. F., Lippold, J.: Inverse response of $^{231}\text{Pa}/^{230}\text{Th}$ to variations of the Atlantic meridional overturning circulation in the North Atlantic intermediate water, *Geo-Mar. Lett.*, *40*, 75–87, <https://doi.org/10.1007/s00367-019-00634-7>, 2020.
- Yang, H.-S., Nozaki, Y., Sakai, H., and Masuda, A.: The distribution of ^{230}Th and ^{231}Pa in the deep-sea surface sediments of the Pacific Ocean, *Geochim. Cosmochim. Acta*, *50*(1), 81–89, [https://doi.org/10.1016/0016-7037\(86\)90050-5](https://doi.org/10.1016/0016-7037(86)90050-5), 1986.
- 770 van Hulten, M., Dutay, J.-C., and Roy-Barman, M.: A global scavenging and circulation ocean model of thorium-230 and protactinium-231 with improved particle dynamics (NEMO-ProThorP 0.1), *Geosci. Model Dev.*, *11*, 3537–3556, <https://doi.org/10.5194/gmd-11-3537-2018>, 2018.
- Vogler, S., Scholten, J., Rutgers Van Der Loeff, M., and Mangini, A.: ^{230}Th in the eastern North Atlantic: The importance of water mass ventilation in the balance of ^{230}Th , *Earth Planet. Sci. Lett.*, *156*(1–2), 61–74, [https://doi.org/10.1016/s0012-821x\(98\)00011-9](https://doi.org/10.1016/s0012-821x(98)00011-9), 1998.
- 775 Walter, H. J., Rutgers Van Der Loeff, M. M., and Hoeltzen, H.: Enhanced scavenging of ^{231}Pa relative to ^{230}Th in the South Atlantic south of the Polar Front: Implications for the use of the $^{231}\text{Pa}/^{230}\text{Th}$ ratio as a paleoproductivity proxy, *Earth Planet. Sci. Lett.*, *149*(1–4), 85–100, [https://doi.org/10.1016/s0012-821x\(97\)00068-x](https://doi.org/10.1016/s0012-821x(97)00068-x), 1997.
- 780 Waelbroeck, C., Pichat, S., Böhm, E., Lougheed, B. C., Faranda, D., Vrac, M., Missiaen, L., Riveiros, N. V., Burckel, P., Lippold, J., Arz, H. W., Dokken, T., Thil, F., and Dapoigny A.: Relative timing of precipitation and ocean circulation changes in the western equatorial Atlantic over the last 45 kyr, *Clim. Past*, *14*, 1315–1330, <https://doi.org/10.5194/cp-14-1315-2018>, 2018.
- Yong Lao, Anderson, R. F., Broecker, W. S., Trumbore, S. E., Hofmann, H. J., and Wolfl, W.: Transport and burial rates of ^{10}Be and ^{231}Pa in the Pacific Ocean during the Holocene period, *Earth Planet. Sci. Lett.*, *113*(1–2), [https://doi.org/10.1016/0012-821X\(92\)90218-K](https://doi.org/10.1016/0012-821X(92)90218-K), 1992.
- 785 Yu, E.-F., François, R., and Bacon, M. P.: Similar rates of modern and last-glacial ocean thermohaline circulation inferred from radiochemical data, *Nature*, *379*(6567), 689–694, <https://doi.org/10.1038/379689a0>, 1996.

790 **Figure 1.** (a) Dissolved ^{231}Pa along 30°W in the Atlantic Ocean and (b) its vertical profile averaged horizontally along 30°W in Siddall_EXP. (c, d) Same as Figs. 1a and 1b except for ^{230}Th . The colored circles in Figs. 1a and 1c represent data from the Atlantic GEOTRACES data (GA02 and GIPY05; Schlitzer et al., 2018). The green and orange circles in Figs. 1b and 1d represent the GA02 data and simulation results.

795 **Figure 2.** (a, c, e) Dissolved ^{231}Pa along 30°W in the Atlantic Ocean and (b, d, f) its vertical profile averaged horizontally along 30°W in BTM_EXP. $K_{\text{bottom}}^{\text{Pa}}$ is set to 5.0×10^4 in Figs. 2a and 2b, 5.0×10^5 in Figs. 2c and 2d, and 5.0×10^6 in Figs. 2e and 2f. The colored circles in Figs. 2a, 2c, and 2e represent data from the Atlantic GEOTRACES data (GA02 and GIPY05; Schlitzer et al., 2018). The green and orange circles in Figs. 2b, 2d, and 2f represent the GA02 data and simulation results.

800 **Figure 3.** (a, c, e, g) Dissolved ^{230}Th along 30°W in the Atlantic Ocean and (b, d, f, h) its vertical profile averaged horizontally along 30°W in BTM_EXP are plotted. $K_{\text{bottom}}^{\text{Th}}$ is set to 5.0×10^5 in Figs. 3a and 3b, 1.0×10^6 in Figs. 3c and 3d, 5.0×10^6 in Figs. 3e and 3f, and 1.0×10^7 in Figs. 3g and 3h. The colored circles in Figs. 3a, 3c, 3e, and 3g represent data from the Atlantic GEOTRACES data (GA02 and GIPY05; Schlitzer et al., 2018). The green and orange circles in Figs. 3b, 3d, 3f, and 3h represent the GA02 data and simulation results.

805 **Figure 4.** The vertical profile of dissolved ^{230}Th averaged horizontally along 30°W in various simulations of KREF_EXP: (a) R2_B20, (b) R2_B10, (c) R2_B5, (d) R4_B10, (e) R4_B5, and (f) R6_B5. The green and orange circles in Figs. 4a–4f represent the Atlantic GEOTRACES data (GA02; Schlitzer et al., 2018) and simulation results. Figure 4g summarizes the choice of parameters (i.e., $K_{\text{ref}}^{\text{Th}}$ and $K_{\text{bottom}}^{\text{Th}}$) in each simulation.

810 **Figure 5.** Dissolved ^{230}Th along 30°W in the Atlantic Ocean in (a) R6_B5 of the KREF_EXP and (b) PCE_EXP. (c) Reference coefficient (K_{ref}) along 30°W in the Atlantic Ocean in PCE_EXP. (d) The vertical profile of dissolved ^{230}Th averaged horizontally along 30°W in R6_B5 of KREF_EXP and PCE_EXP. The colored circles in Figs. 5a and 5b represent data from the Atlantic GEOTRACES data (GA02 and GIPY05; Schlitzer et al., 2018). The green, yellow, and orange circles in Fig. 5d represent the GA02 data and KREF_EXP and PCE_EXP simulation results.

815 **Figure 6.** (a) Dissolved ^{231}Pa , (b) dissolved ^{230}Th , (c) particulate ^{231}Pa , and (d) particulate ^{230}Th along 30°W in the Atlantic Ocean in CTRL_EXP. (e) Sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios normalized by the production ratio of 0.093 in CTRL_EXP. The colored circles represent observational data. Dissolved ^{231}Pa and ^{230}Th data are taken from the Atlantic GEOTRACES data (GA02 and GIPY05; Schlitzer et al., 2018). Particulate ^{231}Pa and ^{230}Th data are taken from the following references (Colley et al., 1995; Moran et al., 1997; Moran et al., 2001; Rutgers van der Loeff and Berger, 1993; Vogler et al., 1998; Walter et al., 1997; Cochran et al., 1987; Moran et al., 2002; Guo et al., 1995). The data of sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios are taken from the following references (Mangianini & Sonntag, 1977; Muller & Mangini, 1980; Anderson et al., 1983; Shimmiel et al., 1986; Schmitz et al., 1986; Yang et al., 1986; Shimmiel & Price, 1988; Yong Lao et al., 1992; François et al., 1993; Frank et al., 1994; Frank, 1996; Bradtmiller et al., 2014, Luo et al., 2010, and their supplemental data).

825 **Figure 7.** The difference between 3D_EXP and 1D_EXP (i.e., 3D_EXP minus 1D_EXP, which represents for ocean transport effect) of (a) dissolved ^{231}Pa , (b) dissolved ^{230}Th , (c) particulate ^{231}Pa , and (d) particulate ^{230}Th along 30°W in the Atlantic Ocean. (e) The difference between 3D_EXP and 1D_EXP of sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios normalized by the production ratio of 0.093.

Figure 8. The difference between CTRL_EXP and 3D_EXP (i.e., CTRL_EXP minus 3D_EXP, which represents for bottom scavenging effect) of (a) dissolved ^{231}Pa , (b) dissolved ^{230}Th , (c) particulate ^{231}Pa , and (d) particulate ^{230}Th along 30°W in the Atlantic Ocean. (e) The difference between CTRL_EXP and 3D_EXP of the sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios normalized by the production ratio of 0.093.

830


Figure 9. Sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios normalized by the production ratio of 0.093 in CTRL_EXP decomposed into contributions from (a) ocean transport solely from ^{231}Pa (i.e., $^{231}\text{Pa}(3\text{D})/^{230}\text{Th}(1\text{D})$), (b) ocean transport solely from ^{230}Th (i.e., $^{231}\text{Pa}(1\text{D})/^{230}\text{Th}(3\text{D})$), (c) bottom scavenging solely from ^{231}Pa (i.e., $^{231}\text{Pa}(\text{CTRL})/^{230}\text{Th}(3\text{D})$ minus $^{231}\text{Pa}(3\text{D})/^{230}\text{Th}(3\text{D})$), and (d) bottom scavenging solely from ^{230}Th (i.e., $^{231}\text{Pa}(3\text{D})/^{230}\text{Th}(\text{CTRL})$ minus $^{231}\text{Pa}(3\text{D})/^{230}\text{Th}(3\text{D})$).

835


Table 1. Parameters of the ^{231}Pa and ^{230}Th model.

Table 2. Equilibrium partition coefficients in experiments Siddall_EXP and CTRL_EXP.

840 **Table 3.** Processes considered in additional experiments. A circle means that the process is considered, and a cross means that it is not considered.



845 Figure 1. (a) Dissolved ^{231}Pa along 30°W in the Atlantic Ocean and (b) its vertical profile averaged horizontally along 30°W in Siddall_EXP. (c, d) Same as Figs. 1a and 1b except for ^{230}Th . The colored circles in Figs. 1a and 1c represent data from the Atlantic GEOTRACES data (GA02 and GIPY05; Schlitzer et al., 2018). The green and orange circles in Figs. 1b and 1d represent the GA02 data and simulation results.



850

Figure 2. (a, c, e) Dissolved ^{231}Pa along 30°W in the Atlantic Ocean and (b, d, f) its vertical profile averaged horizontally along 30°W in BTM_EXP. $K_{\text{bottom}}^{\text{Pa}}$ is set to 5.0×10^4 in Figs. 2a and 2b, 5.0×10^5 in Figs. 2c and 2d, and 5.0×10^6 in Figs. 2e and 2f. The colored circles in Figs. 2a, 2c, and 2e represent data from the Atlantic GEOTRACES data (GA02 and GIPY05; Schlitzer et al., 2018). The green and orange circles in Figs. 2b, 2d, and 2f represent the GA02 data and simulation results.

855












Figure 3. (a, c, e, g) Dissolved ^{230}Th along 30°W in the Atlantic Ocean and (b, d, f, h) its vertical profile averaged horizontally along 30°W in BTM_EXP are plotted. $K_{\text{bottom}}^{\text{Th}}$ is set to 5.0×10^5 in Figs. 3a and 3b, 1.0×10^6 in Figs. 3c and 3d, 5.0×10^6 in Figs. 3e and 3f, and 1.0×10^7 in Figs. 3g and 3h. The colored circles in Figs. 3a, 3c, 3e, and 3g represent data from the Atlantic GEOTRACES data (GA02 and GIPY05; Schlitzer et al., 2018). The green and orange circles in Figs. 3b, 3d, 3f, and 3h represent the GA02 data and simulation results.



865

Figure 4. The vertical profile of dissolved ^{230}Th averaged horizontally along 30°W in various simulations of KREF_EXP: (a) R2_B20, (b) R2_B10, (c) R2_B5, (d) R4_B10, (e) R4_B5, and (f) R6_B5. The green and orange circles in Figs. 4a–4f represent the Atlantic GEOTRACES data (GA02; Schlitzer et al., 2018) and simulation results. Figure 4g summarizes the choice of parameters (i.e., $K_{\text{ref}}^{\text{Th}}$ and $K_{\text{bottom}}^{\text{Th}}$) in each simulation.

870





Figure 5. Dissolved ^{230}Th along 30°W in the Atlantic Ocean in (a) R6_B5 of the KREF_EXP and (b) PCE_EXP. (c) Reference coefficient (K_{ref}) along 30°W in the Atlantic Ocean in PCE_EXP. (d) The vertical profile of dissolved ^{230}Th averaged horizontally along 30°W in R6_B5 of KREF_EXP and PCE_EXP. The colored circles in Figs. 5a and 5b represent data from the Atlantic GEOTRACES data (GA02 and GIPY05; Schlitzer et al., 2018). The green, yellow, and orange circles in Fig. 5d represent the GA02 data and KREF_EXP and PCE_EXP simulation results.




880


Figure 6. (a) Dissolved ^{231}Pa , (b) dissolved ^{230}Th , (c) particulate ^{231}Pa , and (d) particulate ^{230}Th along 30°W in the Atlantic Ocean in CTRL_EXP. (e) Sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios normalized by the production ratio of 0.093 in CTRL_EXP. The colored circles represent observational data. Dissolved ^{231}Pa and ^{230}Th data are taken from the Atlantic GEOTRACES data (GA02 and GIPY05; Schlitzer et al., 2018). Particulate ^{231}Pa and ^{230}Th data are taken from the following references (Colley et al., 1995; Moran et al., 1997; Moran et al., 2001; Rutgers van der Loeff and Berger, 1993; Vogler et al., 1998; Walter et al., 1997; Cochran et al., 1987; Moran et al., 2002; Guo et al., 1995). The data of sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios are taken from the following references (Mangianini & Sonntag, 1977; Muller & Mangini, 1980; Anderson et al., 1983; Shimmield et al., 1986; Schmitz et al., 1986; Yang et al., 1986; Shimmield & Price, 1988; Yong Lao et al., 1992; François et al., 1993; Frank et al., 1994; Frank, 1996; Bradtmiller et al., 2014, Luo et al., 2010, and their supplemental data).

885

890



895 Figure 7. The difference between 3D_EXP and 1D_EXP (i.e., 3D_EXP minus 1D_EXP, which represents for ocean transport effect) of (a) dissolved ^{231}Pa , (b) dissolved ^{230}Th , (c) particulate ^{231}Pa , and (d) particulate ^{230}Th along 30°W in the Atlantic Ocean. (e) The difference between 3D_EXP and 1D_EXP of sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios normalized by the production ratio of 0.093.




900

Figure 8. The difference between CTRL_EXP and 3D_EXP (i.e., CTRL_EXP minus 3D_EXP, which represents for bottom scavenging effect) of (a) dissolved ^{231}Pa , (b) dissolved ^{230}Th , (c) particulate ^{231}Pa , and (d) particulate ^{230}Th along 30°W in the Atlantic Ocean. (e) The difference between CTRL_EXP and 3D_EXP of the sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios normalized by the

905

production ratio of 0.093.



910 Figure 9. Sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratios normalized by the production ratio of 0.093 in CTRL_EXP decomposed into contributions from (a) ocean transport solely from ^{231}Pa (i.e., $^{231}\text{Pa}(3\text{D})/^{230}\text{Th}(1\text{D})$), (b) ocean transport solely from ^{230}Th (i.e., $^{231}\text{Pa}(1\text{D})/^{230}\text{Th}(3\text{D})$), (c) bottom scavenging solely from ^{231}Pa (i.e., $^{231}\text{Pa}(\text{CTRL})/^{230}\text{Th}(3\text{D})$ minus $^{231}\text{Pa}(3\text{D})/^{230}\text{Th}(3\text{D})$), and (d) bottom scavenging solely from ^{230}Th (i.e., $^{231}\text{Pa}(3\text{D})/^{230}\text{Th}(\text{CTRL})$ minus $^{231}\text{Pa}(3\text{D})/^{230}\text{Th}(3\text{D})$).

Variable	Symbol	Value	Units
²³¹ Pa production from ²³⁵ U decay	β^{Pa}	2.33×10^{-3}	dpm m ⁻³ yr ⁻¹
²³⁰ Th production from ²³⁴ U decay	β^{Th}	2.52×10^{-2}	dpm m ⁻³ yr ⁻¹
Decay constant of ²³¹ Pa	λ^{Pa}	2.13×10^{-5}	yr ⁻¹
Decay constant of ²³⁰ Th	λ^{Th}	9.12×10^{-6}	yr ⁻¹
Sinking velocity of particles	w_s	1000	m yr ⁻¹
Thickness of euphotic zone	z_0	100	m
Penetration depth of CaCO ₃	z_p	2000	m
Dissolution constant of opal	B	0.12	°C ⁻¹ yr ⁻¹
Minimum temperature of sea water	T_0	-2	°C
Dissolution rate of POC	ε	0.858	-
Total activity of ²³¹ Pa or ²³⁰ Th	A_{total}	variable	dpm m ⁻³
Activity of dissolved ²³¹ Pa or ²³⁰ Th	A_d	variable	dpm m ⁻³
Activity of particle ²³¹ Pa or ²³⁰ Th	A_p	variable	dpm m ⁻³
Ratio of particle concentration to fluid density	C	variable	-

915

Table 1. Parameters of the ²³¹Pa and ²³⁰Th model.

Experiment	Siddall_EXP		CTRL_EXP	
	²³¹ Pa	²³⁰ Th	²³¹ Pa	²³⁰ Th
K_{ref}	1.0×10^7	1.0×10^7	1.0×10^7	$\left(\frac{C_{\text{total}}}{10^{-7}}\right)^{-0.42} \times 10^7$
K_{CaCO_3}	$K_{\text{ref}} / 40$	K_{ref}	$K_{\text{ref}} / 40$	K_{ref}
K_{opal}	$K_{\text{ref}} / 6$	$K_{\text{ref}} / 20$	$K_{\text{ref}} / 6$	$K_{\text{ref}} / 20$
K_{POC}	K_{ref}	K_{ref}	K_{ref}	K_{ref}
K_{bottom}	0	0	5.0×10^5	5.0×10^5

Table 2. Equilibrium partition coefficients in experiments Siddall_EXP and CTRL_EXP.

Experiment	Water-column reversible scavenging	Bottom scavenging	Ocean transport
CTRL_EXP	0	0	0
3D_EXP	0	X	0
1D_EXP	0	X	X

Table 3. Processes considered in additional experiments. A circle means that the process is considered, and a cross means that it is not considered.