



16 **Abstract.** Most land surface models can, depending on the simulation experiment, calculate the vegetation distribution and  
17 dynamics internally by making use of biogeographical principles or use vegetation maps to prescribe spatial and temporal  
18 changes in vegetation distribution. Irrespective of whether vegetation dynamics are simulated or prescribed, it is not practical  
19 to represent vegetation across the globe at the species level because of its daunting diversity. This issue can be circumvented  
20 by making use of 5 to 20 plant functional types (PFT) by assuming that all species within a single functional type show identical  
21 land-atmosphere interactions irrespective of their geographical location. In this study, we hypothesize that remote-sensing  
22 based assessments of above-ground biomass can be used to constrain the process in which real-world vegetation is discretized  
23 in PFT maps. Remotely sensed biomass estimates for Africa were used in a Bayesian framework to estimate the probability  
24 density distributions of woody, herbaceous, and bare soil fractions for the 15 land cover classes, according to the UN-LCCS  
25 typology, present in Africa. Subsequently, the 2.5<sup>th</sup> and 97.5<sup>th</sup> percentile of the probability density distributions were used to  
26 create 2.5% and 97.5% credible interval PFT maps. Finally, the original and constrained PFT maps were used to drive biomass  
27 and albedo simulations with the ORCHIDEE model. This study demonstrates that remotely sensed biomass data can be used  
28 to better constrain the share of dense forest PFTs but that additional information on bare soil fraction is required to constrain  
29 the share of herbaceous PFTs. Even though considerable uncertainties remain, using remotely sensed biomass data enhances  
30 the objectivity and reproducibility of the process by reducing the dependency on expert knowledge and allows assessing and  
31 reporting the credible interval of the PFT maps which could be used to benchmark future developments.

## 32 **1 Introduction**

33 Degradation, fires and deforestation of tropical forests are responsible for two thirds of the global net deforestation emissions  
34 (Houghton et al., 2012; Le Quéré et al., 2015; Friedlingstein et al., 2020). Although African tropical rainforests represent only  
35 one third of the global tropical rainforests (Lewis et al., 2009), they were responsible for almost all, i.e., 1.48 PgC in 2015 and  
36 1.65 PgC in 2016. of the net carbon (C) emissions of pan-tropical regions, but substantial uncertainty is associated with these  
37 estimates, i.e., 1.15 for 2015 and 1.0 PgC for 2016. mainly driven by fire and land use changes (Palmer et al., 2019). The  
38 uncertainty of model estimates, such as mentioned above, broadly comes from three sources: (1) the vegetation distribution in  
39 the model, (2) the ability of the model to simulate biomass accumulation of undisturbed vegetation, and (3) the ability of the  
40 model to simulate natural and anthropogenic disturbances of the standing biomass. As this study will focus on improving the  
41 description of the vegetation distribution, the first question that needs to be answered is why vegetation distribution remains  
42 so uncertain?

43 Most land surface models can either calculate the vegetation distribution internally by making use of biogeographical principles  
44 (Sitch et al., 2003; Krinner et al., 2005; Clark et al., 2011) or use vegetation maps to prescribe spatial and temporal changes in  
45 vegetation distribution. Where the first approach results in a description of the potential vegetation, the second approach is

46 more suitable when actual vegetation is to be studied. Irrespective of whether potential or actual vegetation is studied, it is not  
47 practical to represent vegetation across the globe at the species level because there are already over 60.000 tree species (Beech  
48 et al., 2017), not to mention the diversity in herbs, forbs and mosses. Land surface models represent this daunting diversity by  
49 making use of 5 to 20 plant functional types (PFT) (Huete et al., 2016). The underlying assumption of plant functional types  
50 is that all species within a single functional type show identical land–atmosphere interactions irrespective of their geographical  
51 location (Huete et al., 2016; Bonan et al., 2002; Brovkin et al., 1997; Chapin et al., 1996). Discretizing real-world vegetation  
52 in PFTs is a first source of uncertainty.

53 When actual vegetation is the focus of a modelling study, the vegetation distribution will have to be prescribed. The  
54 construction of vegetation maps first requires real-world observations, typically through satellite-based remote sensing.  
55 Current remote sensing technology does not enable distinguishing individual tree species; hence, vegetation is observed as  
56 land cover types (Defourny, P., 2019) which group vegetation with similar sensory characteristics. The remote sensing  
57 observations themselves as well as classifying them in land cover types are the second and third source of uncertainties (Hansen  
58 et al., 2013. Mitchard et al., 2014. Hurtt et al., 2004). Because the land surface models require the vegetation to be discretized  
59 in PFTs, which may differ between different land surface models, the land cover types will have to be remapped on PFT maps.  
60 The rules applied in remapping satellite-based land cover types in PFT maps is formalized in so-called “cross-walking tables”  
61 (CWT) (Poulter et al., 2011; Poulter et al., 2015) which are a fourth source of uncertainty (Hartley et al., 2017).

62 Although CWTs have been extensively used to create PFT maps (Wei et al., 2018; Wei et al., 2016; Poulter et al., 2011;  
63 Krinner et al 2005), the process of associating land cover types with specific PFTs remains difficult to reproduce since several  
64 iterations of expert knowledge are required (Poulter et al., 2011; Poulter et al., 2015). Various land cover classifications exist,  
65 the commonly used FAO (Food and Agriculture Organization) Land Cover Classification System (LCCS; Di Gregorio and  
66 Jansen, 2000). Most classes of the LCCS correspond to a mix of PFTs, which fractions are difficult to assess and likely variable  
67 across regions. For example, several classes are labelled as a mosaic of vegetation types (i.e., “Mosaic of natural vegetation  
68 (tree, shrubs, herbs)”); see Table 2 in Poulter et al., 2015). Not surprisingly, efforts have been made to decrease the need of  
69 expert knowledge in favour of more objective and reproducible approaches, e.g., classification rules based on a suite of  
70 improved and standard MODIS products (Wanxiao et al., 2008). Moreover, producing PFT maps from satellite-based land  
71 cover maps needs to become fully automated when the temporal frequency of satellite-based land cover and biomass maps  
72 will increase thanks to the recent GEDI Lidar data (Dubayah et al., 2020) or future SAR missions like the NASA-ISRO  
73 Synthetic Aperture Radar (NiSAR) or BIOMASS missions (Le Toan et al., 2011; Quegan et al., 2019).


74 In this study, we hypothesize that remote sensing-based assessments of above ground biomass (ABG) can decrease the  
75 dependency on expert knowledge when setting up CWTs and as such contribute to the automation of the land cover class  
76 mapping into PFTs for land surface models. The main rationale is that the above-ground biomass content of an ecosystem  
77 provides information on the fraction of tree PFTs of that ecosystem. In this context, the objective of this study are: (1) construct  
78 a framework of data assimilation in which biomass remote sensing products can be routinely used to update an existing or  
79 create a new CWT, (2) constrained a cross-walking table used to convert the ESA-CCI Global Land Cover map into a PFT

80 map, and (3) propagate the credible interval from using a CWT in the production of PFT maps, to the simulation results of  
81 biomass and albedo maps derived from a land surface model. Such a framework will be applied and tested over Africa using  
82 the above ground biomass product derived by (Bouvet et al., 2018) for that continent with the ORCHIDEE land surface model  
83 (Krinner et al., 2005) more specifically tag 2.0 revision 6592 close tag 2.2 used for the recent Climate Modelling  
84 Intercomparison Project - phase 6 (CMIP6) (Boucher et al., 2020).

## 85 **2 Materials and methods**

### 86 **2.1 Overview**

87 Cross-walking tables (CWT) (Poulter et al., 2015) are used to convert the 43 land cover types distinguished on the ESA-CCI  
88 land cover product into generic plant functional types (13 PFTs in (Poulter et al., 2015)) distinguished by large-scale land  
89 surface models such as the ORCHIDEE model (Krinner et al., 2005) used in this study. These generic PFTs are further grouped  
90 and/or divided to match each model-specific PFT classification, using additional grid-cell information to separate grassland  
91 and crop C3 versus C4 photosynthetic pathway (Still et al., 2003) and to split generic PFT according to bioclimatic zones (i.e.,  
92 Koppen Geiger climate classification map) (see more details for the ORCHIDEE model in (Lurton et al., 2020)). In this study,  
93 we provide a proof of concept by creating a new ORCHIDEE PFT map by combining information from the ESA-CCI land  
94 cover product and the AGB product for Africa (Bouvet et al., 2018) to estimate woody, herbaceous and bare soil cover fractions  
95 within each land cover type of the ESA-CCI product. Subsequently, the estimated cover fractions are used to constrain the  
96 existing CWT and create a new ORCHIDEE PFT map applicable primarily for Africa (Fig. 1). Finally, the impact of using  
97 AGB maps to constrain the PFT maps on the skill of the ORCHIDEE model to simulate the contemporary biomass and its  
98 spatial distribution over Africa is quantified. Note that the approach is tested over Africa but is generic enough to be applied  
99 everywhere.



100  
101  
102  
103  
104  
105

*Figure 1: Approach to assimilate the information held by aboveground biomass (AGB) maps into plant functional type (PFT) maps. Remote sensing AGB and land cover products are jointly assimilated to obtain cross-walking tables that can be used to make PFT maps. Owing to the uncertainty analysis in the data assimilation approach, an ensemble of cross-walking tables and PFT cover fraction maps can be produced. Subsequently, the land surface model ORCHIDEE can be run for different PFT maps to quantify the uncertainty from propagation of the uncertainty from remote sensing products into a model simulation.*

106  
107

## 2.2 Dataset products

### 2.2.1 Land cover map

108  
109  
110  
111  
112  
113  
114  
115  
116  
117

ESA’s Climate Change Initiative for Land Cover (CCI-LC) produced consistent global LC maps at 300 m spatial resolution on an annual basis for the year 2015 (Defourny, P., 2019). Only one year (2015) has been used to estimate the new vegetation cover cross-walking table. The typology of CCI-LC maps follows the Land Cover Classification System (LCCS) developed by the United Nations (UN) Food and Agriculture Organization (FAO), to enhance compatibility with similar products such as GLC2000, and GlobCover 2005 and 2009. The UN-LCCS typology was designed as a hierarchical classification, which allows adjusting the thematic detail of the legend. The “level 1” legend, also called “global” legend, counts 22 classes and is globally consistent and thus suitable for global applications such as creating PFT maps for land surface models. The “level 2” or “regional” legend counts 43 classes which are not present all over the world and could be used in this study given its focus on a single continent, i.e., Africa (see section 2.2.3). In addition, the UN-LCCS partly overlaps with the PFTs used in climate models.

118

### 2.2.2 Aboveground biomass map

119  
120

This study also makes use of a continental map of AGB of African savannas and woodlands for the year 2010 (Bouvet et al., 2018). The map has a 25 m resolution and is built from the 2010 L-band data of the Phased Array L-band Synthetic Aperture

121 Radar (PALSAR) on the Advanced Land Observing Satellite (ALOS) satellite. Covering the African continent required about  
122 180 data strips of which 91% were acquired between May and November 2010. The remaining 9% of the domain was filled  
123 with imagery from 2009 and 2008. The data have been processed by the Japan Aerospace Exploration Agency (JAXA) using  
124 the large-scale mosaicking algorithm described in Shimada and Ohtaki (2010), including ortho-rectification, slope correction  
125 and radiometric calibration between neighbouring strips, and multi-image filtering described in Bouvet et al., 2018.

126 The continental AGB map was derived as follows: (1) stratification into wet/dry season areas in order to account for seasonal  
127 effects in the relationship between PALSAR backscatter and AGB, (2) the development of a statistical model relating the  
128 PALSAR backscatter to observed AGB, (3) Bayesian inversion of the direct model, to obtain AGB and its credible interval  
129 for pixels where no observations are available, and (4) masking out non-vegetated areas using the ESA-CCI Land Cover dataset  
130 (but see section 2.1.1). The resulting AGB map was visually compared with existing AGB maps (Saatchi et al., 2011; Baccini  
131 et al., 2012; Avitabile et al., 2016) and cross-validated with AGB estimates obtained from field measurements and LiDAR  
132 datasets (Naidoo et al., 2015). Cross-validation revealed a good accuracy of the dataset, with an RMSD between 8 and 17 t/ha.  
133 For more details on the creation and evaluation of the AGB maps see Bouvet et al., 2018.

### 134 **2.2.3 Pre-processing**

135 One known limitation of the original AGB map (Bouvet et al., 2018) is the signal saturation and in some cases the decrease of  
136 the signal (Mermoz et al., 2015) occurring in L-band SAR for AGB values higher than 85 t/ha. In order to overcome this issue,  
137 a second AGB map was created, based on two other ancillary datasets: a map of tree cover (Hansen et al., 2013) and a map of  
138 tree height (Simard et al., 2011). The AGB was estimated by deriving an empirical relationship between biomass, available  
139 from airborne Lidar estimates, and the product of tree cover and tree height. The second version targets dense forest areas such  
140 as in the Congo basin and is used to adjust the AGB values at locations where signal saturation occurred. Because of a coarser  
141 resolution from the tree height map ( $0.01^\circ \times 0.01^\circ$ , 100 ha) than the original AGB map ( $0.00025^\circ \times 0.00025^\circ$ , 0.0625 ha), the  
142 new biomass map has been rescaled to  $0.01^\circ$  resolution. The rescaling drastically reduced the noise produced by PALSAR  
143 measurement artefacts (personal communication Thuy Le Toan). The original AGB map was downsampled by an average  
144 resampling method, i.e., computing the weighted average of all contributing pixels. To do so, we used the Gdalwarp function  
145 from GDAL (GDAL/OGR). The map used in this study is a composite of the two versions of the biomass map by using the  
146 following rules:

- 147     ▪ For broadleaved evergreen forests (UN-LCCS land cover type 50), flood forests (UN-LCCS 160), and closed  
148       broadleaved deciduous forests (UN-LCCS 61), the map based on tree cover and tree height was used because there  
149       is no AGB estimates in the map based on PALSAR.
- 150     ▪ For broadleaved deciduous forests (UN-LCCS 60) the maximum between the two maps was used because its biomass  
151       ranged around the threshold of  $85 \text{ t ha}^{-1}$  and may create truncated distribution.

- For the other land cover types, which typically have a biomass well below 85t ha<sup>-1</sup> the AGB value from the PALSAR map was used because it is considered more reliable than the statistical relationship between biomass, vegetation cover and vegetation height especially for the lower biomass.

Given the spatial domain of this study, only the 31 land cover types defined on the ESA CCI-LC map and present in Africa were retained. The complexity of the study was further reduced by removing all land types that cover less than 1.0% (304.158 km<sup>2</sup>) of the African surface or that contain less than 1% (i.e., 1.1 Gt) of the total AGB of Africa. Filtering retrained 15 out of the 31 land cover types including bare land. These 15 land cover types (Table 1) represent 96% of the surface of Africa and 98% of its AGB.

One additional issue had to be dealt with: the spatial resolution of the land cover map (9 ha) largely differed from the resolution of the AGB map (0.01°x0.01°, 100 ha). Therefore, each observational point on the AGB map is represented by 11x11 pixels on the land cover map. To simplify the overall data assimilation methodology (see section 3.2), we chose to use only AGB pixels (100 ha) which have a unique land cover type (i.e., pure pixels, in terms of land cover type). To this aim, the variety of land cover types across the 11x11 pixels within each AGB pixel (i.e., the number of present,  $Vlct$ ) was calculated and only pixels where  $Vlct=1$  was retained. Although this criterion resulted in discarding 99% of the pixels, each of the 15 land cover types considered could be represented by at least 2000 pixels. To remove outlier pixels, we choose to pick up the 2000 pixels strictly below the biomass value representing the 97.5<sup>th</sup> percentile of each LCT biomass distribution shown in the figure 2.

## 2.3 Data assimilation

### 2.3.1 Linking land cover fractions and AGB

A linear model was used to relate the satellite-based AGB of a 100-ha pixel to the cover fraction of the satellite-based vegetation types present at the same location. This relationship can be written as:

$$B_p = \sum_{i=1}^{nV} F_{p,i} \cdot B_{ref_i} \quad (1)$$

where  $B_p$ , is the AGB at a given pixel  $p$ ,  $F_{p,i}$  is the cover fraction of the vegetation type  $i$  (i.e., the generic plant functional type (PFT) used for land surface models, see section 2.1 - overview),  $B_{ref_i}$  is the reference AGB for the vegetation type  $i$  and  $nV$  is the number of vegetation types (i.e., number of PFTs) present in the pixel  $p$ . Given the number of unknowns ( $nV$  being usually above 1), equation 1 has many solutions; many of which have no biological meaning. The equifinality of this model can be reduced by arguing that the large difference in biomass between woody, herbaceous and non-vegetated ecosystems combined by their respective cover fraction explains most of the biomass at pixel level. Following this assumption, equation 1 can be simplified as:

$$B_p = F_{p,w} \cdot Bref_w + F_{p,h} \cdot Bref_h + F_{p,b} \cdot Bref_b \quad (2)$$

where  $F_{p,w}$ ,  $F_{p,h}$  and  $F_{p,b}$  are the fractions cover for woody vegetation (i.e., woody PFTs), herbaceous vegetation (i.e., grassland and cropland) and non-vegetated areas, respectively.  $Bref_w$  and  $Bref_h$  are the reference AGB of woody and herbaceous vegetation, respectively, while  $Bref_b = 0$ . Hence,  $F_{p,h}$  in equation 2 can be substituted according to  $F_{p,w} + F_{p,h} + F_{p,b} = 1$  to obtain:

$$B_p = F_{p,w} \cdot Bref_w + (1 - F_{p,w} - F_{p,b}) \cdot Bref_h \quad (3)$$

Although equation 3 no longer details which vegetation types  $i$  (i.e., PFTs) are present on each pixel  $p$ , it still has four unknowns and can, therefore, not be solved analytically. Nevertheless, a statistical solution is within reach if  $F_{p,w}$ ,  $F_{p,b}$ ,  $Bref_w$  and  $Bref_h$  are estimated from a population of AGB observations containing several independent repetitions that largely exceeds the number of unknowns. In this study, over 2000 repetitions were available for each of the 15 land cover types that were retained following filtering (section 2.2.3). The statistical solution will thus consist of four mean parameter values (i.e.,  $F_{p,w}$ ,  $F_{p,b}$ ,  $Bref_w$  and  $Bref_h$ ) for each of these 15 land cover types.

As described in section 2.2.3. the selection of homogeneous AGB pixels, i.e., which have a unique land cover class across the 11.11 underlying land cover sub-pixels allow us to rewrite the equation 3 as follow:

$$\underline{Bp_p} = F_{lc,w} \cdot Bref_{lc,w} + (1 - F_{lc,w} - F_{lc,b}) \cdot Bref_{lc,h} \quad (4)$$

where  $\underline{Bp_p}$  is the average AGB of a specific land cover type  $lc$  and  $F_{lc,w}$ ,  $F_{lc,b}$ ,  $Bref_{lc,w}$ ,  $Bref_{lc,h}$  are the unknowns. The unknown parameters of the regression model (eq. 4) were estimated by using a Bayesian inference method. This approach has been chosen because it helps to synthesize various sources of information as well as to propagate credible intervals in the result of our land surface model (Ellison 2004). Bayesian inference requires, however, setting prior probability distributions for each of the unknowns, i.e., the biomasses and land cover fractions for each of the 16 land cover types. Given these prior probability distributions, Bayesian inference retrieves the posterior probability distribution for each of the unknown parameters.

### 2.3.2 Prior value distributions for $Bref_{lc,w}$ , $Bref_{lc,h}$ and $Bp_p$

The AGB pixels were stratified according to their land cover type and for each land cover type the information contained in the distribution of the satellite based AGB served to estimate the mean and standard deviation of the prior values of  $Bref_{lc,w}$ . To avoid negative  $Bref$  values we used a normal truncated distribution with  $0 < a, b < +\infty$  where  $(a, b)$  are the truncated range:

$$Bref_{lc,w} \sim N(\mu_{lc,w}, \sigma_{lc,w}, a, b) \quad (5)$$

where,  $\mu_{lc,w}$  is calculated as follow:




209  $\mu_{lc,w} = X^{th} \text{ per.} (Bp_{lc})$  (6)

210

211 Where  $Bp_{lc}$  is a vector containing  $Bp_p$  values that belong to the land cover type  $lc$  and  $X^{th} \text{ per}$  denotes the 97.5<sup>th</sup> percentile for  
 212 the woody cover types. This choice assumes that with the 97.5<sup>th</sup> percentile we select the AGB value of a pixel covered only  
 213 by woody vegetation (i.e., woody PFT) for the selected land cover type. In contrast to using a few in-situ observations to define  
 214  $\mu_{lc,w}$ , our approach offers the advantage to rely on a large ensemble of satellite-derived AGB observations and to be coherent  
 215 with the following optimization.

216



217

218 **Figure 2: Probability density distribution of the pure land cover pixel for biomass concentration  $Bp_p$  for 15 selected land cover types.**  
 219 **The dashed line represents the 97.5<sup>th</sup> percentile used as the prior estimate for the reference biomass concentration for trees  $Bref_{lc,w}$ . The**  
 220 **dashed line represents the 50<sup>th</sup> percentile also used as the prior estimate for the reference biomass for herbaceous cover**

221

222

223 Without any information about the variability of  $Bref_{lc,w}$ , we choose to represent  $\sigma_{lc,w}$  as:

224  $\sigma_{lc,w} = \mu_{lc,w} \cdot 0.0375$  (7)

225

226

227

Where 0.0375 accounts for a 30% uncertainty encompassed between the interquartile range of the normally distributed  $Bref_{lc,w}$ . Compared to  $Bref_{lc,w}$ ,  $Bref_{lc,h}$  is more difficult to assess from the satellite-derived data because it often shows bimodal distributions which may stem from biomass degradation or the presence of shrubs which biomass better resembles that of a

228 grassland than a woody ecosystem (Fig. 2). We found that while the 2.5<sup>th</sup> percentile is representing the lowest biomass for  
 229 herbaceous ecosystem, the 50<sup>th</sup> percentile seems to better describe  $Bref_{lc, h}$ , following equation 6. Having no information about  
 230 the variability of  $Bref_{lc, h}$ ,  $\sigma_{lc, w}$  followed equation 7.

231 Finally,  $Bp_p$  which was the 97.5<sup>th</sup> for woody cover types or the 50<sup>th</sup> percentile for herbaceous cover types, comes with a  
 232 measurement uncertainty that was thought to follow a normal truncated distribution with  $0 < a, b < Bref_{lc, w}$  where (a, b) are  
 233 the truncated range. Given that this uncertainty is not known at the pixel level, an uninformative prior was set for the standard  
 234 deviation  $\sigma b_{lc}$  which can vary between 0 and 200 t/ha. We deliberately took a large uncertainty to cover the observation that  
 235 considerable uncertainty remains in satellite-based biomass estimates (Bouvet et al., 2018):

$$236 \quad Bp_p \sim N(\mu, \sigma b_{lc}^2, a, b) \text{ with } \sigma b_{lc} \sim U(0, 200) \text{ and } \mu = Bp_p \quad (8)$$

### 237 2.3.3 Prior value distributions for $F_{lc, w}$ , $F_{lc, b}$ and $F_{lc, h}$

238  $F_{lc, w}$ ,  $F_{lc, b}$  and  $F_{lc, h}$  were defined as fractions of respectively woody vegetation, bare soil and herbaceous vegetation within a  
 239 given land cover type, their values thus range between zero and one and their sum is equal to 1. For this reason, a Dirichlet  
 240 distribution was used to describe the probability distribution of the woody, bare soil and herbaceous cover fractions:

$$241 \quad (F_{lc, w}, F_{lc, b}, F_{lc, h}) \sim Di(\theta_{lc, t}, \theta_{lc, b}, \theta_{lc, h}) \quad (9)$$

242 OpenBUGS (Thomas, 2010), the software that was used in this study, cannot use a Dirichlet distribution as a stochastic node.  
 243 This constraint can be overcome by making the cover fractions dependent on each other:

$$244 \quad F_{lc, w} = q_{lc, 1} \quad (10)$$

$$245 \quad F_{lc, b} = q_{lc, 2} \cdot (1 - q_{lc, 1}) \quad (11)$$

$$246 \quad F_{lc, h} = (1 - q_{lc, 1}) \cdot (1 - q_{lc, 2}) \quad (12)$$

247 Let  $q_{lc, i}$  with  $i = 1, \dots, K - 1$  and  $K$  the number of fractions, be a series of independent beta distributions,  $Be(\alpha_i, \beta_i)$ .

$$248 \quad q_{lc, i} \sim Be(\alpha_{lc, i}, \beta_{lc, i}) \quad (13)$$

249 The parameters of the beta distribution of the cover fraction of bare soil, woody vegetation and herbaceous vegetation (eq. 9)  
 250 can then be estimated as follows:

$$251 \quad \alpha_{lc,i} = \theta_{lc,i} \cdot (\omega_{lc,i} - 2) + 1 \quad (14)$$

$$252 \quad \omega_{lc,i} \sim U(0,1000) \quad (15)$$

253 where  $\theta_{lc,i}$  which represents the fraction of each land cover type taken from expert knowledge used to define the so-called  
 254 cross walking table (CWT) and taken from a recent update of the CWT.  $\omega_{lc,i}$  was described by an uninformative uniform  
 255 distribution and thus reflects the relatively low trust we have in the current CWT. The dependencies between the beta  
 256 distributions comes from  $\beta_{lc,i}$  that is estimated as:

$$257 \quad \beta_{lc,i} = \sum_{u=i+1}^K \alpha_{lc,u} \quad (16)$$

## 258 **2.4 Credible interval propagation**

### 259 **2.4.1 Propagating the credible interval from the CWT into the PFT map**

260 The posterior estimates of the cover fractions ( $F_{lc,w}$ ,  $F_{lc,b}$ ,  $F_{lc,h}$ ) will be directly used to make up a new cross-walking table.  
 261 The posterior estimates of the cover fractions values are then used to recalculate woody and herbaceous fraction of each generic  
 262 PFT of the CWT. In other words, we keep the original split of the different woody PFT defined in the prior CWT and only  
 263 rescale the total woody fraction to  $F_{lc,w}$ . Then we rescale the bare soil fraction based on  $F_{lc,b}$  to finally rescale short vegetation  
 264 PFTs (grass and crop).

265 Given that these posterior estimates come with a probability distribution, a probability distribution of the CWT could be made.  
 266 In this study, the 2.5 and 97.5 percentiles and the mode, i.e., the most common value, of the posterior estimates were used to  
 267 create three cross-walking tables that were then applied on the ESA-CCI-LC product to create two PFT maps that represent  
 268 the 95% interval confidence of the ESA-CC-LC product and one PFT map which represents the one that is used in an  
 269 ORCHIDEE simulation. The impact of the various PFT map was quantified for simulated above ground biomass and simulated  
 270 surface albedo by running three simulations that only differed by the PFT map used to initialize the ORCHIDEE land surface  
 271 model.

272 In the study, the uncertainty propagation index aimed to identify the ecoregions where the AGB and surface albedo estimates  
 273 are most sensitive to uncertainties from the PFT map. This sensitivity was calculated as:

$$274 \quad S_{eco,b} = \frac{ABS(X^{97.5} - X^{2.5})}{ABS(F_{eco,b}^{97.5} - F_{eco,b}^{2.5}) \times 100} \quad (17)$$

275 Where X stands for AGB (t/ha) or surface albedo (unitless),  $S_{eco,b}$  is expressed in the unit of X for a 1% change in bare soil  
 276 fraction.

### 277 **2.4.2 Description of the ORCHIDEE land surface model**

278 ORCHIDEE (Krinner et al., 2005; Boucher et al 2020) is the land surface model of the IPSL (Institut Pierre Simon Laplace)  
 279 Earth system model. Hence, by conception, it can be coupled to a global circulation model. In a coupled setup, the atmospheric

280 conditions affect the land surface which, in turn, affects the atmospheric conditions. However, when a study focuses just on  
281 changes in the land surface ORCHIDEE rather than on the interaction with the atmosphere, it also can be run as a stand-alone  
282 land surface model. The stand-alone configuration receives atmospheric conditions such as temperature, humidity, and wind,  
283 to mention a few, from the so-called meteorological forcing. The resolution of the meteorological forcing determines the spatial  
284 resolution and can cover any area ranging from a single grid point to the entire globe.

285 Although ORCHIDEE does not enforce a spatial or temporal resolution, the model does use a spatial grid and equidistant time  
286 steps. The spatial resolution is an implicit user setting that is determined by the resolution of the meteorological data.  
287 ORCHIDEE can run on any temporal resolution; however, this apparent flexibility is restricted as the processes are nested and  
288 formalised at given time steps: half-hourly (i.e., photosynthesis and energy budget), daily (i.e., net primary production), and  
289 annual (i.e., vegetation dynamics). Hence, meaningful simulations have a temporal resolution of 1 min to 1 h for the energy  
290 balance, water balance, and photosynthesis calculations. In the land-only configuration used in this study, the default time step  
291 for these processes is 30 minutes.

292 When an application requires the land surface to be characterised by its actual vegetation, the vegetation will have to be  
293 prescribed by annual land cover maps. These maps must follow specific rules for the land surface models to be able to read  
294 them. In the case of ORCHIDEE the share of each of the 15 possible plant functional types needs to range between 0 and 1  
295 and be specified for each pixel. When satellite-based land cover maps are used as the basis for an ORCHIDEE-specific PFT  
296 map, the satellite-based land cover classification will need to be converted to match the ORCHIDEE specifications. As  
297 mentioned already above, this involves two steps: i) the derivation of generic PFTs from the satellite land cover classes (in our  
298 case the ESA-CCI-LC product) through the CWT discussed in this paper and ii) the final mapping of the generic PFTs into  
299 the 15 ORCHIDEE-specific PFTs using additional information on the bioclimatic zones and the partition of grassland/crops  
300 into C3 versus C4 photosynthetic pathway (Lurton et al., 2020).

301 In this study, AGB was defined as the sum of leaf biomass, fruit biomass, aboveground sapwood biomass, and aboveground  
302 heartwood biomass which are default output variables of ORCHIDEE. Surface albedo was defined as the albedo in the visible  
303 wavelengths and is a default output variable of ORCHIDEE.

304 **Table 1: Description of the 15 plant functional types (PFT) used in ORCHIDEE to represent global vegetation.**

PFT	Climate	Vegetation type	Phenology class
1	global	NA	Bare soil
2	Tropical	Woody	Broadleaf evergreen
3	Tropical	Woody	Broadleaf deciduous
4	Temperate	Woody	Needleleaf Evergreen
5	Temperate	Woody	Broadleaf Evergreen
6	Temperate	Woody	Broadleaf Summer green
7	Boreal	Woody	Needleleaf Evergreen
8	Boreal	Woody	Broadleaf Summer green
9	Boreal	Woody	Needleleaf Deciduous
10	Temperate	Herbaceous	Natural (C3)
11	global	Herbaceous	Natural (C4)
12	global	Herbaceous	Managed (C3)
13	global	Herbaceous	Managed (C4)
14	Tropical	Herbaceous	Natural (C3)
15	Boreal	Herbaceous	Natural (C3)

305

### 306 **2.4.3 Experimental setup**

307 ORCHIDEE tag 2.0 (rev 6592) was used to run tree simulations that only differed by the PFT map used. Following a 340-  
 308 yearlong spinup to initialise the carbon pools in the model, each simulation consisted of a 110years long simulation between  
 309 1901 to 2010 with the CRU-NCEP v8 climate reconstruction (Viovy, 2017) that matched the simulation years. CO2  
 310 concentration was fixed to 299.16 ppm and thus corresponds to the 2010 concentration.

### 311 **2.4.4 Ecoregions**

312 Results related to the land surface model simulation were presented by subdividing the African continent into ecologically  
 313 homogeneous regions, so-called ecoregions, as defined by Olson et al., (2001).

## 314 **3 Results**

### 315 **3.1 Prior and posterior distributions estimates**

#### 316 **3.1.1 Vegetation cover fraction: prior and reference biomass distributions**


317 Prior distributions for the cover fractions and reference biomasses were determined for all 15 land cover classes separately,  
318 nevertheless, four broadly different groups could be distinguished: (1) The 97.5<sup>th</sup> percentile of biomass distribution for each  
319 land cover belonging in the first group was so high, i.e., from 245 to 416 t/ha, that the land cover types in this group must  
320 correspond to a substantial tree cover, i.e., a woody cover fraction from 0.58 to 0.75. Examples of this group are land cover  
321 types UN-LCCS 50, 61 and 160 (tree cover broadleaf types in Table 2). (2) Contrary to the first group, the 97.5<sup>th</sup> percentile of  
322 biomass distribution for each land cover type of the second group is so low, i.e., from <12 to 42 t/ha, that these land cover  
323 types must be dominated by grasses or bare soil, i.e., a woody cover fraction of 0.1 or less and a substantial bare soil cover  
324 fraction up to 0.71. Examples of this group are UN-LCCS 130, 150 and 153 (grassland and sparse vegetation in Table 2). (3)  
325 The biomass of the third group falls in between these extremes representing mosaic land cover types like the UN-LCCS 10,  
326 11, 30, 40, 100, 110 and 120 (mosaic landscape in Table 2). When taken over the African continent, the biomass distribution  
327 of these land cover types shows bimodal biomass distributions indicating considerable variability within these land cover types  
328 (Fig. 2). (4) The bimodal biomass distribution of the fourth group is backed by a rather high woody reference biomass  
329 associated with a low woody cover fraction which may represent an ecosystem highly disturbed by either silvicultural practice  
330 or a fire regime. UN-LCCS 60, 62 fall into this group which represents the woodland to dry savanna continuum.

#### 331 **3.1.2 Vegetation cover fraction: posterior distributions**

332 Owing to the Bayesian approach, the woody and herbaceous fraction within each land cover type is no longer deterministic  
333 (as was the case with the previous generation of cross-walking table such as in Poulter et al., 2015) but now comes with a  
334 distribution. This distribution is the outcome of propagating the credible interval on the retrieved parameters obtained from  
335 the Bayesian approach into the final product, i.e., the PFT cover fraction map. The 95% credible interval was studied by  
336 comparing the 2.5 and 97.5 percentiles of the distribution of woody, herbaceous and bare soil fractions ( $F_{lc, w}$ ,  $F_{lc, h}$ ,  $F_{lc, b}$ ).

337 The mean change in forest cover fraction between the 2.5 and 97.5 percentiles of the distribution of constrained PFT maps  
338 over Africa was  $1.6 \pm 2.6\%$ . At the ecoregion scale (when averaging the cover fraction over the ecoregion), the largest  
339 uncertainty in forest cover fraction was found in the Congo basin with an average of  $-6.3 \pm 0.5\%$  for the six ecoregions where  
340 LCT 50 is dominant (Fig 3A).

341 The 95% uncertainty interval for bare soil cover fraction is  $13 \pm 8\%$  mainly due to the large uncertainty of the cropland and  
342 mosaic cropland (UN-LCCS 10, 11, 30, 40). In ecoregions where these LCTs are dominant, this credible interval increases to  
343  $24 \pm 7\%$  (Fig. 3B). Moreover, dense forest land cover type i.e., LCT 50, 160 also come with  $15 \pm 4\%$  uncertainty in their bare  
344 soil fraction estimates (Fig. 3B).



345  
 346  
 347  
 348  
 349  
 350

**Figure 3: Uncertainty in CWT constrained by an AGB map. Absolute change in forest (A) and bare soil (B) cover fraction (%) between the 2.5 and 97.5 percentile PFT maps. High values represent a large uncertainty in the estimation of the true cover fraction. C and D represent disagreement estimated as the difference between the CWT based on expert knowledge and the CWT constrained by an AGB map. Disagreement in forest (C) and bare soil (D) is expressed as absolute change (%). High values represent a strong disagreement between the two methods. Black lines delimit the different ecoregion according to Olson et al., 2001.**

351 **Table 2: Surface area (%), share in the continental biomass (%), prior parameters, and posterior median and credible interval values**  
352 **for each of the 15 land cover types considered in this study. The numbering, description and surface area of each land cover type is**  
353 **based on the ESA-CCI product (Defourny, P., 2019), where its share in the continental biomass is based on a compilation of Bouvet et**  
354 **al 2018.  $\theta_{lc}$ ,  $\mu_{lc}$  and  $\sigma_{lc}$  represent the parameters describing the prior distributions of  $F_{lc}$  and  $Bref_{lc}$ . Estimation of these parameters is**  
355 **detailed in section 2.3. For each land cover type and each parameter, the 2.5. the constrained and the 97.5 percentiles are computed.**  
356 **We use the mode for the constrained CWT as an approximation of the posterior  $\theta_{lc, w}$ , since the posterior distributions of  $F_{lc, i}$  may be**  
357 **asymmetric.**

id	Information UN-LCCS short description	Surface area (%)	Biomass (%)	Priors					Posterior fractions used for the CWTs								
				$F_{lc,w}$ $\theta_{lc,w}$	$F_{lc,h}$ $\theta_{lc,h}$	$F_{lc,b}$ $\theta_{lc,b}$	$Bref_{lc,w}$ $\mu_{lc,w} \pm \sigma_{lc,w}$	$Bref_{lc,h}$ $\mu_{lc,h} \pm \sigma_{lc,h}$	CWT 2.5 %			CWT 97.5 %			Refined CWT		
10	Cropland rainfed	7.6	5	0.01	0.98	0.01	83±3.1	9±0.3	0.19	0.34	0.47	0.13	0.86	0.01	0.14	0.83	0.03
11	Cropland rainfed - Herbaceous cover	3.2	3.3	0.01	0.74	0.01	84±3.1	6±0.2	0.13	0.49	0.38	0.09	0.89	0.02	0.11	0.85	0.04
30	Mosaic cropland (>50%) / natural vegetation (tree/shrub/herbaceous cover) (<50%)	2.3	3.1	0.25	0.49	0.01	85±3.2	4±0.2	0.2	0.77	0.03	0.17	0.83	0.00	0.18	0.81	0.01
40	Mosaic natural vegetation (tree/shrub/herbaceous cover) (>50%) / cropland (<50%)	2.2	1.9	0.5	0	0.01	84±3.1	9±0.3	0.26	0.69	0.05	0.22	0.78	0.00	0.24	0.75	0.01
50	Tree cover broadleaved evergreen closed to open (>15%)	6.7	45.1	0.99	0.29	0.01	416±15	9±0.3	0.71	0.02	0.27	0.79	0.1	0.11	0.76	0.01	0.23
60	Tree cover broadleaved deciduous closed to open (>15%)	4.2	8.7	0.7	0.14	0.01	245±9.2	9±0.3	0.27	0.66	0.07	0.23	0.76	0.01	0.25	0.73	0.02
61	Tree cover broadleaved deciduous closed (>40%)	0.4	1.8	0.85	0.44	0.01	252±9.5	9±0.3	0.54	0.44	0.02	0.61	0.3	0.09	0.58	0.4	0.02
62	Tree cover broadleaved deciduous open (15-40%)	10.6	13.1	0.55	0.39	0.01	111±4.2	9±0.3	0.35	0.61	0.04	0.3	0.69	0.01	0.32	0.66	0.02
100	Mosaic tree and shrub (>50%) / herbaceous cover (<50%)	1.8	1.5	0.6	0.59	0.01	85±3.2	9±0.3	0.16	0.77	0.07	0.13	0.86	0.01	0.15	0.84	0.01
110	Mosaic herbaceous cover (>50%) / tree and shrub (<50%)	1.6	1.2	0.4	0.39	0.01	75±2.8	5±0.2	0.07	0.85	0.08	0.05	0.94	0.01	0.06	0.93	0.01
120	Shrubland	13.3	7.7	0.6	0.98	0.01	85±3.2	9±0.3	0.16	0.74	0.1	0.12	0.87	0.01	0.14	0.85	0.01
130	Grassland	6.5	1.5	0.01	0.2	0.01	42±1.6	3±0.1	0.11	0.56	0.33	0.07	0.92	0.01	0.09	0.9	0.01
150	Sparse vegetation (tree/shrub/herbaceous cover) (<15%)	1.6	0.2	0.1	0.29	0.7	12±0.5	3±0.1	0.04	0.45	0.51	0.15	0.01	0.84	0.1	0.19	0.71
153	Sparse herbaceous cover (<15%)	1.1	0.1	0.01	0.24	0.7	22±0.8	3±0.1	0.1	0.29	0.61	0.01	0.97	0.02	0.02	0.81	0.17
160	Tree cover flooded fresh or brackish water	0.7	3.5	0.75	1	0.01	386±14	9±0.3	0.6	0.39	0.01	0.69	0.27	0.04	0.65	0.34	0.01

358


359 Nonetheless, in a classic simulation experiment the most common values of  $F_{lc, w}$ ,  $F_{lc, h}$ ,  $F_{lc, b}$  will be used. The most common  
360 values of  $F_{lc, w}$ ,  $F_{lc, h}$ ,  $F_{lc, b}$  are given by the mode of the posterior distribution (“constrained CWT” in Table 2.a). The mode was  
361 used to show the difference between the original and the constrained PFT maps (Fig. 3C-D). The mean difference in forest  
362 cover fraction between the prior (original) and the constrained PFT maps is  $-15 \pm 12\%$  (Fig. 3C). Largest disagreement between  
363 was observed over the Somali Acacia-Commiphora Bushlands and Thickets and the Kalahari Xeric Savanna where forest  
364 cover fraction was found to be on average  $32 \pm 1\%$  lower in the constrained PFT maps (Fig. 3C). The Bare soil cover fraction  
365 changes on average by  $3.1 \pm 0.5\%$  (Fig. 3D). The constrained PFT map has on average  $16 \pm 4\%$  more bare soil cover fraction  
366 over the Congo Basin than the original map (Fig. 3C).

### 367 3.3 Uncertainty propagation of the PFT maps on the aboveground biomass and visible albedo estimates from 368 ORCHIDEE simulations

369 PFT maps are essential boundary conditions of land surface models because they condition the spatial distribution of various  
370 ecosystem states-properties (i.e., carbon content, albedo, water-carbon-energy fluxes, etc). When tested with ORCHIDEE tags  
371 2.0 (rev 6592), the absolute difference in biomass stock between the 2.5 and 97.5 percentile maps was  $0.5 \pm 5.7$  t/ha (Fig. 4A)  
372 representing 0.2 t/ha/% of cover fraction (Fig 4C). Notable exception is the Congo basin where different PFT maps could result



373 in AGB estimates that differ by 18 t/ha (Fig. 4A) for a 6.5% difference in the forest cover (Fig 3A). Different PFT maps make  
374 the average visible albedo range from  $0.081 \pm 0.055$  to  $0.083 \pm 0.055$ . The largest uncertainty for the visible albedo simulated  
375 with ORCHIDEE was found over the Nigerian lowland forest (0.158) and West Sudanian Savanna (0.107) (Fig. 4B) which  
376 represent a 24% to 11% change in forest cover respectively. The sensitivity is the highest in the western Congo basin with  
377 1.4% of albedo/% of cover faction. In contrast, West Sudanian Savanna possesses a low sensitivity with 0.5%. To summarise,  
378 we found that a smaller forest to bare soil transition uncertainty can drastically change the albedo of an ecoregion than a larger  
379 uncertainty in the grassland/cropland to bare soil transition.



380

381

382

383

*Figure 4: Confident interval propagation of the PFTs maps into AGB and visible albedo simulated by ORCHIDEE. (A) uncertainty propagation into AGB and (B) uncertainty propagation into visible albedo from the difference between the 2.5% and the 97.5% PFT map defined by the optimisation procedure. uncertainty propagation index (eq. 16) for AGB (C) and visible albedo (D).*

## 384 **4 Discussion**

### 385 **4.1 Discretizing vegetation**

386 Irrespective of the data products, the methods, and the model used, discretizing vegetation comes with its own challenges.  
387 Discretizing transitions of ecosystems into land cover type classes (Sankaran et al., 2005), for example, can lead to systematic  
388 uncertainties since all pixels that belong to the same land cover class will get the same vegetation cover fractions in the cross-  
389 walking table (see 4.1.3). This approach articulates a key assumption underlying the PFT-approach, i.e., that only one life form  
390 survives and thus dominates the vegetation due to competition for nutrients, light and water (Hutchinson et al., 1961). However,  
391 the Savanna ecosystem, for example, is characterised by the coexistence of trees, shrubs and grasses which has been explained  
392 by interactions between vegetation, rainfall, fire, and browsing regimes (Eigentler and Sherratt., 2020). This makes savannas  
393 one of the most difficult ecosystems to classify in a land cover type and subsequently convert it into a PFT map.

394 Over Africa, land cover classes such as shrubland (UN-LCCS 120) represent a wide range of ecosystems, from sparse xeric  
395 shrubland composed of small bushes, e.g., *Penzia incana* (Thunb.) Kuntze, grasses, e.g., *Sip agrostis* spp. such as found in  
396 Karoo desert, to dense thicket composed by succulent, e.g., *Portulacaria afra* Jacq. and spinescent shrubs (~3m tall) (Mills, et  
397 al., 2005). Combining land cover types and biomass maps showed that the shrubland pixels in Africa more often resemble  
398 sparse xeric shrubland than dense thickets. Improving the ability to simulate land surface properties of shrublands in a changing  
399 world, especially in Africa where shrub encroachment is an important land cover dynamic (Wigley et al., 2010. Buitenwerf et  
400 al., 2012. O'Connor et al., 2014), is likely to benefit from a more detailed representation of shrublands in land surface models.  
401 A first step could be to represent shrubs as small trees, as was tested with the ORCHIDEE model for arctic ecosystems (Druel  
402 et al., 2017), but ultimately the control of precipitation on plant density (Rietkerk et al., 2002) should also be modelled.

403 Another major challenge with discretizing vegetation is how degraded ecosystems should be classified. From a modelling  
404 point of view, they should be classified as the land cover type that occurred prior to the degradation and the cause of the  
405 degradation. e.g., fire, grazing, erosion, should be explicitly accounted for in the land surface model. This ideal strongly differs  
406 from the current approach in which the degraded vegetation is classified as if it is in its natural state. Even when having the  
407 correct PFTs, the current approach would fail to simulate the observed biomass if degradation occurred. As an alternative, the  
408 PFT map could duplicate all PFTs to distinguish between a PFT in its natural state and in its degraded state. This approach in  
409 which degradation is accounted for in the PFT maps would, however, reduce degradation to a binary problem rather than  
410 addressing its continuous nature.

### 411 **4.2 Knowledge gain from using the AGB map**

412 In the absence of an AGB map, previous efforts to build cross-walking tables (Poulter et al., 2015) had to rely in part on expert  
413 knowledge. That generation of cross-walking tables can be considered as the best-available-knowledge in the absence of AGB  
414 data or other information on the land surface cover. The method developed and demonstrated in this study mostly relies on  
415 data but comes with its own assumptions and statistical complexities. The key assumptions are that: (1) previous cross-walking

416 tables (Poulter et al., 2015) are a reliable source to set the prior distribution for PFT cover, (2) the biomass map (Bouvet et al.,  
417 2018) is a reliable source to set the prior distribution of the reference biomasses, and (3) the land cover classification contains  
418 homogeneous land cover types (Defourny, P. et al., 2019). A key question is thus whether the added complexity justifies the  
419 knowledge gained by jointly assimilating a land cover and a biomass map when producing a CWT?  
420 Ideally this question should be addressed by assessing the reduction of the credible interval associated to the posterior  
421 distribution of the PFT map when using the AGB map to constrain the CWT (in comparison to a prior when no AGB is used).  
422 However, the present generation of CWT without AGB information, does not come with a distribution (except the attempt in  
423 Hartley et al., (2017)), calling for an alternative approach to assess the knowledge gain. Given that the prior distribution of the  
424 cover fraction was based on the previous CWT, the difference between the prior and the posterior distributions can be  
425 considered as the knowledge gained from using AGB information. Following this reason, the question we seek to answer is:  
426 “Is the cover fraction used by the original cross walking table falling outside the 95% credible interval of our posterior  
427 estimate?”  
428 If the answer is no, the biomass map is more likely in agreement with the previous effort to estimate the original cross walking  
429 table. If the answer is yes, adding the information contained in the satellite-based biomass maps is most likely in strong  
430 disagreement with the previous effort to estimate the original cross walking table. The original CWT has a global extent, and  
431 the constrained CWT is only valid for Africa. Therefore, knowledge gains should be carefully interpreted as they may reflect  
432 trade-offs that had to be made previously to construct a global rather than regional CWT. Knowledge gains were assessed for:  
433 “croplands”, “dense evergreen forests”, “woodlands and savannas”, and “xeric shrublands and grasslands” separately.

#### 434 **4.2.1 Croplands (UN-LCCS 10, 11, 30, 40).**

435 Despite the cover fraction of woody vegetation on croplands being close to none in the original CWT, this study found that  
436 the four land cover types associated with croplands, UN-LCCS 10, 11, 30, 40 are in fact covered with 11% to 24% woody  
437 vegetation (Table 2). This large difference in the presence of woody vegetation on croplands is also reflected in the biomass  
438 data, which suggest two distinct but co-existing agricultural systems in Africa, i.e., one system with a low biomass and one  
439 around with a higher biomass.

440 The agricultural system with the low biomasses likely represents annually replanted crops such as millet, sorghum, wheat,  
441 sweet potatoes or cassava (FAO), with a maximum reported biomass between 10 and 15 t/ha for high-input cropping associated  
442 with commercial production of cassava and sweet potatoes. These values are in line with values estimated as reference biomass  
443 (see 2.3.2). Nonetheless, 97% of total cropland area Africa is rainfed (Calzadilla et al., 2009) and most of Africa’s agricultural  
444 land is used for subsistence or small-scale farming associated with low-input cropping which explains why the actual average  
445 biomass estimate from the CESBIO map for cropland is between  $2.0 \pm 0.7$  t/ha (Fig. 2) and thus considerably lower than the  
446 potential production.

447 The high biomass agricultural system which is estimated at  $83 \pm 3$  t/ha in the CESBIO map (Fig. 2) likely includes plantations  
448 for coffee, rubber, fruits as well as shelter trees and forest remnants (FAO). Permanent croplands do not have their own land

449 cover type in the UN-LCCS or in ORCHIDEE. The mixture of bare soil, herbaceous vegetation and woody vegetation, makes  
450 it challenging to discretize African croplands into the current PFTs (Table 1). Moreover, small changes in the woody reference  
451 biomass for high biomass agricultural systems lead to large changes in cover fractions of herbaceous vegetation and bare soil  
452 ratio. Without constraint reference biomass estimates, total biomass alone does not sufficiently constrain the share of woody  
453 vegetation. For the time being, high biomass agricultural systems could be with a woodland fraction ranging from 9.0% to  
454 26% (Table 2). Although this could be an acceptable solution for biomass and albedo simulations, it will underestimate the  
455 agricultural production in the region.

#### 456 **4.2.2 Tropical rainforest (UN-LCCS 50, 160).**

457 The woody cover fraction of tropical rainforest in the original CWT is close to 90% and falls outside the credible interval of  
458 the posterior estimates, i.e., 71 to 79%. This lower cover fraction from many pixels classified as tropical rainforest that do not  
459 reach the reference biomass of  $416 \pm 16$  t/ha (Fig. 2). The reference derived from the biomass map matches the AGB observed  
460 at field plots of intact forests in the Congo basin (Lewis et al., 2013) but the large value in bare soil cover fraction for these  
461 land cover types may thus reflect wide-spread degradation of the forests in the region (Tyukavina et al., 2018) or a too high  
462 reference biomass (Kearsley et al., 2013).

#### 463 **4.2.3 Tropical deciduous forest, woodland, and savanna (UN-LCCS 61, 60 and 62).**

464 The woody cover fraction of the tropical deciduous forest ranged between 45% and 75% in the original CWT. Refining the  
465 CWT using AGB information shifts this range to between 27% and 58%. For savanna (UN-LCCs 62) the original cover  
466 fractions are within the constrained 95% CI. For woody cover, the fraction of deciduous forest (UN-LCCS 61) decreased from  
467 85% to 58%. We observe an overall decrease for the woody cover fraction since the reference biomass is much higher than  
468 the actual biomass of most of the pixels.

469 Although the reference biomasses used in this study are in line with previously reported values (Carreira et al., 2013),  
470 disagreement between the original and the constrained CWT is considerable. The original CWT starts from the view that all  
471 ecosystems (except croplands) are in their natural state. The AGB map, however, does not contain any evidence in support of  
472 this view but rather suggests that 50% of the savanna (UN-LCCs 62) are 65% below their reference biomass. Likewise, 50%  
473 for dry woodland (UN-LCCs 60) are 71% below their reference biomass (Fig. 2). The AGB map thus suggests wide-spread  
474 degradation of these ecosystems which are in a highly anthropized region (Mitchard et al., 2013). Uncertainty coming from  
475 the reference biomasses could be reduced by field observations at the ecoregion or finer spatial scales.

476 For deciduous forest, however, the difference in cover fraction of woody vegetation between the original CWT and the  
477 constrained CWT could also be explained by an inaccurate estimation of the reference biomass due to a too coarse definition  
478 of the deciduous woody vegetation ranging from deciduous forest, over woodlands to savannas which are composed by  
479 different dominant tree species, with different biomasses (Sawadogo et al., 2010).

#### 480 **4.2.4 Xeric shrubland (UN-LCCS 100, 110, 120).**

481 The woody cover fraction of xeric shrublands and grasslands ranged between 40 and 60% in the original CWT. Accounting  
482 for the information contained in the AGB map significantly decreased the woody cover fraction range toward 5.0 and 16%.  
483 Indeed, shrubs which represent a large part of the xeric shrublands were originally classified as woody vegetation for the  
484 ORCHIDEE model (i.e., when moving from the generic PFTs to the ORCHIDEE-specific PFTs; see section 2). This  
485 assumption is true from an ecological point of view but in a simplified world like in land surface models, xeric shrubland has  
486 an aboveground biomass that resembles cropland and grassland (Fig. 2). By overlaying the land cover type and aboveground  
487 biomass maps, 37% of the African shrublands were found to be degraded with a biomass of  $2.7 \pm 1.5$  t/ha, 54% were found to  
488 be intact with a biomass of  $22 \pm 19$  t/ha and 9% of the shrublands are thickets with a biomass of  $68 \pm 11$  t/ha. This is in line with  
489 other aboveground biomass estimates from remote sensing products (Saatchi et al., 2011; Mitchard et al., 2013; Avitabile et  
490 al., 2016) and in situ measurements where shrublands, degraded thicket, and intact thicket in south Africa accumulated 3. 24  
491 and 102 t/ha of biomass respectively (Mills, et al., 2005). These findings suggest that in the model world, xeric shrubland is  
492 best represented by a large fraction of herbaceous plant functional groups, when the overall objective is to model AGB.

#### 493 **4.2.4 Sparse vegetation (UN-LCCS 150, 153).**

494 The constrained cover fraction estimates are in line with the original CWT for UN-LCCS 150 which represent the most  
495 common class of sparse vegetation. The constrained cover fraction for UN-LCCS 153 has a larger herbaceous i.e., 29 to 97%,  
496 then the bare soil cover fraction, i.e., 2.0 to 61% contrary to the original CWT. The herbaceous cover fraction could be  
497 overestimated if a too low reference biomass was used. A reference biomass of 3.0 t/ha was used and is acceptable compared  
498 to the reported biomass for the Succulent and Nama Karoo Biomes ranging from 0.5 to 7.6 t/ha (Rutherford, 1978; Rutherford  
499 and Westfall, 1986). Given the current lack of reference biomass observations, disagreement between the original and  
500 constrained CWT could be resolved by using an independent estimate of bare soil fraction.

### 501 **4.3 Consequences for land surface modelling**

#### 502 **4.3.1 Which land cover types affect the biomass estimate?**

503 The large disagreement in cover fraction estimates (30 to 40%) resulted in small disagreement in biomass, i.e., <1.0 t/ha in  
504 regions with little precipitation like Somali Acacia-Commiphora Bushlands and Thickets and the Kalahari Xeric Savanna. This  
505 counter-intuitive result is explained by the growth processes simulated in ORCHIDEE. Under xeric climate conditions  
506 ORCHIDEE simulates low tree biomasses (< 2.0 t/ha) because the low precipitation and subsequent plant water availability  
507 results in a continuous high tree mortality. Nonetheless, in forest ecoregions like the eastern Guinean forests or in the Congo  
508 basin, where the sensitivity to a change in the cover fractions ranged from 1.0 to 5.0 t/ha/% and had a considerable impact on  
509 the simulation since a 15% uncertainty in the bare soil fraction may lead to a 75 t/ha uncertainty of the biomass in the tropical  
510 forest of the Congo basin. Underestimating the forest cover in humid ecoregions will have a much larger consequence on the

511 simulated AGB than overestimating the forest cover in xeric ecoregions. The uncertainty surrounding the land cover fractions  
512 should thus be further reduced for the land cover types that already come with the lowest uncertainty, i.e., the forests.

### 513 **4.3.2 Which land cover types affect the albedo estimate?**

514 As for AGB, uncertainties in land cover fractions are only partly reflected in the uncertainties of the visible albedo. Dampening  
515 is caused by the fact that the reflectivity of grassland (0.06), cropland (0.06) are close to the leaf reflectivity of a forest (0.03  
516 to 0.04) compared to bare soils reflectivity (0.1 to 0.25 depending on the colour of the soil) in ORCHIDEE. By increasing the  
517 bare soil cover fraction, the albedo will increase accordingly but changing forest into grassland will not drastically change  
518 albedo. The most sensitive area is the western tropical forest in the Congo basin for which a 15% change in bare soil cover  
519 fraction may trigger a 15% change in the visible albedo (fig 3C). Similar as for AGB, the uncertainty surrounding the land  
520 cover fractions of the forested land cover types should be further reduced to reduce the uncertainty of the model simulations.

### 521 **4.4 Outlook**

522 In this study a single biomass map was used as this enabled keeping the focus on the method itself. Nevertheless, other biomass  
523 products are available (Saatchi et al., 2011; Baccini et al., 2012; Avitabile et al., 2016; Santoro et al., 2021) and could have  
524 been used. Repeating this study for each of these biomass products would add another source of uncertainty to the cross-  
525 walking table. Owing to the method presented in this study, this uncertainty could then be propagated into the PFT map and  
526 all the way up to the simulated biomass, albedo -as done in this study for one biomass product- and other land surface  
527 properties. Considering different biomass products would give an insight of the impact of satellite-based biomass estimates on  
528 the discretisation of the vegetation and by extension surface properties as estimated by land surface models. Likewise, a single  
529 land cover map has been used in our analysis, but other products are available as well (Copernicus, UN-spider, Li et al., 2020).  
530 By using different land cover maps, one could quantify the uncertainty in the land cover classification and propagate it to  
531 evaluate its impact on the simulated land surface properties.

532 Compared to other continents, the Africa vegetation has been documented by relatively few quantitative observations (Mills,  
533 et al., 2005; Saatchi et al., 2011; Asner et al., 2012; Réjou-Méchain et al., 2015). Hence, it is the continent where remote  
534 sensing data could largely enhance our knowledge on the issue. Recent high-resolution satellite observations bear the promise  
535 to significantly reduce the credible interval around the aboveground carbon stock to estimate the CO<sub>2</sub> emissions from tropical  
536 forests (Hansen et al., 2013; Bouvet et al., 2018; Defourny et al., 2019; Buchhorn et al., 2020) but land surface models will  
537 need to be ready to routinely assimilate these data to fully benefit from the information contained in biomass maps. This study  
538 demonstrated one way of how satellite-based biomass data can help modellers to constraint the initialization process by means  
539 of refining the cross-walking tables that are used to map land cover classes derived from satellite observations into PFT maps.  
540 Nevertheless, biomass maps could be used for applications other than model initialization (this study), including model  
541 parameterisation and model evaluation.

542 The biomass map could be used to optimise model parameters related to growth, turnover and mortality to better simulate the  
543 vegetation biomass for the different PFTs. The evaluation stage could benefit from the biomass maps by benchmarking the  
544 model results against observed relationships between biomass-climate and biomass-land-use to better distinguish and simulate  
545 the difference between actual and potential biomass (Sankaran et al., 2005). Although the availability of several biomass  
546 products makes it possible to use one product to inform the cross-walking tables and another product to evaluate the simulated  
547 surface properties, the magnitude of present-day differences between biomass products (Mitchard et al., 2013) is expected to  
548 result in major inconsistencies when different biomass products are used for different purposes (e.g., assimilation,  
549 parameterization, evaluation) into a single analysis. In this study, less than 0.01% (see 2.3.1) of the information contained in  
550 the biomass map was used to constraint the cross-walking table and none was used to optimise model parameters. The  
551 simulated biomass remains, therefore, largely independent from the biomass map which implies that a single biomass map can  
552 be used for land cover optimisation (as in this study), and in a second step for parameter optimization or model evaluation.  
553 With an increase in resolution of the land cover map comes a decrease in the reliance on the cross-walking tables. Cross  
554 walking tables will no longer be required once the resolution will be high enough (around 10 x 10 m) such that each pixel  
555 contains a single vegetation type equivalent to a single PFT classification used by land surface models (Li et al., 2020). No  
556 longer having to rely on cross walking tables would likely reduce the width of the credible intervals of the PFT map. As there  
557 would no longer be a need to estimate woody and herbaceous fractions, there would no longer be a need for the information  
558 contained in the biomass map. It will then be feasible to solely use biomass maps to better parameterize the processes that  
559 contribute to simulating the reference biomass. It should be noted, however, that higher resolutions will not solve the basic  
560 challenge of discretizing vegetation. High resolution land cover maps would split structurally complex ecosystems, for  
561 example savannas, into a pure forest fraction and a pure grassland fraction. This would overlook the interactions between the  
562 grasses and the trees which are among the defining ecological characteristics of a savanna.  
563 Finally, we should note that other satellite-derived products than the AGB could be used to constrain the mapping of the land  
564 cover classes into model PFTs (i.e., CWT). For instance, the global tree cover fraction map, at 30-meter resolution, from  
565 Hansen et al., (2013) could also be used to constraint the fraction of bare soil within each land cover class like what was done  
566 in this study with the AGB map.

## 567 **4.5 Conclusion**

568 This study demonstrates how an aboveground biomass map could be used to constrain a cross-walking table that enables  
569 remapping land cover types derived from satellite-observations into plant functional types used as a boundary condition in  
570 land surface models. Given that previous cross-walking tables did not report uncertainties as they were mostly based on expert-  
571 knowledge, it remains unclear how much the use of an additional constraint really improved the cross-walking tables.  
572 Nevertheless, the considerable uncertainties remaining in the cross-walking table that made use of the aboveground biomass  
573 map suggests that total biomass map should be complemented with a bare soil map to better constrain the cross-walking table.  
574 Likewise, the reference biomass for both herbaceous and woody vegetation need to be constrained to at least the ecoregion



575 scale to avoid underestimating or overestimating bare soil fractions. The method developed in this study helped to estimate  
576 the uncertainty of cross-walking tables which can now be used to benchmark further methodological developments. Moreover,  
577 the method identified bare soil cover fraction would be required to reduce the uncertainty of future cross-walking tables and  
578 the plant functional type maps they generate.

## 579 **5 Acknowledgements**

580 This study was primarily financed by the French space agency, Centre National d'Etude Spatiale (CNES), through the  
581 "BIOMASS-Valorisation" project from the TOSCA research program which contributed to the funding of Guillaume Marie  
582 and Cécile Dardel. The Marie Sklodowska Curie Fellowship CLIMPRO (MSCA-Fellowship EU 895455) partly funded  
583 Guillaume Marie.

584

585

586

## 587 **6 Data availability**

- 588 ▪ CESBIO African AGB map. Biomass map of Africa created by CESBIO can be downloaded at [https://www.theia-](https://www.theia-land.fr/en/product/african-biomass-map)  
589 [land.fr/en/product/african-biomass-map](https://www.theia-land.fr/en/product/african-biomass-map). It consists of a GIF file in which Africa is spatially discretized in pixels of  
590 1x1km. The unit is a ton of dry mass per hectare (t/ha). Contact person: alexandre.bouvet@cesbio.cnes.fr
- 591 ▪ The land cover map is freely available from: <http://www.esa-landcover-cci.org>.
- 592 ▪ The ecoregion map used in this study is freely available from:  
593 <https://databasin.org/datasets/68635d7c77f1475f9b6c1d1dbe0a4c4c/>

## 594 **7 Code availability**

- 595 ▪ All R scripts and ORCHIDEE tags 2.0 (rev 6592) source code is available at:  
596 <https://zenodo.org/badge/latestdoi/345907299> or DOI: [10.5281/zenodo.4785328](https://doi.org/10.5281/zenodo.4785328)
- 597 ▪ ORCHIDEE tags 2.0 (rev 6592) code is also available from:  
598 [https://forge.ipsl.jussieu.fr/orchidee/wiki/GroupActivities/CodeAvailabilityPublication/ORCHIDEE\\_tags\\_2.0\\_gmd](https://forge.ipsl.jussieu.fr/orchidee/wiki/GroupActivities/CodeAvailabilityPublication/ORCHIDEE_tags_2.0_gmd_2021_Africabrowser/tags/ORCHIDEE_2_1)  
599 [\\_2021\\_Africabrowser/tags/ORCHIDEE\\_2\\_1](https://forge.ipsl.jussieu.fr/orchidee/wiki/GroupActivities/CodeAvailabilityPublication/ORCHIDEE_tags_2.0_gmd_2021_Africabrowser/tags/ORCHIDEE_2_1)

## 8 Author contribution:

G. Marie, S. Luysaert and P. Peylin designed the experiments and G. Marie carried them out. G. Marie developed the OPENBUGS model code and performed the simulations. G. Marie and S. Luysaert prepared the manuscript with contributions from all co-authors.

## 9 References

- ALOS - EoPortal Directory - Satellite Missions*, earth.esa.int/web/eoportal/satellite-missions/a/alos. *Tools*, orchidas.lsce.ipsl.fr/dev/lccci/tools.php.
- Asner, Gregory P., et al., “A Universal Airborne LiDAR Approach for Tropical Forest Carbon Mapping.” *Oecologia*, vol. 168. no. 4. 2011. pp. 1147–1160., doi:10.1007/s00442-011-2165-z.
- Avitabile, Valerio, et al., “An Integrated Pan-Tropical Biomass Map Using Multiple Reference Datasets.” *Global Change Biology*, vol. 22. no. 4. 2016. pp. 1406–1420., doi:10.1111/gcb.13139.
- Baccini, A., et al., “Estimated Carbon Dioxide Emissions from Tropical Deforestation Improved by Carbon-Density Maps.” *Nature Climate Change*, vol. 2. no. 3. 2012. pp. 182–185., doi:10.1038/nclimate1354.
- Bauer, Peter, et al., “The Quiet Revolution of Numerical Weather Prediction.” *Nature*, vol. 525. no. 7567. 2015. pp. 47–55., doi:10.1038/nature14956.
- Beech, E., et al., “GlobalTreeSearch: The First Complete Global Database of Tree Species and Country Distributions.” *Journal of Sustainable Forestry*, vol. 36. no. 5. 2017. pp. 454–489., doi:10.1080/10549811.2017.1310049.
- Bonan, Gordon B., et al., “Landscapes as Patches of Plant Functional Types: An Integrating Concept for Climate and Ecosystem Models.” *Global Biogeochemical Cycles*, vol. 16. no. 2. 2002. doi:10.1029/2000gb001360.
- Boucher, Olivier, et al., “Presentation and Evaluation of the IPSL-CM6A-LR Climate Model.” *Journal of Advances in Modeling Earth Systems*, vol. 12. no. 7. 2020. p. e2019MS002010. doi:https://doi.org/10.1029/2019MS002010.
- Bouvet, Alexandre, et al., “An Above-Ground Biomass Map of African Savannas and Woodlands at 25 m Resolution Derived from ALOS PALSAR.” *Remote Sensing of Environment*, vol. 206. 2018. pp. 156–173., doi:10.1016/j.rse.2017.12.030.
- Brovkin, Victor, et al., “A Continuous Climate-Vegetation Classification for Use in Climate-Biosphere Studies.” *Ecological Modelling*, vol. 101. no. 2-3. 1997. pp. 251–261., doi:10.1016/s0304-3800(97)00049-5.
- Buchhorn, Marcel, et al., “Copernicus Global Land Cover Layers—Collection 2.” *Remote Sensing*, vol. 12. no. 6. 2020. p. 1044., doi:10.3390/rs12061044.
- Buitenwerf, R., et al., “Increased Tree Densities in South African Savannas: >50 Years of Data Suggests CO<sub>2</sub> as a Driver.” *Global Change Biology*, vol. 18. no. 2. 2011. pp. 675–684., doi:10.1111/j.1365-2486.2011.02561.x.

- Calzadilla, Alvaro, et al., “Climate Change and Agriculture: Impacts and Adaptation Options in South Africa.” *Water Resources and Economics*, vol. 5. 2014. pp. 24–48., doi:10.1016/j.wre.2014.03.001.
- Carreira, Valeria Paula, et al., “Gene-by-Temperature Interactions and Candidate Plasticity Genes for Morphological Traits in *Drosophila Melanogaster*.” *PLoS ONE*, vol. 8. no. 7. 2013. doi:10.1371/journal.pone.0070851.
- Chapin, F. Stuart, et al., “Plant Functional Types as Predictors of Transient Responses of Arctic Vegetation to Global Change.” *Journal of Vegetation Science*, vol. 7. no. 3. 1996. pp. 347–58. doi:10.2307/3236278.
- Clark, D. B., et al., “The Joint UK Land Environment Simulator (JULES), Model Description – Part 2: Carbon Fluxes and Vegetation Dynamics.” *Geoscientific Model Development*, vol. 4. no. 3. 2011. pp. 701–722., doi:10.5194/gmd-4-701-2011.
- Defourny, Pierre, and ESA Land Cover CCI project team. *Dataset Record: ESA Land Cover Climate Change Initiative (Land\_Cover\_cci): Global Land Cover Maps, Version 2.0.7*. 28 Nov. 2019. catalogue.ceda.ac.uk/uuid/b382ebe6679d44b8b0e68ea4ef4b701c.
- “Development Activities.” *Documentation/Forcings – ORCHIDEE*, forge.ipsl.jussieu.fr/orchidee/wiki/Documentation/Forcings.
- Dietze, Michael C., et al., “Iterative Near-Term Ecological Forecasting: Needs, Opportunities, and Challenges.” *Proceedings of the National Academy of Sciences*, vol. 115. no. 7. 2018. pp. 1424–1432., doi:10.1073/pnas.1710231115.
- Di Gregorio, A. (2005). Land cover classification system: classification concepts and user manual: LCCS (Vol. 2). Food & Agriculture Org.
- Druel, A., Peylin, P., Krinner, G., Ciais, P., Viovy, N., Peregon, A., ... & Mironycheva-Tokareva, N. (2017). Towards a more detailed representation of high-latitude vegetation in the global land surface model ORCHIDEE (ORC-HL-VEGv1. 0). *Geoscientific Model Development*, 10(12), 4693-4722.
- Dubayah, Ralph, et al., “The Global Ecosystem Dynamics Investigation: High-Resolution Laser Ranging of the Earth’s Forests and Topography.” *Science of Remote Sensing*, vol. 1. June 2020. p. 100002. doi:10.1016/j.srs.2020.100002.
- Eamus, Derek, et al., “Savannas.” *Vegetation Dynamics*, pp. 383–414., doi:10.1017/cbo9781107286221.017.
- Eigentler, L., and J.a. Sherratt. “Spatial Self-Organisation Enables Species Coexistence in a Model for Savanna Ecosystems.” *Journal of Theoretical Biology*, vol. 487. 2020. p. 110122., doi:10.1016/j.jtbi.2019.110122.
- Ellison, Aaron M. “Bayesian Inference in Ecology.” *Ecology Letters*, vol. 7. no. 6. 2004. pp. 509–520., doi:10.1111/j.1461-0248.2004.00603.x.
- “FAO.org.” *Land Cover Classification System (LCCS)*, Food and Agriculture Organization of the United Nations, [www.fao.org/land-water/land/land-governance/land-resources-planning-toolbox/category/details/en/c/1036361/](http://www.fao.org/land-water/land/land-governance/land-resources-planning-toolbox/category/details/en/c/1036361/).
- Friedlingstein, Pierre, et al., “Global Carbon Budget 2020.” *Earth System Science Data*, vol. 12. no. 4. Dec. 2020. pp. 3269–340. doi:10.5194/essd-12-3269-2020.

- GDAL/OGR contributors (2022). GDAL/OGR Geospatial Data Abstraction Software Library. Open-Source Geospatial Foundation. URL <https://gdal.org>
- Hansen, M. C., et al., “High-Resolution Global Maps of 21st-Century Forest Cover Change.” *Science*, vol. 342. no. 6160. 2013. pp. 850–853., doi:10.1126/science.1244693.
- Hardin, G. “The Competitive Exclusion Principle.” *Science*, vol. 131. no. 3409. 1960. pp. 1292–1297., doi:10.1126/science.131.3409.1292.
- Hartley, A. J., MacBean, N., Georgievski, G., & Bontemps, S. (2017). Uncertainty in plant functional type distributions and its impact on land surface models. *Remote Sensing of Environment*, 203. 71-89.
- “Home.” *ESA Climate Office*, [climate.esa.int/en/](http://climate.esa.int/en/).
- “Home.” *ORCHIDEE*, [orchidee.ipsl.fr/](http://orchidee.ipsl.fr/).
- Houghton, R. A., et al., “Carbon Emissions from Land Use and Land-Cover Change.” *Biogeosciences*, vol. 9. no. 12. 2012. pp. 5125–5142., doi:10.5194/bg-9-5125-2012.
- Huete, Alfredo. “Vegetation’s Responses to Climate Variability.” *Nature*, vol. 531. no. 7593. Mar. 2016. pp. 181–82. doi:10.1038/nature17301.
- Hurt, G. C., et al., “Linking Models and Data on Vegetation Structure.” *Journal of Geophysical Research: Biogeosciences*, vol. 115. no. G2. 2010. doi:10.1029/2009jg000937.
- “Japan Aerospace Exploration Agency.” *JAXA*, [global.jaxa.jp/](http://global.jaxa.jp/).
- Kearsley, Elizabeth, et al., “Conventional Tree Height–Diameter Relationships Significantly Overestimate Aboveground Carbon Stocks in the Central Congo Basin.” *Nature Communications*, vol. 4. no. 1. Aug. 2013. p. 2269. doi:10.1038/ncomms3269.
- Krinner, G., et al., “A Dynamic Global Vegetation Model for Studies of the Coupled Atmosphere-Biosphere System.” *Global Biogeochemical Cycles*, vol. 19. no. 1. 2005. doi:10.1029/2003gb002199.
- Le Toan, T., et al., “The BIOMASS Mission: Mapping Global Forest Biomass to Better Understand the Terrestrial Carbon Cycle.” *Remote Sensing of Environment*, vol. 115. no. 11. Nov. 2011. pp. 2850–60. doi:10.1016/j.rse.2011.03.020.
- Lewis, Simon L., et al., “Increasing Carbon Storage in Intact African Tropical Forests.” *Nature*, vol. 457. no. 7232. 2009. pp. 1003–1006., doi:10.1038/nature07771.
- Lewis, Sophie C., et al., “Modeling Insights into Deuterium Excess as an Indicator of Water Vapor Source Conditions.” *Journal of Geophysical Research: Atmospheres*, vol. 118. no. 2. 2013. pp. 243–262., doi:10.1029/2012jd017804.
- Li, Qingyu, et al., “Mapping the Land Cover of Africa at 10 m Resolution from Multi-Source Remote Sensing Data with Google Earth Engine.” *Remote Sensing*, vol. 12. no. 4. 2020. p. 602., doi:10.3390/rs12040602.
- Li, Wei, et al., “Major Forest Changes and Land Cover Transitions Based on Plant Functional Types Derived from the ESA CCI Land Cover Product.” *International Journal of Applied Earth Observation and Geoinformation*, vol. 47. May 2016. pp. 30–39. doi:10.1016/j.jag.2015.12.006.

- Li, Wei, et al., “Gross and Net Land Cover Changes in the Main Plant Functional Types Derived from the Annual ESA CCI Land Cover Maps (1992–2015).” *Earth System Science Data*, vol. 10. no. 1. Jan. 2018. pp. 219–34. doi:<https://doi.org/10.5194/essd-10-219-2018>.
- Liang, Shunlin. *Advances in Land Remote Sensing: System, Modelling, Inversion and Application*. Springer, 2008.
- Lurton, Thibaut, et al., “Implementation of the CMIP6 Forcing Data in the IPSL-CM6A-LR Model.” *Journal of Advances in Modeling Earth Systems*, vol. 12. no. 4. 2020. p. e2019MS001940. doi:<https://doi.org/10.1029/2019MS001940>.
- Mermoz, Stéphane, et al., “Decrease of L-Band SAR Backscatter with Biomass of Dense Forests.” *Remote Sensing of Environment*, vol. 159. Mar. 2015. pp. 307–17. doi:10.1016/j.rse.2014.12.019.
- Mills, A. J., et al., “Ecosystem Carbon Storage under Different Land Uses in Three Semi-Arid Shrublands and a Mesic Grassland in South Africa.” *South African Journal of Plant and Soil*, vol. 22. no. 3. 2005. pp. 183–190., doi:10.1080/02571862.2005.10634705.
- Mills, A. J., et al., “Ecosystem Carbon Storage under Different Land Uses in Three Semi-Arid Shrublands and a Mesic Grassland in South Africa.” *South African Journal of Plant and Soil*, vol. 22. no. 3. 2005. pp. 183–190., doi:10.1080/02571862.2005.10634705.
- Mitchard, Edward T. A., et al., “Markedly Divergent Estimates of Amazon Forest Carbon Density from Ground Plots and Satellites.” *Global Ecology and Biogeography*, vol. 23. no. 8. 2014. pp. 935–946., doi:10.1111/geb.12168.
- Mitchard, Edward Ta, et al., “Uncertainty in the Spatial Distribution of Tropical Forest Biomass: A Comparison of Pan-Tropical Maps.” *Carbon Balance and Management*, vol. 8. no. 1. 2013. doi:10.1186/1750-0680-8-10.
- Naidoo, Laven, et al., “Savannah Woody Structure Modelling and Mapping Using Multi-Frequency (X-, C- and L-Band) Synthetic Aperture Radar Data.” *ISPRS Journal of Photogrammetry and Remote Sensing*, vol. 105. July 2015. pp. 234–50. doi:10.1016/j.isprsjprs.2015.04.007.
- NiSAR : First global SAR mission, in partnership with ISRO, <https://nisar.jpl.nasa.gov> .
- O'connor, Tim G, et al., “Bush Encroachment in Southern Africa: Changes and Causes.” *African Journal of Range & Forage Science*, vol. 31. no. 2. 2014. pp. 67–88., doi:10.2989/10220119.2014.939996.
- Oke, T. R. *Boundary Layer Climates*. Routledge, 2002.
- Organisation Des Nations Unies Pour L'alimentation Et L'agriculture. *Food and Agriculture Organization of the United Nations*, [www.fao.org/home/fr/](http://www.fao.org/home/fr/).
- Palmer, Paul I., et al., “Net Carbon Emissions from African Biosphere Dominate Pan-Tropical Atmospheric CO<sub>2</sub> Signal.” *Nature Communications*, vol. 10. no. 1. 2019. doi:10.1038/s41467-019-11097-w.
- Poulter, B., et al., “Plant Functional Type Mapping for Earth System Models.” *Geoscientific Model Development*, vol. 4. no. 4. 2011. pp. 993–1010., doi:10.5194/gmd-4-993-2011.
- Pütz, Sandro, et al., “Long-Term Carbon Loss in Fragmented Neotropical Forests.” *Nature Communications*, vol. 5. no. 1. 2014. doi:10.1038/ncomms6037.

- Quegan, Shaun, et al., “The European Space Agency BIOMASS Mission: Measuring Forest Above-Ground Biomass from Space.” *Remote Sensing of Environment*, vol. 227. June 2019. pp. 44–60. doi:10.1016/j.rse.2019.03.032.
- Quéré, C. Le, et al., “Global Carbon Budget 2015.” *Earth System Science Data*, vol. 7. no. 2. 2015. pp. 349–396., doi:10.5194/essd-7-349-2015.
- Réjou-Méchain, Maxime, et al., “Using Repeated Small-Footprint LiDAR Acquisitions to Infer Spatial and Temporal Variations of a High-Biomass Neotropical Forest.” *Remote Sensing of Environment*, vol. 169. 2015. pp. 93–101., doi:10.1016/j.rse.2015.08.001.
- Rietkerk, M., Boerlijst, M. C., van Langevelde, F., HilleRisLambers, R., de Koppel, J. V., Kumar, L., ... & de Roos, A. M. (2002). Self-organization of vegetation in arid ecosystems. *The American Naturalist*, 160(4), 524-530.
- Saatchi, S. S., et al., “Benchmark Map of Forest Carbon Stocks in Tropical Regions across Three Continents.” *Proceedings of the National Academy of Sciences*, vol. 108. no. 24. 2011. pp. 9899–9904., doi:10.1073/pnas.1019576108.
- Sankaran, Mahesh, et al., “Determinants of Woody Cover in African Savannas.” *Nature*, vol. 438. no. 7069. 2005. pp. 846–849., doi:10.1038/nature04070.
- Santoro, Maurizio, et al., “The Global Forest Above-Ground Biomass Pool for 2010 Estimated from High-Resolution Satellite Observations.” *Earth System Science Data Discussions*, August 2021. pp. 1–38. doi:10.5194/essd-13-3927-2021
- Schaaf: MODIS Albedo and Reflectance Anisotropy Products. Accessed 24 Mar. 2021.
- Shimada, Masanobu, and Takahiro Ohtaki. “Generating Large-Scale High-Quality SAR Mosaic Datasets: Application to PALSAR Data for Global Monitoring.” *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, vol. 3. no. 4. 2010. pp. 637–656., doi:10.1109/jstars.2010.2077619.
- Simard, Marc, et al., “Mapping Forest Canopy Height Globally with Spaceborne Lidar.” *Journal of Geophysical Research: Biogeosciences*, vol. 116. no. G4. 2011. doi:https://doi.org/10.1029/2011JG001708.
- Sitch, S., et al., “Evaluation of Ecosystem Dynamics, Plant Geography and Terrestrial Carbon Cycling in the LPJ Dynamic Global Vegetation Model.” *Global Change Biology*, vol. 9. no. 2. 2003. pp. 161–185., doi:10.1046/j.1365-2486.2003.00569.x.
- Still, Christopher J., et al., “Global Distribution of C3 and C4 Vegetation: Carbon Cycle Implications.” *Global Biogeochemical Cycles*, vol. 17. no. 1. 2003. pp. 6-1-6–14. doi:https://doi.org/10.1029/2001GB001807.
- Sun, Wanxiao, et al., “Mapping Plant Functional Types from MODIS Data Using Multisource Evidential Reasoning.” *Remote Sensing of Environment*, vol. 112. no. 3. Mar. 2008. pp. 1010–24. doi:10.1016/j.rse.2007.07.022.
- Thomas, Neal, 2010. "Overview". OpenBUGS website. Retrieved 9 October 2010.
- Tyukavina, Alexandra, et al., “Congo Basin Forest Loss Dominated by Increasing Smallholder Clearing.” *Science Advances*, vol. 4. no. 11. 2018. doi:10.1126/sciadv.aat2993.

UN-spider: “CCI LAND COVER - S2 Prototype Land Cover 20m Map of Africa.” *UN*, [un-spider.org/links-and-resources/data-sources/cci-land-cover-s2-prototype-land-cover-20m-map-africa](http://un-spider.org/links-and-resources/data-sources/cci-land-cover-s2-prototype-land-cover-20m-map-africa).

Viovy, Nicolas, 2017. CRUNCEP data set:

<https://vesg.ipsl.upmc.fr/thredds/fileServer/work/p529viov/cruncep/readme.html>

Wigley, B. J., et al., “Bush Encroachment under Three Contrasting Land-Use Practices in a Mesic South African Savanna.” *African Journal of Ecology*, vol. 47. 2009. pp. 62–70., doi:10.1111/j.1365-2028.2008.01051.x.