

Global seamless tidal simulation using a 3D unstructured-grid model (SCHISM v5.10.0)

Y. Joseph Zhang¹, Tomas Fernandez-MontBlanc², William Pringle³, Hao-Cheng Yu¹, Linlin Cui¹, Saeed Moghimi⁴

5 ¹Center for Coastal Resource Management, Virginia Institute of Marine Science, College of William & Mary, Gloucester Point, 23062, USA

² Earth Sciences Department, University of Cadiz INMAR, Puerto Real, 11519, Spain

³Environmental Science Division, Argonne National Laboratory, Lemont, IL, 60439, USA

⁴NOAA National Ocean Service, Silver Spring, MD, 20910, USA

10

Correspondence to: Y. Joseph Zhang (yjzhang@vims.edu)

Abstract. We present a new 3D unstructured-grid global ocean model to study both tidal and non-tidal processes, with a focus on the total water elevation. Unlike existing global ocean models, the new model resolves estuaries and rivers down to ~8m without the need for grid nesting. The model is validated with both satellite and in-situ observations for elevation, temperature and salinity. Tidal elevation solutions have a mean complex RMSE of 4.2 cm for M2 and 5.4 cm for all 5 major constituents in the deep ocean (the RMSEs for the other 4 constituents (S2, N2, K1, O1) are respectively: 2.05cm, 0.93cm, 2.08cm, 1.34cm). The non-tidal residual assessed by a tide gauge dataset (GESLA) has a mean RMSE of 7 cm. For the first time ever, we demonstrate the potential for seamless simulation, on a single mesh, from the global ocean into several estuaries along the US west coast. The model is able to accurately capture the total elevation, even at some upstream stations. The model can therefore potentially serve as the backbone in a global tide-surge and compound flooding forecasting framework.

15
20

1 Introduction

Global ocean modelling traditionally focuses on large-scale processes but is increasingly looking into the roles played by smaller-scale processes (internal gravity waves (IGW), topographic and lee waves etc) in the global energy budget (see a review by Arbic et al. 2018, hereafter A18). The state-of-the-art global ocean models now boast 1/12° or finer resolution (thus fully eddying resolving) (A18). More and more models are incorporating barotropic and/or baroclinic tides in their simulation, due to their importance in ocean mixing and global energy budget. Table 1 summarizes the major characteristics of several global tide models. Both structured- and unstructured grid (UG) based models have been successfully developed for global tides, starting from the simpler 2D barotropic model, with or without assimilation of altimetry observation. Prominent examples include highly accurate altimetry-informed TPXOv9 (Egbert and Erofeeva 2002) and FES2014 (Lyard et al. 2021) tidal databases. Recently, Pringle et al. (2021) used a 2D model to accurately simulate tide and surge concurrently with high-resolution areas of the mesh focused on hurricane landfall regions. While such a 2D model cannot simulate baroclinic effects,

25
30

it has been suggested that including a baroclinic term derived from a separate 3D ocean circulation model could be used to improve the energy spectrum of modelled sea surface heights, particularly at the low frequency end (Pringle et al., 2019). 3D baroclinic models that include concurrent simulation of eddying circulation and tidal motion are also becoming feasible (Arbic et al. 2010; Savage et al. 2017). Wang et al. (2022) recently proposed a reduced-layer (9 layers) 3D baroclinic model with nudged temperature and salinity fields for improving operational total water level forecasts.

As the global ocean constitutes a quasi-closed system, simulating global ocean processes with both tidal and non-tidal frequencies should include coastal oceans where most of the tidal energy is dissipated; collectively, the shelves dissipate about 70-75% of the tidal energy (A18). Also, as A18 mentioned, improving nearshore tides will have a back-effect that will also improve open-ocean tides, especially in tidally energetic areas (e.g., north Atlantic). Accurately accounting for this dissipation process requires high resolution nearshore to represent the complex geometry and bathymetry found therein, which represents one of the grand challenges in ocean modelling (Holt et al. 2017). In fact, we suspect that the need for several additional types of drags (wave drags etc) in the global ocean modelling in contrast to basin-scale modelling might be related to the inadequate representation of coastal oceans, as even the state-of-the-art resolution of $1/48^\circ$ (Savage et al. 2017) is hardly resolving the coastal features. The need for high resolution in the coastal ocean would inevitably strain the already high computational cost associated with global simulations, and in this regard UG models can effectively mitigate the cost. To this day, however, few UG global models resolve both tidal and non-tidal processes. Logemann et al. (2021) assessed the impact of coastal refinement on tides using an UG global ocean model but did not systematically compare their results with TPXO solution in the deeper ocean, and the reported error metrics appear unsatisfactory.

Despite the tremendous progress made, so far no 3D models for the global ocean exist that can simultaneously include estuaries and rivers without resorting to grid nesting. This is related to the very different characteristics between global and coastal oceans and estuaries (Fringer et al. 2019); chief among those differences are the drastically different spatial scales and force balances (geostrophic vs ageostrophic; weakly vs strongly forced regimes). The stability and efficiency constraints that come with resolving small-scale processes as commonly found in the coastal/estuarine regimes are formidable. For this reason, some global models have or are developing their own versions of ‘coastal model’ components that are intended to be nested into the corresponding global models to better close the energy budget (e.g., Andosov et al. 2019).

In this paper, we present a new 3D baroclinic UG model for the global ocean that incorporates both tidal and non-tidal processes and their interactions and is capable of resolving both ocean basins and the estuaries with a *single* mesh. For topics as large as this, inevitably we have to focus on a subset of interests. Here we focus on the short-term predictability (on the scale of 1 year or shorter) of the *Total Water Level* (TWL) including both tidal and non-tidal components for both large- and small-scales simultaneously, with the ultimate goal of building a global storm tide and compound flood operational model that resolves both eddying motions and some small-scale processes found near the islands and inside estuaries and rivers of interest. This represents a bold approach of ocean modelling that would completely do away with the need for open boundary conditions as in the case of regional models, and can also effectively close the last remaining gap in the energy budget in the global ocean.

65 We will first describe in Section 2 the observational datasets used in this paper, as well as the 3D UG model and its setup for the global ocean simulation. We proceed to model validation and assessment of tidal and non-tidal elevation, temperature and salinity in Section 3. In Section 4, we highlight the importance of representation of the ice shelves near Antarctica in the model bathymetry as well as feedback from shallow areas. Lastly, we demonstrate the model’s potential capability in capturing small-scale processes inside estuaries with the focus on TWL prediction in the shallows (including the challenging upstream area);
70 more detailed quantitative assessment for 3D processes would entail site specific calibration and is left for future study. We also discuss the need for closing the gaps in the theoretical understanding of the cross-scale processes. A short summary is presented in Section 5.

Table 1: List of global ocean tidal models.

Model name/references	Horizontal resolution	2D or 3D	Data assimilation	Ice component
TPXO	1/30°	2D	Y	No
FES2014	1/16° (originally variable)	2D	Y	No
ADCIRC (Pringle et al. 2021)	Variable (1.5 to 25km; 150m minimum)	2D (with special treatments for baroclinicity, internal wave drags etc)	N	No
ICON (Logemann et al. 2021)	Variable (down to 7km along coast)	3D	N	No
HYCOM (Arbic et al. 2010)	1/12.5°	3D	N	No
Schindelegger et al. (2018)	1/12°	2D	Y	No
Pickering et al. (2017)	1/8°	2D	N	N

75

2 Method

2.1 Observation

In this paper we will primarily validate the model using a satellite-derived reanalysis product OSTIA (for sea surface temperature (SST)), an altimetry-informed global tidal model TPXOv9, a global tide gauge dataset (GESLA) for sea-surface
80 height (SSH), and ARGO floats for temperature and salinity profiles. Tide gauges observation will be used sparingly as most

of these are located in complex nearshore regions that require accurate bathymetry information and more mesh work to capture that information. We will use a few tide gauge data for a target study of the US west coast in Section 4. Our ultimate goal is to build a global 3D UG model that is able to seamlessly transition from ocean basin into small creeks; to achieve this goal, however, it's essential that the model has sufficient skills for large-scale processes first. We note that the current 3D UG model
85 has been extensively applied and validated in many coastal regions so the latter does not present fundamental challenges for the model as long as a proper calibration procedure is followed.

2.2 Model description

SCHISM (schism.wiki) is an open-source community model solving 3D hydrostatic form of the Navier-Stokes equation with Boussinesq approximation (Zhang et al. 2016). Major innovative features of SCHISM include: (1) semi-implicit time stepping
90 scheme that bypasses the most stringent stability constraints (and thus allows very fine resolution of $O(1m)$ without the need to reduce the time step); (2) a highly flexible 3D gridding system, with hybrid quadrangular-triangular unstructured mesh in the horizontal dimension and localized sigma coordinates with shaved cells (LSC^2) in the vertical dimension (Zhang et al. 2015). The flexible gridding system enables powerful 'polymorphism' with a single SCHISM grid being able to seamlessly morph between full 3D, 2DH, 2DV and quasi-1D configurations (Zhang et al. 2016); (3) judicious combination of higher- and
95 lower-order schemes to ensure accurate representation of diversity of processes from creek to ocean basin scales (Zhang et al. 2016; Ye et al. 2019). These features have previously allowed a single model to be used for challenging compound flooding studies that involve coastal transition zones between hydrodynamic and hydrologic regimes, forced by ocean, atmosphere and watershed rivers (Ye et al. 2020; Zhang et al. 2020; Huang et al. 2021; Ye et al. 2021). Global compound flooding processes are not the focus of this paper.

100 The global unstructured mesh in the horizontal dimension consists of ~4.6 million nodes, ~9 million triangular elements, with nominal resolution of 10-15km in the ocean basin (Figs. 1-2), thus barely eddy resolving. The mesh resolution generally increases to ~3km at most of the coastline of the continents or islands; higher resolution of ~1-2km is applied in North America and western Pacific due to our interests in those regions. As an illustration, we have also added detailed representations of a few estuaries and rivers along the US west coast (Fig. 1(b-e)). We demonstrate in Section 4.3 the model's potential for seamless
105 cross-scale transition into nearshore and estuaries, with a minimum element size of ~8m found near a coastal highway inside the Columbia River estuary (Fig. 1e). Therefore, the mesh size spans 4 orders of contrast (from ~10km to ~8m). Overall, about 50% of elements have resolution 5km or higher (Fig. 2c).

Consistent with our main goal in this paper, we use Gebco (GEBCO Compilation Group 2019) as the main DEM sources, with a resolution of 500m for global oceans. This resolution is adequate in most of the coastal and deep oceans but is not sufficient
110 for nearshore and estuaries. As shown in Section 4.1, it's important to include the ice shelf effect on the bathymetry in the Southern Ocean, and we therefore use RTOPO for this purpose (Schaffer et al. 2016). To improve the model skill in the target estuaries in the US west coast, we have locally utilized a hierarchy of DEMs from NOAA's Coastal Relief Model (~90m resolution) and CUDEMs (1-10m resolution; CUDEM 2022) and USGS's CoNED (1-3m resolution; CoNED 2022).

The 3D model takes full advantage of the flexible vertical gridding system (LSC²; Zhang et al. 2015). The number of sigma
 115 layers varies from a maximum of 34 to 1 (i.e. 2DH configuration) at mesh nodes, and the average number of layers among all
 nodes is 32. Using 1 layer in shallow and dry areas (where the first master grid depth is less than 0.4m; cf. Zhang et al. 2015)
 greatly improved the efficiency and robustness of the model (Huang et al. 2021). The second master grid depth is set at 10m
 with 23 layers in order to have adequate resolution for the vertical stratification in shallows (cf. Zhang et al. 2015). The vertical
 high resolution is focused on the near-surface zone at the expense of the bottom in order to conserve computational cost. As a
 120 result, the near-bottom vertical layers can be as thick as 1km in the deep ocean; in other words, the logarithmic bottom boundary
 layer at deep depths is not well resolved and therefore, we apply zero friction in the deep depths. Alternatively, using a small
 friction coefficient (10^{-4}) gave similar results. To ensure adequate energy dissipation toward shallows, we use a simple depth-
 dependent bottom friction coefficient (used in the quadratic drag formulation) that linearly increases from 0 at depth 200m to
 0.0025 at 50m (i.e., 0 friction is used at depths deeper than 200m and 0.0025 is used at depths shallower than 50m, with a
 125 linear transition in between the two depths). For the sake of simplicity, no attempt has been made to optimize the friction in
 each region yet, and this is left for future work.

A main advantage of using a 3D baroclinic model is that it accounts for internal tides (ITs), whose production over open-ocean
 topographic features accounts for about 25-30% of the energy lost in the global barotropic tidal energy budget (A18). The
 results from Egbert and Ray (2003) suggest IT dissipation is an important contributor to the mixing that underpins the large-
 130 scale circulation (Munk and Wunsch 1998). According to A18, a horizontal resolution of at least $1/10^\circ$ is needed for a fully
 vigorous low-mode IT field. On the other hand, a horizontal resolution of about $1/24^\circ$ or finer is necessary for simulating a
 vigorous IGWs continuum. Therefore, the current model, because of the limited computational resources available to us, only
 resolves ITs but not IGWs. Although parameterized IGW drag formulation (e.g., Garner 2005) could be included in the model,
 its inclusion in a baroclinic model is tricky as part of the signal is already resolved (A18). Therefore, we neglect IGW drag
 135 here.

The semi-implicit model uses a non-split time step of 120s, a turbulence closure scheme of $k-kl$ of the Generic Length Scale
 Model (Umlauf and Burchard 2003), and a bi-harmonic viscosity (Zhang et al. 2016). Since no bathymetry smoothing was
 done in our mesh, the presence of very steep bottom slopes near numerous islands and ocean trenches requires additional
 momentum stabilization than provided by the bi-harmonic viscosity. Therefore, we add a Smagorinsky-type viscosity that is
 140 designed to ‘penalize’ the steep slopes. In line with the Shapiro filter-like implementation of viscosity inside SCHISM that
 works well with highly distorted UGs (Zhang et al. 2016), the Smagorinsky-type viscosity is implemented as:

$$\gamma = 0.5 \tanh(C\Delta t\Gamma) \quad (1)$$

$$\Gamma = \left[u_x^2 + v_y^2 + \frac{(u_y + v_x)^2}{2} \right]^{1/2}, \quad (2)$$

where γ is the filter strength with a maximum value of 0.5 (Zhang et al. 2016), Δt is the time step, Γ is the deformation rate,
 145 and C is a non-dimensional constant specified by the user. In the global model, we found that $C=1000$ is sufficient to suppress

most spurious noise in the horizontal velocity field. For tracer transport, the 3rd-order WENO scheme (Ye et al. 2019) is used at depths greater than 10m (whereas an upwind scheme is used for shallower depths).

Besides the Smagorinsky-type viscosity scheme, other new features of SCHISM developed in this work include: (1) self-attracting and loading (SAL) tides scheme using the depth-dependent parameterization of Stepanov and Hughes (2004); (2) efficient asynchronous I/O using dedicated ‘scribe’ cores, which is essential for large core counts and significantly improves parallel I/O and scaling.

The primary validation period used in this paper is a 120-day window from June 1, 2011 to September 29, 2011. Atmospheric forcing from ERA5 (with a resolution of 25km) was applied onto the ocean surface, including wind, air pressure, precipitation, and heat fluxes. Relaxation of temperature and salinity near the ocean surface toward their climatologic values, which is commonly utilized in many global ocean models to prevent long-term drift in decadal/century scale simulations (Ringler et al. 2013), was not applied here due to the relatively short duration of the simulation. The tidal potential with 5 constituents (M2,S2,N2,K1,O1) and self-attracting and loading (SAL) tides were included. For the harmonic analysis and comparison with TPXOv9, we used the model results from Days 20 to 60 and turned off the atmospheric forcing, as the 40-day results used were long enough to distinguish the 5 constituents used in the analysis.

The model was initialized with a dynamic flow field interpolated from the global $1/12^\circ$ (~ 10 km) HYCOM. Linear interpolation was used in the interpolation of sea-surface height (SSH), horizontal velocity, salinity, and temperature. Constant extrapolation was used in regions not covered by HYCOM. The HYCOM derived SSH and horizontal velocity represent the non-tidal component; once started, the tidal potential and SAL will initiate the tidal motion in the system. Altogether 868 largest rivers were included along the coast, with monthly mean flow information from Dai (2021). The river temperature input was set to be the ‘ambient’ flow temperature due to the lack of such information. Starting the simulation from a fully dynamically equilibrated flow field allows us to significantly reduce the time required for warming up the model. Still, the discrepancies between HYCOM and SCHISM and initiation of the tides require a short ramp-up period, estimated to be shorter than 20 days.

We remark here on a few limitations of the current model. The need for IGW drag may be reassessed in the future. Alternative implementations of SAL, e.g., interpolation from TPXO or FES should be explored. In the high latitudes, the lack of an ice model is a major gap for the model; even though we have this component inside the SCHISM system, its inclusion would significantly increase the computational cost especially under high mesh resolution. The mesh resolution in the deep ocean is barely eddy resolving; for accurate simulation of the eddy processes, finer mesh resolution would be required. Lastly, atmospheric forcing could use higher spatial and temporal resolution for more accurate simulation of coastal processes especially during storm events.

SCHISM’s good parallel scaling allows efficient simulation given adequate resources. For example, on 5600 cores of an Intel cluster (TACC’s Frontera) the model is able to run ~ 120 times faster than real time. Doubling the core count to 11200 increases this real-time ratio to ~ 223 , which translates to an excellent parallel efficiency of $\sim 95\%$.

180





Fig. 1: (a) Global unstructured mesh, with (b) successive local refinements in the US west coast, including (c,e) the Columbia River and estuary, and (d) San Francisco Bay.



185 Fig. 2: (a) Mesh resolution as measured by equivalent diameter, with (b) zoom-in near the Columbia River; the orange transect will be used to show salt intrusion path. (c) Histogram of resolution.

3 Model validation

We start the model validation with surface elevation, with respect to both tidal (Section 3.1) and non-tidal components (Sections 3.2). For 3D variables, we focus on temperature and salinity as these are the major drivers of the large-scale processes due to their contributions to the uneven oceanic mass distribution. Note that the focus of this study is on the accurate prediction of the total elevation. We follow HYCOM in using NGVD29 (geopotential) as the vertical datum, which is problematic in the assessment of either the altimetry (which is referred to a fixed geoid (Jahanmard et al. 2021)) or some tide gauges that refer to different datums (e.g. NAVD88). More rigorous assessment of the total water level is left out for future studies after a geoid-based datum becomes available, and here we focus on the tidal elevation as well as the *variability* of the non-tidal elevation (i.e., the mean biases are removed when datums do not match or are unknown). Standard error metrics are reported below, including: RMSE (Root-Mean-Square Error), MAE (Mean Absolute Error), correlation coefficient and Wilmot score (Wilmot 1981).

3.1 Co-tidal chart for M2

Globally, the M2 amphidromes correspond to the local minimum of the tidal energy in the open ocean or near islands (e.g., Taiwan, New Zealand, Madagascar etc) where the tides tend to rotate in the form of Kelvin waves (Fig. 3b). The amplitude maxima, on the other hand, are typically found near semi-enclosed basins near resonant modes (e.g., European Seas, Hudson Bay, Bay of Fundy etc), where the tidal transformation is rather complex (Fig. 3b).

A major benefit of the 3D model is that the time varying dissipation of tidal energy due to internal tides is accounted for inside the model. Therefore, with a simple specification of bottom friction (as a function of depths), the simulated M2 distribution already has a good skill as compared to the benchmark TPOXv9 tidal database (Fig. 3).

The complex RMSE for a constituent is defined as (Wang et al. 2012):

$$RMSE_j = \sqrt{\frac{1}{2} VD_j^2}, \quad (3)$$

$$VD_j^2 = [(A_o)_j \cos(\varphi_o)_j - (A_m)_j \cos(\varphi_m)_j]^2 + [(A_o)_j \sin(\varphi_o)_j - (A_m)_j \sin(\varphi_m)_j]^2, \quad (4)$$


where VD stands for vector difference, A is the amplitude, ‘m’ and ‘o’ refer to model and observation, φ is the phase, and j is the constituent index. The area averaged RMSE for a specific area Ω is then defined as:

$$\overline{RMSE}_j = \sqrt{\frac{\int_{\Omega} RMSE_j^2 d\Omega}{\int_{\Omega} d\Omega}}, \quad (5)$$

The \overline{RMSE} can be computed for a single constituent (e.g. M2) or summed up for a group of constituents (e.g., all 5 constituents) to give a single number for those constituents.

215 The averaged complex RMSE for M2 is 4.2cm for depths greater than 1km, and 14.3cm for shallower depths. The averaged total RMSE for all constituents) is 5.4cm / 16.6cm or depths greater/less than 1km. The breakdown of RMSEs for the other 4 constituents (S2, N2, K1, O1) is: 2.05cm, 0.93cm, 2.08cm, and 1.34cm for depths greater than 1km; 6.07cm, 2.60cm, 4.71cm, and 2.84cm for depths shallower than 1km. These results are slightly better than the previous best 3D model results without data assimilation (Schindelegger et al. 2018) but slightly worse than those in Pringle et al. (2021); e.g., the total RMSE from
220 their model is 3.9 cm / 17.2 cm in the deep/shallow ocean respectively. Note that there are differences in the shallow areas included in each model, which makes the numbers for the shallows less reliable than those for the deeper depths. In addition, the sensitivity results using 8 tidal constituents (M2, S2, N2, K2, K1, O1, P1, Q1) are similar (Fig. 3f); the averaged complex RMSE for M2 is essentially the same as using the 5 constituents.

Our elevation results are quite satisfactory given the fact that minimal calibration (with respect to bottom friction etc) was
225 conducted; the more complete physics as incorporated in the 3D baroclinic model reduced the amount of calibration required to achieve good tidal results compared to 2D barotropic models (e.g., Blakely et al., 2022). Our own experience with global SCHISM 2D and with regional studies (Huang et al. 2021, 2022; Ye et al. 2020) also confirmed that an elaborate calibration exercise using spatially variable frictions was necessary to improve the elevation skill in the 2D model. Compared to other global 3D models, our model seems to be able to obtain satisfactory results without the need for some elaborate drag
230 formulations described in A18; one plausible reason is that the higher resolution used in the coastal ocean has provided adequate energy dissipation with improved seafloor representation in areas with complex bathymetry and geometry.



235 **Fig. 3:** comparison of co-tidal chart for M2 between (a) SCHISM and (b) TPXOv9. The complex M2 RMSE is shown in (c); some geographic locations are labelled here also. (d) Sensitivity results without Antarctica ice shelf represented. (e) Sensitivity results with shallow areas (<50m) removed. (f) Sensitivity results using 8 constituents.

3.2 GESLA tide gauges comparison

240 The modelled sea levels are compared with observed sea levels from tide gauge stations in the Global Extreme Sea Level Analysis (GESLA) dataset (Woodworth et al., 2016). The tidal harmonic analysis is performed using the t-tide package

(Pawlowicz et al. 2002). The skills of the model to reproduce tides are assessed using the RMSE for each tidal constituent (cf. Eq. (3)), averaged over all tide gauges. Then the root sum of square (RSS) is used as an overall skill index:


$$RSS = \sqrt{\sum_{j=1}^n RMSE_j^2}, \quad (6)$$

245 where $n=5$ is the total number of constituents. The tidal skill scores are also computed from the FES2012 model, an altimetry-informed model used as a reference to evaluate our model.

The assessment of model performance to reproduce the non-tidal residual (NTR) sea level variation is also conducted. Due to the uncertainty in the vertical datums used in many gauges, NTR time series are obtained by de-meaning for the common period, and de-tiding using the t-tide package. Afterwards skill scores of RMSE and Pearson correlation coefficient are
250 computed. In order to assess the model predictive skill for extreme water levels, RMSE and correlation scores are also computed for the upper tail of the time series, i.e. values exceeding 95th percentile of the observed NTR.

Overall, a satisfactory model performance is observed in coastal areas in comparison with GESLA. The model RMSEs are less than 0.1 m at ~45% of tidal stations for M2 and less than 0.05 m at ~58% of the stations for S2 (Fig. 4). The comparison with FES2012 indicates larger error in SCHISM (+6cm in M2 RMSE and +2cm for S2 RMSE) (Table 2). It is an acceptable
255 performance given the fact that FES2012 incorporates data assimilation from altimetry data (Carrère et al., 2013). Most of the larger errors occur in areas with DEMs of large uncertainty (e.g., Canadian coasts with fjords; southern Chilean/Argentine coasts) or in areas with insufficient resolution (e.g., European Seas) (Fig. 4). The uneven error distribution may guide future priority in mesh development.


The model also accurately reproduces the NTR in coastal waters (Fig. 5). The average RMSE is only 7cm with a median value
260 of 6cm; ~88% of RMSEs are below 10cm and only 1.6% exceed 15cm, mostly at stations located in the North Sea and Northwestern Pacific Ocean. The RMSE increases for extreme conditions (NTR>95th percentile), but the model still adequately reproduces the extreme conditions with an averaged RMSE of 11cm and a median of 9cm. In addition, RMSEs are less than 15cm at ~80% of the tidal stations under extreme conditions. These results are consistent with our previous results using the 3D model (Ye et al. 2020; Huang et al. 2021, 2022), and indeed, the better skill at capturing the NTR is a major
265 advantage of 3D over 2D models, because the NTR is largely driven by the eddying motions and large-scale ocean current systems that originate from the uneven ocean mass distribution.



270 Fig. 4: Histogram and maps of scatter plots of M2 and S2 vector difference error.

Table 2. Summary of model performance to reproduce the main semidiurnal tidal component for SCHISM and FES2012 models against GESLA. The RMSE and RSS are averaged over all tide gauges.

	RMSE M2	RMSE S2	RSS
SCHISM	0.32	0.11	0.35
FES2012	0.26	0.09	0.28



275 Fig. 5. Maps and histogram of RMSEs for non-tidal residual (NTR). ~88% of RMSEs are below 10cm.

3.3 SST

280 The quality of the simulated sea-surface temperature (SST) is assessed against a reanalysis product, OSTIA (Good et al. 2020), which blends different satellites with in-situ data into regular 0.05° resolution sea surface temperature estimation on a daily basis. This dataset provides global SST with an overall analysis error of $\sim 0.4^\circ\text{C}$. Most large errors are distributed in the Arctic region which can reach 3.6°C maximum. In order to isolate the effects from the Arctic, the analysis is done for the global domain with and without the Arctic region (latitude $> 60^\circ\text{N}$). The OSTIA analysis error excluding the Arctic is reduced to 0.3°C on average.

285 The simulated SST is mostly similar to OSTIA (Fig. 6). In particular, the model is able to capture major boundary currents (Kuroshio, Gulf Stream etc) as well as equatorial instabilities. A closer look at the comparison reveals larger warm biases in the Arctic region (Fig. 7), which is not surprising because we did not include the ice component in our model. Excluding the Arctic, the averaged MAE stays much lower at $\sim 0.8^\circ\text{C}$ throughout the simulation period (Fig. 7b). Besides the high latitude regions in the northern hemisphere, relatively higher biases are also found near the boundary currents and equatorial instability regions (Fig. 7a).




Fig. 6: Comparison of SST between (a) OSTIA; (b) SCHISM at the end of 120-day simulation (Sept 29, 2011).




Fig. 7: SST from the 120-day run from June 1, 2011. (a) SST Bias; (b) MAE.

3.4 SSS

295 Compared to SST observations, sea-surface salinity (SSS) observations are scarcer. Although NASA's Aquarius satellite
missions did cover the simulation period, the data may be too coarse for our purpose. Therefore, we use HYCOM (which has
assimilated profile data) as a reference solution in our comparison. The two models are largely comparable, including major
fronts and instabilities near equator, the freshwater plumes and boundary currents, and intrusion from north Atlantic into Arctic
etc (Fig. 8). Freshwater plumes from Amazon and other large rivers appear to be larger in SCHISM. Due to the absence of an
300 ice component in our model, there are also some differences in the Arctic Ocean. Overall, the modelled SSS is satisfactory.




Fig. 8: Comparison of SSS between (a) 1/12° HYCOM and (b) SCHISM at the end of 120-day simulation (Sept 29, 2011).

3.5 ARGO profiles

To assess the model skill in capturing the vertical structures of temperature and salinity, we use all ARGO data in each ocean basins except the Arctic (Fig. 9).
305

Total number of ARGO profiles is around 330 average per day in our simulation period. The model results are first interpolated from surface to 2000m to match the measuring range of ARGO. We follow the original ARGO data structure to divide our analysis into 3 different basins (Atlantic, Pacific and Indian Oceans), each including parts of the Southern Ocean. Due to the relatively small number of profiles (less than 30 per day) available near the surface (< 6m) in all basins, 0-6m data are excluded to produce more reliable statistics.
310

Overall, the modelled temperature and salinity profiles are satisfactory, with a MAE of $\sim 0.6^{\circ}\text{C}$ for temperature and $\sim 0.2\text{PSU}$ for salinity (Fig. 10). Of all ocean basins, the Pacific has the smallest biases (Fig. 11). The simulated temperature in all basins tends to have a cold bias ($\sim -0.4^{\circ}\text{C}$) near the surface, and biases below 200m depth are smaller. The simulated salinity in all basins has a positive bias below 1400m depth (Fig. 11), which is inherited from the initial condition. The good skills for temperature and salinity profiles are also confirmed by the high correlation scores that exceed 0.9 most of the time (Fig. 12).
315




Fig. 9: Locations of ARGO profiles used in this study (Period: Aug/1 ~ Sep/28 2011, 20823 profiles in total)




Fig. 10: Averaged daily MAEs from all ARGO profiles for (a) temperature and (b) salinity.





Fig. 11: Mean biases in each ocean basin. (a-c): temperature biases in Atlantic, Pacific and Indian Oceans respectively; (d-f): salinity biases. Each plot is divided into two parts to clearly show the larger biases in the top 200m depth.



325 Fig. 12: Averaged daily correlation coefficients for all ARGO profiles.

4. Sensitivity tests and discussion

Calibrating a 3D baroclinic model like ours can be an expensive exercise, especially for 3D variables such as temperature, salinity and velocity. However, since the focus of this paper is on the water surface elevation, we found through various sensitivity tests that the elevation results are sensitive to only a few parameters, including the bathymetry and bottom friction, because the thermo-steric contribution to elevation has been accounted for in the model. Compared to 2D models, the more complete physics embedded inside 3D models greatly simplifies the calibration process, e.g. the time-varying IT induced dissipation is already included in the model. This finding is consistent with our previous finding for a sub-domain of the US east coast and Gulf of Mexico coast (Huang et al. 2022). In this section we will show the sensitivity of the simulated elevations to the representation of the ice shelf effects in the Southern Ocean as well as exclusion of shallow areas.

A main novelty of our 3D UG model is its capability to seamlessly traverse scales from global oceans into very localized scales as found in the estuaries and rivers. To the best of our knowledge, this capability has not been demonstrated before without resorting to grid nesting. Therefore, our model represents a major advancement in efficiently simulating global and local scales in a *single* UG model, thus allowing the interaction and connection among scales to be fully explored in our model. We demonstrate this potential here using two estuaries in the US west coast as example.

4.1 Southern Ocean: ice shelf effect

Bathymetry is known to play a pivotal role in the tidal and non-tidal processes (Ye et al. 2018; Huang et al. 2022). As far as the global tide is concerned, one particularly important region is the Southern Ocean, which has an extensive distribution of ice shelves along the Antarctica coast. The existence of ice shelves effectively changes the local bathymetry, which affects tidal propagation locally and beyond (Blakely et al., 2022). Without accounting for those shelves, an erroneous amphidrome appears between Drake Passage and Ross Sea (Fig. 3d vs 3a), and the amphidrome just east of the Ross Sea (Fig. 3a) is displaced westward (Fig. 3d). Other differences are also visible south of Australia and in the Weddell Sea (Fig. 3d vs 3a).

4.2 Feedback from shallow areas

To demonstrate the feedback from the shallow areas to the deeper ocean and its impact on global tidal energy dissipation, we removed all shallow areas with depth less than 50m. Even though the shallow areas only account for ~10% of the total surface area (and even less in terms of volume), excluding the shallows significantly degrades the model skill in the deep ocean (e.g., the total RMSE is more than doubled from 5.4 cm to 11 cm for depths greater than 1km). The lack of dissipation by the shallows is seen in the overestimated M2 amplitudes, especially in tidally energetic areas (e.g., European Seas). The findings here are consistent with previous estimates; e.g. A18 estimated that collectively, the shelves dissipate about 70-75% of the tidal energy. Without the shallows, most of the tidal energy would be reflected back into the deep ocean.

As a proof of concept, we illustrate the model's potential in traversing from large oceanic to small estuarine scales using a few estuaries in the US west coast as an example but leave detailed calibration and validation to future studies. Obviously, the calibration process will be expensive given the large mesh size used here. However, we show that with local mesh refinement and minimal calibration done in the current 3D model, the model is able to capture some small-scale processes.

360 The Columbia River and San Francisco Bay (SFB) are the two largest estuaries in the US west coast (excluding Alaska), and are characterized as meso-tidal systems. The river discharges vary greatly between the two systems and over time. The Columbia River has a long-term mean flow of $2.5\text{--}11 \times 10^3 \text{ m}^3/\text{s}$ over a typical year (Bottom et al., 2005). SFB receives most of the freshwater inputs from the north Bay which is connected to the Sacramento-San Joaquin Delta, and the net Delta outflow is smaller than the Columbia River (from $\sim 500 \text{ m}^3/\text{s}$ to $2000 \text{ m}^3/\text{s}$). Both systems exhibit similar seasonality in the river flow, 365 with the lowest flow occurring in late summer and highest flow during spring freshets. Due to the specificity of the forcings experienced by the two systems, the salt intrusion processes are quite different. The Columbia River estuary shows a strong spring-neap variation in the stratification, and occasionally exhibits salt wedge conditions (Jay and Smith 1990). In the MacCready-Geyer estuarine classification diagram, the Columbia River Estuary (CORIE) is classified as a 'time dependent salt wedge' system (Geyer and MacCready 2004). The shorter shelf width near the CORIE makes it more susceptible to the 370 prevailing coastal upwelling that is more common along the Oregon-Washington shelf. Previous modelling studies (Karna and Baptista 2016) indicate that the Columbia River processes in particular are extremely challenging for numerical models. The SFB, on the other hand, represents a typical partially mixed estuary (Geyer and MacCready 2004).


The total elevations at three NOAA tide gauges in SFB and one gauge in CORIE are assessed in Fig. 13. Fortuitously, the vertical datum used in the model, NGVD29 (inherited from HYCOM) is close to the Local MSL used at these 3 gauges, and 375 therefore, no adjustment of the vertical datum is necessary. For other coastal regions (e.g., US east coast), the datum differences can be substantial, which calls for a geoid-based datum for regional and global tidal models (Jahanmard et al. 2021). With particular attention paid to the mesh representation near these two estuaries, the model is able to accurately capture the tidal and non-tidal elevations in the two systems, as shown in Fig. 13, with RMSEs for the total elevations between 8.3cm (at Richmond) and 18.7cm (at Astoria), and the correlation coefficients and Wilmot scores all exceeding 0.95. As mentioned in 380 Section 3.2, one major advantage of 3D models is their ability to better capture the NTR and thus TWL. Our experience suggests that with similar mesh refinement procedure, reliable DEMs and some calibration with respect to the bottom drag coefficients, similar elevation skills can be obtained for other estuarine systems.

The Columbia River plume is a major coastal feature in the region and can extend 100s of kilometres offshore (Baptista et al. 2005) and has a major impact on the ecosystem (Burla et al. 2010). The plume is highly dynamic and mostly wind driven but 385 modulated by tides and river discharge (Burla et al. 2010). Fig. 14 shows a 'canonical' view of the plume when the wind forcing is weak or relaxed. The combination of the Coriolis and inertial forces turns the plume northward with a coastally trapped jet in the north (in the form of Kelvin waves) and visible recirculation inside the freshwater bulge (Garcia-Berdeal et

al. 2002). The model's ability to qualitatively capture the plume shape and extent is an important first step for further calibration.


390 .

Although a more careful validation against observation (including the vertical salinity profiles) is necessary to ascertain the model skill in capturing the smaller-scale 3D processes in estuaries, which would inevitably involve site-specific parameterization and calibration procedure, the preliminary results shown here are very promising and offer the potential to finally close the gap in simulating the global ocean-estuary-river-lake continuum. Note that the model does allow specification of spatially variable parameterizations such as bottom drag and horizontal mixing scheme, which will be necessary in future calibration process. On the other hand, the physical justifications of these choices, in the context of cross-scale processes from global ocean to estuaries and rivers, warrant further research; e.g., traditionally different horizontal mixing schemes have been used in the two regimes (Fringer et al. 2019).



400

Fig. 13: Comparison of elevation at 4 tide gauges in Columbia River and SF Bay, USA. See Fig. 1(cd) for gauge location. The RMSEs are 18.7, 9.4, 8.3 cm and 15.4 cm; Correlation Coefficients are 0.98, 0.98, 0.98, and 0.90; Wilmot skills are 0.98, 0.99, 0.99, and 0.95 respectively. Both model and data use NGVD29 as the vertical datum.



405 Fig. 14: Surface salinity plume near the Columbia River during a wind relaxation period.

5. Conclusion

We have developed a new 3D unstructured-grid (UG) model (SCHISM) for simulating the global ocean together with coastal ocean and even estuaries in a single mesh, with high resolution applied in the latter. The simulated total elevation (including
410 both tidal and non-tidal components), temperature and salinity have been validated against satellite and in-situ observation data. The simulated tide showed good skill, with a mean complex RMSE of 4.2cm for M2 and 5.4cm for the all major constituents in depths greater than 1km. The non-tidal residual assessed by the global tide gauge dataset (GESLA) had a mean RMSE of 7cm. The mean MAE for SST excluding Arctic is $\sim 0.8^{\circ}\text{C}$.

For the first time ever, we demonstrated the potential for seamless simulation, without the need for grid nesting, from the
415 global ocean into a few estuaries in the US west coast in very high resolution. The model was able to accurately capture the
total elevation, and qualitatively capture the challenging salinity plume dynamics in the Columbia River.

Even though the 3D model is more expensive than 2D models, the improved accuracy and ease of calibration for the total
water levels justify its cost; this advantage is in addition to the obvious benefits of being able to predict other 3D variables
(velocity, temperature and salinity etc). With adequate computational resources, the model can effectively serve as the engine
420 of a global tide-surge and even compound flooding forecasting framework. More meshing work and calibration will be required
to further improve its accuracy in specific regions and the UG nature of the model greatly simplifies the required work. In
addition, a global tidal model would greatly benefit from transitioning from a tidal datum based to geoid-based model to allow
more accurate simulation for the total elevations.

Author contribution

425 All authors contributed to writing of the paper. YJZ and WP worked on the model setup. TF-M did analyses of the GESLA
and altimetry. HY and LC did the analyses for satellite SST and ARGO. SM advised and managed the project.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgement

430 This study was partially supported by NOAA Water Initiative (NA20NOS4200205). WP was supported by the U.S. National
Oceanic and Atmospheric Administration (NOAA) through a Virginia Institute of Marine Science subaward under a Strategic
Partnership Project agreement A21162 to Argonne National Laboratory through U.S. Department of Energy contract DE-
AC02-06CH11357. Simulations used in this paper were conducted using the following computational facilities:

(1) William & Mary Research Computing for providing computational resources and/or technical support (URL:
435 <https://www.wm.edu/it/rc>)

(2) the Extreme Science and Engineering Discovery Environment (XSEDE), which is supported by National Science
Foundation grant number OCI-1053575;

(3) Texas Advanced Computing Center (TACC; Grant EAR21010), The University of Texas at Austin.

The ARGO data used are from the ARGO project (<https://argo.ucsd.edu/data/acknowledging-argo/>).

440 Code availability

Source code of SCHISM v5.10.0 can be accessed at <https://github.com/schism-dev> or <https://doi.org/10.5281/zenodo.6851013>. An example of the 3D model setup can be found at: <http://ccrm.vims.edu/yinglong/TMP/Glb/RUN06a/>

References

- 445 Accad, Y., and C. L. Pekeris: Solution of the tidal equations for the M2 and S2 tides in the world oceans from a knowledge of the tidal potential alone, *Philos. Trans. R. Soc. London*, 290, 235–266, 1978.
- Androsov, A., Fofonova, V., Kuznetsov, I., Danilov, S., Rakowsky, N., Harig, S., Brix, H., and Wiltshire, K. H.: FESOM-C v.2: coastal dynamics on hybrid unstructured meshes, *Geosci. Model Dev.*, 12, 1009–1028, <https://doi.org/10.5194/gmd-12-1009-2019>, 2019.
- 450 Arbic, B. K., Alford, M. H., Ansong, J. K., Buijsman, M. C., Ciotti, R. B., Farrar, J., Hallberg, R. W., Henze, C. E., Hill, C. N., Luecke, C. A., Menemenlis, D., Metzger, E., Müller, M., Nelson, A. D., Nelson, B. C., Ngodock, H. E., Ponte, R. M., Richman, J. G., Savage, A. C., Scott, R. B., Shriver, J. F., Simmons, H. L., Souopgui, I., Timko, P. G., Wallcraft, A. J., Zamudio, L., Zhao, Z.: A Primer on Global Internal Tide and Internal Gravity Wave Continuum Modeling in HYCOM and MITgcm. *New Frontiers In Operational Oceanography*, 307-391, doi:10.17125/gov2018.ch13., 2018.
- 455 Arbic, B.K., A.J. Wallcraft, and E.J. Metzger: Concurrent simulation of the eddying general circulation and tides in a global ocean model. *Ocean Modelling* 32, 175-187, doi:10.1016/j.ocemod.2010.01.007, 2010.
- Baptista, A.M., Zhang, Y., Chawla, A., Zulauf, M.A., Seaton, C., Myers, E.P., Kindle, J., Wilkin, M., Burla, M. and Turner, P.J.: A cross-scale model for 3D baroclinic circulation in estuary-plume-shelf systems: II. Application to the Columbia River. *Continental Shelf Research*, 25, 935-972, 2005.
- 460 Baptista, A. M.: CORIE: the first decade of a coastal-margin collaborative observatory, *Oceans 2006, MTS/ IEEE*, Boston, MA, 2006.
- Blakely, C. P., Ling, G., Pringle, W. J., Contreras, M. T., Wirasaet, D., Westerink, J. J., Moghimi, S., Seroka, G., Shi, L., Myers, E., Owensby, M., & Massey, C.: Dissipation and Bathymetric Sensitivities in an Unstructured Mesh Global Tidal Model. *Journal of Geophysical Research: Oceans*, 127(5), e2021JC018178. <https://doi.org/10.1029/2021JC018178>, 2022.
- 465 Bottom, D. L., C. A. Simenstad, J. Burke, A. M. Baptista, D. A. Jay, K. K. Jones, E. Casillas, and M. H. Schiewe: Salmon at river's end: The role of the estuary in the decline and recovery of Columbia River salmon. NOAA Tech. Memo., NMFS-NWFSC-68, 246 pp., U.S. Dept. of Commerce, 2005.
- Burla, M., Baptista, A.M. Zhang, Y., and Frolov, S.: Seasonal and inter-annual variability of the Columbia River plume: a perspective enabled by multi-year simulation databases. *Journal of Geophysical Research: special issue on NSF RISE*
- 470 project, 115, C00B16, 2010.

- Carrere, L., Lyard, F., Cancet, M., Guillot, A., & Roblou, L.: FES 2012: a new global tidal model taking advantage of nearly 20 years of altimetry. *20 Years of Progress in Radar Altimetry*, 710, 13, 2013.
- CoNED: Coastal National Elevation Database. <https://www.usgs.gov/publications/coastal-national-elevation-database>, 2022.
- CUDEM: Continuously Updated Digital Elevation Model,
475 https://coast.noaa.gov/htdata/raster2/elevation/NCEI_ninth_Topobathy_2014_8483/, 2022.
- Dai, A.: Hydroclimatic trends during 1950–2018 over global land, *Climate Dynamics*, DOI: 10.1007/s00382-021-05684-1, 2021.
- Egbert, G.D., and Erofeeva, S.Y.: Efficient inverse modeling of barotropic ocean tides. *Journal of Atmospheric and Oceanic Technology* 19.2 (2002): 183-204, 2002.
- 480 Egbert, R.D., and Ray, R.D.: Semi-diurnal and diurnal tidal dissipation from TOPEX/Poseidon altimetry. *Geophysical Research Letters* 30, 1907, doi:10.1029/2003GL017676, 2003.
- Fringer, O.B., Dawson, C.N., He, R., Ralston, D.K., and Zhang, Y.: The future of coastal and estuarine modeling: Findings from a workshop, *Ocean Modelling*, 143, 101458. <https://doi.org/10.1016/j.ocemod.2019.101458>, 2019.
- Garcia-Berdeal, I., Hickey, B.M., Kawase, M.: Influence of wind stress and ambient flow on a high discharge river plume. *J. Geophys. Res.* 107, 3130, 2002.
485
- Garner, S.T.: A topographic drag closure built on an analytical base flux. *Journal of the Atmospheric Sciences* 62, 2302-2315, doi:10.1175/JAS3496.1, 2005.
- GEBCO Compilation Group: GEBCO 2019 Grid, <https://doi.org/10.5285/836f016a-33be-6ddc-e053-6c86abc0788e>, 2019.
- Geyer, W.R. and MacCready, P.: The Estuarine Circulation, *Annual Review of Fluid Mechanics* 2004 46:1, 175-197, 2004.
- 490 Good, S.; Fiedler, E.; Mao, C.; Martin, M.J.; Maycock, A.; Reid, R.; Roberts-Jones, J.; Searle, T.; Waters, J.; While, J.; Worsfold, M.: The Current Configuration of the OSTIA System for Operational Production of Foundation Sea Surface Temperature and Ice Concentration Analyses. *Remote Sens.*, 12, 720, doi:10.3390/rs12040720, 2020.
- Huang, W., Ye, F., Zhang, Y., Park, K., Du, J., Moghimi, S., Myers, E., Pe'eri, S., Calzada, J.R., Yu, H.C., Nunez, K., and Liu, Z.: Compounding factors for extreme flooding around Galveston Bay during Hurricane Harvey, *Ocean Modelling*, 158,
495 101735. link <https://doi.org/10.1016/j.ocemod.2020.101735>, 2021.
- Huang, W., Zhang, Y., Wang, Z., Ye, F., Moghimi, S., Myers, E., Yu, H.: Tidal simulation revisited, *Ocean Dynamics*, 72:187-205. <https://doi.org/10.1007/s10236-022-01498-9>, 2022.
- Holt, J., Hyder, P., Ashworth, M., Harle, J., Hewitt, H. T., Liu, H., New, A. L., Pickles, S., Porter, A., Popova, E., Allen, J. I., Siddorn, J., and Wood, R.: Prospects for improving the representation of coastal and shelf seas in global ocean models,
500 *Geosci. Model Dev.*, 10, 499–523, <https://doi.org/10.5194/gmd-10-499-2017>, 2017.
- Jahanmard, V., Delpeche-Ellmann, N., and Ellmann, A.: Realistic dynamic topography through coupling geoid and hydrodynamic models of the Baltic Sea. *Continental Shelf Research* 222, p. 104421. <https://doi.org/10.1016/j.csr.2021.104421>, 2021.

- Jay, D.A., and Smith, J.D.: Circulation, density distribution and nea-spring transitions in the Columbia river estuary. *Prog. Oceanogr.* 25, 81–112, 1990.
- 505 Karna, T. and Baptista, A.M.: Evaluation of a long-term hindcast simulation for the Columbia River estuary, *Ocean Modelling*, 99, 1-14, 2016.
- Logemann, K., Linardakis, L., Korn, P.: Global tide simulations with ICON-O: testing the model performance on highly irregular meshes. *Ocean Dynamics* 71, 43–57. <https://doi.org/10.1007/s10236-020-01428-7>, 2021.
- 510 Lyard, F. H., Allain, D. J., Cancet, M., Carrère, L., and Picot, N.: FES2014 global ocean tide atlas: design and performance, *Ocean Sci.*, 17, 615–649, <https://doi.org/10.5194/os-17-615>, 2021.
- Munk, W.H., and Wunsch, C.: Abyssal recipes II: energetics of tidal and wind mixing. *Deep Sea Research I* 45, 1977–2010, doi:10.1016/S0967-0637(98)00070-3, 1998.
- Pawlowicz, R., Beardsley, R., Lentz, S.: Classical tidal harmonic analysis including error estimates in MATLAB using T_TIDE, *Computers & Geosciences*, 28(8), 929-937, 2002.
- 515 Pickering, M.D., Horsburgh, K.J., Blundell, J.R., Hirschi, J.J.-M. Nicholls, R.J., Verlaan, M., & Wells, N.C.: The impact of future sea-level rise on the global tides, *Continental Shelf Research*, 142, 50-68, 2017.
- Pringle, W. J., Gonzalez-lopez, J., Joyce, B., Westerink, J. J., & van der Westhuysen, A. J.: Baroclinic Coupling Improves Depth-Integrated Modeling of Coastal Sea Level Variations around Puerto Rico and the U.S. Virgin Islands. *Journal of Geophysical Research: Oceans*, 124(3), 2196–2217. <https://doi.org/10.1029/2018JC014682>, 2019.
- 520 Pringle, W. J., Wirasaet, D., Roberts, K. J., and Westerink, J. J.: Global storm tide modeling with ADCIRC v55: unstructured mesh design and performance, *Geosci. Model Dev.*, 14, 1125–1145, <https://doi.org/10.5194/gmd-14-1125-2021>, 2021.
- Ringler, T., Petersen, M., Higdon, R.L., Jacobsen, D., Jones, P.W., Maltrud, M.: A multi-resolution approach to global ocean modeling. *Ocean Modell.* 69, 211– 232. <http://dx.doi.org/10.1016/j.ocemod.2013.04.010>, 2013.
- 525 Savage, A.C., B.K. Arbic, M.H. Alford, J.K. Ansong, J.T. Farrar, D. Menemenlis, A.K. O’Rourke, J.G., Richman, J.F. Shriver, G. Voet, A.J. Wallcraft, and L. Zamudio: Spectral decomposition of internal gravity wave sea surface height in global models. *Journal of Geophysical Research Oceans* 122, 7803-7821, doi:10.1002/2017JC013009, 2017.
- Schaffer, J., Timmermann, R., Erik Arndt, J., Savstrup Kristensen, S., Mayer, C., Morlighem, M., and Steinhage, D.: A global, high-resolution data set of ice sheet topography, cavity geometry, and ocean bathymetry, *Earth System Science Data*, 8, 530 543–557, <https://doi.org/10.5194/essd-8-543-2016>, 2016.
- Schindelegger, M., Green, J. A., Wilmes, S. B., and Haigh, I. D.: Can We Model the Effect of Observed Sea Level Rise on Tides? *Journal of Geophysical Research: Oceans*, 123, 4593–4609, <https://doi.org/10.1029/2018JC013959>, 2018.
- Stepanov, V. N., & Hughes, C. W.: Parameterization of ocean self-attraction and loading in numerical models of the ocean circulation. *Journal of Geophysical Research C: Oceans*, 109(3), 1–11. <https://doi.org/10.1029/2003jc002034>, 2004.
- 535 Umlauf, L., Burchard, H.: A generic length-scale equation for geophysical turbulence models. *J. Mar. Res.* 6, 235–265, 2003.
- Wang, X., Chao, Y., Shum, C. K., Yi, Y., and Fok, H. S.: Comparison of two methods to assess ocean tide models, *Journal of Atmospheric and Oceanic Technology*, 29, 1159–1167, 2012.

- Wang, P., Bernier, N. B., & Thompson, K. R.: Adding baroclinicity to a global operational model for forecasting total water level: Approach and impact. *Ocean Modelling*, 174, 102031. <https://doi.org/10.1016/j.ocemod.2022.102031>, 2022.
- 540 Willmott, C.J.: On the validation of models, *Physical Geography*, 2, 184-194, 1981.
- Woodworth, P. L., Hunter, J. R., Marcos, M., Caldwell, P., Menéndez, M., & Haigh, I.: Towards a global higher-frequency sea level dataset. *Geoscience Data Journal*, 3(2), 50-59, 2016.
- Ye, F., Zhang, Y., Wang, H., Friedrichs, M.A.M., Irby, I.D., Ateljevich, E., Valle-Levinson, A., Wang, Z., Huang, H., Shen, J., Du, J.: A 3D unstructured-grid model for Chesapeake Bay: importance of bathymetry, *Ocean Modelling*, 127, 16-39, 545 2018.
- Ye, F., Zhang, Y., He, R., Wang, Z., Wang, H.V., and Du, J.: Third-order WENO transport scheme for simulating the baroclinic eddying ocean on an unstructured grid, *Ocean Modelling*, 143, 101466. <https://doi.org/10.1016/j.ocemod.2019.101466>, 2019.
- Ye, F., Zhang, Y., Yu, H., Sun, W., Moghimi, S., Myers, E.P., Nunez, K., Zhang, R., Wang, H.V., Roland, A., Martins, K., 550 Bertin, X., Du, J., and Liu, Z.: Simulating storm surge and compound flooding events with a creek-to-ocean model: importance of baroclinic effects, *Ocean Modelling*, 145. <https://doi.org/10.1016/j.ocemod.2019.101526>, 2020.
- Ye, F. and Huang, W. and Zhang, Y. J. and Moghimi, S. and Myers, E. and Pe'eri, S. and Yu, H.-C.: A cross-scale study for compound flooding processes during Hurricane Florence, 21, 1703-1719, *Natural Hazards and Earth System Sciences*, <https://nhess.copernicus.org/articles/21/1703/2021>, 2021.
- 555 Zhang, Y. and Baptista, A.M.: SELFE: A semi-implicit Eulerian-Lagrangian finite-element model for cross-scale ocean circulation, *Ocean Modelling*, 21(3-4), 71-96, 2008.
- Zhang, Y., Ateljevich, E., Yu, H.-C., Wu, C.-H., and Yu, J.C.S.: A new vertical coordinate system for a 3D unstructured-grid model, *Ocean Modelling*, 85, 16-31, 2015.
- Zhang, Y., Ye, F., Stanev, E.V., Grashorn, S.: Seamless cross-scale modeling with SCHISM, *Ocean Modelling*, 102, 64-81. 560 doi:10.1016/j.ocemod.2016.05.002, 2016.
- Zhang, Y., Ye, F., Yu, H., Sun, W., Moghimi, S., Myers, E.P., Nunez, K., Zhang, R., Wang, H.V., Roland, A., Du, J., and Liu, Z.: Simulating compound flooding events in a hurricane, *Ocean Dynamics*, <https://doi.org/10.1007/s10236-020-01351-x>, 2020.