

The CMCC Decadal Prediction System

Dario Nicoli¹, Alessio Bellucci^{1,a}, Paolo Ruggieri^{1,b}, Panos Athanasiadis¹, Stefano Materia¹, Daniele Peano¹, Giusy Fedele¹ and Silvio Gualdi¹

¹Fondazione Centro Euro-Mediterraneo sui Cambiamenti Climatici, Bologna, 40127, Italy

^anow at: Consiglio Nazionale delle Ricerche, Istituto di Scienze dell'Atmosfera e del Clima (CNR-ISAC), Bologna, 40129, Italy

^bnow at: Department of Physics and Astronomy, University of Bologna, Bologna, 40126, Italy.

Correspondence to: Dario Nicoli (dario.nicoli@cmcc.it)

- SUPPLEMENTARY MATERIAL -

Supplementary Figures:




Fig S1: MSLP ACC for lead years 1-9. Stippling denotes points where 95% statistical significance is not reached, according to a one-tailed t test. The actual number of degrees of freedom has been computed following Bretherton et al., 1999. Green lines represent the longitudinal domain considered for the NAO index computation.




Fig S2: Detrended near-surface temperature ACC for lead year 4-10. Stippling denotes points where 95% statistical significance is not reached, according to a one-tailed t test. The actual number of degrees of freedom has been computed following Bretherton et al., 1999.