

The tidal effects in the Finite-volumE Sea ice–Ocean Model (FESOM2.1): a comparison between parameterised tidal mixing and explicit tidal forcing

Pengyang Song¹, Dmitry Sidorenko¹, Patrick Scholz¹, Maik Thomas^{2,3}, and Gerrit Lohmann^{1,4}

¹Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research (AWI), Bremerhaven, Germany

²Helmholtz Centre Potsdam–GFZ German Research Centre for Geosciences, Potsdam, Germany

³Institute of Meteorology, Freie Universität Berlin, Berlin, Germany

⁴MARUM–Centre for Marine Environmental Sciences, University of Bremen, Bremen, Germany

Correspondence: Pengyang Song (pengyang.song@awi.de)

Abstract. Tides are proved to have a significant effect on the ocean and climate. Previous modelling research either adds a tidal mixing parameterisation or an explicit tidal forcing to the ocean models. However, no research compares the two approaches in the same framework. Here we implement both schemes into a general ocean circulation model and assess both methods by comparing the results. The aspects for comparison involve hydrography, sea ice, **mixed layer depth**, Meridional Overturning Circulation (MOC), vertical diffusivity, barotropic streamfunction and energy diagnostics. We conclude that although the mesh resolution is poor in resolving internal tides in most mid-latitude and shelf-break areas, explicit tidal forcing still shows stronger tidal mixing at the Kuril–Aleutian Ridge and the Indonesian Archipelago than the tidal mixing parameterisation. Beyond that, the explicit tidal forcing method leads to a stronger upper cell of the Atlantic MOC by enhancing the Pacific MOC and the Indonesian Throughflow. Meanwhile, the tidal mixing parameterisation leads to a stronger lower cell of the Atlantic MOC due to the tidal mixing in deep oceans. Both methods maintain the Antarctic Circumpolar Current at a higher level than the control run by increasing the meridional density gradient **but with different mechanisms**. We also show several phenomena that are not considered in the tidal mixing parameterisation, for example, the changing of energy budgets in the ocean system, the bottom drag induced mixing on the continental shelves, and the sea ice transport by tidal motions. Due to the limit of computational capacity, an internal-tide-resolving simulation is not feasible for climate studies. However, a high-resolution short-term tidal simulation is still required to improve parameters and parameterisation schemes in climate studies.

1 Introduction

Based on Sandström’s theorem, the ocean is proved not to be a heat engine (e.g., Huang, 2009). That is to mean, the global Thermohaline Circulation (THC) is not driven by the heat difference between the equator and the poles. Lot has been discussed on what is driving the global THC. Munk and Wunsch (1998) concluded that tides and winds maintain the global THC and ocean stratification. They also estimated the global mean vertical diffusivity (the same as diapycnal diffusivity in the whole context) as $10^{-4} \text{ m}^2 \text{ s}^{-1}$, which mismatches $10^{-5} \text{ m}^2 \text{ s}^{-1}$ from observational results in most parts of the global ocean. Thus, some areas with large vertical diffusivities are expected to contribute to the global strength of upwelling. Observational results

(Polzin et al., 1997; Ledwell et al., 2000) present large vertical diffusivities at near-bottom rough topographies in the Mid-Atlantic Ridge, which is believed to be generated by tide–topography interactions. These observational results indicate the significance of tide-induced mixing effects from a climatic point of view.

Tide-induced mixing has long been considered in modelling research. Plenty of research on tuning vertical diffusivity parameters has been conducted to reduce model biases. By testing different vertical diffusivity parameters in an ocean general circulation model (OGCM), Bryan (1987) found that ocean circulation is sensitive to vertical diffusivity. As vertical diffusivity increases, the thickness of ocean thermocline, the strength of meridional overturning circulation (MOC), and poleward heat transport increase. After the bottom-enhanced structure of vertical diffusivities is revealed from the observations (Polzin et al., 1997), the vertical diffusivity in ocean models is now considered as a vertical profile and not anymore as a constant parameter. Tsujino et al. (2000) discussed the influence of different vertical diffusivity profiles on the global ocean circulations, concluding that different profile shapes lead to different strengths of Pacific MOC (PMOC). However, most early research does not consider the horizontal inhomogeneity of tide-induced mixing in the global ocean. Since the raising of a tide-induced mixing parameterisation (St. Laurent et al., 2002; Simmons et al., 2004), a better treatment that considers both vertical variation and horizontal inhomogeneity of the vertical diffusivity is widely used in OGCMs. It should be noted here that the horizontal distribution of the enhanced vertical diffusivities is related to the baroclinic tide conversion rate, which is provided by an analytical or numerical tidal model (e.g., Jayne, 2009). Thus the horizontal distribution is closer to the real ocean. Following the Community Ocean Vertical Mixing (Griffies et al., 2015, CVMIX) library, hereinafter, we name this parameterisation as CVMIX_TIDAL. In the following research, CVMIX_TIDAL, along with parameterisations with similar formulas (e.g., Friedrich et al., 2011; Tatebe et al., 2018), has been adopted in OGCMs to reveal the tidal effect on the ocean and climate. It has been proved that the consideration of tide-induced mixing may have an impact on aspects related to the global THC, for example, the Atlantic MOC (AMOC) (Simmons et al., 2004; Jayne, 2009), the strength and deepness of Antarctic Circumpolar Current (ACC) (Saenko and Merryfield, 2005), the Southern Ocean sea ice, and the Circumpolar Deep Water formation (Tatebe et al., 2018), the ventilation of the North Pacific Intermediate Water and the deep North Pacific circulation (Oka and Niwa, 2013), and the ocean biogeochemistry (Friedrich et al., 2011). Lee et al. (2006) also implemented bottom-enhanced vertical diffusivities in an OGCM. However, the main difference from above is that they enhanced vertical diffusivities on continental shelves, parameterising the tidal shear caused by the bottom drag in shallow areas. Their results show that the mixing caused by tidal shear on continental shelves improves the modelled hydrography at coastal areas and adjacent marginal seas.

Instead of adding parameterisation schemes to OGCMs, another way to consider the tidal effect is to add an explicit tidal forcing to the momentum equations. Global tides have been added to some OGCMs (e.g., Thomas et al., 2001; Shriver et al., 2012; Arbic et al., 2018; Logemann et al., 2021), but little research has been conducted from a climatic point of view. By comparing sensitivity runs with and without tidal motions in an OGCM, Schiller (2004) found that tidal motions can rectify Indonesian Archipelago hydrography and the Indonesian Throughflow (ITF) transport. Similar work via a coupled climate model concludes that the simulation with tidal motions can improve the modelled North Atlantic Current and Western European Climate (Müller et al., 2010). Yu et al. (2016) also implemented a tidal module into an OGCM. But unlike the above research that concentrates on regional circulations, they discuss the influence on global hydrography and circulation. The main finding

is that the tidal motions reduce the strength of wind-induced circulation and the upper cell of AMOC while slightly increasing the strength of the bottom cell of AMOC. In addition, a special kind of explicit forcing representing the tidal effect is the tidal residual circulation. By considering the tidal residual circulation as an external forcing in an OGCM, Bessières et al. (2008) concluded that the ocean state is insensitive to the tidal residual circulation, which can be neglected in climate research.

Generally speaking, there are mainly two approaches to study the tidal effect on the ocean and climate: parameterised tidal mixing and explicit tidal forcing. Even though both approaches have been conducted in previous research, no research has assessed these two approaches in the same framework. Therefore, we implement both approaches in our work. Our objectives are:

- (1) to find out the differences between these two approaches in expressing the effect of internal tide mixing in an OGCM;
- (2) to evaluate the advantages and disadvantages of these two approaches in climate research;
- (3) to see if there are any tidal effects other than internal tide mixing that would change the model results, which is not considered in the CVMIX_TIDAL parameterisation.

In Sect. 2, we first introduce the applied numerical model and its configuration. ~~Then in Sect. A, we verify our newly implemented tidal potential model.~~ The model results are presented in Sect. 3 and are further discussed in Sect. 4. The last section is the conclusion of this work.

2 Model description

The model applied in this work is the Finite-volume Sea ice–Ocean Model (Danilov et al., 2017, FESOM2), which is the successor of the Finite Element Sea ice–Ocean Model (Wang et al., 2014, FESOM). FESOM2 applies unstructured meshes, which are composed of irregular-sized triangles. In FESOM2 meshes, scalars and vectors are located at triangle vertices and triangle centres, respectively. By using multi-resolution meshes, specific regions of the global ocean can be well studied with moderate computational effort. The arbitrary Lagrangian–Eulerian vertical coordinate, which incorporates different vertical coordinates in the same frame, is introduced in FESOM2 (Danilov et al., 2017; Scholz et al., 2019). In this work, we apply the latest release of the model, namely, FESOM2.1. Thus in this paper, all terminologies for FESOM2 refer to FESOM2.1.

2.1 Model setup

In this work, we apply a common mesh for FESOM2, namely, the CORE-II mesh (<https://fesom.de/models/meshesetups/>). The CORE-II mesh features low resolution ($\sim 1^\circ$) in mid-latitude areas and high resolution at coastal regions, the equator, and north of 50° N. It has about 127k horizontal vertices and has 47 progressively thickening vertical layers from top to bottom. Since our work includes the simulation of tidal motions, the instant sea surface displacement can exceed 10 m at some spots, which requires either to thicken the uppermost vertical layer (from 5 m to 20 m) or to select an appropriate vertical coordinate. Here we keep the vertical layer configuration while changing the vertical coordinate to the z-star coordinate. To keep a positive thickness for the uppermost layer, the z-star coordinate distributes ocean surface displacement into changes in layer thicknesses for all vertical layers except the bottom layer. A standard timestep for FESOM2 configuration is 2700 s, which is too long to

90 resolve fast-travelling barotropic tides. Through sensitivity tests, an optimal timestep of 720 s has been determined (see [Seet-Appendix A](#)). In addition, since the model considers tidal motions and some related internal gravity waves, the Courant–Friedrichs–Lewy (CFL) stability condition could be constrained by the vertical advection term (Lemarié et al., 2015). Thus, a vertical velocity splitting method is applied in this work to keep vertical advection and oceanic mixing stable (Shchepetkin, 2015; Danilov et al., 2017). Instead of explicitly treating the whole vertical advection term, when the vertical Courant number
 95 exceeds unity, the excess part of the vertical velocity would be treated implicitly.

Except for the customised model configuration above, we adopt other generic settings for FESOM2 in this work. The model starts from the Polar Science Centre Hydrographic Climatology (Steele et al., 2001, PHC3) and is driven by the Coordinated Ocean-ice Reference Experiments phase II (CORE-II) reanalysis atmospheric forcing (Large and Yeager, 2009) ranging from 1948 to 2009. In addition, the sea surface salinity is restored to climatology values during the simulation. The horizontal
 100 viscosity in the model applies an ocean kinetic energy backscatter parameterisation scheme (Juricke et al., 2019). Constant background mixing is set as 10^{-4} and $10^{-5} \text{ m}^2 \text{ s}^{-1}$ for vertical viscosity and diffusivity. In addition to the constant background mixing, we apply the K-profile parameterisation (Large et al., 1994, KPP) in this work. The Gent–McWilliams scheme (Gent and McWilliams, 1990) and the Redi isoneutral diffusion (Redi, 1982) are applied to the parameterisation of subgrid eddy-induced mixing. Note here that the Gent–McWilliams parameterisation applied is formulated following Ferrari et al. (2010).

105 2.2 The tidal potential module

We implement a global tidal potential module (Thomas et al., 2001) into FESOM2 to consider the real tidal motions. The tidal potential in FESOM2 can be expressed as

$$\Omega = \Omega_g + \Omega_e + \Omega_{SAL} = \alpha\Omega_g + \beta g\eta. \quad (1)$$

The tidal forcing is expressed in the model as the gradient of the tidal potential Ω , which consists of three parts according to Eq.
 110 (1). The lunisolar gravitational potential Ω_g is calculated by considering the real-time position of the Sun and Moon with an ephemeris approach (Duffett-Smith, 1990). Except for the lunisolar gravitational potential term, two correction terms should also be considered. The solid Earth tide Ω_e driven by the Sun and Moon has a counteracting effect to the ocean tides, which is expressed as a portion of the lunisolar gravitational potential. Thus, $\Omega_g + \Omega_e$ can be written as $\alpha\Omega_g$, where α represents the effective Earth elasticity factor. Though α is distinct for diurnal tidal components (Wahr, 1981), an identical factor of 0.69
 115 is applied in this work following Kantha (1995). The self-attraction and loading (SAL) effect Ω_{SAL} considers the seafloor deformation and the changing of the earth gravity field caused by the weight distribution of seawater. To determine the SAL effect, one needs an explicit calculation based on a convolution integral (e.g., Shihora et al., 2022). However, this method is costly, and thus we use a simplified form $\beta g\eta$. η and g represent sea surface elevation and the gravitational acceleration. The scale factor β has been estimated in previous work (e.g., Accad and Pekeris, 1978; Ray, 1998; Stepanov and Hughes, 2004)
 120 and is set to 0.1 here.

2.3 The CVMIX_TIDAL parameterisation

The CVMIX_TIDAL parameterisation is implemented into FESOM2 along with the CVMIX library (Scholz et al., 2022). It is expressed as additional vertical diffusivity and vertical viscosity terms in the model. Take vertical diffusivity (k_v) as an example.

$$125 \quad k'_v = k_v + \frac{\Gamma \epsilon}{N^2} = k_v + \frac{q\Gamma E(x,y)F(z)}{\rho N^2} \quad (2)$$

In Eq. (2), the term k_v represents the original vertical diffusivity, which is the sum of constant background vertical diffusivity and KPP vertical diffusivity in this work. ρ and N denote seawater density and buoyancy frequency, respectively. The mixing efficiency Γ , which is set to 0.2 following Osborn (1980), represents the ~~rate~~ fraction of dissipated mechanical energy that leads to oceanic mixing. ϵ represents the tidal energy dissipation rate, which can be further expanded as $qE(x,y)F(z)/\rho$. The tidal dissipation efficiency q is estimated as 0.33, indicating that one-third of the tidal energy is dissipated locally, while the rest is radiated away and dissipated remotely as propagating low mode internal waves. Note that the remotely dissipated energy is already contained in the background vertical diffusivity in an OGCM (Simmons et al., 2004). The horizontal and vertical distribution functions of tidal energy dissipation are expressed as $E(x,y)$ and $F(z)$, which can be further expanded as below.

$$135 \quad E(x,y) = \frac{1}{2}\rho_0 N_b \kappa h^2 \langle u^2 \rangle \quad (3)$$

$$140 \quad F(z) = \frac{e^{-(H+z)/\zeta}}{\zeta(1 - e^{-H/\zeta})} \quad (4)$$

In Eq. (3), $E(x,y)$ means the parameterised energy conversion rate from barotropic to baroclinic tides. ρ_0 and N_b represent the reference seawater density and bottom buoyancy frequency, respectively. $\langle u^2 \rangle$ is the temporal mean square tidal velocity, representing the barotropic tide energy. κ and h are the wavenumbers and amplitude scales of ocean topography, representing the bottom roughness. It should be noted here that κ is a tuning variable so that the global integral of $E(x,y)$ fits the general value (Munk and Wunsch, 1998, 1 TW). In Eq. (4), the vertical distribution function $F(z)$ is a bottom intensified exponential profile whose depth integral is unity. ζ , which is suggested as 500 m, denotes the e-folding scale of $F(z)$ from the bottom $-H$. The additional vertical viscosity is equal to the additional vertical diffusivity. That means the parameterisation considers the viscosity–diffusivity ratio (also known as the Prandtl number) as unity, which ~~consists~~ agrees with the Prandtl number in the KPP scheme beneath the surface boundary layer.

2.4 Sensitivity runs

We set three experiments with the above configuration to investigate the influence of tide-induced mixing on the ocean state. The control run is named NOTIDE, which does not consider any kind of tidal effect. The first comparison run, LSTIDE, considers the lunisolar tidal potential in the momentum equations. The second comparison run, CVTIDE, does not simulate real ocean tides but applies the parameterisation of tide-induced mixing (CVMIX_TIDAL). ~~The information of the sensitivity~~

runs is listed in Table ???. It should be noted that all other model configurations are identical among the three sensitivity runs. Following the spin-up strategy of Danek et al. (2019), the three sensitivity experiments start from the PHC3 climatology and are periodically forced by the CORE-II forcing. Each experiment is conducted for five consecutive cycles, namely, R1 to R5. Each cycle corresponds with the year 1948 to 2009. The spin-up strategy is demonstrated intuitively in Table ??. Unless otherwise specified, all the model results shown in this paper are the average of the last 50 model years (1960 to 2009 in R5).

3 Validation of the tidal potential module

Though the tidal potential module has been applied in previous work (e.g., Müller et al., 2010; Weber and Thomas, 2017), the performance of this module in FESOM2 still needs to be evaluated. One crucial factor that should be carefully treated is the model timestep. Thus, a set of sensitivity runs is conducted to find the best timestep for the CORE-II mesh. The sensitivity runs are configured with timesteps set to 300, 360, 400, 450, 600, 720, 900, 1200, and 1800, respectively. Each experiment is integrated for one year, and the hourly model output of sea surface height is applied for harmonic analysis using the T_TIDE package (Pawlowicz et al., 2002). Figure A1 shows the co-tidal charts of M2 tides from all sensitivity runs, demonstrating that the amplitudes of M2 tides are getting larger when we reduce the model timestep. However, the Greenwich phase lags are not significantly changed except for the last two runs (with timesteps set to 1200 and 1800). The reason is that the timesteps of the last two runs are too long to resolve the propagation of tidal waves, leading to damped tidal amplitudes and distorted patterns of co-phase lines in, for example, the South Pacific Ocean.

To quantitatively determine the timestep for the best performance of the tidal potential module applying CORE-II mesh, we compare the harmonic tidal constants from the model results with those from tide gauge data. The University of Hawaii Sea Level Centre research quality data (Caldwell et al., 2015) is applied in this work. We obtain 687 tide gauge series at 540 tide gauge stations. Note that some tide gauge series refer to the same station and we integrate those into one. We exclude some tide gauge stations because: (1) these stations are not located in the CORE-II mesh; (2) the water depths at these stations in the model are shallower than 200. We exclude shelf tide gauges because global tide models do not have a good performance in shelf areas inherently. Improving model results in shelf areas may lead to a larger error in the global view. A similar approach is also used in previous work that compares model results with pelagic tide gauges (Müller et al., 2010). Since most tide gauges span over multiple years, each tide gauge series is truncated into several yearly segments according to the calendar year. For the preciseness of analysis, segments with a completeness index (proportion of data span without missing data) lower than 0.8 are discarded. Tidal harmonic analysis is done to each annual segment at each station, and the harmonic tidal constants at each station are thus averaged from multiyear constants. Finally, 229 tide gauge stations are applied for the comparison with model results.

180 A root-mean-square error (RMSE), referring to Cummins and Oey (1997), is defined as below to estimate the deviations of model results:-

$$rmse = \sqrt{\frac{1}{N} \sum_{i=1}^N \left[\frac{1}{2} (A_o^2 + A_m^2) - A_o A_m \cos(\phi_o - \phi_m) \right]}$$

In Eq. (A1), A_o and ϕ_o represent tidal amplitudes and Greenwich phase lags derived from the observational data, while A_m and ϕ_m represent those derived from the model results. It should be noted here that the RMSE defined in Eq. (A1) can be evaluated either for one tidal component throughout all tide gauges or several tidal components at one tide gauge. In the first case, $N = 229$ for one tidal component throughout all 229 tide gauge stations (see Fig. A2a); in the second case, $N = 2$ for two main diurnal/semidiurnal tidal components at one tide gauge station (see Figs. A2b and A2c). Fig. A2a shows the RMSEs at all tide gauge stations for each sensitivity run's four main tidal constituents. It can be found that the last two sensitivity runs present significant errors for M2 tides. As we claim before in Fig. A1, a large timestep may not resolve fast-travelling barotropic tides, leading to the damping of tide amplitudes. By considering all RMSEs of the four main tidal constituents, one can find that an optimal timestep lies between 450 and 720. Considering the computational cost, we finally choose a timestep of 720 to simulate global tides with the CORE-II mesh. With the timestep set to 720, the RMSEs for diurnal (O1 and K1) and semidiurnal (M2 and S2) tidal components at each tide gauge station is further shown in Figs. A2b and A2c, respectively. It can be seen that semidiurnal tides have larger deviations than diurnal ones, and gauges close to the coasts have larger deviations than those in basin centres.

In general, the tidal potential model demonstrates good consistency with tide gauges in most areas. However, we should state that some tide gauges also show considerable inconsistency from series to series and from year to year. For example, the Greenwich phase lags at some stations may have a discrepancy exceeding 120° among different years. That would introduce significant errors in estimating harmonic tidal constants at these stations. We speculate that the inconsistency may come from the changing of observational instruments.

3 Model results

3.1 Hydrography

We compare model results with the World Ocean Atlas 2018 (Locarnini et al., 2018; Zweng et al., 2018, WOA18). The decadal-averaged annual dataset with a resolution of 0.25° is obtained. For comparability, the in situ temperatures from the WOA18 dataset are converted to potential temperatures. Unless otherwise specified, temperatures are referred to as [refer to](#) potential temperatures in the below context.

Figure 1 shows the temperature [biases differences](#) between the sensitivity runs and WOA18. In the upper layer, LSTIDE generally reduces the temperature biases in the low-latitude areas of the three major oceans, indicating lower temperature biases than CVTIDE. However, in the subpolar North Pacific Ocean, CVTIDE shows a better reduction of the warm biases than LSTIDE. In the intermediate layer, LSTIDE generally reduces the temperature biases in most mid- and low-latitude areas.

But CVTIDE has lower temperature biases in the whole Arctic region. In addition, neither CVTIDE nor LSTIDE presents ~~major-significant~~ improvements to the Southern Ocean. In the deep layer, CVTIDE shows a cooling effect in the Southern Ocean, but LSTIDE shows a warming effect in the basins of the three major oceans, showing lower temperature biases in the global view. Figure 2 is the same as Fig. 1 but for salinity. In the upper layer, CVTIDE mainly reduces the salinity biases in the tropical Pacific Ocean. In contrast, LSTIDE reduces the salinity biases in the tropical Pacific Ocean and the tropical Indian Ocean. In the intermediate layer, CVTIDE reduces salinity biases in the tropical Pacific Ocean and the tropical Indian Ocean. In contrast, LSTIDE reduces salinity biases in the North Pacific Ocean and North Atlantic subtropical gyre area. In the deep layer, only LSTIDE shows improvements in the low-latitude area in the Atlantic Ocean.

The tidal effects also demonstrate ~~different-degrees-of-improvements~~ improvements to different extents at different depths of each basin. Figure 3 shows ~~basin-averaged~~ basin-averaged temperature and salinity biases at each vertical layer in the model. In the left panels, LSTIDE shows significant improvements to the temperature biases in the Pacific, Atlantic, and Indian Oceans deeper than 1000 m. CVTIDE only reduces temperature biases in the Arctic Ocean at 500–2000 m in depth. Also, both CVTIDE and LSTIDE reduce temperature biases in the Southern Ocean deeper than 4000 m. In the right panels, ~~improvements can only be found~~ we can only find improvements in LSTIDE. Salinity biases are slightly reduced in the Atlantic and Pacific Oceans, around 1500 m in depth. ~~It should also be noted~~ We should also note that in Fig. 3j, ~~large-significant~~ salinity biases emerge ~~in-on~~ the surface Arctic Ocean. That comes from the differences between the PHC3 and WOA datasets. Our model starts with the PHC3 climatology, which integrates the WOA dataset and the Arctic Ocean Atlas.

~~Both~~ CVTIDE and LSTIDE also greatly influence the hydrography in the continental shelf and slope areas, which cannot be clearly shown in Figs. 1–3. By assembling grids with the same bathymetry and averaging the temperature and salinity biases, the bathymetry averaged biases with respect to depths are shown in Fig. 4. In Fig. 4a, the model results in the control run demonstrate significant temperature biases near continental shelf and slope topographies, and both CVTIDE and LSTIDE decrease the biases in these areas. By comparing Fig. 4c with Fig. 4e, CVTIDE has a better improvement near the bottom of the continental shelf and slope, while LSTIDE ~~has a significant reduction of~~ significantly reduces temperature biases at 1000–4000 m in depth, which is not shown in CVTIDE. In Fig. 4b, significant salinity biases in the control run concentrate in the shallow continental shelf areas. It is shown in Figs. 4d and 4f that CVTIDE decreases salinity biases at the near-bottom continental shelf area, while LSTIDE decreases salinity biases in the whole layer of continental shelf area. Neither of the two comparison runs demonstrates a significant improvement to other areas, except for the slight ~~improvement~~ modification of LSTIDE around 1500 m in depth.

3.2 Sea ice

~~Compared with the control run, both~~ Figure 5 shows the results of sea ice thickness (SIT) in the polar regions. Both tidal effects alter the sea ice results, but the patterns are different. ~~Figure ?? shows the results of sea ice thickness (SIT) in the Arctic region.~~

In the Arctic region, CVTIDE generally ~~shows SIT decrease. In March, a remarkable SIT decrease occurs~~ decreases SIT in the Labrador Shelf ~~(along with the Hudson Strait, the Davis Strait, and the Baffin Bay) and the Okhotsk Sea. In September, an~~

245 ~~SIT decrease occurs in the Baffin Bay, the northeast Greenland Shelf, and the north Barents Shelf. But an SIT increase emerges~~
~~East Greenland Shelf and the Okhotsk Sea but increases SIT in the Canadian Arctic Archipelago (CAA) in CVTIDE. LSTIDE~~
~~also shows an SIT decrease. LSTIDE shows a slighter SIT decrease than CVTIDE in the Okhotsk Sea and Labrador Shelf~~
~~but is weaker than CVTIDE. Unlike CVTIDE, increases SIT in the Greenland Sea. One of the most significant differences~~
~~between LSTIDE and CVTIDE is that LSTIDE shows a remarkable SIT decrease in the CAA. The mean SIT in the CAA~~
250 ~~in the control run is 6.01 m(5.23) in March (September). LSTIDE decreases; LSTIDE decreases by 0.60 m(0.67), while~~
~~CVTIDE increases by 0.03 m(0.03). In addition, SIT slightly increases in the Greenland-Iceland Sea (March), the Baffin Bay,~~
~~the northeast Greenland Shelf, and the north Barents Shelf (September) in LSTIDE, and that is also different from CVTIDE.~~

~~Figure ?? is the same as Fig. ?? but for Antarctica. In March, both tidal effects show similar changes in SIT around~~
~~Antarctica, but CVTIDE is more prominent than LSTIDE. In September, In the Antarctic region, both CVTIDE and LSTIDE~~
255 ~~also show similar patterns, including a general SIT decrease around Antarctica and an SIT increase on the Weddell Sea Shelf~~
~~and the Ross Sea Shelf. But in the Western Antarctica Continental Shelf and Ross Sea Continental Shelf. On the West Antarctic~~
~~Peninsula Continental Shelf, SIT decreases in CVTIDE while but increases in LSTIDE. In addition, in the area ranging from~~
~~30° E to 60° E, SIT increases in CVTIDE but decreases in Generally speaking, CVTIDE shows a more significant impact on~~
~~SIT than LSTIDE.~~

260 ~~It should also be pointed out that in both Fig. ?? and Fig. ??, We should also point out that CVTIDE consistently reduces~~
~~SIT in both regions at shelf-break areas, which are indicated by the grey dashed lines. However, LSTIDE is not showing does~~
~~not show this consistency. The difference is further discussed in Sect. 4.~~

3.3 Mixed layer depth

~~Two definitions of mixed layer depth (MLD) are computed in FESOM2. The first one is defined as the shallowest depth~~
265 ~~where the vertical buoyancy gradient is equal to a local critical buoyancy gradient, following Large et al. (1997). The results~~
~~are shown in Fig. ?. In the North Pacific Ocean, CVTIDE deepens MLD in the Sea of Okhotsk and the Bering Sea, while~~
~~LSTIDE shows significant changes in the Kuroshio-Oyashio front. The pattern features a “blue-to-red” dipole, indicating an~~
~~equatorward displacement of the deep mixed layer area near the Kuroshio. In the North Atlantic Ocean, LSTIDE shows a~~
~~deepening effect on the MLD, while CVTIDE shows a shoaling effect on the MLD in the centre Labrador Sea. In the Southern~~
270 ~~Ocean, both CVTIDE and LSTIDE deepen MLD in the area ranging from 180° to 60° W, but the effect of CVTIDE is stronger~~
~~than that of LSTIDE.~~

~~The second kind of MLD, following Monterey and Levitus (1997), is defined as the depth where the density over depth~~
~~differs by 0.125 sigma units from the surface density. This definition shows larger MLDs in the deep convection area compared~~
~~with the previous one. As shown in Fig. ?, the MLDs in the Labrador Sea and the Weddell Sea can be greater than 3000,~~
275 ~~and both CVTIDE and LSTIDE deepen the MLDs in the deep convection areas. In the Labrador Sea, the deepening effect of~~
~~LSTIDE (500) is stronger than that of CVTIDE (200). In the Weddell Sea, both CVTIDE and LSTIDE extend regions with~~
~~deep mixed layers to the east (ranging from 30° W to 0°), but the deepening effect of CVTIDE (1700) is much stronger than~~
~~that of LSTIDE (460).~~

3.4 Meridional Overturning Circulation

280 ~~Both CVTIDE and LSTIDE influence the MOC. As shown in Meridional Overturning Circulation The left column of Fig. 6~~
~~e and 6e, shows that~~ CVTIDE and LSTIDE ~~show different effects in the~~ affect differently on AMOC. CVTIDE weakens the
AMOC upper cell by 0.50 Sv and strengthens the lower cell by 0.75 Sv, while LSTIDE strengthens the AMOC upper cell by
1.50 Sv and weakens the lower cell by 0.25 Sv. That indicates CVTIDE weakens the North Atlantic Deep Water (NADW)
formation and strengthens the Antarctic Bottom Water (AABW) formation, while LSTIDE ~~shows~~ behaves the opposite.

285 The results of PMOC are shown in the right column of Fig. 6. The main effect of CVTIDE on the PMOC occurs in the south
Indo-Pacific Ocean. The deep water upwelling at 20° S is enhanced ~~for by~~ 4 Sv. ~~The~~ LSTIDE has a ~~much~~ more significant
effect than CVTIDE. Figure 6f features two cells: the south cell upwells deep water in the tropical Indo-Pacific Ocean, while the
north cell upwells deep water in the subarctic North Pacific Ocean. Both cells enhance deep water upwelling ~~for by~~ 5 Sv, and
the upwelled deep waters are released to the Indo-Pacific Ocean's intermediate layer (about 1000–2000 m) ~~of the Indo-Pacific~~
290 ~~Ocean. It should also be noted here that the~~. The north cell in Fig. 6f shows consistency with the North Pacific bathymetry,
which is further discussed in Sect. 4.

3.4 Vertical diffusivity

~~The results of~~ Since tidal mixing is stimulated by topographies, tidal mixing and vertical diffusivity are bottom-enhanced.
Figures 7a–7c show the near-bottom vertical diffusivity in the three sensitivity runs ~~are shown in~~. Compared with the control
295 run, Fig. 7. ~~The middle column shows that~~ b shows observably that the CVMIX_TIDAL parameterisation mainly enhances
vertical diffusivity by a large value in the deep ocean. ~~The difference between CVTIDE and NOTIDE is most significant at~~
~~the bottom layer and is dependent on the~~ In CVTIDE, the vertical diffusivity is enhanced where bottom topography roughness
~~Vertical diffusivity is enhanced in~~ are large, such as seamounts, deep ocean ridges ~~and~~ shelf-break areas. The pattern of
LSTIDE is much different compared with CVTIDE, and we cannot see a coherent ~~increase~~ enhancement of vertical diffusivity
300 in the ~~deep/bottom layer ocean. We should also notice that the differences between CVTIDE/LSTIDE and NOTIDE in the upper~~
~~and intermediate layers show similar patterns with the MLD results~~ near-bottom ocean. The global mean vertical diffusivity
profiles (Fig. ??). ~~That indicates deep mixed layers are related to high vertical diffusivities in the upper~~ 7d) also demonstrate
that compared with CVTIDE, LSTIDE does not enhance vertical diffusivity significantly in the deep ocean.

3.5 Barotropic streamfunction

305 ~~The barotropic streamfunctions in the three sensitivity runs are shown in Fig. 8.~~ Figure 8 demonstrates that CVTIDE and
LSTIDE show similar patterns in changing the global barotropic streamfunction. The patterns of streamfunction differences
in Figs. 8b and 8c mainly consist of two loops. The “blue loop” ~~is mainly located~~ mainly locates in the Indo-Pacific Ocean:
the southward transport enhances the ITF and finally contributes to the ACC, ~~while~~ and its recirculation locates in the western
Pacific Ocean. The “red loop” enhances the ACC separately. The blue and red loops are stronger in LSTIDE than those in
310 CVTIDE. CVTIDE enhances the ITF and the ACC through the Drake Passage by 1.72 Sv and 4.79 Sv, while ~~those values~~

for the values of LSTIDE are 3.92 Sv and 10.34 Sv, respectively. In addition, we can find an important essential difference regarding the blue loop: the blue loop extends to the subpolar Pacific Ocean and the South Atlantic Ocean in LSTIDE but not in CVTIDE. The difference determines the PMOC–AMOC connection, which is ~~mentioned again~~ discussed in Sect. 4.

3.6 Energy

315 ~~Several energy terms are diagnosed~~ Based on the energy budget equation (see Appendix B1), we diagnose the energy terms
from the model results ~~, and and list~~ the global integration values ~~of these energy terms are shown~~ in Table 1. Due to the
additional energy input in LSTIDE (see tidal potential power and total kinetic energy), three energy dissipation terms (bottom
drag, vertical viscous dissipation and horizontal viscous dissipation) are enhanced compared with NOTIDE and CVTIDE,
especially for bottom drag. The additional energy sink in these three terms accounts for about 1 TW. ~~We should also notice~~
320 ~~that~~ The rest 3.5 TW of the total 4.6 TW tidal potential power ~~goes to surface pressure power, which comes from the sea~~
~~surface movement induced by surface tides. Even though the global integration of buoyancy flux shows no difference in the~~
~~three runs, the horizontal distributions show differences. The horizontal distributions of vertical viscous dissipation, bottom~~
~~drag, and buoyancy flux in the three sensitivity runs are further shown in Fig. 9. From Fig. 9c, we can see that LSTIDE~~
~~enhances bottom drag significantly. In most areas, the enhancement is about one order, but it can reach an order of 5 in several~~
325 ~~marginal seas because~~ is converted from kinetic energy to barotropic potential energy.

Figure 9 shows the horizontal distribution of vertically integrated bottom drag and viscous dissipation. Because of the strong
tidal currents ~~. Because of the addition of explicit tidal velocities, the bottom drag and the vertical shear of horizontal velocity~~
~~are enhanced, especially on in marginal seas and~~ continental shelf areas. ~~Thus, LSTIDE also enhances viscous dissipation,~~
~~which is related to vertical shear. It should be noted here that,~~ LSTIDE enhances bottom drag and viscous dissipation
330 significantly in these areas. Like the vertical diffusivity, the vertical viscosity is also enhanced in the CVMIX_TIDAL pa-
rameterisation, but the effect is negligible (~~see Fig. 9d). From the bottom row of Fig. 9, we can find that compared e).~~

Even though the global integration of buoyancy flux shows no difference in the three runs, the horizontal distribution shows
differences (Fig. 10). Regional integration over the Kuril–Aleutian Ridge area (120° E–120° W, 30° N–60° N) in NOTIDE,
CVTIDE and LSTIDE are –1.22, –1.52 and –2.78 TW, respectively. In the Indonesian Archipelago area (90° E–150° E,
335 15° S–15° N), regional integration values in NOTIDE, CVTIDE and LSTIDE are –0.82, –1.13 and –1.19 TW, respectively.
Compared with CVTIDE, LSTIDE ~~has~~ shows a stronger enhancement of buoyancy flux in the Kuril–Aleutian Ridge and
Indonesian Archipelago but weaker in other mid-latitude areas. The lower panels of Fig. 10 indicate that with tidal currents in
LSTIDE, turbulent buoyancy flux plays an essential role in the oceanic mixing and buoyancy flux (see Appendix B2).

4 Discussion

340 4.1 The effect of resolution on tidal mixing

As shown in Munk and Wunsch (1998, their Fig. 4), internal tides are important [linking links](#) between surface tides and oceanic mixing. Indeed, Niwa and Hibiya (2011, 2014) demonstrate that the conversion rate of internal tide energy in a global tide model depends on the model resolution. Thus, if the model mesh cannot resolve internal tides, the internal-tide-induced mixing is missing in the model result. Figure 11 presents the ratio of mode-1 internal tide wavelengths to the mesh resolution.

345 We can find that most of the model areas cannot resolve propagating internal tides in the mid-latitude area, especially for semidiurnal ones in Fig. 11b. In addition, even though the unstructured model mesh has higher resolution in shelf-slope areas, it still cannot resolve internal tides therein because the wavelengths of internal tides are smaller in these areas. So we infer that LSTIDE has a strong dependence on the model resolution.

Evidence can be found in the model results. In Fig. 4, compared with CVTIDE (Figs. 4c and 4d), LSTIDE (Figs. 4e and 4f)

350 shows a weaker reduction of hydrographic biases near the bottom of shelf breaks. The internal tides at shelf-slope areas cannot be resolved in LSTIDE, which weakens the mixing effect. Tidal mixing on shelf breaks also affects SIT. Unlike other [areas, places](#) where the temperature decreases monotonically with depth, the polar regions have colder water near the surface than in the deeper layers. Thus in polar [areasregions](#), the tidal mixing on shelf-slope areas would lead to a warmer surface ocean, which should decrease the SIT. In our result, LSTIDE shows a smaller SIT decrease than CVTIDE on the shelf-slope area, such as

355 the Labrador Shelf ~~in Figs. ??b and ??c. Similar results can also be found near the West Antarctica Shelf (Figs. ??e and ??f and the Antarctic Continental Shelf (Fig. 5).~~ The effect of the resolution is also reflected in the model vertical diffusivity. Both observational results (Polzin et al., 1997) and CVMIX_TIDAL parameterisation suggest larger vertical diffusivities near the bottom, which is caused by the tide-topography interaction. However, vertical diffusivities in the [deep and bottom near-bottom](#) layers are not generally higher in LSTIDE than in NOTIDE (~~Figs. 7i and 7j~~[Fig. 7c](#)). The lack of resolving internal tides causes

360 the underestimation of vertical shear, thus leading to a different result compared with CVTIDE. The buoyancy flux results (~~Figs. 9h and 9i~~[10b and 10c](#)), which are most directly connected to oceanic upwelling, also show less mixing effects in mid-latitude areas in LSTIDE compared with CVTIDE. Figure 11 shows the model mesh can roughly resolve propagating internal tides in tropical areas, where strong mixing can be found in LSTIDE. ~~Figure 6f~~[LSTIDE \(Fig. 6f\)](#) shows strong mixing and upwelling in the equatorial Indo-Pacific Ocean, ~~while this~~ [which](#) is not clearly shown in CVTIDE (Fig. 6d). We think ~~that~~ this

365 comes from the underestimation of tidal dissipation efficiency (q). The tidal dissipation efficiency is set to 0.33 in this work, but previous work indicates higher efficiencies in the Indonesian Archipelago area (Koch-Larrouy et al., 2007; Nagai and Hibiya, 2015).

Figure 11 also implies ~~that there are critical latitudes for propagating internal tides~~ [trapped internal tides beyond critical latitudes](#). According to the dispersion relationship of internal tides, where the inertial frequency is higher than a tidal frequency

370 ($f > \omega$), internal tides at the tidal frequency would not have a wave solution, resulting in trapped internal tides instead of propagating internal tides (~~light grey areas~~ [latitudes higher than the dashed lines](#) in Fig. 11). Unlike propagating internal tides, trapped internal tides can only be dissipated in a small domain, leading to a higher local dissipation rate and thus stronger local

mixing. The Kuril Ridge and Aleutian Ridge, which connect the ~~Sea of Okhotsk~~ Okhotsk Sea and the Bering Sea with the North Pacific Ocean, are the main generation sites for trapped diurnal internal tides. Large dissipation rates by trapped internal tides are validated in the previous modelling research (Niwa and Hibiya, 2011, 2014; Tanaka et al., 2010, 2013; Falahat and Nycander, 2015), and intense mixing is also observed therein (Nakamura et al., 2010; Itoh et al., 2014).

LSTIDE demonstrates intense mixing in the Kuril Ridge and Aleutian Ridge, resulting in higher buoyancy flux than NOTIDE in these areas (Fig. 9i, 10c). LSTIDE does not change the vertical diffusivity significantly, ~~but~~, Still, it enhances the upwelling in the North Pacific. ~~This can be explained by the turbulent buoyancy fluxes (see Appendix ??). By calculating turbulent buoyancy fluxes and the equivalent vertical diffusivity, we can find that LSTIDE generates an equivalent vertical diffusivity reaching 10^{-2} via turbulent buoyancy fluxes (see Fig. ??). We explain this by tide-induced turbulent buoyancy flux (Fig. 10f and Appendix B2).~~ The strong mixing upwells an additional 5 Sv of deep water in the North Pacific (around 50° N) to the intermediate layer and then spreads to the south (Fig. 6f). The ~~intermediate water then~~ strong mixing in the Indonesian Archipelago also results in strong upwelling. The upwelled water from both the Indonesian Archipelago and Kuril–Aleutian Ridge contributes to the ITF (Fig. 8c) and flows into the Indian Ocean. Stronger PMOC helps to reduce temperature biases in the intermediate Indo-Pacific Ocean (Fig. 1f): the upwelling of the deep water reduces the warm biases in the North Pacific Ocean and cold biases in the tropical Indo-Pacific Ocean in NOTIDE. ~~The formation of intermediate water also pushes the Kuroshio–Oyashio front to the south, and that explains why we have “blue-to-red” dipole patterns for MLD (Fig. ??e) and vertical diffusivity (Fig. 7e) in the North Pacific.~~

Trapped internal tides and related mixing can be simulated with LSTIDE, because ~~the CORE-II mesh has higher resolution in the high-latitude areas and the~~, the CORE-II mesh is fine enough to resolve trapped internal tides ~~are not forming as travelling waves (Fig. 11).~~ As a comparison, even though CVTIDE does not show much influence on the PMOC in the North Pacific Ocean, it shows similar but weaker patterns in reducing the temperature biases in the intermediate layer. This probably comes from two reasons: (1) most of the trapped internal tides’ energy is dissipated locally, so the tidal dissipation efficiency is close to unity ($q \approx 1$) in this area instead of 0.33 from CVTIDE; (2) the simulation is not long enough to get stable. It should also be emphasised here that even though the PMOC lower cell in the North Pacific is reported in most previous work (e.g., Simmons et al., 2004; Oka and Niwa, 2013), our results in LSTIDE demonstrate a different shape for the PMOC lower cell in the North Pacific. The significant upwelling occurs in the Aleutian Ridge ~~and~~ the upwelling has great consistency with the topography (take the brown transection in Fig. 6f as an example). ~~It seems that the~~ The diurnal tidal mixing beyond critical latitudes causes an upwelling along the bottom boundary, which is not reported in previous work. Nevertheless, our result supports some previous arguments (Scott and Marotzke, 2002; Ferrari et al., 2016) that the upwelling caused by near boundary mixing is more important than interior mixing in closing the THC.

4.2 The tidal effects on the global THC

The model also shows interesting results in the spin-up. The time series of several main constituents of the global THC is demonstrated in Fig. 12. In Figs. 12a and 12b, we find that both CVTIDE and LSTIDE ~~demonstrate show~~ a stronger PMOC and ITF than the control run since the beginning. In the fourth and fifth cycles, the discrepancies of PMOC and ITF in strength

between CVTIDE/LSTIDE and NOTIDE become stable. In Fig. 12c, it is shown that in the first cycle, the strength of AMOC upper cell in both CVTIDE and LSTIDE is weaker than that in the control run. However, since the second cycle, the strength of the AMOC upper cell in LSTIDE starts to surpass that in NOTIDE. The strength of the AMOC upper cell in CVTIDE is weaker
410 than that in NOTIDE in all cycles. As mentioned in Sect. 2, the CVMIX_TIDAL scheme applies the same formula to the model viscosities, potentially increasing model dissipation and leading to a weaker circulation on shelf-slope areas. Thus, the wind-driven circulation, the poleward heat flux and the AMOC upper cell are weakened, which corresponds with previous work (e.g., Jayne, 2009; Yu et al., 2016). But a major difference between CVTIDE and LSTIDE embodies the subsequent enhancement of the AMOC upper cell in LSTIDE. Results from LSTIDE reveal that the upwelling in the North Pacific and Indonesian
415 Archipelago links to the North Atlantic. Figures 6f and 8c ~~reveal~~indicate that the enhancement of mixing in the North Pacific and Indonesian Archipelago results in a stronger PMOC and ITF. The ITF mass transport finally forms the “blue loop” in the barotropic streamfunction difference (Fig. 8c). The westernmost part of this loop extends to the South Atlantic, which seems to have no connection with the North Atlantic. But note here the ~~barotropic streamfunction has an offset effect in the vertical direction.~~Figure vertical structure of oceanic circulation cannot be revealed by the barotropic streamfunction. Nevertheless,
420 Fig. 6e further shows that the upper cell of AMOC is enhanced, and the effect originates from the South Atlantic. On the contrary, even though CVTIDE also has a stronger ITF and a “blue loop” streamfunction difference, it does not strengthen the upper cell of AMOC because the “blue loop” is restricted in the Indo-Pacific Ocean and does not extend to the South Atlantic (Fig. 8b). We
The connection between the Indo-Pacific Ocean and the Atlantic Ocean is the Agulhas leakage. The Agulhas leakage is strengthened in LSTIDE (Fig. 8c), but the pattern is cut off in CVTIDE (Fig. 8b). And that explains why the
425 strengthening of PMOC and ITF only leads to AMOC increase in LSTIDE. The importance of Agulhas leakage in connecting cross-basin circulation is also reported in Biastoch et al. (2008). In addition, we rule out the possibility that the tidal mixing in the Atlantic itself causes the strengthening of the AMOC upper cell for two reasons: (1) in Fig. 6e, the AMOC cell does not show a closed overturning circulation, which means 1.5 Sv of AMOC is not upwelled in the Atlantic Ocean north to 30° S; (2) the transection plot does not show either stronger vertical diffusivity or turbulent buoyancy flux in the Atlantic Ocean
430 (figure not shown). The cycle revealed in our result corresponds well with ~~the schematic diagram in~~previous paper (Talley et al., 2011, her Fig. S14.1c). The enhancement of ITF due to tidal mixing is also reported in previous research (Schiller, 2004; Sasaki et al., 2018), and the link between stronger PMOC lower cell and stronger AMOC upper cell can also be found in Melet et al. (2016).

As to the AMOC lower cell, LSTIDE shows a weakening effect while CVTIDE shows a strengthening effect (Fig. 12d).
435 It is widely believed that the global tidal mixing and wind-driven upwelling in the Southern Ocean affect the MOC (Ledwell et al., 2000; Marshall and Speer, 2012). Since the model configuration applies the same atmospheric forcing in the three sensitivity runs, the reason concerning wind stress would be excluded. Moreover, considering the vertical advection–diffusion balance (Munk and Wunsch, 1998; Prange et al., 2003), the tidal mixing should strengthen the MOC, as CVTIDE shows. As discussed above, LSTIDE has the disadvantage ~~in of~~demonstrating tide-induced mixing in the vast mid-latitude areas
440 due to the resolution; thus, the AMOC lower cell is weakened instead of strengthened in LSTIDE. We also guess that the enhanced vertical diffusivities may be added to deep convection areas, directly leading to stronger convection. However, the

CVMIX_TIDAL parameterisation implies that the enhanced vertical diffusivities are not added to areas where gravitational instability occurs ($N^2 < 0$). So we finally conclude that the AMOC lower cell is enhanced due to tidal mixing, and more AABW formation in the Weddell Sea is a result instead of a cause of the AMOC lower cell strengthening. A similar conclusion
445 can ~~also~~ be found in Exarchou et al. (2012).

CVTIDE and LSTIDE also demonstrate that the ACC transport through the Drake Passage is larger than that in NOTIDE (Figs. 8 and 12e). We can see from Fig. 12e that the strength of ACC in all three sensitivity runs decreases during the five cycles, but the ~~strength decrease more slowly~~ decreasing rate is lower in the comparison runs, which can be explained ~~with~~ by Fig. 13. Figures 13d and 13f show that both comparison runs have similar effects on changing potential density in the
450 southern hemisphere. In the mid- and low- latitude areas, both tidal effects demonstrate denser seawater in the upper ocean and lighter seawater in the lower ocean, indicating the promotion of ocean potential energy caused by the tidal mixing. In the high-latitude areas, both comparison runs show denser seawater from surface to bottom, indicating the reduction of ocean potential energy due to the deep convection. In depths below 1000 m, both CVTIDE and LSTIDE show potential density decrease at subtropical latitudes. The decrease ~~of~~ in potential density is caused by the mixing and advection effect indicated by
455 the global MOC (GMOC). For example, Fig. 13e shows that strong mixing and upwelling take place in the tropical region, and the upwelled water ~~is further advected~~ further advects to the south (similar to Fig. 6f). The advection effect causes the “blue tongue” pattern in Fig. 13f (1000–2000 m in depth). In addition, the ~~increase of potential density~~ potential density increase at subpolar latitudes can be explained by the AABW formation. In the above context, our results show that the tidal mixing in CVTIDE causes the strengthening of the AABW formation. This is also why the potential density increase at subpolar latitudes
460 is more substantial in CVTIDE than ~~that~~ in LSTIDE. Even though CVTIDE and LSTIDE alter potential density differently, both experiments increase the meridional density gradients in the Southern Ocean, which prevents the strength of ACC from decreasing due to the thermal wind relation. Thus we conclude that both CVTIDE and LSTIDE affect the ACC strength by enhancing the meridional density gradient. Similar results are also shown in Saenko and Merryfield (2005) that with tidal mixing parameterisation, the ACC transport is stronger by 25% compared with ~~a no-tide~~ their control run.

465 4.3 Other tidal effects: what does CVTIDE miss?

The “barotropic tide–baroclinic tide–oceanic mixing” chain is considered the most crucial tidal effect on the ocean and climate. But we are still questioning if there are any effects from tides other than internal tide mixing. Since CVTIDE only considers internal tide mixing effects and LSTIDE considers real tide effects, we can answer this question by comparing LSTIDE with CVTIDE.

470 The most significant difference is the energy in the model. If the tidal potential term is added to the ocean primitive equation, tide energy is thus input to the system (Table 1). The global tidal potential power is ~~estimated as~~ 4.6 TW in our model. However, most of the energy goes to the ~~surface pressure power~~ barotropic potential energy, indicating a lack of internal tides in the system. If the model resolves more internal tides, the surface tide amplitudes and the ~~surface pressure power term~~ conversion to barotropic potential energy would be smaller due to the internal wave drag (e.g., Jayne and St. Laurent, 2001;
475 Sulzbach et al., 2021). This part of energy would be redistributed to the buoyancy flux and energy dissipation terms, depending

on the conversion rate between surface tides and internal tides. External energy input not only leads to higher kinetic energy but also higher energy dissipation. Comparing Fig. 9b with Fig. 9c, we find that the bottom drag is significantly enhanced in LSTIDE than in CVTIDE. Regarding the viscous dissipation, LSTIDE shows remarkable enhancement on continental shelves (Fig. 9f), while CVTIDE does not show significant changes compared with NOTIDE. The reason is that LSTIDE has strong tidal shear caused by ~~bottom drag~~ the bottom drag of barotropic tides on continental shelves, while CVTIDE only increases viscosity and diffusivity on shelf breaks and seamounts, indicating an effect of internal tides.

Additionally, we ~~also~~ find that considering the strong tidal shear on continental shelves can reduce model biases. Both Fig. 4e and Fig. 4f indicate that in LSTIDE, the temperature and salinity biases on continental shelves (e.g., bottom depth less than 100 m) are better reduced compared with CVTIDE. Via a coupled climate model, Lee et al. (2006) also found that local hydrographic biases can be reduced when considering tidal shear effects on continental shelves.

From a mathematical point of view, ~~the addition of~~ adding periodic tidal motions can also affect stress terms, leading to a tidal residual effect. Even though tidal velocities can roughly be eliminated via time-averaging, the stress terms can always leave non-zero cross terms. These effects are more remarkable in regional simulations than global ones. Our results show that only the sea ice distribution in the CAA is significantly affected by the tidal residual effect. Figure ~~??~~ 5 shows that in LSTIDE, the SIT in the CAA is reduced by about 10%. Adding strong tidal currents in the narrow channels can lead to an additional ocean-to-ice stress, and the theoretical basis can be referred to in Hibler (1979, his Eq. 3). Our results also show that the additional stress causes sea ice transports from CAA to the adjacent areas, ~~such as the north Baffin Bay~~ (Fig. ~~??f~~).

5 Conclusions

The crucial effect of tides on the ocean and climate has been widely discussed in previous work (e.g., Polzin et al., 1997; Munk and Wunsch, 1998; Ledwell et al., 2000). Moreover, it is generally implemented in an OGCM via two approaches: parameterised tidal mixing and explicit tidal forcing. In this work, we implement both modules in FESOM2 to assess the two methods in the same framework. By analysing and summarising different aspects from the model results, we draw the conclusions below.

First, the resolution of the CORE-II mesh is not fine enough to simulate internal tides and internal tide mixing. Thus, the real tide simulation (LSTIDE) is not as good as tidal mixing parameterisation (CVTIDE). For instance, our control run shows ~~large~~ significant temperature biases in near-bottom shelf-slope areas, which can be reduced by tidal mixing. Also, tidal mixing in polar regions can lead to lower SIT in the model because of the polar stratificational feature. Additionally, tidal mixing in the deep ocean can lead to a stronger AMOC lower cell, with stronger AABW formation and deeper ~~MLD~~ mixed layer depth in the Weddell Sea. Nevertheless, these phenomena are not significant in LSTIDE ~~than~~ as in CVTIDE because LSTIDE generally underestimates the strength of tidal mixing.

Second, previous research has pointed out that applying 0.33 to the mixing efficiency (q) may cause underestimating tidal mixing in some areas. Our results further show that tidal mixing in the Indonesian Archipelago and Kuril–Aleutian Ridge are stronger in LSTIDE than CVTIDE. That is not contradictory ~~with~~ to the above statement, because the resolution in these areas

is relatively high and trapped internal tides are different from propagating internal tides. Our results show that the tidal mixing
510 in these two areas drives stronger upwelling in the deep Pacific Ocean. In the Kuril–Aleutian Ridge, deep water is upwelled to
the intermediate layer and advected to the tropical region; in the Indonesian Archipelago, ~~the upwelled deep water enhances
the ITF along with the intermediate~~ deep water ~~from the north~~ is also upwelled to the intermediate layer and advected to the
further south. Both upwelled water contributes to the ITF enhancement. The deep water upwelling in the Pacific Ocean and
the ITF enhancement result in lower hydrographic biases in the intermediate and deep layers of the North Pacific and tropical
515 Indo-Pacific Ocean, which is more ~~observably-observable~~ in LSTIDE than CVTIDE. Beyond that, ~~a link between the Agulhas
current connecting the~~ stronger PMOC upwelling ~~and with the~~ stronger AMOC upper cell is only found in LSTIDE. ~~Which
The Agulhas current is not enhanced in CVTIDE, thus the AMOC upper cell does not strengthen along with PMOC upwelling.
What~~ should also be noticed here is that the strong upwelling in the North Pacific Ocean in LSTIDE is not caused by the
vertical diffusivity in the model, but by the turbulent buoyancy flux. ~~Based on the theoretical derivation (Appendix ??), the
520 equivalent vertical diffusivity of turbulent buoyancy flux can reach 10^{-2} .~~

Third, both tidal effects enhance the strength of the ACC. We conclude that this is caused by thermal wind balance: both
CVTIDE and LSTIDE increase meridional density gradient in the Southern Ocean. However, the reason for ~~the increasing
of increasing~~ meridional density gradient is different: CVTIDE increases seawater density in the deep layers of subpolar
regions due to stronger AABW formation; LSTIDE decreases seawater density in the intermediate layers because of the PMOC
525 difference.

Finally, we summarise some other tidal effects which cannot be presented in the CVMIX_TIDAL parameterisation. Our
results show that when we implement real tide motion into the model, both total ocean kinetic energy and energy dissipation
increase. The globally integrated bottom drag is remarkably enhanced by almost one order of magnitude. Real tides also lead
to stronger viscous dissipation on continental shelves, indicating strong tidal shear and mixing in these shelf areas. Our results
530 further prove that the hydrographic biases in shelf areas are less reduced in CVTIDE than LSTIDE because the mixing ~~process
caused by barotropic tides~~ in shelf areas is not considered in the CVMIX_TIDAL parameterisation. In addition, the tidal
residual effect is ~~also~~ significant in ocean-to-ice stress, ~~which~~. This is verified in our results that the SIT in the CAA is reduced
by ~~about~~ 10% and the sea ice is transported to the adjacent areas.

In conclusion, our work shows that both spatial and temporal resolution needs to be refined to simulate real tides in climate
535 research. Previous research indicates that a model usually requires at least 0.1° to carry out a global internal tide simulation
(e.g., Shriver et al., 2012; Li et al., 2015), but no research ~~conduct-conducted~~ a run longer than 10 years with a resolution as
high as 0.1° due to the enormous computational cost. Thus, for long-term climate ~~study-studies~~ (e.g., Shi and Lohmann, 2016;
Ackermann et al., 2020; Lohmann et al., 2020), it is more applicable to use a parameterisation scheme for climate simula-
tions. Nevertheless, the real tide simulation also shows some phenomena that are not well expressed by the CVMIX_TIDAL
540 parameterisation. For example, the underestimation of the mixing efficiency parameter in the Indonesian Archipelago and
Kuril–Aleutian Ridge, the linking between stronger PMOC upwelling and stronger AMOC upper cell, and the tidal effects on
coastal and shelf areas are revealed in our results. The tidal effects, including mixing effects, are revealed from our ocean-only
model results, but ~~it should also be noted that the effects~~ we should also notice that the impact may be different in a coupled

climate model. In addition, vertical mixing might have changed over Earth's history and provide a possible mechanism for past
545 climate changes and climate latitudinal gradients (Lohmann, 2020). Thus, it is important to consider tidal effects in climate
research. Our future work on climate research considering tidal effects is to modify a tidal parameterisation scheme via a
high-resolution short-term real-tide case, as well as to evaluate tidal mixing schemes from other theoretical frameworks (e.g.,
Olbers and Eden, 2013).

Code and data availability. The FESOM2.1 source code applied in this work, the last 50 years averaged model results, as well as the post-
550 processing and visualisation codes, are archived on Zenodo (Song et al., 2021). The whole simulation results can be obtained by contacting
the corresponding author. The tide gauge data can be downloaded from the University of Hawaii Sea Level Centre's website (last access: 27
July 2020). The PHC3 climatology, WOA18 climatology and CORE-II forcing are freely available online.

Appendix A: Validation of the tidal potential module

Though the tidal potential module has been applied in previous work (e.g., Müller et al., 2010; Weber and Thomas, 2017), the
555 performance of this module in FESOM2 still needs to be evaluated. One crucial factor that should be carefully treated is the
model timestep. Thus, a set of sensitivity runs is conducted to find the best timestep for the CORE-II mesh. The sensitivity
runs are configured with timesteps set to 300, 360, 400, 450, 600, 720, 900, 1200, and 1800 s, respectively. Each experiment is
integrated for one year, and the hourly model output of sea surface height is applied for harmonic analysis using the T_TIDE
package (Pawlowicz et al., 2002). Figure A1 shows the co-tidal charts of M2 tides from all sensitivity runs, demonstrating that
560 the amplitudes of M2 tides are getting larger when we reduce the model timestep. However, the Greenwich phase lags are not
significantly changed except for the last two runs (with timesteps set to 1200 and 1800 s). The reason is that the timesteps
of the last two runs are too long to resolve the propagation of tidal waves, leading to damped tidal amplitudes and distorted
patterns of co-phase lines in, for example, the South Pacific Ocean.

To quantitatively determine the timestep for the best performance of the tidal potential module applying CORE-II mesh, we
565 compare the harmonic tidal constants from the model results with those from tide gauge data. The University of Hawaii Sea
Level Centre research quality data (Caldwell et al., 2015) is applied in this work. We obtain 687 tide gauge series at 540 tide
gauge stations. Note that some tide gauge series refer to the same station and we integrate those into one. We exclude some
tide gauge stations because: (1) these stations are not located in the CORE-II mesh; (2) the water depths at these stations in the
model are shallower than 200 m. We exclude shelf tide gauges because global tide models do not have a good performance in
570 shelf areas inherently. Improving model results in shelf areas may lead to a larger error in the global view. A similar approach is
also used in previous work that compares model results with pelagic tide gauges (Müller et al., 2010). Since most tide gauges
span over multiple years, each tide gauge series is truncated into several yearly segments according to the calendar year. For
the preciseness of analysis, segments with a completeness index (proportion of data span without missing data) lower than 0.8
are discarded. Tidal harmonic analysis is done to each annual segment at each station, and the harmonic tidal constants at each

575 station are thus averaged from multiyear constants. Finally, 229 tide gauge stations are applied for the comparison with model results.

A root-mean-square error (RMSE), referring to Cummins and Oey (1997), is defined as below to estimate the deviations of model results.

$$rmse = \sqrt{\frac{1}{N} \sum_{i=1}^N \left[\frac{1}{2} (A_o^2 + A_m^2) - A_o A_m \cos(\phi_o - \phi_m) \right]} \quad (A1)$$

580 In Eq. (A1), A_o and ϕ_o represent tidal amplitudes and Greenwich phase lags derived from the observational data, while A_m and ϕ_m represent those derived from the model results. It should be noted here that the RMSE defined in Eq. (A1) can be evaluated either for one tidal component throughout all tide gauges or several tidal components at one tide gauge. In the first case, $N = 229$ for one tidal component throughout all 229 tide gauge stations (see Fig. A2a); in the second case, $N = 2$ for two main diurnal/semidiurnal tidal components at one tide gauge station (see Figs. A2b and A2c). Figure A2a shows the RMSEs at all tide gauge stations for each sensitivity run's four main tidal constituents. It can be found that the last two sensitivity runs present significant errors for M2 tides. As we claim before in Fig. A1, a large timestep may not resolve fast-travelling barotropic tides, leading to the damping of tide amplitudes. By considering all RMSEs of the four main tidal constituents, one can find that an optimal timestep lies between 450 s and 720 s. Considering the computational cost, we finally choose a timestep of 720 s to simulate global tides with the CORE-II mesh. With the timestep set to 720 s, the RMSEs for diurnal (O1 and K1) and semidiurnal (M2 and S2) tidal components at each tide gauge station is further shown in Figs. A2b and A2c, respectively. It can be seen that semidiurnal tides have larger deviations than diurnal ones, and gauges close to the coasts have larger deviations than those in basin centres.

595 In general, the tidal potential model demonstrates good consistency with tide gauges in most areas. However, we should state that some tide gauges also show considerable inconsistency from series to series and from year to year. For example, the Greenwich phase lags at some stations may have a discrepancy exceeding 120° among different years. That would introduce significant errors in estimating harmonic tidal constants at these stations. We speculate that the inconsistency may come from the changing of observational instruments.

Appendix B: Theoretical background and equation derivation

B1 Calculation of the kinetic energy budget in FESOM2

600 We start from the ocean momentum balance Eq. (B1). \mathbf{u}_h and \mathbf{u} are horizontal and three-dimensional velocity vectors; f is the Coriolis parameter; $\hat{\omega}$ is the unit vector of the Earth angular velocity; ρ_o is a constant reference density; P represents the hydrostatic pressure; g is the gravitational acceleration; η is sea surface height; Ω is the tidal potential, which is also explained in Eq. (1); A_h and A_v are horizontal and vertical viscosity coefficients. ∇ and ∇_h represent three-dimensional and horizontal

605 gradient operator, respectively. Note here our experiment setting does not consider sea ice pressure and air pressure, thus, surface pressure only comes from the sea surface height.

$$\frac{\partial \mathbf{u}_h}{\partial t} + \mathbf{u} \cdot \nabla \mathbf{u}_h + f\hat{\omega} \times \mathbf{u}_h = -\frac{1}{\rho_0} \nabla_h P - g \nabla_h \eta + \nabla_h \Omega + \frac{\partial}{\partial x} \left(A_h \frac{\partial \mathbf{u}_h}{\partial x} \right) + \frac{\partial}{\partial y} \left(A_h \frac{\partial \mathbf{u}_h}{\partial y} \right) + \frac{\partial}{\partial z} \left(A_v \frac{\partial \mathbf{u}_h}{\partial z} \right) \quad (\text{B1})$$

Dot product between Eq. (B1) and $\rho_0 \mathbf{u}_h$ gives Eq. (B2). ρ is seawater density minus the constant reference density ρ_0 . ke means kinetic energy, which is expressed as $\frac{1}{2} \rho_0 \mathbf{u}_h^2$.

$$\begin{aligned} \frac{\partial ke}{\partial t} + \nabla \cdot (\mathbf{u} ke) = & -\nabla \cdot (\mathbf{u} P) - \rho g w - \mathbf{u}_h \cdot \nabla_h (\rho_0 g \eta) + \mathbf{u}_h \cdot \nabla_h (\rho_0 \Omega) + \\ & \frac{\partial}{\partial x} \left(A_h \frac{\partial ke}{\partial x} \right) + \frac{\partial}{\partial y} \left(A_h \frac{\partial ke}{\partial y} \right) + \frac{\partial}{\partial z} \left(A_v \frac{\partial ke}{\partial z} \right) - A_h \left(\frac{\partial \mathbf{u}_h}{\partial x} \right)^2 - A_h \left(\frac{\partial \mathbf{u}_h}{\partial y} \right)^2 - A_v \left(\frac{\partial \mathbf{u}_h}{\partial z} \right)^2 \end{aligned} \quad (\text{B2})$$

610 A global, full-depth integration and long-time mean drops out some minor terms and gives Eq. (B3). Note here that our experiment setting considers free slip lateral boundaries, so the integration of horizontal viscous term does not involve lateral friction. Nevertheless, the integration of vertical viscous term involves surface and bottom stress terms, in which \mathbf{u}_s and \mathbf{u}_b are surface and bottom horizontal velocities, and τ_s and τ_b are surface and bottom stress.

$$0 = -\rho g w - \mathbf{u}_h \cdot \nabla_h (\rho_0 g \eta) + \mathbf{u}_h \cdot \nabla_h (\rho_0 \Omega) - A_h \left(\frac{\partial \mathbf{u}_h}{\partial x} \right)^2 - A_h \left(\frac{\partial \mathbf{u}_h}{\partial y} \right)^2 - A_v \left(\frac{\partial \mathbf{u}_h}{\partial z} \right)^2 + \tau_s \cdot \mathbf{u}_s + \tau_b \cdot \mathbf{u}_b \quad (\text{B3})$$

615 In FESOM2, the horizontal viscosity is not calculated with an explicit form (Juricke et al., 2019). In order to evaluate the horizontal viscous dissipation, we have to compute directly the acceleration caused by horizontal viscosity ($\mathbf{G}_{\text{visch}}$) in the model. Thus the horizontal viscous dissipation term is actually calculated with $\mathbf{u}_h \cdot \mathbf{G}_{\text{visch}}$. The kinetic energy budget in FESOM2 is finally given as Eq. (B4). Note here we introduce buoyancy $b = -g\rho/\rho_0$ into the first term of the right hand side.

$$0 = \rho_0 b w - \mathbf{u}_h \cdot \nabla_h (\rho_0 g \eta) + \mathbf{u}_h \cdot \nabla_h (\rho_0 \Omega) + \mathbf{u}_h \cdot \mathbf{G}_{\text{visch}} - A_v \left(\frac{\partial \mathbf{u}_h}{\partial z} \right)^2 + \tau_s \cdot \mathbf{u}_s + \tau_b \cdot \mathbf{u}_b \quad (\text{B4})$$

620 Thus, on the right hand side of Eq. (B4), we have in order the buoyancy flux term, the ~~surface pressure power~~ barotropic potential energy-kinetic energy conversion term, the tidal potential power term, the horizontal viscous dissipation term, the vertical viscous dissipation term, the wind stress term and the bottom friction term. Theoretically, in a global, full-depth integration and long-term mean, these terms should balance. Residuals may come from: (1) the integral of terms we omit from Eq. (B2) to Eq. (B3); (2) horizontal interpolation between triangle vertices and triangle centroid or vertical interpolation between cell surface and cell centre; (3) the time difference between each term in Eq. (B1) and the velocity applied in the dot product. As to the third point, to reduce the residual term, we suggest to use the central time velocity instead of the updated velocity at each model timestep, especially for the velocity-sea surface height coupled term in a tidal case. Besides, all terms are calculated at each timestep and then averaged over time (take buoyancy flux as an example, $\rho_0 \overline{bw}$ instead of $\rho_0 \bar{b}\bar{w}$).

B2 The relationship between turbulent buoyancy flux and vertical diffusivity

630 Equation (B5) shows the ocean mass balance equation. K_h and K_v are horizontal and vertical diffusivity coefficients. S represents other source and sink terms regarding the mass balance. For the following derivation, we use buoyancy b instead of seawater density ρ here.

$$\frac{\partial b}{\partial t} + \mathbf{u}_h \cdot \nabla_h b + w \frac{\partial b}{\partial z} = \nabla_h \cdot (K_h \nabla_h b) + \frac{\partial}{\partial z} \left(K_v \frac{\partial b}{\partial z} \right) + S \quad (\text{B5})$$

635 Splitting turbulence terms from the advection terms gives Eq. (B6), in which overlines are time mean and primes are turbulence.

$$\frac{\partial \bar{b}}{\partial t} + \overline{\mathbf{u}_h} \cdot \nabla_h \bar{b} + \nabla_h \cdot (\overline{\mathbf{u}'_h b'}) + \bar{w} \frac{\partial \bar{b}}{\partial z} + \frac{\partial}{\partial z} (\overline{w' b'}) = \nabla_h \cdot (K_h \nabla_h \bar{b}) + \frac{\partial}{\partial z} \left(K_v \frac{\partial \bar{b}}{\partial z} \right) + S \quad (\text{B6})$$

A vertical advection–diffusion balance (e.g., Munk and Wunsch, 1998) gives below Eq. (B7).

$$\bar{w} \frac{\partial \bar{b}}{\partial z} = - \frac{\partial}{\partial z} (\overline{w' b'}) + \frac{\partial}{\partial z} \left(K_v \frac{\partial \bar{b}}{\partial z} \right) \quad (\text{B7})$$

640 Vertical diffusivity parameter K_v represents the turbulent buoyancy ~~flux-fluxes~~ $\overline{w' b'}$ that cannot be resolved in the model. In a traditional OGCM, such as NOTIDE/CVTIDE, the ocean currents roughly follow the geostrophic balance, but in LSTIDE we add ageostrophic flows. Tidal flows can easily cross the isobaths, therefore generating upwelling/downwelling around topographies, which alters turbulent buoyancy fluxes $\overline{w' b'}$. Assuming tidal motion as a turbulence of the ocean mean flow, CVTIDE ~~enhances K_v to consider tidal effects. However, LSTIDE already considers real tidal motions as well as the~~ parameterises turbulent buoyancy fluxes ~~caused by tides. This is equal to changing by altering vertical diffusivity parameter K_v in CVTIDE.~~
645 ~~An equivalent vertical diffusivity from the turbulent buoyancy flux is expressed as-. Thus, in Eq. (??). N is the buoyancy frequency.-~~

$$\underline{K_v} = - \frac{\overline{w' b'}}{N^2}$$

~~The presence of turbulent buoyancy flux $\overline{w' b'}$ explains B7), CVTIDE mainly changes the second right-hand-side term, while LSTIDE mainly changes the first right-hand-side term. That is~~ why LSTIDE shows stronger MOC without enhancing the
650 vertical diffusivity in the model.

Note here we use $\overline{w' b'} = \overline{wb} - \bar{w}\bar{b}$ to calculate turbulent buoyancy flux from the model ~~result~~ results.

B3 Estimating internal tide scale in the model

With a stratification $N^2(z)$ given, one can first solve the mode- n eigenvector $\Phi_n(z)$ and the respective mode- n eigenvalue c_n via the Sturm–Liouville equation below. In this work, we only study mode-1 internal tides.

$$655 \quad \frac{d^2 \Phi_n(z)}{dz^2} + \frac{N^2(z)}{c_n^2} \Phi_n(z) = 0 \quad (\text{B8})$$

With eigenvalue speeds c_n , according to Falahat and Nycander (2015) and Song and Chen (2020), the wavenumbers k_n of internal tides in the model are calculated as below.

$$k_n = \begin{cases} \frac{\sqrt{\omega^2 - f^2}}{c_n}, & \text{if } f < \omega; \\ \frac{\sqrt{f^2 - \omega^2}}{c_n}, & \text{if } f > \omega. \end{cases} \quad (\text{B9})$$

660 In Eq. (B9), ω means the angular frequency of a tidal component, and f represents the Coriolis frequency, which changes with latitude. The first case of Eq. (B9) applies to propagating internal tides, occurring below the critical latitudes. The second case of Eq. (B9) applies to trapped internal tides, occurring beyond the critical latitudes. Note that even though trapped internal tides do not propagate, they are confined within a scale related to k_n .

Finally, one can calculate the scale of an internal tide via $L = 2\pi/k_n$.

665 *Author contributions.* P. Song and D. Sidorenko implemented the tidal module into FESOM2. P. Scholz implemented the CVMIX package into FESOM2. P. Song performed all the experiments. All authors contributed to the analysis of the results.

Competing interests. The authors declare that they have no conflict of interest.

670 *Acknowledgements.* The authors thank Nikolay Koldunov for the help on FESOM2 settings and Sergey Danilov for discussing the model results. P. Song is funded by the *China Scholarship Council*. G. Lohmann and M. Thomas received funding through the topic *Ocean and Cryosphere under climate change* in the Program *Changing Earth - Sustaining our Future*. The experiments were performed at the AWI Computing Centre.

References


- Accad, Y. and Pekeris, C. L.: Solution of the tidal equations for the M2 and S2 tides in the world oceans from a knowledge of the tidal potential alone, *Philosophical Transactions of the Royal Society of London. Series A, Mathematical and Physical Sciences*, 290, 235–266, <https://doi.org/10.1098/rsta.1978.0083>, 1978.
- 675 Ackermann, L., Danek, C., Gierz, P., and Lohmann, G.: AMOC Recovery in a Multicentennial Scenario Using a Coupled Atmosphere-Ocean-Ice Sheet Model, *Geophysical Research Letters*, 47, 1–10, <https://doi.org/10.1029/2019GL086810>, 2020.
- Arbic, B. K., Alford, M. H., Ansong, J. K., Buijsman, M. C., Ciotti, R. B., Farrar, J. T., Hallberg, R. W., Henze, C. E., Hill, C. N., Luecke, C. A., Menemenlis, D., Metzger, E. J., Müller, M., Nelson, A. D., Nelson, B. C., Ngodock, H. E., Ponte, R. M., Richman, J. G., Savage, A. C., Scott, R. B., Shriver, J. F., Simmons, H. L., Souopgui, I., Timko, P. G., Wallcraft, A. J., Zamudio, L., and Zhao, Z.: A Primer
680 on Global Internal Tide and Internal Gravity Wave Continuum Modeling in HYCOM and MITgcm, in: *New Frontiers in Operational Oceanography*, pp. 307–392, *GODAE OceanView*, <https://doi.org/10.17125/gov2018.ch13>, 2018.
- Bessières, L., Madec, G., and Lyard, F.: Global tidal residual mean circulation: Does it affect a climate OGCM?, *Geophysical Research Letters*, 35, L03 609, <https://doi.org/10.1029/2007GL032644>, 2008.
- Biastoch, A., Böning, C. W., and Lutjeharms, J.: Agulhas leakage dynamics affects decadal variability in Atlantic overturning circulation,
685 *Nature*, 456, 489–492, <https://doi.org/10.1038/nature07426>, 2008.
- Bryan, F.: Parameter Sensitivity of Primitive Equation Ocean General Circulation Models, *Journal of Physical Oceanography*, 17, 970–985, [https://doi.org/10.1175/1520-0485\(1987\)017<0970:PSOPEO>2.0.CO;2](https://doi.org/10.1175/1520-0485(1987)017<0970:PSOPEO>2.0.CO;2), 1987.
- Caldwell, P., Merrifield, M., and Thompson, P.: Sea level measured by tide gauges from global oceans—the Joint Archive for Sea Level holdings (NCEI Accession 0019568), Version 5.5., NOAA National Centers for Environmental Information, Dataset,
690 <https://doi.org/10.7289/V5V40S7W>, 2015.
- Cummins, P. F. and Oey, L. Y.: Simulation of barotropic and baroclinic tides off northern British Columbia, *Journal of Physical Oceanography*, 27, 762–781, [https://doi.org/10.1175/1520-0485\(1997\)027<0762:SOBAPT>2.0.CO;2](https://doi.org/10.1175/1520-0485(1997)027<0762:SOBAPT>2.0.CO;2), 1997.
- Danek, C., Scholz, P., and Lohmann, G.: Effects of high resolution and spinup time on modeled North Atlantic circulation, *Journal of Physical Oceanography*, 49, 1159–1181, <https://doi.org/10.1175/JPO-D-18-0141.1>, 2019.
- 695 Danilov, S., Sidorenko, D., Wang, Q., and Jung, T.: The finite-volume sea ice-Ocean model (FESOM2), *Geoscientific Model Development*, 10, 765–789, <https://doi.org/10.5194/gmd-10-765-2017>, 2017.
- Duffett-Smith, P.: *Astronomy with your Personal Computer*, Cambridge University Press, 2 edn., <https://doi.org/10.1017/CBO9780511564888>, 1990.
- Exarchou, E., Von Storch, J. S., and Jungclaus, J. H.: Impact of tidal mixing with different scales of bottom roughness on the general
700 circulation, *Ocean Dynamics*, 62, 1545–1563, <https://doi.org/10.1007/s10236-012-0573-1>, 2012.
- Falahat, S. and Nycander, J.: On the generation of bottom-trapped internal tides, *Journal of Physical Oceanography*, 45, 526–545, <https://doi.org/10.1175/JPO-D-14-0081.1>, 2015.
- Ferrari, R., Griffies, S. M., Nurser, A. G., and Vallis, G. K.: A boundary-value problem for the parameterized mesoscale eddy transport, *Ocean Modelling*, 32, 143–156, <https://doi.org/10.1016/j.ocemod.2010.01.004>, 2010.
- 705 Ferrari, R., Mashayek, A., McDougall, T. J., Nikurashin, M., and Campin, J. M.: Turning ocean mixing upside down, *Journal of Physical Oceanography*, 46, 2239–2261, <https://doi.org/10.1175/JPO-D-15-0244.1>, 2016.

- Friedrich, T., Timmermann, A., Decloedt, T., Luther, D. S., and Mouchet, A.: The effect of topography-enhanced diapycnal mixing on ocean and atmospheric circulation and marine biogeochemistry, *Ocean Modelling*, 39, 262–274, <https://doi.org/10.1016/j.ocemod.2011.04.012>, 2011.
- 710 Gent, P. R. and McWilliams, J. C.: Isopycnal Mixing in Ocean Circulation Models, *Journal of Physical Oceanography*, 20, 150–155, [https://doi.org/10.1175/1520-0485\(1990\)020<0150:imiocm>2.0.co;2](https://doi.org/10.1175/1520-0485(1990)020<0150:imiocm>2.0.co;2), 1990.
- Griffies, S. M., Levy, M., Adcroft, A. J., Danabasoglu, G., Hallberg, R. W., Jacobsen, D., Large, W., and Ringler, T.: Theory and numerics of the Community Ocean Vertical Mixing (CVMIX) project, Tech. rep., 2015.
- Hibler, W. D.: A Dynamic Thermodynamic Sea Ice Model, *Journal of Physical Oceanography*, 9, 815–846, [https://doi.org/10.1175/1520-0485\(1979\)009<0815:ADTSIM>2.0.CO;2](https://doi.org/10.1175/1520-0485(1979)009<0815:ADTSIM>2.0.CO;2), 1979.
- 715 Huang, R. X.: Energetics of the oceanic circulation, in: *Ocean Circulation: Wind-Driven and Thermohaline Processes*, p. 149–258, Cambridge University Press, Cambridge, <https://doi.org/10.1017/CBO9780511812293.004>, 2009.
- Itoh, S., Tanaka, Y., Osafune, S., Yasuda, I., Yagi, M., Kaneko, H., Konda, S., Nishioka, J., and N. Volkov, Y.: Direct breaking of large-amplitude internal waves in the Urup Strait, *Progress in Oceanography*, 126, 109–120, <https://doi.org/10.1016/j.pocean.2014.04.014>, 2014.
- 720 Jayne, S. R.: The impact of abyssal mixing parameterizations in an ocean general circulation model, *Journal of Physical Oceanography*, 39, 1756–1775, <https://doi.org/10.1175/2009JPO4085.1>, 2009.
- Jayne, S. R. and St. Laurent, L. C.: Parameterizing tidal dissipation over rough topography, *Geophysical Research Letters*, 28, 811–814, <https://doi.org/10.1029/2000GL012044>, 2001.
- Juricke, S., Danilov, S., Kutsenko, A., and Oliver, M.: Ocean kinetic energy backscatter parametrizations on unstructured grids: Impact on mesoscale turbulence in a channel, *Ocean Modelling*, 138, 51–67, <https://doi.org/10.1016/j.ocemod.2019.03.009>, 2019.
- 725 Kantha, L. H.: Barotropic tides in the global oceans from a nonlinear tidal model assimilating altimetric tides 1. Model description and results, *Journal of Geophysical Research*, 100, 25 283, <https://doi.org/10.1029/95jc02578>, 1995.
- Koch-Larrouy, A., Madec, G., Bouruet-Aubertot, P., Gerkema, T., Bessières, L., and Molcard, R.: On the transformation of Pacific Water into Indonesian Throughflow Water by internal tidal mixing, *Geophysical Research Letters*, 34, L04604, <https://doi.org/10.1029/2006GL028405>, 2007.
- 730 Large, W. G. and Yeager, S. G.: The global climatology of an interannually varying air–sea flux data set, *Climate Dynamics*, 33, 341–364, <https://doi.org/10.1007/s00382-008-0441-3>, 2009.
- Large, W. G., McWilliams, J. C., and Doney, S. C.: Oceanic vertical mixing: A review and a model with a nonlocal boundary layer parameterization, *Reviews of Geophysics*, 32, 363–403, <https://doi.org/10.1029/94RG01872>, 1994.
- 735 Large, W. G., Danabasoglu, G., Doney, S. C., and McWilliams, J. C.: Sensitivity to Surface Forcing and Boundary Layer Mixing in a Global Ocean Model: Annual-Mean Climatology, *Journal of Physical Oceanography*, 27, 2418–2447, [https://doi.org/10.1175/1520-0485\(1997\)027<2418:STSFAB>2.0.CO;2](https://doi.org/10.1175/1520-0485(1997)027<2418:STSFAB>2.0.CO;2), 1997.
- Ledwell, J. R., Montgomery, E. T., Polzin, K. L., St. Laurent, L. C., Schmitt, R. W., and Toole, J. M.: Evidence for enhanced mixing over rough topography in the abyssal ocean, *Nature*, 403, 179–182, <https://doi.org/10.1038/35003164>, 2000.
- 740 Lee, H. C., Rosati, A., and Spelman, M. J.: Barotropic tidal mixing effects in a coupled climate model: Oceanic conditions in the Northern Atlantic, *Ocean Modelling*, 11, 464–477, <https://doi.org/10.1016/j.ocemod.2005.03.003>, 2006.
- Lemarié, F., Debreu, L., Madec, G., Demange, J., Molines, J. M., and Honnorat, M.: Stability constraints for oceanic numerical models: Implications for the formulation of time and space discretizations, *Ocean Modelling*, 92, 124–148, <https://doi.org/10.1016/j.ocemod.2015.06.006>, 2015.

- 745 Li, Z., von Storch, J. S., and Müller, M.: The M2 internal tide simulated by a $1/10^\circ$ OGCM, *Journal of Physical Oceanography*, 45, 3119–3135, <https://doi.org/10.1175/JPO-D-14-0228.1>, 2015.
- Locarnini, M., Mishonov, A., Baranova, O., Boyer, T., Zweng, M., Garcia, H., Seidov, D., Weathers, K., Paver, C., and Smolyar, I.: *World Ocean Atlas 2018, Volume 1: Temperature*, NOAA Atlas NESDIS 82, 1, 1–20, 2018.
- Logemann, K., Linardakis, L., Korn, P., and Schrum, C.: Global tide simulations with ICON-O: testing the model performance on highly
750 irregular meshes, *Ocean Dynamics*, 71, 43–57, <https://doi.org/10.1007/s10236-020-01428-7>, 2021.
- Lohmann, G.: Temperatures from energy balance models: the effective heat capacity matters, *Earth System Dynamics*, 11, 1195–1208, <https://doi.org/10.5194/esd-11-1195-2020>, 2020.
- Lohmann, G., Butzin, M., Eissner, N., Shi, X., and Stepanek, C.: Abrupt Climate and Weather Changes Across Time Scales, *Paleoceanography and Paleoclimatology*, 35, <https://doi.org/10.1029/2019PA003782>, 2020.
- 755 Marshall, J. and Speer, K.: Closure of the meridional overturning circulation through Southern Ocean upwelling, *Nature Geoscience*, 5, 171–180, <https://doi.org/10.1038/ngeo1391>, 2012.
- Melet, A., Legg, S., and Hallberg, R.: Climatic impacts of parameterized local and remote tidal mixing, *Journal of Climate*, 29, 3473–3500, <https://doi.org/10.1175/JCLI-D-15-0153.1>, 2016.
- Monterey, G. and Levitus, S.: *Climatological cycle of mixed layer depth in the world ocean*, US government printing office, NOAA NESDIS,
760 Washington, DC, 1997.
- Müller, M., Haak, H., Jungclaus, J. H., Sündermann, J., and Thomas, M.: The effect of ocean tides on a climate model simulation, *Ocean Modelling*, 35, 304–313, <https://doi.org/10.1016/j.ocemod.2010.09.001>, 2010.
- Munk, W. and Wunsch, C.: Abyssal recipes II: Energetics of tidal and wind mixing, *Deep-Sea Research Part I: Oceanographic Research Papers*, 45, 1977–2010, [https://doi.org/10.1016/S0967-0637\(98\)00070-3](https://doi.org/10.1016/S0967-0637(98)00070-3), 1998.
- 765 Nagai, T. and Hibiya, T.: Internal tides and associated vertical mixing in the Indonesian Archipelago, *Journal of Geophysical Research: Oceans*, 120, 3373–3390, <https://doi.org/10.1002/2014JC010592>, 2015.
- Nakamura, T., Isoda, Y., Mitsudera, H., Takagi, S., and Nagasawa, M.: Breaking of unsteady lee waves generated by diurnal tides, *Geophysical Research Letters*, 37, <https://doi.org/10.1029/2009GL041456>, 2010.
- Niwa, Y. and Hibiya, T.: Estimation of baroclinic tide energy available for deep ocean mixing based on three-dimensional global numerical
770 simulations, *Journal of Oceanography*, 67, 493–502, <https://doi.org/10.1007/s10872-011-0052-1>, 2011.
- Niwa, Y. and Hibiya, T.: Generation of baroclinic tide energy in a global three-dimensional numerical model with different spatial grid resolutions, *Ocean Modelling*, 80, 59–73, <https://doi.org/10.1016/j.ocemod.2014.05.003>, 2014.
- Oka, A. and Niwa, Y.: Pacific deep circulation and ventilation controlled by tidal mixing away from the sea bottom, *Nature Communications*, 4, 2419, <https://doi.org/10.1038/ncomms3419>, 2013.
- 775 Olbers, D. and Eden, C.: A global model for the diapycnal diffusivity induced by internal gravity waves, *Journal of Physical Oceanography*, 43, 1759–1779, <https://doi.org/10.1175/JPO-D-12-0207.1>, 2013.
- Osborn, T. R.: Estimates of the Local Rate of Vertical Diffusion from Dissipation Measurements, *Journal of Physical Oceanography*, 10, 83–89, [https://doi.org/10.1175/1520-0485\(1980\)010<0083:EOTLRO>2.0.CO;2](https://doi.org/10.1175/1520-0485(1980)010<0083:EOTLRO>2.0.CO;2), 1980.
- Pawlowicz, R., Beardsley, B., and Lentz, S.: Classical tidal harmonic analysis including error estimates in MATLAB using TIDE, *Computers and Geosciences*, 28, 929–937, [https://doi.org/10.1016/S0098-3004\(02\)00013-4](https://doi.org/10.1016/S0098-3004(02)00013-4), 2002.
- 780 Polzin, K. L., Toole, J. M., Ledwell, J. R., and Schmitt, R. W.: Spatial variability of turbulent mixing in the abyssal ocean, *Science*, 276, 93–96, <https://doi.org/10.1126/science.276.5309.93>, 1997.

- Prange, M., Lohmann, G., and Paul, A.: Influence of Vertical Mixing on the Thermohaline Hysteresis: Analyses of an OGCM, *Journal of Physical Oceanography*, 33, 1707–1721, [https://doi.org/10.1175/1520-0485\(2003\)033<1707:iovmot>2.0.co;2](https://doi.org/10.1175/1520-0485(2003)033<1707:iovmot>2.0.co;2), 2003.
- 785 Ray, R. D.: Ocean self-attraction and loading in numerical tidal models, *Marine Geodesy*, 21, 181–192, <https://doi.org/10.1080/01490419809388134>, 1998.
- Redi, M. H.: Oceanic isopycnal mixing by coordinate rotation, *Journal of Physical Oceanography*, 12, 1154–1158, [https://doi.org/10.1175/1520-0485\(1982\)012<1154:OIMBCR>2.0.CO;2](https://doi.org/10.1175/1520-0485(1982)012<1154:OIMBCR>2.0.CO;2), 1982.
- Saenko, O. A. and Merryfield, W. J.: On the effect of topographically enhanced mixing on the global ocean circulation, *Journal of Physical*
790 *Oceanography*, 35, 826–834, <https://doi.org/10.1175/JPO2722.1>, 2005.
- Sasaki, H., Kida, S., Furue, R., Nonaka, M., and Masumoto, Y.: An Increase of the Indonesian Throughflow by Internal Tidal Mixing in a High-Resolution Quasi-Global Ocean Simulation, *Geophysical Research Letters*, 45, 8416–8424, <https://doi.org/10.1029/2018GL078040>, 2018.
- Schiller, A.: Effects of explicit tidal forcing in an OGCM on the water-mass structure and circulation in the Indonesian throughflow region,
795 *Ocean Modelling*, 6, 31–49, [https://doi.org/10.1016/S1463-5003\(02\)00057-4](https://doi.org/10.1016/S1463-5003(02)00057-4), 2004.
- Scholz, P., Sidorenko, D., Gurses, O., Danilov, S., Koldunov, N., Wang, Q., Sein, D., Smolentseva, M., Rakowsky, N., and Jung, T.: Assessment of the Finite-volumE Sea ice-Ocean Model (FESOM2.0) - Part 1: Description of selected key model elements and comparison to its predecessor version, *Geoscientific Model Development*, 12, 4875–4899, <https://doi.org/10.5194/gmd-12-4875-2019>, 2019.
- Scholz, P., Sidorenko, D., Danilov, S., Wang, Q., Koldunov, N., Sein, D., and Jung, T.: Assessment of the Finite-VolumE Sea ice–Ocean Model
800 (FESOM2.0) – Part 2: Partial bottom cells, embedded sea ice and vertical mixing library CVMix, *Geoscientific Model Development*, 15, 335–363, <https://doi.org/10.5194/gmd-15-335-2022>, 2022.
- Scott, J. R. and Marotzke, J.: The Location of Diapycnal Mixing and the Meridional Overturning Circulation, *Journal of Physical Oceanography*, 32, 3578–3595, [https://doi.org/10.1175/1520-0485\(2002\)032<3578:TLODMA>2.0.CO;2](https://doi.org/10.1175/1520-0485(2002)032<3578:TLODMA>2.0.CO;2), 2002.
- Shchepetkin, A. F.: An adaptive, Courant-number-dependent implicit scheme for vertical advection in oceanic modeling, *Ocean Modelling*,
805 91, 38–69, <https://doi.org/10.1016/j.ocemod.2015.03.006>, 2015.
- Shi, X. and Lohmann, G.: Simulated response of the mid-Holocene Atlantic meridional overturning circulation in ECHAM6-FESOM/MPIOM, *Journal of Geophysical Research: Oceans*, 121, 6444–6469, <https://doi.org/10.1002/2015JC011584>, 2016.
- Shihora, L., Sulzbach, R., Dobslaw, H., and Thomas, M.: Self-attraction and loading feedback on ocean dynamics in both shallow water equations and primitive equations, *Ocean Modelling*, 169, 101 914, <https://doi.org/10.1016/j.ocemod.2021.101914>, 2022.
- 810 Shriver, J. F., Arbic, B. K., Richman, J. G., Ray, R. D., Metzger, E. J., Wallcraft, A. J., and Timko, P. G.: An evaluation of the barotropic and internal tides in a high-resolution global ocean circulation model, *Journal of Geophysical Research: Oceans*, 117, <https://doi.org/10.1029/2012JC008170>, 2012.
- Sidorenko, D., Danilov, S., Koldunov, N., Scholz, P., and Wang, Q.: Simple algorithms to compute meridional overturning and barotropic streamfunctions on unstructured meshes, *Geoscientific Model Development*, 13, 3337–3345, <https://doi.org/10.5194/gmd-13-3337-2020>,
815 2020.
- Simmons, H. L., Jayne, S. R., St. Laurent, L. C., and Weaver, A. J.: Tidally driven mixing in a numerical model of the ocean general circulation, *Ocean Modelling*, 6, 245–263, [https://doi.org/10.1016/S1463-5003\(03\)00011-8](https://doi.org/10.1016/S1463-5003(03)00011-8), 2004.
- Song, P. and Chen, X.: Investigation of the Internal Tides in the Northwest Pacific Ocean Considering the Background Circulation and Stratification, *Journal of Physical Oceanography*, 50, 3165–3188, <https://doi.org/10.1175/JPO-D-19-0177.1>, 2020.

- 820 Song, P., Sidorenko, D., Scholz, P., Thomas, M., and Lohmann, G.: The tidal effects in the Finite-volume Sea ice–Ocean Model (FESOM2.1): a comparison between parameterized tidal mixing and explicit tidal forcing, <https://doi.org/10.5281/zenodo.6700113>, 2021.
- St. Laurent, L. C., Simmons, H. L., and Jayne, S. R.: Estimating tidally driven mixing in the deep ocean, *Geophysical Research Letters*, 29, 21–1–21–4, <https://doi.org/10.1029/2002GL015633>, 2002.
- Steele, M., Morley, R., and Ermold, W.: PHC: A Global Ocean Hydrography with a High-Quality Arctic Ocean, *Journal of Climate*, 14, 2079–2087, [https://doi.org/10.1175/1520-0442\(2001\)014<2079:PAGOHW>2.0.CO;2](https://doi.org/10.1175/1520-0442(2001)014<2079:PAGOHW>2.0.CO;2), 2001.
- 825 Stepanov, V. N. and Hughes, C. W.: Parameterization of ocean self-attraction and loading in numerical models of the ocean circulation, *Journal of Geophysical Research: Oceans*, 109, <https://doi.org/10.1029/2003jc002034>, 2004.
- Sulzbach, R., Dobslaw, H., and Thomas, M.: High-Resolution Numerical Modeling of Barotropic Global Ocean Tides for Satellite Gravimetry, *Journal of Geophysical Research: Oceans*, 126, <https://doi.org/10.1029/2020JC017097>, 2021.
- 830 Talley, L. D., Pickard, G. L., Emery, W. J., and Swift, J. H.: Chapter S14 - Global Circulation and Water Properties: Supplementary Materials, in: *Descriptive Physical Oceanography (Sixth Edition)*, pp. 1–9, Academic Press, Boston, 6 edn., <https://doi.org/10.1016/B978-0-7506-4552-2.10026-5>, 2011.
- Tanaka, T., Yasuda, I., Tanaka, Y., and Carter, G. S.: Numerical study on tidal mixing along the shelf break in the Green Belt in the south-eastern Bering Sea, *Journal of Geophysical Research: Oceans*, 118, 6525–6542, <https://doi.org/10.1002/2013JC009113>, 2013.
- 835 Tanaka, Y., Hibiya, T., Niwa, Y., and Iwamae, N.: Numerical study of K1 internal tides in the Kuril straits, *Journal of Geophysical Research: Oceans*, 115, C09 016, <https://doi.org/10.1029/2009JC005903>, 2010.
- Tatebe, H., Tanaka, Y., Komuro, Y., and Hasumi, H.: Impact of deep ocean mixing on the climatic mean state in the Southern Ocean, *Scientific Reports*, 8, 14 479, <https://doi.org/10.1038/s41598-018-32768-6>, 2018.
- Thomas, M., Sündermann, J., and Maier-Reimer, E.: Consideration of ocean tides in an OGCM and impacts on subseasonal to decadal polar motion excitation, *Geophysical Research Letters*, 28, 2457–2460, <https://doi.org/10.1029/2000GL012234>, 2001.
- 840 Tsujino, H., Hasumi, H., and Suginohara, N.: Deep Pacific Circulation Controlled by Vertical Diffusivity at the Lower Thermocline Depths, *Journal of Physical Oceanography*, 30, 2853–2865, [https://doi.org/10.1175/1520-0485\(2001\)031<2853:DPCCBV>2.0.CO;2](https://doi.org/10.1175/1520-0485(2001)031<2853:DPCCBV>2.0.CO;2), 2000.
- Wahr, J. M.: Body tides on an elliptical, rotating, elastic and oceanless earth, *Geophysical Journal of the Royal Astronomical Society*, 64, 677–703, <https://doi.org/10.1111/j.1365-246X.1981.tb02690.x>, 1981.
- 845 Wang, Q., Danilov, S., Sidorenko, D., Timmermann, R., Wekerle, C., Wang, X., Jung, T., and Schröter, J.: The Finite Element Sea Ice–Ocean Model (FESOM) v.1.4: Formulation of an ocean general circulation model, *Geoscientific Model Development*, 7, 663–693, <https://doi.org/10.5194/gmd-7-663-2014>, 2014.
- Weber, T. and Thomas, M.: Tidal dynamics and their influence on the climate system from the Cretaceous to present day, *Global and Planetary Change*, 158, 173–183, <https://doi.org/10.1016/j.gloplacha.2017.09.019>, 2017.
- 850 Yu, Y., Liu, H., and Lan, J.: The influence of explicit tidal forcing in a climate ocean circulation model, *Acta Oceanologica Sinica*, 35, 42–50, <https://doi.org/10.1007/s13131-016-0931-9>, 2016.
- Zweng, M., Seidov, D., Boyer, T., Locarnini, M., Garcia, H., Mishonov, A., Baranova, O., Weathers, K., Paver, C., and Smolyar, I.: *World Ocean Atlas 2018, Volume 2: Salinity*, NOAA Atlas NESDIS, 1, 1–20, 2018.



Co-tidal charts of M2 tides from the timestep sensitivity runs. The colours indicate tide amplitudes, and the white lines indicate Greenwich phase lags with an interval of 60° . The timestep setting in each run is labelled on each panel.

(a) RMSE of the four main tidal components (O1, K1, M2, and S2 tides) for all 229 tide gauges; (b) In the sensitivity run with timestep set to 720, RMSE of main diurnal tidal components (O1 and K1 tides) at each tide-gauging station. Scatter points are classified into four levels by the magnitudes of RMSE; (c) The same as panel (b) but for main semidiurnal tidal components (M2 and S2 tides).

Figure 1. Depth averaged temperature biases between the model results and WOA18 in the upper (0–500 m), intermediate (500–1500 m), and deep (1500–5000 m) ocean. The left column shows the biases between NOTIDE and WOA18, while the middle/right column shows the difference between CVTIDE/LSTIDE and NOTIDE. Opposite colours in the left column and the middle/right column indicate CVTIDE/LSTIDE reduces temperature biases in NOTIDE.




Figure 2. The same as Fig. 1, but for salinity.




Figure 3. Basin averaged hydrographic biases between the model results and WOA18 with respect to water depths. The hydrographic biases are absolute values. The left and the right column represent temperature and salinity biases, while each row shows the biases in different basins. **Note that the [The](#)** averaging is nodal area-weighted.




Figure 4. Bathymetry averaged hydrographic biases between the model results and WOA18 with respect to water depths. The hydrographic biases are absolute values, and model mesh grids with the same bathymetry (maximum water depth) are assembled and averaged. The middle and lower rows are presented as the difference with the upper row; thus the blue/red colours indicate tidal effects are decreasing/increasing model biases in the control run. ~~Note that the~~ The averaging is nodal area-weighted, and the axes are scaled with cube root.

Arcic-sea ice thickness in the three sensitivity runs. The left column shows the result from the control run, while the middle/right column shows the biases between CVTIDE/LSTIDE and NOTIDE. The upper and lower rows represent the model results in March and September, respectively. Note that the grey backgrounds mask the ice-free areas, and the grey dashed line represents the 500 isobath.

The same as Fig. ??, but for Antarctica. The grey dashed line represents the 1000 isobath.

Mixed layer depths defined in Large et al. (1997). The left column represents the results from the control run, while the middle/right column represents the difference between CVTIDE/LSTIDE and NOTIDE. The upper and lower rows represent the model results in March and September, respectively.




Figure 5. Polar sea ice thickness in the three sensitivity runs. The same as Fig. ?? but for polar projection. The upper and lower rows show the Arctic (March) and Antarctic (September) results. Here, the grey backgrounds mask the ice-free areas, and the grey dashed line represents the 500/1000 m isobath for the upper/lower panels. The mixed layer depths are defined after Monterey and Levitus (1997).




Figure 6. The MOC results in the three sensitivity runs. The upper panel shows the result from the control run, while the other two panels show the difference between CVTIDE/LSTIDE and NOTIDE differences. The left column and right column represent AMOC and PMOC, respectively. The brown dashed line in panel (f) denotes the North Pacific topography along the 180° transection. Note that All the MOC contains the vertical component of bolus velocities (Gent and McWilliams, 1990). The calculation of MOC in the unstructured mesh applies the algorithm introduced in Sidorenko et al. (2020).




Figure 7. Depth-averaged vertical diffusivity in the upper (0–500 m), intermediate (500–1500 m), deep (1500–5000 m), and the bottom layer of the ocean. The left column shows the global mean vertical diffusivity profile in the control run, while the middle/right column shows the difference between CVTIDE/LSTIDE and NOTIDE three sensitivity runs. Note that vertical diffusivity is shown in a decimal logarithmic scale, and the middle/right column shows the difference of scaled vertical diffusivity logarithmic scales.




Figure 8. The barotropic streamfunction in the three sensitivity runs. The upper panel shows the result in the control run, while the middle/lower panel shows the difference-between-CVTIDE/LSTIDE-and-NOTIDE differences. Note that the The lower panel also represents the tidal residual mean circulation. The calculation of barotropic streamfunction in the unstructured mesh applies the algorithm introduced in Sidorenko et al. (2020).




Figure 9. Horizontal distribution of vertically integrated energy diagnostic dissipation terms in Table 1 (we change the sign so that positive values represent energy dissipation). The upper, middle, and lower rows indicate viscous dissipation, bottom drag, and buoyancy flux, respectively viscous dissipation (including horizontal and vertical components). The left column shows the control run, and the middle and right columns show the differences between CVTIDE/LSTIDE and NOTIDE differences. Note that the upper and middle rows Colorbars are shown in a decimal logarithm scale, which is the same as Fig. 7 logarithmic scales.




Figure 10. The ratio Horizontal distribution of mode-1 internal tide wavelength and mesh resolution vertically integrated buoyancy flux terms. Buoyancy flux in this figure is expressed as $g\rho w$, while turbulent buoyancy flux is $\overline{g\rho'w'}$ (positive values represent upwelling). The upper left column shows the control run, and lower panels represent the ratio for K1 middle and M2 internal tides, respectively. Note here that right columns show the wavelengths are calculated by solving a Sturm-Liouville eigenvalue problem, which can be referred to in Song and Chen (2020, their Eq. 16) differences.




Figure 11. The vertical diffusivity, the turbulent buoyancy flux ratio of mode-1 K1/M2 internal tide scale and its equivalent vertical diffusivity along mesh resolution. The thick grey dashed line indicates the transection-critical latitudes of 65°E internal tides. The calculation of turbulent buoyancy flux and the equivalent vertical diffusivity internal tide scale is introduced in Appendix ???. All panels are shown in a decimal logarithm scale B3.




Figure 12. The time series of (a) 40° N PMOC, (b) ITF, (c) 26° N AMOC, (d) 20° S AMOC, (e) ACC through the Drake Passage from the three sensitivity runs in all five cycles. The light and thin lines are original annual time series, and the dark and thick lines are five-year moving average results for each cycle. All vertical axes have the unit of Sv. Note that 26° N AMOC and 20° S AMOC denote the strength of AMOC upper cell and lower cell, respectively. The calculation of ITF and ACC flux through the transections in the Indonesian Archipelago and the Drake Passage applies the algorithm introduced in Sidorenko et al. (2020).




Figure 13. The GMOC and potential density with reference depth at 4000 m (σ_4) in the southern hemisphere. The upper panels denote results in the control run, while the middle/lower panels denote the difference between CVTIDE/LSTIDE and NOTIDE differences. Note that the The potential density in the figure is globally zonal averaged, with meridional bin set to 1°.

Table 1. Basic information—A global integration of kinetic energy budget from the model results. The surface wind stress, bottom stress, vertical viscous dissipation, horizontal viscous dissipation, buoyancy flux, tidal potential power, the barotropic potential energy–kinetic energy conversion term, and total kinetic energy are in order listed in the table. The expressions of these terms can be found in Appendix B1. All except the sensitivity runs last term have a unit of TW, while kinetic energy has a unit of 10^{18} J.

Experiment ID Vertical mixing scheme Tidal motions Description NOTIDE KPP False FESOM2 control run CVTIDE KPP+CVMIX_TIDAL False FESOM2 considering tide-induced mixing param. LSTIDE KPP True FESOM2 considering lunisolar tidal motions

Model spin-up strategy for each sensitivity run. Cycle ID R1 R2 R3 R4 R5 Forcing year 1948–2009 1948–2009 1948–2009 1948–2009 1948–2009 1948–2009 Model year 1–62 63–124 125–186 187–248 249–310

A global integration of kinetic energy budget from the model results. The surface wind stress, bottom stress, vertical viscous dissipation, horizontal viscous dissipation, buoyancy flux, tidal potential power, surface pressure power terms and total kinetic energy are in order listed in the table. The expressions of these terms can be found in the Appendix ???. Note that all except the last term have a unit of 10^{18} W, while kinetic energy has a unit of 10^{18} J.

Experiment ID	wind stress	bott stress	vert visc	horiz visc	buoy flux	tide	surf pressure-PE2KE	KE
NOTIDE	3.11	−0.18	−2.19	−0.17	−0.58	0.00	−0.02	1.30
CVTIDE	3.11	−0.18	−2.19	−0.18	−0.58	0.00	−0.02	1.31
LSTIDE	3.11	−0.85	−2.29	−0.30	−0.58	4.55	−3.51	1.53




Figure A1. Co-tidal charts of M2 tides from the timestep sensitivity runs. The colours indicate tide amplitudes, and the white lines indicate Greenwich phase lags with an interval of 60° . The timestep setting in each run is labelled on each panel.




Figure A2. (a) RMSE of the four main tidal components (O1, K1, M2, and S2 tides) for all 229 tide gauges; (b) In the sensitivity run with timestep set to 720 s, RMSE of main diurnal tidal components (O1 and K1 tides) at each tide-gauging station. Scatter points are classified into four levels by the magnitudes of RMSE; (c) The same as panel (b) but for main semidiurnal tidal components (M2 and S2 tides).