

Objective identification of meteorological fronts: climatologies from ERA-Interim and ERA5

Philip G. Sansom^{1,2} and Jennifer L. Catto¹

¹Faculty of Environment, Science and Economy, University of Exeter, North Park Road, Exeter, EX4 4QE, UK

²Met Office, FitzRoy Road, Exeter, EX1 3PB, UK

Correspondence: Jennifer L. Catto (j.catto@exeter.ac.uk)

Abstract. Meteorological fronts are important for their associated surface impacts, including extreme precipitation and extreme winds. Objective identification of fronts is therefore of interest in both operational weather prediction and research settings. We have implemented a number of changes to a previous implementation of an objective front identification algorithm, applied these to reanalysis datasets, and present the improvements associated with these changes. The previous implementation used a different order of operations when identifying fronts compared to the the original algorithm. First, we show that the originally proposed order of operations yields smoother fronts with fewer breaks. Next, we propose the selection of the front identification thresholds in terms of climatological quantiles of the threshold fields. This allows for comparison between datasets of differing resolutions. Finally, we include a number of numerical improvements in the implementation of the algorithm, such as more accurate finite differencing, direct calculation of the wet-bulb potential temperature, and better handling of short fronts, which yield further benefits in smoothness and number of breaks. This updated version of the algorithm has been made fully portable and scalable to different datasets in order to enable future climatological studies of fronts and their impacts.

1 Introduction

Atmospheric fronts are of great importance for the day-to-day variability of weather in the mid-latitudes. They are associated with a large proportion of total and extreme precipitation, as demonstrated with numerous case studies (Browning, 2004), modelling, and more recently, long-term climatologies (Berry et al., 2011b; Parfitt et al., 2017b; Schemm et al., 2017). They are also strongly linked to extreme wind events (Dowdy and Catto, 2017; Catto et al., 2019; Raveh-Rubin and Catto, 2019; Catto and Dowdy, 2021), and are key for air-sea interaction (Parfitt et al., 2017b). With a wealth of global gridded observationally-constrained and model-produced data, there is a desire to be able to objectively identify these frontal features in the gridded data. This avoids the huge time requirements of a manual analysis, and allows the features to be linked to high impact weather, such as extreme precipitation or winds (Catto et al., 2012; Catto and Pfahl, 2013; Dowdy and Catto, 2017). The application of

the methods to model data of historical and future climate also allows the models to be evaluated for their ability to capture the dynamical features and their connection to precipitation events (Leung et al., 2022), and to investigate the future of such features and how they may impact water resources and natural hazards (Catto et al., 2014).

A number of methods have been developed to perform such objective identification in recent years. Hewson (1998) compiled a summary of methods used to identify frontal features in gridded data, and further developed the methods based on a thermal front parameter. Thomas and Schultz (2019) highlighted the three main factors required in identifying fronts with such a thermal front parameter: first, the thermal variable and vertical level to be considered, e.g., temperature, potential temperature, or equivalent (or wet-bulb) potential temperature at 850 hPa; second, a function of the variable, e.g., the gradient, or some second or third derivative; and finally, some thresholds. They found that different thermal variables each had pros and cons, and could be selected depending on the purpose of the study. The study by Jenkner et al. (2010) used equivalent potential temperature and its second derivative to place the frontal lines. This results in the fronts lying in the centre of a frontal zone, rather than at the leading edge as a synoptic meteorologist would typically put them. Berry et al. (2011b) directly applied the methods of Hewson (1998) to gridded data at $2.5^\circ \times 2.5^\circ$ resolution, placing fronts on the warm side of the strong temperature gradient. This also included the addition of a numerical line-joining algorithm, which is the final piece of the front identification puzzle.

Other methods have used dynamical information to identify fronts. Simmonds et al. (2012) used information solely on wind shifts. This method was found to work better in the Southern Hemisphere than the Northern Hemisphere by Schemm et al. (2015). A combination of this and the thermal method was used by Bitsa et al. (2021) to identify cold fronts in the Mediterranean, with the method tailored to suit the smaller spatial scale of fronts in this region. Parfitt et al. (2017b) used a combination of vorticity and temperature, requiring both a thermal gradient and a wind shift. While each method has its advantages and disadvantages, many of the methods typically identify many of the same features (Hope et al., 2014).

A major difficulty in applying objective front identification is the many datasets and differing resolutions. This is particularly an issue when using gradients of thermal properties, since the resolution of the data will have a large impact on these gradients. The thresholds used to define fronts need to be varied depending on the resolution. For example, the threshold used in Berry et al. (2011b) was $-8 \times 10^{-12} \text{ K m}^{-2}$, while in Dowdy and Catto (2017) on the higher resolution data was $-5 \times 10^{-11} \text{ K m}^{-2}$. Recently, Soster and Parfitt (2022) investigated the sensitivity of results to the use of different datasets and found a large difference in front frequency between the datasets. Higher resolution datasets consistently show higher frequency of frontal points, with the differences reduced when re-gridded to a common grid. This was shown lead to large differences between datasets in the proportion of precipitation attributed to fronts.

Despite the many methods of identifying fronts, and issues and uncertainties associated with each of them, the thermal front parameter method of Hewson (1998) has been successful in identifying the key climatological features of front frequency and the link to other variables in a number of studies (e.g., Berry et al., 2011b, a; Catto et al., 2012; Catto and Pfahl, 2013; Catto et al., 2014; Dowdy and Catto, 2017). The code developed by Berry et al. (2011b) and applied in those subsequent studies was originally developed on the European Centre for Medium-Range Weather Forecasts' (ECMWF) ERA-40 reanalysis (Uppala et al., 2005) at $2.5^\circ \times 2.5^\circ$ resolution, and later to the ECMWF ERA-Interim (Dee et al., 2011) reanalysis at $0.75^\circ \times 0.75^\circ$

resolution. However, that implementation was not easily portable due to being written in a mixture of NCL and Fortran, and would not scale to the ECMWF ERA5 reanalysis at $0.25^\circ \times 0.25^\circ$ or other high resolution datasets. The aim of this study is to create a portable implementation of the front identification method of (Hewson, 1998), as implemented by Berry et al. (2011b), that is able to scale to contemporary high resolution (re-)analyses with horizontal grid-spacings of 0.25° or less. We demonstrate a quantile based method of tuning the thresholds. First the data used are described in Section 2. Section 3 gives a description of the thermal front parameter method, and the improvements over the previous implementation of the algorithm. In Section 4 we compare the front climatology using the new method with previous methods and different datasets. We finish in Section 5 with a discussion of the benefits and challenges associated with such objective identification methods.

2 Data

The updated front identification procedure is applied to the ECMWF ERA-Interim reanalysis (ECMWF Reanalysis - Interim, Dee et al., 2011). The data used here have a resolution of $0.75^\circ \times 0.75^\circ$ on a regular longitude-latitude grid. The 6-hourly instantaneous air temperature and specific humidity fields at the 850 hPa level were used to compute the wet-bulb potential temperature θ_W , using the direct method of Davies-Jones (2008, Equation 3.8), in order to identify fronts. The 6-hourly eastward and northward wind components at 850 hPa were used to compute the front speed using the method of Hewson (1998, Equation 13), allowing classification into cold, warm or quasi-stationary fronts. ERA-Interim was chosen over the more recent ERA5 reanalysis (ECMWF Reanalysis v5, Hersbach et al., 2020) for the primary analysis since the updated procedure is of greatest benefit in middle and low resolution models, and the resolution of ERA-Interim is equal to that of the highest resolution among standard CMIP6 GCMs. Our baseline for comparison is the global climatology of fronts in ERA-Interim at $0.75^\circ \times 0.75^\circ$ produced by Dowdy and Catto (2017) using the method of Berry et al. (2011b). We also present a high resolution climatology based on applying the updated front identification procedure to the ERA5 reanalysis using the same 6-hourly fields as ERA-Interim but with a grid spacing of $0.25^\circ \times 0.25^\circ$.

3 Methodology

Following Hewson (1998) and Berry et al. (2011b), fronts are identified in the wet-bulb potential temperature field θ_W at 850 hPa. As described in Hewson (1998, their Equation 5), and implemented in Berry et al. (2011b), fronts are located as the zero contour of:

$$\nabla \cdot \nabla |\nabla \theta_W| = 0 \quad \text{or} \quad \nabla^2 |\nabla \theta_W| = 0. \quad (1)$$

For a one-dimensional front (Type 1 front in Hewson (1998)), this is simply the third derivative of the wet-bulb potential temperature θ_W (see Figure 3 of Hewson (1998) for an intuitive explanation). We will refer to Equation 1 as the thermal front locator (TFL). In practice, most atmospheric fronts are curved and not simple one-dimensional objects. Hewson (1998) derived an alternative (their Equation 6) to Equation 1, based on the computation of “five-point mean axes”, designed to

mitigate the effects of frontal curvature on the computation of Equation 1, which can lead to noise and exaggerated frontal curvature. Although the alternative definition was preferred by Hewson (1998), we keep the definition in Equation 1 primarily for compatibility with Berry et al. (2011b) and the numerous studies which have utilised that implementation. However, the option to use the alternative definition may be included in a future version of the code documented by this study.

90 Hewson (1998) defined two additional criteria that must be met in order for a zero contour of the Equation 1 to be considered a front. First, the rate of change of θ_W across the front in the direction of cold air must exceed some threshold value K_1 . This criterion was formalised in Equation 9 of Hewson (1998) as:

$$\nabla |\nabla \theta_W| \cdot \frac{\nabla \theta_W}{|\nabla \theta_W|} < K_1, \quad \text{where } K_1 \leq 0 \text{ K m}^{-2}. \quad (2)$$

This is the thermal front parameter (TFP) defined by Renard and Clarke (1965). For a one-dimensional front, this criterion simply states that the second derivative of θ_W must be negative, placing the front on the warm side of the gradient. Second, the gradient of θ_W in the adjacent baroclinic zone (ABZ) must be greater than some threshold value K_2 . This criterion was formalised in Equation 11 of Hewson (1998) as:

$$|\nabla \theta_W|_{ABZ} > K_2, \quad \text{where } K_2 \geq 0 \text{ K m}^{-1}, \quad (3)$$

with

$$100 \quad |\nabla \theta_W|_{ABZ} = |\nabla \theta_W| + m\chi |\nabla |\nabla \theta_W||,$$

where $m = 1/\sqrt{2}$ and χ is the grid length. For a one-dimensional front, this criterion simply states that the magnitude of the gradient of θ_W must be greater than K_2 a fraction m of a grid length in the direction of greatest increase in the gradient of θ_W , i.e., inside the adjacent baroclinic zone. The value of $m = 1/\sqrt{2}$ was suggested by Hewson (1998) and we found it to be effective at the resolution of ERA-Interim (0.75°) and ERA5 (0.25°), but it may require additional tuning in very high resolution data sets.

105 Fronts are identified as warm, cold or quasi-stationary using the front speed defined by Equation 13 of Hewson (1998), which is given here as:

$$\frac{\mathbf{V} \cdot \nabla |\nabla \theta_W|}{|\nabla |\nabla \theta_W||}, \quad (4)$$

where $\mathbf{V} = (u, v)$ is the vector wind field at 850 hPa. Following Berry et al. (2011b), we adopt a threshold of $K_3 = 1.5 \text{ m s}^{-1}$ such that front points are defined as belonging to warm fronts if they have speed exceeding 1.5 m s^{-1} , and as belonging to cold fronts if they have speed less than -1.5 m s^{-1} . All other front points are defined as belonging to quasi-stationary fronts.

The automatic front identification method described by Equations 1–4 has been re-implemented in the R statistical computing language (R Core Team, 2021). The new implementation includes one key methodological change described in Section 3.1, as well as number of numerical updates compared to that of Berry et al. (2011b).

115 3.1 Methodological changes

The intention of this study was to create a portable and scalable implementation of the front identification method of Hewson (1998) as implemented by Berry et al. (2011b), since that implementation has been successfully used in a number of other studies (e.g., Berry et al., 2011a; Catto and Pfahl, 2013; Dowdy and Catto, 2017). However, one key methodological change was implemented regarding the order of operations when locating front objects as lines. Berry et al. (2011b) take what we will
120 call a “mask-then-join” approach. First they locate all those grid boxes that satisfy Equation 2 to form a mask (the criterion in Equation 3 is not used). Zero points of Equation 1 are located by an exhaustive search using linear interpolation between only those grid boxes included in the mask defined Equation 2. Finally, a line-joining algorithm is used to join the zero points of Equation 1 into lines representing fronts. Points are joined to their nearest neighbour if the euclidean distance calculated in degrees of longitude and latitude between two points is less than a specified threshold. This requires the repeated calculation
125 of the distance between the current point and all remaining un-joined points, making the algorithm computationally expensive. Berry et al. (2011b) also apply a minimum front length criteria of 250 km.

In contrast, Hewson (1998) originally proposed a “contour-then-mask” approach, which we adopt here. We identify zero points in the complete field defined by Equation 1 and join them into lines using a contouring algorithm, specifically the `contourLines()` function in R. Zero points are again located by linear interpolation, but only zero points located in adjacent
130 grid boxes are considered for joining into lines, reducing the computational expense compared to an exhaustive search and avoiding the need for repeatedly calculating the distance between large numbers of points. We then interpolate the values of the fields defined by Equations 2 and 3 onto the points located by the contouring algorithm. Only points that meet the criteria defined by Equations 2 and 3 are retained, leaving a set of pre-joined line segments representing fronts.

The two approaches are compared in Figure 1. Zero points in Equation 1 usually occur between grid points. That means
135 that adjacent grid boxes meeting the criteria in Equation 2 are required in order to find zero points using Equation 1 by the mask-then-join approach. At or below the $0.75^\circ \times 0.75^\circ$ resolution of ERA-Interim, the region that satisfies Equation 2 is often narrow, frequently only one grid box wide. Therefore, the mask-then-join approach frequently fails to locate front points. This behaviour can be seen in Figure 1a where no front points are identified between 44.25N and 45.75N since two zonally adjacent grid boxes would be required for successful interpolation of a zero point between two masked points, given the orientation of
140 the front. This may result in some features not being identified at all or, more frequently, gaps in what should be continuous features. The line-joining algorithm used by Berry et al. (2011b) attempts to mitigate this by using a search radius larger than one grid length, but this is only partially effective. In Figure 1a, the search radius is effective in joining the southern-most located point, but fails to bridge the gap between 44.25N and 45.75N to the region between 46.5N and 48.0N where multiple adjacent grid points might once again enable the location of zero points. The number of points located in that northern region
145 is then too small to meet the minimum front length criteria on their own. The contour-then-mask approach originally




Figure 1. Front identification in ERA-Interim at 00:00 on 2001-01-01. (a) Using the mask-then-join approach, and (b) using the contour-then-mask approach. Black contours show $\nabla \cdot \nabla |\nabla \theta_W| = 0$ and red shading indicates regions where $\text{TFP} \leq -5 \times 10^{-11} \text{ K m}^{-2}$ by masking in (a) and interpolation in (b). Circles indicate front points located by each algorithm.

proposed by Hewson (1998) and demonstrated in Figure 1b is able to successfully identify the whole front as a single object. The masked region is shown for illustration only, in practice the masking variables are interpolated directly on to the potential front points located on the zero contour. Overall, the contour-then-mask approach results in more fronts and front points identified, and fewer breaks, as can be seen in the examples in Figures 3a and 3b, and the climatologies in Figures 4b and 4c.

150 The expected decrease in the number of fronts due to there being fewer breaks is compensated by the number of new fronts located due to the increased sensitivity of the contour-then-mask approach to identifying potential front points. In some cases these new fronts were missed completely by the mask-then-join approach, in others they fail to meet the length criteria without additional points located by the contour-then-mask approach.

3.2 Choosing the thresholds and level of smoothing

155 Although automated methods offer the promise of objective feature identification, it is still usually necessary to set some key parameters. For front identification there are three parameters that require tuning: the amount of smoothing applied to the θ_W field, the TFP threshold K_1 , and the gradient threshold K_2 . Some studies have compared outputs with manual analyses by meteorologists to calibrate the parameters. This is difficult, time consuming and calibrates the algorithm to the subjective judgement of a single meteorologist. Also, all three parameters depend on the resolution of the data. Therefore, the calibration
160 must be repeated for each new dataset, or datasets brought to a common resolution for comparison. Instead, we offer some suggestions for objective calibration criteria.

We first address the smoothing problem, since the amount of smoothing applied to θ_W affects the choice of K_1 and K_2 . The purpose of smoothing is to remove local minima and maxima that might break up otherwise continuous features. We particularly wish to avoid local extrema in the TFL field defined by Equation 1, which will appear as short closed contours of
165 $TFL = 0 \text{ K m}^{-3}$. Therefore, it makes sense to examine the effect of smoothing on the average length of these contours. Previous studies applying the method of Berry et al. (2011b) to ERA-Interim used $n = 2$ passes of a simple five-point average to smooth the θ_W field. In testing on ERA-Interim data, it was found that the average length of the contours of $TFL = 0 \text{ K m}^{-3}$ initially increases rapidly with the number of passes of the five-point smoother, but after 6–10 passes, the effect of further smoothing diminishes (see Figure 1 of Supplementary Material). Therefore, we settled on $n = 8$ passes of a five-point smoother.

170 The noise in the TFL field may in part be due to the choice to use Equation 1, as implemented by Berry et al. (2011b), to define the location of the fronts, rather than the method preferred by Hewson (1998, their Equation 6) designed to mitigate the effects of frontal curvature. Equation 1 was retained for its simplicity, and compatibility with Berry et al. (2011b) and subsequent studies. However, the alternative method preferred by Hewson (1998) may be made available as an option in future versions of the code associated with this study. Jenkner et al. (2010) classify all closed contours in the front locating field
175 encircling an area smaller than a given threshold as being associated with (potential) local, rather than synoptic fronts. Such a criterion introduces additional subjectivity, but would effectively reduce the noise when identifying synoptic fronts, possibly allowing less smoothing to be used, and further distinguish fronts associated with orography and other local features. The issue of noise and surface driven gradients was also discussed in Hewson (2001).

It is common to define weather phenomena as events exceeding some percentile of the climatological distribution. Therefore,
180 we propose a quantile based approach to setting the thresholds K_1 and K_2 . The advantage of setting thresholds in terms of climatological quantiles is that the thresholds should be comparable between datasets of differing resolution, while the actual values can differ quite widely. For example, Berry et al. (2011b) used a threshold of $-8 \times 10^{-12} \text{ K m}^{-2}$ at 2.5° resolution in ERA-40, compared to the threshold of $-5 \times 10^{-11} \text{ K m}^{-2}$ at 0.75° resolution used in ERA-Interim by Dowdy and Catto (2017). In order to compute quantiles, we require climatologies of the TFP and the magnitude of the gradient, obtained by evaluating
185 Equations 2 and 3, respectively, over an extended time period for the region of interest. The time period considered was 1979–2018 in ERA-Interim. Since most fronts occur in the extra-tropical regions, we will focus our attention there. We seek quantiles of the TFP and the magnitude of the gradient that produce continuous fronts in good agreement with published charts for the




Figure 2. Choosing the thresholds K_1 and K_2 . (a) The 25th percentile of the TFP, and (b) the 50th percentile of $|\nabla\theta_W|$ by latitude and month of the year. Each coloured line represents a different month: blue for DJF, yellow for MAM, red for JJA, and orange for SON. Horizontal dotted lines represent the major circles of latitude. Vertical dashed lines indicate the thresholds chosen in the text: $-1.6 \times 10^{-11} \text{ K m}^{-2}$ in (a); and $7.5 \times 10^{-6} \text{ K m}^{-1}$ in (b).

North Atlantic and Europe, focusing on January and July 2020. Combinations of quantiles of both the TFP and the magnitude of the gradient were systematically compared (see Supplementary Material for examples). We set the first threshold K_1 to the 25th percentile (0.25 quantile) of the climatological distribution of the TFP. In the Northern Hemisphere extra-tropics this is around $-1.6 \times 10^{-11} \text{ K m}^{-2}$. We set the second threshold K_2 equal to the 50th percentile (0.50 quantile) of the climatological distribution of the magnitude of the gradient of θ_W . In the Northern Hemisphere extra-tropics (23.4°N – 66.6°N) this is around $7.5 \times 10^{-6} \text{ K m}^{-1}$. These choices are subjective, and an operational meteorologist might make other choices. However, in the absence of strong physical reasoning, these quantiles have a simple symmetry, i.e., each is approximately the 50th percentile of the allowed range (since $K_1 < 0 \text{ K m}^{-2}$ and globally the 50th percentile of TFP is approximately 0 K m^{-2}), and produce continuous fronts in good agreement with published charts.

Figure 2 illustrates the monthly and latitudinal climatological variation in the chosen quantiles of TFP and the magnitude of the gradient of θ_W . The distributions of both TFP and the magnitude of the gradient are very different in the tropics compared to the extra-tropics. The value of TFP chosen for K_1 is biased towards the upper latitudes of the northern hemisphere extra-tropics where fronts are frequently observed and associated with extra-tropical cyclones. The chosen value of the magnitude of gradient lies in the middle of the seasonal variation in the extra-tropics, which is fairly constant between around 35°N–65°N and 30°S–50°S, with greater spread in the northern hemisphere. The chosen values are broadly representative of the quantiles across the seasons in both the northern and southern hemisphere extra-tropics. Given the relative insensitivity to reasonable values of the K_1 and K_2 shown in the Appendix, the chosen values should be representative across the seasons and both hemispheres for both criteria.

For comparison, previous studies applying the method of Berry et al. (2011b) to ERA-Interim used a threshold of $K_1 = -5 \times 10^{-11} \text{ K m}^{-2}$ after $n = 2$ smoothing passes. The second threshold K_2 in Equation 3 was not implemented by Berry et al. (2011b), equivalent to setting $K_2 = 0 \text{ K m}^{-1}$ since $|\nabla\theta_W| \geq 0$ by definition. Our threshold K_1 is higher primarily due to the additional smoothing, but the exclusion of the second threshold K_2 may have caused Berry et al. (2011b) to choose a lower threshold for K_1 in order to remove unwanted features that could more effectively have been eliminated by implementing the second threshold K_2 .

3.3 Comparing fronts from different datasets

When comparing analyses from different weather and climate datasets, the most common approach is to interpolate all the datasets to a common resolution, usually the lowest resolution among them. For some features such as fronts that are more easily identified in higher resolution data, this can be limiting. The objective calibration criteria described in Section 3.2 provide one route by which fronts could be identified at the native resolution of each dataset and then compared. The quantile based criteria will identify the same fraction of grid boxes potentially containing front points for any reasonable resolution and number of smoothing passes. However, computing the required climatologies is time consuming. An alternative is to keep the thresholds K_1 and K_2 constant, and adjust the number of smoothing passes such that the climatological distributions of the TFP and the magnitude of the gradient are similar between datasets. Specifically, the quantiles used to set the thresholds should be similar. In testing, it was found that matching the threshold quantile of the TFP field provided a more consistent comparison than that of the gradient field. It is sufficient to compare the quantiles for only a small subset of the data, provided the same subset is used for each dataset, avoiding the need to compute a long climatology in order to determine. In testing, various lengths and spatial extents of training data were considered for comparing ERA-Interim and ERA5, from one month, to 30 years, for the Northern Hemisphere extra-tropics, Southern Hemisphere extra-tropics, or the whole globe. One month of data was found to be sufficient to consistently determine an appropriate number of smoothing passes. The procedure is not sensitive to either the month of the year or the spatial extent, among those considered. In practice, we used January 2000 for the Northern Hemisphere extra-tropics, consistent with the examples in Figures 1 and 3.

3.4 Numerical updates

230 Berry et al. (2011b) used repeated applications of a simple central finite difference approximation to the first derivative to evaluate all the derivatives in Equations 1–4 at each grid box. The simple approximation uses one grid box on either side of the box in question to approximate the first derivative to second order accuracy. The zonal and meridional components of the derivatives are evaluated separately using one box to the left and right, or above and below, respectively. However, repeated applications of the approximation to the first derivative degrades the accuracy for higher derivatives. In contrast, we use an
235 explicit central finite difference approximation to the second derivatives required to evaluate ∇^2 when computing the TFL in Equation 1, avoiding the need for repeated applications of the first derivative, and maintaining second order accuracy. The explicit approximation uses two grid boxes either side of the grid box in question. The zonal and meridional components are evaluated separately. The computation of both the first and second derivatives was also updated to maintain second order accuracy at the edges of the domain using forward and backward differences. The increased accuracy comes at minimal
240 computation cost, but is valuable due to the small scales of the quantities of interest, i.e., the thresholds K_1 and K_2 are very small, order 10^{-11} and 10^{-6} , indicative of scale of the fields of TFP and $|\nabla\theta_W|$ themselves, therefore accurate and stable numerical schemes are required to accurately locate fronts.

Other numerical differences include using the mixed-phase parametrization of relative humidity from the ECMWF Integrated Forecasting System (ECMWF, Section 7.4.2) rather than the table based approach from the NCAR Command Language (NCL,
245 2019). In the new implementation, the wet-bulb potential temperature (θ_W) is computed using the direct method of Davies-Jones (2008, Equation 3.8), rather than an iterative method. The final numerical difference between the two implementations is how short fronts are handled. In the original application, Berry et al. (2011b) reject any fronts less than three points long. In later applications this was updated to a great-circle distance based criterion where fronts whose end points are less than 250 km apart are rejected. In our implementation, we sum the great-circle distance between all adjacent points in each front and reject
250 fronts whose total length is less than 250 km.

4 Results

4.1 Comparison with previous implementations

Figure 3 illustrates the difference between the mask-then-join and contour-then-mask methods, and the effect of the updated parameter choices (i.e., n , K_1 , and K_2) in ERA-Interim at 00:00 UTC on 2001-01-01. The mask-then-join approach using the
255 original parameters (Figure 3a) is clearly identifying fronts, but they are fractured with frequent gaps. The contour-then-mask (Figure 3b) results in much smoother front features with fewer gaps, and more fronts identified. Figure 3c shows the results of the updated parameters with more smoothing cycles and stronger thresholds. Figure 3d shows the fronts identified in ERA5, and will be discussed further in Section 4.3. Compared to the original parameters, the front features are smoother, with fewer breaks and many spurious local fronts have been removed. One feature that can be seen is the warm front directly to the
260 southeast of the cold front that crosses the south of the UK. Such features were noted by Hewson (1998), and are associated




Figure 3. Comparison of methods in ERA-Interim at 00:00 on 2001-01-01. (a) mask-then-join with $n = 2$, $K_1 = -5 \times 10^{-11} \text{ K m}^{-2}$ and $K_2 = 0 \text{ K m}^{-1}$, (b) contour-then-mask with $n = 2$, $K_1 = -5 \times 10^{-11} \text{ K m}^{-2}$ and $K_2 = 0 \text{ K m}^{-1}$, (c) contour-then-mask with $n = 8$, $K_1 = -1.6 \times 10^{-11} \text{ K m}^{-2}$ and $K_2 = 7.5 \times 10^{-6} \text{ K m}^{-1}$, and (d) in ERA5 using contour-then-mask with $n = 96$, $K_1 = -1.6 \times 10^{-11} \text{ K m}^{-2}$ and $K_2 = 7.5 \times 10^{-6} \text{ K m}^{-1}$. Thin black lines indicate contours of wet-bulb potential temperature θ_W . Thick blue lines indicate cold fronts, thick red lines indicate warm fronts and thick black lines indicate quasi-stationary fronts. All fronts were classified using a threshold of $K_3 = 1.5 \text{ ms}^{-1}$

with a warm conveyor belt running adjacent to the front. Hewson and Tittley (2010) suggest a third masking criteria based on potential temperature rather than wet-bulb potential temperature that may be implemented in a future version of the code documented in this study.

Figure 4 compares the front frequency climatologies from three different implementations of the Hewson (1998) algorithm applied to ERA-Interim with identical parameters (i.e., the same number of smoothing passes, and thresholds K_1 and K_2): the implementation of Berry et al. (2011b) used by Dowdy and Catto (2017) (Figure 4a); a version incorporating our numerical updates but using the original mask-then-join approach (Figure 4b); and our final version using the contour-then-mask approach (Figure 4c). Figure 4d shows that our numerical updates result in slightly lower numbers of fronts identified in across most of the northern and southern hemisphere extra-tropics, and slightly higher numbers of fronts identified in the tropics. Figures 4e and 4f compare our final version with the implementation of Berry et al. (2011b) and the version incorporating only the numerical updates. The greatest increases in the number of fronts identified are seen in or adjacent to regions where fronts were already common, highlighting the effectiveness of the contour-then-mask approach at reducing the breaks in the fronts. In the densest region of the North Atlantic storm track, the number of fronts identified increases by almost 100 %.




Figure 4. Comparison of global climatologies of front frequency as a percentage of 6-hourly times. (a) Dowdy & Catto (2017), (b) updated implementation using the mask-then-join approach (c) updated implementation using the contour-then-mask approach, (d) percentage difference between mask-then-join and Dowdy & Catto (2017), (e) percentage difference between contour-then-mask and Dowdy & Catto (2017), and (f) percentage difference between contour-then-mask and mask-then-join. All climatologies were computed with $n = 2$ smoothing cycles, $K_1 = -5 \times 10^{-11} \text{ K m}^{-2}$ and $K_2 = 0 \text{ K m}^{-1}$

While changing the implementation of the front identification leads to an increase in the number of fronts identified, as shown
 275 in Figure 4, the next aspect of the updated method is a change to the parameters used. Figure 5 compares the climatology of
 front frequency of our final version with updated parameters (i.e., smoothing passes, and thresholds K_1 and K_2) applied to
 ERA-Interim against the implementation by Berry et al. (2011b) with the original parameters. The updated parameters result
 in slightly fewer fronts identified in almost all regions, due to the increased smoothing, making the climatology more similar
 to earlier estimates, but with the smoother individual fronts given by the contour-then-mask method. The greatest decreases
 280 are seen on the edge of the tropics, adjacent to regions with high front activity. This pattern is to be expected due to the rapid
 drop-off in the climatological quantile values of the masking parameters in the tropics in Figure 2.




Figure 5. Updated parameters. Percentage difference between ERA-Interim climatology of front frequency computed using updated parameters $n = 8$, $K_1 = -1.6 \times 10^{-11} \text{ K m}^{-2}$ and $K_2 = 7.5 \times 10^{-6} \text{ K m}^{-1}$, and the original parameters $n = 2$, $K_1 = -5 \times 10^{-11} \text{ K m}^{-2}$ and $K_2 = 0 \text{ K m}^{-1}$.




Figure 6. Updated global climatologies of front frequency as a percentage of times. (a) All fronts, (b) cold fronts, (c) warm fronts, (d) quasi-stationary fronts. All climatologies were computed with $n = 8$ smoothing cycles, $K_1 = -1.6 \times 10^{-11} \text{ K m}^{-2}$ and $K_2 = 7.5 \times 10^{-6} \text{ K m}^{-1}$ and $K_3 = 1.5 \text{ m s}^{-1}$.

4.2 Front climatology from ERA-Interim

Figure 6 shows the climatology of front frequency of our final version with updated parameters applied to ERA-Interim, including the breakdown into cold, warm and quasi-stationary fronts. Figure 6a allows for a direct comparison with the climatologies

285 in Figure 4, showing that while the updated parameters reduce the number of fronts identified compared to the updated numerical implementation only, overall more fronts are still identified in almost all regions than in earlier versions. Figure 6b and 6c show that cold and warm fronts occur with similar frequencies in most extra-tropical regions, as previously shown in Berry et al. (2011b). Figure 6d shows that quasi-stationary fronts occur most often where winds are weaker, particularly in the horse latitudes and inter-tropical convergence zone, and adjacent to high orography, as expected.

290 Figure 7 breaks the classification of fronts down still further, reporting cold and warm fronts by season. Unsurprisingly, cold fronts in the Northern Hemisphere are most common at the beginning of the storm track regions of both the Atlantic and Pacific oceans in northern winter (DJF, Figure 7a). In contrast, warm fronts in northern summer (JJA, Figure 7g) tend to outnumber cold fronts (Figure 7c). In agreement with Berry et al. (2011b), the seasonal distribution of fronts in the Southern Hemisphere is much more stable. Cold fronts are slightly more common though less widely distributed in the Southern Hemisphere during
295 southern summer (DJF, Figure 7a) than in southern winter (JJA, Figure 7c). When the storm track moves poleward in Southern winter there are larger numbers of warm fronts near Antarctica (JJA, Figure 7g).

4.3 Front climatology from ERA5

The ERA5 reanalysis has a higher resolution than ERA-Interim, with grid spacing of $0.25^\circ \times 0.25^\circ$ compared to $0.75^\circ \times 0.75^\circ$ for ERA-Interim. For ERA5, a total of $n = 96$ smoothing cycles were required to make the climatologies of the TFP and
300 gradient similar to ERA-Interim. Figure 3d illustrates fronts identified over Europe and the North Atlantic at 00:00 UTC on 2001-01-01. As expected, the features are very similar to those identified in ERA-Interim against which it was calibrated (Figure 3c). Figure 8 compares the frequency of fronts identified in ERA5 with that in ERA-Interim when fronts are identified in ERA5 at $0.25^\circ \times 0.25^\circ$ grid spacing with $n = 96$ smoothing cycles but identical thresholds to those used for ERA-Interim, then aggregated to $0.75^\circ \times 0.75^\circ$ grid spacing for comparison with ERA-Interim. Aggregation is performed by counting individual
305 fronts identified at the higher resolution passing through the lower resolution grid. When aggregated to the same resolution, more fronts are identified almost everywhere in ERA5 than in ERA-Interim. Decreases in frequency are primarily associated with areas of high orography. The pattern of increase broadly follows the general distribution of fronts, with more fronts seen where they were already common, particularly in the storm tracks where front frequency increases by 20% to 40%. The greatest percentage increases are seen in the inter-tropical convergence zone (ITCZ) east and west of South America where
310 very few fronts were identified in ERA-Interim.

Figure 9 shows the climatology of fronts by type identified in ERA5 at its native $0.25^\circ \times 0.25^\circ$ resolution. Due to the smaller grid boxes, the frequency is necessarily lower than for ERA-Interim in Figure 6 and the aggregated data in Figure 8. One ERA-Interim grid box contains nine ERA5 grid boxes. A perfectly straight front passing through one ERA-Interim grid box would pass through only three of the nine associated ERA5 grid boxes. Therefore, one might expect the front frequency in ERA5 at
315 its native resolution to be approximately one third of the frequency in ERA-Interim. Comparing Figures 6 and 9 shows that this is approximately the case. Figure 9d suggests that the large increases in the fronts in the ITCZ seen in Figure 8b is primarily associated quasi-stationary fronts, due to the light winds associated with the ITCZ.




Figure 7. Updated seasonal climatologies of front frequency. (a) Cold fronts in DJF, (b) Cold fronts in MAM, (c) Cold fronts in JJA, (d) Cold fronts in SON, (e) Warm fronts in DJF, (f) Warm fronts in MAM, (g) Warm fronts in JJA, (h) Warm fronts in SON. All climatologies were computed with $n = 8$ smoothing cycles, $K_1 = -1.6 \times 10^{-11} \text{ K m}^{-2}$ and $K_2 = 7.5 \times 10^{-6} \text{ K m}^{-1}$ and $K_3 = 1.5 \text{ m s}^{-1}$.

Figure 10 shows the seasonal breakdown of cold and warm fronts in ERA5, which is provided to be able to compare the most up-to-date climatology from ERA5 with previous studies. In general the maximum warm front frequency occurs at higher




Figure 8. ERA5 compared to ERA-Interim. (a) ERA5 climatology of all fronts at $0.75^\circ \times 0.75^\circ$, and (b) percentage difference between ERA5 and ERA-Interim. The ERA5 climatology were computed with $n = 96$ smoothing cycles, $K_1 = -1.6 \times 10^{-11} \text{ K m}^{-2}$ and $K_2 = 7.5 \times 10^{-6} \text{ K m}^{-1}$ and $K_3 = 1.5 \text{ m s}^{-1}$. ERA5 fronts were identified at $0.25^\circ \times 0.25^\circ$ then regridded to $0.75^\circ \times 0.75^\circ$ for comparison with ERA-Interim.

320 latitudes than the maximum cold front frequency, due to the structure of extratropical cyclones and the associated poleward transport of warm air. During DJF especially, the sea surface temperature (SST) fronts associated with the Gulf Stream in the North Atlantic and Kuroshio current in the North Pacific are clearly visible. The influence of the SST on the atmosphere is more marked for higher resolution ocean and atmosphere (Parfitt et al., 2016, 2017a).

5 Discussion

325 In this paper, we have presented an updated implementation of the automatic front identification method of Berry et al. (2011b), based on Hewson (1998). The updated implementation was designed specifically to scale to modern high resolution data sets. It is open source and does not require compilation, making it extremely portable. Despite not requiring compilation, computational performance is improved over earlier versions that were implemented in NCL with compiled components. Performance improvements come primarily from three areas: (i) the improved efficiency of the contouring algorithm compared
 330 to the line joining algorithm; (ii) vectorization of many calculations to avoid unnecessary loops; (iii) reduced memory usage by avoiding pre-allocating unnecessarily large arrays. One month of global ERA-Interim data at 6-hourly intervals and $0.75^\circ \times$




Figure 9. ERA5 global climatologies. (a) All fronts, (b) cold fronts, (c) warm fronts, (d) quasi-stationary fronts. All climatologies were computed with $n = 96$ smoothing cycles, $K_1 = -1.6 \times 10^{-11} \text{ K m}^{-2}$ and $K_2 = 7.5 \times 10^{-6} \text{ K m}^{-1}$ and $K_3 = 1.5 \text{ ms}^{-1}$.

0.75° resolution can be processed in around 6 minutes using a single core of an Intel i7-8565U based laptop with a theoretical maximum speed of 4.6 GHz. The same amount of global ERA5 data at $0.25^\circ \times 0.25^\circ$ can be processed in around 1 hour. Memory requirements are minimal since only one time step is processed at once. The improved scalability enables us to present high resolution climatologies of cold, warm and quasi-stationary fronts for all seasons from the ERA5 reanalysis.

In addition to several numerical improvements, the revised implementation uses the contour-then-mask approach originally proposed by Hewson (1998) rather than the mask-then-join approach used by Berry et al. (2011b). The advantages of the contour-then-mask approach are demonstrated by example and by comparison of climatologies which show increased numbers of fronts identified almost everywhere. Gaps in what should be continuous fronts are reduced in ERA-Interim, and greater improvements are expected in lower resolution datasets for the reasons demonstrated in Figure 1. This improvement will be useful when linking frontal features to precipitation or winds (or compound extreme events) as in Catto and Dowdy (2021), or when using more object-based connections such as Papritz et al. (2014). Computational performance is improved since the contouring algorithm is more efficient than the line joining algorithm used by Berry et al. (2011b).

Most automatic feature detection algorithms require a calibration or training step involving comparison to analyses by a meteorologist. While this step cannot be neglected, we propose a quantile based approach to setting thresholds for front identification. Setting thresholds in terms of climatological quantiles makes the thresholds more easily comparable between data sets of differing resolution. By considering the climatological distributions of the masking variables, we have demonstrated




Figure 10. ERA5 seasonal climatologies of front frequency. (a) Cold fronts in DJF, (b) Cold fronts in MAM, (c) Cold fronts in JJA, (d) Cold fronts in SON, (e) Warm fronts in DJF, (f) Warm fronts in MAM, (g) Warm fronts in JJA, (h) Warm fronts in SON. All climatologies were computed with $n = 96$ smoothing cycles, $K_1 = -1.6 \times 10^{-11} \text{ K m}^{-2}$ and $K_2 = 7.5 \times 10^{-6} \text{ K m}^{-1}$ and $K_3 = 1.5 \text{ m s}^{-1}$.

for the first time the regional and seasonal variation of the TFP and gradient fields. Subsequent analyses may consider adopting latitudinally or seasonally varying thresholds in order to capture features that may be missed by or eliminate spurious features

350 included by the used constant thresholds. We also demonstrate how the quantile based method can also be used to compare analyses from models or analyses at different resolutions by smoothing until the climatological quantiles are similar. In ERA5 this results in greater numbers of fronts identified even after smoothing, similar to the results of Parfitt et al. (2017b) after interpolation to lower resolution. Smoothing has the advantage of allowing feature identification to be conducted at the native resolution of each data set.

355 In addition to the various numerical and methodological improvements presented in this study, further numerical improvements, methodological choices, and alternative choices of meteorological field are possible. In addition to improving the accuracy of the finite difference approximations of the second derivative fields to second order, more accurate finite difference schemes could be used for both the first and second order derivatives. In testing, moving to fourth order accuracy for both first and second derivative fields did yield modest increases in both the number of fronts and front points identified, for minimal
360 computational expense. However, the increased accuracy of the derivatives is difficult to justify given the coarse resolution of the fields in question. Following Berry et al. (2011b), we identify fronts as zero contours in the field defined by Equation 5 of Hewson (1998), effectively the third derivative of the wet-bulb potential temperature field at 850 hPa. Firstly, meteorological fields other than wet-bulb potential temperature could be considered, see Hewson (1998) for a list of previously considered fields. Secondly, Hewson (1998) derived an alternative expression for the front locator field, designed to mitigate the effects
365 of frontal curvature which can otherwise lead to noise and excessive curvature in identified fronts. We retained the simpler definition for compatibility with Berry et al. (2011b) and subsequent studies, but the alternative definition preferred by Hewson (1998) may be included as an option in a future version of the code associated with this study. Additional diagnostics such as distinguishing between local and synoptic fronts suggested by Jenkner et al. (2010), or the additional criteria proposed by Hewson and Titley (2010) designed to eliminate spurious features associated with proximity to the warm conveyor
370 belt, could also be implemented. Furthermore, while all distance calculations are carried out on a sphere in the updated implementation, contouring and interpolation still take place on a regular longitude-latitude grid. Greater accuracy could be achieved at high latitudes by also carrying out these operations on a sphere.

While cyclone identification algorithms routinely include the ability to track cyclonic features over subsequent time steps, similar feature tracking algorithms are almost absent for fronts. Front tracking is inherently more complex than cyclone tracking
375 since fronts are complex line objects whereas cyclones can be reduced to simple point objects or point objects with an associated area. Hewson and Titley (2010) proposed a sophisticated tracking scheme for cyclonic features developing on fronts, which relies on accurate identification and classification of fronts in order to identify cyclones early in their life cycle, but is limited to tracking point objects associated with cyclones rather than fronts themselves. To the authors' knowledge, only Rüdüsühli et al. (2020) have documented a front tracking algorithm. An openly available front tracking algorithm would offer new possibilities
380 in terms of attributing and analysing impacts of individual fronts, e.g., precipitation or wind events, or understanding biases in weather and climate models.

Code availability. Code for revised method detailed in this paper are available from <https://doi.org/10.5281/zenodo.7278068> and future developments will be available at https://github.com/phil-sansom/front_id.

Author contributions. PGS developed and tested the software, produced the results, and wrote the paper. JLC led the project, interpreted
385 results, and wrote the paper.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. This research was supported by Natural Environment Research Council (NERC) grant NE/V004166/1. The authors thank Dr. Duncan Ackerley for comments on the manuscript.

References

- 390 Berry, G., Jakob, C., and Reeder, M.: Recent global trends in atmospheric fronts, *Geophysical Research Letters*, 38, 1–6, <https://doi.org/10.1029/2011GL049481>, 2011a.
- Berry, G., Reeder, M. J., and Jakob, C.: A global climatology of atmospheric fronts, *Geophysical Research Letters*, 38, 1–5, <https://doi.org/10.1029/2010GL046451>, 2011b.
- Bitsa, E., Flocas, H. A., Kouroutzoglou, J., Galanis, G., Hatzaki, M., Latsas, G., Rudeva, I., and Simmonds, I.: A Mediterranean cold front identification scheme combining wind and thermal criteria, *International Journal of Climatology*, 41, 6497–6510, <https://doi.org/https://doi.org/10.1002/joc.7208>, 2021.
- 395 Browning, K. A.: The sting at the end of the tail: Damaging winds associated with extratropical cyclones, *Quarterly Journal of the Royal Meteorological Society*, 130, 375–399, <https://doi.org/10.1256/qj.02.143>, 2004.
- Catto, J. L. and Dowdy, A. J.: Understanding compound hazards from a weather system perspective, *Weather and Climate Extremes*, 32, <https://doi.org/10.1016/j.wace.2021.100313>, 2021.
- 400 Catto, J. L. and Pfahl, S.: The importance of fronts for extreme precipitation, *Journal of Geophysical Research Atmospheres*, 118, 10,791–10,801, <https://doi.org/10.1002/jgrd.50852>, 2013.
- Catto, J. L., Jakob, C., Berry, G., and Nicholls, N.: Relating global precipitation to atmospheric fronts, *Geophysical Research Letters*, 39, 1–6, <https://doi.org/10.1029/2012GL051736>, 2012.
- 405 Catto, J. L., Nicholls, N., Jakob, C., and Shelton, K. L.: Atmospheric fronts in current and future climates, *Geophysical Research Letters*, 41, 7642–7650, <https://doi.org/10.1002/2014GL061943>, 2014.
- Catto, J. L., Ackerley, D., Booth, J. F., Champion, A. J., Colle, B. A., Pfahl, S., Pinto, J. G., Quinting, J. F., and Seiler, C.: The Future of Midlatitude Cyclones, *Current Climate Change Reports*, 5, 407–420, <https://doi.org/10.1007/s40641-019-00149-4>, 2019.
- Davies-Jones, R.: An efficient and accurate method for computing the wet-bulb temperature along pseudoadiabats, *Monthly Weather Review*, 410 136, 2764–2785, <https://doi.org/10.1175/2007MWR2224.1>, 2008.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J. J., Park, B. K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J. N., and Vitart, F.: The ERA-Interim reanalysis: 415 Configuration and performance of the data assimilation system, *Quarterly Journal of the Royal Meteorological Society*, 137, 553–597, <https://doi.org/10.1002/qj.828>, 2011.
- Dowdy, A. J. and Catto, J. L.: Extreme weather caused by concurrent cyclone, front and thunderstorm occurrences, *Scientific Reports*, 7, 1–8, <https://doi.org/10.1038/srep40359>, 2017.
- ECMWF: Part IV: Physical processes, in: IFS Documentation CY47R3, ECMWF, <https://doi.org/10.21957/eyrpir4vj>, 2021.
- 420 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J. N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, 425 <https://doi.org/10.1002/qj.3803>, 2020.

- Hewson, T. D.: Objective fronts, *Meteorological Applications*, 5, 37–65, <https://doi.org/10.1017/S1350482798000553>, 1998.
- Hewson, T. D.: Objective identification of fronts, frontal waves and potential waves, in: *Cost Action 78 Final Report – Improvement of Nowcasting Techniques*, edited by Lagouvardos, K., Liljas, E., Conway, B., and Sunde, J., pp. 285–290, European Commission EUR 19544. Cambridge University Press, Luxembourg, 2001.
- 430 Hewson, T. D. and Tittley, H. A.: Objective identification, typing and tracking of the complete life-cycles of cyclonic features at high spatial resolution, *Meteorological Applications*, 17, 355–381, <https://doi.org/10.1002/met.204>, 2010.
- Hope, P., Keay, K., Pook, M., Catto, J. L., Simmonds, I., Mills, G., McIntosh, P., Risbey, J., and Berry, G.: A comparison of automated methods of front recognition for climate studies: A case study in southwest Western Australia, *Monthly Weather Review*, 142, 343–363, <https://doi.org/10.1175/MWR-D-12-00252.1>, 2014.
- 435 Jenkner, J., Sprenger, M., Schwenk, I., Schwierz, C., Dierer, S., and Leuenberger, D.: Detection and climatology of fronts in a high-resolution model reanalysis over the Alps, *Meteorological Applications*, 17, 1–18, <https://doi.org/10.1002/met.142>, 2010.
- Leung, L. R., Boos, W. R., Catto, J. L., A. DeMott, C., Martin, G. M., Neelin, J. D., O’Brien, T. A., Xie, S., Feng, Z., Klingaman, N. P., Kuo, Y., Lee, R. W., Martinez-Villalobos, C., Vishnu, S., Priestley, M. D. K., Tao, C., and Zhou, Y.: Exploratory Precipitation Metrics: Spatiotemporal Characteristics, Process-Oriented, and Phenomena-Based, *Journal of Climate*, 35, 3659–3686, <https://doi.org/10.1175/JCLI-D-21-0590.1>, 2022.
- 440 NCL: The NCAR Command Language, UCAR/NCAR/CISL/TDD, Boulder, Colorado, <https://doi.org/10.5065/D6WD3XH5>, 2019.
- Papritz, L., Pfahl, S., Rudeva, I., Simmonds, I., Sodemann, H., and Wernli, H.: The Role of Extratropical Cyclones and Fronts for Southern Ocean Freshwater Fluxes, *Journal of Climate*, 27, 6205–6224, <https://doi.org/10.1175/JCLI-D-13-00409.1>, 2014.
- Parfitt, R., Czaja, A., Minobe, S., and Kuwano-Yoshida, A.: The atmospheric frontal response to SST perturbations in the Gulf Stream region, *Geophysical Research Letters*, 43, 2299–2306, <https://doi.org/https://doi.org/10.1002/2016GL067723>, 2016.
- 445 Parfitt, R., Czaja, A., and Kwon, Y.-O.: The impact of SST resolution change in the ERA-Interim reanalysis on wintertime Gulf Stream frontal air-sea interaction, *Geophysical Research Letters*, 44, 3246–3254, <https://doi.org/https://doi.org/10.1002/2017GL073028>, 2017a.
- Parfitt, R., Czaja, A., and Seo, H.: A simple diagnostic for the detection of atmospheric fronts, *Geophysical Research Letters*, 44, 4351–4358, <https://doi.org/10.1002/2017GL073662>, 2017b.
- 450 R Core Team: R: A Language and Environment for Statistical Computing, R Foundation for Statistical Computing, Vienna, Austria, <https://www.r-project.org>, 2021.
- Raveh-Rubin, S. and Catto, J. L.: Climatology and dynamics of the link between dry intrusions and cold fronts during winter, Part II: Front-centred perspective, *Climate Dynamics*, 53, 1893–1909, <https://doi.org/10.1007/s00382-019-04793-2>, 2019.
- Renard, R. J. and Clarke, L. C.: Experiments in Numerical Objective Frontal Analysis 1, *Monthly Weather Review*, 93, 547–556, [https://doi.org/10.1175/1520-0493\(1965\)093<0547:einofa>2.3.co;2](https://doi.org/10.1175/1520-0493(1965)093<0547:einofa>2.3.co;2), 1965.
- 455 Rüdüsühli, S., Sprenger, M., Leutwyler, D., Schär, C., and Wernli, H.: Attribution of precipitation to cyclones and fronts over Europe in a kilometer-scale regional climate simulation, *Weather and Climate Dynamics*, 1, 675–699, <https://doi.org/10.5194/wcd-1-675-2020>, 2020.
- Schemm, S., Rudeva, I., and Simmonds, I.: Extratropical fronts in the lower troposphere-global perspectives obtained from two automated methods, *Quarterly Journal of the Royal Meteorological Society*, 141, 1686–1698, <https://doi.org/10.1002/qj.2471>, 2015.
- 460 Schemm, S., Sprenger, M., Martius, O., Wernli, H., and Zimmer, M.: Increase in the number of extremely strong fronts over Europe? A study based on ERA-Interim reanalysis (1979–2014), *Geophysical Research Letters*, 44, 553–561, <https://doi.org/10.1002/2016GL071451>, 2017.

- Simmonds, I., Keay, K., and Bye, J. A. T.: Identification and climatology of Southern Hemisphere mobile fronts in a modern reanalysis, *Journal of Climate*, 25, 1945–1962, <https://doi.org/10.1175/JCLI-D-11-00100.1>, 2012.
- 465 Soster, F. and Parfitt, R.: On Objective Identification of Atmospheric Fronts and Frontal Precipitation in Reanalysis Datasets, *Journal of Climate*, 35, 4513–4534, <https://doi.org/10.1175/JCLI-D-21-0596.1>, 2022.
- Thomas, C. and Schultz, D.: What are the Best Thermodynamic Quantity and Function to Define a Front in Gridded Model Output?, *Bulletin of the American Meteorological Society*, 100, <https://doi.org/10.1175/BAMS-D-18-0137.1>, 2019.
- Uppala, S. M., Kallberg, P. W., Simmons, A. J., Andrae, U., Bechtold, V. D. C., Fiorino, M., Gibson, J. K., Haseler, J., Hernandez, A., Kelly,
470 G. A., Li, X., Onogi, K., Saarinen, S., Sokka, N., Allan, R. P., Andersson, E., Arpe, K., Balmaseda, M. A., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Caires, S., Chevallier, F., Dethof, A., Dragosavac, M., Fisher, M., Fuentes, M., Hagemann, S., Holm, E., Hoskins, B. J., Isaksen, L., Janssen, P. A. E. M., Jenne, R., McNally, A. P., Mahfouf, J. F., Morcrette, J. J., Rayner, N. A., Saunders, R. W., Simon, P., Sterl, A., Trenberth, K. E., Untch, A., Vasiljevic, D., Viterbo, P., and Woollen, J.: The ERA-40 re-analysis, *Quarterly Journal of the Royal Meteorological Society*, 131, 2961–3012, <https://doi.org/10.1256/qj.04.176>, 2005.