

# A Sub-Grid Parameterization Scheme for Topographic Vertical Motion in CAM5-SE

Yaqi Wang<sup>1</sup>, Lanning Wang<sup>1,2\*</sup>, Juan Feng<sup>1</sup>, Zhenya Song<sup>3</sup>, Qizhong Wu<sup>1</sup>, Huaqiong Cheng<sup>1</sup>

5 <sup>1</sup>College of Global Change and Earth System Science (GCESS), Beijing Normal University, Beijing 100875, China

<sup>2</sup>Laboratory for Regional Oceanography and Numerical Modeling, Pilot National Laboratory for Marine Science and Technology, Qingdao 266237, China

10 <sup>3</sup>First Institute of Oceanography, and Key Laboratory of Marine Science and Numerical Modeling, Ministry of Natural Resources, Qingdao 266061, China

*Corresponding to: Lanning Wang (wangln@bnu.edu.cn)*

## Abstract:

15 Overestimation of precipitation over steep mountains is always a common bias of atmospheric general circulation models (AGCMs). One basic reason is the imperfection of parameterization scheme. Sub-grid topography has a non-negligible role in the dynamics of the actual atmosphere, and therefore the sub-grid topographic parameterization schemes have been the focus of model development. This study  
20 [proposes a sub-grid parameterization scheme for topographic vertical motion in CAM5-SE \(Community Atmospheric Model version 5 with spectral element dynamical core\)](#) to revise the original vertical velocity by adding the topographic vertical motion and then resulting a significant improvement of simulation in precipitation over steep mountains. The results show a better improvement in precipitation simulation in steep  
25 mountains, such as the steep edge of the Tibetan Plateau and the Andes. The positive deviations of the precipitation on the mountain tops and the negative deviations in the windward slope are revised. The improved scheme of topographic vertical motion reduces the model biases of summer mean precipitation simulations by up to 48% (6.23 mm day<sup>-1</sup>) on the mountain tops. The improvement of convective precipitation (4.83  
30 mm day<sup>-1</sup>) contributes the most to the improvement of the total precipitation simulation. In addition, we extend the dynamic lifting effect of topography from the lowest layer [\(Single experiment\)](#) to multiple layers, approaching the [surface-layer, bottom model layers \(Multi experiment\)](#). Moreover, the water vapor transport in low-altitude regions in front of the windward slope is also considerably improved, leading to simulations of  
35 much more realistic circulation patterns in the multi-layer scheme. Since the sub-grid parameterization scheme addresses the more detailed problem caused by topography, the water vapor is transported further to the northwest in the multi-layer scheme. The topographic vertical motion schemes in both the Single- and Multi-experiments can improve the model performance in simulating precipitation in all regions with complex  
40 terrain.

## 1 Introduction

Numerical models have been widely used and become an essential tool to predict and simulate the weather and climate. However, there are still large deviations compared with observations, especially for precipitation simulation and prediction. It is of great  
45 scientific and social relevance to accurately simulate precipitation by using atmospheric general circulation models (AGCMs). In particular, the Coupled Model Intercomparison Project Phase 5 (CMIP5) and Phase 6 (CMIP6) models always overestimate the precipitation in regions with steep topography, which have been investigated in previous studies (Liu et al. 2014; Akinsanola et al. 2021; Cui et al. 2021).  
50 Jia et al. (2019) found that all CMIP5 models overestimate the monthly precipitation over the Tibetan Plateau by an average of 48.2 mm (~150%), with larger biases during spring and summer. Zhu and Yang (2020) also found that the [systematic-model biases \(overestimation of precipitation\) in the over the Tibetan Plateau still exist, in the CMIP6 models were even performed larger \(more poorly based on the performance of 23](#)

55 ~~models-positive) than~~ in ~~CMIP6~~ the CMIP5 models. Similar problems also exist in  
precipitation simulations in other mountain regions with steep terrains, such as the  
Andes in South America, the Rocky Mountains of North America, and Indonesia.  
Excessive precipitation was simulated in both weather/climate models and  
global/regional models in regions with steep and high mountains, but less precipitation  
60 ~~in windward slopes~~ before the foothills of the steep slope (Done et al., 2004; Kunz and  
Kottmeier, 2006; Alpert et al., 2012; Chao 2012; Navale and Singh, 2020).

The reasons for excessive precipitation simulated by numerical models over steep  
mountains are complex, involving the horizontal resolution, dynamical ~~cores~~ framework,  
physical processes, and their complicated interactions (Liang et al., 2021). There is  
65 plenty of evidence of a close relationship between orography and precipitation patterns  
at spatial scales of a few kilometers, even in climatological precipitation rates. Thus,  
improving model resolution is a possible way to improve the biases of precipitation  
simulations. Kimoto et al. (2005) found that higher-resolution versions of General  
Circulation Models (GCMs) can better characterize the frequency distributions of  
70 different precipitation patterns. Similar results can be found in regional models. Lin et  
al. (2018) compared the simulations with resolutions of 30 km, 10 km and 2 km based  
on the Weather Research and Forecasting model, and they found that higher-resolution  
simulations can reduce positive precipitation biases over the Tibetan Plateau. However,  
increasing spatial resolution does not always improve precipitation simulations in some  
75 areas, for example, in lowlands of southeastern England (Chan et al. 2013; Wang et al.  
2017). The relationship between the spatial resolution of models and the quality of  
precipitation simulation remains elusive. Additionally, high-resolution climate models  
require a large amount of computation and storage. Some parameterization schemes are  
also proposed to improve the accuracy of precipitation simulation, which mainly focus  
80 on the parameterization schemes for physical processes. For example, in the past 20  
years, much effort has been made to develop stochastic convection schemes and apply  
them to numerical models, resulting in some substantial improvements in precipitation  
simulation (Chen et al. 2010; Fonseca et al. 2015; Wang and Zhang, 2016; Attada et al.  
2020).

85 The simulation bias of topographic precipitation has been a challenge for numerical  
models. Most studies are based on improving model resolution and the parameterization  
schemes of physical processes, but few studies focused on the modification of the  
dynamic ~~core~~ framework for numerical models, especially the dynamic lifting. At  
spatial scales greater than approximately 40 km and for mountain ranges exceeding  
90 approximately 1.5 km in height, the maximum condensation is generated over low,  
steep and windward slopes due to upslope flow (Roe 2005). An important quantity of  
orographic precipitation is water vapor flux. In numerical models, Yu et al. (2015)  
replaced the semi-Lagrangian method with a finite-difference approach for the trace  
transport algorithm to restrain the "overshoot" of water vapor to the high-altitude region  
95 of the windward slopes. Codron and Sadourny (2002) tested the advected water vapor  
with respect to saturation values and redistributed it accordingly over the grid points

found along the advecting path. Actually, these two schemes add the limitation of ~~oversaturation~~ supersaturation for water vapor advection, which may cause partial precipitation when the water vapor advects upward mountain slopes along terrain-following coordinates. Less water vapor is transported to summits and plateaus and settles in windward slopes and foothills in advance, thus improving precipitation simulations in steep mountains. These studies only ~~improve~~ improved the scheme of water vapor advection scheme, ~~and only~~. Shen et al. (2007) proposed a sub-grid correction parameterization scheme for pressure tendency by considering slope and orientation according to the disturbance lifting caused by each fine grid. Based on this, the precipitation simulation in the ~~P-σ~~ regional climate model of Nanjing University over complex terrain areas was improved. But it is only a case study of precipitation simulation in East China.

As mentioned above, sufficient water vapor and dynamic lifting are the necessary conditions for precipitation (Shen et al. 2021). Considering the shortcomings of the current dynamic lifting studies for numerical models, ~~in this study~~, we propose a sub-grid parameterization scheme of topographic vertical motion and apply in CAM5, one of global atmosphere general circulation models, to improve precipitation simulation in areas with complex terrain. In particular, we extend the dynamic lifting effect of topography on airflow from the lowest model layer to multiple layers and consider the influence of the decay of vertical airflow.

The remainder of this paper is organized as follows. Section 2 describes the modeling context and the data used in this research and details the sub-grid parameterization scheme for topographic vertical velocity. Section 3 analyzes and compares the precipitation simulated by two topographic vertical velocity experiments. The main conclusions and discussion are presented in section 4.

## 2 Model, methodology and experiments

### 2.1 CAM5-SE

The models used in this study are the Community Earth System Model (CESM; Hurrell et al. 2013) version 1.2.1. from the National Center for Atmospheric Research (NCAR) and the Community Atmospheric Model version 5 (CAM5; Neale et al. 2010) with the new spectral element dynamical core (CAM-SE). The CAM-SE is based on the High-Order Method Modeling Environment spectral element method (HOMME, Dennis et al. 2012) and adopts a conventional vector-invariant form of the moist primitive equations. Noted that the CAM-SE uses the vector-invariant form of the momentum equation instead of the vorticity-divergence equation. The pressure vertical velocity can be expressed by  $\omega = D_p/D_t$ , as shown in Eq. (1).

$$\omega = \frac{\partial p}{\partial t} + \vec{u} \cdot \nabla p + \dot{\eta} \frac{\partial p}{\partial \eta} = \vec{u} \cdot \nabla p - \int_{\eta_{\text{top}}}^{\eta} \nabla \cdot \left( \frac{\partial p}{\partial \eta} \vec{u} \right) d\eta', \quad (1)$$

Given description of the coordinate in CAM5-SE, the continuous system of equations can be written following the first law of thermodynamics, Kasahara (1974) and Simmons and Strüfing (1981). The prognostic equations are as shown in Eq. (2) (Neale et al., 2010).

$$\begin{aligned} \frac{\partial \zeta}{\partial t} &= \mathbf{k} \cdot \nabla \times \left( \frac{\mathbf{n}}{\cos \phi} \right) + F_{\zeta_H}, \\ \frac{\partial \delta}{\partial t} &= \nabla \cdot \left( \frac{\mathbf{n}}{\cos \phi} \right) - \nabla^2 (E + \Phi) + F_{\delta_H}, \\ \frac{\partial T}{\partial t} &= \frac{-1}{a \cos^2 \phi} \left[ \frac{\partial}{\partial \lambda} (UT) + \cos \phi \frac{\partial}{\partial \phi} (VT) \right] + T\delta - \dot{\eta} \frac{\partial T}{\partial \eta} + \frac{R}{c_p^*} T_v \frac{\omega}{p} \\ &\quad + Q + F_{T_H} + F_{F_H}, \\ \frac{\partial q}{\partial t} &= \frac{-1}{a \cos^2 \phi} \left[ \frac{\partial}{\partial \lambda} (Uq) + \cos \phi \frac{\partial}{\partial \phi} (Vq) \right] + q\delta - \dot{\eta} \frac{\partial q}{\partial \eta} + S, \\ \frac{\partial \pi}{\partial t} &= \int_1^{\eta_t} \nabla \cdot \left( \frac{\partial p}{\partial \eta} \mathbf{V} \right) d\eta \end{aligned} \quad (2)$$

The third equation in Eqs. (2) above shows that in the SE dynamic framework, vertical velocity affects the tendency of temperature  $\frac{\partial T}{\partial t}$  directly, and affects pressure P through the equation of state  $P = \rho RT$  indirectly. Thus, the correction of vertical velocity can change the atmospheric circulation and precipitation.

The major model physics of CAM5-SE include: (1) the separate deep convection scheme is ZM (Zhang and McFarlane 1995; Richter and Rasch 2008), the (2) The shallow convection scheme is University of Washington shallow convection schemes ((UW, Park and Bretherton 2009) and a). (3) The cloud microphysics scheme is MG1.0 (Morrison and Gettelman 2008; Gettelman et al. 2010). (4) The moist turbulence scheme (Gettelman et al. 2013) for calculating sub-grid vertical transport of heat and moisture. The cloud microphysics (Morrison and Gettelman 2008; Gettelman et al. 2010) includes both the direct and indirect effect for sulfate and black and organic carbon. For tracer advection, CAM-SE is modeled closely on the finite volume core. It uses the same conservation form of the transport equation— is diag\_TKE (Turbulent Kinetic Energy, Bretherton and the same vertically Lagrangian discretization (Lin, 2004); Park, 2009a). (5) The radiation scheme is Rapid Radiative Transfer Model for GCM (RRTMG) package (Mlawer et al. 1997).

## 2.2 Topographic vertical motion and sub-grid topography parameterization scheme

Alpert and Shafir (1989) found that orographic precipitation at micro/meso scales is highly predictable with the adiabatic assumption that the lifting is determined by  $\mathbf{V} \cdot \nabla Z_s$ . The surface vertical velocity caused by the forced lifting of topography can be expressed by Eq. (23).

$$\omega_s = \mathbf{V}_s \cdot \nabla Z_s, \quad (23)$$

In the P-coordinate system, Eq. (23) can be rewritten as Eq. (34):

$$\omega = \frac{dp_s}{dt} = \frac{\partial p_s}{\partial t} + \vec{V}_s \cdot \nabla p_s, \quad (34)$$

Where  $\vec{V}_s$  and  $p_s$  indicate the surface wind velocity and the surface pressure, respectively. After considering the topographic vertical velocity, Eq. (3) can be rewritten as Eq. (45).

$$\omega = \omega_0 + \omega_s, \quad (45)$$

$$\omega_s = -\rho g \vec{V}_s \cdot \nabla Z_s = -\rho g \cdot |\vec{V}_s| \cdot \tan\theta_N \cdot \cos(\theta - \varphi_N)$$

$$= -\rho g \sqrt{u^2 + v^2} \cdot \tan\theta_N \cdot (\cos\theta \cdot \cos\varphi_N + \sin\theta \cdot \sin\varphi_N), \quad (56)$$

where  $\omega_s$  denotes the topographic vertical velocity of the lowest model layer,  $\theta$  is the wind direction,  $\theta_N$  is the slope, and  $\varphi_N$  is the aspect,  $\rho$  is air density and  $g$  are constants is gravitational acceleration. It can be seen that the surface topographic vertical velocity is proportional to the surface wind speed, the tangent of the slope and the cosine of the angle between the mountain aspect and the wind direction. Figure 1a shows the distribution of surface topographic vertical velocity with the slope and the angle between the wind direction and aspect under unit wind speed. In fact, the angle between the mountain aspect and the wind direction ranges from  $0^\circ$  to  $360^\circ$ . When the angle in the range of  $0^\circ$ – $90^\circ$  or  $270^\circ$ – $360^\circ$ , it indicates an ascending motion, while the angle of  $90^\circ$ – $270^\circ$ , it represents a descending motion. The angle range of  $0^\circ$ – $90^\circ$  is chosen just because it can cover the range of cosine values and is adequately representative. This study only focuses on the simulation of precipitation caused by blocking uplift in windward slopes. At the current model resolution, the maximum slope captured by Digital Elevation Model (DEM) data is  $61^\circ$ , indicating that the maximum surface topographic vertical velocity is about 22Pa/s, and is positively correlated with slope. That is, when the mountain is the steepest and the angle between the wind direction and aspect is the smallest, the topographic vertical velocity reaches the maximum. However, when the slope is less than  $\sim 5^\circ$ , the topographic vertical velocity is so small that it can be ignored.

Generally, only the single layer, that is the lowest layer, is considered Shen et al. (2007)

proposed a sub-grid correction parameterization scheme for pressure tendency in numerical models. [regional climate model of Nanjing University](#). However, in the actual atmospheric movement, the topographic vertical motion not only affects the surface layer lowest model level, but also affects near surface layers. ~~Although it is realized in theory, it has not been applied in weather/climate prediction.~~ Thus, we extend the topographic vertical velocity from single layer to multi layers, as shown in Eq. (67):

$$\omega = \omega_0 + \omega_s \times \gamma, \quad (67)$$

where  $\gamma$  indicates the attenuation coefficient of topographic vertical velocity  $\omega_s$  and it increases with the elevation, as shown in Eq. (78):

$$\gamma = \frac{\text{sh}\left(\sqrt[2]{2} \frac{2\pi}{L} \sqrt{\frac{\sigma}{f^2}} \times p\right)}{\text{sh}\left(\sqrt[2]{2} \frac{2\pi}{L} \sqrt{\frac{\sigma}{f^2}} \times p_0\right)}, \quad (7)$$

$$\gamma = \frac{\sinh\left(\sqrt[2]{2} \frac{2\pi}{L} \sqrt{\frac{\sigma}{f^2}} \times p\right)}{\sinh\left(\sqrt[2]{2} \frac{2\pi}{L} \sqrt{\frac{\sigma}{f^2}} \times p_0\right)}, \quad (8)$$

where  $f$  represents the Coriolis term,  $p_0$  is the reference pressure,  $p$  is the actual pressure,  $\sigma = -\frac{T}{\theta} \frac{\partial \theta}{\partial p}$  is a constant, and  $L$  is the wavelength. Because the complexity of hyperbolic sine function calculation and the fact that the initial pressure in complex terrain areas actually does not start from the sea level but from the surface layer, we simplify Eq. (89) according to [Taylor's formula Taylor series](#) to make  $\gamma$  become an exponential function that varies only with latitude and pressure difference  $\Delta p$ :

$$\gamma \approx e^{\left(\frac{\sqrt{\sigma}}{2dl \times f \times \sin(lat)}\right) \times (-\Delta p)}, \quad (89)$$

where  $\Delta p$  indicates the difference between the surface pressure and the pressure on a certain model layer,  $dl$  is model horizontal resolution.  $\frac{\sqrt{\sigma}}{2dl \times f \times \sin(lat)}$  is static variable which can be preprocessed at each integration step without calculation. After simplification, the divergence of  $\gamma$  between Eq (78) and Eq (89) is only  $10^{-10}$ . Thus, the simplified Eq. (89) can be applied in numerical models to calculate the multi-layer topographic vertical velocity.

Figure 1b shows the linear variation of the unit topographic vertical velocity intensity with altitude at the given model resolution. The results indicated that with the increase of model resolution, the topographic vertical velocity decreases rapidly with altitude.

When  $L=10\text{km}$ , that is high-resolution numerical models and  $\Delta p=10\text{hPa}$ , there is almost only one layer in the model, and the topographical vertical velocity decreases to be negligible. In low-resolution numerical models ( $L=150\text{ km}$ ), when  $\Delta p \geq 150\text{hPa}$  10hPa above the surface which is lower than the next layer of the lowest model vertical layer in CAM5-SE, so a single layer parameterization scheme is enough. For  $L=150\text{km}$ , the influence of topography on the vertical motion can be negligible. Therefore, the influence reaches up to 150hPa above the attenuation of surface, so multi-layer topographic vertical velocity can be ignored in high-resolution numerical models but can be considered in low-resolution numerical models parameterization scheme is necessary. It can provide some new information for numerical simulations. Notably, preprocessing the sub-grid topographic data before the model integration may increase a small amount of computation compared with CAM5-SE simplify calculation.

[Insert Figure 1]

The trigonometric function of slope and aspect calculated by Eq. (56) is parameterized to the model dynamic processes to evaluate the topographic vertical motion. A realistic statistical method based The topography data used in this study is from the United States Geological Survey (USGS) DEM with a resolution of  $1\text{ km} \times 1\text{ km}$  (Sub Grid). The simulations are performed at the horizontal resolution of different model grid (Coarse Grid). Thus, the coarse grid contains several sub grids. We define a coarse grid as a terrestrial grid when the number of sub-grids on land occupies more than 10% of the total number of sub-grids, otherwise it is a marine grid. If the number of sub-grids with slope  $\geq 5^\circ$  in the terrestrial grid exceeds 10%, the terrestrial grid is considered as a complex topographic area coarse grid and needs to be parameterized. After that, the product of the trigonometric function transformation to calculate sub-grid functions of the slope and aspect for describing the orographic characteristics of complex areas over the globe has been proposed and tested in of each sub-grid in complex topographic area coarse grid is calculated, that is  $\tan\theta_N \times \cos\varphi_N$  (TC) and  $\tan\theta_N \times \sin\varphi_N$  (TS). According to Wang et al. (2022). Since  $\tan(\text{slope}) \times \cos(\text{aspect})$  ( $\tan\theta_N \cos\varphi_N$  in Eq. (5)) and  $\tan(\text{slope}) \times \sin(\text{aspect})$  ( $\tan\theta_N \sin\varphi_N$  in Eq (5)) conform to the (2022), it was found that the sub grids contained in the coarse grids of all topographic areas follow Gaussian distribution, it is transformed into the standard normal distribution (standardized transformation) before processing. According to the probability P table, the value of  $Z_p$  corresponds to Z during Z-Scores ( $Z = \frac{x-\mu}{\sigma}$ ,  $\mu$  denotes the average, and  $\sigma$  indicates the standard deviation). Then at the representative value of several sub-grid topography values at the model coarse grid scale is selected ( $y_p = \mu + Z_p * \sigma$ ) and can be easily described and applied. (Wang et al., 2022). Finally, bring the representative value into Eq.(6) to calculate  $\omega_s$ . Before the experiments, advanced preprocessing is used to calculate the probability densities of the trigonometric function and grid weights.



## 2.3 Experimental design and data

The CAM stand-alone model can be run using CESM scripts, which is coupled to a data ocean model, a thermodynamic sea ice model and an active land model, when one of “F” component sets of CESM is chosen. We choose the F\_2000\_CAM5 component set of CESM to conduct numerical experiments. The simulations are performed at the horizontal resolution of ne30 (about 1°) and 30 hybrid sigma-pressure levels, with an integration time step of 1800 s. Three 6-year simulations are forced by the prescribed current sea surface temperature and sea ice range with seasonal variations and are recycled yearly (Stone et al. 2018). The one without any modification is the control experiment (Ctl experiment). The others are the sensitivity experiments, which are the same as the control experiment but consider the lowest topographic vertical velocity (Single experiment) and the decrease of multi-layer topographic vertical velocity (Multi experiment). All the three cases are carried out for 6 years, and the first year of simulation is discarded ~~to avoid any~~ spin-up..

~~The topography data used is the United States Geological Survey (USGS) DEM data with the resolution of 1km×1km.~~ The Global Precipitation Climatology Project (GPCP) Level 3 Monthly 0.5-Degree V3.0 beta (Huffman et al. 2019) from 1987 to 2016 is used to evaluate the simulated precipitation. Monthly mean atmospheric data, comprising surface pressure, specific humidity, zonal and meridional wind ((at 11 vertical levels from 1000 to 700 hPa) during 1991–2021, are from the European Centre for Medium-Range Weather Forecasts Reanalysis 5 data set (ERA5) on a 0.25° × 0.25° grid (~~Hans et al., 2020~~). used for comparison with model results (Hans et al., 2020). And the lowest model layer wind is derived from the ERA-Interim at a 0.25° horizontal grid spacing and 60 model levels.

## 2.4 Improvement or divergence ratio

Divergence ratio is an indicator used to measure the difference ratio between simulation results and observation results. Improvement ratio is an indicator used to measure the improvement ratio between Single (Multi) and Ctl experiments. In mountain meteorology, the precipitation enhancement ratio (PER) is the ratio of the precipitation  $P$  at mountain peak or some other selected points to the precipitation at the reference point or in the reference region  $P_{REF}$ , as presented in Eq. (910).

$$PER = \frac{P}{P_{REF}}, \quad (910).$$

The reference region should be far enough removed that it is unaffected by the mountain, but still in the same climate zone (Smith 2019). We extend Eq. (910) to any physical quantity to obtain Eq. (4011).

$$PER = \frac{\Delta P}{P_{REF}}, \quad (4011)$$

where  $\Delta P$  indicates the difference in simulations between the sensitivity and control experiments or the difference between the simulations from the control experiment and observation data.  $P_{REF}$  represents simulations from the control experiment. Then, the PER reflects the improvement ratio or divergence ratio.

### 3 Results and discussion

#### 3.1 Precipitation simulation over the Tibetan Plateau

A region of 22°N–45°N and 70°E–105°E is selected to cover the Tibetan Plateau. The Tibetan Plateau is influenced by the plateau monsoon and has a distinct seasonal pattern of wet summer and dry winter (Su et al. 2013). The precipitation reaches its annual maximum in summer, accounting for 60%–70% of the annual accumulated precipitation (Yanai and Wu 2006; Wang et al. 2018). Therefore, summer precipitation is of great significance for this study in the region.

The geographical distributions of boreal summer (June–August, JJA) mean precipitation amount from GPCP, Ctl, Single and Multi experiments are shown in Fig. 2. In summer, most precipitation over East Asia is related to the Indian summer monsoon and the East Asian summer monsoon (Tao and Chen 1987). The results indicate that for the GPCP (Fig. 2a), a large rainfall amount is concentrated in the Bay of Bengal and the southeastern periphery of the Tibetan Plateau, but for the simulations from the Ctl (Fig. 2b), Single (Fig. 2c) and Multi (Fig. 2d) experiments, little rainfall is received in these areas. However, the precipitation increase appears on the southern slope of the Tibetan Plateau in model experiments, but there is little rainfall in this region in GPCP.

[Insert Figure 2]

In order to illustrate the biases of the model simulation and the improvement of the topographic vertical motion scheme, the differences in the summer precipitation between sensitivity experiments, Ctl experiment and GPCP are shown in Fig. 3. The most striking feature of the bias distribution is its close relation with topography. Positive precipitation bias controlling the Tibetan Plateau has been a common error in many climate models for a long time (Yu et al., 2015). The largest overestimations of the Ctl experiment (Fig. 3c) are found over the eastern and southern edges of the Tibetan Plateau, mostly in the regions with altitudes of 500 m and 4000 m. According to Eq. (4011), the divergence ratio is about 80% (Fig. 3f). In addition, the larger underestimations of precipitation can be found in front of the southern slope of the Tibetan Plateau, mostly in the region below the altitude of 500m. The region with the

largest underestimation is located in the area of 22°N, 90–98°E, with an underestimation ratio of about 100%. However, underestimation ratios in other regions are 20–40%. This result indicates that the southwesterly wind transports the water vapor from the ocean to the southern slope of the Tibetan Plateau. Due to the mountains, the airflow climbs upward and produces plenty of precipitation. The simulation bias is that the condensate that should have been generated in the Bay of Bengal is brought to the southern slope of the Tibetan Plateau. It is noteworthy that after considering the topographic vertical velocity, the simulation results are remarkably improved. The positive precipitation deviations in the southern and eastern edges of the Tibetan Plateau and the negative deviations in the low-altitude region of the windward slope are obviously improved. Moreover, the Multi experiment (Fig. 3b) performs better than the Single experiment (Fig. 3a), and the improvement ratios of positive deviations for the Single and Multi experiments are both 20%–30% (Fig. 3d and 3e). The results above indicate that the modification of topographic vertical velocity plays a vital role in topographic precipitation simulations.

[Insert Figure 3]

More details of model performance and precipitation variations are revealed by the meridional and latitudinal averages of precipitation over the Tibetan Plateau. The meridional average precipitation though the Tibetan Plateau over 87°E–95°E (Fig. 4b) suggests that the precipitation peak for the Ctl (green line) is located north of the GPCP (black line), but more precipitation than GPCP. The precipitation distribution for the Single (blue line) experiment is the same as that for the Ctl experiment. However, the peak in Multi experiment (red line) is located north of GPCP, but the rain intensity is nearly equal. This result indicates that considering the decaying of multi-layer vertical velocity can significantly reduce the overestimation of precipitation over the south foot of the Tibetan Plateau. Fig. 4a shows the latitudinal average of precipitation over 22°N–25°N. Compared with the GPCP (black line), the Ctl experiment (green line) considerably underestimates the rainfall in front of the southern edge of the Tibetan Plateau. At the eastern peak 91°E, the difference between Ctl and GPCP is about -8.41 mm day<sup>-1</sup>, and the maximum value of Multi experiment (14.51 mm day<sup>-1</sup>) presents similar magnitude to that of GPCP (17 mm day<sup>-1</sup>). At the windward peak 26°N, the difference between Ctl and GPCP is about 12.5 mm day<sup>-1</sup>, and the value of Single experiment (14.1 mm day<sup>-1</sup>) presents similar magnitude to that of GPCP (14.22 mm day<sup>-1</sup>).

[Insert Figure 4]

~~To further investigate the impact of vertical circulation on precipitation simulations, Figure 5 displays the vertical pressure velocity, meridional vertical circulation and their difference averaged over 87°E–95°E. It can be found that for the Single, Multi and Ctl experiments (Fig. 5a–5c), there is strong southerly wind near 27°N–38°N, but the Ctl experiment does not simulate the variability of the vertical velocity. The vertical motion~~

for the Single and Multi experiments appears at 28°N, which is an essential factor of orographic precipitation. Fig. 5d and 5e visually show the differences between the vertical pressure velocity and meridional-vertical circulation among the Single, Multi and Ctl experiments. Compared with Ctl experiment, the mountain blocking for the affects the Indian summer monsoon, weakening the southerly wind component. Due to the stronger vertical motion, the vertical and southerly wind components for the Multi experiment are stronger than those for the Single experiment.

[Insert Figure 5]

In terms of the biases of model simulations, Fig. 65 presents differences in convective precipitation, large-scale precipitation, shallow convective precipitation and ZM convective precipitation between the simulations and GPCP. The deviations in the convective precipitation present almost the same spatial pattern (Fig. 6a5a and 6e5e) as the total precipitation (Figs. 3a–3b), especially along the southern and eastern edges of the Tibetan Plateau. The deviation in the spatial pattern of large-scale precipitation is slightly different (Fig. 6b5b and 6f5f). The Single and Multi experiments only revise the positive deviations of precipitation in the middle region of the southern slope (28°N–32°N, 82°E–88°E), and the simulations of Multi experiment are slightly higher than those from the Single experiment. However, both Single and Multi experiments greatly improve the negative deviations of precipitation in front of the southern slope (22°N–25°N, 90°E–97°E). The deviations in the spatial pattern of shallow convective precipitation (Fig. 6e5c and 6g) show almost the same between Single and Ctl experiments and between Multi and Ctl experiments, and the most negative deviations are both located at altitudes above 500 m. In the regions with altitudes below 500 m, the deviation of the ZM convective precipitation (Fig. 6d5d and 6h5h) presents almost the same spatial pattern as that of the convective precipitation (Fig. 6a5a and 6e5e).

[Insert Figure 65]

To further analyze which type of precipitation improvement is dominant, we investigate the contributions of convective precipitation, large-scale precipitation, ZM convective precipitation and shallow convective precipitation to the improvement of total precipitation simulations (Fig. 76). The results suggest that for the improvement of the overestimation of total precipitation at altitudes from 500m to 4000 m (pink shaded areas in Fig. 3c), the Multi experiment performs better than the Single experiment. The total precipitation overestimation of 12.9 mm day<sup>-1</sup> is improved by 6.23 mm day<sup>-1</sup> for the Multi experiment and 3.23 mm day<sup>-1</sup> for the Single experiment (Fig. 7a6a). For the improvement of the total precipitation simulations in the Multi experiment, the improvement of convective precipitation (4.83 mm day<sup>-1</sup>) accounts for the largest proportion part, while the improvement of large-scale precipitation is only 1.4 mm day<sup>-1</sup>. This is due to the fact that the water vapor is lifted higher by the topographic vertical motion in the Multi experiment, which is favorable for triggering convective precipitation. In terms of convective precipitation, there is little difference in the

improvement between the shallow convective and ZM convective precipitation, and the improvements of precipitation simulations are both about  $2 \text{ mm day}^{-1}$ . The improvement of precipitation simulation for the Single experiment is similar to that for the Multi experiment, but the large-scale precipitation negatively contributes to the improvement of total precipitation in the Single experiment. Below 500 m, the underestimation of the total precipitation is about  $3 \text{ mm day}^{-1}$ , and the Single and Multi experiments both improve  $\sim 1.2 \text{ mm day}^{-1}$ , but the composition of precipitation types contributing to the improvement is different (Fig. 7b6b). In the Single experiment, the decrease of biases comes mainly from the improvement of large-scale precipitation simulation, and the improvement of convective precipitation can be negligible. This is because in the Single experiment, the water vapor of the whole layer is lifted, and therefore the improvement of total precipitation simulation is dominated by the improvement of large-scale precipitation simulation. However, the contribution of convective precipitation to the improvement of total precipitation simulation is greater than that of the large-scale precipitation in the Multi experiment. Moreover, ZM convective precipitation is the dominant precipitation type in convective precipitation, and shallow convective precipitation makes a negative contribution to the improvement of total precipitation simulation.

[Insert Figure 6]

### 3.2 Circulation simulation over the Tibetan Plateau

To further investigate the impact of vertical circulation on precipitation simulations, Figure 5 displays the vertical pressure velocity, meridional vertical circulation and their difference averaged over  $87^{\circ}\text{E}$ – $95^{\circ}\text{E}$ . It can be found that for the Single, Multi and Ctl experiments (Fig. 7a–7c), there is strong southerly wind near  $27^{\circ}\text{N}$ – $38^{\circ}\text{N}$ , but the Ctl experiment does not simulate the variability of the vertical velocity. The vertical motion for the Single and Multi experiments appears at  $28^{\circ}\text{N}$ , which is an essential factor of orographic precipitation. Fig. 7d and 7e visually show the differences between the vertical pressure velocity and meridional-vertical circulation among the Single, Multi and Ctl experiments. Mountain blocking has an impact on the Indian summer monsoon, reducing the southerly wind component in Single and Multi experiments compared to Ctl experiment. Due to the stronger vertical motion, the vertical and southerly wind components for the Multi experiment are stronger than those for the Single experiment.

[Insert Figure 7]

Since the differences in the total precipitable water (TPW) and 10m wind are related to precipitation, we analyze the distributions of the spatial differences of the 10m wind and TPW for the Single, Multi and Ctl experiments over the Tibetan Plateau (Fig. 8). Compared with Ctl experiment, the TPW shows negative deviations in the southern and eastern edges of the Tibetan Plateau in both the Single and Multi experiments. In front

445 of the southern slope ~~(windward)~~, the TPW presents positive deviations in the Multi  
experiment (Fig. 8a) but negative deviations in the Single (Fig. 8b), indicating that the  
Multi experiment improves the precipitation simulation in front of the windward slope  
and allows the water vapor transported to the front of the southern slope of the Tibetan  
Plateau with the Asian monsoon. This result is consistent with the precipitation  
450 distribution in Fig. 3. Also, the 10m wind can prove this result. In the Single and Multi  
experiments, the wind speed in high altitude regions decreases. However, only in the  
Multi experiment, there are positive deviations at the southern foot of the Tibetan  
Plateau, i.e., low-altitude windward-slope regions (Fig. 8a–8b).

[Insert Figure 8]

455 Water vapor transport is a critical factor in determining precipitation distribution and  
an essential quantity for the orographic precipitation is the horizontal water vapor flux.  
As shown in Fig. 9, the water vapor transported from the northern Indian Ocean reaches  
the coast of the Asian continent along the Indian peninsula and the Bay of Bengal in the  
Ctl (Fig. 9c), Single (Fig. 9a), Multi (Fig. 9b) experiments and ERA5 (Fig. 9d). After  
460 that, the water vapor is separated into two branches, one of which reaches the southern  
slope of the Tibetan Plateau and flows eastward after being blocked by the plateau. The  
other branch transports eastward. Compared with the Ctl experiment, more water vapor  
is transported from the northern Indian Ocean in the Multi experiment, and more water  
vapor converges in front of the southern slope of the Tibetan Plateau (80°E–87°E,  
465 24°N–26°N), but less water vapor climbs the slope. Additionally, the water vapor  
transported eastward weakens due to the blocking of the plateau, forming a weakened  
"water belt". It can be explained by Yu et al. (2015), i.e., the altitude of land surface  
jumps from lower than 200 m to more than 4000 m within approximately 4 model grids,  
and the CAM5 (Ctl experiment) allows the multi-grid transport and spurious  
470 accumulated water vapor at cold and high-altitude regions. In contrast, the scheme of  
multi-layer topographic vertical motion implemented in the Multi experiment considers  
the climbing and bypassing of airflow. Thus, in the Multi experiment, water vapor is  
more in low-altitude regions and less in high-altitude regions. As a result, the  
precipitation is more in front of the slope and less in the southern slope of the Tibetan  
475 Plateau, which is consistent with the previous conclusion of total precipitation (Fig. 3).  
When the water vapor transports northward, there is a branch of water vapor in East  
Asia, which moves northwestward after bypassing westward and weakens markedly.  
This leads to a decrease in precipitation on the eastern edge of the Tibetan Plateau.  
Therefore, the differences between the simulations and observations, the excessive  
480 precipitation on higher slopes and less precipitation on lower slopes are considerably  
improved. In terms of the Single experiment, the variation of water vapor presents  
almost the same spatial pattern as that in the Multi experiment but less than in the Multi  
experiment. The only difference is that there is no noticeable increase in water vapor in  
lower slopes due to less pronounced variation in precipitation. Rahimi et al. (2019)  
485 investigated the relationship between the location of precipitation peak along slopes  
and horizontal resolution, and they found that finer resolution could allow the peak

location to move northward. Previous studies found that the orographic drag of complex topography may only be resolved at horizontal resolutions of a few kilometers or even finer resolutions (Sandu et al., 2016; Wang and Zhang, 2020). However, our research demonstrates that considering the sub-grid parameterization scheme of slope gradient and surface and adding the topographic vertical motion in the CAM5-SE can address the impacts of topographic complexity on precipitation. It significantly improves the underestimation of precipitation over the windward slope of the Tibetan Plateau and the overestimation of precipitation over the steep edge of high mountains at the horizontal resolutions of hundred kilometers, which is equivalent to the horizontal resolutions of a few kilometers or ~~the integration step of~~ a few months simulation in climate models (Li et al. 2022).-

[Insert Figure 9]

Upslope flow is critical for orographic precipitation, which allows air to climb over mountains more easily (Smith 2019). Figure 10 presents the meridional-vertical cross-section of water vapor transport along 90°E. The results suggest that for the Single and Multi experiments (Fig. 10a and 10b), the vertical water vapor transport considerably enhances from 27°N, and even the lifting height in the Multi experiment is higher than that in the Single experiment. Compared with the Ctl experiment, the lifting height of water vapor reaches about 700 hPa in the Single experiment (Fig. 10d), while it reaches about 650 hPa in the Multi experiment (Fig. 10e). The upslope flow supplies the water vapor to the windward slope, and the airflow blocking reduces the precipitation over the region above 500 m.

[Insert Figure 10]

### 3.3 Precipitation simulation in other complex terrain areas

A similar precipitation response can be found in other high mountains, such as the Andes in South America. Figure 11 shows the biases of precipitation simulated in the Single, Multi and Ctl experiments in South America during austral summer (December to February). It can be found that in December–February, there is strong southerly wind at 850 hPa (Figs. 11a–11b) on the western edge of the Andes (from west of 30°S to 10°S), and large positive precipitation biases can be found in front of the foot of the Andes (Fig. 11c). In the Ctl experiment, the precipitation is overestimated on ridges above 1000 m and is underestimated in some low-altitude regions on the eastern slope. These biases are closely associated with the strong wind at 850 hPa on the eastern edge of the Andes. In both Single and Multi experiments (Figs. 11a and 11b), the overestimation of precipitation decreases on ridges above 1000 m and increases in the windward slope at the eastern region of the Andes.

[Insert Figure 11]

525 The distributions of spatial differences in the specific humidity and TPW in South  
America for the Single, Multi and Ctl experiments are shown in Fig. 12. Similar to on  
the Tibetan Plateau, compared with the Ctl experiment, the TPW shows negative  
deviations in mountain tops in both the Single and Multi experiments, which is in  
agreement with the precipitation distribution in Fig. 11. However, the TPW on the foot  
530 of the northeastern slope (windward) only displays positive deviations in the Multi  
experiment but negative deviations in the Single experiment (Fig. 12a and b). This  
result suggests that the Multi experiment improves the precipitation simulation in front  
of the windward slope, and in both the Multi and Single experiments, the water vapor  
is transported to the eastern slope. Thus, the TPW accumulates in this area to form large  
positive deviations. The results for the specific humidity (Fig.12c–12d) and TPW are  
535 consistent. In the Single and Multi experiments, there are dry deviations in high-altitude  
regions. However, only in the Multi experiment, there are wet deviations at the southern  
foot of the Tibetan Plateau, i.e., the low-altitude windward-slope regions.

[Insert Figure 12]

540 Table 1 presents the root mean square error (RMSE) of precipitation simulations in  
several typical areas with complex terrain during boreal summer or winter (figure  
omitted). The results indicate that in the Tibetan Plateau, (70°E–105°E, 22°N–45°N,  
boreal summer precipitation), Equatorial New Guinea and Indonesia (100°E–150°E,  
10°S–10°N) ~~and~~, boreal summer precipitation), South America (30°W–90°W, 60°S–  
5°N), boreal winter precipitation), and North America (155°W–122°W, 30°N–65°N,  
545 boreal winter precipitation) the RMSE values of precipitation simulations in the  
sensitivity experiments are smaller than those in the Ctl experiment. For the Ctl  
experiment, the RMSE is the largest ~~in the over~~ Tibetan Plateau- (5.44) and the smallest  
over North America (1.57). Almost all GCMs have large deviations in precipitation  
simulations on the Tibetan Plateau. Therefore, after considering the dynamic lifting of  
550 topography, the improvement of biases in this area is the most pronounced- followed  
by Equatorial New Guinea (26.3%) and the smallest in North America (9.55%).  
Moreover, the improvement of the Multi experiment is better than that of the Single  
experiment, reaching about 29.23%, which indicates that the steeper the mountains are,  
the more obvious the influence of lifting condensation on multi-layer vertical velocity  
555 is. The impact of single topographic vertical motion is limited to low-altitude areas.  
However, in Africa, the surface is relatively flat, and the slope gradient is small- (Wang  
et al., 2022). Thus, the method in this research may not be as effective- so it is no longer  
mentioned in Table 1.

[Insert Table 1]

560 Notably, the improvement of precipitation simulations is noticeable over the Tibetan  
Plateau but not in the Rocky Mountain region in North America (figure omitted). The  
main reason is that in the Rocky Mountain region, the wind direction is parallel to the  
mountain range, and the angle between the prevailing wind direction on the western



565 side of the mountain (steep slope) and the slope surface is close to 90°. Thus, there can  
be no lifting motion caused by topography. The topographic vertical motion is not only  
dependent on the slope gradient, but also associated with the angle between the wind  
direction and the slope surface. Therefore, the large amount of water vapor from the  
ocean cannot be transported to the mountains. In order to understand and solve these  
570 remaining problems, more numerical experiments and more detailed analyses should  
be further conducted. Moreover, when we only consider the steep slope of mountains,  
it would greatly impact the precipitation simulation of the regional climate. Future  
research is also needed to investigate the possibility of applying the topographic vertical  
motion scheme to extreme precipitation simulation in local areas, allowing weather  
models to more accurately simulate extreme precipitation caused by topography.

## 575 **4 Conclusions and discussion**

A common bias of the AGCMs is the overestimation of orographic precipitation. One  
primary reason for this bias is the imperfection of the sub-grid terrain parameterization  
scheme. One critical reason is that the influence of topographic lifting on airflow and  
water vapor transport is not considered in numerical models. In this study, we  
580 investigate whether such excessive precipitation simulation can be improved by  
considering the topographic vertical velocity in the CAM5-SE. The results show that  
the simulated precipitation in steep regions is sensitive to topographic vertical velocity.  
In the Multi experiment, the underestimated total precipitation is remarkably improved  
at lower layers on steep windward slopes. A common bias of the AGCMs is the  
585 ~~overestimation of orographic precipitation. One primary reason for this bias is the~~  
~~imperfection of the sub-grid terrain parameterization scheme. One critical reason is that~~  
~~the influence of topographic lifting on airflow and water vapor transport is not~~  
~~considered in numerical models. In this study, we investigate whether such excessive~~  
~~precipitation simulation can be improved by considering the topographic vertical~~  
590 ~~velocity in the CAM5-SE. The results show that the simulated precipitation in steep~~  
~~regions is sensitive to topographic vertical velocity. In the Multi experiment, the~~  
~~underestimated total precipitation is remarkably improved at lower layers on steep~~  
~~windward slopes.~~ However, in the Ctl experiment, there are large dry biases, **andbut** the  
overestimation of precipitation in high-altitude areas of steep mountains is markedly  
595 reduced **in Multi experiment**. The increase of precipitation on steep windward slopes  
and the decrease of precipitation in high-altitude areas of mountains are mainly due to  
the contribution of convective precipitation, which is greater in the Multi experiment  
than in the Single experiment. The improvement of precipitation simulations is closely  
related to dynamic lifting. If the dynamic uplifting effect is not considered, every grid  
600 is flat without considering the slope gradient and slope surface. In this case, a large  
amount of water vapor accumulates in high-altitude areas on the top of mountains. This  
is partially responsible for the excessive water vapor and precipitation in high-altitude  
regions of steep mountains in the Ctl experiment.

Moreover, in this study, the sub-grid parameterization scheme of the topographic vertical motion performs well in precipitation simulations over complex terrains, such as the Tibetan Plateau and the Andes in South America. Moreover, the improvement of precipitation simulations for the Multi experiment is better than that for the Single experiment. As shown in Fig. 1a, with increasing numerical model resolution, the influence of topography on multi-layer vertical velocity weakens. Therefore, it is necessary to use high-resolution numerical experiments to verify whether the dynamic lifting effect of sub-grid topography on airflow still exists.

~~Notably, the improvement of precipitation simulations is noticeable over the Tibetan Plateau but not in the Rocky Mountain region in North America (figure omitted). The main reason is that in the Rocky Mountain region, the wind direction is parallel to the mountain range, and the angle between the prevailing wind direction on the western side of the mountain (steep slope) and the slope surface is close to 90°. Thus, there can be no lifting motion caused by topography. The topographic vertical motion is not only dependent on the slope gradient, but also associated with the angle between the wind direction and the slope surface. Therefore, the large amount of water vapor from the ocean cannot be transported to the mountains. In order to understand and solve these remaining problems, more numerical experiments and more detailed analyses should be further conducted. Moreover, when we only consider the steep slope of mountains, it would greatly impact the precipitation simulation of the regional climate. Future research is also needed to investigate the possibility of applying the topographic vertical motion scheme to extreme precipitation simulation in local areas, allowing weather models to more accurately simulate extreme precipitation caused by topography.~~

**Acknowledgments:** The authors would like to thank the administrator of Beijing Normal University High Performance Computing for providing the high-performance computing (HPC) environment and technical support.

**Author contributions:** YQW and LNW designed the experiments and the scope and structure of the manuscript. YQW carried out the simulations and analyzed the results with help from JF, ZYS, QZW and HQC. All authors gave comments and contributed to the development of the paper.

**Funding:** The research work presented in this paper was supported by the Marine S&T Fund of Shandong Province for Pilot National Laboratory for Marine Science and Technology (Qingdao) (2022QNL010202) and the National Natural Science Foundation of China (41830536).

**Code and Data availability:** GPCP V3.0 data is available from <https://doi.org/10.5067/TTO0VJF2FSYR> (last access: 29 June 2022). DEM data can be found at <http://usgs.gov/>. The ERA5 atmospheric datasets used in this study is available from: <https://doi.org/10.24381/cds.6860a573> (last access: 29 June 2022). The source code of CAM-SE5.3 is available from <http://www.cesm.ucar.edu/models/cesm1.2/> (last

access: 20 April 2022). The dataset related to this paper is available online via Zenodo:  
<https://doi.org/10.5281/zenodo.7256923>.

645 **Competing interests.** The authors declare that they have no conflict of interest.

## Reference


- 650 Akinsanola A A, Ongoma V, Kooperman G J.: Evaluation of CMIP6 models in  
simulating the statistics of extreme precipitation over Eastern Africa. *Atmos. Res.*  
254: 105509 , 2021.
- Alpert P, Shafir H. Mesoy-Scale Distribution of Orographic Precipitation: Numerical  
Study and Comparison with Precipitation Derived from Radar Measurements. *J*  
*Appl Meteorol*, 28(10):1105-1117 , 1989.
- 655 Alpert, P., F. Jin, and H. Shafir.: Orographic precipitation simulated by a super-high  
resolution global climate model over the Middle East, *National Security and*  
*Human Health Implications of Climate Change*, 1, 301–306, doi:10.1007/978-94-  
007-2430-3\_26 , 2012.
- Attada R, Dasari H P, Kunchala R K, et al.: Evaluating cumulus parameterization  
schemes for the simulation of Arabian Peninsula winter rainfall. *Journal of*  
660 *Hydrometeorology*. 21(5): 1089-1114 , 2020.
- [Bretherton, C. S., and S. Park, A new moist turbulence parameterization in the  
community atmosphere model, \*J. Climate\*, 22, 3422–3448, 2009a.](#)
- Chan, S. C., E. J. Kendon, H. J. Fowler, S. Blenkinsop, C. A. T. Ferro, and D. B.  
Stephenson.: Does increasing the spatial resolution of a regional climate model  
665 improve the simulated daily precipitation? *Clim. Dyn.* 41, 1475–1495 , 2013.
- Chao, W. C.: Correction of excessive precipitation over steep and high mountains in a  
GCM. *J. Atmos. Sci.*, 69, 1547–1561 , 2012.
- Chen HM, Zhou TJ, Neale RB, Wu XQ, Zhang GJ.: Performance of the new NCAR  
CAM3.5 in East Asian summer monsoon simulations: sensitivity to modifications  
670 of the convection scheme . *J. Clim.* 23(13): 3657– 3675, 2010.
- Codron F & Sadourny R.: Saturation limiters for water vapour advection schemes:  
impact on orographic precipitation. *Tellus A* 54(4):338–349 , 2002.
- Cui T, Li C, Tian F.: Evaluation of temperature and precipitation simulations in CMIP6  
models over the Tibetan Plateau. *Earth and Space Science*. 8(7): e2020EA001620 ,  
675 2021.
- Dennis, J., Edwards, K., Evans, J., Guba, O., Lauritzen, P. H., Mirin, A. A., St-Cyr, A.,  
Taylor, M. A., & Worley, P. H.: CAM-SE: A scalable spectral element dynamical  
core for the Community Atmosphere Model. *International Journal of High*  
*Performance Computing Applications*, 26(1), 74– 89. 2012.
- 680 Done, J. M., L. R. Leung, C. A. Davis, and B. Kuo.: Regional climate simulation using  
the WRF model. Preprints, Fifth WRF/14th MM5 Users' Workshop, Boulder, CO,  
National Center for Atmospheric Research, P8, 2004.
- Fonseca, R. M., Zhang, T., and Yong, K.T.: Improved simulation of precipitation in the  
tropics using a modified BMJ scheme in the WRF model, *Geosci. Model Dev.*, 8,  
685 2915–2928 , 2015.
- Gettelman A., and Coauthors.: Global simulations of ice nucleation and ice  
supersaturation with an improved cloud scheme in the community atmosphere  
model. *J. Geophys. Res.*, 115, D18216 , 2010.
- [Gettelman, A., J. E. Kay, and J. T. Fasullo.: Spatial decomposition of climate feedbacks  
in the Community Earth System Model. \*J. Climate\*, 26, 3544– 3561 , 2013.](#)
- 690

- Hans, H., Bell, B., Berrisford, P., Hirahara, S., Horanyi, A., Munoz-Sabater, J., et al.: The ERA5 global reanalysis. *Journal of the Meteorological Society of Japan*, 146(730), 1999–2049. <https://doi.org/10.1002/qj.3803>, 2020.
- 695 Huffman, G.J., D.T. Bolvin, E.J. Nelkin.: GPCP Precipitation Level 3 Monthly 0.5-Degree V3.0 Beta. Greenbelt, MD: NASA GES DISC, 2019.
- Hurrell J W, Holland M M, Gent P R, et al.: The community earth system model: a framework for collaborative research. *Bull Am Meteorol Soc* 94:1339–1360 , 2013.
- 700 Jia K, Ruan Y, Yang Y, Zhang C.: Assessing the Performance of CMIP5 Global Climate Models for Simulating Future Precipitation Change in the Tibetan Plateau. *Water*. 11(9):1771, 2019.
- [Kasahara A \(1974\) Various vertical coordinate systems used for numerical weather prediction, \*Mon. Wea. Rev.\*, 102, 509–522.](#)
- 705 Kimoto, M., N. Yasutomi, C. Yokoyama, and S. Emori.: Projected changes in precipitation characteristics around Japan under the global warming. *SOLA*, 1, 85–88 , 2005.
- Kunz, M., and C. Kottmeier.: Orographic enhancement of precipitation over low mountain ranges. Part II: Simulations of heavy precipitation events over Southwest Germany, *J. Appl. Meteorol. Climatol.*, 45, 1041–1055 , 2006.
- 710 Liang Y, Yang B, Wang M, et al.: Multiscale simulation of precipitation over East Asia by variable resolution CAM-MPAS. *Journal of Advances in Modeling Earth Systems*, 13(11): e2021MS002656, 2021.
- Li, G., Chen, H., Xu, M. et al.: Impacts of Topographic Complexity on Modeling Moisture Transport and Precipitation over the Tibetan Plateau in Summer. *Adv. Atmos. Sci.* 39, 1151–1166 , 2022.
- 715 Lin C, Chen D, Yang K, et al.: Impact of model resolution on simulating the water vapor transport through the central Himalayas: implication for models’ wet bias over the Tibetan Plateau. *Clim. Dyn.* 51(9): 3195-3207, 2018.
- [Lin SJ.: A vertically Lagrangian finite volume dynamical core for global models. \*Mon Wea Rev\* 132: 2293–2397 , 2004.](#)
- 720 Liu Z, Mehran A, Phillips T J, et al.: Seasonal and regional biases in CMIP5 precipitation simulations. *Clim. Res.* 60(1): 35-50, 2014.
- Mlawer EJ, Taubman SJ, Brown PD, Iacono MJ, Clough SA.: Radiative transfer for inhomogeneous atmospheres: RRTM, a validated correlated-k model for the longwave. *J Geophys Res* 102(D14):16,663–16, 1997.
- 725 Morrison, H., and A. Gettelman.: A new two-moment bulk stratiform cloud microphysics scheme in the NCAR Community Atmosphere Model (CAM3). Part I: Description and numerical tests. *J. Climate*, 21, 3642–3659, 2008.
- Navale A, Singh C.: Topographic sensitivity of WRF-simulated rainfall patterns over the North West Himalayan region. *Atmos Res* 242:105003, 2020.
- 730 Neale RB, Chen C, Gettelman A, Lauritzen PH, Park S, Williamson DL, Conley AJ, Carcia R, Kinnison D, Lamarque J, Marsh D, Mills M, Smith AK, Tilmes S, Morrison H, Cameron-Smith P, Collins WD, Iacono MJ, Easter RC, Ghan SJ, Liu

- 735 X, Rasch PJ, Tayloy MA.: Description of the NCAR community atmosphere  
model (CAM 5.0). NCAR tech note TN-486 , 2010.
- Park S and Bretherton CS.: The University of Washington Shallow Convection and  
Moist Turbulence Schemes and Their Impact on Climate Simulations with the  
Community Atmosphere Model, *J. Clim.* 22, 3449–3469, 2009.
- 740 Rahimi, S. R., C. L. Wu, X. H. Liu, and H. Brown.: Exploring a variable-resolution  
approach for simulating regional climate over the Tibetan Plateau using VR-  
CESM. *J. Geophys. Res.*, 124, 4490–4513, 2019.
- Richter JH, Rasch PJ.: Effects of convective momentum transport on the atmospheric  
circulation in the community atmosphere model, version 3. *J. Clim.* 21(7):1487–  
1499 , 2008.
- 745 Roe, G. H.: Orographic precipitation, *Annu. Rev. Earth Planet. Sci.*, 33, 645– 671, 2005.
- Sandu, I., P. Bechtold, A. Beljaars, A. Bozzo, F. Pithan, T. G. Shepherd, and A. Zadra.:  
Impacts of parameterized orographic drag on the Northern Hemisphere winter  
circulation. *Journal of Advances in Modeling Earth Systems*, 8, 196–211, 2016.
- 750 Shen S, Xiao H, Yang H, et al.: Variations of water vapor transport and water vapor-  
hydrometeor-precipitation conversions during a heavy rainfall event in the Three-  
River-Headwater region of the Tibetan Plateau. *Atmospheric Research*. 264:  
105874 , 2021.
- Shen Y, Zhang Y, Qian Y.: A Parameterization Scheme for the Dynamic Effects of  
Subgrid Topography and Its Impacts on Rainfall Simulation. *Plateau Meteorology*.  
755 26(4): 655-665 (in Chinese, 2007).
- [Simmons AJ and Strüfing R \(1981\) An energy and angular-momentum conserving  
finite difference scheme, hybrid coordinates and medium-range weather prediction,  
Technical Report ECMWF Report No. 28, European Centre for Medium–Range  
Weather Forecasts, Reading, U.K.,68 pp.](#)
- 760 Smith, R. B.: 100 Years of progress on mountain meteorology research. A Century of  
Progress in Atmospheric and Related Sciences: Celebrating the American  
Meteorological Society Centennial, *Meteor. Monogr.*, No. 59, Amer. Meteor. Soc.,  
<https://doi.org/10.1175/AMSMONOGRAPHS-D-18-0022.1>, 2019.
- 765 Stone, D., Risser, M. D., Angelil, O., Wehner, M., Cholia, S., Keen, N., Krishnan, H.,  
Obrien, T. A., and Collins, W. D.: A basis set for exploration of sensitivity to  
prescribed ocean conditions for estimating human contributions to extreme  
weather in CAM5.1–1degree, *Weather Clim. Extremes*, 19, 10–19, 2018.
- Su F, Duan X, Chen D, Hao Z, Cuo L.: Evaluation of the global climate models in the  
CMIP5 over the Tibetan Plateau. *J. Clim.* 26: 3187–3208, 2013.
- 770 Tao, S. Y., and L. X. Chen.: A review of recent research on the East Asian summer  
monsoon in China. *Monsoon Meteorology*, C. P. Chang and T. N. Krishnamurti,  
Eds., Oxford University Press, 60–92, 1987.
- Wang X., Pang G., Yang M.: Precipitation over the Tibetan Plateau during recent  
decades: a review based on observations and simulations. *Int. J. Climatol*, 38, pp.  
775 1116-1131,2018.

- Wang, Y. and Zhang, G. J.: Global climate impacts of stochastic deep convection parameterization in the NCAR CAM5, *J. Adv. Model. Earth Sy.*, 8, 1641–1656, 2016.
- 780 Wang, Y., and Coauthors.: Synergy of orographic drag parameterization and high resolution greatly reduces biases of WRF-simulated precipitation in central Himalaya. *Climate Dyn.*, 54, 1729–1740 , 2020.
- Wang, Y., G. J. Zhang, and Y.-J. He.: Simulation of precipitation extremes using a stochastic convective parameterization in the NCAR CAM5 under different resolutions. *J. Geophys. Res. Atmos.*, 122, 12 875–12 891 , 2017.
- 785 Wang, Y., Wang, L., Feng, J. et al.: A statistical description method of global sub-grid topography for numerical models. *Clim Dyn.* <https://doi.org/10.1007/s00382-022-06447-2>,2022.
- Yanai, M., and Wu G. X. Effects of the Tibetan Plateau. *The Asian Monsoon*, B. Wang, Ed., Springer, 513–549 , 2006.
- 790 Yu, R., Li, J., Zhang, Y. et al.: Improvement of rainfall simulation on the steep edge of the Tibetan Plateau by using a finite-difference transport scheme in CAM5. *Clim Dyn* 45, 2937–2948 , 2015.
- Zhang GJ, McFarlane NA.: Sensitivity of climate simulations to the parameterization of cumulus convection in the Canadian Climate Centre general circulation model. *Atmos-Ocean* 33(3):407–446,1995.
- 795 Zhu YY and Yang S.: Evaluation of CMIP6 for historical temperature and precipitation over the Tibetan Plateau and its comparison with CMIP5. *Advances in Climate Change Research*. 11(3): 239-251, 2020.


800




**Figure 1.** (a) Distribution of surface topographic vertical velocity (Pa/s) at different slope and aspect in 10m/s wind speed; (b) the decreasing of the unit topographic vertical velocity with height at different grid scales.

805






810 **Figure 2.** Spatial distributions of summer (June–August) average precipitation amount ( $\text{mm day}^{-1}$ ) from (a) the GPCP data and simulation in (b) Ctl, (c) Single and (d) Multi experiments. Vectors in Fig. 2a represent the summer wind at [925 hPa](#) the lowest model level in ERA-Interim, vectors in Figs. 2b–2d represent the summer wind simulation at the lowest model level, and the black contour indicate the altitude of 3000 m.




815 **Figure 3.** Differences of summer average precipitation amount ( $\text{mm day}^{-1}$ ) (a) between Single and Ctl experiments, (b) between Multi and Ctl experiments and (c) between Ctl experiment and GPCP, improvement ratio of (d) Single experiment and (e) Multi experiment, (f) divergence ratio of Ctl. Black contours indicate the altitudes of 500 m and 4000 m.



820


**Figure 4.** Summer precipitation averaged over (a) 22°N–25°N and (b) 87°E–95°E. Green, red, blue and black lines represent the simulated precipitation in the Ctl, Multi, Single experiments and from the GPCP data, respectively. The grey dotted lines indicate the altitudes (km).

825




Figure

5.



830

**Figure 5.** Meridional-vertical circulation (vectors) and vertical velocity (shading) averaged over 87°E–95°E in (a) Single, (b) Multi and (c) Ctl experiments, and their differences (d) between the Single and Ctl experiments and (e) between the Multi and Ctl experiments.




835


**Figure 6.** Difference of (a) convective precipitation, (b) large-scale precipitation, (c) shallow convection precipitation, (d) precipitation from ZM convection between Single and Ctl experiments. (e-h) As in (a-d) but between Multi and Ctl experiments. Black contours indicate the altitudes of 500 m and 4000 m. Dotted areas are statistically significant at the 90% confidence level


840

(a) precipitation between 500 and 4000 m



(b) precipitation below 500 m






**Figure 76.** Difference of the precipitation types between the sensitivity and control experiments. (a) Positive deviations of precipitation simulations over the region with altitudes within 500–4000 m and (b) negative deviations of precipitation simulations over the region below 500 m.


845





850


**Figure 7.** Meridional-vertical circulation (vectors) and vertical velocity (shading) averaged over 87°E–95°E in (a) Single, (b) Multi and (c) Ctl experiments, and their differences (d) between the Single and Ctl experiments and (e) between the Multi and Ctl experiments.



**Figure 8.** As in Fig. 6 but for Difference of (a–b) total precipitable water ( $\text{kg m}^{-2}$ ) and (c–d) 10-m wind speed ( $\text{m s}^{-1}$ ) between Single, Multi and Ctl experiments. Dotted areas are statistically significant at the 90% confidence level.


855

860




**Figure 9.** Distribution of the composite whole-layer water vapor flux (from the lowest model level to the seventh model level) in the (a) Single, (b) Multi, (c) Ctl experiments and (d) ERA5 over East Asia. Black contours indicate the altitudes of 3000 m.

865




**Figure 10.** As in Fig. 5, but along 90°E meridional-vertical water vapor transport (vectors) and meridional water transport (shading) along 90°E in (a) Single, (b) Multi and (c) Ctl experiments, and their differences (d) between the Single and Ctl experiments and (e) between the Multi and Ctl experiments.

870



875

**Figure 11.** As in Fig.3a–c, but **Figure 11.** Differences of winter (December–February) average precipitation amount ( $\text{mm day}^{-1}$ ) (a) between Single and Ctl experiments, (b) between Multi and Ctl experiments and (c) between Ctl experiment and GPCP over South America. Vectors in Fig. 11a and 11b represent the 850 hPa wind in the Single and Multi experiments, respectively. Black contours indicate the altitudes of 1000 m and 2000 m.



880

**Figure 12.** Difference of (a)–(b) Same as Fig.8 total precipitable water (kg m<sup>-2</sup>) and (c)–(d), (e)–(d) same as (a)–(b) but for the lowest model level specific humidity (g/kg) between Single, Multi and Ctl experiments over South America. Black contours indicate the altitudes of 1000 m and 2000 m.

885

**Table 4R1.** RMSE in different regions.

Regions	Ctl experiment	Single experiment	Multi experiment
Tibetan Plateau ( <a href="#">70°E–105°E, 22°N–45°N</a> )	5.44	4.88 (10.3%)	3.85 (29.23%)
Equatorial New Guinea ( <a href="#">100°E–150°E, 10°S–10°N</a> )	2.55	2.2 (13.73%)	1.88 (26.3%)
South America ( <a href="#">30°W–90°W, 60°S–5°N</a> )	2.13	2.04 (4.23%)	1.91 (10.33%)
North America ( <a href="#">155°W–122°W, 30°N–65°N</a> )	<a href="#">1.57</a>	<a href="#">1.46 (7%)</a>	<a href="#">1.42(9.55%)</a>