

Can TROPOMI-NO₂ satellite data be used to track the drop and resurgence of NO_x emissions between 2019 - 2021 using the multi-source plume method (MSPM)?

Enrico Dammers¹, Janot Tokaya¹, Christian Mielke², Kevin Hausmann², Debora Griffin³, Chris McLinden³, Henk Eskes⁴, and Renske Timmermans¹

¹Netherlands Organisation for Applied Scientific Research (TNO), Utrecht, Princetonlaan 6, 3584 CB, The Netherlands

²Umweltbundesamt (UBA), Dessau-Roßlau, Wörlitzer Pl. 1, 06844, Germany

³Air Quality Research Division, Environment and Climate Change Canada, 4905 Dufferin St, M3H 5T4, Toronto, Canada

⁴KNMI, De Bilt, Utrechtseweg 297, 3731 GA, The Netherlands

Correspondence: Enrico Dammers (enrico.dammers@tno.nl)

Abstract. NO_x is an important primary air pollutant ~~-, dominantly- of major environmental concern which is predominantly~~ produced by anthropogenic ~~-, mostly combustion-based, activities from sectors such as industry, traffic and transport~~ combustion activities. NO_x ~~is directly linked to negative health and environmental impacts. Currently, the construction of emission inventories to~~ keep track of NO_x emissions is based on official national reported emissions and proxies such as activity data as well as ~~direct measurements. The effort to properly construct an accurate inventory is significant and time-consuming which causes a~~ reporting offset between one and five years with respect to the current date. Next to this temporal lag difficulties in composed ~~inventories~~ needs to be accounted for in national emission inventories, according to international treaties. Constructing accurate inventories requires substantial time and effort, resulting in reporting delays of one to five years. In addition to this, difficulties can arise from temporal and country specific legislative and protocol differences ~~between countries and over time in reporting~~ of emissions. ~~Satellite based-~~ To address these issues, satellite-based atmospheric composition measurements ~~provide offer~~ a unique opportunity ~~to fill this gap and independently estimate emissions on a large scale~~ for independent and large-scale estimation of emissions in a consistent, transparent ~~and comprehensible way. They give the possibility to check for compliance~~ with emission reduction targets in a timely manner as well as to observe rapid emission reductions such as experienced during the COVID-19 lock-downs. In this study we apply a consistent methodology to derive ~~-, and comprehensible manner. Here we~~ test the multi-source plume method (MSPM) to assess the NO_x emissions over Germany ~~for the years of in the Corona period~~ from 2019-2021. For the years where reporting is available, the differences between satellite estimates and inventory totals were within ~~400kt~~ 75-100 kt (NO₂) NO_x (<10% of inventory values). The large reduction of NO_x emissions (~15%) related to the COVID-19 lock-downs ~~were was~~ observed in both the inventory and satellite derived emissions. The recent projections for the inventory emissions ~~of 2021~~ pointed to a recovery of the ~~emissions towards pre-COVID19 levels~~ this 2021 emissions towards pre-COVID-19 levels. In the satellite derived emissions however, ~~such an~~ increase was not observed. While emissions from the larger power-plants did rebound to ~~earlier levels, others pre-COVID-19 levels, other~~ sectors such as road transport ~~and shipping did not and could be linked~~ did not, likely due to a reduction in the number of heavier transport trucks. This again

illustrates the value of having a consistent satellite based methodology for faster ~~projections~~ emission estimates to guide and check the conventional emission inventory reporting. The method described in this manuscript also ~~meet~~ meets the demand for independent verification of the official emission inventories, which will enable inventory compilers to detect potentially problematic reporting issues. ~~Transparency and comparability~~, bolstering transparency and comparability: two key values for emission reporting, ~~are thus bolstered by this technique~~.

1 Introduction

Nitrogen monoxide (NO) and dioxide (NO₂) play an important role in the atmospheric chemistry as they influence the abundance of tropospheric ozone (~~?~~) (Seinfeld and Pandis, 2006), and lead to aerosol formation. These primary air pollutants are collectively called nitrogen oxides (NO_x ≡ NO + NO₂). Since NO₂ is for the most part formed primarily through rapid oxidation of NO, their concentrations are strongly related. NO₂ is a major source of air pollution and exposure ~~to high concentrations~~ can result in ~~nausea, dyspnoea, headache and respiratory disease~~ significant health problems that cause an association between long-term exposure and reduced live-life expectancy (Atkinson et al., 2018; Belch et al., 2021). Hence ~~objective~~, objective concentration limits are set by the European ~~union~~ Union on the hourly (200 $\mu\text{g m}^{-3}\text{m}^{-3}$) and yearly (40 $\mu\text{g m}^{-3}\text{m}^{-3}$) NO₂ exposure levels ~~with recent WHO limits reducing this level further~~, with recent World Health Organization (WHO, 2021) limits reducing the annual mean limit to 10 $\mu\text{g m}^{-3}$ (yearly, WHO (2021)). ~~Next to these adverse health effect NO₂ causes m⁻³. As well as adverse health effects, NO₂ also places~~ a strain on the environment through soil and water acidification as well as eutrophication (Galloway et al., 2003).

Many anthropogenic activities contribute to the atmospheric NO₂ concentration ~~because~~ since NO₂ is formed in combustion processes where air (being about 80% nitrogen) is the oxidant. Natural sources of NO_x include lightning and soil emissions. The main sources of NO₂ emissions are the internal combustion engines that burn fossil fuels in motor vehicles and industry. The overall atmospheric evolution and budget of NO_x in the atmosphere has been determined with ever-increasing accuracy over the last decades. National environmental agencies are required to monitor the level of NO_x and the contribution of human activity on it according to international agreements, such as the ~~Geneva~~ Convention on Long-Range Transboundary Air Pollution (UNECE)(CLRTAP, <https://unece.org/environment-policy/air>) by the United Nations Economic Commission for Europe (UNECE). Efforts undertaken to limit NO_x emissions have resulted in strong reductions of ambient NO₂ concentration in many parts of the world (Jamali et al., 2020).

Inventories of NO_x emissions are commonly compiled using a bottom-up approach based on proxies as well as direct emission measurements, for example in stacks. Retrieving data at detailed levels as well as creation of representative emission factors that translate an activity into emissions is, however, a very labour intensive task. ~~Emissions~~ For example, emissions from road transport for example depend on several factors such as: fleet composition, type of fuel, engine maintenance and design, outside temperature, usage profile and road conditions. New technology standards, reported numbers, and real life measures (or lack thereof compared to ~~projections~~ emission estimates) are slow to be incorporated in the emission inventories, as they need to fulfill the good practice guidelines of the respective protocol commonly agreed upon by the EU member states.

Therefore, inventories cannot reflect the latest actual emission trends in "near-real-time". This is problematic especially when large deviations from business as usual scenarios occur, which are then only reflected in the inventories with a great time lag. For example air quality applications such as forecasts depend on accurate emission inventories to represent these changes. A recent example is the large changes in emissions following the ~~2020-COVID~~ COVID-19 lock-downs and the post lockdown recovery phase of the emissions, which both are poorly represented in current air quality applications (Goldberg et al., 2020; Griffin et al., 2020; Barré et al., 2021).

A potential solution to speed up the creation of up to date emissions from inventories, in a harmonized way, is the usage of satellite observations of air pollutants (Beirle et al., 2011; Fioletov et al., 2011; Mijling et al., 2009; Miyazaki et al., 2012; McLinden et al., 2016; Goldberg et al., 2019; Griffin et al., 2020; Dammers et al., 2019; Ding et al., 2020), which can be used to verify the reported emissions, constrain emission sources, and analyse trends. Furthermore, methods that allow for independent verification, can potentially be used to trace and reveal significant discrepancies in the current emission inventories and have proven to be accurate. An example would be the Diesel-Gate Scandal (Jonson et al., 2017), which revealed that diesel cars had been emitting at least four times more NO_x in on-road driving than in type approval tests. Timely verification of the inventories could potentially more rapidly identify such discrepancies.

Over the past decade the data availability of satellite-based atmospheric composition measurements has increased tremendously. Furthermore, due to an increase in instrument sensitivity, spatial and temporal resolution these satellite-based measurements are becoming more and more attractive for air quality monitoring and emission studies. Recent scientific developments have shown the viability of various methods in estimating emissions based on satellite observations. In the case of NO_x, the earliest methods were mostly ~~aimed to~~ developed to estimate the emissions of individual point sources (Beirle et al., 2011) followed by regional estimates at lower spatial resolutions (Mijling et al., 2009; Miyazaki et al., 2012). The more recent Tropospheric Monitoring Instrument (TROPOMI) with its unprecedented ~~horizontal-resolution~~ spatial resolution of 3.5 × 5.5 km² improved the resolvability of individual and clusters of emission sources (Ding et al., 2020; McLinden et al., 2022).

~~Here,~~ The TROPOMI-NO₂ product offers an inventory independent source to verify NO_x emissions. These observations of spatio-temporal trends offer the possibility for inventory agencies to independently check their findings on, for example, emission reduction of NO_x throughout the country without having to rely on bottom-up inventory data products, such as the Emission Database for Global Atmospheric Research (EDGAR) (Crippa et al., 2019), the Copernicus Atmospheric Monitoring Service (CAMS) database (Granier et al., 2019; Kuenen et al., 2022), or other country specific gridded data products like from the Gridding Emission Tool for ArcGis (GRETA) (Schneider et al., 2016). Fast changing spatio-temporal patterns may only be captured by space-borne data in a timely manner in comparison to the above mentioned gridded data products.

A major aim behind the research work presented here is the provision of a tool, developed for the Umweltbundesamt (UBA, German Environment Agency), to compare satellite-derived emissions to inventory emissions for air pollutants, in order to verify the bottom-up computed emissions with independent data from space-borne measurements. This should help inventory compilers to build trust in their work and identify potentially problematic issues, in case large deviations between inventory data and space-borne data trends are present. Furthermore, the tool should allow for fast checks if a country is compliant with

90 its national air pollutant reduction targets, which have been initiated by the EU (EU, 2022) or if adjustments need to be made (Dore, 2022).

In this study, we apply one of the more recently developed methods (Fioletov et al., 2017) to TROPOMI-NO₂ observations to derive the NO_x emissions for Germany for the period of 2019-2021. The plume based fitting method relies on wind data and a parameterization of multiple Gaussian plumes originating from corresponding point sources to estimate the strength of
95 the emissions at these point source location. These estimates are then compared to the emissions in the current inventories for 2019 and 2020, as well as the projected emissions (~~latest draft~~) of 2021 to assess their validity and analyse the expected variations due to the ~~COVID-COVID-19~~ lock-downs. The plume based fitting routine is part of an open-access standalone ~~offline tool~~¹. Besides the plume based fitting routine two further methods were implemented during the development phase, a simple mass-balance approach ~~for~~ for which we coined the term "naive method", and the divergence approach as described by
100 (~~Beirle et al., 2019~~)Beirle et al. (2019). Furthermore, the simple mass-balanced method was employed into an online web-tool (~~space-emissions, described further in the manuscript~~),² geared towards emission inventory agencies that are interested to compare their national total emissions to an independent, yet easily comprehensible space-borne measurement. More details on the implementation and comparison of these (~~other~~) methods can be found in (~~Dammers et al., 2023~~)Dammers et al. (2023).
In this study we focus on the plume based fitting method.

105 2 Methodology and Datasets

~~A major aim behind the research work presented here was the provision of a tool, developed for the Umweltbundesamt (UBA), to compare satellite-derived emissions to inventory emissions for air pollutants to verify the bottom-up computed emissions with independent data from space-borne measurements. This should help inventory compilers to build trust in their work and identify potentially problematic issues, should there be large deviations between inventory data and space-borne data trends. Furthermore, the tool should allow for fast checks if a country is compliant with its national air pollutant reduction targets, which have been initiated by EU decisions e.g.: (EU, 2022) or if adjustments need to be made as exemplified and discussed in (Dore, 2022). The national inventory data is reported through the informative inventory report (IR): For the case of Germany it is publicly available from its original source: . The TROPOMI-NO₂ product offers an inventory independent source to verify NO_x emissions. These space-borne observations of spatio-temporal trends offer the possibility for inventory agencies to independently check their findings on e.g. emission reduction of NO_x throughout the country without having to rely on bottom-up inventory data products, such as the Emission Database for Global Atmospheric Research (EDGAR) (Crippa et al., 2019), the Copernicus Atmospheric Monitoring Service (CAMS) database (Granier et al., 2019; Kuenen et al., 2022) , or other country specific gridded data products e.g. from the Gridding Emission Tool for ArcGis (GRETA) (Schneider et al., 2016) . Fast changing spatio-temporal patterns may only be captured by space-borne data in a timely manner in comparison to the
115 above named gridded data products.~~

¹<https://github.com/UBA-DE-Emissionsituation/space-emissions>

²<https://space-emissions.net/>

2.1 Data sets

2.1.1 Emission inventory

The reporting of the national air pollutants follows international guidelines that are available via the EMEP centre on Emission Inventories and Projections³. The reported inventory data for Germany, in form of the detailed informative inventory report (IIR), ~~detailing describing~~ the technical methodology, may be found here:
125 <https://iir.umweltbundesamt.de/2022/>.

The data is arranged in time series per gas species considering the different emission sources of NO_x in sectors such as; e.g. public power, industry and traffic in very detailed, ~~dis-aggregated~~ ~~disaggregated~~ form, at the national level. The bottom up creation of these ~~data inventories~~ is driven by statistical data ~~, provided by e.g. provided by~~ the German statistical authority (DESTASIS), and complex models that use this data to compile the emissions for a specific gas (or aerosol) for a specific emission source in a specific sector and year. The uncertainties for each reported emission source depend on the availability of the data used for the emission calculation, and may vary considerably. As an example, uncertainties in emissions from sectors, which are quite accurately described by statistical data and models ~~, such as emissions from large power plants, show much lower uncertainties than sectors that are governed by a great complexity such as the natural variability of NO_x from agricultural emissions in soils (e.g., uncertainties can be more than 300 percent % for Agricultural Soils).~~ ~~Though TROPOMI emission estimates provided by this project may not be suitable to resolve these issues of specific sectors they may, however, contribute to identify problems with reported emission totals e.g. should the temporal trend between the inventory data and TROPOMI data diverge significantly. Otherwise the tool is able to build confidence and trust into the emission inventory, should both trends (the TROPOMI-based emission estimates and the national total) match.~~ ~~UBA (2023)).~~
135 ~~UBA (2023)).~~

In this study both the gridded (CLRTAP, 2021) and non-gridded inventory (CLRTAP, 2022) reported emission data sets are ~~used~~, retrieved directly from the respective Convention on Long-range Transboundary Air Pollution (CLRTAP) inventories, which follow the nomenclature for reporting (NFR) standard. The gridded data set is only available for 2019 ~~while~~, ~~whilst~~ the non-gridded data is available for both 2019 and 2020. The 2021 data is a prognosis based on the trends observed between 2012-2019 for all emission classes, under the assumption that the patterns in most emission sectors rebound after the 2020 ~~COVID-COVID-19~~ lock-downs. An overview of relative contributions of individual sectors to the total gridded emissions is shown by sector in Fig. A1.
145

All emissions except the MEMO items (~~MEMO items are~~ additional reported emissions on non-standard emission such as volcanoes, forest fires etc.) are selected from the CLRTAP inventories. ~~To this set two~~ ~~Two~~ natural sources are added ~~to this set~~, namely non-agricultural soils and lightning. Globally, the lightning NO constitutes about 3% of the total NO_x emission budget. ~~Please note that according~~ ~~According~~ to the guidebook (EEA, 2019) only 20% of the lightning NO is formed in the lowest 1000 meter of the atmosphere and the remaining 80% at higher altitude (all inter cloud lightning above 5 km height) ~~(EEA, 2019), (EEA, 2019))~~. A rough estimate for the lightning emissions can be made on the basis of the number of flashes per km² and the expected ~~mol-NO_x-NO_x emissions~~ released per flash. A study by Anderson and Klugmann (2014) gives an
150

³<https://www.ceip.at/reporting-instructions>

average of about 2 flashes per km² throughout Germany, with fewer flashes in the central and northern parts. Assuming that
155 on average the number of lightning flashes did not increase significantly in combination with a production of about 180 mol
NO_x/flash (Bucsela et al., 2019), and German surface area of about ~~357.000~~ 357,000 km², gives us a German lightning NO_x
emission total of about 5 kt ~~NO(NO₂)~~ NO_x per year. This emission total is very minor and spread out over a large domain,
and not expected to be a significant source of error when comparing the satellite derived emission estimates with the emission
inventory. From this point forward in the manuscript kt (NO₂) NO_x is written as kt (NO_x).

160 Due to widespread nitrogen pollution and deposition in Germany it is complicated to make an estimate of purely non-
anthropogenic and non-agricultural soil emissions. There are several studies that looked at soil NO_x emissions for the European
domain, mostly based on the anthropogenic emissions Yienger and Levy II (1995), but fewer that focus on purely natural
emissions. Simpson et al. (1999) gave an estimate of 3-90 kt NO_x of forest emissions and 20 kt NO_x for grassland soils.
This estimate was more recently updated by Simpson and Darras (2021) and is available as the CAMS-GLOB-SOIL inventory
165 (Simpson, 2022), with a reported 2018 German emission total of about 160 kt NO_x. Within the inventory the emissions are split
between fertilizer induced, biome, deposition related, and pulsed soil emissions. There is always a danger of double counting
such emissions but the fertilizer induced emissions of 100 kt NO_x match closely to those included within the 2019 GNFR
data of ~~about~~ approximately 110 kt NO_x (classed under L. AgriOther sources). The remaining ~~60kt~~ 60 kt of NO_x per year is
a combination of biome, deposition related, and pulsed soil emissions. The distribution of non-agricultural source emissions
170 are quite uniformly distributed throughout Germany, peaking somewhat towards the northeastern part of the country. Note
that Simpson and Darras (2021) stress that the derived soil emissions still have a large uncertainty range, mostly related to a
lack of observations, missing data for some biomes, and the uncertainty of input parameters such as soil temperatures. Annual
variations are expected to be large depending on variations in soil temperatures. Simpson and Darras (2021) do not provide an
upper and lower range of the emissions.

175 Additionally, we use the European Pollutant Release and Transfer Register (E-PRTR) for the ~~locations and emission strength~~
emission locations and strengths of the largest industrial emission sources within Germany. The latest data set (v18) can be
accessed via: <https://www.eea.europa.eu/data-and-maps/data/industrial-reporting-under-the-industrial-6>. Only sources with an
emission strength above 0.25 kt NO_x per year are selected for later comparison to the satellite derived emissions. Note, that the
most recent data available is based on reported emissions of the year 2017, and thus we only use the data as a rough indication
180 of source strength.

2.1.2 TROPOMI-NO₂

The TROPOMI instrument, on ~~board of~~ the S5P satellite platform, was launched on the 13th of October 2017. The satellite
instrument achieves almost full daily coverage of the globe through a sun-synchronous orbit with a local overpass at around
13:30 (Veefkind et al., 2012). TROPOMI has an unprecedented horizontal resolution of 3.5 × 5.5 km² for the NO₂ product.
185 Details on the retrieval are described in the Copernicus user manuals (Algorithm Theoretical Basis Document (ATBD)) as well
as in earlier publications such as van Geffen et al. (2022). The TROPOMI-NO₂ standard product has several data streams, with
the near-real-time product available within 3 hours (NRTI), the offline(OFFL) (2019-2021) reprocessed version following one




Figure 1. TROPOMI-NO₂ (PAL [product](#), v2.3.1) ~~yearly-averages-year averaged vertical column density concentrations over Germany~~ for ~~the years~~ 2019-2021.

day later. A complete reprocessed version is provided at more irregular intervals (RPRO, April 2018-November 2018). Finally, a reanalysis of the full data set became available at the end of 2021, named the PAL product, which is currently available until
 190 the end of November 2021, connecting seamlessly to OFFL v2.3.1 from November 2021 to July 2022. The TROPOMI-NO₂ product went through several upgrades concerning its product versions over the years, with the most recent three upgrades from version v1.3.2 to version v1.4.0, then v2.2.0, then v2.3.1, and v2.4.0 taking place in [respectively](#) November 2020, July 2021, November 2021 and July 2022. The most recent upgrade to version 2 involved a more major overhaul greatly improving the overall quality of the retrieval (van Geffen et al., 2022; Zhao et al., 2022).

195 The TROPOMI-NO₂-PAL product includes a reanalysis of the earlier data and provides a consistent version throughout (v2.3.1). This product is recommended to be used for any longer time series analysis and has been used in this study. We combine this product with 2 months of the newest OFFL data (v2.3.1) to complete the data series for 2021. The PAL product is available through the PAL data portal ⁴.

The quality of the TROPOMI-NO₂ PAL and OFFL products based on the v2.3.1 processor version is discussed by (~~van Geffen et al., 2022~~
 200 [van Geffen et al. \(2022\)](#)). Furthermore, the previous data set versions 1.2.x and 1.3.x were relatively well validated (Verhoelst et al., 2021). The TROPOMI NO₂ data ~~are typically well correlated but show an underestimation of the tropospheric column correlate well~~ when compared to ground-based MAX-DOAS and PANDORA instruments (Verhoelst et al., 2021), ~~but tend to show an underestimation of the tropospheric column~~. The median negative bias ranges from -15% to -35% in most clean to slightly polluted regions ~~and~~ up to -50% over highly polluted regions for versions 1.2.x and 1.3.x. This bias is reduced
 205 in the PAL data set (van Geffen et al., 2022) with reported improvements for the tropospheric columns from an average low bias of -32% to -23%. The range of the differences for individual sites are ~~however-, however,~~ quite wide with for

⁴<https://data-portal.s5p-pal.com/products/no2.html>

example the MAX-DOAS in de Bilt, the Netherlands, showing a range of around -75% up to around +50% (25 and 75 % percentiles) with a median of around -20%. The negative bias can be explained by the low spatial resolution of the a-priori profiles as well as the treatment of clouds and aerosols in the retrieval. As for the TROPOMI data quality criteria, the recommended set were used, which means only observations with a cloud fraction below 0.3 were used, based on the cloud_fraction_crb_nitrogen_dioxide_window variable in the data files. Furthermore, observations with a quality value (qa_value) below 0.75 were filtered from the data set.

Figure 1 shows yearly averages of the ~~TROPOMI-NO2~~ TROPOMI-NO₂ (PAL, v2.3.1) data. Here the impact of the COVID-19 lockdown measures in 2020 is clearly visible: The industrialized Ruhr valley at the western border of Germany shows far reduced levels of NO₂, if compared to 2019. The same is also observed in the industrial centers further to the South-South-West, which almost vanishes in 2020 and show only a very slow recovery of emissions in 2021.

2.1.3 Meteorology

The methodology in this study makes use of the wind rotation approach as explained in detail in Pommier et al. (2013), Fioletov et al. (2015), and Dammers et al. (2019). The required meteorology is taken from ECMWF's ERA-5 dataset (~~(Hersbach et al., 2020, 2018)~~) (Hersbach et al., 2020, 2018), which was downloaded at a ~~resolution of~~ 0.25°x0.25° resolution and 1-hour temporal resolution. To match each of the satellite footprints the meteorological fields are interpolated (spatially and temporally) to each of the observations. We assume that the majority of the NO_x mass from local emissions is located in the lower boundary layer (Beirle et al., 2019; McLinden et al., 2022; Griffin et al., 2020) and for the transport of NO_x an average is taken of the wind fields of the first 100 hPa (around the first kilometre) above the surface. These are approximately the levels between ~1000-900 hPa for a typical sea level location, and for a location with a surface pressure of 800 hPa, winds between 800 and 700 hPa are averaged. The surface pressure at the location of the satellite observations are used to determine the 100hPa layer.

2.2 Emission estimates tooling

The plume based fitting routine presented here was developed together with two other methods to form the core of the space-emissions tool as developed for the ~~Umweltbundeamt (UBA)~~ UBA. The other two methods are a simple mass-balance approach, which was coined the "naive method", and a third method, the divergence approach as described by (Beirle et al., 2019) Beirle et al. (2019). The tooling is available in two forms, an open-access standalone offline tool, which contains all three methods and of which the source-code can be found at <https://github.com/UBA-DE-Emissionsituation/space-emissions>, and an online web-tool, which currently only allows for the use of the naïve method. The focus in this study is on the plume based fitting method. More details on the other methodologies and inter-comparison of these (~~other~~) other methods can be found in (Dammers et al., 2023) Dammers et al. (2023). The tool is offered as a web-based application available at: <https://space-emissions.net/>. The data processing is hosted at the German national Copernicus data service initiative⁵, which offers a direct

⁵<https://code-de.org/de/>

link to the required Sentinel-5P data products especially tailored for governmental agencies of Germany. This web-based application tool is directly targeted at users from the emission inventory community and, therefore, uses the TEMIS monthly L3 data product available ~~from in the naive method processing at~~ <https://www.temis.nl/>. The design of the tool is based on a modular structure that encourages later additions of other compatible air pollutants to the tool chain such as SO₂ or to add more technical and computational more demanding methods, other than the here employed mass-balance technique in later development steps. This was necessary as it offers a concise development framework to which more advanced techniques may be added later on, driven by the emission inventory community.


The ~~workflow of both the offline and online tools are~~ [online tool works](#) as follows, (also shown in Fig. 2: ~~A~~): a user is required to select the country of the world they want to target with their analysis, as well as the desired air pollutant (in this case NO₂), from the respective pull down menus of the processing options (online) or by providing a shapefile of the region of choice (offline). After that, the time span for the observations needs to be selected, ~~where data is available from the onset of the S5P NO₂ data product (February 2018). At last~~ [covering the period for which the data is available](#). Finally, the user initiates the computation, which returns the analysis results to the graphical user interface. The user may then download the graphical results as well as the analysis results as a comma separated value (.csv) file and/or other ancillary data using the post-processing options (netCDF4 files). Advanced users and software developers are also encouraged to visit: <https://github.com/UBA-DE-Emissionsituation/space-emissions> for the source code of the project.

2.2.1 Multi-source plume method (MSPM)


Emissions were derived using the Multi-source plume method (MSPM) (Fioletov et al., 2017; McLinden et al., 2022; Dammers et al., 2022), which was originally developed by Fioletov et al. (2017) and can be used for an assessment of emissions from both area and point sources. For a more detailed explanation we refer readers to those publications. In short, the method relates observations and emission sources by creating a linear system of plume functions, which effectively establish a system of source and receptor relations, ~~Eq. 1~~, in which the total column density of each observation is described as a combination of ~~the~~ total column densities of all sources plume functions. ~~This is expressed as:~~

$$\mathbf{Ax} = \mathbf{B} \tag{1}$$

~~In Eq. 1 where~~ ~~A~~ ~~represent a is the~~ linear system of source-receptor relations, ~~x~~ the emission sources, and ~~B~~ the satellite observed vertical column density ~~;~~ (in our case the TROPOMI-NO₂ tropospheric columns). Several additional terms can be incorporated in ~~x~~ to account for regional product biases and for background concentrations. While the TROPOMI-NO₂ product does have local biases, the small number of validation stations hampers an accurate determination and correction for the product bias. To account for the bias we apply a correction on an overpass to overpass basis, following Beirle et al. (2019), removing the lowest 5% of the observed total column density within the larger domain. The short lifetime of NO₂ ensures that further corrections for background concentrations ~~is are~~ not needed.



A



B

Figure 2. Screenshots from the space-emissions tool <https://space-emissions.net/> for Germany. A illustrates the interface and the visualisation of the result. Whist B illustrates the result in context to the reported NO_x values.

270 Any (~~Gaussian~~) plume function can be used to represent the relations in matrix A, here we use the ~~EMG~~ exponentially modified Gaussian (EMG) plume function, which has been successfully applied in previous studies (de Foy et al., 2014; Fioletov et al., 2017; McLinden et al., 2020; Dammers et al., 2019). Using this method, observations are rotated around a single point, the emission source, so that each is positioned in a similar upwind-downwind frame (Pommier et al., 2013) with respect to the wind direction. This enables us to describe the position of each observation as a point within a downwind plume.

275 For more details on the plume rotation method see, Fig. S4 in Pommier et al. (2013). The EMG plume function describes

the vertical column density (VCD) concentrations downwind of a source. The VCD at each position x, y near a source can be described by Eq. 2, where ~~a represents the peak~~ a represents the emission enhancement, $f(x, y)$ the crosswind diffusion (Eq. 4), and $g(y, s)$ (Eq. 6) a convolution of the downwind advection and diffusion, ~~within all functions~~, Within all functions x represents the crosswind position, y the downwind position, and s the wind-speed.

$$280 \quad \underline{\text{Column}_{NO_2, j}} V(x, y, s) = a \cdot f(x, y) \cdot g(y, s) + B \quad (2)$$

$$\underline{f(x, y) = \frac{1}{\sigma_1 \sqrt{2\pi}} \exp\left(-\frac{x^2}{2\sigma_1^2}\right)}$$

$$\underline{g(y, s) = \frac{\lambda_1}{2} \exp\left(\frac{\lambda_1(\lambda_1 \sigma^2 + 2y)}{2}\right) \operatorname{erfc}\left(\frac{\lambda_1 \sigma^2 + y}{\sqrt{2}\sigma}\right)}$$

$$285 \quad \sigma_1 = \begin{cases} \sqrt{\sigma^2 - 1.5y} & , y < 0 \\ \sigma & , y \geq 0 \end{cases} \quad (3)$$

$$\underline{f(x, y) = \frac{1}{\sigma_1 \sqrt{2\pi}} \exp\left(-\frac{x^2}{2\sigma_1^2}\right)} \quad (4)$$

$$\lambda_1 = \frac{\lambda}{s} \quad (5)$$

$$\underline{g(y, s) = \frac{\lambda_1}{2} \exp\left(\frac{\lambda_1(\lambda_1 \sigma^2 + 2y)}{2}\right) \operatorname{erfc}\left(\frac{\lambda_1 \sigma^2 + y}{\sqrt{2}\sigma}\right)} \quad (6)$$

Parameters, σ and λ represent the plume-spread and decay-rate of NO_2 , with $\tau = 1/\lambda$ being the decay time or life time. The parameters σ_1 and λ_1 shown in Eqs. 3 and 5 represent the adjusted form of a plume upwind of the source (σ_1), and the ~~the~~ decay-rate divided by the wind-speed (λ_1). Each observation j can then be described by the sum of the enhancements of all sources i forming, Eq. 7.

$$\text{Column}_{NO_2, j}(\psi_j, \theta_j, \mathbf{s}_j) = \sum_i a_i f(x_{i,j}, y_{i,j}) g(y_{i,j}, s_j) + B_{i,j} \quad (7)$$

The emission rate of each source i can then be calculated by dividing the emission enhancement a_i by the decay-rate λ ;
 295 $E = a/\tau = a\lambda$.

In this work, a grid with a resolution of 0.1 x 0.1 degrees is used to describe the emission, covering the full domain of Germany with ~~two~~ 2 degrees of padding added to the ~~domains~~ edges to reduce any edge effects (Dammers et al., 2022). ~~Following Beirle et al. (2019) The effective lifetime of NO_x depends on both the chemical decay rate and loss to surfaces (dry deposition). Within our domain of interest, the chemical decay will be the dominant factor. Earlier studies using the EMG plume functions derived lifetimes between 2-5 hours based on the decay downwind of major sources worldwide (Beirle et al., 2011; de Foy et al., 2015; Gol~~
 300 ~~. Following those results~~ we assume a ~~lifetime of about mean lifetime of~~ 4 hours (~~±25%~~) ~~± 1 hour to account for local and seasonal variations~~. The plume-spread can be seen as a combination of the diffusion, satellite footprint size, and the spatial size of the sources (McLinden et al. (2022)). Taking into account the effective TROPOMI foot-print as well as the added diffusion we use a value of 7 km for the plume-spread. A dampening factor is added to the linear system in equation 1, forming Eq.
 305 8 to reduce oscillation effects within the solution. The resulting sparse linear system can be solved efficiently with the ~~Scipy sparse.linalg.lsqr~~ ~~Scipy sparse.linalg.lsqr~~ package (Paige and Saunders, 1982; Virtanen et al., 2020).

$$\begin{bmatrix} \mathbf{A} \\ \gamma\mathbf{C} \end{bmatrix} \mathbf{x} = \begin{bmatrix} \mathbf{B} \\ \mathbf{0} \end{bmatrix} \quad (8)$$

Satellite observations of short lived species are only representative of emissions near the overpass time. A correction factor should be applied to the satellite-based estimated emissions to account for the diurnal and seasonal variability. To account for
 310 this we can use a basic box-model to approximate the mass over time and apply a posterior correction. Assuming a mass $m(t)$, and a constant lifetime ($\tau = 1/lifetime$) and E the emission at time t , the mass can be calculated with,

$$m(t) = m(t-1)e^{-\tau} + E(t) \quad (9)$$

This equation is applied to the domain-wide emissions that are injected into the domain for a whole year, including a few days of spin-up, and averaged and normalised for a selection of expected lifetimes. For the temporal distribution we use the
 315 average NO_x emission profile for all NO_x sources within the German domain as used in the LOTOS-EUROS model (Manders et al., 2017). A lifetime of about 4 hours and an overpass time of around 13:00 LST results in a correction factor of 1.24, meaning that the estimated emissions can be expected to ~~bias-high-be overestimated~~ by around 24%.

Depending on the source location and time of year this value is expected to vary due to variations in the temporal emission profile. However, as actual measurements of diurnal cycles of nitrogen-dioxide emissions are rare and only exist for larger
 320 power-plants, only the variability in the model emissions can be used to create a regional adjustment parameter. Surface concentration observations should in turn be used to analyse and optimize the modelled diurnal emission profiles for individual sectors. To calculate the viability of such a regional factor the adjustment parameter was calculated for each cell. The standard deviation of the regional parameters is around ~0.05. Therefore, to reduce complexity, the value of 1.24 is assumed for the entire domain. A similar parameter is derived to account for the seasonal variability of the emissions in combination with the
 325 variable availability of ~~TROPOMI-NO₂~~ ~~TROPOMI-NO₂~~ observations passing the data quality filters. The correction parameter

is calculated as the weighted mean of the number of observations per month and the mean correction factor for each month. Using this approach, a value of 1.05 is found. Combined with the diurnal parameter this gives a factor of approximately 1.30.

TROPOMI is only capable ~~in of~~ observing NO_2 , ~~therefore~~. ~~Therefore~~, an additional correction is needed to account for the NO mass. ~~In previous studies this is commonly done by using a NO_x/NO_2 concentration ratio~~. ~~Values commonly~~
330 ~~lie within the 1.3-1.4 range (Beirle et al., 2011, 2019) depending~~ depends on the local chemistry ~~with values commonly falling~~
~~within the 1.2-1.5 range for polluted regions (Beirle et al., 2011, 2019, 2021; Lange et al., 2022)~~. In this study we apply the
1.32 \pm 0.26 factor as used by Beirle et al. (2019) ~~and include the standard deviation of 0.26 further into the uncertainty budget~~
~~account for the variations~~.

2.2.2 Method uncertainties

335 ~~Based on the methods and choice of parameters described in the previous sections a summary can be made of the total expected~~
~~uncertainty of our method. An overview of the uncertainty parameters with a short summary of the chosen parameter values~~
~~and impact on the emissions is given in table 1.~~

~~One of the major parameters of uncertainty is the TROPOMI- NO_2 data product. As stated earlier, the current TROPOMI- NO_2~~
~~product overestimates concentrations in background/low emission regions (+a few %) while having a negative bias in source~~
~~regions: -35%, up to -50% in extreme cases (Verhoelst et al., 2021), which according to van Geffen et al. (2022) adds up to a~~
340 ~~potential mean bias of around -32% to -23%. Assuming -23% for a larger industrialized region such as Germany, we end up~~
~~with an underestimation of the emissions by a factor of -30%. Locally these values can decrease further (high-emission zones)~~
~~or increase (low-emission zones, up to a positive percentage). The main cause for the bias can be found in the inaccuracies~~
~~of the air mass factor (AMF) which come from uncertainties in the underlying modelled concentration fields and missing~~
345 ~~variations in the stratospheric NO_2 concentrations. Local variations due to errors in the AMF cannot be corrected for without~~
~~the use of a chemistry transport model (CTM) and lead to under or overestimation of emissions in high source and background~~
~~regions. A recent approach using the modelled CAMS-Europe profiles (Douros et al., 2023) shows that the large negative bias~~
~~can be resolved with the help of higher resolution a-priori profiles. Beside this systematic uncertainty the VCDs will also~~
~~have a random uncertainty (of up to 30-50% for individual observations). Due to the large number of observations used to~~
350 ~~constrain each source, the impact of those uncertainties are expected to be minor. Furthermore, there is the detection limit of~~
~~the TROPOMI instrument, which limits the ability to detect smaller sources. The study by Beirle et al. (2019) gives a limit~~
~~of about 0.11kg/s, based on the Divergence method. An emission source of 0.11kg/s equals about 3.5 kt NO_x per year. This~~
~~is based on a peak fit which typically has a radius of 25km, which roughly gives us a 2500km² area, which when divided~~
~~the detection limit over the area, results in an detection limit of around 1.4 tonnes km⁻². To summarize the total expected~~
355 ~~uncertainty in the emissions due to the TROPOMI product will add up to around -30%.~~

~~The second major parameter with a large uncertainty is the choice of lifetime. An underestimation of the chemical losses~~
~~could lead to an overall overestimation of the emissions, and vice versa an overestimation of the lifetime can lead to an~~
~~underestimation of the emissions. A doubling of the lifetime roughly halves the emissions, which shows the importance of the~~
~~parameter. Lifetimes as stated are location-dependent and to more accurately estimate will require further detailed plume and~~

360 chemistry (model) studies. Examples of recent studies (Beirle et al., 2011, 2019, 2021; Lange et al., 2022) give an indication
of the typical ranges for the NO_x (chemical) lifetime and give a range of 2-5 hours, with the study by Lange et al. (2022) giving
a value of 3-5 hours representative of the Germany domain. The 4±1 hour results result in an -33 to +20% under/overestimation
of the emissions.

365 The NO_x to NO₂ ratio can also have local variations, which affect the total emissions. At source level the majority NO_x
seems to be emitted as NO, which can rapidly turn into NO₂ after which an equilibrium is reached, the speed of which depends
on the availability of O₃. Beirle et al. (2021) recently gave a modelled estimate of the ratio, which was very close to the factor
1.32(±20%) given in his original study, with values moving towards 1.0 for industrial areas just north of the equator while
values tended towards higher ranges (1.6) for less industrialized and high-latitude regions.

370 Next up, there is the influence of the wind-speed and direction, for which we assume an uncertainty of up to 1ms⁻¹
(McLinden et al., 2022). The effect can move in both directions and translates into an uncertainty of around 15-20% for average
conditions over Germany (based on the matched wind-fields) which matches earlier uncertainty estimates by Griffin et al. (2021); McLinden

375 Finally, the diurnal and seasonal variations show some variations in the order of a few percent (<5%). Note that a fixed
parameter was determined for the whole German domain, but locally the diurnal correction factor can be lower/larger for the
more continuous/strongly varying emissions. For example, in the case of power-plants, which run more continuously than road
transport, this can result into a negative bias for the emissions.

380 Taken together, these error terms result into a Germany averaged error range between -50 to +35% for the Gaussian plume
method. The low error estimate corresponds to source regions where the low bias of the TROPOMI VCDs, effectively low
biasing the emissions, are counteracted by the potentially high bias in the emissions of the NO_x:NO₂ ratio and effective
lifetimes. Both values should be seen as conservative estimates, which would occur in the unlikely case that the inaccuracy in
NO_x:NO₂ ratio, lifetime, AMF and wind-fields all nudge the estimate in the same direction for all locations in the domain of
interest. In reality not all errors point in a similar direction, like the product bias pointing in opposite directions for background
and source regions.

2.2.3 Sector specific emissions

385 A direct sector based attribution of emissions is not feasible using the satellite data only. Therefore, additional data needs to be
taken into account to attempt to estimate a potential sectoral attribution of the emission. We used the GNFR/CLRTAP sector
outputs to create a spatial index filter for the emission data. The gridded-NFR-GNFR data is used as a basis and summed to
and regridded for all the NFR classes, to match the 0.1°x0.1° grid for each of the NFR classes used in this study. A Gaussian
filter (*scipy.ndimage*, Virtanen et al. (2020)) is applied to the data with a sigma of 1 grid cell, applying a light smoothing to
390 account for the maximum resolvability of each source and emission methodology. The resulting masks are divided by the total
emissions of all sectors to derive each sector's fraction of all emissions (emission fraction), and shown in the supplement Fig.
A1 for the non smoothed version and Fig. A2 for the Gaussian smoothed version. For further sectoral emissions analysis only
locations with an emission fraction above 50 % are selected, the resulting mask is shown in Fig. A3.

<u>Parameter</u>	<u>Summary</u>	<u>Impact on final emissions(%)</u>
<u>TROPOMI: AMF/other bias</u>	<u>-23% mean bias</u>	<u>-30%</u>
<u>TROPOMI: Noise</u>	<u>30-50% for individual observations depending on VCD range</u>	<u>Minor, large number of observations reduces uncertainty.</u>
<u>TROPOMI: Detection limit</u>	<u>3.5 kt NO_x for isolated individual sources</u>	<u>±1.4 tonnes km⁻²</u>
<u>Total: TROPOMI</u>		<u>-30%</u>
<u>Lifetime</u>	<u>4 hours±25%</u>	<u>-33% to +20%</u>
<u>NO_x:NO₂ ratio</u>	<u>1.32±0.26</u>	<u>±20% (a factor of 1.41 gives an increase of +7%)</u>
<u>Wind Fields</u>	<u>±1m s⁻¹</u>	<u>±15-20%</u>
<u>Diurnal & Seasonal emission cycles</u>	<u>1.3±0.05</u>	<u>±5%</u>
<u>Total uncertainty</u>		<u>-50% to +35%</u>

Table 1. Summary of uncertainty parameters for emission estimates

3 Results

395 3.1 Inter-comparison with the emission inventory

For a comparison with the gridded inventory data, we used the 2019 data from the satellite derived emissions and the respective NO_x data from the GNFR inventory (CLRTAP, 2021). Figure 3 gives a visual comparison of the 2019 data sets. Both sets were compared using the Structural Similarity Index Measure (SSIM) (Wang et al., 2004) for a quantitative comparison of the images. The SSIM operator is a metric, which was developed to evaluate the image quality of video frames . It uses a window based comparison analysis to track subtle differences between two images so that the spatial structure of both images is also taken into account when calculating the SSIM score. This way similarity and dissimilarity between two 2D data sets may be quantified with the SSIM score in a way – which assesses image similarity in a more human-vision-based mode. Since its introduction SSIM has become a standard comparison operator for computing the similarity between 2d data sets and is now also available in standard open-source data analysis packages such as scikit-image (van der Walt et al., 2014).

405 The resulting SSIM analysis for the 2019 GNFR and TROPOMI derived emission data shows a SSIM score value of 0.6 between both data sets. However, to consider the different approaches of both data sets and to harmonize effects of different baseline resolution, a Gaussian filter is applied to both sets of data that compensates the effect of the larger point spread function




Figure 3. The Structural Similarity Index Measure SSIM of 0.6 was calculated between the gridded inventory data (top-left panel, A) and the emissions derived with the TROPOMI data for 2019 (top-right panel, B). Please note that the details in the (image-) data structure (major road networks and urban areas) are very similar between both sets of data. This is highlighted by a SSIM score of 0.6, which quantifies as the similarity between the data as highly significant. If the data is Gaussian filtered effects of spatially sharper GNFR data (panel C), compared to TROPOMI data (panel D) are compensated and yield a SSIM score of 0.79

(PSF) of the sensor. If both images are Gaussian filtered and compared, the resultant SSIM score is 0.79 and now closer to a score of 1.0, which would depict spatial structure identity between the two sets of data. This illustrates that the spatial structure of both data sets show a high similarity and space-borne derived emissions from the method presented here capture similar large emission sources such as major cities, road networks and industrial areas as the GNFR data set.

3.2 Multi-year emissions




Figure 4. From left to right; NO_x satellite derived yearly emissions for 2019-2021, and the GNFR inventory emissions of 2019. Rightmost figure H: Hamburg B: Berlin, C: Cologne, LU: Lusatia, LE: Leipzig, M: Munich, S: Stuttgart, F: Frankfurt.

The satellite derived emissions for the individual years between 2019-2021 are shown in Figure 4. The rightmost plot shows the emissions as part of the GNFR inventory emissions. A comparison between the satellite derived emissions of individual years and the inventory emissions for 2019 are shown in Fig. 5. Figure 6 shows the change in the spatial satellite derived emission distribution of the largest sources in Germany between 2019 and 2020 or 2021. As the gridded inventory emissions are only available for 2019, we can only compare the individual years to that years inventory emissions.

~~Fig4 again illustrates the exceptional qualities of the TROPOMI instrument over previous sensors. Fig. 4~~ shows that the spatial data from the 2019 space-borne emission estimates have elevated NO_x emission values of around 5-7t/km² outlining the major motorway network in Germany. Most notably is the motorway A2 westwards from Berlin, via Magdeburg and Hannover towards the Rhine-Ruhr region. Another interesting motorway is the A1 from the Ruhr area of North-Rhine-Westphalia towards Bremen and Hamburg. The motorway ring and spider-alike road networks and settlements fanning out from around Berlin are also nicely visible in 2019.

While the TROPOMI instrument represents a huge step in the capabilities to spot individual emission sources, there are still limits to the spatial resolvability. The top row of Fig. 5 shows a direct comparison while the bottom row shows the same results but now with the application of the Gaussian filter, as previously used in Fig.3. The main difference between these two rows are the large positive/negative swings around the more localized emissions and/or major point source like emitters such




Figure 5. Difference between the satellite derived and inventory emissions (2019) for years 2019-2021 (A-C). The red values indicate a higher value for the satellite derived emissions compared to the inventory emissions. Top row shows the original difference between both emission sets, the bottom row shows the same sets but now with the Gaussian filter applied to both sets before subtracting the 2019 inventory emissions (D-F). The letters in the figure indicate the following: H: Hamburg B: Berlin, C: Cologne, LU: Lusatia, LE: Leipzig, M: Munich, S: Stuttgart, F: Frankfurt.

as power-plants. Such variations are ~~however~~, however, not observed around emitters with a large spatial footprints such as cities. This is an excellent showcase of the limit of the method and satellite instrument. Through the size of the satellite pixel's footprint and the misrepresentations of the wind fields (i.e. artefacts) there is an actual limit to the overall spatial resolvability of individual sources. This limit was reported by McLinden et al. (2022) to be around 5-10km for TROPOMI, which matches well to the size of the source grid used here. The 0.1x0.1 degree spatial resolution used in this study is thus at the limit of the methods capabilities to constrain individual neighbouring sources and some smearing is thus expected around the strongest point like sources. The Gaussian (smearing) filter can be used as a first order correction, which results in the lower row of plots. Compared to the inventory emissions of 2019, the Figs. 5 D, E and F show similar patterns between the years, with strong negative differences observed around the major sources, while the background regions (i.e., regions with emissions below 2 tonnes km²) show a consistent positive difference of around 0-1 tonnes km². There are several potential causes for these systematic patterns which will be evaluated in the discussion section.




Figure 6. Difference between 2020-2019 and 2021-2019 satellite derived emissions. A Gaussian filter has been applied to both derived emission sets. The red dots indicate the locations of the largest NO_x emitters within Germany, with the size of the dots a reflection of the individual source strength. The red triangles indicate the larger Power-Plants, with the letter combinations indicating the names of the powerplants: NEU (Neurath), NIE (Niederaussem), WW (Weisweiler), LD (Lippendorf), JAN (Janschwalde), SP (Schwarze Pump), and BB (Boxberg).

Outside of the systematic patterns there are several variations visible between the years. The year 2020 shows a noticeable drop in NO_x emissions around ~~the before mentioned~~ industrial, cities, and highways (Fig. 6). The largest reduction of NO_x

emissions between 2019 and 2020 are in the industrial areas in the Rhine Ruhr region and the upper Rhine area. The rise in emissions from 2020 to 2021 in the TROPOMI data in Fig. 4 and Fig. 6 is most noticeable in the larger urban areas. However, the 2021 NO_x emissions are still lower than in 2019, for example in the industrial centers of the Rhine-Ruhr region (note the red dots indicating the major industrial emitters) and further south along the Rhine. Only the A1 motorway ([the line of emissions between the major emissions clusters at C and H](#)) is still clearly visible in the 2020 and 2021 emission estimates (Fig. 4), whilst the data from other settlement and road networks (e.g., around Berlin) is much less obvious than in the 2019 emission estimates. This is also visible in Fig. 6 in the area with roads leading away from Berlin, where the difference between the 2021 and 2019 estimated emissions still show negative difference of the order 0-1 tonnes km².

Two of the most prominently visible changes (2019-2020) shown in Fig. 6 are the industrial "Ruhr" region, the largest and oldest industrial core of Germany in the westernmost part of the country, and the area of Lusatia in the eastern part of the country with ~~its big lignite mining for power generation in a large-scale lignite mining industry to supply~~ coal fired power plants. These two areas are shown as detailed maps in Fig. 7. Compared to 2019 the emissions have dropped substantially in 2020 and 2021 (up to <-5t/km²). The power generation in Germany has seen an upsurge in the usage of coal fired power stations for power generation in 2021 compared to the COVID-19 year 2020, as reported by the DESTASIS in its press briefing ([link](#)) stating that coal had been the most important source of electricity generation in Germany in 2021. This can be seen in Fig. 7, where there is an increase near one of the large emission centers right at the eastern border of Germany. The Schwarze Pump (SP) and Lippendorf (~~LPLD~~) power-plants even show an increase in emissions compared to 2019. Meanwhile the emissions from the Janschwalde (JAN) power-plant show a strong reduction in 2020 that continues into 2021, which was expected as the power-plant reduced its operation capacity as planned (Vattenfall, 2015; EPH, 2022). The three large power-plants in the west show similar patterns with the Neurath (NEU) and Niederaussem (NIE) plants showing a strong decrease in 2020 rebounding upwards moving into 2021. The Weisweiler power-plant reduces into 2019 while reducing further into 2021. This drop can be explained by two potential causes. Firstly, a ~~reduction of operation capacity was planned.~~ [planned reduction in operation capacity](#) and secondly by the ~~floods caused by flooding from~~ exceptional rainfall in mid July 2021, which also affected the nearby lignite mining pits (RWE statement / [Link to news item](#)).

465 3.3 Sector-specific emissions

An aggregated version of the spatially distributed results is shown in the bar plot of Fig.8, in which the country wide fitted emissions are compared to the country-wide sector specific emission totals. Note, that we added the ~~60kt~~ [60 kt NO_x](#) from natural soil emissions and ~~5kt~~ [5 kt NO_x](#) from lightning emissions to the N_Natural class. These emissions were not included in the previous spatial plots. In line with the previously discussed results, both the satellite and inventory emissions show a large drop from 2019-2020, of comparable size. The slight increase in the projected inventory emissions from 2020-2021 is however not matched by a change in the satellite emissions.

Emissions sources that have a strong spatio-temporal imprint on TROPOMI data should show independent patterns for regions where the sources cause the majority of emissions. To find out what type of source is causing this mismatch, we make




Figure 7. Difference between 2020-2019 and 2021-2019 satellite derived emissions. Upper row depicting the industrial "Ruhr" region, whilst the lower 3 panels show Lusatia at the eastern border of Germany. A Gaussian filter has been applied to all data sets prior to subtraction. The red dots indicate the locations of the largest NO_x emitters within Germany, with the size of the dots a reflection of the individual source strength. The black diamonds indicate the larger Power-Plants, with the letter combinations indicating the names of the powerplants: NEU (Neurath), NIE (Niederaussem), WW (Weisweiler), LD (Lippendorf), JAN (Janschwalde), SP (Schwarze Pump), and BB (Boxberg).




Figure 8. Emission changes over the years as reported by the national inventory of Germany and observed by the TROPOMI instrument. Black error bars indicate the uncertainties on the inventory emissions while the red error bars show the uncertainty in the satellite derived emissions. Note the slight rise in the reported emissions of 2021, compared to the year 2020 due to COVID-19 lockdown measures.

use of the sectoral masks (e.g., fig. A3) to derive sector specific patterns from the spatio-temporal data from the satellite derived
475 and inventory emission data.

Only 5 sectors have locations which are dominated (e.g., above 50% of the total emissions) by a single emission sector, of which the PublicPower sector has the largest emissions in a single location while the road transport ~~and agricultural~~ emissions are more spread out over roads and pastures throughout the country. Note, that the PublicPower, Shipping and Industrial emissions cover a very limited area with only PublicPower showing very high emissions. Figure 9 shows the sector specific
480 emissions as indexed by the 2019 emissions for the ~~non-agricultural~~ Public Power, Industry, Road Transport and Shipping sources. Based on earlier projections, and trends over the previous years, 2021 inventory emissions are expected to be at just over 90% of the 2019 emissions. The emissions related to power generation have bounced back almost to the ~~pre-COVID~~ pre-COVID-19 levels even though the Janschwalde power plant in the east reduced its operation capacity as planned (Vattenfall, 2015; EPH, 2022). Further resurgences are to be expected (for 2022) by the plans of the reactivation of old coal fired power
485 plants in the wake of the European energy crisis and the potential fears of a blackout in Germany. While Road transport emissions were expected to show a recovery this is not matched by patterns in the satellite derived emissions. The slow recovery can potentially be explained by the reduced number of kilometers by trucks (>3500kg) which is almost down by 10% in 2021 compared to 2019 (KBA, 2022). Shipping emissions have continued their trend with no ~~show-of-any-sign-of~~ recovery. While this reduction was expected based on past trends, the cause can be found in the global shipping crisis and disrupted
490 supply chains.




Figure 9. Satellite derived and inventory emissions for each source sector as indexed by the 2019 emissions. A clear downward trend is visible for most sectors. Dotted lines indicate the inventory emissions, the other line the satellite derived emissions.

4 Discussion

As the results showed, the captured spatial variability within satellite derived emissions are very similar to those in inventory emissions. The values for the NO_x emissions retrieved from the TROPOMI observations diverge on average ~~100kt from the reported emissions by 75-100 NO_x (<10%) from the emissions reported for Germany~~. There are some variations observed
495 between the years but the difference between both series fall within the uncertainty range of both emission totals (Fig. 8). The uncertainties of both sets are quite large compared to the yearly variations, which hampers stronger conclusions on the quality of the inventory and satellite based estimates. We can however discuss the various causes of uncertainty and how these can be reduced. The uncertainty range of the reported inventory emissions is estimated to fall between -9.2 and +15.8% (see <https://iir.umweltbundesamt.de/2022/>), which translates to about -100 to +180 kt NO_x in 2019 for the inventory. Note that this
500 range does not include potentially missed sources, such as stronger than expected natural emissions (e.g., [Soil emissions](#)) and any of the MEMO items. ~~As discussed in section 2.1.1 especially soil emissions show large range within the literature, and some of the spatial patterns show possible hints a smaller but widespread source throughout Germany (Fig. 5).~~

Besides the above discussed items it should be noted that the emissions from road transport are required to be based on the fuels sold approach. This approach does not account for all the emissions which occur in Germany from vehicles which
505 were fueled abroad and are driving in Germany (this might constitute an underestimation in the inventory). On the other hand the emissions from foreign vehicles (for instance from the Netherlands) which bought their fuel in Germany and were not driving in Germany are in this fuel sold approach allocated to Germany (this might constitute an overestimation of the German emissions). However, it is not known how much emissions are associated with these cross border phenomena's for Germany. Data from The Netherlands show this might be a significant difference; the ~~NO_x~~ NO_x emission based on fuel used
510 is approximately 5.5 % less than the emissions based on fuel sold as reported in the GNFR total. However, as fuel prices in Belgium and Germany are cheaper than in The Netherlands, Dutch drivers fuel frequently in those countries thus the Dutch case represents the higher end of the difference between fuel sold and fuel used approach, only surpassed by Luxembourg with one of the lowest fuel prices in Europe.

Another source of uncertainty are the emissions near the border regions. Emissions within the first 10 to 20km outside of the
515 border can be expected to be smeared out in the inversions due to the limited resolvability of the instrument and methodology. The stronger the source is, the better is the resolvability. So for the larger sources 10 km can be assumed. Making a loop around the German borders there are a few areas of interest. Starting at the border of the Netherlands and moving clockwise on both sides of the border, there are several larger sources, such as the Weisweiler power plant in Eschweiler, the Dolna Odra power station in Poland, and several power plants near the border in the Czech Republic, but also several smaller and larger cities.
520 By taking a polygon that is 10 km wider and narrower in shape than the existing German borders the smeared emissions near the borders can be approximated. Based on the European CAMS-REG v5.1 inventory (emissions 2018, based on the in 2020 reported emissions, (Kuenen et al., 2022)) we find that around 120 kt NO_x of the German emissions take place within Germany and within 10 km of the borders, and around 75 kt NO_x just outside of Germany within 10 km of the border. Assuming that at most half of the full amount of these emissions smear out past the border, the smeared loss in emissions is about 22kt on

525 the total emissions. This should be seen as an upper limit. Furthermore, of these emissions a large majority takes place in the western part of Germany, where the most common wind direction is wind coming from the west. In effect it can be expected that the smearing of those emissions will be reduced further.

A more probable source for the ~~100kt~~ 100 kt NO_x mismatch however can be found in the satellite derived estimates. ~~The As stated in section 2.2.2 the TROPOMI based emission estimates on average can have an uncertainty in the range of 30-40~~ 35-50% translating to about ~~+400kt and based on various uncertainty parameters. Based on the multi-source methodology we can point out several important parameters of uncertainty. Firstly, there is the TROPOMI-NO₂ product. As stated earlier, the current TROPOMI-NO₂ product overestimates concentrations in background/low emission regions (+a few %) while having a negative bias in source regions (-35%, up to -50% in extreme cases (Verhoelst et al., 2021), although improved from a mean bias of around -32% to -23% (van Geffen et al., 2022)). The cause for the bias can be found in the inaccuracies of the air mass factor (AMF) which comes from uncertainties in the underlying modelled concentration fields and missing variations in the stratospheric NO₂ concentrations. Local variations due to errors in the AMF cannot be corrected for without the use of a chemistry transport model (CTM) and can lead to under and overestimation of emissions in high source and background regions. A recent approach using the modelled CAMS-Europe profiles show that the large negative bias can be resolved with the help of higher resolution a-priori profiles. Beside this systematic uncertainty the VCDs will also have a random uncertainty,~~ but due to the large number of observations used to constrain each source, the impact of those uncertainties is expected to be minor. Finally, there is the detection limit of the TROPOMI instrument, which limits the detect ability of sources. The study by Beirle et al. (2019) gives a limit of about 0.1 Hkg/s, although based on the Divergence method. An emission source of 0.1 Hkg/s equals about 3.5kt NO_x per year. This is based on a peak fit which typically has a radius of 25km, which roughly gives us a 2500km² area, which when dividing the detection limit over the area, results in an detection limit of around 1.4 tonnes km⁻². To put that number in perspective, ~~400 kt NO_x. A more complete error analysis based on simulated observations with controlled conditions and a subsequent Monte-Carlo analysis of error propagation could give a more accurate estimate, but falls outside of the scope of this study. A study by Dammers et al. (2022) however did perform such an analysis. While using a very similar set of input parameters, the study derived a mean uncertainty between 15-20% which increases when close to large mountains. Two important differences between this manuscript and that study (Dammers et al., 2022), are the uncertainties and bias in the smaller differences observed in Fig5 are in many cases above that limit except for the background regions in southern Germany. Secondly, while the assumption of a 4 (±25%) hour the effective lifetime seems to be correct for 2019 Germany as a whole. It is important to note that an underestimation of the chemical losses could lead to an overall overestimation of satellite product (which only shows a minor negative bias) and the emissions, and vice-versa an overestimation of the lifetime can lead to lack of a NO_x:NO₂ ratio. Without both parameters the 15-20% uncertainty would translate into an uncertainty of around ±150-200~~ kt NO_x.

555 For the regional emission mismatches, of the uncertainties studied in Dammers et al. (2022) and in this study, only the lifetime, the satellite product bias, and NO_x:NO₂ parameters can have a large enough systematic effect on the estimated emissions to explain the observed differences. Additionally, one could argue that the wind fields around larger hills and mountains can have a systematic effect. However, throughout our region of interest most of the mismatches (Fig. 5) are observed

560 away from the main mountainous regions. The negative bias observed around the major emitters (up to and over 50%) can be explained by both the product bias, an increase of $\text{NO}_x:\text{NO}_2$ ratios near the source (Lange et al., 2022; Griffin et al., 2021), and a mismatch in the NO_x lifetime. The product bias by itself can be expected to cause an underestimation of the emissions. A doubling of the lifetime roughly halves the emissions which shows the importance of the parameter. Lifetimes however are location-dependent and require more detailed plume and chemistry studies. Examples of recent other studies show the large ranges for the NO_x (chemical) lifetime. Examples are the study by Lorente et al. (2019) give a range of 11 hours in winter, to 1-4 hours in spring for the region around Paris. Previous studies specifically using the EMG plume function found effective lifetime in the order of 2-5 hours for very strong emitters. Furthermore, other possible causes for the over or underestimation of derived emissions are the diurnal and seasonal variations (expected to be several % for the seasonal variations). Currently a fixed parameter was determined for the whole German domain, but locally the diurnal correction factor can be lower for the more continuous emissions, for example in the case of power plants, and thus in effect create a negative bias for emissions. The NO_x at least -20% (van Geffen et al., 2022). A higher NO_x to $\text{NO}:\text{NO}_2$ ratio can also have local variations, which affect the total emissions. For the majority NO_x seems to be emitted as NO , which is then rapidly turned into NO_2 after which an equilibrium is reached, of which the speed depends on the availability of O_3 . Beirle et al. (2021) recently gave a modelled estimate of the ratio, which was very close to the factor 1.32 (\pm ratio of 1.5 at the upper end of the literature values (Lange et al., 2022; Beirle et al., 2021; Griffin et al., 2021) would result in an additional underestimation of about -15%. The lifetime values reported in literature show a more random variation. Assuming the 3.3 hour estimate from Fioletov et al. (2022) is more accurate for emission zones, this would add an additional 20% given in his original study, with values moving towards 1.0 for industrial areas just north of the equator while values tended towards higher ranges (1.6) for less industrialized and high-latitude regions. Finally, there is the influence of the wind fields, for which we estimate an uncertainty of up to 1m/s over flat to 2m/s in mountainous and hilly areas. The effect can move in both directions and translates into an uncertainty of around 15-20% for average conditions over Germany (based on the matched wind fields). Taken together, these error terms result into a Germany averaged error range between 10-30% for the Gaussian plume method. The low error estimate corresponds to source regions where the low bias of the TROPOMI VCDs, effectively low biasing the emissions, are counteracted by the potentially high bias in the emissions of the $\text{NO}_x:\text{NO}_2$ ratio and effective lifetimes. Both values should be seen as conservative estimates, which would occur in the extremely unlikely case that % low bias to our estimates. Taken all together these values add up to an underestimation of about -45%, which is close to the inaccuracy in observed difference. The positive difference observed away from the major emitters, and especially in regions with intensive use of arable land⁶, could potentially hint at underestimation of soil emissions throughout Germany (Fig. 5). As discussed in section 2.1.1, soil emissions show a large range within the literature, with strong variations due to the availability of nitrogen, soil type, humidity and temperature. Some variations are observed between the years which could reflect changes in any of these parameters. The detection limit of the instrument does not seem like a likely candidate as it would result in low bias, similarly the slight high bias of the product cannot explain the larger differences observed in the northwest. The other two parameters that can cause a systematic offset, the $\text{NO}_x:\text{NO}_2$ ratio, lifetime, AMF and wind fields all nudge the estimate in the same direction for all locations in the domain

⁶<https://www.eea.europa.eu/data-and-maps/figures/agricultural-land-use-intensity-1>

of interest. In reality not all errors point in a similar direction, with the AMF/concentration term pointing in opposite directions for background and source regions. A complete error analysis based on simulated observations with controlled conditions and a subsequent Monte-Carlo analysis of error propagation can give a more accurate estimate, but that falls outside of the scope of this study. Ratios and lifetime, also do not seem to be a logical suspect. The $\text{NO}_x:\text{NO}_2$ ratio can at most cause a few percent of a positive bias (i.e. a ratio of 1.25 would only result in a few percent difference) while the lifetime would need to double or triple to explain the difference observed in the north-west.

The year to year variations in the TROPOMI- NO_2 derived emissions are of the order of a few to ten % (Fig 8). While the estimated errors of individual years are larger than those variations, most error components will point in similar directions stay consistent between the years. For example the A similar conclusion was made by Fioletov et al. (2022) who performed various experiments to test the impact of a common offset, in lifetime and plume-width, on the emission estimates of several years. In our case the consistency of the TROPOMI product version ensures that negative bias in the TROPOMI product can be expected to stay stable between the years over the high VCD regions while staying slightly positive over background regions. The only terms that are expected to slightly change are the $\text{NO}_x:\text{NO}_2$ ratio, the effective lifetime, and changes in wind-patterns. The changes in wind-patterns will only matter for regions in the border regions as misinterpretation of the wind-fields will typically results in the wrongful attribution of emissions within Germany. These This leaves the $\text{NO}_x:\text{NO}_2$ and effective lifetime as the main source of uncertainty, both related to the timing of emissions and the chemistry. A potential method to constrain this effect is performing a CTM run over the same period but with fixed yearly emissions over the whole period. The emission estimate methodology of this study can then be used to estimate the emissions of the individual years and thus derive the influence of changing chemistry and meteorology. This however again falls outside of the scope of this study.

5 Conclusions and Outlook

This work has shown that TROPOMI can be used as a verification tool for emission inventories, even for those inventory compilers, which are unfamiliar with remote sensing data, given the here presented web-tool. Emission inventory compilers may monitor near-real-time trends in NO_x emissions with the tool via top-down space-borne data without the need to wait for the completion of the statistical data required for the classic statistical "bottom-up" approach for the calculation of emissions. This is of particular importance for the quantification of unforeseen events such as the outbreak of the COVID-19 pandemic, which has been shown in this paper by comparing the 2019 emission data to the COVID-COVID-19 (2020) and post-COVID post-COVID-19 year (2021). Individual sectors are, however, difficult to assess given the low spatial resolution of TROPOMI. However, if we look at single large contributors to emissions such as the public power sector shown in Fig. 9 it is possible to track the rebound in emissions after the corona year 2020. Which has been due to the increased usage of coal fired power plants for power generation in 2021 compared to 2020. Similar trends and changes in NO_2 concentration may now be assessed by the emission inventory community worldwide as they are now able to compare their countries results to others using the here presented fully transparent methods. This has previously not been possible in a convenient way for inventory compilers. As at least comprehensive data science knowledge is required to access and query other data products e.g. from the ECMWF

atmospheric data ~~store (adsstorage (ADS⁷))~~, the web tool is complemented by the source code offer, which specifically invites other developers to extend the space-borne emissions code-base and web-tool by their own contributions. ~~This truly offers a substantial value to the international emission inventory community.~~

630 Space-borne data from TROPOMI and other satellites contain valuable information that can be used as a verification tool for emission inventories. NO_x retrievals from space-borne sensors such as OMI and TROPOMI can be used to monitor the quite dramatically decreasing evolution of NO_x emissions over the years with new emission estimation methods such as (Fioletov et al., 2017). Although, sub-sector and facility related data still is difficult to assess, ~~given the 3.5x5.5km² resolution of TROPOMI, the~~ the data still delivers valuable insight into the coarser spatial distribution of emission clusters, such as the
635 chemical industry parks around Halle and Leipzig or large coal fired power stations in the east of Germany. This may help to monitor emission reductions directly of these large industrial clusters. This satellite-based emission estimates, based on a single consistent methodology applied to several countries, can be used to verify the compliance towards meeting the air pollution reduction targets throughout the whole of the European Union, which ensures a maximum of transparency for all stakeholders. This ultimately values the principals of the European Green Deal initiative⁸, which tries to leverage new technology for a
640 sustainable EU.

With the presented space-emissions tool other emission inventory compilers without remote sensing expertise are encouraged to employ space-emissions for verification of their inventories. This would make the space-emissions tool a critical building block of emission compliance reporting thanks to the Copernicus Sentinel data set (i.e., TROPOMI) that are provided by ESA. We are looking forward to the feedback from the emission inventory community and their results using the online and
645 offline tools. ~~While the~~ The here developed methodology and online (and offline) tool ~~is~~ was initially focused on NO_x emission estimates from TROPOMI observations. In future the incorporation of OMI data would also extend the time series to the year 2005, which is of great importance for the verification of a more complete time-series of the inventory. The additions of other pollutants should also be envisioned for future work under the reservation that the respective method is applicable to the selected pollutant. In the near future the geostationary Sentinel-4 satellite is scheduled for launch and will provide hourly data
650 on tropospheric constituents over Europe. This will allow tooling such as those used in this manuscript to explore additional functionality such as the measurement of time profiles and might allow for emission estimates on a weekly or even daily basis and provide information on the diurnal emission cycle. While the methodology was only applied on a yearly basis in this study, TROPOMI has enough spatio-temporal coverage to move to seasonal or monthly estimates, potentially trading the spatial resolution of the emission fields for an increase in temporal resolvability.

655 Future improvements to the methodology should focus on updating the AMF with the help of higher resolution modelled fields, the addition of a location dependent lifetime (for example based on concentration of NO₂, O₃ and OH), the addition of local NO_x:NO₂ ratios and local corrections for diurnal and seasonal cycles) which all three would make sense from a physical perspective and form the largest uncertainty in the method outside satellite bias. Some of these improvements require simulated model fields, of which some are available in the form of (open-access) CAMS ensemble runs(~~link website, model fields~~).

⁷<https://ads.atmosphere.copernicus.eu>

⁸https://commission.europa.eu/strategy-and-policy/priorities-2019-2024/european-green-deal_en

660 Other required variables such as temperature, UV radiation, precipitation and humidity, which would be used for adjusted lifetimes, are also available at the various ECMWF data storage. These quantities and/or estimates can be downloaded with the ERA download tool and already make a relatively easy improvement to the lifetime estimates and thereby reduce the overall uncertainty of those terms.

Code and data availability. An offline version of the emission code is available at <https://github.com/UBA-DE-Emissionsituation/space-emissions>. All code used to produce further results figures etc, can be provided on request. The TROPOMI L2 data product versions (OFFL / PAL) can be accessed through the ESA Sentinel-5P data hub (<https://s5phub.copernicus.eu>) and the PAL data-portal (<https://data-portal.s5p-pal.com/>). The emission inventory data-sets can be accessed via <https://ir.umweltbundesamt.de/2022/>, and the GNFR/NFR data sets via Link to NFR and Link to GNFR.

Appendix A: Additional figures




Figure A1. Fraction of NO_x emissions emitted by each emission sector for each grid cell within the German domain. Yellow indicates locations with emissions dominated (>50%) by an individual source sector. The displayed data is based on gridded GNFR inventory emissions of 2019.




Figure A2. Fraction of NO_x emissions emitted by each emission sector for each grid cell within the German domain, smoothed with gaussian. The displayed data is based on gridded GNFR inventory emissions of 2019.



Figure A3. Emission source locations selected to produce sectoral trends. The produced masks are based on the results shown in Fig. A2, for all locations with an emission fraction above 50%. used to distinguish different source sectors in the emissions derived from satellite data.




Figure A4. Difference between the satellite derived and inventory emissions (2019) for years 2019-2021 over two zoom regions. The red values indicate a higher value for the satellite derived emissions compared to the inventory emissions. Upper row depicting the industrial "Ruhr" region, whilst the lower 3 panels show Lusatia at the eastern border of Germany.

670 *Author contributions.* Enrico Dammers devised and implemented the methods presented in this paper and carried out the data analysis and writing of this publication together with Janot Tokaya, who made critical contributions to the data stream handling of the method (retrieval of the correct scenes, and CAMS meteorology data). Renske Timmermans coordinated the scientific work from the TNO side and gave critical input to the scientific work with respect to the atmospheric chemistry of NO_x. Christian Mielke interpreted the spatial data of the algorithm runs and wrote the parts of the publication that deal with the emission inventory relevant topics. Kevin Hausmann designed and implemented
675 the webtool and the level-0 emission estimation on the Code-DE Platform and provided the critical scientific environment for the success of this project. Debora Griffin and Chris McLinden helped devise the methods presented in this paper and gave critical input to the manuscript. Henk Eskes provided information on the satellite product. Finally, all authors discussed the results and reviewed the manuscript.

Competing interests. The authors declare that there are no competing interests present in carrying out the work presented in this publication. The work has been financed by the Umweltbundesamt in the Leitplankenproject (FKZ: 3720515010).

680 *Acknowledgements.* We acknowledge the hard work done by KNMI, ESA, the team behind the PAL data portal, and the TROPOMI teams for making TROPOMI a success and providing easy access to the NO₂ data. ECMWF ERA5 data Hersbach et al. (2020) was downloaded from the Copernicus Climate Change Service (C3S) Climate Data Store (<https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-pressure-levels?tab=overview>). We thank Stefan Feigenspan for providing the 2019 GRETA data and for sharing his knowledge on emission gridding.

685 **References**

- Anderson, G. and Klugmann, D.: A European lightning density analysis using 5 years of ATDnet data, *Natural Hazards and Earth System Sciences*, 14, 815–829, <https://doi.org/10.5194/nhess-14-815-2014>, 2014.
- Atkinson, R. W., Butland, B. K., Anderson, H. R., and Maynard, R. L.: Long-term concentrations of nitrogen dioxide and mortality: a meta-analysis of cohort studies, *Epidemiology (Cambridge, Mass.)*, 29, 460, 2018.
- 690 Barré, J., Petetin, H., Colette, A., Guevara, M., Peuch, V.-H., Rouil, L., Engelen, R., Inness, A., Flemming, J., Pérez García-Pando, C., Bowdalo, D., Meleux, F., Geels, C., Christensen, J. H., Gauss, M., Benedictow, A., Tsyro, S., Friese, E., Struzewska, J., Kaminski, J. W., Douros, J., Timmermans, R., Robertson, L., Adani, M., Jorba, O., Joly, M., and Kouznetsov, R.: Estimating lockdown-induced European NO₂ changes using satellite and surface observations and air quality models, *Atmospheric Chemistry and Physics*, 21, 7373–7394, <https://doi.org/10.5194/acp-21-7373-2021>, 2021.
- 695 Beirle, S., Boersma, K. F., Platt, U., Lawrence, M. G., and Wagner, T.: Megacity emissions and lifetimes of nitrogen oxides probed from space, *Science*, 333, 1737–1739, 2011.
- Beirle, S., Borger, C., Dörner, S., Li, A., Hu, Z., Liu, F., Wang, Y., and Wagner, T.: Pinpointing nitrogen oxide emissions from space, *Science Advances*, 5, eaax9800, <https://doi.org/10.1126/sciadv.aax9800>, 2019.
- Beirle, S., Borger, C., Dörner, S., Eskes, H., Kumar, V., de Laat, A., and Wagner, T.: Catalog of NO_x emissions from point sources as derived from the divergence of the NO₂ flux for TROPOMI, *Earth System Science Data*, 13, 2995–3012, <https://doi.org/10.5194/essd-13-2995-2021>, 2021.
- 700 Belch, J. J., Fitton, C., Cox, B., and Chalmers, J. D.: Associations between ambient air pollutants and hospital admissions: more needs to be done, *Environmental Science and Pollution Research*, 28, 61 848–61 852, <https://doi.org/10.1007/s11356-021-16544-0>, 2021.
- Bucsela, E. J., Pickering, K. E., Allen, D. J., Holzworth, R. H., and Krotkov, N. A.: Midlatitude Lightning NO_x Production Efficiency Inferred From OMI and WLLN Data, *Journal of Geophysical Research: Atmospheres*, 124, 13 475–13 497, <https://doi.org/https://doi.org/10.1029/2019JD030561>, 2019.
- CLRTAP: National gridded data of emissions (CLRTAP), <https://cdr.eionet.europa.eu/de/un/clrtap/gridded/envyizg6q/>, 2021.
- CLRTAP: LRTAP Convention - National emission inventories, https://cdr.eionet.europa.eu/de/un/clrtap/inventories/envygjjnq/index_html?, 2022.
- 710 Crippa, M., Guizzardi, D., Muntean, M., Schaaf, E., Lo Vullo, E., Solazzo, E., Monforti-Ferrario, F., Olivier, J., and Vignati, E.: EDGAR v5.0 global air pollutant emissions, European Commission, Joint Research Centre (JRC)[Dataset] PID: <http://data.europa.eu/89h/377801afb094-4943-8fdc-f79a7c0c2d19>, 2019.
- Dammers, E., McLinden, C. A., Griffin, D., Shephard, M. W., Van Der Graaf, S., Lutsch, E., Schaap, M., Gainairu-Matz, Y., Fioletov, V., Van Damme, M., Whitburn, S., Clarisse, L., Cady-Pereira, K., Clerbaux, C., Coheur, P. F., and Erismann, J. W.: NH₃ emissions from large point sources derived from CrIS and IASI satellite observations, *Atmospheric Chemistry and Physics*, 19, 12 261–12 293, <https://doi.org/10.5194/acp-19-12261-2019>, 2019.
- 715 Dammers, E., Shephard, M., Griffin, D., Chow, E., White, E., Hickman, J., Tokaya, J., Lutsch, E., Kharol, S., van der Graaf, S., Cady-Pereira, K., Bittman, S., McLinden, C., Erismann, J., and Schaap, M.: County-level ammonia emissions monitored worldwide, *Nature Geoscience*, a, a, https://assets.researchsquare.com/files/rs-1752718/v1_covered.pdf?c=1656451952, 2022.
- 720 Dammers, E., Tokaya, J., Timmermans, R., Schaap, M., Coenen, P., Mielke, C., and Hausmann, K.: Satellite-based Emission Verification, Pilot Study, <https://www.umweltbundesamt.de/publikationen/satellite-based-emission-verification>, 2023.

- de Foy, B., Wilkins, J. L., Lu, Z., Streets, D. G., and Duncan, B. N.: Model evaluation of methods for estimating surface emissions and chemical lifetimes from satellite data, *Atmospheric Environment*, 98, 66 – 77, <https://doi.org/https://doi.org/10.1016/j.atmosenv.2014.08.051>, 2014.
- 725 de Foy, B., Lu, Z., Streets, D. G., Lamsal, L. N., and Duncan, B. N.: Estimates of power plant NO_x emissions and lifetimes from OMI NO₂ satellite retrievals, *Atmospheric Environment*, 116, 1–11, <https://doi.org/https://doi.org/10.1016/j.atmosenv.2015.05.056>, 2015.
- Ding, J., van der A, R. J., Eskes, H. J., Mijling, B., Stavrou, T., van Geffen, J. H. G. M., and Veefkind, J. P.: NO_x Emissions Reduction and Rebound in China Due to the COVID-19 Crisis, *Geophysical Research Letters*, 47, e2020GL089912, <https://doi.org/https://doi.org/10.1029/2020GL089912>, e2020GL089912 2020GL089912, 2020.
- 730 Dore, C.: Technical Guidance for Emission Inventory Adjustments under the Amended Gothenburg Protocol: Inventory adjustments in context of ERCs, CEIP, a, a, <https://doi.org/2514/06AL22/2310>, 2022.
- Douros, J., Eskes, H., van Geffen, J., Boersma, K. F., Compennolle, S., Pinardi, G., Blechschmidt, A.-M., Peuch, V.-H., Colette, A., and Veefkind, P.: Comparing Sentinel-5P TROPOMI NO₂ column observations with the CAMS regional air quality ensemble, *Geoscientific Model Development*, 16, 509–534, <https://doi.org/10.5194/gmd-16-509-2023>, 2023.
- 735 EEA: MEP/EEA air pollutant emission inventory guidebook 2019: Technical guidance to prepare national emission inventories', EEA Technical report, (12/2019), <https://www.eea.europa.eu/publications/emep-eea-guidebook-2019>, 2019.
- EPH: Jänschwalde(Accessed on < 05-10-2022 >), <https://www.eppowereurope.cz/en/companies/janschwalde/>, 2022.
- EU: Directive 2001/81/EC of the European Parliament and of the Council on national emission ceilings for certain atmospheric pollutants, <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2001:309:0022:0030:EN:PDF>, 2022.
- 740 Fioletov, V., McLinden, C., Krotkov, N., Moran, M., and Yang, K.: Estimation of SO₂ emissions using OMI retrievals, *Geophysical Research Letters*, 38, 2011.
- Fioletov, V., McLinden, C. A., Kharol, S. K., Krotkov, N. A., Li, C., Joiner, J., Moran, M. D., Vet, R., Visschedijk, A. J., van der Gon, D., et al.: Multi-source SO₂ emission retrievals and consistency of satellite and surface measurements with reported emissions, *Atmospheric Chemistry and Physics*, 17, 12 597–12 616, 2017.
- 745 Fioletov, V., McLinden, C. A., Griffin, D., Krotkov, N., Liu, F., and Eskes, H.: Quantifying urban, industrial, and background changes in NO₂ during the COVID-19 lockdown period based on TROPOMI satellite observations, *Atmospheric Chemistry and Physics*, 22, 4201–4236, <https://doi.org/10.5194/acp-22-4201-2022>, 2022.
- Fioletov, V. E., McLinden, C. A., Krotkov, N., and Li, C.: Lifetimes and emissions of SO₂ from point sources estimated from OMI, *Geophysical Research Letters*, 42, 1969–1976, <https://doi.org/https://doi.org/10.1002/2015GL063148>, 2015.
- 750 Galloway, J. N., Aber, J. D., Erisman, J. W., Seitzinger, S. P., Howarth, R. W., Cowling, E. B., and Cosby, B. J.: The nitrogen cascade, *Bioscience*, 53, 341–356, 2003.
- Goldberg, D. L., Lu, Z., Streets, D. G., de Foy, B., Griffin, D., McLinden, C. A., Lamsal, L. N., Krotkov, N. A., and Eskes, H.: Enhanced Capabilities of TROPOMI NO₂: Estimating NO_x from North American Cities and Power Plants, *Environmental science & technology*, 53, 12 594–12 601, 2019.
- 755 Goldberg, D. L., Anenberg, S. C., Griffin, D., McLinden, C. A., Lu, Z., and Streets, D. G.: Disentangling the impact of the COVID-19 lockdowns on urban NO₂ from natural variability, *Geophysical Research Letters*, 47, e2020GL089269, 2020.
- Goldberg, D. L., Anenberg, S. C., Kerr, G. H., Moheg, A., Lu, Z., and Streets, D. G.: TROPOMI NO₂ in the United States: A Detailed Look at the Annual Averages, Weekly Cycles, Effects of Temperature, and Correlation With Surface NO₂ Concentrations, *Earth's Future*, 9, e2020EF001 665, <https://doi.org/https://doi.org/10.1029/2020EF001665>, e2020EF001665 2020EF001665, 2021.

- 760 Granier, C., Darras, H., Denier van der Gon, J., Doubalova, N., Elguindi, B., Galle, M., Gauss, M., Guevara, J., Jalkanen, J., and Kuenen, C.: The Copernicus Atmosphere Monitoring Service Global and Regional Emissions, Report April 2019 version (Research Report), ECMWF, Reading, UK, Reading, UK,[data set], 10, 2019.
- Griffin, D., McLinden, C. A., Racine, J., Moran, M. D., Fioletov, V., Pavlovic, R., Mashayekhi, R., Zhao, X., and Eskes, H.: Assessing the impact of Corona-Virus-19 on nitrogen dioxide levels over Southern Ontario, Canada, *Remote Sensing*, 12, 4112, 2020.
- 765 Griffin, D., McLinden, C. A., Dammers, E., Adams, C., Stockwell, C. E., Warneke, C., Bourgeois, I., Peischl, J., Ryerson, T. B., Zarzana, K. J., Rowe, J. P., Volkamer, R., Knote, C., Kille, N., Koenig, T. K., Lee, C. F., Rollins, D., Rickly, P. S., Chen, J., Fehr, L., Bourassa, A., Degenstein, D., Hayden, K., Mihele, C., Wren, S. N., Liggio, J., Akingunola, A., and Makar, P.: Biomass burning nitrogen dioxide emissions derived from space with TROPOMI: methodology and validation, *Atmospheric Measurement Techniques*, 14, 7929–7957, <https://doi.org/10.5194/amt-14-7929-2021>, 2021.
- 770 Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J.-N.: ERA5 hourly data on pressure levels from 1979 to present. Copernicus Climate Change Service (C3S) Climate Data Store (CDS). (Accessed on < 18-01-2022 >), <https://doi.org/10.24381/cds.bd0915c6>, 2018.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., et al.: The ERA5 global reanalysis, 2020.
- 775 Jamali, S., Klingmyr, D., and Tagesson, T.: Global-Scale Patterns and Trends in Tropospheric NO₂ Concentrations, 2005–2018, *Remote Sensing*, 12, <https://doi.org/10.3390/rs12213526>, 2020.
- Jonson, J. E., Borken-Kleefeld, J., Simpson, D., Nyíri, A., Posch, M., and Heyes, C.: Impact of excess NO_x emissions from diesel cars on air quality, public health and eutrophication in Europe, *Environmental Research Letters*, 12, 094017, <https://doi.org/10.1088/1748-9326/aa8850>, 2017.
- 780 Kuenen, J., Dellaert, S., Visschedijk, A., Jalkanen, J.-P., Super, I., and Denier van der Gon, H.: CAMS-REG-v4: a state-of-the-art high-resolution European emission inventory for air quality modelling, *Earth System Science Data*, 14, 491–515, <https://doi.org/10.5194/essd-14-491-2022>, 2022.
- Lange, K., Richter, A., and Burrows, J. P.: Variability of nitrogen oxide emission fluxes and lifetimes estimated from Sentinel-5P TROPOMI observations, *Atmospheric Chemistry and Physics*, 22, 2745–2767, <https://doi.org/10.5194/acp-22-2745-2022>, 2022.
- 785 Lorente, A., Boersma, K., Eskes, H., Veefkind, J., Van Geffen, J., De Zeeuw, M., Denier Van Der Gon, H., Beirle, S., and Krol, M.: Quantification of nitrogen oxides emissions from build-up of pollution over Paris with TROPOMI, *Scientific reports*, 9, 1–10, 2019.
- Manders, A. M., Builtjes, P. J., Curier, L., van der Gon, D., Hugo, A., Hendriks, C., Jonkers, S., Kranenburg, R., Kuenen, J. J., Segers, A. J., et al.: Curriculum vitae of the LOTOS–EUROS (v2. 0) chemistry transport model, *Geoscientific Model Development*, 10, 4145–4173, <https://doi.org/https://doi.org/10.5194/gmd-10-4145-2017>, 2017.
- 790 McLinden, C., Griffin, D., Fioletov, V., Zhang, J., Dammers, E., Adams, C., Loria, M., Krotkov, N., and Lamsal, N.: Monitoring of total and off-road NO_x emissions from Canadian oil sands surface mining using the Ozone Monitoring Instrument, To be submitted, *Atmospheric Chemistry and Physics*, can be shared upon request, 2022.
- McLinden, C. A., Fioletov, V., Shephard, M. W., Krotkov, N., Li, C., Martin, R. V., Moran, M. D., and Joiner, J.: Space-based detection of missing sulfur dioxide sources of global air pollution, *Nature Geoscience*, 9, 496–500, 2016.
- 795 McLinden, C. A., Adams, C. L., Fioletov, V., Griffin, D., Makar, P. A., Zhao, X., Kovachik, A., Dickson, N., Brown, C., Krotkov, N., et al.: Inconsistencies in sulfur dioxide emissions from the Canadian oil sands and potential implications, *Environmental Research Letters*, 16, 014012, <https://doi.org/https://doi.org/10.1088/1748-9326/abcbbb>, 2020.

- Mijling, B., Van Der A, R., Boersma, K., Van Roozendaal, M., De Smedt, I., and Kelder, H.: Reductions of NO₂ detected from space during the 2008 Beijing Olympic Games, *Geophysical Research Letters*, 36, 2009.
- 800 Miyazaki, K., Eskes, H., and Sudo, K.: Global NO_x emission estimates derived from an assimilation of OMI tropospheric NO₂ columns, *Atmospheric Chemistry and Physics*, 12, 2263–2288, 2012.
- Paige, C. C. and Saunders, M. A.: LSQR: An Algorithm for Sparse Linear Equations and Sparse Least Squares, *ACM Trans. Math. Software*, pp. 43–71, 1982.
- Pommier, M., McLinden, C. A., and Deeter, M.: Relative changes in CO emissions over megacities based on observations from space, *Geophysical research letters*, 40, 3766–3771, <https://doi.org/https://doi.org/10.1002/grl.50704>, 2013.
- 805 Schneider, C., Pelzer, M., Toenges-Schuller, N., Nacken, M., and Niederau, A.: ArcGIS basierte Lösung zur detaillierten, deutschlandweiten Verteilung (Gridding) nationaler Emissionsjahreswerte auf Basis des Inventars zur Emissionsberichterstattung: Forschungskennzahl 3712 63 240 2, *Texte*, 71, 5, 2016.
- Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics*, A Wiley-Inter Science Publication, John Wiley & Sons Inc, Hoboken, New Jersey, 2006.
- 810 Simpson, D.: Copernicus Atmosphere Monitoring Service soil global NO_x emissions (CAM5-GLOB-SOIL v2.2), <https://doi.org/https://doi.org/10.24380/kz2r-fe18>, 2022.
- Simpson, D. and Darras, S.: Global soil NO emissions for Atmospheric Chemical Transport Modelling: CAM5-GLOB-SOIL v2.2, *Earth System Science Data Discussions*, 2021, 1–35, <https://doi.org/10.5194/essd-2021-221>, 2021.
- 815 Simpson, D., Winiwarter, W., Börjesson, G., Cinderby, S., Ferreira, A., Guenther, A., Hewitt, C. N., Janson, R., Khalil, M. A. K., Owen, S., et al.: Inventorying emissions from nature in Europe, *Journal of Geophysical Research: Atmospheres*, 104, 8113–8152, <https://doi.org/https://doi.org/10.1029/98JD02747>, 1999.
- UBA: German Informative Inventory Report, https://iir.umweltbundesamt.de/2023/general/uncertainty_evaluation/start, 2023.
- UNECE: Convention on Long-Range Transboundary Air Pollution, <https://unece.org/environment-policy/air>.
- 820 van der Walt, S., Schönberger, J. L., Nunez-Iglesias, J., Boulogne, F., Warner, J. D., Yager, N., Gouillart, E., Yu, T., and the scikit-image contributors: scikit-image: image processing in Python, *PeerJ*, 2, e453, <https://doi.org/10.7717/peerj.453>, 2014.
- van Geffen, J., Eskes, H., Compernelle, S., Pinardi, G., Verhoelst, T., Lambert, J.-C., Sneep, M., ter Linden, M., Ludewig, A., Boersma, K. F., and Veefkind, J. P.: Sentinel-5P TROPOMI NO₂ retrieval: impact of version v2.2 improvements and comparisons with OMI and ground-based data, *Atmospheric Measurement Techniques*, 15, 2037–2060, <https://doi.org/10.5194/amt-15-2037-2022>, 2022.
- 825 Vattenfall: Vattenfall to phase-out 1,000 MW lignite capacity (Accessed on < 05-10-2022 >), <https://group.vattenfall.com/press-and-media/pressreleases/2015/vattenfall-to-phase-out-1000-mw-lignite-capacity>, 2015.
- Veefkind, J. P., Aben, I., McMullan, K., Förster, H., de Vries, J., Otter, G., Claas, J., Eskes, H. J., de Haan, J. F., Kleipool, Q., van Weele, M., Hasekamp, O., Hoogeveen, R., Landgraf, J., Snel, R., Tol, P., Ingmann, P., Voors, R., Kruizinga, B., Vink, R., Visser, H., and Levelt, P. F.: TROPOMI on the ESA Sentinel-5 Precursor: A GMES mission for global observations of the atmospheric composition for climate, air quality and ozone layer applications, *rse*, 120, 70–83, <https://doi.org/10.1016/j.rse.2011.09.027>, 2012.
- 830 Verhoelst, T., Compernelle, S., Pinardi, G., Lambert, J.-C., Eskes, H. J., Eichmann, K.-U., Fjæraa, A. M., Granville, J., Niemeijer, S., Cede, A., Tiefengraber, M., Hendrick, F., Pazmiño, A., Bais, A., Bazureau, A., Boersma, K. F., Bogner, K., Dehn, A., Donner, S., Elokhorov, A., Gebetsberger, M., Goutail, F., Grutter de la Mora, M., Gruzdev, A., Gratsea, M., Hansen, G. H., Irie, H., Jepsen, N., Kanaya, Y., Karagkiozidis, D., Kivi, R., Kreher, K., Levelt, P. F., Liu, C., Müller, M., Navarro Comas, M., Piters, A. J. M., Pommereau, J.-P., Portafaix, T., Prados-Roman, C., Puentedura, O., Querel, R., Remmers, J., Richter, A., Rimmer, J., Rivera Cárdenas, C., Saavedra de Miguel, L.,
- 835

- Sinyakov, V. P., Stremme, W., Strong, K., Van Roozendaal, M., Veeffkind, J. P., Wagner, T., Wittrock, F., Yela González, M., and Zehner, C.: Ground-based validation of the Copernicus Sentinel-5P TROPOMI NO₂ measurements with the NDACC ZSL-DOAS, MAX-DOAS and Pandonia global networks, *Atmospheric Measurement Techniques*, 14, 481–510, <https://doi.org/10.5194/amt-14-481-2021>, 2021.
- 840 Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D., Burovski, E., Peterson, P., Weckesser, W., Bright, J., van der Walt, S. J., Brett, M., Wilson, J., Millman, K. J., Mayorov, N., Nelson, A. R. J., Jones, E., Kern, R., Larson, E., Carey, C. J., Polat, İ., Feng, Y., Moore, E. W., VanderPlas, J., Laxalde, D., Perktold, J., Cimrman, R., Henriksen, I., Quintero, E. A., Harris, C. R., Archibald, A. M., Ribeiro, A. H., Pedregosa, F., van Mulbregt, P., and SciPy 1.0 Contributors: SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python, *Nature Methods*, 17, 261–272, <https://doi.org/10.1038/s41592-019-0686-2>, 2020.
- 845 Wang, Z., Bovik, A. C., Sheikh, H. R., and Simoncelli, E. P.: Image quality assessment: from error visibility to structural similarity, *IEEE transactions on image processing*, 13, 600–612, 2004.
- WHO: WHO global air quality guidelines: particulate matter (PM_{2.5} and PM₁₀), ozone, nitrogen dioxide, sulfur dioxide and carbon monoxide, World Health Organization, <https://www.who.int/publications/i/item/9789240034228>, 2021.
- Yienger, J. J. and Levy II, H.: Empirical model of global soil-biogenic NO_x emissions, *Journal of Geophysical Research: Atmospheres*, 100, 11 447–11 464, <https://doi.org/https://doi.org/10.1029/95JD00370>, 1995.
- 850 Zhao, X., Fioletov, V., Alwarda, R., Su, Y., Griffin, D., Weaver, D., Strong, K., Cede, A., Hanisco, T., Tiefengraber, M., McLinden, C., Eskes, H., Davies, J., Ogyu, A., Sit, R., Abboud, I., and Lee, S. C.: Tropospheric and Surface Nitrogen Dioxide Changes in the Greater Toronto Area during the First Two Years of the COVID-19 Pandemic, *Remote Sensing*, 14, <https://doi.org/10.3390/rs14071625>, 2022.