

The Earth system model CLIMBER-X v1.0 – Part 2: The global carbon cycle

Matteo Willeit¹, Tatiana Ilyina², Bo Liu², Christoph Heinze³, Mahé Perrette¹, Malte Heinemann⁴, Daniela Dalmonech⁵, Victor Brovkin^{2,6,1}, Guy Munhoven⁷, Janine Boerker⁸, Jens Hartmann⁸, Gibran Romero Mujalli⁸, and Andrey Ganopolski¹

¹Potsdam Institute for Climate Impact Research (PIK), Member of the Leibniz Association, P.O. Box 6012 03, D-14412 Potsdam Germany

²Max Planck Institute for Meteorology, Hamburg, Germany

⁶CEN, University of Hamburg, Germany; also a guest at PIK, Potsdam, Germany

³University of Bergen, Bergen, Norway

⁴Christian-Albrechts-Universität zu Kiel, Kiel, Germany

⁵Institute for Agriculture and Forestry Systems in the Mediterranean, National Research Council of Italy (CNR-ISAFOM), Perugia, Italy

⁷Dépt. d'Astrophysique, Géophysique et Océanographie, Université de Liège, Liège, Belgium

⁸Universität Hamburg, Hamburg, Germany

Correspondence: Matteo Willeit (willeit@pik-potsdam.de)

Abstract. The carbon cycle component of the newly developed Earth System Model of intermediate complexity CLIMBER-X is presented. The model represents the cycling of carbon through atmosphere, vegetation, soils, seawater and marine sediments. Exchanges of carbon with geological reservoirs occur through sediment burial, rock weathering and volcanic degassing. The state-of-the-art HAMOCC6 model is employed to simulate ocean biogeochemistry and marine sediments processes. The land model PALADYN simulates the processes related to vegetation and soil carbon dynamics, including permafrost and peatlands. The dust cycle in the model allows for an interactive determination of the input of the micro-nutrient iron into the ocean. A rock weathering scheme is implemented into the model, with the weathering rate depending on lithology, runoff and soil temperature. CLIMBER-X includes a simple representation of the methane cycle, with explicitly modelled natural emissions from land and the assumption of a constant residence time of CH₄ in the atmosphere. Carbon isotopes ¹³C and ¹⁴C are tracked through all model compartments and provide a useful diagnostic for model-data comparison.

A comprehensive evaluation of the model performance for present-day and the historical period shows that CLIMBER-X is capable of realistically reproducing the historical evolution of atmospheric CO₂ and CH₄, but also the spatial distribution of carbon on land and the 3D structure of biogeochemical ocean tracers. The analysis of model performance is complemented by an assessment of carbon cycle feedbacks and model sensitivities compared to state-of-the-art CMIP6 models.

Enabling interactive carbon cycle in CLIMBER-X results in a relatively minor slow-down of model computational performance by ~20%, compared to a throughput of ~10,000 simulation years per day on a single node with 16 CPUs on a high performance computer in a climate-only model setup. CLIMBER-X is therefore well suited to investigate the feedbacks between climate and the carbon cycle on temporal scales ranging from decades to >100,000 years.

1 Introduction

20 Atmospheric CO₂ exerts a profound control on the state of the Earth system. Although it is present only in tiny concentrations in the present-day atmosphere, by absorbing radiation in the longwave spectral range it has a substantial effect on the energy balance of the Earth. In the present day atmosphere, CO₂ is the second most important greenhouse gas, after water vapor. CO₂ is also a fundamental molecule for life on Earth, as it serves as 'food' in the photosynthesis process. The atmospheric CO₂ concentration is hence a main control on the growth rate of plants on land.

25 From ice core data it is well known that atmospheric CO₂ concentrations showed pronounced variations over the last million years (e.g. Petit et al., 1999; Augustin et al., 2004) that played an important role for the climate evolution over the Pleistocene (last ~2.6 million years) by amplifying the variations associated with glacial-interglacial cycles (e.g. Ganopolski and Calov, 2011; Abe-Ouchi et al., 2013). Furthermore, on even longer time scales, a secular decrease in CO₂ is thought to have been the main driver of the gradual cooling over the Cenozoic (last 66 million years) (e.g. Raymo and Ruddiman, 1992).

30 Over the last few centuries, human activities have strongly disrupted the natural CO₂ balance, by directly emitting CO₂ from fossil sources into the atmosphere. The resulting increase in atmospheric CO₂ has been the main factor for the observed rapid climate warming since the preindustrial period (e.g. Gulev et al., 2021).

Modelling the atmospheric CO₂ concentration is thus fundamental both for understanding past climate changes and for predicting the future evolution of the Earth system under different anthropogenic emission scenarios. However, it is far from
35 trivial, because atmospheric CO₂ is the result of complex biogeochemical processes on land, in the ocean, in marine sediments and in the lithosphere. Additionally, because of the long time scales involved in some of the carbon cycle processes, the interactive simulation of atmospheric CO₂ has been, and still is, a challenge for state-of-the-art Earth system models. Fast Earth system models of intermediate complexity have therefore been extensively employed for investigating carbon cycle-climate feedbacks (e.g. Bern3D (Müller et al., 2008; Tschumi et al., 2011; Stocker et al., 2013), cGENIE (Ridgwell et al.,
40 2007; Cao et al., 2009), CLIMBER-2 (Brovkin et al., 2002, 2007, 2012), iLOVECLIM (Bouttes et al., 2015), LOVECLIM (Goosse et al., 2010) and Uvic (Eby et al., 2009; Zickfeld et al., 2011; Mengis et al., 2020)). Among these, CLIMBER-2 has successfully reproduced glacial-interglacial variations in CO₂ (Ganopolski and Brovkin, 2017; Willeit et al., 2019), but some of the processes involved remain uncertain. CLIMBER-X builds on the past experience in modelling the global carbon cycle with CLIMBER-2, but adds an improved and more detailed representation of carbon cycle processes both on land and in the
45 ocean. Improvements include a generally higher spatial resolution, a 3D ocean model, a state-of-the-art ocean biogeochemistry and marine sediment model, a more comprehensive description of vegetation and soil carbon processes, including permafrost and peatlands, and a new chemical weathering scheme.

In the following, the biogeochemistry components of CLIMBER-X are presented. The climate core of CLIMBER-X is described in detail in Willeit et al. (2022).




Figure 1. Schematic illustration of the natural biogeochemical cycles in the CLIMBER-X model.

50 2 Model description

CLIMBER-X represents the cycling of carbon through atmosphere, vegetation, soils, seawater and marine sediments. Through sediment burial, chemical weathering of rocks and volcanic degassing, carbon is also exchanged with geological reservoirs. A schematic illustration of the carbon cycle in the model is shown in [FigureFig. 1](#). The carbon cycle component of CLIMBER-X consists of the ocean biogeochemistry and marine sediment models from HAMOCC6 (Maier-Reimer and Hasselmann, 1987; Ilyina et al., 2013; Heinze et al., 1999; Mauritsen et al., 2019) and the land model PALADYN (Willeit and Ganopolski, 2016), which includes dynamic vegetation, a soil carbon model and the weathering model of Hartmann (2009) and Börker et al. (2020). The atmospheric CO₂ concentration is determined interactively by the exchange of carbon between atmosphere, seawater, land and lithosphere. The model includes a representation of the dust cycle, with simulated dust deposition determining the input of the micro-nutrient iron into the ocean. CLIMBER-X also includes a simple representation of the methane cycle, with explicitly modelled natural emissions from land and the assumption of a constant residence time of CH₄ in the atmosphere. The model is enabled with the carbon isotopes ¹³C and ¹⁴C, which are tracked through all model compartments.

The different model components are described in more detail in the following sections.

2.1 Ocean biogeochemistry and marine sediments: HAMOCC

HAMOCC (Maier-Reimer and Hasselmann, 1987; Maier-Reimer et al., 1993; Ilyina et al., 2013) is a state-of-the-art ocean biogeochemistry model, which is part of the MPI-ESM, the Earth system model of the Max Planck Institute for Meteorology

(MPI). The latest version (Mauritsen et al., 2019), which is the version employed by the MPI in the ~~latest model intercomparison project~~ [Coupled Model Intercomparison Project Phase 6](#) (CMIP6) ~~for the Sixth Assessment Report of the IPCC~~, has been the starting point for the implementation of the model into CLIMBER-X. As a first step, the original HAMOCC6 code has been adapted to the CLIMBER-X structure. Notably, for easier parallelisation, it has been transformed from a 3D model into a 1D
70 vertical column model in which each water column is independent from the others. This is possible because the biogeochemical processes in the model are restricted to local vertical interactions. The different columns are interacting only through horizontal advection by ocean currents, which takes place in the ocean model.

HAMOCC represents the biogeochemical processes in the water column, the sediments, and at the air-sea interface. Marine biology dynamics is based on an extended NPZD (nutrients, phytoplankton, zooplankton, and detritus) approach (Six and
75 Maier-Reimer, 1996). The carbonate chemistry in the model follows the latest OMIP protocol (Orr et al., 2017), which uses the robust and safe pH calculation routines from SolveSAPHE-r1 (Munhoven, 2013). In the water column, the following biogeochemical tracers are simulated: dissolved inorganic carbon (DIC), total alkalinity (TA), phosphate (PO_4), nitrate (NO_3), nitrous oxide (N_2O), dissolved nitrogen gas (N_2), silicate (SiO_2), dissolved bioavailable iron (Fe), dissolved oxygen (O_2), phytoplankton (Phy), zooplankton (Zoo), dissolved organic matter (DOC), particulate organic matter (POC), opal shells, calcium carbonate shells (CaCO_3), terrigenous material (dust) and hydrogen sulfide (H_2S). The composition of organic material
80 follows a constant Redfield ratio ($\text{C:N:P:O}_2 = 122:16:1:-172$) after Takahashi et al. (1985), and for the micronutrient iron ($\text{Fe:C} = 4 \times 10^{-6}:1$).

The marine sediment module, which is part of HAMOCC, is based on Heinze et al. (1999). It essentially simulates the same processes between dissolved tracers (DIC, TA, PO_4 , NO_3 , O_2 , Fe, SiO_2 , H_2S and N_2) in pore water and solid sediment
85 constituents (POC, opal, CaCO_3 and dust) as in the water column. Pore water tracers are exchanged with the overlying water column via diffusion. Sedimentation fluxes of POC, CaCO_3 , opal and dust are added to the solid components of the sediment. Accumulation of solid sediment material will lead to active sediment layer content being shifted to the burial layer and back if boundary conditions changes lead to chemical erosion of previously buried sediment.

Next we describe the changes introduced into HAMOCC as part of its implementation into CLIMBER-X.

90 N_2 fixation is represented by a diagnostic formulation, whereby the nitrate influx into the surface layer is a function of the nitrate deficit relative to phosphate, multiplied by a constant fixation rate (Ilyina et al., 2013). Prognostic N_2 fixers have recently been included in HAMOCC (Paulsen et al., 2017), based on the physiological characteristics of the cyanobacterium *Trichodesmium*. However, for simplicity and because uncertainties in nitrogen fixation remain large (e.g. Zehr and Capone, 2020), in CLIMBER-X cyanobacteria are disabled by default.

95 Following Heinemann et al. (2019), we have implemented a representation of aggregates in the model. Particulate organic carbon is assumed to form aggregates with the denser calcite and opal built during phytoplankton and zooplankton growth, as well as with dust particles. The sinking speed of these aggregates depends on their excess density (Gehlen et al., 2006; Heinemann et al., 2019). Note that this approach neglects the effects of, e.g., aggregate size distribution and porosity on the sinking speed (Maerz et al., 2020), and it does not, like other numerically more expensive schemes (e.g., Kriest and Evans,

100 2000) explicitly resolve the biological and physical aggregation and disaggregation processes. [The introduction of the ballasting scheme required a retuning of the dissolution rates of calcite and opal as shown in Table 1.](#)

Following recent evidence that the remineralisation of organic carbon depends on temperature (e.g. Laufkötter et al., 2017), we have introduced a Q10 temperature dependence for the remineralisation of POC and DOC (Segschneider and Bendtsen, 2013; Crichton et al., 2021), with a default Q10 value of 2. [The complete set of remineralisation parameters is listed in Table 1.](#)

105

In the original HAMOCC, iron complexation by organic substances is assumed when the iron concentration exceeds a given threshold and dissolved iron is then removed from the water column at a fixed rate. In CLIMBER-X, we explicitly model iron complexation, differentiating between free and complexed iron forms following Archer and Johnson (2000) and Parekh et al. (2004). The complexed iron is associated with an organic ligand and only the free iron is available for scavenging. The ligand concentration is assumed to be constant at 1 nmol kg^{-1} with a ligand stability constant of $1 \times 10^{11} \text{ kg mol}^{-1}$. The speciation of iron is then determined by equilibrium kinetics. The scavenging rate of free iron is a combination of a minimum scavenging rate and a scavenging rate that is proportional to the POC, calcite and opal concentrations following Aumont et al. (2015) and Hauck et al. (2013). Compared to HAMOCC we have also increased the stoichiometric iron ratio in organic compounds from $\text{Fe:C} = 3 \times 10^{-6}:1$ to $\text{Fe:C} = 4 \times 10^{-6}:1$. [The parameters related to the iron cycle are also reported in Table 1.](#)

115 The carbon 13 isotope has been recently implemented in HAMOCC by Liu et al. (2021). In CLIMBER-X we extended this approach to also include radiocarbon.

Since the ocean model in CLIMBER-X is a rigid lid model, following the OMIP protocol (Orr et al., 2017), we explicitly take into account the local concentration-dilution effect of the net surface freshwater flux, which changes surface DIC concentration and alkalinity. [Two options are available in the model to implement the dilution effect on DIC and alkalinity. The first one ensures that the net global surface tracer flux is zero by applying deviations from the global average freshwater flux to the global average surface tracer concentration. The second \(default\) option applies the actual local surface freshwater flux to compute a new virtual top ocean layer thickness and then dilutes the tracers accordingly. In this case, the conservation of tracer inventories is ensured by compensating disbalances over the global ocean. Additionally, during times when ocean volume is changing because of buildup or melt of land ice, concentrations of all tracers are globally adjusted while conserving tracer inventories. This is a reasonable simplification, considering that land ice volume changes occur on multi-millennial time scales, over which the ocean can be considered well mixed.](#)

125 Based on scale analysis, we have excluded fast sinking tracers (CaCO_3 , opal, POC and dust) from advection, as these particles have sinking speeds which are large enough so that vertical transfer between different grid cells is more rapid than horizontal transfer by advection would be, considering the relatively coarse resolution of the ocean model. Following a similar line of thought also short-lived tracers like phytoplankton and zooplankton are excluded from oceanic transport. However, convection and wind-driven surface vertical mixing are applied to all biogeochemical tracers.

In CLIMBER-X, HAMOCC is integrated with a time step of one day, which is also the time step of the physical ocean model.

Table 1. Modified HAMOCC parameters used in CLIMBER-X compared to HAMOCC6 (i.e. Table 2 in Ilyina et al. (2013)).

Parameter	Description	CLIMBER-X	HAMOCC6	Unit
<i>Nutrients</i>				
μ_{cyan}	N_2 fixation rate	0.0025	0.005	d^{-1}
$\lambda_{\text{det}}^{\text{ref}}$	POC remineralisation rate at temperature T_{ref}	0.05	0.025	d^{-1}
$\lambda_{\text{N}}^{\text{ref}}$	denitrification rate at temperature T_{ref}	0.15	0.07	d^{-1}
$\lambda_{\text{S}}^{\text{ref}}$	sulfate reduction rate at temperature T_{ref}	0.005	0.005	d^{-1}
Q_{10}	Q10 for temperature dependence of remineralisation rate	2	1	
T_{ref}	reference temperature for remineralisation rate	20		$^{\circ}\text{C}$
<i>Iron cycle</i>				
$f_{\text{Fe}}^{\text{dust}}$	fraction of iron mass in dust	0.025	0.035	kgkg^{-1}
d_{Fe}	iron solubility in surface water	0.01	0.01	
$R_{\text{Fe:C}}$	stoichiometric iron ratio in organic compounds	4×10^{-6}	3×10^{-6}	molFemolC^{-1}
L	total ligand concentration	1×10^{-9}		kmolm^{-3}
k_{L}	ligand stability constant	1×10^{11}		$\text{m}^3\text{kmol}^{-1}$
$l_{\text{scav}}^{\text{min}}$	minimum free Fe scavenging rate	3×10^{-5}		d^{-1}
$l_{\text{scav}}^{\text{POC}}$	slope of free Fe scavenging rate by POC	0.002		$(\text{mmolCm}^{-3})^{-1}\text{d}^{-1}$
$l_{\text{scav}}^{\text{shells}}$	slope of free Fe scavenging rate by shells	0.002		$(\text{mmol(C/Si)m}^{-3})^{-1}\text{d}^{-1}$
<i>Shell material</i>				
K_{SiO}	half-saturation constant for Si(OH)_4 uptake	5×10^{-6}	1×10^{-6}	kmolSim^{-3}
$R_{\text{Ca:P}}$	CaCO_3 :P uptake ratio	10	20	molCmolP^{-1}
$R_{\text{Si:P}}$	opal:P uptake ratio	50	25	molSimolP^{-1}
λ_{calc}	dissolution rate of calcite shells	0.3	0.075	d^{-1}
λ_{opal}	dissolution rate of opal shells	0.0025	0.01	d^{-1}
<i>Sediments</i>				
$\lambda_{\text{det}}^{\text{sed,ref}}$	sediment POC remineralisation rate at temperature $T_{\text{ref}}^{\text{sed}}$	0.025	0.01	$(\text{kmolO}_2\text{m}^{-3})^{-1}\text{d}^{-1}$
$\lambda_{\text{N}}^{\text{sed,ref}}$	sediment denitrification rate at temperature $T_{\text{ref}}^{\text{sed}}$	0.1	0.01	d^{-1}
$\lambda_{\text{S}}^{\text{sed,ref}}$	sediment sulfate reduction rate at temperature $T_{\text{ref}}^{\text{sed}}$	2.5×10^{-5}	2.5×10^{-5}	d^{-1}
Q_{10}^{sed}	Q10 for temperature dependence of remineralisation rate	2	1	
$T_{\text{ref}}^{\text{sed}}$	reference temperature for remineralisation rate in sediments	5		$^{\circ}\text{C}$
$\lambda_{\text{calc}}^{\text{sed}}$	sediment dissolution rate constant of CaCO_3	0.02	0.0086	$(\text{kmolCaCO}_3\text{m}^{-3})^{-1}\text{d}^{-1}$
$\lambda_{\text{opal}}^{\text{sed}}$	sediment dissolution rate constant of opal	0.005	0.0026	$(\text{kmolSi(OH)}_4\text{m}^{-3})^{-1}\text{d}^{-1}$

2.2 Land carbon cycle: PALADYN

135 PALADYN is a comprehensive land surface-vegetation-carbon cycle model designed specifically for the use in CLIMBER-X (Willeit and Ganopolski, 2016). It includes a detailed representation of the land carbon cycle. Photosynthesis is computed following the Farquhar model (Farquhar et al., 1980; Collatz et al., 1991) and depends on absorbed shortwave radiation, air temperature, vapor pressure deficit between leaf and ambient air, atmospheric CO₂ and soil moisture. Carbon assimilation by vegetation is coupled to the transpiration of water through stomatal conductance. The model includes a dynamic vegetation
140 module with 5 plant functional types (PFTs) competing for the gridcell share based on their respective net primary productivity. The model distinguishes between mineral soil carbon, peat carbon, buried carbon and shelf carbon. Each soil carbon 'type' has its own soil carbon pools generally represented by a litter, a fast and a slow carbon pool in each of the five soil layers. Carbon can be redistributed between the layers by vertical diffusion. For the vegetated macro surface type, decomposition is a function of soil temperature and soil moisture. Carbon in permanently frozen layers is assigned a long turnover time which
145 effectively locks carbon in permafrost. Carbon buried below ice sheets and on ocean shelves is treated separately. The land model also includes a dynamic peat module. PALADYN includes carbon isotopes ¹³C and ¹⁴C, which are tracked through all carbon pools in vegetation and soil. Isotopic discrimination is modelled only during the photosynthetic process. A simple methane module is implemented to represent methane emissions from anaerobic carbon decomposition in wetlands and peatlands. The integration of PALADYN into the coupled CLIMBER-X framework and subsequent sensitivity analyses of the land carbon
150 cycle feedbacks, which were not performed with the offline PALADYN setup in Willeit and Ganopolski (2016), highlighted the need to improve certain aspects of the model. These improvements are described next.

We have updated the parameterisation of the roughness length for heat and moisture. Originally, it was simply taken to be proportional to the roughness length for momentum, but there is ample evidence from observations that the roughness length for scalars can be orders of magnitude lower than that for momentum when the surface roughness is large (e.g. Zilitinkevich, 1995; Chen and Z
155 . We have therefore implemented the parameterisation from Zilitinkevich (1995), which includes a dependence of the surface roughness length for heat and moisture on the roughness Reynolds number. With this new parameterisation the exchange coefficient for the turbulent surface fluxes shows a much weaker dependence on the roughness of the surface, which has an impact on the vegetation feedback.

We have introduced a topographic erodibility factor for dust emissions following Ginoux et al. (2001). It assumes that a
160 basin with pronounced topographic variations contains large amount of sediments which have accumulated in the valleys and depressions and which can easily be mobilized by wind. The following topographic factor is then applied to scale dust emissions:

$$f_{\text{topo}} = \left(\frac{\max(0, z_{\text{max}} - z)}{z_{\text{max}} - z_{\text{min}}} \right)^5, \quad (1)$$

where z is the grid-cell mean elevation, and z_{max} and z_{min} are the maximum and minimum surface elevations computed from
165 the high-resolution topography in the surrounding $15 \times 15^\circ$. The exponent 5 is taken from Zender et al. (2003).

The rubisco-limited photosynthesis rate [in](#) the version of PALADYN model described in Willeit and Ganopolski (2016) was based on the 'strong optimality' hypothesis of Haxeltine and Prentice (1996), which assumes that rubisco activity and the nitrogen content of leaves vary with canopy position and seasonally so as to maximize net assimilation at the leaf level (Schaphoff et al., 2018). However, we found that this formulation led to a relatively small increase in gross primary production
 170 over the historical period, which resulted into an overestimation of atmospheric CO₂ in coupled historical simulations. We therefore introduced a new formulation for the maximum rubisco capacity, with dependencies on PFT-specific, constant foliage nitrogen concentration, specific leaf area and leaf temperature following Thornton and Zimmermann (2007) as implemented in CLM4.5 (Oleson et al., 2010).

In the original PALADYN formulation, the internal leaf CO₂ concentration used for photosynthesis was computed based
 175 on the Cowan–Farquhar optimality hypothesis (Medlyn et al., 2011). In the new model version, for C3 plants, we have implemented an alternative scheme following the more general least-cost optimality model (Prentice et al., 2014; Lavergne et al., 2019) with the moisture dependence proposed by Lavergne et al. (2020).

In the isotopic discrimination during photosynthesis (Δ) we included an explicit fractionation term for photorespiration as recommended by several recent studies (Ubierna and Farquhar, 2014; Schubert and Jahren, 2018; Lavergne et al., 2019):

$$180 \quad \Delta = 4.4 \frac{c_a - c_i}{c_a} + 27 \cdot \frac{c_i}{c_a} - 12 \frac{\Gamma_*}{p_a}, \quad (2)$$

where c_a and c_i are the ambient and leaf internal CO₂ concentrations, p_a is the ambient partial pressure of CO₂ and Γ_* is the CO₂ compensation point.

[For the distinction between evergreen and summergreen trees, in addition to a threshold on coldest month temperature we have introduced a PFT-specific threshold on the growing degree days above 5°C, which is set to 600 for needleleaf trees and
 185 900 for shrubs following Sitch et al. \(2003\).](#)

In the dynamic vegetation model a parameter (λ) is used to partition the net primary production (NPP) between local growth of existing vegetation and lateral expansion ('spreading') of vegetation coverage within the grid cell, with all of the NPP being used for growth for small leaf area index (LAI) values, and all the NPP being used for 'spreading' for large LAI values. λ is assumed to be a piecewise linear function of the leaf area index between a minimum and maximum LAI. For small leaf area
 190 indices, all of the NPP is used for local growth ($\lambda = 0$); for LAI above a critical value LAI_{\min} , a fraction ($\lambda > 0$) is used for 'spreading':

$$\lambda = \frac{LAI - LAI_{\min}}{LAI_{\max} - LAI_{\min}}. \quad (3)$$

However, since the simulated leaf area index depends strongly on NPP, which in turn has a pronounced dependence on atmospheric CO₂, this formulation results in a strong dependence of λ on CO₂, with an increasingly larger fraction of NPP being
 195 used for 'spreading' as CO₂ increases. We have therefore implemented a CO₂ dependence in the maximum leaf area index to reduce this effect:

$$LAI_{\max} = LAI_{\max}^{\text{ref}} \cdot \left(1 + 0.5 \cdot \log\left(\text{CO}_2/\text{CO}_2^{\text{ref}}\right)\right). \quad (4)$$

The fraction of decomposed litter respired directly as CO₂ to the atmosphere has been reduced from 0.7 to 0.6 and the fraction of decomposed litter transferred to the slow soil carbon pool has been doubled from 0.015 to 0.03. Together these
200 changes result in more carbon ~~being accumulated~~ accumulating into the soil.

A simple representation of land use change has been introduced into the model following Burton et al. (2019) as described in Willeit et al. (2022). A fraction of each grid cell is prescribed as being used for agriculture and land use is then represented as a limitation to the space available for the woody PFTs to expand into. When forests and shrubs are affected by land use change, an additional disturbance rate of 1 yr⁻¹ is prescribed on top of the standard background disturbance, leading to vegetation
205 dying. The resulting dead vegetation carbon is then added as litter to the soil carbon pools, and a large part will be respired directly to the atmosphere within a few years. ~~Soil Storage of wood from deforestation into products such as paper or wood for construction is not accounted for in the model and soil~~ carbon is assumed to not be directly affected by land use practices. Following deforestation, the model will grow C3 or C4 grasses, depending on climate conditions.

The partitioning of the soil carbon decomposed under anaerobic conditions into CO₂ and CH₄ ~~was used~~ a prescribed
210 constant ratio in Willeit and Ganopolski (2016). We modified this by making the fraction released as CH₄ dependent on temperature with a Q10 of 1.8, following Riley et al. (2011) and Kleinen et al. (2020).

We implemented a chemical weathering model to compute the riverine fluxes of bicarbonate ions (HCO₃⁻), (and therefore dissolved inorganic carbon and alkalinity) to the ocean and the consumption of atmospheric CO₂. The weathering rate depends on the lithology and on the climate variables temperature and runoff. The lithological map of Hartmann and Moosdorf (2012)
215 distinguishing 16 different lithologies is used to describe the spatial distribution of rocks. The parameters for the chemical weathering equations for all lithologies, except for carbonate sedimentary rocks and loess, are based on a spatially explicit runoff-dependent model of chemical weathering, which was calibrated for 381 catchments in Japan (Hartmann, 2009), with the additional temperature dependence of Hartmann et al. (2014). The effect of soil shielding on the weathering rate suggested by Hartmann et al. (2014) has not been considered since information on soil shielding is not readily available for periods
220 beyond the recent past. For carbonate sedimentary rocks, the weathering rate follows the approach of Amiotte Suchet and Probst (1995) with a dependence on runoff. Alternatively, the temperature dependent formulation of Romero-Mujalli et al. (2019) is available for use in the model. The weathering rate for loess sediments depends on runoff following Börker et al. (2020). The global distribution of loess cover for present-day and for the last glacial maximum, as well as the lithologies of the continental shelves that were exposed at the last glacial maximum, ~~is are~~ taken from Börker et al. (2020). The weathering fluxes are transferred from the land to the ocean in the same way as water runoff, following the runoff routing scheme.
225

The carbon isotopes fluxes from chemical weathering are computed assuming a δ¹³C of 1.8 permil for carbon originating from carbonate minerals (Derry and France-Lanord, 1996).

Equations describing silicate and phosphorus weathering fluxes are also available as part of the weathering model. However, silicate and phosphorus riverine fluxes are not considered in the default model setup, as they would result in further complica-
230 tions related to the conservation of nutrients in the ocean. Instead, as discussed in ~~section~~ Sect. 2.1, the silicate and phosphorus budgets are closed by assuming that the sediment burial flux is ~~prescribed~~ returned as input at the ocean surface.

2.3 Atmospheric CO₂

The atmospheric CO₂ concentration in CLIMBER-X is a globally uniform value. It can either be prescribed (as constant or time-dependent) or interactively computed by the model from the following prognostic equation for the total carbon content stored as CO₂ in the atmosphere (C_{atm}):

$$\frac{dC_{\text{atm}}}{dt} = F_{\text{ocn}} + F_{\text{land}} + F_{\text{anth}} - F_{\text{weath}} + F_{\text{volc}} + F_{\text{CH}_4\text{ox}}. \quad (5)$$

The source and sink terms on the right hand side represent, from left to right, the net sea-air carbon flux, the global net land to atmosphere carbon flux, the anthropogenic carbon emissions (excluding land-use change), the CO₂ consumption by silicate and carbonate weathering, the volcanic degassing flux and the CO₂ flux from the oxidation of atmospheric methane originating from non-agricultural sources. The CO₂ consumption by weathering is computed assuming that all carbon in the HCO₃⁻ originating from the weathering of silicate rocks ($F_{\text{HCO}_3^-}^{\text{sil}}$) comes from the atmosphere, while only half of the carbon in the HCO₃⁻ originating from the weathering of carbonate rocks and sediments ($F_{\text{HCO}_3^-}^{\text{carb}}$) comes from the atmosphere:

$$F_{\text{weath}} = F_{\text{HCO}_3^-}^{\text{sil}} + 0.5 \cdot F_{\text{HCO}_3^-}^{\text{carb}}. \quad (6)$$

The constant volcanic degassing rate is set to half the silicate weathering rate (e.g. Munhoven and François, 1994) as determined by an equilibrium spinup simulation:

$$F_{\text{volc}} = 0.5 \cdot F_{\text{HCO}_3^-}^{\text{sil}}. \quad (7)$$

The flux from the oxidation of methane, $F_{\text{CH}_4\text{ox}}$, is computed by the CH₄ model as described in [See Sect. 2.4](#) below. The atmospheric CO₂ concentration is then computed from C_{atm} using a conversion factor of 2.12 PgC ppm⁻¹ (Denman et al., 2007).

Equations similar to eq. 5 are used also for the carbon isotopes ¹³C and ¹⁴C. The prognostic equation for the stable isotope ¹³C in atmospheric CO₂ is:

$$\frac{d^{13}C_{\text{atm}}}{dt} = F_{\text{ocn}}^{13} + F_{\text{land}}^{13} + F_{\text{anth}}^{13} - F_{\text{weath}}^{13} + F_{\text{volc}}^{13}. \quad (8)$$

The ¹³C fluxes from land and ocean are explicitly computed by the land and ocean carbon cycle models as described in detail in Willeit and Ganopolski (2016) and Liu et al. (2021). The $\delta^{13}\text{C}$ of anthropogenic carbon emissions is ~~taken to be~~ [prescribed as time-dependent from historical data of Andres et al. \(2016\)](#) and the ¹³C flux from CO₂ consumption by weathering, assuming no fractionation, is simply computed as:

$$F_{\text{weath}}^{13} = F_{\text{weath}} \frac{^{13}C_{\text{atm}}}{C_{\text{atm}}}. \quad (9)$$

The ¹³C of volcanic degassing is computed assuming a $\delta^{13}\text{C}$ of -5 permil.

The prognostic equation for radiocarbon ¹⁴C in atmospheric CO₂ reads:

$$\frac{d^{14}C_{\text{atm}}}{dt} = F_{\text{ocn}}^{14} + F_{\text{land}}^{14} - F_{\text{weath}}^{14} + F_{\text{prod}}^{14} - \frac{^{14}C_{\text{atm}}}{\tau_{^{14}\text{C}}}. \quad (10)$$

Carbon sources originating from geological reservoirs, i.e. volcanic degassing, are assumed to contain no radiocarbon. Similarly, radiocarbon is assumed to be absent in anthropogenic carbon emissions from fossil fuel burning, because the age of fossils far exceeds the half-life of ^{14}C . The production rate of radiocarbon in the atmosphere ($F_{\text{prod}}^{14}\text{C}$) is prescribed in the model and the radiocarbon decay time is $\tau_{14\text{C}} = 8267$ yr.

265 2.4 Atmospheric CH_4

Similarly to CO_2 , atmospheric CH_4 is also considered to be well-mixed in the atmosphere and is therefore represented as a globally uniform value. The atmospheric CH_4 concentration can be prescribed, or it can be interactively computed by the model from:

$$\frac{d\text{CH}_4}{dt} = F_{\text{ind}}^{\text{emis}} + F_{\text{anth}}^{\text{emis}} - \frac{\text{CH}_4}{\tau_{\text{CH}_4}}. \quad (11)$$

270 Methane sources include natural emissions from wetlands and peatlands ($F_{\text{ind}}^{\text{emis}}$), which are explicitly simulated by the model as originating from anaerobic decomposition processes of carbon in soils (Willeit and Ganopolski, 2016). Other natural sources of methane are generally smaller (e.g. Saunio et al., 2020; Kleinen et al., 2020) and are neglected here for simplicity. Anthropogenic methane emissions ($F_{\text{anth}}^{\text{emis}}$) are prescribed in the model. The sink of methane from oxidation in the atmosphere is computed using a constant residence time of CH_4 , $\tau_{\text{CH}_4} = 9.5$ years, which is a reasonable first approximation at least for climate conditions ranging between the last glacial maximum and present-day (Kleinen et al., 2020; Levine et al., 2011; Hopcroft et al., 2017).

3 Closed and open carbon cycle model configurations and model spinup

Two different configurations of the carbon cycle model are available and can be chosen according to the specific needs.

The first (and simplest) setup consists of ocean, land and atmosphere carbon cycle components only. In this setup marine
 280 sediments are disabled and particulate fluxes that reach the ocean floor are completely remineralised/dissolved in the bottom ocean grid cell. Rock weathering from land is also switched off, so that the carbon exchange between ocean, land and atmosphere occurs only through air-sea fluxes and through land-atmosphere exchanges. In this setup the carbon system is closed, in the sense that there are no natural sources and sinks from and to geological reservoirs. As a response to an external climate perturbation, carbon is then simply redistributed between atmosphere, ocean and land, with the total carbon in the system being
 285 conserved. This setup is equivalent to what is used in many state-of-the-art Earth System models for climate change projections on centennial time scales (e.g. Séférian et al., 2020). The model spinup for this simple setup is straightforward and requires only to run the model to steady state with a prescribed atmospheric CO_2 concentration for $\approx 10,000$ years. The slowest time scale in this setup is given by the slow decomposition rate of organic carbon in frozen soils, which is limited to a maximum value set by default to 5000 years. The initial state for the spinup run is given by observed present-day 3D concentrations of
 290 different tracers in the ocean (Lauvset et al., 2016; Olsen et al., 2016; Garcia et al., 2013b), while the land surface is assumed to be covered by bare soil and with no carbon stored on land.

The closed carbon cycle setup is applicable to simulations of up to 1000 yr. On longer time scales, sediment and weathering processes become important and need to be accounted for when performing long-term transient simulations with interactive CO₂. Although it is unlikely that in reality the slow carbon cycle processes related to marine sediments, peatlands and permafrost carbon are in equilibrium at any specific point in time, for practical reasons we assume that such an equilibrium is a reasonable first approximation. Assuming that the preindustrial is an equilibrium state of the climate-carbon cycle system allows to run perturbation experiments with the interactive carbon cycle without having to deal with possible long-term drifts in atmospheric CO₂. However, the long time scale of ~100,000 years involved in ocean sediment processes represents a challenge in running the model into equilibrium, even for a high-throughput model like CLIMBER-X. We therefore implemented a scheme to run the physical ocean and ocean biogeochemistry models in an offline setup with prescribed climatological daily input fields at the ocean surface. This setup results in a speedup of a factor >2 relative to running the fully coupled climate-carbon cycle model, meaning that ocean carbon cycle and marine sediments can be run into equilibrium in about a week of computing time on a high performance computer. In detail, the spinup procedure of the full carbon cycle configuration comprises two different stages. Atmospheric CO₂ is prescribed to a constant value throughout the process, at 280 ppm for the pre-industrial case. The first ~~spinup~~-stage aims at spinning up the sediment model. For ~~that~~ this purpose the full carbon-cycle climate model is run for 5,000 years and every 300 years the sediment model is run ~~in-stand-alone-offline~~ for 1000 years. During this stage all net fluxes into the sediments are compensated for and ~~prescribed-as-input~~ returned as inputs at the ocean surface in order to approximately conserve water column tracer inventories while the sediments are filling up. In the second stage we switch to simulated DIC and alkalinity weathering fluxes from land and at the same time also switch to the more efficient offline ocean-biogeochemistry setup described above and run the model until an approximate equilibrium is reached after ~100,000 years (Fig. 2). A simplification that is made in the open carbon cycle setup is that organic carbon and opal that are buried into the sediments, and are therefore effectively leaving the system, are returned in remineralized form to the surface ocean so that phosphorus and silica inventories of the ocean-sediment system are conserved throughout the simulation.

The carbon fluxes among the different model components in the open setup for equilibrium pre-industrial conditions are schematically illustrated in Fig. 3. The volcanic degassing rate is equal to half the atmospheric CO₂ consumption by silicate weathering, in accordance with theory (Munhoven and François, 1994). Note that not only the carbon budget of the different compartments (atmosphere, ocean, lithosphere) is well balanced, but also the ocean alkalinity budget is.

4 Model evaluation for historical period and present day

Here we present results from a CLIMBER-X simulation with interactive CO₂ and CH₄ in the open carbon cycle setup for the historical period (1850–2015) and provide a comprehensive evaluation of model performance against various observational datasets. The forcings for this simulation include variations in solar radiation (Matthes et al., 2017), radiative forcing of volcanic eruptions (Prather et al., 2013), globally uniform N₂O concentrations from Köhler et al. (2017), globally uniform CFC11 and CFC12 concentrations from Meinshausen et al. (2016), 3D O₃ concentrations and 2D SO₄²⁻ load from the ensemble mean of CMIP6 models and land use change (pasture and cropland fractions) from Ma et al. (2020). The model is initialized from an




Figure 2. Open versus closed carbon cycle spinup. The figure shows surface input of (a) DIC and (b) alkalinity, the evolution of (c) DIC and (d) alkalinity inventories in the ocean and (e) the air-sea CO_2 flux. The grey vertical lines indicate the switch between first and second spinup phase, as described in the text.




Figure 3. CLIMBER-X carbon fluxes and reservoirs in equilibrium with pre-industrial conditions for the open carbon cycle setup.

325 80,000-year equilibrium simulation with the open carbon cycle setup for pre-industrial boundary conditions and a prescribed atmospheric CO_2 of 280 ppm, as described in section 3 and shown in Fig. 2.

4.1 Present day

In the following, different simulated climatological characteristics are compared to observations to assess the model performance for present day. Unless stated otherwise the comparison with observations is for the time interval from 1981 to 2010. To
 330 give an overview of how CLIMBER-X compares to state-of-the-art Earth system models based on general circulation models, we also include results from model simulations from the recent coupled model intercomparison project CMIP6 (Eyring et al., 2016). ~~In particular, for~~ The following CMIP6 models are included for ocean biogeochemistry: CESM2, IPSL-CM6A-LR, MRI-ESM2-0, MIROC-ES2L, MPI-ESM1-2-LR, UKESM1-0-LL and CanESM5. For the land carbon cycle, the following models are used for comparison: ACCESS-ESM1-5, BCC-CSM2-MR, CanESM5, CNRM-ESM2-1, GFDL-ESM4, IPSL-CM6A-LR,
 335 MIROC-ES2L, MPI-ESM1-2-LR, MRI-ESM2-0, NorESM2-LM, UKESM1-0-LL. For ocean biogeochemistry, we highlight how the model compares with results from the MPI-ESM1-2-LR employing the original marine carbon cycle model HAMOCC6.

4.1.1 Ocean biogeochemistry and marine sediments

An overview of simulated global variables characterising the ocean carbon cycle are presented and compared to observation-based estimates in Table 2, providing a summary of model performance for the present day.

340 Representing ocean ventilation time scale reasonably well is a prerequisite for simulating biogeochemical tracers in the ocean. The ocean uptake of CFCs of anthropogenic origin over the historical period is often used to probe the ventilation

Table 2. Ocean biogeochemistry [Global values of the main ocean biogeochemical variables for the present-day.](#)

	CLIMBER-X	Estimated range	Unit	
<i>Ocean-Atmosphere Fluxes</i>				
Pre-industrial CO ₂ flux	0.2	0.2–0.6	PgCyr ⁻¹	Jacobson et al. (2007);Regnier
N ₂ O flux	5.0	1.9–9.4	TgNyr ⁻¹	Buitenhuis
<i>Surface Nutrients and Alkalinity</i>				
Surface alkalinity	2324 <u>2410</u>	2250–2300 <u>2355</u>	mmolm ⁻³	GLODAPv2, (Lauvset et al., 2016; Olsen
Surface nitrate	6.15 <u>6.3</u>	5.0 <u>5.2</u>	mmolNm ⁻³	WOA 2013, Garcia e
Surface phosphate	0.46 0.51	<u>0.53</u>	mmolPm ⁻³	WOA 2013, Garcia e
Surface silicate	12.8 <u>8.2</u>	8 <u>7.5</u>	mmolSim ⁻³	WOA 2013, Garcia e
<i>Primary Production</i>				
Net primary production	53	47-60	PgCyr ⁻¹	Johnson and Bif (2021);Carr
N-fixation	122 <u>88</u>	51-200	TgNyr ⁻¹	Karl et al. (2002);Großkopf
<i>Export production</i>				
POC export at 100 m	6.0 <u>5.9</u>	5.8–12.9	PgCyr ⁻¹	Dunne
CaCO ₃ export at 100 m	0.65 <u>0.62</u>	0.38–1.8	PgCyr ⁻¹	Dunne
Opal export at 100 m	113.7 <u>105</u>	94.5–155.5	TmolSiy ⁻¹	Tréguer and De La F
<i>Sediments</i>				
POC sediment deposition	0.55 <u>0.57</u>	0.93–3.2	PgCyr ⁻¹	Dunne
CaCO ₃ sediment deposition	0.26 <u>0.27</u>	0.16–0.4	PgCyr ⁻¹	Battaglia et al. (2016); Milliman and Dr
Opal sediment deposition	56 <u>79</u>	22–79 <u>79–84</u>	PgCyr ⁻¹	Tréguer and De La Rocha (2013); Dunne et al. (2007) Tréguer
POC burial	0.23 <u>0.09</u>	0.07–0.7	PgCyr ⁻¹	Cartapanis
CaCO ₃ burial	0.2 <u>0.19</u>	0.13–0.45	PgCyr ⁻¹	Cartapanis
Opal burial	6.3 <u>5.3</u>	2.7–9.9	TmolSiy ⁻¹	Tréguer and De La F

of the ocean on decadal time scales, while the pre-industrial radiocarbon concentration in the ocean provides information on the age distribution of the water masses in an approximate equilibrium state. We therefore start by comparing how well the model reproduces the CFC11 and radiocarbon distributions in the ocean. The inventory of CFC11 in the ocean starts to increase after ≈ 1950 , as a consequence of its increase in the atmosphere (Fig. 4). Estimates for CFC11 inventory in the year ≈ 1994 are available from models from the OCMIP model intercomparison (Dutay et al., 2002) and from direct observations (Willey et al., 2004). CLIMBER-X results are generally consistent with these estimates (Fig. 4), indicating that, at least at the global scale, the decadal ventilation time scale in CLIMBER-X is well in line with observations and other models. In terms of spatial distribution, the CFC11 uptake is overestimated in the North Pacific, the northern Indian Ocean and around Antarctica, while too small CFC11 concentrations are simulated at mid-latitudes in all basins at depths between 500 and 1000 m (Fig. 5).




Figure 4. Historical global cumulative ocean uptake of CFC11 in CLIMBER-X compared to observations (Willey et al., 2004) and OCMIP models (Dutay et al., 2002).




Figure 5. Zonally averaged CFC11 concentration in CLIMBER-X (left column) and GLODAP (Key et al., 2004) (middle column) for different basins: Atlantic (top), Pacific (middle) and Southern Ocean (bottom). The model bias is shown in the right column.




Figure 6. Zonally averaged pre-industrial radiocarbon ventilation age in CLIMBER-X (left column) and GLODAP Key et al. (2004) (middle column) for different basins: Atlantic (top), Pacific (middle) and Southern Ocean (bottom). The model bias is shown in the right column.

The radiocarbon ventilation age in the pre-industrial gives additional insights into the ocean ventilation in quasi-equilibrium conditions, an information which is complementary to CFC11. The radiocarbon ventilation age of the deep ocean is nicely reproduced by CLIMBER-X, while radiocarbon age is systematically overestimated in the upper kilometer across all ocean basins (Fig. 6). The too old (in terms of radiocarbon age) sub-surface waters could be a result of the model not explicitly resolving synoptic processes in the atmosphere and therefore not representing the non-linear effects of synoptic variability on vertical mixing of tracers. For instance, a one-time mixing down to 200 m depth by a wind storm could have a large effect on some tracers, which cannot be resolved by using climatological mean winds. We would expect this non-linear effect to be much more important for radiocarbon than for nutrients. The analyses of CFC11 and radiocarbon provide important insights on the ocean ventilation in the model and will be useful when discussing model biases in the distribution of other biogeochemical tracers below.

The spatial pattern of the air-sea CO_2 exchange is well captured by the model (Fig. 7), with outgassing generally taking place in the tropics and CO_2 being taken up in mid- to high northern latitudes and in mid-latitudes of the Southern Hemisphere. The main difference compared to other models is observed around the Equator, with a less pronounced peak in CO_2 release simulated by CLIMBER-X (Fig. 8), which is likely related to deficiencies in the simulated ocean circulation close to the equator, where the geostrophic approximation employed in CLIMBER-X reaches its limit of applicability. In the Southern Ocean, most CMIP6 models tend to overestimate the CO_2 uptake compared to observations (e.g. Gruber et al., 2009) (Fig. 8),




Figure 7. Air–sea CO₂ flux in (a) CLIMBER-X compared to (b) observations from Landschützer et al. (2016).

while CLIMBER-X is apparently more consistent with recent estimates, although with substantial differences in the spatial distribution of the CO₂ flux (Fig. 7). Notably, in the Southern Ocean the CLIMBER-X air-sea CO₂ exchange diverges from that simulated by the MPI-ESM1-2-LR model (Fig. 8), which employs the original HAMOCC6 ocean biogeochemistry model. This is possibly related to the lower simulated net primary production in the Southern Ocean in CLIMBER-X compared to MPI-ESM (Fig. 9a). However, the MPI-ESM seems to be an outlier in the simulated primary production in the Southern Ocean, possibly because of biases in climate, which are unrelated to the HAMOCC ocean carbon cycle model.

The export of particulate organic carbon from the euphotic layer drives the biological pump and generally follows the primary productivity pattern, with modifications due to varying sinking speeds and remineralisation rates of POC in the water column. While the net primary productivity in CLIMBER-X is in line with CMIP6 models (Fig. 9a) and the globally integrated value of 55 PgCyr⁻¹ agrees well with observations (Fab. Table 2), the export production in the model is generally at the lower end of the CMIP6 model range (Fig. 9b). CaCO₃ and opal export are compared to CMIP6 models in Fig. 9c,d.

Primary production in the ocean is limited by the availability of nutrients. Over large parts of the surface ocean nitrogen concentrations constitute the main limiting factor for photosynthesis in CLIMBER-X (Fig. 10). However, over the Southern Ocean, in the equatorial Pacific and in the North Pacific production is limited by the availability of iron (Fig. 10). This is in accordance with observations showing that iron limitation is usually important where subsurface nutrient supply is enhanced, such as in oceanic upwelling regions (e.g. Moore et al., 2013). Since one of the main iron sources in the ocean is from mineral




Figure 8. Zonal mean air-sea CO_2 flux in CLIMBER-X compared to observations from Landschuetzer et al. (2016) and selected CMIP6 models.




Figure 9. ~~Zonal~~ Global zonal mean (a) net primary production, (b) particulate organic carbon export at 100 m depth, (c) CaCO_3 and (d) opal export at 100 m depth. Results from CLIMBER-X are compared to CMIP6 models.




Figure 10. Nutrient limitation of marine net primary productivity in CLIMBER-X.




Figure 11. (a) CLIMBER-X annual dust deposition flux compared to model based products of (b) Albani et al. (2016), (c) Lambert et al. (2015) and (d) Hopcroft et al. (2015). The respective globally integrated deposition values are given in brackets in the panel titles.

dust deposited at the ocean surface (e.g. Tagliabue et al., 2016), iron limitation is confined to regions with low dust deposition. The dust cycle is an integral part of CLIMBER-X, and the dust deposition is therefore explicitly modelled. The simulated
 385 dust deposition compares reasonably well with estimates from complex ESMs for the present-day (Fig. 11), although they are relatively poorly constrained. A comparison of dust deposition fluxes with observations over land further indicates that the model is able to capture the general pattern of dust deposition rate (Fig. 12).

The simulated dissolved iron concentration in surface water is closely related to the dust deposition shown in Fig. 11. It is therefore high in the Atlantic and Indian oceans, lower in the Southern Ocean and very small over large parts of the Pacific
 390 (Fig. 13). This is broadly consistent with observations (e.g. Tagliabue et al., 2012), but measurements of iron concentration in ocean water are still relatively sparse. ~~The surface iron concentration in CLIMBER-X is systematically higher than in CMIP6 models at mid-latitudes in the Southern Hemisphere, which is a result of excessive dust emissions from subtropical SH land (Fig. 11).~~




Figure 12. Simulated versus observed dust deposition fluxes at different locations available from the AeroCom dataset (Huneus et al. (2011) and references therein). The dashed lines indicate one order of magnitude deviation from the 1:1 line.




Figure 13. Surface dissolved iron concentration in CLIMBER-X.




Figure 14. Zonal-Global zonal mean surface concentrations of the nutrients (a) nitrate, (b) phosphate, (c) silicate and (d) dissolved iron. CLIMBER-X is compared to observations (Garcia et al., 2013b) and CMIP6 model results.

The main features of the surface nitrate concentration are well reproduced by CLIMBER-X, with large concentrations in the Southern Ocean, moderate values in the upwelling region of the Eastern equatorial Pacific and in the North Atlantic and North Pacific and low values elsewhere (Fig. 15, Fig. 14a). The most pronounced model biases are found in too high nitrate concentrations in the Arctic and too low values in the North Pacific. The simulated basin-wide vertical distribution of nitrate is in very good agreement with observations (Fig. 16). ~~The simulated basin-wide vertical distribution of different ocean tracers compares well with observations (Fig. 16), and generally lies within the relatively large range of values returned by CMIP6 models.~~

The 3D phosphate distribution in the global ocean is nicely captured by the model (Fig. 17, Fig. 14b, Fig. 16), except for too low concentrations simulated in the surface ocean of the North Pacific and the northern Indian Ocean. The negative bias in the North Pacific is consistent with the too low simulated surface nitrate concentrations, both originating from a too vigorous ventilation of water masses in the upper kilometer in the physical ocean model.

As a result of reduced primary productivity in the Southern Ocean in CLIMBER-X compared to MPI-ESM1-2-LR, both surface nitrate and phosphate concentrations are consistently higher in CLIMBER-X (Fig. 14a,b), as less nutrients are assimilated during photosynthesis.

Silicate concentration is generally overestimated in the sub-surface ocean and underestimated in the deep North Pacific and North Indian oceans (Fig. 18). ~~A positive surface silicate concentration bias in the Southern Ocean is seen in most CMIP6 models, similarly to other nutrients~~ (Fig. 14c,17).

The large scale patterns of oxygen concentration in ocean waters simulated by CLIMBER-X is largely consistent with observations (Fig. 19), but the extent and depth of the oxygen minimum zones, in particular in the Eastern equatorial Pacific,




Figure 15. Surface NO_3 concentration in (a) CLIMBER-X compared to (b) observations from the World Ocean Atlas 2013 (WOA13, Garcia et al. (2013b)). The model bias is shown in (c).

is overestimated. This bias is common to many ~~EMP~~CMIP5 models (e.g. Cabré et al., 2015). Other biases include a too oxygen depleted Southern Ocean and too high oxygen concentrations in the upper North Pacific and North Indian oceans, again resulting from the excessive water mass ventilation in those regions as discussed above.

Both DIC and alkalinity are generally overestimated in the upper ocean (Fig. 16), particularly in Antarctic intermediate water masses, and underestimated in the deep ocean (Fig. 20,21). ~~The~~ These biases in the simulated vertical distribution of DIC and alkalinity could be due to a relatively low CaCO_3 export from the euphotic layer (Table 2), which leads to a too weak vertical redistribution. Additionally, the simulated DIC concentration is generally too low in the ~~upper~~ North Pacific and ~~Indian~~ Oceans, while alkalinity is underestimated in the deep Pacific/northern Indian Oceans.

Carbon isotopes ~~The carbon 13 isotope~~ in the ocean ~~help~~helps to track the distribution of different water masses. The higher $\delta^{13}\text{C}$ values in the Atlantic compared to the Pacific Ocean, originating mainly from the pronounced overturning circulation in




Figure 16. Global and basin-wide average profiles of different biogeochemical tracers in the ocean, from top to bottom: DIC, alkalinity, nitrate, phosphate, silicate and dissolved iron. CLIMBER-X results (black) are compared to observations (blue) (Lauvset et al., 2016; Olsen et al., 2016; Garcia et al., 2013a,b) and CMIP6 model results (grey). Results from the MPI-ESM1-2-LR are shown by the green dashed lines.

The boundary of the Southern Ocean is set at 35°S and the Southern Ocean section is not included in the profiles of the Atlantic, Pacific and Indian ocean.




Figure 17. Zonally averaged PO_4 concentration in CLIMBER-X (left column) and WOA13 (Garcia et al., 2013b) (middle column) for different basins: Atlantic (top), Pacific (middle) and Southern and Indian Oceans (bottom). The model bias is shown in the right column.

the Atlantic, which is absent in the Pacific, are generally captured by the model (Fig. 22). The main bias is a too low $\delta^{13}\text{C}$ associated with Antarctic intermediate waters and too high $\delta^{13}\text{C}$ values in the upper North Pacific and Indian oceans. negative biases at 500-1500 m depth are associated with the 'nutrient trapping' problem (Aumont et al., 1999; Dietze and Loeptien, 2013) that is often seen in ESMs. This problem is characterised by high concentrations of remineralised nutrients and carbon and, therefore, low $\delta^{13}\text{C}$ is also overestimated at all depths in the Arctic. The radiocarbon ventilation age of the deep ocean is nicely reproduced by CLIMBER-X, while radiocarbon age is systematically overestimated in the upper kilometer across all ocean basins (Fig. 6). The too old (in terms of radiocarbon age) sub-surface waters could be a result of the model not explicitly resolving synoptic processes in the atmosphere, leading to an underestimation of the wind-driven mixing of passive carbon tracers in the upper ocean. C (Liu et al., 2021). The positive biases through the whole water column in the North Atlantic, North Pacific and northern Indian Ocean are possibly the result from too strong ventilation in these regions in the model.

Zonally averaged radiocarbon ventilation age in CLIMBER-X (left column) and GLODAP Key et al. (2004) (middle column) for different basins: Atlantic (top), Pacific (middle) and Southern and Indian Oceans (bottom). The model bias is shown in the right column.

In the Atlantic and Indian Ocean CaCO_3 dominates the sediment composition, in accordance with observations (Fig. 23a,d). However, little CaCO_3 is simulated in large parts of the sediment in the eastern Pacific Ocean, where observations indi-




Figure 18. Zonally averaged Si concentration in CLIMBER-X (left column) and WOA13 (Garcia et al., 2013b) (middle column) for different basins: Atlantic (top), Pacific (middle) and Southern and Indian Oceans (bottom). The model bias is shown in the right column.




Figure 19. Oxygen concentration in CLIMBER-X (left column) and WOA13 (Garcia et al., 2013a) (middle column) at different ocean depths: from top to bottom, 100 m, 500 m, 1000 m and 3000 m. The model bias is shown in the right column.




Figure 20. Zonally averaged dissolved inorganic carbon in CLIMBER-X (left column) and GLODAPv2 (Lauvset et al., 2016; Olsen et al., 2016) (middle column) for different basins: Atlantic (top), Pacific (middle) and Southern and Indian Oceans (bottom). The model bias is shown in the right column.

cate widespread CaCO_3 content in the Southern Hemisphere (Fig. 23a,d). The underestimation of calcite weight fractions in sediments of the eastern South Pacific Ocean is caused by water being undersaturated with respect to calcite in this area.
 440 This leads to dissolution of most of the calcite produced at the surface before it can even reach the sediments. The strongly undersaturated water is ultimately a result of deficiencies in the simulated ocean circulation. Some other models show similar deficiencies in the simulated calcite fraction in Pacific sediments (e.g. Kurahashi-Nakamura et al., 2022). Global CaCO_3 sediment deposition and burial are in line with observational underestimates (Tab. Table 2), with around 25% of the deposited CaCO_3 undergoing dissolution. The opal content in sediments in CLIMBER-X is ~~substantially~~ overestimated (Fig. 23b,e),
 445 even though the global opal sedimentation and burial fluxes are fully consistent with observational estimates (Tab. Table 2). Opal is particularly abundant in the eastern equatorial Pacific, simply as a result of missing CaCO_3 in the sediments in that area. Organic carbon is found mainly on the continental margins and in the equatorial east Pacific, in agreement with observations (Fig. 23c,f), although CLIMBER-X tends to underestimate the organic carbon content in sediments, possibly because of a too small sediment deposition flux of POC (Tab. Table 2).




Figure 21. Zonally averaged total alkalinity in CLIMBER-X (left column) and GLODAPv2 (Lauvset et al., 2016; Olsen et al., 2016) (middle column) for different basins: Atlantic (top), Pacific (middle) and Southern and Indian Oceans (bottom). The model bias is shown in the right column.

450 4.1.2 Land carbon cycle

A detailed evaluation of the land carbon cycle component has already been presented in the original PALADYN description paper (Willeit and Ganopolski, 2016). However, here we partly repeat the analysis to show the model performance in the coupled climate model setup and with the additional modifications to the model described above.

A selection of simulated global variables characterising the land carbon cycle is presented and compared to observation-
455 based estimates in Table 3, providing a summary of model performance for the present day.

Photosynthesis is the basic process by which carbon enters the land domain. The simulated gross primary production (GPP), which quantifies this process, is in good agreement with observational estimates, both in terms of global integral (Tab:Table 3) and in terms of spatial distribution (Fig. 24a,b,c).

The total carbon stored in the vegetation, both above ground and below ground, is slightly overestimated in the model
460 (Tab:Table 3), but the meridional distribution, mainly originating from large-scale differences in precipitation, is well reproduced (Fig. 24d,e,f). Most of soil carbon in CLIMBER-X is stored in cold soils of the NH high-latitudes, in agreement with observations (Fig. 24g,h,i). However, compared to estimates from Carvalhais et al. (2014) the soil carbon distribution is too skewed towards high northern latitudes and there is too little carbon in the tropics. Most CMIP6 models underestimate soil carbon in the tropics as well (Fig. 24j).




Figure 22. Zonally averaged $\delta^{13}\text{C}$ in CLIMBER-X (left column) and Eide et al. (2017) (middle column) for different basins: Atlantic (top), Pacific (middle) and Southern ~~and Indian Oceans~~ Ocean (bottom). The model bias is shown in the right column.




Figure 23. Weight fraction of calcite, opal and organic carbon in marine sediments as simulated by CLIMBER-X (top), compared to observations (~~Archer, 1996; Seiter et al., 2004~~) (Hayes et al., 2021) (bottom).

Table 3. Land-Global values for the main variables of the land carbon cycle.

	CLIMBER-X	Estimated range	Unit	Source
<i>Primary production</i>				
Gross primary production	120	115–131	PgCyr ⁻¹	Beer et al. (2010)
Net primary production	67	42–70	PgCyr ⁻¹	Ito (2011)
<i>Land carbon pools</i>				
Vegetation carbon	490 <u>472</u>	392–437	PgC	Fan et al. (2020)
Soil carbon	2187 <u>2145</u>	3300–4800	PgC	Fan et al. (2020)
Soil carbon top 1 m	1532 <u>1521</u>	1200–2000	PgC	Varney et al. (2022)
Soil carbon top 1 m 60–90°N	441 <u>436</u>	314–526	PgC	Varney et al. (2022)
Permafrost area	19.8 <u>19.1</u>	18.7	10 ⁶ × km ²	Brown et al. (1998); Tarnocai et al. (2009)
Carbon in permafrost area	861 <u>796</u>	1100–1500	PgC	Hugelius et al. (2014)
Peatland area	2.5 <u>2.4</u>	4.4	10 ⁶ × km ²	Yu et al. (2010)
Carbon in peatlands	347 <u>340</u>	530–694	PgC	Yu et al. (2010)
<i>CH₄</i>				
Maximum monthly wetland area	5	5.1	10 ⁶ × km ²	Prigent et al. (2007); Papa et al. (2010)
Total CH ₄ emissions	214	100–217	TgCH ₄ yr ⁻¹	Saunois et al. (2020)
Tropical CH ₄ emissions	182	71–155	TgCH ₄ yr ⁻¹	Saunois et al. (2020)
Extratropical CH ₄ emissions	33	12–64	TgCH ₄ yr ⁻¹	Saunois et al. (2020)
<i>Weathering (pre-industrial)</i>				
CO ₂ consumption	21.4 <u>22.6</u>	17–27	TmolCyr ⁻¹	Munhoven (2002)
Carbonate weathering	25.0 <u>20.1</u>	10–25.4	TmolCyr ⁻¹	Munhoven (2002)
Silicate weathering	8.9 <u>12.6</u>	10.8–19.7	TmolCyr ⁻¹	Munhoven (2002)
Alkalinity flux to ocean	33.9 <u>32.7</u>	30–40	Tmolyr ⁻¹	Amiotte Suchet et al. (2003); Gaillardet et al. (1999)

465 ~~Total soil carbon estimates vary considerably, in particular when considering the full soil depth (Fan et al., 2020, e.g.).~~ In
CLIMBER-X, ~~three quarters (~1500 PgC) of the total ~2200 in soils of carbon~~ are stored in the top soil meter ~~,~~ in good
agreement with different estimates (~~Tab. Table 3~~). ~~The~~ ~~However, with ~2150 PgC, the total soil carbon content seems to be~~
~~underestimated compared to observations, which suggest >3000 PgC. This indicates that too little carbon is simulated in~~
~~soil below 1 m depth. However, total soil carbon content estimates vary widely between datasets (e.g. Fan et al., 2020), with~~
470 ~~e.g. 1952±198 PgC in WISE30sec (Batjes, 2016) and 3141±893 PgC in Sanderman et al. (2017) in the top 2 m of soil. Most~~
~~of carbon in mineral soil layers below one meter is recalcitrant and its response to changes in environmental conditions is~~
~~uncertain. In Earth System models, total soil carbon storage is usually much lower (1206±445 PgC, Varney et al. (2022)) as~~
~~these models account for active carbon responding on centennial time scale. In CLIMBER-X, one possible explanation for~~

underestimated carbon content in deeper soil layers is that the maximum turnover time scale of soil carbon is set to 5000
475 years in the model, which limits the amount of carbon that can be accumulated in cold, frozen soil layers. Other possible
reasons include: (i) a general underestimation of vertical carbon transport by diffusion, particularly into perennally frozen
soil layers, (ii) a possible depth dependence of soil carbon turnover due to processes other than temperature and moisture
(e.g. Koven et al., 2013) and that are not included in the model. Consistently, the carbon contained in areas affected by per-
mafrost is ~ 860800 PgC, which is also a bit lower than the $\sim 1100\text{--}1500$ PgC suggested by observations (Tab. Table 3). The
480 Let us note that even when models are initialized with the observed permafrost carbon stock of ≈ 1300 PgC, remapping on
model resolution and accounting for differences in soil temperatures between models and observations generally leads to a
reduction of permafrost carbon stocks (e.g. Kleinen and Brovkin, 2018). The CLIMBER-X simulated peatland extent is lower
than estimated (Yu et al., 2010), and consistently also the peat carbon is underestimated accordingly (Tab. Table 3).

The turnover time of terrestrial ecosystem carbon is an integrated quantitative measure of the residence time of carbon on
485 land, from the time it is fixed by photosynthesis to the time it is returned to the atmosphere through respiration processes. It
is computed as the ratio between land carbon stocks (vegetation+soil) and gross primary production. The ecosystem carbon
turnover time simulated by CLIMBER-X is in line with CMIP6 models, while it is underestimated compared to observation-
based estimates from Fan et al. (2020) (Fig. 24j,k,l). However, it should be noted that the large uncertainties in soil carbon
content result in a rather uncertain estimated ecosystem carbon turnover time (Fan et al., 2020).

490 The global maximum monthly wetland extent in CLIMBER-X agrees well with observations (Tab. Table 3), although with
substantial differences in the geographic distribution (Fig. 25). Compared to the multi-satellite product from GIEMS (Global
Inundation Extent from Multi-Satellites) (Prigent et al., 2007; Papa et al., 2010) the model simulates larger wetland extent
in tropical forest areas. However, if compared to other wetland products based on data other than from satellite, GIEMS is
underestimating wetlands below dense forests (e.g. the Amazon forest) (e.g. Melack and Hess, 2010) In south-east Asia, the
495 GIEMS wetland extent also includes extensive rice cultivation areas, which are not represented in the model.

In CLIMBER-X methane is emitted exclusively from wetlands. However, because of the dependence of methane emissions
on soil carbon decomposition rates and because of the temperature dependence of the fraction of wetland carbon respired as
methane, wetland methane emissions are dominated by tropical sources (Tab. Table 3, Fig. 26), in agreement with observations
(e.g. Saunio et al., 2020). The total CH_4 emissions from wetlands are at the high end of recent estimates, which is a result of
500 tuning the emissions in the model to match the observed emissions from all natural sources.

Chemical weathering fluxes are generally high where runoff is high, with the separation between silicate and carbonate
weathering being modulated by lithological properties (Fig. 27). The global CO_2 consumption rate by weathering and the
alkalinity flux to the ocean in form of bicarbonate produced by rock weathering are in good agreement with observational esti-
mates (Tab. Table 3), while the partitioning between carbonate and silicate weathering is skewed toward carbonate weathering
505 (Tab. Table 3).




Figure 24. (a) Simulated GPP compared to (b) observations (Jung et al., 2011). (c) Comparison of zonally integrated GPP. (d) Simulated vegetation carbon compared to (e) observations (Carvalhais et al., 2014). (f) Comparison of zonally integrated vegetation carbon. (g) Simulated soil carbon compared to (h) observations (Carvalhais et al., 2014). (i) Comparison of zonally integrated soil carbon. (j) Simulated ecosystem carbon turnover time compared to (k) observations (Fan et al., 2020). (l) Comparison of zonal mean ecosystem carbon turnover time. In panels (c), (f), (i) and (l) results from CLIMBER-X are shown in black, observations in blue and CMIP6 models in grey.

4.2 Historical period

As shown by Willeit et al. (2022), the historical climate evolution is well simulated by CLIMBER-X. Here we extend this analysis by focusing on the carbon cycle response.

The historical atmospheric CO_2 concentration is well reproduced by the model, with CO_2 at the year 2015 being within a few ~ 5 ppm of direct measurements (Fig. 28). Biases in simulated CO_2 of ~ 10 ppm are quite common in state-of-the-art ESMs (e.g. Hoffman et al., 2014; Friedlingstein et al., 2014).

The partitioning of the anthropogenic carbon emitted over the historical period among the different spheres is compared with recent estimates of the Global Carbon Budget (GCB) (Friedlingstein et al., 2022) by the Global Carbon Project in Fig. 29. The amount of fossil carbon emitted from anthropogenic activities is prescribed from empirical data and therefore by definition




Figure 25. Maximum monthly wetland fraction (a) in CLIMBER-X compared to (b) the GIEMS dataset (Papa et al., 2010; Prigent et al., 2007).




Figure 26. Natural methane emission simulated by CLIMBER-X for the present-day.




Figure 27. CLIMBER-X (a) silicate and (b) carbonate weathering flux distribution for the present-day.




Figure 28. Historical atmospheric CO_2 concentration from a coupled CLIMBER-X simulation compared to observations (Köhler et al., 2017).




Figure 29. Historical global carbon budget in CLIMBER-X. The dashed lines are estimates from Friedlingstein et al. (2022).

515 matches with estimates from Friedlingstein et al. (2022). The carbon emissions resulting from land use change practices are underestimated in CLIMBER-X compared to the GCB, although the actual values remain uncertain (e.g. Gasser et al., 2020). A substantial fraction of this anthropogenic CO₂ emission is absorbed by the ocean and the land, while the rest remains in the atmosphere. In CLIMBER-X, the ocean carbon uptake is a bit lower and the land carbon uptake a bit higher than GCB estimates, but the net effect is a realistic airborne fraction of carbon remaining in the atmosphere. The ocean carbon uptake

520 is driven by the chemical disequilibrium between surface air CO₂ concentrations and the concentration of dissolved CO₂ in the surface ocean water and is relatively well understood, as also indicated by the narrow uncertainty range obtained from different CMIP6 models (Fig. 30a). The CLIMBER-X ocean carbon uptake falls within this narrow range although it tends to be at the lower end. The land carbon uptake is largely driven by an increase in gross primary productivity as a response to increasing atmospheric CO₂. The net primary productivity increase simulated by CLIMBER-X over the historical period is

525 in agreement with what is shown by most CMIP6 models (Fig. 30b). However, the effect of this NPP increase on vegetation carbon varies widely among models (Fig. 30c), also because of the confounding factor of land use change. In CLIMBER-X the net effect is a vegetation carbon stock decrease by ~50 PgC. The historical evolution of soil carbon additionally depends on the response of microbial decomposition to changing environmental conditions, particularly soil temperatures. The increasing

530 NPP, and consequently larger input of litter carbon into the soil, dominates over the negative effect of increasing temperatures in CLIMBER-X, leading to an increase in soil carbon by ~50 PgC (Fig. 30d).

Since CLIMBER-X is enabled with carbon isotopes, it also allows for a comparison of isotopic signatures to observations, thereby providing additional constraints on processes involved in carbon cycle exchanges. As an example, the historical $\delta^{13}C$ of atmospheric CO₂ is compared to observations in Fig. 31.

The general historical trend in atmospheric CH₄ is captured by the model (Fig. 32a). Prescribed anthropogenic methane emissions are the dominant source for the increase of the atmospheric methane burden, but natural emissions from land are also increasing due to the increase in NPP and soil temperature (Fig. 32b).

535




Figure 30. Historical anomalies of (a) air-sea CO₂ flux, (b) net primary production on land, (c) vegetation carbon and (d) soil carbon in CLIMBER-X compared to CMIP6 models. The anomalies are computed relative to the time interval 1850–1880 CE.




Figure 31. Historical δ¹³C of atmospheric CO₂ in CLIMBER-X compared with observations (Eggleston et al., 2016).




Figure 32. Historical (a) atmospheric CH₄ concentration in CLIMBER-X compared to observations (Köhler et al., 2017) and (b) prescribed anthropogenic CH₄ emissions and natural land emissions as simulated in CLIMBER-X.

5 Carbon cycle feedbacks

Carbon cycle processes both on land and in the ocean are sensitive to changes in climate and atmospheric CO₂. This implies that carbon fluxes between ocean and atmosphere and between land and atmosphere will respond to changes in climate and CO₂ concentration, which will in turn affect CO₂ and consequently climate. Quantifying the strength of these feedbacks is important to understand how the climate will respond to anthropogenic CO₂ emissions. For that, a linear feedback decomposition analysis has been proposed by Friedlingstein et al. (2006) to estimate these feedbacks in Earth system models. The analysis relies on a set of model simulations under idealized 1 %yr⁻¹ CO₂ increase experiments, whereby in one simulation the CO₂ increase is seen only by the radiative code in the atmosphere (radiatively coupled), in another one the CO₂ increase is seen only by the carbon cycle (biogeochemically coupled) and in a final one both radiation and carbon cycle see the increase in atmospheric CO₂ (fully coupled). This set of simulations allows to roughly separate the effect of changes in climate and changes in CO₂ on land and ocean carbon fluxes. To a first approximation, the carbon cycle feedback to climate is usually quantified using global temperature changes, while in reality climate obviously influences the carbon cycle in more complex ways than just through (global) temperature.

The carbon cycle feedback parameters have been estimated for ~~several phases of the coupled model intercomparison projects C4MIP, CMIP5 and CMIP6 models~~ (Friedlingstein et al., 2006; Arora et al., 2013, 2020). In ~~Figure~~Fig. 33 the carbon cycle–climate (γ) and the carbon cycle–concentration (β) feedbacks in CLIMBER-X are compared to CMIP6 model results separately for land and ocean. An increase in CO₂ has a positive effect on the uptake of carbon by both land and ocean, resulting in a lowering of CO₂ and therefore a negative feedback on climate (implying positive β , Fig.33a,c). Conversely, a climate warming has a generally negative impact on the ability of ocean and land to store carbon, leading to a positive feedback loop (implying negative γ , Fig.33b,d). The β and γ ~~are generally in good agreement with~~ values obtained here are well within the range of estimates from CMIP6 models (Arora et al., 2020), although in CLIMBER-X the CO₂ fertilisation effect on land is rather high (Fig.33a) and the ocean carbon uptake as a response to an increase in atmospheric CO₂ is at the lower end of the CMIP6 models (Fig.33c).

560 6 The zero emissions commitment

The zero emissions commitment (ZEC) is a measure of the amount of additional future temperature change following a complete cessation of CO₂ emissions (e.g. Matthews and Solomon, 2013). A model intercomparison project has been established in an effort to analyze and compare the ZEC in different Earth system models (Jones et al., 2019). Here we use this standardized and idealized experimental setup to compare the carbon cycle response in CLIMBER-X with results from the ZECMIP models for the 1000 PgC emission pulse (MacDougall et al., 2020). The experiment branches off from a 1 % per year CO₂ increase run with CO₂ emissions set to zero at the point of 1000 PgC of total carbon emissions. We performed this experiment with both the open and closed carbon cycle setups.

The results of the CLIMBER-X simulations are generally well within the large range of results from state-of-the-art ESMs and EMICs participating in ZECMIP (Fig. 34). Atmospheric CO₂ concentration is rapidly decreasing after stopping the carbon




Figure 33. Carbon cycle feedbacks in CLIMBER-X compared to CMIP6 models (Arora et al., 2020). The (a,c) β and (b,d) γ parameters are shown at the time of CO_2 doubling (year ~ 70) and at the time of CO_2 quadrupling (year ~ 140) in idealized $1\% \text{yr}^{-1}$ CO_2 increase experiments.

570 emissions (Fig. 34b), while global temperature shows a more modest decrease (Fig. 34a). The ocean continues to take up carbon throughout the simulation (Fig. 34c), while the land turns from a sink to a source of carbon roughly at the time of the peak in CO_2 (Fig. 34d). CLIMBER-X initially shows a relatively weak ocean carbon uptake compared to ZECMIP models, while the land carbon uptake is at the high end of the ZECMIP model ensemble.

The difference between the experiments with open and closed carbon cycle setup are negligible for the first few centuries
 575 but continue to grow over time, with CO_2 decreasing faster in the open setup (Fig. 34b).

7 Conclusions

We have described the major features of the carbon-cycle components of the newly developed CLIMBER-X Earth System model. The model includes a detailed representation of carbon cycle processes on land, in the ocean and in marine sediments, thus allowing to investigate the complex interactions between climate and the carbon cycle. Two setups of the global carbon cycle, closed and open, are available in CLIMBER-X, allowing both a proper comparison with CMIP6 type models on centennial
 580 scale and multi-millennia simulations. We have evaluated the model performance for the historical period against observations and state-of-the-art CMIP6 models, showing that many characteristics and feedbacks of the carbon cycle are reasonably well




Figure 34. Comparison of CLIMBER-X simulations with ZECMIP model results in terms of (a) global temperature, (b) atmospheric CO₂ concentration, (c) cumulative ocean carbon uptake and (d) cumulative land carbon uptake for the standard ZECMIP experiment with 1000 PgC cumulative carbon emissions. For CLIMBER-X, results with both the open (solid lines) and closed (dashed lines) carbon cycle setups are shown.

captured by the model. Biases in the simulated distribution of ocean biogeochemical tracers exist and can mostly be related to deficiencies in the simulated ocean circulation changes, which can at least partly be attributed to the comparatively coarse resolution of the ocean model and to the frictional-geostrophic approximation that it rests upon. On land, the carbon in soil layers below 1 m is underestimated, particularly in the permafrost zone, with possible implications for the land carbon cycle response to global warming.

Some possible directions for future model developments include the extension of the organic carbon cycle allowing for burial on land and in sediments and fluxes from land to ocean, the refinement of the carbonate chemistry on shelves, including coral growth, and possibly the addition of the nitrogen cycle on land, which could be important for nitrogen limitation of photosynthesis and would allow to have interactive atmospheric N₂O, considering that N₂O fluxes from the ocean are already available from the ocean biogeochemistry model HAMOCC.

The computational efficiency of CLIMBER-X will enable it to be used for systematic explorations of the coupled climate-carbon cycle evolution over timescales ranging from decades up to ~100,000 years, while also allowing a quantification of related uncertainties.

Code and data availability. The source code of CLIMBER-X v1.0 is archived on Zenodo (<https://doi.org/10.5281/zenodo.7898797>), with the exception of the HAMOCC model, which is covered by the Max Planck Institute for Meteorology software licence agreement as part of the MPI-ESM (https://code.mpimet.mpg.de/attachments/download/26986/MPI-ESM_SLA_v3.4.pdf). CMIP6 model data are licensed under a Creative Commons Attribution-ShareAlike 4.0 International License (<https://creativecommons.org/licenses>) and can be accessed through the ESGF nodes (for instance esgf-data.dkrz.de/search/cmip6-dkrz/).

Author contributions. M.W. and A.G. designed the model. T.I. and B.L. provided the HAMOCC model code and assisted in the implementation of the model into CLIMBER-X. C.H. helped with the sediment model setup and configuration. M.P. re-arranged HAMOCC into a column model for the purpose of integration into CLIMBER-X. M.H. implemented and tuned the particle ballasting scheme. D.D. contributed to the improvements in the land carbon cycle model. V.B. and G.M. assisted in the implementation and setup of the open carbon cycle. J.B., J.H. and G.R.M. developed the weathering model and contributed to its implementation into CLIMBER-X. M.W. coupled the different model components and tuned and tested the model. M.W. performed the model simulations, prepared the figures and wrote the paper, with contributions from all authors.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. M.W., B.L., M.H. and J.B. are funded by the German climate modeling project PalMod supported by the German Federal Ministry of Education and Research (BMBF) as a Research for Sustainability initiative (FONA) (grant nos. 01LP1920B, 01LP1917D, 01LP1506C (is this correct Janine?) Malte? Bo? 01LP1919B, 01LP1919C, 01LP1920C). G.M. is a Research Associate with the Belgian Fund for Scientific Research – F.R.S.-FNRS. We thank Irene Stemmler for technical support in implementing HAMOCC into CLIMBER-X and Thomas Kleinen for discussions on the methane cycle. We acknowledge the World Climate Research Programme, which, through its Working Group on Coupled Modelling, coordinated and promoted CMIP6. We thank the climate modeling groups for producing and making available their model output, the Earth System Grid Federation (ESGF) for archiving the data and providing access, and the multiple funding agencies who support CMIP6 and ESGF. We thank Nuno Carvalhais for providing the soil and vegetation carbon dataset. [The authors gratefully acknowledge the European Regional Development Fund \(ERDF\), the German Federal Ministry of Education and Research and the Land Brandenburg for supporting this project by providing resources on the high performance computer system at the Potsdam Institute for Climate Impact Research.](#)

620 References

- Abe-Ouchi, A., Saito, F., Kawamura, K., Raymo, M. E., Okuno, J., Takahashi, K., and Blatter, H.: Insolation-driven 100,000-year glacial cycles and hysteresis of ice-sheet volume., *Nature*, 500, 190–3, <https://doi.org/10.1038/nature12374>, <http://dx.doi.org/10.1038/nature12374>, 2013.
- Albani, S., Mahowald, N. M., Murphy, L. N., Raiswell, R., Moore, J. K., Anderson, R. F., McGee, D., Bradtmiller, L. I., Delmonte, B.,
625 Hesse, P. P., and Mayewski, P. A.: Paleodust variability since the Last Glacial Maximum and implications for iron inputs to the ocean, *Geophysical Research Letters*, 43, 3944–3954, <https://doi.org/10.1002/2016GL067911>, 2016.
- Amiotte Suchet, P. and Probst, J. L.: A global model for present-day atmospheric/soil CO₂ consumption by chemical erosion of continental rocks (GEM-CO₂), *Tellus B: Chemical and Physical Meteorology*, 47, 273–280, <https://doi.org/10.3402/tellusb.v47i1-2.16047>, 1995.
- Amiotte Suchet, P., Probst, J.-L., and Ludwig, W.: Worldwide distribution of continental rock lithology: Implications for the atmospheric/soil CO₂ uptake by continental weathering and alkalinity river transport to the oceans, *Global Biogeochemical Cycles*, 17, n/a–n/a, <https://doi.org/10.1029/2002GB001891>, [http://dx.doi.org/10.1029/2002GB001891](http://dx.doi.org/10.1029/2002GB001891%5Cnhttp://doi.wiley.com/10.1029/2002GB001891), 2003.
- Andres, R., Boden, T., and Marland, G.: Annual Fossil-Fuel CO₂ Emissions: Global Stable Carbon Isotopic Signature, <https://doi.org/10.3334/CDIAC/FFE.DB1013.2017>, <https://data.ess-dive.lbl.gov/datasets/doi:10.3334/CDIAC/FFE.DB1013.2017>, 2016.
- 635 Archer, D. E.: An atlas of the distribution of calcium carbonate in sediments of the deep sea, *Global Biogeochemical Cycles*, 10, 159–174, <https://doi.org/10.1029/95GB03016>, <http://doi.wiley.com/10.1029/95GB03016>, 1996.
- Archer, D. E. and Johnson, K.: A model of the iron cycle in the ocean, *Global Biogeochemical Cycles*, 14, 269–279, <https://doi.org/10.1029/1999GB900053>, <http://doi.wiley.com/10.1029/1999GB900053>, 2000.
- Arora, V. K., Boer, G. J., Friedlingstein, P., Eby, M., Jones, C. D., Christian, J. R., Bonan, G., Bopp, L., Brovkin, V., Cadule, P., Hajima, T.,
640 Ilyina, T., Lindsay, K., Tjiputra, J. F., and Wu, T.: Carbon-concentration and carbon-climate feedbacks in CMIP5 earth system models, *Journal of Climate*, 26, 5289–5314, <https://doi.org/10.1175/JCLI-D-12-00494.1>, 2013.
- Arora, V. K., Katavouta, A., Williams, R. G., Jones, C. D., Brovkin, V., Friedlingstein, P., Schwinger, J., Bopp, L., Boucher, O., Cadule, P., Chamberlain, M. A., Christian, J. R., Delire, C., Fisher, A. R. A., Hajima, T., Ilyina, T., Joetzjer, E., Kawamiya, M., Koven, C. D., Krasting, J. P., Law, R. M., Lawrence, D. M., Lenton, A., Lindsay, K., Pongratz, J., Raddatz, T., Séférian, R., Tachiiri, K., Tjiputra, J. F., Wiltshire,
645 A., Wu, T., and Ziehn, T.: Carbon-concentration and carbon-climate feedbacks in CMIP6 models and their comparison to CMIP5 models, *Biogeosciences*, 17, 4173–4222, <https://doi.org/10.5194/bg-17-4173-2020>, 2020.
- Augustin, L., Barbante, C., Barnes, P. R. F., Barnola, J. M., Bigler, M., Castellano, E., Cattani, O., Chappellaz, J., Dahl-Jensen, D., Delmonte, B., Dreyfus, G., Durand, G., Falourd, S., Fischer, H., Flückiger, J., Hansson, M. E., Huybrechts, P., Jugie, G., Johnsen, S. J., Jouzel, J., Kaufmann, P., Kipfstuhl, J., Lambert, F., Lipenkov, V. Y., Littot, G. C., Longinelli, A., Lorrain, R., Maggi, V., Masson-Delmotte, V.,
650 Miller, H., Mulvaney, R., Oerlemans, J., Oerter, H., Orombelli, G., Parrenin, F., Peel, D. a., Petit, J.-R., Raynaud, D., Ritz, C., Ruth, U., Schwander, J., Siegenthaler, U., Souchez, R., Stauffer, B., Steffensen, J. P., Stenni, B., Stocker, T. F., Tabacco, I. E., Udisti, R., Van De Wal, R. S. W., Van Den Broeke, M., Weiss, J., Wilhelms, F., Winther, J.-G., Wolff, E. W., and Zucchelli, M.: Eight glacial cycles from an Antarctic ice core., *Nature*, 429, 623–8, <https://doi.org/10.1038/nature02599>, <http://www.ncbi.nlm.nih.gov/pubmed/15190344>, 2004.
- Aumont, O., Orr, J. C., Monfray, P., Madec, G., and Maier-Reimer, E.: Nutrient trapping in the equatorial Pacific: The ocean circulation solution, *Global Biogeochemical Cycles*, 13, 351–369, <https://doi.org/10.1029/1998GB900012>, <http://doi.wiley.com/10.1029/1998GB900012>, 1999.

- Aumont, O., Ethé, C., Tagliabue, A., Bopp, L., and Gehlen, M.: PISCES-v2: An ocean biogeochemical model for carbon and ecosystem studies, *Geoscientific Model Development*, 8, 2465–2513, <https://doi.org/10.5194/gmd-8-2465-2015>, 2015.
- 660 Batjes, N.: Harmonized soil property values for broad-scale modelling (WISE30sec) with estimates of global soil carbon stocks, *Geoderma*, 269, 61–68, <https://doi.org/10.1016/j.geoderma.2016.01.034>, <https://linkinghub.elsevier.com/retrieve/pii/S0016706116300349>, 2016.
- Battaglia, G., Steinacher, M., and Joos, F.: A probabilistic assessment of calcium carbonate export and dissolution in the modern ocean, *Biogeosciences*, 13, 2823–2848, <https://doi.org/10.5194/bg-13-2823-2016>, 2016.
- Beer, C., Reichstein, M., Tomelleri, E., Ciais, P., Jung, M., Carvalhais, N., Rodenbeck, C., Arain, M. A., Baldocchi, D., Bonan, G. B., Bondeau, A., Cescatti, A., Lasslop, G., Lindroth, A., Lomas, M., Luysaert, S., Margolis, H., Oleson, K. W., Rouspard, O., Veenendaal, E., 665 Viovy, N., Williams, C., Woodward, F. I., and Papale, D.: Terrestrial Gross Carbon Dioxide Uptake: Global Distribution and Covariation with Climate, *Science*, 329, 834–838, <https://doi.org/10.1126/science.1184984>, <http://www.sciencemag.org/cgi/doi/10.1126/science.1184984>, 2010.
- Börker, J., Hartmann, J., Amann, T., Romero-Mujalli, G., Moosdorf, N., and Jenkins, C.: Chemical Weathering of Loess and Its Contribution to Global Alkalinity Fluxes to the Coastal Zone During the Last Glacial Maximum, Mid-Holocene, and Present, *Geochemistry, Geophysics, Geosystems*, 21, <https://doi.org/10.1029/2020GC008922>, 2020.
- 670 Bouttes, N., Roche, D. M., Mariotti, V., and Bopp, L.: Including an ocean carbon cycle model into iLOVECLIM (v1.0), *Geoscientific Model Development*, 8, 1563–1576, <https://doi.org/10.5194/gmd-8-1563-2015>, 2015.
- Brovkin, V., Bendtsen, J., Claussen, M., Ganopolski, A., Kubatzki, C., Petoukhov, V., and Andreev, A.: Carbon cycle, vegetation, and climate dynamics in the Holocene: Experiments with the CLIMBER-2 model, *Global Biogeochemical Cycles*, 16, 86– 675 1–86–20, <https://doi.org/10.1029/2001GB001662>, <http://doi.wiley.com/10.1029/2001GB001662><http://www.agu.org/pubs/crossref/2002/2001GB001662.shtml>, 2002.
- Brovkin, V., Ganopolski, A., Archer, D., and Rahmstorf, S.: Lowering of glacial atmospheric CO₂ in response to changes in oceanic circulation and marine biogeochemistry, *Paleoceanography*, 22, n/a–n/a, <https://doi.org/10.1029/2006PA001380>, <http://doi.wiley.com/10.1029/2006PA001380>, 2007.
- 680 Brovkin, V., Ganopolski, A., Archer, D., and Munhoven, G.: Glacial CO₂ cycle as a succession of key physical and biogeochemical processes, *Climate of the Past*, 8, 251–264, <https://doi.org/10.5194/cp-8-251-2012>, <http://www.clim-past.net/8/251/2012/>, 2012.
- Brown, J., Ferrians, O., Heginbottom, J. A., and Melnikov, E.: Circum-Arctic Map of Permafrost and Ground-Ice Conditions, 1998.
- Buitenhuis, E. T., Suntharalingam, P., and Le Quéré, C.: Constraints on global oceanic emissions of N₂O from observations and models, *Biogeosciences*, 15, 2161–2175, <https://doi.org/10.5194/bg-15-2161-2018>, 2018.
- 685 Burton, C., Betts, R., Cardoso, M., Feldpausch, R. T., Harper, A., Jones, C. D., Kelley, D. I., Robertson, E., and Wiltshire, A.: Representation of fire, land-use change and vegetation dynamics in the Joint UK Land Environment Simulator vn4.9 (JULES), *Geoscientific Model Development*, 12, 179–193, <https://doi.org/10.5194/gmd-12-179-2019>, 2019.
- Cabré, A., Marinov, I., Bernardello, R., and Bianchi, D.: Oxygen minimum zones in the tropical Pacific across CMIP5 models: Mean state differences and climate change trends, *Biogeosciences*, 12, 5429–5454, <https://doi.org/10.5194/bg-12-5429-2015>, 2015.
- 690 Cao, L., Eby, M., Ridgwell, A., Caldeira, K., Archer, D., Ishida, A., Joos, F., Matsumoto, K., Mikolajewicz, U., Mouchet, A., Orr, J. C., Plattner, G. K., Schlitzer, R., Tokos, K., Totterdell, I., Tschumi, T., Yamanaka, Y., and Yool, A.: The role of ocean transport in the uptake of anthropogenic CO₂, *Biogeosciences*, 6, 375–390, <https://doi.org/10.5194/bg-6-375-2009>, 2009.
- Carr, M.-E., Friedrichs, M. A., Schmeltz, M., Noguchi Aita, M., Antoine, D., Arrigo, K. R., Asanuma, I., Aumont, O., Barber, R., Behrenfeld, M., Bidigare, R., Buitenhuis, E. T., Campbell, J., Ciotti, A., Dierssen, H., Dowell, M., Dunne, J., Esaias, W., Gentili, B.,

- 695 Gregg, W., Groom, S., Hoepffner, N., Ishizaka, J., Kameda, T., Le Quéré, C., Lohrenz, S., Marra, J., Mélin, F., Moore, K., Morel, A., Reddy, T. E., Ryan, J., Scardi, M., Smyth, T., Turpie, K., Tilstone, G., Waters, K., and Yamanaka, Y.: A comparison of global estimates of marine primary production from ocean color, *Deep Sea Research Part II: Topical Studies in Oceanography*, 53, 741–770, <https://doi.org/10.1016/j.dsr2.2006.01.028>, <https://linkinghub.elsevier.com/retrieve/pii/S0967064506000555>, 2006.
- Cartapanis, O., Galbraith, E. D., Bianchi, D., and Jaccard, S. L.: Carbon burial in deep-sea sediment and implications for oceanic inventories of carbon and alkalinity over the last glacial cycle, *Climate of the Past*, 14, 1819–1850, <https://doi.org/10.5194/cp-14-1819-2018>, <https://www.clim-past-discuss.net/cp-2018-49/><https://cp.copernicus.org/articles/14/1819/2018/>, 2018.
- 700
- Carvalho, N., Forkel, M., Khomik, M., Bellarby, J., Jung, M., Migliavacca, M., Mu, M., Saatchi, S., Santoro, M., Thurner, M., Weber, U., Ahrens, B., Beer, C., Cescatti, A., Randerson, J. T., and Reichstein, M.: Global covariation of carbon turnover times with climate in terrestrial ecosystems, *Nature*, 514, 213–217, <https://doi.org/10.1038/nature13731>, 2014.
- 705
- Chen, F. and Zhang, Y.: On the coupling strength between the land surface and the atmosphere: From viewpoint of surface exchange coefficients, *Geophysical Research Letters*, 36, 1–5, <https://doi.org/10.1029/2009GL037980>, 2009.
- Collatz, G., Ball, J., Grivet, C., and Berry, J. A.: Physiological and environmental regulation of stomatal conductance, photosynthesis and transpiration: a model that includes a laminar boundary layer, *Agricultural and Forest Meteorology*, 54, 107–136, [https://doi.org/10.1016/0168-1923\(91\)90002-8](https://doi.org/10.1016/0168-1923(91)90002-8), <http://www.sciencedirect.com/science/article/pii/0168192391900028><http://linkinghub.elsevier.com/retrieve/pii/0168192391900028>, 1991.
- 710
- Crichton, K. A., Wilson, J. D., Ridgwell, A., and Pearson, P. N.: Calibration of temperature-dependent ocean microbial processes in the cGENIE.muffin (v0.9.13) Earth system model, *Geoscientific Model Development*, 14, 125–149, <https://doi.org/10.5194/gmd-14-125-2021>, 2021.
- Denman, K. L., Brasseur, G., Chidthaisong, A., Ciais, P., Cox, P. M., Dickinson, R. E., Hauglustaine, D., Heinze, C., Holland, E., Jacob, D., Lohmann, U., Ramachandran, S., da Silva Dias, P. L., Wofsy, S. C., and Zhang, X.: Couplings Between Changes in the Climate System and Biogeochemistry, in: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Solomon, S., Qin, D., Manning, M., Marquis, M., Averyt, K., Tignor, M. M. B., Jr., H. L. M., and Chen, Z., chap. 7, pp. 499–587, Cambridge University Press, Cambridge (UK), <https://www.ipcc.ch/site/assets/uploads/2018/02/ar4-wg1-chapter7-1.pdf>, 2007.
- 715
- Derry, L. A. and France-Lanord, C.: Neogene growth of the sedimentary organic carbon reservoir, *Paleoceanography*, 11, 267–275, <https://doi.org/10.1029/95PA03839>, <http://doi.wiley.com/10.1029/95PA03839>, 1996.
- 720
- Dietze, H. and Loeptien, U.: Revisiting "nutrient trapping" in global coupled biogeochemical ocean circulation models, *Global Biogeochemical Cycles*, 27, 265–284, <https://doi.org/10.1002/gbc.20029>, 2013.
- Dunne, J. P., Sarmiento, J. L., and Gnanadesikan, A.: A synthesis of global particle export from the surface ocean and cycling through the ocean interior and on the seafloor, *Global Biogeochemical Cycles*, 21, 1–16, <https://doi.org/10.1029/2006GB002907>, 2007.
- 725
- Dutay, J.-C., Bullister, J., Doney, S., Orr, J., Najjar, R., Caldeira, K., Campin, J.-M., Drange, H., Follows, M., Gao, Y., Gruber, N., Hecht, M., Ishida, A., Joos, F., Lindsay, K., Madec, G., Maier-Reimer, E., Marschall, J., Matear, R., Monfray, P., Mouchet, A., Plattner, G.-K., Sarmiento, J., Schlitzer, R., Slater, R., Totterdell, I., Weirig, M.-F., Yamanaka, Y., and Yool, A.: Evaluation of ocean model ventilation with CFC-11: comparison of 13 global ocean models, *Ocean Modelling*, 4, 89–120, [https://doi.org/10.1016/S1463-5003\(01\)00013-0](https://doi.org/10.1016/S1463-5003(01)00013-0), <http://linkinghub.elsevier.com/retrieve/pii/S1463500301000130>, 2002.
- 730

- Eby, M., Zickfeld, K., Montenegro, A., Archer, D., Meissner, K. J., and Weaver, A. J.: Lifetime of anthropogenic climate change: Millennial time scales of potential CO₂ and surface temperature perturbations, *Journal of Climate*, 22, 2501–2511, <https://doi.org/10.1175/2008JCLI2554.1>, 2009.
- Eggleston, S., Schmitt, J., Bereiter, B., Schneider, R., and Fischer, H.: Evolution of the stable carbon isotope composition of atmospheric CO₂ over the last glacial cycle, *Paleoceanography*, 31, 434–452, <https://doi.org/10.1002/2015PA002874>, 2016.
- Eide, M., Olsen, A., Ninnemann, U. S., and Johannessen, T.: A global ocean climatology of preindustrial and modern ocean $\delta^{13}\text{C}$, *Global Biogeochemical Cycles*, 31, 515–534, <https://doi.org/10.1002/2016GB005473>, 2017.
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geoscientific Model Development*, 9, 1937–1958, <https://doi.org/10.5194/gmd-9-1937-2016>, 2016.
- Fan, N., Koirala, S., Reichstein, M., Thurner, M., Avitabile, V., Santoro, M., Ahrens, B., Weber, U., and Carvalhais, N.: Apparent ecosystem carbon turnover time: Uncertainties and robust features, *Earth System Science Data*, 12, 2517–2536, <https://doi.org/10.5194/essd-12-2517-2020>, 2020.
- Farquhar, G. D., von Caemmerer, S., and Berry, J. A.: A biochemical model of photosynthetic CO₂ assimilation in leaves of C₃ species, *Planta*, 149, 78–90, <https://doi.org/10.1007/BF00386231>, 1980.
- Friedlingstein, P., Cox, P., Betts, R., Bopp, L., von Bloh, W., Brovkin, V., Cadule, P., Doney, S., Eby, M., Fung, I., Bala, G., John, J., Jones, C., Joos, F., Kato, T., Kawamiya, M., Knorr, W., Lindsay, K., Matthews, H. D., Raddatz, T., Rayner, P., Reick, C., Roeckner, E., Schnitzler, K.-G., Schnur, R., Strassmann, K., Weaver, A. J., Yoshikawa, C., and Zeng, N.: Climate–Carbon Cycle Feedback Analysis: Results from the C₄ MIP Model Intercomparison, *Journal of Climate*, 19, 3337–3353, <https://doi.org/10.1175/JCLI3800.1>, <http://journals.ametsoc.org/doi/abs/10.1175/JCLI3800.1>, 2006.
- Friedlingstein, P., Meinshausen, M., Arora, V. K., Jones, C. D., Anav, A., Liddicoat, S. K., and Knutti, R.: Uncertainties in CMIP5 climate projections due to carbon cycle feedbacks, *Journal of Climate*, 27, 511–526, <https://doi.org/10.1175/JCLI-D-12-00579.1>, 2014.
- Friedlingstein, P., Jones, M. W., O’Sullivan, M., Andrew, R. M., Bakker, D. C. E., Hauck, J., Le Quéré, C., Peters, G. P., Peters, W., Pongratz, J., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S. R., Anthoni, P., Bates, N. R., Becker, M., Bellouin, N., Bopp, L., Chau, T., T. T., Chevallier, F., Chini, L. P., Cronin, M., Currie, K. I., Decharme, B., Djeutchouang, L. M., Dou, X., Evans, W., Feely, R. A., Feng, L., Gasser, T., Gilfillan, D., Gkritzalis, T., Grassi, G., Gregor, L., Gruber, N., Gürses, Ö., Harris, I., Houghton, R. A., Hurtt, G. C., Iida, Y., Ilyina, T., Luijkx, I. T., Jain, A., Jones, S. D., Kato, E., Kennedy, D., Klein Goldewijk, K., Knauer, J., Korsbakken, J. I., Körtzinger, A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lienert, S., Liu, J., Marland, G., McGuire, P. C., Melton, J. R., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S.-I., Niwa, Y., Ono, T., Pierrot, D., Poulter, B., Rehder, G., Resplandy, L., Robertson, E., Rödenbeck, C., Rosan, T. M., Schwinger, J., Schwingshackl, C., Séférian, R., Sutton, A. J., Sweeney, C., Tanhua, T., Tans, P. P., Tian, H., Tilbrook, B., Tubiello, F., van der Werf, G. R., Vuichard, N., Wada, C., Wanninkhof, R., Watson, A. J., Willis, D., Wiltshire, A. J., Yuan, W., Yue, C., Yue, X., Zaehle, S., and Zeng, J.: Global Carbon Budget 2021, *Earth System Science Data*, 14, 1917–2005, <https://doi.org/10.5194/essd-14-1917-2022>, <https://essd.copernicus.org/articles/14/1917/2022/>, 2022.
- Gaillardet, J., Dupré, B., Louvat, P., and Allègre, C.: Global silicate weathering and CO₂ consumption rates deduced from the chemistry of large rivers, *Chemical Geology*, 159, 3–30, [https://doi.org/10.1016/S0009-2541\(99\)00031-5](https://doi.org/10.1016/S0009-2541(99)00031-5), <https://linkinghub.elsevier.com/retrieve/pii/S0009254199000315>, 1999.
- Ganopolski, A. and Brovkin, V.: Simulation of climate, ice sheets and CO₂ evolution during the last four glacial cycles with an Earth system model of intermediate complexity, *Climate of the Past*, 13, 1695–1716, <https://doi.org/10.5194/cp-13-1695-2017>, 2017.

- Ganopolski, A. and Calov, R.: The role of orbital forcing, carbon dioxide and regolith in 100 kyr glacial cycles, *Climate of the Past*, 7, 1415–1425, <https://doi.org/10.5194/cp-7-1415-2011>, [http://www.clim-past.net/7/1415/2011/](http://www.clim-past.net/7/1415/2011/cp-7-1415-2011.pdf), 2011.
- Garcia, H. E., Boyer, T. P., Locarnini, R. A., Antonov, J. I., Mishonov, A. V., Baranova, O. K., Zweng, M. M., Reagan, J. R., and Johnson, D. R.: *World Ocean Atlas 2013. Volume 3: dissolved oxygen, apparent oxygen utilization, and oxygen saturation*, NOAA Atlas NESDIS 75, 3, 27, 2013a.
- Garcia, H. E., Locarnini, R., Boyer, T. P., Antonov, J. I., Baranova, O. K., Zweng, M. M., Reagan, J. R., and Johnson, D. R.: *World Ocean Atlas 2013 Volume 4 : Nutrients (phosphate , nitrate , silicate)*, NOAA Atlas NESDIS 76, 4, 396, 2013b.
- Gasser, T., Crepin, L., Quilcaille, Y., Houghton, R. A., Ciais, P., and Obersteiner, M.: Historical CO₂ emissions from land use and land cover change and their uncertainty, *Biogeosciences*, 17, 4075–4101, <https://doi.org/10.5194/bg-17-4075-2020>, 2020.
- Gehlen, M., Bopp, L., Emprin, N., Aumont, O., Heinze, C., and Ragueneau, O.: Reconciling surface ocean productivity, export fluxes and sediment composition in a global biogeochemical ocean model, *Biogeosciences*, 3, 521–537, <https://doi.org/10.5194/bg-3-521-2006>, 2006.
- Ginoux, P., Chin, M., Tegen, I., Prospero, J. M., Holben, B., Dubovik, O., and Lin, S. J.: Sources and distributions of dust aerosols simulated with the GOCART model, *Journal of Geophysical Research Atmospheres*, 106, 20 255–20 273, <https://doi.org/10.1029/2000JD000053>, 2001.
- Goosse, H., Brovkin, V., Fichefet, T., Haarsma, R., Huybrechts, P., Jongma, J., Mouchet, A., Selten, F., Barriat, P. Y., Campin, J. M., Deleersnijder, E., Driesschaert, E., Goelzer, H., Janssens, I., Loutre, M. F., Morales Maqueda, M. A., Opsteegh, T., Mathieu, P. P., Munhoven, G., Pettersson, E. J., Renssen, H., Roche, D. M., Schaeffer, M., Tartinville, B., Timmermann, A., and Weber, S. L.: Description of the Earth system model of intermediate complexity LOVECLIM version 1.2, *Geoscientific Model Development*, 3, 603–633, <https://doi.org/10.5194/gmd-3-603-2010>, 2010.
- Großkopf, T., Mohr, W., Baustian, T., Schunck, H., Gill, D., Kuypers, M. M. M., Lavik, G., Schmitz, R. A., Wallace, D. W. R., and LaRoche, J.: Doubling of marine dinitrogen-fixation rates based on direct measurements, *Nature*, 488, 361–364, <https://doi.org/10.1038/nature11338>, <http://www.nature.com/articles/nature11338>, 2012.
- Gruber, N., Gloor, M., Mikaloff Fletcher, S. E., Doney, S. C., Dutkiewicz, S., Follows, M. J., Gerber, M., Jacobson, A. R., Joos, F., Lindsay, K., Menemenlis, D., Mouchet, A., Müller, S. A., Sarmiento, J. L., and Takahashi, T.: Oceanic sources, sinks, and transport of atmospheric CO₂, *Global Biogeochemical Cycles*, 23, 1–21, <https://doi.org/10.1029/2008GB003349>, 2009.
- Gulev, S., Thorne, P., Ahn, J., Dentener, F., Domingues, C., Gerland, S., Gong, D., Kaufman, D., Nnamchi, H., Quaas, J., Rivera, J., Sathyendranath, S., Smith, S., Trewin, B., von Schuckmann, K., and Vose, R.: Chapter 2: Changing state of the climate system, in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J., Maycock, T., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., pp. 287–422, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/9781009157896.004>, 2021.
- Hartmann, J.: Bicarbonate-fluxes and CO₂-consumption by chemical weathering on the Japanese Archipelago - Application of a multi-lithological model framework, *Chemical Geology*, 265, 237–271, <https://doi.org/10.1016/j.chemgeo.2009.03.024>, <http://dx.doi.org/10.1016/j.chemgeo.2009.03.024>, 2009.
- Hartmann, J. and Moosdorf, N.: The new global lithological map database GLiM: A representation of rock properties at the Earth surface, *Geochemistry, Geophysics, Geosystems*, 13, 1–37, <https://doi.org/10.1029/2012GC004370>, 2012.

- Hartmann, J., Moosdorf, N., Lauerwald, R., Hinderer, M., and West, A. J.: Global chemical weathering and associated p-release - the role of lithology, temperature and soil properties, *Chemical Geology*, 363, 145–163, <https://doi.org/10.1016/j.chemgeo.2013.10.025>, <http://dx.doi.org/10.1016/j.chemgeo.2013.10.025>, 2014.
- 810 Hauck, J., Völker, C., Wang, T., Hoppema, M., Losch, M., and Wolf-Gladrow, D. A.: Seasonally different carbon flux changes in the Southern Ocean in response to the southern annular mode, *Global Biogeochemical Cycles*, 27, 1236–1245, <https://doi.org/10.1002/2013GB004600>, 2013.
- Haxeltine, A. and Prentice, I. C.: A General Model for the Light-Use Efficiency of Primary Production, <https://doi.org/10.2307/2390165>, <http://www.jstor.org/stable/2390165>, 1996.
- 815 Hayes, C. T., Costa, K. M., Anderson, R. F., Calvo, E., Chase, Z., Demina, L. L., Dutay, J. C., German, C. R., Heimbürger-Boavida, L. E., Jaccard, S. L., Jacobel, A., Kohfeld, K. E., Kravchishina, M. D., Lippold, J., Mekik, F., Missiaen, L., Pavia, F. J., Paytan, A., Pedrosa-Pamies, R., Petrova, M. V., Rahman, S., Robinson, L. F., Roy-Barman, M., Sanchez-Vidal, A., Shiller, A., Tagliabue, A., Tessin, A. C., van Hulten, M., and Zhang, J.: Global Ocean Sediment Composition and Burial Flux in the Deep Sea, *Global Biogeochemical Cycles*, 35, 1–25, <https://doi.org/10.1029/2020GB006769>, 2021.
- 820 Heinemann, M., Segschneider, J., and Schneider, B.: CO₂ drawdown due to particle ballasting by glacial aeolian dust: An estimate based on the ocean carbon cycle model MPIOM/HAMOCC version 1.6.2p3, *Geoscientific Model Development*, 12, 1869–1883, <https://doi.org/10.5194/gmd-12-1869-2019>, 2019.
- Heinze, C., Maier-Reimer, E., Winguth, A. M. E., and Archer, D.: A global oceanic sediment model for long-term climate studies, *Global Biogeochemical Cycles*, 13, 221–250, <https://doi.org/10.1029/98GB02812>, <http://doi.wiley.com/10.1029/98GB02812>, 1999.
- 825 Hoffman, F. M., Randerson, J. T., Arora, V. K., Bao, Q., Cadule, P., Ji, D., Jones, C. D., Kawamiya, M., Khatiwala, S., Lindsay, K., Obata, A., Shevliakova, E., Six, K. D., Tjiputra, J. F., Volodin, E. M., and Wu, T.: Causes and implications of persistent atmospheric carbon dioxide biases in Earth System Models, *Journal of Geophysical Research: Biogeosciences*, 119, 141–162, <https://doi.org/10.1002/2013JG002381>, <https://onlinelibrary.wiley.com/doi/10.1002/2013JG002381>, 2014.
- Hopcroft, P. O., Valdes, P. J., Woodward, S., and Joshi, M. M.: Last glacial maximum radiative forcing from mineral dust aerosols in an Earth system model, *Journal of Geophysical Research: Atmospheres*, 120, 8186–8205, <https://doi.org/10.1002/2015JD023742>, 2015.
- 830 Hopcroft, P. O., Valdes, P. J., O'Connor, F. M., Kaplan, J. O., and Beerling, D. J.: Understanding the glacial methane cycle, *Nature Communications*, 8, <https://doi.org/10.1038/ncomms14383>, 2017.
- Hugelius, G., Strauss, J., Zubrzycki, S., Harden, J. W., Schuur, E. a. G., Ping, C.-L. L., Schirrmeister, L., Grosse, G., Michaelson, G. J., Koven, C. D., O'Donnell, J. A., Elberling, B., Mishra, U., Camill, P., Yu, Z., Palmtag, J., and Kuhry, P.: Estimated stocks of circumpolar permafrost carbon with quantified uncertainty ranges and identified data gaps, *Biogeosciences*, 11, 6573–6593, <https://doi.org/10.5194/bg-11-6573-2014>, <http://www.biogeosciences.net/11/6573/2014/>, 2014.
- 835 Huneeus, N., Schulz, M., Balkanski, Y., Griesfeller, J., Prospero, J., Kinne, S., Bauer, S., Boucher, O., Chin, M., Dentener, F., Diehl, T., Easter, R., Fillmore, D., Ghan, S., Ginoux, P., Grini, A., Horowitz, L., Koch, D., Krol, M. C., Landing, W., Liu, X., Mahowald, N., Miller, R., Morcrette, J. J., Myhre, G., Penner, J., Perlwitz, J., Stier, P., Takemura, T., and Zender, C. S.: Global dust model intercomparison in AeroCom phase i, *Atmospheric Chemistry and Physics*, 11, 7781–7816, <https://doi.org/10.5194/acp-11-7781-2011>, 2011.
- 840 Ilyina, T., Six, K. D., Segschneider, J., Maier-Reimer, E., Li, H., and Núñez-Riboni, I.: Global ocean biogeochemistry model HAMOCC: Model architecture and performance as component of the MPI-Earth system model in different CMIP5 experimental realizations, *Journal of Advances in Modeling Earth Systems*, 5, 287–315, <https://doi.org/10.1029/2012MS000178>, 2013.

- Ito, A.: A historical meta-analysis of global terrestrial net primary productivity: Are estimates converging?, *Global Change Biology*, 17, 845 3161–3175, <https://doi.org/10.1111/j.1365-2486.2011.02450.x>, 2011.
- Jacobson, A. R., Fletcher, S. E., Gruber, N., Sarmiento, J. L., and Gloor, M.: A joint atmosphere-ocean inversion for surface fluxes of carbon dioxide: 1. Methods and global-scale fluxes, *Global Biogeochemical Cycles*, 21, <https://doi.org/10.1029/2005GB002556>, 2007.
- Johnson, K. S. and Bif, M. B.: Constraint on net primary productivity of the global ocean by Argo oxygen measurements, *Nature Geoscience*, 14, 769–774, <https://doi.org/10.1038/s41561-021-00807-z>, <http://dx.doi.org/10.1038/s41561-021-00807-z>, 2021.
- 850 Jones, C. D., Frölicher, T. L., Koven, C., MacDougall, A. H., Damon Matthews, H., Zickfeld, K., Rogelj, J., Tokarska, K. B., Gillett, N. P., Ilyina, T., Meinshausen, M., Mengis, N., Séférian, R., Eby, M., and Burger, F. A.: The Zero Emissions Commitment Model Intercomparison Project (ZECMIP) contribution to C4MIP: Quantifying committed climate changes following zero carbon emissions, *Geoscientific Model Development*, 12, 4375–4385, <https://doi.org/10.5194/gmd-12-4375-2019>, 2019.
- Jung, M., Reichstein, M., Margolis, H. a., Cescatti, A., Richardson, A. D., Arain, M. A., Arneth, A., Bernhofer, C., Bonal, D., Chen, J., 855 Gianelle, D., Gobron, N., Kiely, G., Kutsch, W., Lasslop, G., Law, B. E., Lindroth, A., Merbold, L., Montagnani, L., Moors, E. J., Papale, D., Sottocornola, M., Vaccari, F., and Williams, C.: Global patterns of land-atmosphere fluxes of carbon dioxide, latent heat, and sensible heat derived from eddy covariance, satellite, and meteorological observations, *Journal of Geophysical Research: Biogeosciences*, 116, 2–4, <https://doi.org/10.1029/2010JG001566>, 2011.
- Karl, D., Michaels, A., Bergman, B., Capone, D., Carpenter, E., Letelier, R., Lipschultz, F., Paerl, H., Sigman, D., and Stal, L.: Dinitro- 860 gen fixation in the world's oceans, *Biogeochemistry*, 57, 47–98, <https://doi.org/10.1023/A:1015798105851>, <https://doi.org/10.1023/A:1015798105851>, 2002.
- Key, R. M., Kozyr, A., Sabine, C. L., Lee, K., Wanninkhof, R., Bullister, J. L., Feely, R. A., Millero, F. J., Mordy, C., and Peng, T. H.: A global ocean carbon climatology: Results from Global Data Analysis Project (GLODAP), *Global Biogeochemical Cycles*, 18, 1–23, <https://doi.org/10.1029/2004GB002247>, 2004.
- 865 Kleinen, T. and Brovkin, V.: Pathway-dependent fate of permafrost region carbon, *Environmental Research Letters*, 13, <https://doi.org/10.1088/1748-9326/aad824>, 2018.
- Kleinen, T., Mikolajewicz, U., and Brovkin, V.: Terrestrial methane emissions from the Last Glacial Maximum to the preindustrial period, *Climate of the Past*, 16, 575–595, <https://doi.org/10.5194/cp-16-575-2020>, 2020.
- Köhler, P., Nehrbass-Ahles, C., Schmitt, J., Stocker, T. F., and Fischer, H.: A 156 kyr smoothed history of the atmospheric greenhouse 870 gases CO₂, CH₄ and N₂O and their radiative forcing, *Earth System Science Data*, 9, 363–387, <https://doi.org/10.5194/essd-9-363-2017>, <https://www.earth-syst-sci-data.net/9/363/2017/https://essd.copernicus.org/articles/9/363/2017/>, 2017.
- Koven, C. D., Riley, W. J., Subin, Z. M., Tang, J. Y., Torn, M. S., Collins, W. D., Bonan, G. B., Lawrence, D. M., and Swenson, S. C.: The effect of vertically resolved soil biogeochemistry and alternate soil C and N models on C dynamics of CLM4, *Biogeosciences*, 10, 7109–7131, <https://doi.org/10.5194/bg-10-7109-2013>, <http://www.biogeosciences.net/10/7109/2013/>, 2013.
- 875 Kriest, I. and Evans, G. T.: A vertically resolved model for phytoplankton aggregation, *Journal of Earth System Science*, 109, 453–469, <https://doi.org/10.1007/BF02708333>, <http://link.springer.com/10.1007/BF02708333>, 2000.
- Kurahashi-Nakamura, T., Paul, A., Merkel, U., and Schulz, M.: Glacial state of the global carbon cycle: time-slice simulations for the last glacial maximum with an Earth-system model, *Climate of the Past*, 18, 1997–2019, <https://doi.org/10.5194/cp-18-1997-2022>, 2022.
- Lambert, F., Tagliabue, A., Shaffer, G., Lamy, F., Winckler, G., Farias, L., Gallardo, L., and De Pol-Holz, R.: Dust fluxes and iron fertilization 880 in Holocene and Last Glacial Maximum climates, *Geophysical Research Letters*, 42, 6014–6023, <https://doi.org/10.1002/2015GL064250>, <http://doi.wiley.com/10.1002/2015GL064250>, 2015.

- Landschützer, P., Gruber, N., and Bakker, D. C.: Decadal variations and trends of the global ocean carbon sink, *Global Biogeochemical Cycles*, 30, 1396–1417, <https://doi.org/10.1002/2015GB005359>, 2016.
- 885 Laufkötter, C., John, J. G., Stock, C. A., and Dunne, J. P.: Temperature and oxygen dependence of the remineralization of organic matter, *Global Biogeochemical Cycles*, 31, 1038–1050, <https://doi.org/10.1002/2017GB005643>, 2017.
- Lauvset, S. K., Key, R. M., Olsen, A., Van Heuven, S., Velo, A., Lin, X., Schirnick, C., Kozyr, A., Tanhua, T., Hoppema, M., Jutterström, S., Steinfeldt, R., Jeansson, E., Ishii, M., Perez, F. F., Suzuki, T., and Watelet, S.: A new global interior ocean mapped climatology: The $1^\circ \times 1^\circ$ GLODAP version 2, *Earth System Science Data*, 8, 325–340, <https://doi.org/10.5194/essd-8-325-2016>, 2016.
- 890 Lavergne, A., Voelker, S., Csank, A., Graven, H., de Boer, H. J., Daux, V., Robertson, I., Dorado-Liñán, I., Martínez-Sancho, E., Battipaglia, G., Bloomfield, K. J., Still, C. J., Meinzer, F. C., Dawson, T. E., Camarero, J. J., Clisby, R., Fang, Y., Menzel, A., Keen, R. M., Roden, J. S., and Prentice, I. C.: Historical changes in the stomatal limitation of photosynthesis: empirical support for an optimality principle, *New Phytologist*, n/a, nph.16314, <https://doi.org/10.1111/nph.16314>, <https://onlinelibrary.wiley.com/doi/abs/10.1111/nph.16314>, 2019.
- Lavergne, A., Sandoval, D., Hare, V. J., Graven, H., and Prentice, I. C.: Impacts of soil water stress on the acclimated stomatal limitation of photosynthesis: Insights from stable carbon isotope data, *Global Change Biology*, 26, 7158–7172, <https://doi.org/10.1111/gcb.15364>, 895 2020.
- Levine, J. G., Wolff, E. W., Jones, A. E., Sime, L. C., Valdes, P. J., Archibald, A. T., Carver, G. D., Warwick, N. J., and Pyle, J. A.: Reconciling the changes in atmospheric methane sources and sinks between the Last Glacial Maximum and the pre-industrial era, *Geophysical Research Letters*, 38, 2–7, <https://doi.org/10.1029/2011GL049545>, 2011.
- Liu, B., Six, K. D., and Ilyina, T.: Incorporating the stable carbon isotope ^{13}C in the ocean biogeochemical component of the Max Planck 900 Institute Earth System Model, *Biogeosciences*, 18, 4389–4429, <https://doi.org/10.5194/bg-18-4389-2021>, 2021.
- Ma, L., Hurtt, G. C., Chini, L. P., Sahajpal, R., Pongratz, J., Frohling, S., Stehfest, E., Klein Goldewijk, K., O’Leary, D., and Doelman, J. C.: Global rules for translating land-use change (LUH2) to land-cover change for CMIP6 using GLM2, *Geoscientific Model Development*, 13, 3203–3220, <https://doi.org/10.5194/gmd-13-3203-2020>, <https://gmd.copernicus.org/articles/13/3203/2020/>, 2020.
- 905 MacDougall, A. H., Frölicher, T. L., Jones, C. D., Rogelj, J., DamonMatthews, H., Zickfeld, K., Arora, V. K., Barrett, N. J., Brovkin, V., Burger, F. A., Eby, M., Eliseev, A. V., Hajima, T., Holden, P. B., Jeltsch-Thömmes, A., Koven, C., Mengis, N., Menviel, L., Michou, M., Mokhov, I. I., Oka, A., Schwinger, J., Séférian, R., Shaffer, G., Sokolov, A., Tachiiri, K., Tjiputra, J., Wiltshire, A., and Ziehn, T.: Is there warming in the pipeline? A multi-model analysis of the Zero Emissions Commitment from CO_2 , *Biogeosciences*, 17, 2987–3016, <https://doi.org/10.5194/bg-17-2987-2020>, 2020.
- 910 Maerz, J., Six, K. D., Stemmler, I., Ahmerkamp, S., and Ilyina, T.: Microstructure and composition of marine aggregates as co-determinants for vertical particulate organic carbon transfer in the global ocean, *Biogeosciences*, 17, 1765–1803, <https://doi.org/10.5194/bg-17-1765-2020>, 2020.
- Maier-Reimer, E. and Hasselmann, K.: Transport and storage of CO_2 in the ocean — an inorganic ocean-circulation carbon cycle model, *Climate Dynamics*, 2, 63–90, <https://doi.org/10.1007/BF01054491>, 1987.
- Maier-Reimer, E., Mikolajewicz, U., and Hasselmann, K.: Mean Circulation of the Hamburg LSG OGCM and Its Sensitivity to the Thermo- 915 haline Surface Forcing, [https://doi.org/10.1175/1520-0485\(1993\)023<0731:MCOTHL>2.0.CO;2](https://doi.org/10.1175/1520-0485(1993)023<0731:MCOTHL>2.0.CO;2), 1993.
- Matthes, K., Funke, B., Andersson, M. E., Barnard, L., Beer, J., Charbonneau, P., Clilverd, M. A., Dudok De Wit, T., Haberleiter, M., Hendry, A., Jackman, C. H., Kretzschmar, M., Kruschke, T., Kunze, M., Langematz, U., Marsh, D. R., Maycock, A. C., Misios, S., Rodger, C. J., Scaife, A. A., Seppälä, A., Shangguan, M., Sinnhuber, M., Tourpali, K., Usoskin, I., Van De Kamp, M., Verronen, P. T., and Versick, S.: Solar forcing for CMIP6 (v3.2), *Geoscientific Model Development*, 10, 2247–2302, <https://doi.org/10.5194/gmd-10-2247-2017>, 2017.

- 920 Matthews, H. D. and Solomon, S.: Irreversible does not mean unavoidable, *Science*, 340, 438–439, <https://doi.org/10.1126/science.1236372>, 2013.
- Mauritsen, T., Bader, J., Becker, T., Behrens, J., Bittner, M., Brokopf, R., Brovkin, V., Claussen, M., Crueger, T., Esch, M., Fast, I., Fiedler, S., Fläschner, D., Gayler, V., Giorgetta, M., Goll, D. S., Haak, H., Hagemann, S., Hedemann, C., Hohenegger, C., Ilyina, T., Jahns, T., Jimenéz-de-la Cuesta, D., Jungclaus, J., Kleinen, T., Kloster, S., Kracher, D., Kinne, S., Kleberg, D., Lasslop, G., Kornbluh, L., Marotzke, 925 J., Matei, D., Meraner, K., Mikolajewicz, U., Modali, K., Möbis, B., Müller, W. A., Nabel, J. E., Nam, C. C., Notz, D., Nyawira, S. S., Paulsen, H., Peters, K., Pincus, R., Pohlmann, H., Pongratz, J., Popp, M., Raddatz, T. J., Rast, S., Redler, R., Reick, C. H., Rohrschneider, T., Schemann, V., Schmidt, H., Schnur, R., Schulzweida, U., Six, K. D., Stein, L., Stemmler, I., Stevens, B., von Storch, J. S., Tian, F., Voigt, A., Vrese, P., Wieners, K. H., Wilkenskield, S., Winkler, A., and Roeckner, E.: Developments in the MPI-M Earth System Model version 1.2 (MPI-ESM1.2) and Its Response to Increasing CO₂, *Journal of Advances in Modeling Earth Systems*, 11, 998–1038, 930 <https://doi.org/10.1029/2018MS001400>, 2019.
- Medlyn, B. E., Duursma, R. a., Eamus, D., Ellsworth, D. S., Prentice, I. C., Barton, C. V. M., Crous, K. Y., De Angelis, P., Freeman, M., and Wingate, L.: Reconciling the optimal and empirical approaches to modelling stomatal conductance, *Global Change Biology*, 17, 2134–2144, <https://doi.org/10.1111/j.1365-2486.2010.02375.x>, <http://doi.wiley.com/10.1111/j.1365-2486.2010.02375.x>, 2011.
- Meinshausen, M., Vogel, E., Nauels, A., Lorbacher, K., Meinshausen, N., Etheridge, D., Fraser, P., Montzka, S., Rayner, P., Trudinger, C., 935 Krummel, P., Beyerle, U., Cannadell, J., Daniel, J., Enting, I., Law, R., O'Doherty, S., Prinn, R., Reimann, S., Rubino, M., Velders, G., Vollmer, M., and Weiss, R.: Historical greenhouse gas concentrations, *Geoscientific Model Development Discussions*, 1, 1–122, <https://doi.org/10.5194/gmd-2016-169>, 2016.
- Melack, J. M. and Hess, L. L.: Remote Sensing of the Distribution and Extent of Wetlands in the Amazon Basin, in: *Amazonian floodplain forests: Ecophysiology, ecology, biodiversity and sustainable management*, edited by Junk, W. J., Piedade, M. T., Wittmann, 940 F., Schöngart, J., and Parolin, P., pp. 43–59, Springer, https://doi.org/10.1007/978-90-481-8725-6_3, http://link.springer.com/10.1007/978-90-481-8725-6_3, 2010.
- Mengis, N., Keller, D. P., MacDougall, A. H., Eby, M., Wright, N., Meissner, K. J., Oschlies, A., Schmittner, A., MacIsaac, A. J., Matthews, H. D., and Zickfeld, K.: Evaluation of the University of Victoria Earth System Climate Model version 2.10 (UVic ESCM 2.10), *Geoscientific Model Development*, 13, 4183–4204, <https://doi.org/10.5194/gmd-13-4183-2020>, <https://gmd.copernicus.org/articles/13/4183/2020/>, 945 2020.
- Milliman, J. D. and Droxler, A. W.: Neritic and pelagic carbonate sedimentation in the marine environment: Ignorance is not bliss, *Geologische Rundschau*, 85, 496–504, <https://doi.org/10.1007/BF02369004>, 1996.
- Moore, C. M., Mills, M. M., Arrigo, K. R., Berman-Frank, I., Bopp, L., Boyd, P. W., Galbraith, E. D., Geider, R. J., Guieu, C., Jaccard, S. L., Jickells, T. D., La Roche, J., Lenton, T. M., Mahowald, N. M., Marañón, E., Marinov, I., Moore, J. K., Nakatsuka, T., Oschlies, A., Saito, 950 M. A., Thingstad, T. F., Tsuda, A., and Ulloa, O.: Processes and patterns of oceanic nutrient limitation, *Nature Geoscience*, 6, 701–710, <https://doi.org/10.1038/ngeo1765>, 2013.
- Müller, S. A., Joos, F., Plattner, G. K., Edwards, N. R., and Stocker, T. F.: Modeled natural and excess radiocarbon: Sensitivities to the gas exchange formulation and ocean transport strength, *Global Biogeochemical Cycles*, 22, 1–14, <https://doi.org/10.1029/2007GB003065>, 2008.
- 955 Munhoven, G.: Glacial - Interglacial changes of continental weathering: Estimates of the related CO₂ and HCO₃⁻ flux variations and their uncertainties, *Global and Planetary Change*, 33, 155–176, [https://doi.org/10.1016/S0921-8181\(02\)00068-1](https://doi.org/10.1016/S0921-8181(02)00068-1), 2002.

- Munhoven, G.: Mathematics of the total alkalinity-pH equation-Pathway to robust and universal solution algorithms: The SolveSAPHE package v1.0.1, *Geoscientific Model Development*, 6, 1367–1388, <https://doi.org/10.5194/gmd-6-1367-2013>, 2013.
- 960 Munhoven, G. and François, L. M.: Glacial-interglacial changes in continental weathering~: Possible implications for atmospheric CO₂, in: *Carbon Cycling in the Glacial Ocean~: Constraints on the Ocean's Role in Global Change*, edited by Zahn, R., Pedersen, T. F., Kaminski, M. A., and Labeyrie, L., vol. 17 of *NATO ASI Series~I~: Global Environmental Change*, pp. 39–58, Springer-Verlag, Berlin, https://doi.org/10.1007/978-3-642-78737-9_3, 1994.
- Oleson, K. W., Lawrence, D. M., Gordon, B., Flanner, M. G., Kluzek, E., Peter, J., Levis, S., Swenson, S. C., Thornton, E., and Feddema, J.: Technical description of version 4.5 of the Community Land Model (CLM), NCAR/TN-478+STR NCAR Technical Note, p. 266, <https://doi.org/10.5065/D6RR1W7M>, <http://citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.172.7769%5Cnpapers3://publication/uuid/E8E12D50-5C26-4DF4-A67C-753D8AC5D002>, 2010.
- 965 Olsen, A., Key, R. M., Van Heuven, S., Lauvset, S. K., Velo, A., Lin, X., Schirnack, C., Kozyr, A., Tanhua, T., Hoppema, M., Jutterström, S., Steinfeldt, R., Jeansson, E., Ishii, M., Pérez, F. F., and Suzuki, T.: The global ocean data analysis project version 2 (GLODAPv2) - An internally consistent data product for the world ocean, *Earth System Science Data*, 8, 297–323, <https://doi.org/10.5194/essd-8-297-2016>, 970 2016.
- Orr, J. C., Najjar, R. G., Aumont, O., Bopp, L., Bullister, J. L., Danabasoglu, G., Doney, S. C., Dunne, J. P., Dutay, J. C., Graven, H., Griffies, S. M., John, J. G., Joos, F., Levin, I., Lindsay, K., Matear, R. J., McKinley, G. A., Mouchet, A., Oschlies, A., Romanou, A., Schlitzer, R., Tagliabue, A., Tanhua, T., and Yool, A.: Biogeochemical protocols and diagnostics for the CMIP6 Ocean Model Intercomparison Project (OMIP), *Geoscientific Model Development*, 10, 2169–2199, <https://doi.org/10.5194/gmd-10-2169-2017>, 2017.
- 975 Papa, F., Prigent, C., Aires, F., Jimenez, C., Rossow, W. B., and Matthews, E.: Interannual variability of surface water extent at the global scale, 1993–2004, *Journal of Geophysical Research*, 115, D12 111, <https://doi.org/10.1029/2009JD012674>, <http://doi.wiley.com/10.1029/2009JD012674>, 2010.
- Parekh, P., Follows, M. J., and Boyle, E.: Modeling the global ocean iron cycle, *Global Biogeochemical Cycles*, 18, n/a–n/a, <https://doi.org/10.1029/2003GB002061>, <http://doi.wiley.com/10.1029/2003GB002061>, 2004.
- 980 Paulsen, H., Ilyina, T., Six, K. D., and Stemmler, I.: Incorporating a prognostic representation of marine nitrogen fixers into the global ocean biogeochemical model HAMOCC, *Journal of Advances in Modeling Earth Systems*, 9, 438–464, <https://doi.org/10.1002/2016MS000737>, 2017.
- Petit, J. R., Jouzel, J., Raynaud, D., Barkov, N. I., Barnola, J.-M., Basile, I., Bender, M., Chappellaz, J., Davis, M., Delaygue, G., Delmotte, M., Kotlyakov, V. M., Legrand, M., Lipenkov, V. Y., Lorius, C., Pépin, L., Ritz, C., Saltzman, E., and Stievenard, M.: Climate and atmospheric history of the past 420,000 years from the Vostok ice core, Antarctica, *Nature*, 399, 429–436, <https://doi.org/10.1038/20859>, <http://www.nature.com/nature/journal/v399/n6735/abs/399429a0.html><https://www.nature.com/articles/20859>, 1999.
- 985 Prather, M., Flato, G., Friedlingstein, P., Jones, C., Lamarque, J., Liao, H., and Rasch, P.: IPCC, 2013: Annex II: Climate System Scenario Table, in: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Stocker, T., Qin, D., Plattner, G.-K., Tignor, M., Allen, S., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgale, P., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
- 990 Prentice, I. C., Dong, N., Gleason, S. M., Maire, V., and Wright, I. J.: Balancing the costs of carbon gain and water transport: testing a new theoretical framework for plant functional ecology., *Ecology letters*, 17, 82–91, <https://doi.org/10.1111/ele.12211>, <http://www.ncbi.nlm.nih.gov/pubmed/24215231>, 2014.

- 995 Prigent, C., Papa, F., Aires, F., Rossow, W. B., and Matthews, E.: Global inundation dynamics inferred from multiple satellite observations, 1993–2000, *Journal of Geophysical Research*, 112, D12 107, <https://doi.org/10.1029/2006JD007847>, <http://doi.wiley.com/10.1029/2006JD007847>, 2007.
- Raymo, M. and Ruddiman, W.: Tectonic Forcing of Late Cenozoic Climate, *Nature*, 359, 117–122, 1992.
- 1000 Regnier, P., Friedlingstein, P., Ciais, P., Mackenzie, F. T., Gruber, N., Janssens, I. A., Laruelle, G. G., Lauerwald, R., Luysaert, S., Andersson, A. J., Arndt, S., Arnosti, C., Borges, A. V., Dale, A. W., Gallego-Sala, A., Godd ris, Y., Goossens, N., Hartmann, J., Heinze, C., Ilyina, T., Joos, F., Larowe, D. E., Leifeld, J., Meysman, F. J., Munhoven, G., Raymond, P. A., Spahni, R., Suntharalingam, P., and Thullner, M.: Anthropogenic perturbation of the carbon fluxes from land to ocean, *Nature Geoscience*, 6, 597–607, <https://doi.org/10.1038/ngeo1830>, 2013.
- 1005 Ridgwell, A., Hargreaves, J. C., Edwards, N. R., Annan, J. D., Lenton, T. M., Marsh, R., Yool, A., and Watson, A.: Marine geochemical data assimilation in an efficient Earth System Model of global biogeochemical cycling, *Biogeosciences*, 4, 87–104, <https://doi.org/10.5194/bg-4-87-2007>, <https://bg.copernicus.org/articles/4/87/2007/>, 2007.
- Riley, W. J., Subin, Z. M., Lawrence, D. M., Swenson, S. C., Torn, M. S., Meng, L., Mahowald, N. M., and Hess, P.: Barriers to predicting changes in global terrestrial methane fluxes: Analyses using CLM4Me, a methane biogeochemistry model integrated in CESM, *Biogeosciences*, 8, 1925–1953, <https://doi.org/10.5194/bg-8-1925-2011>, 2011.
- 1010 Romero-Mujalli, G., Hartmann, J., and B rker, J.: Temperature and CO₂ dependency of global carbonate weathering fluxes – Implications for future carbonate weathering research, *Chemical Geology*, 527, 118 874, <https://doi.org/10.1016/j.chemgeo.2018.08.010>, <https://doi.org/10.1016/j.chemgeo.2018.08.010>, 2019.
- Sanderman, J., Hengl, T., and Fiske, G. J.: Soil carbon debt of 12,000 years of human land use, *Proceedings of the National Academy of Sciences of the United States of America*, 114, 9575–9580, <https://doi.org/10.1073/pnas.1706103114>, 2017.
- 1015 Saunois, M., R. Stavert, A., Poulter, B., Bousquet, P., G. Canadell, J., B. Jackson, R., A. Raymond, P., J. Dlugokencky, E., Houweling, S., K. Patra, P., Ciais, P., K. Arora, V., Bastviken, D., Bergamaschi, P., R. Blake, D., Brailsford, G., Bruhwiler, L., M. Carlson, K., Carrol, M., Castaldi, S., Chandra, N., Crevoisier, C., M. Crill, P., Covey, K., L. Curry, C., Etiope, G., Frankenberg, C., Gedney, N., I. Hegglin, M., H glund-Isaksson, L., Hugelius, G., Ishizawa, M., Ito, A., Janssens-Maenhout, G., M. Jensen, K., Joos, F., Kleinen, T., B. Krummel, P., L. Langenfelds, R., G. Laruelle, G., Liu, L., MacHida, T., Maksyutov, S., C. McDonald, K., McNorton, J., A. Miller, P., R. Melton, J., Morino, I., M ller, J., Murguia-Flores, F., Naik, V., Niwa, Y., Noce, S., O’Doherty, S., J. Parker, R., Peng, C., Peng, S., P. Peters, G., Prigent, C., 1020 Prinn, R., Ramonet, M., Regnier, P., J. Riley, W., A. Rosentretter, J., Segers, A., J. Simpson, I., Shi, H., J. Smith, S., Paul Steele, L., F. Thornton, B., Tian, H., Tohjima, Y., N. Tubiello, F., Tsuruta, A., Viovy, N., Voulgarakis, A., S. Weber, T., Van Weele, M., R. Van Der Werf, G., F. Weiss, R., Worthy, D., Wunch, D., Yin, Y., Yoshida, Y., Zhang, W., Zhang, Z., Zhao, Y., Zheng, B., Zhu, Q., Zhu, Q., and Zhuang, Q.: The global methane budget 2000–2017, *Earth System Science Data*, 12, 1561–1623, <https://doi.org/10.5194/essd-12-1561-2020>, 2020.
- 1025 Schaphoff, S., von Bloh, W., Rammig, A., Thonicke, K., Biemans, H., Forkel, M., Gerten, D., Heinke, J., J germeyr, J., Knauer, J., Langerwisch, F., Lucht, W., M ller, C., Rolinski, S., and Waha, K.: LPJmL4 – a dynamic global vegetation model with managed land – Part 1: Model description, *Geoscientific Model Development*, 11, 1343–1375, <https://doi.org/10.5194/gmd-11-1343-2018>, <https://www.geosci-model-dev-discuss.net/gmd-2017-146/><https://gmd.copernicus.org/articles/11/1343/2018/>, 2018.
- Schubert, B. A. and Jahren, A. H.: Incorporating the effects of photorespiration into terrestrial paleoclimate reconstruction, *Earth-Science Reviews*, 177, 637–642, <https://doi.org/10.1016/j.earscirev.2017.12.008>, <https://doi.org/10.1016/j.earscirev.2017.12.008>, 2018.
- 1030 S f rian, R., Berthet, S., Yool, A., Palmi ri, J., Bopp, L., Tagliabue, A., Kwiatkowski, L., Aumont, O., Christian, J., Dunne, J., Gehlen, M., Ilyina, T., John, J. G., Li, H., Long, M. C., Luo, J. Y., Nakano, H., Romanou, A., Schwinger, J., Stock, C., Santana-Falc n, Y., Takano, Y.,

- Tjiputra, J., Tsujino, H., Watanabe, M., Wu, T., Wu, F., and Yamamoto, A.: Tracking Improvement in Simulated Marine Biogeochemistry Between CMIP5 and CMIP6, *Current Climate Change Reports*, 6, 95–119, <https://doi.org/10.1007/s40641-020-00160-0>, 2020.
- 1035 Segschneider, J. and Bendtsen, J.: Temperature-dependent remineralization in a warming ocean increases surface pCO₂ through changes in marine ecosystem composition, *Global Biogeochemical Cycles*, 27, 1214–1225, <https://doi.org/10.1002/2013GB004684>, 2013.
- Seiter, K., Hensen, C., Schröter, J., and Zabel, M.: Organic carbon content in surface sediments—defining regional provinces, *Deep Sea Research Part I: Oceanographic Research Papers*, 51, 2001–2026, <https://doi.org/10.1016/j.dsr.2004.06.014>, <https://linkinghub.elsevier.com/retrieve/pii/S0967063704001384>, 2004.
- 1040 Sitch, S., Smith, B., Prentice, I. C., Arneth, A., Bondeau, A., Cramer, W., Kaplan, J. O., Levis, S., Lucht, W., Sykes, M. T., Thonicke, K., and Venevsky, S.: Evaluation of ecosystem dynamics, plant geography and terrestrial carbon cycling in the LPJ dynamic global vegetation model, *Global Change Biology*, 9, 161–185, <https://doi.org/10.1046/j.1365-2486.2003.00569.x>, <http://doi.wiley.com/10.1046/j.1365-2486.2003.00569.x>, 2003.
- Six, K. D. and Maier-Reimer, E.: Effects of plankton dynamics on seasonal carbon fluxes in an ocean general circulation model, *Global Biogeochemical Cycles*, 10, 559–583, <https://doi.org/10.1029/96GB02561>, <http://doi.wiley.com/10.1029/96GB02561>, 1996.
- 1045 Stocker, B. D., Roth, R., Joos, F., Spahni, R., Steinacher, M., Zaehle, S., Bouwman, L., and Prentice, I. C.: Multiple greenhouse-gas feedbacks from the land biosphere under future climate change scenarios, *Nature Climate Change*, 3, 666–672, <https://doi.org/10.1038/nclimate1864>, <http://www.nature.com/doi/10.1038/nclimate1864>, 2013.
- Tagliabue, A., Mtshali, T., Aumont, O., Bowie, A. R., Klunder, M. B., Roychoudhury, A. N., and Swart, S.: A global compilation of dissolved iron measurements: Focus on distributions and processes in the Southern Ocean, *Biogeosciences*, 9, 2333–2349, <https://doi.org/10.5194/bg-9-2333-2012>, 2012.
- 1050 Tagliabue, A., Aumont, O., DeAth, R., Dunne, J., Dutkiewicz, S., Galbraith, E., Misumi, K., Moore, J., Ridgeway, A., Sherman, E., Stock, C., Vichi, M., Volker, C., and Yool, A.: How well do global ocean biogeochemistry models simulate dissolved iron distributions?, *Global Biogeochemical Cycles*, 30, 149–174, <https://doi.org/10.1002/2015GB005289>.Received, 2016.
- Takahashi, T., Broecker, W. S., and Langer, S.: Redfield ratio based on chemical data from isopycnal surfaces., *Journal of Geophysical Research*, 90, 6907–6924, <https://doi.org/10.1029/JC090iC04p06907>, 1985.
- 1055 Tarnocai, C., Canadell, J. G., Schuur, E. a. G., Kuhry, P., Mazhitova, G., and Zimov, S.: Soil organic carbon pools in the northern circumpolar permafrost region, *Global Biogeochemical Cycles*, 23, n/a–n/a, <https://doi.org/10.1029/2008GB003327>, <http://doi.wiley.com/10.1029/2008GB003327>, 2009.
- Thornton, P. E. and Zimmermann, N. E.: An improved canopy integration scheme for a Land Surface Model with prognostic canopy structure, *Journal of Climate*, 20, 3902–3923, <https://doi.org/10.1175/JCLI4222.1>, 2007.
- 1060 Tréguer, P. J. and De La Rocha, C. L.: The World Ocean Silica Cycle, *Annual Review of Marine Science*, 5, 477–501, <https://doi.org/10.1146/annurev-marine-121211-172346>, <https://www.annualreviews.org/doi/10.1146/annurev-marine-121211-172346>, 2013.
- Tréguer, P. J., Sutton, J. N., Brzezinski, M., Charette, M. A., Devries, T., Dutkiewicz, S., Ehlert, C., Hawkings, J., Leynaert, A., Liu, S. M., 1065 Monferrer, N. L., López-Acosta, M., Maldonado, M., Rahman, S., Ran, L., and Rouxel, O.: Reviews and syntheses: The biogeochemical cycle of silicon in the modern ocean, *Biogeosciences*, 18, 1269–1289, <https://doi.org/10.5194/bg-18-1269-2021>, 2021.
- Tschumi, T., Joos, F., Gehlen, M., and Heinze, C.: Deep ocean ventilation, carbon isotopes, marine sedimentation and the deglacial CO₂ rise, *Climate of the Past*, 7, 771–800, <https://doi.org/10.5194/cp-7-771-2011>, 2011.

- Ubierna, N. and Farquhar, G. D.: Advances in measurements and models of photosynthetic carbon isotope discrimination in C3 plants, *Plant, Cell and Environment*, 37, 1494–1498, <https://doi.org/10.1111/pce.12346>, 2014.
- 1070 Varney, R. M., Chadburn, S. E., Burke, E. J., and Cox, P. M.: Evaluation of soil carbon simulation in CMIP6 Earth system models, *Biogeosciences*, 19, 4671–4704, <https://doi.org/10.5194/bg-19-4671-2022>, 2022.
- Willeit, M. and Ganopolski, A.: PALADYN v1.0, a comprehensive land surface-vegetation-carbon cycle model of intermediate complexity, *Geoscientific Model Development*, 9, 3817–3857, <https://doi.org/10.5194/gmd-9-3817-2016>, 2016.
- 1075 Willeit, M., Ganopolski, A., Calov, R., and Brovkin, V.: Mid-Pleistocene transition in glacial cycles explained by declining CO₂ and regolith removal, *Science Advances*, 5, 1–9, <https://doi.org/10.1126/sciadv.aav7337>, 2019.
- Willeit, M., Ganopolski, A., Robinson, A., and Edwards, N. R.: The Earth system model CLIMBER-X v1.0 – Part 1: Climate model description and validation, *Geoscientific Model Development*, 15, 5905–5948, <https://doi.org/10.5194/gmd-15-5905-2022>, <https://gmd.copernicus.org/articles/15/5905/2022/>, 2022.
- 1080 Willey, D. A., Fine, R. A., Sonnerup, R. E., Bullister, J. L., Smethie, W. M., and Warner, M. J.: Global oceanic chlorofluorocarbon inventory, *Geophysical Research Letters*, 31, <https://doi.org/10.1029/2003GL018816>, 2004.
- Yang, K., Koike, T., Ishikawa, H., Kim, J., Li, X., Liu, H., Liu, S., Ma, Y., and Wang, J.: Turbulent flux transfer over bare-soil surfaces: Characteristics and parameterization, *Journal of Applied Meteorology and Climatology*, 47, 276–290, <https://doi.org/10.1175/2007JAMC1547.1>, 2008.
- 1085 Yu, Z., Loisel, J., Brosseau, D. P., Beilman, D. W., and Hunt, S. J.: Global peatland dynamics since the Last Glacial Maximum, *Geophysical Research Letters*, 37, n/a–n/a, <https://doi.org/10.1029/2010GL043584>, <http://doi.wiley.com/10.1029/2010GL043584>, 2010.
- Zehr, J. P. and Capone, D. G.: Changing perspectives in marine nitrogen fixation, *Science*, 368, <https://doi.org/10.1126/science.aay9514>, 2020.
- Zender, C. S., Newman, D., and Torres, O.: Spatial heterogeneity in aeolian erodibility: Uniform, topographic, geomorphic, and hydrologic hypotheses, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/10.1029/2002jd003039>, 2003.
- 1090 Zheng, W., Wei, H., Wang, Z., Zeng, X., Meng, J., Ek, M., Mitchell, K., and Derber, J.: Improvement of daytime land surface skin temperature over arid regions in the NCEP GFS model and its impact on satellite data assimilation, *Journal of Geophysical Research Atmospheres*, 117, 1–14, <https://doi.org/10.1029/2011JD015901>, 2012.
- Zickfeld, K., Eby, M., Damon Matthews, H., Schmittner, A., and Weaver, A. J.: Nonlinearity of carbon cycle feedbacks, *Journal of Climate*, 1095 24, 4255–4275, <https://doi.org/10.1175/2011JCLI3898.1>, 2011.
- Zilitinkevich, S.: Non-local turbulent transport: pollution dispersion aspects of coherent structure of convective flows, *International Conference on Air Pollution - Proceedings*, 1, 53–60, <https://doi.org/10.2495/air950071>, 1995.