CLM5-FruitTree: A new sub-model for deciduous fruit trees in the **Community Land Model (CLM5)**

Olga Dombrowski¹, Cosimo Brogi¹, Harrie-Jan Hendricks Franssen¹, Damiano Zanotelli², Heye Bogena¹

¹Agrosphere (IBG-3), Forschungszentrum Jülich GmbH, Jülich, 52425, Germany 5

²Faculty of Science and Technology, Free University of Bolzano, Bolzano, 39100, Italy

Correspondence to: Olga Dombrowski (o.dombrowski@fz-juelich.de)

Abstract. The inclusion of perennial, woody crops in land surface models is crucial to address their role in carbon (C) sequestration, food production, and water requirements under climate change. To help quantifying the biogeochemical and biogeophysical processes associated with these agro-ecosystems, we developed and tested a new sub-model, CLM5-FruitTree, for deciduous fruit orchards within the framework of the Community Land Model version 5 (CLM5). The model development 10 included (1) a new perennial crop phenology description, (2) an adapted C and nitrogen allocation scheme, considering both storage and photosynthetic growth of annual and perennial plant organs, (3) typical management practices associated with fruit orchards, and (4) the parameterization of an apple plant functional type. CLM5-FruitTree was tested using extensive field measurements from an apple orchard in South Tyrol, Italy. Growth and partitioning of biomass to the individual plant components was well represented by CLM5-FruitTree and average yield was predicted within 2.3 % of the observed values despite low simulated inter-annual variability compared to observations. The simulated seasonal course of C, energy, and water

- fluxes was in good agreement with the eddy covariance (EC) measurements owing to the accurate representation of the prolonged growing season and typical leaf area development of the orchard. We found that gross primary production, net radiation, and latent heat flux were highly correlated (r > 0.94) with EC measurements and showed little bias ($< \pm 5$ %). Simulated respiration components, sensible heat, and ground heat flux were less consistent with observations. This was 20
- attributed to simplifications in the orchard structure and to the presence of additional management practices that are not yet represented in CLM5-FruitTree. Finally, the results suggested that the representation of microbial and autotrophic respiration, and energy partitioning in complex, heterogeneous discontinuous canopies in CLM5 requires further attention. The new submodel CLM5-FruitTree improved the representation of agricultural systems in CLM5 and can be used to study land surface

processes in fruit orchards at the local, regional or larger scale. 25

15

Kommentiert [m1]: Name of sub-model was changed to CLM5-FruitTree throughout the manuscript in response to a comment from reviewer 2

1 Introduction

Orchards and other perennial fruit crops are a major component of the global agricultural production with significant coverage and yield in China, the United States, south-western Africa, and some parts of Europe (Fao, 2021). In the European region, perennial crops are a key economic element of Mediterranean agro-ecosystems as they provide 45 % of the local agricultural output (Lobianco and Roberto, 2006). Apples are the most important fruit tree crop as one third of European orchards is devoted to their production. With a coverage of 984²₂509 ha, they provide a yearly harvest of over 17 million tons which is one fifth of the overall European fruit production in terms of output value (Fao, 2021).

- 35 In contrast to annual crops, fruit trees can be productive for several decades before rotation is needed. Their prolonged growing season, standing biomass, and low respiratory losses can support carbon (C) storage and promote higher <u>carbon-C</u> use efficiencies (Wünsche and Lakso, 2000; Zanotelli et al., 2013). The transport of C stored in biomass into the soil in addition to reduced soil tillage and disturbances under fruit orchards compared to annual crops further promote C sequestration (Bwalya, 2012; Wu et al., 2012; Ledo et al., 2020). The FAO has therefore suggested perennial agriculture as a possible measure to
- 40 mitigate climate change and enhance food security (Glover et al., 2010), and many studies have recently investigated this potential for various fruit orchards (Wu et al., 2012; Scandellari et al., 2016; Hammad et al., 2020; Yasin et al., 2021). The study of water and irrigation requirements in fruit orchards has become another field of intense research due to the need for a more resilient agriculture in the context of climate change and water supply shortages (Maestre-Valero et al., 2017; El Jaouhari et al., 2018; O'connell and Scalisi, 2019; Segovia-Cardozo et al., 2022). In order to answer questions related to C sequestration,
- 45 water requirements, and sustainable food production of fruit orchards, a better understanding of the related ecosystem processes is vital (Fader et al., 2015).

Models with a comprehensive description of the carbon, water, and energy fluxes, such as global land surface models (LSMs), are a powerful tool to explore complex ecosystems like the abovementioned fruit orchards. The use of LSMs was recently extended to not only model the processes at the land–atmosphere interface, but also to study the response of ecosystems and

- 50 water resources to climate change (Prentice et al., 2015; Fisher and Koven, 2020; Blyth et al., 2021). To quantify these effects, LSMs need to represent a wide range of land use and vegetation types. However, most LSMs consider only perennials such as deciduous and coniferous trees, as well as major annual crops such as wheat, soy, or maize (Lawrence et al., 2018). Recently, some LSMs additionally included bioenergy crops (Schaphoff et al., 2018), while others group crops into a few generic crop types (Noilhan and Mahfouf, 1996; Krinner et al., 2005; Balsamo et al., 2009). Despite their significance, perennial crops,
- 55 such as fruit trees, are rarely considered in LSMs and attempts of including them in global and regional modelling environments are scarce (Fader et al., 2015; Cheng et al., 2020). An example of such an attempt is the inclusion of agricultural trees (e.g., grapes, cotton, and apple trees) in the Lund-Potsdam-Jena managed Land (LPJmL) model to improve the representation of Mediterranean agro-ecosystems (Fader et al., 2015). Here, agricultural trees were modelled as small trees and fruit harvest was determined as the product of a plant specific harvest index and the net primary productivity (NPP). Other authors parameterized
- 60 oil palm trees, a perennial evergreen crop, in the Community Land Model (CLM) version 4.5 (Fan et al., 2015). Palm trees

were represented by a new phenology where large palm leaves with fruit bunches emerge successively, leaves are pruned regularly, and harvest occurs once a month. Recently, two perennial grasses for energy production were parameterized in the latest version of the model, CLM5 (Cheng et al., 2020). Parameters for bioenergy crops were tuned using sensitivity analysis and observations, while harvest was represented by removing around 70 % of the aboveground biomass.

- 65 While the above-mentioned studies describe some common features of perennial plants, they do not, or only partially, represent the seasonal deciduous phenology of fruit trees or the explicit modelling of fruit growth. Furthermore, key aspects such as C reserve <u>deposition accumulation</u> and mobilization in the following spring are generally not considered, possibly due to necessary simplifications or because the drivers of these processes are still not fully understood (Le Roux et al., 2001; Neumann, 2020). The absence of perennial crops in land surface models introduces a significant bias in the representation of
- 70 biogeophysical and biogeochemical processes in agro-ecosystems where this type of cultivation is prevalent. As a result, the response to climate change in terms of C sequestration, water requirements, or food production cannot be assessed adequately in regions such as the Mediterranean, where perennial, woody crops are very common and play a vital role for food security and economy (Fader et al., 2015; Lobianco and Roberto, 2006).

Although deciduous fruit trees share certain characteristics with natural vegetation and annual crops in LSMs such as CLM5, several particularities in their growth dynamics and management practices still prevent a meaningful simulation using currently

- available representations of vegetation. In this study, we therefore provide CLM5 with the ability of modelling perennial fruit trees and the associated processes. For this purpose, we developed a new sub-model named CLM5-FruitTree within the existing model framework of CLM5. CLM5-FruitTree combines elements of the broadleaf deciduous tree subroutine such as growth and C turnover of woody components, with distinctive phenological stages and a harvestable organ similar to the annual crop subroutine. We first describe the model conceptualization including the new phenology, carbon and nitrogen (CN) allocation,
- and management options. We further demonstrate the applicability of CLM5-FruitTree by parameterizing a new apple plant functional type (PFT). Finally we evaluate and discuss the model performance using extensive field data from an apple orchard in South Tyrol, Italy.

2 Methods

85 2.1 Vegetation characterizations in CLM5

The latest version of the Community Land Model, CLM5, simulates the exchange of water, energy, C, and nitrogen (N) between land and atmosphere as well as their storage and transport on the land surface and in the subsurface, driven by climate variability and modulated by soil and vegetation states and characteristics. The land surface in CLM5 is characterized by one of five land units namely glacier, lake, urban, vegetated, and crop. These units are further divided to capture the variability in

90 soil, vegetation, and management options (i.e., irrigated or non-irrigated). Compared to previous model versions, CLM5 features various improvements in the representation of land use and vegetation modelling such as plant CN cycling, soil and plant hydrology and crop modelling (Lawrence et al., 2018; Lombardozzi et al., 2020).

Kommentiert [m2]: Section 2.1 was restructured in response to a comment from reviewer 1

Kommentiert [m3]: Reference to Lombardozzi was added in response to a comment of reviewer 2 Many of the C and N cycle components of CLM5 were originally derived from the Biome BioGeochemical Cycles (Biome-BGC) model (Thornton et al., 2002). Here, vegetation is represented conceptually by three different plant C and N pools that

- 95 are maintained separately for the individual plant organs (leaf, live/dead stem, fine root, live/dead coarse root, and grain). The storage pools represent C and N reserves, the transfer pools serve as intermediate pools to separate fluxes in and out of the storage pools, and the display pools represent the actual growth of a given organ (Fig. 1). C made available through photosynthesis is first used to support maintenance respiration of live organs based on organ N content, temperature, and a constant base rate as proposed by Atkin et al. (2015). Dead stem and dead coarse root components are assumed to consist of
- 100 dead xylem cells, without metabolic function (no C cost for maintenance). The remaining C can then be allocated to the growth of new tissue considering associated growth respiration costs. Maintenance respiration together with growth respiration and C cost of N uptake from the soil comprise the autotrophic respiration component (R_a) in CLM5. Plant material reaching the end of its lifespan feeds into different litter pools from where it progressively decomposes to soil organic matter under C losses through heterotrophic respiration (R_b).
- 105 For the simulation of fruit orchards, a module for perennial deciduous crops is needed which is currently missing in CLM5. Such module must account for the perennial deciduous nature of fruit trees, which is similar to the existing representation of broadleaf deciduous trees (BDT) included in Biome-BGC but with differences in phenological triggers, vegetation structure, and C partitioning. In addition, it must represent growth and harvest of the fruits and typical management practices, of which some are already conceptualized in the prognostic Biogeochemistry Crop Module (BGC-crop) while others are not yet 110 implemented.

The inclusion of the Biome BGC (Biome BioGeochemical Cycles) model into CLM4 allowed the simulation of 15 different PFTs of natural, woody and non-woody, vegetation including broadleaf deciduous trees (BDT). The latter <u>The</u> is described by an algorithm for the seasonal phenology of <u>deciduous treesBDT</u> that controls changes in phenology, as well as the C and N fluxes between different plant and litter pools. Here, initial leaf <u>developmentonset</u> and <u>senescenceoffset periods that</u> mark the

- 115 beginning and end of a growing season based on temperature and day length thresholds. Once a new onset growth period is initiated, C and corresponding N fluxes accumulated in the previous season, occur out of the storage pools into the transfer pools, (intermediate pool to separate fluxes in and out of the storage pool) from where they are gradually transferred sent to the displayed growth-pools (Fig. 1). All plant organs are maintained separately in each of the three pools. During the active growth period, C and corresponding N storage pools of the individual plant organs are replenished based on fixed
- 120 stoichiometric relationships and specified C:N ratios of each plant organ. During At leaf offsetsenescence, C and N pools feed the litter or coarse woody debris pool except for live stem and live coarse roots that are mostly retained as structural woody tissue (dead stem and dead coarse roots).

In addition, CLM5 now features the Biogeochemistry Crop Module (BGC-crop), adopted from the prognostic crop module of the Agro-Ecosystem Integrated Biosphere Simulator (Agro-IBIS), <u>currently features which provides</u> eight different annual crop species with interactive crop management options (i.e., irrigation and fertilization). Another 23 currently inactive crop types

125

Formatiert: Tiefgestellt

Kommentiert [m4]: Reference to Lombardozzi was added in response to a comment of reviewer 2

allocation follow three phenological phases: (1) from planting to leaf emergence, (2) from leaf emergence to the start of grain fill, and (3) from grain fill to grain maturity and harvest, that are controlled by temperature and growing degree-day (GDD) thresholds. Different to natural vegetation, <u>crops have a grain pool representing the harvestable organ but no structural woody</u> <u>tissue. Furthermore</u>, all assimilates are directed to the displayed pools while the storage pools remain unused. At harvest, C

- and N from the grain pool are transferred to a grain product pool while a small amount is kept to reseed the crop in the following year. All remaining plant parts feed the litter cycle_-(Fig. 1). The reader is referred to Lombardozzi et al. (2020) <u>and the</u> technical documentation of CLM5 for a more detailed description of the <u>BDTvegetation</u> and crop <u>description representation</u> (Lawrence et al., 2018).
- 135 From the above description of the existing model vegetation componentsmodules, the following limitations for the application of CLM5 to deciduous fruit trees arise: (1) the current CLM-BGC-crop algorithm does not allow the simulation of perennial and/or woody crops; (2) the seasonal deciduousBDT phenology algorithm although describing some characteristics common to fruit trees, lacks the capability of simulating a harvestable organ, individualependent development of different plant parts, and the separation of growth from C reserves of the previous year and photosynthetic growth of active growth in the current
- 140 season; (3) typical management practices of fruit orchards such as transplanting of tree seedlings and pruning are currently not represented in CLM5; and (4) there is no parameterized fruit tree PFT in the default parameter set of CLM5.

2.2 Model conceptualization and technical implementation

130

To resolve the model limitations discussed in Sect. 2.1, we have developed a new sub-model CLM5-FruitTree to model the ecosystem processes and exchanges of energy and matter of deciduous fruit trees grown in commercial orchards systems with

- 145 a focus on the simulation of biomass growth and yieldas well as the associated ecosystem processes. More specifically, for the implementation of CLM5-FruitTree, we introduced a new phenology subroutine that describes the main phenological development of fruit trees and includes including triggers for seasonal orcharderop management optionspractices typical under organic or conventional production next to the existing subroutines of natural vegetation and crops.-_In addition, the CN allocation module as well as corresponding modules (including_C and N state and flux updates, vegetation structure, and
- 150 respiration) were modified to reproduce the growth dynamics of fruit trees and to model the fates of C and N in the orchard system. The sub-model development does not include any changes to the existing calculation schemes for radiative transfer or momentum, heat, and water fluxes to explicitly account for the discontinuous canopy structure of tree rows and vegetated or non-vegetated alleys in fruit orchards. In-row and between-row planting distances and alley vegetation are not defined directly. Instead, the orchard structure and the area covered by the canopy is accounted for through parameterization of the leaf and
- 155 stem area indices, the planting density, maximum canopy height, and aerodynamic parameters, similar to the implementation of crops and forest in CLM5. The modification of sub-routines and creation of new parameters followed CLM5 standards and the given model structure. Where necessary, new history and restart variables were added to restart simulations with bit-forbit continuity of the output (Lawrence et al., 2018). Furthermore, to test the new sub-model, an apple PFT was parameterized

Kommentiert [m5]: Section 2.2 was restructured and renamed in response to the comment of reviewer 1

5

and added to the active crops using one of the existing but currently inactive crop types in CLM5, type 35 and 36 (rain fed and

160 irrigated citrus).

CLM5-FruitTree combines characteristics of both BDT and annual crops to simulate a perennial woody crop with a harvestable organ <u>making use of the existing concepts of storage, transfer, and display vegetation pools described in section 2.1 (Fig. 1)</u>. Similar to the existing BDT phenology algorithm in CLM5, the fruit tree algorithm uses a perennial deciduous phenology with standing woody biomass and annual leaf shedding. During the active growth period however, the phenology and CN allocation

165 of vegetative and harvestable organs are described by distinct growth phases and are driven by a GDD summation similar to the crop phenology.

An orchard is established by transplanting small tree seedlings from nursery, a typical planting method for this type of cultivation (Wheaton et al., 1990; Corelli-Grappadelli and Marini, 2008). Once planted, the orchard remains productive according to a user-defined lifespan which, depending on fruit tree type and production system, typically ranges between for

- 170 2105 and 30–40 years (Demestihas et al., 2017; Cerutti et al., 2014) as defined by the user. The sub-model makes no specific assumptions on the rootstock, but the effect of different rootstocks in terms of tree height and rooting depth can be set by the user via the respective parameters, *ztopmx* and *root_dmx* (Table C1). In CLM5-FruitTree, both stored C and current photosynthesis contribute to the growth of the fruit tree, as leaf and shoot development at the beginning of a growing season utilize carbohydrate reserves and nitrogenous compounds that were accumulated during the previous season (Tromp, 1983;
- Oliveira and Priestley, 1988; Loescher et al., 1990). Deciduous fruit trees are dormant in winter and resume growth in spring after meeting species- and cultivar-specific chilling and heat requirements (Anderson et al., 1985; Faust et al., 1997; Zavalloni et al., 2006), which is represented in CLM5-FruitTree using the chilling and forcing model suggestedproposed by Cesaraccio et al. (2004). Early in the season, the canopy develops rapidly until it reaches maturity typically by midsummer, while leaf shedding occurs when temperatures drop in autumn (Kozlowski, 1992; Loescher et al., 1990; Lakso et al., 1990). Fruit trees
- 180 usually start flowering 3–4 weeks after bud burstbreak, which is not specifically represented by CLM5-FruitTree which instead assumes that fruit growth begins at the end of floweringafter full bloom (Lakso et al., 1999). The implementation of flowering to include effects of non-optimal pollination, frost during flowering, or hormonal processes affecting fruit set and development is outside of the scope of this development and of minor importance for large scale simulations and processes at ecosystem level that are typically the focus of land surface models such as CLM5. Consequently, CLM5-FruitTree does not produce
- 185 information on fruit size or number but only on total yield which we consider adequate for most applications of the sub-model development. Fruit growth is described by two stages, cell division and cell expansion that together form a sigmoid growth curve <u>observed</u> for many fruit tree species such as apple, pear, and orange (Corelli-Grappadelli and Lakso, 2002; Jackson, 2011).

In the following, the new developments to account for the distinct phenology, CN allocation, and management practices of a fruit orchard are described in more detail. Other biochemical and biophysical processes such as photosynthesis, water and litter cycles, and fixation and uptake of N were not modified except for minor adaptations to the retranslocation of N and respiration to enable the use of certain parts of these scripts for the fruit tree PFT. The technical implementation of some features of the Formatiert: Schriftart: Kursiv Formatiert: Schriftart: Kursiv new phenology routine (transplanting, pruning, harvest, and final rotation) was based on CLM-Palm, a previous model development for palm trees in CLM4.5 (Fan et al. 2015 and unpublished code). References where code elements were directly

195 reused or modified based on CLM-Palm are made in the published source code of CLM5-FruitTree (Dombrowski, 2022). Along with the new sub-model, an apple PFT was parameterized using one of the existing but thus far inactive crop types in CLM5, type 35 and 36 (rain fed and irrigated citrus). **Kommentiert [m6]:** Reference to CLM-Palm was added following an email exchange with Yuanchao Fan. Specific references are also added in the source code on zenodo.



Figure 1: Schematic of the main phenology and C allocation features of the broadleaf deciduous tree and annual crop representations in CLM5 as well as 200 the new sub-model CLM5-FruitTree. C pools within the dashed boxes are the individual components that make up the displayed C pool (the same components can be found for the other main plant pools: storage and transfer pools respectively). Carbon pools and fluxes in green were reused for CLM5-FruitTree while pools and fluxes in other-brown were modified or newly added.

2.2.1 Phenology

A new orchard lifecycle is initialized by transplanting seedlings at the beginning of the year during dormancy. An initial biomass of 5 gC m⁻² is assigned to leaf and fine root transfer pools, and 0.5 gC m⁻² is assigned to the dead stem pool. Each pool is also assigned the corresponding amount of N. AppleTree growth thereafter is described by six post-planting phenological stages, namely: (1) bud break, (2) fruit growth, (3) fruit ripening, (4) canopy maturity, (5) fruit maturity and harvest, and (6) start of leaf senescence (Fig. 2).

Bud break is predicted by a sequential model that first accumulates chill days followed by anti-chill days based on a predefined temperature threshold and chilling requirement (Cesaraccio et al., 2004). More information on the sequential model and the calibration of model parameters can be found in Appendix A. At bud break, a portion of the C in the storage pool of all plant components, except fruits, is transferred to the actively growing C pools over a period of 50 days as controlled by the newly added parameter *ndays_stor*. (Oliveira and Priestley, 1988; Loescher et al., 1990)Outside the dormant period, leaf and fruit development occur independentlyin parallel but with a time shift as fruit growth typically starts 4–5 weeks after bud break while canopy development continues until mid-season and leaf senescence does not occur until after the fruits are harvested (Wünsche and Lakso, 2000; Goldschmidt and Lakso, 2005) (Fig. 2).

The thermal thresholds to reach phases (2)–(5) (Appendix C) are defined as accumulated GDDs since bud break and can be adjusted by the user via the parameter file which applies to all parameters listed in Table C1 of the Appendix. GDDs are determined as the difference between the average daily air temperature and a base temperature of 4 °C with a maximum daily

- 220 increment of 26 °days (Eq. (1)). Different to the existing deciduous phenology, leaf senescence is triggered not by day length but by the drop of the daily mean temperature below a critical temperature threshold, in this case the base temperature. This approach was selected since many fruit trees that belong to the Rosaceae family (e.g., apple, pear, plum, and cherry) are unaffected by photoperiod and instead controlled by temperature (Heide and Prestrud, 2005). The last day of the leaf_- offsetsenescence period marks the beginning of dormancy. The new phenology subroutine of CLM5-FruitTree also controls
- 225 C reserve dynamics, stem and root turnover, and final rotation, which involves removing and replanting trees when the maximum orchard lifespan is reached.



250

Figure 2: Fruit tree phenological stages of (1) bud break at the end of dormancy, (2) the start of fruit growth, (3) fruit ripening, (4) canopy maturity, (5) harvest, and (6) the start of leaf senescence. The lengths of phenological stages (2)-(5) are determined by their respective growing degree-day thresholds (GDD) starting from bud break (GDD_{leaf}=0), while stage (6) is determined by a critical temperature threshold (T_{crit}). Coloured bars correspond to the time any plant organ is present on the field throughout a year.

2.2.2 Carbon and nitrogen allocation

CN allocation to the growth of new tissue (display pools) and to storage pools follows the phenological stages described in Sect. 2.2.1 (Fig. 2). A coupled CN allocation subroutine determines the fate of newly assimilated C from photosynthesis. A 235 user-defined initial biomass can be assigned to leaf and fine root transfer pools via the transplant parameter (Table C1), while additionally 10 % of this biomass is assigned to the dead stem pool to define an initial stem area index > 0. Each pool is also assigned the corresponding amount of N. Adjustments to this parameter have only little effect on the biomass growth and yield of the adult trees as the trees reach their maximum canopy height and develop their full LAI within the first couple of years after transplanting. Thereafter, the potential allocation to the different plant components is based on allocation coefficients and 240 allometric relationships between dead and live parts of stem and coarse root. Throughout the growing period until harvest, 5 % of the newly assimilated C is allocated to the storage pools, as defined by the fcur parameter₂. The remainder is allocated to the displayed C pools except for fruits where all allocated C is assigned to the displayed pool. For all other organs, the remaining C is also allocated to the displayed C pools. At bud break, a fraction of the C in the storage pool of all plant components, except fruits, is transferred to the actively growing C pools over a period that can be specified by the newly added 245 parameter *ndays_stor*. This is based on the assumption that resources are partially mobilized to support growth of new tissue (Oliveira and Priestley, 1988; Loescher et al., 1990). Lacking more specific knowledge of the exact fraction, the default of 0.5 used by the seasonal deciduous phenology in CLM5 is adopted for fruit trees. Before the start of fruit growth, phase (1), newly assimilated available C and corresponding N are partitioned between leaves,

stem_a and root pools. The allocation coefficients are calculated according to a set of equations that were adapted from the AgroIBIS crop phenology algorithm used in CLM5-BGC-crop (Lawrence et al., 2018):

Formatiert: Schriftart: Kursiv

$$GDD_{T2m} = GDD_{T2m} + T_{2m} - T_f - 4 \qquad \text{where} \quad 0 \le T_{2m} - T_f - 4 \le 26 \,^{\circ} days \,, \tag{1}$$

$$a_{repr} = 0 , \qquad (2)$$

$$a_{froot} = a_{froot}^{i} - \left(a_{froot}^{i} - a_{froot}^{f}\right) * \frac{GDD_{T2m} - GDD_{leaf}}{GDD_{fruit} - GDD_{leaf}},$$
(3)

$$a_{leaf} = (1 - a_{froot}) * \frac{a_{leaf}^{i} * (e^{-b} - e^{-b^* GDD_{I}mat - GDD_{leaf}})}{e^{-b - 1}},$$
(4)

$$a_{livestem} = 1 - a_{repr} - a_{froot} - a_{leaf} ,$$
⁽⁵⁾

where GDD_{T2m} are the accumulated growing degree-days for the 2-m air temperature with maximum increments of 26 °days;
T_{2m} is the simulated 2-m air temperature in K; T_f is the freezing temperature of water and equals 273.15 K; GDD_{leaf},
GDD_{fruit}, and GDD_{lfmat} are thermal thresholds for bud break, start of fruit growth, and canopy maturity, respectively; b is an
exponential factor; aⁱ_{leaf}, aⁱ_{froot}, and a^f_{froot} are initial and final values for the allocation coefficients to leaf (a_{leaf}) and fine root (a_{froot}), respectively; and a_{repr} and a_{livestem} are the allocation coefficients to fruit and live stem, respectively.

Once fruit growth begins in phase (2), an increasing proportion of the assimilated C and corresponding N is allocated to this organ, causing leaf allocation to decline and fruit allocation to plateau at a high value once canopy maturity is reached. Allocation to fFine roots and stem continues to decline and then settles at a constant value until harvest.

$$a_{livestem} = a_{livestem} * \left(1 - \frac{(GDD_{T2m} - GDD_{leaf}) - (GDD_{fruit} - GDD_{leaf})}{(GDD_{mat} - GDD_{leaf})^{d_L - (GDD_{fruit} - GDD_{leaf})}} \right)^{d_{alloc}^{ASim}}, \tag{6}$$

$$a_{repr} = 1 - a_{froot} - a_{livestem} - a_{leaf},\tag{(1)}$$

260 where GDD_{mat} is the thermal threshold for fruit maturity and harvest, while d_L and d_{alloc}^{stem} are stem allocation decline factors.

After harvest and until the start of dormancy, all of the newly assimilated C is sent to the storage pools following the notion that late in the season, assimilates are used mostly to fill up reserves that can be mobilized to resume growth in the following spring (Le Roux et al., 2001). Fruit trees store C in the perennial woody parts of the tree from where it is remobilized to support the growth of new shoots, leaves, and fine roots (Oliveira and Priestley, 1988; Millard, 1996; Le Roux et al., 2001). Since in CLM5 separate storage pools are assigned to each plant organ, the newly added *aleafstor* parameter (Table C1) defines the

fraction of allocatable C going to the leaf storage pool while the remainder is split equally between roots and stem.

265

Fruit trees, similar to other deciduous species, have been observed to translocate N out of senescent leaves to be reused by other tree organs (Millard, 1996; Malaguti et al., 2001; Millard et al., 2006). Therefore, CLM5-FruitTree adopts the same N retranslocation strategy as used in the BDT phenology during which N is removed from falling litter based on leaf and litter

270 <u>C:N ratios and the available C to pay for the extraction of N from increasingly recalcitrant litter pools. Subsequently it is transferred to the plant N pool from where it can be used for the growth of new plant tissue (Lawrence et al., 2018).</u>

2.2.3 Representation Parameterization of other processes and of management practices

- Fruit trees, similar to other deciduous species, have been observed to translocate N out of senescent leaves to be reused by
 other tree organs (Murneek, 1942; Bollard, 1960). Therefore, CLM FruitTree adopts the same N retranslocation strategy as
 used in the BDT phenology (Lawrence et al., 2018). Furthermore, management practices such as fertilization and stem pruning
 are represented in the new sub-model. Fertilization is performed on a yearly basis after the occurrence of bud break, as N
 fertilization in early spring is still the most common practice in fruit orchards even though autumn fertilization or multiple
 applications via fertigation are also in use to increase fertilizer N use efficiency and reduce N losses (Sanchez et al., 1995;
 Carranca et al., 2018). We use the existing fertilization scheme of the crop phenology that adds fertilizer directly to the soil
- mineral N pool. A user-defined fertilization rate or amount can be applied as synthetic fertilizer or manure respectively, although there currently is no difference in model behaviour for these two fertilizer types (Lawrence et al., 2018). Winter pruning is a common practice in fruit orchards and may be performed throughout the winter to control the shape and
- size of fruit trees, and partially to manage crop load (Grechi et al., 2008). Pruning is performed as the tree enters dormancy by removing a user defined fraction, *prune_fr* (Table C1), of the dead stem from both storage and displayed C pools. Pruning residues are treated in one of two ways: (1) residues are added to the wood harvest pool and exported from the field or (2) residues are mulched into the soil thus feeding the litter cycle. The former is typically adopted in conventional agriculture were residues are exported and treated as waste (Benyei et al., 2018) or utilized for energy production (Kazimierski et al., 2021).

The latter is more common in orchards uIn organic production. In many intensive orchard production systems, pruning residues

- 290 are mulched into the soil, possiblyto increasinge C sequestration in soils (Montanaro et al., 2010; Aguilera et al., 2015). Alternatively, residues may also be exported and treated as waste (Benyei et al., 2018) or utilized for energy production, (Kazimierski et al., 2021).During the first three years after planting, trees are not pruned to allow some initial stem biomass to grow. In practice, winter pruning in fruit orchards may be performed throughout the winter to control the shape and size of fruit trees, and partially to manage crop load (Grechi et al., 2008). In CLM5-FruitTree, pruning is performed as the tree enters
- 295 dormancy by removing a user-defined fraction, *prune_fr* (Table C1), of the dead stem from both storage and displayed C pools. We remove C from the dead stem pool instead of the live stem pool since the former is the main wood pool in CLM5 that receives 85 % of the C allocated to total new wood. Furthermore, the implemented live wood turnover in CLM5 converts live stem to dead stem at the end of the growing season to account for differences in maintenance respiration and C:N ratios between these tissue types (Lawrence et al., 2018). Hence the live stem C pool remains rather small and stable over the years so that

 Formatiert: Schriftfarbe: Automatisch

 Formatiert: Schriftfarbe: Automatisch

- 300 applying pruning to this pool would have little effect on total tree biomass. The pruning implemented in CLM5-FruitTree affects only the dead stemtree biomass and tree-height that is calculated based on this biomass pool, which in turn eaffects the calculation of turbulent fluxes of sensible and latent heat. However, this effect is small, and since turbulent fluxes are generally low in winter, the exact timing of pruning does not play a significant role in the magnitude of these fluxes. During the first three years after planting, trees are not pruned to allow some initial stem biomass to grow. The sub-model treats pruning 305 residues in one of two ways to account for their possible difference in fate: (1) residues are added to the wood harvest pool
- and exported from the field or (2) residues are added to the woody debris pool thus feeding the litter cycle. When the orchard reaches the end of its lifespan, C of all biomass pools (storage, transfer, and display) is sent to either the

litter pool for leaves and fine roots, or the wood harvest pool for live and dead stem and coarse roots, while any remaining C in the fruit pool is harvested. The orchard can then be replanted in the following year. Lastly, the standard irrigation routine 310 implemented in CLM5 can be used for irrigated orchards by selecting the irrigated crop PFT.

2.3 Model implementation and testing

2.3.1 Site data

- Extensive field measurements from an apple-growing region in the Adige River valley, South Tyrol, Italy (46°21' N, 11°16' E; 240 m a.s.l.) were used to parameterize and test the new sub-model CLM5-FruitTree along with the new apple PFT
 315 (Zanotelli et al., 2013; Zanotelli et al., 2015; Zanotelli et al., 2019). Measurements were obtained from an approximately 0.5 ha irrigated apple orchard planted in 2000 with the Fuji apple cultivar grafted on M9 dwarfing rootstock. The apple trees were planted at a row and tree spacing of 3x1 m (3333 trees ha⁻¹). A 1.8 m wide grass strip was grown between the tree rows, which was mowed 3 times a year. Other management practices included regular pruning, spring fertilization of 7.5 gN m⁻² yr⁻¹, and tillage of the soil directly underneath the trees (Zanotelli et al., 2013). Stand-related data included general stand characteristics
- 320 and phenology observations, leaf area index, C:N ratios, rooting distribution at three depth ranges (0–20 cm, 20–40 cm, and 40–60 cm), measurements of the biomass growth of different tree organs at monthly or seasonal interval, and fruit harvest information (Table 1). Furthermore, daily soil respiration measurements from a control and a trenching plot (with (R_s) and without root respiration (R_h), respectively) were performed in 2010. Additionally, an eddy covariance (EC) station provided measurements of the turbulent exchange of trace gases and energy at the studied apple orchard between 2013 and 2015. The
- 325 quality check, gap filling, and flux partitioning of collected data followed the procedure outlined in Reichstein et al. (2005). The average closure of the energy balance was 60 %. To correct for the closure failure, the missing energy was assigned to the latent (LE) and sensible (H) heat fluxes based on the daily Bowen ratio (Zanotelli et al., 2019). Measured or derived fluxes included net ecosystem CO₂ exchange (NEE), ecosystem respiration (R_{eco}), gross primary production (GPP), LE, H, and evapotranspiration (ET) at half-hourly intervals. Furthermore, soil heat flux (G) measured at 5cm depth as well as soil moisture
- 330 measurements up to a depth of 60 cm of soil are available. Table 1 gives a summary of the available data and measurement

periods. A complete description of the measurement procedures and instruments can be found in Zanotelli et al. (2013), Zanotelli et al. (2015), and Zanotelli et al. (2019).

Meteorological data, recorded partly at the EC tower and at the Laimburg meteorological station located 4 km from the site (46°23' N, 11°17' E; 224 m a.s.l.), were used at an hourly time step to force the model. Measured data included precipitation, solar radiation, net radiation (R_n , only at EC tower), air temperature, air pressure (only at Laimburg), relative humidity, and

wind speed. Measurements of incoming longwave radiation (LW_{in}) were available for 2010 only, but additional calculations following Konzelmann et al. (1994) and Sedlar and Hock (2009) were produced and used as forcing for the remaining years 2011–2019 (Appendix B). This was necessary since the use of the internally calculated LW_{in} in CLM5 resulted in unrealistic underestimations compared to the available measurements of LW_{in} leading to a significant bias in R_n.

340 Table 1: Summary of available data from an apple orchard in the Adige River valley, South Tyrol, Italy between 2010 and 2019. Solid lines represent continuous <u>and</u>, dotted lines monthly measurements, <u>whileand</u> diamonds represent single measurements.

Data	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
Weather data EC tower		•		-			•			
Weather data Laimburg										
Biomass (NPP) components	• • • • • •	•	•							
C:N ratios of biomass components	• • • • • •									
Leaf area index (LAI)	• • • • • •	•••••								
Fruit production (yield)	•	• •	*	•	*	•				
Root distribution (0-60 cm)	*									
Soil respiration (Rs, Rh)		•								
Soil heat flux (G)							•			
Soil water content (SWC, 0-60 cm)							•			
EC data: carbon (GPP, R_{eco} , NEE), energy (R_n , LE, H), water (ET)										

2.3.2 Model setup

345

335

The model was set up in point mode to simulate the apple orchard in the Adige valley using available sand, clay, and organic matter fractions. The model was spun-up for 200 years, first in accelerated decomposition and then in normal decomposition mode, until all state variables, such as total ecosystem soil C and soil water, reached equilibrium (Lawrence et al., 2018). For the model spin-up, the CRUNCEPv7 atmospheric forcing data set from 1986 to 2016 was used (Viovy, 2018). The apple orchard was then initiated using the newly developed sub-model and the apple PFT selecting the site-specific management (i.e., fertilization with 7.5 gN m⁻² yr⁻¹, irrigation, mulching of pruning material). Simulations were performed for a period of

350 10 years to mirror the time from orchard establishment in 2000 up to the start of the measurements in 2010 using 10 years (2010–2019) of the available meteorological data from Laimburg meteorological station. Simulations were then extended for another 6 years from 2010 to 2015 for model parameterization and performance evaluation purposes.

2.3.3 Parameterization

Key parameters of the new sub-model as well as other PFT-specific parameters were parameterized using the first 3 years of

- 355 simulations between 2010 and 2012. The length of phenological stages and associated parameters were determined based on field observations of bud <u>burstbreak</u>, full bloom, and harvest as well as non-cultivar specific apple phenology descriptions that were found in the literature (Appendix C). The length of the period where growth is supported out of reserves (*pdays_stor*) was calibrated based on the biomass measurements and the estimate by Zanotelli et al. (2013) that apple trees use stored carbohydrates in the first two months after bud break. C allocation coefficients were calculated based on the monthly
- 360 measurements in 2010, by dividing the biomass growth of the individual plant organs by the total biomass increment. Subsequently, model parameters associated with the CN allocation subroutine (Eq. (2) – (7)) were calibrated manually to match the coefficients obtained from the observations and the overall biomass partitioning on a yearly basis. Parameter values for C:N ratios of all plant organs and maximum LAI were based on field observations in 2010 and 2010–2012, respectively. The specific leaf area index was calculated by dividing monthly measurements of LAI by leaf biomass and taking the average
- 365 of the obtained values. Plant Structural and morphological parameters such as maximum tree height (<u>topmx</u>), plantingstocking density (<u>nstem</u>), the ratio of stem height to radius at breast height (<u>taper</u>), or rooting depth (<u>toot dmx</u>) were adjusted based on site-specific information (Zanotelli et al., 2013). Initial biomass at transplanting was assumed 5 gC m²/₂ resulting in an initial tree height of around 100 cm and a stem diameter of 16 mm. As seedlings are dormant at the time of transplanting, their LAI is 0. The CLM5 root distribution parameter (<u>tootprof beta</u>), which sets the root ratios at different depths, was calibrated by
- 370 least squares regression of the measured root ratios at 0–20 cm, 20–40 cm, and 40–60 cm depths and the calculated ratios. Optical parameters for leaf transmittance and reflectance in the visible and near-IR were set to average values reported for apple by Bastías and Corelli-Grappadelli (2012). Stem reflectance and transmittance were assumed to be similar to other woody species and therefore set to the values used for BDT in CLM5, similar to the assumptions made by Fan et al. (2015) for the palm tree development in CLM4.5. The ratio of momentum roughness length to canopy top height (<u>z0mr</u>) was set to the average
- 375 value of the ranges reported for apple and citrus orchards to account for the differences in canopy structure compared to annual crops and forest (Tanny and Cohen, 2003; La Fuente-Sáiz et al., 2017). No specific values could be found for the ratio of displacement to top of canopy height (*displar*), the leaf orientation index (xl), or the intercept to calculate the top of canopy maintenance respiration base rate (*lmr_intercept_atkin*). These values were assumed to be comparable to other deciduous trees
- and thus set to the values used for BDT in CLM5. Parameters related to C reserve dynamics (e.g., *fcur*) and photosynthesis
 (e.g., the slope of the relationship between leaf N per unit area and the maximum rate of carboxylation at 25 °C (<u>s vcad</u>)) were adjusted to match observed LAI and productivity data. All parameters with their values and references to the literature are summarized in Table C1 of the Appendix.

Formatiert: Schriftart: Kursiv

-	Formatiert: Schriftart: Kursiv
-{	Formatiert: Schriftart: Kursiv
\neg	Formatiert: Schriftart: Kursiv
X	Formatiert: Schriftart: Kursiv
X	Formatiert: Hochgestellt
-(Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv

Formatiert: Schriftart: Kursiv	
Formatiert: Schriftart: Kursiv	
Formatiert: Schriftart: 10 Pt., Kursiv	
Formatiert: Schriftart: Kursiv	

2.3.4 Sensitivity Analysis

A simple one-by-one sensitivity analysis was performed to further tune model parameters and assess the influence of newly 385 added parameters on the simulation results. As a complete sensitivity analysis of all PFT-related parameters would have exceeded the scope of this study, the analysis focused on key parameters of the new phenology and CN allocation subroutines. Other potentially influential parameters were selected based on previously performed sensitivity analyses by Göhler et al. (2013) for CLM3.5, and by Cheng et al. (2020) and Dagon et al. (2020) for CLM5 taking into account differences between previous and current model versions. Parameters selected for the analysis were perturbed by varying a parameter by ± 30 %, ± 20 %, and ± 10 % while keeping the others fixed to the value of the control simulation (after initial parameterization). The

390

goal here was not to perform an in-depth analysis covering the full range of possible parameter values, but rather to provide a first indication of influential parameters in the new sub-model similar to the approach of Fan et al. (2015). As a measure of sensitivity, the parameter effect (PE) was calculated using the average of 3 years of simulations between 2013 and 2015 of the control and the perturbed simulations for selected output variables and the following formula adjusted from Luo et al. (2020):

$$\Delta X_{i,j} = \sum_{k=1}^{n} \frac{|\overline{X_{i,j,k}} - \overline{X_{i,control}}|}{|\overline{X_{i,control}}|} , \qquad (8)$$

$$PE_{i,j} = \frac{\Delta X_{i,j}}{max \left[\left(\Delta X_{i,j} \right)_{1 \le i \le n; 1 \le j \le m} \right]},$$
(9)

395

400

where Where X is a simulated value of the control or a perturbation run, ΔX is the summed absolute difference between the <u>control and the perturbation run across all perturbations</u>, k is the parameter perturbation factor, i is the ith variable across n =6 selected output variables including: GPP, NEE, R_a , LE, maximum LAI, and yield, *i* is the *j*th parameter across *m* selected parameters. $PE_{i,j}$ is a number between 0 and 1 that represents the sensitivity of an output variable *i* to the parameter *j*, with 1 meaning high and 0 meaning low sensitivity. The parameters selected for sensitivity analysis are indicated in Table C1 of the Appendix.

2.3.5 Model performance evaluation

Modelling results are compared to observed biomass, yield, and LAI data as well as ecosystem fluxes retrieved from the EC measurements. Statistical indices for model performance evaluation include the Pearson's coefficient of correlation (r), the 405 root mean square error (RMSE) and the percent bias error (%bias):

$$r = \frac{\left(\frac{1}{n}\sum_{i=1}^{n} (x_{i}^{o} - \mu^{o})_{*}(x_{i} - \mu)\right)}{\sigma^{*}\sigma^{o}},\tag{10}$$

$$RMSE = \sqrt{\frac{1}{n} \sum_{i=1}^{n} (X_i - X_i^o)^2},$$

$$\% bias = \frac{\sum_{i=1}^{n} (x_i - x_i^o)}{\sum_{i=1}^{n} (x_i^o)},$$
(12)

(11)

where *i* is the time step, *n* is the total number of time steps, and X_i and X_i^o are simulated and observed values at each time step respectively, μ and μ^o are simulated and observed mean values, respectively, and σ and σ^o are simulated and observed standard deviations.

410

3 Results and discussion

3.1 Sensitivity analysis

A total of 34 parameters were initially considered for the sensitivity analysis of which the 13 most influential parameters (PE > 0.1 for at least one of the selected output variables) are shown in Fig. 3. GPP, NEE, R_a, and yield have similar sensitivity

- 415 patterns and are most sensitive to the leaf C:N ratio (*leafcn*) and the relationship between leaf N and the maximum rate of carboxylation at 25 °C (*s_vcad*). Together with the specific leaf area (*slatop*) and other constants, they control the maximum photosynthetic capacity in the photosynthesis calculation and thus largely influence total C assimilation. As expected, LAI is most influenced by parameters that control the CN allocation to leaves such as the initial leaf allocation coefficient (*fleafi*), the GDDs needed to reach canopy maturity (*lfinat*), the maximum LAI (*laimx*), photosynthetic parameters, and, to a smaller extent,
- 420 the fraction of C allocated to the leaf storage pool to refill C reserves (*aleafstor*). The first three parameters influence leaf biomass and thus show a considerable effect on GPP, NEE, R_a, and yield. The same output variables are affected in a similar fashion by the GDDs needed until fruit harvest (*hybgdd*) that control the amount of C allocated to fruits. LE is influenced largely by the parameter controlling stomatal conductance (*medlynslope*), and the photosynthetic parameters (*leafcn*, *s_vcad*). Overall, photosynthetic parameters play a key role in determining the magnitude of the studied output variables with an average
- 425 PE value close to 0.7 across all six variables. Phenological parameters (top seven parameters in Fig. 3) are generally less influential for the same output variables with average PE values up to 0.43. These findings are largely consistent with earlier studies of parameter sensitivity (Göhler et al., 2013; Cheng et al., 2020; Dagon et al., 2020; Luo et al., 2020). In contrast to Luo et al. (2020) we did not find a strong effect of the root distribution parameter (*rootprof_beta*) on LE, which can be attributed to the low water stress due to the irrigation management of the studied orchard.
- 430 While the one-at-a-time sensitivity analysis provides some insight into model sensitivity, the ranking of influential parameters is strongly influenced by the choice of parameters and output variables, the parameter perturbation strategy (i.e. percent change,

linear sampling), and the index chosen as the sensitivity measure. Parameter tuning based on this analysis is further complicated since this approach does not consider parameter covariation that is particularly strong for plant parameters that influence photosynthesis (Göhler et al., 2013). Selecting parameter values based on the individual best simulation hence does not necessarily yield the best overall result (Luo et al., 2020). We therefore decided to first adjust *s_vcad* to best match the observed average GPP. In the following, we further adjusted *fleafi*, *hybgdd* and *medlynslope* to improve the simulated biomass components as well as the LE flux, respectively.



Figure 3: Parameter effect (PE) as a measure of sensitivity of selected output variables to the most influential model parameters.
Output variables include gross primary production (GPP), net ecosystem exchange (NEE), autotrophic respiration (R_a), maximum leaf area index (LAI_{max}), latent heat flux (LE) and yield. Parameters are: Post-harvest leaf allocation coefficient to storage (aleafstor), initial leaf allocation coefficient (fleafi), GDD to canopy maturity (lfmat), root allocation coefficients at the start of fruit development (arootf) and until harvest (arootf2), GDD needed until harvest (hybgdd), maximum LAI (laimx), fraction of allocation that goes to currently displayed growth (fcur), C:N ratios of fruits (graincn) and leaves (leafcn), specific leaf area at top of canopy (slatop), slope of the relationship between leaf N per unit area and the maximum rate of carboxylation at 25 °C (s_vcad), and the medlyn slope of the conductance–photosynthesis relationship (medlynslope). For more details on the parameters, see Appendix C.

3.2 Modelling results

450

435

In the following, we present the modelling results according to the initial parameterization and the updated parameters values from the sensitivity analysis. Daily simulations or yearly sums are compared to observed biomass, yield, and LAI data as well as ecosystem fluxes retrieved from the EC measurements aggregated to daily mean values.

3.2.1 Biomass growth and yield

The patterns in seasonal biomass allocation simulated by CLM<u>5</u>-FruitTree show good agreement with the monthly observations from 2010 (Fig. 4a). The beginning and end of the growing season are well captured. After bud break in the beginning of March, biomass is primarily allocated to the vegetative organs of leaves, fine roots, and woody organs, and growth is supported by C and N reserves until the start of fruit growth in early May (50 days according to the *pdays_stor* parameter). In the following months, fruit biomass grows rapidly until harvest takes place in mid-October, following the typical sigmoidal growth curve that is well captured by the new phenology and CN allocation. <u>LSimulated leaf</u> biomass peaks in early July mid-June and remains constant thereafter, with leaf <u>senescenceoffset</u> starting later October when temperatures drop below 4 °C. Pruning is performed when the tree enters dormancy by removing 85 % of the stem biomass assimilated over the season according to the observed pruning amounts in the studied apple orchard (Zanotelli et al., 2013; Zanotelli et al., 2015). From

according to the observed pruning amounts in the studied apple orchard (Zanotelli et al., 2013; Zanotelli et al., 2015). From 2010 to 2012, the modelled percentage of biomass allocation to plant organs was generally in agreement with the observations (Zanotelli et al., 2015) with differences ranging between 1 and 5 % for fruits, leaves, aboveground wood, and roots (Fig. 4b).
Penzel et al. (2020) stated that different studies reported biomass allocation to fruits ranging from 50 to 85 % depending on apple cultivar suggesting considerable variability in allocation coefficients. This emphasizes the benefit of a cultivar specific
calibration in order to obtain realistic modelling results. On the other hand, it suggests that a more general parameterization,

that reflects an average apple tree, may be necessary to apply CLM5-FruitTree at larger scales and across multiple cultivars.



Formatiert: Schriftart: Kursiv

Figure 4: (a) Observed and simulated growth of leaves, fruits, fine roots, aboveground (live and dead stem), and belowground biomass (live and dead coarse roots) during 2010. (b) Observed and simulated biomass components between 2010 and 2012 as percentage of total biomass.

The timing for initial leaf development in spring and leaf offsetsenescence in late autumn are sufficiently well captured by the implemented bud burstbreak prediction algorithm and the simple temperature threshold for leaf abscission respectively (Fig. 5). Observed maximum LAI varied between 2.8 and 3.3 m² m⁻² and occurred during the first half of July. The simulations reached similar values in 2010 and 2012 matching the observations, while the lower simulated LAI in 2011 underestimated

- 475 the measurements due to a smaller C transfer from storage and lower solar radiation early in the growing season. Such The discrepancy between the low simulated LAI and the high observed LAI in 2011 could have been further exacerbated bybe further related to a lighter pruning performed in the previous winter, compared to other years (Zanotelli et al., 2013). Such management practice is sometimes performed in an attempt to counteract the strongwell known alternate bearing behaviour of the Fuji applevariety, which causes a substantial drop in yield following a high yielding year (Belleggia et al., 2009; Atay
- 480 et al., 2013; Pasa et al., 2021). <u>As alm</u> consequence toof the light pruning, a larger amount of vegetative and flower buds remained on the tree leading to <u>a higher yield more growth</u> and, possibly <u>explainingcontributing to</u> the larger discrepancy between relatively high observed LAI and relatively low simulated <u>valuesLAI</u>. <u>The adjusted pruning is however based on a somewhat subjective assessment of the farmer and information about the exact amount is hardly available. Thus CLM5-FruitTree currently adopts a simplified pruning practice based on the removal of a fixed portion of the seasonal stem growth which manages tree size and total woody biomass without affecting LAI.</u>

Measured LAI showed a slow decline soon after maximum LAI was reached, while simulated values in contrast are assumed to remain constant until-the leaf senescence-offset period is initiated. The observed early decline may be an artefact of the sampling strategy used to determine LAI that extrapolated individual leaf area measurements to the whole tree, assuming a constant leaf distribution within the tree (Zanotelli et al., 2013). Another reason could be some premature-a limited leaf fall in occurring during the summer at the expense of the inner shadowed leaves, as notedobserved during field sampling. Other studies suggest that the LAI of fruit trees generally stays constant until a rapid decline with the startonset of senescence (Lakso et al., 1999; Pallas et al., 2016), supporting the simulated LAI dynamic.



Kommentiert [m7]: Number and naming of x-ticks was adapted

Figure 5:-Observed (±standard error) and simulated leaf area index (LAI) between 2010 and 2012. Simulated daily leaf area index (LAI) between 2010 and 2012 together with observations (±standard error) of LAI that were made once a month for the same period. Ticks on the x-axis refer to the beginning of the month.

Simulated yield averaged-at 70 t ha⁻¹ between 2010 and 2015 and was within 2.3 % of the observed average yield. While simulated yield varied between 61 and 76 t ha⁻¹, the observations showed a greater inter-annual variability (IAV), as exemplified in the case of the year 2012 (low yield of 51 t ha⁻¹) and 2015 (high yield of 101 t ha⁻¹) (Fig. 6). Low <u>IAVinterannual variability</u> of yield has also been observed in previous crop simulations with CLMS for winter wheat (Boas et al., 2021)

suggesting that additional certain drivers of IAVyield variability such as extreme environmental conditions (e.g., frost, heat, and hail) or plant pests and diseases and -the resulting plant physiological responses (e.g., stress-induced leaf shedding or failure to flower) (Charrier et al., 2021), as specific management practices are missing or not represented with sufficient detail in CLM5. In the case of apple trees, yield is also tightly linked to the number of flowers and early fruit growth, which in turn 505 depend on a complex interaction of the environmental conditions during winter dormancy and the start of the new growing season (Chmielewski et al., 2012; Corelli-Grappadelli and Lakso, 2002). Additionally, C reserves accumulated in the previous year (Greer et al., 2002) and crop load management play an important role in determining the final harvest (Penzel et al., 2020). The latter includes pruning or fruit thinning to ensure optimal fruit growth and to reduce the effect of prevent alternate bearing. 510 The low observed yield in 2012 may be a result of such behaviour. -This phenomenon and the processes involved are not universal so that different fruit trees may be bearing regularly, irregularly or biannually (Hoblyn et al., 1937; Monselise and Goldschmidt, 1982). As such, alternate bearing and its treatment through pruning or fruit thinning cannot easily be generalized and are thus not currently implemented in CLM5-FruitTree, which could have further reduced simulated IAV. Storage growth is considered in CLM5-FruitTree and exhibited an impact on the final yield of the following season as shown by the sensitivity 515 analysis of the *aleafstor* and *fcur* parameters (Fig. 3). However, its effect on fruit growth in CLM5-FruitTree is indirect since it supports leaf development in the early growth stage but does not directly contribute to fruit growth. Identifying the driving forces of reserve deposition and mobilization, and their quantification remains an unsolved issue and there is yet no consistent formulation of this process in tree modelling (Le Roux et al., 2001; Allen et al., 2005). Predicting final yield in fruit orchards is further complicated by the fact that harvest is usually based on certain fruit quality traits such as firmness, or soluble solids 520 and can occur successively as fruits may not mature at the same time (Corelli-Grappadelli and Lakso, 2002; Musacchi and

- Serra, 2018). Within this context, the proposed simplifications of the C reserve dynamics and fruit harvest are likely reasons contributingfor to the difference in observed and simulated yields while the missing representation of specific crop load management practices regarding the Fuji apple contributed to the low variability in simulated yields. Considering the many specific challenges in modelling this apple cultivar, we believe that the yield predictions are satisfactory enough in the context 525 of the sub-model development.





530 3.2.2 Ecosystem fluxes and soil moisture variation

Carbon fluxes

As shown in Fig. 7, CLM5-FruitTree was able to capture the overall patterns of GPP, NEE, and R_{eco}, particularly during the transition between dormancy periods and growing seasons (April to November). Simulated C fluxes are highly correlated with observations (r ≥ 0.84) while the RMSE ranges between 1.12 and 1.53 gC m⁻² d⁻¹. Observed and simulated peak C fixation occurred mid-July-June (Fig. 7a), corresponding to the maximum (negative) NEE (Fig. 7c) and maximum LAI (Fig. 5). Simulated NEE becomes negative (net carbon sink) around April and returns to positive (net carbon source) around November in agreement with the observed dynamic (Fig. 7c). Observed yearly sums of GPP (NEE) were 1.60 (-0.49), 1.43 (-0.48), and 1.65 (-0.76) kgC m⁻² yr⁻¹ for 2013, 2014, and 2015, respectively. Simulated yearly sums of GPP (NEE) were 1.58 (-0.53), 1.56 (-0.51), and 1.53 (-0.57) kgC m⁻² yr⁻¹ for the same years showing a negligible positive bias of on average 0.17 % for GPP (Fig. 540) and a small underestimation (less negative) of on average 3.8 % for NEE (Fig. 7d). Simulated and observed R_{eco} (Fig. 7e)

- generally increased until July, because of the increase in air temperature and respiratory costs of the developing canopy, and declined thereafter as air temperature started to drop. Simulations of R_{eco} tend to slightly underestimate observations between April and late August and to overestimate observations during winter, although discrepancies are relatively small. Observed yearly sums of R_{eco} were 1.13 (2013), 0.98 (2014), and 0.94 (2015) kgC m⁻² yr⁻¹ while simulated values were 1.08, 1.08, and
- 545 0.99 kgC m⁻² yr⁻¹, respectively. CLM<u>5</u>-FruitTree overestimated yearly R_{eco} by on average 3.3 % explaining most of the difference in observed and simulated NEE in 2013, while differences in 2014 and 2015 are owed to a combination of small biases in both GPP and R_{eco} . Measured R_{eco} showed irregular fluctuations in the early part of the growing season 2013, and mid- to late-season 2014 and 2015 that are not reproduced well by the model. These fluctuations mostly correspond to the

observed temperature dynamics (not shown) as a result of the applied gap filling that is based on an air (or soil) temperature-

- 550 R_{eco} relationship (Reichstein et al., 2005). Such discrepancies between observed and simulated dynamics could be further explained by the occurrence of field management practices such as mowing of the grassed alleys or soil tillage under the tree rows, which are currently not represented in CLM5-FruitTree. Such practices could have led to a temporary rise in soil respiration (R_s) due to increased heterotrophic respiration (R_h) as discussed in Zanotelli et al. (2013). Indeed soil tillage experiments performed in an apple orchard located in a Loess plateau in Shaanxi Province in China were found to increase R_s 14-57 % depending on the tillage method (Hou et al., 2021).
 - Zanotelli et al. (2013) measured a total R_s of 801±95 gC m⁻² in 2010 contributing around 90 % to R_{eco} , based on soil chamber measurements within the orchard (total soil respiration). The comparison with parallel measurements in a trenched plot produced a high ratio R_b/R_s of 0.77 for the apple orchard. In contrast, simulated R_g was 510 gC m⁻² contributing merely 45 % to R_{eco} for the same year with a ratio R_b/R_s of 0.87. Simulated R_{gco} was instead dominated by autotrophic respiration (R_a) due
- 560 to high C costs for maintenance, mainly of leaf biomass (<u>data</u> not shown). Total simulated R_s-contributed merely 45 % (510 gC m⁻²) to R_{eco} with a ratio R_h/R_s of 0.87. Other studies found that R_s contributed 56-67 % to R_{eco} in irrigated citrus orchards of different ages <u>that share</u> common management practices (i.e., use of heavy machinery, irrigation, fertilization, tree pruning, and mulching) as well as structural similarities (e.g. planting in tree rows) with the studied apple orchard. Both aspects have a strong influence on soil respiration components in orchards (Martin-Gorriz et al., 2020). In forest ecosystems-and >60 % in
- 565 forest ecosystems, where the magnitude of ecosystem fluxes was found to be be senerally somewhat comparable to orchards, R_{g} contributed > 60 % to R_{gco} (Lasslop et al., 2012; Zanotelli et al., 2013).

In addition to the missing representation of certain management practices, CLM_{5} -FruitTree currently does not account for an active ground cover in the orchard which has shown to enhance R_{s} in an Italian olive orchard through increased fine root and microbial biomass (Turrini et al., 2017). Furthermore, the simplified representation of microbial activity in CLM5, through

570 fixed respiration fractions for litter and soil organic matter pools, may limit the ability of CLM5-FruitTree to accurately represent soil respiration processes. Not accounting for mycorrhizal respiration may fail to adequately represent R_{eco} of the orchard, as measurements suggested a substantial contribution of 11±6 % to total R_s in an apple orchard (Tomè et al., 2016). Lastly, biases in simulated soil temperature, or soil moisture organic matter content, and fine root density could partially further contribute to explain the above-discussed differences as these factors have a major effect on R_g in apple orchards (Ceccon et al., 2011).

In contrast to the underestimation of R_{g} in the model, the simulated R_{g} of 693 gC m_{1}^{-2} was almost twice the measured value of 372±195 gC m_{1}^{-2} . In our simulations R_{g} in CLM5 is the sum of maintenance and growth respiration for the different plant organs as well as the C cost associated with N uptake from the soil, whereby-maintenance respiration compriseds the main part of R_{g} (with on average 78 %-on average in this study). CLM5-The calculationes of maintenance respiration in CLM5 (see

580 section 2.1)based on tissue N content, temperature, and a constant base maintenance rate. This however, does not account for a lower or varying maintenance cost observed in mature apple orchard canopies compared to annual crops (Bepete and Lakso, 1996; Lakso et al., 1999). It therefore seems likely that the cost for leaf tissue maintenance costs forin the orchard areis Formatiert: Tiefgestellt
Formatiert: Tiefgestellt

Formatiert: Tiefgestellt

Formatiert: Tiefgestellt

Formatiert: Tiefgestellt

-{	Formatiert: Tiefgestellt
1	Formatiert: Tiefgestellt
Υ	Formatiert: Hochgestellt
Υ	Formatiert: Hochgestellt

overestimated unrealistically high in CLM5-FruitTree, accounting for on average 45 % of R_a or (28 % of R_{eco}). In comparison, in citrus orchards, with an evergreen canopy, leaf respiration was found to contribute 18–20 % to annual R_{eco} (Martin Gorriz et al., 2020). This could also explain the lower simulated carbon use efficiency (NPP/GPP) of 0.59 compared to 0.71 found by Zanotelli et al. (2013). Further work and more experimental data are needed to better understand the differences in modelled and observed respiration partitioning and to improve the performance of CLM5-FruitTree to adequately simulate the respiration components in fruit orchards.



Kommentiert [m8]: Number and naming of x-ticks was adapted

590 Figure 7: Daily instantaneous (a, c, e) and cumulative (b, d, f) observed and simulated fluxes of gross primary productivity (GPP), net ecosystem exchange (NEE), and ecosystem respiration (R_{eco}) for the studied apple orchard between 2013 and 2015. Pearson's coefficient of correlation (r), the root mean square error (RMSE) and the percent bias (%bias) are displayed as statistical indices.

Energy and water fluxes

585

The simulated seasonal course of the energy balance components R_n , G, LE and H agrees well with observed dynamics in the orchard (Fig. 8). CLM<u>5</u>-FruitTree shows a high performance in reproducing R_n and LE with $r \ge 0.97$ and RMSE of 15.98 and 17.85 W m⁻², respectively (Fig. 8a and c). Due to the lack of LW_{in} measurements, the CLM5-internal LW_{in} calculation based on a clear-sky parameterization after Idso (1981) was used initially. This resulted in a significant underestimation of 5 % (511 MJ) for LW_{in} and 18 % (471 MJ) for R_n compared to the observations in 2010. The R_n bias could be reduced by 14 % for the observed time series when LW_{in} was calculated by considering cloud cover as described in Appendix B. This stresses the

- 600 necessity to account for cloud cover, ideally combined with locally calibrated parameters, for an accurate calculation of LW_{in} . The remaining small negative bias of 4.48 % in R_n is due to negative simulated R_n during the winter months (Fig. 8b), which may be a result of the higher reflectance of solar radiation from bare soil compared to a grass surface (Bryś et al., 2019). The model assumes a bare soil (except for stem area) during the dormancy period, as the grass-covered alleys in the orchard are not considered explicitly.
- 605 The simulated LE (Fig. 8c) shows similar dynamics and variability as the observations following the increase and decrease of GPP (Fig. 7a) and LAI (Fig. 5). Similarly to LE, modelled ET shows a high correlation coefficient of 0.97 and small RMSE of 0.62 mm d⁻¹ (Fig. 8i). Simulated ET exceeds observed ET by 1.1 mm d⁻¹ on average during its peak in July but the overall bias is almost negligible (Fig. 8j). Total observed ET is 901 (2013), 858 (2014), and 883 (2015) mm while corresponding simulated values are 916, 877, and 925 mm, respectively. When examining the order of magnitudes of the ET components,
- canopy transpiration takes up around 85 % of ET, followed by soil evaporation and canopy evaporation (data not shown). Typically, apple orchard ET represents a combined flux from the apple trees and the grassed alley system, which is not explicitly represented in CLM5-FruitTree since CLM5 does currently not consider inter-row grass coverage or intercropping. Ntshidi et al. (2021) found that the contribution of understory transpiration is high in young, non-bearing apple orchards, but contributes less than 10 % to whole orchard ET in mature orchards with high canopy cover, which may explain the good model
 performance despite not considering the grass cover.
- Simulated H and G are less consistent with the observations with r values of 0.54 and 0.64, respectively, and large %bias (Fig. 8e and 8g), which is partially due to the much smaller magnitudes of the two fluxes compared to R_n and LE. A possible reason for the lower amplitude of observed G (Fig. 8h) compared to simulated values may be the dampening effect of the grass cover providing additional shading during summer and insolation during winter (Bryś et al., 2019; Oorthuis et al., 2021). Observed
- 620 H was rather constant throughout the year with slightly higher values at the start and end of the growing season when the canopy was not yet fully developed, or leaves were shedding. CLM5-FruitTree simulated a clear rise of H until April, closely following the observations, but H thereafter declined steeply in May with negative values in August 2013 and 2015. Negative H during August corresponds with maximum LE and the main simulated irrigation season (June to September) that added 357 (2013), 281 (2014), and 517 mm (2015) of water to the orchard (Fig. 9a). In a study conducted with CLM4.5, intense irrigation
- 625 was found to strongly influence the convective heat fluxes by increasing LE and decreasing H (Zeng et al., 2017). Although precise measurements of the irrigation amount in the orchard are not available for the studied period, the average yearly irrigation was estimated around 200 mm with no irrigation in 2014 due to sufficient rainfall (Montagnani et al., 2018). The difference in irrigation amounts may in part explain why the described phenomenona is not observed in the measurements. Indeed negative simulated H in the summer months occurred as a result of strong evaporative cooling of ground and vegetation temperature through energy absorption by LE following irrigation, that irrigation that caused simulated LE to exceed simulated
 - $R_{\rm n}.$ This behaviour was not observed in the measurements where LE rarely exceeded $R_{\rm n}$ and was mostly owed to an

overestimation of simulated LE compared to the measurements. Persisting model weaknesses in the partitioning of the energy balance were pointed out by a recent study examining land surface processes over a tropical rainforest using CLM4.5 and CLM5, and were linked to missing detail in the representation of the canopy and an oversensitivity of vegetation temperature

635 to incoming solar radiation among others (Song et al., 2020). As a result, the authors observed an overestimation of LE and unrealistically high day-to-night changes in G, which was also observed in this study when examining the model output at an hourly time step (results not shown).

Energy partitioning in orchards is strongly influenced by the positioning and pruning of branches to optimize tree architecture for higher productivity, planting density, tree height, and LAI distribution (López-Olivari et al., 2016). Consequently, the
 640 contribution of H and LE can significantly differ in patchythe discontinuous orchard canopy (grass-covered alleys between

tree rows) compared to the closed canopies of annual crops (La Fuente-Sáiz et al., 2017). Currently CLM5 is still limited to the assumption of a closed canopy structure that is uniform in space Next to the aspects discussed above, hence biases in the simulated energy balance components are hence are likely to arise from this model limitation, s to represent such heterogeneous canopies. Future developments towards integrating multi-layer schemes for canopy processes and the explicit representation of the canopy to improve the related processes are desirable for a more realistic representation of the orchard canopy structure.

Soil moisture variation

Simulated mean soil moisture (SM) at 5 cm depth was within 1.6 % vol of the observed value during the three observed growing seasons, despite the higher simulated irrigation amount (Fig. 9b). Simulated daily values show a greater variability than the measured data in response to precipitation and to frequent irrigation (Fig. 9a–b). In contrast, observed SM in the deeper soils

- 650 (30–60 cm), was 3–11 % vol higher during the growing season compared to simulated values (Fig. 9c–d). Considering the total investigated soil depth, simulations exhibit a larger variability in SM throughout the year, with a general overestimation in winter and underestimation during the growing season (especially in the deeper soils). However, the collected SM data was limited to a single soil profile that may not adequately reflect the average soil moisture of the apple orchard, which should be considered when comparing measurements and simulations. Even though the measurements are incomplete, the constant high
- 655 observed SM in the deeper soils suggests an ample supply of water due to capillary rise from the shallow ground water table that typically ranges between 1.2 and 1.85 m in the area (Montagnani et al., 2018). This process replenishes the water removed by ET processes and may explain the reduced need for irrigation compared to the simulations. Despite the shallow simulated ground water table (generally 1.2 m depth), groundwater could not be used for root water uptake in the simulation as the rooting depth of the orchard was restricted to 0.8 m according to local measurements and capillary rise is currently not for a simulation of the orchard was restricted to 0.8 m according to local measurements and capillary rise is currently not
- 660 implemented in CLM5.



Kommentiert [m9]: Number and naming of x-ticks was adapted

Figure 8: Daily and cumulative observed and simulated fluxes of net radiation (R_n) , ground heat (G), latent heat (H), sensible heat (LE) and evapotranspiration (ET) for the studied apple orchard between 2013 and 2015. The coefficient of determination (r), the root mean square error (RMSE) and the percent bias (%bias) are displayed as statistical indices.

Kommentiert [m10]: SWC was changed to SM



4 Conclusions

670 The novel CLM5-FruitTree was developed to model perennial deciduous fruit orchards and thus extended the representation of agricultural systems in CLM5. The development included a new phenology subroutine to account for the perennial nature, prolonged growing season, and the distinct phenological development of fruit trees compared to annual crops. Furthermore, C reserve dynamics of perennial deciduous trees were considered by adapting the CN allocation and typical management

Figure 9: Precipitation and simulated irrigation (a), and observed and simulated soil water content<u>moisture</u> (S<u>MWC</u>) at 0.05 m (b), 0.3 m (c), and 0.6 m (d) depth from 2013-2015.

practices associated with fruit orchards were represented, such as transplanting of seedlings and winter pruning. To evaluate 675 the development, a new apple PFT was parameterized, and the model was set up and tested using extensive site data of a mature apple orchard in Northern Italy.

One-by-one parameter sensitivity analysis revealed that photosynthetic parameters and parameters associated with canopy conductance have the highest influence on GPP, NEE, LE and yield, while phenological parameters were more influential in biomass partitioning to the different plant organs. Due to the high number of model parameters and parameter covariation, future studies could propose a more comprehensive sensitivity analysis with a training dataset consisting of multiple sites,

which would give more insight into model sensitivity and could further improve the parameterization.

680

CLM5-FruitTree was able to capture the seasonal biomass development as well as the average relative partitioning of the total biomass into the different plant organs. The inclusion of C reserves next to photosynthetic growth was imperative to enable regrowth at the end of a dormancy period and influenced LAI development, total seasonal biomass, and yield. Average
 685 simulated yield was within 2.3 % of observation even though CLM5-FruitTree showed a lower inter annual variabilityIAV likely due to the simplification of C reserve dynamics, specific management practices, and the alternate bearing behaviour exhibited by the Fuji apple cultivar.

The new phenology and CN allocation algorithms well represented the seasonal course of carbon, water, and energy fluxes of the orchard. The magnitude of ecosystem fluxes was particularly well captured for GPP, R_n, LE, and ET with correlation

- 690 coefficients > 0.94 and % bias < \pm 5 %. The model exhibited small biases in NEE and R_{eco} that were most likely caused by the overestimation of R_a , especially leaf maintenance respiration, and an underestimation of R_g . Possible reasons for the smaller simulated contribution of R_s to R_{eco} could be the missing representation of the grass-covered alleys, differences in simulated and actual soil temperature or organic matter content, and oversimplification of microbial respiration processes. Additionally, large negative biases in simulated H were found over most of the main irrigation season during summer as the model simulated
- 695 a strong evaporative cooling of the surface temperature. Further model developments should consider the improvement of canopy processes related to energy partitioning and the inclusion of an active ground cover in the orchard representation to improve the yearly energy budget calculations and possibly soil respiration. An explicit representation of the microbial community and a more flexible calculation of R_a, i.e. considering tissue age, should also be the focus of future model improvements. While the particular alternate bearing of the Fuji variety
- 700 posed a challenge in this specific study, the pruning routine that is currently implemented may be sufficient for most other apple cultivars and fruit tree species for which this behaviour is less pronounced or not exhibited. However, future developments could be envisioned once the model is further tested and applied. In addition, management practices such as mowing or soil tillage could further enhance the model capability of capturing the dynamics and fate of assimilated C. Fruit thinning is another common practice in orchards, but its implementation would be more challenging, as the current model
- 705
 structure does not represent individual fruits. This process could however be implicitly accounted for through parameterization

 of the C allocation to fruits. Finally, the application of the newly developed sub-model to different geographical regions and

Formatiert: Tiefgestellt

other types of fruit trees or apple cultivars is needed to further validate the model and give more insight into the transferability of the development to different types of orchards.

Overall, our results demonstrate the ability of the newly developed CLM5-FruitTree sub-model to represent the seasonal dynamics and magnitudes of growth and ecosystem fluxes in a deciduous fruit orchard. As such, this development constitutes an important contribution towards a more comprehensive representation of the agricultural land surface in CLM5 by adding a perennial, woody crop to the existing annual crop types. This will allow for a more realistic evaluation of land use and climate change effects, or water availability at regional scale such as the Mediterranean or parts of China and the US, where perennial agriculture such as fruit orchards cover large parts of the agricultural landscape.

715

710

Appendix

Appendix A: Sequential model for bud burstbreak prediction

720

The bud <u>burstbreak</u> prediction in CLM<u>5</u>-FruitTree is based on the sequential model developed by Cesaraccio et al. (2004). Negative chill days (C_d) are accumulated from the 1st of November followed by positive anti-chill days (C_a) to overcome the different stages of tree dormancy, rest and quiescence. The chilling requirement (C_R) defines the threshold for the accumulation of C_d and is reached when $\sum C_d \leq C_R$. Thereafter C_a accumulation begins until $C_R + \sum C_a \geq 0$ at which bud <u>burstbreak</u> occurs. The accumulation of C_d and C_a on a given day is calculated from maximum (T_x) and minimum (T_n) daily air temperature as well as a temperature threshold for chill accumulation (T_C), and varies depending on five possible temperature cases that relate T_x, T_n, T_C, and 0 °C with the daily mean air temperature (Table A1). The optimal values for C_R and T_C were calibrated based on bud <u>burstbreak</u> observations from 2010–2013 for the Adige site by minimizing the RMSE between observations and predicted bud <u>burstbreak</u>. The optimal value for C_R was -68 while T_c was 4 °C resulting in an RMSE of 7.2 days.

725

Table A1: Chill day (C _d) and anti-chill day (C _a) calculation for five different temperature cases relating maximum (T _x) and minimum
(T_n) air temperature to the air temperature threshold (T_C) and 0 °C, T_M is the air mean temperature.

Temperature Cases	Chill days	Anti-chill days
$0 \le T_C \le T_n \le T_x$	$C_d = 0$	$C_a = T_M - T_C$
$0 \le T_n \le T_C < T_x$	$C_{d} = -\left[(T_{M} - T_{n}) - \frac{(T_{x} - T_{C})^{2}}{2(T_{x} - T_{n})} \right]$	$C_{a} = \frac{(T_{x} - T_{C})^{2}}{2(T_{x} - T_{n})}$
$0 \le T_n \le T_x \le T_C$	$C_d = -(T_M - T_n)$	$C_a = 0$
$T_n < 0 \le T_x \le T_C$	$C_d = -\left[\frac{T_x^2}{2(T_x - T_n)}\right]$	$C_a = 0$
$T_n < 0 < T_C < T_x$	$C_d = -\frac{T_x^2}{2(T_x - T_n)} - \frac{(T_x - T_C)^2}{2(T_x - T_n)}$	$C_{a} = \frac{(T_{x} - T_{C})^{2}}{2(T_{x} - T_{n})}$

730 Appendix B: Calculation of incoming longwave radiation

Incoming longwave radiation (LWin) can be expressed based on the Stefan Boltzmann Law as:

$$LW_{in} = \varepsilon_{eff} * \sigma * T^4 = \varepsilon_{cs} * F * \sigma * T^4 , \qquad (B1)$$

where ε_{eff} is the effective emissivity that can be expressed by multiplying the clear-sky atmospheric emissivity ε_{cs} with a cloud factor *F* (always ≥ 1) that expresses the increase of *LW*_{in} under cloudy conditions, σ is the Stefan Boltzmann constant (5.67 x 10⁻⁸ W m⁻² K⁻¹) and *T* is the 2m air temperature in K.

735 Clear-sky emissivity was obtained using the Konzelmann et al. (1994) parameterization as follows:

$$\varepsilon_{cs} = 0.23 + 0.484 * \left(\frac{e}{\tau}\right)^{\frac{1}{8}},\tag{B2}$$

where e is the vapour pressure in Pa at 2m.

Equation (B1) can be rearranged to obtain F as follows:

$$F = \frac{LW_{in}}{\varepsilon_{cs^*\sigma^*T^4}} \tag{B3}$$

F was calculated at hourly interval using measured LW_{in} data from 2010 and ϵ_{cs} calculated using the above Eq. (B2).

As proposed by Sedlar and Hock (2009), in the absence of cloud data, the cloud factor F can be parameterized as a function of 740 the atmospheric transmissivity index τ , which is defined as follows:

$$\tau = \frac{SW_{in}}{SW_{toa}} , \qquad (B4)$$

where SW_{in} is the incoming shortwave radiation, and SW_{toa} is the theoretical shortwave radiation received at the top of the atmosphere.

Figure B1 shows the linear equation that was fitted to the relationship of F and τ for the year 2010. For the calculation of clearsky emissivity, all data where τ was greater than 0.7 (N=3863) was considered based on the suggestion by Campbell (1985).

745

Kommentiert [m11]: Increased font size of the figure following a comment of reviewer 2



Figure B1: Cloud factor F as a function of atmospheric emissivity τ for hourly observations. The black line represents the linear equation for $F(\tau)$ and $F \ge 1$. Clear-sky emissivity is parameterized based on Konzelmann et al. (1994).

For the nighttime values and for very low incoming shortwave radiation $(SW_{in} < 15 \text{ W m}^2)$, τ was gap-filled with the mean of 750 the two surrounding values to obtain a complete time series of LW_{in} data. Figure B2 shows the results of the LW_{in} parameterization compared to LW_{in} calculated by CLM5 and to the observed data for the year 2010. As performance statistics the Pearson's r (r), the root mean square error (RMSE) and percent bias (%bias) are given.



Figure B2: Comparison of observed LW_{in} with the parameterization using (a) Konzelmann et al. (1994) according to Eq. (B2) and

755 the cloud factor parameterization F(τ), and (b) the calculation procedure used in CLM5, as well as (c) cumulative observed and calculated LW_{in} for 2010. Pearson's r, RMSE and %bias are given as performance statistics.

Kommentiert [m12]: Increased font size of the figure following a comment of reviewer 2

Appendix C: Parameters used in CLM5-FruitTree and for the apple PFT

760

Table C1: Parameters adapted or added in the new CLM5-FruitTree sub-model and the apple PFT including phenology, CN allocation, photosynthesis, vegetation structure as well as optical and respiration parameters. Parameters were adjusted based on field observations or literature values and are listed with their definition, unit, value, and references to the literature.

Parameter	Definition	Unit	Value	Reference
Phenological paramet	ers			
baset	Base temperature for GDD	°C	4	Based on commonly used values for apple trees (Reyes et al.,
	accumulation			2016; Díez-Palet et al., 2019; Penzel et al., 2020)
crequ	Chilling requirements for bud	unitless	-68	Calibrated using bud break dates from Zanotelli et al. (2013) and
	burstbreak of fruit tree crops			Zanotelli et al. (2015), and the sequential model (Cesaraccio et al.,
				2004)
crit_temp	Critical temperature to	Κ	278.15	Adjusted based on LAI measurements (Zanotelli et al., 2013)
	initiate leaf offsetsenescence			
	for fruit tree crops			
grnfill* (GDDfruit)	GDD needed from bud break	°days	400	Based on observed and commonly used values for apple trees
	to beginning of fruit			(Zanotelli et al., 2013; Lakso et al., 2000; Neumann, 2020; Penzel
	development			et al., 2020)
grnrp*(GDDripe)	GDD needed from bud break	°days	1100	Based on observed and commonly used values for apple trees
	to the fruit ripening phase			(Lakso et al., 2000; Zanotelli et al., 2013; Neumann, 2020; Penzel
				et al., 2020)
huileaf (GDDleaf)	GDD accumulated at the	°days	-	Calculated based on sequential model for bud burstbreak
	moment of bud break (end of			prediction (Cesaraccio et al., 2004)
	dormancy period)			
hybgdd* (GDDmat)	GDD needed from bud break	°days	2880	Based on observed and commonly used values for apple trees
	until fruit harvest			(Lakso et al., 2000; Zanotelli et al., 2013; Neumann, 2020; Penzel
				et al., 2020)
laimx*	Maximum leaf area index	m^2/m^2	3	Based on observed and commonly used values for apple trees
				(Valancogne et al., 1999; Li et al., 2002; Zanotelli et al., 2013)
lfmat*(GDDlfmat)	GDD needed from bud break	°days	1350	Based on observed and commonly used values for apple trees
	to canopy maturity			(Lakso et al., 2000; Zanotelli et al., 2013; Neumann, 2020; Penzel
				et al., 2020)

max_NH_harvest_date	maximum harvest date for	date (md)	1015	Based on typical harvest dates in NH
	northern hemisphere (NH)			
max_NH_planting_date	maximum planting date for	date (md)	101	Only needed for orchard establishment and initiation of sequential
	NH			model for bud break, tree is still dormant
min_NH_planting_date	minimum planting date for	date (md)	101	Only needed for orchard establishment and initiation of sequential
	NH			model for bud break, tree is still dormant
mxmat	Maximum orchard age	days	9125	Based on common values for apple orchards (Lakso et al., 2000;
				Zanotelli et al., 2013; Penzel et al., 2020)
ndays_stor	Length of period for storage	days	50	Based on common values for fruit orchards (Kozlowski, 1992;
	growth of fruit tree crops			Dejong and Grossman, 1994; Wünsche and Lakso, 2000)
perennial	Binary flag for perennial crop	unitless	1	
	phenology			
root_dmx	Maximum rooting depth of	m	0.8	Based on observed rooting depth (Zanotelli et al., 2013)
	crops			
rootprof_beta*	Rooting beta parameter, for C	unitless	0.964	Calibrated based on root sampling campaign of root mass up to
	and N vertical discretization			60cm (Zanotelli 2010, unpublished data)
woody	Binary flag for woody	unitless	1	
woody	Binary flag for woody lifeform	unitless	1	
woody C and N allocation para	Binary flag for woody lifeform meters	unitless	1	
woody C and N allocation para aleaff*	Binary flag for woody lifeform meters Final leaf allocation	unitless	0.01	Adjusted based on monthly biomass measurements (Zanotelli et
woody C and N allocation para aleaff*	Binary flag for woody lifeform meters Final leaf allocation coefficient	unitless	0.01	Adjusted based on monthly biomass measurements (Zanotelli et al., 2013)
woody C and N allocation para aleaff* aleafstor*	Binary flag for woody lifeform meters Final leaf allocation coefficient Leaf allocation coefficient to	unitless unitless	1 0.01 0.3	Adjusted based on monthly biomass measurements (Zanotelli et al., 2013) Adjusted based on monthly biomass measurements (Zanotelli et
woody C and N allocation para aleaff* aleafstor*	Binary flag for woody lifeform meters Final leaf allocation coefficient Leaf allocation coefficient to storage post-harvest used in	unitless unitless	1 0.01 0.3	Adjusted based on monthly biomass measurements (Zanotelli et al., 2013) Adjusted based on monthly biomass measurements (Zanotelli et al., 2013)
woody C and N allocation para aleaff* aleafstor*	Binary flag for woody lifeform meters Final leaf allocation coefficient Leaf allocation coefficient to storage post-harvest used in CNAllocation	unitless unitless	1 0.01 0.3	Adjusted based on monthly biomass measurements (Zanotelli et al., 2013) Adjusted based on monthly biomass measurements (Zanotelli et al., 2013)
woody C and N allocation para aleaff* aleafstor* allconss*	Binary flag for woody lifeform meters Final leaf allocation coefficient Leaf allocation coefficient to storage post-harvest used in CNAllocation Power to control the shape of	unitless unitless unitless	1 0.01 0.3 1.5	Adjusted based on monthly biomass measurements (Zanotelli et al., 2013) Adjusted based on monthly biomass measurements (Zanotelli et al., 2013) Adjusted based on monthly biomass measurements (Zanotelli et
woody C and N allocation para aleaff* aleafstor* allconss*	Binary flag for woody lifeform meters Final leaf allocation coefficient Leaf allocation coefficient to storage post-harvest used in CNAllocation Power to control the shape of the stem allocation curve	unitless unitless unitless	1 0.01 0.3 1.5	Adjusted based on monthly biomass measurements (Zanotelli et al., 2013) Adjusted based on monthly biomass measurements (Zanotelli et al., 2013) Adjusted based on monthly biomass measurements (Zanotelli et al., 2013)
woody C and N allocation para aleaff* aleafstor* allconss* arootf*	Binary flag for woody lifeform meters Final leaf allocation coefficient Leaf allocation coefficient to storage post-harvest used in CNAllocation Power to control the shape of the stem allocation curve Root allocation coefficient at	unitless unitless unitless unitless unitless	1 0.01 0.3 1.5 0.2	Adjusted based on monthly biomass measurements (Zanotelli et al., 2013) Adjusted based on monthly biomass measurements (Zanotelli et al., 2013) Adjusted based on monthly biomass measurements (Zanotelli et al., 2013) Adjusted based on monthly biomass measurements (Zanotelli et
woody C and N allocation para aleaff* aleafstor* allconss* arootf*	Binary flag for woody lifeform meters Final leaf allocation coefficient Leaf allocation coefficient to storage post-harvest used in CNAllocation Power to control the shape of the stem allocation curve Root allocation coefficient at start of fruit development	unitless unitless unitless unitless unitless	1 0.01 0.3 1.5 0.2	Adjusted based on monthly biomass measurements (Zanotelli et al., 2013) Adjusted based on monthly biomass measurements (Zanotelli et al., 2013) Adjusted based on monthly biomass measurements (Zanotelli et al., 2013) Adjusted based on monthly biomass measurements (Zanotelli et al., 2013)
woody C and N allocation para aleaff* aleafstor* allconss* arootf* arootf2*	Binary flag for woody lifeform meters Final leaf allocation coefficient Leaf allocation coefficient to storage post-harvest used in CNAllocation Power to control the shape of the stem allocation curve Root allocation coefficient at start of fruit development Final root allocation	unitless unitless unitless unitless unitless unitless	1 0.01 0.3 1.5 0.2 0.08	Adjusted based on monthly biomass measurements (Zanotelli et al., 2013) Adjusted based on monthly biomass measurements (Zanotelli et
woody C and N allocation para aleaff* aleafstor* allconss* arootf* arootf2*	Binary flag for woody lifeform meters Final leaf allocation coefficient Leaf allocation coefficient to storage post-harvest used in CNAllocation Power to control the shape of the stem allocation curve Root allocation coefficient at start of fruit development Final root allocation coefficient until harvest	unitless unitless unitless unitless unitless unitless	1 0.01 0.3 1.5 0.2 0.08	Adjusted based on monthly biomass measurements (Zanotelli et al., 2013) Adjusted based on monthly biomass measurements (Zanotelli et al., 2013)
woody C and N allocation para aleaff* aleafstor* allconss* arootf* arootf2* arooti*	Binary flag for woody lifeform meters Final leaf allocation coefficient Leaf allocation coefficient to storage post-harvest used in CNAllocation Power to control the shape of the stem allocation curve Root allocation coefficient at start of fruit development Final root allocation coefficient until harvest Initial root allocation	unitless unitless unitless unitless unitless unitless unitless	1 0.01 0.3 1.5 0.2 0.08 0.7	Adjusted based on monthly biomass measurements (Zanotelli et al., 2013) Adjusted based on monthly biomass measurements (Zanotelli et al., 2013)
woody C and N allocation para aleaff* aleafstor* allconss* arootf* arootf2* arooti*	Binary flag for woody lifeform meters Final leaf allocation coefficient Leaf allocation coefficient to storage post-harvest used in CNAllocation Power to control the shape of the stem allocation curve Root allocation coefficient at start of fruit development Final root allocation coefficient until harvest Initial root allocation coefficient	unitless unitless unitless unitless unitless unitless unitless	1 0.01 0.3 1.5 0.2 0.08 0.7	Adjusted based on monthly biomass measurements (Zanotelli et al., 2013) Adjusted based on monthly biomass measurements (Zanotelli et al., 2013)

astemf*	Final stem allocation	unitless	0.22	Adjusted based on monthly biomass measurements (Zanotelli et
	coefficient]			al., 2013)
bfact*	Exponential factor used for	unitless	-0.5	Adjusted based on monthly biomass measurements (Zanotelli et
	fraction allocated to leaf			al., 2013)
declfact*	Decline factor to control the	unitless	4	Adjusted based on monthly biomass measurements (Zanotelli et
	shape of the stem allocation			al., 2013)
	curve			
fcur*	Fraction of C and N allocated	unitless	0.95	Tuned based on observed LAI and yield data (Zanotelli et al.,
	to the displayed pools			2013)
fleafi*	Initial leaf allocation	unitless	0.85	Adjusted based on monthly biomass measurements (Zanotelli et
	coefficient			al., 2013)
flivewd	Fraction of new wood that is		0.15	Same as BDT in CLM5
	live			
frootCN	Fine root C:N ratio	gC/gN	32	Average of 6 measurements (Zanotelli 2010, unpublished data)
grainCN*	Fruit C:N ratio	gC/gN	139	Average of 6 measurements (Zanotelli 2010, unpublished data)
leafCN*	Leaf C:N ratio	gC/gN	19.7	Average of 6 measurements (Zanotelli 2010, unpublished data)
lflitCN	Litter C:N ratio	gC/gN	60	Average of 4 measurements (Zanotelli 2010, unpublished data)
livewdCN	Livewood C:N ratio	gC/gN	60	Average of 6 measurements (Zanotelli 2010, unpublished data)
transplant	Initial carbon for crops	gC	5	
	transplanted from nursery			
Photosynthetic parameter	ers			
i_vcad*	Intercept of the relationship	$\mu molCO_2/m^2/s$	5.2	Adjusted in between BDT and crop
	between leaf N per unit area			
	and Vcmax25top			
medlynslope*	Medlyn slope of	$\mu mol H_2O/\mu mol CO_2$	8.2	Tuned based observed GPP and ET data (Zanotelli et al., 2015)
	conductance-photosynthesis			
	relationship			
s_vcad*	Slope of the relationship	µmolCO2/s/gN	34	Tuned based on observed LAI and yield data (Zanotelli et al.,
	between leaf N per unit area			2013)
	and Vcmax25top			
slatop*	Specific leaf area at top of	m^2/gC	0.028	Mean value for the growing season based on LAI and leaf biomass
	canopy			measurements (Zanotelli et al., 2013)

Vegetation structure and	d management				
displar	Ratio of displacement height	unitless	0.67	Same as BDT in CLM5	
	to canopy top height				
mulch_pruning	Binary flag for mulching (1)	unitless	1	Based on reported organic farming practices (Zanotelli et al.,	
	or export (0) of pruning			2013)	
	material				
prune_fr	Fraction of dead stem that is	unitless	0.85	Based on reported pruning quantity (Zanotelli et al., 2015)	
	pruned				
stockingnstem	Planting densityNumber of	trees#/ham2	3333<u>0.3</u>	Based on reported planting density (Zanotelli et al., 2013)	Formatiert: Hochgestellt
	trees per hectare		<u>3</u>		
taper	Ratio of stem height to radius		120	Based on reported stem diameter and tree allometry and height	
	at breast height			(Zanotelli et al., 2013)	
xl*	Leaf/stem orientation index	unitless	0.25	Same as BDT in CLM5	
z0mr*	Ratio of momentum	unitless	0.06	Based on average values reported for apple (La Fuente-Sáiz et al.,	
	roughness length to canopy			2017) and citrus (Tanny and Cohen, 2003) orchards	
	top height				
ztopmx	Maximum canopy height for	m	<u>3.6</u> 4	Based on reported tree heights (Zanotelli et al., 2013)	
	crops				
Optical parameters					
rholnir*	Leaf reflectance: near-IR	fraction	0.5	Based on average values for apple trees (Bastías and Corelli-	
				Grappadelli, 2012)	
rholvis*	Leaf reflectance: visible	fraction	0.1	Based on average values for apple trees (Bastías and Corelli-	
				Grappadelli, 2012)	
rhosnir*	Stem reflectance: near-IR	fraction	0.39	Same as BDT in CLM5	
rhosvis*	Stem reflectance: visible	fraction	0.16	Same as BDT in CLM5	
taulnir*	Leaf transmittance: near-IR	fraction	0.3	Based on average values for apple trees (Bastías and Corelli-	
				Grappadelli, 2012)	
taulvis*	Leaf transmittance: visible	fraction	0.04	Based on average values for apple trees (Bastías and Corelli-	
				Grappadelli, 2012)	
tausnir*	Stem transmittance: near-IR	fraction	0.001	Same as BDT in CLM5	
tausvis*	Stem transmittance: visible	fraction	0.001	Same as BDT in CLM5	

Respiration				
FUN_fracfixers*	The maximum fraction of	fraction	0.25	Same as BDT in CLM5
	assimilated carbon that can be			
	used to pay for N fixation			
lmr_intercept_atkin	Intercept in the calculation of	$\mu molCO_2/m^2/s$	1.756	Same as BDT in CLM5
	the top of canopy leaf			
	maintenance respiration base			
	rate.			

* parameters included in the sensitivity analysis

Code availability: The new sub-model CLM5-FruitTree is freely available via Zenodo, https://doi.org/10.5281/zenodo.6595378025014 (Dombrowski, 2022).

Author contribution: OD developed and modified the code for the sub-model; designed, performed, and analysed the simulations; and prepared the original draft of the manuscript. HB, HJHF, and CB supervised the research, and, together with DZ contributed to the manuscript writing through review & editing.

Competing interests: The authors declare no conflict of interest.

Acknowledgements: The authors gratefully acknowledge Damiano Zanotelli and his team for providing the field data of the apple orchard in the South Tyrol, Italy. The authors thank Yuanchao Fan for sharing the source files for his development of CLM-Palm (Fan et al., 2015) that aided the development of the new sub-model CLM5-FruitTree. Furthermore, the authors

17/0 CLM-Fain (ran et al., 2013) that adde the development of the new sub-inder CLM2-Future: Future: Future index, the additions thank Martin Thalheimer for providing the meteorological data from Laimburg meteorological station. This research was supported by the ATLAS project funded by the European Union's Horizon 2020 research and innovation programme under grant agreement No. 857125. Lastly, Harrie-Jan Hendricks Franssen acknowledges support from the Deutsche Forschungsgemeinschaft under Germany's Excellence Strategy, EXC-2070–390732324–PhenoRob.

775 References

Aguilera, E., Guzmán, G., and Alonso, A.: Greenhouse gas emissions from conventional and organic cropping systems in Spain. II. Fruit tree orchards, Agronomy for Sustainable Development, 35, 725-737, <u>https://doi.org/10.1007/s13593-014-0265-v,</u> 2015. Allen, M., Prusinkiewicz, P., and DeJong, T.: Using L-systems for modeling source-sink interactions, architecture and physiology of growing trees: the L-PEACH model, New phytologist, 166, 869-880, <u>https://doi.org/10.1111/j.1469-8137.2005.01348.x</u>, 2005.

- 780 Anderson, J., Richardson, E., and Kesner, C.: Validation of chill unit and flower bud phenology models for 'Montmorency' sour cherry, I International Symposium on Computer Modelling in Fruit Research and Orchard Management 184, 71-78, https://doi.org/10.17660/ActaHortic.1986.184.7, 1985.
- Atay, A. N., Koyuncu, F., and Atay, E.: Relative susceptibility of selected apple cultivars to alternate bearing, Journal of Biological and Environmental Sciences, 7, 81-86, 2013.
- 785 Atkin, O. K., Bloomfield, K. J., Reich, P. B., Tjoelker, M. G., Asner, G. P., Bonal, D., Bönisch, G., Bradford, M. G., Cernusak, L. A., and Cosio, E. G.: Global variability in leaf respiration in relation to climate, plant functional types and leaf traits, New Phytologist, 206, 614-636, <u>https://doi.org/10.1111/nph.13253</u>, 2015. Balsamo, G., Beliaars, A., Scipal, K., Viterbo, P., van den Hurk, B., Hirschi, M., and Betts, A. K.: A revised hydrology for the ECMWF

model: Verification from field site to terrestrial water storage and impact in the Integrated Forecast System, Journal of hydrometeorology, 10, 623-643, <u>https://doi.org/10.1175/2008JHM1068.1</u>, 2009.

Bastías, R. M. and Corelli-Grappadelli, L.: Light quality management in fruit orchards: physiological and technological aspects, Chilean Journal of Agricultural Research, 72, 574, 2012.

Belleggia, A., Murri, G., and Neri, D.: Crop load control of 'Fuji' apple in central Italy, XI International Symposium on Plant Bioregulators in Fruit Production 884, 399-405, https://doi.org/10.17660/ActaHortic.2010.884.48, 2009.

795 Benyei, P., Cohen, M., Gresillon, E., Angles, S., Araque-Jiménez, E., Alonso-Roldán, M., and Espadas-Tormo, I.: Pruning waste management and climate change in Sierra Mágina's olive groves (Andalusia, Spain), Regional environmental change, 18, 595-605, https://doi.org/10.1007/s10113-017-1230-5, 2018.

 Bepete, M. and Lakso, A.: Apple fruit respiration in the field: relationships to fruit growth rate, temperature, and light exposure, VI International Symposium on Integrated Canopy, Rootstock, Environmental Physiology in Orchard Systems 451, 319-326, https://doi.org/10.17660/ActaHortic.1997.451.37, 1996.

Blyth, E. M., Arora, V. K., Clark, D. B., Dadson, S. J., De Kauwe, M. G., Lawrence, D. M., Melton, J. R., Pongratz, J., Turton, R. H., and Yoshimura, K.: Advances in land surface modelling, Current Climate Change Reports, 7, 45-71, https://doi.org/10.1007/s40641-021-00171-5, 2021.

Boas, T., Bogena, H., Grünwald, T., Heinesch, B., Ryu, D., Schmidt, M., Vereecken, H., Western, A., and Hendricks Franssen, H.-J.:
 Improving the representation of cropland sites in the Community Land Model (CLM) version 5.0, Geoscientific Model Development, 14, 573-601, https://doi.org/10.5194/gmd-14-573-2021, 2021.

Bryś, K., Bryś, T., and Ojrzyńska, H.: Long-wave radiation balances of grassy surface and bare soil in Wrocław, E3S Web of Conferences, 1-8, https://doi.org/10.1051/e3sconf/201911600012, 2019.

Formatiert: Englisch (Vereinigtes Königreich)	
Formatiert: Englisch (Vereinigtes Königreich)	
Feldfunktion geändert	
Feldfunktion geändert	
Formatiert: Englisch (Vereinigtes Königreich)	
Formatiert: Englisch (Vereinigtes Königreich)	
Feldfunktion geändert	
Formatiert: Englisch (Vereinigtes Königreich)	
Formatiert: Englisch (Vereinigtes Königreich)	
Feldfunktion geändert	
Formatiert: Englisch (Vereinigtes Königreich)	
Formatiert: Englisch (Vereinigtes Königreich)	
Feldfunktion geändert	
Formatiert: Englisch (Vereinigtes Königreich)	
Formatiert: Englisch (Vereinigtes Königreich)	
Feldfunktion geändert	
Formatiert: Englisch (Vereinigtes Königreich)	
Formatiert: Englisch (Vereinigtes Königreich)	
Feldfunktion geändert	
Formatiert: Englisch (Vereinigtes Königreich)	
Formatiert: Englisch (Vereinigtes Königreich)	
Feldfunktion geändert	
Formatiert: Englisch (Vereinigtes Königreich)	
Formatiert: Englisch (Vereinigtes Königreich)	
Feldfunktion geändert	
Formatiert: Englisch (Vereinigtes Königreich)	
Formatiert: Englisch (Vereinigtes Königreich)	
Feldfunktion geändert	
Formatiert: Englisch (Vereinigtes Königreich)	
Formatiert: Englisch (Vereinigtes Königreich)	
Feldfunktion geändert	
Formatiert: Englisch (Vereinigtes Königreich)	
Formatiert: Englisch (Vereinigtes Königreich)	

Formatiert: Englisch (Vereinigtes Königreich)

		F	ormatiert	
1	Bwalya, J. M.: Estimation of net carbon sequestration potential of citrus under different management systems using the life cycle approach,	F	eldfunktion geändert	
810	M.S. thesis, University of Zambia, <u>http://thesisbank.jhia.ac.ke/id/eprint/8503</u> , 2012.	F	ormatiert	
	Carranca, C., Brunetto, G., and Tagliavini, M.: Nitrogen nutrition of fruit trees to reconcile productivity and environmental concerns, Plants,	F	eldfunktion geändert	
	7, 1-12, https://doi.org/10.3390/plants7010004, 2018.	F	eldfunktion geändert	
015	Ceccon, C., Panzacchi, P., Scandellari, F., Prandi, L., Ventura, M., Russo, B., Millard, P., and Tagliavini, M.: Spatial and temporal effects	/ }	armatiant	
815	of soil temperature and moisture and the relation to fine root density on root and soil respiration in a mature apple orchard, Plant and Soil, 342, 195-206, <u>https://doi.org/10.1007/s11104-010-0684-8</u> , 2011.		ormatiert	
	Cerutti, A. K., Beccaro, G. L., Bruun, S., Bosco, S., Donno, D., Notarnicola, B., and Bounous, G.: Life cycle assessment application in the	/		
	fruit sector: state of the art and recommendations for environmental declarations of fruit products, Journal of cleaner production, 73, 125- 125, https://dx.doi.org/10.1016/j.doi.org/10.0017_2014		eldfunktion geandert	
820	Cesaracio, C., Spano, D., Styder, R. L., and Duce, P.: Chilling and forcing model to predict hud-hurst of crop and forest species.	F	eldfunktion geändert	
	Agricultural and Forest Meteorology, 126, 1-13, https://doi.org/10.1016/j.agrformet.2004.03.002, 2004.	- (F	ormatiert	
	Charrier, G., Martin-Stpaul, N., Damesin, C., Delpierre, N., Hänninen, H., Torres-Ruiz, J. M., and Davi, H.: Interaction of drought and frost		eldfunktion geändert	
	in tree ecophysiology: rethinking the timing of risks, Annals of Forest Science, 78, 1-15, <u>https://doi.org/10.1007/s13595-021-01052-5, 2021.</u>	F	ormatiert	
825	Cheng, F., Huang, M., Chen, M., Guan, K., Bernacch, C., Peng, B., and Tan, Z.: raraneterizing perennia bioenergy crops in Version 5 of the Community Land Model based on site-level observations in the Central Midwestern United States, Journal of Advances in Modeling Earth Systems, 12, 1-24, <u>https://doi.org/10.1029/2019MS001719</u> , 2020. Chmielewski, FM., Blümel, K., and Pálešová, I.: Climate change and shifts in dormancy release for deciduous fruit crops in Germany,		eldfunktion geändert	
			ormatiert	
	Climate research, 54, 209-219, https://doi.org/10.3354/cr01115_2012.	- [F	ormatiert	
830	conditions (kevnote). XXVI International Horicultural Congress: Kev Processes in the Growth and Cropping of Deciduous Fruit and Nut	F	eldfunktion geändert	
	Trees 636, 425-441, https://doi.org/10.17660/ActaHortic.2004.636.52, 2002.	F	eldfunktion geändert	
	Corelli-Grappadelli, L. and Marini, R.: 11 Orchard Planting Systems, in: The peach: botany, production and uses, CABI, Pennsylvania, U.S., 264, 2008	~ (F	ormatiert	
	Dagon, K., Sanderson, B. M., Fisher, R. A., and Lawrence, D. M.: A machine learning approach to emulation and biophysical parameter estimation with the Community Land Model, version 5, Advances in Statistical Climatology, Meteorology and Oceanography, 6, 223-244, https://doi.org/10.5104/acemp.6.223.2020_2020		eldfunktion geändert	
835			ormatiert	
	https://doi.org/10.179/astructure/2219/2020. DeJong, T. M. and Grossman, Y. L.: A supply and demand approach to modeling annual reproductive and vegetative growth of deciduous	F	eldfunktion geändert	
	fruit trees, HortScience, 29, 1435-1442, 1994.	F	ormatiert	
0.40	Demestihas, C., Plénet, D., Génard, M., Raynal, C., and Lescourret, F.: Ecosystem services in orchards. A review, Agronomy for Sustainable		aldfunktion goöndort	
840	Development, 37, 1-21, https://doi.org/10.100//s13593-017-0422-1_2017.	/ -		
	Diez-ratet, L, Fuiles, L, Save, K., Biel, C., de renaue, F., Mianau, A., Vaigas, F., Avia, G., Caloo, J., and Alanda, A., Biooning unice Mediterratena climate: Estimating cultivar-specific chill and heat requirements of almond and apple trees using a statistical approach.		ormatiert	
	Agronomy, 9, 760-781, https://doi.org/10.3390/agronomy9110760, 2019.	F	eldfunktion geändert	
845	Dombrowski, O.: odombro/TSW: CLMS-FruitTree: A new sub-model for deciduous fruit trees, Zenodo, https://doi.org/10.5281/zenodo.6025014_2022	F	ormatiert	
	El Jaouhari, N., Abouabdillah, A., Bouabid, R., Bourioug, M., Aleya, L., and Chaoui, M.: Assessment of sustainable deficit irrigation in a	F	eldfunktion geändert	
	Moroccan apple orchard as a climate change adaptation strategy, Science of the total environment, 642, 574-581,	F	ormatiert	
	https://doi.org/10.1016/j.scitotenv.2018.06.108_2018.	F	eldfunktion geändert	
850	in the LPJmL model, Geoscientific Model Development, 8, 3545-3561, <u>https://doi.org/10.5194/gmd-8-3545-2015</u> , 2015.	F	ormatiert	
	Fan, Y., Roupsard, O., Bernoux, M., Le Maire, G., Panferov, O., Kotowska, M. M., and Knohl, A.: A sub-canopy structure for simulating	F	ormatiert	
	https://doi.org/10.5194/gmd-8-3785-2015, 2015.	F	eldfunktion geändert	
055	FAO Crops and livestock products: https://www.fao.org/faostat/en/#data/QCL_last access: 22 December 2021.	F	eldfunktion geändert	
800	Faust, M., Erez, A., Kowland, L. J., Wang, S. Y., and Norman, H. A.: Bud dormancy in perennial fruit frees: physiological basis for dormancy induction, maintenance, and release, HortScience, 32, 623-629, 1997.	F	ormatiert	
	Fisher, R. A. and Koven, C. D.: Perspectives on the future of land surface models and the challenges of representing complex terrestrial	F	eldfunktion geändert	
	systems, Journal of Advances in Modeling Earth Systems, 12, 1-24, <u>https://doi.org/10.1029/2018MS001453</u> , 2020. Glover, J. D., Reganold, J., Bell, L., Borevitz, J., Brummer, E., Buckler, E. S., Cox, C., Cox, T. S., Crews, T., and Culman, S.: Increased	(F	ormatiert	
860	food and ecosystem security via perennial grains, Science, 328, 1638-1639, https://doi.org/10.1126/science.1188761, 2010.	F	eldfunktion geändert	
	Göhler, M., Mai, J., and Cuntz, M.: Use of eigendecomposition in a parameter sensitivity analysis of the Community Land Model, Journal of Geophysical Research: Biogeosciences, 118, 904-921, https://doi.org/10.1002/jgrg.20072, 2013.	F	ormatiert	 ()
	Goldschmidt, E. E. and Lakso, A. N.: Fruit tree models: scope and limitations, Information and Communication Technology (ICT)	F	eldfunktion geändert	
I	Development and Adoption: Perspectives of Technological Innovation, 1-19, 2005.	F	 ormatiert	
		Ċ		()
	39			

865	Grechi, I., Sauge, M. H., Sauphanor, B., Hilgert, N., Senoussi, R., and Lescourret, F.: How does winter pruning affect peach tree-Myzus		Feldfunktion geändert	
	persicae interactions?, Entomologia Experimentalis et Applicata, 128, 369-379, https://doi.org/10.1111/j.1570-7458.2008.00720.x, 2008.		Formatiert	
	Greer, D. H., Wünsche, J. N., and Halligan, E. A.: Influence of postharvest temperatures on leaf gas exchange, carbohydrate reserves and			
	anocations, subsequent buobreak, and irrut yield of Braeburn appre (Mausu domesuica) irress, New Zealand Journal of Crop and Horiticultural Science 30 (175,185, bttps://doi.org/10.108/001140/671.2002.9514213.2002		Formatiert	
870	Hammad, H. M., Nauman, H. M. F., Abbas, F., Ahmad, A., Bakhat, H. F., Saeed, S., Shah, G. M., Ahmad, A., and Cerdà, A.: Carbon		Feldfunktion geändert	
070	sequestration potential and soil characteristics of various land use systems in arid region, Journal of environmental management, 264,		Feldfunktion geändert	
	110254, https://doi.org/10.1016/j.jenvman.2020.110254, 2020.		Formatiert	
	Heide, U. M. and Prestrud, A. K.: Low temperature, but not photoperiod, controls growth cessation and dormancy induction and release in annle and pear. Tree physicalogy 25, 109-114. https://doi.org/10.1093/treenbw/25, 1.109.2005		Formatiert	
875	hoblyn, T., Grubb, N., Painter, A., and Wates, B.: Studies in biennial bearing, —I. Journal of pomology and horticultural science, 14, 39-76,		Foldfunktion goöndort	
	https://doi.org/10.1080/03683621.1937.11513464, 1937.			
	Hou, T., Wang, Y., Guo, F., Jia, Q., Wu, X., Wang, E., and Hong, J.: Soil Respiration Characteristics and Influencing Factors for Apple		Feldfunktion geändert	
	Orchards in Different Regions on the Loess Plateau of Shaahxi Province, Sustainability, 15, $1-21$, <u>https://doi.org/10.3590/su15094780, 2021</u> .		Formatiert	
880	research (7, 295-304, https://doi.org/10.1029/WR017i002p00295, 1981.	$\langle \backslash \uparrow$	Feldfunktion geändert	
	Jackson, D.: Flowers and fruit, in: Temperate and subtropical fruit production., CABI, Cambridge, U.K., 34-43, 2011.	$\langle \rangle \rangle$	Formatiert	
	Kazimierski, P., Hercel, P., Suchocki, T., Smoliński, J., Pladzyk, A., Kardaš, D., Łuczak, J., and Januszewicz, K.: Pyrolysis of Pruning		Formatiert	
	Residues from various types of Orchards and Pretreatment for Energetic Use of Biochar, Materials, 14, 1-14, https://doi.org/10.3390/ma14112969.2021		Feldfunktion geändert	
885	Konzelmann, T., van de Wal, R. S., Greuell, W., Bintanja, R., Henneken, E. A., and Abe-Ouchi, A.: Parameterization of global and longwave			
	incoming radiation for the Greenland Ice Sheet, Global and Planetary change, 9, 143-164, https://doi.org/10.1016/0921-8181(94)90013-2	\mathbb{N}	Formatiert	<u>[</u>
	1994. V zaloveli T. Cashabudata courses and sinks in mandu plants. The hotonical ration, 58, 107, 222, 1002	\langle / \rangle	Feldfunktion geändert	
	Kolzowski, L. Caroonyunae sources and sinks in woody plants, the botanical review, 56, 107-222, 1592.		Formatiert	
890	global vegetation model for studies of the coupled atmosphere-biosphere system, Global Biogeochemical Cycles, 19, 1-33,		Feldfunktion geändert	
	https://doi.org/10.1029/2003GB002199_2005.	L Y	Formatiert	
	Ia ruente-Saiz, D., Ortega-rarias, S., Fonseca, D., Ortega-Saiazar, S., Kilic, A., and Alien, K.: Calibration of METRIC model to estimate energy halance over a drin, irrigated annle orchard Remate Sensing 9, 1,18, bitrs://doi.org/10.3300/rs6070670, 2017		Feldfunktion geändert	
	Lakso, A., White, M., and Tustin, D.: Simulation modeling of the effects of short and long-term climatic variations on carbon balance of		Formatiert	
895	apple trees, VII International Symposium on Orchard and Plantation Systems 557, 473-480,	$\langle \rangle$		
	https://doi.org/10.17660/ActaHortic.2001.557.63, 2000.		Feidfunktion geandert	
	Latso, A., winische, J., Fahler, J., and Grappaden, E. C., Weastrement and modering of carbon balance of the apple tree, fortiscience, 54, 1040-1047, 1999.	\sim	Formatiert	
	Lasslop, G., Migliavacca, M., Bohrer, G., Reichstein, M., Bahn, M., Ibrom, A., Jacobs, C., Kolari, P., Papale, D., and Vesala, T.: On the		Feldfunktion geändert	
900	choice of the driving temperature for eddy-covariance carbon dioxide flux partitioning, Biogeosciences, 9, 5243-5259,	Ì	Formatiert	
	Jauper, 2010. S194: 095-5243-2012, 2012.	<	Feldfunktion geändert	
	L.: Technical description of version 5.0 of the Community Land Model (CLM), Natl. Cent. Atmospheric Res. (NCAR), available at:		Formatiert	
005	http://www.cesm.ucar.edu/models/cesm2/land/CLM50_Tech_Note.pdf (last access: 2 May 2021), 2018.		Feldfunktion geändert	
905	Le Roux, X., Lacointe, A., Escobar-Gutterrez, A., and Le Dizes, S.: Carbon-based models of individual tree growth: a critical appraisal, Annals of Ecorest Science 58, 469-506 https://doi.org/10.1051/fcoast-2001140.2001	\searrow	Formatiert	
	Ledo, A., Smith, P., Zerihun, A., Whitaker, J., Vicente-Vicente, J. L., Qin, Z., McNamara, N. P., Zinn, Y. L., Llorente, M., and Liebig, M.:	$\langle \cdot \rangle$		
	Changes in soil organic carbon under perennial crops, Global change biology, 26, 4158-4168, https://doi.org/10.1111/gcb.15120, 2020.		Feidfunktion geandert	
010	Li, F., Cohen, S., Naor, A., Shaozong, K., and Erez, A.: Studies of canopy structure and water use of apple trees on three rootstocks,	\swarrow	Formatiert	
910	ricultural water Management, 55, 1-14, <u>https://doi.org/10.1016/S05/8-5/74(01)00184-6</u> , 2002.		Feldfunktion geändert	
	Germany, IDEMA DELIVERABLE No. 17, 2006.	$\langle \rangle$	Formatiert	
	Loescher, W. H., McCamant, T., and Keller, J. D.: Carbohydrate reserves, translocation, and storage in woody plant roots, HortScience, 25,		Feldfunktion geändert	
915	81, 1990. ardazzi D.I. Lu V. Laurence P.I. Laurence D.M. Swenson S. Oleson K.W. Wieder W.R. and Ainsworth F.A. Simulating		Formatiert	
15	agriculture in the Community Land Model version 5, Journal of Geophysical Research: Biogeosciences, 125, 1-19		Feldfunktion geändert	
	https://doi.org/10.1029/2019JG005529, 2020.		Formation	
	López-Olivari, R., Ortega-Farías, S., and Poblete-Echeverría, C.: Partitioning of net radiation and evapotranspiration over a superintensive			<u>l</u>
I	anp-imgated on ve orchard, imgation science, 54, 17-51, <u>https://doi.org/10.1007/s00271-015-0484-2</u> 2016.	\leq	Feldfunktion geändert	
			Formatiert	

1		
920	Luo, Q., Wen, J., Hu, Z., Lu, Y., and Yang, X.: Parameter sensitivities of the Community Land Model at two alpine sites in the three-river	
	source region, Journal of Meteorological Research, 34, 851-864, https://doi.org/10.100//s13351-020-9205-8, 2020.	/
	Maestre-Valero, J. F., Testi, L., Jimenez-Bello, M. A., Castel, J. R., and Intrigliolo, D. S.: Evapotranspiration and carbon exchange in a citrus	/
	orchard using eduy covariance, irrigation Science, 55, 597-406, <u>mitps://doi.org/10.100//S00271-017-0548-6</u> , 2017.	
025	Matagui, D., Minard, P., Wendier, R., Helpouri, A., and Tagnavini, M.: Translocation of amino actos in the xytem of apple (Matus domestica Backb) tracks in spring as a consequence of both N remebilization and not unreal of Europingate Retarguing 22 1665 1671	
923	bitski, just sin spring as a consequence of both is remoonization and foot uptake, Journal of Experimental Botany, 52, 1005-1071,	/
	https://doi.org/10.1095/jfeX00/32.301.1005_2001.	/
	matun-odniz, B., Gonzalez-Real, M. M., Egea, G., and Banne, A.: Ecosystem respiration of old and young irrigated circus orchards in a carsing the second and second se	
	semant children Agricultural and Forest Meteorology, 200, 1-15, $\frac{1}{2}$ (the <i>S</i> -doctograd-torological control agricultural and Forest Meteorology, 200, 1-15, $\frac{1}{2}$ (the <i>S</i> -doctograd-torological control agricultural and Forest Meteorology, 200, 1-15, $\frac{1}{2}$ (the <i>S</i> -doctograd-torological control agricultural and Forest Meteorology, 200, 1-15, $\frac{1}{2}$ (the <i>S</i> -doctograd-torological control agricultural agr	_
030	Minaut, F., Ecophysiology of the memara cycing of introgen for the growth, Zenschift für Frianzenerhandung und Bodenkunde, 159, 1- 10. https://doi.org/10.1002/1016.1091.0906.3581500102.1006	
250	10, antips.//doi.org/10.1002/jfni.1790.301320102 1790. Millard D. Wandler R. Grassi G. Grelet G. A. and Tagliavini M. Translocation of nitrogen in the xylem of field-grown cherry and	~
	milau, 1., weinter, R., Orassi, O., Oreret, OA., and Tagnavini, M.: Hanstocation of inflogen in the xytem of neur-grown cherry and poplar transmission reambility and the state of the s	
	Monselies S and Foldochundit F - Alternate beging in fuit trace Horticultural reviews 4, 1982	
	Montagnani L. Zanotelli D. Tagliavini M. and Tomelleri F.: Timescale effects on the environmental control of carbon and water fluxes	
935	of an apple orchard Ecology and evolution 8, 416-434, https://doi.org/10.1002/ece3.3633, 2018	
155	Montanaro G Celano G Dichio B and Xilovannis C. Effects of soil-protecting activity Loro.	~
	productivity in fruit tree orchards L and Degradation & Development 21 132-138 https://doi.org/10.1002/dr 917-2010	/
	Musachi S, and Serra S: Apple fruit quality: Overview on pre-baryest factors Scientia Horizoluturae 234 409-430	_
	https://doi.org/10.1016/j.scienta.2017.2.057.2018	
940	Neumann, L.: Improving the apple carbon balance model MaluSim by integrating and testing water deficit and fruit drop effects. M.A. thesis,	_
	Hannover: Institutionelles Repositorium der Leibniz Universität Hannover, Hannover, Germany, 147 pp., https://doi.org/10.15488/10194	
	2020.	~
	Noilhan, J. and Mahfouf, JF.: The ISBA land surface parameterisation scheme, Global and planetary Change, 13, 145-159,	
	https://doi.org/10.1016/0921-8181(95)00043-7, 1996.	
945	Ntshidi, Z., Dzikiti, S., Mazvimavi, D., and Mobe, N.: Contribution of understorey vegetation to evapotranspiration partitioning in apple	
	orchards under Mediterranean climatic conditions in South Africa, Agricultural Water Management, 245, 1-11,	
	https://doi.org/10.1016/j.agwat.2020.106627, 2021.	_
	O'Connell, M. and Scalisi, A.: Sensing fruit and tree performance under deficit irrigation in 'September Bright' nectarine, International	
	Symposium on Precision Management of Orchards and Vineyards 1314, 9-16, https://doi.org/10.17660/ActaHortic.2021.1314.2, 2019.	_
950	Oliveira, C. M. and Priestley, C. A.: Carbohydrate reserves in deciduous fruit trees, Hort. Rev, 10, 403-430, 1988.	
	Oorthuis, R., Vaunat, J., Hürlimann, M., Lloret, A., Moya, J., Puig-Polo, C., and Fraccica, A.: Slope orientation and vegetation effects on	
	soil thermo-hydraulic behavior. An experimental study, Sustainability, 13, 1-14, <u>https://doi.org/10.3390/su13010014</u> , 2021.	~
	Pallas, B., Da Silva, D., Valsesia, P., Yang, W., Guillaume, O., Lauri, PE., Vercambre, G., Génard, M., and Costes, E.: Simulation of	
055	carbon allocation and organ growth variability in apple tree by connecting architectural and source–sink models, Annals of Botany, 118,	
955	317-330, https://doi.org/10.1093/a0b/mcw085, 2016.	~
	Pasa, M. d. S., Carra, B., Brignenti, A. F., Pinto, F. A. M. F., Meilo-Farlas, P. C. d., and Herter, F. G.: Emephon as a potential tool to manage alternet hearing of E-bill employable targe Residence of the second seco	
	auternate bearing of Full apple trees, Revisita Ceres, 68, 100-184, <u>https://doi.org/10.159/0054-757A202188050005</u> , 2021.	~
	Penzel, M., Lakso, A. N., Isounas, N., and Zude-Sasse, M.: Carbon consumption of developing trutt and the fruit bearing capacity of individual Pollo 215 and Bingua apple trace. Lit A graphic, 24, 400, 422, https://doi.org/10.21545/integr/10.21545/	
960	mututual Korto 5015 and Finova apple trees, int. Agrophiss, 34 , 405 - 425 , $\frac{1015}{M}$, $\frac{1015}{M}$, $\frac{1012}{M}$, $\frac{1012}{M}$, $\frac{102}{M}$, $\frac{1012}{M}$, 1012	~
200	modelling Atmospheric Chamistry and Wang 1.51, S087-6005 https://doi.org/10.510//gop.15.5087.2015.2015	1
	Reichstein M Falder E Baldocchi D Panale D Aubinet M Berbiere P Bernhofer C Buchmann N Gilmanov T and Granier	
	A · On the separation of net ecosystem exchange into assimilation and ecosystem respiration: review and improved algorithm Global changes	1
	biology 11 1424-1439 https://doi.org/10.1111/j.1365-24486.2005.001002 x 2005	
965	Reves, F., Delong, T., Franceschi, P., Tagliavini, M., and Gianelle, D.: Maximum growth potential and periods of resource limitation in	
	apple tree. Frontiers in Plant Science, 7, 1-12, https://doi.org/10.3389/fpls.2016.00233, 2016.	1
	Sanchez, E. E., Khemira, H., Sugar, D., and Righetti, T. L.: Nitrogen management in orchards, in: Nitrogen Fertilization in the Environment,	
	Marcel Dekker, Inc., New York, U.S., 1995.	
	Scandellari, F., Caruso, G., Liguori, G., Meggio, F., Palese, A. M., Zanotelli, D., Celano, G., Gucci, R., Inglese, P., and Pitacco, A.: A survey	
970	of carbon sequestration potential of orchards and vineyards in Italy, European Journal of Horticultural Science, 81, 106-114,	
	https://doi.org/10.17660/eJHS.2016/81.2.4, 2016.	
	Schaphoff, S., von Bloh, W., Rammig, A., Thonicke, K., Biemans, H., Forkel, M., Gerten, D., Heinke, J., Jägermeyr, J., and Knauer, J.:	1
	LPJmL4-a dynamic global vegetation model with managed land-Part 1: Model description, Geoscientific Model Development, 11, 1343-	
	1375, https://doi.org/10.5194/gmd-2017-145, 2018.	_
		1

	/	Feldfunktion geändert	
ites in the three river		Formatiert	
sites in the three-fiver		Formatiert	
n exchange in a citrus		Feldfunktion geändert	
pple (Malus domestica		Feldfunktion geändert	
tany, 52, 1665-1671,		Formatiert	
d citrus orchards in a		Formatiert	
020.		Feldfunktion geändert	
Bodenkunde, 159, 1-		Feldfunktion geändert	$\overline{\Box}$
eld-grown cherry and		Formatiert	Ē
0 0		Feldfunktion geändert	$\overline{\square}$
rbon and water fluxes		Formatiert	Ĩ
roon and water maxes		Feldfunktion geändert	H
l organic carbon and		Formatiert	H
irae, 234, 409-430,		Formatiert	H
		Feldfunktion geändert	H
p effects, M.A. thesis, org/10 15488/10194		Feldfunktion geändert	<u> </u>
	<	Formationt	<u> </u>
hange, 13, 145-159,			<u> </u>
partitioning in apple	<		<u> </u>
ement, 245, 1-11,		Formatiert	<u> </u>
ctarine. International	<	reidfunktion geandert	<u> </u>
<u>021.1314.2</u> , 2019.		Formatiert	<u> </u>
vegetation effects on	\searrow	Feldfunktion geandert	<u></u>
2021.		Formatiert	<u> </u>
tes, E.: Simulation of		Feldfunktion geändert	<u></u>
nnais of Botany, 118,		Formatiert	
tential tool to manage		Feldfunktion geändert	
2021. t bearing capacity of		Formatiert	
020.	\sim	Feldfunktion geändert	
neration land-surface	K,	Formatiert	
nov, T., and Granier,	$\langle \rangle$	Formatiert	<u></u>
orithm, Global change	$\langle \rangle$	Feldfunktion geändert	
resource limitation in		Feldfunktion geändert	
	\mathbb{N}	Formatiert	
n in the Environment,	//	Feldfunktion geändert	
Pitacco, A.: A survey	//)	Formatiert	
cience, 81, 106-114,		Feldfunktion geändert)
yr, J., and Knauer. J.:		Formatiert	$\overline{\Box}$
velopment, 11, 1343-		Feldfunktion geändert	
		Formatiert	
		Feldfunktion geändert	
		Formatiert	Ä
			<u> </u>

975	Sedlar, J. and Hock, R.: Testing longwave radiation parameterizations under clear and overcast skies at Storglaciären, Sweden, The				
	Cryosphere, 3, 75-84, https://doi.org/10.5194/tc-3-75-2009, 2009.		Forn	natiert	
	Segovia-Cardozo, D. A., Franco, L., and Provenzano, G.: Detecting crop water requirement indicators in irrigated agroecosystems from soil water content profiles: An application for a citrus orchard, Science of The Total Environment, 806,		Feld	lfunktion geändert	
	https://doi.org/10.1016/j.scitotenv.2021.150492, 2022.		Forn	natiert	
980	Song, J., Miller, G. R., Cahill, A. T., Aparecido, L. M. T., and Moore, G. W.: Modeling land surface processes over a mountainous rainforest		Feld	lfunktion geändert	
	in Costa Rica using CLM4. 5 and CLM5, Geoscientific Model Development, 13, 5147-5173, https://doi.org/10.5194/gmd-13-5147-2020_ 2020	<u></u>	Feld	lfunktion geändert	
	Tanny, J. and Cohen, S.: The effect of a small shade net on the properties of wind and selected boundary layer parameters above and within	\sim	-		
	a citrus orchard, Biosystems Engineering, 84, 57-67, https://doi.org/10.1016/S1537-5110(02)00233-7, 2003.		Forn	natiert	
985	Thornton, P. E., Law, B. E., Gholz, H. L., Clark, K. L., Falge, E., Ellsworth, D. S., Goldstein, A. H., Monson, R. K., Hollinger, D., and Falk,		Forn	natiert	
	M.: Modeling and measuring the effects of disturbance history and climate on carbon and water budgets in evergreen needlelear forests, Arricultural and forest metagenove 113, 185-222, 2002		Feld	lfunktion geändert]
	Tomè, E., Ventura, M., Folegot, S., Zanotelli, D., Montaenani, L., Mimmo, T., Tonon, G., Tagliavini, M., and Scandellari, F.: Mycorrhizal				
	contribution to soil respiration in an apple orchard, Applied Soil Ecology, 101, 165-173, https://doi.org/10.1016/j.apsoil.2016.01.016, 2016.		Feld	lfunktion geändert	
990	Tromp, J.: Nutrient reserves in roots of fruit trees, in particular carbohydrates and nitrogen, Plant and Soil, 71, 401-413,		Form	nationt	
	pttps://doi.org/10.1007/BF02182682, 1983.				
	Turrini, A., Caruso, G., Avio, E., Gennai, C., Palla, M., Agnolucci, M., Iomei, P. E., Giovannetti, M., and Gucci, R.: Protective green cover anhances scil aceptization and native mycorrhizal participation of the high design of the high design of the scilar participation of the sci	\backslash	Feld	lfunktion geändert	
	emances son respiration and narve mycontract potential compared with son unage in a night-density on ve of chard in a long term study, Applied Soil Ecology, 116, 70–78, https://doi.org/10.1016/i.apsoil.2017.04.001.2017		Forn	natiert	
995	Valancogne, C., Dayau, S., Pieri, P., Ferreira, M. I., Silvestre, J., and Angelocci, L.: Influence of orchard and vineyard characteristics on		Feld	lfunktion geändert	
	maximal plant transpiration, 3. International symposium on irrigation of Horticultural crops, Lisbonne, Portugal, <u>https://hal.inrae.fr/hal-02770605_1000</u>		Forn	natiert	
	UZ/10005 1999. Vioux N - CRUNCEP Version 7 - Atmospheric Forcing Data for the Community Land Model Research Data Archive at the National Center	$\langle \rangle$	Form	matiert: Englisch (Vereinigtes Königreich)	
	for Atmospheric Research, Computational and Information Systems Laboratory, https://doi.org/10.5065/PZ8F-F017, 2018.		Cald		
1000	Wheaton, T., Castle, W., Whitney, J., Tucker, D., and Muraro, R.: A high density citrus planting, Proc. Fla. State Hort. Soc, Florida, U.S.,	//	reiu		
	55-59, 1990.		Forn	matiert: Englisch (Vereinigtes Konigreich)	
	Wu, I., Wang, Y., Yu, C., Chiarawipa, R., Zhang, X., Han, Z., and Wu, L.: Carbon sequestration by fruit trees-Chinese apple orchards as an example PLoS one 7, 1-13 https://doi.org/10.1371/journal.neu.003883.2012		Feld	lfunktion geändert	
	winsche J. N. and Lakso, A. N.: Apple the physiology-implications for orchard and tree management, Compact Fruit Tree, 33, 82-88,	\checkmark '	Forn	natiert	
1005	2000.	$\langle \rangle$	Feld	lfunktion geändert	
	Yasin, G., Farrakh Nawaz, M., Zubair, M., Qadir, I., Saleem, A. R., Ijaz, M., Gul, S., Amjad Bashir, M., Rehim, A., and Rahman, S. U.: Assessing the Contribution of Citrus Orchards in Climate Change Mitigation through Carbon Sequestration in Sargodha District, Pakistan, Sustainability, 13, 1-10, https://doi.org/10.3300/su132212412, 2021		Forn	natiert	
			Fold	lfunktion geändert	
	Zanotelli, D., Montagnani, L., Andreotti, C., and Tagliavini, M.: Evapotranspiration and crop coefficient patterns of an apple orchard in a	\langle	-		
1010	sub-humid environment, Agricultural Water Management, 226, 1-11, https://doi.org/10.1016/j.agwat.2019.105756, 2019.		Forn	natiert	<u> (</u>)
	Zanotelli, D., Montagnani, L., Manca, G., and Tagliavini, M.: Net primary productivity, allocation pattern and carbon use efficiency in an apple orthogened assessed by integrating eddy coursing and continuous soil chamber measurements. Biogenetic and continuous soil chamber measurements.		Feld	lfunktion geändert	
	appresentation appresentation of the second and a second and a second appresentation of the second appr		Form	natiert	
	Zanotelli, D., Montagnani, L., Manca, G., Scandellari, F., and Tagliavini, M.: Net ecosystem carbon balance of an apple orchard, European		Feld	lfunktion geändert	
1015	Journal of Agronomy, 63, 97-104, http://dx.doi.org/10.1016/j.eja.2014.12.0002, 2015.		Form	natiert	
	Based on Growing Degree-day Accumulation, Journal of the American Society for Horticultural Science, 131, 601-607,		Feld	lfunktion geändert	
	https://doi.org/10.21273/JASHS.131.5.601, 2006.		Form	natiert	
1020	Leng, I., Are, Z., and Liu, S.: Seasonal effects of irrigation on land-atmosphere latent neat, sensible neat, and carbon fluxes in semiarid basin, Earth System Dynamics, 8, 113-127, https://doi.org/10.5194/esd-8-113-2017, 2017.		Feld	lfunktion geändert	
	an a	//	Forn	natiert	
			Feld	lfunktion geändert	
			Form	natiert	
				nation	