

Improving snow albedo modeling in E3SM land model (version 2.0) and assessing its impacts on snow and surface fluxes over the Tibetan Plateau

5 Dalei Hao¹, Gautam Bisht¹, Karl Rittger², Edward Bair³, Cenlin He⁴, Huilin Huang¹, Cheng Dang⁵, Timbo Stillinger³, Yu Gu⁶, Hailong Wang¹, Yun Qian¹ and L. Ruby Leung¹

¹Atmospheric Sciences and Global Change Division, Pacific Northwest National Laboratory, Richland, WA, USA

²Institute for Arctic and Alpine Research, University of Colorado, Boulder, CO, USA

³Earth Research Institute, University of California, Santa Barbara, CA, USA

10 ⁴Research Applications Laboratory, National Center for Atmospheric Research, Boulder, CO, USA

⁵Joint Center for Satellite Data Assimilation, University Corporation for Atmospheric Research, Boulder, CO, USA

⁶Joint Institute for Regional Earth System Science and Engineering and Department of Atmospheric and Oceanic Sciences, University of California, Los Angeles, CA, USA

15 *Correspondence to:* Dalei Hao (dalei.hao@pnnl.gov)

Abstract. With the highest albedo of the land surface, snow plays a vital role in Earth's surface energy budget and water cycle. Snow albedo is primarily controlled by snow grain properties (e.g., size and shape) and light absorbing particles (LAPs) such as black carbon (BC) and dust. The mixing state of LAPs in snow also has impacts on LAP-induced snow albedo reduction and surface radiative forcing (RF). However, most land surface models assume that snow grain shape is spherical and LAPs are externally mixed with the snow grains. This study improves the snow radiative transfer model in the Energy Exascale Earth System Model version 2.0 (E3SM v2.0) Land Model (ELM v2.0) by considering non-spherical snow grain shapes (i.e., spheroid, hexagonal plate and Koch snowflake) and internal mixing of dust-snow and systematically evaluates the impacts on surface energy budget and water cycle over the Tibetan Plateau (TP). A series of ELM simulations with different treatments of snow grain shape, mixing state of BC-snow and dust-snow, and sub-grid topographic effects (TOP) on solar radiation are performed. Compared with two remote sensing snow products derived from the Moderate Resolution Imaging Spectroradiometer, both the control ELM simulation (ELM_Control) with the default configurations of spherical snow grain shape, internal mixing of BC-snow, external mixing of dust-snow and without TOP and the ELM simulation with new model features (ELM_New) can capture the overall snow distribution reasonably. Additionally, ELM_New overall shows smaller biases in snow cover fraction than ELM_Control in spring when snow melt is important for water managers. The estimated LAP-induced RF in ELM_New ranges from 0 to 19.3 W m⁻² with the area-weighted average value of 1.5 W m⁻² that is comparable to the reported values in the existing studies. The Koch snowflake shape, among other non-spherical shapes, shows the largest difference from spherical shape in spring when snow processes related to surface energy budget and water cycle have high importance. The impacts of the mixing state of LAP in snow are smaller than the shape effects and depend on snow grain shape. Compared to external mixing, internal mixing of LAP-snow can lead to larger snow albedo reduction and

35 snowmelt, which further affect surface energy budget and water cycle. The individual contributions of non-spherical snow
shape, mixing state of LAP-snow, and local topography impacts on the snow and surface fluxes have different signs and
magnitudes, and their combined effects may be negative or positive due to complex and non-linear interactions among the
factors. Overall, the changes of net solar radiation in spring due to individual and combined effects range from -28.6 to 16.9
40 W m⁻² and -29.7 to 12.2 W m⁻², respectively. This study advances understanding of the role of snow grain shape and mixing
state of LAP-snow in land surface processes and offers guidance for improving snow simulations and RF estimates in Earth
system models under climate change.

1 Introduction

With the highest broadband albedo (defined as the ratio of reflected to incident radiative flux at the surface from $0.2\mu\text{m}$ to $4.0\mu\text{m}$) of all land surfaces, snow plays an important role in Earth's radiation budget (Flanner et al., 2011), water cycle (Barnett et al., 2005), and regional/global climate change via positive snow-albedo feedback (Hall, 2004; Riihelä et al., 2021). Snow albedo (α_{sno}) is regulated by solar zenith angle (SZA), atmospheric conditions, snow properties, and snow impurities such as light absorbing particles (LAPs) (He and Flanner, 2020). Snow properties such as snow grain size and shape have large influences on single scattering optical properties of snow (Dang et al., 2016; Tanikawa et al., 2020; Warren, 1982). LAPs include black carbon (BC), light-absorbing organic carbon (also known as brown carbon, BrC), and mineral dust can darken the snow-covered surface (Wu et al., 2018; Zhao et al., 2014), increasing the amount of energy absorbed by snow and accelerating snowmelt (Kang et al., 2020; Sarangi et al., 2019). The surface radiative forcing (RF, a measure of the change in surface radiative flux) from LAPs has contributed to rapid glacier retreat (Xu et al., 2009). BC is the optically absorbing component of soot and has been identified as one of the most important anthropogenic emissions affecting global climate due to its large RF (Bond et al., 2013; Flanner et al., 2007; Hadley and Kirchstetter, 2012). BrC absorbs more light at shorter wavelengths compared to BC (Andreae and Gelencsér, 2006). The impurity effects of BrC in snow on modelling α_{sno} and its RF have not been widely investigated due to its large variabilities and uncertainties in the optical properties (Brown et al., 2021; Skiles et al., 2018). Dust has smaller mass absorption efficiency compared to BC, but the impurity effects of dust on snow albedo reduction (SAR) and its RF are found to be comparable to or even greater than BC in the high-mountain Asia (HMA) and the Sierra Nevada due to the larger dust deposition (Kaspari et al., 2014; Sarangi et al., 2020; Sterle et al., 2013; Usha et al., 2020).

Considering the high sensitivity of α_{sno} to snow grain shape, the assumptions of spherical snow grain shape may be inappropriate to realistically model α_{sno} . Radiative transfer models or snow albedo parameterizations in the Earth System Models (ESMs), e.g., the Snow, Ice, and Aerosol Radiative (SNICAR) model (Flanner et al., 2007), explain the dependence of α_{sno} on snow grain size, snow depth, SZA, underlying surface, atmospheric conditions, and spectral wavelength. However, the snow grain shape in these models/parameterizations is generally assumed to be spherical. In reality, snow grains are usually irregular and non-spherical (Kokhanovsky et al., 2005), and depend on snow age and meteorological conditions (Dominé et al., 2003; LaChapelle, 1969). Compared to spherical grains, non-spherical snow grains show weaker forward scattering and thus a smaller asymmetry factor, resulting in higher albedo (Dang et al., 2016; Libois et al., 2013; Räisänen et al., 2015; Räisänen et al., 2017). He et al. (2017, 2018b) developed parameterizations to explicitly represent the impacts of three typical non-spherical grain shapes (i.e., spheroid, hexagonal plate and Koch snowflake) on single-scattering properties of snow based on the geometric-optics surface wave (GOS) approach. These parameterizations have been implemented in the standalone version of the SNICAR model (He et al., 2018a) and have further been incorporated into the latest SNICAR with Adding-

Doubling solver, version 3.0 (SNICAR-AD v3) (Flanner et al., 2021). However, these new parameterizations have not been incorporated into ESMs and applied at regional or global scales yet.

The mixing state of LAPs in snow (i.e., external mixing and internal mixing) has large impacts on α_{sno} , and only accounting for external mixing of LAP-snow could lead to underestimations of modeled SAR and RFs (Flanner et al., 2012; Liou et al., 2014). LAP-snow internal mixing can occur through wet deposition or dry deposition followed by successive snow events over high elevation regions (Liou et al., 2014). Compared to external mixing, internal mixing of LAP-snow can enhance SAR and thus lead to larger RF (Flanner et al., 2012; He et al., 2018b; Liou et al., 2014; Tanikawa et al., 2020). However, snow albedo models/parameterizations in ESMs often assume that LAPs are only externally mixed with snow (Flanner et al., 2007). Flanner et al. (2012) showed that the internal mixing of BC-snow for a spherical snow grain shape can be accurately described by the dynamic effective medium approximation (DEMA) (Chýlek and Srivastava, 1983). He et al. (2017, 2018a, 2018b) implemented a series of computationally efficient polynomial-based parameterizations to represent the effects of BC-snow internal mixing on α_{sno} based on GOS simulations for both spherical and non-spherical snow grain shapes. He et al. (2019) further developed parameterizations to quantify the effects of dust-snow internal mixing for different snow grain shapes. This recent progress provides an opportunity to account for the LAP-snow internal mixing in ESMs and quantify their effects on land surface processes.

Realistically representing the impacts of snow grain properties and mixing state of LAPs on α_{sno} in ESMs is required for improving the simulations of snow processes and surface energy budget and water cycle. However, the non-spherical grain shape and internal mixing of dust-snow are still not represented in the snow radiative transfer model of the land model (ELM v2.0) of the Energy Exascale Earth System Model version 2.0 (E3SM v2.0). E3SM, supported by the U.S. Department of Energy (DOE), is a state-of-the-art fully coupled ESM aimed at improving projections of future changes of the water cycle, biogeochemistry, and cryosphere systems (Golaz et al., 2019; Leung et al., 2020). The land component of E3SM v2.0, ELM v2.0 was originally developed from the Community land model version 4.5 (CLM4.5), which uses SNICAR to model the snow radiative transfer processes. Dang et al. (2019) incorporated a two-stream radiative transfer scheme with delta-Eddington approximation and adding-doubling technique into SNICAR of ELM v2.0 (named as SNICAR-AD). This new SNICAR-AD model improves and unifies the treatment of snow shortwave radiative transfer in the land and sea ice components of E3SM (Dang et al., 2019). The internal mixing of BC-snow has been incorporated in SNICAR-AD of ELM v2.0 using a new look-up table derived from the DEMA approach (Wang et al., 2020a). However, the non-spherical snow grain shape and internal mixing of dust-snow are not included in the SNICAR-AD of ELM v2.0. The impact of non-spherical grain shape and internal mixing of LAP-snow on α_{sno} has been well studied using snow albedo models uncoupled with land surface models (Dang et al., 2016; He et al., 2018a; Räisänen et al., 2015). Inclusion of parameterizations for non-spherical grain shape and LAP-snow internal mixing into the snow radiative transfer model of ELM v2.0 offers a feasible way to analyze the sensitivity of surface energy budget and water cycle to such changes and quantify the corresponding uncertainties. Moreover, an improved

110 parameterization of sub-grid topographic effects on solar radiation has been recently implemented in ELM v1.0 (Hao et al.,
2021a), which makes it possible to study the interactions between improved snow radiative transfer model and sub-grid
topographic effects on solar radiation.

115 This study aims to improve the SNICAR-AD in ELM v2.0 by accounting for non-spherical snow grain shape and internal
mixing of LAP-snow and systematically evaluate their impacts on surface energy budget and water cycle over the Tibetan
Plateau. The analytical parameterizations of He et al. (2017, 2018a, 2018b) for three non-spherical grain shape (i.e., spheroid,
hexagonal plate and Koch snowflake) and He et al. (2019) for internal mixing of dust-snow are implemented in SNICAR-AD
of ELM v2.0. A series of ELM simulations with different assumptions of snow grain shape and mixing state of LAP-snow are
carried out. Then the control ELM simulation (ELM_Control) with the default configurations in the original ELM v2.0 and
120 the ELM simulation with new model features (ELM_New) implemented in this study are compared with two snow surface
property remote sensing products derived from the Moderate Resolution Imaging Spectroradiometer (MODIS). The RFs from
LAPs' impurity effects in the ELM_Control and ELM_New are also compared to the values reported in existing studies. The
impacts of snow grain shape and mixing state of LAP-snow on the snow-related processes, energy budget, and water cycle are
analyzed. The interactions between the improved snow radiative transfer model and sub-grid topographic effects on solar
125 radiation are also investigated.

2 Model Improvements in ELM v2.0

2.1 Snow albedo modeling in ELM and SNICAR-AD v3

130 E3SM v2.0 (including ELM v2.0) was released on September 29, 2021 (<https://github.com/E3SM-Project/E3SM/releases/tag/v2.0.0>). The default settings for hydrologic and biogeochemistry models in ELM v2.0 are the same
as ELM v1.0 and ELM v1.1. ELM v1.0 incorporated new features for better representing soil hydrology and carbon cycle
dynamics (Golaz et al., 2019). New options for modeling α_{sno} are implemented in ELM v2.0. The SNICAR-AD model is also
adopted in ELM v2.0 to more accurately describe the multi-layer snow radiative transfer processes (Dang et al., 2019).
SNICAR-AD can prognostically simulate the evolution of the snow grain size and LAP concentration via deposition and
redistribution. SNICAR-AD of ELM showed better performance in simulating α_{sno} both visible (VIS) and near-infrared (NIR)
135 wavelengths at various SZAs under all sky conditions than the original SNICAR (Dang et al., 2019). The internal mixing of
hydrophilic BC-snow is represented in SNICAR-AD of ELM v2.0 using a new look-up table derived using the DEMA
approach (Wang et al., 2020a). In parallel, Hao et al. (2021a) updated the two-stream radiative transfer scheme of ELM v1.0
to account for the sub-grid topographic effects on solar radiation (TOP), which showed improved performance on simulating
surface energy balance and snow processes over the Tibetan Plateau (TP) compared to the original plane-parallel (PP) scheme.

140

SNICAR-AD v3 includes numerous recent improvements and extensions in SNICAR (Flanner et al., 2021). Specifically, it adopts the Adding-Doubling solver for the radiative transfer equations; adds new representations of carbon dioxide snow, snow algae, and new types of dust, BrC, and volcanic ash; contains options for ice refractive indices; parameterizes the impacts of non-spherical snow grain shape; and considers the SZA dependence of surface spectral irradiances under different atmospheric profiles. However, internal mixing of LAP-snow is not accounted for in SNICAR-AD v3.

As part of this study, we implement the following new features into the snow albedo model in ELM v2.0: 1) SZA dependence of clear-sky surface irradiance (the original model assumes that clear-sky surface irradiance doesn't change with SZA); 2) the parameterization of He et al. (2017, 2018a, 2018b) for non-spherical snow grain shape (i.e., spheroid, hexagonal plate, and Koch snowflake); and 3) the parametrization of He et al. (2019) for internal mixing of dust-snow. For the first enhancement, new look-up tables of SNICAR-AD v3 are incorporated in ELM v2.0, which considers the surface irradiance dependence on SZA for six types of atmospheric profiles under both clear-sky and cloudy conditions: mid-latitude winter, mid-latitude summer, sub-Arctic winter, sub-Arctic summer, summit Greenland, and high mountain. The remaining two enhancements are described in Sections 2.2 and 2.3.

155 2.2 Parameterizing the impacts of snow grain shape

Snow grain shape affects the single scattering optical properties, such as single-scattering albedo (ω , which describes the probability that a photon experiencing an extinction event is scattered as opposed to absorbed), extinction efficiency (Q_{ext} , which is the sum of scattering efficiency and absorption efficiency and represents the light attenuation ability of the particle), and asymmetry factor (g , which represents the average cosine of the scattering phase angle and determines the fractions of backscattered and forward-scattered light). Previous studies represented non-spherical grains as spheres of an equivalent radius, R_e , such that the specific surface area (i.e., surface area to mass ratio) is equal to that of the non-spherical grain. Although the equivalent sphere performs well in computing ω and Q_{ext} (Grenfell and Warren, 1999), it is not sufficiently accurate in estimating g (Dang et al., 2016). He et al. (2017) defined a specific-projected-area-equivalent radius, R_s , of an equivalent sphere with the same orientation-averaged projected area to volume ratio as that of the non-spherical grain:

$$165 \quad R_s = \frac{3V_{snow}}{4A_{snow}} \quad (1)$$

where V_{snow} is the volume of snow grain and A_{snow} is the projected area of a snow grain averaged over all directions. For convex shapes (e.g., spheroid and hexagonal plate), $R_s=R_e$, whereas for concave shape (e.g. Koch snowflake), $R_s>R_e$ generally (He et al., 2017). He et al. (2017) developed new parameterizations to further consider the effects of non-spherical grain shape on g , by introducing two additional factors for describing the degree of non-sphericity: (1) aspect ratio (AR) which is the ratio of grain width to length, and (2) shape factor (SF) which is the ratio of R_s of a non-spherical grain to that of an equal-volume sphere. Specifically, three typical non-spherical grain shapes (Figure S1) are considered: spheroid, hexagonal plate and Koch snowflake, which capture the major morphological characteristics of snow grain structures observed by scanning electron

microscopy (Dominé et al., 2003; Erbe et al., 2003; Liou et al., 2014) Simulated snow albedo under the assumptions of the three non-spherical snow shapes show better agreement with field measurements (He et al., 2018a). The recommended AR

175 and SF values for the three non-spherical grain shapes are listed in Table 1 of He et al. (2017).

According to (Fu, 2007), the asymmetry factor for hexagonal plate/column, g_{hex} , can be derived by

$$g_{hex} = \frac{1-g'}{2\omega} + g' \quad (2)$$

where g' is as

$$g' = \begin{cases} a_0 + a_1 \cdot AR + a_2 \cdot AR^2, & 0.1 \leq AR \leq 1.0 \\ b_0 + b_1 \cdot \ln(AR) + b_2 \cdot \ln^2(AR), & 1 < AR \leq 20.0 \end{cases} \quad (3)$$

180 where both a_i and b_i are wavelength-dependent fitted coefficients provided in Tables 1 and 2 of Fu (2007). ω is the snow single-scattering albedo, and AR is the snow grain aspect ratio. The asymmetry factor for other non-spherical shapes (i.e., spheroid and Koch snowflake), g_{ns} , can be calculated by

$$g_{ns} = g_{hex} \cdot C_g \quad (4)$$

where C_g , an empirical correction factor, is parameterized as the function of shape factor (SF) and R_s :

$$C_g = c_0 \cdot \left(\frac{SF_{ns}}{SF_{hex}}\right)^{c_1} \cdot (2R_s)^{c_2} \quad (5)$$

where c_i are wavelength-dependent fitted coefficients provided in Table 3 of He et al. (2017). These parameterizations have shown good performance when compared to the sophisticated geometric optics ray-tracing simulations (Flanner et al., 2021; He et al., 2017).

2.3 Parameterizing the impacts of internal mixing of dust-snow

190 The mixing state of dust-snow (Figure S1) mainly contributes to the change of ω (He et al., 2019; Liou et al., 2014). He et al. (2019) proposed a parameterization to account for the internal mixing of dust-snow by modifying ω . Specifically, compared to pure snow, the snow single-scatter co-albedo ($1-\omega$) enhancement ($E_{1-\omega}$) caused by dust-snow internal mixing was defined as

$$E_{1-\omega} = \frac{1-\omega_{dust}}{1-\omega_p} \quad (6)$$

195 Where ω_p and ω_{dust} are single-scattering albedo of pure snow and dirty snow internally mixed by snow, respectively. $E_{1-\omega}$ can be empirically calculated by

$$E_{1-\omega} = d_0 + d_1 \cdot (C_{dust})^{d_2} \quad (7)$$

where C_{dust} is the dust mass concentration and d_i with $i = 1,2$ are wavelength-dependent fitted coefficients listed in Table 1 of He et al. (2019). This parameterization has shown good consistencies with the Monte Carlo ray-tracing simulations (He et

200 al., 2019).

3 Model Simulations and Remote Sensing Data

3.1 Experimental Design

Tibetan Plateau and its surrounding regions (named TP in the study), characterized by complex topography and frequent snow cover, are selected as a testbed in the study. Some regions of TP include areas where LAPs have large impacts on snowmelt 205 (Sarangi et al., 2020). This study area has a large elevation gradient ranging from below 1500 m to above 6000 m (Figure 1). Only model results for grid cells with elevation above 1500 m are considered in the study.




Figure 1: Spatial distribution of elevation and the contour lines of 1.5 and 4.5 km over the Tibetan Plateau.

A series of ten ELM simulations with different configurations at 0.125° spatial resolution are conducted to investigate the 210 sensitivity to the new improvements in the snow albedo model (Table 1). For all cases, the prescribed satellite phenology (SP) mode is used, where the 8-day 500 m MODIS leaf area index (LAI) data is upscaled to derive the climatological monthly data with 0.125° spatial resolution (Myneni et al., 2002). The Global Soil Wetness Project Phase 3 (GSWP3) v1 forcing data (Dirmeyer et al., 2006) with 0.5° spatial resolution and 3-hourly temporal resolution is used to drive ELM. The prescribed 215 climatological monthly aerosol deposition data at $1.9^\circ \times 2.5^\circ$ is used from the Community Atmosphere Model version 5 coupled with chemistry (Lamarque et al., 2010; Liu et al., 2012), which provides the dry and wet deposition rate of LAPs. The bilinear interpolation technique is used to spatially downscale the GSWP3v1 and aerosol deposition data to 0.125° spatial resolution. The temporal downscaling of the solar data, precipitation data, and all the other atmospheric data are performed using the 220 cosine of the SZA-based, nearest-neighbour, and linear interpolation methods, respectively. The configurations of the snow albedo model include (1) mid-latitude winter atmosphere profile, (2) SZA-dependence of solar irradiance, (3) external mixing of hydrophobic BC with snow, (4) four different snow grain shapes (i.e. sphere, spheroid, hexagonal plate, Koch snowflake), (5) internal and external mixing of hydrophilic BC and dust with snow, and (6) two different solar radiation parameterizations (i.e., TOP and PP). The parameterization of He et al. (2019) for the effects of internally-mixed dust in snow assumes no BC present in the snow. Thus, we do not include a model configuration that simultaneously considered the internal mixing of BC and dust in snow. The control simulation with the default settings in the original ELM is named ELM_Control, while the case

225 with all the model features is named ELM_New (Table 1). For each case, ELM is first run from 1950 to 2000 for model spin-up, followed by another 10 years from 2001-2010 for model analysis. The output variables are stored at a half-hourly interval to match the satellite observations and then aggregated to multi-year averaged seasonal scales.

230 **Table 1. Model configurations with different snow grain shapes, mixing states of LAP-snow and solar radiation parameterizations.**

Case ID	Snow Grain Shape	Mixing of hydrophilic	Mixing of	dust-	Solar	radiation
		BC-snow	snow		parameterization	
Sph_BCI nt_DExt_PP (ELM_Control)	Sphere	Internal	External		PP	
Sphd_BCI nt_DExt_PP	Spheroid	Internal	External		PP	
Hex_BCI nt_DExt_PP	Hexagonal Plate	Internal	External		PP	
Koc_BCI nt_DExt_PP	Koch Snowflake	Internal	External		PP	
Sph_BCE nt_DExt_PP	Sphere	External	External		PP	
Sph_BCE nt_DInt_PP	Sphere	External	Internal		PP	
Koc_BCE nt_DExt_PP	Koch Snowflake	External	External		PP	
Koc_BCE nt_DInt_PP	Koch Snowflake	External	Internal		PP	
Sph_BCE nt_DInt_TOP	Sphere	External	Internal		TOP	
Koc_BCE nt_DInt_TOP (ELM_New)	Koch Snowflake	External	Internal		TOP	

3.2 MODIS data

Two snow surface property products derived from MODIS onboard Terra are used to evaluate both the ELM_Control and ELM_New. The first set are the spatially and temporally complete (STC) Snow-Covered Area and Grain Size (MODSCAG) 235 and MODIS Dust and Radiative Forcing in Snow (MODDRFS) products (Painter et al., 2012; Painter et al., 2009; Rittger et al., 2020) that we hereafter refer to as “STC-MODSCAG” and “STC-MODDRFS”. STC-MODSCAG uses a physically-based spectral mixture algorithm to estimate the snow cover fraction (f_{sno}) and grain size and has been shown to have a better performance than MODIS MOD10A1 snow cover products (Rittger et al., 2013) especially in the melt season. STC-MODDRFS is used to calculate LAP-induced SAR based on the relative difference between satellite measured dirty α_{sno} and 240 modeled pure α_{sno} , which is derived from the grain size. The snow surface properties are interpolated and smoothed to reduce the errors and uncertainties caused by off-nadir views, cloud contamination and data noise, resulting in improved quality and spatio-temporal consistency (Rittger et al., 2020). Snow albedo estimates combining STC-MODSCAG and STC-MODDRFS

have a 4 to 6 percent RMSE and negligible bias (Bair et al., 2019). The Snow Property Inversion from Remote Sensing (SPIReS) product is the second MODIS dataset used for evaluating ELM simulations. The SPIReS product estimates LAP 245 concentrations and snow grain size simultaneously with optimized data processing and integrated spatio-temporal interpolation and smoothing procedures (Bair et al., 2021b). Then LAP-induced SAR can be calculated by the albedo look-up table derived from a radiative transfer model (Bair et al., 2019). Spherical snow grain shape is assumed in the STC-MODSCAG/STC-MODDRFS and SPIReS.

250 Both the daily 500 m f_{sno} and SAR in STC-MODSCAG/STC-MODDRFS and SPIReS products from 2001 to 2010 are used to compare with the ELM control simulation. SAR estimated by STC-MODDRFS covers the spectral range from 0.350 to 0876 μm and is converted to broadband via a scaling factor of 0.63 (Bair et al., 2019). Both products are aggregated to 0.125° spatial resolution and multi-year averaged seasonal scales.

255 **3.3 Evaluation and analysis**

Both ELM_Control and ELM_New in Table 1 are compared to the MODIS data. The MODIS data represents the snow states at the overpass time of about 10:30 AM (local solar time). Thus, ELM simulated snow-related outputs are extracted from 10:00 am to 11:00 AM (local solar time) to match the satellite observations and then aggregated to multi-year averaged seasonal scales: winter (DJF), spring (MAM), summer (JJA) and autumn (SON). Considering that the snow cover in summer and 260 autumn is relatively low, this study focuses on winter and spring.

The impacts of snow grain shape and mixing state of LAP in snow on energy budget and water cycles in winter and spring are investigated from the various model configurations as listed in Table 1. Specifically, the impacts on the snow-related processes, energy budget, and water cycle are analyzed. The contributions of different influencing factors including snow grain shape, 265 mixing state of LAP-snow, and sub-grid topographic effects are separated and compared based on the differences between various model configurations (see Table 2). Only the model grids with f_{sno} larger than 0.01 in ELM_Control are used in the analysis. Note that RFs over all model grids are averaged (weighted by areas) to calculate the area-weighted average RFs over the TP with elevation above 1500 m (i.e., zero is involved in the calculation for any grids when snow is not present). The maximum absolute difference (AD_{max} , calculated as the maximum of the absolute value of the difference between two cases), 270 maximum relative difference (RD_{max} , calculated as the maximum of the absolute value of the relative difference between two cases), mean absolute difference (AD_{mean} , calculated as the mean of the absolute value of the difference between two cases), and mean relative difference (RD_{mean} , calculated as the mean of the absolute value of the relative difference between two cases) are used to evaluate the effects of snow grain shape and mixing state. One-way analysis of variance (ANOVA), which compares the means between different groups and determines whether they are statistically significant different from each other is used

275 to determine whether the effects of snow grain shape and mixing state are statistically significant (if the levels of statistical significance (p values) are less than 0.05) or not.

Table 2. Separating the contributions of different influencing factors based on the model experiments in Table 1.

Influencing Factor	Abbreviation	Snow grain shape	Difference between cases
Non-spherical shape (compared to spherical shape)	Non-spherical effect	Spheroid	Sphd_BCInt_DExt_PP – Sph_BCInt_DExt_PP
		Hexagonal Plate	Hex_BCInt_DExt_PP – Sph_BCInt_DExt_PP
		Koch Snowflake	Koc_BCInt_DExt_PP – Sph_BCInt_DExt_PP
Mixing state of BC-snow (Internal-External)	BC mixing state effect	Sphere	Sph_BCInt_DExt_PP – Sph_BCExt_DExt_PP
		Koch Snowflake	Koc_BCInt_DExt_PP – Koc_BCExt_DExt_PP
Mixing state of dust-snow (Internal-External)	Dust mixing state effect	Sphere	Sph_BCExt_DInt_PP – Sph_BCExt_DExt_PP
		Koch Snowflake	Koc_BCExt_DInt_PP – Koc_BCExt_DExt_PP
Sub-grid topographic effects	TOP effect	Sphere	Sph_BCExt_DInt_TOP – Sph_BCExt_DInt_PP
		Koch Snowflake	Koc_BCExt_DInt_TOP – Koc_BCExt_DInt_PP
Combined effect	Combined effect	Koch Snowflake	Koc_BCExt_DInt_TOP – Sph_BCInt_DExt_PP

280 4. Results

4.1 Comparison with remote sensing snow data


Compared to the MODIS data, ELM simulations (i.e., ELM_Control and ELM_New) show less spatial variability (Figure S2), which may be caused by the relatively coarse spatial resolution of the GSWP3 atmospheric forcing data and aerosol deposition data. The simulated f_{sno} in both ELM_Control and ELM_New shows similar magnitudes to the two MODIS products in winter

285 (Figure S2). ELM and MODIS data show higher f_{sno} in the western regions of the TP in winter (Figure S2). ELM_Control overestimates f_{sno} in the western regions of the TP for winter, and ELM_New shows similar bias to ELM_Control (Figure 2a-b) for f_{sno} . The area-weighted average f_{sno} for winter in ELM_New is within the range of the uncertainty bounds in STC-MODSCAG and SPIReS, while ELM_Control slightly underestimates the area-weighted average of f_{sno} for winter (Figure 3a). In spring, both ELM_Control and ELM_New underestimate f_{sno} compared to STC-MODSCAG and SPIReS, however,

290 ELM_New has a smaller bias than ELM_Control in the western regions of the TP (Figure 2c-d). Compared to the mean value of the two MODIS datasets, ELM_New reduces 0.014 (13.6%) of the bias of ELM_Control in the area-weighted average of f_{sno} for spring (Figure 3a). Overall, both ELM_Control and ELM_New capture the overall elevation gradients of f_{sno} for winter

(Figure 4a). At different elevation intervals, ELM shows similar mean values and ranges of f_{sno} with STC-MODSCAG and SPIReS for winter. In spring, ELM_New reduces the underestimation of f_{sno} in ELM_Control for each elevation interval 295 (Figure 4b).

Both ELM_Control and ELM_New show large differences in the spatial distribution of LAP-induced SAR, compared to two remote sensing datasets (Figure 5). In winter, both ELM_Control and ELM_New have a similar magnitude of SAR to STC-MODDRFS, while SPIReS has lower values. The area-weighted average SAR for winter in both ELM_Control and ELM_New 300 is within the range of the uncertainty bounds in SPIReS (Figure S3b). In spring, ELM_Control and ELM_New show a similar change along the elevation gradient as compared to the STC-MODDRFS (Figure 4d), but the estimated SAR in ELM_Control, ELM_New and STC-MODDRFS is greater than SPIReS. These differences may be due to the overestimation of snow grain 305 size (Figure S3) and associated underestimation of α_{sno} in ELM (Sarangi et al., 2020). The two MODIS products show large differences, probably due to different assumptions used in their retrieval algorithm, limitations of multi-spectral sensor, and/or persistent cloud cover over the TP necessitating interpolation. The coarse resolution and different time periods of the aerosol deposition data used in the ELM simulations can contribute to the inconsistencies among the three datasets. Overall, ELM_New 310 better represents the snow cover distribution than ELM_Control, while LAP-induced SAR in both ELM_Control and ELM_New has poor consistency with the MODIS estimates.



310 **Figure 2:** (a,c) The f_{sno} bias ($\delta_{\text{ELM_Control}}$) of ELM_Control compared to the mean value of STC-MODSCAG and SPIReS, and (b,d) the difference ($|\delta_{\text{ELM_New}}| - |\delta_{\text{ELM_Control}}|$) between the absolute values of the biases of ELM_New ($\delta_{\text{ELM_New}}$) and ELM_Control ($\delta_{\text{ELM_Control}}$) for winter (a-b) and spring (c-d). The negative values (blue color) in (b,d) show that ELM_New has the smaller bias than ELM_Control. The areas with f_{sno} smaller than 0.01 are masked.




Figure 3: The area-weighted average of (a) f_{sno} and (b) SAR induced by LAPs for winter and spring from ELM, STC-MODSCAG/STC-MODDRFS, and SPIReS. The bar width represents the uncertainty bounds of STC-MODSCAG and SPIReS from Bair et al., (2021a).





Figure 4: Statistical distributions of (a,b) f_{sno} and (c,d) SAR induced by LAPs under different elevation ranges for winter and spring from ELM, STC-MODSCAG/STC-MODDRFS and SPIReS.




325

Figure 5: Spatial distributions of SAR estimated from: (a,e) ELM_Control, (b,f) ELM_New, (c,g) STC-MODSCAG/STC-MODDRFS and (d,h) SPIReS for winter (a-d) and spring (e-h). The areas with f_{sno} smaller than 0.01 are masked..

4.2 Radiative forcing induced by LAPs in snow

330 The RFs induced by different LAPs in both ELM_Control and ELM_New show divergent spatial distributions with seasonal variations (Figures 6 and S4). Western regions show larger RFs induced by all LAPs than the eastern regions (Figure 6a,c), while BC-induced RFs are larger in the southwestern regions (Figure 6b,e) and dust-induced RFs are larger in the northwestern regions (Figure 6c,f). Overall, spring has larger RFs than winter in ELM_New, because LAP-induced SAR is larger in spring (Figure 5a-b,c-f) caused by accumulated LAP concentration and resurfacing as snow melts and the larger incident solar
 335 radiation over TP in spring than in winter. Table S1 summarizes the statistical maximum and mean values of RFs induced by different LAPs.

The LAP-induced RF values estimated in both ELM_Control and ELM_New case are within the range of the RF values reported by other studies. Table 3 summarizes the LAP-induced RF values for spring or non-monsoon season over the TP
 340 reported by previous studies based on different climate models and different snow radiative transfer models. In spring, RFs by all LAPs in ELM_New are 0-19.3 W m⁻², which is close to the results in Qian et al. (2011) that showed RFs within 5-25 W m⁻² during spring. RFs by BC in the control simulation are about 0-5.4 W m⁻², which is similar to Ji (2011). Gertler et al. (2016) also reviewed different studies in the Himalaya and showed that BC-induced RFs range from 0 to 28.0 W m⁻². RFs by dust in ELM_New are about 0-11.9 W m⁻², which are similar to the results in Xie et al. (2018a). The area-weighted average RFs by
 345 all LAPs, BC and dust in ELM_New are 1.5, 0.3 and 0.9 W m⁻², respectively in spring. Our results are also consistent with (Zhang et al., 2015), who found that snow LAPs induced RF has a seasonal peak in spring and a spatial maximum over the northwest TP with an average RF of 5 and 3.5 W m⁻², respectively, for BC and dust.



350 **Figure 6. Spatial distributions of RF values from (a,d) all LAPs (BC+dust), (b,e) BC and (c,f) dust for (a-c) winter and (d-f) spring in ELM_New. The areas with f_{sno} smaller than 0.01 are masked.**

Table 3. A summary of RF values of LAPs in snow at seasonal scales over the TP reported by the existing studies.

Model Name	Snow radiative transfer model	Aerosol type	RF value (W m ⁻²)	Period	Reference
Community Atmosphere Model, version 3 (CAM 3) CAM 3.1	SNICAR	BC	>20 in some parts of TP	Spring	(Flanner et al., 2007)
CAM 3.1	SNICAR	BC+Dust	0-9.5 or even >9.5	Spring	(Flanner et al., 2009)
CAM 3.1	SNICAR	BC+Dust	5-25	Spring	(Qian et al., 2011)
GEOS-Chem	Four-stream radiative transfer model	BC	Approximately 8-16	Spring	(Kopacz et al., 2011)
GEOS-Chem	Geometric-optics surface-wave (GOS) approach	BC	5-10	Spring	(He et al., 2014)
CAM 5	SNICAR	BC	5 over the northwest TP	Spring	(Zhang et al., 2015)
		Dust	3 over the northwest TP	Spring	
Regional Climate Model version 4.3.4	SNICAR	BC	0-6	November-April	(Ji, 2016)
CAM 4	Bulk Aerosol Model parameterizations of the dust size distribution (BAM)	Dust	0-20 or even >20	Spring	(Xie et al., 2018a)
ELM_Control	SNICAR-AD	BC+Dust	0-21.9	Spring	This study
ELM_New	Improved SNICAR-AD	BC+Dust	0-19.3	Spring	This study

355 4.3 Impacts of snow grain shape and mixing state of LAP-snow

4.3.1 Impacts on snow related processes

Compared to the spherical shape, all three non-spherical grain shapes show larger α_{sno} and higher f_{sno} in spring (Figure 7) and

all the differences in α_{sno} and f_{sno} between non-spherical and spherical shapes are significant (ANOVA: $p < 0.05$), because non-spherical grain shapes have a smaller asymmetry factor and their forward-scattering is weaker than the spherical shape. Among

360 them, spheroid shape has the smallest differences from spherical shape and Koch snowflake has the largest differences from spherical shape (Figure 7). For instance, when $f_{sno} \geq 0.5$, AD_{mean} in α_{sno} and f_{sno} between Koch snowflake and spherical shape are 0.08 and 0.09, respectively (Table 4). The simulated snow water equivalent (SWE) also changes with snow grain shape and AD_{mean} in SWE between Koch snowflake and spherical shape is 56.65 mm (Table 4). Similar results are obtained in winter (Figure S5), but they generally show smaller differences than those in spring. Similar results are obtained for spring and winter, 365 thus only the results in spring are reported in the subsequent sections and the corresponding results for winter are reported in the supplementary material.

Compared to the external mixing of LAP-snow, internal mixing leads to smaller α_{sno} and lower f_{sno} (Figure 8), although all the differences in α_{sno} and f_{sno} between internal and external mixing are not significant (ANOVA: $p > 0.1$). For instance, when

370 $f_{sno} \geq 0.5$, under spherical shape, AD_{mean} in α_{sno} between internal and external mixing of BC-snow and dust-snow is

approximately 0.009, while under Koch snowflake shape, AD_{mean} caused by mixing state of BC-snow and dust-snow are 0.007 and 0.005, respectively. Similar results can be obtained for f_{sno} from Figure 8e-h.

Both SAR and RFs induced by LAPs are sensitive to snow grain shape and mixing state of LAP-snow (Figure 9). The effects of snow grain shape and mixing state are more significant for regions with higher f_{sno} . Non-spherical shapes have smaller SAR and RFs induced by different LAPs, compared to the spherical shape. For instance, when $f_{sno} \geq 0.5$, all LAP-induced mean SAR and RF under spherical shape (Sph_BCI nt_DExt_PP case) are 0.065 and 14.6 W m^{-2} , while those under Koch snowflake (Koc_BCI nt_DExt_PP case) are 0.051 and 12.1 W m^{-2} . Compared to external mixing, internal mixing of LAP-snow leads to larger SAR and higher RF values especially under spherical shape. For instance, when $f_{sno} \geq 0.5$, the dust-induced SAR and RF in the Sph_BCE nt_DExt_PP case have mean values of 0.034 and 7.5 W m^{-2} , while the corresponding SAR and RF are larger in the Sph_BCE nt_DInt_PP case with the mean values of 0.041 and 9.1 W m^{-2} , respectively. The effects of internal mixing become smaller for non-spherical grain shapes. For instance, the Koc_BCE nt_DInt_TOP case has smaller SAR and RFs compared to the Sph_BCE nt_DInt_PP case.

385




Figure 7: The differences in α_{sno} (a-c) and f_{sno} (d-f) between different snow grain shapes: (a,d) spheroid - sphere, (b,e) hexagonal plate - sphere and (c,f) Koch snowflake - sphere, for spring. Here the differences between non-spherical and spherical grain shapes are positive. The specific calculation methods are listed in Table 2. The areas with f_{sno} smaller than 0.01 are masked.

390




Figure 8: The differences in α_{sno} (a-d) and f_{sno} (e-h) between different mixing states (internal - external) of snow-LAP: (a,e) BC (sphere), (b,f) dust (sphere), (c,g) BC (Koch snowflake) and (d,h) dust (Koch snowflake) for spring. Here the differences between internal and external mixing are negative. The specific calculation methods are listed in Table 2. The corresponding results for winter are shown in Figure S6. The areas with f_{sno} smaller than 0.01 are masked.





Figure 9: Boxplots of SAR (a-b) and RF (c-d) from all LAPs, BC, and dust for different f_{sno} values: (a,c) < 0.5 and (b,d) ≥ 0.5 in spring under different cases listed in Table 1. For the case ID labelled in x-axis, the '_PP' suffix is omitted to keep them simplified. Red circles represent the mean values. The corresponding results for winter are shown in Figure S7.

4.3.2 Impacts on energy budget and water cycle


The impacts of snow grain shape and mixing state of LAP-snow on surface energy balance are large especially when f_{sno} is high (Figures 10 and S8-S10). Generally as f_{sno} increases, non-spherical grain shape has larger effects on surface energy balance terms due to a larger change in α_{sno} and thus land surface albedo (α_{sur}) (Figures 10 and S8). Overall, Koch snowflake shape has the largest differences from spherical shape in all surface energy budget terms (Figure 10). Due to the largest change of α_{sur} , net solar radiation (R_{net}^s) of Koch snowflake has the largest difference from that of sphere (Table 4). For instance, AD_{mean} and RD_{mean} of R_{net}^s are 17.6 W m^{-2} and 0.17, respectively when $f_{\text{sno}} \geq 0.5$. The change of R_{net}^s further leads to the change of surface temperature (T_{sur}), latent heat (F_{lat}) and sensible heat (F_{sen}) fluxes. Similarly, the effect of the mixing state of LAP-snow on surface energy balance overall increases with f_{sno} (Figures S9-S10). The effects of mixing state on surface energy balance have an opposite sign with those of non-spherical shape, because internal mixing leads to smaller α_{sno} than external mixing. Overall, the magnitude of mixing state effects is smaller than the effects of non-spherical grain shape (Figures 10 and S8-S10). Snow grain shape also affects the differences in the energy balance terms between the internal and external mixing of LAP-snow, because non-spherical grains tend to have larger α_{sno} (Figure 7) and could affect SAR induced by LAPs. For instance, when $f_{\text{sno}} \geq 0.5$, under spherical shape, AD_{mean} between internal and external mixing of BC-snow in R_{net}^s is 2.3 W m^{-2} , while under Koch snowflake shape, the value is 2.1 W m^{-2} (Table 4).



420 **Figure 10:** The boxplots of the differences (Δ) in surface energy budget terms: (a) surface albedo (α_{sur}), (b) net solar
 425 radiation (R_{net}^s), (c) surface temperature (T_{sur}), (d) latent heat flux (F_{lat}) and (e) sensible heat flux (F_{sen}), between
 different snow grain shapes: spheroid – sphere, hexagonal plate – sphere and Koch snowflake – sphere under different
 snow cover fractions (f_{sno}) for spring. See Table 2 for the specific calculation methods.

The water cycle is also affected by snow grain shape and mixing state of LAP-snow especially over the regions with high snow
 425 cover, due to snowmelt and altered F_{lat} and F_{sen} (Table 4 and Figures S11-S14). Due to the increased α_{sur} and reduced R_{net}^s ,
 non-spherical grain shape shows reduced snowmelt, evapotranspiration (ET), and runoff, and Koch snowflake shape has the
 largest difference from spherical shape (Figures 11 and S11). As f_{sno} increases, the non-spherical effects on these water cycle
 related terms generally become larger. When $f_{\text{sno}} \geq 0.5$, Δ_{mean} in snowmelt, ET and runoff between Koch snowflake and
 430 spherical shape are 0.43, 0.25, and 0.47 mm/day, respectively. The mixing state (i.e., the difference between internal and
 external mixing) of LAP-snow has smaller but opposite effects on water cycle than the effects from non-spherical shape

(Figures S12-S13). The effects of mixing state also increase with f_{sno} . The snow grain shape also affects the magnitude of mixing state effects. For instance, when $f_{\text{sno}} \geq 0.5$, AD_{mean} between internal and external mixing of BC-snow in snowmelt is 0.04 mm/day, while it is 0.07 mm/day under Koch snowflake shape.



435 **Figure 11:** Same as Figure 10, except for hydrological terms: (a) snowmelt, (b) ET, and (c) runoff.

Table 4. Statistics of the impacts of snow grain shape and mixing state of LAP-snow on snow-related processes, surface energy and water cycles in spring for the grid cells of $f_{\text{smo}} \geq 0.5$ in the control simulation.

Category	Variable	Non-spherical effect (Koch)				BC mixing state effect for sphere				BC mixing state effect for Koch				Dust mixing state effect for Sphere				Dust mixing state effect for Koch			
		AD _{max}	RD _{max}	AD _{mean}	RD _{mean}	AD _{max}	RD _{max}	AD _{mean}	RD _{mean}	AD _{max}	RD _{max}	AD _{mean}	RD _{mean}	AD _{max}	RD _{max}	AD _{mean}	RD _{mean}	AD _{max}	RD _{max}	AD _{mean}	RD _{mean}
Snow-related processes	\mathcal{C}_{smo}	0.096	0.15	0.076	0.11	0.023	0.03	0.009	0.01	0.018	0.03	0.007	0.01	0.016	0.02	0.009	0.01	0.010	0.02	0.005	0.01
	f_{smo}	0.16	0.32	0.09	0.15	0.03	0.07	0.01	0.02	0.04	0.07	0.01	0.02	0.02	0.04	0.01	0.02	0.01	0.03	0.01	0.01
	SWE (mm)	101.65	0.54	56.65	0.26	18.33	0.11	7.04	0.03	17.48	0.13	6.45	0.03	10.20	0.05	4.83	0.02	8.66	0.05	3.81	0.02
	SAR	0.027	0.36	0.014	0.22	0.014	0.24	0.007	0.11	0.013	0.23	0.006	0.10	0.013	0.17	0.007	0.11	0.008	0.14	0.005	0.07
	RF (W m ⁻²)	5.82	0.36	2.82	0.19	3.23	0.21	1.51	0.10	3.02	0.23	1.43	0.10	2.59	0.17	1.51	0.10	1.85	0.13	1.07	0.07
	\mathcal{C}_{sur}	0.060	0.08	0.043	0.06	0.012	0.02	0.005	0.01	0.014	0.02	0.005	0.01	0.007	0.010	0.004	0.01	0.006	0.01	0.003	0.01
Energy cycle	R_{net}^s (W m ⁻²)	28.60	0.28	17.63	0.17	5.89	0.05	2.28	0.02	7.32	0.05	2.10	0.02	3.41	0.04	1.62	0.02	2.60	0.03	1.30	0.01
	T_{sur} (K)	1.92	-	1.01	-	0.42	-	0.13	-	0.43	-	0.12	-	0.17	-	0.09	-	0.16	-	0.07	-
	F_{lat} (W m ⁻²)	13.41	0.74	7.16	0.27	2.79	0.15	0.97	0.04	2.75	0.08	0.86	0.03	1.71	0.12	0.67	0.03	1.37	0.06	0.50	0.02
	F_{sen} (W m ⁻²)	12.50	0.98	4.93	0.56	2.82	0.23	0.71	0.08	2.23	0.16	0.54	0.06	1.11	0.17	0.49	0.06	0.86	0.09	0.32	0.04
	Snowmelt (mm/day)	2.07	0.42	0.43	0.08	0.21	0.04	0.04	0.01	0.41	0.05	0.07	0.01	0.27	0.06	0.04	0.01	0.39	0.05	0.06	0.01
Water cycle	ET (mm/day)	0.46	0.74	0.25	0.27	0.10	0.15	0.03	0.04	0.10	0.08	0.03	0.03	0.06	0.12	0.02	0.03	0.05	0.06	0.02	0.02
	Runoff (mm/day)	2.75	0.51	0.47	0.16	0.83	0.14	0.06	0.02	0.96	0.23	0.07	0.02	0.63	0.16	0.05	0.02	0.37	0.11	0.05	0.02

4.4 Combined effects of snow grain shape, mixing state of LAP-snow and sub-grid topography

The snow grain shape, mixing state of LAP-snow, and sub-grid topography can all affect α_{sur} and thus the surface energy balance and water cycle. Taking R_{net}^S as an example, their combined effects on R_{net}^S may be negative or positive and vary from -29.7 to 12.2 W m^{-2} , depending on the sign and magnitude of the effect of individual influencing factors (Figure 12). The non-spherical shape effects over TP are negative (Figure 12b), while the effects of mixing state of LAPs (BC and dust) are positive for both sphere and Koch snowflake with a smaller magnitude (Figure 12e-h). Different from the snow grain shape and mixing state, topography can affect the surface radiation budget over both the snow-covered and snow-free complex terrain (Figure 12c,d) due to the shadowing effects and multi-scattering from adjacent terrain. The TOP effects can be positive or negative, depending on the local topographic features. Thus interactions among the three influencing factors are complex and nonlinear, and their effects can be supplementary or cancelling. For instance, the TOP effects can be different under different snow grain shapes and they vary from -5.2 to 14.1 W m^{-2} under spherical shape and from -7.5 to 16.9 W m^{-2} under Koch snowflake shape (Figure 12c-d).

450




Figure 12: Spatial distributions of the change of net solar radiation (R_s^{net}) contributed by individual influencing factors and their combined effects in spring, which are derived based on Table 2. The corresponding results in winter are shown in Figure S14.

455

5. Discussion

Snow grain shape and mixing state of LAP-snow play an important role in snow processes, surface energy balance and water cycle. Snow grain shape has a large effect on α_{sno} and the Koch snowflake shows the largest differences from sphere, as reported in (He et al., 2018a). The reduction of snow albedo induced by LAP can be comparable to or even larger than the 460 impacts of snow grain shape when the LAP concentration in snow is high (Figures 5e-f and 7a-c). Mixing state of LAP-snow

also has impacts on SAR induced by BC and dust (He et al., 2018a; He et al., 2019), but overall the impacts of the mixing state of LAP-snow are smaller than the non-spherical shape effects (Sections 4.3 and 4.4). With the treatment of Koch snowflake shape, internal mixing of LAP-snow has smaller effects on α_{sno} than that of the spherical shape (Figure 9; He et al., 2018a). Besides, both the impacts of non-spherical snow grain shape and mixing states are sensitive to snow grain size (He et al., 465 2018a). However, there are still uncertainties in modeling the evolution of snow grain size in ELM (Figure S3), which needs further improvements. Their impacts on α_{sno} further affect the surface energy balance and water cycles (Section 4.3.2). For example, the Koch-snowflake shape decreases R_{net}^s by up to 30 W m⁻² compared to the spherical shape (Table 4). Topography also changes the apparent A_{sur} and solar radiation absorbed by the surface, and in turn affects snow dynamics (Hao et al., 2021a; Hao et al., 2021b; Hao et al., 2018). Compared to the negative effects of non-spherical shape on R_{net}^s , topography generally 470 has positive effects on R_{net}^s with a larger magnitude than mixing state of LAP-snow (Figure 12). Their effects can be additive or cancelling out each other and thus the combined effects may be cooling or warming (Figure 12), which depend on the specific snow grain shape, mixing state, and local topographic features. A novel topography-based sub-grid structure (topounit) (Tesfa and Leung, 2017) has been added in ELM. Furthermore, jointly including the improved snow albedo model from this 475 study, TOP solar radiation parameterization of Hao et al. (2021b) within the topounit structure, and considering the downscaling of snowfall based on topounit from (Tesfa et al., 2020) will be promising to improve the simulation of snow dynamics.

There remain some uncertainties in parameterizing the effects of LAPs on α_{sno} in ELM. A mono-disperse (uniform) snow grain 480 size distribution is assumed in the new treatments of Section 2.2, although non-uniform (e.g., lognormal) snow size distributions were observed and may have significant impacts on α_{sno} (Saito et al., 2019). The typical properties of BC and dust, such as refractive index, size distribution, and particle morphology from observations were adopted in the parameterizations of ELM (He et al., 2019; He et al., 2017). The maximum allowable BC and dust concentrations are 1000 ppb and 1000 ppm, respectively (He et al., 2019; He et al., 2017). The parameterizations assume that BC/dust particles 485 randomly distribute internally in the snow grains, but a non-random distribution of BC/dust in snow grains may affect the magnitude of SAR (Dombrovsky and Kokhanovsky, 2020; Shi et al., 2021). The parameterization for the effects of internally-mixed BC (or dust) assumes no presence of dust (or BC) in the snow. In order to avoid the possible overestimations of SAR, the simultaneous internal mixing of BC and dust in snow grains is not included in this study, which requires future 490 investigations. It is also assumed that different types of LAPs are externally mixed with each other. However, different types can also be internally mixed (i.e., coating or attachment) with each other (Chung et al., 2012; Pu et al., 2021), leading to an enhanced absorption through the so-called lensing effects (Bond et al., 2006; Lack et al., 2012). For instance, BC absorption enhancement is strongly relevant to the amount of coating material and is also source and region dependent (Liu et al., 2015).. The effects of BrC with high uncertainties are neglected in our study. In contrast to BC, sources, chemical composition and optical properties of BrC are still poorly understood (Wu et al., 2016) and its effects on RF and its contributions to SAR remain 495 highly uncertain (Beres et al., 2020). With some assumptions of the photochemical and optical properties of BrC in SNICAR

495 within the Community Land Model (version 4), Brown et al. (2021) found that the global (and TP) mean RF of BrC in snow can be comparable to that of BC. A more realistic representation of the impacts of snow grain properties and diverse LAPs on α_{sno} that is validated against field measurements and remote sensing data in land surface models (e.g., ELM) is required for climate change studies.

500 The magnitude and spatial distribution of snow cover in the ELM simulations are similar to the MODIS estimates in winter. Although ELM_New reduces the f_{sno} biases in ELM_Control, there are still underestimations in spring compared to the MODIS estimates (Figure S2). This underestimation is also found in CLM4.5 as reported by (Xie et al., 2018b), which used the same snow cover parameterizations of (Swenson and Lawrence, 2012) as in ELM v2.0. The underestimation may be caused by the use of a constant snow accumulation ratio, empirical snowmelt shape factor, and the underrepresentation of the complex 505 vegetation-snow interaction processes of snow interception and dynamical removal from canopy (Xie et al., 2018b). The canopy-gap effects on f_{sno} are not accounted for in ELM v2.0. Additionally, uncertainties of the atmospheric forcing data can affect the accuracy of the ELM snow simulations (Wang et al., 2020b). Although ELM v2.0 is run at a 0.125° spatial resolution, 0.5° GSWP3v1 atmospheric data is used in our simulations, which leads to a smaller spatial variability of snow simulated by ELM v2.0 compared to MODIS data (Figures 2 and 5). The wind-blown of snow and the subsequent sublimation process 510 also frequently affect the snow dynamics and are unaccounted for in ELM v2.0 (Orsolini et al., 2019; Xie et al., 2019).

515 There are large differences in SAR between ELM and MODIS data. ELM shows a similar magnitude to STC-MODDRFS in winter, but is larger than STC-MODDRFS in spring, while SPIRES shows lower values than ELM v2.0 in both winter and spring. The ELM results are in agreement with He et al. (2018a), showing that BC-induced SAR can be above 0.1 in the non-monsoon season, and the magnitude of SAR is comparable to the field measurements-based analysis in the central and western 520 Himalayas of (Gul et al., 2021). However, a few field measurements showed that the snowpacks in the Indus Basin are clean in winter (Negi et al., 2010). Spring snow has larger RF values from different LAPs than winter snow (Figure 6; Qian et al., 2011), due to larger LAP concentrations accumulated during snowmelt process (Kang et al., 2020) and larger incident solar radiation. Dust has a larger RF value than BC in spring over the northwestern regions of the TP (Figure 6f), as reported in 525 (Sarangi et al., 2020), and is related to the regional difference of the source, transport, and deposition of BC and dust. The use of prescribed climatological aerosol deposition data may in part contribute to the inconsistencies between model results and remote sensing data. The simulation biases of snow accumulation, melting, refreezing, compaction, aging, and water transport across snow layers in ELM can affect the SAR estimates. Note that there are large differences in SAR between STC-MODDRFS and SPIRES (Bair et al., 2021a), because the two models use different approaches to estimating SAR and interpolating SAR estimates. Snow estimates from MODIS are affected by frequent cloud cover over TP, coarse spectral resolution of MODIS, uncertainty in atmospheric correction, reflectance measurement errors, and the impacts of canopy cover (Bair et al., 2021b; Stillinger et al., 2022). Although f_{sno} estimated by STC-MODDRFS and SPIReS is relatively reliable, there 530 are still large uncertainties in the estimated snow grain size and LAP content and thus SAR due to the limited capability of

multi-spectral sensors to distinguish between darkening from micro-scale topography and LAPs in low concentrations (Bair 530 et al., 2022). The ongoing and upcoming hyperspectral satellites such as PRISMA (PRRecursore IperSpettrale della Missione Applicativa) mission, the Surface Biology and Geology (SBG) led by NASA and the Copernicus Hyperspectral Imaging Mission for the Environment (CHIME) led by European Space Agency (ESA) will be promising for improving the remotely sensed estimates of LAP-induced SAR. Long-term field measurements of snow grain characteristics, α_{sno} and LAP concentrations over the TP are needed to evaluate the model simulations and advance our understanding of LAP effects. Our 535 study focuses on the relative sensitivity rather than absolute accuracy, and thus the abovementioned uncertainties are expected to have only a small influence on our results.

There are some limitations of this study. Our sensitivity tests are based on assumptions of a spatiotemporally homogenous snow grain shape and mixing state of LAP-snow and currently there is no knowledge about the specific snow grain shape and 540 mixing state of dust-snow regionally or globally. Only four typical types of snow grain shapes are included in the analysis, but the real snow grain shape is more complicated and irregular (Kokhanovsky et al., 2005). Besides, in reality, both the snow grain shape and mixing state of LAP-snow are spatially-inhomogeneous and time-varying (Räisänen et al., 2015). For instance, dust may be partially internally and externally mixed with snow grains and simply assuming external mixing of dust-snow could underestimate the dust effects, while assuming fully dust-snow internal mixing will overestimate the dust effects (Shi et 545 al., 2021). The land-atmosphere interaction is neglected by performing offline ELM simulations over the TP. Further efforts are needed to better evaluate the global impacts of LAP by better coupling ELM and the atmospheric aerosol model. The impacts of BrC and snow algae are also excluded in this study, which may be comparable to BC and dust effects (Brown et al., 2021; Dang and Hegg, 2014; Di Mauro, 2020; Ganey et al., 2017).

6. Conclusions

550 Snow albedo is sensitive to snow grain shape and mixing state of LAP-snow. This study implemented the computationally efficient parameterizations for non-spherical grain shape (i.e., spheroid, hexagonal plate, and Koch snowflake) and the internal mixing of dust-snow in the snow radiative transfer model (i.e., SNICAR-AD) in ELM v2.0. Both ELM_Control and ELM_New show similar snow distribution in winter as two MODIS snow products (STC-MODSCAG/STC-MODDRFS and SPIReS). There are some noticeable statistical differences between ELM simulations and MODIS estimates in spring, and ELM_New 555 reduces the bias of f_{sno} in ELM_Control. The LAP-induced RFs in both ELM_Control and ELM_New in spring are also within the range of the reported values in previous studies, though considerably higher than MODIS snow products in spring. All the snow-related processes, surface energy balance and water cycles are sensitive to the treatment of snow grain shape. The Koch snowflake shape shows the largest difference from the default spherical shape treatment in ELM. The impacts of the mixing state of LAP-snow are smaller than the non-spherical shape effects, which also depend on snow grain shape. The effects of 560 non-spherical shape, mixing state of LAP-snow, and sub-grid topography on snow and surface fluxes have different signs and

magnitudes. Their combined effects are complex, non-linear and may be negative or positive depending on the specific snow grain shape, mixing state, and local topographic features. Overall, the changes of net solar radiation in spring due to individual and combined effects range from -28.6 to 16.9 W m⁻² and -29.7 to 12.2 W m⁻², respectively. This study advances our understanding of uncertainties in snow albedo modeling and its effects on surface energy and water cycles, and offers guidance
565 for improving the simulations of snow processes and RF estimates in ESMs. Future efforts are needed to couple the impacts of BrC and snow algae and investigate the climate effects of LAPs in snow via land-atmosphere coupling.

Code and data availability

The description and codes of E3SM v2.0 (including ELM v2.0) are publicly available at
570 <https://doi.org/10.11578/E3SM/dc.20210927.1> and <https://github.com/E3SM-Project/E3SM/releases/tag/v2.0.0> (last access: 12 April 2022), respectively. Starting from ELM 2.0, the model codes for snow albedo modeling improvements described in this paper is available at <https://doi.org/10.5281/zenodo.6324131>, and code to reproduce all results and plot all figures is publicly available at https://github.com/daleihao/Snow_Albedo_Parameterization_in_ELM and
<https://doi.org/10.5281/zenodo.6321316>.

575

Author contributions

DH designed the study, implemented the parameterization, performed the simulations, analyzed the results, and drafted the original manuscript. GB designed the study, discussed the results, and edited the manuscript. CH and CD helped with the snow albedo parameterization and edited the manuscript. EB, KR, and TS provided the remote sensing data and edited the
580 manuscript. HH, YG, HW, YQ, and LRL discussed the results and edited the manuscript.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

This research was conducted at Pacific Northwest National Laboratory, which is operated for the U.S. Department of Energy
585 by Battelle Memorial Institute under contract DEAC05-76RL01830. This research used resources of the National Energy Research Scientific Computing Center (NERSC), a DOE Office of Science User Facility supported by the Office of Science of the U.S. Department of Energy under contract no. DE-AC02-15 05CH11231. The reported research also used DOE's Biological and Environmental Research Earth System Modeling program's Compy computing cluster located at Pacific

Northwest National Laboratory. We thank Jeff Dozier for his help with snow albedo modeling. The STC-MODSCAG/STC-590 MODDRFS data is available at <ftp://snowserver.colorado.edu/pub/fromRittger/products/Indus>. The SPIReS data is available at <ftp://ftp.snow.ucsb.edu/pub/org/snow/products/SPIRES/Indus/>.

Financial support

595 This research has been supported by the U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research, Earth System Model Development program area, as part of the Climate Process Team projects and the U.S. National Oceanic and Atmospheric Administration (NOAA, grant no. NOAA-OAR-CPO-2019-2005530 and NA19OAR4310243). This research was also supported by NASA awards: 80NSSC21K0997, 80NSSC20K1349, 80NSSC20K1722, & 80NSSC18K1489. Y. Gu acknowledges the support by (while serving at) the National Science Foundation.

600

References

Andreae, M. O. and Gelencsér, A.: Black carbon or brown carbon? The nature of light-absorbing carbonaceous aerosols, *Atmos. Chem. Phys.*, 6, 3131-3148, 2006.

605 Bair, E., Stillinger, T., Rittger, K., and Skiles, M.: COVID-19 lockdowns show reduced pollution on snow and ice in the Indus River Basin, *Proceedings of the National Academy of Sciences*, 118, e2101174118, 2021a.

Bair, E. H., Dozier, J., Stern, C., LeWinter, A., Rittger, K., Savagian, A., Stillinger, T., and Davis, R. E.: Divergence of apparent and intrinsic snow albedo over a season at a sub-alpine site with implications for remote sensing, *The Cryosphere*, 16, 1765-1778, 2022.

610 Bair, E. H., Rittger, K., Skiles, S. M., and Dozier, J.: An Examination of Snow Albedo Estimates From MODIS and Their Impact on Snow Water Equivalent Reconstruction, *Water Resources Research*, 55, 7826-7842, 2019.

Bair, E. H., Stillinger, T., and Dozier, J.: Snow Property Inversion From Remote Sensing (SPIReS): A Generalized Multispectral Unmixing Approach With Examples From MODIS and Landsat 8 OLI, *IEEE Transactions on Geoscience and Remote Sensing*, 59, 7270-7284, 2021b.

615 Barnett, T. P., Adam, J. C., and Lettenmaier, D. P.: Potential impacts of a warming climate on water availability in snow-dominated regions, *Nature*, 438, 303-309, 2005.

Beres, N. D., Sengupta, D., Samburova, V., Khlystov, A. Y., and Moosmüller, H.: Deposition of brown carbon onto snow: changes in snow optical and radiative properties, *Atmos. Chem. Phys.*, 20, 6095-6114, 2020.

620 Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M. G., Ghan, S., Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schulz, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo, T., Warren, S. G., and Zender, C. S.: Bounding the role of black carbon in the climate system: A scientific assessment, *Journal of Geophysical Research: Atmospheres*, 118, 5380-5552, 2013.

Bond, T. C., Habib, G., and Bergstrom, R. W.: Limitations in the enhancement of visible light absorption due to mixing state, *Journal of Geophysical Research: Atmospheres*, 111, 2006.

625 Brown, H., Wang, H., Flanner, M., Liu, X., Singh, B., Zhang, R., Yang, Y., and Wu, M.: Brown Carbon Fuel and Emission Source Attributions to Global Snow Darkening Effect, *Earth and Space Science Open Archive*, doi: doi:10.1002/essoar.10507836.1, 2021. 37, 2021.

Chung, C., Lee, K., and Müller, D.: Effect of internal mixture on black carbon radiative forcing, *Tellus B: Chemical and Physical Meteorology*, 64, 10925, 2012.

630 Chýlek, P. and Srivastava, V.: Dielectric constant of a composite inhomogeneous medium, *Physical Review B*, 27, 5098, 1983.

Dang, C., Fu, Q., and Warren, S. G.: Effect of Snow Grain Shape on Snow Albedo, *Journal of the Atmospheric Sciences*, 73, 3573-3583, 2016.

Dang, C. and Hegg, D. A.: Quantifying light absorption by organic carbon in Western North American snow by serial chemical extractions, *Journal of Geophysical Research: Atmospheres*, 119, 10,247-210,261, 2014.

635 Dang, C., Zender, C. S., and Flanner, M. G.: Intercomparison and improvement of two-stream shortwave radiative transfer schemes in Earth system models for a unified treatment of cryospheric surfaces, *The Cryosphere*, 13, 2325-2343, 2019.

Di Mauro, B.: A darker cryosphere in a warming world, *Nature Climate Change*, 10, 979-980, 2020.

Dirmeyer, P. A., Gao, X., Zhao, M., Guo, Z., Oki, T., and Hanasaki, N.: GSWP-2: Multimodel Analysis and Implications for Our Perception of the Land Surface, *Bulletin of the American Meteorological Society*, 87, 1381-1398, 2006.

640 Dombrovsky, L. A. and Kokhanovsky, A. A.: Light absorption by polluted snow cover: Internal versus external mixture of soot, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 242, 106799, 2020.

Dominé, F., Lauzier, T., Cabanes, A., Legagneux, L., Kuhs, W. F., Techmer, K., and Heinrichs, T.: Snow metamorphism as revealed by scanning electron microscopy, *Microscopy Research and Technique*, 62, 33-48, 2003.

645 Erbe, E. F., Rango, A., Foster, J., Josberger, E. G., Pooley, C., and Wergin, W. P.: Collecting, shipping, storing, and imaging snow crystals and ice grains with low-temperature scanning electron microscopy, *Microscopy research and technique*, 62, 19-32, 2003.

Flanner, M. G., Arnheim, J., Cook, J. M., Dang, C., He, C., Huang, X., Singh, D., Skiles, S. M., Whicker, C. A., and Zender, C. S.: SNICAR-AD v3: A Community Tool for Modeling Spectral Snow Albedo, *Geosci. Model Dev. Discuss.*, 2021, 1-49, 2021.

650 Flanner, M. G., Liu, X., Zhou, C., Penner, J. E., and Jiao, C.: Enhanced solar energy absorption by internally-mixed black carbon in snow grains, *Atmos. Chem. Phys.*, 12, 4699-4721, 2012.

Flanner, M. G., Shell, K. M., Barlage, M., Perovich, D. K., and Tschudi, M. A.: Radiative forcing and albedo feedback from the Northern Hemisphere cryosphere between 1979 and 2008, *Nature Geoscience*, 4, 151-155, 2011.

655 Flanner, M. G., Zender, C. S., Hess, P. G., Mahowald, N. M., Painter, T. H., Ramanathan, V., and Rasch, P. J.: Springtime warming and reduced snow cover from carbonaceous particles, *Atmos. Chem. Phys.*, 9, 2481-2497, 2009.

Flanner, M. G., Zender, C. S., Randerson, J. T., and Rasch, P. J.: Present-day climate forcing and response from black carbon in snow, *Journal of Geophysical Research: Atmospheres*, 112, 2007.

Fu, Q.: A New Parameterization of an Asymmetry Factor of Cirrus Clouds for Climate Models, *Journal of the Atmospheric Sciences*, 64, 4140-4150, 2007.

660 Ganey, G. Q., Loso, M. G., Burgess, A. B., and Dial, R. J.: The role of microbes in snowmelt and radiative forcing on an Alaskan icefield, *Nature Geoscience*, 10, 754-759, 2017.

Golaz, J.-C., Caldwell, P. M., Van Roekel, L. P., Petersen, M. R., Tang, Q., Wolfe, J. D., Abeshu, G., Anantharaj, V., Asay-Davis, X. S., Bader, D. C., Baldwin, S. A., Bisht, G., Bogenschutz, P. A., Branstetter, M., Brunke, M. A., Brus, S. R., Burrows, S. M., Cameron-Smith, P. J., Donahue, A. S., Deakin, M., Easter, R. C., Evans, K. J., Feng, Y., Flanner, M., Foucar, J. G., Fyke, J. G., Griffin, B. M., Hannay, C., Harrop, B. E., Hoffman, M. J., Hunke, E. C., Jacob, R. L., Jacobsen, D. W., Jeffery, N., Jones, P. W., Keen, N. D., Klein, S. A., Larson, V. E., Leung, L. R., Li, H.-Y., Lin, W., Lipscomb, W. H., Ma, P.-L., Mahajan, S., Maltrud, M. E., Mametjanov, A., McClean, J. L., McCoy, R. B., Neale, R. B., Price, S. F., Qian, Y., Rasch, P. J., Reeves Eyre, J. E. J., Riley, W. J., Ringler, T. D., Roberts, A. F., Roesler, E. L., Salinger, A. G., Shaheen, Z., Shi, X., Singh, B., Tang, J., Taylor, M. A., Thornton, P. E., Turner, A. K., Veneziani, M., Wan, H., Wang, H., Wang, S., Williams, D. N., 665 Wolfram, P. J., Worley, P. H., Xie, S., Yang, Y., Yoon, J.-H., Zelinka, M. D., Zender, C. S., Zeng, X., Zhang, C., Zhang, K., Zhang, Y., Zheng, X., Zhou, T., and Zhu, Q.: The DOE E3SM Coupled Model Version 1: Overview and Evaluation at Standard Resolution, *Journal of Advances in Modeling Earth Systems*, 11, 2089-2129, 2019.

Grenfell, T. C. and Warren, S. G.: Representation of a nonspherical ice particle by a collection of independent spheres for scattering and absorption of radiation, *Journal of Geophysical Research: Atmospheres*, 104, 31697-31709, 1999.

675 Gul, C., Kang, S., Puppala, S. P., Wu, X., He, C., Xu, Y., Koch, I., Muhammad, S., Kumar, R., and Dubache, G.: Measurement of Light absorbing particles in surface snow of central and western Himalayan glaciers: spatial variability, radiative impacts, and potential source regions, *Atmos. Chem. Phys. Discuss.*, 2021, 1-21, 2021.

Hadley, O. L. and Kirchstetter, T. W.: Black-carbon reduction of snow albedo, *Nature Climate Change*, 2, 437-440, 2012.

Hall, A.: The Role of Surface Albedo Feedback in Climate, *Journal of Climate*, 17, 1550-1568, 2004.

680 Hao, D., Bisht, G., Gu, Y., Lee, W. L., Liou, K. N., and Leung, L. R.: A parameterization of sub-grid topographical effects on solar radiation in the E3SM Land Model (version 1.0): implementation and evaluation over the Tibetan Plateau, *Geosci. Model Dev.*, 14, 6273-6289, 2021a.

685 Hao, D., Bisht, G., Huang, M., Ma, P.-L., Tesfa, T. K., Lee, W.-L., Gu, Y., and Leung, L. R.: Impacts of sub-grid topographic representations on surface energy balance and boundary conditions in the E3SM Land Model, *Earth and Space Science Open Archive*, doi: doi:10.1002/essoar.10508175.1, 2021b. 37, 2021b.

690 Hao, D., Wen, J., Xiao, Q., Wu, S., Lin, X., Dou, B., You, D., and Tang, Y.: Simulation and Analysis of the Topographic Effects on Snow-Free Albedo over Rugged Terrain, *Remote Sensing*, 10, 278, 2018.

695 He, C. and Flanner, M.: Snow Albedo and Radiative Transfer: Theory, Modeling, and Parameterization. In: *Springer Series in Light Scattering: Volume 5: Radiative Transfer, Remote Sensing, and Light Scattering*, Kokhanovsky, A. (Ed.), Springer International Publishing, Cham, 2020.

700 He, C., Flanner, M. G., Chen, F., Barlage, M., Liou, K. N., Kang, S., Ming, J., and Qian, Y.: Black carbon-induced snow albedo reduction over the Tibetan Plateau: uncertainties from snow grain shape and aerosol–snow mixing state based on an updated SNICAR model, *Atmos. Chem. Phys.*, 18, 11507-11527, 2018a.

705 He, C., Li, Q., Liou, K.-N., Takano, Y., Gu, Y., Qi, L., Mao, Y., and Leung, L. R.: Black carbon radiative forcing over the Tibetan Plateau, *Geophysical Research Letters*, 41, 7806-7813, 2014.

710 He, C., Liou, K.-N., and Takano, Y.: Resolving Size Distribution of Black Carbon Internally Mixed With Snow: Impact on Snow Optical Properties and Albedo, *Geophysical Research Letters*, 45, 2697-2705, 2018b.

715 He, C., Liou, K.-N., Takano, Y., Chen, F., and Barlage, M.: Enhanced Snow Absorption and Albedo Reduction by Dust-Snow Internal Mixing: Modeling and Parameterization, *Journal of Advances in Modeling Earth Systems*, 11, 3755-3776, 2019.

720 He, C., Takano, Y., Liou, K.-N., Yang, P., Li, Q., and Chen, F.: Impact of Snow Grain Shape and Black Carbon?Snow Internal Mixing on Snow Optical Properties: Parameterizations for Climate Models, *Journal of Climate*, 30, 10019-10036, 2017.

Ji, Z.-M.: Modeling black carbon and its potential radiative effects over the Tibetan Plateau, *Advances in Climate Change Research*, 7, 139-144, 2016.

Kang, S., Zhang, Y., Qian, Y., and Wang, H.: A review of black carbon in snow and ice and its impact on the cryosphere, *Earth-Science Reviews*, 210, 103346, 2020.

725 Kaspari, S., Painter, T. H., Gysel, M., Skiles, S. M., and Schwikowski, M.: Seasonal and elevational variations of black carbon and dust in snow and ice in the Solu-Khumbu, Nepal and estimated radiative forcings, *Atmos. Chem. Phys.*, 14, 8089-8103, 2014.

Kokhanovsky, A. A., Aoki, T., Hachikubo, A., Hori, M., and Zege, E. P.: Reflective properties of natural snow: approximate asymptotic theory versus in situ measurements, *IEEE Transactions on Geoscience and Remote Sensing*, 43, 1529-1535, 2005.

Kopacz, M., Mauzerall, D. L., Wang, J., Leibensperger, E. M., Henze, D. K., and Singh, K.: Origin and radiative forcing of black carbon transported to the Himalayas and Tibetan Plateau, *Atmos. Chem. Phys.*, 11, 2837-2852, 2011.

LaChapelle, E. R.: *Field Guide to Snow Crystals*, 1969.

Lack, D. A., Langridge, J. M., Bahreini, R., Cappa, C. D., Middlebrook, A. M., and Schwarz, J. P.: Brown carbon and internal mixing in biomass burning particles, *Proceedings of the National Academy of Sciences*, 109, 14802, 2012.

Lamarque, J. F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Liouesse, C., Mieville, A., Owen, B., Schultz, M. G., Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne, J., Cooper, O. R., Kainuma, M., Mahowald, N., McConnell, J. R., Naik, V., Riahi, K., and van Vuuren, D. P.: Historical (1850–2000) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols: methodology and application, *Atmos. Chem. Phys.*, 10, 7017-7039, 2010.

730 Leung, L. R., Bader, D. C., Taylor, M. A., and McCoy, R. B.: An Introduction to the E3SM Special Collection: Goals, Science Drivers, Development, and Analysis, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS001821, 2020.

Libois, Q., Picard, G., France, J. L., Arnaud, L., Dumont, M., Carmagnola, C. M., and King, M. D.: Influence of grain shape on light penetration in snow, *The Cryosphere*, 7, 1803-1818, 2013.

735 Liou, K. N., Takano, Y., He, C., Yang, P., Leung, L. R., Gu, Y., and Lee, W. L.: Stochastic parameterization for light absorption by internally mixed BC/dust in snow grains for application to climate models, *Journal of Geophysical Research: Atmospheres*, 119, 7616-7632, 2014.

Liu, S., Aiken, A. C., Gorkowski, K., Dubey, M. K., Cappa, C. D., Williams, L. R., Herndon, S. C., Massoli, P., Fortner, E. C., Chhabra, P. S., Brooks, W. A., Onasch, T. B., Jayne, J. T., Worsnop, D. R., China, S., Sharma, N., Mazzoleni, C., Xu, L.,

730 Ng, N. L., Liu, D., Allan, J. D., Lee, J. D., Fleming, Z. L., Mohr, C., Zotter, P., Szidat, S., and Prévôt, A. S. H.: Enhanced light absorption by mixed source black and brown carbon particles in UK winter, *Nature Communications*, 6, 8435, 2015.

735 Liu, X., Easter, R. C., Ghan, S. J., Zaveri, R., Rasch, P., Shi, X., Lamarque, J. F., Gettelman, A., Morrison, H., Vitt, F., Conley, A., Park, S., Neale, R., Hannay, C., Ekman, A. M. L., Hess, P., Mahowald, N., Collins, W., Iacono, M. J., Bretherton, C. S., Flanner, M. G., and Mitchell, D.: Toward a minimal representation of aerosols in climate models: description and evaluation in the Community Atmosphere Model CAM5, *Geosci. Model Dev.*, 5, 709-739, 2012.

740 Myneni, R. B., Hoffman, S., Knyazikhin, Y., Privette, J. L., Glassy, J., Tian, Y., Wang, Y., Song, X., Zhang, Y., Smith, G. R., Lotsch, A., Friedl, M., Morisette, J. T., Votava, P., Nemani, R. R., and Running, S. W.: Global products of vegetation leaf area and fraction absorbed PAR from year one of MODIS data, *Remote Sensing of Environment*, 83, 214-231, 2002.

Negi, H. S., Singh, S. K., Kulkarni, A. V., and Semwal, B. S.: Field-based spectral reflectance measurements of seasonal snow cover in the Indian Himalaya, *International Journal of Remote Sensing*, 31, 2393-2417, 2010.

745 Orsolini, Y., Wegmann, M., Dutra, E., Liu, B., Balsamo, G., Yang, K., de Rosnay, P., Zhu, C., Wang, W., Senan, R., and Arduini, G.: Evaluation of snow depth and snow cover over the Tibetan Plateau in global reanalyses using in situ and satellite remote sensing observations, *The Cryosphere*, 13, 2221-2239, 2019.

Painter, T. H., Bryant, A. C., and Skiles, S. M.: Radiative forcing by light absorbing impurities in snow from MODIS surface reflectance data, *Geophysical Research Letters*, 39, 2012.

750 Painter, T. H., Rittger, K., McKenzie, C., Slaughter, P., Davis, R. E., and Dozier, J.: Retrieval of subpixel snow covered area, grain size, and albedo from MODIS, *Remote Sensing of Environment*, 113, 868-879, 2009.

Pu, W., Shi, T., Cui, J., Chen, Y., Zhou, Y., and Wang, X.: Enhancement of snow albedo reduction and radiative forcing due to coated black carbon in snow, *The Cryosphere*, 15, 2255-2272, 2021.

755 Qian, Y., Flanner, M. G., Leung, L. R., and Wang, W.: Sensitivity studies on the impacts of Tibetan Plateau snowpack pollution on the Asian hydrological cycle and monsoon climate, *Atmos. Chem. Phys.*, 11, 1929-1948, 2011.

Räisänen, P., Kokhanovsky, A., Guyot, G., Jourdan, O., and Nousiainen, T.: Parameterization of single-scattering properties of snow, *The Cryosphere*, 9, 1277-1301, 2015.

Räisänen, P., Makkonen, R., Kirkevåg, A., and Debernard, J. B.: Effects of snow grain shape on climate simulations: sensitivity tests with the Norwegian Earth System Model, *The Cryosphere*, 11, 2919-2942, 2017.

760 Riihelä, A., Bright, R. M., and Anttila, K.: Recent strengthening of snow and ice albedo feedback driven by Antarctic sea-ice loss, *Nature Geoscience*, 14, 832-836, 2021.

Rittger, K., Painter, T. H., and Dozier, J.: Assessment of methods for mapping snow cover from MODIS, *Advances in Water Resources*, 51, 367-380, 2013.

Rittger, K., Raleigh, M. S., Dozier, J., Hill, A. F., Lutz, J. A., and Painter, T. H.: Canopy Adjustment and Improved Cloud Detection for Remotely Sensed Snow Cover Mapping, *Water Resources Research*, 56, e2019WR024914, 2020.

765 Saito, M., Yang, P., Loeb, N. G., and Kato, S.: A Novel Parameterization of Snow Albedo Based on a Two-Layer Snow Model with a Mixture of Grain Habits, *Journal of the Atmospheric Sciences*, 76, 1419-1436, 2019.

Sarangi, C., Qian, Y., Rittger, K., Bormann, K. J., Liu, Y., Wang, H., Wan, H., Lin, G., and Painter, T. H.: Impact of light-absorbing particles on snow albedo darkening and associated radiative forcing over high-mountain Asia: high-resolution WRF-Chem modeling and new satellite observations, *Atmos. Chem. Phys.*, 19, 7105-7128, 2019.

770 Sarangi, C., Qian, Y., Rittger, K., Ruby Leung, L., Chand, D., Bormann, K. J., and Painter, T. H.: Dust dominates high-altitude snow darkening and melt over high-mountain Asia, *Nature Climate Change*, 10, 1045-1051, 2020.

Shi, T., Cui, J., Chen, Y., Zhou, Y., Pu, W., Xu, X., Chen, Q., Zhang, X., and Wang, X.: Enhanced light absorption and reduced snow albedo due to internally mixed mineral dust in grains of snow, *Atmos. Chem. Phys.*, 21, 6035-6051, 2021.

Skiles, S. M., Flanner, M., Cook, J. M., Dumont, M., and Painter, T. H.: Radiative forcing by light-absorbing particles in snow, *Nature Climate Change*, 8, 964-971, 2018.

Sterle, K. M., McConnell, J. R., Dozier, J., Edwards, R., and Flanner, M. G.: Retention and radiative forcing of black carbon in eastern Sierra Nevada snow, *The Cryosphere*, 7, 365-374, 2013.

775 Stillinger, T., Rittger, K., Raleigh, M. S., Michell, A., Davis, R. E., and Bair, E. H.: Landsat, MODIS, and VIIRS snow cover mapping algorithm performance as validated by airborne lidar datasets, *The Cryosphere Discuss.*, 2022, 1-37, 2022.

Swenson, S. C. and Lawrence, D. M.: A new fractional snow-covered area parameterization for the Community Land Model and its effect on the surface energy balance, *Journal of Geophysical Research: Atmospheres*, 117, 2012.

780 Tanikawa, T., Kuchiki, K., Aoki, T., Ishimoto, H., Hachikubo, A., Niwano, M., Hosaka, M., Matoba, S., Kodama, Y., Iwata, Y., and Stammes, K.: Effects of Snow Grain Shape and Mixing State of Snow Impurity on Retrieval of Snow Physical Parameters From Ground-Based Optical Instrument, *Journal of Geophysical Research: Atmospheres*, 125, e2019JD031858, 2020.

785 Tesfa, T. K., Leung, L. R., and Ghan, S. J.: Exploring Topography-Based Methods for Downscaling Subgrid Precipitation for Use in Earth System Models, *Journal of Geophysical Research: Atmospheres*, 125, e2019JD031456, 2020.

790 Tesfa, T. K. and Leung, L. Y. R.: Exploring new topography-based subgrid spatial structures for improving land surface modeling, *Geosci. Model Dev.*, 10, 873-888, 2017.

795 Usha, K. H., Nair, V. S., and Babu, S. S.: Modeling of aerosol induced snow albedo feedbacks over the Himalayas and its implications on regional climate, *Climate Dynamics*, 54, 4191-4210, 2020.

800 Wang, H., Easter, R. C., Zhang, R., Ma, P.-L., Singh, B., Zhang, K., Ganguly, D., Rasch, P. J., Burrows, S. M., Ghan, S. J., Lou, S., Qian, Y., Yang, Y., Feng, Y., Flanner, M., Leung, L. R., Liu, X., Shrivastava, M., Sun, J., Tang, Q., Xie, S., and Yoon, J.-H.: Aerosols in the E3SM Version 1: New Developments and Their Impacts on Radiative Forcing, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS001851, 2020a.

805 Wang, Y., Xie, Z., Jia, B., Wang, L., Li, R., Liu, B., Chen, S., Xie, J., and Qin, P.: Sensitivity of Snow Simulations to Different Atmospheric Forcing Data Sets in the Land Surface Model CAS-LSM, *Journal of Geophysical Research: Atmospheres*, 125, e2019JD032001, 2020b.

810 Warren, S. G.: Optical properties of snow, *Reviews of Geophysics*, 20, 67-89, 1982.

815 Wu, G.-M., Cong, Z.-Y., Kang, S.-C., Kawamura, K., Fu, P.-Q., Zhang, Y.-L., Wan, X., Gao, S.-P., and Liu, B.: Brown carbon in the cryosphere: Current knowledge and perspective, *Advances in Climate Change Research*, 7, 82-89, 2016.

820 Wu, L., Gu, Y., Jiang, J. H., Su, H., Yu, N., Zhao, C., Qian, Y., Zhao, B., Liou, K. N., and Choi, Y. S.: Impacts of aerosols on seasonal precipitation and snowpack in California based on convection-permitting WRF-Chem simulations, *Atmos. Chem. Phys.*, 18, 5529-5547, 2018.

825 Xie, X., Liu, X., Che, H., Xie, X., Li, X., Shi, Z., Wang, H., Zhao, T., and Liu, Y.: Radiative feedbacks of dust in snow over eastern Asia in CAM4-BAM, *Atmos. Chem. Phys.*, 18, 12683-12698, 2018a.

830 Xie, Z., Hu, Z., Ma, Y., Sun, G., Gu, L., Liu, S., Wang, Y., Zheng, H., and Ma, W.: Modeling Blowing Snow Over the Tibetan Plateau With the Community Land Model: Method and Preliminary Evaluation, *Journal of Geophysical Research: Atmospheres*, 124, 9332-9355, 2019.

835 Xie, Z., Hu, Z., Xie, Z., Jia, B., Sun, G., Du, Y., and Song, H.: Impact of the snow cover scheme on snow distribution and energy budget modeling over the Tibetan Plateau, *Theoretical and Applied Climatology*, 131, 951-965, 2018b.

840 Xu, B., Cao, J., Hansen, J., Yao, T., Joswia, D. R., Wang, N., Wu, G., Wang, M., Zhao, H., Yang, W., Liu, X., and He, J.: Black soot and the survival of Tibetan glaciers, *Proceedings of the National Academy of Sciences*, 106, 22114, 2009.

845 Zhang, R., Wang, H., Qian, Y., Rasch, P. J., Easter, R. C., Ma, P. L., Singh, B., Huang, J., and Fu, Q.: Quantifying sources, transport, deposition, and radiative forcing of black carbon over the Himalayas and Tibetan Plateau, *Atmos. Chem. Phys.*, 15, 6205-6223, 2015.

850 Zhao, C., Hu, Z., Qian, Y., Ruby Leung, L., Huang, J., Huang, M., Jin, J., Flanner, M. G., Zhang, R., Wang, H., Yan, H., Lu, Z., and Streets, D. G.: Simulating black carbon and dust and their radiative forcing in seasonal snow: a case study over North China with field campaign measurements, *Atmos. Chem. Phys.*, 14, 11475-11491, 2014.