

# An EnKF-based ocean data assimilation system improved by adaptive observation error inflation (AOEI)

Shun Ohishi<sup>1,2,3</sup>, Takemasa Miyoshi<sup>1,2,4</sup>, and Misako Kachi<sup>5</sup>

<sup>1</sup>RIKEN Center for Computational Science, Kobe, 6500047, Japan

5 <sup>2</sup>RIKEN Cluster for Pioneering Research, Kobe, 6500047, Japan

<sup>3</sup>Institute for Space-Earth Environmental Research, Nagoya University, Nagoya, 4648601, Japan

<sup>4</sup>RIKEN Interdisciplinary Theoretical and Mathematical Sciences Program (iTHEMS), Kobe, 6500047, Japan

<sup>5</sup>Earth Observation Research Center, Japan Aerospace Exploration Agency, Tsukuba, 3058505, Japan

10 *Correspondence to:* Shun Ohishi ([shun.ohishi@riken.jp](mailto:shun.ohishi@riken.jp)) and Takemasa Miyoshi ([takemasa.miyoshi@riken.jp](mailto:takemasa.miyoshi@riken.jp))

## Abstract.

A previous study proposed an adaptive observation error inflation (AOEI) method for an ensemble Kalman filter-based atmospheric data assimilation system to assimilate all-sky infrared brightness temperatures. Brightness temperature differences between clear- and cloudy-sky radiances are large, and observation-minus-forecast differences or innovations are therefore likely to be large around boundaries between clear- and cloudy-sky regions. The AOEI method mitigates these discrepancies by adaptively inflating observation errors. Ocean frontal regions have similar characteristics to the borders between clear- and cloudy-sky regions with large innovations. Consequently, we have implemented the AOEI with an EnKF-based regional ocean data assimilation system, in which the assimilation interval is set to one day to utilize frequent satellite observations. We conducted sensitivity experiments to investigate the impacts of the AOEI on salinity structure, geostrophic balance, and accuracy. A control run, in which the AOEI is not applied, shows the degradation of low-salinity North Pacific Intermediate Water around the Kuroshio Extension region, where the innovation amplitude and forecast ensemble spread are large in association with the fronts and eddies. The resulting large temperature and salinity increments weaken the density stratification, leading to large vertical diffusivity. As a result, the low salinity water in the intermediate layer is lost through strong vertical diffusion. When the AOEI is used, the salinity structure in the ocean interior is preserved because the AOEI suppresses the salinity degradation by reducing the temperature and salinity increments. We also demonstrate that the AOEI provides significant improvement of the geostrophic balance and the accuracy of temperature, salinity, and surface flow fields.

## Short summary (487/500 character)

An adaptive observation error inflation (AOEI) method was proposed for atmospheric data assimilation to mitigate erroneous analysis updates caused by large observation-minus-forecast differences for satellite brightness temperature around clear- and cloudy-sky boundaries. This study implemented the AOEI with an ocean data assimilation system, leading to an improvement of analysis accuracy and dynamical balance around the frontal regions with large meridional temperature differences.

## 1. Introduction

The ensemble Kalman filter (EnKF) estimates flow-dependent forecast errors from an ensemble of model forecasts, and calculates the best estimates (i.e., analyses) by combining forecasts and observations with their error covariances (Evensen, 1994, 2003). The EnKF has the advantage of being easy to implement for various models (See table 1 of Ohishi et al. in review), but it has been used in only two ocean reanalysis datasets thus far (Balmaseda et al., 2015; Martin et al., 2015): the Predictive Ocean Atmosphere Model for Australia (PAOMA) Ensemble Ocean Data Assimilation System (PEODAS; Yin et al., 2011) and TOPAZ4 (Sakov et al., 2012). In contrast, the three-dimensional variational method (3D-VAR) is the most widely used in ocean analysis datasets (e.g. Miyazawa et al., 2017; Zuo et al., 2019).

With the enhancement of in-situ and satellite observations, the number of observations has increased dramatically. Argo profiling float observations since the 2000s provide a large number of in-situ temperature and salinity data in the ocean interior. Although satellite sea surface salinity (SSS) data since 2010 are relatively inaccurate, particularly in coastal and high-latitude regions (Abe and Ebuchi, 2014), previous studies have demonstrated the positive effects of SSS assimilation on the analyses of ocean interior structures such as mixed and barrier layers (Chakraborty et al., 2015), low salinity water caused by river discharge (Toyoda et al., 2015), and El Niño-Southern Oscillation prediction (Hackert et al., 2011). A Japanese geostationary satellite, Himawari-8 (Bessho et al., 2016; Kurihara et al., 2016), has observed sea surface temperatures (SSTs) in the Pacific region at a high spatiotemporal resolution of 2 km and 10 minutes since July 2015. The Surface Water and Ocean Topography (SWOT) satellite is scheduled to be launched in 2022 and will provide high-resolution and two-dimensional sea surface height (SSH) anomalies (SSHAs).

For effective use of dense and frequent satellite observations, Ohishi et al. (in review) performed sensitivity experiments using an EnKF-based ocean data assimilation system with an assimilation interval of one day, which is more frequent than the five- and seven-day intervals in the existing EnKF-based systems (PEODAS and TOPAZ4, respectively). They demonstrated that the combination of incremental analysis updates (IAU; Bloom et al., 1996) and relaxation-to-prior perturbations (RTPP; Kotsuki et al., 2017; Zhang et al., 2004) to restore the forecast ensemble perturbations toward the analysis by 80%–90% produced optimal results in terms of both dynamic balance and accuracy. However, their system contained several tuning parameters such as observation errors, ensemble size, and localization scale. Previous studies have prescribed observation errors in various ways, e.g., using spatiotemporally fixed constants (Miyazawa et al., 2012; Xu and Oey, 2014); assuming them to be standard deviations calculated from historical observations (Miyazawa et al., 2009; Usui et al., 2006); estimating them from other assimilation datasets (Penny et al., 2013); and assuming that observation error covariance matrices are proportional to the forecast error covariance matrices (Carton et al., 2018; Yin et al., 2011). A technique to adaptively inflate the observation errors based on the innovation statistics (Desroziers et al., 2005), which is called as an adaptive observation error inflation (AOEI) method, was recently proposed for assimilating all-sky infrared satellite brightness temperatures in an atmospheric data assimilation system (Minamide and Zhang, 2017; Zhang et al., 2016). Since the brightness temperature differences between clear- and cloudy-sky radiances are large, there are large observation-minus-forecast differences or innovations around boundaries between clear- and cloudy-sky regions even for the tiny boundary differences

Deleted: is

Deleted: and

70 between forecasts and observations. This results in erroneous analysis increments and degrades the analysis. The AOEI  
mitigates the large discrepancies between forecasts and observations by adaptively inflating the observation errors. Ocean  
frontal regions such as Kuroshio and Kuroshio Extension (KE) regions have large spatiotemporal variations, and the innovations  
around the frontal regions also tend to be large even for small differences of frontal positions between the forecasts and  
observations. Therefore, ocean fronts have similar characteristics to the borders between clear- and cloudy-sky regions with  
large innovations, and the AOEI method is therefore expected to be useful for improving EnKF-based ocean data assimilation  
75 systems.

This study aims to investigate the causes of the salinity degradation around the KE region, and to evaluate the impacts  
of the AOEI on the salinity structure, dynamical balance, and accuracy. The remainder of this paper is organized as follows:  
details of the AOEI method, the experiment design, [the temperature and salinity budget equations](#), and the methods to evaluate  
geostrophic balance and accuracy in sensitivity experiments are presented in Sect. 2; Sect. 3 describes the causes of the salinity  
80 degradation in the intermediate layer and the positive impacts of the AOEI on the geostrophic balance and accuracy for  
temperature, salinity, and surface flow. A summary is provided in Sect. 4.

## 2. Methods

### 2.1. Adaptive observation error inflation (AOEI)

85 Manual tuning of observation errors is computationally expensive, and several studies have proposed adaptive estimation  
methods using the innovation statistics of Desroziers et al. (2005):

$$\langle d_b^o (d_b^o)^T \rangle \approx \mathbf{H} \mathbf{P}^b \mathbf{H}^T + \mathbf{R}. \quad (1)$$

90 Here,  $\langle \cdot \rangle$  denotes the statistical expectation,  $d_b^o (= \mathbf{y} - \mathbf{H} \bar{\mathbf{x}}^b)$  is an innovation vector, where  $\mathbf{y}$ ,  $\mathbf{H}$ , and  $\bar{\mathbf{x}}^b$  denote an  
observation vector, linear observation operator, and forecast ensemble mean state vector, respectively.  $\mathbf{P}^b$  and  $\mathbf{R}$  are the  
forecast and observation error covariance matrices, respectively. Expressing Eq. (1) in a scalar form, the observation error  
 $\sigma_{est-o}$  may be estimated by

$$95 \sigma_{est-o}^2 = (d_b^o)^2 - \sigma_{H(x^b)}^2, \quad (2)$$

where  $\sigma_{H(x^b)}$  is the forecast ensemble spread in observation space. [Here, the forecast ensemble spreads are assumed to be  
accurate, and  \$\(d\_b^o\)^2\$  is assumed to be equivalent to  \$\langle \(d\_b^o\)^2 \rangle\$ . To avoid underestimation of the observation errors, in the AOEI  
method \(Minamide and Zhang, 2017; Zhang et al., 2016\), larger observation errors  \$\sigma\_o\$  between the estimated and prescribed  
100 errors are used:](#)

Deleted: I

$$\sigma_o^2 = \max\{\sigma_{pre-o}^2, \sigma_{est-o}^2\}, \quad (3)$$

105 where  $\sigma_{pre-o}$  is the prescribed observation error. As described in Sect. 1, the AOEI suppresses erroneous analysis increments associated with systematic errors, biases, and representation errors by adaptively inflating the observation errors when the squared innovation is larger than the sum of the prescribed observation and ensemble-based forecast error variances.

**Deleted:** the forecast ensemble spreads are assumed to be correct, and that  $(d_B^f)^2$  is assumed to be equivalent to  $((d_B^f)^2)$

**Deleted:** We note that in the AOEI.

## 2.2. Experimental design

110 This study uses an EnKF-based regional ocean data assimilation system known as sbPOM-LETKF (Ohishi et al. in review), comprising a sigma-coordinate regional ocean model, the Stony Brook Parallel Ocean Model version 1.0 (sbPOM; Jordi and Wang, 2012; Ohishi et al. in review), and a three-dimensional local ensemble transform Kalman filter (3D-LETKF; Hunt et al., 2007; Miyoshi and Yamane, 2007). The sbPOM is configured for the northwestern Pacific region (117°–180°E, 15°–50°N) with horizontal resolution of 0.25° and 50 sigma layers. The bottom topography is taken from a 1 arc-minute global relief  
 115 model of Earth’s surface (ETOPO1; Amante and Eakins, 2009) and is smoothed by a Gaussian filter with a 200 km e-folding scale to reduce pressure gradient errors at steep bottom slopes (Mellor et al., 1994). Monthly (seasonal) temperature and salinity climatologies from the World Ocean Atlas 2018 (WOA18; Locarnini et al., 2019; Zweng et al., 2019) with horizontal resolution of 1° and 57 (103) layers are used for the initial conditions over depths shallower (deeper) than 1500 m. Lateral boundary conditions for temperature, salinity, and horizontal velocity are derived from the Simpler Ocean Data Assimilation (SODA)  
 120 version 3.7.2 (Carton et al., 2018) with horizontal resolution of 0.5° and 50 layers. The Japanese 55-year Reanalysis (JRA55; Kobayashi et al., 2015) with horizontal and temporal resolution of 1.25° and 6 hours, respectively, is adopted for the atmospheric boundary conditions including air temperature and specific humidity at 2 m, wind velocity at 10 m, shortwave radiation, total cloud fraction, sea level pressure, and precipitation. River discharge is obtained from the Japan Aerospace Exploration Agency (JAXA)’s land surface and river simulation system, Today’s Earth (TE)-Global  
 125 (<https://www.eorc.jaxa.jp/water/>), with horizontal and temporal resolution of 0.25° and 3 hours, respectively. To avoid filter divergence, the atmospheric and lateral boundary conditions other than rainfall and river discharge are perturbed in the same way as in Ohishi et al. (in review). The model with 100 ensemble members is spun up from 1 January 2011 to 6 July 2015, using the initial conditions with no motion. During the spin-up period, simulated temperature and salinity are nudged towards the monthly climatology from the WOA18 with a 90-day timescale to prevent northward overshoot of the Kuroshio along the  
 130 east coast of Japan.

The LETKF with 100 ensemble members is used to assimilate the following observations on a one-day assimilation interval: satellite SSTs from Himawari-8 (Bessho et al., 2016; Kurihara et al., 2016) and the Global Change Observation Mission–Water (GCOM-W: <https://gportal.jaxa.jp/gpr/?lang=en>); satellite SSS from Soil Moisture and Ocean Salinity (SMOS: [http://www.esa.int/Applications/Observing\\_the\\_Earth/SMOS](http://www.esa.int/Applications/Observing_the_Earth/SMOS)) and Soil Moisture Active Passive (SMAP) version 4.3  
 135 (Meissner et al., 2018); SSH estimated by summing satellite SSH anomalies from the Copernicus Marine Environment Monitoring Service (CMEMS: <https://marine.copernicus.eu/>) and mean dynamic ocean topography obtained by averaging the

140 simulated SSH over 2012–14; and in-situ temperatures and salinity from the Global Temperature and Salinity Profile Programme (GTSP; Sun et al., 2010) and Advance automatic QC (AQC) Argo Data version 1.2a (AQC: [http://www.jamstec.go.jp/ARGO/argo\\_web/argo/?page\\_id=100&lang=en](http://www.jamstec.go.jp/ARGO/argo_web/argo/?page_id=100&lang=en)).

145 **Covariance localization in the observation space is applied using the Gaussian function with horizontal and vertical localization scales  $LS = 300$  km and  $100$  m, respectively, following Miyazawa et al. (2012) and Penny et al. (2013). We assume that the localization function becomes zero beyond  $2\sqrt{10/3}LS \approx 1100$  km (370 m) in the horizontal (vertical) direction (Miyoshi et al., 2007). Following Miyazawa et al. (2012), the prescribed observation errors for temperature, salinity, and SSH are set to  $1.0$  °C,  $0.3$ , and  $0.2$  m, respectively. Here, we set the larger salinity observation errors than those from Miyazawa et al. (2012), because the SSS satellite observations are relatively noisy and the measurement errors would be large (Meissner et al., 2018). We adopt the combination of the IAU (Bloom et al., 1996; Ohishi et al. in review) and RTPP (Kotsuki et al., 2017; Zhang et al., 2004), in which the analysis ensemble perturbations are relaxed toward the forecast ensemble perturbations by 90 % while maintaining the analysis ensemble mean, because the sensitivity experiments in Ohishi et al. (in review) demonstrated that it results in the best dynamical balance and accuracy. Although this may not be optimal, our computational resources are limited, and thus the RTPP relaxation parameter is fixed at 90%.**

155 To highlight the impacts of the AOEI on the ocean salinity structure and dynamical balance, we conduct AOEI and control (CTL) runs with and without applying the AOEI, respectively, from the start date of the Himawari-8 observation (7 July 2015) to 31 December 2015. Furthermore, we perform the 1.5Terr run with the same setting as the CTL run but with the temperature observation error of  $1.5$  °C, and compare the accuracy between the CTL, AOEI, and 1.5Terr runs. During the assimilation period, the SSS nudging with a 90-day timescale is applied to prevent a surface freshening drift as in the spin-up period.

Commented [SO1]: Minor comment #1

Deleted: T

Commented [SO2]: Minor comment #2

Commented [SO3]: Major Comment #3

Deleted: evaluate

Deleted: ,

Deleted: , and accuracy

160

### 2.3. Temperature and salinity budget equations in the ocean interior

To quantitatively investigate ocean interior temperature and salinity differences between the AOEI and CTL runs, respectively, we use the temperature  $T$  and salinity  $S$  budget equations:

$$165 \frac{\partial T}{\partial t} = \nabla \cdot (\kappa \circ \nabla T) - \mathbf{v} \cdot \nabla T + \frac{q_{sw}}{\rho_0 c_p} + (\text{increment}) \text{ and} \quad (4)$$

$$\frac{\partial S}{\partial t} = \nabla \cdot (\kappa \circ \nabla S) - \mathbf{v} \cdot \nabla S + (\text{increment}). \quad (5)$$

170 Here,  $\nabla = (\partial/\partial x, \partial/\partial y, \partial/\partial z)$  denotes the three-dimensional gradient operator.  $\kappa = (\kappa_x, \kappa_y, \kappa_z)$  is a diffusivity vector.  $\circ$  indicates a Schur product.  $\mathbf{v} = (u, v, w)$  is three-dimensional velocity.  $\rho_0 = 1025 \text{ kg m}^{-3}$  is the reference density.  $c_p = 4190 \text{ J kg}^{-1} \text{ }^\circ\text{C}^{-1}$  is the specific heat of the seawater, and  $q_{sw}$  is downward shortwave radiation parameterized by

$$q_{sw} = Q_{sw} \left\{ R_{sw} \exp\left(-\frac{|z|}{\gamma_1}\right) + (1 - R_{sw}) \exp\left(-\frac{|z|}{\gamma_2}\right) \right\} \quad (6)$$

(Paulson and Simpson, 1977), where  $Q_{sw}$  is shortwave radiation at the sea surface,  $R_{sw} = 0.62$  is a separation constant, and  $\gamma_1 = 0.60 \text{ m}$  and  $\gamma_2 = 20.0 \text{ m}$  are attenuation length scales. These values are set to the case of Type IA from Jerlov (1976). (increment) indicates the analysis increments. Equations (4) and (5) do not include a residual term because the system conserves each term of the temperature and salinity budget equations, respectively.

#### 2.4. Evaluation method

As in Ohishi et al. (in review), this study evaluates geostrophic balance and accuracy using the nonlinear balance equation (NBE) and root mean square deviations (RMSDs) relative to observations, respectively (see subsections 2.4.1 and 2.4.2).

##### 2.4.1. Nonlinear balance equation

For the analysis fields, the geostrophic balance equation is represented as

$$f \mathbf{k} \times \delta \mathbf{u} = -g \nabla_h \delta \eta, \quad (7)$$

where  $f$  is the vertical component of the Coriolis parameter,  $\mathbf{k}$  is a unit vector in the vertical direction,  $\delta$  is the analysis increment,  $\mathbf{u} = (u, v)$  denotes horizontal velocity at the sea surface,  $g = 9.8 \text{ m s}^{-2}$  is gravitational acceleration,  $\nabla_h = (\partial/\partial x, \partial/\partial y)$  is the horizontal gradient operator, and  $\eta$  denotes SSH. By  $\partial/\partial x$  of the  $x$ -component and  $\partial/\partial y$  of the  $y$ -component of Eq. (7), the geostrophic equation can be reduced to the nonlinear balance equation (NBE; Shibuya et al., 2015; Zhang et al., 2001):

$$-f \delta \zeta + \beta \delta u + g \nabla_h^2 \delta \eta = 0, \quad (8)$$

where  $\zeta = \partial v/\partial x - \partial u/\partial y$  is the relative vorticity at the sea surface and  $\beta = \partial f/\partial y$  is the planetary vorticity gradient. If the analysis fields do not satisfy geostrophic balance, there is an absolute NBE residual:

$$\Delta NBE \equiv \text{abs}(-f \delta \zeta + \beta \delta u + g \nabla_h^2 \delta \eta), \quad (9)$$

where  $\text{abs}$  indicates taking the absolute value. A smaller (larger)  $\Delta NBE$  indicates more (less) geostrophic balance.

##### 2.4.2. RMSD

Deleted: 2.3

Deleted: 2.3.1

Deleted: 2.3.2

Deleted: 2.3.1

Deleted: 4

Deleted: 4

Deleted: 5

Deleted: 6

Deleted: 2.3.2

We evaluate the accuracy of temperature, salinity, horizontal velocities, and SSH using the RMSDs calculated relative to the following observations: in-situ temperature and salinity over 1–525 m depth and in-situ horizontal velocity over 8–36 m depth at (144.6°E, 32.3°N) south of the KE from the Kuroshio Extension Observatory (KEO) buoy (<https://www.pmel.noaa.gov/ocs/>;

See Fig. 11), SSH and SSH anomaly gridded datasets with horizontal resolution of 0.25° from Archiving, Validation and Interpretation of Satellite Oceanographic data (AVISO; Ducet et al., 2000), in-situ surface horizontal velocity from surface drifter buoys of the Global Drifter Program (Elipot et al., 2016), and Himawari-8 SSTs. We note that the AVISO and Himawari-8 are not independent because the satellite SSH anomalies and SSTs are used in this system, respectively, whereas the KEO and surface drifter buoys are independent observations. The validation in the ocean interior in this study is limited due to the paucity of available independent observations.

In this study, we calculate the  $\Delta NBE$  (RMSDs) using daily outputs from the CTL and AOEI runs (the CTL, AOEI, and 1.5Terr runs). To compare the AOEI run (AOEI and 1.5Terr runs) with the CTL run, we also calculate improvement ratios  $IR$  for  $\Delta NBE$  (RMSD):

$$IR_{NBE} = \frac{(\Delta NBE)_{CTL} - (\Delta NBE)_{AOEI}}{(\Delta NBE)_{CTL}} \times 100 \text{ and} \quad (10)$$

$$IR_{RMSD} = \frac{(RMSD)_{CTL} - (RMSD)_{AOEI/1.5Terr}}{(RMSD)_{CTL}} \times 100. \quad (11)$$

Here, the subscripts  $CTL$ ,  $AOEI$ ,  $1.5Terr$  indicate the CTL, AOEI, and 1.5Terr runs, respectively. Using the bootstrap method with 10,000 cycles, we detect significant improvement and degradation in the AOEI and 1.5Terr runs relative to the CTL run at the 99% confidence level.

### 3. Results

In subsection 3.1, the degradation of the low-salinity structure in the CTL run is described. Subsection 3.2 presents how the AOEI is applied to SST, SSS, and SSH fields. The detail of the improvement of the low-salinity structure by the AOEI is provided in subsection 3.3, and the results of the geostrophic balance and accuracy are described in subsection 3.4.

#### 3.1. Salinity degradation of the NPIW around the KE region in the CTL run

As shown in Fig. 1, the SST field in the CTL run agrees well with the satellite observations. Although the satellite-derived SSS has large errors, especially in coastal and high-latitude regions (Abe and Ebuchi, 2014), the SSS spatial pattern appears to be reproduced well in the analysis field (Fig. 2). However, the CTL run has noisier signals in the latter half of the experiment period, particularly in the SSS analysis fields. We also assess the monthly-mean temperature, salinity, and potential density  $\sigma_\theta$  along 150°E and 35°N sections across the KE. During the initial stages of the experiment period, the North Pacific Intermediate Water (NPIW), characterized by low minimum salinity, is distributed within  $\sigma_\theta = 26.5\text{--}27.25 \text{ kg m}^{-3}$  (Talley, 1993; Yasuda, 1997) (Fig. 3a, b). However, as the assimilation period progresses, the low salinity structure in the intermediate

Deleted: 15

Deleted: and

Deleted: is

Deleted: an

Deleted: dataset

Deleted: and

Deleted:

Deleted: and

Deleted: 7

Deleted: AOEI

Deleted: ,

Deleted: →

Deleted: 8

Deleted: where

Deleted: and

Deleted: and

Deleted: run

Deleted: ¶

Moved (insertion) [3]

Deleted: Here,  $\rho_0 = 1025 \text{ kg m}^{-3}$  is the reference density,  $c_p = 4190 \text{ J kg}^{-1} \text{ }^\circ\text{C}^{-1}$  is the specific heat of the seawater, and  $q_{sw}$  is downward shortwave radiation parameterized by ¶

$$q_{sw} = Q_{sw} \left\{ R \exp\left(-\frac{|z|}{\gamma_1}\right) + (1 - R) \exp\left(-\frac{|z|}{\gamma_2}\right) \right\} \rightarrow \rightarrow \rightarrow \rightarrow \rightarrow (17) ¶$$

(Paulson and Simpson, 1977), where  $Q_{sw}$  is shortwave radiation at the sea surface,  $R = 0.62$  is a separation constant, and  $\gamma_1 = 0.60 \text{ m}$  and  $\gamma_2 = 20.0 \text{ m}$  are attenuation length scales. These values are set to the case of Type IA from Jerlov (1976).

Deleted: mechanism

Deleted:

Deleted: that of

Deleted: is

Deleted: in

Deleted: T

Deleted: S

Deleted: (Figs. 3, 4)





salinity budget analysis indicates that this salinity improvement results from the reduction of the vertical salinity diffusion in the AOEI run relative to the CTL run (See the detail in Appendix B). Figure 8 shows the vertical profile of the vertical diffusivity  $\kappa_z$  averaged over the KE region (140°–160°E, 30°–40°N) for the whole experiment period and the maximum of the averaged diffusivity over the whole experiment period within 300–1000 m depth. As is consistent with the results of the salinity budget analysis, there is exceedingly large vertical diffusivity at 300–800 m depth around the KE region, which results in salinity degradation induced by strong vertical diffusion in the CTL run. In contrast, the low salinity water in the intermediate layer persists in the AOEI run because the vertical diffusivity is smaller.

Weak density stratification and strong vertical shear are favorable conditions for the generation of large vertical diffusivity (Davis et al., 2016; Pacanowski and Philander, 1981). To gain dynamical insight into the vertical diffusivity difference between the AOEI and CTL runs, the temporal tendency of the vertical diffusivity  $\kappa_z$ , the squared buoyancy frequency  $N^2 = -g/\sigma_\theta \partial\sigma_\theta/\partial z$ , and the squared vertical shear  $\mathbf{u}_z^2 = |\partial\mathbf{u}/\partial z|^2 (\partial\kappa_z/\partial t, \partial N^2/\partial t, \text{ and } \partial\mathbf{u}_z^2/\partial t, \text{ respectively})$  are summed during the positive vertical diffusivity tendency ( $\partial\kappa_z/\partial t > 0$ ), and are then averaged in the KE region (140°–160°E, 30°–40°N). The buoyancy frequency can be represented as the sum of the contributions from the temperature and salinity vertical gradients ( $N_T^2$  and  $N_S^2$ , respectively):

$$N^2 = -\frac{g}{\sigma_\theta} \frac{\partial\sigma_\theta}{\partial z} = g \left( \alpha_T \frac{\partial T}{\partial z} - \beta_S \frac{\partial S}{\partial z} \right) \equiv N_T^2 + N_S^2. \quad (12)$$

Here,  $\alpha_T$  ( $\beta_S$ ) is the thermal (salinity) expansion coefficient.

Figure 9a, b shows that the total vertical diffusivity tendency is smaller in the AOEI run than the CTL run, which agrees qualitatively with the diffusivity averaged over the whole period (Fig. 8a, b). As is clear from Fig. 9c, d, the total shear tendency is almost zero in both the CTL and AOEI runs. The total buoyancy frequency tendency makes substantial contributions in the CTL and AOEI runs, and its amplitude is smaller in the AOEI run than the CTL run. Since the negative values indicate weakening of the density stratification, the density stratification is less weakened in the AOEI run than the CTL run. The difference in the total buoyancy frequency tendency between the AOEI and CTL runs is caused by the differences of both the total  $\partial N_T^2/\partial t$  and  $\partial N_S^2/\partial t$  (Fig. 9e). Although  $\partial N_T^2/\partial t$  ( $\partial N_S^2/\partial t$ ) can be decomposed into the temporal tendency terms of the vertical temperature (salinity) gradient and the thermal (salinity) expansion coefficient, we confirmed that the latter terms have almost no impact on  $\partial N_T^2/\partial t$  and  $\partial N_S^2/\partial t$  (figure not shown). Therefore, the differences in the temperature and salinity vertical gradient tendencies result in less weakening of the density stratification in the AOEI run than the CTL run.

To investigate the causes of the differences in the temperature and salinity vertical gradient tendencies between the AOEI and CTL runs, we derive the temperature and salinity stratification tendency equations, respectively, by taking the vertical derivatives of the temperature and salinity budget equations [Eqs. (4) and (5)] in the ocean interior;

$$\frac{\partial}{\partial t} \left( \frac{\partial T}{\partial z} \right) = \frac{\partial}{\partial z} \{ \nabla \cdot (\boldsymbol{\kappa} \circ \nabla T) \} - \frac{\partial}{\partial z} (\mathbf{v} \cdot \nabla T) + \frac{1}{\rho_0 c_p} \frac{\partial q_{sw}}{\partial z} + \frac{\partial}{\partial z} (\text{increment}) \text{ and} \quad (13)$$

Deleted: 12

Moved down [4]:  $\alpha_T$  ( $\beta_S$ ) is the thermal (salinity) expansion coefficient.

Deleted: The density vertical gradient can be decomposed as

$$\frac{\partial\sigma_\theta}{\partial z} = -\sigma_\theta \alpha_T \frac{\partial T}{\partial z} + \sigma_\theta \beta_S \frac{\partial S}{\partial z}, \rightarrow \rightarrow \rightarrow \rightarrow \rightarrow \rightarrow \rightarrow \rightarrow (11)$$

where  $\alpha_T$  ( $\beta_S$ ) is the thermal (salinity) expansion coefficient. Consequently,

Deleted: t

Deleted: 12

Moved (insertion) [4]

Deleted: 13

Formatted: Indent: First line: 1.27 cm

Deleted: 12

Deleted: 13

Deleted: 13

Deleted: We note

Deleted: that

Deleted: :

Deleted: ¶

$$\frac{\partial N_T^2}{\partial t} = g \alpha_T \frac{\partial}{\partial z} \frac{\partial T}{\partial z} + g \frac{\partial T}{\partial z} \frac{\partial \alpha_T}{\partial z} \text{ and} \rightarrow \rightarrow \rightarrow \rightarrow \rightarrow \rightarrow \rightarrow \rightarrow (13)$$

$$\frac{\partial N_S^2}{\partial t} = -g \beta_S \frac{\partial}{\partial z} \frac{\partial S}{\partial z} - g \frac{\partial S}{\partial z} \frac{\partial \beta_S}{\partial z} \rightarrow \rightarrow \rightarrow \rightarrow \rightarrow \rightarrow \rightarrow \rightarrow (14)$$

W

Deleted: e

Deleted: to obtain the temperature and salinity stratification tendency equations

Deleted: 15

$$425 \quad \frac{\partial}{\partial t} \left( \frac{\partial S}{\partial z} \right) = \frac{\partial}{\partial z} \{ \nabla \cdot (\kappa \circ \nabla S) \} - \frac{\partial}{\partial z} (v \cdot \nabla S) + \frac{\partial}{\partial z} (\text{increment}). \quad (14)$$

Deleted: .

Deleted: 16

As in the total vertical diffusivity tendency calculated the above, the terms in Eqs. (13) and (14) are summed when  $\partial \kappa_z / \partial t > 0$ , and then averaged in the KE region (140°–160°E, 30°–40°N) (Fig. 10). We note that positive values in Eqs. (13) and (14) indicate opposite effects on the density stratification: a positive temperature (salinity) vertical gradient tendency strengthens (weakens) the density stratification.

**Moved up [3]:** Here,  $\rho_0 = 1025 \text{ kg m}^{-3}$  is the reference density,  $c_p = 4190 \text{ J kg}^{-1} \text{ }^\circ\text{C}^{-1}$  is the specific heat of the seawater, and  $q_{sw}$  is downward shortwave radiation parameterized by

$$q_{sw} = Q_{sw} \left\{ R \exp\left(-\frac{|z|}{\gamma_1}\right) + (1-R) \exp\left(-\frac{|z|}{\gamma_2}\right) \right\} \quad (17)$$

(Paulson and Simpson, 1977), where  $Q_{sw}$  is shortwave radiation at the sea surface,  $R = 0.62$  is a separation constant, and  $\gamma_1 = 0.60 \text{ m}$  and  $\gamma_2 = 20.0 \text{ m}$  are attenuation length scales. These values are set to the case of Type IA from Jerlov (1976).

Deleted: respectively.

**Deleted:** Here,  $\rho_0 = 1025 \text{ kg m}^{-3}$  is the reference density,  $c_p = 4190 \text{ J kg}^{-1} \text{ }^\circ\text{C}^{-1}$  is the specific heat of the seawater, and  $q_{sw}$  is downward shortwave radiation parameterized by

$$q_{sw} = Q_{sw} \left\{ R \exp\left(-\frac{|z|}{\gamma_1}\right) + (1-R) \exp\left(-\frac{|z|}{\gamma_2}\right) \right\} \quad (17)$$

(Paulson and Simpson, 1977), where  $Q_{sw}$  is shortwave radiation at the sea surface,  $R = 0.62$  is a separation constant, and  $\gamma_1 = 0.60 \text{ m}$  and  $\gamma_2 = 20.0 \text{ m}$  are attenuation length scales. These values are set to the case of Type IA from Jerlov (1976).

Deleted: A

Deleted: 15

Deleted: 16

Deleted: 14

Deleted: 15

Deleted: 16

Deleted: 15

Deleted: 14

Deleted: (Fig. 14a, b)

Deleted: 14

Deleted: 15

Deleted: 15

Deleted: 15

Deleted: 16

Deleted: 14

Deleted: 14

Deleted: 14

Deleted: 16

Deleted: 14

Deleted: Fig. 5

Deleted: a

In the CTL and AOEI runs, the temperature gradient tendency term [the left-hand side (LHS) term of Eq. (13)] is negative and indicates that the temperature and density stratification are weakened at all depths (Fig. 10a, b). Compared with the CTL run, the amplitude of this term is smaller, and thus the temperature and density stratification is less weakened in the AOEI run. As shown in Fig. 10c, the difference in the temperature gradient tendency between the AOEI and CTL runs is due mainly to the analysis increment gradient term [the last term on the right-hand side (RHS) of Eq. (13)] and in part to the advection gradient term [the second term on the RHS of Eq. (13)], whereas the diffusion and shortwave penetration gradient terms [the first and third terms on the RHS of Eq. (13), respectively] make almost no contribution.

In the CTL run, the salinity gradient tendency term [the LHS term of Eq. (14)] indicates that the salinity (density) stratification is strengthened (weakened) at all depths (Fig. 10d). In the AOEI run, the salinity (density) stratification is weakened (strengthened) at 200–400 m depth and slightly strengthened (weakened) at 400–1000 m depth (Fig. 10e). The salinity gradient tendency term is smaller in the AOEI run than the CTL run at all depths, and thus the salinity (density) stratification is less strengthened (weakened) in the AOEI run relative to the CTL run (Fig. 10f). The difference in the analysis increment gradient terms [the last term on the RHS of Eq. (14)] between the AOEI and CTL runs dominates that in the salinity gradient tendency term, whereas the differences between the diffusion and advection gradient terms [the second and third terms on the RHS of Eq. (14), respectively] have little influence. This indicates that less strengthening (weakening) of the salinity (density) stratification in the AOEI run relative to the CTL run is due to the smaller analysis increment in the AOEI run. The impacts of the SST, SSS, and SSH assimilation are limited to between the surface and about 370 m depth because of the prescribed vertical localization scale of 100 m described in subsection 2.2, and consequently only in-situ temperature and salinity assimilation generates analysis increments in the intermediate layer.

The AOEI contributes to maintaining the density stratification by reducing the temperature and salinity increments and preventing the occurrence of large vertical diffusivity that degrades low salinity water in the intermediate layer around the KE region. In the CTL run, the salinity increments restore the degraded low salinity water (See Appendix A) but lead to degradation through the formation of large vertical diffusivity at the same time. Thus, it seems that positive feedback exists that may degrade the salinity structure.

### 3.4 Improvement of geostrophic balance and accuracy by the AOEI

In this subsection, we investigate the impacts of the AOEI on the geostrophic balance and accuracy. Figure 11 shows  $\Delta NBE$  averaged over the whole period in the CTL and AOEI runs. In the CTL run,  $\Delta NBE$  is large in the mid-latitude regions, especially along the KE (Fig. 11a). In the AOEI run,  $\Delta NBE$  is smaller than the CTL run for the entire domain (Fig. 11b). The spatiotemporal averaged  $\Delta NBE$  over the whole experiment period and domain is  $0.57 \times 10^{-10} \text{ s}^{-2}$  and  $0.35 \times 10^{-10} \text{ s}^{-2}$  for the CTL and AOEI runs, respectively, and the balance is significantly improved in the AOEI run relative to the CTL run. This is probably because the analysis increments are smaller in the AOEI run than the CTL run.

To investigate the accuracy in the ocean interior, we calculate the RMSDs of CTL, AOEI, and 1.5Terr runs relative to in-situ temperature, salinity, and horizontal velocity observations from the KEO buoy south of the KE (Figs. 11a, 12). Results are only presented for the temperature and salinity because no significant results are obtained for the horizontal velocity. The RMSDs for both temperature and salinity are smaller in the AOEI run than the CTL run, and the AOEI run provides significant temperature (salinity) improvements at 0–150 m (50–400 m) depth relative to the CTL run. This is probably because the AOEI suppresses the development of the strong vertical diffusion that leads to the salinity degradation and because of the improvement in the balance. We have confirmed that the AOEI run also has smaller temperature (salinity) RMSDs than the 1.5Terr run throughout the depth except for two observation points at 225 and 275 m (150 and 525 m) depth. Therefore, among the experiments, the AOEI run is the best for the accuracy of temperature and salinity south of the KE.

We also investigate the accuracy of the SSH, surface flow, and SST fields, respectively, calculating the spatiotemporally averaged RMSDs relative to the SSH and SSHA datasets from the AVISO, to in-situ surface horizontal velocity observations from the drifter buoys, and to Himawari-8 SSTs (Fig. 13). Although the AOEI run slightly degrades the SST accuracy relative to the CTL run (Fig. 13e), the RMSDs in the AOEI run are smaller for all other variables, and indicate significant improvements with the exception of surface meridional velocity (Fig. 13a–d). The improvement of the surface flow field in the AOEI run would result from the better balance and accuracy of the density structure in the ocean interior. Relative to the 1.5Terr run, the AOEI run improves the accuracy for all variables.

Kurihara et al. (2016), for example, show that the RMSDs of the Himawari-8 SSTs relative to the buoys are about  $0.5^\circ\text{C}$  and are larger in the higher latitude regions with larger zenith angle, and observation error variances have substantial contributions to the RMSDs. However, the ensemble spreads are much smaller than the RMSDs for all variables (Fig. 13). We have found that the ensemble spreads in the subtropical region appear to be underdispersive even if the perturbed atmospheric and lateral boundary conditions are applied (cf. Figs. 5e, 6e). Methods to inflate the ensemble spread more would be required for further improvements of the accuracy, but this will be a future issue.

#### 4. Summary

We have implemented the AOEI with the sbPOM-LETKF ocean data assimilation system and conducted sensitivity experiments to investigate the impacts on the low-salinity NPIW around the KE region, geostrophic balance, and accuracy in the analysis field. In the CTL run, the large analysis increments by in-situ temperature and salinity assimilation weaken the density stratification. The resulting exceedingly large vertical diffusivity induces the strong vertical diffusion that breaks the

Deleted: 15

Deleted: 15

Deleted: 15

Deleted: 15

Deleted: 16

Deleted: , and to

Deleted:

Commented [SO4]: Major comment #2

Deleted: 17

Deleted: T

Deleted: in the AOEI run,

Deleted: relative to the CTL run,

Deleted:

Deleted: The SSHA RMSDs averaged over the experiment period also show that the AOEI run may represent an improvement relative to the CTL run in some high RMSD regions; for example, along the Kuroshio (130–140°E, 30°–35°N) and downstream of the KE (160°–165°E, 30°N) (Fig. 17a, b). Here, we note that these improvements are not necessarily caused by the larger ensemble spread in the AOEI run than the CTL run, as the AOEI decreases the analysis increments by inflating the observation errors. Instead, the

Commented [SO5]: Major comment #1

Commented [SO6]: Minor comment #7

Commented [SO7]: Major comment #3

Deleted: in the AOEI run would result in an improvement in the surface flow field. We have confirmed that the AOEI run is more accurate with the experiment with larger temperature observation error of  $1.5^\circ\text{C}$  than the CTL run (not shown).

Formatted: Highlight

low salinity structure in the NPIW around the KE region. The salinity increment contributes to restoring the low salinity water, but at the same time causes the salinity degradation by generating strong vertical diffusion. Therefore, the positive feedback appears to occur, degrading the salinity structure.

The AOEI decreases the temperature and salinity increments around the KE region by adaptively inflating the temperature and salinity observation errors, respectively. As a result, the AOEI mitigates the salinity degradation seen in the CTL run, and therefore, the low salinity water is maintained in the AOEI run. In addition, the AOEI significantly improves the geostrophic balance probably because of the reduction of the analysis increments. Moreover, the AOEI prevents the development of strong vertical diffusion and improves the accuracy of temperature and salinity in the ocean interior. Furthermore, the improvements of the balance and density structure in the ocean interior contribute to more accurate SSH and surface flow fields. In summary, this study demonstrates the positive impacts of the AOEI on the balance and accuracy of the temperature, salinity, and surface flow fields.

Since our available computational resources were limited, we fixed the tuning parameter of the RTPP, perturbed atmospheric forcing, ensemble size, localization scale, and prescribed observation errors. Further experiments to explore more optimal settings are required, and this will be investigated in the future. Since low salinity water is distributed in the intermediate layer in western boundary current regions in all ocean basins, we would expect that this study will be helpful for improving existing EnKF-based ocean data assimilation systems. Minamide and Zhang (2017) noted that the AOEI has the advantage of being easily implemented with various EnKF-based systems, and this study serves as a good example for the usefulness of the AOEI. We are currently constructing high-resolution reanalysis datasets in the western North Pacific and Maritime Continent regions based on this system, and plan to develop (near) real-time ensemble forecast systems.

Commented [S08]: Minor comment #3

Deleted: d

## Appendices

### Appendix A: Cause of salinity degradation in the CTL run

To investigate the cause of the salinity degradation in the CTL run, we calculate the salinity budget equation [Eq. (5)] in the intermediate layer around the KE region ( $140^{\circ}$ – $160^{\circ}$ E,  $30^{\circ}$ – $40^{\circ}$ N, 500–1000 m depth; Fig. 3g, h). Figure A1a indicates that the salinity tendency term [the LHS term of Eq. (5)] is positive and corresponds to the salinity increase shown in left and middle panels of Fig. 3. The positive salinity tendency term is caused mainly by the diffusion term [the first term on the RHS of Eq. (5)] and partly by the advection term [the second term on the RHS of Eq. (5)]. The diffusion term is dominated only by the vertical diffusion, and the horizontal diffusion makes almost no contribution (Fig. A1b). The advection term consists of different components in different months during the experiment period (Fig. A1c): meridional advection in July, zonal and meridional advection in August, and zonal and vertical advection in September–December 2015. In contrast, the analysis increment term [the last term on the RHS of Eq. (5)] has only a minor impact but plays a role in restoring the low salinity water. Therefore, the vertical diffusion is the main cause of the salinity degradation in the intermediate layer around the KE region.

## Appendix B: Cause of salinity improvement in the AOEI run

To investigate the cause of the salinity improvement in the AOEI run relative to the CTL run, we calculate the salinity budget equation [Eq. (5)] difference between the AOEI and CTL runs (Fig. B1):

$$\Delta \left( \frac{\partial S}{\partial t} \right) = \Delta \{ \nabla \cdot (\kappa \circ \nabla S) \} - \Delta \{ v \cdot \nabla S \} + \Delta(\text{increment}), \quad (\text{B1})$$

where  $\Delta$  indicates the AOEI run minus the CTL run. The salinity tendency difference term [the LHS term of Eq. (B1)] indicates that the salinity structure is maintained in the AOEI run by suppressing the salinity increase throughout the experiment period (Fig. B1a). The diffusion and advection difference terms [the first and second terms on the RHS of Eq. (B1), respectively] contribute almost equally to the salinity tendency difference term. The diffusion difference term is dominated by only the vertical diffusion difference, whereas the advection difference term is dominated by different components in different months: by the meridional advection difference in July; by all advection differences in August–September; and by vertical and partly zonal advection differences in October–December 2015. The reduction in the vertical diffusion is therefore the main cause of the improvement for low salinity water in the AOEI run relative to the CTL run.

### Code and data availability

The source code for sbPOM version 1.0 and LETKF are available from <https://zenodo.org/record/6482744> (last access: 27 June 2022, Jordi and Wang, 2012; Miyoshi and Yamane, 2007; Ohishi et al. in review). The source code for COARE version 3.5 (Brodeau et al., 2017; Edson et al., 2013) was downloaded from <https://github.com/brodeau/aerobulk> (last access: 13 April 2021).

We thank Dr. Kenshi Hibino for providing us with an earlier version of TE-Global, before the official release of the latest version (<https://www.eorc.jaxa.jp/water/>, last access: 13 April 2021). The observation datasets are: the surface drifter buoy data ([https://www.aoml.noaa.gov/phod/gdp/hourly\\_data.php](https://www.aoml.noaa.gov/phod/gdp/hourly_data.php), last access: 13 April 2021, Elipot et al., 2016); the KEO buoy data (<https://www.pmel.noaa.gov/ocs/>, last access: 13 April 2021); ETOPO1 (<https://www.ngdc.noaa.gov/mgg/global/>, last access: 13 April 2021, Amante and Eakins, 2009); WOA18 (<https://www.ncei.noaa.gov/access/world-ocean-atlas-2018/>, last access: 13 April 2021, Locarnini et al., 2019; Zweng et al., 2019); the satellite SSTs from Himawari-8 (<https://www.eorc.jaxa.jp/ptree/index.html>, last access: 13 April 2021; Bessho et al., 2016; Kurihara et al., 2016) and GCOM-W (<https://gportal.jaxa.jp/gpr/?lang=en>, last access: 13 April 2021); the satellite-derived SSS from SMOS ([http://www.esa.int/Applications/Observing\\_the\\_Earth/SMOS](http://www.esa.int/Applications/Observing_the_Earth/SMOS), last access: 13 April 2021) and SMAP version 4.3 (<https://podaac.jpl.nasa.gov/>, last access: 13 April 2021, Meissner et al., 2018); the satellite-derived SSHA and AVISO (Ducet et al., 2000) from CMEMS (<https://marine.copernicus.eu/>, last access: 13 April 2021); and in-situ temperatures and salinity from GTSP (<https://www.ncei.noaa.gov/products/global-temperature-and-salinity-profile-programme>, last access: 13 April 2021, Sun et al., 2010) and AQC Argo version 1.2a

**Deleted:** To investigate the cause of this salinity improvement, we estimate the salinity budget equation [Eq. (9)] difference between the AOEI and CTL runs (Fig. 11):

$$\Delta \left( \frac{\partial S}{\partial t} \right) = \Delta \{ \nabla \cdot (\kappa \circ \nabla S) \} - \Delta \{ v \cdot \nabla S \} + \Delta(\text{increment}), \quad (\text{10})$$

where  $\Delta$  indicates the AOEI run minus the CTL run. The salinity tendency difference term [the LHS term of Eq. (10)] indicates that the salinity structure is maintained in the AOEI run by suppressing the salinity increase throughout the experiment period (Fig. 11a). The diffusion and advection difference terms [the first and second terms on the RHS of Eq. (10), respectively] contribute almost equally to the salinity tendency difference term. The diffusion difference term is dominated by only the vertical diffusion difference, whereas the advection difference term is dominated by different components in different months: by the meridional advection difference in July; by all advection differences in August–September; and by vertical and partly zonal advection differences in October–December 2015. The reduction in the vertical diffusion is therefore the main cause of the improvement for low salinity water in the AOEI run relative to the CTL run.

650 ([http://www.jamstec.go.jp/ARGO/argo\\_web/argo/?page\\_id=100&lang=en](http://www.jamstec.go.jp/ARGO/argo_web/argo/?page_id=100&lang=en), last access: 13 April 2021). The global JRA55  
atmosphere and SODA 3.7.2 ocean reanalysis datasets are from <http://search.diasjp.net/en/dataset/JRA55> (last access: 13 April  
2021, Kobayashi et al., 2015) and [https://www.soda.umd.edu/soda3\\_readme.htm](https://www.soda.umd.edu/soda3_readme.htm) (last access: 13 April 2021, Carton et al.,  
2018), respectively.

#### Author contribution

655 SO developed the code for the ocean data assimilation system, conducted the sensitivity experiments, and analyzed the outputs.  
SO and TM prepared the paper with contributions from MK.

#### Competing interests

The authors declare that they have no conflict of interest.

660

#### Acknowledgements

665 *We thank two anonymous reviewers for constructive comments.* Numerous comments from Drs. Nariaki Hirose, Takahiro  
Toyoda, Yosuke Fujii, and Norihisa Usui at the Meteorological Research Institute, Yoichi Ishikawa at JAMSTEC, Katsumi  
Takayama at IDEA Consultants, Inc., Naoki Hirose at Kyushu University, and participants in the ocean data assimilation  
summer school helped us to develop the presented system. This work used computational resources of the JAXA  
Supercomputer System Generation 2 and 3 (JSS2 and JSS3, respectively) and the supercomputer Fugaku provided by RIKEN  
through the HPCI Research Project (Project ID: hp210166, [hp220167](https://www.hpci.go.jp/portal/entry/20210401), ra000007).

#### Financial support

670 This work was supported by JST AIP Grant Number JPMJCR19U2, Japan; MEXT (JPMXP1020200305) as "Program for  
Promoting Researches on the Supercomputer Fugaku" (Large Ensemble Atmospheric and Environmental Prediction for  
Disaster Prevention and Mitigation); the COE research grant in computational science from Hyogo Prefecture and Kobe City  
through Foundation for Computational Science; JST, SICORP Grant Number JPMJSC1804, Japan; JSPS KAKENHI Grant  
Number JP19H05605; the JAXA; JST, CREST Grant Number JPMJCR20F2, Japan; Cabinet Office, Government of Japan,  
675 Cross-ministerial Moonshot Agriculture, Forestry and Fisheries Research and Development Program, "Technologies for Smart  
Bio-industry and Agriculture"(funding agency: Bio-oriented Technology Research Advancement Institution) No. JPJ009237;  
RIKEN Pioneering Project "Prediction for Science"; RIKEN Engineering Network Project.

#### References

680 Abe, H. and Ebuchi, N.: Evaluation of sea-surface salinity observed by Aquarius, J. Geophys. Res. Ocean., 119, 8109–8121,  
doi:10.1002/2014JC010094, 2014.  
Amante, C. and Eakins, B. W.: ETOPO1 1 Arc-Minute Global Relief Model: Procedures, Data Sources and Analysis., 2009.

Deleted: ¶  
Review statement¶

- 685 Balmaseda, M. A., Hernandez, F., Storto, A., Palmer, M. D., Alves, O., Shi, L., Smith, G. C., Toyoda, T., Valdivieso, M., Barnier, B., Behringer, D., Boyer, T., Chang, Y. S., Chepurin, G. A., Ferry, N., Forget, G., Fujii, Y., Good, S., Guinehut, S., Haines, K., Ishikawa, Y., Keeley, S., Köhl, A., Lee, T., Martin, M. J., Masina, S., Masuda, S., Meyssignac, B., Mogensen, K., Parent, L., Peterson, K. A., Tang, Y. M., Yin, Y., Vernieres, G., Wang, X., Waters, J., Wedd, R., Wang, O., Xue, Y., Chevallier, M., Lemieux, J. F., Dupont, F., Kuragano, T., Kamachi, M., Awaji, T., Caltabiano, A., Wilmer-Becker, K. and
- 690 Gaillard, F.: The ocean reanalyses intercomparison project (ORA-IP), *J. Oper. Oceanogr.*, 8(0), s80–s97, doi:10.1080/1755876X.2015.1022329, 2015.
- Bessho, K., Date, K., Hayashi, M., Ikeda, A., Imai, T., Inoue, H., Kumagai, Y., Miyakawa, T., Murata, H., Ohno, T., Okuyama, A., Oyama, R., Sasaki, Y., Shimazu, Y., Shimoji, K., Sumida, Y., Suzuki, M., Taniguchi, H., Tsuchiyama, H., Uesawa, D., Yokota, H. and Yoshida, R.: An introduction to Himawari-8/9 – Japan’s new-generation geostationary
- 695 meteorological satellites, *J. Meteorol. Soc. Japan*, 94(2), 151–183, doi:10.2151/jmsj.2016-009, 2016.
- Bloom, S. C., Takacs, L. L., da Silva, A. M. and Ledvina, D.: Data assimilation using incremental analysis updates, *Mon. Weather Rev.*, 124(6), 1256–1271, doi:10.1175/1520-0493(1996)124<1256:DAUIAU>2.0.CO;2, 1996.
- Brodeau, L., Barnier, B., Gulev, S. K. and Woods, C.: Climatologically significant effects of some approximations in the bulk parameterizations of turbulent air–sea fluxes, *J. Phys. Oceanogr.*, 47(1), 5–28, doi:10.1175/JPO-D-16-0169.1, 2017.
- 700 Carton, J. A., Chepurin, G. A. and Chen, L.: SODA3: A new ocean climate reanalysis, *J. Clim.*, 31(17), 6967–6983, doi:10.1175/JCLI-D-17-0149.1, 2018.
- Chakraborty, A., Sharma, R., Kumar, R. and Basu, S.: Joint assimilation of Aquarius-derived sea surface salinity and AVHRR-derived sea surface temperature in an ocean general circulation model using SEEK filter: Implication for mixed layer depth and barrier layer thickness, *J. Geophys. Res. Ocean.*, 120(10), 6927–6942, doi:10.1002/2015JC010934, 2015.
- 705 Davis, P. E. D., Lique, C., Johnson, H. L. and Guthrie, J. D.: Competing effects of elevated vertical mixing and increased freshwater input on the stratification and sea ice cover in a changing Arctic Ocean, *J. Phys. Oceanogr.*, 46(5), 1531–1553, doi:10.1175/JPO-D-15-0174.1, 2016.
- Desroziers, G., Berre, L., Chapnik, B. and Poli, P.: Diagnosis of observation, background and analysis-error statistics in observation space, *Q. J. R. Meteorol. Soc.*, 131(613), 3385–3396, doi:10.1256/qj.05.108, 2005.
- 710 Ducet, N., Le Traon, P. Y. and Reverdin, G.: Global high-resolution mapping of ocean circulation from TOPEX/Poseidon and ERS-1 and -2, *J. Geophys. Res.*, 105(C8), 19477–19498, doi:10.1029/2000JC900063, 2000.
- Edson, J. B., Jampana, V., Weller, R. A., Bigorre, S. P., Plueddemann, A. J., Fairall, C. W., Miller, S. D., Mahrt, L., Vickers, D. and Hersbach, H.: On the exchange of momentum over the open ocean, *J. Phys. Oceanogr.*, 43(8), 1589–1610, doi:10.1175/JPO-D-12-0173.1, 2013.
- 715 Elipot, S., Lumpkin, R., Perez, R. C., Lilly, J. M., Early, J. J. and Sykulski, A. M.: A global surface drifter data set at hourly resolution, *J. Geophys. Res. Ocean.*, 121(5), 2937–2966, doi:10.1002/2016JC011716, 2016.
- Evensen, G.: Sequential data assimilation with a nonlinear quasi-geostrophic model using Monte Carlo methods to forecast error statistics, *J. Geophys. Res.*, 99, 10143–10162, 1994.


- Evensen, G.: The Ensemble Kalman Filter: Theoretical formulation and practical implementation, *Ocean Dyn.*, 53, 343–367, doi:10.1007/s10236-003-0036-9, 2003.
- Hackert, E., Ballabrera-Poy, J., Busalacchi, A. J., Zhang, R. H. and Murtugudde, R.: Impact of sea surface salinity assimilation on coupled forecasts in the tropical Pacific, *J. Geophys. Res. Ocean.*, 116, C05009, doi:10.1029/2010JC006708, 2011.
- Hunt, B. R., Kostelich, E. J. and Szunyogh, I.: Efficient data assimilation for spatiotemporal chaos: A local ensemble transform Kalman filter, *Phys. D*, 230, 112–126, doi:10.1016/j.physd.2006.11.008, 2007.
- Isoguchi, O., Kawamura, H. and Oka, E.: Quasi-stationary jets transporting surface warm waters across the transition zone between the subtropical and the subarctic gyres in the North Pacific, *J. Geophys. Res. Ocean.*, 111(10), C10003, doi:10.1029/2005JC003402, 2006.
- Jerlov, N. G.: *Marine Optics.*, 1976.
- Jordi, A. and Wang, D. P.: sbPOM: A parallel implementation of Princeton Ocean Model, *Environ. Model. Softw.*, 38, 59–61, doi:10.1016/j.envsoft.2012.05.013, 2012.
- Kida, S., Mitsudera, H., Aoki, S., Guo, X., Ito, S., Kobashi, F., Komori, N., Kubokawa, A., Miyama, T., Morie, R., Nakamura, H., Nakamura, T., Nakano, H., Nishigaki, H., Nonaka, M., Sasaki, H., Sasaki, Y. N., Suga, T., Sugimoto, S., Taguchi, B., Takaya, K., Tozuka, T., Tsujino, H. and Usui, N.: Oceanic fronts and jets around Japan: a review, *J. Oceanogr.*, 71, 469–497, doi:10.1007/s10872-015-0283-7, 2015.
- Kobayashi, S., Ota, Y., Harada, Y., Ebata, A., Moriya, M., Onoda, H., Onogi, K., Kamahori, H., Kobayashi, C., Endo, H., Miyaoka, K. and Takahashi, K.: The JRA-55 reanalysis: General specifications and basic characteristics, *J. Meteorol. Soc. Japan*, 93(1), 5–48, doi:10.2151/jmsj.2015-001, 2015.
- Kotsuki, S., Ota, Y. and Miyoshi, T.: Adaptive covariance relaxation methods for ensemble data assimilation: experiments in the real atmosphere, *Q. J. R. Meteorol. Soc.*, 143(705), 2001–2015, doi:10.1002/qj.3060, 2017.
- Kurihara, Y., Murakami, H. and Kachi, M.: Sea surface temperature from the new Japanese geostationary meteorological Himawari-8 satellite, *Geophys. Res. Lett.*, 43(3), 1234–1240, doi:10.1002/2015GL067159, 2016.
- Locarnini, R. A., Mishonov, A. V., Baranova, O. K., Boyer, T. P., Zweng, M. M., Garcia, H. E., Reagan, J. R., Seidov, D., Weathers, K. W., Paver, C. R. and Smolyar, I. V.: *World Ocean Atlas 2018, Volume 1: Temperature*. A. Mishonov, Technical Editor., NOAA Atlas NESDIS, 81(July), 52, 2019.
- Martin, M. J., Balmaseda, M., Bertino, L., Brasseur, P., Brassington, G., Cummings, J., Fujii, Y., Lea, D. J., Lellouche, J. M., Mogensen, K., Oke, P. R., Smith, G. C., Testut, C. E., Waagbø, G. A., Waters, J. and Weaver, A. T.: Status and future of data assimilation in operational oceanography, *J. Oper. Oceanogr.*, 8(0), s28–s48, doi:10.1080/1755876X.2015.1022055, 2015.
- Meissner, T., Wentz, F. J. and Vine, D. M. Le: The salinity retrieval algorithms for the NASA Aquarius version 5 and SMAP version 3 releases, *Remote Sens.*, 10(7), 1121, doi:10.3390/rs10071121, 2018.




- Mellor, G. L., Ezer, T. and Oey, L.-Y.: The pressure gradient conundrum of sigma coordinate ocean models, *J. Atmos. Ocean. Technol.*, 11(4), 1126–1134, doi:10.1175/1520-0426(1994)011<1126:TPGCOS>2.0.CO;2, 1994.
- Minamide, M. and Zhang, F.: Adaptive observation error inflation for assimilating all-sky satellite radiance, *Mon. Weather Rev.*, 145(3), 1063–1081, doi:10.1175/MWR-D-16-0257.1, 2017.
- Miyazawa, Y., Zhang, R., Guo, X., Tamura, H., Ambe, D., Lee, J. S., Okuno, A., Yoshinari, H., Setou, T. and Komatsu, K.: Water mass variability in the western North Pacific detected in a 15-year eddy resolving ocean reanalysis, *J. Oceanogr.*, 65(6), 737–756, doi:10.1007/s10872-009-0063-3, 2009.
- Miyazawa, Y., Miyama, T., Varlamov, S. M., Guo, X. and Waseda, T.: Open and coastal seas interactions south of Japan represented by an ensemble Kalman filter, *Ocean Dyn.*, 62(4), 645–659, doi:10.1007/s10236-011-0516-2, 2012.
- Miyazawa, Y., Varlamov, S. M., Miyama, T., Guo, X., Hihara, T., Kiyomatsu, K., Kachi, M., Kurihara, Y. and Murakami, H.: Assimilation of high-resolution sea surface temperature data into an operational nowcast/forecast system around Japan using a multi-scale three-dimensional variational scheme, *Ocean Dyn.*, 67(6), 713–728, doi:10.1007/s10236-017-1056-1, 2017.
- Miyoshi, T. and Yamane, S.: Local ensemble transform Kalman filtering with an AGCM at a T159/L48 resolution, *Mon. Weather Rev.*, 135(11), 3841–3861, doi:10.1175/2007MWR1873.1, 2007.
- Miyoshi, T., Yamane, S. and Enomoto, T.: Localizing the error covariance by physical distances within a local ensemble transform Kalman filter (LETKF), *SOLA*, 3, 89–92, doi:10.2151/sola.2007-023, 2007.
- [Ohishi, S., Hihara, T., Aiki, H., Ishizaka, J., Miyazawa, Y., Kachi, M., and Miyoshi, T.: An ensemble Kalman filter system with the Stony Brook Parallel Ocean Model v1.0, \*Geosci. Model Dev. Discuss.\* \[preprint\], doi: 10.5194/gmd-2022-40, in review, 2022.](#)
- Pacanowski, R. C. and Philander, S. G. H.: Parameterization of vertical mixing in numerical models of tropical oceans, *J. Phys. Oceanogr.*, 11, 1443–1451, doi:10.1175/1520-0485(1981)011<1443:POVMIN>2.0.CO;2, 1981.
- Paulson, C. A. and Simpson, J. J.: Irradiance measurements in the upper ocean, *J. Phys. Oceanogr.*, 7, 952–956, doi:10.1175/1520-0485(1977)007<0952:IMITUO>2.0.CO;2, 1977.
- Penny, S. G., Kalnay, E., Carton, J. A., Hunt, B. R., Ide, K., Miyoshi, T. and Chepurin, G. A.: The local ensemble transform Kalman filter and the running-in-place algorithm applied to a global ocean general circulation model, *Nonlinear Process. Geophys.*, 20(6), 1031–1046, doi:10.5194/npg-20-1031-2013, 2013.
- Sakov, P., Counillon, F., Bertino, L., Lisæter, K. A., Oke, P. R. and Korabely, A.: TOPAZ4: an ocean-sea ice data assimilation system for the North Atlantic and Arctic, *Ocean Sci.*, 8(4), 633–656, doi:10.5194/os-8-633-2012, 2012.
- Shibuya, R., Sato, K., Tomikawa, Y., Tsutsumi, M. and Sato, T.: A study of multiple tropopause structures caused by inertia-gravity waves in the antarctic, *J. Atmos. Sci.*, 72(5), 2109–2130, doi:10.1175/JAS-D-14-0228.1, 2015.
- Sun, C., Thresher, A., Keeley, R., Hall, N., Hamilton, M., Chinn, P., A. Tran, Goni, G., Villeon, L. P. de la, Carval, T., Cowen, L., Manzella, G., Gopalakrishna, V., Guerrero, R., Reseghetti, F., Kanno, Y., Klein, B., Rickard, L., Baldoni, A., Lin, S., Ji, F. and Nagaya, Y.: The data management system for the global temperature and salinity profile programme, in

Formatted: English (US)

- Proceedings of OceanObs'09: Sustained Ocean Observations and Information for Society, pp. 931–938, European Space Agency., 2010.
- Talley, L. D.: Distribution and formation of North Pacific intermediate water, *J. Phys. Oceanogr.*, 23(3), 517–537, doi:10.1175/1520-0485(1993)023<0517:DAFONP>2.0.CO;2, 1993.
- 790 Toyoda, T., Fujii, Y., Kuragano, T., Matthews, J. P., Abe, H., Ebuchi, N., Usui, N., Ogawa, K. and Kamachi, M.: Improvements to a global ocean data assimilation system through the incorporation of Aquarius surface salinity data, *Q. J. R. Meteorol. Soc.*, 141(692), 2750–2759, doi:10.1002/qj.2561, 2015.
- Usui, N., Tsujino, H., Fujii, Y. and Kamachi, M.: Short-range prediction experiments of the Kuroshio path variabilities south of Japan, *Ocean Dyn.*, 56(5–6), 607–623, doi:10.1007/s10236-006-0084-z, 2006.
- 795 Xu, F. H. and Oey, L. Y.: State analysis using the Local Ensemble Transform Kalman Filter (LETKF) and the three-layer circulation structure of the Luzon Strait and the South China Sea, *Ocean Dyn.*, 64(6), 905–923, doi:10.1007/s10236-014-0720-y, 2014.
- Yasuda, I.: The origin of the North Pacific intermediate water, *J. Geophys. Res. Ocean.*, 102(C1), 893–909, doi:10.1029/96JC02938, 1997.
- 800 Yin, Y., Alves, O. and Oke, P. R.: An ensemble ocean data assimilation system for seasonal prediction, *Mon. Weather Rev.*, 139(3), 786–808, doi:10.1175/2010MWR3419.1, 2011.
- Zhang, F., Davis, C. A., Kaplan, M. L. and Koch, S. E.: Wavelet analysis and the governing dynamics of a large-amplitude mesoscale gravity-wave event along the east coast of the United States, *Q. J. R. Meteorol. Soc.*, 127(577), 2209–2245, doi:10.1002/qj.49712757702, 2001.
- 805 Zhang, F., Snyder, C. and Sun, J.: Impacts of initial estimate and observation availability on convective-scale data assimilation with an ensemble Kalman filter, *Mon. Weather Rev.*, 132(5), 1238–1253, doi:10.1175/1520-0493(2004)132<1238:IOIEAO>2.0.CO;2, 2004.
- Zhang, F., Minamide, M. and Clothiaux, E. E.: Potential impacts of assimilating all-sky infrared satellite radiances from GOES-R on convection-permitting analysis and prediction of tropical cyclones, *Geophys. Res. Lett.*, 43(6), 2954–2963, doi:10.1002/2016GL068468, 2016.
- 810 Zuo, H., Balmaseda, M. A., Tietsche, S., Mogensen, K. and Mayer, M.: The ECMWF operational ensemble reanalysis–analysis system for ocean and sea ice: a description of the system and assessment, *Ocean Sci.*, 15(3), 779–808, doi:10.5194/os-15-779-2019, 2019.
- Zweng, M. M., Reagan, J. R., Seidov, D., Boyer, T. P., Antonov, J. I., Locarnini, R. A., Garcia, H. E., Mishonov, A. V., 815 Baranova, O. K., Weathers, K. W., Paver, C. R. and Smolyar, I. V.: World Ocean Atlas 2018, Volume 2: Salinity. A. Mishonov, Technical Editor, NOAA Atlas NESDIS, 82(July), 50, 2019.



820 Figure 1: Monthly-mean SST for (a) July, (c) October, and (e) December 2015 in the CTL run. (b), (d), and (f) Same as in (a), (c), and (e), but for assimilated satellite SSTs. Thin (thick) black contour intervals are 2 (10) °C. White lines in (a) indicate 150°E (35°N) used for the zonal (meridional) sections in Fig. 3.



Deleted:

Deleted: (4)





Figure 2: Same as Fig.1 but for SSS. Thin (thick) black contour intervals are 0.25 (2). In (b), (d), and (f), contour intervals are not shown because the satellite observations are noisy.



Deleted:





Figure 3: The zonal section of monthly-mean salinity (color) and potential density (contour) along 35°N, in (a) July, (d) October, and (g) December 2015 in the CTL run. (b), (e), and (h) Same as (a), (d), and (g), but for the meridional section along 150°E, (c), (f), (i) Same as (b), (e), (h) but for AOEI run. Thin (thick) contour intervals are 0.25 (2) kg m<sup>-3</sup>. The white box in (g) [(h) and (i)], encloses a longitude (latitude)-depth section of 140°–160°E (30°–40°N) and 500–1000 m depth.



835



Deleted:

Deleted: meridional... section of monthly-mean salinity/temperature... (color) and potential density (contour) along 35°N150°E... in (a) July, (dc... October, and (ge... 2015 in the CTL run. (b), (ed..., and (hf... Same as (a), (dc..., and (ge..., but for the meridional section along 150°E salin' (... [1]

Deleted: (f)



Moved down [1]:






Figure 4: Monthly-mean ratio of the area where the AOEI is applied to the system domain in the (a) SST, (b) SSS, and (c) SSH fields in the AOEI run (black bars). Red (blue) lines indicate when the innovation is positive (negative).




**Figure 5:** (a) Ratio of the period when the AOEI is applied to the SST compared with the whole experiment period in the AOEI run. (b) and (c) Same as (a) but for when the innovation is positive and negative, respectively. (d) and (e) innovation amplitude and ensemble spread averaged during the period when the AOEI is applied, respectively. **White** contours indicate SST averaged over the whole period. Thin (thick) contour intervals are 2 (10) °C.




**Deleted:**

**Deleted:** 7

**Deleted:** Black



910 Figure 6: Same as Fig. 5 but for SSS. Thin (thick) contour intervals are 0.25 (1).



Deleted:

Deleted: 8

Deleted: 7



915






Figure 7: (a) SST and (b) SSS forecast biases (color) and averages (contour) over the whole experiment period. Thin (thick) contour intervals are 2 (10) °C in (a) and 0.25 (2) in (b).




Deleted:  
Deleted: 9



Moved down [2]:

Deleted: 1

... [4]



955 **Figure 8:** Vertical diffusivity (black), and squared buoyancy frequency (red) and shear (blue) averaged over the KE region (140°–160°E, 30°–40°N) for the whole experiment period in the (a) CTL and (b) AOEI runs. Maxima of the averaged vertical diffusivity for the whole experiment period within 300–1000 m depth in the (c) CTL and (d) AOEI runs. Black contours show SSH averaged over the whole period. Thin (thick) contour intervals are 0.2 (1) m.

960


Deleted:

Deleted: 12

Deleted: lines


Deleted: lines

Deleted: lines




**Figure 9:** Total vertical diffusivity tendency during the positive vertical diffusivity tendency period averaged over the KE region (140°–160°E, 30°–40°N) in the (a) CTL and (b) AOEI runs. (c) and (d) Same as (a) but for the squared buoyancy frequency (black) and shear (gray) tendency. (e) Same as (c) and (d), but for the AOEI minus CTL run. In (c)–(e), cyan (orange) lines indicate contributions from  $\partial N_T^2 / \partial t$  ( $\partial N_S^2 / \partial t$ ).

970



Deleted:

Deleted: 13



975 Figure 10: (a)–(c) Same as Fig. 9 but for each term of the temperature stratification budget equation [Eq. (13)]:  
 Temperature gradient tendency term (the LHS term; black), temperature diffusion gradient term (the first term on  
 the RHS; red), temperature advection gradient term (the second term on the RHS; blue), shortwave penetration  
 gradient term (the third term on the RHS; orange), and temperature increment gradient term (the last term on the  
 RHS; cyan). The shortwave penetration gradient term is almost zero and overlaps with the temperature diffusion  
 gradient term. (d)–(f) Same as (a)–(c) but for the salinity stratification budget equation [Eq. (14)]. We note that the  
 shortwave penetration gradient term is almost zero and is behind the diffusion gradient term in (a)–(c).

Deleted: 14

Deleted: 13

Deleted: 15

Deleted: 16

Commented [S09]: Minor comment #6







Figure 11:  $\Delta NBE$  (color) and SSH (contour) averaged over the whole experiment period for the (a) CTL and (b) AOEI runs. Thin (thick) contour intervals are 0.2 (1) m. Spatiotemporally averaged  $\Delta NBE$  over the whole period and domain is shown in the lower right corners. The black star in (a) denotes the position of the KEO buoy.

990





995 **Figure 12:** (a) Temperature and (b) salinity RMSDs averaged over the whole experiment period at the KEO buoy in the CTL (black), AOEI (orange), and 1.5Terr (cyan) runs. Open circles indicate significant improvement in the AOEI and 1.5Terr runs relative to the CTL run.



Deleted:

Deleted: 16

Deleted: and

Deleted: red






Figure 13: RMSDs for (a) SSH and (b) SSHA relative to the AVISO surface (c) zonal and (d) meridional velocity relative to the drifter buoys, and (e) SSTs relative to the Himawari-8 over the whole domain and period. Black dots indicate the ensemble spread in the observation space. We note that the ranges of the vertical axis are different between the RMSDs and ensemble spreads.

Deleted: 17

Deleted: and for

Commented [SO10]: Major Comments #1, #2

Commented [SO11]: Minor comment #7




**Figure A1:** (a) Monthly mean for each term in the salinity budget equation [Eq. (5)] averaged over the KE region in the intermediate layer (140°–160°E, 30°–40°N, and 500–1000 m depth) in the CTL run: the salinity tendency term (the LHS term; black bars), salinity diffusion term (the first term on the RHS; red line), salinity advection term (the second term on the RHS, blue line), and salinity increment term (the last term on the RHS; gray line). (b) and (c) Same as (a), but for salinity diffusion and advection terms (black bars), and zonal (orange lines), meridional (cyan lines), and vertical (green lines) components.

Moved (insertion) [1]

Deleted: 5

Deleted: 9





020 **Figure B1:** (a) Monthly mean for each term for the difference between the AOEI and CTL runs in the salinity budget equation [Eq. (B1)]: Salinity tendency difference term (the LHS term; black bars), salinity diffusion difference term (the first term on the RHS; red line), salinity advection difference term (the second term on the RHS; blue line), and salinity increment difference term (the last term on the RHS; gray line). (b) and (c) Same as (a) but for the salinity diffusion and advection difference terms (black bars), respectively, and zonal (orange lines), meridional (cyan lines), and vertical (green lines) components.

025

Moved (insertion) [2]

Deleted: 11

Deleted: 10

Deleted: 9




Figure 18: RMSDs for SSHA relative to the AVISO averaged over the whole experiment period in the (a) CTL and (b) AOEI runs. (c) and (d) Same as (a) and (b) but for the SSHA ensemble spread.

**Page 21: [1] Deleted      Shun Ohishi      8/26/22 6:31:00 PM**

**Page 21: [1] Deleted      Shun Ohishi      8/26/22 6:31:00 PM**

**Page 21: [1] Deleted      Shun Ohishi      8/26/22 6:31:00 PM**

**Page 21: [1] Deleted      Shun Ohishi      8/26/22 6:31:00 PM**

**Page 21: [1] Deleted      Shun Ohishi      8/26/22 6:31:00 PM**

**Page 21: [1] Deleted      Shun Ohishi      8/26/22 6:31:00 PM**

**Page 21: [1] Deleted      Shun Ohishi      8/26/22 6:31:00 PM**

**Page 21: [1] Deleted      Shun Ohishi      8/26/22 6:31:00 PM**

**Page 21: [1] Deleted      Shun Ohishi      8/26/22 6:31:00 PM**

**Page 21: [1] Deleted      Shun Ohishi      8/26/22 6:31:00 PM**

**Page 22: [2] Deleted      Shun Ohishi      8/26/22 6:35:00 PM**

**Page 22: [3] Deleted      Shun Ohishi      8/26/22 6:52:00 PM**

**Page 22: [3] Deleted      Shun Ohishi      8/26/22 6:52:00 PM**

**Page 25: [4] Deleted      Shun Ohishi      8/29/22 5:44:00 PM**