

# Impacts of Ice-Particle Size Distribution Shape Parameter on Climate Simulations with the Community Atmosphere Model Version 6 (CAM6)

Wentao Zhang<sup>1</sup>, Xiangjun Shi<sup>1,\*</sup>, and Chunsong Lu<sup>2</sup>

<sup>1</sup> School of Atmospheric Sciences, Nanjing University of Information Science and Technology, Nanjing 210044, China.

<sup>2</sup> Key Laboratory for Aerosol-Cloud-Precipitation of China Meteorological Administration, Nanjing University of Information Science and Technology, Nanjing 210044, China.

*Correspondence to:* Xiangjun Shi (shixj@nuist.edu.cn)

**Abstract.** The impacts of the ice crystal size distribution shape parameter ( $\mu_i$ ) were considered in the two-moment bulk cloud microphysics scheme of the Community Atmosphere Model Version 6 (CAM6). The  $\mu_i$ 's impact on the statistical mean radii of ice crystals can be analysed based on their calculating formulas. Under the same mass ( $q_i$ ) and number ( $N_i$ ), the ratios of the mass-weighted radius ( $R_{qi}$ , not related to  $\mu_i$ ) to other statistical mean radii (e.g., effective radiative radius) are completely determined by  $\mu_i$ . Off-line tests show that  $\mu_i$  has a significant impact on the cloud microphysical processes owing to the  $\mu_i$ -induced changes in ice crystal size distribution and statistical mean radii (excluding  $R_{qi}$ ). Climate simulations show that increasing  $\mu_i$  would lead to higher  $q_i$  and lower  $N_i$  in most regions, and these impacts can be explained by the changes in cloud microphysical processes. After increasing  $\mu_i$  from 0 to 5, the longwave cloud radiative effect increases (stronger warming effect) by  $5.58 \text{ W m}^{-2}$  (25.11%), and the convective precipitation rate decreases by  $-0.12 \text{ mm day}^{-1}$  (7.64%). In short, the impacts of  $\mu_i$  on climate simulations are significant and the main influence mechanisms are also clear. This suggests that the  $\mu_i$ -related processes deserve to be parameterized in a more realistic manner.

## 1 Introduction

Clouds are an integral part of the Earth's radiation budget and global water cycle (Liou, 1986; Luo and Rossow, 2004; Bony et al., 2015; Zhou et al., 2016). Since cloud microphysical processes occur at scales that are much smaller than the resolution of commonly-used atmospheric models, it remains a significant challenge for atmospheric models to represent cloud-related processes, especially ice-phase cloud microphysical processes (Mitchell et al., 2008; Spichtinger and Gierens, 2009; Wang and Penner, 2010; Erfani and Mitchell, 2016; Paukert et al., 2019; Morrison et al., 2020; Proske et al., 2021). Because it is impossible for commonly-used atmospheric models (excluding the ideal model with the recently developed Lagrangian particle-based scheme) to individually describe cloud particles (e.g., cloud droplets or ice crystals), only the macrostatistical features of cloud particles are represented in cloud microphysics schemes. From the outset, the development of cloud microphysics schemes has resulted in two distinct categories: bulk microphysics parameterization and spectral (bin)

microphysics (Milbrandt and Yau, 2005; Khain et al., 2015). The spectral (bin) approach explicitly represents the cloud 30 particle size distributions (PSDs) by using tens to hundreds of bins. The computational cost of this approach is very high because of the massive interactions among different bins. The bulk microphysics scheme represents the PSDs by a semiempirical distribution function. Compared to the spectral (bin) scheme, the bulk microphysics scheme has high computational efficiency and has been widely used in climate models (Morrison et al., 2005; Lohmann et al., 2007; Salzmann et al., 2010; Gettelman and Morrison, 2015).

35 In the bulk cloud microphysics schemes used for climate models, the PSD is usually described by the gamma distribution function with three parameters, namely, the intercept parameter ( $N_0$ ), the slope parameter ( $\lambda$ ), and the spectral shape parameter ( $\mu$ ) (Khain et al., 2015; Morrison et al., 2020). Note that the commonly used two-moment bulk microphysics scheme predicts only the mass and number of cloud particles, which cannot constrain these three parameters (i.e.,  $N_0$ ,  $\lambda$ , and  $\mu$ ). Therefore, one of these three parameters (typically  $\mu$ ) must be determined from an empirical formula or set to a given 40 value (e.g., Morrison and Gettelman, 2008; Barahona et al., 2014; Eidhammer et al., 2017). For instance, the  $\mu$  ( $\mu_i$ ) of ice crystals (ICs, only represents cloud ice in this study) in the two-moment bulk stratiform cloud microphysics scheme developed by Morrison and Gettelman 2008 (hereafter MG scheme) is set to zero (i.e., the  $\mu_i$  is ignored). In recent years, off-line tests and short-term simulations (a few days or less) with high-resolution atmospheric models (e.g., cloud-resolving models and mesoscale models) have shown that  $\mu_i$  has a significant impact on cloud microphysical processes and synoptic 45 systems (Milbrandt and Yau, 2005; Milbrandt and McTaggart-Cowan, 2010; Loftus et al., 2014; Khain et al., 2015; Milbrandt et al., 2021). Unlike short-term simulations, climate simulations pay more attention to the equilibrium states or quasi-equilibrium states because the feedback processes become important (Sherwood et al., 2015; King et al., 2020). However, in terms of climate simulations, few studies have focused on the influence of  $\mu_i$ .

In this study, in order to investigate the impacts of  $\mu_i$  on climate simulations with the Community Atmosphere Model version 50 6 (CAM6) model, the impacts of  $\mu_i$  were considered in the MG scheme by a tunable parameter. There were two major motivations behind this work. First, are the impacts of  $\mu_i$  notable? If yes, it's necessary for climate models to represent the  $\mu_i$  and  $\mu_i$ -related processes in a more realistic manner. And second, what are the main mechanisms for these impacts? These would be helpful to understand the climate simulations with the impacts of  $\mu_i$ . This paper is organized as follows: the modified MG scheme and experimental setup are described in Section 2; cloud microphysical process off-line tests and 55 CAM6 model simulation results are analyzed in Section 3; and finally, the summary and conclusions are provided in Section 4.

## 2 Model and experiments

### 2.1 The modified MG scheme

The CAM6 model, which is the atmospheric component of the Community Earth System Model Version 2.1.3 (CAM6, 60 Bogenschutz et al., 2018; CESM2, Danabasoglu et al., 2020), was used in this study. It is noteworthy that the treatments of clouds in climate models are usually divided into two categories: convective cloud schemes with simplified cloud microphysics and larger-scale stratiform cloud schemes with relatively detailed cloud microphysics. In the CAM6 model, the convective cloud scheme does not consider the PSD of ICs (Zhang and McFarlane, 1995; Zhang et al., 1998; Bogenschutz et al., 2013; Larson, 2017). The stratiform cloud microphysics was represented by the updated MG scheme with prognostic 65 precipitation (Gettelman and Morrison, 2015). In both versions of the MG scheme, the ICs are assumed to be spherical, and the PSD of ICs is described by the gamma distribution function:

$$N'_i(D) = N_{0i} D^{\mu_i} e^{-\lambda_i D} \quad (1)$$

where  $N'_i(D)$  is the number density (i.e.,  $\delta N_i / \delta D$ ) of the ICs with diameter  $D$ .  $N_{0i}$ ,  $\lambda_i$ , and  $\mu_i$  (nonnegative values) are the intercept parameter, the slope parameter, and the spectral shape parameter, respectively. Given that  $\mu_i$  is known,  $N_{0i}$  and  $\lambda_i$  can be determined by the local in-cloud IC mass and number mixing ratio ( $q_i$  and  $N_i$ , prognostic variables in units of  $\text{kg kg}^{-1}$  70 and  $\text{kg}^{-1}$ , respectively).

$$\lambda_i = \left[ \frac{\pi \rho_i N_i \Gamma(4 + \mu_i)}{6 q_i \Gamma(1 + \mu_i)} \right]^{1/3} \quad (2)$$

$$N_{0i} = \frac{N_i \lambda_i^{(1+\mu_i)}}{\Gamma(1 + \mu_i)} \quad (3)$$

where the IC bulk density ( $\rho_i$ ) is  $500 \text{ kg m}^{-3}$  and  $\mu_i$  is zero in the default MG scheme.  $\Gamma(x) = \int_0^\infty t^{x-1} e^{-t} dt$  is the gamma function. It is noteworthy that the  $k$ th moment of this size distribution ( $M_k$ ) is found by integrating the distribution in this form:  $M_k = \int_0^\infty N_{0i} D^{\mu_i+k} e^{-\lambda_i D} dD = N_{0i} \Gamma(k + \mu_i + 1) / \lambda_i^{(k+\mu_i+1)}$  (Eidhammer et al., 2014). Furthermore, the recursive property of the gamma function (i.e.,  $\Gamma(x + 1) = x\Gamma(x)$ ) is also used for the following formula derivation.

75 Eqs. (2), (3) also indicate that, under the same  $q_i$  and  $N_i$ , changes in  $\mu_i$  could impact the other two parameters regarding the PSD of ICs (i.e.,  $N_{0i}$  and  $\lambda_i$ ). Meanwhile, the number-weighted radius ( $R_{ni}$ ) related to the IC deposition/sublimation process, the effective radiative radius ( $R_{ei}$ ) used for the radiative transfer scheme, and other statistical mean radii might be influenced. To better understand the influence of  $\mu_i$  on the ice-phase cloud microphysical processes, the equations for calculating the statistical mean radii are introduced first. The mass-weighted radius ( $R_{qi}$ ) is calculated from Eq. (4). The number-weighted 80 radius ( $R_{ni}$ ), which is the so-called mathematical mean value, is calculated from Eq. (5). The area-weighted radius ( $R_{ai}$ ) is calculated from Eq. (6).  $R_{ei}$ , which is defined as the cross-section weighted radius (Schumann et al., 2011; Wyser, 1998), is calculated from  $R_{qi}^3 / R_{ai}^2$  (Eq. 7). Note that  $R_{qi}$  can be calculated by  $q_i$  and  $N_i$  (the last term of Eq. 4, without  $\mu_i$ ), and the other

statistical mean radii (e.g.,  $R_{ni}$ ,  $R_{ai}$ , and  $R_{ei}$ ) can be calculated by  $R_{qi}$  and  $\mu_i$  (Eqs. 5-7). In other words, the ratios of the other statistical mean radii (e.g.,  $R_{ni}$ ,  $R_{ai}$ , and  $R_{ei}$ ) to  $R_{qi}$  are functions of  $\mu_i$ . For nonnegative  $\mu_i$  values, Eqs. (5), (6) indicate that  $R_{ni}$  and  $R_{ai}$  are always less than  $R_{qi}$ . This can be explained by the physical reason that larger ICs contribute more to  $R_{qi}$  than to  $R_{ni}$  and  $R_{ai}$ . Similarly,  $R_{ei}$  is always greater than  $R_{qi}$  (Eq. 7). Furthermore, Eqs. (5), (6), (7) also indicate that with increasing  $\mu_i$ ,  $R_{ni}$ ,  $R_{ai}$ , and  $R_{ei}$  approach  $R_{qi}$ . In Section 3.1, more analyses are provided by off-line tests.

$$R_{qi} = \frac{1}{2} \left[ \frac{\int_0^\infty D^3 N_i(D) dD}{\int_0^\infty N_i(D) dD} \right]^{1/3} = \frac{1}{2\lambda_i} \left[ \frac{\Gamma(\mu_i + 4)}{\Gamma(\mu_i + 1)} \right]^{1/3} = \left( \frac{3}{4\pi\rho_i} \frac{q_i}{N_i} \right)^{1/3} \quad (4)$$

$$R_{ni} = \frac{1}{2} \frac{\int_0^\infty D N_i(D) dD}{\int_0^\infty N_i(D) dD} = \frac{1}{2\lambda_i} \frac{\Gamma(\mu_i + 2)}{\Gamma(\mu_i + 1)} = R_{qi} \left[ \frac{\Gamma(1 + \mu_i)}{\Gamma(4 + \mu_i)} \right]^{\frac{1}{3}} \frac{\Gamma(\mu_i + 2)}{\Gamma(\mu_i + 1)} = R_{qi} \frac{(\mu_i + 1)}{[(\mu_i + 3)(\mu_i + 2)(\mu_i + 1)]^{\frac{1}{3}}} \quad (5)$$

$$R_{ai} = \frac{1}{2} \left[ \frac{\int_0^\infty D^2 N_i(D) dD}{\int_0^\infty N_i(D) dD} \right]^{1/2} = \frac{1}{2\lambda_i} \left[ \frac{\Gamma(\mu_i + 3)}{\Gamma(\mu_i + 1)} \right]^{1/2} = R_{qi} \left[ \frac{\Gamma(1 + \mu_i)}{\Gamma(4 + \mu_i)} \right]^{1/3} \left[ \frac{\Gamma(\mu_i + 3)}{\Gamma(\mu_i + 1)} \right]^{1/2} \\ = R_{qi} \frac{[(\mu_i + 2)(\mu_i + 1)]^{\frac{1}{2}}}{[(\mu_i + 3)(\mu_i + 2)(\mu_i + 1)]^{\frac{1}{3}}} \quad (6)$$

$$R_{ei} = \frac{1}{2} \frac{\int_0^\infty D^3 N_i(D) dD}{\int_0^\infty D^2 N_i(D) dD} = \frac{(R_{qi})^3}{(R_{ai})^2} = \frac{1}{2\lambda_i} \frac{\Gamma(\mu_i + 4)}{\Gamma(\mu_i + 3)} = R_{qi} \left[ \frac{\Gamma(1 + \mu_i)}{\Gamma(4 + \mu_i)} \right]^{1/3} \frac{\Gamma(\mu_i + 4)}{\Gamma(\mu_i + 3)} \\ = R_{qi} \frac{(\mu_i + 3)}{[(\mu_i + 3)(\mu_i + 2)(\mu_i + 1)]^{\frac{1}{3}}} \quad (7)$$

Because  $\mu_i$  is zero in the default MG scheme, the equations for the cloud microphysical processes are simplified by omitting  $\mu_i$  (Morrison and Gettelman, 2008; Gettelman and Morrison, 2015). In this study, these equations are modified to consider the impact of  $\mu_i$  (i.e., nonzero  $\mu_i$ ). In the default MG scheme, there are three cloud microphysical processes, which are related to the PSD of ICs. They consist of the deposition/sublimation of ICs, the autoconversion of IC to snow, and the mass-weighted and number-weighted IC fall velocities ( $V_{qi}$  and  $V_{ni}$ ), respectively. Table 1 shows the original and modified equations for these cloud microphysical processes. The  $dq_i/dt$  (i.e., the time derivative of  $q_i$ ) caused by the deposition/sublimation process (including the Wegener-Bergeron process in mixed-phase clouds) is calculated from  $dq_i/dt = S_i/(T_p\tau_i)$ , where  $S_i$ ,  $T_p$ , and  $\tau_i$  are the ice supersaturation, a psychrometric correction to account for the release of latent heat, and the supersaturation relaxation time scale, respectively (Morrison and Gettelman, 2008). Among them,  $\tau_i$  is related to  $\mu_i$ . In the original equation of  $\tau_i$  (Table 1, left column),  $N_{0i} = \lambda_i N_i$  (Eq. 3) and  $\lambda_i^{-1} = 2R_{ni}$  at  $\mu_i = 0$  (Eq. 5). Therefore, the original equation for  $\tau_i$  can be rewritten as the modified equation (Table 1, right column). The modified equation indicates that  $\tau_i$  is inversely proportional to  $N_i R_{ni}$ , which is consistent with the equation obtained by Korolev et al. (2003). This modified equation also indicates that, under the same  $q_i$  and  $N_i$  ( $R_{qi}$  is also fixed),  $\mu_i$  can affect  $\tau_i$  (i.e., the IC deposition/sublimation process) via the influence on  $R_{ni}$ . In the MG scheme, ICs with radii greater than the threshold ( $R_{cs}$ ) are considered to be snow.

Correspondingly, the mass and number of ICs converted to snow ( $q_{i\text{auto}}$  and  $N_{i\text{auto}}$ ) are represented by the integration of those ICs with radii greater than  $R_{cs}$ . Therefore, the incomplete gamma function, ( $\Gamma(s, x) = \int_x^\infty t^{s-1}e^{-t} dt$ ), is used to calculate  $q_{i\text{auto}}$  and  $N_{i\text{auto}}$  (right column). It is necessary to note that, at  $\mu_i = 0$ , the modified equations for  $q_{i\text{auto}}$  and  $N_{i\text{auto}}$  can be 105 rewritten as the original equations (i.e., omitting  $\mu_i$ , left column) based on a property of the incomplete gamma function (i.e.,  $\Gamma(s, x) = (s-1)! e^{-x} \sum_{k=0}^{s-1} \frac{x^k}{k!}$ , where  $s$  is a positive integer). Based on the diameter-fall speed relationship,  $V = aD^b$  ( $a$  and  $b$  are empirical coefficients), and the properties of the gamma function,  $\mu_i$  is considered in the equations for mass-weighted and number-weighted terminal fall speeds ( $V_{qi}$  and  $V_{ni}$ , Table 1).

**Table 1.** Equations for calculating the  $\mu_i$ -related cloud microphysical processes\*.

	Original ( $\mu_i = 0$ )	Modification (nonzero $\mu_i$ )
$\tau_i$	$\frac{1}{2\pi\rho_a D_v N_{0i} \lambda_i^{-2}}$	$\frac{1}{2\pi\rho_a D_v N_i 2R_{ni}}$
$N_{i\text{auto}}$	$\frac{N_{0i}}{\lambda_i} e^{-\lambda_i 2R_{cs}}$	$N_{0i} \frac{\Gamma(1 + \mu_i, \lambda_i 2R_{cs})}{\lambda_i^{1+\mu_i}}$
$q_{i\text{auto}}$	$\frac{\pi\rho_i N_{0i}}{6} \left[ \frac{(2R_{cs})^3}{\lambda_i} + \frac{3(2R_{cs})^2}{\lambda_i^2} + \frac{6(2R_{cs})}{\lambda_i^3} + \frac{6}{\lambda_i^4} \right] e^{-\lambda_i 2R_{cs}}$	$\frac{\pi\rho_i N_{0i}}{6} \frac{\Gamma(4 + \mu_i, \lambda_i 2R_{cs})}{\lambda_i^{4+\mu_i}}$
$V_{ni}$	$\frac{\left(\frac{\rho_{a850}}{\rho_a}\right)^{0.35} a \Gamma(1 + b)}{\lambda_i^b \Gamma(1)}$	$\frac{\left(\frac{\rho_{a850}}{\rho_a}\right)^{0.35} a \Gamma(1 + b + \mu_i)}{\lambda_i^b \Gamma(1 + \mu_i)}$
$V_{qi}$	$\frac{\left(\frac{\rho_{a850}}{\rho_a}\right)^{0.35} a \Gamma(4 + b)}{\lambda_i^b \Gamma(4)}$	$\frac{\left(\frac{\rho_{a850}}{\rho_a}\right)^{0.35} a \Gamma(4 + b + \mu_i)}{\lambda_i^b \Gamma(4 + \mu_i)}$

110 \*Where  $D_v$  is the diffusivity of water vapor in air ( $D_v$  is calculated as a function of temperature and pressure,  $D_v = 8.794 \times 10^{-5} \times T^{1.81}/P$ );  $R_{cs}$  is the threshold radius for the autoconversion of IC to snow ( $R_{cs} = 100 \mu\text{m}$ );  $\rho_a$  is the air density;  $\rho_{a850}$  is the reference air density at 850 hPa, and  $a$  and  $b$  are empirical coefficients ( $a = 700 \text{ m}^{1-b} \text{ s}^{-1}$ ,  $b = 1$ ).

## 2.2 CAM6 experimental design

Observational studies have shown that  $\mu_i$  is less than 5 under most conditions (Heymsfield, 2003; McFarquhar et al., 2015). 115 This study focuses only on investigating the influence of  $\mu_i$ . There are four  $\mu_i$ -related processes (i.e., the radiative transfer process and three cloud microphysical processes) in the modified CAM6 model. Note that  $\mu_i$  can be set to different values for different processes with the advantage of model simulations. Seven experiments were conducted in this study (Table 2). The Mu0 experiment is considered to be the reference experiment because  $\mu_i$  is set to zero for all of the  $\mu_i$ -related processes. The  $\mu_i$  is set to 2 for all of the  $\mu_i$ -related processes in the Mu2 experiment, and the  $\mu_i$  is set to 5 for all of the  $\mu_i$ -related processes in

120 the Mu5 experiment. The comparison between the Mu2 (or Mu5) and Mu0 experiments shows the influence of  $\mu_i$  on climate simulations. Furthermore, to investigate the influence of each  $\mu_i$ -related process, an additional four experiments, namely, Tao5, Auto5, Fall5, and Rei5, were conducted. In this study, for ease of expression, “ $\Delta$ ” is used to denote the difference from the Mu0 experiment (e.g.,  $\Delta\text{Tao5} = \text{Tao5} - \text{Mu0}$ ). Without specification, the comparisons between model simulations are relative to the Mu0 experiment. When analyzing a cloud property variable (e.g.,  $q_i$ ), it is necessary to know which 125 experiment the variable comes from. To show this information, the experiment name is added as a superscript. For example, the  $q_i$  from the Mu5 experiment is denoted as  $q_i^{\text{Mu5}}$ , the difference in  $q_i$  between the Mu5 and Mu0 experiments is denoted as  $q_i^{\Delta\text{Mu5}}$ , and the relative change of  $q_i$  from the Mu5 experiment is denoted as  $q_i^{\Delta\text{Mu5/Mu0}}$ .

In this study, all experiments were atmosphere-only simulations (i.e., sea surface temperature and sea ice are prescribed) with a horizontal resolution of  $1.9^\circ$  latitude  $\times 2.5^\circ$  longitude and 32 vertical layers. All experiments run for 11 model years 130 and the last 10 years were used for the analyses. In addition, the standard deviation calculated from the averages of each year (i.e., 10 averages) was used to check the statistical significance of the multiyear average (i.e., 10-year average).

**Table 2.** The values of  $\mu_i$  in all experiments conducted in this study.  $\mu_{i\_tao}$ ,  $\mu_{i\_auto}$ ,  $\mu_{i\_fall}$ , and  $\mu_{i\_rei}$  indicate the  $\mu_i$  used for calculating the IC deposition/sublimation (*tao*), autoconversion of IC to snow (*auto*), IC fall velocity (*fall*), and the  $R_{ei}$  used for the radiation scheme (*rei*), respectively.

Names	$\mu_{i\_tao}$	$\mu_{i\_auto}$	$\mu_{i\_fall}$	$\mu_{i\_rei}$
Mu0	0	0	0	0
Mu2	2	2	2	2
Mu5	5	5	5	5
Tao5	5	0	0	0
Auto5	0	5	0	0
Fall5	0	0	5	0
Rei5	0	0	0	5


### 135 3 Results and analysis

#### 3.1 Off-line tests

To better understand the impact of  $\mu_i$  on climate simulations, the impacts of  $\mu_i$  on the IC PSD and  $\mu_i$ -related cloud microphysical processes are first illustrated by off-line tests. In these off-line tests, the impact of  $\mu_i$  was analyzed at a given  $R_{qi}$  (i.e., the ratio of  $q_i$  to  $N_i$  is fixed).

140 Eqs. (1), (2), (3) indicate that the normalized IC size distribution (i.e., the relative contributions of each bin) can be calculated from  $R_{qi}$  and  $\mu_i$ . Fig. 1 shows the impact of  $\mu_i$  on the normalized PSDs of ICs. Under the same  $\mu_i$ , the shapes of the PSDs (i.e., the relative number or mass contributions of each bin) with  $R_{qi} = 20 \mu\text{m}$  (small IC scenario) are the same as those

with  $R_{qi} = 60 \mu\text{m}$  (large IC scenario). In other words, the shape of the PSD is completely determined by  $\mu_i$  (i.e., spectral shape parameter). As expected, the PSDs move toward larger radii with increasing  $R_{qi}$ . As introduced in the study of Milbrandt et al. (2021), the PSD becomes narrow with increasing  $\mu_i$ . Note that, in terms of number, the contributions of the smaller size bins significantly decrease with increasing  $\mu_i$ . Unlike the number contributions, the mass contributions of the larger size bins significantly decrease with increasing  $\mu_i$  because the mass contribution is more sensitive to the IC radius. Under the large IC scenario (i.e.,  $R_{qi} = 60 \mu\text{m}$ ), the mass contribution of the ICs with radii greater than  $R_{cs}$  is significantly decreased with increasing  $\mu_i$ . The above analyses suggest that the cloud microphysical processes that depend on the PSD of ICs (e.g., autoconversion of IC to snow) might be significantly influenced by  $\mu_i$ .



**Figure 1.** The relative number (upper panel) and mass (lower panel) contributions from each radius bin of ICs. Each bin width is the same based on the logarithm of the particle radius.  $N_i$  and  $q_i$  are the total number and mass of ICs, respectively. A total of 100 bins were used here. The solid lines indicate the normal IC scenario (i.e.,  $R_{qi} = 20 \mu\text{m}$ ), and the dotted lines indicate the large IC scenario (i.e.,  $R_{qi} = 60 \mu\text{m}$ ). The vertical black line indicates the  $R_{cs}$  that was used for the autoconversion of IC to snow ( $R_{cs} = 100 \mu\text{m}$ ).

The off-line tests were performed for the  $\mu_i$ -related cloud microphysical processes and statistical mean radii (Table 3). As introduced in Section 2.1,  $R_{ni}$ ,  $R_{ai}$ , and  $R_{ei}$  can be calculated from  $R_{qi}$  and  $\mu_i$ . Both  $R_{ni}$  and  $R_{ai}$  significantly increase with increasing  $\mu_i$  (Table 3). This is in agreement with their calculation equations (Eqs. 5, 6).  $R_{ni}$  is approximately half of  $R_{qi}$  at  $\mu_i = 0$  (i.e., 11.00/20 and 33.02/60), while  $R_{ni}$  is close to  $R_{qi}$  at  $\mu_i = 5$  (i.e., 17.26/20 and 51.78/60). According to the calculation equation of  $R_{ei}$  (Eq. 7),  $R_{ei}$  decreases with increasing  $\mu_i$ . The ratios of  $R_{ei}$  to  $R_{qi}$  at  $\mu_i = 0$ , 2, and 5 are 1.65 (i.e., 33.02/20 and 99.06/60), 1.28 (i.e., 25.54/20 and 76.63/60), and 1.15 (i.e., 23.01/20 and 69.04/60), respectively. It is necessary to point out that with increasing  $\mu_i$ , both  $R_{ni}$ ,  $R_{ai}$ , and  $R_{ei}$  approach  $R_{qi}$  (Table 3) because the PSD of ICs becomes narrow (Fig.1). As expected,  $\tau_i$  decreases with increasing  $\mu_i$  (Table 3) because  $\tau_i$  is inversely proportional to  $R_{ni}$  (Table 2). The decrease in  $\tau_i$

suggests that the  $dq_i/dt$  caused by the deposition/sublimation process is accelerated (Morrison and Gettelman, 2008).

165 Compared to the  $1/\tau_i$  with  $\mu_i = 0$  (i.e.,  $3.35 \times 10^{-4} \text{ s}^{-1}$  and  $10.04 \times 10^{-4} \text{ s}^{-1}$ ), the  $1/\tau_i$  with  $\mu_i = 2$  ( $4.66 \times 10^{-4} \text{ s}^{-1}$  and  $13.98 \times 10^{-4} \text{ s}^{-1}$ ) and  $\mu_i = 5$  ( $5.25 \times 10^{-4} \text{ s}^{-1}$  and  $15.74 \times 10^{-4} \text{ s}^{-1}$ ) increase by 39.10% and 56.72%, respectively. This is consistent with the previous finding (ICs vapor deposition process is obviously accelerated by increasing  $\mu_i$ ) reported by Mitchell (1991). In Table 3,  $N_{i\text{auto}}/N_i$  and  $q_{i\text{auto}}/q_i$  indicate the portion of ICs that convert to snow in terms of number and mass, respectively. Under the small IC scenario (i.e.,  $R_{qi} = 20 \mu\text{m}$ ), regardless of the value of  $\mu_i$ , both  $N_{i\text{auto}}/N_i$  and  $q_{i\text{auto}}/q_i$  are very small (< 2%, 170 Table 3) because there are few ICs with radii greater than  $R_{cs}$  (Fig. 1). Under the large IC scenario (i.e.,  $R_{qi} = 60 \mu\text{m}$ ), there is a considerable portion of ICs with radii greater than  $R_{cs}$ , especially the mass contribution (Fig. 1). The  $q_{i\text{auto}}/q_i$  at  $\mu_i = 0, 2$ , and 5 are 64.08%, 36.54%, and 18.40%, respectively (Table 3). This suggests that the autoconversion of IC to snow becomes difficult with increasing  $\mu_i$ . Compared with the considerable values for  $q_{i\text{auto}}/q_i$ , the  $N_{i\text{auto}}/N_i$  is relatively small (i.e., 4.84% at  $\mu_i = 0$ , 4.23% at  $\mu_i = 2$ , and 2.63% at  $\mu_i = 5$ ). Therefore, the  $R_{qi}$  of the residual ICs ( $R_{qi\_afterauto}$ ;  $43.36 \mu\text{m}$  at  $\mu_i = 0$ ,  $52.31 \mu\text{m}$  at 175  $\mu_i = 2$ , and  $56.57 \mu\text{m}$  at  $\mu_i = 5$ ) is obviously lower than the original  $R_{qi}$  ( $60 \mu\text{m}$ ). During the falling process, it is inevitable that  $V_{qi}$  is greater than  $V_{ni}$  because larger ICs with faster falling contribute more in the  $V_{qi}$ . Thus, larger ICs appear preferentially in the lower model layers. This is called the size-sorting mechanism (Milbrandt and Yau, 2005).  $V_{qi}$  decreases with increasing  $\mu_i$ , while  $V_{ni}$  increases with increasing  $\mu_i$  (Table 3). This could also be explained by their calculation equations (the corresponding derivations are similar to those for  $R_{ni}$ ,  $R_{ai}$ , and  $R_{ei}$ , not shown). With increasing  $\mu_i$ , the difference between  $V_{qi}$  180 and  $V_{ni}$  decreases (Table 3) because the PSD of ICs becomes narrow (Fig.1). As a result, the size-sorting process becomes slow. For instance, there are many ICs with  $R_{qi} = 60 \mu\text{m}$  in a model layer. The height of each model layer is 200 m. After one model time step (10 min), some ICs fall into the lower layer. For  $\mu_i = 0$ , the  $R_{qi}$  of the ICs that are still in the model layer ( $R_{qi\_leftover}$ ) is  $42.11 \mu\text{m}$ , and the  $R_{qi}$  of the ICs in the lower layer ( $R_{qi\_lowlayer}$ ) is  $95.24 \mu\text{m}$ . For  $\mu_i = 2$ ,  $R_{qi\_leftover}$  is  $52.45 \mu\text{m}$ , and  $R_{qi\_lowerlayer}$  is  $75.60 \mu\text{m}$ . For  $\mu_i = 5$ ,  $R_{qi\_leftover}$  is  $55.81 \mu\text{m}$ , and  $R_{qi\_lowerlayer}$  is  $68.68 \mu\text{m}$ . It is clear that the difference in  $R_{qi}$  185 between these two adjacent layers that is caused by the sedimentation process (i.e., the difference between  $R_{qi\_leftover}$  and  $R_{qi\_lowerlayer}$ ) becomes small with increasing  $\mu_i$ . In short, the above analyses clearly suggest that  $\mu_i$  has a significant impact on the cloud microphysical processes and statistical mean radii of ICs.


**Table 3.** Off-line tests\* for the cloud microphysical processes and statistical mean radii at  $R_{qi} = 20 \mu\text{m}$  (left) and  $R_{qi} = 60 \mu\text{m}$  (right).

	$R_{qi} = 20 \mu\text{m}$			$R_{qi} = 60 \mu\text{m}$		
	$\mu_i = 0$	$\mu_i = 2$	$\mu_i = 5$	$\mu_i = 0$	$\mu_i = 2$	$\mu_i = 5$
$R_{ni} (\mu\text{m})$	11.00	15.33	17.26	33.02	45.98	51.78
$R_{ai} (\mu\text{m})$	15.57	17.70	18.64	46.70	53.09	55.93
$R_{ei} (\mu\text{m})$	33.02	25.54	23.01	99.06	76.63	69.04
$I/\tau_i (10^{-4} \text{ s}^{-1})$	3.35	4.66	5.25	10.04	13.98	15.74
$\tau_i (\text{s})$	2989.21	2146.68	1906.05	996.40	715.56	635.35
$q_{iauto}/q_i (\%)$	2.00	0.01	0	64.08	36.54	18.40
$N_{iauto}/N_i (\%)$	0.01	0	0	4.84	4.23	2.63
$R_{qi\_afterauto} (\mu\text{m})$	19.87	20.00	20.00	43.36	52.31	56.57
$V_{qi} (\text{cm s}^{-1})$	7.96	5.54	4.68	23.87	16.62	14.04
$V_{ni} (\text{cm s}^{-1})$	1.99	2.77	3.12	5.97	8.31	9.36

190 \*Note  $\tau_i$  is calculated at  $T = 220 \text{ K}$ ,  $P = 330 \text{ hPa}$ , and  $N_i = 10^5 \text{ kg}^{-1}$  ( $\sim 50 \text{ L}^{-1}$ ).  $V_{ni}$  and  $V_{qi}$  are calculated at  $T = 220 \text{ K}$  and  $P = 330 \text{ hPa}$ .

### 3.2 CAM6 simulations

During the evolution of stratiform clouds, the properties of ice clouds (e.g.,  $q_i$ ,  $N_i$ , and  $R_{ni}$ , including mixed-phase clouds) largely determine the ice-phase cloud microphysical processes. Meanwhile, these cloud microphysical processes in turn change the cloud properties. They interact as both cause and effect and finally reach equilibrium climate states. To facilitate 195 the subsequent analyses, the cloud properties and  $\mu_i$ -related cloud microphysical processes are shown together in one figure. For ease of expression, “ $\delta$ ” is used to denote the changes in cloud properties that are caused by the cloud microphysical process during one model time step (tendency  $\times$  one time step). For example, the changes in  $q_i$  and  $N_i$  that are caused by the sedimentation process during one model time step are denoted as  $\delta q_{ised}$  and  $\delta N_{ised}$ , respectively.



**Figure 2.** Annual zonal mean in-cloud variables from the Mu0 (first and fourth columns),  $\Delta\text{Mu2}$  (second and fifth columns) and  $\Delta\text{Mu5}$  (third and sixth columns) experiments. Shown are the ICs mass mixing ratio ( $q_i$ ) and number density ( $N_i$ ), mass-weighted and number-weighted radii ( $R_{qi}$  and  $R_{ni}$ ), effective radius ( $R_{ei}$ ), reciprocal of the supersaturation relaxation time scale ( $1/\tau_i$ ), ice supersaturation ( $S_i$ ), change in  $q_i$  caused by deposition/sublimation process ( $\delta q_{idep}$ ), portion of ICs that are converted to snow in terms of mass and number ( $q_{iauto}/q_i$  and  $N_{iauto}/N_i$ ), newly formed IC number density from the nucleation process ( $\delta N_{inuc}$ ), the updated mass-weighted radius ( $R_{qi}^*$ ) used for calculating the sedimentation process, mass-weighted and number-weighted fall velocities ( $V_{qi}$  and  $V_{ni}$ ), and changes in  $q_i$  and  $N_i$  that are caused by the sedimentation process ( $\delta q_{ised}$  and  $\delta N_{ised}$ ). Except for  $\delta q_{idep}$ ,  $\delta q_{ised}$ ,  $\delta N_{inuc}$  and  $\delta N_{ised}$ , the other variables are shown as their relative changes (i.e.,  $\Delta\text{Mu2}/\text{Mu0}$  and  $\Delta\text{Mu5}/\text{Mu0}$ ). The Y-axis indicates the atmospheric pressure (unit: hPa). The two black dashed curves are the 0 and  $-37^\circ\text{C}$  isotherms. All results are sampled from model grids where the ice cloud fraction is greater than 1%. The shadows indicate that the differences between two experiments are not significant at the 95% level based on the Student's  $t$  test.

Fig. 2 shows the model results from the Mu0, Mu2, and Mu5 experiments. The  $q_i^{\text{Mu0}}$  is larger in the upper tropical troposphere ( $> 3 \mu\text{g L}^{-1}$ ) and relatively larger in the lower troposphere over middle latitudes ( $> 1 \mu\text{g L}^{-1}$ ). The spatial pattern of  $q_i^{\text{Mu0}}$  is generally in agreement with the satellite retrieval data (Li et al., 2012). Higher  $N_i^{\text{Mu0}}$  ( $> 200 \text{ L}^{-1}$ ) can be found in the tropopause region, where homogeneous freezing produces a large number of ICs (not shown) due to sufficient soluble aerosol particles, higher subgrid vertical velocity and lower temperature (Shi et al., 2015). All statistical

mean radii (i.e.,  $R_{qi}$ ,  $R_{ni}$ , and  $R_{ei}$ ) decrease with altitude increasing. One possible reason is that it is hard for ICs to grow big in the upper troposphere because the water vapor density is very small over there (lower temperature). Furthermore, the size-sorting effect (i.e., sedimentation process) could also be a contributor to this phenomenon (Milbrandt and Yau, 2005; Khain et al., 2015). As expected,  $R_{ni}$  is less than  $R_{qi}$ , and  $R_{ei}$  is larger than  $R_{qi}$ . After considering the impact of  $\mu_i$  (i.e.,  $\mu_i = 2$  or 5),

220 the  $\Delta\text{Mu}2$  and  $\Delta\text{Mu}5$  experiments show that  $q_i$  is significantly increased while  $N_i$  is significantly decreased. The  $q_i^{\Delta\text{Mu}2/\text{Mu}0}$  is 30-100% in nearly all regions, and the  $q_i^{\Delta\text{Mu}5/\text{Mu}0}$  reaches even higher levels ( $> 100\%$ ) in most regions. Both  $N_i^{\Delta\text{Mu}2/\text{Mu}0}$  and  $N_i^{\Delta\text{Mu}5/\text{Mu}0}$  are  $< -20\%$  above the  $-37^\circ\text{C}$  isotherm and even reach  $-50\%$  in the upper tropical troposphere. Consistent with the increase in  $q_i$  and the decrease in  $N_i$ , the  $R_{qi}$  significantly increases. The  $R_{qi}^{\Delta\text{Mu}2/\text{Mu}0}$  is 30-100% above the  $-37^\circ\text{C}$  isotherm, and the  $R_{qi}^{\Delta\text{Mu}5/\text{Mu}0}$  is 30-100% in most regions and even reaches 100% in a few regions of the upper tropical troposphere.

225 Because  $R_{ni}$  increases with increasing  $\mu_i$  at a fixed  $R_{qi}$  value (Section 3.1), the relative increases in  $R_{ni}$  from the  $\Delta\text{Mu}2$  and  $\Delta\text{Mu}5$  experiments (i.e.,  $R_{ni}^{\Delta\text{Mu}2/\text{Mu}0}$  and  $R_{ni}^{\Delta\text{Mu}5/\text{Mu}0}$ ) are obviously higher than the relative increases in  $R_{qi}$  (i.e.,  $R_{qi}^{\Delta\text{Mu}2/\text{Mu}0}$  and  $R_{qi}^{\Delta\text{Mu}5/\text{Mu}0}$ ). The  $R_{ni}^{\Delta\text{Mu}2/\text{Mu}0}$  is  $> 100\%$  in some regions, and the  $R_{ni}^{\Delta\text{Mu}5/\text{Mu}0}$  is  $> 100\%$  in most regions. Compared with the relative increases in  $R_{qi}$ , the relative increases in  $R_{ei}$  from the  $\Delta\text{Mu}2$  and  $\Delta\text{Mu}5$  experiments are obviously reduced or even negative because  $R_{ei}$  decreases with increasing  $\mu_i$  at a fixed  $R_{qi}$  value (Section 3.1). Overall, the impacts of  $\mu_i$  on  $q_i$  and  $N_i$  are notable. The changes in the statistical mean radii (i.e.,  $R_{qi}$ ,  $R_{ni}$ , and  $R_{ei}$ ) can be explained by the changes in  $q_i$ ,  $N_i$ , and  $\mu_i$ .

230 This paragraph analyzes the interaction between the ice cloud properties ( $q_i$ ,  $N_i$ , and  $R_{ni}$ ) and the IC deposition/sublimation process, and the influence of  $\mu_i$  on this interaction. Since  $1/\tau_i$  is proportional to  $N_i R_{ni}$  (Table 1), the  $1/\tau_i^{\text{Mu}0}$  is larger in the upper tropical troposphere ( $> 20 \times 10^{-4} \text{ s}^{-1}$ ) due to the high  $N_i^{\text{Mu}0}$  ( $> 200 \text{ L}^{-1}$ ). Both the  $\Delta\text{Mu}2$  and  $\Delta\text{Mu}5$  experiments show that the  $1/\tau_i$  increases in most regions because the relative increase in  $R_{ni}$  (i.e.,  $R_{ni}^{\Delta\text{Mu}2/\text{Mu}0}$  and  $R_{ni}^{\Delta\text{Mu}5/\text{Mu}0}$ ) is stronger than the 235 relative decrease in  $N_i$  (i.e.,  $N_i^{\Delta\text{Mu}2/\text{Mu}0}$  and  $N_i^{\Delta\text{Mu}5/\text{Mu}0}$ ). However, the  $1/\tau_i$  is slightly decreased in some regions of the upper tropical troposphere because the relative decrease in  $N_i$  is remarkable ( $< -50\%$ ) in these regions. The  $\delta q_{idep}$  which indicates the change in  $q_i$  caused by the deposition/sublimation process is mainly determined by the  $1/\tau_i$  and in-cloud ice supersaturation ( $S_i$ ) (Morrison and Gettelman, 2008). Except for a very small region, the annual zonal mean  $S_i^{\text{Mu}0}$  is positive. This is consistent with the deposition events being much more frequent than sublimation events (not shown). When  $S_i > 0$ , 240 ice-supersaturation (i.e.,  $S_i > 0$ ) towards ice-saturation (i.e.,  $S_i = 0$ ) occurs because the water vapor is consumed by  $\delta q_{idep}$  (Korolev et al., 2003; Krämer et al., 2009). The  $S_i^{\text{Mu}0}$  is lower ( $< 3\%$ ) in the upper tropical troposphere due to the high  $1/\tau_i^{\text{Mu}0}$  ( $> 20 \times 10^{-4} \text{ s}^{-1}$ ). Both the  $\Delta\text{Mu}2$  and  $\Delta\text{Mu}5$  experiments show that  $S_i$  is increased in the upper tropical troposphere due to the decreasing  $1/\tau_i$ , and  $S_i$  is decreased in the other regions due to the increasing  $1/\tau_i$ . It is noteworthy that the  $S_i^{\Delta\text{Mu}2}$  and  $S_i^{\Delta\text{Mu}5}$  in the mixed-phase cloud layers are obviously weaker than those in the pure ice cloud layers (i.e., above the  $-37^\circ\text{C}$  isotherm). 245 This is consistent with that the  $S_i$  is relatively stable in mixed-phase clouds because liquid droplets are often present. The  $\delta q_{idep}^{\text{Mu}0}$  is generally decreased with the altitude because the saturated vapor pressure significantly decreases with decreasing air temperature. The comparison between  $\delta q_{idep}$  and  $q_i$  suggests that  $\delta q_{idep}$  is an important source of  $q_i$ . The  $\delta q_{idep}^{\Delta\text{Mu}2}$  and  $\delta q_{idep}^{\Delta\text{Mu}5}$  are greater than  $0.1 \mu\text{g L}^{-1}$  in most mixed-phase cloud layers due to the strongly increasing  $1/\tau_i$  and relatively stable


$S_i$  values. This suggests that the increasing  $\mu_i$  could lead to a higher equilibrium state of  $q_i$  in the mixed-phase cloud layers via the deposition process. The  $\delta q_{idep}^{\Delta Mu2}$  and  $\delta q_{idep}^{\Delta Mu5}$  are negative between 200 hPa and 300 hPa mainly because the  $S_i^{\Delta Mu2}$  and  $S_i^{\Delta Mu5}$  are negative and the  $1/\tau_i^{\Delta Mu2/Mu0}$  and  $1/\tau_i^{\Delta Mu5/Mu0}$  are relatively small. The  $\delta q_{idep}^{\Delta Mu2}$  and  $\delta q_{idep}^{\Delta Mu5}$  are positive above 100 hPa mainly because the  $S_i^{\Delta Mu2}$  and  $S_i^{\Delta Mu5}$  are positive. These results indicate that the impact of  $\mu_i$  on  $\delta q_{idep}$  becomes complex above the  $-37^{\circ}\text{C}$  isotherm, where  $S_i$  is more susceptible to  $1/\tau_i$  and  $\delta q_{idep}$ . Meanwhile, the impact of  $\mu_i$  on  $\delta q_{idep}$  also becomes weak above the  $-37^{\circ}\text{C}$  isotherm because the feedback processes (i.e., the interaction between  $S_i$  and  $\delta q_{idep}$ ) become important. In short, the  $\mu_i$ -induced changes in the deposition/sublimation process (i.e.,  $1/\tau_i$  and  $\delta q_{idep}$ ) can be largely explained by the changes in  $N_i$  and  $R_{ni}$ . One reason for the higher  $q_i$  in the mixed-phase cloud layers from the Mu2 and Mu5 experiments is that  $\delta q_{idep}$  increases with increasing  $\mu_i$ .

This paragraph analyzes the interaction between the ice cloud properties ( $q_i$ ,  $N_i$ , and  $R_{qi}$ ) and the autoconversion of IC to snow process (hereafter the autoconversion process), and the influence of  $\mu_i$  on this interaction. Both  $q_{iauto}/q_i^{\text{Mu0}}$  and  $N_{iauto}/N_i^{\text{Mu0}}$  are decreased with the altitude because the  $R_{qi}^{\text{Mu0}}$  is decreased with the altitude. As expected, the  $q_{iauto}/q_i^{\text{Mu0}}$  is considerable and much larger than the  $N_{iauto}/N_i^{\text{Mu0}}$ . It is clear that the autoconversion process is an important sink of  $q_i$ . However, the autoconversion process is not an important sink of  $N_i$  because the  $N_{iauto}/N_i$  is very small. Both the  $\Delta\text{Mu2}$  and  $\Delta\text{Mu5}$  experiments show that the  $q_{iauto}/q_i$  is significantly decreased because the autoconversion process obviously becomes difficult at higher  $\mu_i$  values (off-line tests, Section 3.1). The difficult autoconversion process leads to an equilibrium state with higher  $q_i$  and larger  $R_{qi}$ . Because of the larger  $R_{qi}$ , the  $N_{iauto}/N_i$  from the Mu2 and Mu5 experiments are significantly increased. The increasing  $N_{iauto}/N_i$  from the Mu2 and Mu5 experiments might be a main reason for the decrease in  $N_i$  in the mixed-phase cloud layers. However, the remarkable decrease in  $N_i$  (mostly in the pure ice cloud layers) from the Mu2 and Mu5 experiments is mainly due to the ice nucleation process. In the MG scheme, the newly formed IC number density (excluding the ICs in mixed-phase clouds) is calculated by a physically based ice nucleation parameterization (Liu and Penner, 2005). Because the autoconversion process becomes difficult in the Mu2 and Mu5 experiments, the in-cloud ICs should have longer lifetimes and larger radii. As a result,  $\delta N_{inuc}$ , which denotes the newly formed IC number density from the nucleation process, significantly decreases in the Mu2 and Mu5 experiments (Fig. 2). The main reason is that the pre-existing ICs would hinder the subsequent ice nucleation process (especially for homogeneous freezing) owing to the depletion of water vapor via deposition growth (Barahona et al., 2014; Shi et al., 2015).  $\delta N_{inuc}$  is the main source of  $N_i$ . Therefore, both the  $\Delta\text{Mu2}$  and  $\Delta\text{Mu5}$  experiments show that  $N_i$  is significantly decreased. In short, the increase in  $\mu_i$  causes the autoconversion process to be difficult and then leads to a higher equilibrium state of  $q_i$  and  $R_{qi}$ . Meanwhile,  $N_i$  is significantly decreased due to the higher equilibrium state of  $q_i$  and  $R_{qi}$  (i.e., the stronger suppression effect of the pre-existing ICs on the ice nucleation process).

This paragraph analyzes the interaction between the ice cloud properties and the IC sedimentation process, and the influence of  $\mu_i$  on this interaction. The sedimentation process is the last cloud microphysical process in the MG scheme. The IC fall velocity is calculated based on the updated cloud properties (i.e., the other cloud microphysical processes at this model time

step have been considered). Here,  $R_{qi}^*$  denotes the updated  $R_{qi}$ , which includes the changes caused by the deposition/sublimation and autoconversion processes at this model time step. In the mixed-phase cloud layers, the  $R_{qi}^{*,\text{Mu}0}$  is slightly less than the  $R_{qi}^{\text{Mu}0}$  because the sedimentation process has not occurred. After considering the impacts of  $\mu_i$  on the cloud microphysical processes introduced above, the relative increases in  $R_{qi}^*$  from the  $\Delta\text{Mu}2$  and  $\Delta\text{Mu}5$  experiments (i.e.,  $R_{qi}^{*,\Delta\text{Mu}2/\text{Mu}0}$  and  $R_{qi}^{*,\Delta\text{Mu}5/\text{Mu}0}$ ) are higher than the relative increase in  $R_{qi}$ . As expected, both  $V_{qi}^{\text{Mu}0}$  and  $V_{ni}^{\text{Mu}0}$  decrease with altitude increasing, and the  $V_{qi}^{\text{Mu}0}$  is obviously larger than the  $V_{ni}^{\text{Mu}0}$ . Although the  $R_{qi}^{*,\Delta\text{Mu}2/\text{Mu}0}$  and  $R_{qi}^{*,\Delta\text{Mu}5/\text{Mu}0}$  are positive, the  $V_{qi}^{\Delta\text{Mu}2/\text{Mu}0}$  and  $V_{qi}^{\Delta\text{Mu}5/\text{Mu}0}$  are negative in most regions because  $V_{qi}$  decreases with increasing  $\mu_i$  (off-line tests, Section 3.1). However, both  $V_{qi}^{\Delta\text{Mu}2/\text{Mu}0}$  and  $V_{qi}^{\Delta\text{Mu}5/\text{Mu}0}$  are positive in some layers over the tropics and subtropics, where the  $R_{qi}^{*,\Delta\text{Mu}2/\text{Mu}0}$  and  $R_{qi}^{*,\Delta\text{Mu}5/\text{Mu}0}$  are relatively higher. Because  $V_{ni}$  increases with increasing  $\mu_i$  at a fixed  $R_{qi}$  value (off-line tests, Table 3) and the  $R_{qi}^*$  from the Mu2 and Mu5 experiments are increased, the relative increases in  $V_{ni}$  from the  $\Delta\text{Mu}2$  and  $\Delta\text{Mu}5$  experiments are remarkable. The  $V_{ni}^{\Delta\text{Mu}2/\text{Mu}0}$  is  $> 100\%$  in some regions, and the  $V_{ni}^{\Delta\text{Mu}5/\text{Mu}0}$  is  $> 100\%$  in most regions.  $\delta q_{ised}$  is mainly determined by the gradient of  $V_{qi}q_i$  in the vertical direction. Actually, the newly updated  $q_i$  between the substeps of the sedimentation process is used for calculating  $\delta q_{ised}$ . Similarly,  $\delta N_{ised}$  is mainly determined by the gradient of  $V_{ni}N_i$  in the vertical direction. Furthermore, the ICs that fall into the clear portions of the lower model layer sublimate instantly. Therefore, both  $\delta q_{ised}^{\text{Mu}0}$  and  $\delta N_{ised}^{\text{Mu}0}$  are negative in most regions. This is consistent with sedimentation being a sink of clouds. The  $\delta q_{ised}$  from the Mu2 and Mu5 experiments (i.e.,  $\delta q_{ised}^{\Delta\text{Mu}2}$  and  $\delta q_{ised}^{\Delta\text{Mu}5}$ ) decrease (negative, stronger sink) in most regions mainly because of the increasing  $V_{qi}$  and higher  $q_i$ . The  $\delta N_{ised}$  from the Mu2 and Mu5 experiments (i.e.,  $\delta N_{ised}^{\Delta\text{Mu}2}$  and  $\delta N_{ised}^{\Delta\text{Mu}5}$ ) increase (positive, weaker sink) in a few layers over the tropics. This is mainly due to the changes 300 in the vertical gradient of  $N_i$ . Both the  $\Delta\text{Mu}2$  and  $\Delta\text{Mu}5$  experiments show that the changes in  $\delta q_{ised}$  and  $\delta N_{ised}$  are generally weaker than the changes in  $\delta q_{idep}$ ,  $\delta q_{iauto}$  (i.e.,  $q_i \times q_{iauto}/q_i$ ), and  $\delta N_{inuc}$ . In short, the fall velocities (i.e.,  $V_{qi}$  and  $V_{ni}$ ) and their impacts on ice clouds (i.e.,  $\delta q_{ised}$  and  $\delta N_{ised}$ ) are mainly determined by the cloud properties (i.e.,  $q_i$ ,  $N_i$ ,  $R_{qi}$ , and  $R_{qi}^*$ ). Although the sedimentation process is also a main factor that determines the cloud properties, the changes in the sedimentation process that are caused by the increasing  $\mu_i$  are not as strong as those in the deposition/sublimation, 305 autoconversion, and nucleation processes.

Based on the analyses presented above, it can be concluded that increasing  $\mu_i$  would lead to a climate equilibrium state with higher  $q_i$  and lower  $N_i$  in most regions. The changes in the statistical mean radii (i.e.,  $R_{qi}$ ,  $R_{ni}$ , and  $R_{ei}$ ) and ice-phase cloud microphysical processes (i.e.,  $\delta q_{idep}$ ,  $q_{iauto}/q_i$ ,  $N_{iauto}/N_i$ ,  $\delta N_{inuc}$ ,  $\delta q_{ised}$  and  $\delta N_{ised}$ ) are mainly determined by the higher  $q_i$ , lower  $N_i$ , and increasing  $\mu_i$ . On the other hand, the higher  $q_i$  and lower  $N_i$  can largely be explained by the changes in the ice-phase 310 cloud microphysical processes (i.e.,  $\delta q_{idep}$ ,  $q_{iauto}/q_i$ , and  $\delta N_{inuc}$ ) that are caused by the increasing  $\mu_i$ . Furthermore, the  $\Delta\text{Mu}2$  and  $\Delta\text{Mu}5$  experiments show very similar spatial patterns for the  $\mu_i$ -induced changes. This suggests that the impact of  $\mu_i$  on the simulated climate equilibrium state is stable.




315 **Figure 3.** Similar to Fig. 2 but for the  $\Delta\text{Tao5}$ ,  $\Delta\text{Auto5}$ ,  $\Delta\text{Fall5}$  and  $\Delta\text{Rei5}$  experiments. Except for the  $R_{ei}^{\text{Rei5}}$ , all statistical mean radii are calculated with  $\mu_i = 0$ .

Fig. 3 shows the changes in the simulated climate equilibrium states that are caused by each individual  $\mu_i$ -related process. The  $1/\tau_i$  from the Tao5 experiment is significantly increased ( $1/\tau_i^{\Delta\text{Tao5}/\text{Mu}0}$ ). Similar to the  $\Delta\text{Mu2}$  and  $\Delta\text{Mu5}$  experiments, this increasing  $1/\tau_i$  could lead to a higher equilibrium state of  $q_i$  in the mixed-phase cloud layers ( $q_i^{\Delta\text{Tao5}/\text{Mu}0}$ ) via the deposition/sublimation process ( $\delta q_{i\text{dep}}^{\Delta\text{Tao5}}$ ). However, this increasing  $1/\tau_i$  leads to lower  $q_i$  and lower  $N_i$  in most of the pure ice cloud layers. The main reason might be that the ICs grow faster and their lifetimes become shorter (Mitchell, 1991; DeMott et al., 2010; Storelvmo et al., 2013). The  $q_{i\text{auto}}/q_i$  from the Auto5 experiment is significantly decreased ( $q_{i\text{auto}}/q_i^{\Delta\text{Tao5}/\text{Mu}0}$ ). This could lead to a higher  $q_i$  in nearly all regions ( $q_i^{\Delta\text{Auto5}/\text{Mu}0}$ ) and a lower  $N_i$  in the pure ice cloud layers ( $N_i^{\Delta\text{Auto5}/\text{Mu}0}$ ). The mechanism is the same as that introduced based on the  $\Delta\text{Mu2}$  and  $\Delta\text{Mu5}$  experiments. It is noteworthy that the  $N_i$  from the Auto5 experiment is slightly increased in some mixed-phase cloud layers. The main reason might be that the accretion of  $N_i$  by snow is significantly decreased in the mixed-phase cloud layers (not shown) due to the difficult autoconversion process. The  $V_{qi}$  from the Fall5 experiment is significantly decreased ( $V_{qi}^{\Delta\text{Fall5}/\text{Mu}0}$ ), and the sink term of  $q_i$  due to sedimentation becomes weaker (i.e., positive  $\delta q_{i\text{sed}}^{\Delta\text{Fall5}}$ ) in most regions. Unlike the  $V_{qi}^{\Delta\text{Fall5}/\text{Mu}0}$ , the  $V_{ni}$  from the Fall5 experiment obviously increase ( $V_{ni}^{\Delta\text{Fall5}/\text{Mu}0}$ ), and the sink term of  $N_i$  due to sedimentation becomes stronger (i.e., negative  $\delta N_{i\text{sed}}^{\Delta\text{Fall5}}$ ).

330  $\delta N_{ised}^{\Delta Fall5}$ ) in the pure ice cloud layers. These might be the main reasons for the increase in  $q_i$  and the decrease in  $N_i$  in the pure ice cloud layers over the tropics ( $q_i^{\Delta Fall5/Mu0}$  and  $N_i^{\Delta Fall5/Mu0}$ ). It is interesting to note that the  $R_{qi}$ ,  $R_{ni}$ , and  $R_{ei}$  from the Fall5 experiment all increase in pure ice cloud layers (i.e., upper layers) and decrease in mixed-phase cloud layers (i.e., lower layers). This can be explained by the  $\mu_i$ -induced weaker size-sorting mechanism (Section 3.1). The  $R_{ei}$  from the Rei5 experiment ( $R_{ei}^{\Delta Rei5/Mu0}$ ) is significantly decreased. Because the change of  $R_{ei}$  does not directly affect the cloud microphysical processes, the changes in cloud properties from the  $\Delta Rei5$  experiment are not statistically significant in most regions. Taken overall, the above analyses clarify the mechanism of  $\mu_i$ 's impacts. Increasing  $\mu_i$  in autoconversion impacts pure ice clouds the most (i.e., significantly increased  $q_i$  and significantly decreased  $N_i$  in the pure ice cloud layers). Furthermore, increasing  $\mu_i$  in autoconversion also leads to a much higher  $q_i$  in the mixed-phase cloud layers. Increasing  $\mu_i$  in deposition/sublimation can also lead to a higher  $q_i$  in the mixed-phase cloud layers. Increasing  $\mu_i$  in sedimentation can lead to a higher IC radius in the 340 upper layers and lower IC radius in the lower layers. The impacts from sedimentation and deposition/sublimation are obviously weaker than those from autoconversion. The changes caused by increasing  $\mu_i$  in the radiative process (i.e.,  $R_{ei}$ ) are relatively chaotic.

The above analyses focus on cloud properties and cloud microphysical processes (i.e., in-cloud variables). This paragraph 345 discusses the impacts of  $\mu_i$  on radiation and precipitation. The annual zonal mean distributions of the ice water path (IWP), column  $N_i$  (ColNi), longwave (CRELW) and shortwave (CREsw) cloud radiative effects, and convective (RainC) and large-scale (RainL) precipitation rates are shown in Fig. 4, and the corresponding global annual mean values are listed in Table 4. The comparison of the Mu0, Mu2 and Mu5 experiments shows that the zonal mean IWPs over all latitudes clearly increase with increasing  $\mu_i$ . This is consistent with the changes in in-cloud  $q_i$  (Fig. 2). The comparison of the  $\Delta Mu5$ ,  $\Delta Tao5$ ,  $\Delta Auto5$ ,  $\Delta Fall5$  and  $\Delta Rei5$  experiments shows that the  $\mu_i$ -induced increases in IWP are mainly provided by the autoconversion 350 process. Compared to the Mu0 experiment, the ColNi from the Mu2 and Mu5 experiments obviously decrease over tropical regions. It is clear that the autoconversion process is also the main contributor to the decreases in ColNi (Fig. 4, right column). Compared to the Mu0 experiment, both CRELW and CREsw from the Mu2 and Mu5 experiments are obviously enhanced mainly because of the increasing IWPs. It is clear that the enhancements of CRELW and CREsw are also mainly contributed to by the autoconversion process (Fig. 4). Both the CRELW and CREsw from the Rei5 experiment are also 355 obviously enhanced in terms of their zonal mean values (Fig. 4) and global mean values (Table 4,  $CRE_{LW}^{\Delta Rei5} = 1.29 \text{ W m}^{-2}$  and  $CRE_{sw}^{\Delta Rei5} = -1.79 \text{ W m}^{-2}$ ). This suggests that the impact of  $\mu_i$  on  $R_{ei}$  could lead to considerable changes in the Earth's radiation budget. Compared to the impacts of  $\mu_i$  on radiation, the impact on large-scale precipitation (i.e., RainL) is not statistically significant (Fig. 4, left column). However, the convective precipitation from the  $\Delta Mu5$  experiment (i.e., RainC $^{\Delta Mu5}$ ) is significantly reduced over the tropics and subtropics (Fig. 4, right column). The reason is that the increase in 360 ice clouds (i.e.,  $q_i$ ) increases atmospheric stability via the radiative budget and then leads to weaker convective precipitation (Andrews et al., 2010; Wang et al., 2014). Overall, the impacts of  $\mu_i$  on radiation and precipitation are considerable. The global mean  $CRE_{LW}^{\Delta Mu5}$ ,  $CRE_{sw}^{\Delta Mu5}$  and  $RainC^{\Delta Mu5}$  are  $5.58 \text{ W m}^{-2}$ ,  $-5.34 \text{ W m}^{-2}$  and  $-0.12 \text{ mm day}^{-1}$ , respectively. These

changes are mainly contributed to by the autoconversion process. Furthermore, the comparisons between the  $\Delta\text{Mu}2$  and  $\Delta\text{Mu}5$  experiments (Fig. 4 and Table 4) show that, in most cases, the  $\mu_i$ -induced changes are enhanced with increasing  $\Delta\mu_i$ .  
 365 This suggests that, in terms of the zonal mean and global mean values, the impacts of  $\mu_i$  are relatively stable.



**Figure 4.** Annual zonal mean distributions of the ice water path (IWP), column  $N_i$  ( $\text{ColN}_i$ ), longwave and shortwave cloud radiative effects ( $\text{CRE}_{\text{LW}}$  and  $\text{CRE}_{\text{sw}}$ ), and convective and large-scale precipitation rates ( $\text{RainC}$  and  $\text{RainL}$ ). In the right panel (i.e.,  $\Delta$ ), the vertical bars overlying these lines indicate the ranges of two standard deviations from the difference of each of 10 years at different latitudes.

370 **Table 4.** The global annual mean variables are shown in Fig. 4. The corresponding standard deviations calculated from the difference of each year for 10 years are shown in brackets.

Experiments	Mu0	$\Delta\text{Mu}2$	$\Delta\text{Mu}5$	$\Delta\text{Tao}5$	$\Delta\text{Auto}5$	$\Delta\text{Fall}5$	$\Delta\text{Rei}5$
IWP ( $\text{g m}^{-2}$ )	4.74	1.37(0.10)	2.97(0.06)	-0.20(0.05)	3.64(0.06)	-0.01(0.04)	-0.09(0.06)
$\text{ColN}_i$ ( $10^8 \text{ m}^{-2}$ )	1.89	-0.41(0.06)	-0.56(0.04)	-0.15(0.03)	-0.37(0.03)	-0.06(0.04)	-0.07(0.05)
$\text{CRE}_{\text{LW}}$ ( $\text{W m}^{-2}$ )	22.22	3.40(0.12)	5.58(0.13)	-0.52(0.07)	3.40(0.11)	0.49(0.09)	1.29(0.10)
$\text{CRE}_{\text{sw}}$ ( $\text{W m}^{-2}$ )	-49.25	-3.00(0.43)	-5.34(0.31)	0.54(0.38)	-2.52(0.46)	-0.20(0.36)	-1.79(0.24)
$\text{RainC}$ ( $\text{mm day}^{-1}$ )	1.57	-0.08(0.01)	-0.12(0.01)	0 (0.01)	-0.07(0.01)	-0.02(0.01)	-0.02(0.01)
$\text{RainL}$ ( $\text{mm day}^{-1}$ )	1.39	0.01(0.01)	0.01(0.01)	0 (0.01)	0.01(0.01)	0 (0)	0 (0.01)

#### 4 Summary and conclusions

This paper investigates the impacts of  $\mu_i$  on climate simulations with the CAM6 model. To achieve this, the two-moment bulk cloud microphysics scheme used in CAM6 was modified to consider the  $\mu_i$ 's impacts by a tunable parameter. After that,

375 the impacts of  $\mu_i$  on the IC size distribution and  $\mu_i$ -related cloud microphysical processes are illustrated first by calculation equations and off-line tests, and the impacts of  $\mu_i$  on the climate simulations are then analyzed with the CAM6 model.

The impacts of  $\mu_i$  on the IC size distribution and various statistical mean radii are clearly explained by the calculation equations.  $R_{qi}$  can be calculated from  $q_i$  and  $N_i$ , and the normalized IC size distribution (i.e., the relative contributions from each bin) can be calculated from  $R_{qi}$  and  $\mu_i$ . The impacts of  $\mu_i$  on mass-weighted size distribution is obviously different from

380 that on the commonly used number-weighted size distribution (Fig.1). Unlike the number contributions, the mass contributions of the larger size bins significantly decrease with increasing  $\mu_i$  because the mass contribution is more sensitive to the IC radius. In the bulk cloud microphysics scheme, the physical processes are calculated based on various statistical mean radii. The ratios of the other statistical mean radii (i.e.,  $R_{ni}$ ,  $R_{ai}$ , and  $R_{ei}$ ) to  $R_{qi}$  are functions of  $\mu_i$ . At  $\mu_i \geq 0$ ,  $R_{ni}$  and  $R_{ai}$  are always less than  $R_{qi}$ , whereas  $R_{ei}$  is always greater than  $R_{qi}$ . The differences among these statistical mean radii become

385 small with increasing  $\mu_i$ , which is consistent with the narrower size distribution determined by higher  $\mu_i$  values.

The impacts of  $\mu_i$  on the  $\mu_i$ -related cloud microphysical processes are clearly illustrated by the off-line tests. Under the same  $q_i$  and  $N_i$  ( $R_{qi}$  is also fixed), the IC deposition/sublimation process is considerably accelerated with increasing  $\mu_i$  because the

390  $R_{ni}$  used for calculating deposition/sublimation is increasing with increasing  $\mu_i$ . Under the same  $R_{qi}$  ( $R_{qi}$  is much less than the snow radius), the autoconversion of IC to snow obviously becomes difficult with increasing  $\mu_i$  because the portion of ICs with radii greater than the threshold ( $> R_{cs}$ ) decreases under the narrow size distribution. A major effect of IC sedimentation

is size-sorting because  $V_{qi}$  is greater than  $V_{ni}$ , and  $\mu_i$  plays an important role in determining the rate of size-sorting (Milbrandt and Yau, 2005). In this study, the off-line tests clearly show that the difference in IC radius between two adjacent model layers caused by sedimentation becomes small with increasing  $\mu_i$  because the difference between  $V_{qi}$  and  $V_{ni}$  becomes small (i.e., the size-sorting rate becomes slow).

395 The climate simulations show that the impacts of  $\mu_i$  on the ice cloud properties are notable, and the main corresponding mechanisms are clear. After increasing  $\mu_i$  from 0 to 2 and 5,  $q_i$  significantly increases, while  $N_i$  significantly decreases. The accelerated deposition process contributes to the higher  $q_i$  in the mixed-phase cloud layers where the ice supersaturation ( $S_i > 0$ ) is relatively stable. The difficult autoconversion process leads to longer IC lifetime and higher  $q_i$ . Meanwhile,  $N_i$

400 significantly decreases because the newly formed IC number density is significantly decreased owing to the longer IC lifetime and higher  $q_i$ . The experiments with only one modified  $\mu_i$ -related process make the mechanisms of  $\mu_i$ 's impacts more clear. Autoconversion contributes the most. The  $\mu_i$  also has considerable impacts on radiation and precipitation. After increasing  $\mu_i$  from 0 to 5, the global mean  $CRE_{LW}$  is increased (stronger warming effect) by  $5.58 \text{ W m}^{-2}$  (25.11%). Meanwhile, the  $CRE_{sw}$  is decreased (less cooling effect) by  $-5.34 \text{ W m}^{-2}$  (10.84%). The enhancement of the cloud radiative

405 effects is largely provided by the higher equilibrium state of  $q_i$ . The considerably stronger CRE<sub>LW</sub> could increase the atmospheric stability and then lead to weaker convective precipitation (Andrews et al., 2010; Wang et al., 2014). As expected, after increasing  $\mu_i$  from 0 to 5, the global mean RainC is decreased by  $-0.12 \text{ mm day}^{-1}$  (7.64%). In short, the impacts of  $\mu_i$  on climate simulations are significant. This suggests that the  $\mu_i$  (i.e., the PSD of ICs) and  $\mu_i$ -related cloud microphysical processes deserve a more realistic representation in climate models, especially for cloud schemes with autoconversion. Fortunately, there have been some studies that can help to address this issue. Such as,  $\mu_i$  is described by an empirical formula (Eidhammer et al., 2017),  $\mu_i$  is prognosed in a three-moment cloud scheme (Milbrandt et al., 2021), and single-ice-category cloud schemes could obviate the need for autoconversion process (e.g., Morrison and Milbrandt, 2015; Eidhammer et al., 2017; Zhao et al., 2017).

410 This study only focuses on the impacts of  $\mu_i$ , the default tunable parameters (except for  $\mu_i$ ) are used in all the simulations. After improving the representation of  $\mu_i$ -related processes, further model tuning and analyses are required based on the updated cloud scheme. Therefore, this study does not estimate which value of  $\mu_i$  could lead to a better simulation. Finally, it is necessary to point out that the main mechanism for  $\mu_i$ 's impacts introduced in this study (i.e., autoconversion becomes difficult with increasing  $\mu_i$ ) is not applicable to climate simulations with the single-ice-category cloud scheme. However, similar to the  $\mu_i$ 's impact on autoconversion, the interaction between small ICs and large ICs (e.g., the accretion of small ICs by large ICs) should become weaker with increasing  $\mu_i$  (i.e., narrower size distribution). Therefore, we can speculate that the impacts of  $\mu_i$  on climate simulations with the single-ice-category cloud scheme may still be worth noting.

## Code and data availability

420 The CAM6 model used in this study is the atmospheric component of the Community Earth System Model version 2.1.3 (CESM2.1.3). The CESM2.1.3 is a release version of CESM2. The model code, scripts, and input data are freely available through a public GitHub repository ([https://escomp.github.io/CESM/versions/cesm2.1/html/downloading\\_cesm.html](https://escomp.github.io/CESM/versions/cesm2.1/html/downloading_cesm.html); last access: 9 May 2022). More details about model workflow can be found in the corresponding quick-start guide. The modified model code, model run control scripts, and simulation results postprocessing scripts covering every data processing action for all the model results reported in the paper are available online at DOI: 10.5281/zenodo.6409156. The FORTRAN code for offline tests is also archived with the same DOI. Furthermore, the NCL scripts and data for making every figure are also available at the above DOI.

## 430 Competing interests

The authors declare that they have no conflict of interest.

## Author contribution

XS designed this study. WZ and XS designed the CAM6 model experiments and developed the modified model code. XS and WZ analyzed the results and wrote the original paper. All authors contributed to improving and reviewing the  
435 manuscript.

## Acknowledgments

This study was supported by the National Key Research and Development Program of China (grant no. 2017YFA0604001) and the National Natural Science Foundation of China (grant no. 41775095). The authors would like to thank Yuxi Zeng and Jiaojiao Liu for checking the English expressions of this manuscript. The model simulation was conducted in the High  
440 Performance Computing Center of Nanjing University of Information Science & Technology.

## References

Andrews, T., Forster, P. M., Boucher, O., Bellouin, N., and Jones, A.: Precipitation, radiative forcing and global temperature change, *Geophysical Research Letters*, 37, L14701, <https://doi.org/10.1029/2010GL043991>, 2010.

Barahona, D., Molod, A., Bacmeister, J., Nenes, A., Gettelman, A., Morrison, H., Phillips, V., and Eichmann, A.:  
445 Development of two-moment cloud microphysics for liquid and ice within the NASA Goddard Earth Observing System Model (GEOS-5), *Geosci. Model Dev.*, 7, 1733-1766, <https://doi.org/10.5194/gmd-7-1733-2014>, 2014.

Bogenshutz, P. A., Gettelman, A., Morrison, H., Larson, V. E., Craig, C., and Schanen, D. P.: Higher-order turbulence closure and its impact on climate simulations in the Community Atmosphere Model, *Journal of Climate*, 26, 9655-9676, <https://doi.org/10.1175/jcli-d-13-00075.1>, 2013.

450 Bogenshutz, P. A., Gettelman, A., Hannay, C., Larson, V. E., Neale, R. B., Craig, C., and Chen, C.-C.: The path to CAM6: coupled simulations with CAM5.4 and CAM5.5, *Geoscientific Model Development*, 11, 235-255, <https://doi.org/10.5194/gmd-11-235-2018>, 2018.

Bony, S., Stevens, B., Frierson, D. M. W., Jakob, C., Kageyama, M., Pincus, R., Shepherd, T. G., Sherwood, S. C., Siebesma, A. P., Sobel, A. H., Watanabe, M., and Webb, M. J.: Clouds, circulation and climate sensitivity, *Nature Geoscience*, 8, 261-  
455 268, <https://doi.org/10.1038/ngeo2398>, 2015.

Danabasoglu, G., Lamarque, J.-F., Bacmeister, J., Bailey, D. A., DuVivier, A. K., Edwards, J., Emmons, L. K., Fasullo, J., Garcia, R., Gettelman, A., Hannay, C., Holland, M. M., Large, W. G., Lauritzen, P. H., Lawrence, D. M., Lenaerts, J. T. M., Lindsay, K., Lipscomb, W. H., Mills, M. J., Neale, R., Oleson, K. W., Otto-Btiesner, B., Phillips, A. S., Sacks, W., Tilmes, S., van Kampenhout, L., Vertenstein, M., Bertini, A., Dennis, J., Deser, C., Fischer, C., Fox-Kemper, B., Kay, J. E.,  
460 Kinnison, D., Kushner, P. J., Larson, V. E., Long, M. C., Mickelson, S., Moore, J. K., Nienhouse, E., Polvani, L., Rasch, P.

J., and Strand, W. G.: The community earth system model version 2 (CESM2), *Journal of Advances in Modeling Earth Systems*, 12, e2019MS001916, <https://doi.org/10.1029/2019MS001916>, 2020.

DeMott, P. J., Prenni, A. J., Liu, X., Kreidenweis, S. M., Petters, M. D., Twohy, C. H., Richardson, M. S., Eidhammer, T., and Rogers, D. C.: Predicting global atmospheric ice nuclei distributions and their impacts on climate, *P. Natl. Acad. Sci. USA*, 107, 11217–11222, <https://doi.org/10.1073/pnas.0910818107>, 2010.

Eidhammer, T., Morrison, H., Bansemer, A., Gettelman, A., and Heymsfield, A. J.: Comparison of ice cloud properties simulated by the Community Atmosphere Model (CAM5) with in-situ observations, *Atmospheric Chemistry and Physics*, 14, 10103-10118, <https://doi.org/10.5194/acp-14-10103-2014>, 2014.

Eidhammer, T., Morrison, H., Mitchell, D., Gettelman, A., and Erfani, E.: Improvements in global climate model microphysics using a consistent representation of ice particle properties, *Journal of Climate*, 30, 609-629, <https://doi.org/10.1175/jcli-d-16-0050.1>, 2017.

Erfani, E. and Mitchell, D. L.: Developing and bounding ice particle mass- and area-dimension expressions for use in atmospheric models and remote sensing, *Atmospheric Chemistry and Physics*, 16, 4379-4400, <https://doi.org/10.5194/acp-16-4379-2016>, 2016.

Gettelman, A. and Morrison, H.: Advanced two-moment bulk microphysics for global models. Part I: Off-line tests and comparison with other schemes, *Journal of Climate*, 28, 1268-1287, <https://doi.org/10.1175/jcli-d-14-00102.1>, 2015.

Heymsfield, A. J.: Properties of tropical and midlatitude ice cloud particle ensembles. Part II: Applications for mesoscale and climate models, *Journal of the Atmospheric Sciences*, 60, 2592-2611, [https://doi.org/10.1175/1520-0469\(2003\)060<2592:Potami>2.0.Co;2](https://doi.org/10.1175/1520-0469(2003)060<2592:Potami>2.0.Co;2), 2003.

Khain, A. P., Beheng, K. D., Heymsfield, A., Korolev, A., Krichak, S. O., Levin, Z., Pinsky, M., Phillips, V., Prabhakaran, T., Teller, A., van den Heever, S. C., and Yano, J. I.: Representation of microphysical processes in cloud-resolving models: Spectral (bin) microphysics versus bulk parameterization, *Reviews of Geophysics*, 53, 247-322, <https://doi.org/10.1002/2014rg000468>, 2015.

King, A. D., Lane, T. P., Henley, B. J., and Brown, J. R.: Global and regional impacts differ between transient and equilibrium warmer worlds, *Nature Climate Change*, 10, 42-47, <https://doi.org/10.1038/s41558-019-0658-7>, 2020.

Korolev, A. V. and Mazin, I. P.: Supersaturation of water vapor in clouds, *Journal of the Atmospheric Sciences*, 60, 2957-2974, [https://doi.org/10.1175/1520-0469\(2003\)060<2957:Sowvic>2.0.Co;2](https://doi.org/10.1175/1520-0469(2003)060<2957:Sowvic>2.0.Co;2), 2003.

Krämer, M., Schiller, C., Afchine, A., Bauer, R., Gensch, I., Mangold, A., Schlicht, S., Spelten, N., Sitnikov, N., Borrmann, S., de Reus, M., and Spichtinger, P.: Ice supersaturations and cirrus cloud crystal numbers, *Atmos. Chem. Phys.*, 9, 3505-3522, <https://doi.org/10.5194/acp-9-3505-2009>, 2009.

Larson, V. E.: CLUBB-SILHS: A parameterization of subgrid variability in the atmosphere, *arXiv preprint arXiv:1711.03675*, 2017.

Li, J. L. F.: An observationally based evaluation of cloud ice water in CMIP3 and CMIP5 GCMs and contemporary reanalyses using contemporary satellite data, *J Geophys Res*, 117, <https://doi.org/10.1029/2012JD017640>, 2012.

495 Liou, K.: Influence of cirrus clouds on weather and climate processes: A Global Perspective, *Monthly Weather Review*, 114, 1167-1199, [https://doi.org/10.1175/1520-0493\(1986\)114<1167:IOCCOW>2.0.CO;2](https://doi.org/10.1175/1520-0493(1986)114<1167:IOCCOW>2.0.CO;2), 1986.

Liu, X. H. and Penner, J. E.: Ice nucleation parameterization for global models, *Meteorologische Zeitschrift*, 14, 499-514, <https://doi.org/10.1127/0941-2948/2005/0059>, 2005.

500 Loftus, A. M., Cotton, W. R., and Carrió, G. G.: A triple-moment hail bulk microphysics scheme. Part I: Description and initial evaluation, *Atmospheric Research*, 149, 35-57, <https://doi.org/10.1016/j.atmosres.2014.05.013>, 2014.

Lohmann, U., Stier, P., Hoose, C., Ferrachat, S., Kloster, S., Roeckner, E., and Zhang, J.: Cloud microphysics and aerosol indirect effects in the global climate model ECHAM5-HAM, *Atmos. Chem. Phys.*, 7, 3425-3446, <https://doi.org/10.5194/acp-7-3425-2007>, 2007.

505 Luo, Z. and Rossow, W. B.: Characterizing tropical cirrus life cycle, evolution, and interaction with upper-tropospheric water vapor using Lagrangian trajectory analysis of satellite observations, *Journal of Climate*, 17, 4541-4563, <https://doi.org/10.1175/3222.1>, 2004.

McFarquhar, G. M., Hsieh, T.-L., Freer, M., Mascio, J., and Jewett, B. F.: The characterization of ice hydrometeor gamma size distributions as volumes in  $N_0 - \lambda - \mu$  phase space: Implications for microphysical process modeling, *Journal of the Atmospheric Sciences*, 72, 892-909, <https://doi.org/10.1175/jas-d-14-0011.1>, 2015.

510 Milbrandt, J. A. and McTaggart-Cowan, R.: Sedimentation-induced errors in bulk microphysics schemes, *Journal of the Atmospheric Sciences*, 67, 3931-3948, <https://doi.org/10.1175/2010jas3541.1>, 2010.

Milbrandt, J. A. and Yau, M. K.: A multimoment bulk microphysics parameterization. Part I: Analysis of the role of the spectral shape parameter, *Journal of the Atmospheric Sciences*, 62, 3051-3064, <https://doi.org/10.1175/jas3534.1>, 2005.

515 Milbrandt, J. A., Morrison, H., Dawson II, D. T., and Paukert, M.: A triple-moment representation of ice in the Predicted Particle Properties (P3) microphysics scheme, *Journal of the Atmospheric Sciences*, 78, 439-458, <https://doi.org/10.1175/jas-d-20-0084.1>, 2021.

Mitchell, D. L.: Evolution of Snow-Size Spectra in Cyclonic Storms. Part II: Deviations from the Exponential Form, *J. Atmos. Sci.*, 48, 1885-1899, [https://doi.org/10.1175/1520-0469\(1991\)048<1885:EOSSII>2.0.CO;2](https://doi.org/10.1175/1520-0469(1991)048<1885:EOSSII>2.0.CO;2), 1991.

Mitchell, D. L., Rasch, P., Ivanova, D., McFarquhar, G., and Nousiainen, T.: Impact of small ice crystal assumptions on ice sedimentation rates in cirrus clouds and GCM simulations, *Geophys. Res. Lett.*, 35, L09806, <https://doi.org/10.1029/2008gl033552>, 2008.

520 Morrison, H. and Gettelman, A.: A new two-moment bulk stratiform cloud microphysics scheme in the community atmosphere model, version 3 (CAM3). Part I: Description and numerical tests, *Journal of Climate*, 21, 3642-3659, <https://doi.org/10.1175/2008jcli2105.1>, 2008.

Morrison, H. and Milbrandt, J. A.: Parameterization of cloud microphysics based on the prediction of bulk ice particle properties. Part I: Scheme description and idealized tests, *Journal of the Atmospheric Sciences*, 72, 287-311, <https://doi.org/10.1175/jas-d-14-0065.1>, 2015.

Morrison, H., Curry, J. A., and Khvorostyanov, V. I.: A new double-moment microphysics parameterization for application in cloud and climate models. Part I: Description, *Journal of the Atmospheric Sciences*, 62, 1665-1677, <https://doi.org/10.1175/jas3446.1>, 2005.

530 Morrison, H., van Lier-Walqui, M., Fridlind, A. M., Grabowski, W. W., Harrington, J. Y., Hoose, C., Korolev, A., Kumjian, M. R., Milbrandt, J. A., Pawlowska, H., Posselt, D. J., Prat, O. P., Reimel, K. J., Shima, S.-I., van Diedenhoven, B., and Xue, L.: Confronting the challenge of modeling cloud and precipitation microphysics, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS001689, <https://doi.org/10.1029/2019MS001689>, 2020.

Paukert, M., Fan, J., Rasch, P. J., Morrison, H., Milbrandt, J. A., Shpund, J., and Khain, A.: Three-moment representation of rain in a bulk microphysics model, *Journal of Advances in Modeling Earth Systems*, 11, 257-277, <https://doi.org/10.1029/2018MS001512>, 2019.

535 Proske, U., Ferrachat, S., Neubauer, D., Staab, M., and Lohmann, U.: Assessing the potential for simplification in global climate model cloud microphysics, *Atmos. Chem. Phys. Discuss.*, 2021, 1-40, <https://doi.org/10.5194/acp-2021-801>, 2021.

Salzmann, M., Ming, Y., Golaz, J. C., Ginoux, P. A., Morrison, H., Gettelman, A., Krämer, M., and Donner, L. J.: Two-  
540 moment bulk stratiform cloud microphysics in the GFDL AM3 GCM: description, evaluation, and sensitivity tests, *Atmos. Chem. Phys.*, 10, 8037-8064, <https://doi.org/10.5194/acp-10-8037-2010>, 2010.

Schumann, U., Mayer, B., Gierens, K., Unterstrasser, S., Jessberger, P., Petzold, A., Voigt, C., and Gayet, J.-F.: Effective radius of ice particles in cirrus and contrails, *Journal of the Atmospheric Sciences*, 68, 300-321, <https://doi.org/10.1175/2010jas3562.1>, 2011.

545 Sherwood, S. C., Bony, S., Boucher, O., Bretherton, C., Forster, P. M., Gregory, J. M., and Stevens, B.: Adjustments in the forcing-feedback framework for understanding climate change, *Bulletin of the American Meteorological Society*, 96, 217-228, <https://doi.org/10.1175/bams-d-13-00167.1>, 2015.

Shi, X., Liu, X., and Zhang, K.: Effects of pre-existing ice crystals on cirrus clouds and comparison between different ice nucleation parameterizations with the Community Atmosphere Model (CAM5), *Atmospheric Chemistry and Physics*, 15, 550 1503-1520, <https://doi.org/10.5194/acp-15-1503-2015>, 2015.

Spichtinger, P. and Gierens, K. M.: Modelling of cirrus clouds – Part 1a: Model description and validation, *Atmos. Chem. Phys.*, 9, 685-706, <https://doi.org/10.5194/acp-9-685-2009>, 2009.

Storelvmo, T., Kristjansson, J. E., Muri, H., Pfeffer, M., Barahona, D., and Nenes, A.: Cirrus cloud seeding has potential to cool climate, *Geophys. Res. Lett.*, 40, 178–182, <https://doi.org/10.1029/2012GL054201>, 2013.

555 Wang, M. and Penner, J. E.: Cirrus clouds in a global climate model with a statistical cirrus cloud scheme, *Atmos. Chem. Phys.*, 10, 5449-5474, <https://doi.org/10.5194/acp-10-5449-2010>, 2010.

Wang, M., Liu, X., Zhang, K., and Comstock, J. M.: Aerosol effects on cirrus through ice nucleation in the Community Atmosphere Model CAM5 with a statistical cirrus scheme, *Journal of Advances in Modeling Earth Systems*, 6, 756-776, <https://doi.org/10.1002/2014MS000339>, 2014.

560 Wyser, K.: The effective radius in ice clouds, *Journal of Climate*, 11, 1793-1802, [https://doi.org/10.1175/1520-0442\(1998\)011<1793:Teriic>2.0.Co;2](https://doi.org/10.1175/1520-0442(1998)011<1793:Teriic>2.0.Co;2), 1998.

Zhang, G. J. and McFarlane, N. A.: Sensitivity of climate simulations to the parameterization of cumulus convection in the Canadian climate centre general circulation model, *Atmosphere-Ocean*, 33, 407-446, <https://doi.org/10.1080/07055900.1995.9649539>, 1995.

565 Zhang, G. J., Kiehl, J. T., and Rasch, P. J.: Response of climate simulation to a new convective parameterization in the National Center for Atmospheric Research Community Climate Model (CCM3), *Journal of Climate*, 11, 2097-2115, [https://doi.org/10.1175/1520-0442\(1998\)011<2097:Rocsta>2.0.Co;2](https://doi.org/10.1175/1520-0442(1998)011<2097:Rocsta>2.0.Co;2), 1998.

Zhao, X., Lin, Y., Peng, Y., Wang, B., Morrison, H., and Gettelman, A.: A single ice approach using varying ice particle properties in global climate model microphysics, *Journal of Advances in Modeling Earth Systems*, 9, 2138-2157, 570 <https://doi.org/10.1002/2017MS000952>, 2017.

Zhou, C., Zelinka, M. D., and Klein, S. A.: Impact of decadal cloud variations on the Earth's energy budget, *Nature Geosci*, 9, 871-874, <https://doi.org/10.1038/ngeo2828>, 2016.