

1 **Assessment of Climate Biases in OpenIFS Version 43R3 across** 2 **Model Horizontal Resolutions and Time Steps**

3
4 Abhishek Savita¹, Joakim Kjellsson^{1,2}, Robin Pilch Kedzierski^{3,1}, Mojib Latif^{1,2}, Tabea Rahm^{1,2},
5 Sebastian Wahl¹ and Wonsun Park^{4,5}

6
7 ¹GEOMAR Helmholtz Centre for Ocean Research Kiel, Kiel, Germany

8 ²Faculty of Mathematics and Natural Sciences, Christian Albrechts University of Kiel, Kiel, Germany

9 ³Meteorological Institute, Universität Hamburg, Hamburg, Germany

10 ⁴Center for Climate Physics, Institute for Basic Science (IBS), Busan, Republic of Korea

11 ⁵Department of Climate System, Pusan National University, Busan, Republic of Korea

12 *Correspondence to:* Abhishek Savita (asavita@geomar.de)

13 **Abstract.**

14 We examine the impact of horizontal resolution and model time step on climate of the OpenIFS version 43R3 atmosphere
15 general circulation model. A series of simulations for the period 1979-2019 are conducted with various horizontal resolutions
16 (i.e., ~100, ~50, and ~25 km) while maintaining the same time step (i.e., 15 minutes) and using different time steps (i.e., 60,
17 30 and 15 minutes) at 100 km horizontal resolution. We find that the surface zonal wind bias reduces significantly over certain
18 regions such as the Southern Ocean, the Northern Hemisphere mid-latitudes, and in tropical and subtropical regions at high
19 horizontal resolution (i.e., ~25 km). Similar improvement is evident too when using a coarse resolution model (~100 km) with
20 a smaller time step (i.e., 30 and 15 minutes). We also find improvements in Rossby wave amplitude and phase speed as well
21 as weather regime patterns when a smaller time step or higher horizontal resolution is used. The improvement in the wind bias
22 when using the shorter time step is mostly due to an increase in shallow and mid-level convection that enhances vertical mixing
23 in the lower troposphere. The enhanced mixing allows frictional effects to influence a deeper layer and reduces wind and wind
24 speed throughout the troposphere. However, precipitation biases generally increase with higher horizontal resolution or smaller
25 time step, whereas the surface-air temperature bias exhibits a small improvement over North America and the Eastern Eurasian
26 continent. We argue that the bias improvement in the highest horizontal resolution (i.e., ~25 km) configuration benefits from
27 a combination of both the enhanced horizontal resolution and the shorter time step. In summary, we demonstrate that by
28 reducing the time step in the coarse resolution (~100 km) OpenIFS model, one can alleviate some climate biases at a lower
29 cost than by increasing the horizontal resolution.

30

31

32

33

34

36 1. Introduction

37 In the last few decades, Atmospheric-Ocean General Circulation Model (AOGCM) simulations from the Coupled Model
38 Intercomparison Project (CMIP) have been widely used to study the internal climate variability and the climate response to
39 external forcing such as increasing atmospheric greenhouse gas concentrations causing global warming. These simulations,
40 however, suffer from long-standing biases (**Bayr et al., 2018; Flato et al., 2014; Gates et al., 1999; Kim et al., 2014; Zhou**
41 **et al., 2020**), which lead to significant uncertainties in short-term and long-term climate projections and potential ecosystem
42 impacts (**Athanasiadis et al., 2022; Couldrey et al., 2021; Meehl and Teng, 2014; Meng et al., 2022**). These biases can
43 arise from a variety of sources, including inaccurate representation of physical processes, poor initialization of model
44 conditions, or inadequate representation of the Earth's topography and land cover.

45

46 Simulations using Atmospheric General Circulation Models (AGCMs) from the Atmosphere Model Intercomparison Project
47 (AMIP), a part of CMIP, are used to study the internal variability of the atmosphere. The AGCMs are less complex than the
48 AOGCMs as the former are constrained by observed Sea Surface Temperature (SST) and Sea Ice Concentration (SIC). Despite
49 being constrained by the observations, the AGCMs also exhibit biases (e.g., **Gates et al., 1999**), and some of these biases have
50 persisted for over several phases of AMIP (**He and Zhou, 2014**). The biases in AGCMs are largely due to the fact that many
51 unresolved processes, such as atmospheric convection, precipitation, clouds, cloud-microphysical and aerosol processes,
52 boundary layer processes, and interactions between the land surface and hydrologic processes, have to be included in a
53 parameterized form in the coarse resolution model (**Ma et al., 2022**). The treatment of unresolved gravity waves and the
54 relatively large model time step also contribute to the biases in AGCMs (**Flato et al., 2014; Gates et al., 1999**).

55

56 Recently, **Liu et al. (2022)** analyzed AOGCM simulations and reported that increasing the horizontal resolution of the ocean
57 component one can reduce SST and precipitation biases in the equatorial Pacific, whereas increasing the horizontal resolution
58 of the atmospheric component did not have the same effect. However, other studies found that a high-horizontal resolution
59 atmosphere model better simulates the main features of tropical precipitation, tropical atmospheric circulation, and extra-
60 tropical cyclones while increasing from 125 km to 40 km horizontal resolution with relatively small improvements for further
61 enhanced horizontal resolution (**Branković and Gregory, 2001; Jung et al., 2012; Williamson et al., 1995**). Similarly,
62 **Roberts et al. (2018)** found that there was not much improvement in the Integrated Forecasting System (IFS) from the
63 European Centre for Medium-Range Weather Forecasting (ECMWF) when increasing horizontal resolution from 50 to 25 km.

64

65 **Jung et al. (2012)** and **Roberts et al. (2018)** demonstrated a time step sensitivity in the coarse and high horizontal resolution
66 model simulations using the IFS model. **Jung et al. (2012)** found that the precipitation and wind biases were reduced at the

67 coarse horizontal resolution when shortening the model time step from 60 to 15 minutes. **Roberts et al. (2018)** did not find
68 such a significant improvement when reducing the model time step from 20 to 15 minutes in their high-resolution (~25 km)
69 configuration. However, both studies did not investigate the model's sensitivity to changes in the model time step in detail.
70 While the semi-implicit semi-Lagrangian scheme, as used in OpenIFS, is unconditionally stable and the time step can be
71 chosen to be very long, a shorter time step generally leads to a decrease in truncation error in the finite differences and thus a
72 more accurate representation of the model dynamics. The physics parameterisations, which are computed independently of
73 each other in OpenIFS, also benefit from a shorter time step as it will allow the various parameterisations to be coupled at a
74 higher frequency (**Beljaars et al., 2018**). However, model parameters for e.g., convection or diffusion may be tuned for a
75 specific time step and shortening the time step can therefore, in some cases, increase model error. Hence, a shorter model time
76 step is expected to reduce biases in model dynamics, e.g., winds, while the results for parameterised processes, e.g.,
77 precipitation, may be mixed.

78

79 In the research community, there is no standard definition for coarse horizontal resolution, as one study considered 200 km as
80 a coarse resolution (~2°) configuration (**Branković and Gregory, 2001**), whereas another study considered 50 km (0.5°) as a
81 coarse resolution (**Roberts et al., 2018**). Likewise, there is no unique rule for setting the model time step dependent on model
82 resolution. Groups using either the IFS or OpenIFS model at horizontal resolutions of ~100 km have used a relatively long
83 time step of 1 hour (**Hazeleger et al., 2012; Kjellsson et al., 2020; Streffing et al., 2022**) or 45 minutes (**Döscher et al.,**
84 **2022**), while other groups using the ARPEGE-Climat with a similar dynamical core use 15 minutes (**Voldoire et al., 2019**).
85 The model's horizontal resolution and time steps are rather chosen on what can be afforded computationally, and their relative
86 contributions to biases in the model's climate are not well documented.

87

88 In this study, we systematically investigate the sensitivity of the OpenIFS model version 43r3v2 to the model time step and
89 horizontal resolution. We mostly focus on the surface zonal winds since they play a crucial role for the ocean circulation in
90 the AOGCMs. We also study the representation of the synoptic-scale variability such as Rossby waves and weather regimes.
91 The paper is structured as follows: section 2 describes the model, experimental design, data and methodology; section 3
92 describes the results and section 4 summarizes the conclusions of this work.

93 **2. Model, Experimental design, Data and Methodology**

94 We conducted a series of experiments with the OpenIFS model. The OpenIFS model is derived from the Integrated Forecasting
95 System at the European Centre for Medium-range Weather Forecasting (ECMWF-IFS) cycle 43 release 3 (43r3). The
96 dynamical core is the same as ECMWF-IFS that uses a two-time-level semi-implicit time stepping with semi-Lagrangian
97 advection (**Temperton et al., 2001**) on a reduced Gaussian grid with a hybrid-sigma vertical coordinate (**Simmons and**
98 **Burridge, 1981**). Likewise, the OpenIFS uses the same model physics as the ECMWF-IFS (cf. **Forbes and Tompkins, 2011**;

99 **Hogan and Bozzo, 2018; Tiedtke, 1993**) but does not include the tangent-linear code or 4D-VAR capabilities. Our version,
100 OpenIFS, is similar to cy43r1 used in **Roberts et al., (2018)**, with the main difference being the new radiation scheme, ecRad
101 (**Hogan and Bozzo, 2018**), introduced in cy43r3.

102

103 Our study is partly motivated by evaluating the suitability of various OpenIFS configurations for coupled climate simulations
104 with FOCI-OpenIFS (**Kjellsson et al., 2020**) with an atmosphere horizontal resolution higher than that of ECHAM6 Tq63/N48
105 (~200km) in FOCI (**Matthes et al., 2020**). Our choices thus fall on three different horizontal resolutions: a low-resolution
106 (T_{co95} , ~100 km), a medium-resolution (T_{co199} , ~50 km), and a high resolution (T_{co399} , ~25 km). The T_{co95} grid is the lowest
107 acceptable resolution since the supported lower-resolution grids, e.g., T195/N48 and Tq42/F32, are either similar to Tq63 in
108 ECHAM6 or coarser. The T_{co399} grid was chosen as an upper limit of what is computationally feasible for AMIP integrations
109 and century-long coupled integrations given our computer resources. All the configurations share the same vertical L91 grid.
110 We did not modify any other model parameters when changing the model horizontal resolutions or model time steps, but we
111 note that some parameters such as launch momentum flux for non-orographic gravity waves scales with resolution in the
112 model. We performed 5 experiments in total (**Table 1**). For simplicity, we now refer now OpenIFS as OIFS in the rest of the
113 sections. We note that exploring the effect of different time steps was only done for the lowest horizontal resolution (T_{co95} ,
114 ~100 km). We did not run similar sensitivity experiments for the high-resolution configuration (T_{co399} , ~25 km) for two
115 reasons. First, the high-resolution configuration is very computationally expensive. Second, it was deemed more important to
116 explore time step sensitivity at low resolution since this configuration (and other similar resolutions) is often used for coupled
117 climate simulations. The potential time-step sensitivity at high-resolution is discussed in the Discussion section.

118

119

120 The lower boundary conditions, i.e., SST and SIC, are taken the from Atmospheric Model Intercomparison Project (AMIP)
121 version 1.1.6 (**Eyring et al., 2016; Taylor et al., 2012**), which are available as monthly means on a $1^\circ \times 1^\circ$ horizontal grid. The
122 external forcing is identical to that used in the CMIP6 AMIP simulations except for the aerosol and ozone concentrations,
123 which are taken from monthly mean climatology. SST and SIC are interpolated from monthly to daily frequency and from
124 $1^\circ \times 1^\circ$ horizontal resolution to the OIFS horizontal grid using bilinear interpolation. All the simulations are run for the period
125 1979–2019. We extend the simulations beyond the AMIP protocol for 1979-2014 up to 2019 by using SST and SIC from the
126 ERA5 reanalysis and the Shared Socioeconomic Pathways 5 (SSP5-8.5) emission scenario. Ozone concentrations are taken
127 from monthly photochemical equilibrium state and aerosol concentrations from monthly CAMS climatology of 11 species.

128

129 Amplitude and phase speed of Rossby wave were computed by performing a Fourier decomposition analysis on 300 hPa daily
130 meridional winds. First, we interpolated both ERA5 and OIFS simulation datasets onto a $2.5^\circ \times 2.5^\circ$ grid using bilinear
131 interpolation. We then applied the Fourier decomposition analysis to determine amplitude and position for each Rossby wave
132 number at each latitude as a function of time. Phase speed is computed as the difference in the daily position of each wave,

133 and stored at the midpoints in the time dimension. For consistency, wave amplitudes are interpolated to the midpoints in time
134 as well. Lastly, seasonal averages are computed from the daily data for the boreal and austral winter seasons over the time
135 period 1979–2019. In the case of phase speed, it is weighed by the corresponding daily (midpoint) amplitude squared when
136 computing the seasonal averages in order to account for the impact of higher-amplitude events. The results are presented in
137 wavenumber-latitude diagrams similar to previous studies (e.g., [Pilch Kedzierski et al., 2020](#); [Wolf and Wirth, 2017](#)). Our
138 wavenumber-latitude analysis is not directly comparable to both studies mentioned above, because we did not apply any high-
139 pass filtering in time before the Fourier decomposition. While the previous literature had similar diagrams with varying
140 measures of wave amplitude, our detailed analysis of phase speed in such a manner is novel in literature to our knowledge and
141 a strong addition as a model performance diagnostic.

142

143 The Weather Regime Patterns (WRPs) were calculated using daily 500-hPa geopotential height (z500) anomalies over the
144 Euro-Atlantic region (30°–90 °N, 80 °W–40 °E) for the boreal winter season during the period 1979-2019. The daily z500
145 daily anomalies were computed by subtracting the daily climatology smoothed by a 20-day running mean from the raw z500
146 data. We calculated the first four Empirical Orthogonal Functions (EOFs) from the ERA5 dataset. In the next step, the OIFS-
147 simulated z500 anomalies were projected on the ERA5 EOFs to obtain Pseudo-Principal Components (Pseudo-PCs). We then
148 applied a K-means clustering algorithm to the individual model pseudo-PCs and observation PCs using four clusters. We chose
149 four clusters because these give the most of the significant clustering. Spatial WRPs are obtained by compositing over all daily
150 z500 anomalies for each regime. More information about the methodology can be found in [Fabiano et al. \(2020\)](#), section 3.1.
151 In order to evaluate the WRPs simulated by the OIFS across configurations more quantitatively, we have additionally estimated
152 the Pearson’s Pattern Correlation Coefficient (PCC) between the WRPs identified in the model and ERA5.

153

154 We compare the climate of OIFS to observational and reanalysis datasets. Precipitation is validated against the Global
155 Precipitation Climatology Project ([GPCP](#); [Huffman et al., 1997](#)), and the surface air temperature (SAT) against the
156 CRUTEM4 ([Harris et al., 2014](#); [Osborn and Jones, 2014](#)). We have used the ERA5 reanalysis ([Hersbach et al., 2020](#)) to
157 evaluate 10-meter surface wind as well as the zonal wind at 300 hPa for the Rossby wave analysis. We use z500 from ERA5
158 to validate the OIFS-simulated weather regimes. We also compare our results with the MERRA2 reanalysis ([Gelaro et al.,
159 2017](#)) and find similar results. Therefore, the comparison with MERRA2 is not shown. The bootstrapping (in total 2000
160 iterations) method is used to compute the 95% confidence interval for the RMSE and the WRPs correlation.

161 3. Results

162 3.1.1 Global and regional surface bias and deriving processes

163 The annual mean 10m zonal wind (surface wind hereafter) bias during the period 1979–2019 for the different OIFS
164 configurations is shown in [Fig. 1](#). We find that the OIFS-LRA-1h configuration has a large surface wind bias over most of the

165 world ocean, with positive biases in the mid-latitudes (the Southern Ocean, North Atlantic and North Pacific) and negative
166 surface wind biases over the tropical oceans (Tropical Pacific, Tropical Indian and Atlantic Ocean) (**Fig. 1b**). Thus, the OIFS-
167 LRA-1h configuration simulates too strong surface westerly winds (and wind speed) over the mid-latitude oceans, which, if
168 coupled to an ocean model, may cause biases in upper-ocean mixing and oceanic uptake of heat and carbon.

169

170 The surface wind bias in the OIFS-HRA-15m configuration is reduced significantly (**Fig. 1f**) over most of the world ocean
171 compared to the OIFS-LRA-1h configuration (**Fig. 1b**), indicating that increasing the horizontal resolution from 100 km to 25
172 km and shortening the time step from 1h to 15-min improves the representation of the surface winds. The surface wind bias
173 also significantly reduces everywhere in the OIFS-MRA-15m configuration (**Fig. 1e**) compared to the OIFS-LRA-1h
174 configuration (**Fig. 1b**). The surface wind bias in OIFS-MRA-15m is larger than that in the OIFS-HRA-15m configuration but
175 smaller than that in the OIFS-LRA-1h configuration. Similar conclusions are obtained by performing Root Mean Square Error
176 (RMSE) analysis, which shows that the OIFS-HRA-15m configuration has the lowest annual and global mean RMSE of
177 surface wind while the OIFS-LRA-1h configuration has the highest RMSE (**Fig. 2a, black line**). Though we have found a
178 significant improvement in the wind bias in the OIFA-HRA-15m configuration, it is not clear yet whether the improvement is
179 due to the increased horizontal resolution or the shorter time step.

180

181 Surface wind bias is also reduced in both the OIFS-LRA-30m (**Fig. 1c**) and OIFS-LRA-15m (**Fig. 1d**) configurations compared
182 to the OIFS-LRA-1h configuration (**Fig. 1b**), and the bias improvement is mostly observed at the same places as in the OIFS-
183 HRA-15m configuration (**Fig. 1f**). The surface-wind bias improvement is similar in the OIFS-LRA-30m and OIFS-LRA-15
184 configurations, except over the North Pacific and Southern Ocean where the OIFS-LRA-15m configuration has a smaller wind
185 bias than the OIFS-LRA-30m configuration. However, we have not seen a large difference between the OIFS-LRA-30m and
186 OIFS-LRA-15 configurations in the global average RMSE analysis (**Fig. 2a**).

187

188 The surface-wind bias improvement in the OIFS-HRA-15m and OIFS-LRA-15m configurations not only exists in the annual
189 average but also in boreal winter (DJF) and summer (JJA) (**Fig. 2a blue and red lines**, respectively). Our results are consistent
190 with **Jung et al. (2012)**, as they found a reduction in wind bias in the tropical Pacific region when they shortened the time step
191 in their coarse resolution configuration. However, this study and the **Jung et al. (2012)** study are not consistent with that of
192 **Robert et al. (2020)** who did not find much time-step sensitivity. We speculate that in **Robert et al. (2020)**, the reduction
193 from 20 to 15 minutes in their high horizontal resolution (25 km) may be too small. Alternatively, the 20-minute time step
194 could be the optimal time step for the 25 km configuration.

195

196 The surface wind bias in the OIFS-HRA-15m and OIFS-LRA-15m configurations looks similar in pattern, but they differ in
197 magnitude. The OIFS-HRA-15m configuration has a smaller bias in the North Pacific, Peru upwelling and Agulhas Bank
198 regions compared to the OIFS-LRA-15m configuration. We hypothesize that the reduction in surface-wind bias in the OIFS-

199 HRA-15m configuration (**Fig. 1f**) compared to the OIFS-LRA-1h configuration (**Fig. 1b**) is a combination of the enhanced
200 horizontal resolution and shorter time step. The improvement in the OIFS-HRA-15m configuration (**Fig. 1f**) compared to the
201 OIFS-LRA-15m configuration (**Fig. 1d**) is due to only the enhanced horizontal resolution as both configurations use the same
202 time step.

203

204 The zonal-wind bias improvement in the OIFS-LRA-15m is further explored using the online zonal wind tendencies from
205 OIFS which are split into dynamics and physics that includes turbulent diffusion, gravity-wave drag and convection:

206

$$207 \quad du/dt = du/dt_{Dyn} + du/dt_{Turb} + du/dt_{Gwd} + du/dt_{Conv} \quad (1)$$

208

209 where du/dt_{Dyn} is the sum of the tendencies from advection, pressure gradient and Coriolis force, du/dt_{Turb} includes tendencies
210 from surface processes, vertical diffusion and orography drag, du/dt_{Gwd} includes gravity-wave drag and non-orographic drag,
211 and du/dt_{Conv} is the tendency from convection. The individual tendencies on the right-hand side of equation (1) are referred to
212 as Dyn, Turb, Gwd and Conv, respectively. They were stored for each model level in the OIFS-LRA-1h and OIFS-LRA-15m
213 configurations. The lowest model level is at 10m height (assuming surface pressure of 1013hPa), so the 10m wind will behave
214 very similarly to the wind at level k=91.

215

216 The averaged zonal wind and zonal wind tendencies over the Southern Ocean (40° S – 60° S and all longitude) in the OIFS-
217 LRA-1h and OIFS-LRA-15m configurations are shown in **Fig. 3a & b**, respectively. The zonal wind tendency (i.e., du/dt) in
218 both OIFS-LRA-15m and OIFS-LRA-1h configurations is very small (~ 2 to 0.04 m/s^{-1}) compared to the other processes (**Fig.**
219 **3b, black lines**). Conv provides westward acceleration between the 700 and 900 hPa pressure levels and eastward acceleration
220 below, indicating a downward transport of westward momentum. Dyn acts to accelerate the flow eastward from 700 hPa and
221 below, likely via momentum advection, pressure-gradient and Coriolis forces, while Turb has the opposite effect, likely via
222 surface friction and vertical mixing processes. In the OIFS-LRA-15m configuration, we find a similar balance as in the OIFS-
223 LRA-1h, but the westward acceleration above and eastward acceleration below is enhanced by Conv, likely by increased
224 downward momentum transport, in agreement with the increased shallow and mid-level convection (**Fig. 3d**). The vertical
225 momentum mixing by shallow and mid-level convection reduces the vertical wind shear, making the westerly winds more
226 barotropic. As a result, the westerly winds weaken throughout the troposphere and even in the stratosphere (**Fig. 3a**). We note
227 similar changes in the Northern Hemisphere mid-latitudes, suggesting similar mechanisms are acting. Gwd has a negligible
228 role for the winds in the lower stratosphere and troposphere, and the Gwd term does not appear sensitive to model time step
229 (**Fig. 3b, orange lines**).

230

231 **Fig. 3c** shows the zonal average of the zonal wind tendencies at the lowest level of the model, as a function of the latitude. In
232 the OIFS-LRA-1h configuration, Conv and Dyn accelerate the surface westerly wind in the mid-latitudes ($\sim 40^\circ \text{ N}$ to $\sim 60^\circ \text{ N}$)

233 in both hemispheres, and these westerly winds are partly balanced by Turb (**Fig. 3c, solid lines**). Dyn has a larger contribution
234 to accelerating the surface westerly winds than Conv (**Fig. 3c, solid lines**). However, the Conv contribution is enhanced in the
235 OIFS-LRA-15m configuration, while the Dyn contribution reduces (**Fig. 3c, dashed lines**). We also find that the contribution
236 to slowing the westerly wind is reduced by Turb in the OIFS-LRA-15m configuration (**Fig. 3c, dashed lines**).

237

238 It is also noteworthy that the individual wind tendencies are significantly larger in the Southern Hemisphere (and Southern
239 Ocean) than in the Northern Hemisphere (Fig. 3c). The larger magnitudes of the tendencies over the Southern Ocean compared
240 to similar latitudes on the Northern Hemisphere is likely due to the Southern Hemisphere having fewer continents in
241 midlatitudes than the Northern Hemisphere and thus the surface is less rough and allows for stronger winds. In the low latitudes,
242 both Dyn and Conv contribute to accelerating the easterly winds, which is partly balanced by Turb in the OIFS-LRA-1h
243 configuration (**Fig. 3c, solid lines**). There are no discernible changes in Conv, Dyn or Turb from OIFS-LRA-1h to OIFS-LRA-
244 15m, indicating that the tropical surface winds are relatively insensitive to model time step (**Fig. 3c, dashed lines**).

245

246 In addition to surface wind, we also investigated the sensitivity of model time step and horizontal resolution for SAT and
247 precipitation. The RMSEs for SAT and precipitation are shown in Figure 2b and 2c, respectively. We find that the OIFS-HRA-
248 15m has the lowest SAT RMSE of all model experiments in both annual and seasonal means, although the RMSE difference
249 across the configurations is not significant (**Fig. 2b**). The reduced SAT RMSE in OIFS-HRA-15m configuration is primarily
250 due to the lowered SAT bias over North America and the eastern part of Russia. Compared to the OIFS-LRA-1h, the SAT
251 RMSE decreases with increased horizontal resolution (OIFS-HRA-15m and OIFS-MRA-15m), and there is no notable
252 improvement when shortened the time step (OIFS-LRA-30m and OIFS-LRA-15m) (Fig 2b).

253

254 We have computed the SAT and precipitation biases with a 3-point smoothing, i.e., approximately 3x3 degree spatial
255 smoothing, which eliminates the wiggles near steep topography arising from the Gibbs' phenomenon in the model spectral
256 fields. We find that smoothing the fields does not change the main result that precipitation biases increase with shorter time
257 step in Tco95 and then decreases somewhat with higher horizontal resolution. Hence, the wiggles are not the main source of
258 precipitation biases and their presence does not impact the findings in this study.

259

260 The OIFS-LRA-1h experiment exhibits the lowest precipitation RMSE of all experiments, with RMSE increasing with shorter
261 time step (OIFS-LRA-15m) and increased horizontal resolution (OIFS-HRA-15m) for both the annual and boreal winter means
262 (**Fig. 2c, black and blue lines**). The patterns of regional precipitation biases are similar across the configurations in the mid-
263 and high-latitudes, whereas the precipitation biases increase in the tropics at the high horizontal resolution or in the smaller
264 time step configuration (not shown). The results suggest that some of the cloud and/or convection parameters may be dependent
265 on resolution or time step and need retuning for each configuration.

Deleted: contribute

Deleted: more

268 3.1.2 Wind and temperature bias in upper-atmosphere

269 We examined the zonal mean u wind bias at different model levels, and it is shown in Figures 4. We find that zonal mean u
270 wind bias over the tropical region (40°S and 40°N) is positive and independent of model horizontal resolution and model time
271 step (Fig. 4b-f). The OIFS-HRA-15m configuration has a relatively large negative bias in the Northern Hemisphere compared
272 to the other configurations. The OIFS-LRA-15m and OIFS-HRA-15m zonal mean u wind bias is similar to that in Robert et
273 al. (2018). However, the zonal mean u wind bias in the Southern Hemisphere is not consistent across horizontal resolution or
274 model time step. The zonal mean u wind bias in midlatitudes (i.e., 70°S to 50°S) is positive and large in the OIFS-LRA-1h
275 configuration and reduces throughout the pressure levels by shortening the model time step in the coarse resolution OpenIFS
276 configuration (i.e., OIFS-LRA-30m and OIFS-LRA-15m). Whereas, the negative zonal wind bias south 70°S in the coarse
277 resolution configuration is consistent across the different time steps (Fig. 3b-d). It is also interesting to note that both OIFS-
278 MRA-15m and OIFS-HRA-15m configurations exhibit a negative bias over the Southern Ocean (SO) at most of the pressure
279 levels, which is not seen in the standard OIFS-LRA-1h, nor in the OIFS-LRA-30m or OIFS-LRA-15m configurations. Overall,
280 we conclude that by reducing the model time step in the coarse resolution configuration, we improve winds not only at the
281 surface but also at higher model levels mostly over the SO. A similar conclusion does not hold for the OIFS-MRA-15m and
282 OIFS-HRA-15m configurations, as both suffer from large negative bias over the SO.

283

284 We also examined the zonal mean temperature bias at different pressure levels. We find a cold bias (1.5 to 6 °C) in the
285 troposphere and lower stratosphere, and a warm bias (1.5 to 6 °C) above the stratosphere across the configurations (Figure not
286 shown). This indicates that OpenIFS simulations (independent of model time step and horizontal resolution) are colder than
287 observations in the lower stratosphere and warmer above. The cold bias in the lower stratosphere is larger in the high resolution
288 (i.e., OIFS-HRA-15m), and the warm bias above the stratosphere is smaller compared to the other configurations. Robert et
289 al. (2018) noticed a similar zonal mean temperature bias and speculated that the zonal mean temperature bias is linked with
290 the sensitivity of spurious mixing due to convection and diffusion.

291 3.2 Rossby wave analysis

292 **Fig. 5** shows the Rossby wave amplitude (gray and black contours) for ERA5 and the individual OIFS simulations for the
293 boreal winter (**Fig. 5A, DJF, Northern Hemisphere; NH**) and austral winter (**Fig. 5B, JJA, Southern Hemisphere; SH**).
294 The color in **Fig. 5** denotes the wave amplitude bias relative to ERA5 (model – ERA5). We focus only on those wave numbers
295 and latitudes that have the highest wave amplitude, because these waves explain most of the variability. The region where the
296 wave amplitude is larger than 5 ms⁻¹ is termed “core region”, which mostly covers the area that is occupied by the thick black
297 contours in **Fig. 5**. In DJF (NH), at north of 70° N, the Rossby wave numbers k=1 and k=2 have the largest amplitude in ERA5
298 whereas at the mid-latitudes (30° N to 60° N), the wave numbers between about k=3 and k=9 have large amplitude with the
299 largest amplitude amounting to 8 ms⁻¹ at about 40° N for the wave number k=6 (**Fig. 5Aa**). During JJA (SH), the wave

300 amplitude is located in a similar core region (**Fig. 5Ba**) as that in DJF (NH). The amplitude is largest south of 70° S for the
301 wave numbers $k=1$ and $k=2$ whereas at the mid-latitudes (45° S to 65° S), the wave numbers between about $k=3$ to 5 have large
302 amplitude with the largest amplitude amounting to 9 ms^{-1} is found at 57.5° S for the wave number $k=4$ (**Fig. 5Ba**).

303

304 In DJF (NH) the OIFS-LRA-1h configuration exhibits a positive bias of $\sim 1 \text{ ms}^{-1}$ in Rossby wave amplitude (i.e., the waves
305 amplitude bias in OIFS-LRA-1h is larger than the ERA-5) in the core region, in particular for wave numbers $k=3-8$ at latitudes
306 between 25° N to 55° N and a negative bias at latitudes between 60° N to 80° N for waver number 2 (**Fig. 5Af**). The wave
307 amplitude biases around the core region in OIFS-LRA-1h in the midlatitudes (20° N to 40° N) are small (~ 0.2) for the higher
308 wave numbers and get better with a shorter time step configuration (OIFS-LRA-15m).

309

310 The Rossby wave amplitude biases in the OIFS-HRA-15m configuration are strongly reduced compared to the OIFS-LRA-1h
311 configuration over the core region (**Fig. 5Ab and 5Af**). The Rossby wave amplitude bias reduction in the OIFS-MRA-15m
312 configuration is mostly similar to that in the OIFS-HRA-15m configuration except for the wave number $k=7$ at 45° N, where
313 the wave amplitude bias is larger in the OIFS-HRA-15m configuration (**Fig. 5Ab and 5Ac**). The OIFS-HRA-15 m and OIFS-
314 MRA-15m configurations also exhibit a positive bias for wave number 2 at high-latitudes 60° N to 80° N. The OIFS-MRA-
315 15m configuration also show a negative bias for the wave number 3 at latitudes between 60° N to 65° N in the core region,
316 which is not present in the other configurations. The OIFS-HRA-15m and OIFS-MRA-15m configurations show similar bias
317 around the core region as in the OIFS-LRA-1h configuration, i.e., high resolution and OIFS-LRA-1h configurations
318 overestimate wave amplitudes for the higher wave numbers. The Rossby wave amplitude biases are progressively reduced
319 from the OIFS-LRA-1h configuration to the OIFS-LRA-30m and OIFS-LRA-15m configurations (**Fig. 5Ad-Af**), indicating a
320 sensitivity of model bias to the time step. The wave amplitude bias for wave number $k=7$ at 45° N exists in all the
321 configurations, and it is smaller in the OIFS-LRA-15m and OIFS-MRA-15m configuration than in the other configurations.
322 Overall, both OIFS-LRA-15m and OIFS-HRA-15m configurations are able to reproduce the observed Rossby-wave
323 amplitudes in DJF (NH) better than OIFS-LRA-1h.

324

325 In JJA (SH), the Rossby wave amplitude bias in the core region is smaller than in DJF (NH) for all the configurations (**Fig.**
326 **5A and 5B**). OIFS-LRA-1h exhibits a positive bias of $\sim 0.5 \text{ ms}^{-1}$ in JJA (SH) for the wave number $k=2$ at latitude between $\sim 50^\circ$
327 S and $\sim 62.5^\circ$ S and for wave numbers $k=4$ to 5 between 30° S and 40° S (**Fig. 5Bf**). The OIFS-LRA-30m configuration shows
328 a positive bias for the wave number $k=2$ to 5 at latitudes between 40° S and 70° S, which is larger than other configurations.

329 The OIFS-HRA-15m and OIFS-MRA-15m configurations exhibits a positive bias $\sim 0.5 \text{ ms}^{-1}$ around the core region and
330 latitude 50° S to 70°S, which does not exist in the other coarse resolution configurations (**Fig. 5Bb-Bf**). The Rossby wave
331 amplitude biases around the core region at the midlatitudes in the high-resolution simulations are consistent and large in the
332 SH than the NH (Fig. 5Ab-c and 5Bb-c).

333

334 We also analyze the phase speed of Rossby waves for ERA5 and across the OIFS' configurations for DJF (NH) and JJA (SH)
335 seasons (**Fig. 6**). In the ERA5 dataset (**Fig. 6Aa**), the Rossby wave phase speed is positive (i.e., eastward moving, solid contour)
336 for wave numbers greater than 2 (i.e., $k > 2$) at most latitudes. The wave numbers $k=1$ to 2 have a positive wave phase speed
337 from the equator to 55° N and a negative wave phase speed (i.e., westward moving, dashed contours) between 60° N and 80°
338 N in DJF (NH) (**Fig. 6Aa**). The maximum phase speed is found at wave number $k=8$ at 40° N, while the minimum is found at
339 wave number $k=1$ at 60° N (**Fig. 6Aa**). In JJA (SH) (**Fig. 6Ba**), the wave phase speeds are mostly positive and large for all the
340 wave numbers and at each latitude, with the maximum phase speed is observed for the wave numbers between $k=6$ and $k=8$
341 and latitudes between 40° S and 60° S, and these waves are moving faster than that in DJF (NH).

342

343 The OIFS-LRA-1h configuration suffers from positive phase speed bias for wave numbers $k=4$ to 8 at latitudes between 42.5°
344 N and 60° N, i.e., waves move faster eastward than in ERA5, and the bias is larger than 1 ms^{-1} . The bias of $\sim 1 \text{ ms}^{-1}$ for wave
345 number $k = 6$ to 8 at 40° N and 60° N is of particular concern as it is near the maximum wave amplitudes in DJF (**Fig. 6Af**).
346 In general, phase speed biases in the OIFS-LRA-1h configuration are strongly reduced as either horizontal resolution is
347 increased or time step is shortened (**Fig. 6Ab-5Af**). In JJA (SH), the OIFS-LRA-1h configuration exhibits a very large (between
348 ~ 1.5 - 2 ms^{-1}) Rossby wave phase speed bias for most of the wave numbers, which is largest for the wave numbers $k=2$ to 8
349 between 15° S to 55° S (**Fig. 6Bf**). Large biases can be found between 15° S and 25° S ($\sim 1.5 \text{ ms}^{-1}$) for most of the wave
350 numbers, but the wave activity is low there (**Fig. 6Bf**). The large phase speed biases are strongly reduced in the OIFS-LRA-
351 30m and OIFS-LRA-15m configurations (**Fig. 6Bd-Bf**), indicating a strong sensitivity to the reduced biases in mean winds
352 and wind speeds (**Fig. 1**). Overall, the Rossby wave speed bias in the OIFS-HRA-15m configuration is smaller than in the
353 OIFS-LRA-1h configuration (**Fig. 6Bb and 6Bf**). However, we note that both the OIFS-MRA-15m and OIFS-HRA-15m
354 configurations exhibit negative biases south of 55° S for wave numbers $k= 1$ to 5, that is, the eastward moving waves are
355 slower than in the ERA5 (**Fig. 6Bb**).

356

357 The wave phase speed analysis reveals a clear improvement in the representation of the Rossby waves in the boreal winter
358 (i.e., NH) when increasing the horizontal resolution and shortening the model time step compared to OIFS-LRA-1h
359 configuration. In austral winter, however, the representation of Rossby wave amplitudes and phase speeds are the most realistic
360 in OIFS-LRA-15m configuration, with longer time steps introducing too fast phase speeds and higher horizontal resolution
361 introducing too slow phase speeds at wave number less than 6 (i.e., $k < 6$).

362 3.3 Weather regimes pattern

363 We derive the four weather regimes patterns (WRPs) over NH in the Euro-Atlantic region from ERA5. The patterns resemble
364 the positive and negative phases of the North Atlantic Oscillation (NAO+ and NAO-, respectively), Scandinavian blocking

365 (Sc. Blocking), and the North Atlantic ridge (Atl. Ridge) pattern (**Fig. 7, bottom row**). These WRPs are consistent with the
366 previous findings (**Dawson et al., 2012; Fabiano et al., 2020; Fabiano et al., 2021**).

367

368 The OIFS-HRA-15m configuration produces WRPs that are more visually similar to those in ERA-5 than does OIFS-LRA-1h
369 (Fig. 6), a result confirmed by the higher pattern correlation coefficient (PCC) between OIFS-HRA-15m and ERA5 compared
370 to the OIFS-LRA-1h and ERA-5 (**Fig. 7 and 8**). The PCCs for NAO+, NAO- and Sc. Blocking all exceed 0.8 in OIFS-HRA-
371 1h while OIFS-LRA-1h does not achieve PCC above 0.8 for any WRP (**Fig. 7**).

372

373 The OIFS-MRA-15m configuration shows smaller PCCs than both the OIFS-HRA-15m and OIFS-LRA-1h configurations
374 (**Fig. 8**), i.e., the improvement from OIFS-LRA-1h to OIFS-HRA-15m does not have a linear relationship with model
375 horizontal resolution or time step. Compared to other configurations and ERA5, OIFS-MRA-15m the z500 anomaly in the
376 NAO+ pattern is too elongated in the southwest-northeast direction, and an unrealistic negative z500 anomaly over the North
377 Atlantic appears in the Sc. Blocking regime (**Fig 7**). Furthermore, OIFS-MRA-15m shows an Atl. Ridge pattern with neither
378 the right structure nor amplitude.

379

380 There is an improvement in the representation of the NAO- regime in the OIFS-LRA-30m configuration over the OIFS-LRA-
381 1h configuration (**Fig. 7**) while the Sc. Blocking regime becomes worse due to the ridge shifting westward. These changes are
382 also reflected in the PCCs (**Fig 8**). Similarly, the OIFS-LRA-15m better represents NAO- and Atl. Ridge than OIFS-LRA-1h
383 while NAO+ and Sc. Blocking worsened. The westward shift of the Sc. Blocking is similar in OIFS-LRA-15m and OIFS-
384 LRA-30m, and the worse NAO+ is related to a northward shift of both the positive and negative z500 anomalies. We note that
385 all experiments use the same SST and sea-ice conditions and that OIFS-LRA-1h, 30m and 15m share the same horizontal
386 resolution, i.e., the changes from OIFS-LRA-1h to OIFS-LRA-15m are not due to SST biases or representation of orography.
387 There does not seem to be a clear improvement as time step is shortened, despite the reduction in mean state biases and Rossby-
388 wave amplitudes and phase speeds.

389

390 The PCC is greater than 0.8 for three out of four WRPs in the OIFS-HRA-15m configuration, hence we argue that the OIFS-
391 HRA-15m has the most realistic representation of the weather regimes pattern out of all experiments here. Large improvement
392 in OIFS-HRA-15m over the other configurations could be due to better resolved topography and land-sea contrasts.

393

394 **3.4 Discussion and Conclusions**

395 We have investigated the sensitivity of the climate biases in the OpenIFS atmosphere model to changes in horizontal resolution
396 and time step by analyzing AMIP simulations for the period 1979-2019 (**Table 1**). The strong positive surface zonal wind bias

397 over the Southern Ocean and Northern Hemisphere mid-latitudes and the negative bias in the tropical and subtropical regions
398 have significantly improved in the high horizontal resolution configuration with a short time step (~25km, OIFS-HRA-15m).
399 A similar improvement is observed at the coarse horizontal resolution version with a shorter time step (~100 km with 30 or
400 15-minutes). The zonal wind bias over the mid-latitudes in both hemispheres is reduced throughout the air column when a
401 smaller time step is used in the coarse resolution version, and we find that the changes in the surface winds are largely due to
402 enhanced shallow and mid-level convection which increases vertical momentum transport. Biases in the surface westerlies in
403 midlatitudes are common in CMIP-class climate models (**Bracegirdle et al., 2020**) and a sensitivity to friction has been noted
404 in idealized model studies (**Chen and Plumb, 2009**). We hypothesize that the enhanced shallow and mid-level convection
405 with a shorter model time step and/or increased horizontal resolution deepened the layer over which friction acts in the lower
406 troposphere so that the frictional effects on the barotropic jet increased, leading to a poleward shift in the jet and reduced biases
407 in zonal wind.

408

409 We also find a notable improvement in the representation of the Rossby wave amplitude and phase speed with increased
410 horizontal resolution and shorter time step at least for the waves accounting for most variability in both austral and boreal
411 winter seasons. The reduced zonal wind throughout the troposphere with a shorter time step (**Fig. 3**) would decrease the
412 eastward phase speed of Rossby waves, which may explain part of the reduced phase speeds (**Fig. 6**) and reduced biases.
413 However, changes in air-sea interactions or eddy-mean flow interactions may also play a role. In particular, we note that a very
414 large reduction in phase speed biases in austral winter in OIFS-LRA-15m compared to OIFS-LRA-1h were concurrent with
415 very large reduction in zonal surface wind biases.

416

417 The weather regime patterns are also more realistic in the high horizontal resolution and short time step configuration OIFS-
418 HRA-15m than OIFS-LRA-1h, but we note that there is no consistent improvement from OIFS-LRA-1h to OIFS-HRA-15m
419 as either horizontal resolution is increased or time step is shortened. For example, both OIFS-MRA-15m and OIFS-LRA-15m
420 are worse than OIFS-LRA-1h. The improvements in the weather regime patterns and Rossby wave amplitude and speed could
421 very well be related to each other as e.g. variations in Rossby wave breaking have been linked to the onset of NAO phases
422 (**Strong and Magnúsdóttir, 2008**) but this would require further and more targeted analysis. The overall good representation
423 of weather regimes in OIFS-LRA-1h compared to simulations with shorter time steps (OIFS-LRA-30m, OIFS-LRA-15m) may
424 be due to compensation of errors. For example, it is possible that improving the wave amplitudes and phase speeds in OIFS-
425 LRA-30m compared to OIFS-LRA-1h exposes the effect of a biases caused by both the coarse resolution in both
426 configurations, e.g., weak interactions with topography, leading to an overall worse representation of weather regimes.

427 We found a gradual reduction in SAT biases in OpenIFS with increased resolution or shorter time steps. The improvements
428 were largely driven by improvements over North America and eastern Russia. **Roberts et al. (2018)** noted similar SAT biases
429 and linked them to surface albedo, which is thus likely the cause here as well. The improvement with increased resolution
430 and/or shorter time step may be a result of improved snow cover. Systematic improvements in the precipitation biases were

431 not observed. Instead, precipitation biases generally increased with finer horizontal resolution or shorter time step, suggesting
432 that some tuning may be required in the physics parameters when changing horizontal resolution and time step.

433

434 We stress that the results presented in this study are specific to the OpenIFS atmosphere model and are crucial for the modeling
435 community that uses the OpenIFS in their climate models such as EC-Earth (Haarsma et al., 2020; Döscher et al., 2022),
436 CNRM (Volodire et al., 2019), AWI (Streffing et al., 2022), and GEOMAR (Kjellsson et al., 2020). However, the results
437 may also have implications for other climate modeling communities, at least for those that use a semi-Lagrangian scheme
438 similar to the IFS (e.g., Walters et al., 2019) in the atmospheric component where long time steps are both possible and often
439 desirable to reduce the computational cost of the model.

440

441 The zonal wind bias improvement in the OpenIFS is important for research questions linked with the Southern Ocean climate
442 dynamics that plays a crucial role in both the global atmosphere and ocean circulation. We propose that the model time step
443 not be longer than 30 minutes at any horizontal resolution to minimize surface wind biases over the ocean. The computational
444 cost increases linearly with dt (time step), whereas the cost scales with horizontal resolution as dx^3 as the number of grid
445 points increases in both dimensions and the time step is likely shortened as well. Hence, reducing the model time step from 45
446 or 60 minutes to 20 or 30 minutes may double the computational cost, but lead to significant improvements in the simulated
447 climate. The optimal model time step for the OpenIFS coarse resolution model (1°) is suggested to be 30-minute, but should
448 likely be somewhat shorter, e.g., 15 min, for higher resolutions. In this study, we have not investigated sensitivity of extreme
449 events to the model time step as our focus is mostly on mean state biases. The effect of model horizontal resolution and time
450 step on precipitation extremes is the topic of another manuscript currently in preparation.

451

452 Another limitation of this study is that the time step sensitivity was only tested for the low-resolution configuration, OIFS-
453 LRA, and not the higher resolutions, e.g., OIFS-HRA. We found that much of the surface wind biases were alleviated by a
454 shorter time step due to increased shallow and mid-level convection (Fig. 3). We therefore speculate that a similar sensitivity
455 should be present at high horizontal resolution (~25 km), i.e., a simulation with OIFS-HRA using a 1h time step would most
456 likely exhibit a much larger surface wind biases than the OIFS-HRA simulation with 15min time step

457

458 **Code and data availability**

459 The OpenIFS model requires a software license agreement with ECMWF, and OpenIFS' license is easily given free of charge
460 to any academic or research institute. The details of the different versions of the OpenIFS model, including the OpenIFS
461 version used in this study, i.e., 43R3, can be found at <https://confluence.ecmwf.int/display/OIFS/About+OpenIFS>. The
462 OpenIFS model source code has been made available for the editor and reviewers.

463 The input datasets (both initial and boundary conditions) needed to run the OpenIFS model, run scripts, the model output, and
464 the Jupiter notebook that support the finding of this study are available at (Savita, 2023). The source code for XIOS 2.5,
465 revision 1910, is available from the official repository at <https://forge.ipsl.jussieu.fr/ioserver/> under CeCILL_V2 license.
466 OpenIFS experiments were made using ESM-Tools (https://github.com/esm-tools/esm_tools/). The OASIS coupler is
467 available at <https://oasis.cerfacs.fr/en/>. The XIOS, ESM-Tools and OASIS coupler used in this study can be downloaded from
468 <https://doi.org/10.5281/zenodo.8189718>.

469 The observational datasets used to validate OpenIFS model results in this study are downloaded from the ERA5
470 (<https://cds.climate.copernicus.eu/>), GPCP (<https://psl.noaa.gov/data/gridded/data.gpcp.html>) and CRUTEM4
471 (<http://badc.nerc.ac.uk/data/cru/>) websites. Total model output exceeds 10 Tb and it not publicly available, but is available
472 from the authors upon reasonable requests.

473 **Author contributions**

474 All the model simulations were conducted by AS and JK. Analysis of the output and the writing of text for this paper
475 coordinated by Savita with substantial contributions from JK, RPK, ML, TR, SW and WP.

476 **Competing interest**

477 The authors declare that they have no conflict of interest.

478 **Acknowledgement**

479 AS, JK and MK are supported by JPI Climate/Ocean (ROADMAP project grant 01LP2002C). We wish to thank the OpenIFS
480 team at ECMWF for the technical support. All simulations were performed on the HLRN machine under shk00018 project
481 resources. All analyses were performed on computer clusters at GEOMAR and Kiel University Computing Center (NESH).
482 We thanks to Anton Beljaars for discussion on ECMWF model physics.

483

484

485 **References**

- 486 Athanasiadis, P. J., Ogawa, F., Omrani, N.-E., Keenlyside, N., Schiemann, R., Baker, A. J., Vidale, P. L., Bellucci, A.,
487 Ruggieri, P., and Haarsma, R.: Mitigating climate biases in the midlatitude North Atlantic by increasing model resolution: SST
488 gradients and their relation to blocking and the jet, *Journal of Climate*, 35, 3385-3406, 2022.
- 489 Bayr, T., Latif, M., Dommenges, D., Wengel, C., Harlaß, J., and Park, W.: Mean-state dependence of ENSO atmospheric
490 feedbacks in climate models, *Climate Dynamics*, 50, 3171-3194, 2018.
- 491 Beljaars, A., Balsamo, G., Bechtold, P., Bozzo, A., Forbes, R., Hogan, R. J., Köhler, M., Morcrette, J.-J., Tompkins, A. M.,
492 and Viterbo, P.: The numerics of physical parametrization in the ECMWF model, *Frontiers in Earth Science*, 6, 137, 2018.
- 493 Bracegirdle, T., Holmes, C., Hosking, J., Marshall, G., Osman, M., Patterson, M., and Rackow, T.: Improvements in
494 circumpolar Southern Hemisphere extratropical atmospheric circulation in CMIP6 compared to CMIP5, *Earth and Space
495 Science*, 7, e2019EA001065, 2020.
- 496 Branković, C. and Gregory, D.: Impact of horizontal resolution on seasonal integrations, *Climate Dynamics*, 18, 123-143,
497 2001.
- 498 Chen, G. and Plumb, R. A.: Quantifying the eddy feedback and the persistence of the zonal index in an idealized atmospheric
499 model, *Journal of the atmospheric sciences*, 66, 3707-3720, 2009.
- 500 Couldrey, M. P., Gregory, J. M., Boeira Dias, F., Dobrohotoff, P., Domingues, C. M., Garuba, O., Griffies, S. M., Haak, H.,
501 Hu, A., and Ishii, M.: What causes the spread of model projections of ocean dynamic sea-level change in response to
502 greenhouse gas forcing?, *Climate Dynamics*, 56, 155-187, 2021.
- 503 Dawson, A., Palmer, T., and Corti, S.: Simulating regime structures in weather and climate prediction models, *Geophysical
504 Research Letters*, 39, 2012.
- 505 Döscher, R., Acosta, M., Alessandri, A., Anthoni, P., Arsouze, T., Bergman, T., Bernardello, R., Boussetta, S., Caron, L. P.,
506 Carver, G., Castrillo, M., Catalano, F., Cvijanovic, I., Davini, P., Dekker, E., Doblas-Reyes, F. J., Docquier, D., Echevarria,
507 P., Fladrich, U., Fuentes-Franco, R., Gröger, M., v. Hardenberg, J., Hieronymus, J., Karami, M. P., Keskinen, J. P., Koenigk,
508 T., Makkonen, R., Massonnet, F., Ménégos, M., Miller, P. A., Moreno-Chamarro, E., Nieradzki, L., van Noije, T., Nolan, P.,
509 O'Donnell, D., Ollinaho, P., van den Oord, G., Ortega, P., Prims, O. T., Ramos, A., Reerink, T., Rousset, C., Ruprich-Robert,
510 Y., Le Sager, P., Schmith, T., Schrödner, R., Serva, F., Sicardi, V., Sloth Madsen, M., Smith, B., Tian, T., Tourigny, E., Uotila,
511 P., Vancoppenolle, M., Wang, S., Wärlind, D., Willén, U., Wyser, K., Yang, S., Yepes-Arbós, X., and Zhang, Q.: The EC-

512 Earth3 Earth system model for the Coupled Model Intercomparison Project 6, *Geosci. Model Dev.*, 15, 2973-3020,
513 10.5194/gmd-15-2973-2022, 2022.

514 Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled
515 Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geoscientific Model Development*,
516 9, 1937-1958, 2016.

517 Fabiano, F., Meccia, V. L., Davini, P., Ghinassi, P., and Corti, S.: A regime view of future atmospheric circulation changes in
518 northern mid-latitudes, *Weather and Climate Dynamics*, 2, 163-180, 2021.

519 Fabiano, F., Christensen, H., Strommen, K., Athanasiadis, P., Baker, A., Schiemann, R., and Corti, S.: Euro-Atlantic weather
520 Regimes in the PRIMAVERA coupled climate simulations: impact of resolution and mean state biases on model performance,
521 *Climate Dynamics*, 54, 5031-5048, 2020.

522 Flato, G., Marotzke, J., Abiodun, B., Braconnot, P., Chou, S. C., Collins, W., Cox, P., Driouech, F., Emori, S., and Eyring, V.:
523 Evaluation of climate models, in: *Climate change 2013: the physical science basis. Contribution of Working Group I to the*
524 *Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, 741-866, 2014.

525 Forbes, R. and Tompkins, A.: An improved representation of cloud and precipitation, *ECMWF Newsletter*, 129, 13-18, 2011.

526 Gates, W. L., Boyle, J. S., Covey, C., Dease, C. G., Doutriaux, C. M., Drach, R. S., Fiorino, M., Gleckler, P. J., Hnilo, J. J.,
527 and Marlais, S. M.: An overview of the results of the Atmospheric Model Intercomparison Project (AMIP I), *Bulletin of the*
528 *American Meteorological Society*, 80, 29-56, 1999.

529 Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G.,
530 and Reichle, R.: The modern-era retrospective analysis for research and applications, version 2 (MERRA-2), *Journal of*
531 *climate*, 30, 5419-5454, 2017.

532 Haarsma, R., Acosta, M., Bakhshi, R., Bretonnière, P.-A. B., Caron, L.-P., Castrillo, M., Corti, S., Davini, P., Exarchou, E.,
533 and Fabiano, F.: HighResMIP versions of EC-Earth: EC-Earth3P and EC-Earth3P-HR. Description, model performance, data
534 handling and validation, (No Title), 2020.

535 Harris, I., Jones, P. D., Osborn, T. J., and Lister, D. H.: Updated high-resolution grids of monthly climatic observations—the
536 CRU TS3.10 Dataset, *International journal of climatology*, 34, 623-642, 2014.

537 Hazeleger, W., Wang, X., and Severijns, C.: SS tef anescu, R Bintanja, A Sterl, Klaus Wyser, T Semmler, S Yang, B Van den
538 Hurk, et al. Ec-earth v2. 2: description and validation of a new seamless earth system prediction model, *Climate dynamics*, 39,
539 2611-2629, 2012.

540 He, C. and Zhou, T.: The two interannual variability modes of the western North Pacific subtropical high simulated by 28
541 CMIP5–AMIP models, *Climate dynamics*, 43, 2455-2469, 2014.

542 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., and
543 Schepers, D.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999-2049, 2020.

544 Hogan, R. J. and Bozzo, A.: A flexible and efficient radiation scheme for the ECMWF model, *Journal of Advances in Modeling
545 Earth Systems*, 10, 1990-2008, 2018.

546 Huffman, G. J., Adler, R. F., Arkin, P., Chang, A., Ferraro, R., Gruber, A., Janowiak, J., McNab, A., Rudolf, B., and Schneider,
547 U.: The global precipitation climatology project (GPCP) combined precipitation dataset, *Bulletin of the american
548 meteorological society*, 78, 5-20, 1997.

549 Jung, T., Miller, M., Palmer, T., Towers, P., Wedi, N., Achuthavarier, D., Adams, J., Althshuler, E., Cash, B., and Kinter Iii, J.:
550 High-resolution global climate simulations with the ECMWF model in Project Athena: Experimental design, model climate,
551 and seasonal forecast skill, *Journal of Climate*, 25, 3155-3172, 2012.

552 Kim, S. T., Cai, W., Jin, F.-F., and Yu, J.-Y.: ENSO stability in coupled climate models and its association with mean state,
553 *Climate dynamics*, 42, 3313-3321, 2014.

554 Kjellsson, J., Streffing, J., Carver, G., and Köhler, M.: From weather forecasting to climate modelling using OpenIFS, ECMWF
555 *Newsletter*, 164, 38-41, 2020.

556 Liu, B., Gan, B., Cai, W., Wu, L., Geng, T., Wang, H., Wang, S., Jing, Z., and Jia, F.: Will increasing climate model resolution
557 be beneficial for ENSO simulation?, *Geophysical Research Letters*, 49, e2021GL096932, 2022.

558 Ma, P.-L., Harrop, B. E., Larson, V. E., Neale, R. B., Gettelman, A., Morrison, H., Wang, H., Zhang, K., Klein, S. A., and
559 Zelinka, M. D.: Better calibration of cloud parameterizations and subgrid effects increases the fidelity of the E3SM Atmosphere
560 Model version 1, *Geoscientific Model Development*, 15, 2881-2916, 2022.

561 Matthes, K., Biastoch, A., Wahl, S., Harlaß, J., Martin, T., Brücher, T., Drews, A., Ehlert, D., Getzlaff, K., and Krüger, F.:
562 The flexible ocean and climate infrastructure version 1 (FOCI1): Mean state and variability, *Geoscientific Model
563 Development*, 13, 2533-2568, 2020.

564 Meehl, G. A. and Teng, H.: CMIP5 multi-model hindcasts for the mid-1970s shift and early 2000s hiatus and predictions for
565 2016–2035, *Geophysical Research Letters*, 41, 1711-1716, 2014.

566 Meng, Y., Hao, Z., Feng, S., Guo, Q., and Zhang, Y.: Multivariate bias corrections of CMIP6 model simulations of compound
567 dry and hot events across China, *Environmental Research Letters*, 17, 104005, 2022.

568 Osborn, T. and Jones, P.: The CRUTEM4 land-surface air temperature data set: construction, previous versions and
569 dissemination via Google Earth, *Earth System Science Data*, 6, 61-68, 2014.

570 Pilch Kedzierski, R., Matthes, K., and Bumke, K.: New insights into Rossby wave packet properties in the extratropical UTLS
571 using GNSS radio occultations, *Atmospheric Chemistry and Physics*, 20, 11569-11592, 2020.

572 Roberts, C. D., Senan, R., Molteni, F., Boussetta, S., Mayer, M., and Keeley, S. P.: Climate model configurations of the
573 ECMWF Integrated Forecasting System (ECMWF-IFS cycle 43r1) for HighResMIP, *Geoscientific model development*, 11,
574 3681-3712, 2018.

575 Savita, A.: Assessment of Climate Biases in OpenIFS Version 43R3 across Model Horizontal Resolutions and Time Steps
576 [dataset], <https://hdl.handle.net/20.500.12085/c74887dc-e609-4392-9faf-48c67276d5d1>, 2023.

577 Simmons, A. J. and Burridge, D. M.: An energy and angular-momentum conserving vertical finite-difference scheme and
578 hybrid vertical coordinates, *Monthly Weather Review*, 109, 758-766, 1981.

579 Streffing, J., Sidorenko, D., Semmler, T., Zampieri, L., Scholz, P., Andrés-Martínez, M., Koldunov, N., Rackow, T., Kjellsson,
580 J., and Goessling, H.: AWI-CM3 coupled climate model: description and evaluation experiments for a prototype post-CMIP6
581 model, *Geoscientific Model Development*, 15, 6399-6427, 2022.

582 Strong, C. and Magnusdottir, G.: Tropospheric Rossby wave breaking and the NAO/NAM, *Journal of the atmospheric
583 sciences*, 65, 2861-2876, 2008.

584 Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An overview of CMIP5 and the experiment design, *Bulletin of the American
585 meteorological Society*, 93, 485-498, 2012.

586 Temperton, C., Hortal, M., and Simmons, A.: A two-time-level semi-Lagrangian global spectral model, *Quarterly Journal of
587 the Royal Meteorological Society*, 127, 111-127, 2001.

588 Tiedtke, M.: Representation of clouds in large-scale models, *Monthly Weather Review*, 121, 3040-3061, 1993.

589 Voltaire, A., Saint-Martin, D., S n si, S., Decharme, B., Alias, A., Chevallier, M., Colin, J., Gu r my, J. F., Michou, M., and
590 Moine, M. P.: Evaluation of CMIP6 deck experiments with CNRM-CM6-1, *Journal of Advances in Modeling Earth Systems*,
591 11, 2177-2213, 2019.

592 Walters, D., Baran, A. J., Boutle, I., Brooks, M., Earnshaw, P., Edwards, J., Furtado, K., Hill, P., Lock, A., and Manners, J.:
593 The Met Office Unified Model global atmosphere 7.0/7.1 and JULES global land 7.0 configurations, *Geoscientific Model*
594 *Development*, 12, 1909-1963, 2019.

595 Williamson, D. L., Kiehl, J. T., and Hack, J. J.: Climate sensitivity of the NCAR Community Climate Model (CCM2) to
596 horizontal resolution, *Climate Dynamics*, 11, 377-397, 1995.

597 Wolf, G. and Wirth, V.: Diagnosing the horizontal propagation of Rossby wave packets along the midlatitude waveguide,
598 *Monthly Weather Review*, 145, 3247-3264, 2017.

599 Zhou, S., Huang, G., and Huang, P.: Excessive ITCZ but negative SST biases in the tropical Pacific simulated by CMIP5/6
600 models: The role of the meridional pattern of SST bias, *Journal of Climate*, 33, 5305-5316, 2020.

601

602

Table

603

Experiment Name	Horizontal resolution	Vertical grid	Time step	CHPSY	SYPD
OIFS-LRA-15m	Tco95/100km	L91	15m	3.3 k	11
OIFS-MRA-15m	Tco199/50km	L91	15m	13.3 k	4
OIFS-HRA-15m	Tco399/25km	L91	15m	19.2 k	2
OIFS-LRA-30m	Tco95/100km	L91	30m	845	21
OIFS-LRA-1h	Tco95/100km	L91	1h	256	36

604


605 **Table 1.** List of the experiments performed across different horizontal resolutions and model time steps using OIFS model.

606

In the above table CHPSY is core hours per simulation year, and SYPD is simulation year per day.

607

608




610


611

612 **Figure 1.** (a) Annual mean ERA5 surface zonal wind [ms^{-1}]. (b-d) Annual mean zonal wind [ms^{-1}] bias for different model
 613 time steps (1h (b), 30m (c), and 15m (d)) using ~ 100 km resolution, and (e-f) with different horizontal resolutions, ~ 50 (e) and
 614 ~ 25 km (f), respectively. Biases are computed with respect to ERA5 over the period 1979–2019.

615



616
 617 **Figure 2.** Root mean squared error of surface zonal wind (a), SAT (b), and precipitation (c) over the period 1979-2019 for all
 618 the configurations: annual (black) and seasonal mean (DJF: blue, JJA: red). The error bars represent a 95% confidence interval.




620

621

622 **Figure 3.** (a) Averaged zonal wind (u) [ms^{-1}] and zonal wind tendencies [$\text{ms}^{-2}/\text{hour}$] over the Southern Ocean ($40^{\circ}\text{S} - 60^{\circ}\text{S}$,
 623 all longitude) as a function of height for OIFS-LRA-1h and OIFS-LRA-15m. Model levels (y-axis left) and pressure levels (y-
 624 axis right). (c) Zonal and time average of zonal wind tendencies at the lowest level of the model as a function of latitude. (d)
 625 Zonal and time average convection difference [$\text{Kgm}^{-2}/\text{hour}$] between OIFS-LRA-15m and OIFS-LRA-1h configurations. The
 626 solid lines in panels (b) and (c) show the wind tendency for OIFS-LRA-1h configuration whereas the dashed lines are for
 627 OIFS-LRA-15m configuration. Shown are averages over 1979-2019.


628



629


630 **Figure 4.** (a) Annual zonal mean ERA5 zonal wind [ms^{-1}]. (b-d) Annual zonal mean zonal wind [ms^{-1}] bias for different model
 631 time steps (1h (b), 30m (c), and 15m (d)) using ~ 100 km resolution, and (e-f) with different horizontal resolutions, ~ 50 (e) and
 632 ~ 25 km (f), respectively. Biases are computed with respect to ERA5 over the period 1979–2019.

633




634

635 **Figure 5.** (A) The Rossby wave amplitude (contours) for different wave numbers in the Northern Hemisphere at 300 hPa (a)
 636 in ERA5 observation and (b-f) in the OIFS model simulations during 1979-2019 in DJF (i.e., boreal winter). The color shows
 637 difference of wave amplitude between the model and ERA5 where it is significant on the 95 % confidence level. The wave
 638 amplitude and contour interval are shown in ms^{-1} . The grey contours start from 2 ms^{-1} and the black contours from 5 ms^{-1} and
 639 the contour interval is 1 ms^{-1} . (B) is similar to (A), but for JJA (i.e., austral winter).



640

641 **Figure 6.** (A) The Rossby wave phase speed (contours) for different wave numbers at 300 hPa in the Northern Hemisphere in
 642 ERA5 (a) observation and (b-f) in the OIFS model simulations during 1979-2019 in DJF (i.e., boreal winter). The color shows
 643 the difference of wave phase speed between model and ERA5 where it is significant on the 95 % confidence level. The wave
 644 phase speed and contour interval are shown in ms^{-1} . The black contours start from 1 ms^{-1} and the contour interval is 1 ms^{-1} .
 645 Panel (B) is similar to panel (A), but for JJA (i.e., austral winter). The dashed contours show a negative phase speed and a gray
 646 contour shows a zero-phase speed.




647

648

649 **Figure. 7.** Weather regime patterns over the Euro–Atlantic regions from ERA5 observation (bottom row) and the individual
 650 OIFS model simulations (1st to 5th row) over the time period 1979–2019 for DJF (boreal winter season).

651



652

653 **Figure 8.** Pattern correlation coefficient of the individual weather regime between OIFS model configurations and ERA5 for
654 the period 1979-2019 for the DJF season. The error bars represent a 95% confidence interval.

655

656

657

658