

# Deep learning applied to CO<sub>2</sub> power plant emissions quantification using simulated satellite images

Joffrey Dumont Le Brazidec<sup>1</sup>, Pierre Vanderbecken<sup>1</sup>, Alban Farchi<sup>1</sup>, Grégoire Broquet<sup>2</sup>, Gerrit Kuhlmann<sup>3</sup>, and Marc Bocquet<sup>1</sup>

<sup>1</sup>CEREA, École des Ponts and EDF R&D, Île-de-France, France

<sup>2</sup>Laboratoire des Sciences du Climat et de l'Environnement, LSCE/IPSL, CEA-CNRS-UVSQ, Université Paris-Saclay, 91198 Gif-sur-Yvette, France

<sup>3</sup>Swiss Federal Laboratories for Materials Science and Technology (Empa), Dübendorf, Switzerland

**Correspondence:** Joffrey Dumont Le Brazidec (joffrey.dumont@enpc.fr)

## Abstract.

The quantification of emissions of greenhouse gases and air pollutants through the inversion of plumes in satellite images remains a complex problem that current methods can only assess with significant uncertainties. The anticipated launch of the CO<sub>2</sub>M ([Copernicus Anthropogenic Carbon Dioxide Monitoring](#)) satellite constellation in 2026 is expected to provide high-resolution images of CO<sub>2</sub> ([carbon dioxide](#)) column-averaged mole fractions (XCO<sub>2</sub>), opening up new possibilities. However, the inversion of future CO<sub>2</sub> plumes from CO<sub>2</sub>M will encounter various obstacles. A challenge is the CO<sub>2</sub> plume low signal-to-noise ratio, due to the variability of the background and instrumental errors in satellite measurements. Moreover, uncertainties in the transport and dispersion processes further complicate the inversion task.

To address these challenges, deep learning techniques, such as neural networks, offer promising solutions for retrieving emissions from plumes in XCO<sub>2</sub> images. Deep learning models can be trained to identify emissions from plume dynamics simulated using a transport model. It then becomes possible to extract relevant information from new plumes and predict their emissions.

In this paper, we employ convolutional neural networks (CNN) to estimate the emission fluxes from a plume in a pseudo XCO<sub>2</sub> image. Our dataset used to train and test such methods includes pseudo images based on simulations of hourly XCO<sub>2</sub>, NO<sub>2</sub> and wind fields near various power plants in Eastern Germany, tracing plumes from anthropogenic and biogenic sources. CNN models are trained to predict emissions from three power plants that exhibit diverse characteristics. The power plants used to assess the deep learning model's performance are not used to train the model. We find that the CNN model outperforms state of the art plume inversion approaches, achieving highly accurate results with an absolute error about half of that of the cross-sectional flux method [and a absolute relative error of ~ 20% when only the XCO<sub>2</sub> and wind fields are used as inputs](#). Furthermore, we show that our estimations are only slightly affected by the absence of NO<sub>2</sub> ([nitrogen dioxide](#)) fields or a detection mechanism as additional information. Finally, interpretability techniques applied to our models confirm that the CNN automatically learns to identify the XCO<sub>2</sub> plume and to assess emissions from the plume concentrations. These promising results suggest a high potential of CNNs in estimating local CO<sub>2</sub> emissions from satellite images.

# 1 Introduction

25 The burning of fossil fuels, such as coal and oil, in power plants (PP), is a primary source of anthropogenic CO<sub>2</sub> emissions. Approximately 50% of worldwide fossil fuel CO<sub>2</sub> emissions originate from large facilities, which encompass PPs (??). As a result, maintaining regular monitoring of these emissions and possessing the capacity to control their reporting is crucial.

Observations from satellites like OCO-2 provide valuable data that can be utilised to estimate CO<sub>2</sub> emissions (???????). Specifically, observations of CO<sub>2</sub> plumes, such as the plume transects obtained from OCO2-2 and OCO-3 satellites, offer a  
30 direct means of quantifying their source. The upcoming launch of the CO<sub>2</sub>M satellites in 2026 is anticipated to capture high-resolution images of CO<sub>2</sub> column-averaged mole fractions (XCO<sub>2</sub>), aiming to have a much larger swath and more accurate CO<sub>2</sub> estimations, further advancing our capabilities in this area. Leveraging these images, however, will present significant challenges (?).

CO<sub>2</sub> plumes are notoriously difficult to invert due to various factors, including 1) image integrity issues caused by cloud  
35 cover or satellite overpasses, which result in missing data in the images used for analysis. Additionally, the estimation of the emissions associated with a plume is further complicated by 2) the measurements low signal-to-noise (SNR) ratio. The noise component encompasses variations in the background as well as errors in the satellite measurements. The SNR problem stands as the main hurdle in the detection of the plume, a crucial step for inversion. Recent research conducted by ? has illustrated the remarkable ability of Convolutional Neural Networks (CNNs) to effectively overcome this obstacle. Lastly, another challenge  
40 stems from 3) the uncertainties in the transport and dispersion processes, specifically, when it comes to estimating the effective wind driving the plume and determining its shape (?).

This paper addresses the second and third problems (except for the errors in the satellite measurements) by employing deep learning techniques to perform inverse modelling of CO<sub>2</sub> plumes. In particular, we focus on developing techniques for inverting CO<sub>2</sub> plumes from PPs of different emission levels.

45 To assert the effectiveness of the method, the predictions of the deep learning model are compared against a state-of-the-art ~~technique, the cross-sectional flux (CSF) approach~~techniques. Plume inversion methods include approaches that use an atmospheric transport model to simulate the plume and compare it to observation (e.g., ??). They also include techniques that quantify emissions from a hotspot based on plume detection in satellite observations (?). These methods can be based on time-averaged plumes, such as the divergence method (??), or on instantaneous images. ? compared several of these approaches,  
50 namely the Gaussian plume inversion, the integrated mass enhancement and the ~~CSF method. Among these approaches, the CSF method is regularly diagnosed as being among the most robust and viable options (??), although recent developments suggest that alternative cross-sectional flux method~~. In the CoCO<sub>2</sub> project, several of these methods were compared using synthetic CO<sub>2</sub>M CO<sub>2</sub> and NO<sub>2</sub> observations (?). Here, we use the cross-sectional flux (CSF) method for comparison, which showed similar accuracy as other well performing methods such as Gaussian plume inversion and the light cross-sectional  
55 flux or based on Gaussian plume (LCSF) method. It should be noted that the CoCO<sub>2</sub> project identified several potential improvements of the methods that may yield superior performance in the future (?).


In this paper, the proposed plume inversion approach is based on convolutional neural networks. This research builds on earlier work in the field of remote sensing image analysis leveraging machine learning techniques (????). Here, plume inversion involves the analysis of an image to extract scalar or vector emissions data at different time steps. Therefore, this task can be framed as an image regression problem, where relying on CNNs can offer significant advantages (?). CNNs, a type of supervised learning method, can be trained on a comprehensive dataset, where all input variables (images) and associated output variables (like emissions) are known. Once trained, these CNNs can effectively process and draw conclusions from unseen observational imagery. CNNs employ convolutional layers to extract essential features from images. Each filter is automatically trained to detect specific patterns, such as edges, corners, or other shapes within the image. By stacking multiple convolutional layers, CNNs become capable of learning intricate patterns, enabling them to capture increasingly complex features. The ability of CNNs to capture and learn spatial features in images makes them a popular choice for various image-related tasks, including image recognition, classification, and regression. Given the nature of our plume inversion task, they are particularly well-suited due to their ability to identify spatial features in images, such as plume shapes or intensity, that correspond to specific emissions. This feature extraction approach effectively harnesses the knowledge embedded in transport models, enabling this automatic capture of plumes dynamics. This ability to capture such features has already been demonstrated by ~~?, ?, or ?~~ ?, ?, or ? which study the segmentation of plumes in CO<sub>2</sub> images.

To train and test the CNN models, this paper relies on a synthetic dataset as CO<sub>2</sub>M data will not be available until 2026. This dataset has been designed to possess similar key features as the forthcoming CO<sub>2</sub>M satellite, such as resolution and the availability of NO<sub>2</sub> data. ~~This study specifically focuses on clear sky conditions, and~~ However, the influence of clouds or systematic error patterns is not considered in the analysis.

Before introducing the inversion methodology and results, we briefly describe the physical fields used to train and evaluate the CNNs. This includes the presentation of the simulated satellite fields in section 2 and of the model used to produce segmentation masks of the plumes used as inputs of the inversion model in section 3. The inversion methodology using CNN is described in section 4, specifying the problem statement, the model, the training process, and the alternative method employed for comparison. The subsequent section 5 delves into the application of the model to three specific PPs. In particular, subsection 5.3 places special emphasis on the interpretability of the trained CNNs. Discussions on the limitations and future directions of this study can be found in section 6, while the conclusions are outlined in section 7.

## 2 Dataset of XCO<sub>2</sub>, winds and NO<sub>2</sub> images

XCO<sub>2</sub> and NO<sub>2</sub> images are taken from the concentration fields simulated for the SMARTCARB project (?). Using the COSMO-GHG model, the SMARTCARB simulations were performed in a region centred around Berlin and ~~covered several nearby~~ covering several nearby coal-fired PPs. These simulations were used to produce synthetic observations of CO<sub>2</sub>M and to evaluate various plume detection and inversion approaches (?????). The data are hourly and cover an entire year. Their spatial resolution is 0.01° and sixty vertical layers ranging spanning from an altitude of 0 to 24 km are used. More information can be found in ?, which presented more extensively a dataset very similar to the one used in this study.



**Figure 1.** Examples of inputs used by the CNN model. The first, second, third and fourth columns represent the XCO<sub>2</sub> images, NO<sub>2</sub> images, vertically averaged *u* and *v* winds, respectively.

90 Images used to train and evaluate the CNN inversion models consist of  $64 \times 64$  pixels, with each pixel covering an area of  $2 \times 2 \text{ km}^2$ . Each image is extracted from the SMARTCARB COSMO-GHG simulated fields so that one hotspot is located in the centre. In addition, the selected size guarantees the inclusion of the majority of the central hotspot plume within the image. The mapping from the original SMARTCARB fields resolution to the 2 km resolution is performed by cubic spline interpolation (?). The 2 km resolution was chosen to be consistent with the resolution expected for CO<sub>2</sub>M observations.

95 It is necessary to take into account the expected noise associated with the satellite instruments. For this, a Gaussian random noise of standard deviation 0.7 ppm, characteristic of the CO<sub>2</sub>M (?), is added to the XCO<sub>2</sub> images.

In addition to XCO<sub>2</sub>, ancillary data can be used to assist in the inversion of XCO<sub>2</sub> plumes. Considering that CO<sub>2</sub>M will provide measurements of NO<sub>2</sub> and the observed strong correlation between NO<sub>2</sub> and CO<sub>2</sub> plumes, noisy NO<sub>2</sub> fields are used in this study. The noise associated with a NO<sub>2</sub> field is implemented as the standard normal distribution multiplied by the NO<sub>2</sub> field values. The median standard value for NO<sub>2</sub> fields surpasses  $1e15 \text{ molec.cm}^{-2}$ , leading to an average noise level in the NO<sub>2</sub> field that exceeds the CO<sub>2</sub>M NO<sub>2</sub> requirement (less than  $1e15 \text{ molec.cm}^{-2}$ ). Furthermore, ERA5 winds are used: the original resolution of 28 km is mapped to 2 km to be consistent with the CO<sub>2</sub> and NO<sub>2</sub> images. To overcome the circular data limitation of statistical methods, the *u* and *v* wind fields are used instead of the direction and magnitude components. This limitation corresponds to the statistical model's inability to correctly interpret wind directions where the value of 360 degrees is equivalent to 0 degrees. More precisely, we use 2D *u* and *v* wind fields which are calculated as the average of the zonal and meridional wind fields over the 37 lower ERA5 vertical levels, respectively, which corresponds roughly to the lowest 4000 km of the atmosphere. Figure 1 presents a series of potential inputs to the CNN.

This paper only addresses the retrieval of PP emissions, although the training dataset includes the city of Berlin. More precisely, depending on the PP evaluated, the training dataset might be composed of any hotspot in {Berlin, Jänschwalde, Schwarze Pumpe, Boxberg, Turow, Patnow, Lippendorf, Opole, Dolna Odra}. The primary rationale behind prioritising the training of the model on PPs is the scarcity of cities in the dataset, which poses a challenge for the model to effectively learn and generalise for cities. However, Berlin is included in the training dataset as supplementary data to aid the model

Hotspot	min	max	mean	std
Berlin	4.8	34.7	16.8	7.2
Jänschwalde	16.4	52.7	33.3	7.7
Boxberg	9.4	30.1	19.0	4.4
Lippendorf	7.5	24.1	15.2	3.5
Turow	4.3	13.8	8.7	2.0
Schwarze Pumpe	4.0	13.0	8.2	1.9
Dolna Odra	3.7	12.5	7.9	1.9
Opole	3.5	11.8	7.5	1.8
Patnow	2.9	9.2	5.8	1.3

**Table 1.** Emission statistics for the considered PPs and the city of Berlin. Fluxes are in Mt CO<sub>2</sub> /yr.


in its learning process. In the SMARTCARB dataset, the modelling of anthropogenic emissions, incorporating fixed diurnal, weekly, and seasonal cycles, was performed using the TNO-MACC III inventory (?). The emissions range, mean, and standard deviation of each hotspot are given in Table 1 [and locations of considered PPs and Berlin are described in 2](#). Moreover, data augmentation techniques are employed to expand the database, as detailed in section 4.2.1.

### 3 Application of the segmentation model

Utilising a segmentation algorithm to incorporate plume contours as ~~critical~~[additional](#) prior information in plume inversion may yield significant benefits. In this section, we provide a brief description and application of the CNN-based method developed in ? that predicts plume contours in XCO<sub>2</sub> images. The methodology of ? involves employing an image-to-image U-net model, which generates images that are subsequently used as inputs for the CNN inversion model, as outlined in section 4.

Apart from a few specific points, the training and model choices are similar to those of ?. A simpler encoder, with fewer neurons, is chosen since the NO<sub>2</sub> fields are used as inputs to the CNN. This simplification of the problem reduces the need for a complex encoder. In addition to NO<sub>2</sub> and XCO<sub>2</sub> fields, winds are also used to assist in the XCO<sub>2</sub> plume contour prediction, although experiments show that the addition of this data has very little influence on the predictions. Finally, the U-net models were designed to make predictions beyond the geographical region of their training data. Specifically, the model that learns to predict the mask of the Boxberg PP plume from an image centred at Boxberg PP is trained on a dataset excluding the images centred at Boxberg.

In Fig. 3, we show the application of a model trained to predict the positions of Turow plumes. It was trained on pairs of fields in the regions of {Boxberg, Berlin, Lippendorf, Patnow, Jänschwalde, Dolna Odra, Schwarze Pumpe, Opole}. The first




**Figure 2.** XCO<sub>2</sub> concentration map with the locations of Berlin and each considered PP within the complete SMARTCARB domain. The map consists only of the concentrations stemming from the major anthropogenic sources. Furthermore, to enhance plume visibility, as fluxes of power plants such as Janschwalde are vastly superior to other fluxes, concentrations exceeding 2 ppmv have been capped at 2 ppmv.

and second columns of the figure show the XCO<sub>2</sub> and NO<sub>2</sub> fields, inputs to the CNN. The third column shows the target plume as a reference point, while the fourth column shows the output of the CNN.

## 4 Deep learning method for the inversion of XCO<sub>2</sub>

### 4.1 Inversion based on supervised learning

135 The inverse problem addressed here is the estimation of the CO<sub>2</sub> emissions accountable for the target plume observed in a given XCO<sub>2</sub> field image. To do so, a CNN is used, processing as input a given XCO<sub>2</sub> field and other additional fields and resulting in a scalar output representing the emission rate of CO<sub>2</sub> in MtCO<sub>2</sub>.yr<sup>-1</sup> at the hour corresponding to the image. This flux unique scalar representation choice was made for the sake of simplification, as the quality of the results is minimally impacted by the choice between targeting average, instantaneous fluxes, or a vector of instantaneous fluxes over the last  $N$  hours. This is due to the relatively slow hourly variation in the CO<sub>2</sub> emission rate of the PP. All future flux quantities are expressed in Mt.yr<sup>-1</sup>. The image-to-scalar, or image regression, problem is depicted in Fig. 4. The CNN is trained using pairs of XCO<sub>2</sub> fields, ancillary data fields (winds, segmentation results and NO<sub>2</sub>) and associated emission fluxes ranging from 3 to 53 Mt.yr<sup>-1</sup> across various times and targets.



**Figure 3.** Examples of the U-net CNN model application on images centred at Turow. The first, second, third and fourth columns represent the XCO<sub>2</sub> images, NO<sub>2</sub> images, weighted Boolean plumes and U-net predictions as probability maps, respectively. All times are in UTC.

## 4.2 CNN model and training parametrisation


145 In this section, we present and discuss the architecture of the model, the hyperparameters, and the learning methodology for the inversion task. In particular, the model is built from preprocessing layers and a core model. The preprocessing layers are used to augment/transform, construct, add noise, and normalise the input data before feeding it into the core model. The core model is designed to extract features from these transformed input data.

### 4.2.1 Description of the preprocessing layers

150 The preprocessing layers consist of a six steps sequence, as presented in Fig. 5. The purpose of steps 1-2-3 is to extend the initial database, thereby enhancing the model’s ability to generalise to unseen data. A pair of input/output data for training is constituted in this way:

1. a target CO<sub>2</sub> plume corresponding to a PP at a time  $t$  is chosen randomly. A background must be constructed to form the XCO<sub>2</sub> image with this plume. This background is chosen randomly and therefore does not necessarily correspond





**Figure 4.** The  $XCO_2$ -Various fields:  $XCO_2$ , ancillary data winds, segmentation results and  $NO_2$ , and emissions (of the central plume) emissions are used as inputs by a CNN that learns to estimate the emissions associated to the central plume concealed under the background. Note that the combination of  $NO_2$  and segmentation results as inputs is never assessed in this manuscript.

155 (geographically and temporally) to the chosen target plume. It may correspond to another PP as well as to another time. Furthermore, in SMARTCARB, the background is partitioned into multiple segments. In our case, the background is constructed from two randomly and independently drawn fields: a field containing the major part of the fluxes including the biogenic fluxes and a field containing a part of the anthropogenic fluxes of the SMARTCARB domain. Finally, potential additional fields (wind,  $NO_2$ , segmentations) corresponding geographically and temporally to the chosen  $CO_2$  plume are selected;

160

2. then,

- the target  $XCO_2$  plume is multiplied by a random uniform scaling factor  $p \sim U(0.25, 2)$ . The corresponding true emissions are also multiplied:  $y_{truth}^{scaled\ plume} = p \times y_{truth}^{plume}$ ;
- a random number  $b \sim U(-3.5, 3.5)$  (in ppmv) is added uniformly to the main background. Extrema of this uniform distribution are chosen as approximately the standard deviation of an average  $XCO_2$  background;
- a random uniform scaling factor  $a \sim U(0.33, 3)$  is applied to multiply the field containing a part of the alternate anthropogenic fluxes.

165

Anthropogenic fluxes scaling factors  $p$  and  $a$  are chosen so that the resultant emissions still correspond to reasonable anthropogenic fluxes;



- 170 3. the XCO<sub>2</sub> field is built as the sum of the target plume, background, and alternate anthropogenic fluxes field components;
4. a Gaussian noise matrix of shape 64 × 64 (equal to the image shape)  $\mathbf{G}_{N_x \times N_y} \sim N(0, 0.7)$  (in ppmv) is added to the XCO<sub>2</sub> field to simulate the satellite observational noise;
5. the noisy XCO<sub>2</sub> field is concatenated with the additional fields. If added, the NO<sub>2</sub> field is noised beforehand;
- 175 6. standardisation (i.e., Z-score normalisation) is performed independently on each channel (each physical field) of the concatenated input data.

These steps are carried out exclusively during the model training phase. The different operations in this process are performed to create a more robust and diverse training dataset. To ensure an accurate assessment of the performance of the trained model, the test dataset used for the evaluation consists only of pre-constructed, physically consistent simulated data. Specifically, no scaling factors are applied and the XCO<sub>2</sub> fields used for testing are always constructed from geographically and temporally  
180 consistent plume and background components.


#### 4.2.2 Description of the core model

The chosen core CNN model, described in Fig. 6 is designed for image regression. It was chosen by comparing its performance with state-of-the-art models such as EfficientNet or Squeezenet (?). These two models are deep neural networks designed primarily for image classification tasks. They incorporate modified versions of CNN, including features such as residual connections and depth-separable convolutions, in order to improve efficiency, speed and ease of implementation. As their initial  
185 implementation tended to overfit, we considered a smaller version with a reduced number of neurons in each layer. But even after tuning, the simpler model depicted below outperformed these more advanced models.

The chosen model takes ~~4 to 5~~ 3 to 4 images of 64×64 pixels as input (which correspond to the XCO<sub>2</sub> field and ancillary data such as the winds). It is constructed as a succession of convolutional, max pooling, batch normalisation, and dropout layers  
190 where:

- convolutional layers aim to identify and extract relevant features by applying a set of learnable filters to the previous feature map. The 2D convolutional operations are applied with a filter size of 32 and a kernel size of 3×3;
- max pooling layers play a key role in reducing the resolution of feature maps while retaining the essential information. This reduces the computational complexity of the network and leads to the extraction of more complex features;
- 195 – batch normalisation layers are used to improve the stability of the network and speed up its learning process;
- dropout layers randomly exclude a certain percentage of the neurons in the previous layer at each iteration to reduce overfitting. They are only activated during the training phase.

The output of the core model is flattened and fed into a fully connected ~~terminal~~ final dense layer with a single output unit, which is activated by a leaking rectified linear unit (ReLU). In total, the CNN has approximately ~ 186,000 ~~parameters are trained by the CNN parameters.~~  
200 parameters.



**Figure 5.** Description of the preprocessing layers as a sequence of six steps: 1) random choice of the XCO<sub>2</sub> field components, 2) scaling transformation of the components, 3) sum of the components, 4) satellite noise simulation, 5) concatenation with the additional data, and 6) normalisation.



parameters. This approach enabled us to produce a range of predictions that were subsequently averaged into a single estimate. To ensure sufficient convergence of these estimates, ~~at least a minimum of~~ five individual models were used for each considered configuration. Through these steps, we significantly reduced the variability observed across multiple models runs, thereby  
220 improving the accuracy and confidence in our system.

### 4.3 Geographical separation between the training and test datasets

We consider a strict geographical separation between the PPs of the training/validation and those of the test dataset. For example, to train a model to predict emissions from Boxberg plumes, we consider a training dataset consisting of images centred at all other available PPs except Boxberg. In this way, the model predicting Boxberg emissions never had access to the  
225 Boxberg emissions inventory.

During the training phase, to avoid overfitting or underfitting, a separate validation dataset is used to follow the model's performance on new data. In our case, the validation dataset is chosen as unseen data from the same geographical area as the training dataset.

Three distinct models are created and trained to predict emissions from three specific PPs. The target PPs are Lippendorf, Boxberg and Turow and are selected ~~to obtain~~:  
230

- to obtain a diverse range of emission rates: between 7 and 24 Mt.yr<sup>-1</sup>, with a mean of 15 Mt.yr<sup>-1</sup> and a std of 3 Mt.yr<sup>-1</sup> for Lippendorf, between 4 and 13 Mt.yr<sup>-1</sup>, with a mean of 9 Mt.yr<sup>-1</sup> and a std of 2 Mt.yr<sup>-1</sup> for Turow, and between 9 to 30 Mt.yr<sup>-1</sup> with a mean of 19 Mt.yr<sup>-1</sup> and a std of 4. Mt.yr<sup>-1</sup> for Boxberg;
- to account for the potential presence of multiple plumes in the same image: the images centred at Boxberg also include  
235 plumes from Jänschwalde, Schwarze Pumpe, and Turow which sometimes overlap;
- because their emission rates are neither at the highest nor the lowest extremes (Boxberg was selected over Jänschwalde because Jänschwalde has the highest emission levels in our dataset, potentially placing it outside the training distribution);
- due to their positioning away from the boundaries of the SMARTCARB domain. These PPs are less affected by border  
240 conditions. As shown in Fig. 1, Patnow or Opole are close to the borders and are influenced by border conditions.

Although these models differ in terms of the training datasets used, they all share the same architectural framework, including hyperparameters, CNN structure, and preprocessing layers. The objective behind creating these models is to demonstrate the effectiveness of the architectural framework, laying the groundwork for a universal model based on this architecture and capable of generalising across future PPs.

### 245 4.4 Alternative method for comparison

~~In order to assess the~~ The accuracy of our inversion method ~~, we compare its predictions with those obtained from~~ is compared to the cross-sectional ~~fluxes method-~~ flux (CSF) method implemented in the Python library for data-driven emission quantification

(ddeg<sup>1</sup>). The CSF method was recently compared with other state-of-the-art methods and shows similar accuracy than other well performing methods (?). This method consists in

- 250
- detecting the plume and extracting it from the background. NO<sub>2</sub> fields are used to help in the detection;
  - dividing the plume into a series of horizontal slices of known areas and heights;
  - estimating the line densities of CO<sub>2</sub> by fitting Gaussian curves to the CO<sub>2</sub> and NO<sub>2</sub> concentrations within each slice;
  - inferring the CO<sub>2</sub> fluxes as the product of the line densities and the wind speed at the sources;
  - deriving the total emission rate by multiplying the flux estimation and the area of each slice, and then averaging all
- 255 downstream fluxes.

The CSF method is limited by the need for accurate estimation of the effective wind (?). Two separate wind estimates have been considered: the first derived from an average of the 37 lower levels of ERA 5 data, and the second corresponding to the wind at 100 meters. The first estimate was chosen because of its superior performance. ~~We implement the method with the Python package for data-driven emission quantification (ddeg<sup>2</sup>).~~

## 260 5 Application: inversion of three power plants and model interpretation

In this section, we study the performance of a trained CNN under the conditions exposed in section 4. ~~Specifically, we analyse the potential of CNNs in the inversion of three different PPs:~~

- 265
- firstly, ~~the model is various ensembles of models with different sets of inputs are~~ evaluated on the Lippendorf PP in section ~~??~~. ~~The training dataset is composed of {Berlin, Jänschwalde, Schwarze Pumpe, Boxberg, Turow, Patnow, Opole, Dolna Odra}~~. ~~Furthermore~~ 5.1.1, the Turow PP in section 5.1.2 and the Boxberg PP in section 5.1.3. In each configuration, the training, validation, and test datasets involve 25152, 4608, and 6289 images, respectively;
  - secondly, we investigate how the assimilation of ~~NO<sub>2</sub> and segmentation fields~~ segmentation fields or NO<sub>2</sub> affects the CNNs. ~~Afterwards, since overfitting arises in certain configurations, discussions and partial solutions to this issue are proposed;~~
  - 270 – ~~secondly, in section ??, the model is investigated on the Turow PP, and is~~ thirdly, we propose to interpret the CNNs using a gradient-based technique and by permuting the input features.

### 5.1 Inversion of plumes performance

We study the performance of various CNN model ensembles in predicting the emissions of the Lippendorf, Turow, or Boxberg PPs. More precisely:

---

<sup>1</sup><https://gitlab.com/empa503/remote-sensing/ddeg>

<sup>2</sup>

- 275 – three model ensembles are trained on {Berlin, Jänschwalde, Schwarze Pumpe, Boxberg, Patnow, Lippendorf, Turow, Patnow, Opole, Dolna Odra} and tested on Lippendorf;
- ~~finally, the model is evaluated on the Boxberg PP in section ??.~~ The training dataset includes three model ensembles are trained on {Berlin, Jänschwalde, Schwarze Pumpe, Turow, Patnow, Boxberg, Lippendorf, Patnow, Opole, Dolna Odra} - Since issues of overfitting arise in certain configurations, discussions and partial solutions to these issues are proposed;
- 280 ~~In each configuration, the training, validation, and test datasets involve 25152, 4608, and 6289 images, respectively.~~ Performance evaluation tested on Turow;
- three model ensembles are trained on {Berlin, Jänschwalde, Schwarze Pumpe, Turow, Lippendorf, Patnow, Opole, Dolna Odra} and tested on Boxberg.

~~We study the performance of three models trained with different sets of inputs to predict the emissions of the Lippendorf~~

285 ~~PP.~~ Lippendorf typical emissions fall between those of low-emission PPs (such as Dolna Odra or Turow) and those of high-emission (Boxberg, Jänschwalde). All three models are trained on a dataset composed of {Berlin, Jänschwalde, Schwarze Pumpe, Boxberg, Turow, Patnow, Opole, Dolna Odra} Turow emissions range between 4 and 14 Mt.yr<sup>-1</sup>, similar to PPs like Opole. This implies that most Turow CO<sub>2</sub> plumes are hidden under the background. The last studied PP is Boxberg which is characterised by high emissions and the presence of other PPs in the vicinity, which entails the presence of other high SNR

290 plumes.

For each of the three target PPs, three model ensembles are trained. The input data for the first model ensemble are the XCO<sub>2</sub> field and the wind fields  $u$  and  $v$ . The second and third ~~models~~ model ensembles use the same three base fields and the output of the segmentation model, or the NO<sub>2</sub> field, respectively, as ~~the~~ fourth input. In the following, for simplicity, these ensembles of CNN models will be referred to simply as "models".

295 Kernel Density Estimation (KDE) plots over the 6289 images of the test dataset are drawn in Fig. ~~??~~ 7 comparing with relative and absolute metrics the true emission rates and the predictions of ~~four approaches~~ the four models for each PP. While Fig. 7 employs signed relative error to assess prediction bias in the CNNs, Table 2 and the majority of the analysis rely on absolute relative error. KDE is a non-parametric statistical technique that estimates the probability density function of a continuous random variable by smoothing its observed data points using a kernel function. The three first ensembles of


300 predictions consist of those of the trained CNNs, and the fourth corresponds to the CSF method application. The main statistics corresponding to these KDE plots are summarised in ~~table ??~~ Table 2.

~~The~~

### 5.1.1 Lippendorf plumes inversion

As reported in Table 2, the utilisation of the CNN approach yields remarkably accurate predictions of Lippendorf emissions

305 compared to the CSF method. The performance of all three CNN models shows a median absolute relative error of approximately 20% and a median absolute error of around 3 Mt.yr<sup>-1</sup> (~~where~~ the average emissions for Lippendorf are 15.2 Mt.yr<sup>-1</sup>).



**Figure 7.** Density plots of the signed relative and absolute error between the predicted and the true Lippendorf emissions. Four sets of predictions are considered, corresponding to the three CNN models with three different ensembles-sets of inputs and the CSF method. Each CNN model is trained with the XCO<sub>2</sub> field and the winds as inputs. Two of the models additionally assimilate the NO<sub>2</sub> field or the predictions of the segmentation model. Predictions with relative errors greater (in absolute) than 200%-150% or absolute errors greater than 30 Mt.yr<sup>-1</sup> were set to 200±150 or 30 to increase visibility. 3% of the CSF method predictions are missing. Those predictions correspond to Lippendorf plumes superimposed on other plumes, where the CSF method can not be applied.



	Absolute relative error [%]			Absolute error [Mt.yr <sup>-1</sup> ]		
	25 %	Median	75 %	25 %	Median	75 %
<b><u>Lippendorf</u></b>						
CNN with no additional input	9.8	21.3	35.6	1.4	3.1	5.2
CNN with segmentation	11.4	23.3	36.5	1.6	3.4	5.6
CNN with NO <sub>2</sub>	8.6	18.1	30.3	1.3	2.7	4.5
CSF	21.0	42.8	70.1	3.1	6.3	10.5
<b><u>Turow</u></b>						
<u>CNN with no additional input</u>	<u>12.3</u>	<u>25.9</u>	<u>43.2</u>	<u>1.0</u>	<u>2.2</u>	<u>3.8</u>
<u>CNN with segmentation</u>	<u>11.0</u>	<u>23.7</u>	<u>41.3</u>	<u>0.9</u>	<u>2.0</u>	<u>3.5</u>
<u>CNN with NO<sub>2</sub></u>	<u>10.9</u>	<u>22.9</u>	<u>38.0</u>	<u>0.9</u>	<u>1.9</u>	<u>3.2</u>
<u>CSF</u>	<u>26.3</u>	<u>52.1</u>	<u>92.0</u>	<u>2.2</u>	<u>4.5</u>	<u>8.1</u>
<b><u>Boxberg</u></b>						
<u>CNN with no additional input</u>	<u>11.7</u>	<u>23.5</u>	<u>37.2</u>	<u>2.1</u>	<u>4.4</u>	<u>7.1</u>
<u>CNN with segmentation</u>	<u>24.2</u>	<u>36.9</u>	<u>48.2</u>	<u>4.2</u>	<u>6.8</u>	<u>9.4</u>
<u>CNN with NO<sub>2</sub></u>	<u>26.4</u>	<u>36.9</u>	<u>45.4</u>	<u>4.5</u>	<u>6.8</u>	<u>9.3</u>
<u>CSF</u>	<u>21.7</u>	<u>41.5</u>	<u>63.5</u>	<u>3.9</u>	<u>7.7</u>	<u>12.3</u>

**Table 2.** Relative ~~Absolute relative~~ and absolute error statistics between predicted and true ~~Lippendorf~~ emissions for ~~three~~ CNN models (~~assimilating three different sets of~~ ~~with various~~ inputs )~~and applied to various target PPs~~ and the CSF method.

In comparison, the CSF method exhibits a higher median absolute relative error performance of around 40%, and the absolute error performance is approximately double, at  $6 \text{ Mt.yr}^{-1}$ . ~~These~~ The CNN results are reliable, as the majority of errors are concentrated below  $10 \text{ Mt.yr}^{-1}$  or 50%, with very few exceeding 100%. This indicates that the models provide trustworthy estimates, with a relatively small margin of error.

### 5.1.2 Turow plumes inversion

The CNN approach produces highly accurate predictions for the low SNR Turow plumes, exhibiting similar performance to the results obtained in the Lippendorf case. The three models yield a median absolute relative error performance of approximately 25% and a median absolute error performance of around  $2 \text{ Mt.yr}^{-1}$ . The results can be considered reliable: 75% of the results fall below a threshold of  $4 \text{ Mt.yr}^{-1}$ . By contrast, the CSF method exhibits a median absolute relative error performance of approximately 50 – 55% and an absolute error more than two times larger. The inclusion of segmentation or  $\text{NO}_2$  fields has a noticeable, albeit not significant, impact on the model's performance, resulting in an improvement on the order of a tenth of a  $\text{Mt.yr}^{-1}$ . Notably, the addition of the  $\text{NO}_2$  field appears to have a slightly greater impact compared to the inclusion of segmentation fields. This implies that when applying the model that assimilates the segmented  $\text{XCO}_2$  fields (with the assistance of the  $\text{NO}_2$  fields), there is a potential loss of information on the  $\text{NO}_2$  fields compared to using the model directly assimilating the  $\text{NO}_2$  fields. The observed phenomenon could be a result of potential overfitting when employing the segmentation model, as it is trained on the same dataset as the inversion model. Consequently, the segmentation predictions on the training dataset are typically superior to those on the test dataset. This discrepancy can lead to an over-reliance on the segmentation fields, subsequently causing overfitting.

### 5.1.3 Boxberg plumes inversion

In contrast to the inversion results obtained for Lippendorf and Turow PPs, there is a significant variation in Boxberg plume inversions performance. A CNN only trained with  $\text{XCO}_2$ ,  $u$  and  $v$  winds demonstrates strong performance, comparable to those of the models estimating Lippendorf and Turow emissions. The median absolute relative error performance is approximately 20 – 25%, and the median absolute error performance is around  $4 \text{ Mt.yr}^{-1}$ . The CSF method exhibits a median absolute relative error performance of around 40% and an absolute error performance close to  $8 \text{ Mt.yr}^{-1}$ . But both models with segmentation or  $\text{NO}_2$  fields show a significant decline in performance, which contradicts our expectations. This phenomenon can be attributed to overfitting and is examined in detail in the section 5.2.2.

### 5.1.4 Cross-sectional fluxes method results

The CSF results align with the findings reported in (?) and (?). Other methods, such as the light cross sectional fluxes technique, might yield predictions with absolute relative errors reduced by 5 to 10% (?).

A notable constraint of the CSF method is that it hinges on the estimation of the effective wind speed inside the plume. Under ideal circumstances, the effective wind speed should be estimated using the wind profile, weighted by the  $\text{CO}_2$  concentration

profile, but in practical applications, estimating this profile presents a substantial challenge. In a separate experiment, the CSF method was utilised on a reduced dataset (compared with that used to study the CNN results). Among other changes, this dataset considers only the CO<sub>2</sub>M overpass, or excludes the plume overlaps processed by the CNN. In this experiment, the exact emission profile used to simulate the COSMO-GHG fields of SMARTCARB was used to compute the effective wind. This methodology resulted in a median absolute error around 4Mt.yr<sup>-1</sup>, thereby indicating that the effective wind estimation significantly contributes to the errors associated with the CSF method. ~~The CNN results are reliable, as the majority of errors are concentrated below 10Mt.yr<sup>-1</sup> or 50%, with very few exceeding 100%. This indicates that the models provide trustworthy estimates, with a relatively small margin of error~~


## 5.2 Results analysis

This section presents an analysis of the results from the preceding section 5.1. It involves examining the trained models, or new models trained under the same configuration, across various datasets, including the test dataset. It is important to note that these analyses were conducted after obtaining the previous section's results and thus did not influence, e.g., the choice of model architecture or hyperparameters.

### 5.2.1 Study of the addition of the segmentation and NO<sub>2</sub> fields

~~Surprisingly, the addition of the segmentation results in the inputs does not improve the inversion process~~Interestingly, the model designed to invert the Lippendorf plumes does not yield more accurate emission estimates when using segmentations as inputs, compared to the model without segmentations. One possible reason for this lack of improvement is that the Lippendorf segmentation fields are inadequate. The segmentation model assimilates NO<sub>2</sub> fields to segment the CO<sub>2</sub> fields and the presence of multiple alternative NO<sub>2</sub> plumes in the NO<sub>2</sub> fields hinders the segmentation model's ability to accurately delineate the contour of the Lippendorf CO<sub>2</sub> plume. However, the incorporation of NO<sub>2</sub> field as inputs slightly increase the quality of the predictions made by the new CNN model. One hypothesis to explain the error discrepancy between the model utilising NO<sub>2</sub> and the one using segmentation fields is due to the segmentation model not capturing NO<sub>2</sub> or CO<sub>2</sub> plume amplitude variations. Precisely, the segmentation model does not discriminate between plume pixels with high amplitude and those with low amplitude. Consequently, if the NO<sub>2</sub> images are of poor quality due to the presence of numerous alternative NO<sub>2</sub> plumes, the segmentation model will struggle to distinguish the main plume accurately. Taking into account these segmentation fields, which do not distinguish the background plumes in the test dataset, results in a degradation of the predictions.

Regarding the benefits brought by the integration of NO<sub>2</sub> fields, their essential contribution is their facilitation of the plume segmentation, rather than a direct enablement of inversion based on the NO<sub>2</sub> levels alone. Indeed, as stated in section 4.2.1, the NO<sub>2</sub> plumes are not scaled like CO<sub>2</sub> plumes. To further investigate this, it is possible to verify this hypothesis by modifying the scaling of the plume in the NO<sub>2</sub> fields. In Fig. ??-8 we draw the relative absolute error of the model at testing stage when taking as inputs scaled NO<sub>2</sub> fields by a constant factor in [0, 3]. The CNN model exhibits its highest performance within the scaling range of 1.0 to 1.3. As expected, performances gradually decrease for scalings-scaling below 1 and above 1.3. However, intriguingly, the model still achieves remarkably satisfactory scores for scalings ranging from 0.5 to 1.75. Notably, these scores



**Figure 8.** Effect of scaling the NO<sub>2</sub> plume of Lippendorf inputs on the performance of the CNN model. The x-axis corresponds to the scaling of the NO<sub>2</sub> plume: 1 corresponds to the original plume, 0 to no plume. The y-axis corresponds to the median absolute relative error of the CNN model evaluated for the given scaling of the NO<sub>2</sub> plumes. The CNN model is the same for each scaling (each dot of the x-axis) and corresponds to the CNN model having obtained the best absolute relative error score. The red dotted line approximately corresponds to the median absolute relative error of a model not learning with NO<sub>2</sub> fields.

outperform those of the ~~ensemble of~~ CNN models without NO<sub>2</sub> field as input. If the model were utilising the amplitude of the NO<sub>2</sub> field as a predictor, deeply inaccurate results could be expected for scalings of 0.5 or 1.5. However, the graph contradicts this assumption, strongly suggesting that the amplitude of the NO<sub>2</sub> field is not employed as a predictor by the model. Instead, it is likely that only the contour of the NO<sub>2</sub> plume and the ratios between different parts of the plume serve as predictors. In essence, the model's reliance on the NO<sub>2</sub> field for predictions appears to be based on the contour of the plume and the relative proportions of its various components, rather than on the absolute amplitude of the NO<sub>2</sub> field values. Finally, the model's inability to accurately estimate emissions for scaling factors exceeding 2 can be attributed to the unprecedented and extreme values that NO<sub>2</sub> plumes can reach, which lie beyond the range of what the model has been trained on.

~~Performance evaluation~~In this section, we study the performance of three models with input similar to those of section ??, with this time a focus on a low-emission PP: Turow. Turow emissions range between 4 and 14 Mt.yr<sup>-1</sup>, similar to PPs like Dolna Odra or Opole. This implies that most Turow CO<sub>2</sub> plumes are hidden under the background. All three models considered are trained on a dataset consisting of {Berlin, Janschwalde, Schwarze Pumpe, Boxberg, Lippendorf, Patnow, Opole, Dolna Odra}.

In a similar manner to section ??, KDE plots are presented in Fig. ?? to compare the true emission rates with the predictions of the four different approaches using relative and absolute metrics and corresponding statistics are summarised in table ??.

Density plots of the relative and absolute error between the predicted and the true Turow emissions. Four sets of predictions are considered, corresponding to the three CNN models with three different ensembles of inputs and the CSF method. Each CNN model is trained with the XCO<sub>2</sub> field and the winds as inputs. Two of the models additionally assimilate the NO<sub>2</sub> field or the predictions of the segmentation model. Predictions with relative errors greater than 200% or absolute errors greater than 30 Mt.yr<sup>-1</sup> were set to 200 or 30 to increase visibility.

25% Median 75% 25% Median 75%

CNN with no additional input 12.3 25.9 43.2 1.0 2.2 3.8 CNN with segmentation 11.0 23.7 41.3 0.9 2.0 3.5 CNN with NO<sub>2</sub> 10.9 22.9 38.0 0.9 1.9 3.2 CSF 26.3 52.1 92.0 2.2 4.5 8.1


Relative and absolute error statistics between predicted and true Turow emissions for three CNN models (assimilating three different sets of inputs) and the CSF method.

The CNN approach produces highly accurate predictions for these low SNR plumes, exhibiting similar performance to the results obtained in the Lippendorf case. The three models yield a median relative error performance of approximately 25% and a median absolute error performance of around 2 Mt.yr<sup>-1</sup>. The results can be considered reliable: 75% of the results fall below a threshold of 4 Mt.yr<sup>-1</sup>. By contrast, the CSF method exhibits a median relative error performance of approximately 50–55% and an absolute error performance more than two times larger.

The inclusion of segmentation or NO<sub>2</sub> fields has a noticeable, albeit not significant, impact on the model's performance, resulting in an improvement on the order of a tenth of a Mt.yr<sup>-1</sup>. Notably, the addition of the NO<sub>2</sub> field appears to have a slightly greater impact compared to the inclusion of segmentation fields. This implies that when applying the model that assimilates the segmented XCO<sub>2</sub> fields (with the assistance of the NO<sub>2</sub> fields), there is a potential loss of information on the NO<sub>2</sub> fields compared to using the model directly assimilating the NO<sub>2</sub> fields. The observed phenomenon could be a result of potential overfitting when employing the segmentation model, as it is trained on the same dataset as the inversion model. Consequently, the segmentation predictions on the training dataset are typically superior to those on the test dataset. This discrepancy can lead to an over-reliance on the segmentation fields, subsequently causing overfitting.

Performance evaluation The last studied PP is Boxberg which is characterised by high emissions and the presence of other PPs in the vicinity, which entails the presence of other high SNR plumes. We study the performance of three models, all trained on a dataset consisting of {Berlin, Jänschwalde, Schwarze Pumpe, Lippendorf, Turow, Patnow, Opole, Dolna Odra}. The three models differ in their inputs, mirroring the approach taken in sections ?? and ?. Figure ?? and table ?? present a comparison between the true emission rates and the corresponding predictions.

Density plots of the relative and absolute error between the predicted and the true Boxberg emissions. Four sets of predictions are considered, corresponding to the three CNN models with three different ensembles of inputs and the CSF method. Each CNN model is trained with the XCO<sub>2</sub> field and the winds as inputs on a dataset composed of {Berlin, Jänschwalde, Schwarze Pumpe, Lippendorf, Turow, Patnow, Opole, Dolna Odra}. Two of the models additionally assimilate the NO<sub>2</sub> field or the predictions of the segmentation model. Predictions with relative errors greater than 200% or absolute errors greater than 30 Mt.yr<sup>-1</sup> were set to 200 or 30 to increase visibility. 15% of the CSF method predictions are missing. Those predictions correspond to Boxberg plumes superimposed on other plumes, where the CSF method cannot be applied.



**Figure 9.** Residuals density between the true emissions of Boxberg PP and the predictions of the CNNs.

~~25% Median 75% 25% Median 75%~~


~~CNN with no additional input 11.7 23.5 37.2 2.1 4.4 7.1 CNN with segmentation 24.2 36.9 48.2 4.2 6.8 9.4 CNN with NO<sub>2</sub> 26.4 36.9 45.4 4.5 6.8 9.3 CSF 21.7 41.5 63.5 3.9 7.7 12.3~~

~~Relative and absolute error statistics between predicted and true Boxberg emissions for three CNN models (assimilating three different sets of inputs) and the CSF method.~~

~~In contrast to the inversion results obtained for Lippendorf and Turow PPs, there is a significant variation in performance among the three CNN models. A CNN only trained with XCO<sub>2</sub>, *u* and *v* winds demonstrates strong performance, comparable to those of the models estimating Lippendorf and Turow emissions in sections ?? and ?. The median relative error performance is approximately 20–25%, and the median absolute error performance is around 4Mt.yr<sup>-1</sup>. By contrast, the CSF method exhibits a median relative error performance of around 40% and an absolute error performance close to 8Mt.yr<sup>-1</sup>. But both models with segmentation or NO<sub>2</sub> fields show a significant decline in performance, which contradicts our expectations. This phenomenon can be attributed to overfitting and is examined in detail in the subsequent sub-section.~~

### 5.2.2 Overfitting investigation

To understand the reasons behind the deviations between predictions and actual values, we conducted an analysis of the residuals in Fig. ???. This examination, which focuses on the disparities between predicted and true emissions, reveals a substantial underestimation of the actual Boxberg emissions by the model incorporating NO<sub>2</sub> or segmentation fields.




**Figure 10.** Evolution during training of the validation and test relative errors between the true emissions of Boxberg and the predictions of the CNNs. Specifically, for ~~each epoch~~, the need of this experiment, new models being trained are trained and employed to predict the emissions corresponding to the validation and test fields at each epoch. Three models are considered: each is trained with the XCO<sub>2</sub> field and the winds as inputs. Two of the models additionally assimilate the NO<sub>2</sub> field or the predictions of the segmentation model. The validation error decreases monotonically with the number of epochs, while the test error does not, which suggests overfitting of the model.

This observation suggests an overfitting issue (?), since the majority of PPs used to train the model exhibit lower emission rates compared to Boxberg (6 out of 7: Schwarze Pumpe, Lippendorf, Turow, Patnow, Opole, and Dolna Odra). The low absolute relative error observed on the training dataset of the models with NO<sub>2</sub> or segmentation fields, as depicted in Fig. 440 ??10, further substantiates concerns regarding overfitting. An overfitting-overfitted model tends to learn features that are overly tailored to the specifics of the training dataset. Consequently, when presented with images from the test dataset with slightly different features, the model struggles to generate accurate predictions.

Our hypothesis is that the model specifically overfits the low emissions PPs, which can be attributed to the information gained through the use of NO<sub>2</sub> or segmentation fields. When provided with new images, the model fails to recognise the new 445 features and consequently yields predictions that align more closely with the training dataset, which predominantly consists of emissions from the low emissions PPs. The failure to recognise new features is due to the incorporation of NO<sub>2</sub> field as inputs. The model's ability to learn highly specific features is limited when no additional input is provided. Conversely, when the model incorporates the NO<sub>2</sub> field, it gains access to more information and can acquire more intricate features. Consequently, the model's capacity to generalise worsens in the latter case.





**Figure 11.** Density plots of the relative and absolute error between the predicted and the true emissions. Four sets of predictions are considered, corresponding to the three CNN models with three different ensembles-sets of inputs and the CSF method. Each CNN model is trained with the  $XCO_2$  field and the winds as inputs on a dataset composed of {Berlin, Janschwalde, Lippendorf, Turow, Opole}. Two of the models additionally assimilate the  $NO_2$  field or the predictions of the segmentation model. Predictions with absolute relative errors greater than ~~200~~150% or absolute errors greater than  $30 \text{ Mt.yr}^{-1}$  were set to ~~200~~150 or 30 to increase visibility. 15% of the CSF method predictions are missing. Those predictions correspond to Boxberg plumes superimposed on other plumes, where the CSF method cannot be applied. A modified dataset is used to avoid overfitting.

450 ~~It is worth noting that some CNN models that we trained incorporating  $NO_2$  or segmentation fields do not exhibit overfitting and provide highly accurate predictions. However, these models do not perform better on the validation dataset, rendering them unsuitable for selection.~~

Next, we examine the performance of a new ensemble of models trained on a more balanced dataset, achieved by removing three out of the five low-emissions PP from the original dataset. The new dataset is composed of {Berlin, Janschwalde, Lippendorf, Turow, Opole}, i.e. of two medium or high emissions PPs, two low emissions PPs and Berlin. Furthermore, the choice of the factor scaling the plumes during training (see section 4.2.1) varies depending on the ensemble member considered (see section 4.2.4). Specifically, the uniform distribution is defined with a minimum scaling factor of 0.25 or 0.5 and a maximum scaling factor of 2 or 3. Once more, we examine the outcomes for three models obtained from ensembling, as depicted in Fig. ~~??~~11, and summarised in ~~table ??~~Table 3.

460 The three CNN models demonstrate very good performance, although the inclusion of  $NO_2$  or segmentation fields still leads to a degradation in results, albeit to a lesser extent than in section ~~??~~5.1.3. For example, the median absolute relative error for

	Absolute relative error [%]			Absolute error [Mt.yr <sup>-1</sup> ]		
	25 %	Median	75 %	25 %	Median	75 %
CNN with no additional input	9.5	20.4	33.8	1.8	3.7	6.3
CNN with segmentation	14.1	26.6	40.6	2.5	4.8	7.6
CNN with NO <sub>2</sub>	13.2	23.8	34.2	2.3	4.4	6.6
CSF	21.7	41.5	63.5	3.9	7.7	12.3

**Table 3.** Relative and absolute error statistics between predicted and true Boxberg emissions for three CNN models (assimilating three different sets of inputs) and the CSF method. A modified dataset is used to avoid overfitting.

[the CNN with NO<sub>2</sub> as additional input is 23.8%, comparing to 36.9% in section 5.1.3.](#) On the one hand, the median [absolute](#) relative error of the CNN model trained on NO<sub>2</sub> and wind fields stands at approximately 20%, while the median absolute error remains below 4Mt.yr<sup>-1</sup>. On the other hand, the degradation of the results when adding the NO<sub>2</sub> or segmentation fields still  
465 can be regarded as overfitting. The model learns features from NO<sub>2</sub> or segmentation fields that are not general enough to cover the case of Boxberg. Furthermore, it fails to acquire compensatory generalisable features such as in the case of Turow, where the model probably gains information about the plume contour from the NO<sub>2</sub> field, which is not straightforwardly apparent in the Turow XCO<sub>2</sub> field.

### 5.3 Interpretation of the CNN inversion models

470 In the two following sections, we introduce and apply two methods to gain a deeper understanding of the behaviour and decision-making processes of the CNNs discussed in this paper. These methods offer valuable insights into the significance of input features in the predictions made by the CNNs:

- the integrated-gradient method allows us to examine the importance of individual pixels across channels;
- the feature permutation method enables us to assess the importance of the channels, i.e. the fields used as inputs.

#### 475 5.3.1 Gradient-based study of the pixels

Integrated Gradients is a gradient-based method for the interpretability of neural networks that enables the assessment of pixel importance in CNN prediction. It calculates the sensitivity of a model’s predictions to input features (here pixels), assigning relevance scores to them. By analysing how changes in the input pixels affect the model’s output, the method provides insights into the importance of each pixel in the decision-making process.

480 One unique aspect of this gradient-based method is that the scores it assigns are relative to a baseline. More precisely, these scores are computed as integrated gradients along a linear interpolation from a blank image to the input.

In Fig. 12, we apply the Integrated Gradient approach to study four different models specific to various sources. The first and second models are built to invert the emissions from Lippendorf (see section [??5.1.1](#)) and Boxberg plumes (see section [??5.1.3](#)), respectively, whereas the two last models target Turow plumes (see section [??5.1.2](#)). The first three models only use  
485 the XCO<sub>2</sub> field and the winds as inputs, whereas the fourth model considers an additional input, the NO<sub>2</sub> field. To apply the Integrated Gradient method, a random plume from the target PP is selected for each model. Columns of Fig. 12 represent the XCO<sub>2</sub> field, the corresponding XCO<sub>2</sub> plume and the Integrated Gradient between the model predictions and the inputs.


In order to simplify the analysis, we choose the model that exhibits the best performance on the test dataset, rather than using the ensemble of models. It is worth noting that such a model yields similar performances to the ensemble of models.

490 The Integrated Gradient technique reveals that the CNN model learns to estimate the emissions of a source based on the pixels of the plume from this source. In the first row, we examine a plume from Lippendorf PP. The Integrated Gradient technique identifies the most important pixels, which correspond to the plume pixels. This indicates that if the pixels associated with the plume were to deviate, the estimated flux rate would be significantly affected. This demonstrates that the CNN model effectively makes inference from the crucial parts of the image. In the second row, we focus on an image centred on Boxberg.  
495 The gradients reveal that the model concentrates exclusively on the Boxberg plume in the centre, disregarding the other plumes when inferring the emissions of Boxberg PP. In the third row of the figure, we examine a plume from Turow PP. Although the precision is lower compared to the Lippendorf or Boxberg cases, the pixels in the general direction of the Turow plume are the main ones used to estimate the emissions. In the fourth row, we display the gradients associated with the same Turow image but for a model trained with NO<sub>2</sub> fields as additional inputs. In this case, the model clearly identifies the pixels corresponding to the  
500 plume as critical, as indicated by the amplitude and contour of the gradients. This reinforces the hypothesis that the improved estimation of Turow emissions when the model is trained with NO<sub>2</sub> fields, can be attributed to the enhanced assessment of the plumes.

### 5.3.2 Feature permutation analysis

Feature permutation is a technique used to determine the importance of input channels used in a model (?). As input variables used here are not independent, the interpretation of the following permutation analysis should be taken with caution.  
505 The principle is to i) permute the values of a feature (e.g. exchange the  $u$  wind field corresponding to an XCO<sub>2</sub> image with another random  $u$  wind field), and ii) use the model to predict emissions for the given input, which includes the XCO<sub>2</sub> field, other associated inputs and the random  $u$  wind field. By comparing the performance of the model on the original dataset with the performance on the permuted dataset, we can measure the impact of each feature on the performance of the model. The more  
510 the permutation of a feature affects the performance of the model, the more important that feature is. In [table-Table 4](#), we present the outcomes of the permutation of the features for nine ensembles of models (for each PP and each ensemble of inputs considered). Each entry in the table represents the degradation in the average [absolute](#) relative error of the model (associated with a specific column) when the corresponding feature is permuted (related to the respective row).

We can formulate hypotheses on several groups of cells in [table-Table 4](#):



**Figure 12.** Evaluation of four CNN models using the Integrated Gradients method on four input sets. Columns 1, 2, and 3 represent the XCO<sub>2</sub> field, the corresponding XCO<sub>2</sub> plume and the Integrated Gradient between the model predictions and the inputs, in that order. The rows represent the four models and corresponding test fields.

PP Feature	Lippendorf			Boxberg			Turow		
	None	Seg.	NO <sub>2</sub>	None	Seg.	NO <sub>2</sub>	None	Seg.	NO <sub>2</sub>
XCO <sub>2</sub>	18.3	27.6	32.7	19.6	29.5	34.5	12.6	9.8	22.7
u-wind	11.6	10.4	3.3	12.4	9.7	4.2	6.6	5.4	0.9
v-wind	4.9	3.7	1.8	5.3	4.9	3.0	3.6	1.7	0.6
<del>Fourth feature</del> <u>Add. input</u>	n/a	22.7	30.3	n/a	28.9	34.6	n/a	6.9	21.0

**Table 4.** Evaluation of the degradation in the average absolute relative error of the model when the corresponding feature is permuted. Each column corresponds to a model, and each row corresponds to a permuted feature. For example, we estimate the emissions corresponding to Boxberg plumes with the CNN model trained with the XCO<sub>2</sub> field, the winds and no additional input. We then estimate the degradation of the predictions of this model on the test dataset with the u-wind fields permuted: the degradation in absolute relative error is 12.4%. ~~Several cells of importance of this table are coloured and discussed thereafter~~ Seg. is for Segmentation. Add. input corresponds to either None, NO<sub>2</sub>, or Segmentation fields depending on the considered column.

- 515 – ~~in the highlighted yellow instances, we verify the anticipated observation~~ comparison of the first row with the others shows that the XCO<sub>2</sub> feature consistently holds (or shares) the highest importance among other features ~~(here, in comparison to u-wind) or shares the highest level of importance;~~
- ~~in the highlighted orange section, we emphasise a trend indicating that~~ examining the second and third rows reveals a pattern where the u-wind ~~appears seems~~ to have a greater impact more significant influence on the inversion ~~than process compared to the~~ v-wind. It might be due to the bigger variance of  $u$  ( $\sim 51 \text{ m}^2 \cdot \text{s}^{-2}$ ) comparing to  $v$  ( $\sim 19 \text{ m}^2 \cdot \text{s}^{-2}$ ):  $u$  is the dominant wind in this situation;
- 520 – ~~purple cells are examples of the XCO<sub>2</sub> field inputs having less importance~~ input permutation seems to have a lesser impact for the CNN targeting Turow. ~~The observed behaviour is due to the low SNR of Turow plumes. Due to inconsistent ancillary (winds, ...) data and Turow plume, the model tends to estimate emissions from points in the image other than the plume. These points are less distinct from the plume than in the scenario with the Boxberg plume. As a result, the relative error is lower~~ (12.6% in absolute relative error degradation comparing to 18.3 and 19.6% for Lippendorf and Boxberg);
- 525 – ~~in the highlighted red section, we underline the increase~~ the importance of the XCO<sub>2</sub> feature ~~importance differs for models with different inputs: specifically, the XCO<sub>2</sub> feature importance increases~~ when additional data are used. ~~For example, the XCO<sub>2</sub> feature permutation degrades the performance of the model by 32.7% when NO<sub>2</sub> is used, and by 18.3% when no additional input is used.~~ This observation is consistent with the overfitting tendency of the CNN models when trained with ~~for example, NO<sub>2</sub> field~~ NO<sub>2</sub> fields. When confronted with inconsistent data (non-corresponding
- 530

XCO<sub>2</sub> and NO<sub>2</sub> fields), the acquired complex features of the model exhibit a total absence of correspondence with these inputs. As a consequence, the model predicts nonsensical emissions;

535 – the ~~brown cells reveal that the wind systematically holds~~ wind inputs systematically hold greater importance in the model without additional input, followed by a relatively diminished importance in the model with segmentation fields, and finally, it exhibits the least significance in the model with NO<sub>2</sub> fields. This suggests that the inclusion of NO<sub>2</sub> fields is more advantageous for inversion compared to segmentation fields. Furthermore, it ~~implies indicates~~ that the winds ~~are used to compensate for the absence or deficiency~~ compensate the absence of additional data.

## 540 6 Discussions and limitations

The approach developed in this paper carries certain limitations: firstly, it focuses exclusively on European PPs, which questions its generalisability to PPs in other regions with different climatic conditions. Secondly, the study emphasises the importance of a balanced dataset, as highlighted by the ~~??-section~~ section 5.2.2, and the need to be able to identify and address potential overfitting issues. It should also be noted that the study did not specifically investigate outliers such as PPs with exceptionally

545 high emission rates. Despite incorporating plume scaling strategies, the model may struggle to generalise to such outliers.

In terms of future research, several areas should be explored such as the challenge posed by clouds. In this respect, CNNs can be trained to ignore missing data caused by cloud cover and to make effective use of the available data. Another aspect to consider is the presence of noise in CO<sub>2</sub>M data. While Gaussian noise may not pose significant issues, if the satellite noise exhibits structured patterns, it would become crucial to develop robust noise modelling techniques to enable CNNs to

550 accurately distinguish and remove such noise. Finally, in real-world applications, training models trained on synthetic datasets ~~and applying the trained models may face challenges when applied~~ to real datasets ~~may encounter distributional differences due to differences in data distribution~~. Strategies such as importance weighting, specific data augmentation techniques, transfer learning, or active learning methods may be necessary to account for these differences and ensure reliable performance.

## 7 Conclusions and perspectives

555 The future availability of CO<sub>2</sub> satellite imagery, through missions such as the Copernicus CO<sub>2</sub> Monitoring (CO<sub>2</sub>M) mission, heralds new ~~opportunities~~ opportunities and challenges for the evaluation of local CO<sub>2</sub> emissions. Emissions can certainly be retrieved from CO<sub>2</sub> plumes of hotspots in the satellite images. But the emission estimation is hindered by two primary obstacles: low signal-to-noise ratio plumes which cannot be extracted straightforwardly from the background, and uncertainties in the transport or dispersion processes which hampers the assessment of the emissions from the plumes.

560 In this work, we assess the ability of Convolutional Neural Networks (CNNs) to invert a plume from satellite imagery using simulated XCO<sub>2</sub>, NO<sub>2</sub>, and wind fields with similar characteristics to the future CO<sub>2</sub>M images. The fields used to train and evaluate the CNNs are based on the SMARTCARB dataset and possess the same resolution, satellite noise level and ancillary data availability to the future CO<sub>2</sub>M images. Each synthetic XCO<sub>2</sub> image encompasses the anthropogenic plume from at least

one power plant, a background arising from other biogenic and anthropogenic fluxes, and a random Gaussian noise intended  
565 to simulate the errors inherent in satellite instruments. But the model evaluation was conducted using simulated data. This  
approach does not account for all the challenges that real satellite images present, specifically issues related to cloud cover and  
systematic error patterns due to surface reflectance and the aerosol dependency of retrievals.

Our source emission estimation model is an image-to-scalar CNN model, which infers from the full XCO<sub>2</sub> field a flux rate,  
estimation of the anthropogenic emission corresponding to the plume. The first layers of the model consist of preprocessing  
570 steps which transform the input data at training time, in particular by adding noise and by scaling the plume and emission rate.  
The core model is a CNN composed of approximately 200,000 neurons divided into convolutional, max pooling, dropout and  
batch normalisation layers.

We ~~demonstrated~~ highly suggest that the design of a "universal" CNN, trained on a small power plant subset and highly  
accurate on all of them, is possible. To do so, we ~~evaluated~~ evaluate the model's ability to generalise to unobserved images  
575 from another region. Explicitly, three CNN models ~~were trained on different datasets but sharing the exact same structure~~  
(hyperparametrisation, architecture, ...) are tested on plumes from three sources: Boxberg, Lippendorf and Turow. The training/validation  
dataset for each CNN is restricted to a dataset consisting of all other power plants except their target. The CNNs  
are highly accurate in each case and the addition of NO<sub>2</sub> fields often improves the results slightly. Precisely, the median absolute  
relative errors for the CNN models are on average close to 20–25%. Moreover, the median absolute error is generally half that  
580 obtained with the CSF method: an alternative and state of the art inversion approach. This highly suggests that it is possible to  
build a universal neural network (which can generalise to all targets) using this methodology.

By using interpretability tools, we demonstrate that the predictions made by the CNNs are grounded in the physically  
meaningful components of the features. The Integrated Gradient method shows that the CNNs learn to predict the emissions  
corresponding to a plume from the pixels making up the plume. The feature permutation technique highlighted several aspects  
585 of the models, such as the expected high importance of the XCO<sub>2</sub> fields compared to the used ancillary data.

Future prospects of the CNN plume inversion method from satellite images encompass the challenges of clouds, cities and  
real satellite images. Concretely, the method should be able to handle missing data caused by clouds. Additionally, the CNN  
approach should incorporate the second important category of hotspots: cities. Finally, the method should be tested on real  
satellite data once it becomes available.

590 *Data availability.* The datasets used in this paper are available on a compliant repository on <https://zenodo.org/record/8096616> and originate  
from <https://zenodo.org/record/4048228>. The weights of the CNNs are available on <https://zenodo.org/record/8095487>. The algorithms are  
available on Zenodo (<https://zenodo.org/record/10100338>) and Github at <https://github.com/cerea-daml/co2-images-inv-dl>.

*Author contributions.* Joffrey Dumont Le Brazidec: Conceptualisation, Methodology, Software, Investigation, Formal analysis, Visualisation,  
Resources, Project administration, Writing - Original Draft; Pierre Vanderbecken: Investigation, Formal analysis, Writing - Review;



595 Alban Farchi: Conceptualisation, Methodology, Project administration, Writing - Review; Marc Bocquet: Conceptualisation, Methodology,  
Project administration, Funding acquisition, Writing - Review; Grégoire Broquet: Conceptualisation, Writing - Review; Gerrit Kuhlmann:  
Resources, Writing - Review.

*Competing interests.* The authors declare that they have no conflict of interest.

600 *Acknowledgements.* This project has been funded by the European Union's Horizon 2020 research and innovation programme under grant  
agreement N° 958927 (Prototype system for a Copernicus CO<sub>2</sub> service). CEREIA is a member of Institut Pierre-Simon Laplace (IPSL). We  
would like to acknowledge Tobias Finn for the valuable thought-provoking discussions, as well as Evan D. Sherwin and our second reviewer  
for their insightful comments.

## References

- Beirle, S., Borger, C., Dörner, S., Li, A., Hu, Z., Liu, F., Wang, Y., and Wagner, T.: Pinpointing nitrogen oxide emissions from space, *Science Advances*, 5, eaax9800, <https://doi.org/10.1126/sciadv.aax9800>, publisher: American Association for the Advancement of Science, 2019.
- 605 Broquet, G., Bréon, F.-M., Renault, E., Buchwitz, M., Reuter, M., Bovensmann, H., Chevallier, F., Wu, L., and Ciais, P.: The potential of satellite spectro-imagery for monitoring CO<sub>2</sub> emissions from large cities, *Atmospheric Measurement Techniques*, 11, 681–708, <https://doi.org/10.5194/amt-11-681-2018>, publisher: Copernicus GmbH, 2018.
- Brunner, D., Kuhlmann, G., Marshall, J., Clément, V., Fuhrer, O., Broquet, G., Löscher, A., and Meijer, Y.: Accounting for the vertical distribution of emissions in atmospheric CO<sub>2</sub> simulations, *Atmos Chem Phys*, 19, 4541–4559, <https://doi.org/10.5194/acp-19-4541-2019>, publisher: Copernicus GmbH, 2019.
- 610 Chevallier, F., Remaud, M., O'Dell, C. W., Baker, D., Peylin, P., and Cozic, A.: Objective evaluation of surface- and satellite-driven carbon dioxide atmospheric inversions, *Atmospheric Chemistry and Physics*, 19, 14 233–14 251, <https://doi.org/10.5194/acp-19-14233-2019>, publisher: Copernicus GmbH, 2019.
- 615 Chevallier, F., Broquet, G., Zheng, B., Ciais, P., and Eldering, A.: Large CO<sub>2</sub> Emitters as Seen From Satellite: Comparison to a Gridded Global Emission Inventory, *Geophysical Research Letters*, 49, e2021GL097 540, <https://doi.org/10.1029/2021GL097540>, \_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2021GL097540>, 2022.
- Chollet, F.: *Deep Learning with Python*, Manning Publications Co., USA, 1st edn., 2017.
- Dumont Le Brazidec, J., Vanderbecken, P., Farchi, A., Bocquet, M., Lian, J., Broquet, G., Kuhlmann, G., Danjou, A., and Lauvaux, T.: Segmentation of XCO<sub>2</sub> images with deep learning: application to synthetic plumes from cities and power plants, *Geoscientific Model Development Discussions*, pp. 1–29, <https://doi.org/10.5194/gmd-2022-288>, publisher: Copernicus GmbH, 2022.
- 620 Finch, D., Palmer, P., and Zhang, T.: Automated detection of atmospheric NO<sub>2</sub> plumes from satellite data: a tool to help infer anthropogenic combustion emissions, *Atmos. Meas. Tech.*, pp. 1–21, <https://doi.org/10.5194/amt-2021-177>, 2021.
- Hakkarainen, J., Szelağ, M. E., Ialongo, I., Retscher, C., Oda, T., and Crisp, D.: Analyzing nitrogen oxides to carbon dioxide emission ratios from space: A case study of Matimba Power Station in South Africa, *Atmospheric Environment: X*, 10, 100 110, <https://doi.org/10.1016/j.aeaoa.2021.100110>, 2021.
- 625 Hakkarainen, J., Ialongo, I., Koene, E., Szelağ, M., Tamminen, J., Kuhlmann, G., and Brunner, D.: Analyzing Local Carbon Dioxide and Nitrogen Oxide Emissions From Space Using the Divergence Method: An Application to the Synthetic SMARTCARB Dataset, *Frontiers in Remote Sensing*, 3, <https://doi.org/10.3389/frsen.2022.878731>, 2022.
- 630 Hakkarainen, J., Tamminen, J., Nurmela, J., Santaren, D., Broquet, G., Chevallier, F., Koene, E., Kuhlmann, G., and Brunner, D.: D4.4 Benchmarking of plume detection and quantification methods | CoCO<sub>2</sub>: Prototype system for a Copernicus CO<sub>2</sub> service, Technical Report 4.4, <https://coco2-project.eu/node/366>, 2023.
- IEA, I. E. A.: Fuel share of CO<sub>2</sub> emissions from fuel combustion, 2019 – Charts – Data & Statistics, <https://www.iea.org/data-and-statistics/charts/fuel-share-of-co2-emissions-from-fuel-combustion-2019>, 2019.
- 635 Jongaramrungruang, S., Matheou, G., Thorpe, A. K., Zeng, Z.-C., and Frankenberg, C.: Remote sensing of methane plumes: instrument tradeoff analysis for detecting and quantifying local sources at global scale, *Atmospheric Measurement Techniques*, 14, 7999–8017, <https://doi.org/10.5194/amt-14-7999-2021>, publisher: Copernicus GmbH, 2021.

- Joyce, P., Ruiz Villena, C., Huang, Y., Webb, A., Gloor, M., Wagner, F. H., Chipperfield, M. P., Barrio Guilló, R., Wilson, C., and Boesch, H.: Using a deep neural network to detect methane point sources and quantify emissions from PRISMA hyperspectral satellite images, *Atmospheric Measurement Techniques*, 16, 2627–2640, <https://doi.org/10.5194/amt-16-2627-2023>, publisher: Copernicus GmbH, 2023.
- 640 Koene, E., Brunner, D., and Kuhlmann, G.: Documentation of plume detection and quantification methods, Technical Report 4.3, Empa, <https://www.coco2-project.eu/node/329>, 2021.
- Kuener, J. J. P., Visschedijk, A. J. H., Jozwicka, M., and Denier van der Gon, H. a. C.: TNO-MACC\_II emission inventory; a multi-year (2003&ndash;2009) consistent high-resolution European emission inventory for air quality modelling, *Atmos Chem Phys*, 14, 10963–10976, <https://doi.org/10.5194/acp-14-10963-2014>, 2014.
- 645 Kuhlmann, G., Broquet, G., Marshall, J., Clément, V., Löscher, A., Meijer, Y., and Brunner, D.: Detectability of CO<sub>2</sub> emission plumes of cities and power plants with the Copernicus Anthropogenic CO<sub>2</sub> Monitoring (CO2M) mission, *Atmos. Meas. Tech.*, 12, 6695–6719, <https://doi.org/10.5194/amt-12-6695-2019>, 2019.
- Kuhlmann, G., Brunner, D., Broquet, G., and Meijer, Y.: Quantifying CO<sub>2</sub> emissions of a city with the Copernicus Anthropogenic CO<sub>2</sub> Monitoring satellite mission, *Atmos. Meas. Tech.*, 13, 6733–6754, <https://doi.org/10.5194/amt-13-6733-2020>, 2020.
- 650 Kuhlmann, G., Henne, S., Meijer, Y., and Brunner, D.: Quantifying CO<sub>2</sub> Emissions of Power Plants With CO<sub>2</sub> and NO<sub>2</sub> Imaging Satellites, *Front. remote sens.*, 2, <https://www.frontiersin.org/article/10.3389/frsen.2021.689838>, 2021.
- Lary, D. J., Alavi, A. H., Gandomi, A. H., and Walker, A. L.: Machine learning in geosciences and remote sensing, *Geoscience Frontiers*, 7, 3–10, <https://doi.org/10.1016/j.gsf.2015.07.003>, 2016.
- 655 Meijer, Y.: Copernicus CO<sub>2</sub> Monitoring Mission Requirements Document, Earth and Mission Science Division, 84, [https://esamultimedia.esa.int/docs/EarthObservation/CO2M\\_MRD\\_v3.0\\_20201001\\_Issued.pdf](https://esamultimedia.esa.int/docs/EarthObservation/CO2M_MRD_v3.0_20201001_Issued.pdf), 2020.
- Molnar, C.: *Interpretable Machine Learning: A Guide For Making Black Box Models Explainable*, Independently published, Munich, Germany, 2022.
- Nassar, R., Hill, T. G., McLinden, C. A., Wunch, D., Jones, D. B. A., and Crisp, D.: Quantifying CO<sub>2</sub> Emissions From Individual Power Plants From Space, *Geophysical Research Letters*, 44, 10,045–10,053, <https://doi.org/10.1002/2017GL074702>, <https://onlinelibrary.wiley.com/doi/pdf/10.1002/2017GL074702>, 2017.
- 660 Nassar, R., Moeini, O., Mastrogiacomo, J.-P., O’Dell, C. W., Nelson, R. R., Kiel, M., Chatterjee, A., Eldering, A., and Crisp, D.: Tracking CO<sub>2</sub> emission reductions from space: A case study at Europe’s largest fossil fuel power plant, *Frontiers in Remote Sensing*, 3, <https://www.frontiersin.org/articles/10.3389/frsen.2022.1028240>, 2022.
- 665 Pillai, D., Buchwitz, M., Gerbig, C., Koch, T., Reuter, M., Bovensmann, H., Marshall, J., and Burrows, J. P.: Tracking city CO<sub>2</sub> emissions from space using a high-resolution inverse modelling approach: a case study for Berlin, Germany, *Atmospheric Chemistry and Physics*, 16, 9591–9610, <https://doi.org/10.5194/acp-16-9591-2016>, publisher: Copernicus GmbH, 2016.
- Reuter, M., Buchwitz, M., Schneising, O., Krautwurst, S., O’Dell, C. W., Richter, A., Bovensmann, H., and Burrows, J. P.: Towards monitoring localized CO<sub>2</sub> emissions from space: co-located regional CO<sub>2</sub> and NO<sub>2</sub> enhancements observed by the OCO-2 and S5P satellites, *Atmos Chem Phys*, 19, 9371–9383, <https://doi.org/10.5194/acp-19-9371-2019>, 2019.
- 670 Tan, M. and Le, Q. V.: EfficientNet: Rethinking Model Scaling for Convolutional Neural Networks, arXiv:1905.11946 [cs, stat], <http://arxiv.org/abs/1905.11946>, 2020.
- Varon, D. J., Jacob, D. J., McKeever, J., Jervis, D., Durak, B. O. A., Xia, Y., and Huang, Y.: Quantifying methane point sources from fine-scale satellite observations of atmospheric methane plumes, *Atmospheric Measurement Techniques*, 11, 5673–5686, <https://doi.org/10.5194/amt-11-5673-2018>, publisher: Copernicus GmbH, 2018.
- 675

- Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D., Burovski, E., Peterson, P., Weckesser, W., Bright, J., van der Walt, S. J., Brett, M., Wilson, J., Millman, K. J., Mayorov, N., Nelson, A. R. J., Jones, E., Kern, R., Larson, E., Carey, C. J., Polat, I., Feng, Y., Moore, E. W., VanderPlas, J., Laxalde, D., Perktold, J., Cimrman, R., Henriksen, I., Quintero, E. A., Harris, C. R., Archibald, A. M., Ribeiro, A. H., Pedregosa, F., van Mulbregt, P., SciPy 1.0 Contributors, Vijaykumar, A., Bardelli, A. P., Rothberg, A., Hilboll, A., Kloeckner, A., Scopatz, A., Lee, A., Rokem, A., Woods, C. N., Fulton, C., Masson, C., Häggström, C., Fitzgerald, C., Nicholson, D. A., Hagen, D. R., Pasechnik, D. V., Olivetti, E., Martin, E., Wieser, E., Silva, F., Lenders, F., Wilhelm, F., Young, G., Price, G. A., Ingold, G.-L., Allen, G. E., Lee, G. R., Audren, H., Probst, I., Dietrich, J. P., Silterra, J., Webber, J. T., Slavič, J., Nothman, J., Buchner, J., Kulick, J., Schönberger, J. L., de Miranda Cardoso, J. V., Reimer, J., Harrington, J., Rodríguez, J. L. C., Nunez-Iglesias, J., Kuczynski, J., Tritz, K., Thoma, M., Newville, M., Kümmerer, M., Bolingbroke, M., Tartre, M., Pak, M., Smith, N. J., Nowaczyk, N., Shebanov, N., Pavlyk, O., Brodtkorb, P. A., Lee, P., McGibbon, R. T., Feldbauer, R., Lewis, S., Tygier, S., Sievert, S., Vigna, S., Peterson, S., More, S., Pudlik, T., Oshima, T., Pingel, T. J., Robitaille, T. P., Spura, T., Jones, T. R., Cera, T., Leslie, T., Zito, T., Krauss, T., Upadhyay, U., Halchenko, Y. O., and Vázquez-Baeza, Y.: SciPy 1.0: fundamental algorithms for scientific computing in Python, *Nature Methods*, 17, 261–272, <https://doi.org/10.1038/s41592-019-0686-2>, 2020.
- 680
- Wang, Y., Broquet, G., Bréon, F.-M., Lespinas, F., Buchwitz, M., Reuter, M., Meijer, Y., Loescher, A., Janssens-Maenhout, G., Zheng, B., and Ciais, P.: PMIF v1.0: assessing the potential of satellite observations to constrain CO<sub>2</sub> emissions from large cities and point sources over the globe using synthetic data, *Geoscientific Model Development*, 13, 5813–5831, <https://doi.org/10.5194/gmd-13-5813-2020>, publisher: Copernicus GmbH, 2020.
- 690
- Wu, D., Lin, J. C., Oda, T., and Kort, E. A.: Space-based quantification of per capita CO<sub>2</sub> emissions from cities, *Environmental Research Letters*, 15, 035004, <https://doi.org/10.1088/1748-9326/ab68eb>, publisher: IOP Publishing, 2020.
- 695
- Zhang, A., Lipton, Z. C., Li, M., and Smola, A. J.: Dive into Deep Learning, <https://doi.org/10.48550/arXiv.2106.11342>, 2022.
- Zheng, B., Chevallier, F., Ciais, P., Broquet, G., Wang, Y., Lian, J., and Zhao, Y.: Observing carbon dioxide emissions over China's cities and industrial areas with the Orbiting Carbon Observatory-2, *Atmospheric Chemistry and Physics*, 20, 8501–8510, <https://doi.org/10.5194/acp-20-8501-2020>, publisher: Copernicus GmbH, 2020.