

The computational and energy cost of simulation and storage for climate science: lessons from CMIP6

Mario C. Acosta¹, Sergi Palomas¹, Stella Paronuzzi¹, Gladys Utrera¹, Joachim Biercamp², Pierre-Antoine Bretonniere¹, Reinhard Budich³, Miguel Castrillo¹, Arnaud Caubel⁴, Francisco Doblas-Reyes¹, Italo Epicoco⁵, Uwe Fladrich⁶, Sylvie Joussaume⁴, Alok Kumar Gupta⁷, Bryan Lawrence⁸, Philippe Le Sager⁹, Grenville Lister⁸, Marie-Pierre Moine¹⁰, Jean-Christophe Rioual¹¹, Sophie Valcke¹⁰, Niki Zadeh¹², and Venkatramani Balaji¹³

¹Barcelona Supercomputing Center, Plaça d'Eusebi Güell, 1-3, 08034 Barcelona, Spain

²German Climate Computing Centre, Bundesstraße 45a, 20146 Hamburg, Germany

³Max Planck Institute, Hofgartenstr. 8, 80539 Munich, Germany

⁴Institut Pierre-Simon Laplace, 11 Bd d'Alembert, 78280 Guyancourt, France

⁵Euro-Mediterranean Center on Climate Change, Via della Libertà, 12, 30121 Venezia VE, Italy

⁶Swedish Meteorological and Hydrological Institute, SE-601 76 Norrköping, Sweden

⁷Norwegian Research Centre, Nygårdsgaten 112, 5008 Bergen, Norway

⁸National Centre for Atmospheric Science, Fairbairn House, 71-75 Clarendon Rd, Woodhouse, Leeds LS2 9PH, United Kingdom

⁹Royal Netherlands Meteorological Institute, Utrechtseweg 297. NL-3731 GA De Bilt, Netherlands

¹⁰European Center for Advanced Research and Training in Scientific Computing, 42 Av. Gaspard Coriolis, 31100 Toulouse, France

¹¹Meteorological Office, Fitzroy road, Exeter, Devon, EX1 3PB, United Kingdom

¹²National Oceanic and Atmospheric Administration, 1401 Constitution Avenue NW, Room 5128, Washington, DC 20230, USA

¹³High Meadows Environmental Institute, Princeton University, Guyot Hall, Room 129, Princeton, NJ 08544-1003, USA

Correspondence: Mario C. Acosta (mario.acosta@bsc.es), Sergi Palomas (sergi.palomas@bsc.es)

Abstract. The Coupled Model Intercomparison Project (CMIP) is one of the biggest international efforts aimed at better understanding past, present, and future of climate changes in a multi-model context. A total of 21 Model Intercomparison Projects (MIPs) were endorsed in its 6th phase (CMIP6), which included 190 different experiments that were used to simulate 40000 years and produced around 40 PB of data in total. This paper presents the main findings obtained from the CPMIP (the Computational Performance Model Intercomparison Project) a collection of a common set of metrics, specifically designed for assessing climate model performance. These metrics were exclusively collected from the production runs of experiments used in CMIP6 and primarily from institutions within the IS-ENES3 consortium. The document presents the full set of CPMIP metrics per institution per experiment, including a detailed analysis and discussion of each of the measurements. During the analysis, we found a positive correlation between the core hours needed, the Complexity of the models, and the Resolution used. Likewise, we show that between 5-15% of the execution cost is spent in the coupling between independent components, and it only gets worse increasing the number of resources. From the data, it is clear that queue times have a great impact on the actual speed achieved, and have a huge variability across different institutions, ranging from none to up to 78% execution

overhead. Furthermore, we evaluated that the estimated carbon footprint of running such big simulations within the IS-ENES3 consortium is 1692 tons of CO₂ equivalent.

15 As a result of the collection, we contribute to the creation of a comprehensive database for future community reference, establishing a benchmark for evaluation and facilitating the multi-model, multi-platform comparisons crucial for understanding climate modelling performance. Given the diverse range of applications, configurations, and hardware utilised, further work is required for the standardisation and formulation of general rules. The paper concludes with recommendations for future exercises aimed at addressing the encountered challenges which will facilitate more collections of a similar nature.

20 *Copyright statement.* TEXT

1 Introduction

Earth System Models (ESMs) are an essential tool for understanding the Earth's climate and the consequences of climate change, which are crucial to the design of response policies to address the current climate emergency resulting from anthropogenic emissions. Modelling the Earth is inherently complex. ESMs are among the most challenging applications that the
25 High-Performance Computing (HPC) industry has had to face, requiring the most powerful computers available, consuming vast amounts of energy in computer power, and producing massive amounts of data in the process (Wang and Yuan 2020; Wang et al. 2010; Fuhrer et al. 2014; McGuffie and Henderson-Sellers 2001; Dennis et al. 2012).

Virtually all models are designed to exploit the parallelism of HPC machines so that we can get the results in a reasonable amount of time while trying to make the best use of the HPC platform. While the technology underneath keeps improving
30 every year (in Petaflops/s, memory bandwidth, I/O speed, etc.) climate software evolves much more slowly. Balaji (2015) and Liu et al. (2013) show how challenging is to adapt multi-scale multi-physics climate models to new hardware or programming paradigms. These models, often community-developed software, are very complex, inherently chaotic, and subject to numerical stability. All of which contribute to a slower evolution of the codes. Bauer et al. (2021) illustrate how climate science did take advantage of Moore's law (Bondyopadhyay, 1998) and Dennard scaling (Frank et al., 2001) without much
35 pressure to fundamentally revise numerical methods and programming paradigms, leading to huge legacy codes mostly driven by scientific concerns. Consequently, such codes achieve notably poor sustained floating-point performance in present-day CPU architectures. Enhancing the performance of these models is crucial to boost the rate at which they can grow (in the resolution, complexity, and features represented). In a context where energy cost is rising, running faster and more cost-effective simulations is key to contributing to the advancement of climate research.

40 The performance of ESMs is hardly limited only by one but by multiple bottlenecks that depend on the model itself and on the properties of the HPC platform on which they run, for instance: models using higher resolutions may benefit from (or be limited by) the speed of the network as the data is split into many nodes, memory-bound models will benefit from having more memory available per core and with faster transmission speed while compute-bound models will perform better in faster CPUs,

models that produce more output will run faster on infrastructures with higher capacities for I/O operations, models that include
45 more individual components will be limited by the load-balance achieved between them and by the coupler performance.

Balaji et al. (2017) proposed a set of 12 performance metrics that define the Computation Performance for Model Intercom-
parison Project (CPMIP) which were specifically designed for climate science by taking into account the structure of ESMs
and how they are executed in real experiments. This set of metrics include the climate experiment and platform properties,
the computational speed and cost (core-hours and energy), measures for the coupling and I/O overhead, and for the memory
50 consumption. Each one is described in detail in Table 1 and Section 3.

In this paper, we present in Section 2 the collection of CPMIP metrics from 33 experiments used for climate projections in
the Coupled Model Intercomparison Project phase 6 (CMIP6). The collection effort has been predominantly led by institutions
affiliated with the IS-ENES3 (Joussaume, 2010), a consortium founded by a Horizon 2020¹ project composed of the most
important weather and climate centres in Europe and devoted to improving the infrastructure to make the Earth System Grid
55 Federation (ESGF) and CMIP publication easier. This compilation is the first of its kind and constitutes a representative part
of the total 124 CMIP6 experiments, involving 45 institutions². Our data encompasses 33 different experiments that were
used to simulate almost 500 000 years during CMIP6 on 14 different HPC machines and involving 14 independent modelling
institutions. All experiments are listed in Table 2, along with the institution in charge, the experiment name, HPC platform,
ocean and atmosphere resolutions, and the main reference to the experiment configuration. In addition, Table 3 shows the
60 complete collection of CPMIP metrics for each one of the models, and Table 4 lists the HPC machines that have been used
to run these models. Furthermore, in Section 3, we include the analysis of the metrics to underscore the most significant
insights derived from this data collection. We study in detail the measurements reported by each institution, grouping them
based on experiment configurations, establishing relationships between intertwined metrics, and discussing the strengths and
difficulties encountered during the analysis of each metric. For instance, our analysis reveals that institutions tend to increase
65 the number of resources used in higher-resolution experiments, thereby mitigating the expected increase in execution time at
the expense of increasing the core-hours required. Similarly, the addition of extra components simulated increases the core-
hours needed, and the cost of coupling interactions and synchronisations between models as well. Institutions reported that
the Coupling cost entails an execution cost overhead typically ranging between 5-15%, and it tends to be more problematic
higher processor counts. Additionally, the numbers indicate that the volume of data generated by an experiment does not
70 correspond to increases in Resolution or core-hours needed, contrary to expectations. We observed very different queue times
for HPC resources across institutions, ranging from instantaneous access to introducing an execution time overhead of up to
78%. Furthermore, we present an initial approximation of the carbon footprint generated from executing these experiments,
totaling 1692 tons of CO₂ equivalent.

Our study emphasises the significance of developing standardised metrics for assessing climate model performance. This
75 contribution will serve to establish a database for future reference and multiple institutions-modellers will be able to use for

¹https://research-and-innovation.ec.europa.eu/funding/funding-opportunities/funding-programmes-and-open-calls/horizon-2020_en, retrieved February 6, 2024.

²<http://esgf-ui.cmcc.it/esgf-dashboard-ui/data-archiveCMIP6.html>, retrieved February 6, 2024.

Table 1. List of CPMIP metrics collected

Metric	Used to evaluate
Resolution (Resol)	number of grid points NXxNYxNZ per component
Complexity (Cmplx)	number of prognostic variables per component
Platform	machine measurements: core count, clock frequency, and double-precision op. per clock cycle
Simulation Years Per Day (SYPD)	number of simulated years per day (24h) of execution time
Core-hours per Simulated Year (CHSY)	cost, measured in core-hours per simulated year
Actual SYPD (ASYPD)	how queue time and interruptions affect the complete experiment duration
Parallelisation (Paral)	total number of cores allocated for the run
Joules Per Simulated Year (JPSY)	energy needed per year of simulation
Memory Bloat (Mem B)	ratio between actual and ideal memory size
Data Output cost (DO)	computing cost for performing I/O
Data Intensity (DI)	amount of data produced after 1 year of simulation divided by the CHSY
Coupling Cost (Cpl C)	computing cost of the coupling algorithm and load imbalance

comparison, which we believe to be essential for evaluating climate modeling performance. The noise and variability present in the dataset are the results of the diversity of the applications represented and the hardware under study. This obscures any attempt to make a general rule formulation. Despite this difficulty, our paper concludes with recommendations for future exercises aimed at addressing these challenges.

80 2 Data collection

The collection process was coordinated and supervised to get the metric results, including meetings, reporting, and surveys conducted at different stages of the CMIP6 simulations (before, during, and after the simulation runs). All the partners listed in Table 2 were invited to participate in the tracking process. The coordination, meetings, and reporting were useful to evaluate the state of the collection from the partners, and we provided support to those institutions that required it during the collection process.

The data collection was divided into two steps: the initial phase comprehends the collection up to March 2020, coinciding with the first IS-ENES3 general assembly where the first results were presented; the second includes the data collected up to the end of 2020 when all the institutions had finished the CMIP6 runs. Finally, IS-ENES3 completed the last update to the Earth System Documentation ³ (ES-DOC) in the middle of 2021, publishing CPMIP along with the other CMIP6 results.

90 As the reader can see, not all institutions managed to provide the full set of CPMIP performance metrics. The metrics frequently missing are the *Coupling Cost*, *Memory Bloat*, and *Data Output cost*. This is primarily attributed to the challenges involved in their collection compared to metrics like *SYPD* or *Parallelisation*, which are well-known within the community and

³<https://es-doc.org/>

Table 2. List of institutions and models that provided the metrics from their CMIP6 executions. Also listed are the HPC platform and resolution used for the atmosphere (ATM) and ocean (OCN) components. Note that "resol" in Table 1 is defined as the number of gridpoints. For better readability, we present here this information using more conventional measure of degrees of latitude and longitude.

Institution	Experiment	HPC machine	Atmosphere resol	Ocean resol	Reference	
BSC	EC-Earth3	MareNostrum4	0.7	1.0	Döscher et al. (2022)	
	EC-EarthVeg		0.7	1.0		
CMCC	CM2-SR5	Zeus	1.0	1.0	Lovato et al. (2022)	
CNRM-CERFACS	CNRM-CM6-1-atm	Beaufix2	1.4	1.0	Voldoire et al. (2019)	
	CNRM-CM6-1		1.4			
	CNRM-CM6-1-HR-atm		0.5	0.25		
	CNRM-CM6-1-HR		0.5			
	CNRM-ESM2-1-atm		1.4	1.0		Séférian et al. (2019)
	CNRM-ESM2-1		1.4			
DKRZ	MPI-ESM1-HR	Mistral	1.0	0.4	Müller et al. (2018)	
GFDL	OM4-p5	Gaea	1.0	0.5	Dunne et al. (2020)	
	ESM4-piC			0.5		
	CM4-piC		1.0	0.25		
	OM4-p25			0.25		
IITM	IITM-ESM	Intel AADITYA	1.875	1.0	Krishnan et al. (2021)	
IMPE	BESM	xc50	1.875	1.0	Veiga et al. (2019)	
IPSL	IPSL-CM6A	Irene-SKL/Curie	2.5	1.0	Boucher et al. (2020)	
KNMI	EC-Earth3	Rhino	0.7	1.0	Döscher et al. (2022)	
	EC-Earth3-AerChem		0.7	1.0		
MPI	MPI-ESM1-LR-ATM	Mistral	4.0	1.5	Müller et al. (2018)	
	MPI-ESM1-LR-LAND		1.875			
	MPI-ESM1-LR		1.875			
NERC	UKESM1-AMIP	Archer xc30	4.0	1.0	Sellar et al. (2020)	
	UKESM1-0-LL		1.875			
	HadGEM3-GC3.1-LL		1.875	1.0		
	HadGEM3-GC3.1-HM		0.8	0.25		Williams et al. (2018),
	HadGEM3-GC3.1-HH		0.8	0.08		
NorESM2	NorESM2-LM	Fram	2.5	1.0	Seland et al. (2020)	
	NorESM2-MM		1.0	1.0		
SMHI	EC-EarthVeg	Tetralith/Beskow	0.7	1.0	Döscher et al. (2022)	
UKMO	UKESM1-0-LL	xce xc40	1.875	1.0	Sellar et al. (2020)	
	HadGEM3-GC3.1-LL		1.875	1.0	Williams et al. (2018)	
	HadGEM3-GC3.1-MM		0.8	0.25		

relatively easier to obtain. Other impediments to collect the CPMIP metrics include time and resources constraints, particularly considering that the focus of the simulations are leans more towards science aspects than to the computational realm during CMIP6 runs. Additionally, some institutions reported that changes in the underlying computational infrastructure have made the collection process more difficult.

Table 3. List of institutions with the model and CPMIP metrics. We also include the Useful Simulated Years (Useful SY), which accounts for the number of years simulated by each experiment that generated data with scientific value

Institution	Experiment	Resol	Cmplx	SYPD	ASYPD	CHSY	Paral	JPSY	Cpl C	Mem B	DO	DI	Useful SY
BSC	EC-Earth3	1.99E+07	34	15.20	9.87	1213	768	4.41E+07	0.080	59.5	1.12	0.041	14020
	EC-EarthVeg	1.99E+07		12.36	7.42	1491	768	4.87E+07	0.100	68.48	1.13	0.059	252
CMCC	CM2-SR5	6.94E+06	397	6.68	6.50	2069	576	1.67E+09	0.074	17.8	1.04	0.050	965
CNRM-CERFACS	CNRM-CM6-1-atm	2.98E+06	128	7.30	6.10	1292	393	3.50E+07					5723
	CNRM-CM6-1	1.02E+07	181	8.10	7.30	1352	400	3.38E+07					22241
	CNRM-CM6-1-HR-atm	2.36E+07	128	2.20	1.80	1541	520	4.80E+07					1190
	CNRM-CM6-1-HR	1.37E+08	181	1.50	1.48	4289	840	1.07E+08					1642
	ESM2-1-atm	2.98E+06	335	7.10	6.40	8520	781	2.28E+08					1759
	ESM2-1	1.10E+07	393	4.70	4.40	21552	1347	5.28E+08					11761
DKRZ	MPI-ESM1-HR	2.00E+07		13.33	11.00	4710	2616	3.21E+08					1864
GFDL	OM4-p5	3.32E+07	13	15.90	12.22	1962	1300	7.50E+07	0.140	33.61		0.039	300
	ESM4-piC	3.76E+07	140	8.65	7.46	13576	4893	5.19E+08	0.270	40.57		0.003	1124
	CM4-piC	1.28E+08	31	9.98	8.16	15388	6399	3.72E+08	0.130	47.64		0.018	657
	OM4-p25	1.26E+08	11	11.50	7.05	9748	4671	5.88E+08	0.260	16.09		0.006	300
IITM	IESM	1.83E+06	168	8.00	7.00	996	332	3.81E+07		36.7			845
IMPE	BESM	6.88E+06	132	3.60	3.40	1853	278					0.020	360
IPSL	IPSL-CM6A	1.06E+07	750	12.00	11.50	1900	950	1.16E+08	0.050	10.00	1.20	0.070	53000
KNMI	EC-Earth3	1.99E+07	34	16.20	16.20	1286	868						1009
	EC-Earth3-AerChem	2.06E+07		3.03	3.03	3549	448						730
MPI	MPI-ESM1-LR-ATM	8.66E+05		45.90	25.20	163	312	1.11E+07					991
	MPI-ESM1-LR-LAND	8.33E+05		282.80	265.40	3	36	1.39E+05					2460
	MPI-ESM1-LR	3.12E+06		55.60	22.70	379	878	2.56E+07					18860
NERC	UKESM1-AMIP	2.35E+06	202	1.64	1.41	7376	504	1.04E+08		52.50	1.31	0.003	45
	UKESM1-0-LL	1.14E+07	252	2.02	1.10	8554	720	3.18E+08	0.078	28.00	1.19	0.005	195
	HadGEM3-GC3.1-LL	1.14E+07	150	4.25	1.06	12198	2160	4.33E+08	0.047	56.80	1.41	0.016	70
	HadGEM3-GC3.1-HM	1.99E+08	54	0.58	0.46	192662	4656	7.70E+09	0.210	154.00		0.001	65
	HadGEM3-GC3.1-HH	1.26E+09	54	0.49	0.34	588931	12024	2.30E+10		183.00	1.41	0.0004	65
NorESM	NorESM2-LM	1.01E+07		13.84	3.03	1665	960	5.60E+07	0.035			0.065	5463
	NorESM2-MM	1.14E+07		8.96	6.14	4886	1824	1.65E+08	0.32			0.060	1021
SMHI	EC-EarthVeg	1.99E+07		12.44	6.65	1667	864					0.028	6337
UKMO	HadGEM3-GC3.1-LL	1.14E+07	228	4.00	3.55	13392	2232	4.97E+08	0.061	46.00	1.03	0.074	5610
	UKESM1-0-LL	1.14E+07	372	4.30	3.60	16074	2880	5.97E+08	0.098	4.60	1.03	0.019	15435
	HadGEM3-GC3.1-MM	1.44E+08	236	1.65	1.32	62836	4320	2.33E+09	0.105	120.00	1.02	0.050	2386

Table 4. List of HPC machines used for the experiments under study, detailing hardware specifications, benchmark results (Linpack and High-Performance Conjugate Gradient, HPCG), theoretical performance (Rpeak), power consumption and Power Usage Effectiveness (PUE) for each system

Institution	Machine	total cores	cores per node	Mem node (GB)	Mem per core (GB)	network	CPU family	CPU freq (GHz)	Rpeak (PFlop/s)	Linpack (PFlop/s)	Power (kW)	HPCG (TFlop/s)	PUE
BSC	MN4	155520	48	96	2.00	Intel Omni-Path	Platinum Skylake	2.10	10.300	6.22	1632	122.24	1.35
CMCC	Zeus	12528	36	96	2.67	InfiniBand	Gold Skylake	3.00	1.202				1.84
CNRM-CERFACS	Beaufix2	73440	40	64	1.60	InfiniBand	E5 Broadwell	2.20	2.590	2.16	830	35.34	
DKRZ/MPI	Mistral	100200	30	68	2.25	InfiniBand	E5 Haswell	2.29	3.960	3.01	1116	44.11	1.19
IITM	AADITYA	38144	16	64	4.00	InfiniBand	E5 Haswell	2.60	0.790	0.72	790		
INPE	xc50	4080	40	192	4.80	Aries Interconnect	Gold Skylake	2.40	0.313				
IPSL	Curie	80640	16	64	4.00	InfiniBand	E5 Sandy Bridge	2.70	1.670	1.36	2132	50.99	1.43
IPSL	Irene	79488	48	192	4.00	InfiniBand	Platinum Skylake	2.70	6.640	4.07	917	52.68	
KNMI	Rhino	4752	28	128	4.57	InfiniBand	Nehalem	3.06	0.058				
NERC	Archer xc30	118080	24	64	2.67	Aries Interconnect	E5 Ivy	2.70	2.550	1.64		80.79	1.10
NorESM	Fram	32256	32	64	2.00	InfiniBand	E5 Broadwell	2.10	1.100	0.95			
SMHI	Beskow	65920	32	64	2.00	Aries Interconnect	E5 Haswell	2.30	2.440	1.80	842		
SMHI	Tetralith	61056	32	96	3.00	Intel Omni-Path	Gold Skylake	2.10	4.340	2.97		65.24	
UKMO	xc40	241920	36	192	5.33	Aries Interconnect	E5 Broadwell	2.10	8.130	7.04			

2.1 Additional data collected

The CPMIP metrics serve not only as a means of computational evaluation but also provide valuable insights for broader analysis. In light of this, we collaborated with the Carbon Footprint Group created within the IS-ENES3 consortium, which was responsible for evaluating the Total Energy Cost associated with the CMIP6 experiments

$$\text{Total Energy Cost} = \text{Useful Simulated Years} \times \text{JPSY} \quad (1)$$

The Total Energy Cost of an experiment is defined as the product of the Useful Simulated Years, defined as years of simulation that produced data with a scientific value that was either shared between the groups or kept within the producer group for scientific analysis, and the Jules per Simulated Year (JPSY). This collaboration enabled us to provide for the first time an estimation of the carbon footprint related to those experiments. The Carbon Footprint was calculated following Equation 2.

$$\text{Carbon Footprint} = \text{Total Energy Cost} \times CF \times PUE \quad (2)$$

where the *Total Energy Cost* is in MWh, *CF* is the greenhouse gas conversion factor from MWh to CO₂ kilogram according to the supplier bill or the country energy mix, and *PUE* (*Power Usage Effectiveness*) accounts for other costs sustained from the data-center, such as cooling. The results for all the institutions that participated in the study during the CPMIP collection are shown during the analysis section in Table 10.

2.2 Uncertainty in the measurements

Understanding measurement uncertainty and machine variability has a significant role in any performance analysis, particularly when comparing models running across different platforms without advanced performance tools or methods like tracing or sampling. Before starting the collection of the metrics, we asked each institution to indicate the machine variability, which was reported to be below 10% for all machines used. This provides an initial rough estimation, subject to future refinement efforts like the usage of benchmarking codes for climate science like the one proposed by van Werkhoven et al. (2023).

It is important to note that not all metrics exhibit the same variability range. Certain metrics, such as Parallelisation, Resolution, platform, and model Complexity, are constant values determined just by the experimental configurations, the HPC infrastructure, and model characteristics. These are considered *static* metrics.

The rest of the metrics are related to the execution speed and therefore subject to different sources of variability. On the one hand metrics like the SYPD or CHSY are well-known by the community and straightforward to collect: this results in less margin of error during collection and any variability should be attributed solely to the machine. On the other hand metrics like the Actual SYPD, JPSY, Coupling Cost, Memory Bloat, Data Intensity, and Data Output Cost are less common to collect and this can lead to confusion and human errors (e.g. whether the Actual SYPD should include system interruptions or only queue time can bring a to systematic misreporting). This represents a second source of variability, difficult to assess and estimate.

Identifying and understanding this uncertainty is key for accurately interpreting and comparing the performance of models across different centers. Special effort has been made to ensure the quality and correctness of the metrics presented in this

work through continuous support of the groups during the collection and double-checking of the reported numbers with the responsible for each institution whenever needed.

130 Future collections like this one will contribute to better identify and address metrics uncertainty, while detailed analysis of individual metrics will enhance our understanding of their characteristics and exhibit variability. For instance, studies like Acosta et al. (2023) focus mainly on the Coupling Cost and offer valuable lessons for understanding and measuring this metric. Therefore, mitigating possible uncertainties arising from misconception or lack of the appropriate tools to collect them in the future.

135 **3 Analysis**

Analysing metrics derived from diverse models, executed on multiple platforms, and managed by independent institutions presents a non-trivial challenge. Moreover, the presence of missing values further complicates the analysis, making it difficult to substitute them with estimations or interpolations, particularly given the relatively limited size of the dataset.

Our approach consisted of first: validating the metrics provided by the institutions. We have sometimes found that the metrics
140 reported for some models were orders of magnitude apart from the rest. In this case, we started actively communicating with the institutions asking them to double-check the values and assisting them in the re-computation process. After going through this process for each one of the metrics and models we came up with the values reported in Section 2: in Table 2 and Table 3 the reader can find the complete list of models for which the CPMIP metrics were collected, with the name of the institution that was in charge for the run, the resolution used for the OCN and ATM, the reference for the experiment configuration,
145 and the CPMIP metrics. Additionally, we include in Table 4 the most relevant information on the HPC platforms used by the institutions and some supplementary metrics in Table 10 related to the execution costs in CO₂ emissions.

Later, and for each of the metrics analysed in detail in the following sections, we filtered by model selecting those where the metric was provided, sorting and/or grouping them by the reported value. Finally, to uncover possible relations among the metrics, we have used both statistical approaches (e.g. Pearson's correlation, Freedman et al., 2007) and qualitative analysis.

150 **3.1 Resolution**

The first attempt to extract valuable information from Table 3 was done by grouping the experiments by resolution, since for the moment we want to compare the performance achieved by ESMs whose target is similar. We are ignoring here the fact that for some simulations the set-up has fewer grid points (e.g. reduced Gaussian in the atmosphere or removal of land points in the ocean) and we are using the total size of the corresponding regular grid. The resolution of a component is measured
155 as the number of grid points it has ($N_X \times N_Y \times N_Z$), and the total resolution is given by the sum of the resolutions of their constituents. There is not a strict consensus on the connection between the number of grid points and the categorisation of low, medium, and high resolutions. Thus, for the grouping we have used both the naming provided by the institution in charge of the experiment and the total number of grid points used for each model configuration. Most configurations have been categorised as low resolution and use up to $2.10E+07$ grid-points in total, or no less than 0.7 degrees latitude-longitude grid spacing for any

160 of the components (see Figure 1 and Table 2). On the other hand, only those experiments with an Ocean/Atmosphere resolution under 0.5 degree are treated as medium-high resolution configurations (see Figure 2).




Figure 1. Atmosphere and ocean grid-points for low-resolution experiments. The yellow color refers to components that are contributing to the atmosphere or the ocean but can not be counted as a general circulation model per se (e.g. land-surface, sea ice, vegetation, etc.).

We see in Figure 1 the low-resolution experiments. The number of grid points for the ATM (red) and OCN (blue) components for each model/institution has been listed in ascending order. We observe that except for EC-Earth, all other models run the OCN at a higher resolution than the ATM. More precisely, the OCN resolution is between 3 to 5 times bigger for MPI-ESM, BESM, CM2-SR2, CNRM-CM6, HadGEM3-LL, UKESM-LL and NorESM-MM. While in EC-Earth, it only accounts for 1/3 of the total model resolution (the remaining 2/3 are used for the ATM). Remarkably, the LM configuration used at NorESM uses a grid for the OCN which is 22 times bigger than the one for the ATM. As one would expect, the total number of grid points of an experiment can be explained solely by the ATM and OCN resolution used, but we will show later how adding more components/features (in yellow in Figure 1) can have quite an impact on the performance anyway.

170 Figure 2 shows the number of ocean and atmosphere grid points for the medium-high resolution experiments. We observe that like most of the low-resolution ones, all experiments use more grid points for the oceanic component than for the atmospheric one (notably, GFDL CM4-piC experiment use 55x more grid points in the OCN component). The ATM resolutions range between 1 and 0.4 degree, while OCN ones mostly run at 1/4 of a degree, except for the NERC-HadGEM3-GC3.1-HH experiment which runs the oceanic component at 1/12.




Figure 2. Atmosphere and ocean grid-points for medium-high resolution experiments

175 3.2 Complexity

The complexity of a coupled model, as defined in Table 1, is the number of prognostic variables among all components. Here, "prognostic" refers to variables that the model directly predicts, such as temperature, atmospheric humidity, salinity, etc. In other words, variables that can be obtained directly as outcomes of the model. This metric is not well-known by the community and never collected before, leading to confusion in some cases. Therefore, the values reported here are to be seen as approximations. Only by continuously measuring these metrics in future collections will we improve our understanding of model complexity and its implications on model performance. The data in Table 5 reveals a wide variability in *Complexity* (*Cmplx*) across the models, with most models reporting a value that ranges between 100 to 400. Notably, GFDL (OM4 and CM4) and EC-Earth have considerably lower *Cmplx*. IPSL-CM6A model stands out in this context with a *Cmplx* of 750, which is markedly higher than the other models, potentially due to its representation of the carbon cycle. Likewise, we were expecting a much higher value for the EC-Earth-Veg experiment, but it was impossible to get this metric for the vegetation component (LPJ-Guess) even after contacting the developers. This highlights the challenge of obtaining this metric with accuracy, partly due to a lack of awareness of the number of prognostic variables of the components among users of the ESMs, leading to an overestimation for this metric, and also because the approximation based on the size of the restart files (Balaji et al., 2017, p. 25) is not always accurate. For instance, LPJ-Guess restart file size can measure tens of GB and depends on the Parallelisation used for this component. What's more, explaining why NERC HadGEM3-GC31 *Cmplx* is almost 3 times larger for the lower resolution configuration (LL) than for the same experiment using more grid points (MM, HM and HH configurations) represents

Table 5. Resolution, SYPD, CHSY, Parallelisation and Coupling Cost for experiments that reported the Complexity metric

Institution	Experiment	Resolution	SYPD	CHSY	Parallelisation	Complexity	Coupling Cost
BSC	EC-Earth3	1.99E+07	15.20	1491	768	34	0.080
CNRM-CERFACS	CNRM-CM6-1-atm	2.98E+06	7.30	1292	393	128	
	CNRM-CM6-1	1.10E+07	8.10	1541	520	181	
	CNRM-CM6-1-HR-atm	2.36E+07	2.20	8520	781	128	
	CNRM-CM6-1-HR	1.37E+08	1.50	21552	1347	181	
	ESM2-1-atm	2.98E+06	7.10	1352	400	335	
	ESM2-1	1.10E+07	4.70	4289	840	393	
GFDL	OM4-p25	1.26E+08	11.50	9748	4671	11	0.130
	OM4-p5	3.32E+07	15.90	1962	1300	13	0.140
	CM4	1.28E+08	9.98	15388	6399	31	0.260
	ESM4	3.76E+07	8.65	13576	4893	140	0.270
IITM	IESM	1.83E+06	8.00	996	332	168	
IMPE	BESM	6.88E+06	3.60	1853	278	132	
IPSL	IPSL-CM6A	1.06E+07	12.00	1900	950	750	0.050
KNMI	EC-Earth3	1.99E+07	16.20	1286	868	34	
NERC	HadGEM3-GC3.1-HM	1.99E+08	0.58	192662	4656	54	0.210
	HadGEM3-GC3.1-HH	1.26E+09	0.49	588931	12024	54	
	HadGEM3-GC3.1-LL	1.14E+07	4.25	12198	2160	150	0.047
	UKESM1-AMIP	2.35E+06	1.64	7376	504	202	
	UKESM1-0-LL	1.14E+07	2.02	8554	720	252	0.078
UMKO	HadGEM3-GC31-LL	1.14E+07	4.00	13392	2232	228	0.061
	HadGEM3-GC31-MM	1.44E+08	1.65	62836	4320	236	0.105
	UKESM1-0-LL	1.14E+07	4.30	16074	2880	372	0.098

a challenge. Similarly, the notable differences between NERC and UMKO measurements, despite both running HadGEM-GC3.1 and UKESM1 models but on different platforms, raise questions about their source, which requires further investigation.

Nonetheless, the data from CNRM-CERFACS provides evidence supporting the idea that the *Cmplx* of a model should remain consistent regardless of the resolution, and only increase as additional features are simulated by the ESM. For instance, the *Cmplx* of CNRM-CM6 ATM standalone runs (CNRM-CM6-1-atm and CNRM-CM6-1-HR-atm) is 128 and grows up to 181 when the OCN component is included for the coupled configuration (CNRM-CM6-1 and CNRM-CM6-1-HR). The same is also observed for the CNRM-ESM2 model, where the *Cmplx* increases from 335 to 393 when adding the OCN component. Furthermore, in both cases, the ESMs require more processing elements when running the coupled version. This shows a clear interconnection between the Parallelisation and *Cmplx* as both will grow when comparing standalone and coupled simulations, other examples are: NERC standalone execution UKESM1-AMIP and UKESM1-LL coupled version, GFDL standalone OM4 (OCN only) runs and the coupled configurations ESM4 and CM4, and CNRM-CM6-atm (ATM only), CNRM-CM6-1 (ATM and OCN) and IPSL-CM6A (ATM, OCN and chemistry).

Therefore, *Cmplx* usually reduces the *SYPD* achieved and/or increases the *CHSY* given that adding a new component will, at best, only increase the latter. Maintaining the same throughput when increasing the *Cmplx* requires the use of more parallel resources, which translates into more costly executions and is usually correlated to parallel efficiency loss due to the need for

Table 6. Experiments that reported the Data Output cost (DO) and Data Intensity (DI) metrics

Institution	Experiment	Resolution	Complexity	SYPD	CHSY	Parallelisation	DO	DI
BSC	EC-Earth3	1.99E+07	34	15.20	1213	768	1.12	0.0410
	EC-EarthVeg	1.99E+07		12.36	1491	768	1.13	0.0590
CMCC	CM2-SR5	6.94E+06	397	6.68	2069	576	1.04	0.0500
GFDL	OM4-p5	3.32E+07	13	15.90	1962	1300		0.0392
	OM4-p25	1.26E+08	11	11.50	9748	4671		0.0178
	ESM4-piC	3.76E+07	140	8.65	13576	4893		0.0032
	CM4-piC	1.28E+08	31	9.98	15388	6399	1.24	0.0058
IMPE	IMPE-BESM	6.88E+06	132	3.60	1853	278		0.0200
IPSL	IPSL-CM6A	1.06E+07	750	12.00	1900	950	1.20	0.0700
NERC	HadGEM3-GC3.1-LL	1.14E+07	150	4.25	12198	2160	1.41	0.0160
	HadGEM3-GC3.1-HM	1.99E+08	54	0.58	192662	4656		0.0006
	HadGEM3-GC3.1-HH	1.26E+09	54	0.49	588931	12024	1.41	0.0004
	UKESM1-AMIP	2.35E+06	202	1.64	7376	504	1.31	0.0030
NorESM	UKESM1-0-LL	1.14E+07	252	2.02	8554	720	1.19	0.0050
	NorESM2-LM	1.01E+07		13.84	1665	960		0.0650
	NorESM2-MM	1.14E+07		8.96	4886	1824		0.0600
SMHI	EC-EarthVeg	1.99E+07		12.44	1667	864		0.0280
UKMO	UKESM1-0-LL	1.14E+07	372	4.30	16074	2880	1.03	0.0190
	HadGEM3-GC31-LL	1.14E+07	228	4.00	13392	2232	1.03	0.0740
	HadGEM3-GC31-MM	1.44E+08	236	1.65	62836	4320	1.02	0.0500

coupling synchronisations and interpolations (e.g. see GFDL results in Table 5). The relation between *Cmplx* and the Coupling Cost is further discussed in subsection 3.5.

3.3 Data output

210 ESMs generate a large amount of output data, including model results, diagnostics, and intermediate variables, which need to be written to storage. Writing and saving this massive amount of data to disk or other storage mediums is time-consuming and can affect the overall performance of the model. Concurrent access to storage resources by multiple processes or multiple model instances can create contention, may represent an I/O bottleneck, and eventually degrade performance and scalability. CPMIP metrics add two metrics to quantify and evaluate the I/O workload: the *Data Output Cost (DO)*, which reflects the cost of performing I/O and is determined as the ratio of *CHSY* with and without I/O; and the *Data Intensity (DI)*, which measures the data production efficiency in terms of data generated per compute hour (i.e. GB/Core-hour).

Data Output Cost

From Table 6, we see that all the experiments conducted by UKMO and CMCC reported a *DO* below 1.05, even though the *DI* varies considerably between the different experiments. Moreover, we observe that the *DO* is much higher for the same ESM (HadGEM-GC31-LL and UKESM1-0-LL) when executed by NERC, reaching 1.19 for UKESM1-0-LL and 1.41 for HadGEM3-GC31-LL. It is not possible to know, however, if this is due to the difference between the HPC platform used or to differences in the model I/O configuration. This underscores the importance of the specific model's I/O configuration in

influencing the *DO* metric. Besides, neither the metrics collected from UKMO nor the ones reported from NERC show that the *DO* should increase when running higher-resolution experiments (HadGEM3-GC31-MM and HH configurations). Moreover, EC-Earth and EC-Earth-Veg *DO* measurements are almost the same, suggesting that adding the vegetation model to EC-Earth does not increase the cost of the I/O, while UKESM runs conducted by NERC show that the *DO* is much higher when running the ATM standalone configuration, UKESM-AMIP, that the coupled run, UKESM-1-LL. Thus, the increase in Complexity or Resolution does not increase the cost of the I/O but the cost of the whole ESM simulation, which can diminish the *DO* metric if I/O workload stays constant.

230 Data Intensity

As seen in Table 6, the *DI* is generally of the order of MB per core-hour and gets smaller as we move to higher-resolution experiments (i.e. higher *CHSY*), meaning that the amount of data generated does not grow proportionally with the number of grid points nor with the execution cost. For instance, the *DI* reported for NERC-HadGEM, UKMO-HadGEM, NorESM2 and GFLD-OM4 experiments decreases when increasing the resolution. Thus, we observe a positive correlation between the *SYPD* and the *DI*.

3.4 Workflow and infrastructure costs

The real execution time of climate experiments can not be explained only by the speed at which a model can run. Queue times before having access to the HPC resources (usually managed by an external scheduler), service disruption, errors in the model/workflow manager, etc. can heavily extend the time-to-solution of ESMs. From the data in Table 1, we see that the difference between the *SYPD* and *ASYPD* reported varies a lot between institutions. Some claim that they had no overhead in their runs (KNMI), while for others it can account for up to 78% (NorESM2-LR). The histogram in Figure 3 helps illustrate the spread of the *ASYPD* overhead: it rarely surpasses 50% and half of the institutions reported it to be less than 20%. Judging from the spread of this metric and from the discussions after the collection, we consider that there are two groups: 1) Institutions that included solely the queue time, which reported an overhead under 20%, and 2) Institutions including not only the queue time but also the system interruptions and/or workflow management, which reported much higher values.

The results support the idea that queuing time represents an increment of around 10-20% to the speed of the ESM. On the other hand, adding interruptions and workflow management the total execution time could increase up to 40-50% compared to the simulation time alone. We do not have enough supporting data to draw any definitive conclusions, so we believe that it would be essential to add finer granularity to the *ASYPD* metric to be able to differentiate both factors. BSC CMIP6 results using the same configuration on two different platforms (Marenostrum and CCA) proved that the percentage of each part (queue time, interruptions or post-processing) could change among platforms even though the CMIP6 experiment is the same⁴. From the metrics listed in Table 3, we see that the difference between *SYPD* and *ASYPD* for the same model can significantly vary depending on the machine used for execution. For EC-Earth3 (standard and vegetation experiments), the overhead ranges from 0% at KNMI to 0.35-0.40% at BSC and up to 0.47% at SMHI. However, it is important to note that the value provided

⁴<https://shorturl.at/lzAHO>, retrieved February 6, 2024




Figure 3. Histogram of the Actual SYPD overhead

255 by KNMI only accounts for the queue time, and they reported having instant access to the HPC resources. Furthermore, for HadGEM3-GC3.1-LL, we observe that NERC and UKMO runs are similar in the model execution speed, achieving approx. 4 SYPD, but totally different in the ASYPD. The overhead due to the workflow at UKMO is just 11%, whereas at NERC it takes 75%. We see something similar when comparing the same institutions for the UKESM-LL execution, where the overhead in UKMO is almost the same as before (16%), but it has drastically decreased at NERC. As we expected, the ASYPD overhead is
 260 related to the model SYPD, but more importantly to the workload of the platform used for the runs. Furthermore, we observed that for UKMO and MPI the smaller the Parallelisation, the smaller the overhead due to the workflow.

3.5 Coupling Cost

Coupling Cost ($Cpl\ C$, Equation 3) is an essential metric evaluated in this study. It quantifies the overhead introduced by coupling within an Earth System Model (ESM). This overhead encompasses various factors, including the coupling algorithms
 265 used for grid interpolations and calculations for conservative coupling. Additionally, it incorporates the impact of the load imbalance, which arises when different independent components of the ESM finish their computations at varying rates, potentially leaving processing elements idle. It is defined as follows:

$$Cpl_Cost \equiv \frac{T_M P_M - \sum_c T_C P_C}{T_M P_M} \quad (3)$$

Where T_M and P_M are the runtime and Parallelisation for the whole coupled model, and T_C and P_C the same for each
 270 individual component it uses.

Figure 4 shows the list of institutions ordered from lower to higher $Cpl\ C$. Most institutions reported that the cost increase due to the coupling accounts for around 5-15% of the total. Only 4 (over the 16 that reported this metric) show an increase of over 20%. The data from GFDL (OM4-p5, OM4-p25, ESM4-piC, and CM4-piC) and UKMO (UKESM-LL and UKESM-AMIP) suggests that the increase in Complexity leads to higher $Cpl\ C$ and lower SYPD. This aligns with the expectations,
 275 as the addition of a new component to the ESM will likely slow down the model and make the load balancing harder. It is

noteworthy that a similar trend is observed in EC-Earth experiments. Even though we don't know the exact value for EC-EarthVeg *Cmplx*, it is known to be higher than in the standard EC-Earth (ATM-OCN) configuration due to the inclusion of vegetation and chemistry models. When comparing the performance of these two runs, we see a decrease in the *SYPD* and a concurrent increase in the *Cmplx* and *Cpl. C*, as discussed in more detail in subsection 3.2.

280 In general, the *Cpl C* tends to rise when running experiments that use a higher Parallelisation. This could reflect a problem in the coupling phase. It can occur that the coupling algorithm is not scaling correctly or that the higher resolution configuration is not well-balanced. It is also likely that since the computing cost of running configurations in lower resolutions is smaller and less time-consuming, institutions can afford to run more spin-up tests and come up with a better distribution of processes among the coupled components to obtain a better load balance. In comparison, the contrary will happen for higher resolutions.

285 Since there are no specific tools to balance a coupled model, these institutions are forced to use a trial-and-error approach, which is not trivial for complex configurations with several components and/or differences in the time-stepping among them.

For these cases, a finer granularity in the *Cpl C* metric and new ways to achieve a well-balanced configuration could be needed, splitting interpolation algorithm and waiting time in different sub-metrics or providing some of the CPMIPs (*SYPD*, *CHSY*...) not only for the coupled version but also per component.




Figure 4. Coupling cost for all the institutions that provided the metric

Table 7. Metrics for models available on different HPC platforms

Experiment	Institution	Resolution	SYPD	CHSY	Parallelisation
EC-Earth3	BSC	1.99E+07	15.20	1213	768
	KNMI	1.99E+07	16.20	1286	868
EC-Earth3Veg	BSC	1.99E+07	12.36	1491	768
	SMHI	1.99E+07	12.44	1667	864
HadGEM3-GC3.1-LL	NERC	1.14E+07	4.25	12198	2160
	UKMO	1.14E+07	4.00	13392	2232
UKESM1-0-LL	NERC	1.14E+07	2.02	8554	720
	UKMO	1.14E+07	4.30	16074	2880

290 **3.6 Speed, cost, and Parallelisation**

The *speed* of execution (*SYPD*) of a model is a fundamental metric that requires careful consideration. However, taken alone, it may not be enough to shed light on the model’s performance itself. The meaning of a model’s speed can only be fully understood when correlated to other important metrics. Among these the *Parallelisation* (*Paral*, i.e. the number of parallel resources allocated) stands out as a factor closely related to the speed and that, at the same time, directly influences the computational *cost* (*CHSY*) of the model execution. In this section, we show a detailed analysis of these three interconnected metrics. Contrary to what one would expect, the *SYPD* achieved by the models in this study is not always related to the resolution used nor to the *Paral* allocated. Although if we analyse how the same model performs on different HPC machines (Table 7), we note that higher values of *Paral* usually correspond to faster but more energy-consuming simulations.




Figure 5. Comparison between CHSY and Parallelisation for both low and medium-high resolution experiments. Experiment configurations are arranged from left to right in ascending number of gridpoints. Note that vertical axis use a logarithmic scale for better visualisation




Figure 6. Parallelisation for low (grey) and medium-high (black) resolution models

As seen in Figure 5, the *Paral* and the *CHSY* are closely correlated in low-resolution models (e.g. CMCC-CM2-SR5, NorESM2-LM, IPSL-CM6A, NERC-HadGEM3-GC3.1-LL, UKMO-HadGEM3-GC3.1-LL, UKMO-UKESM1-0-LL, NorESM2-MM, BSC-EC-Earth3, BSC-EC-EarthVeg, KNMI-EC-Earth3, SMHI-EC-EarthVeg), showing that models do not scale in the current generation of HPC platforms. Otherwise, one would see that the *CHSY* of ESMs with similar *Resol* do not increase when using more processors given that the models run faster (i.e. higher *SYPD*). From the data, it is also clear which models are under-performing. Take for instance KNMI-EC-Earth3-AerChem, despite using a smaller Parallelisation compared to its family counterparts (BSC-EC-Earth3, BSC-EC-EarthVeg, KNMI-EC-Earth3 and SMHI-EC-EarthVeg), exhibits a higher *CHSY*. Similarly, NERC-UKESM1-AMIP and NERC-UKESM1-0-LL employ less Parallelisation compared to UKMO-UKESM1-0-LL, yet the *CHSY* does not decrease proportionally. Also, as illustrated in Figure 6, the level of Parallelisation tends to increase as we move to higher-resolution experiments. Thus, and given that we do not observe a relation between the Resolution and the *SYPD* achieved, we conclude that most institutions try to maintain at high-medium resolution the same *SYPD* achieved when running lower-resolution configurations, at the cost of increasing the *CHSY*. Future collections that include more medium-high resolution experiments will help creating further relationships for these experiments.

In addition, and already discussed in subsection 3.5, the *Cpl C* grows together with the Parallelisation, although there isn't any sign that it limits the speed of the models.

Table 8. Resolution, SYPD, CHSY, Paral and Memory Bloat results for UKESM, EC-Earth and HadGEM3-GC31 experiments

Experiment	Resolution	SYPD	CHSY	Parallelisation	Memory Bloat
BSC-EC-Earth3	1.99E+07	15.2	1213	768	59.50
BSC-EC-EarthVeg	1.99E+07	12.4	1491	768	68.48
NERC-HadGEM3-GC3.1-LL	1.14E+07	4.3	12198	2160	56.80
NERC-HadGEM3-GC3.1-HM	1.99E+08	0.6	192662	4656	154.00
NERC-HadGEM3-GC3.1-HH	1.26E+09	0.5	588931	12024	183.00
UKMO-HadGEM3-GC31-LL	1.14E+07	4.0	13392	2232	46.00
UKMO-HadGEM3-GC31-MM	1.44E+08	1.7	62836	4320	120.00
NERC-UKESM-AMIP	2.35E+06	1.6	7376	504	52.50
NERC-UKESM-LL	1.14E+07	2.0	8554	720	28.00

3.7 Memory Bloat

315 The *Memory Bloat* (*Mem B*, Equation 4) is the only CPMIP metric to evaluates models' memory usage by computing the ratio between the real to the ideal memory size. It is defined as:

$$Memory_Bloat \equiv \frac{M - Parallelisation \cdot X}{M_i} \quad (4)$$

Where M is the actual memory size, X is the binary file size and M_i the ideal memory size. The ideal memory size represents the size of the complete model state, which can be obtained by exploring the restart file size. This ratio is typically falls between 10-100. Large *Mem B* values signal of excessive buffering. As an example (Balaji et al. (2017)), for a rectangular grid with a halo size of 2 in X and Y directions, and a 20x20 domain decomposition, the 2-D array including halos is 576 (24x24) instead of 400 (20x20), resulting in a bloat factor of 1.44. Similarly, a 10x10 decomposition would yield an array area of 196, and a bloat ratio of 1.96.

Table 8 presents the *Mem B* values reported for various models along other CPMIP metrics. We observe how the *Mem B* increases with the resolution (e.g. NERC-HadGEM31), likely due to larger subdomains assigned to each compute unit in higher resolutions if the Parallelisation does not increase proportionally. Additionally, *Mem B* also increases when Complexity grows but the Parallelisation remains constant (e.g. BSC-EC-Earth3 with and without the vegetation model) as it requires keeping more data in memory. It is important to acknowledge the challenges in obtaining accurate memory usage for such applications, and the authors are aware that institutions faced difficulties in providing this data. Therefore, the reliability of the reported values varies between sources, and should be contrasted by future measurements (e.g. CPMIP collection for CMIP7). Precise memory measurements, however, can only achievable with more advanced tools and approaches (memory profilers, MPI environment variables, etc.).

Table 9. Other CMIP6 measurements. The "Useful" metric, whenever used, accounts only for experiments that led to scientific value. The Power Usage Effectiveness (PUE) depends on the HPC machine used (Table 4)

Institution	Useful	Total	Useful Data	Total Data	Useful core	Total core	Total	Total Energy	PUE	Conversion Factor (MWh - kg CO ₂ eq)	Carbon Footprint (tons CO ₂)
	Simulated Years*	Simulated Years	Produced (PB)	Produced (PB)	hours (millions)	hours (millions)	Person/Months	Cost (TeraJoules)			
CMCC	965		0.097		1.99		7	1.61	1.84	408	329
CNRM-CERFACS	47,000		1.350	2.48	160.00	365.00	450	6.18	1.43	40	97
DKRZ	1,276	1,321	0.600		5.52	5.90		0.41	1.19	184	24
EC-Earth	28,105	38,854	0.800	1.41	31.13	46.36	115	1.24	1.35	357	165
IPSL	75,000	165,000	1.800	7.60	150.00	320.00	200	8.72	1.43	50	172
MPI-M	24,175	35,000	1.930		16.31			0.62	1.19	184	37
NCC-NorESM2	23,096		0.600		27.23	80.00	150	1.69			
NERC	640		0.460		55.50			2.17	1.10	0	0
UKMO	37,237		10.400		683.00			26.70	1.35	87	868

*The Useful Simulated Years column values can differ from Table 1 given that some of the experiment runs were not shown in that table

3.8 Carbon footprint

In addition to the CPMIP collection, we have also gathered the general metrics shown in Table 10. These metrics provide both useful (only accounting for simulations that produced data with scientific value) and total (encompassing all simulations, including spin-up and any runs that were finally discarded) numbers for the complete execution of CMIP6 experiments at the different institutions. They can be used to provide an idea about the total and useful number of years simulated, data produced and core hours consumed to finish the European community CMIP6 experiments. Although we did our best to collect the most updated data, we are aware that these numbers could have changed since the data collection was finished. We know that some institutions were doing some minor and final executions and updating databases such as ESGF. However, we consider Table 10 a very good representation of the effort done for the collection during CMIP6. In any case and taking into account the previous reasons, we do not analyse the results themselves and we will use this information to evaluate the Carbon Footprint associated with running models for large-scale projects like CMIP6, which is also a very interesting example for the community. By considering the useful Simulated Years, the HPC machine efficiency, and the KWH to CO₂ conversion rates provided by each energy supplier, we calculated the Carbon Footprint (in tons of CO₂) using Equation 2. As the reader can see, NERC reported a zero Carbon Footprint due to their green tariff power supplier. Among other institutions, CMCC is the one with the highest CF, followed by EC-Earth. Both significantly surpass the emissions of the other institutions: CERFACS, MPI, and UKMO have very small CO₂ emissions per kWh. Regarding machine efficiency, CMCC reported that Zeus is the least power-efficient machine, with a Power Usage Effectiveness (PUE) of 1.84. CERFACS, IPSL, EC-Earth and UKMO reported similar values for their machines, while DKRZ, MPI-M and NERC have reported a PUE under 1.2. We believe that CMCC's Carbon Footprint may be overestimated, considering they simulated fewer than 1000 years yet reported nearly double the CO₂ emissions compared to EC-Earth or IPSL, despite these institutions having simulated longer experiments (in SY). The Total Energy Cost of UKMO seems too big compared to their reported Useful Simulated Years. However, this can be attributed to the cost of maintaining the useful data produced, which amounts to 10.4 PB of disk space. The total Carbon Footprint is 1692 tCO₂, even when

355 accounting for the experiments executed by only 8 out of the 49 institutions that are enlisted in CMIP6⁵. Based on a 2018
study by Acosta et al. Acosta and Bretonnière (2018), the Earth science group at the BSC, comprising around 80 people, had
a CO2 equivalent of commuting (29 tCO2eq/yr), computing infrastructure (397 tCO2eq/yr), building and infrastructure (117
tCO2eq/yr), and travel (255 tCO2eq/yr). The total budget was, therefore, estimated to be near 800 tCO2eq/yr. Consequently,
the carbon footprint from the execution of only this small subset of experiments more than doubles our budget in a single year.
360 This finding is consistent with observations from other groups within the community, such as a similar study conducted by
CERFACS between 2019 and 2021, which reported a total budget of around 700 tCO2eq/yr. Nonetheless, the contributions
that CMIP6 has to climate science are invaluable and beyond the immediate costs associated to running the simulations.

4 Drawbacks and actions recommended

Thanks to the experience learned from the data collection and analysis done, we recognise the importance of highlighting the
365 specific drawbacks we have found during this first collection as well as our recommendations to improve the collection and
analysis for future iterations of multi-model climate research projects, such as CMIP7. The authors will continue working on
this topic in the future not only to provide new approaches to facilitate the collection, but also in fostering the collaboration of
the weather and climate science community to address the computational challenges of Earth modelling. Table 10 shows a list
of the main drawbacks along suggested actions for improvement.

⁵https://wcrp-cmip.github.io/CMIP6_CVs/docs/CMIP6_institution_id.html

Table 10. Drawbacks and Recommended Actions for CMIP6 Metrics

Drawbacks	Recommended Actions
CPMIPs are not enough to compare the performance of different ESMs running on different HPC platforms.	Multi-model comparisons will be better grounded once more data is available. Integrating the CPMIPs in the High-Performance Climate and Weather (HPCW, van Werkhoven et al., 2023) benchmark to evaluate the performance of the different machines used by the community.
Lack of resources and time to collect metrics after CMIP experiments.	Perform metric collection before or during CMIP experiments. Develop portable and automated processes for efficient collection.
Inconsistencies in metric collection hinder inter-model comparisons.	Normalise metric collection methods across institutions before multi-model runs. Develop tools to automatise the collection (e.g. integrated into the workflow manager).
Difficulty in identifying computational bottlenecks due to limited information.	Split sensitive metrics into sub-metrics for finer analysis. For instance, the Coupling Cost should separate interpolation from load-imbalance cost, and the ASYPD should differentiate between queue time and system interruptions.

370 5 Conclusions

One of the limiting factors for climate science is the computational performance that Earth System Models (ESM) can achieve on modern High-Performance Computing (HPC) platforms. This limitation imposes constraints on the number of years that can be simulated, the number of ensembles that can be used, the resolution used by the models, the number of features simulated in one experiment, I/O intensity, data diagnostics calculated during the run, etc. Evaluating the performance of an ESM is a tremendous amount of work that generally requires: profiling the application, using tools to visualise and understand the profiling information, and developing and applying solutions based on the bottlenecks found. This process becomes even more complex when dealing with models used in large-scale multi-model projects like CMIP6, where multiple ESM are executed by different institutions that have access to diverse HPC platforms. To address these challenges, the Computational Performance Model Intercomparison Project (CPMIP) metrics were designed to be: universally available, easy to collect, and representative of the actual performance of ESMs and of the entire life-cycle of modeling (i.e. simulation and workflow costs).

This paper presents, for the first time, the results obtained from the CPMIP collection during the CMIP6 exercise. It provides the list of 14 institutions involved, primarily from the IS-ENES3 consortium, along with the 33 CMIP6 experiment configurations and the CPMIP metrics collected for each experiment. Furthermore, it goes well beyond mere data presentation and offers in-depth analysis for each metric collected to demonstrate the broader utility of the CPMIP collection. For instance, this

385 study investigates the resolution used by each model on the oceanic and atmospheric components, explores the relationship
between execution speed and cost with the other metrics, assesses the impact of running models with higher processor counts,
complexity, or I/O requirements, examines the overhead caused by queuing and workflow management, explores the coupling
cost across different configurations, etc.

Besides the CPMIP metrics analysis, this paper highlights results obtained from collaborations with other groups, such as
390 the Carbon Footprint Group. This collaboration underscores the shared concern of multiple institutions regarding computational
performance in climate science and the joint effort to estimate the carbon footprint of the simulations conducted during the
CMIP6 exercise.

Finally, the paper addresses the main issues and drawbacks encountered during the collection and analysis of the metrics,
including the heterogeneity of the models and HPC machines used, as well as uncertainty in the metric measurements reported.
395 These points should be of particular interest to the partners, aiming to improve and facilitate future collections. The paper
also proposes recommendations to confront these challenges, which can be adopted by the community for the development
of novel tools and more finely-grained metrics that would facilitate upcoming similar works. Moreover, the improvement
and development of benchmarks specially designed for climate science will significantly enhance multi-platform performance
comparisons. Continuous collection of these metrics in future multi-model projects (e.g. CMIP7) will facilitate the development
400 of a shared database for the community and vendors.

Data availability. The original data used during this work can be found here: <http://bit.ly/3Y6XhHM>

Author contributions. M.C. Acosta led the data collection process, ensuring that multiple institutions provided the necessary data and pro-
viding technical assistance throughout. S. Palomas conducted the data analysis and validation, and took the lead in writing the manuscript.
S.P. Ticco contributed to data analysis and played a key role in revising and correcting the manuscript. Other co-authors were responsible for
405 data collection from their respective models and institutions

Competing interests. The contact author declares that Sophie Valcke, a co-author of this article, serves as a member of the editorial board of
the Geoscientific Model Development (GMD) journal. The authors have no other competing interests to declare.

Acknowledgements. The research leading to these results has received funding from the EU H2020 ISENES3, under grant agreement n°
824084 and co-funding from the Spanish National Research Council through OEMES (PID2020-116324RA-I00)

410 References

- Acosta, M. and Bretonnière, P.-A.: Towards Minimising Carbon Footprint of Climate Modelling: Modelling Centre Perspective, C report, 2018.
- Acosta, M. C., Palomas, S., and Tourigny, E.: Balancing EC-Earth3 Improving the Performance of EC-Earth CMIP6 Configurations by Minimizing the Coupling Cost, *Earth and Space Science*, 10, e2023EA002912, <https://doi.org/https://doi.org/10.1029/2023EA002912>,
415 e2023EA002912 2023EA002912, 2023.
- Balaji, V.: Climate Computing: The State of Play, *Computing in Science & Engineering*, 17, 9–13, <https://doi.org/10.1109/MCSE.2015.109>, 2015.
- Balaji, V., Maisonnave, E., Zadeh, N., Lawrence, B. N., Biercamp, J., Fladrich, U., Aloisio, G., Benson, R., Caubel, A., Durachta, J., Foujols, M.-A., Lister, G., Mocavero, S., Underwood, S., and Wright, G.: CPMIP: measurements of real computational performance of Earth
420 system models in CMIP6, *Geoscientific Model Development*, 10, 19–34, <https://doi.org/10.5194/gmd-10-19-2017>, 2017.
- Bauer, P., Dueben, P. D., Hoefler, T., Quintino, T., Schulthess, T. C., and Wedi, N. P.: The digital revolution of Earth-system science, *Nature Computational Science*, 1, 104–113, <https://doi.org/10.1038/s43588-021-00023-0>, 2021.
- Bondyopadhyay, P.: Moore’s law governs the silicon revolution, *Proceedings of the IEEE*, 86, 78–81, <https://doi.org/10.1109/5.658761>, 1998.
- Boucher, O., Servonnat, J., Albright, A. L., Aumont, O., Balkanski, Y., Bastrikov, V., Bekki, S., Bonnet, R., Bony, S., Bopp, L., Bracon-
425 not, P., Brockmann, P., Cadule, P., Caubel, A., Cheruy, F., Codron, F., Cozic, A., Cugnet, D., D’Andrea, F., Davini, P., de Lavergne, C., Denvil, S., Deshayes, J., Devilliers, M., Ducharne, A., Dufresne, J.-L., Dupont, E., Éthé, C., Fairhead, L., Falletti, L., Flavoni, S., Foujols, M.-A., Gardoll, S., Gastineau, G., Ghattas, J., Grandpeix, J.-Y., Guenet, B., Guez, Lionel, E., Guilyardi, E., Guimberteau, M., Hauglustaine, D., Hourdin, F., Idelkadi, A., Joussaume, S., Kageyama, M., Khodri, M., Krinner, G., Lebas, N., Levvasseur, G., Lévy, C., Li, L., Lott, F., Lurton, T., Luysaert, S., Madec, G., Madeleine, J.-B., Maignan, F., Marchand, M., Marti, O., Mellul, L., Meur-
430 desoif, Y., Mignot, J., Musat, I., Otlé, C., Peylin, P., Planton, Y., Polcher, J., Rio, C., Rochetin, N., Rousset, C., Sepulchre, P., Sima, A., Swingedouw, D., Thiéblemont, R., Traore, A. K., Vancoppenolle, M., Vial, J., Vialard, J., Viovy, N., and Vuichard, N.: Presentation and Evaluation of the IPSL-CM6A-LR Climate Model, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS002010, <https://doi.org/https://doi.org/10.1029/2019MS002010>, 2020.
- Dennis, J. M., Vertenstein, M., Worley, P. H., Mirin, A. A., Craig, A. P., Jacob, R., and Mickelson, S.: Computational performance of
435 ultra-high-resolution capability in the Community Earth System Model, *The International Journal of High Performance Computing Applications*, 26, 5–16, 2012.
- Döscher, R., Acosta, M., Alessandri, A., Anthoni, P., Arsouze, T., Bergman, T., Bernardello, R., Boussetta, S., Caron, L.-P., Carver, G., Castrillo, M., Catalano, F., Cvijanovic, I., Davini, P., Dekker, E., Doblás-Reyes, F. J., Docquier, D., Echevarria, P., Fladrich, U., Fuentes-Franco, R., Gröger, M., v. Hardenberg, J., Hieronymus, J., Karami, M. P., Keskinen, J.-P., Koenigk, T., Makkonen, R., Massonnet, F.,
440 Ménégos, M., Miller, P. A., Moreno-Chamarro, E., Nieradzick, L., van Noije, T., Nolan, P., O’Donnell, D., Ollinaho, P., van den Oord, G., Ortega, P., Prims, O. T., Ramos, A., Reerink, T., Rousset, C., Ruprich-Robert, Y., Le Sager, P., Schmith, T., Schrödner, R., Serva, F., Sicardi, V., Sloth Madsen, M., Smith, B., Tian, T., Tourigny, E., Uotila, P., Vancoppenolle, M., Wang, S., Wårlind, D., Willén, U., Wyser, K., Yang, S., Yepes-Arbós, X., and Zhang, Q.: The EC-Earth3 Earth system model for the Coupled Model Intercomparison Project 6, *Geoscientific Model Development*, 15, 2973–3020, <https://doi.org/10.5194/gmd-15-2973-2022>, 2022.
- 445 Dunne, J. P., Horowitz, L. W., Adcroft, A. J., Ginoux, P., Held, I. M., John, J. G., Krasting, J. P., Malyshev, S., Naik, V., Paulot, F., Shevliakova, E., Stock, C. A., Zadeh, N., Balaji, V., Blanton, C., Dunne, K. A., Dupuis, C., Durachta, J., Dussin, R., Gauthier, P. P. G., Griffies, S. M.,

- Guo, H., Hallberg, R. W., Harrison, M., He, J., Hurlin, W., McHugh, C., Menzel, R., Milly, P. C. D., Nikonov, S., Paynter, D. J., Ploshay, J., Radhakrishnan, A., Rand, K., Reichl, B. G., Robinson, T., Schwarzkopf, D. M., Sentman, L. T., Underwood, S., Vahlenkamp, H., Winton, M., Wittenberg, A. T., Wyman, B., Zeng, Y., and Zhao, M.: The GFDL Earth System Model Version 4.1 (GFDL-ESM 4.1): Overall
450 Coupled Model Description and Simulation Characteristics, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS002015, <https://doi.org/https://doi.org/10.1029/2019MS002015>, 2020.
- Frank, D., Dennard, R., Nowak, E., Solomon, P., Taur, Y., and Wong, H.-S. P.: Device scaling limits of Si MOSFETs and their application dependencies, *Proceedings of the IEEE*, 89, 259–288, <https://doi.org/10.1109/5.915374>, 2001.
- Freedman, D., Pisani, R., and Purves, R.: *Statistics (international student edition)*, Pisani, R. Purves, 4th edn. WW Norton & Company, New
455 York, 2007.
- Fuhrer, O., Osuna, C., Lapillonne, X., Gysi, T., Cumming, B., Bianco, M., Arteaga, A., and Schulthess, T. C.: Towards a performance portable, architecture agnostic implementation strategy for weather and climate models, *Supercomputing Frontiers and Innovations*, 1, 45–62, <https://doi.org/10.14529/jsfi140103>, 2014.
- Joussaume, S.: IS-ENES: Infrastructure for the European Network for Earth System Modelling, in: EGU General Assembly Conference
460 Abstracts, EGU General Assembly Conference Abstracts, p. 6039, 2010.
- Krishnan, R., Swapna, P., Choudhury, A. D., Narayansetti, S., Prajeesh, A., Singh, M., Modi, A., Mathew, R., Vellore, R., Jyoti, J., et al.: The IITM earth system model (IITM ESM), arXiv preprint arXiv:2101.03410, 2021.
- Liu, Z., Wang, B., Wang, T., Tian, Y., Xu, C., Wang, Y., Yu, W., Cruz, C. A., Zhou, S., Clune, T., and Klasky, S.: Profiling and Improving I/O
465 Performance of a Large-Scale Climate Scientific Application, in: 2013 22nd International Conference on Computer Communication and Networks (ICCCN), pp. 1–7, <https://doi.org/10.1109/ICCCN.2013.6614174>, 2013.
- Lovato, T., Peano, D., Butenschön, M., Materia, S., Iovino, D., Scoccimarro, E., Fogli, P. G., Cherchi, A., Bellucci, A., Gualdi, S., Masina, S., and Navarra, A.: CMIP6 Simulations With the CMCC Earth System Model (CMCC-ESM2), *Journal of Advances in Modeling Earth Systems*, 14, e2021MS002814, <https://doi.org/https://doi.org/10.1029/2021MS002814>, 2022.
- McGuffie, K. and Henderson-Sellers, A.: Forty years of numerical climate modelling, *International Journal of Climatology*, 21, 1067–1109,
470 <https://doi.org/https://doi.org/10.1002/joc.632>, 2001.
- Müller, W. A., Jungclaus, J. H., Mauritsen, T., Baehr, J., Bittner, M., Budich, R., Bunzel, F., Esch, M., Ghosh, R., Haak, H., Ilyina, T., Kleine, T., Kornbluh, L., Li, H., Modali, K., Notz, D., Pohlmann, H., Roeckner, E., Stemmler, I., Tian, F., and Marotzke, J.: A Higher-resolution Version of the Max Planck Institute Earth System Model (MPI-ESM1.2-HR), *Journal of Advances in Modeling Earth Systems*, 10, 1383–1413, <https://doi.org/https://doi.org/10.1029/2017MS001217>, 2018.
- 475 Seland, Ø., Bentsen, M., Olivié, D., Toniazzo, T., Gjermundsen, A., Graff, L. S., Debernard, J. B., Gupta, A. K., He, Y.-C., Kirkevåg, A., Schwinger, J., Tjiputra, J., Aas, K. S., Bethke, I., Fan, Y., Griesfeller, J., Grini, A., Guo, C., Ilicak, M., Karset, I. H. H., Landgren, O., Liakka, J., Moseid, K. O., Nummelin, A., Spensberger, C., Tang, H., Zhang, Z., Heinze, C., Iversen, T., and Schulz, M.: Overview of the Norwegian Earth System Model (NorESM2) and key climate response of CMIP6 DECK, historical, and scenario simulations, *Geoscientific Model Development*, 13, 6165–6200, <https://doi.org/10.5194/gmd-13-6165-2020>, 2020.
- 480 Sellar, A. A., Walton, J., Jones, C. G., Wood, R., Abraham, N. L., Andrejczuk, M., Andrews, M. B., Andrews, T., Archibald, A. T., de Mora, L., Dyson, H., Elkington, M., Ellis, R., Florek, P., Good, P., Gohar, L., Haddad, S., Hardiman, S. C., Hogan, E., Iwi, A., Jones, C. D., Johnson, B., Kelley, D. I., Kettleborough, J., Knight, J. R., Köhler, M. O., Kuhlbrodt, T., Liddicoat, S., Linova-Pavlova, I., Mizieliński, M. S., Morgenstern, O., Mulcahy, J., Neiningner, E., O’Connor, F. M., Petrie, R., Ridley, J., Rioual, J.-C., Roberts, M., Robertson, E., Rumbold, S., Seddon, J., Shepherd, H., Shim, S., Stephens, A., Teixeira, J. C., Tang, Y., Williams, J., Wiltshire, A., and Griffiths,

- 485 P. T.: Implementation of U.K. Earth System Models for CMIP6, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS001946, <https://doi.org/https://doi.org/10.1029/2019MS001946>, 2020.
- Séférian, R., Nabat, P., Michou, M., Saint-Martin, D., Voldoire, A., Colin, J., Decharme, B., Delire, C., Berthet, S., Chevallier, M., Sénési, S., Franchisteguy, L., Vial, J., Mallet, M., Joetzjer, E., Geoffroy, O., Guérémy, J.-F., Moine, M.-P., Msadek, R., Ribes, A., Rocher, M., Roehrig, R., Salas-y Mélia, D., Sanchez, E., Terray, L., Valcke, S., Waldman, R., Aumont, O., Bopp, L., Deshayes, J., Éthé, C., and
490 Madec, G.: Evaluation of CNRM Earth System Model, CNRM-ESM2-1: Role of Earth System Processes in Present-Day and Future Climate, *Journal of Advances in Modeling Earth Systems*, 11, 4182–4227, <https://doi.org/https://doi.org/10.1029/2019MS001791>, 2019.
- van Werkhoven, B., van den Oord, G., Sclocco, A., Heldens, S., Azizi, V., Raffin, E., Guibert, D., Lucido, L., Moulard, G.-E., Giuliani, G., van Stratum, B., and van Heerwaarden, C.: To make Europe’s Earth system models fit for exascale - Deliverable D3.5, <https://doi.org/10.5281/zenodo.7671032>, ESiWACE2 stands for Centre of Excellence in Simulation of Weather and Climate in Europe
495 Phase 2. ESiWACE2 is funded by the European Union’s Horizon 2020 research and innovation programme (H2020-INFRAEDI-2018-1 call) under grant agreement 823988., 2023.
- Veiga, S. F., Nobre, P., Giarolla, E., Capistrano, V., Baptista Jr., M., Marquez, A. L., Figueroa, S. N., Bonatti, J. P., Kubota, P., and Nobre, C. A.: The Brazilian Earth System Model ocean–atmosphere (BESM-OA) version 2.5: evaluation of its CMIP5 historical simulation, *Geoscientific Model Development*, 12, 1613–1642, <https://doi.org/10.5194/gmd-12-1613-2019>, 2019.
- 500 Voldoire, A., Saint-Martin, D., Sénési, S., Decharme, B., Alias, A., Chevallier, M., Colin, J., Guérémy, J.-F., Michou, M., Moine, M.-P., Nabat, P., Roehrig, R., Salas y Mélia, D., Séférian, R., Valcke, S., Beau, I., Belamari, S., Berthet, S., Cassou, C., Cattiaux, J., Deshayes, J., Douville, H., Éthé, C., Franchistéguy, L., Geoffroy, O., Lévy, C., Madec, G., Meurdesoif, Y., Msadek, R., Ribes, A., Sanchez-Gomez, E., Terray, L., and Waldman, R.: Evaluation of CMIP6 DECK Experiments With CNRM-CM6-1, *Journal of Advances in Modeling Earth Systems*, 11, 2177–2213, <https://doi.org/https://doi.org/10.1029/2019MS001683>, 2019.
- 505 Wang, D. and Yuan, F.: High-Performance Computing for Earth System Modeling, pp. 175–184, Springer International Publishing, Cham, https://doi.org/10.1007/978-3-030-47998-5_10, 2020.
- Wang, D., Post, W., and Wilson, B.: Climate change modeling: Computational opportunities and challenges, *Computing in Science & Engineering*, 13, 36–42, 2010.
- Williams, K. D., Copsey, D., Blockley, E. W., Bodas-Salcedo, A., Calvert, D., Comer, R., Davis, P., Graham, T., Hewitt, H. T., Hill, R., Hyder, P., Ineson, S., Johns, T. C., Keen, A. B., Lee, R. W., Megann, A., Milton, S. F., Rae, J. G. L., Roberts, M. J., Scaife, A. A., Schiemann, R., Storkey, D., Thorpe, L., Watterson, I. G., Walters, D. N., West, A., Wood, R. A., Woollings, T., and Xavier, P. K.: The Met Office Global Coupled Model 3.0 and 3.1 (GC3.0 and GC3.1) Configurations, *Journal of Advances in Modeling Earth Systems*, 10, 357–380, <https://doi.org/https://doi.org/10.1002/2017MS001115>, 2018.