

# Ocean biogeochemistry in the coupled ocean-sea ice-biogeochemistry model FESOM2.1-REcoM3

Özgür Gürses<sup>1</sup>, Laurent Oziel<sup>1</sup>, Onur Karakuş<sup>1</sup>, Dmitry Sidorenko<sup>1</sup>, Christoph Völker<sup>1</sup>, Ying Ye<sup>1</sup>, Moritz Zeising<sup>1</sup>, Martin Butzin<sup>1,2</sup>, and Judith Hauck<sup>1</sup>

<sup>1</sup>Alfred-Wegener-Institut, Helmholtz-Zentrum für Polar- und Meeresforschung

<sup>2</sup>MARUM-Center for Marine Environmental Sciences

**Correspondence:** Özgür Gürses (ozgur.gurses@awi.de)

## Abstract.

The cycling of carbon in the oceans is affected by feedbacks driven by changes in climate and atmospheric CO<sub>2</sub>. Understanding these feedbacks is therefore an important prerequisite for projecting future climate. Marine biogeochemical models are a useful tool, but as any model, are a simplification and need to be continually improved. In this study, we coupled the Finite-volumE Sea ice-Ocean Model (FESOM2.1) to the Regulated Ecosystem Model version 3 (REcoM3). FESOM2.1 is an update of the Finite Element Sea ice-Ocean Model (FESOM1.4) and operates on variable mesh resolution. Unlike standard structured-mesh ocean models, the mesh flexibility allows for a realistic representation of small-scale dynamics in key regions at affordable computational cost. Compared to the previous coupled model version FESOM1.4-REcoM2, the model FESOM2.1-REcoM3 utilizes a new dynamical core based on a finite volume discretization instead of finite elements, but retains central parts of the biogeochemistry model. As a new feature, carbonate chemistry including water vapor correction is computed by mocsy-2.0. Moreover, REcoM3 has an extended food web that includes macrozooplankton and fast-sinking detritus. Dissolved oxygen is also added as a new tracer. In this study we assess the ocean and biogeochemical state simulated with FESOM2.1-REcoM3 in a global setup at relatively low spatial resolution forced with JRA55-do atmospheric reanalysis. The focus is on the recent period 1958-2021 to assess how well the model can be used for present-day and future climate change scenarios on decadal to centennial timescales. A bias in global ocean-atmosphere preindustrial CO<sub>2</sub> flux present in the previous model version FESOM1.4-REcoM2 could be significantly reduced. In addition, the computational efficiency is 2–3 times higher than that of FESOM1.4-REcoM. Overall, it is found that FESOM2.1-REcoM3 is a skillful tool for ocean biogeochemical modelling applications.

## 1 Introduction

There is an unequivocal consensus and concern about the effects of increasing greenhouse gases in the atmosphere due to human activities. Since the beginning of the industrial era (year 1750) the concentration of carbon dioxide (CO<sub>2</sub>) in the air has substantially risen from 277 ppm to 417.2 ppm (year 2022, Friedlingstein et al., 2022b). The ocean has taken up a remarkably constant fraction of 25-30% of human CO<sub>2</sub> emissions from fossil fuel burning and land-use change throughout time (Crisp

et al., 2022). For the recent decade 2012-2021, the rate of ocean anthropogenic carbon uptake (including effects of climate  
25 change) amounted to  $2.9 \pm 0.4 \text{ PgC yr}^{-1}$  (26% of total  $\text{CO}_2$  emissions) (Friedlingstein et al., 2022b). A similar proportion  
was taken up by the terrestrial biosphere  $3.1 \pm 0.6 \text{ PgC yr}^{-1}$  (2012-2021), but the total air-to-land  $\text{CO}_2$  flux is substantially  
lower because of emissions from land-use change, mainly deforestation, that amounted to  $1.2 \pm 0.7 \text{ PgC yr}^{-1}$  (2012-2021). The  
ocean carbon sink has grown over the past decades in response to the near-exponential rise in  $\text{CO}_2$  emissions (Friedlingstein  
et al., 2022b). While the global ocean carbon sink estimate is assigned an uncertainty of  $0.4 \text{ PgC yr}^{-1}$  and medium confidence,  
30 regional patterns of the sink differ more strongly. This points to the balance between physical and biological processes, which  
are more difficult to model as also illustrated in model deficiencies in accurately representing the seasonal cycle of  $\text{pCO}_2$   
and  $\text{CO}_2$  fluxes (Mongwe et al., 2018). Both climate change and rising atmospheric  $\text{CO}_2$  feed back on the fraction of  $\text{CO}_2$   
emissions that will end up in the ocean over the next century (Friedlingstein et al., 2003; Canadell et al., 2021). Models are  
important tools in estimating how large these feedbacks are.

35 The flux of  $\text{CO}_2$  between atmosphere and ocean is controlled by two main mechanisms: the solubility pump and the bio-  
logical pump. The solubility pump describes the air-sea  $\text{CO}_2$  exchange that occurs to satisfy a thermodynamic equilibrium,  
and the subsequent transport of carbon from the surface to the deep ocean with the overturning circulation. This leads to  $\text{CO}_2$   
uptake at mid- to high latitudes through high solubility in cold waters and large vertical motion in deep water formation re-  
gions. In contrast, warm ocean regions in the tropics and subtropics as well as upwelling regions lose carbon to the atmosphere  
40 (Takahashi et al., 2009; Wanninkhof et al., 2013). The solubility pump is responsible for anthropogenic carbon uptake. The  
biological carbon pump comprises the fixation of  $\text{CO}_2$  into biomass by phytoplankton and the subsequent downward transfer  
of dead organic material (Boyd et al., 2019). The biological carbon pump is responsible for 75% of the natural vertical carbon  
gradient and for the large-interbasin gradient between the deep Pacific and Atlantic (Sarmiento and Gruber, 2006). Without the  
biological carbon pump, atmospheric  $\text{CO}_2$  would be higher by 200 ppm (Maier-Reimer et al., 1996), and perturbations thereof  
45 can have large effects on atmospheric  $\text{CO}_2$  (Kwon et al., 2009; Lauderdale and Cael, 2021) as also known from paleo evidence  
(Galbraith and Skinner, 2020).

Global ocean biogeochemistry models (GOBMs, Fennel et al., 2022) are used to assess the global ocean carbon sink (Hauck  
et al., 2020), its regional patterns (Fay and McKinley, 2021), and effects of climate change and variability on the ocean carbon  
sink (Le Quéré et al., 2010; Hauck et al., 2013; DeVries et al., 2019; Bunsen, 2022). By their representation of pH, the marine  
50 oxygen cycle and phytoplankton primary production as the base of the marine food web, they also offer information about the  
environmental conditions for marine life and how these will develop under climate change (Bopp et al., 2013; Laufkötter et al.,  
2015; Kwiatkowski et al., 2020). However, modeling the marine biogeochemistry of the ocean is subject to several sources  
of uncertainties. First, GOBMs are expensive with respect to the computational cost due to the advection of a large number  
of tracers and therefore, often demand low spatial resolution. This leads to deficiencies in the representation of significant  
55 physical processes such as (sub)mesoscale currents (McWilliams, 2016) which can have large impacts on transport and mixing  
processes that strongly affect biological productivity (Lévy et al., 2018; Keerthi et al., 2022). Second, the descriptions of  
ecological interactions and of the physiology of primary and secondary producers in GOBMs are still mostly based on empirical  
or semi-empirical mathematical descriptions, such as, the dependency of zooplankton grazing rates on prey abundance (Doney

et al., 2001; Rohr et al., 2022). These contain a large number of parameters that are only partly constrained from observations, making it necessary to tune these parameters in GOBMs to some extent. Choices in these parameters can have strong effects on the biological carbon pump (e.g. Lauderdale and Cael, 2021). It has been demonstrated that the largest source of uncertainty for projections of net primary production (NPP, Tagliabue et al., 2021) comes from model uncertainty, not scenario uncertainty (Frölicher et al., 2016).

Ocean circulation models formulated on unstructured meshes have become an alternative to existing structured global ocean models (Danilov, 2013). The Finite-Element Sea ice–Ocean Model (from now on FESOM1.4, Wang et al., 2014) is one of the first global models with multiple resolutions designed to simulate the large-scale ocean circulation. While it has already been used in numerous applications (Sidorenko et al., 2015; Wekerle et al., 2017), another dynamical core, the Finite-volume Sea ice–Ocean Model version 2.1 (FESOM2.1), has been developed (Danilov et al., 2017). The advantages of a finite volume formulation are (a) better throughput and scalability as a result of a more efficient data structure (Koldunov et al., 2019), (b) the availability of clearly defined fluxes, and (c) the possibility to choose from a selection of transport algorithms, which was very limited before (Danilov et al., 2017). Furthermore, the arbitrary Lagrangian Eulerian (ALE) vertical coordinate is introduced which provides different types of vertical coordinates (Scholz et al., 2019). The Regulated Ecosystem Model (REcoM) is an ocean biogeochemistry model that describes the lower trophic levels of the marine ecosystem using the plankton functional type approach, and that bases its description of primary production on a physiological model for phytoplankton growth that takes into account nutrient availability effects on photoacclimation (Geider et al., 1998) and, for diatoms, on the relative frustule weight (Hohn, 2009). One specificity of REcoM is the representation of flexible stoichiometry, which leads to a description of elemental fluxes that can deviate from the fixed Redfield ratios often used in models (Redfield et al., 1963).

Here, we document the ocean biogeochemistry in the Regulated Ecosystem Model version 3 (REcoM3) coupled to the ocean and sea ice model FESOM2.1, and assess its performance in reproducing carbon and nutrient biogeochemical fluxes as well as the distribution of phytoplankton and zooplankton. Our aim is to analyze the new setup regarding the coupled model state under historical atmospheric CO<sub>2</sub> forcing and the associated model bias and drift from the experiment with constant preindustrial (PI) CO<sub>2</sub> level. We thus focus on evaluating model aspects with regard to the effects of climate change and CO<sub>2</sub> increase on carbon fluxes on century-scale time-scales. We exclude in our analysis the deep-sea distribution of carbon and nutrients. This would require model runs over at least 500 to two thousand years (Séférian et al., 2020), which will be done in follow-up work.

## 85 **2 Methods**


### **2.1 Model Description**

We present the coupled ocean-sea ice-biogeochemistry model FESOM2.1-REcoM3. The previous model version FESOM1.4-REcoM2 has been described by Schourup-Kristensen et al. (2014). Unlike its predecessor FESOM1.4, which uses a finite element formulation, the ocean model is now based on a finite volume discretization, which makes tracer conservation much easier to achieve. FESOM2.1 was described by Danilov et al. (2017) and evaluated in Scholz et al. (2019, 2021). The ocean biogeochemistry is simulated by the Regulated Ecosystem Model version 3 (REcoM3), which builds upon the previous version

REcoM2 (Hauck et al., 2013; Schourup-Kristensen et al., 2014). The advection and diffusion of 28 passive biogeochemical tracers is handled by FESOM2.1, whereas REcoM3 calculates sources and sinks, driven by biological interactions or biogeochemical exchange processes.

### 95 2.1.1 Ocean Model FESOM2.1

FESOM2.1 solves the hydrostatic primitive equations under the Boussinesq approximation (Danilov et al., 2017). This equation set in differential form is discretized on a finite set of points (nodes). As a first step of mesh generation, a 2-dimensional grid is created by combining these nodes in triangular shapes (elements). At this stage, mesh resolution (i.e., the size of triangles) can be adjusted in areas of interest without requiring a nesting approach. A 3-dimensional mesh is produced by projecting triangles in vertical direction forming prisms. The scalar quantities (tracers, pressure) are located at nodes while the horizontal velocities are defined at centroids of elements (See Figs. 1 and 2 in Danilov et al., 2017). A pair of control volumes are defined: The vector control volumes are the prisms based on elements. The scalar control volumes are formed by connecting cell centroids with edge midpoints (Fig. 1). Integration is carried out on a staggered Arakawa B type of mesh (Scholz et al., 2019).



**Figure 1.** Scheme of the cell-vertex discretization in 3-dimensional space. Blue dots correspond to scalar quantities including REcoM3 state variables, located at the mid-layer vertices of triangles. Red dots represent horizontal velocities located at mid-layer cell centers of the triangles. Yellow dots depict vertical transfer velocities, placed at the layer boundaries aligned with scalar quantities in vertical.

We use FESOM2.1, an updated version of FESOM2.0. The updated model features include several developments, such as parallel and asynchronous output writing. An important new feature that we applied is the kinematic backscatter parameterisation. This method takes into account the scales at which energy is scattered back to the resolved flow by introducing a negative viscosity term (Juricke et al., 2020). This greatly improves the simulation of eddy effects in coarse resolution mesh setups (Juricke et al., 2020). The model code also includes representation of ice-shelf cavities (Timmermann et al., 2012), which has been used in regional studies with FESOM1.4-REcoM2 (Nissen et al., 2022). Ice-shelf cavities are, however, not used in this study. Isonutral tracer diffusion (Redi, 1982) and the Gent-McWilliams (GM, Gent and McWilliams, 1990; Griffies, 1998) eddy stirring parameterization are applied. Both GM and Redi are scaled with horizontal resolution with a maximum value of

2000 m<sup>2</sup> s<sup>-1</sup> at 100 km horizontal resolution, and decrease linearly below a resolution of 40 km to reach zero at 30 km resolution effectively switching the parameterization off. As vertical mixing parameterization, the K-profile scheme is used (KPP, Large et al., 1994) with a background vertical diffusivity of 1 x 10<sup>-4</sup> m<sup>2</sup> s<sup>-1</sup> for momentum and 1 x 10<sup>-5</sup> m<sup>2</sup> s<sup>-1</sup> for tracers. 115 Furthermore, the Monin-Obukhov length dependent vertical mixing parameterization is applied in the surface boundary layer south of 50°S (Timmermann and Beckmann, 2004).


Regarding the vertical discretization, FESOM2.1 is formulated with an arbitrary Lagrangian-Eulerian (ALE) scheme, a synthesis of different types of vertical coordinates. In the model configuration used here, we apply a full free-surface formulation and thus permit the vertical movement of the surface and of all other layers (referred to as *zstar*, Scholz et al. (2019)). This drastically improves tracer conservation properties (Campin et al., 2004). Partially filled cells are used at the ocean floor resulting 120 in a smoother representation of the bathymetry.

The sea ice component (Finite-Element Sea Ice Model, FESIM version 2) solves for sea ice concentration, ice and snow thickness, as well as ice drift velocity (Danilov et al., 2015). It is discretized on the same unstructured horizontal mesh as the ocean model. The elastic-viscous-plastic solver and flux corrected transport scheme are used for sea ice advection (Danilov 125 et al., 2015). The formulation of sea ice thermodynamics follows the work of Timmermann et al. (2009).

### 2.1.2 Biogeochemistry Model REcoM3

REcoM3 is a water column biogeochemistry and ecosystem model which incorporates cycles of carbon and nutrients (nitrogen, iron, and silicon) with varying intracellular stoichiometry in phytoplankton, zooplankton and detritus (see Appendix for detailed description and equations). Starting from the work by Schartau et al. (2007), REcoM was first used to describe 130 carbon overconsumption in mesocosm experiments. After coupled to the ocean and sea ice model MITgcm (Marshall et al., 1997), the previous version (REcoM2) with two phytoplankton classes, one zooplankton and one detritus class was applied to study the cycling of marine carbon on present (Hauck et al., 2013, 2018) and glacial time-scales (Du et al., 2022; Völker and Köhler, 2013), as well as the marine iron cycle (e.g., Völker and Tagliabue, 2015; Tagliabue et al., 2016; Ye and Völker, 2017; Pagnone et al., 2019). Moreover, REcoM2 was employed in assessments on the efficiency of ocean alkalinity enhance- 135 ment (Köhler et al., 2013; Hauck et al., 2016), in data assimilation studies (Pradhan et al., 2019) and as a test bed for model development, e.g., for development of a parameterization of iron-ligand binding based on pH (Ye et al., 2020) among others. Simultaneously, REcoM2 was coupled to FESOM1.4 (Schourup-Kristensen et al., 2014). These coupled model set-ups were used either in a global configuration (e.g., Schourup-Kristensen et al., 2014; Hauck et al., 2020) with a regional focus on the Arctic or the Antarctic (Hauck et al., 2015; Schourup-Kristensen et al., 2018; Oziel et al., 2022; Nissen et al., 2022) or in 140 regional configurations (Taylor et al., 2013; Losch et al., 2014). Recently, the model has matured to include two groups of each classes of phytoplankton, zooplankton and detritus (REcoM3, Fig. 2).

Marine primary production is computed through representation of two phytoplankton functional types (PFTs), namely diatoms and small phytoplankton. The diverse group of small phytoplankton comprises a wide range of taxa, including, for instance, non-silicifying, calcifying, and non-calcifying haptophytes and green algae. The model allows PFTs to adapt their 145 internal stoichiometry (C:N:Chl:CaCO<sub>3</sub> ratios for small phytoplankton and C:N:Chl:Si for diatoms) to nutrient levels, ambi-



**Figure 2.** Schematic diagram of the components and interactions in the REcoM3 model. Small phytoplankton (phy) and diatoms (dia) take up inorganic nutrients (nut). Small zooplankton (small zoo.) and macrozooplankton (macrozoo.) consume phytoplankton and particles. Macrozooplankton feeds on small zooplankton. Phytoplankton aggregation, zooplankton sloppy feeding, mortality and fecal pellets generate sinking detritus (slow and fast). Sinking detritus degrades to dissolved organic carbon and nitrogen. Dissolved organic material (DOM) then remineralizes to dissolved inorganic carbon and nitrogen. The number of tracers and the other processes are shown in the appendix (Fig. A2).

ent light and temperature, based on the photoacclimation model by Geider et al. (1998). Si uptake by diatoms is regulated as well, based on the internal Si:N quota, following Hohn (2009). This parameterization takes into account the strong decoupling between Si and N metabolism (e.g., Claquin et al., 2002), and prescribes the observed change in Si:N ratios under Fe and N limitation. The intracellular iron pool is derived from intracellular nitrogen with a fixed Fe:N ratio, based on the fact that  
 150 intracellular iron is mostly associated with the photosynthetic electron transport chain and nitrogen metabolism (Geider and La Roche, 1994; Raven, 1988). REcoM3 also includes the photo-damage parameterization by Álvarez et al. (2018). Calcium carbonate production is assumed to be linearly dependent on the gross small phytoplankton production. CaCO<sub>3</sub> dissolution is described by a depth-dependent dissolution rate.

Zooplankton is represented by two groups, small zooplankton and polar macrozooplankton (Karakuş et al., 2021) and each  
 155 group has a carbon and a nitrogen tracer. The small zooplankton group in the model is associated with relatively higher grazing rates compared to macrozooplankton and is widely spread in the global ocean. The polar macrozooplankton is mainly present in

the Southern Ocean and northern high latitudes. The respiration rate is described mechanistically for macrozooplankton taking into account reduced metabolism in winter and increased metabolism at high grazing rates (Karakuş et al., 2021). For small zooplankton, respiration is calculated with a fixed respiration rate constant and biomass in contrast to the previous version REcoM2 where respiration was used to drive zooplankton C:N back towards Redfield ratio (Hauck et al., 2013; Schourup-Kristensen et al., 2014). Grazing is computed by applying a sigmoidal function with variable preferences on both phytoplankton and detritus (Fasham et al., 1990).

Particulate organic matter (detritus) is split into two groups. The sinking speed of the first detritus group increases linearly with depth (from 20 m day<sup>-1</sup> from the surface to 192 m day<sup>-1</sup> at 6000 m depth; Kriest and Oschlies, 2008). The sinking speed of the second group (fast-sinking detritus) is constant throughout the water column (200 m day<sup>-1</sup>, Karakuş et al., 2021). Remineralisation of carbon and nitrogen occurs in two steps. Detrital material is first degraded to dissolved organic matter and then remineralised to the inorganic forms (dissolved inorganic carbon and nitrogen). For iron, it is assumed that the organic form is directly bioavailable, so it enters the dissolved iron pool in one step.

REcoM3 comprises a single-layer sediment pool for nitrogen, silicon, dissolved inorganic carbon and calcium carbonate. The sinking detritus and associated minerals are accumulated in this layer when they reach the ocean floor. This material is subsequently returned back to the water column to the pools of dissolved inorganic nitrogen, carbon and silicon, as well as alkalinity with a fixed remineralisation rate. The release of iron to the bottom layer of the ocean is assumed to be proportional to the release of inorganic nitrogen (Elrod et al., 2004).

### 2.1.3 Updates to previous REcoM version coupled to FESOM1.4

There are numerous improvements relative to the previously documented version FESOM1.4-REcoM2 (Schourup-Kristensen et al., 2014), and the main changes are listed below:

#### REcoM

1. The routines for calculating carbonate chemistry and air-sea CO<sub>2</sub>-exchange used in FESOM1.4-REcoM2 which followed the guidelines provided by the Ocean Carbon Model Intercomparison Project (Orr, 1999) were replaced by the mocsy-2.0 scheme of Orr and Epitalon (2015). While both use the same thermodynamic equilibrium to calculate surface pCO<sub>2</sub> and CO<sub>2</sub> flux, mocsy-2.0 uses the faster and more accurate algorithm SolveSAPHE (Munhoven, 2013). Among other differences, it follows best practice guides and uses recommended equilibrium constants. The gas exchange formulation is updated to Wanninkhof (2014), which is largely equivalent to Ho et al. (2006). The computed fluxes are scaled with the ice-free area.
2. Dissolved oxygen was added as a new tracer in REcoM3. The air-sea O<sub>2</sub> flux is calculated using the mocsy-2.0 routines (Orr and Epitalon, 2015). Photosynthesis, respiration and remineralisation change oxygen with a fixed O<sub>2</sub>:C ratio, and remineralisation does not depend on O<sub>2</sub> levels in the current model version.

- 190 3. A second zooplankton group and a fast-sinking detritus class were added. The second zooplankton group represents a slow-growing polar macrozooplankton with a feeding preference on diatoms which produces fast-sinking and carbon-rich fecal pellets (Karakuş et al., 2021).
4. Intracellular iron concentration is connected to intracellular nitrogen via a constant ratio Fe:N leading to some variation in the Fe:C ratio, as briefly presented in Tagliabue et al. (2016) and Pagnone et al. (2019).
5. Sedimentary release of iron was added to the model (Tagliabue et al., 2016), in addition to the previously considered Fe input with dust deposition.

## 195 FESOM

Biogeochemical fluxes returned back to the ocean from the benthos are treated with a specific bottom boundary condition. Variable bottom topography leads to a smaller scalar control volume located at the lowermost level. This is because scalar control volumes are obtained by connecting the areas from the elements they are attached a constant level (see Fig. 1 in Danilov et al., 2017). Therefore, the number of elements around a single surface node may vary with depths when it meets  
200 non-flat topography. We thus computed the control volume and associated fluxes for each node by considering all surrounding elements at different depth levels.

### Forcing

Our simulation was forced by the atmospheric reanalysis JRA55-do data set (Tsujino et al., 2018) instead of the CORE-II data set (Large and Yeager, 2009) that was used in previous assessments (Schourup-Kristensen et al., 2014). JRA55-do is a blend of  
205 reanalysis data and satellite observations and has the advantage to provide regularly updated near real time data up to present day with higher temporal (3-hourly) resolution. The freshwater supplied by rivers is a climatology and provided by Large and Yeager (2004) as part of the CORE forcing. Nutrient, carbon and alkalinity supply via river discharge is not included in the experiments described here.

## 2.2 Experimental setup and data

210 In this study, we used a mesh with a nominal resolution of 1 degree as a background. The horizontal resolution is enhanced on the equatorial belt and in the region north of 50°N to match 1/3 degree and 25 km, respectively. The mesh has 48 unevenly spaced vertical layers where the layer thickness ranges from 5 m in the surface to 250 m in the deep ocean (Scholz et al., 2019).

Initial fields for temperature and salinity were taken from the winter statistical fields of Polar Science Center Hydrographic Climatology (PHC3, updated from Steele et al., 2001) that ingests observations from the period 1900-1994. Total alkalinity  
215 (Alk) and preindustrial dissolved inorganic carbon (DIC) were initialized from version 2 of the Global Ocean Data Analysis Project (GLODAPv2) climatology centered to the year 2002 (Lauvset et al., 2016) based on data collected between 1972 and 2013. Dissolved inorganic nitrogen (DIN) and dissolved silicic acid (DSi) were started with values from the Levitus World Ocean Atlas climatology of 2013 (Garcia et al., 2014) occupied between 1955 and 2012. We used the Levitus World Ocean Atlas climatology of 2018 for dissolved oxygen (Garcia et al., 2019a) (See Table 2) based on data for the time span 1955-2017.



220 Due to scarcity of observations, the iron field (DFe) was initialized with output from the Pelagic Interaction Scheme for Carbon and Ecosystem Studies (PISCES) model (Aumont et al., 2003) which was corrected using observed profiles for the Southern Ocean (de Baar et al., 1999; Boye et al., 2001). Sensitivity tests indicated that high values stemming from a hydrothermal vent in the Eastern Equatorial Pacific lead to unreasonably large values in the interior Pacific Ocean due to advective fluxes. Therefore, the region spanning the latitudes of 12.5°S - 9.5°N, longitudes 72°W - 106°W was masked to a maximum value of  
 225  $0.3 \mu\text{mol m}^{-3}$  (below 2000 m). All other tracers were initialized with small values.

Iron was supplied to the ocean by dust deposition and from sediments. The sedimentary flux was assumed to scale with organic nitrogen flux into the sediment, as found in Elrod et al. (2004). REcoM3 used monthly averages of dust deposition (Albani et al., 2014). We assumed that 3.5% of the dust field consists of iron of which 1.5% dissolves into a bio-available form when deposited in the surface ocean. We did not include aeolian nitrogen deposition in our simulations.

**Table 1.** List of simulations performed in this study.

Experiment	Period	Atmospheric CO <sub>2</sub>	Atmospheric Forcing
pre-spinup	1611 – 1799	constant (278 ppm)	RYF61
<i>A<sub>spinup</sub></i>	1800 – 1957	increasing	RYF61
<i>B<sub>spinup</sub></i>	1800 – 1957	constant (278 ppm)	RYF61
A	1958 – 2021	increasing	JRA55-do
B	1958 – 2021	constant (278 ppm)	RYF61
D	1958 – 2021	constant (278 ppm)	JRA55-do

230 The atmospheric reanalysis data sets of JRA55-do v.1.5.0 (Tsuji no et al., 2018) were used to force the model for the period 1958-2021 (hereafter, JRA55-do). A single repeating annual cycle of all forcing fields (year 1961) was used to perform the spinup simulations and a control experiment. This is referred to as Repeat Year Forcing (hereafter called RYF61). We have deliberately chosen the year 1961 as it had rather neutral El Niño conditions and further contained a low amount of anthropogenic perturbation as compared to the years 1990 and 1991 recommended by Stewart et al. (2020).

235 A series of experiments was carried out in a global setup to investigate the performance of the coupled FESOM2.1-REcoM3 model. The experiments follow the definitions used in the Global Carbon Budget (Friedlingstein et al., 2022a) and in the RECCAP (Regional Carbon Cycle Assessment and Processes, <https://reccap2-ocean.github.io/>) projects and are summarized in Table 1. Our first experiment was forced with varying climate from the JRA55-do data set, and varying atmospheric CO<sub>2</sub> levels (hereafter referred as A). Atmospheric CO<sub>2</sub> mixing ratio (xCO<sub>2</sub>) values are taken from the Global Carbon Budget (Joos and Spahni, 2008; Ballantyne et al., 2012; Friedlingstein et al., 2022a). A second simulation was forced by RYF61 atmospheric reanalysis fields and a preindustrial atmospheric CO<sub>2</sub> mixing ratio of 278 ppm. This configuration, here termed as B, is considered as the control run. Our last simulation was forced with varying climate from the JRA55-do data set and a preindustrial atmospheric CO<sub>2</sub> mixing ratio of 278 ppm. This experiment is referred as D and is used to separate the effects of rising atmospheric CO<sub>2</sub> and of climate change on the DIC inventory. Using the simulations A and B, the global ocean anthropogenic CO<sub>2</sub>  
 240 sink was estimated by taking the model biases and drift from the control run into account. We used a coupled system spinup  
 245


(i.e., a direct strategy, Séférian et al., 2016). Before starting simulations A, B and D, we performed spinup experiments in two stages. In the first stage, a 189-year long (equivalent to three cycles of JRA55-do forcing) preindustrial spinup simulation (named as pre-spinup) was conducted using RYF61 atmospheric forcing and a preindustrial atmospheric CO<sub>2</sub> mixing ratio of 278 ppm until the air-sea CO<sub>2</sub> reaches a quasi-equilibrium state. The  $A_{spinup}$  and  $B_{spinup}$  simulations are a continuation of the pre-spinup simulation with either increasing ( $A_{spinup}$ ) or constant ( $B_{spinup}$ ) atmospheric CO<sub>2</sub> and run from 1800–1957. From the spinup simulations, A, B and D were branched off in 1958 and run until the end of 2021. FESOM1.4-REcoM2 and FESOM2.1-RECOM3 reach a throughput of 6 simulated years per day (SYPD) and 16 SYPD using the same mesh configuration and the same experimental setup (See Table 1) on 288 cores with time steps of 15 min and 45 min, respectively. All modelled mean fields shown in this work are averaged over the period 2012–2021 unless stated otherwise.

**Table 2.** List of the observational data sets used to initialize the biogeochemistry model and assess its performance.

Data set	Variable name	Unit	Reference
Dissolved inorganic carbon	DIC	mmol m <sup>-3</sup>	Global Ocean Data Analysis Project version 2 (Lauvset et al., 2016)
Total Alkalinity	Alk	mmol m <sup>-3</sup>	Global Ocean Data Analysis Project version 2 (Lauvset et al., 2016)
Dissolved inorganic nitrogen	DIN	mmol m <sup>-3</sup>	World Ocean Atlas (Garcia et al., 2014)
Dissolved inorganic silicon	DSi	mmol m <sup>-3</sup>	World Ocean Atlas (Garcia et al., 2014)
Oxygen	O <sub>2</sub>	mmol m <sup>-3</sup>	World Ocean Atlas (Garcia et al., 2019b)
Chlorophyll <i>a</i> concentration	Chl	mg m <sup>-3</sup>	OC-CCI (Sathyendranath et al., 2019) and Southern Ocean (Johnson et al., 2013)
Net primary production	NPP	mmol m <sup>-3</sup>	CPBM (Westberry et al., 2008) and VGPM (Behrenfeld and Falkowski, 1997)


In this section we assess the performance of FESOM2.1-REcoM3 in simulating the observed mean state of nutrients, chlorophyll  $a$ , net primary production, and export production in the near-surface ocean as well as air-sea  $\text{CO}_2$  flux primarily under elevating  $\text{CO}_2$ . Before assessing the biogeochemical variables, we analyze key features of the ocean model.

### 3.1 Modelled hydrography, mixed layer and Atlantic meridional overturning




**Figure 3.** Maps of simulated FESOM2.1-REcoM3 (Sim A) surface temperature [ $^{\circ}\text{C}$ ] (A), practical salinity [ $\text{g kg}^{-1}$ ] (D) with observations from the Polar Science Center Hydrographic Climatology (PHC3, updated from Steele et al., 2001) and corresponding differences (C, F) averaged over the time period 2012-2021.

260 An extended analysis of analogous FESOM runs is presented in Scholz et al. (2019, 2021). Here we analyze only a few relevant diagnostics to prove the validity of the presented research. We start the analysis by inspecting the spatial distribution of the model bias in surface hydrography, presented in Fig. 3 as the difference between modeled mean 2012-2021 and the PHC3 Climatology (Steele et al., 2001). For temperature and salinity respectively, we found a global spatial correlation coefficient ( $r$ ) of 0.99 and 0.99, with a Root Mean Squared Error (RMSE) of  $0.82^{\circ}\text{C}$  and  $0.43$  psu. In the northern North Atlantic the bias is expressed by the cold ( $\sim 4^{\circ}\text{C}$  colder) and fresh ( $\sim 1$  psu fresher) anomalies around Newfoundland which is the typical bias for standalone and climate models at coarse resolutions (see e.g., Scaife et al., 2011). Further south, the bias depicts a  
265




**Figure 4.** Vertical representation of the Atlantic Meridional Overturning Circulation [Sv] in simulations A, B and their difference [Sv].



**Figure 5.** Time-series of annual mean Atlantic Meridional Overturning Circulation [Sv] maxima in simulations A and B.

dipole anomaly associated with the Gulf Stream going too far north, which is a commonly addressed shortcoming for non-eddy-permitting models (see e.g., Zhang and Vallis, 2007; Storkey et al., 2018). Similar issues are found in comparable current systems, such as the Kuroshio and Malvina systems. It is, however, surprising that in general FESOM is far too saline at the surface being on average 0.3 psu saltier than the climatology. The reason for this bias could be the imperfections in the river discharge from CORE forcing and the relatively low surface salinity restoring, using a piston velocity of 50m/300 days in the simulations. In most of the ocean the SST and SSS differences act in an opposite manner at buoyancy. Hence increase or decrease in SST is accompanied with an increase or decrease in SSS. The only exception is the Indian Ocean, where East and West in sim A become less and more buoyant, respectively (Fig. 3).

In Fig. 4, we augment the diagnostic by inspecting the Atlantic meridional overturning circulation (AMOC) which provides the most general characteristic of water mass transformation and production. The mean AMOC in both runs is expressed with the basinwide mid depth cell showing a maximum of ~ 15 Sv at ca 40°N. The bottom cell, induced by the circulation of the Antarctic Bottom water, is also well reproduced with a minimum of ~ -5 Sv. Even though the runs depict large differences in temperature and salinity from the observed climatology, the simulated AMOC shows the canonical picture as known from




**Figure 6.** Maps of simulated FESOM2.1-REcoM3 (Sim A) maximum Mixed Layer Depth [m] in March (A), and September (D) averaged over the time period 2012-2021, observation-based maximum Mixed Layer Depth from Sallée et al. (2021, B and E) occupied between 1970 and 2018 and corresponding differences (C, F).

280 other stand-alone ocean and coupled climate models (Griffies et al., 2009; Jungclaus et al., 2013; Danabasoglu et al., 2014). This indicates that although biases in the representation of water mass properties and ventilation mechanisms are present, they still result in a reasonable density distribution which maintains realistic transports.

The difference between runs A and B shows that the mid depth and bottom cells are stronger in simulation B. Consequently, the difference A-B is expressed by a basinwide positive anomaly with a maximum of  $\sim 3$  Sv. We also show the time-series, for both runs, of AMOC maxima for the years 1958-2021 (Fig. 5). In run A, the time-series depicts a multidecadal variability with a minimum of  $\sim 9.5$  Sv and a maximum of  $\sim 13.5$  Sv. Concurrently the reference run B depicts nearly constant value, but with a small increase, between 9.5 Sv and 10 Sv, which is a result of the repeated year forcing.

290 Finally in Fig. 6, we present the simulated and observed (Sallée et al., 2021, referred to as Atlas) annual maximum mixed layer depth (MLD) pattern for March and September following the same methods (the depth at which the potential density referenced to the surface exceeds the density of the water at 10 m by a threshold of  $0.03 \text{ kg m}^{-3}$ ). Overall, the modelled MLD fits well with the observations although some common discrepancies remained in the deep mixing areas. In the Northern Hemisphere, the deepest MLD ( $>1000$  m) is found in the Labrador as well as in Greenland-Iceland-Norwegian Seas. The magnitude is larger than in Sallée et al. (2021), but is in the same range as other modelling studies (Griffies et al., 2009; Sidorenko et al., 2011). In the Southern Hemisphere, winter deep mixing in high-latitudes is also overestimated compared to 295 the observations, especially in the Pacific sector of the Southern Ocean. From inspecting the model runs and their differences we conclude that FESOM2.1 simulated a reasonable ocean state which can be used for further analysis.

### 3.2 Nutrients, ocean productivity and ecosystem




**Figure 7.** Maps of simulated FESOM2.1-REcoM3 (Sim A) surface [0-100m] concentration of dissolved inorganic nitrogen [ $\text{mmol m}^{-3}$ ] (A), dissolved inorganic silicon [ $\text{mmol m}^{-3}$ ] (D) with observations from the World Ocean Atlas 2018 climatology (B and E, Garcia et al., 2019b) and corresponding differences (C, F) averaged over the time period 2012-2021.

#### 3.2.1 Modeled versus in situ nutrients

We first compared the spatial distribution of surface (averaged over the top 100m depth layer) ocean dissolved inorganic nitrogen (DIN) and dissolved silicate (DSi) from REcoM3 with the World Ocean Atlas 2018 (Garcia et al., 2019b) climatologies (Fig. 7). While simulated surface DIN concentrations were lower than observations in the subpolar regions, a large positive DIN bias of up to  $20 \text{ mmol m}^{-3}$  was found in the subtropical South Pacific Ocean. The simulated DSi was overestimated in the Southern Ocean and underestimated in the northern Pacific. Exceptions are the Pacific and Atlantic sectors of the coastal Southern Ocean where the modeled DSi concentrations are lower than the observations. These patterns were already present in FESOM1.4-REcoM2 (Schourup-Kristensen et al., 2014), however, two recent improvements should be noted. First, the large and positive DIN bias in the northern subtropical Pacific (Schourup-Kristensen et al., 2014) disappeared. This is caused by replacing the dust deposition input forcing field from Mahowald et al. (2003) with Albani et al. (2014), which results in more realistic (i.e., less strong) iron limitation. Second, the silicate bias in the Southern Ocean is reduced in magnitude and extent

compared to Schourup-Kristensen et al. (2014). This is related to tuning experiments (not shown), which resulted in a larger  
 310 share of diatoms in the Southern Ocean (Fig. 11) compared to Schourup-Kristensen et al. (2014), thus drawing down more  
 silicic acid. Along with the increased share of diatoms, also the Southern Ocean and global opal export increased. For the  
 global ocean, the opal export increased from 74.5 Tmol Si yr<sup>-1</sup> in Schourup-Kristensen et al. (2014) to 168 Tmol Si yr<sup>-1</sup>  
 in the present study and is thus at the upper instead of the lower end of the range of 69-185 Tmol Si yr<sup>-1</sup> (Dunne et al.,  
 2007) and higher than the best estimate of Tréguer et al. (2021) (Table 3). In the Southern Ocean, opal export increased  
 315 from 21.5 Tmol Si yr<sup>-1</sup> in Schourup-Kristensen et al. (2014) to 85.5 Tmol Si yr<sup>-1</sup>, which is higher than the range of 21-  
 54 Tmol Si yr<sup>-1</sup> reported by (Dunne et al., 2007). The silicic acid bias is rather insensitive to formulation and parameter  
 choice of opal dissolution, but very sensitive to the share of diatoms in the Southern Ocean. The correlation coefficient (*r*) and  
 root mean squared error (RMSE) between simulated and observed annual mean were 0.88 and 0.86 mmol m<sup>-3</sup> for DIN, and  
 0.47 and 0.54 mmol m<sup>-3</sup> for DSi. The correlation with observed DIN is higher than in Schourup-Kristensen et al. (2014, 0.75),  
 320 which we relate to the disappearance of the DIN bias in the northern subtropical Pacific. The correlation with observed DSi  
 is lower than in FESOM-1.4-REcoM2, despite the reduction in magnitude and extent in the Southern Ocean DSi bias. Moderately  
 high silicic acid values in the northern high latitudes are not reproduced. This may be related to mixing that is too sluggish, or  
 to overly strong silicic acid draw-down by diatoms (Figs. 9 and A1), possibly linked to iron limitation that may be too weak  
 (Fig. 12).



**Figure 8.** Maps of simulated FESOM2.1-REcoM3 (Sim A) surface [0-50m] concentration of dissolved iron [ $\mu\text{mol m}^{-3}$ ] (A), and of the  
 AI-based global reconstruction by Huang et al. (2022) (B). Note the different colorscale for the two plots.

325 Despite the enormous increase in the number of observations of dissolved iron with the GEOTRACES project, observations  
 have not reached a global coverage that makes it possible to construct a global climatology. Therefore the modeled dissolved  
 iron is compared here to the global surface pattern of dissolved iron by Huang et al. (2022), which uses an artificial intelligence  
 method (random forest) to construct a near global iron field, based on the observations in the second intermediate GEOTRACES

data product (Schlitzer et al., 2018), plus some older in-situ iron observations compiled in Tagliabue et al. (2012), and on co-located hydrographic observations. The pattern of modeled dissolved iron (Fig. 8, averaged over the top 50 m) shows the expected pattern of high concentrations in regions with high dust deposition, mainly in the tropical Atlantic Ocean and the eastern part of the Arabian Sea, but also to some extent in the southern subtropical Atlantic and Indian Oceans. Concentrations are extremely low in the subpolar Southern Ocean, and almost the whole Equatorial and South Pacific. Iron concentrations are also low in the subpolar North Pacific, and — less so, but still noticeable — in the subpolar North Atlantic. Oceanic regions adjacent to extended shelves, especially in the Arctic, show somewhat elevated iron concentrations. If we compare this to the AI-generated global pattern of dissolved iron from Huang et al. (2022) we find qualitatively similar patterns, like the elevated iron concentration in the equatorial and subtropical Atlantic and the Arabian Sea, or the low concentrations in the subpolar Southern Ocean, the equatorial Pacific, and the subpolar North Pacific, but the amplitude of the patterns is overall smaller. The largest discrepancy in amplitude is found under the Saharan dust plume in the tropical Atlantic, where the model produces maximal dissolved iron values that are almost three times as high as the reconstruction from Huang et al. (2022). Direct observations in the tropical Atlantic also show dissolved iron concentrations that reach  $1.2 \text{ nmol L}^{-1}$  (e.g. Hatta et al., 2015), while modeled maxima are  $> 3 \text{ nmol L}^{-1}$ . A further important difference is that the distribution by Huang et al. (2022) shows slightly elevated iron concentrations in the center of the subtropical South Pacific, where the model in contrast has extremely low values. This discrepancy causes a too strong iron limitation in this region in the model, probably explaining the overly high DIN concentrations in the model South Pacific. The too strong amplitude of the patterns in modeled dissolved iron, which is also found in other models, likely has a number of causes. The most important one is the assumption of a constant solubility in dust-deposited iron. Dust deposition close to the main source regions is on average coarser and has experienced less chemical processing during its transport, which both would lead to a lower solubility. The opposite is true for particles deposited far from the source regions, such as in the South Pacific. A second contribution might be the missing source from pyrogenic aerosols, which are far more soluble. Also, the effect of dust particles as iron scavengers, which has not been included in this simulation, has been shown to reduce the overly high dissolved iron concentrations often found in models under the main dust deposition regions (Ye and Völker, 2017; Pagnone et al., 2019). Furthermore, the intensity and extension of dust plumes vary between modelled dust deposition fluxes (e.g. Myriokefalitakis et al., 2018). The field of dust deposition by Albani et al. (2014), used in our model to calculate aeolian iron input, is within the range of modern estimates but surely contains some uncertainties. Despite the overall too strong amplitude of the patterns in dissolved iron, especially in the regions of high dust deposition, the model is able to reproduce the main regions where iron availability limits phytoplankton productivity (Moore et al., 2013), namely the subpolar Southern Ocean, the equatorial and North Pacific, and to some extent also the seasonal iron limitation in the subpolar North Atlantic (Nielsdóttir et al., 2009), but overestimates iron limitation in the subtropical South Pacific.

### 3.2.2 Modeled versus satellite-based phytoplankton biomass and productivity

We first evaluated spatial distributions of modelled chlorophyll *a* concentration (Fig. 9), an indicator for phytoplankton biomass, and vertically integrated Net Primary Production (NPP, Fig. 10). We compared chlorophyll *a* concentrations obtained from FESOM2.1-REcoM3 simulations, averaged from 2012 to 2021, with an ocean color remote sensing merged data



set (from the Ocean Colour Climate Change Initiative, OC-CCI, (Sathyendranath et al., 2019), averaged from 1998 to 2019. We also compared the modeled NPP with satellite estimations such as the Vertically Generalized Production Model (VGPM, Behrenfeld and Falkowski, 1997, Fig. 10) and the updated Carbon-based Productivity Model (CbPM, Westberry et al., 2008, see Appendix Fig. A1). VGPM is a chlorophyll-based algorithm that can be considered as a standard NPP estimation from ocean color for the last 20 years (Lee and Marra, 2022). CbPM uses spectrally resolved light attenuation and is based on a semi-analytical algorithm (Garver-Siegel-Maritorena, GSM, Maritorena et al., 2002). All data sets were also compared along a latitudinal distribution together with an improved chlorophyll *a* algorithm for the Southern Ocean (Johnson et al., 2013, Fig. 11).

The results for chlorophyll *a* and NPP obtained here are comparable to those presented by Schourup-Kristensen et al. (2014). Over large parts of the global ocean, the mean surface chlorophyll *a* concentrations are in agreement with observations (Fig. 9 panels C and D). Yet, there are regional differences. In temperate latitudes, the modeled chlorophyll *a* concentrations are somewhat higher than observed while the subtropical gyres show concentrations slightly lower than observations. The comparison of modeled and observational-based satellite estimates of chlorophyll *a* yielded a correlation of 0.66 and an RMSE of 0.38 mg m<sup>-3</sup>. FESOM2.1-REcoM3 also shows a reasonably well simulated latitudinal variation of chlorophyll *a* compared to satellite estimations (Fig. 11A). The model underestimates chlorophyll *a* concentrations in most of the coastal regions, especially in the high latitude regions. In the southern high latitudes, FESOM2.1-REcoM3 follows the Southern Ocean adjusted chlorophyll *a* data set (Johnson et al., 2013) quite well, except for the coastal regions close to Antarctica (approximately south of 70°S). In the Arctic Ocean, the model strongly underestimates chlorophyll *a* concentrations which is driven by negative biases reaching up to 3 mg chlorophyll *a* m<sup>-3</sup> on the continental shelves (Fig. 9). Although FESOM2.1-REcoM3 did reproduce well NPP distribution at low latitudes (Fig. 10,11), it also strongly under-estimated NPP at higher latitudes when compared to VGPM ( $r = 0.43$ ,  $RMSE = 0.34 \text{ mgC m}^{-2} \text{ d}^{-1}$ ), in particular in productive areas north of 50°N and coastal areas (Fig. 10). For regional applications, further analysis and possibly tuning may be needed. When compared with VGPM, the model simulation generally underestimated the remotely sensed NPP estimations (Table 3), especially in the subtropical Pacific. Yet, with a value of 35.9 PgC yr<sup>-1</sup> the modeled global total NPP is slightly above the range of earlier modeling studies (23.7 - 30.7 PgC yr<sup>-1</sup>, Schneider et al., 2008), and within the range of recent Earth System Models (24.5 - 57.3 PgC yr<sup>-1</sup>, S  f  rian et al., 2020). It is lower than other satellite-based estimates of 47.3 PgC yr<sup>-1</sup> (Behrenfeld and Falkowski, 1997), 52 PgC yr<sup>-1</sup> (Westberry et al., 2008) and 48.7 - 52.5 PgC yr<sup>-1</sup> reported by Kulk et al. (2020).

Both simulated chlorophyll *a* concentrations and NPP from FESOM2.1-REcoM3 seemed to be under-estimated in coastal regions. Too low primary production and chlorophyll *a* levels were particularly evident in coastal regions, which could be linked to deficiencies in either the chlorophyll *a* data set and/or in the model. For the former, the chlorophyll *a* OC-CCI data set and the CbPM primary production data set uses the GSM algorithm. GSM tries to distinguish optical signatures from phytoplankton, particles and dissolved organic matter, but still requires regional tuning in coastal regions, where the presence of non-biotic optically active material (i.e. yellow substances and sediments) makes chlorophyll *a* retrieval challenging (Blondeau-Patissier et al., 2014). The over-estimation of chlorophyll *a* in coastal waters is even more pronounced with standard global chlorophyll algorithms used in the VGPM primary production data set, such as OC4 that are only adapted to CASE-I waters (low influence


of dissolved organic matter and non-algal particles). Therefore, both remotely sensed NPP estimations carry uncertainties related to the global algorithms. For example, turbid waters over the Arctic shelves are known to artificially increase both chlorophyll *a* and NPP estimates from remote sensing (Matsuoka et al., 2012; Mitchell, 1992; Mustapha et al., 2012). Some recent advances used local parametrizations with in situ data, which resulted in much lower productivity levels in those coastal areas (Lewis et al., 2020; Lewis and Arrigo, 2020). Generally, the NPP and chlorophyll differences to satellite-based estimates could also be linked to model deficiencies, such as coarse model resolution and associated weak upwelling, missing complexity in simulated phytoplankton classes, but also the so far unconsidered nutrient input from terrigenous sources.

The low values of primary production could be caused by several top-down and/or bottom-up effects. The nutrient dynamics that partly control NPP, are the result of a delicate balance between physical (mixing, stratification and upwelling systems) and biogeochemical processes. To investigate bottom-up controls on regional NPP dynamics, we derived the most limiting factor (either light or nutrients) of growth of diatom and small phytoplankton. This factor ranges between 0 (most limiting) and 1 (no limitation) and is based on the nutrient uptake Michaelis–Menten kinetics of REcoM. The Michaelis–Menten coefficient (MM) is computed as  $MM = [Nut]/([Nut] + KNut)$ , with [Nut] being the nutrient concentration, and KNut a nutrient and phytoplankton dependent half-saturation constant. The light limitation is defined as the carbon-specific photosynthesis rate divided by the maximum photosynthetic rate. We derived maps showing the most limiting factor (factor closest to zero, either nutrients DIN, DSi, or DFe, or light) in the annual mean (Fig. 12).


The spatial distribution of the dominant growth-limiting factor for diatoms and small phytoplankton over the time period 2012-2021 is shown in Fig. 12. Over large areas of the Southern Pacific and almost the entire Southern Ocean diatoms were limited by iron availability. Elsewhere, except for the Arctic Ocean where light was the most limiting factor, diatom growth was controlled by the abundance of dissolved silicic acid. Nutrient uptake of small phytoplankton was limited by iron in the South Pacific, DIN within the band of 45°S-45°N in the Atlantic and Indian Oceans and insufficient light at high latitudes (south of 45°S and north of 45°N).

The large-scale patterns of limitation were in general agreement with observations (Moore et al., 2013) and other modelling studies (Long et al., 2021a), although the degree of silicic acid limitation for diatoms (outside the iron-limited Southern Ocean) varied across models (Laufkötter et al., 2015). The severe iron limitation in most of the Pacific might contribute to the lower productivity levels than observed in the same regions (Fig. 10).


In addition to bottom-up explanations, one can also raise a too high grazing pressure from zooplankton due to the choice of grazing formulations and parameter values as a reason for low primary production (Anderson et al., 2010; Prowe et al., 2012; Karakuş et al., 2021). In fact, Karakuş et al. (2022) demonstrated that a separation of the small zooplankton group in REcoM into micro- and mesozooplankton leads not only to a 25% increase in NPP but also to a reduction of overly strong iron limitation in the South Pacific, due to nutrient recycling by zooplankton. Further, REcoM does not explicitly represent picophytoplankton (e.g., non N<sub>2</sub>-fixing cyanobacteria such as *Synechococcus* and *Prochlorococcus*) and nitrogen fixers, and this might contribute to an underestimation of NPP.



**Figure 9.** Maps of simulated FESOM2.1-REcoM3 (Sim A) surface (log10 transformed) chlorophyll *a* concentration [ $\text{mg Chl } a \text{ m}^{-3}$ ] of small phytoplankton (A), diatoms (B) and the sum of both phytoplankton groups (C). The satellite-based merged dataset OC-CCI is shown in D (Sathyendranath et al., 2019) with corresponding differences between FESOM2.1-REcoM3 and OC-CCI (E). Note the different time periods of the simulation (2012-2021) and OC-CCI (1998-2019).



**Figure 10.** Maps of simulated FESOM2.1-REcoM3 (Sim A) vertically integrated net primary production [ $\text{mg C m}^{-2} \text{ d}^{-1}$ ] of small phytoplankton (A), diatoms (B), and the sum of both phytoplankton groups (C). The satellite-based Vertically Generalized Production Model (VGPM) is shown in D (Behrenfeld and Falkowski, 1997) with corresponding differences between FESOM2.1-REcoM3 and VGPM (E). All fields are averaged over the time period 2012 to 2021.




**Figure 11.** Latitudinal distribution of vertically integrated and zonally averaged (A) chlorophyll  $a$  [ $\text{mg Chl m}^{-3}$ ] and (B) net primary production [ $\text{mg C m}^{-2} \text{d}^{-1}$ ] simulated by FESOM2.1-REcoM3 (blue line). The satellite-based merged chlorophyll  $a$  datasets of OC-CCI (Sathyendranath et al., 2019, orange line) and the improved chlorophyll  $a$  algorithm for the Southern Ocean (Johnson et al., 2013, green line) are shown in (A). The satellite-based dataset of the Vertically Generalized Production Model (VGPM) (Behrenfeld and Falkowski, 1997, purple line) and the Carbon-based Productivity Model (CbPM) (Westberry et al., 2008, red line) are shown in (B).

### 3.2.3 Modeled versus MAREDAT zooplankton biomass

In REcoM3, the small zooplankton group is widely spread in the global ocean and the highest biomass occurs in high-productivity regions (Fig. 13A). The macrozooplankton is present in the high latitudes (Fig. 13B) since it is parametrized as a polar macrozooplankton group (Karakuş et al., 2021). We compare the latitudinal distribution of integrated modeled zooplankton biomass with gridded global zooplankton biomass data from MAREDAT (Buitenhuis et al., 2010; Moriarty et al., 2013; Moriarty and O'Brien, 2013). The simulated biomass of small and total zooplankton reproduces MAREDAT-derived biomass reasonably well in low to mid latitudes, but underestimates biomass in the polar regions (Fig. 13C). The underestimation of zooplankton biomass in the northern high latitudes may be related to an underestimation of primary production in the same region. In agreement with the MAREDAT data set (Moriarty et al., 2013), macrozooplankton is not present in low latitudes.

### 3.2.4 Synthesis

The modeled biogeochemical fluxes were compared to the previous version FESOM1.4-REcoM2 and observational studies (Table 3). Modelled global NPP is higher in FESOM2.1-REcoM3 than in FESOM1.4-REcoM2, but still lower than in satellite-




**Figure 12.** Maps showing the spatial distribution of the most limiting factor in the model’s surface water for diatoms (A) and small phytoplankton (B). Hatching denotes areas of weak limitation (all limiting factors  $>0.5$ ). Fe: iron, DIN: dissolved inorganic nitrogen, DSi: dissolved silicic acid.

based estimates. The estimate is comparable to other global modelling studies (Schneider et al., 2008; Séférian et al., 2020).

445 Export production (EP) is slightly higher in FESOM2.1-REcoM3 than in the previous version, and falls within the observational range previously documented in the literature for both the global ocean and the Southern Ocean. For the global ocean, FESOM2.1-REcoM3 NPP and EP estimations remained at the lower end of the range despite a slight increase in NPP. A more detailed description of zooplankton results in more efficient nutrient recycling and can thus increase NPP by 25% (see also explanation in section 3.2.2, Karakuş et al., 2022). In the Southern Ocean, estimations of NPP and EP remained very close

450 to observation-based estimates. Maybe the most noticeable change between the two model versions is the substantial increase in opal export which increased by a factor of four in the Southern Ocean, passing from the lower to the higher end of the observational range of an earlier review (Dunne et al., 2007), and lies above an updated estimate (Tréguer et al., 2021). This is due to an increase in diatoms relative contribution to the total NPP in high latitudes (Fig. 11).



**Figure 13.** Maps of annual mean surface (A) small zooplankton and (B) macrozooplankton concentrations in FESOM2.1-REcoM3 (Sim A). Latitudinal distribution of vertically integrated zooplankton biomass  $[\text{mg C m}^{-2}]$  of (C) modeled small zooplankton (solid blue line) and sum of microzooplankton and mesozooplankton from MAREDAT (orange dots for individual observations and solid brown line for the zonal mean of the observations, Buitenhuis et al., 2010; Moriarty and O'Brien, 2013) and (D) modeled macrozooplankton (solid blue line) and macrozooplankton from MAREDAT (orange dots, Moriarty et al., 2013). Modeled zooplankton biomass is averaged over the time period 2012 to 2021. The zonal mean of macrozooplankton is not calculated due to the low number of observations.


**Table 3.** Global and Southern Ocean net primary production (NPP) and export production (EP) in FESOM2.1-REcoM3 and estimates from the literature. The Southern Ocean is considered as the region south of 50°S. The numbers for VGPM and CbPM are recalculated after interpolation to the model mesh over the years 2012-2019

	Unit	FESOM1.4-REcoM2 (Sim. A)	FESOM2.1-REcoM3 (Sim. A)	Range from literature
NPP global	PgC yr <sup>-1</sup>	32.5	35.8	50.5 (VGPM, this study) 68.9 (CbPM, this study) 47.3 (Behrenfeld and Falkowski, 1997) 52 (Westberry et al., 2008) 23.7 – 30.7 (Schneider et al., 2008) 48.7 - 52.5 (Kulk et al., 2020) 24.5 - 57.3 (CMIP6) (Séférian et al., 2020)
EP global	PgC yr <sup>-1</sup>	6.1	6.3	9.6 (Schlitzer, 2004) 5.8-13 (Dunne et al., 2007) 5 (Henson et al., 2011) 5.9 (Siegel et al., 2014)
Opal export global	Tmol Si yr <sup>-1</sup>	74.5	168	69-185 (review in Dunne et al., 2007) 112 (Tréguer et al., 2021)
CaCO <sub>3</sub> export global	PgC yr <sup>-1</sup>	1.2	0.89	0.1 - 4.7 (Seifert et al., 2022)
NPP Southern Ocean	PgC yr <sup>-1</sup>	3.1	3.2	3.48 (VGPM, this study) 3.92 (CbPM, this study) 1.1-4.9 (Carr et al., 2006) 5.7 (Behrenfeld and Falkowski, 1997)
EP Southern Ocean	PgC yr <sup>-1</sup>	1.1	1.5	1.0 (Schlitzer, 2002; Nevison et al., 2012)
Opal export Southern Ocean	Tmol Si yr <sup>-1</sup>	21.5	85.5	21-54 (Dunne et al., 2007)
CaCO <sub>3</sub> export Southern Ocean	PgC yr <sup>-1</sup>		0.31	0.018 (Dunne et al., 2007)



### 3.3 Carbon cycle

#### 455 3.3.1 Dissolved inorganic carbon and alkalinity



**Figure 14.** Maps of simulated FESOM2.1-REcoM3 (Sim A) surface [0-100 m] concentration of dissolved inorganic carbon [ $\text{mmol m}^{-3}$ ] (A) and alkalinity [ $\text{mmol m}^{-3}$ ] (D); corresponding data from GLODAPv2 (B, E; Lauvset et al., 2016) and model-data differences (C, F). The comparison is for the period 1998-2006 which is centered at year 2002 to be comparable to GLODAP.


Insight into the carbonate system can be obtained by inspecting surface maps of modeled dissolved inorganic carbon and alkalinity and the corresponding observational GLODAPv2 climatologies (Fig. 14). Global patterns of simulated concentrations resemble the observed fields reasonably well ( $r = 0.99$ ,  $\text{RMSE} = 36.5 \text{ mmol m}^{-3}$ , calculated from annual means) with highest DIC values in the subtropical gyres of the Atlantic and south Pacific, as well as the subpolar North Atlantic and the Southern Ocean. Similar to GLODAP, highest alkalinity values are found in the subtropical gyres of the Atlantic and south Pacific with a good agreement with global spatial features ( $r = 0.99$ ,  $\text{RMSE} = 33.9 \text{ mmol m}^{-3}$ ). Yet, simulated surface surface DIC and alkalinity concentrations were slightly overestimated in most of the global ocean. Two major exceptions are the Arctic Ocean and the North Atlantic where the concentrations were underestimated, as well as the tropical upwelling regions and the northern Indian Ocean for DIC. The departure from observations differ in their patterns relative to FESOM-1.4-REcoM (which was too low in DIC and alkalinity in the tropics and subtropics, too high in high latitudes, not shown), which indicates that different realisations in circulation or mixing may drive these bias patterns. This is in line with an overestimation of surface salinity in

460

465

most of the global ocean with the exception of the North Atlantic and the Arctic Ocean (see Fig. 3). Also, surface alkalinity biases are generally attributed to a dominant physical (preformed) signal with a smaller contribution from the calcium carbonate cycle and a negligible contribution from organic matter remineralization (Koeve et al., 2014). However, tuning the model to result in a higher  $\text{CaCO}_3$  production could possibly also counteract the positive alkalinity bias. Similarly, a higher NPP in the South Pacific could regionally ameliorate the high DIC bias. A positive bias in alkalinity at constant atmospheric  $\text{CO}_2$  in the spin-up (not shown) and simulation A (Fig. 14) leads to a positive bias in DIC as surface water with a higher alkalinity can hold more  $\text{CO}_2$  in equilibrium than a low-alkalinity surface ocean. The range of biases is similar ( $\pm 100 \text{ mmol m}^{-3}$ ) as in other ocean biogeochemical models (e.g., Tjiputra et al., 2020; Long et al., 2021a).

### 475 3.3.2 Surface ocean $\text{pCO}_2$ and air-sea $\text{CO}_2$ flux



**Figure 15.** Maps of surface ocean  $\text{pCO}_2$  [ $\mu\text{atm}$ ]. The top row compares the (A) simulated FESOM2.1-REcoM3 (Sim A) surface partial pressure of  $\text{CO}_2$  to the (B)  $\text{pCO}_2$ -based data-product (Chau et al., 2022), both averaged over 2012-2020. The bottom panel (C) shows model-data differences. Note that simulation and observations are masked every month with sea-ice concentration  $> 15\%$


We compare the pattern of the temporal mean (2012-2021) surface ocean partial pressure of  $\text{CO}_2$  ( $\text{pCO}_2$ , Fig. 15) and air-sea  $\text{CO}_2$  flux (Fig. 16) to the  $\text{pCO}_2$ -based data-product of Chau et al. (2022) with a seamless coverage from open ocean

to the coasts (Fig. 16). Different pCO<sub>2</sub>-products largely agree with each other in terms of spatial patterns, although they may differ with respect to amplitude and timing of variability of regionally or globally integrated fluxes (Fay et al., 2021; Fay and McKinley, 2021). Therefore we chose one of them (Chau et al., 2022) and focus on the comparison of the spatial pattern with our model. We further evaluate the temporal evolution of pCO<sub>2</sub> in FESOM2.1-REcoM3 with a direct comparison to surface ocean pCO<sub>2</sub> observations from the Surface Ocean CO<sub>2</sub> Atlas (SOCAT, Bakker et al., 2016), where we subsampled the model output for spatio-temporal locations where observations exist, following Hauck et al. (2020) and Friedlingstein et al. (2022b) in Fig. 17.

The large-scale spatial patterns of pCO<sub>2</sub> are well reproduced (Fig. 15) with high values in the tropics that are typically higher than atmospheric values (red colors), and lower values in the subpolar Southern and Pacific Ocean and the high-latitude North Atlantic ( $r = 0.75$  and  $RMSE = 29.2 \mu\text{atm}$ , ice-free areas). However, compared to the pCO<sub>2</sub>-product of Chau et al. (2022), model pCO<sub>2</sub> values are overestimated in the subtropical gyres (Fig. 15C). Further, the North Atlantic pCO<sub>2</sub> is on average lower than the pCO<sub>2</sub>-product, and the two data sets also differ on over- versus undersaturation of pCO<sub>2</sub> relative to the atmosphere in the polar Southern Ocean (higher values in FESOM2.1-REcoM3). The latter may well be explained by a known summer bias in Southern Ocean pCO<sub>2</sub> observations (e.g., Metzl et al., 2006; Gregor et al., 2019). FESOM2.1-REcoM3 also simulates very high pCO<sub>2</sub> values on the Russian shelves in the Arctic, where hardly any observations exist. Similarly high pCO<sub>2</sub> values were reported for this region by Anderson et al. (2009), but missing repeat observations prevent a conclusion on whether this is a robust signal and what its extent in time and space is.

FESOM2.1-REcoM3 reproduced the temporal evolution of surface ocean pCO<sub>2</sub> reasonably well compared to SOCAT when accounting for where and when pCO<sub>2</sub> sampling took place (Fig. 17). The annual correlation coefficient and RMSE between simulated and observed global mean pCO<sub>2</sub> are 0.93 and 4.6  $\mu\text{atm}$ , respectively. The subsampled model follows the SOCAT time-series closely, including its variability due to sampling distribution in space and time. The global mismatch with SOCAT pCO<sub>2</sub> as measured by the RMSE is comparable or slightly below the value for FESOM-1.4-REcoM2 (see supplementary Figure S9, 1985-2018, in Hauck et al., 2020) and comparable, but at the high end of the range of other models in GCB2022 (1990-2021, Friedlingstein et al., 2022b). On a monthly scale, the RMSE is higher (38  $\mu\text{atm}$ ), as the models capture the large-scale patterns better than smaller-scale variability according to a previous assessment (Hauck et al., 2020). An analysis of large-scale regional patterns (North, Tropics, South, Fig. 17) reveals that the model reproduced well the trend, but overestimates the mean of pCO<sub>2</sub> in the tropics and underestimates pCO<sub>2</sub> in the northern extra-tropics and to a lesser extent in the southern extra-tropics in recent decades, as also indicated in the maps (Fig. 15).


The air-sea CO<sub>2</sub> flux spatial pattern was reasonably reproduced by FESOM2.1-REcoM3 with CO<sub>2</sub> uptake in the subpolar regions of both hemispheres, and outgassing in the tropics and north Pacific (Fig. 16,  $r = 0.72$ ,  $RMSE = 1.45 \text{ mol C m}^{-2} \text{ yr}^{-1}$ ). Generally, the CO<sub>2</sub> flux patterns mirror the pCO<sub>2</sub> patterns (Fig. 15), but with the additional imprint of spatial variability of wind speed. Hence, the CO<sub>2</sub> uptake in the subpolar Southern Ocean may appear large compared to pCO<sub>2</sub>, which is not as strongly undersaturated in the South as in the North Atlantic. Regions of mean outgassing in the Southern Ocean are of smaller extent in the model than in the pCO<sub>2</sub>-product. While it is well established that outgassing of CO<sub>2</sub> in the polar Southern Ocean occurs in winter (e.g., Bakker et al., 1997), its magnitude and timing varies between estimates and is under debate (Gruber




**Figure 16.** Maps of air-sea CO<sub>2</sub> fluxes [mol C m<sup>-2</sup> yr<sup>-1</sup>]. The top row compares the (A) simulated FESOM2.1-REcoM3 (Sim A) CO<sub>2</sub> flux to the (B) pCO<sub>2</sub>-based data-product (Chau et al., 2022), both averaged over 2012–2020. The bottom panel (C) shows model-data differences. Negative numbers indicate a flux into the ocean. Note that simulation and observations are masked every month with sea-ice concentration > 15%

et al., 2009; Lenton et al., 2013; Gray et al., 2018; Bushinsky et al., 2019; Sutton et al., 2021; Long et al., 2021b). The misfit between the annual mean modelled CO<sub>2</sub> flux and the pCO<sub>2</sub>-based data-product generally mimic pCO<sub>2</sub> biases and thus shows small positive differences (less uptake or more outgassing) in the subtropical gyres and small negative differences (stronger uptake or less outgassing) in the equatorial Pacific, and the Southern Ocean (Fig. 16, bottom panel). The strongest biases were found in the northern high latitudes (negative bias) and the upwelling zone of the eastern tropical Pacific (positive bias). FESOM2.1-REcoM also generally captures the large-scale patterns of coastal CO<sub>2</sub> fluxes with CO<sub>2</sub> uptake in the mid- and high-latitudes (poleward of 25°N/S) and outgassing in the tropical coastal ocean, as described in a recent synthesis based on low- and high-resolution models and pCO<sub>2</sub>-products (Resplandy et al., 2023). The large mismatch in pCO<sub>2</sub> on the Siberian shelves does not show up in CO<sub>2</sub> flux as sea ice prevents CO<sub>2</sub> outgassing throughout most of the year.

We continue our investigation with the analysis of the global ocean-atmosphere CO<sub>2</sub> flux time-series (Fig. 18). In 1800, the first year of spinup after the first 189 years of pre-spinup of simulation B (not shown), the global ocean-atmosphere CO<sub>2</sub> flux




**Figure 17.** Comparing annual mean  $p\text{CO}_2$  [ $\mu\text{atm}$ ] from FESOM2.1-REcoM3 (Sim A) (subsamped for spatiotemporal locations of observations in SOCAT, red) with observations from SOCATv2022 (light blue, updated from Bakker et al., 2016). Results are shown spatially averaged for (A) the global ocean, (B) the northern hemisphere ( $>30^\circ\text{N}$ ), (C) the Tropics ( $30^\circ\text{S}$ - $30^\circ\text{N}$ ), and (D) the southern hemisphere ( $<30^\circ\text{S}$ ). The time-series are shown for all observations in SOCAT (since 1970), but correlation coefficient  $r$  [unitless] and Root Mean Squared Error RMSE [ $\mu\text{atm}$ ] are indicated in the panels for the time period 1990–2021.



**Figure 18.** Time-series of simulated annual mean global ocean-atmosphere  $\text{CO}_2$  flux in  $\text{PgC yr}^{-1}$  in the experiments conducted in this study. FESOM2.1-REcoM3 spinup was conducted for 347 years (including 189 years of pre-spinup, not shown in the plot) under repeat year forcing taken from the year 1961 (RYF61). Here we show the spin-up since 1800 that is continued as the control simulation B after 1958 for FESOM-1.4-REcoM2 (yellow) and FESOM2.1-REcoM3 (magenta) with a constant  $\text{CO}_2$  concentration of 278 ppm (dashed lines) and the spin-up under increasing  $\text{CO}_2$  that is continued as simulation A after 1958 (solid lines). The control simulation B started in the year 1958 and was conducted for 64 years with RYF61 (dashed lines). Simulation A also started in 1958 and was forced with inter-annual varying forcing JRA55-do-1.4.0 (solid lines). Please note that spinup period for FESOM1.4-REcoM2 and FESOM2.1-REcoM3 differ from each other, the latter being longer than the former.

was already in a stable state and converged towards a value close to zero. Under the assumption that the ocean and atmosphere  
 525 were in equilibrium at constant preindustrial  $\text{CO}_2$  and without riverine carbon transported into the ocean (Aumont et al., 2001; Resplandy et al., 2018; Regnier et al., 2022), an equilibrium flux of zero is expected for simulation B. Any deviation from this can be considered a bias (Hauck et al., 2020). The global bias of the annual air-sea  $\text{CO}_2$  flux in the FESOM2.1-REcoM3 control simulation amounts to  $-0.12 \text{ PgC yr}^{-1}$ . The control simulation conducted with the older model version FESOM1.4 had a larger bias with a positive flux of around  $0.4 \text{ PgC yr}^{-1}$  at the end of the simulation. In addition to the bias, the drift is reduced  
 530 from  $0.00264 \text{ PgC yr}^{-2}$  in FESOM1.4-REcoM2 to  $-0.00011 \text{ PgC yr}^{-2}$  in FESOM2.1-REcoM3 with longer spin-up. Despite different spinup procedures (FESOM1.4 has a shorter spin-up period), sim A with both FESOM2.1 and FESOM1.4 reveals similar  $\text{CO}_2$  fluxes under interannually varying forcing after 1980, which indicates a dominance of the forcing over the initial conditions. This also questions the common assumption that the same bias occurs in the control and historical simulations.

We next assess the model performance of the interannually varying simulation (A) by comparison with the Global Carbon  
 535 Budget's ensemble of  $\text{pCO}_2$ -based data-products and other ocean biogeochemistry models (Fig. 19). Note that all model time series shown in Fig. 19 are referenced relative to their control simulations with a constant atmospheric  $\text{CO}_2$  concentration and without climate-change forcing (simulation B). Although being consistent with the interannual variability, air-sea  $\text{CO}_2$  fluxes of FESOM1.4 are at the lower end of the range compared to other Global Ocean Biogeochemistry Models and  $\text{pCO}_2$ -



**Figure 19.** Globally integrated annual air-sea CO<sub>2</sub> flux from ten Global Ocean Biogeochemistry Models (GOBMs) and seven pCO<sub>2</sub>-based data-products used in the Global Carbon Budget 2022 (Table 4 in Friedlingstein et al., 2022a), namely (Wright et al., 2021; Schwinger et al., 2016; Lacroix et al., 2021; Berthet et al., 2019; Hauck et al., 2020; Liao et al., 2020; Doney et al., 2009; Aumont et al., 2015; Nakano et al., 2011; Urakawa et al., 2020; Long et al., 2021a; Landschützer et al., 2016; Rödenbeck et al., 2022; Chau et al., 2022; Gloege et al., 2021; Zeng et al., 2014; Iida et al., 2015; Gregor and Gruber, 2021) after applying bias correction to the models and river flux adjustment of 0.65 PgC yr<sup>-1</sup> (Regnier et al., 2022) to the pCO<sub>2</sub>-products. The thick black line indicates the model ensemble mean and the thick blue line shows the mean of the pCO<sub>2</sub>-product ensemble. Thin dashed lines are from individual models and pCO<sub>2</sub>-products. FESOM2.1-REcoM3 (magenta) shows the ocean carbon flux for the period of 1959-2021 whereas FESOM1.4-REcoM2 (yellow) covers the period of 1959-2019. Positive numbers indicate a flux into the ocean.

based estimates. In contrast, starting from the mid-1960s, FESOM2.1 shows a higher CO<sub>2</sub> flux in comparison to FESOM1.4. Considering the fact that both model versions do not depart much from each other in simulation A, the increase in net CO<sub>2</sub> flux is mostly attributed to the level of CO<sub>2</sub> fluxes in their control simulations (Fig. 18).

After accounting for the bias in simulation B, the simulated ocean carbon sink (1990-1999) is  $1.74 \pm 0.11$  PgC yr<sup>-1</sup> and  $2.17 \pm 0.13$  PgC yr<sup>-1</sup> for FESOM1.4-REcoM2 and FESOM2.1-REcoM3, respectively. Hence, FESOM2.1-REcoM3 is closer to the best estimate for the 1990s ( $2.2 \pm 0.4$  PgC yr<sup>-1</sup>, IPCC, based on seven different methodologies Denman et al., 2007; Ciais et al., 2014) than FESOM1.4-REcoM2. The cumulative uptake over the period of 1959-2019 amounts to 93.4 PgC (FESOM1.4-REcoM2) and 116.6 PgC (FESOM2.1-REcoM3) which is a 25% increase in CO<sub>2</sub> flux. Yet, the FESOM2.1-REcoM3 CO<sub>2</sub> fluxes are lower than the mean of the pCO<sub>2</sub>-based data-products since about 2008 and thus affirm the growing discrepancy between global ocean biogeochemistry models and pCO<sub>2</sub>-products (Friedlingstein et al., 2022a). It is likely that the models underestimate the mean ocean carbon uptake (Friedlingstein et al., 2022a), linked to biases in ventilation (Goris

550 et al., 2018; Terhaar et al., 2021, 2022; Bourgeois et al., 2022) and surface ocean buffer capacity (Vaittinada Ayar et al., 2022; Terhaar et al., 2022), and it is thus encouraging that FESOM2.1-REcoM3 has a comparatively high mean flux (Fig. 19). The pCO<sub>2</sub>-products are statistical models that inter- and extrapolate sparse pCO<sub>2</sub> observations and have substantial uncertainties themselves. In particular, they are sensitive to sparse and unevenly distributed observations (e.g., Gloege et al., 2021; Hauck et al., 2023) and it was shown that two of these methods overestimate the decadal CO<sub>2</sub> flux trend (2000-2018) by 20-35%  
555 based on the current pCO<sub>2</sub> observation distribution using a synthetic data set (Hauck et al., 2023).

### 3.3.3 DIC Inventory Changes

**Table 4.** FESOM2.1-REcoM3 DIC inventory for simulation A (in PgC) in 1994, and change in DIC inventory between 1800-1994 and 1994-2007 calculated from simulation A minus simulation B to account for model drift. Thus, the FESOM2.1-REcoM3 numbers encompass anthropogenic carbon cycle processes and the effect of climate change on the natural carbon cycle. Gruber et al. (2019) estimate the anthropogenic carbon inventory change, which is equivalent to simulation A minus simulation D (constant atmospheric CO<sub>2</sub>, variable climate). We have given the Gruber et al. (2019) anthropogenic plus back-of-the-envelope natural carbon inventory changes in parenthesis, which is roughly comparable to simulation A minus simulation B (only available for the global ocean).

	Year	Global [PgC]	North [PgC]	Tropics [PgC]	South [PgC]
<i>Total DIC inventory</i>					
FESOM2.1-REcoM3 (sim A)	1994	38167.4	5259.8	21108.1	11799.5
<i>DIC inventory change</i>					
FESOM2.1-REcoM3 (sim A minus sim B)	1800 to 1994	121	19.5	54.4	47.2
FESOM2.1-REcoM3 (sim A minus sim D)	1800 to 1994	119	22.3	52.5	44.3
Sabine et al. (2004), Gruber et al. (2019)	1800 to 1994	118±19 (111±21)	25.1	46.6	48.0
FESOM2.1-REcoM3 (sim A minus sim B)	1994 to 2007	29.9	5.4	12.6	12.0
FESOM2.1-REcoM3 (sim A minus sim D)	1994 to 2007	30.9	5.8	13.2	11.9
Gruber et al. (2019)	1994 to 2007	34±4 (29±5)	5.9	17.5	10.4

The interior ocean DIC inventory in FESOM2.1-REcoM3 amounts to about 38,200 PgC, which is in the reported range of 37,200±200 to 39,000 PgC (Sundquist, 1985; Keppler et al., 2020). The DIC inventory is thought to change primarily in response to the rise in atmospheric CO<sub>2</sub> and the resulting DIC inventory change is often referred to as anthropogenic carbon.  
560 Effects of climate change (warming, circulation changes) are thought to be one order of magnitude smaller, have the opposite sign, and affect both the natural carbon cycle and the anthropogenic carbon uptake and inventory (see Hauck et al. (2020); Friedlingstein et al. (2022a); Crisp et al. (2022) for more details on the different simulations and carbon components). The observation-based estimates of anthropogenic DIC inventory change use back-calculation techniques to separate anthropogenic carbon changes from the vast natural carbon reservoir. Thus, we here analyze the DIC inventory change in FESOM-REcoM




565 for simulation A minus B, which quantifies the total DIC inventory change while accounting for model drift. In addition, to derive the comparable DIC inventory change component as in observation-based studies, we make use of a third simulation (called simulation D), which is forced by interannual varying climate and preindustrial atmospheric CO<sub>2</sub>. Quantifying the DIC inventory change over a specific period from simulation A minus D is then coherent with the anthropogenic carbon definition used in Gruber et al. (2019).

570 The DIC inventory grew over time in accordance with observation-based estimates (Table 4, Sabine et al., 2004; Gruber et al., 2019). The simulated anthropogenic DIC inventory change 1800-1994 (119 PgC) is in good agreement with the observation-based anthropogenic DIC inventory change (118±19 PgC). For the total DIC inventory change, FESOM2.1-REcoM3 estimates a somewhat higher number (121 PgC, compared to 111±21 PgC), but is well within the reported uncertainty of the observation-based estimate. For the period 1994-2007, the total DIC increase is 29.9 PgC (simulation A minus simulation B, 575 i.e., drift-corrected) and thus slightly lower than the estimate by Gruber et al. (2019). However, Gruber et al. (2019) only quantifies the ocean anthropogenic DIC inventory increase (34±4 PgC) and neglects the counter effect by climate change. When considering the poorly constrained response of the natural carbon inventory to climate change, the model estimate falls within the uncertainty range of Gruber et al. (2019, 29±5 PgC). Alternatively, estimating the anthropogenic DIC inventory change from simulations A minus D leads to 30.9 PgC, which is within the uncertainty range of Gruber et al's 34±4 PgC. In GCB 580 2022, only four models simulated an anthropogenic DIC inventory change (1994-2007, simulation A minus D) ≥30 PgC, i.e. within the Gruber et al. uncertainty range. The other six models ranged between 25.5 and 28.3 PgC, and the model ensemble mean was 28.3±2.6 PgC (Friedlingstein et al., 2022a). FESOM2.1-REcoM3 is thus one of the few ocean biogeochemistry models that falls within the range of interior ocean anthropogenic carbon accumulation that is also supported by O<sub>2</sub>/N<sub>2</sub> ratios (Tohjima et al., 2019) and atmospheric inversions (see also discussion in Friedlingstein et al., 2022b). Notably, FESOM2.1- 585 REcoM3 reproduces the latitudinal distribution of anthropogenic carbon accumulation 1994-2007 with the maximum in the tropics (30°S-30°N), followed by the Southern Ocean (south of 30°S), and the North (north of 30°N). However, it also underestimates the accumulation in the tropics, as most other models do (Friedlingstein et al., 2022a). If the observation-based assessment of DIC inventory changes in North, Tropics and South is correct, this may indicate a transport of anthropogenic carbon from the Southern Ocean into the tropics that is too weak, or an air-sea CO<sub>2</sub> flux in the tropics, with too little ocean 590 uptake (or too much release) of CO<sub>2</sub>.

### 3.4 Oxygen

The simulated global O<sub>2</sub> concentration distribution at the surface ocean and intermediate depths was consistent with observed patterns in WOA2018 (Fig. 20, with  $r = 0.98/0.91$  and  $RMSE = 19.6/38.4$  mmol m<sup>3</sup> for surface (0-10 m) and intermediate depths (300-500 m) respectively). The model successfully reproduced the typical spatial patterns (Schmidtke et al., 2017): (1) Oxygen 595 Minimum Zones in the western boundary upwelling systems where old deoxygenated waters are brought to the surface, (2) high concentrations in the high latitude regions where cold temperature increases oxygen solubility (Arctic and Southern Oceans), and (3) moderate oxygen concentrations in the more stratified tropical gyres. Nevertheless, there were regional discrepancies. At the surface, the model slightly underestimated O<sub>2</sub> concentrations in the high latitude surface ocean. At intermediate depth,



**Figure 20.** Maps of surface (0-10m, top row) and intermediate (300-500m, bottom row) concentration of simulated FESOM2.1-REcoM3 (Sim A) dissolved  $O_2$  [ $mmol\ m^{-3}$ ] (A, D), World Ocean Atlas 2018 climatology of dissolved  $O_2$  (B, E; (Garcia et al., 2019b)) and corresponding differences (C, F) over the time period 2012-2021.

the model generally overestimated oxygen levels, especially in the Pacific Ocean and the sub-polar Southern Ocean with biases  
 600 exceeding  $100\ mmol\ m^{-3}$ . Within the 100-600 m layer, FESOM2.1-REcoM3 performed remarkably well with simulated  
 values of about  $160 \pm 105\ mmol\ m^{-3}$ , which is very close to the observations from the WOA2018 ( $158 \pm 103\ mmol\ m^{-3}$ ). A  
 previous model intercomparison study of oxygen concentrations within the 100-600 m layer (Cocco et al., 2013) showed that  
 such performances are common as all evaluated models fell within the error range of observations.

#### 4 Conclusions and Outlook

605 We have presented a new coupled ocean biogeochemistry model FESOM2.1-REcoM3. Building upon finite volumes for the  
 ocean component improves the numerical efficiency and leads to higher numerical throughput of the coupled model (Danilov  
 et al., 2017). Furthermore, the biogeochemistry component was extended to incorporate state of the art carbonate chemistry  
 routines, a second zooplankton and detritus group and simulates the cycling of oxygen in the ocean. In its present configuration,  
 the overall realism of FESOM2.1-REcoM3 in simulating the observed mean biogeochemical state is comparable to that of most  
 610 GOBMs, while being among the more realistic models for estimating global ocean anthropogenic carbon uptake. There are

still a number of model shortcomings, such as a lower simulated NPP and regional misfit between the annual mean CO<sub>2</sub> flux of the model simulation and the pCO<sub>2</sub>-based data-product that will be addressed in the future.


This model set-up provides the basis for further model development, e.g., the inclusion of coccolithophores as an additional phytoplankton functional type and the sensitivity of phytoplankton growth to rising CO<sub>2</sub> (Seifert et al., 2022), as well as the  
615 separation of the generic small zooplankton group into micro- and mesozooplankton that reduces model biases in nutrient fields, increases net primary production and better captures the top-down control on phytoplankton bloom phenology (Karakuş et al., 2022). We further plan to incorporate more detailed iron biogeochemistry as developed in REcoM coupled to MITgcm (e.g., Ye et al., 2020), and the explicit representation of the effects of viscosity and ballasting on particle sinking speed, as well as oxygen-dependent remineralization, following Cram et al. (2018) to address knowledge gaps in carbon export and transfer  
620 to depth (Henson et al., 2022). Other on-going work addresses the role of rivers for carbon and nutrient transport into the ocean and the remineralization time-scale of this river-derived organic material (Aumont et al., 2001; Lacroix et al., 2020; Regnier et al., 2022), and thus tackles a major uncertainty in the ocean carbon cycle and comparison of ocean carbon sink estimates based on pCO<sub>2</sub>-products and ocean biogeochemistry models (e.g., Hauck et al., 2020).

*Code availability.* The FESOM2.1-REcoM3 source code is available at [https://github.com/FESOM/fesom2/tree/fesom2.1\\_recom](https://github.com/FESOM/fesom2/tree/fesom2.1_recom) (last access: 31 December 2022); the version of FESOM2.1-REcoM3 used for this paper can be found at <https://doi.org/10.5281/zenodo.7502419>.  
625 A manual is available at: <https://recom.readthedocs.io/en/latest/>.


*Author contributions.* Conceptualization was done by OG, LO, CV and JH. Data were prepared by OG, JH, and OK. Analysis of simulations and visualization was done by all authors. All authors contributed to writing of the paper.

*Competing interests.* The authors declare no competing interests.

630 *Acknowledgements.* This research was funded by the Initiative and Networking Fund of the Helmholtz Association (Helmholtz Young Investigator Group Marine Carbon and Ecosystem Feedbacks in the Earth System [MarESys], grant number VH-NG-1301) and the ERC-2022-STG OceanPeak (grant agreement 101077209). LO and YY received funding from the European Union's Horizon 2020 research and innovation program under grant agreement number 820989 (project COMFORT). The work reflects only the authors' view; the European Commission and their executive agency are not responsible for any use that may be made of the information the work contains. LO acknowledges funding from the Germany Ministry for Education and Research (BMBF) project nuArctic (grant no. 03F0918A). MZ is funded by the  
635 Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Projektnummer 268020496 – TRR 172, within the Transregional Collaborative Research Center “Arctic Amplification: Climate Relevant Atmospheric and SurfaCe Processes, and Feedback Mechanisms (AC)<sup>3</sup>. YY and MB received funding by the German Federal Ministry of Education and Research (BMBF) through the project PalMod



**Figure A1.** Maps of simulated FESOM2.1-REcoM3 (Sim A) vertically integrated net primary production [ $\text{mgC m}^{-2} \text{d}^{-1}$ ] of small phytoplankton (A), diatoms (B), and the sum of both phytoplankton groups (C). The satellite-based Carbon-based Productivity Model (CbPM) is shown in (D; Westberry et al., 2008) with corresponding differences between FESOM2.1-REcoM3 and VGPM (E). All fields are averaged over the time period 2012 to 2021.



**Figure A2.** Conceptual diagram of the ocean biogeochemical model REcoM3. The 28 tracers can be grouped (indicated by boxes) into dissolved nutrients, carbonate system parameters and oxygen (upper left), phytoplankton functional types (center), zooplankton functional types (upper right), two detritus classes (lower right), and dissolved organic material (lower left). Source and sink terms are depicted by arrows. For reasons of diagrammatic clarity, connections of dissolved oxygen (Oxy) to other state variables are omitted here. Similarly, the release of alkalinity, dissolved inorganic nutrients and organic matter from the sediment are not shown.

## Appendix A: Equations

This appendix provides an overview of the underlying model equations and lists all biogeochemical variables of FESOM2.1-REcoM3. Changes in state variables in REcoM3 are controlled by biological and chemical processes, in addition to the changes induced by ocean circulation, mixing, diffusion and advection computed by FESOM2.1. While some variables exchange across the ocean surface and/or the sea floor, others, like dead organic matter (detritus) sink through the water column. The concentration change for a state variable  $S$  is formulated as follows:

$$\frac{\partial S}{\partial t} = -\mathbf{U} \cdot \nabla S + \nabla \cdot (\kappa \cdot \nabla S) + SMS(S), \quad (\text{A1})$$

where  $S$  is the volumetric concentration of a state variable,  $\mathbf{U}$  is the three-dimensional advection velocity and  $\kappa$  is the diffusivity. The term  $SMS(S)$  represents the biogeochemical sources minus sinks. The slow-sinking detritus class is assumed to sink with a velocity, which increases linearly with depth as a first-order description of the shift to larger and faster-sinking particles with increasing depth (Kriest and Oschlies, 2008). A constant sinking rate is applied to the fast-sinking detritus class. REcoM3 has 28 oceanic and four explicit benthic state variables (Tables A1 and A2).

### A1 Sources minus sinks

#### A1.1 Nutrients

##### 655 A1.1.1 Dissolved inorganic nitrate (DIN)

The simulated DIN conceptually represents the concentrations of nitrate, nitrite and ammonia, while in practice only nitrate is considered. The concentration of DIN in the water column rises when DON is remineralized and diminishes as a consequence of assimilation by small phytoplankton and diatoms:

$$SMS(\text{DIN}) = \underbrace{\rho_{\text{DON}} \cdot f_T \cdot \text{DON}}_{\text{DON remineralization}} - \underbrace{V_{\text{small}}^{\text{N}} \cdot \text{PhyC}_{\text{small}}}_{\text{N-assimilation, small phytoplankton}} - \underbrace{V_{\text{dia}}^{\text{N}} \cdot \text{PhyC}_{\text{dia}}}_{\text{N-assimilation, diatoms}}. \quad (\text{A2})$$

660 The state variables  $\text{DON}$ ,  $\text{PhyC}_{\text{small}}$  and  $\text{PhyC}_{\text{dia}}$  are listed in Table A1. The value of the remineralization rate constant ( $\rho_{\text{DON}}$ ) is given in Table A8. The temperature dependency of remineralization ( $f_T$ ) is calculated in Eq. A43. See Section A3.4 for details on the carbon-specific nitrogen-assimilation rates  $V_{\text{small}}^{\text{N}}$  and  $V_{\text{dia}}^{\text{N}}$  (Table A5).

##### A1.1.2 Dissolved silicic acid (DSi)

silicon assimilation (Si-assimilation) and increases when biogenic silica from one of the two detritus classes dissolves.

$$665 \quad SMS(\text{DSi}) = \underbrace{\rho_{\text{Si}}^T \cdot \text{DetSi}}_{\text{Remineralization, slow-sinking detritus}} + \underbrace{\rho_{\text{Si}}^T \cdot \text{DetZ2Si}}_{\text{Remineralization, fast-sinking detritus}} - \underbrace{V^{\text{Si}} \cdot \text{PhyC}_{\text{dia}}}_{\text{Si-assimilation, diatoms}} \quad (\text{A3})$$

The state variables  $\text{PhyC}_{\text{dia}}$ ,  $\text{DetSi}$  and  $\text{DetZ2Si}$  are listed in Table A1. The temperature dependent remineralization rate of silicon ( $\rho_{\text{Si}}^T$ ) and the carbon-specific Si-assimilation rate ( $V^{\text{Si}}$ ) are calculated in Eqs. A45 and A51, respectively (Table A5).

### A1.1.3 Dissolved iron (DFe)

Excretion of phyto- and zooplankton and remineralization of detritus release iron with a fixed iron:nitrate ratio ( $q^{\text{Fe:N}}$ ). Unlike  
 670 for nitrogen, which is released as dissolved organic nitrogen and needs to be remineralized further to become available as nu-  
 trient again, the released iron is directly put into the dissolved pool iron, basically assuming that all dissolved iron is ultimately  
 bio-available. Iron assimilation (again assumed to be proportional to nitrogen assimilation, from now on N-assimilation) by  
 both phytoplankton classes lower the level of dissolved iron. In addition, free inorganic iron  $\text{Fe}'$  is scavenged onto sinking  
 675 particles, with a rate that is proportional to particle concentration. We take detrital carbon as a proxy for the mass of sinking  
 particles.

$$\begin{aligned}
 \text{SMS}(\text{DFe}) = & q^{\text{Fe:N}} \cdot \left( \underbrace{\epsilon_{\text{phy}}^{\text{N}} \cdot f_{\text{lim, small}}^{\text{N:Cmax}} \cdot \text{PhyN}_{\text{small}}}_{\text{Excretion, small phytoplankton}} + \underbrace{\epsilon_{\text{phy}}^{\text{N}} \cdot f_{\text{lim, dia}}^{\text{N:Cmax}} \cdot \text{PhyN}_{\text{dia}}}_{\text{Excretion, diatoms}} \right) \\
 & + \underbrace{\rho_{\text{DetN}} \cdot f_T \cdot \text{DetN}}_{\text{Remineralization, slow-sinking detritus}} + \underbrace{\rho_{\text{DetN}} \cdot f_T \cdot \text{DetZ2N}}_{\text{Remineralization, fast-sinking detritus}} \\
 & + \underbrace{\epsilon_{\text{zoo}}^{\text{N}} \cdot \text{ZooN}}_{\text{Excretion, small zooplankton}} + \underbrace{\epsilon_{\text{zoo2}}^{\text{N}} \cdot \text{Zoo2N}}_{\text{Excretion, macrozooplankton}} \\
 & - \underbrace{V_{\text{small}}^{\text{N}} \cdot \text{PhyC}_{\text{small}}}_{\text{N-assimilation, small phytoplankton}} - \underbrace{V_{\text{dia}}^{\text{N}} \cdot \text{PhyC}_{\text{dia}}}_{\text{N-assimilation, diatom}} \\
 & - \underbrace{\kappa_{\text{Fe}} \cdot \text{DetC} \cdot \text{Fe}'}_{\text{Scavenging, slow-sinking detritus}} - \underbrace{\kappa_{\text{Fe}} \cdot \text{DetZ2C} \cdot \text{Fe}'}_{\text{Scavenging, fast-sinking detritus}}
 \end{aligned} \tag{A4}$$

The state variables  $\text{PhyC}_{\text{small}}$ ,  $\text{PhyC}_{\text{dia}}$ ,  $\text{PhyN}_{\text{small}}$ ,  $\text{PhyN}_{\text{dia}}$ ,  $\text{DetC}$ ,  $\text{DetN}$ ,  $\text{DetZ2C}$ ,  $\text{DetZ2N}$ ,  $\text{ZooN}$ ,  $\text{Zoo2N}$  are listed in  
 Table A1. Intracellular Fe:N ratio ( $q^{\text{Fe:N}}$ ) and scavenging rate of iron ( $\kappa_{\text{Fe}}$ ) are given in Table A4. Excretion rates ( $\epsilon_{\text{phy}}^{\text{N}}$ ,  
 $\epsilon_{\text{zoo}}^{\text{N}}$  and  $\epsilon_{\text{zoo2}}^{\text{N}}$ ) and the degradation rate for detritus N ( $\rho_{\text{DetN}}$ ) are listed in Table A8. The temperature dependency ( $f_T$ ) is  
 680 calculated in Eq. A43. The limitation by intracellular nitrogen ( $f_{\text{lim, small}}^{\text{N:Cmax}}$ ,  $f_{\text{lim, dia}}^{\text{N:Cmax}}$ ) is described in Eq. A55. Scavenging is  
 calculated following Parekh et al. (2004). The total concentration of dissolved iron ( $\text{Fe}_T$ ) is separated into free iron ( $\text{Fe}'$ ) and  
 iron complexed with organic ligands ( $\text{Fe}_L$ ), which is not scavenged. Complexation reactions are fast (Tagliabue and Völker,  
 2011), so we assume instantaneous equilibrium between free iron and free ligand ( $L'$ ) which is computed using a constant  
 $K_{\text{Fe}_L} = \frac{[\text{Fe}'] \cdot [L']}{[\text{Fe}_L]}$ , by solving

$$685 \quad \text{Fe}_T = \text{Fe}' + \text{Fe}_L \quad L_T = \text{Fe}_L + L' \tag{A5}$$

For simplicity we assume here a constant total ligand concentration  $L_T$ , unlike in Völker and Tagliabue (2015). Variable ligand  
 concentration, like in Misumi et al. (2011) or Völker and Tagliabue (2015), or variable ligand binding strength, like in Ye et al.  
 (2020) will be explored in the future. The values for  $K_{\text{Fe}_L}$  and  $L_T$  are listed in Table A4.

## A1.2 Carbon cycle

### 690 A1.2.1 Dissolved inorganic carbon (DIC)

DIC concentration increases with respiration of phyto- and zooplankton, remineralization of semi-labile dissolved organic carbon, dissolution of calcitic detritus and dissolution of  $\text{CaCO}_3$  in zooplankton guts. Loss terms are carbon fixation by primary producers and the formation of calcium carbonate. In addition, sea–air flux of  $\text{CO}_2$  leads to an exchange of carbon with the atmosphere, depending on the partial pressure difference of  $\text{CO}_2$  between ocean and atmosphere. This exchange is  
 695 treated separately as a boundary condition. The partial pressure of surface ocean  $\text{CO}_2$  is computed using the mocsy-2.0 routines (Orr and Epitalon, 2015).

$$\begin{aligned}
 \text{SMS(DIC)} = & \underbrace{(r_{\text{small}} - P_{\text{small}}) \cdot \text{PhyC}_{\text{small}}}_{\text{Net respiration, small phytoplankton}} + \underbrace{(r_{\text{dia}} - P_{\text{dia}}) \cdot \text{PhyC}_{\text{dia}}}_{\text{Net respiration, diatom}} + \underbrace{\rho_{\text{DOC}} \cdot f_T \cdot \text{DOC}}_{\text{Remineralization of DOC}} \\
 & + \underbrace{r_{\text{zoo}} \cdot \text{ZooC}}_{\text{Respiration, small zoo}} + \underbrace{r_{\text{zoo2}} \cdot \text{Zoo2C}}_{\text{Respiration, macrozoo}} + \underbrace{\text{Diss}_{\text{calc}} \cdot \text{DetCalc}}_{\text{Calcite dissolution, slow-sinking detritus}} + \underbrace{G_{\text{small}}^{\text{zoo}} \cdot q_{\text{small}}^{\text{CaCO}_3:\text{N}} \cdot \text{Diss}_{\text{calc\_guts}}}_{\text{CaCO}_3 \text{ dissolution in guts, small zoo}} \quad (\text{A6}) \\
 & - \underbrace{\psi \cdot P_{\text{small}} \cdot \text{PhyC}_{\text{small}}}_{\text{Calcification}} + \underbrace{\text{Diss}_{\text{calc2}} \cdot \text{DetZ2Calc}}_{\text{Calcite dissolution, fast-sinking detritus}} + \underbrace{G_{\text{small}}^{\text{zoo2}} \cdot q_{\text{small}}^{\text{CaCO}_3:\text{N}} \cdot \text{Diss}_{\text{calc\_guts}}}_{\text{CaCO}_3 \text{ dissolution in guts, macrozoo}}
 \end{aligned}$$

The state variables  $\text{PhyC}_{\text{small}}$ ,  $\text{PhyC}_{\text{dia}}$ ,  $\text{DOC}$ ,  $\text{ZooC}$ ,  $\text{Zoo2C}$ ,  $\text{DetCalc}$ ,  $\text{DetZ2Calc}$  are listed in Table A1. Respiration rate constants of small phytoplankton ( $r_{\text{small}}$ ), diatoms ( $r_{\text{dia}}$ ) and zooplankton groups ( $r_{\text{zoo}}$  and  $r_{\text{zoo2}}$ ) are computed in Sections A3.2  
 700 and A4.1, respectively. Photosynthesis terms ( $P_{\text{small}}$  and  $P_{\text{dia}}$ ) are calculated in Eq. A46. The remineralization rate constant ( $\rho_{\text{DOC}}$ ) is listed in Table A8 and the temperature dependency ( $f_T$ ) is given in Eq. A43. Calcite dissolution by detritus ( $\text{Diss}_{\text{calc}}$ ,  $\text{Diss}_{\text{calc2}}$ ) is calculated in Eq. A38. The constant for dissolution of calcium carbonate in zooplankton guts ( $\text{Diss}_{\text{calc\_guts}}$ ) is listed in Table A5.  $G_{\text{small}}^{\text{zoo}}$  and  $G_{\text{small}}^{\text{zoo2}}$  are grazing terms and explained in Section A4.2. The value of the calcite production ratio ( $\psi$ ) is given in Table A3.

### 705 A1.2.2 Total alkalinity (Alk)

The balance of alkalinity is affected by primary production, remineralization of dissolved organic matter, dissolution of calcitic detritus and dissolution of  $\text{CaCO}_3$  in zooplankton guts. Alkalinity increases when nitrogen is assimilated and when  $\text{CaCO}_3$  is dissolved (Wolf-Gladrow et al., 2007). Simultaneously, it is reduced by calcification as well as remineralization of dissolved organic nitrogen. The effect of phosphate assimilation and remineralization onto alkalinity is taken into account assuming a  
 710 constant N:P Redfield ratio (16:1).



$$\text{SMS}(\text{Alk}) = (1 + 1/16) \cdot \underbrace{V_{\text{small}}^{\text{N}} \cdot \text{PhyC}_{\text{small}}}_{\text{N-assimilation, small phytoplankton}} + (1 + 1/16) \cdot \underbrace{V_{\text{dia}}^{\text{N}} \cdot \text{PhyC}_{\text{dia}}}_{\text{N-assimilation, diatom}} \quad (\text{A7})$$

$$\begin{aligned} & - (1 + 1/16) \cdot \underbrace{\rho_{\text{DON}} \cdot f_T \cdot \text{DON}}_{\text{Remineralization of DON}} - 2 \cdot \underbrace{\psi \cdot P_{\text{small}} \cdot \text{PhyC}_{\text{small}}}_{\text{Calcification}} \\ & + 2 \cdot \underbrace{\text{Diss}_{\text{calc}} \cdot \text{DetCalc}}_{\text{Calcite dissolution, slow-sinking detritus}} + 2 \cdot \underbrace{G_{\text{small}}^{\text{zoo}} \cdot q_{\text{small}}^{\text{CaCO}_3:\text{N}} \cdot \text{Diss}_{\text{calc\_guts}}}_{\text{CaCO}_3 \text{ dissolution in guts, small zoo}} \\ & + 2 \cdot \underbrace{\text{Diss}_{\text{calc}2} \cdot \text{DetZ2Calc}}_{\text{Calcite dissolution, fast-sinking detritus}} + 2 \cdot \underbrace{G_{\text{small}}^{\text{zoo}2} \cdot q_{\text{small}}^{\text{CaCO}_3:\text{N}} \cdot \text{Diss}_{\text{calc\_guts}}}_{\text{CaCO}_3 \text{ dissolution in guts, macrozoo}} \end{aligned} \quad (\text{A8})$$

715 The state variables  $\text{PhyC}_{\text{small}}$ ,  $\text{PhyC}_{\text{dia}}$ ,  $\text{DON}$ ,  $\text{DetCalc}$ ,  $\text{DetZ2Calc}$  are listed in Table A1. The N-assimilation ( $V_{\text{small}}^{\text{N}}$  and  $V_{\text{dia}}^{\text{N}}$ ) is calculated in Section A3.4. The remineralization rate constant ( $\rho_{\text{DON}}$ ) is given in Table A8. The temperature dependency ( $f_T$ ) is calculated in Eq. A43. The value of the calcite production ratio ( $\psi$ ) is given in Table A3. The photosynthesis term ( $P_{\text{small}}$ ) is calculated in Eq. A46. The calcite dissolution by detritus ( $\text{Diss}_{\text{calc}}$ ,  $\text{Diss}_{\text{calc}2}$ ) is calculated in Eq. A38. Dissolution of calcium carbonate in guts ( $\text{Diss}_{\text{calc\_guts}}$ ) is listed in Table A5.  $G_{\text{small}}^{\text{zoo}}$  and  $G_{\text{small}}^{\text{zoo}2}$  are grazing terms and explained in Section A4.2.

## 720 A1.3 Phytoplankton

### A1.3.1 Nitrogen

The phytoplankton nitrogen pools increase through N-assimilation. The assimilation process is assumed to be proportional to carbon biomass, with a carbon-specific uptake rate that depends on the C:N ratio of phytoplankton and the external DIN concentration (Geider et al., 1998). Excretion of biogenic nitrogen to semi-labile DON drains the pool. At high intracellular C:N  
725 ratio, excretion is downregulated. Aggregation and grazing by the two zooplankton groups transfer nitrogen to the zooplankton and detritus pools.

$$\text{SMS}(\text{PhyN}_{\text{small}}) = \underbrace{V_{\text{small}}^{\text{N}} \cdot \text{PhyC}_{\text{small}}}_{\text{N-assimilation}} \quad (\text{A9})$$

$$- \underbrace{\epsilon_{\text{phy}}^{\text{N}} \cdot f_{\text{lim,small}}^{\text{N:Cmax}} \cdot \text{PhyN}_{\text{small}}}_{\text{DON excretion}} - \underbrace{\text{Agg} \cdot \text{PhyN}_{\text{small}}}_{\text{Aggregation loss}} - \underbrace{G_{\text{small}}^{\text{zoo}}}_{\text{Grazing loss by small zoo}} - \underbrace{G_{\text{small}}^{\text{zoo}2}}_{\text{Grazing loss by macrozoo}} \quad (\text{A10})$$

$$\text{SMS}(\text{PhyN}_{\text{dia}}) = \underbrace{V_{\text{dia}}^{\text{N}} \cdot \text{PhyC}_{\text{dia}}}_{\text{N-assimilation}} \quad (\text{A11})$$

$$- \underbrace{\epsilon_{\text{phy}}^{\text{N}} \cdot f_{\text{lim,dia}}^{\text{N:Cmax}} \cdot \text{PhyN}_{\text{dia}}}_{\text{DON excretion}} - \underbrace{\text{Agg} \cdot \text{PhyN}_{\text{dia}}}_{\text{Aggregation loss}} - \underbrace{G_{\text{dia}}^{\text{zoo}}}_{\text{Grazing loss by small zoo}} - \underbrace{G_{\text{dia}}^{\text{zoo}2}}_{\text{Grazing loss by macrozoo}} \quad (\text{A12})$$

The state variables  $\text{PhyC}_{\text{small}}$ ,  $\text{PhyN}_{\text{small}}$ ,  $\text{PhyC}_{\text{dia}}$ , and  $\text{PhyN}_{\text{dia}}$  are listed in Table A1. The N-assimilation ( $V_{\text{small}}^{\text{N}}$  and  $V_{\text{dia}}^{\text{N}}$ ) is explained in Section A3.4. The constant excretion rate constant ( $\epsilon_{\text{phy}}^{\text{N}}$ ) is given in Table A8. When the C:N ratio of the cells becomes too high, excretion of DON is downregulated by the limiter function ( $f_{\text{lim,small}}^{\text{N:Cmax}}$ ,  $f_{\text{lim,dia}}^{\text{N:Cmax}}$ ) that is described in Eq. A55.

Phytoplankton aggregation (Agg) defines the transfer of nitrogen into the detritus pools which depends quadratically on detritis  
 735 and phytoplankton concentrations (Eq. A52). Grazing loss terms ( $G_{small}^{zoo}$ ,  $G_{small}^{zoo2}$ ,  $G_{dia}^{zoo}$  and  $G_{dia}^{zoo2}$ ) are explained in Section A4.2.

### A1.3.2 Carbon

The carbon biomass of small phytoplankton and diatoms increases as a result of carbon assimilation during photosynthesis. Loss terms include excretion of DOC, which is limited by the availability of proteins as in the nitrogen pool, respiration, aggregation, and grazing.

$$740 \text{ SMS}(\text{PhyC}_{small}) = \underbrace{(P_{small} - r_{small}) \cdot \text{PhyC}_{small}}_{\text{Net photosynthesis}} \quad (\text{A13})$$

$$- \underbrace{\text{Agg} \cdot \text{PhyC}_{small}}_{\text{Aggregation loss}} - \underbrace{\epsilon_{phy}^C \cdot f_{lim,small}^{N:Cmax} \cdot \text{PhyC}_{small}}_{\text{Excretion of DOC}} - \underbrace{q_{small}^{C:N} \cdot G_{small}^{zoo}}_{\text{Grazing loss by small zoo}} - \underbrace{q_{small}^{C:N} \cdot G_{small}^{zoo2}}_{\text{Grazing loss by macrozoo}}$$

$$\text{SMS}(\text{PhyC}_{dia}) = \underbrace{(P_{dia} - r_{dia}) \cdot \text{PhyC}_{dia}}_{\text{Net photosynthesis}} \quad (\text{A14})$$

$$- \underbrace{\text{Agg} \cdot \text{PhyC}_{dia}}_{\text{Aggregation loss}} - \underbrace{\epsilon_{phy}^C \cdot f_{lim,dia}^{N:Cmax} \cdot \text{PhyC}_{dia}}_{\text{Excretion of DOC}} - \underbrace{q_{dia}^{C:N} \cdot G_{dia}^{zoo}}_{\text{Grazing loss by small zoo}} - \underbrace{q_{dia}^{C:N} \cdot G_{dia}^{zoo2}}_{\text{Grazing loss by macrozoo}}$$

The state variables  $\text{PhyC}_{small}$  and  $\text{PhyC}_{dia}$  are listed in Table A1. The photosynthesis terms ( $P_{small}$  and  $P_{dia}$ ) are calculated  
 745 in Eq. A46. Rates of respiration by small phytoplankton ( $r_{small}$ ), diatoms ( $r_{dia}$ ) are explained in Section A3.2. The constant for DOC excretion rate of phytoplankton ( $\epsilon_{phy}^C$ , Table A8) is downregulated by the limiter factor ( $f_{lim,small}^{N:Cmax}$ ,  $f_{lim,dia}^{N:Cmax}$ ) when the N:C ratio becomes too high (Eq. A55). Phytoplankton aggregation (Agg) is calculated in Eq. A52. Grazing terms ( $G_{small}^{zoo}$ ,  $G_{small}^{zoo2}$ ,  $G_{dia}^{zoo}$  and  $G_{dia}^{zoo2}$ ) are explained in Section A4.2.  $q^{C:N} = \text{PhyC}/\text{PhyN}$ , is used to convert the grazing units from mmol N to mmol C.

### 750 A1.3.3 CaCO<sub>3</sub>

The formation of biogenic calcium carbonate in our model is limited to coccolithophores only, which are assumed to form a constant fraction of the non-diatom phytoplankton. Formation of CaCO<sub>3</sub> by heterotrophs, such as foraminifera or pteropods is neglected. Biogenic CaCO<sub>3</sub> produced by coccolithophores is transformed into detritus CaCO<sub>3</sub> with all forms of organic carbon loss, i.e. organic matter excretion, respiration, aggregation and grazing. Calcifiers are assumed to comprise a certain  
 755 fraction of the total small phytoplankton concentration, specified by the parameter  $\psi$  (Table A3), tying the calcite production of calcifiers to the growth of small phytoplankton.

$$\text{SMS}(\text{PhyCalc}) = \underbrace{\psi \cdot P_{small} \cdot \text{PhyC}_{small}}_{\text{Calcification}} - \underbrace{r_{small} \cdot \text{PhyCalc}}_{\text{Respiration}} - \underbrace{G_{small}^{zoo} \cdot q_{small}^{CaCO_3:N}}_{\text{Grazing loss, small zoo}} - \underbrace{G_{small}^{zoo2} \cdot q_{small}^{CaCO_3:N}}_{\text{Grazing loss, macrozoo}} \quad (\text{A15})$$

$$- \underbrace{\epsilon_{phy}^C \cdot f_{lim,small}^{N:Cmax} \cdot \text{PhyCalc}}_{\text{Excretion loss}} - \underbrace{\text{Agg} \cdot \text{PhyCalc}}_{\text{Aggregation loss}}$$

The state variables  $\text{PhyC}_{\text{small}}$  and  $\text{PhyCalc}$  are listed in Table A1. The value of the calcite production ratio ( $\psi$ ) is given in Table A3. The constant excretion rate ( $\epsilon_{\text{phy}}^{\text{C}}$ , Table A8) is downregulated by the limiter factor  $f_{\text{lim,small}}^{\text{N:Cmax}}$  (Eq. A55) when the N:C ratio becomes too high. Photosynthesis ( $P_{\text{small}}$ ), respiration ( $r_{\text{small}}$ ) and the aggregation of phytoplankton (Agg) rates are calculated in Eqs. A46, A48 and A52, respectively. Grazing terms ( $G_{\text{small}}^{\text{zoo}}$  and  $G_{\text{small}}^{\text{zoo2}}$ ) are explained in Section A4.2.  $q_{\text{small}}^{\text{CaCO}_3:\text{N}} = \text{PhyCalc}/\text{PhyN}_{\text{small}}$  is used to convert the grazing units from mmol N to mmol  $\text{CaCO}_3$ .

### A1.3.4 Diatom silicon

The silica frustule of diatoms is built through Si-assimilation, which we assume to be carbon-specific, and regulated by cellular quotas (see below). Any decrease in N-biomass through excretion, grazing or aggregation leads to a corresponding transfer of silica to the detritus silica pool.

$$\begin{aligned} \text{SMS}(\text{PhySi}) = & \underbrace{V^{\text{Si}} \cdot \text{PhyC}_{\text{dia}}}_{\text{Diatom Si-assimilation}} \\ & - \underbrace{\epsilon_{\text{phy}}^{\text{N}} \cdot f_{\text{lim,dia}}^{\text{N:Cmax}} \cdot \text{PhySi}_{\text{dia}}}_{\text{Excretion to detritus}} - \underbrace{\text{Agg} \cdot \text{PhySi}_{\text{dia}}}_{\text{Aggregation loss}} - \underbrace{G_{\text{dia}}^{\text{zoo}} \cdot q^{\text{Si:N}}}_{\text{Grazing loss, small zoo}} - \underbrace{G_{\text{dia}}^{\text{zoo2}} \cdot q^{\text{Si:N}}}_{\text{Grazing loss, macrozoo}} \end{aligned} \quad (\text{A16})$$

The state variables  $\text{PhyC}_{\text{dia}}$  and  $\text{PhySi}_{\text{dia}}$  are described in Table A1. Si-assimilation ( $V^{\text{Si}}$ ) and aggregation rates (Agg) are calculated in Eqs. A51 and A52, respectively. The constant excretion rate ( $\epsilon_{\text{phy}}^{\text{N}}$ , Table A8) is downregulated by the limiter factor  $f_{\text{lim,dia}}^{\text{N:Cmax}}$  (Eq. A55) when the N:C ratio becomes too high. Grazing terms ( $G_{\text{dia}}^{\text{zoo}}$  and  $G_{\text{dia}}^{\text{zoo2}}$ ) are explained in Section A4.2. The intracellular ratio between diatom silicon and nitrate is defined as  $q^{\text{Si:N}} = \text{PhySi}_{\text{dia}}/\text{PhyN}_{\text{dia}}$ .

### A1.3.5 Chlorophyll *a*

Chlorophyll *a* synthesis is structured as a function of irradiance and of N-assimilation, following Geider et al. (1998). Chlorophyll *a* is degraded at a light-dependent rate (See Álvarez et al. (2018)), and lost via aggregation and grazing. The grazing losses in terms of nitrogen biomass are converted to chlorophyll loss using the intracellular Chl:N ratio.

$$\text{SMS}(\text{PhyChl}_{\text{small}}) = \underbrace{S_{\text{small}}^{\text{chl}} \cdot \text{PhyC}_{\text{small}}}_{\text{Chlorophyll } a \text{ synthesis}} \quad (\text{A17})$$

$$- \underbrace{G_{\text{small}}^{\text{zoo}} \cdot q_{\text{small}}^{\text{Chl:N}}}_{\text{Grazing loss, small zoo}} - \underbrace{G_{\text{small}}^{\text{zoo2}} \cdot q_{\text{small}}^{\text{Chl:N}}}_{\text{Grazing loss, macrozoo}} - \underbrace{\text{deg}_{\text{small}}^{\text{chl}} \cdot \text{PhyChl}_{\text{small}}}_{\text{Degradation loss}} - \underbrace{\text{Agg} \cdot \text{PhyChl}_{\text{small}}}_{\text{Aggregation loss}} \quad (\text{A18})$$

$$\text{SMS}(\text{PhyChl}_{\text{dia}}) = \underbrace{S_{\text{dia}}^{\text{chl}} \cdot \text{PhyC}_{\text{dia}}}_{\text{Chlorophyll } a \text{ synthesis}} \quad (\text{A19})$$

$$- \underbrace{G_{\text{dia}}^{\text{zoo}} \cdot q_{\text{dia}}^{\text{Chl:N}}}_{\text{Grazing loss, small zoo}} - \underbrace{G_{\text{dia}}^{\text{zoo2}} \cdot q_{\text{dia}}^{\text{Chl:N}}}_{\text{Grazing loss, macrozoo}} - \underbrace{\text{deg}_{\text{dia}}^{\text{chl}} \cdot \text{PhyChl}_{\text{dia}}}_{\text{Degradation loss}} - \underbrace{\text{Agg} \cdot \text{PhyChl}_{\text{dia}}}_{\text{Aggregation loss}} \quad (\text{A20})$$

The state variables  $\text{PhyC}_{\text{small}}$ ,  $\text{PhyC}_{\text{dia}}$ ,  $\text{PhyChl}_{\text{small}}$  and  $\text{PhyChl}_{\text{dia}}$  are listed in Table A1. The chlorophyll *a* synthesis ( $S_{\text{small}}^{\text{chl}}$ ,  $S_{\text{dia}}^{\text{chl}}$ ) and the aggregation (Agg) terms are calculated in Eqs. A49 and A52, respectively. The degradation parameters ( $\text{deg}_{\text{small}}^{\text{chl}}$ ,  $\text{deg}_{\text{dia}}^{\text{chl}}$ )

deg<sub>dia</sub><sup>chl</sup>) are given in Table A8. Grazing terms ( $G_{\text{small}}^{\text{zoo}}$ ,  $G_{\text{small}}^{\text{zoo2}}$ ,  $G_{\text{dia}}^{\text{zoo}}$ , and  $G_{\text{dia}}^{\text{zoo2}}$ ) are explained in Section A4.2. The conversion factor from mmol N to mg Chl *a* is defined as  $q^{\text{Chl:N}} = \text{PhyChl}/\text{PhyN}$ .

## 785 A1.4 Zooplankton

### A1.4.1 Nitrogen

Both zooplankton classes increase their nitrogen biomass via grazing on phytoplankton and detritus while mortality and excretion of DON reduce it. Macrozooplankton further feeds on small zooplankton and releases nitrogen via fecal pellet production.

$$\text{SMS}(\text{ZooN}) = \underbrace{\gamma_{\text{zoo}} \cdot G_{\text{tot}}^{\text{zoo}}}_{\text{Grazing}} - \underbrace{G_{\text{zoo}}}_{\text{Grazing loss, macrozoo}} - \underbrace{m_{\text{zoo}} \cdot \text{ZooN}^2}_{\text{Mortality}} - \underbrace{\epsilon_{\text{zoo}}^{\text{N}} \cdot \text{ZooN}}_{\text{Excretion of DON}} \quad (\text{A21})$$

$$790 \quad \text{SMS}(\text{Zoo2N}) = \underbrace{\gamma_{\text{zoo2}} \cdot G_{\text{tot}}^{\text{zoo2}}}_{\text{Grazing}} - \underbrace{m_{\text{zoo2}} \cdot \text{Zoo2N}^2}_{\text{Mortality}} - \underbrace{\epsilon_{\text{zoo2}}^{\text{N}} \cdot \text{Zoo2N}}_{\text{Excretion of DON}} - \underbrace{f_{\text{n}} \cdot G_{\text{tot}}^{\text{zoo2}}}_{\text{Fecal pellet}} \quad (\text{A22})$$

The state variables ZooN and Zoo2N are listed in Table A1. Only a fraction of the grazed phytoplankton ( $\gamma_{\text{zoo}}$ ,  $\gamma_{\text{zoo2}}$ , Table A3) enters the zooplankton biomass. The rest is transferred to detritus due to sloppy feeding. The grazing terms ( $G_{\text{tot}}^{\text{zoo}}$ ,  $G_{\text{tot}}^{\text{zoo2}}$ ) are calculated in Section A4.2. The mortality parameter ( $m_{\text{zoo}}$ ,  $m_{\text{zoo2}}$ ) and fecal pellet production rate constant ( $f_{\text{n}}$ ) are listed in Table A3. The DON excretion terms ( $\epsilon_{\text{zoo}}^{\text{N}}$ ,  $\epsilon_{\text{zoo2}}^{\text{N}}$ ) are given in Table A8.

### 795 A1.4.2 Carbon

The zooplankton carbon biomass increases with carbon uptake via grazing and decreases through carbon losses through mortality, respiration and carbon excretion to the semi-labile DOC pool. Macrozooplankton further gains carbon by grazing on small zooplankton and loses it via fecal pellet production.

$$\text{SMS}(\text{ZooC}) = \underbrace{\gamma_{\text{zoo}} \cdot (G_{\text{small}}^{\text{zoo}} \cdot q_{\text{small}}^{\text{C:N}} + G_{\text{dia}}^{\text{zoo}} \cdot q_{\text{dia}}^{\text{C:N}})}_{\text{Grazing on phytoplankton}} + \underbrace{\gamma_{\text{zoo}} \cdot (G_{\text{det}}^{\text{zoo}} \cdot q_{\text{det}}^{\text{C:N}} + G_{\text{detZ2}}^{\text{zoo}} \cdot q_{\text{detZ2}}^{\text{C:N}})}_{\text{Grazing on detritus}} - \underbrace{G_{\text{zoo}} \cdot q_{\text{zoo}}^{\text{C:N}}}_{\text{Grazing loss by macrozoo}} \quad (\text{A23})$$

$$800 \quad - \underbrace{m_{\text{zoo}} \cdot \text{ZooN}^2 \cdot q_{\text{zoo}}^{\text{C:N}}}_{\text{Zooplankton mortality}} - \underbrace{r_{\text{zoo}} \cdot \text{ZooC}}_{\text{Respiration loss}} - \underbrace{\epsilon_{\text{zoo}}^{\text{C}} \cdot \text{ZooC}}_{\text{Excretion of DOC}} \quad (\text{A24})$$

$$\text{SMS}(\text{Zoo2C}) = \underbrace{\gamma_{\text{zoo2}} \cdot (G_{\text{small}}^{\text{zoo2}} \cdot q_{\text{small}}^{\text{C:N}} + G_{\text{dia}}^{\text{zoo2}} \cdot q_{\text{dia}}^{\text{C:N}})}_{\text{Grazing on phytoplankton}} + \underbrace{\gamma_{\text{zoo2}} \cdot (G_{\text{det}}^{\text{zoo2}} \cdot q_{\text{det}}^{\text{C:N}} + G_{\text{detZ2}}^{\text{zoo2}} \cdot q_{\text{detZ2}}^{\text{C:N}})}_{\text{Grazing on detritus}} + \underbrace{\gamma_{\text{zoo2}} \cdot (G_{\text{zoo}} \cdot q_{\text{zoo}}^{\text{C:N}})}_{\text{Grazing on small zoo}} \quad (\text{A25})$$

$$- \underbrace{m_{\text{zoo2}} \cdot \text{Zoo2N}^2 \cdot q_{\text{zoo2}}^{\text{C:N}}}_{\text{Zooplankton mortality}} - \underbrace{r_{\text{zoo2}} \cdot \text{Zoo2C}}_{\text{Respiration loss}} - \underbrace{\epsilon_{\text{zoo2}}^{\text{C}} \cdot \text{Zoo2C}}_{\text{Excretion of DOC}} - \underbrace{f_{\text{c}} \cdot G_{\text{cflux}}}_{\text{Fecal pellet}} \quad (\text{A26})$$

The state variables ZooN, ZooC, Zoo2N and Zoo2C are listed in Table A1. A fraction of the grazed phytoplankton ( $\gamma_{\text{zoo}}$  and  $\gamma_{\text{zoo2}}$ , Table A3) is kept in the zooplankton biomass while the remainder is returned back to detritus pool as a consequence of sloppy feeding. Grazing terms ( $G_{\text{small}}^{\text{zoo}}$ ,  $G_{\text{dia}}^{\text{zoo}}$ ,  $G_{\text{det}}^{\text{zoo}}$ ,  $G_{\text{detZ2}}^{\text{zoo}}$ ,  $G_{\text{small}}^{\text{zoo2}}$ ,  $G_{\text{dia}}^{\text{zoo2}}$ ,  $G_{\text{det}}^{\text{zoo2}}$ ,  $G_{\text{detZ2}}^{\text{zoo2}}$  and  $G_{\text{zoo}}$ ) are calculated in Section A4.2. The respiration terms of zooplankton ( $r_{\text{zoo}}$  and  $r_{\text{zoo2}}$ ) are calculated in Eqs. A60 and A61. Mortality parameters ( $m_{\text{zoo}}$ ,  $m_{\text{zoo2}}$ ) are listed in Table A3. The DOC excretion terms ( $\epsilon_{\text{zoo}}^{\text{C}}$ ,  $\epsilon_{\text{zoo2}}^{\text{C}}$ ) are in Table A8. The grazing flux in terms of

nitrogen biomass is converted to carbon biomass using the respective intracellular C:N ratios ( $q_{\text{small}}^{\text{C:N}}$ ,  $q_{\text{dia}}^{\text{C:N}}$ ,  $q_{\text{det}}^{\text{C:N}}$ ,  $q_{\text{detZ2}}^{\text{C:N}}$ ,  $q_{\text{zoo}}^{\text{C:N}}$  and  $q_{\text{zoo2}}^{\text{C:N}}$ ) where,  $q_{\text{small}}^{\text{C:N}} = \text{PhyC}_{\text{small}}/\text{PhyN}_{\text{small}}$ ,  $q_{\text{dia}}^{\text{C:N}} = \text{PhyC}_{\text{dia}}/\text{PhyN}_{\text{dia}}$ ,  $q_{\text{det}}^{\text{C:N}} = \text{DetC}/\text{DetN}$ ,  $q_{\text{detZ2}}^{\text{C:N}} = \text{DetZ2C}/\text{DetZ2N}$ ,  $q_{\text{zoo}}^{\text{C:N}} = \text{ZooC}/\text{ZooN}$  and  $q_{\text{zoo2}}^{\text{C:N}} = \text{Zoo2C}/\text{Zoo2N}$ . Total grazed carbon biomass ( $G_{\text{flux}}$ ) and the fecal pellet production rate constant ( $f_c$ , Table A3) together determine the fraction of carbon being lost to the large detritus carbon pool via fecal pellets.

## A1.5 Detritus

### A1.5.1 Nitrogen

Detrital nitrogen pool increases as a result of sloppy feeding and mortality. Sloppy feeding is outlined as a function of grazing fluxes and grazing efficiency of macrozooplankton. In other words, the grazed phytoplankton partly goes to the macrozooplankton biomass depending on the grazing efficiency. The phytoplankton aggregation contributes only to slow-sinking detritus. Fecal pellet production is defined only for macrozooplankton group. Detritus is degraded to DON based on temperature and a remineralisation rate.

$$\begin{aligned} \text{SMS}(\text{DetN}) = & \underbrace{(G_{\text{small}}^{\text{zoo}} + G_{\text{dia}}^{\text{zoo}}) \cdot (1 - \gamma_{\text{zoo}})}_{\text{Sloppy feeding}} + \underbrace{m_{\text{zoo}} \cdot \text{ZooN}^2}_{\text{Zooplankton mortality}} - \underbrace{\gamma_{\text{zoo}} \cdot (G_{\text{det}}^{\text{zoo}} + G_{\text{detZ2}}^{\text{zoo}})}_{\text{Grazing loss, small zoo}} \\ & + \underbrace{\text{Agg} \cdot (\text{PhyN}_{\text{small}} + \text{PhyN}_{\text{dia}})}_{\text{Phytoplankton aggregation}} - \underbrace{\rho_{\text{DetN}} \cdot f_T \cdot \text{DetN}}_{\text{Degradation to DON}} \end{aligned} \quad (\text{A27})$$

$$\text{SMS}(\text{DetZ2N}) = \underbrace{(G_{\text{small}}^{\text{zoo2}} + G_{\text{dia}}^{\text{zoo2}} + G_{\text{zoo}}) \cdot (1 - \gamma_{\text{zoo2}})}_{\text{Sloppy feeding}} - \underbrace{\gamma_{\text{zoo2}} \cdot (G_{\text{det}}^{\text{zoo2}} + G_{\text{detZ2}}^{\text{zoo2}})}_{\text{Grazing loss, macrozoo}} \quad (\text{A28})$$

$$+ \underbrace{m_{\text{zoo2}} \cdot \text{Zoo2N}^2}_{\text{Mortality}} + \underbrace{f_n \cdot G_{\text{tot}}}_{\text{Fecal pellet}} - \underbrace{\rho_{\text{DetN}} \cdot f_T \cdot \text{DetZ2N}}_{\text{Degradation to DON}} \quad (\text{A29})$$

The state variables  $\text{PhyN}_{\text{small}}$ ,  $\text{PhyN}_{\text{dia}}$ ,  $\text{ZooN}$ ,  $\text{DetN}$ ,  $\text{Zoo2N}$  and  $\text{DetZ2N}$  are listed in Table A1. The grazing efficiency ( $\gamma_{\text{zoo}}$  and  $\gamma_{\text{zoo2}}$ ), mortality ( $m_{\text{zoo}}$ ,  $m_{\text{zoo2}}$ ) and fecal pellet production rate constant ( $f_n$ ) are listed in Table A3. Grazing terms ( $G_{\text{small}}^{\text{zoo}}$ ,  $G_{\text{dia}}^{\text{zoo}}$ ,  $G_{\text{det}}^{\text{zoo}}$ ,  $G_{\text{detZ2}}^{\text{zoo}}$ ,  $G_{\text{small}}^{\text{zoo2}}$ ,  $G_{\text{dia}}^{\text{zoo2}}$ ,  $G_{\text{det}}^{\text{zoo2}}$ ,  $G_{\text{detZ2}}^{\text{zoo2}}$  and  $G_{\text{zoo}}$ ) are calculated in Section A4.2. The remineralisation rate constant of DON ( $\rho_{\text{DetN}}$ ) is listed in Table A8. The temperature dependency  $f_T$  is calculated in Eq. A43. The aggregation (Agg) term is calculated in Eq. A52.

### A1.5.2 Carbon

Detrital carbon sources are associated with sloppy feeding, aggregation of phytoplankton, mortality of small zooplankton and fecal pellet production by macrozooplankton. Degradation of DetC and DetZ2C to DOC is the only loss term.

$$\text{SMS}(\text{DetC}) = \underbrace{(G_{\text{small}}^{\text{zoo}} \cdot q_{\text{small}}^{\text{C:N}} + G_{\text{dia}}^{\text{zoo}} \cdot q_{\text{dia}}^{\text{C:N}})}_{\text{Sloppy feeding}} \cdot (1 - \gamma_{\text{zoo}}) + \underbrace{m_{\text{zoo}} \cdot \text{ZooN}^2 \cdot q_{\text{zoo}}^{\text{C:N}}}_{\text{small zoo mortality}} - \underbrace{\gamma_{\text{zoo}} \cdot (G_{\text{det}}^{\text{zoo}} \cdot q_{\text{det}}^{\text{C:N}} + G_{\text{detZ2}}^{\text{zoo}} \cdot q_{\text{detZ2}}^{\text{C:N}})}_{\text{Grazing loss by macro zoo}} \quad (\text{A30})$$

$$+ \underbrace{\text{Agg} \cdot (\text{PhyC}_{\text{small}} + \text{PhyC}_{\text{dia}})}_{\text{Phytoplankton aggregation}} - \underbrace{\rho_{\text{DetC}} \cdot f_T \cdot \text{DetC}}_{\text{Degradation to DOC}}$$

$$\text{SMS}(\text{DetZ2C}) = \underbrace{(G_{\text{small}}^{\text{zoo2}} \cdot q_{\text{small}}^{\text{C:N}} + G_{\text{dia}}^{\text{zoo2}} \cdot q_{\text{dia}}^{\text{C:N}} + G_{\text{zoo}} \cdot q_{\text{zoo}}^{\text{C:N}})}_{\text{Sloppy feeding}} \cdot (1 - \gamma_{\text{zoo2}}) - \underbrace{\gamma_{\text{zoo2}} \cdot (G_{\text{det}}^{\text{zoo2}} \cdot q_{\text{det}}^{\text{C:N}} + G_{\text{detZ2}}^{\text{zoo2}} \cdot q_{\text{detZ2}}^{\text{C:N}})}_{\text{Grazing loss by macro zoo}} \quad (\text{A31})$$

$$+ \underbrace{m_{\text{zoo2}} \cdot \text{Zoo2N}^2 \cdot q_{\text{zoo2}}^{\text{C:N}}}_{\text{Mortality}} + \underbrace{f_c \cdot G_{\text{flux}}}_{\text{Fecal pellet}} - \underbrace{\rho_{\text{DetC}} \cdot f_T \cdot \text{DetZ2C}}_{\text{Degradation to DOC}} \quad (\text{A32})$$

835 The state variables  $\text{PhyC}_{\text{small}}$ ,  $\text{PhyC}_{\text{dia}}$ ,  $\text{ZooN}$ ,  $\text{DetC}$ ,  $\text{Zoo2N}$  and  $\text{DetZ2C}$  are listed in Table A1. The grazing efficiency ( $\gamma_{\text{zoo}}$  and  $\gamma_{\text{zoo2}}$ ) and mortality ( $m_{\text{zoo}}$   $m_{\text{zoo2}}$ ) parameters are listed in Table A3. Grazing terms ( $G_{\text{small}}^{\text{zoo}}$ ,  $G_{\text{dia}}^{\text{zoo}}$ ,  $G_{\text{det}}^{\text{zoo}}$ ,  $G_{\text{detZ2}}^{\text{zoo}}$ ,  $G_{\text{small}}^{\text{zoo2}}$ ,  $G_{\text{dia}}^{\text{zoo2}}$ ,  $G_{\text{det}}^{\text{zoo2}}$  and  $G_{\text{zoo}}$ ) are calculated in Section A4.2. The remineralisation rate of DOC ( $\rho_{\text{DetC}}$ ) is listed in Table A8. Temperature dependency  $f_T$  is calculated in Eq. A43. The aggregation (Agg) term is calculated in Eq. A52. Total grazed carbon biomass ( $G_{\text{flux}}$ ) and the fecal pellet production rate constant ( $f_c$ , Table A3) together determine the fraction of carbon  
840 being lost to the large detritus carbon pool via fecal pellets. The quotas  $q_{\text{small}}^{\text{C:N}} = \text{PhyC}_{\text{small}}/\text{PhyN}_{\text{small}}$ ,  $q_{\text{dia}}^{\text{C:N}} = \text{PhyC}_{\text{dia}}/\text{PhyN}_{\text{dia}}$ ,  $q_{\text{zoo}}^{\text{C:N}} = \text{ZooC}/\text{ZooN}$ ,  $q_{\text{zoo2}}^{\text{C:N}} = \text{Zoo2C}/\text{Zoo2N}$ ,  $q_{\text{det}}^{\text{C:N}} = \text{DetC}/\text{DetN}$  and  $q_{\text{detZ2}}^{\text{C:N}} = \text{DetZ2C}/\text{DetZ2N}$  are used to convert the units from mmol N to mmol C.

### A1.5.3 Silica

Biogenic detrital silica increases with excretion fluxes from diatoms to detritus, aggregation and grazing and decreases with  
845 silica dissolution from  $\text{DetSi}$  and  $\text{DetZ2Si}$ .

$$\text{SMS}(\text{DetSi}) = \underbrace{(\epsilon_{\text{phy}}^{\text{N}} \cdot f_{\text{lim,dia}}^{\text{N:Cmax}})}_{\text{Diatom excretion}} + \underbrace{\text{Agg}}_{\text{Aggregation}} \cdot \text{DiaSi} + \underbrace{G_{\text{dia}}^{\text{zoo}} \cdot q_{\text{dia}}^{\text{Si:N}}}_{\text{Sloppy feeding}} - \underbrace{\rho_{\text{Si}}^{\text{T}} \cdot \text{DetSi}}_{\text{Remineralization to DSi}} \quad (\text{A33})$$

$$\text{SMS}(\text{DetZ2Si}) = \underbrace{G_{\text{dia}}^{\text{zoo2}} \cdot q_{\text{dia}}^{\text{Si:N}}}_{\text{Sloppy feeding}} - \underbrace{\rho_{\text{Si}}^{\text{T}} \cdot \text{DetZ2Si}}_{\text{Remineralization to DSi}} \quad (\text{A34})$$

The state variables  $\text{DiaSi}$ ,  $\text{DetSi}$  and  $\text{DetZ2Si}$  are listed in Table A1. The constant excretion rate ( $\epsilon_{\text{phy}}^{\text{N}}$ , Table A8) is down-regulated by the limiter factor  $f_{\text{lim,dia}}^{\text{N:Cmax}}$  (Eq. A55) when the N:C ratio becomes too high. The remineralization rates ( $\rho_{\text{Si}}^{\text{T}}$ ),  
850 the aggregation (Agg) and the grazing on diatoms ( $G_{\text{dia}}^{\text{zoo}}$ ,  $G_{\text{dia}}^{\text{zoo2}}$ ) are calculated in Eqs. A45, A52 and A65, respectively. The intracellular ratio between diatom silicon and carbon is defined as  $q_{\text{dia}}^{\text{Si:N}} = \text{PhySi}_{\text{dia}}/\text{PhyN}_{\text{dia}}$ .

### A1.5.4 $\text{CaCO}_3$

The coccolithophore fraction of small phytoplankton loses biogenic  $\text{CaCO}_3$  to the detrital  $\text{CaCO}_3$  pool along with excretion, aggregation, respiration and grazing. Dissolution of  $\text{CaCO}_3$  leads to an increase in DIC and alkalinity.

$$855 \quad \text{SMS}(\text{DetCalc}) = \underbrace{\epsilon_C^{\text{phy}} \cdot f_{\text{lim,small}}^{\text{N:Cmax}} \cdot \text{PhyCalc}}_{\text{Small phytoplankton, excretion}} + \left( \underbrace{\text{Agg}}_{\text{Aggregation}} + \underbrace{r_{\text{small}}}_{\text{Respiration}} \right) \cdot \text{PhyCalc} + \underbrace{G_{\text{small}}^{\text{zoo}} \cdot q_{\text{small}}^{\text{CaCO}_3:\text{N}}}_{\text{Grazing loss}} \quad (\text{A35})$$

$$- \underbrace{G_{\text{small}}^{\text{zoo}} \cdot q_{\text{small}}^{\text{CaCO}_3:\text{N}} \cdot \text{Diss}_{\text{calc-guts}}}_{\text{CaCO}_3 \text{ dissolution in guts}} - \underbrace{\text{Diss}_{\text{calc}} \cdot \text{DetCalc}}_{\text{CaCO}_3 \text{ dissolution, slow-sinking detritus}}$$

$$\text{SMS}(\text{DetZ2Calc}) = \underbrace{G_{\text{small}}^{\text{zoo2}} \cdot q_{\text{small}}^{\text{CaCO}_3:\text{N}}}_{\text{Grazing loss}} \quad (\text{A36})$$

$$- \underbrace{G_{\text{small}}^{\text{zoo2}} \cdot q_{\text{small}}^{\text{CaCO}_3:\text{N}} \cdot \text{Diss}_{\text{calc-guts}}}_{\text{CaCO}_3 \text{ dissolution in guts}} - \underbrace{\text{Diss}_{\text{calc2}} \cdot \text{DetZ2Calc}}_{\text{CaCO}_3 \text{ dissolution, fast-sinking detritus}} \quad (\text{A37})$$

The state variables PhyCalc, DetCalc and DetZ2Calc are listed in Table A1. The constant excretion rate ( $\epsilon_C^{\text{phy}}$ , Table A8) is downregulated by the limiter factor  $f_{\text{lim,small}}^{\text{N:Cmax}}$  (Eq. A55) when the N:C ratio becomes too high. The respiration ( $r_{\text{small}}$ ), the aggregation (Agg) and the grazing on small pytoplankton ( $G_{\text{small}}^{\text{zoo}}$  and  $G_{\text{small}}^{\text{zoo2}}$ ) are calculated in Eqs. A48, A52 and A64, respectively. The ratio  $q_{\text{small}}^{\text{CaCO}_3:\text{N}} = \text{PhyCalc}/\text{PhyN}_{\text{small}}$ .

**Calcite dissolution:** As the detritus calcite sinks through the water column it is subject to dissolution. We follow Yamanaka and Tajika (1996) assuming an exponential decrease of the  $\text{CaCO}_3$  flux with depth. As we also assume an increasing sinking speed of small detritus with depth, following Kriest and Oschlies (2008), the dissolution rate is scaled with the sinking velocity.

$$\text{Diss}_{\text{calc}} = \text{Diss}_{\text{calc\_rate}} \cdot w_{\text{det}} \quad \text{Diss}_{\text{calc2}} = \text{Diss}_{\text{calc\_rate}} \quad (\text{A38})$$

$\text{Diss}_{\text{calc}}$  and  $\text{Diss}_{\text{calc2}}$  are the dissolution rate constants for slow- and fast-sinking detritus classes (Table A5). The reference dissolution rate ( $\text{Diss}_{\text{calc\_rate}}$ , Table A8) is based on a length scale of 3500 m and velocity of  $20 \text{ m d}^{-1}$ . The sinking speed at depth  $z$  ( $w_{\text{det}}$ , Table A5) is calculated as follows:

$$870 \quad w_{\text{det}} = 0.0288 \cdot z + w_0 \quad (\text{A39})$$

Here,  $z$  denotes the depth and  $w_0$  is the sinking speed at the ocean surface (Table A3). The dissolution rate for fast-sinking detritus class ( $\text{Diss}_{\text{calc2}}$ ) is assumed to be constant throughout the water column and is set to the value of  $\text{Diss}_{\text{calc\_rate}}$  (Table A8).

## A1.6 Dissolved oxygen (Oxy)

Oxy concentration increases with carbon fixation by primary producers. It decreases with respiration of phyto- and zooplank- tons, remineralization of dissolved organic carbon. In addition, sea-air flux of  $\text{O}_2$  leads to an exchange of oxygen with the atmosphere, depending on the partial pressure difference of  $\text{O}_2$  between ocean and atmosphere. This exchange is treated separately as a boundary condition. The partial pressure of surface ocean  $\text{O}_2$  is computed using the mocsy-2.0 routines (Orr and

Epitalon, 2015).

$$\begin{aligned}
\text{SMS}(\text{Oxy}) = & \underbrace{(P_{\text{small}} - r_{\text{small}}) \cdot \text{PhyC}_{\text{small}}}_{\text{Net production, small phytoplankton}} + \underbrace{(P_{\text{dia}} - r_{\text{dia}}) \cdot \text{PhyC}_{\text{dia}}}_{\text{Net production, diatom}} \\
& - \underbrace{\rho_{\text{DOC}} \cdot f_T \cdot \text{DOC}}_{\text{Remineralization of DOC}} - \underbrace{r_{\text{zoo}} \cdot \text{ZooC}}_{\text{Respiration, small zoo}} - \underbrace{r_{\text{zoo2}} \cdot \text{Zoo2C}}_{\text{Respiration, macrozoo}}
\end{aligned} \tag{A40}$$

880 The state variables  $\text{PhyC}_{\text{small}}$ ,  $\text{PhyC}_{\text{dia}}$ ,  $\text{DOC}$ ,  $\text{ZooC}$  and  $\text{Zoo2C}$  are listed in Table A1. Respiration rate constants of small phytoplankton ( $r_{\text{small}}$ ), diatoms ( $r_{\text{dia}}$ ) and zooplankton groups ( $r_{\text{zoo}}$  and  $r_{\text{zoo2}}$ ) are computed in Sections A3.2 and A4.1, respectively. Photosynthesis terms ( $P_{\text{small}}$  and  $P_{\text{dia}}$ ) are calculated in Eq. A46. The remineralization rate constant  $\rho_{\text{DOC}}$  is listed in Table A8 and the temperature dependency ( $f_T$ ) is given in Eq. A43.

### A1.7 Dissolved organic material

885 Dissolved organic matter in our model is a representation of the semi-labile fraction only, the refractory and labile fractions are not included.

#### A1.7.1 Dissolved organic nitrogen (DON)

DON is produced via nitrogen excretion by phytoplankton, zooplankton and by degradation of detrital nitrogen. DON is turned into DIN by remineralization which is the only sink term.

$$\begin{aligned}
890 \text{ SMS}(\text{DON}) = & \underbrace{\epsilon_{\text{phy}}^{\text{N}} \cdot f_{\text{lim,small}}^{\text{N:Cmax}} \cdot \text{PhyN}_{\text{small}}}_{\text{Excretion, small phytoplankton}} + \underbrace{\epsilon_{\text{dia}}^{\text{N}} \cdot f_{\text{lim,dia}}^{\text{N:Cmax}} \cdot \text{PhyN}_{\text{dia}}}_{\text{Excretion, diatom}} \\
& + \underbrace{\epsilon_{\text{zoo}}^{\text{N}} \cdot \text{ZooN}}_{\text{Excretion, small zoo}} + \underbrace{\epsilon_{\text{zoo2}}^{\text{N}} \cdot \text{Zoo2N}}_{\text{Excretion, macrozoo}} + \underbrace{\rho_{\text{DetN}} \cdot f_T \cdot \text{DetN}}_{\text{Detritus degradation, slow-sinking}} + \underbrace{\rho_{\text{Det2ZN}} \cdot f_T \cdot \text{Det2ZN}}_{\text{Detritus degradation, fast sinking}} - \underbrace{\rho_{\text{DON}} \cdot f_T \cdot \text{DON}}_{\text{Remineralization}}
\end{aligned} \tag{A41}$$

The state variables  $\text{PhyN}_{\text{small}}$ ,  $\text{PhyN}_{\text{dia}}$ ,  $\text{ZooN}$ ,  $\text{DetN}$ ,  $\text{Zoo2N}$ ,  $\text{Det2ZN}$  and  $\text{DON}$  are listed in Table A1. The constant excretion rate of nitrogen from phytoplankton and zooplankton classes ( $\epsilon_{\text{phy}}^{\text{N}}$ ,  $\epsilon_{\text{dia}}^{\text{N}}$ ,  $\epsilon_{\text{zoo}}^{\text{N}}$  and  $\epsilon_{\text{zoo2}}^{\text{N}}$ ), the degradation rate of detritus ( $\rho_{\text{DetN}}$ ,  $\rho_{\text{Det2ZN}}$ ) and the remineralization rate of DON ( $\rho_{\text{DON}}$ ) are listed in Table A8. The constant excretion rate of phytoplankton  
895 is downregulated by the limiter function ( $f_{\text{lim,small}}^{\text{N:Cmax}}$ ,  $f_{\text{lim,dia}}^{\text{N:Cmax}}$ , Eq. A55) when the N : C ratio becomes too high. The temperature dependency  $f_T$  is calculated in Eq. A43.

#### A1.7.2 Dissolved organic carbon (DOC)

DOC is produced via carbon excretion by phytoplankton and zooplankton and by degradation of detrital carbon. DOC is turned into DIC by remineralization which is the only sink term.

$$\begin{aligned}
900 \text{ SMS}(\text{DOC}) = & \underbrace{\epsilon_{\text{phy}}^{\text{C}} \cdot f_{\text{lim,small}}^{\text{N:Cmax}} \cdot \text{PhyC}_{\text{small}}}_{\text{Excretion, small phytoplankton}} + \underbrace{\epsilon_{\text{dia}}^{\text{C}} \cdot f_{\text{lim,dia}}^{\text{N:Cmax}} \cdot \text{PhyC}_{\text{dia}}}_{\text{Excretion, diatom}} \\
& + \underbrace{\epsilon_{\text{zoo}}^{\text{C}} \cdot \text{ZooC}}_{\text{Excretion, small zoo}} + \underbrace{\epsilon_{\text{zoo2}}^{\text{C}} \cdot \text{Zoo2C}}_{\text{Excretion, macrozoo}} + \underbrace{\rho_{\text{DetC}} \cdot f_T \cdot \text{DetC}}_{\text{Detritus degradation, slow sinking}} + \underbrace{\rho_{\text{Det2C}} \cdot f_T \cdot \text{Det2C}}_{\text{Detritus degradation, fast sinking}} - \underbrace{\rho_{\text{DOC}} \cdot f_T \cdot \text{DOC}}_{\text{Remineralization}}
\end{aligned} \tag{A42}$$



The state variables PhyC<sub>small</sub>, PhyC<sub>dia</sub>, ZooC, DetC, Zoo2C, Det2C and DOC are listed in Table A1. The constant excretion rate of nitrogen from phytoplankton and zooplankton classes ( $\epsilon_{\text{phy}}^{\text{C}}$ ,  $\epsilon_{\text{dia}}^{\text{C}}$ ,  $\epsilon_{\text{zoo}}^{\text{C}}$  and  $\epsilon_{\text{zoo2}}^{\text{C}}$ ), the degradation rate of detritus ( $\rho_{\text{DetC}}$ ,  $\rho_{\text{Det2C}}$ ) and the remineralization rate of DOC ( $\rho_{\text{DOC}}$ ) are listed in Table A8. The constant excretion rate of phytoplankton is downregulated by the limiter factor ( $f_{\text{lim,small}}^{\text{N:Cmax}}$ ,  $f_{\text{lim,dia}}^{\text{N:Cmax}}$ , Eq. A55) when the N:C ratio becomes too high. Temperature dependency  $f_T$  is calculated in Eq. A43.

## A2 Temperature dependence of rates

**Arrhenius function:** Most metabolic processes are faster at higher temperatures. This temperature dependence is defined relative to a reference temperature.

$$f_T = \exp\left(-4500 \cdot \left(\frac{1}{T} - \frac{1}{T_{\text{ref}}}\right)\right) \quad (\text{A43})$$

$T$  and  $T_{\text{ref}}$  are the local and reference temperature in K, respectively (Table A6).

**Macrozooplankton grazing:** Macrozooplankton grazing is temperature dependent. A dimensionless exponential temperature function (Butzin and Pörtner, 2016) is used for the parameterization of the temperature dependency ( $f_{T_{\text{zoo2}}}$ , Table A5). Specifically, the following parameterization provides an optimum curve with a maximum at 0.5°C as described in Karakuş et al. (2021).

$$f_{T_{\text{zoo2}}} = \frac{\exp\left(\frac{Q_a}{T_r} - \frac{Q_a}{T}\right)}{1 + \exp\left(\frac{Q_h}{T_h} - \frac{Q_h}{T}\right)} \quad (\text{A44})$$

$T_r$  is the intrinsic optimum temperature for development and  $T_h$  is the temperature above which inhibitive processes dominate.  $Q_a$  and  $Q_h$  are the temperatures for the uninhibited and inhibited reaction kinetics, respectively (Table A9).  $T$  is the local temperature in K.

**Silicon dissolution:** The temperature dependent dissolution rate of silicon ( $\rho_{\text{Si}}^T$ , Table A5) is calculated following Maerz et al. (2020), but with a minimum dissolution rate.

$$\rho_{\text{Si}}^T = \max\left(0.023 \cdot 2.6^{\frac{T-10}{10}}, \rho_{\text{Si}}\right) \quad (\text{A45})$$

$T$  is the local temperature in °C. The minimum dissolution rate ( $\rho_{\text{Si}}$ ) is listed in Table A8.

## A3 Phytoplankton processes

Phytoplankton growth equations are based on Geider et al. (1998) with small modifications for diatom silicon uptake, following Hohn (2009).

### A3.1 Photosynthesis

The rate of the carbon specific (C-specific from now on) photosynthesis for phytoplankton ( $P_{\text{small}}, P_{\text{dia}}$ ) is parameterized as follows:

$$930 \quad P_{\text{small}} = P_{\text{max}}^{\text{small}} \cdot \left( 1.0 - \exp \left( \frac{-\alpha_{\text{small}} \cdot q^{\text{Chl:C}} \cdot \text{PAR}}{P_{\text{max}}^{\text{small}}} \right) \right), \quad P_{\text{dia}} = P_{\text{max}}^{\text{dia}} \cdot \left( 1.0 - \exp \left( \frac{-\alpha_{\text{dia}} \cdot q^{\text{Chl:C}} \cdot \text{PAR}}{P_{\text{max}}^{\text{dia}}} \right) \right) \quad (\text{A46})$$

The light harvesting efficiency ( $\alpha_{\text{small}}, \alpha_{\text{dia}}$ ) per chlorophyll is listed in Table A7. PAR is the photosynthetically available radiation (Table A5). The intracellular Chl to C ratio ( $q^{\text{Chl:C}}$ ) is defined as PhyChl/PhyC and varies as a result of photoacclimation. The apparent maximum photosynthetic rate ( $P_{\text{max}}^{\text{small}}, P_{\text{max}}^{\text{dia}}$ ) is defined below.

$$P_{\text{max}}^{\text{small}} = \mu_{\text{C,small}}^{\text{max}} \cdot \min (f_{\text{lim,small}}^{\text{Fe}}, f_{\text{lim,small}}^{\text{N:Cmin}}) \cdot f_T, \quad P_{\text{max}}^{\text{dia}} = \mu_{\text{C,dia}}^{\text{max}} \cdot \min (f_{\text{lim,dia}}^{\text{Fe}}, f_{\text{lim,dia}}^{\text{N:Cmin}}, f_{\text{lim,dia}}^{\text{Si:Cmin}}) \cdot f_T \quad (\text{A47})$$

935 The value of  $\mu_{\text{C,small}}^{\text{max}}, \mu_{\text{C,dia}}^{\text{max}}$  is listed in Table A7. The limitation terms ( $f_{\text{lim,small}}^{\text{N:Cmin}}, f_{\text{lim,dia}}^{\text{N:Cmin}}, f_{\text{lim,dia}}^{\text{Si:Cmin}}, f_{\text{lim,small}}^{\text{Fe}}$  and  $f_{\text{lim,dia}}^{\text{Fe}}$ ) are presented in Section A3.6 and the temperature dependency ( $f_T$ ) is calculated in Eq. A43.

### A3.2 Respiration

The phytoplankton respiration rate ( $r_{\text{small}}$  and  $r_{\text{dia}}$ , Table A5) is calculated as a base respiration plus a second term proportional to N-assimilation, as a measure of biosynthesis:

$$940 \quad r_{\text{small}} = \underbrace{\text{res}_{\text{small}} \cdot f_{\text{lim,small}}^{\text{N:Cmax}}}_{\text{Maintenance}} + \underbrace{\zeta \cdot V_{\text{small}}^{\text{N}}}_{\text{N-assim}}, \quad r_{\text{dia}} = \underbrace{\text{res}_{\text{dia}} \cdot f_{\text{lim,dia}}^{\text{N:Cmax}}}_{\text{Maintenance}} + \underbrace{\zeta \cdot V_{\text{dia}}^{\text{N}}}_{\text{N-assim}} \quad (\text{A48})$$

The values for the maintenance respiration rate ( $\text{res}_{\text{small}}, \text{res}_{\text{dia}}$ ) and the cost of biosynthesis ( $\zeta$ ) are listed in Table A7. Si-assimilation is assumed to be inexpensive, so it is not included as additional cost in the respiration (Hohn, 2009). The limiter function ( $f_{\text{lim,small}}^{\text{N:Cmax}}$  and  $f_{\text{lim,dia}}^{\text{N:Cmax}}$ ) is described in Eq. A55 and the N-assimilation rate ( $V_{\text{small}}^{\text{N}}, V_{\text{dia}}^{\text{N}}$ ) is calculated in Eq. A50.

### A3.3 Chlorophyll *a* synthesis

945 The chlorophyll synthesis rate ( $S_{\text{small}}^{\text{chl}}, S_{\text{dia}}^{\text{chl}}$ , Table A5) is proportional to N-assimilation, with the proportionality factor varying as a function of the C-specific photosynthesis rate, relative to the maximum possible photosynthetic rate at the current Chl:C ratio of the cell, which depends on photosynthetically available radiation and light harvesting efficiency.

$$S_{\text{small}}^{\text{chl}} = V_{\text{small}}^{\text{N}} \cdot q_{\text{max,small}}^{\text{Chl:N}} \cdot \min \left( 1, \frac{P_{\text{small}}}{\alpha_{\text{small}} \cdot q^{\text{Chl:C}} \cdot \text{PAR}} \right), \quad S_{\text{dia}}^{\text{chl}} = V_{\text{dia}}^{\text{N}} \cdot q_{\text{max,dia}}^{\text{Chl:N}} \cdot \min \left( 1, \frac{P_{\text{dia}}}{\alpha_{\text{dia}} \cdot q^{\text{Chl:C}} \cdot \text{PAR}} \right) \quad (\text{A49})$$

950 The N-assimilation ( $V_{\text{small}}^{\text{N}}, V_{\text{dia}}^{\text{N}}$ ) is computed in Eq. A50. The conversion factor of the maximum Chl : N ratio ( $q_{\text{max,small}}^{\text{Chl:N}}, q_{\text{max,dia}}^{\text{Chl:N}}$ ) and the light harvesting efficiency ( $\alpha_{\text{small}}, \alpha_{\text{dia}}$ ) are listed in Table A7. The C-specific photosynthesis ( $P_{\text{small}}, P_{\text{dia}}$ ) is given in Eq. A46. PAR is the photosynthetically available radiation (Table A5) and the intracellular Chl to C ratio ( $q^{\text{Chl:C}}$ ) is defined as PhyChl/PhyC.

### A3.4 N- and Si-assimilation

**Nitrogen:** The C-specific N-assimilation rate is a function of the maximum rate of C-specific photosynthesis and DIN concentration. N-assimilation depends on the DIN concentration in seawater via Michaelis–Menten kinetics. The N : C uptake ratio and a function of the intracellular quota between N and C further, which downregulates uptake under high N:C ratio further modify the N-assimilation.

$$V_{\text{small}}^{\text{N}} = V_{\text{cm}}^{\text{small}} \cdot P_{\text{max}}^{\text{small}} \cdot \sigma_{\text{N:C}}^{\text{small}} \cdot f_{\text{lim,small}}^{\text{N:Cmax}} \cdot \frac{\text{DIN}}{K_{\text{small}}^{\text{N}} + \text{DIN}}, \quad V_{\text{dia}}^{\text{N}} = V_{\text{cm}}^{\text{dia}} \cdot P_{\text{max}}^{\text{dia}} \cdot \sigma_{\text{N:C}}^{\text{dia}} \cdot f_{\text{lim,dia}}^{\text{N:Cmax}} \cdot \frac{\text{DIN}}{K_{\text{dia}}^{\text{N}} + \text{DIN}} \quad (\text{A50})$$

$V_{\text{cm}}^{\text{small}}, V_{\text{cm}}^{\text{dia}}, \sigma_{\text{N:C}}^{\text{small}}, \sigma_{\text{N:C}}^{\text{dia}}, K_{\text{small}}^{\text{N}}$  and  $K_{\text{dia}}^{\text{N}}$  are listed in Table A7. The maximum rate of photosynthesis ( $P_{\text{max}}^{\text{small}}$  and  $P_{\text{max}}^{\text{dia}}$ ) is given in Eqs. A47.  $f_{\text{lim,small}}^{\text{N:Cmax}}$  and  $f_{\text{lim,dia}}^{\text{N:Cmax}}$  are described in Eq. A55. DIN corresponds to insitu concentration.

**Silicon:** The building of a silica frustule of diatoms requires silicate uptake. The C-specific Si-assimilation rate is a function of a factor for C-specific N-uptake, a rate constant of C-specific photosynthesis, maximum uptake ratio N : C for small phytoplankton and DSi concentration. The maximum Si : C ratio, temperature, and the scaling factor for the maximum nitrogen uptake further regulate the N-assimilation.

$$V_{\text{Si}}^{\text{Si}} = V_{\text{cm}}^{\text{dia}} \cdot \mu_{\text{C,dia}}^{\text{max}} \cdot f_T \cdot \sigma_{\text{Si:C}} \cdot f_{\text{lim}}^{\text{Si:Cmax}} \cdot f_{\text{lim,dia}}^{\text{N:Cmax}} \cdot \frac{\text{DSi}}{K_{\text{Si}} + \text{DSi}} \quad (\text{A51})$$

The scaling factor for the N-uptake ( $V_{\text{cm}}^{\text{dia}}$ ), the maximum Rate constant of C-specific photosynthesis ( $\mu_{\text{C,dia}}^{\text{max}}$ ), the uptake ratio of the maximum Si : C ( $\sigma_{\text{Si:C}}$ ) and half-saturation constant for silicate uptake ( $K_{\text{Si}}$ ) are listed in Table A7. The temperature dependency ( $f_T$ ) is computed in Eq. A43. The limitation by the intracellular ratios N : C and Si : C ( $f_{\text{lim,dia}}^{\text{N:Cmax}}, f_{\text{lim}}^{\text{Si:Cmax}}$ ) are described in Eqs. A55 and A56, respectively. DSi corresponds to in situ concentration.

### 970 A3.5 Aggregation loss

The aggregation rate (Agg, Table A5) is proportional to the concentration of small phytoplankton, diatoms and detritus. The effect of increased stickiness of diatoms under nutrient limitation (Waite et al., 1992; Aumont et al., 2015) is taken into account by multiplying the diatom biomass with  $(1 - q_{\text{lim}}^{\text{dia}})$ . When the nutrient limitation is high (i.e, low  $q_{\text{lim}}^{\text{dia}}$ ), the aggregation rate increases in the model.

$$\text{Agg} = \phi_{\text{phy}} \cdot (\text{PhyN}_{\text{small}} + (1 - q_{\text{lim}}^{\text{dia}}) \cdot \text{PhyN}_{\text{dia}}) + \phi_{\text{det}} \cdot (\text{DetN} + \text{DetZ2N}) \quad (\text{A52})$$

$$q_{\text{lim}}^{\text{dia}} = \min(f_{\text{lim,dia}}^{\text{Fe}}, f_{\text{lim,dia}}^{\text{N:Cmin}}, f_{\text{lim,dia}}^{\text{Si:Cmin}}) \quad (\text{A53})$$

The state variables  $\text{PhyN}_{\text{small}}, \text{PhyN}_{\text{dia}}, \text{DetN}$  and  $\text{DetZ2N}$  are described in Table A1. The values of the maximum aggregation loss parameters ( $\phi_{\text{phy}}$  and  $\phi_{\text{det}}$ ) are listed in Table A3. The limitation terms ( $f_{\text{lim,dia}}^{\text{N:Cmin}}, f_{\text{lim,dia}}^{\text{Si:Cmin}}$  and  $f_{\text{lim,dia}}^{\text{Fe}}$ ) are presented below (Section A3.6).

### A3.6 Nutrient limitation

The metabolic processes such as C-specific photosynthesis, respiration rate and excretion losses are treated as functions of the intracellular nitrogen status (i.e., N : C ratios  $q$ ) following Geider et al. (1998). Intracellular ratios between nutrients and carbon limit uptake of nitrogen and silicon which is modeled via a non-linear function as in Schourup-Kristensen et al. (2014).

$$985 \quad f_{\text{lim}}(\theta, q_1, q_2) = 1 - \exp(-\theta(|\Delta q| - \Delta q)^2) \quad (\text{A54})$$

Here,  $\Delta q = q_1 - q_2$  is the difference between the current intracellular nutrient:C quota and a prescribed maximum or minimum quota. The dimensionless constant  $\theta$  controls the limitation.

#### A3.6.1 $f_{\text{lim}}^{\text{N:Cmax}}$

The limiter  $f_{\text{lim}}^{\text{N:Cmax}}$  downregulates the metabolic processes such as nitrogen and Si-assimilation, excretion and maintenance  
990 respiration of phytoplankton when the intracellular nitrogen quota ( $q^{\text{N:C}}$ ) becomes too high.  $f_{\text{lim}}^{\text{N:Cmax}}$  is one when the current  $q^{\text{N:C}} < 0.151$  (i.e., Redfield ratio, 16N:106C) and zero for  $q^{\text{N:C}} > 0.2$  (i.e., 21.2N:106C). It determines the end of the uptake of nitrogen and silicon in assimilation processes as well as the cease of carbon and nitrogen release during the respiration and excretion of DON/DOC and  $\text{CaCO}_3$  processes of phytoplankton (See Section A1.5.4).

$$f_{\text{lim,small}}^{\text{N:Cmax}} = f_{\text{lim}}(\theta_{\text{max}}^{\text{N}}, q_{\text{small}}^{\text{N:C}}, q_{\text{small}}^{\text{N:Cmax}}), \quad f_{\text{lim,dia}}^{\text{N:Cmax}} = f_{\text{lim}}(\theta_{\text{max}}^{\text{N}}, q_{\text{dia}}^{\text{N:C}}, q_{\text{dia}}^{\text{N:Cmax}}) \quad (\text{A55})$$

995 The limitation function for quota regulation is calculated with Eq. A54.  $q_{\text{small}}^{\text{N:C}}$  and  $q_{\text{dia}}^{\text{N:C}}$  are the current intracellular nitrogen quota for small phytoplankton and diatoms, respectively. Dimensionless constants  $\theta_{\text{max}}^{\text{N}}$ ,  $q_{\text{small}}^{\text{N:Cmax}}$  and  $q_{\text{dia}}^{\text{N:Cmax}}$  are listed in Table A6.

#### A3.6.2 $f_{\text{lim}}^{\text{Si:Cmax}}$

The limiter  $f_{\text{lim}}^{\text{Si:Cmax}}$  downregulates the Si-assimilation of diatoms when the intracellular silicon quota (Si : C) becomes too  
1000 high.  $f_{\text{lim}}^{\text{Si:Cmax}}$  is one when the current  $q^{\text{Si:C}} < 0.76$  and zero for  $q^{\text{Si:C}} > 0.8$ . It determines the end of the uptake of silicon in assimilation processes. The limiter function is described in Eq. A54 and is calculated as follows:

$$f_{\text{lim}}^{\text{Si:Cmax}} = f_{\text{lim}}(\theta_{\text{max}}^{\text{Si}}, q^{\text{Si:C}}, q^{\text{Si:Cmax}}) \quad (\text{A56})$$

Dimensionless constants  $\theta_{\text{max}}^{\text{Si}}$  and  $q^{\text{Si:Cmax}}$  are listed in Table A6.

#### A3.6.3 $f_{\text{lim}}^{\text{Si:Cmin}}$

1005 Carbon fixation and aggregation loss in diatoms are further downregulated by a factor ( $f_{\text{lim,dia}}^{\text{Si:Cmin}}$ , see Eq. A54) when the intracellular silicon quota ( $q^{\text{Si:C}}$ ) approaches a minimum value ( $q^{\text{Si:Cmin}}$ ), mimicking the arrest of cellular division at low cellular Si (Claquin et al., 2002).  $f_{\text{lim,dia}}^{\text{Si:Cmin}}$  is zero when the current  $q^{\text{Si:C}} < 0.04$  and one for  $q^{\text{Si:C}} > 0.08$ .

$$f_{\text{lim,dia}}^{\text{Si:Cmin}} = f_{\text{lim}}(\theta_{\text{min}}^{\text{Si}}, q^{\text{Si:Cmin}}, q^{\text{Si:C}}) \quad (\text{A57})$$

Dimensionless constants  $\theta_{\text{min}}^{\text{Si}}$  and  $q^{\text{Si:Cmin}}$  are listed in Table A6.

1010 **A3.6.4**  $f_{\text{lim}}^{\text{Fe}}$

Growth-limitation by iron is modeled with Michaelis–Menten kinetics, implicitly assuming that all dissolved iron is ultimately bioavailable.

$$f_{\text{lim,small}}^{\text{Fe}} = \frac{\text{DFe}}{K_{\text{small}}^{\text{Fe}} + \text{DFe}}, \quad f_{\text{lim,dia}}^{\text{Fe}} = \frac{\text{DFe}}{K_{\text{dia}}^{\text{Fe}} + \text{DFe}} \quad (\text{A58})$$

Stat variable DFe is listed in Table A1. The half saturation constants ( $K_{\text{small}}^{\text{Fe}}$  and  $K_{\text{dia}}^{\text{Fe}}$ ) are given in Table A6.

1015 **A3.6.5**  $f_{\text{lim}}^{\text{N:Cmin}}$

In addition to iron limitation, photosynthesis is limited by nitrogen in small phytoplankton and diatoms using the Eq. A54. Nitrogen limitation ( $f_{\text{lim,small}}^{\text{N:Cmin}}$ ,  $f_{\text{lim,dia}}^{\text{N:Cmin}}$ ) is described as a function of the intracellular nitrogen quota ( $q_{\text{small}}^{\text{N:C}}$ ,  $q_{\text{dia}}^{\text{N:C}}$ ) with growth ending at a minimum quota ( $q_{\text{small}}^{\text{N:Cmin}}$ ,  $q_{\text{dia}}^{\text{N:Cmin}}$ ).

$$f_{\text{lim,small}}^{\text{N:Cmin}} = f_{\text{lim}}(\theta_{\text{min}}^{\text{N}}, q_{\text{small}}^{\text{N:Cmin}}, q_{\text{small}}^{\text{N:C}}), \quad f_{\text{lim,dia}}^{\text{N:Cmin}} = f_{\text{lim}}(\theta_{\text{min}}^{\text{N}}, q_{\text{dia}}^{\text{N:Cmin}}, q_{\text{dia}}^{\text{N:C}}) \quad (\text{A59})$$

1020 Dimensionless constants  $\theta_{\text{min}}^{\text{N}}$ ,  $q_{\text{small}}^{\text{N:Cmin}}$  and  $q_{\text{dia}}^{\text{N:Cmin}}$  are listed in Table A6.

## A4 Zooplankton processes

### A4.1 Zooplankton respiration

Small zooplankton: When the intracellular C:N ratio in zooplankton exceeds the Redfield ratio, a temperature dependent respiration ( $r_{\text{zoo}}$ , Table A5) is assumed to drive it back with a time scale  $\tau$ .

$$1025 \quad r_{\text{zoo}} = \frac{q_{\text{zoo}}^{\text{C:N}} - q_{\text{standard}}^{\text{C:N}}}{\tau} \cdot f_T \quad (\text{A60})$$

The time scale for respiration ( $\tau$ ) is listed in Table A7. The temperature dependence ( $f_T$ ) is calculated in Eq. (A43). The ratios are defined as  $q_{\text{zoo}}^{\text{C:N}} = \text{ZooC}/\text{ZooN}$  and  $q_{\text{standard}}^{\text{C:N}} = 106\text{C}/16\text{N}$ .

**Macrozooplankton:** The daily respiration rate constant of macrozooplankton ( $r_{\text{zoo}2}$ , Table A5) is modeled following Karakuş et al. (2021).

$$1030 \quad r_{\text{zoo}2} = R_s \cdot (1 + R_f + R_a) \quad (\text{A61})$$

The standard respiration rate ( $R_s$ ) is listed in Table A3. The feeding activity factor ( $R_f$ , Table A5) is defined as the ratio of grazing flux to carbon biomass of macrozooplankton which increases linearly from 0 to 1 for ratio between 0% and 10% and is 1 otherwise. The respiration activity factor ( $R_a$ , Table A5) defines reduced macrozooplankton respiration rate in austral/boreal winter with the value of  $-0.5$ .

## 1035 A4.2 Grazing

In REcoM3, there are two zooplankton classes, small zooplankton (< 2cm) and macrozooplankton (2-20 cm). The small zooplankton group grazes on small phytoplankton and diatoms as well as on fast- and slow-sinking detrital particles. While macrozooplankton grazes on similarly both phytoplankton classes and detritus groups, it further grazes on small zooplankton. Total grazing of both zooplankton groups is based on the Holling type III ingestion function as follows:

$$1040 \quad G_{\text{tot}}^{\text{zoo}} = \xi_{\text{zoo}} \cdot \frac{(\sum_i p_i \cdot N_i)^2}{\sigma_{\text{zoo}} + (\sum_i p_i \cdot N_i)^2} \cdot f_T \cdot \text{ZooN} \quad (\text{A62})$$

$G_{\text{tot}}^{\text{zoo}}$  ( $G_{\text{tot}}^{\text{zoo}2}$ ) is the total grazing flux which is calculated for small (macro) zooplankton. ZooN (Zoo2N) is listed in Table A1. The maximum grazing rate ( $\xi_{\text{zoo}}$ ,  $\xi_{\text{zoo}2}$ ) and the half saturation constants ( $\sigma_{\text{zoo}}$ ,  $\sigma_{\text{zoo}2}$ ) are listed in Table A10. The temperature dependency terms ( $f_T$ ,  $f_{T_{\text{zoo}2}}$ ) are given in Eqs. A43 and A44. In the model, relative grazing preferences are implemented following Fasham et al. (1990). Variable relative grazing preferences ( $p_i$ ) are calculated using the nominal preferences for

1045 small phytoplankton, diatoms, slow-/fast-sinking detritus and small zooplankton (Table A10) as follows:

$$p_i = \frac{p'_i \cdot N_i}{\sum_i p'_i \cdot N_i} \quad (\text{A63})$$

. Here, summation  $i$  is done over each food source to calculate the relative proportion of the food. Total grazing is used to calculate the grazing of zooplankton groups on individual food source, i.e., small phytoplankton ( $i=1$ ,  $\text{PhyN}_{\text{small}}$ ), diatoms ( $i=2$ ,  $\text{PhyN}_{\text{dia}}$ ), both detritus classes ( $i=3$ ,  $\text{DetN}$  and  $i=4$ ,  $\text{DetZ2N}$ ) and ( $i=5$ ,  $\text{ZooN}$ ) in the case of macrozooplankton as the ratio

1050 of each food source to total food source ( $G_{\text{small}}$ ,  $G_{\text{dia}}$ ,  $G_{\text{det}}$ ,  $G_{\text{detZ2}}$  and  $G_{\text{zoo}}$ ).

$$G_{\text{small}}^{\text{zoo}} = G_{\text{tot}}^{\text{zoo}} \cdot \frac{p_{\text{small}} \cdot \text{PhyN}_{\text{small}}}{\sum_i p_i \cdot N_i}, \quad G_{\text{small}}^{\text{zoo}2} = G_{\text{tot}}^{\text{zoo}2} \cdot \frac{p_{\text{small}} \cdot \text{PhyN}_{\text{small}}}{\sum_i p_i \cdot N_i} \quad (\text{A64})$$

$$G_{\text{dia}}^{\text{zoo}} = G_{\text{tot}}^{\text{zoo}} \cdot \frac{p_{\text{dia}} \cdot \text{PhyN}_{\text{dia}}}{\sum_i p_i \cdot N_i}, \quad G_{\text{dia}}^{\text{zoo}2} = G_{\text{tot}}^{\text{zoo}2} \cdot \frac{p_{\text{dia}} \cdot \text{PhyN}_{\text{dia}}}{\sum_i p_i \cdot N_i} \quad (\text{A65})$$

$$1055 \quad G_{\text{det}}^{\text{zoo}} = G_{\text{tot}}^{\text{zoo}} \cdot \frac{p_{\text{det}} \cdot \text{DetN}}{\sum_i p_i \cdot N_i}, \quad G_{\text{det}}^{\text{zoo}2} = G_{\text{tot}}^{\text{zoo}2} \cdot \frac{p_{\text{det}} \cdot \text{DetN}}{\sum_i p_i \cdot N_i} \quad (\text{A66})$$

$$G_{\text{detZ2}}^{\text{zoo}} = G_{\text{tot}}^{\text{zoo}} \cdot \frac{p_{\text{detZ2}} \cdot \text{DetZ2N}}{\sum_i p_i \cdot N_i}, \quad G_{\text{detZ2}}^{\text{zoo}2} = G_{\text{tot}}^{\text{zoo}2} \cdot \frac{p_{\text{detZ2}} \cdot \text{DetZ2N}}{\sum_i p_i \cdot N_i} \quad (\text{A67})$$

$$G_{\text{zoo}} = G_{\text{tot}}^{\text{zoo}2} \cdot \frac{p_{\text{zooN}} \cdot \text{ZooN}}{\sum_i p_i \cdot N_i} \quad (\text{A68})$$

where  $G_{\text{zoo}}$  is associated with macrozooplankton grazing on small zooplankton.  $\text{PhyN}_{\text{small}}$ ,  $\text{PhyN}_{\text{dia}}$ ,  $\text{ZooN}$ ,  $\text{DetN}$  and

1060  $\text{DetZ2N}$  are listed in Table A1.

## A5 Bottom boundary fluxes

The model contains a benthic layer at the sea floor. Within this benthic layer, the total amounts of organic carbon, organic nitrogen, biogenic silica and CaCO<sub>3</sub> are modeled.

**Loss to benthos:** When the slow- and fast-sinking detritus reach the ocean bottom, they continue to sink into the benthic layer with the speed  $w_{\text{det}}$  (Eq. A39) and  $w_{\text{detZ2}} = 200 \text{ m d}^{-1}$ , respectively. This results in a detrital flux ( $\text{BenF}_{\text{DetN}}$ ,  $\text{BenF}_{\text{DetZ2N}}$ ,  $\text{BenF}_{\text{DetC}}$ ,  $\text{BenF}_{\text{DetZ2C}}$ ,  $\text{BenF}_{\text{DetSi}}$ ,  $\text{BenF}_{\text{DetZ2Si}}$ ,  $\text{BenF}_{\text{DetCalc}}$  and  $\text{BenF}_{\text{DetZ2Calc}}$ , Table A11) from the water column to the benthos.

$$\text{BenF}_{\text{DetN}} = -w_{\text{det}} \cdot \text{DetN} \quad (\text{A69})$$

$$\text{BenF}_{\text{DetC}} = -w_{\text{det}} \cdot \text{DetC} \quad (\text{A70})$$

$$1070 \quad \text{BenF}_{\text{DetSi}} = -w_{\text{det}} \cdot \text{DetSi} \quad (\text{A71})$$

$$\text{BenF}_{\text{DetCalc}} = -w_{\text{det}} \cdot \text{DetCalc} \quad (\text{A72})$$

$$\text{BenF}_{\text{DetZ2N}} = -w_{\text{detZ2}} \cdot \text{DetZ2N} \quad (\text{A73})$$

$$\text{BenF}_{\text{DetZ2C}} = -w_{\text{detZ2}} \cdot \text{DetZ2C} \quad (\text{A74})$$

$$\text{BenF}_{\text{DetZ2Si}} = -w_{\text{detZ2}} \cdot \text{DetZ2Si} \quad (\text{A75})$$

$$1075 \quad \text{BenF}_{\text{DetZ2Calc}} = -w_{\text{detZ2}} \cdot \text{DetZ2Calc} \quad (\text{A76})$$

These fluxes increase the total amount of the different benthic state variables. The state variables DetN, DetC, DetSi, DetCalc, DetZ2N, DetZ2C, DetZ2Si and DetZ2Calc are described in Table A1.

**Input from benthos:** The lowermost ocean layer located next to the benthic layer receives remineralized inorganic matter back from the benthos. These fluxes, at the same time reduce the amount of the benthic variables. In addition, a sediment flux of Fe from the sediment is calculated from the nitrogen flux, but assuming a Fe:N ratio that is higher than in biomass. This parameterization models that the release of iron from the sediment is driven by redox processes, which are ultimately tied to their remineralization of organic matter.

$$\text{BenF}_{\text{DIN}} = \rho_{\text{ben}}^{\text{N}} \cdot \text{BenthosN} \quad (\text{A77})$$

$$\text{BenF}_{\text{DSi}} = \rho_{\text{ben}}^{\text{Si}} \cdot \text{BenthosSi} \quad (\text{A78})$$

$$1085 \quad \text{BenF}_{\text{DIC}} = \rho_{\text{ben}}^{\text{C}} \cdot \text{BenthosC} + \text{Diss}_{\text{calc}} \cdot \text{BenthosCalc} + \text{Diss}_{\text{calc2}} \cdot \text{BenthosCalc2} \quad (\text{A79})$$

$$\text{BenF}_{\text{Alk}} = (1 + 1/16) \cdot \rho_{\text{ben}}^{\text{N}} \cdot \text{BenthosN} + 2 \cdot \text{Diss}_{\text{calc}} \cdot \text{BenthosCalc} \quad (\text{A80})$$

$\text{BenF}_{\text{DIN}}$ ,  $\text{BenF}_{\text{DSi}}$ ,  $\text{BenF}_{\text{DIC}}$  and  $\text{BenF}_{\text{Alk}}$  (Table A11) denote the fluxes of DIN, DSi, DIC and Alk returned into the bottom layer of the ocean. Constant remineralization rates ( $\rho_{\text{ben}}^{\text{N}}$ ,  $\rho_{\text{ben}}^{\text{Si}}$  and  $\rho_{\text{ben}}^{\text{C}}$ ) are listed in Table A8. The calcite dissolution rates  $\text{Diss}_{\text{calc}}$  and  $\text{Diss}_{\text{calc2}}$  are calculated in Eq. (A38). BenthosN, BenthosSi, BenthosC and BenthosCalc denote the vertically integrated benthos concentration of dissolved nitrogen, silicate, carbon and calcium carbonate, respectively (Table A2). The

1090

alkalinity of the lowermost ocean layer located next to the benthic layer is changed by the remineralization of DIN, dissolved inorganic phosphate converted from DIN with Redfield ratio) and dissolution of calcite from the benthos.

**Table A1.** List of oceanic state variables in REcoM3.

Variable	Description	Unit
DIN	Dissolved Inorganic Nitrogen	[mmol N m <sup>-3</sup> ]
DSi	Dissolved Inorganic Silicon	[mmol N m <sup>-3</sup> ]
DFe	Dissolved Inorganic Iron	[μmol Fe m <sup>-3</sup> ]
DIC	Dissolved Inorganic Carbon	[mmol C m <sup>-3</sup> ]
Alk	Alkalinity	[mmol C m <sup>-3</sup> ]
PhyN <sub>small</sub>	Intracellular nitrogen concentration in small phytoplankton	[mmol N m <sup>-3</sup> ]
PhyC <sub>small</sub>	Intracellular carbon concentration in small phytoplankton	[mmol C m <sup>-3</sup> ]
PhyC <sub>calc</sub>	Intracellular calcite concentration in small phytoplankton	[mmol CaCO <sub>3</sub> m <sup>-3</sup> ]
PhyChl <sub>small</sub>	Intracellular chl <i>a</i> concentration in small phytoplankton	[mg Chl m <sup>-3</sup> ]
PhyN <sub>dia</sub>	Intracellular nitrogen concentration in diatoms	[mmol N m <sup>-3</sup> ]
PhyC <sub>dia</sub>	Intracellular carbon concentration in diatoms	[mmol C m <sup>-3</sup> ]
PhySi <sub>dia</sub>	Intracellular silicon concentration in diatoms	[mmol Si m <sup>-3</sup> ]
PhyChl <sub>dia</sub>	Intracellular chl <i>a</i> concentration in diatoms	[mg Chl m <sup>-3</sup> ]
ZooN	small zooplankton nitrogen concentration	[mmol N m <sup>-3</sup> ]
Zoo2N	Macrozooplankton nitrogen concentration	[mmol N m <sup>-3</sup> ]
ZooC	small zooplankton carbon concentration	[mmol C m <sup>-3</sup> ]
Zoo2C	Macrozooplankton carbon concentration	[mmol C m <sup>-3</sup> ]
DetN	Slow-sinking detritus nitrogen concentration	[mmol N m <sup>-3</sup> ]
Det2N	Fast-sinking detritus nitrogen concentration	[mmol N m <sup>-3</sup> ]
DetC	Slow-sinking detritus carbon concentration	[mmol C m <sup>-3</sup> ]
Det2C	Fast-sinking detritus carbon concentration	[mmol C m <sup>-3</sup> ]
DetCalc	Slow-sinking detritus calcite concentration	[mmol CaCO <sub>3</sub> m <sup>-3</sup> ]
Det2Calc	Fast-sinking detritus calcite concentration	[mmol CaCO <sub>3</sub> m <sup>-3</sup> ]
DetSi	Slow-sinking detritus silicon concentration	[mmol Si m <sup>-3</sup> ]
Det2Si	Fast-sinking detritus silicon concentration	[mmol Si m <sup>-3</sup> ]
DON	Extracellular dissolved organic nitrogen	[mmol N m <sup>-3</sup> ]
DOC	Extracellular dissolved organic carbon	[mmol C m <sup>-3</sup> ]
Oxy	Dissolved oxygen concentration	[mmol O m <sup>-3</sup> ]



**Table A2.** List of benthic state variables in REcoM3.

Variable	Description	Unit
BenthosN	Vertically integrated N concentration	[mmol N m <sup>-2</sup> ]
BenthosC	Vertically integrated C concentration	[mmol C m <sup>-2</sup> ]
BenthosSi	Vertically integrated Si concentration	[mmol Si m <sup>-2</sup> ]
BenthosCalc	Vertically integrated calcite concentration	[mmol CaCO <sub>3</sub> m <sup>-2</sup> ]

**Table A3.** Parameters for sources-minus-sinks equations.

Parameter	Value	Description	Unit
$\psi$	0.02	Calcite production ratio	[dimensionless]
$\gamma_{zoo}$	0.4	Fraction of grazing flux to small zooplankton pool	[dimensionless]
$\gamma_{zoo2}$	0.8	Fraction of grazing flux to macrozooplankton pool	[dimensionless]
$m_{zoo}$	0.05	Small zooplankton mortality rate	[m <sup>3</sup> mmol N <sup>-1</sup> d <sup>-1</sup> ]
$m_{zoo2}$	0.003	Macrozooplankton mortality rate	[m <sup>3</sup> mmol N <sup>-1</sup> d <sup>-1</sup> ]
$\phi_{phy}$	0.015	Max aggregation loss parameter for phytoplankton N	[m <sup>3</sup> mmol N <sup>-1</sup> d <sup>-1</sup> ]
$\phi_{det}$	0.165	Max aggregation loss parameter for detritus N	[m <sup>3</sup> mmol N <sup>-1</sup> d <sup>-1</sup> ]
$w_0$	20.0	Detritus sinking speed at surface	[m d <sup>-1</sup> ]
$f_n$	0.104	N fecal pellet production rate constant	[m <sup>3</sup> mmol N <sup>-1</sup> d <sup>-1</sup> ]
$f_c$	0.236	C fecal pellet production rate constant	[m <sup>3</sup> mmol C <sup>-1</sup> d <sup>-1</sup> ]

**Table A4.** Parameters for iron calculations.

Parameter	Value	Description	Unit
$q^{Fe:N}$	0.033	Intracellular Fe : N ratio	[ $\mu$ mol Fe mmol N <sup>-1</sup> ]
$K_{FeL}$	100.0	Iron stability constant	[m <sup>-3</sup> $\mu$ mol]
$L_T$	1.0	Total ligand concentration	[ $\mu$ mol m <sup>-3</sup> ]
$\kappa_{Fe}$	0.07	Scavenging rate of iron	[m <sup>3</sup> mmol C <sup>-1</sup> d <sup>-1</sup> ]
$q^{Fe:N}$	0.033	Intracellular Fe : N ratio	[ $\mu$ mol Fe mmol N <sup>-1</sup> ]

**Table A5.** Model variables.

Variable	Description	Unit
Agg	Aggregation rate constant [ $\text{d}^{-1}$ ]	
$\text{Diss}_{\text{calc}}$	The dissolution rate constant for slow-sinking detritus	$[\text{d}^{-1}]$
$\text{Diss}_{\text{calc}2}$	The dissolution rate constant for fast-sinking detritus	$[\text{d}^{-1}]$
$\text{Diss}_{\text{calc\_guts}}$	Dissolution of calcium carbonate in guts constant	$[\text{d}^{-1}]$
$w_{\text{det}}$	Sinking velocity of detritus	$[\text{m d}^{-1}]$
$f_T$	Temperature dependence of rates	[dimensionless]
$f_{T_{\text{zoo}2}}$	Temperature dependence of macrozooplankton grazing rates	[dimensionless]
$G_{\text{tot}}$	Total zooplankton grazing rate	$[\text{mmol N m}^{-3} \text{d}^{-1}]$
$G_{\text{small}}$	Small phytoplankton specific zooplankton grazing rate	$[\text{mmol N m}^{-3} \text{d}^{-1}]$
$G_{\text{dia}}$	Diatom specific zooplankton grazing rate	$[\text{mmol N m}^{-3} \text{d}^{-1}]$
PAR	Photosynthetically Available Radiation	$[\text{W m}^{-2}]$
$P_{\text{small}}, P_{\text{dia}}$	C-specific actual rate constant of photosynthesis [ $\text{d}^{-1}$ ]	
$P_{\text{max}}$	C-specific light saturated rate constant of photosynthesis	$[\text{d}^{-1}]$
$r_{\text{small}}$	Small phytoplankton respiration rate constant	$[\text{d}^{-1}]$
$r_{\text{dia}}$	Diatoms respiration rate constant	$[\text{d}^{-1}]$
$r_{\text{zoo}}$	small zooplankton respiration rate constant	$[\text{d}^{-1}]$
$r_{\text{zoo}2}$	Macrozooplankton respiration rate constant	$[\text{d}^{-1}]$
$R_f$	Macrozooplankton feeding activity factor	$[\text{d}^{-1}]$
$R_a$	Macrozooplankton respiration activity factor	$[\text{d}^{-1}]$
$S_{\text{small}}^{\text{chl}}, S_{\text{dia}}^{\text{chl}}$	Rate of chlorophyll <i>a</i> synthesis	$[\text{mg Chl mmol C}^{-1} \text{d}^{-1}]$
$T$	Local temperature	[K]
$V_{\text{small}}^{\text{N}}, V_{\text{dia}}^{\text{N}}$	N-assimilation	$[\text{mmol N mmol C}^{-1} \text{d}^{-1}]$
$\rho_{\text{Si}}^T$	Temperature dependent remineralization rate constant of Si	$[\text{d}^{-1}]$
$V^{\text{Si}}$	Si-assimilation	$[\text{mmol Si mmol C}^{-1} \text{d}^{-1}]$

**Table A6.** Parameters for limitation functions.

Parameter	Value	Description	Unit
$K_{\text{small}}^{\text{Fe}}$	0.04	Half saturation constant for small phytoplankton Fe uptake	$[\mu\text{mol Fe m}^{-3}]$
$K_{\text{dia}}^{\text{Fe}}$	0.12	Half saturation constant for diatom Fe uptake	$[\mu\text{mol Fe m}^{-3}]$
$q_{\text{small}}^{\text{N:Cmin}}$	0.04	Minimum intracellular N : C ratio for small phytoplankton	$[\text{mmol N mmol C}^{-1}]$
$q_{\text{dia}}^{\text{N:Cmin}}$	0.04	Minimum intracellular N : C ratio for diatoms	$[\text{mmol N mmol C}^{-1}]$
$q_{\text{small}}^{\text{N:Cmax}}$	0.2	Maximum intracellular N : C ratio for small phytoplankton	$[\text{mmol N mmol C}^{-1}]$
$q_{\text{dia}}^{\text{N:Cmax}}$	0.2	Maximum intracellular N : C ratio for diatoms	$[\text{mmol N mmol C}^{-1}]$
$q_{\text{dia}}^{\text{Si:Cmin}}$	0.04	Minimum intracellular Si : C ratio for diatoms	$[\text{mmol Si mmol C}^{-1}]$
$q_{\text{dia}}^{\text{Si:Cmax}}$	0.8	Maximum intracellular Si : C ratio for diatoms	$[\text{mmol Si mmol C}^{-1}]$
$\theta_{\text{min}}^{\text{N}}$	50	Minimum limiter regulator for N	$[\text{mmol C mmol N}^{-1}]$
$\theta_{\text{max}}^{\text{N}}$	1000	Maximum limiter regulator for N	$[\text{mmol C mmol N}^{-1}]$
$\theta_{\text{min}}^{\text{Si}}$	1000	Minimum limiter regulator for Si	$[\text{mmol C mmol N}^{-1}]$
$\theta_{\text{max}}^{\text{Si}}$	1000	Maximum limiter regulator for Si	$[\text{mmol C mmol N}^{-1}]$
$T_{\text{ref}}$	288.15	Reference temperature for Arrhenius function	[K]

**Table A7.** Parameters for phytoplankton processes.

Parameter	Value	Description	Unit
$\alpha_{\text{small}}$	0.14	Light harvesting efficiency for small phytoplankton	$[\text{mmol C m}^2 (\text{mg Chl W d})^{-1}]$
$\alpha_{\text{dia}}$	0.19	Light harvesting efficiency for diatoms	$[\text{mmol C m}^2 (\text{mg Chl W d})^{-1}]$
$\mu_{\text{C,small}}^{\text{max}}$	3.0	Rate constant of C-specific photosynthesis	$[\text{d}^{-1}]$
$\mu_{\text{C,dia}}^{\text{max}}$	3.5	Rate constant of C-specific photosynthesis	$[\text{d}^{-1}]$
$\text{res}_{\text{small}}$	0.01	Maintenance respiration rate constant	$[\text{d}^{-1}]$
$\text{res}_{\text{dia}}$	0.01	Maintenance respiration rate constant	$[\text{d}^{-1}]$
$\zeta$	2.33	Cost of biosynthesis of N	$[\text{mmol C mmol N}^{-1}]$
$q_{\text{max,small}}^{\text{Chl:N}}$	3.15	Maximum Chl:N ratio for phytoplankton	$[\text{mg Chl mmol N}^{-1}]$
$q_{\text{max,dia}}^{\text{Chl:N}}$	4.2	Maximum Chl:N ratio for phytoplankton	$[\text{mg Chl mmol N}^{-1}]$
$K_{\text{small}}^{\text{N}}$	0.55	Half saturation constant for small phytoplankton N uptake	$[\text{mmol N m}^{-3}]$
$K_{\text{dia}}^{\text{N}}$	1.00	Half saturation constant for diatom N uptake	$[\text{mmol N m}^{-3}]$
$V_{\text{cm}}^{\text{small}}$	0.7	scaling factor for C-specific N-uptake for small phytoplankton	[dimensionless]
$V_{\text{cm}}^{\text{dia}}$	0.7	scaling factor for C-specific N-uptake for diatoms	[dimensionless]
$\sigma_{\text{N:C}}^{\text{small}}$	0.2	Maximum uptake ratio N : C for small phytoplankton	$[\text{mmol N mmol C}^{-1}]$
$\sigma_{\text{N:C}}^{\text{dia}}$	0.2	Maximum uptake ratio N : C for diatoms	$[\text{mmol N mmol C}^{-1}]$
$K_{\text{Si}}$	4.00	Half saturation constant for diatom Si uptake	$[\text{mmol Si m}^{-3}]$
$\sigma_{\text{Si:C}}$	0.2	Maximum uptake ratio Si : C	$[\text{mmol Si mmol C}^{-1}]$

**Table A8.** Degradation parameters for sources-minus-sinks equations.

Parameter	Value	Description	Unit
$\epsilon_{\text{phy}}^{\text{N}}$	0.05	Small phytoplankton excretion constant of organic N	[d <sup>-1</sup> ]
$\epsilon_{\text{dia}}^{\text{N}}$	0.05	Diatoms excretion constant of organic N	[d <sup>-1</sup> ]
$\epsilon_{\text{phy}}^{\text{C}}$	0.1	Small phytoplankton excretion constant of organic C	[d <sup>-1</sup> ]
$\epsilon_{\text{dia}}^{\text{C}}$	0.1	Diatoms excretion constant of organic C	[d <sup>-1</sup> ]
$\epsilon_{\text{zoo}}^{\text{N}}$	0.15	small zooplankton excretion constant of organic N	[d <sup>-1</sup> ]
$\epsilon_{\text{zoo}2}^{\text{N}}$	0.02	Macrozooplankton excretion constant of organic N	[d <sup>-1</sup> ]
$\epsilon_{\text{zoo}}^{\text{C}}$	0.15	small zooplankton excretion constant of organic C	[d <sup>-1</sup> ]
$\epsilon_{\text{zoo}2}^{\text{C}}$	0.02	Macrozooplankton excretion constant of organic C	[d <sup>-1</sup> ]
$\rho_{\text{ben}}^{\text{N}}$	0.005	Remineralization rate constant for benthos N	[d <sup>-1</sup> ]
$\rho_{\text{ben}}^{\text{Si}}$	0.005	Remineralization rate constant for benthos Si	[d <sup>-1</sup> ]
$\rho_{\text{ben}}^{\text{C}}$	0.005	Remineralization rate constant for benthos C	[d <sup>-1</sup> ]
$\rho_{\text{DON}}$	0.11	Remineralization constant of DON	[d <sup>-1</sup> ]
$\rho_{\text{DOC}}$	0.1	Remineralization constant of DOC	[d <sup>-1</sup> ]
$\rho_{\text{DetN}}$	0.165	Degradation constant of DetN	[d <sup>-1</sup> ]
$\rho_{\text{DetZ2N}}$	0.165	Degradation constant of DetZ2N	[d <sup>-1</sup> ]
$\rho_{\text{DetC}}$	0.15	Degradation constant of DetC	[d <sup>-1</sup> ]
$\rho_{\text{DetZ2C}}$	0.15	Degradation constant of DetZ2C	[d <sup>-1</sup> ]
$\text{deg}_{\text{small}}^{\text{chl}}$	0.2	Small phytoplankton chlorophyll <i>a</i> degradation rate constant	[d <sup>-1</sup> ]
$\text{deg}_{\text{dia}}^{\text{chl}}$	0.2	Diatom chlorophyll <i>a</i> degradation rate constant	[d <sup>-1</sup> ]
$\text{DisS}_{\text{calc\_rate}}$	0.005714	Dissolution of calcium carbonate constant	[d <sup>-1</sup> ]

**Table A9.** Parameters for macrozooplankton grazing.

Parameter	Value	Description	Unit
$Q_a$	28145	Temperatures for the uninhibited reaction kinetics	[°K]
$Q_h$	105234	Temperatures for the inhibited reaction kinetics	[°K]
$T_r$	272.5	Intrinsic optimum temperature	[°K]
$T_h$	274.5	Temperature above which inhibitive processes dominate	[°K]

**Table A10.** Parameters for zooplankton processes.

Parameter	Value	Description	Unit
$\xi_{zoo}$	2.4	Maximum grazing rate constant, small zooplankton	[d <sup>-1</sup> ]
$\xi_{zoo2}$	0.1	Maximum grazing rate constant, macrozooplankton	[d <sup>-1</sup> ]
$\sigma_{zoo}$	0.35	Half saturation constant, small zooplankton	[(mmol N m <sup>-3</sup> ) <sup>2</sup> ]
$\sigma_{zoo2}$	0.0144	Half saturation constant, macrozooplankton	[(mmol N m <sup>-3</sup> ) <sup>2</sup> ]
$\tau$	0.01	Time scale constant for zooplankton respiration	[d <sup>-1</sup> ]
$R_s$	0.0107	Standard respiration rate constant	[d <sup>-1</sup> ]
small zooplankton			
$p'_{small}$	1.0	Initial grazing preference for small phytoplankton	[dimensionless]
$p'_{dia}$	0.5	Initial grazing preference for diatoms	[dimensionless]
$p'_{det}$	0.5	Initial grazing preference for slow-sinking detritus	[dimensionless]
$p'_{detZ2}$	0.5	Initial grazing preference for fast-sinking detritus	[dimensionless]
Macrozooplankton			
$p'_{small}$	0.5	Initial grazing preference for small phytoplankton	[dimensionless]
$p'_{dia}$	1.0	Initial grazing preference for diatoms	[dimensionless]
$p'_{zoo}$	0.8	Initial grazing preference for zooplankton	[dimensionless]
$p'_{det}$	0.5	Initial grazing preference for slow-sinking detritus	[dimensionless]
$p'_{detZ2}$	0.5	Initial grazing preference for fast-sinking detritus	[dimensionless]

**Table A11.** Benthos variables.

Variable	Description	Unit
BenF <sub>Alk</sub>	Flux of alkalinity from benthos to bottom water	[mmol m <sup>-2</sup> d <sup>-1</sup> ]
BenF <sub>DIC</sub>	Flux of C from benthos to bottom water	[mmol C m <sup>-2</sup> d <sup>-1</sup> ]
BenF <sub>DIN</sub>	Flux of N from benthos to bottom water	[mmol N m <sup>-2</sup> d <sup>-1</sup> ]
BenF <sub>DSi</sub>	Flux of Si from benthos to bottom water	[mmol Si m <sup>-2</sup> d <sup>-1</sup> ]
BenF <sub>DetCalc</sub>	Flux of slow-sinking detritus calcite from the water to the benthos	[mmol CaCO <sub>3</sub> m <sup>-2</sup> d <sup>-1</sup> ]
BenF <sub>DetC</sub>	Flux of slow-sinking detritus C from the water to the benthos	[mmol C m <sup>-2</sup> d <sup>-1</sup> ]
BenF <sub>DetN</sub>	Flux of slow-sinking detritus N from the water to the benthos	[mmol N m <sup>-2</sup> d <sup>-1</sup> ]
BenF <sub>DetSi</sub>	Flux of slow-sinking detritus Si from the water to the benthos	[mmol Si m <sup>-2</sup> d <sup>-1</sup> ]
BenF <sub>DetZ2Calc</sub>	Flux of fast-sinking detritus calcite from the water to the benthos	[mmol CaCO <sub>3</sub> m <sup>-2</sup> d <sup>-1</sup> ]
BenF <sub>DetZ2C</sub>	Flux of fast-sinking detritus C from the water to the benthos	[mmol C m <sup>-2</sup> d <sup>-1</sup> ]
BenF <sub>DetZ2N</sub>	Flux of fast-sinking detritus N from the water to the benthos	[mmol N m <sup>-2</sup> d <sup>-1</sup> ]
BenF <sub>DetZ2Si</sub>	Flux of fast-sinking detritus Si from the water to the benthos	[mmol Si m <sup>-2</sup> d <sup>-1</sup> ]

## References

- Albani, S., Mahowald, N. M., Perry, A. T., Scanza, R. A., Zender, C. S., Heavens, N. G., Maggi, V., Kok, J. F., and Otto-Bliesner, B. L.:  
1095 Improved dust representation in the Community Atmosphere Model, *Journal of Advances in Modeling Earth Systems*, 6, 541–570,  
<https://doi.org/10.1002/2013MS000279>, 2014.
- Álvarez, E., Thoms, S., and Völker, C.: Chlorophyll to Carbon Ratio Derived From a Global Ecosystem Model With Photodamage, *Global  
Biogeochemical Cycles*, 32, 799–816, <https://doi.org/10.1029/2017GB005850>, 2018.
- Anderson, L. G., Jutterström, S., Hjalmarsson, S., Wählström, I., and Semiletov, I. P.: Out-gassing of CO<sub>2</sub> from Siberian Shelf seas by  
1100 terrestrial organic matter decomposition, *Geophysical Research Letters*, 36, <https://doi.org/https://doi.org/10.1029/2009GL040046>, 2009.
- Anderson, T. R., Gentleman, W. C., and Sinha, B.: Influence of grazing formulations on the emergent properties of  
a complex ecosystem model in a global ocean general circulation model, *Progress in Oceanography*, 87, 201–213,  
<https://doi.org/https://doi.org/10.1016/j.pocean.2010.06.003>, 3rd GLOBEC OSM: From ecosystem function to ecosystem prediction,  
2010.
- 1105 Aumont, O., Orr, J. C., Monfray, P., Ludwig, W., Amiotte-Suchet, P., and Probst, J.-L.: Riverine-driven interhemispheric transport of carbon,  
*Global Biogeochemical Cycles*, 15, 393–405, <https://doi.org/https://doi.org/10.1029/1999GB001238>, 2001.
- Aumont, O., Maier-Reimer, E., Blain, S., and Monfray, P.: An ecosystem model of the global ocean including Fe, Si, P colimitations, *Global  
Biogeochemical Cycles*, 17, <https://doi.org/https://doi.org/10.1029/2001GB001745>, 2003.
- Aumont, O., Ethé, C., Tagliabue, A., Bopp, L., and Gehlen, M.: PISCES-v2: an ocean biogeochemical model for carbon and ecosystem  
1110 studies, *Geoscientific Model Development*, 8, 2465–2513, <https://doi.org/10.5194/gmd-8-2465-2015>, 2015.
- Bakker, D., De Baar, H., and Bathmann, U.: Changes of carbon dioxide in surface waters during spring in the Southern Ocean, *Deep Sea  
Research Part II: Topical Studies in Oceanography*, 44, 91–127, [https://doi.org/https://doi.org/10.1016/S0967-0645\(96\)00075-6](https://doi.org/https://doi.org/10.1016/S0967-0645(96)00075-6), 1997.
- Bakker, D. C. E., Pfeil, B., Landa, C. S., Metzl, N., O'Brien, K. M., Olsen, A., Smith, K., Cosca, C., Harasawa, S., Jones, S. D., Nakaoka, S.,  
1115 Nojiri, Y., Schuster, U., Steinhoff, T., Sweeney, C., Takahashi, T., Tilbrook, B., Wada, C., Wanninkhof, R., Alin, S. R., Balestrini, C. F.,  
Barbero, L., Bates, N. R., Bianchi, A. A., Bonou, F., Boutin, J., Bozec, Y., Burger, E. F., Cai, W.-J., Castle, R. D., Chen, L., Chierici, M.,  
Currie, K., Evans, W., Featherstone, C., Feely, R. A., Fransson, A., Goyet, C., Greenwood, N., Gregor, L., Hankin, S., Hardman-Mountford,  
N. J., Harlay, J., Hauck, J., Hoppema, M., Humphreys, M. P., Hunt, C. W., Huss, B., Ibáñez, J. S. P., Johannessen, T., Keeling, R., Kitidis,  
V., Körtzinger, A., Kozyr, A., Krasakopoulou, E., Kuwata, A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lo Monaco, C., Manke, A.,  
Mathis, J. T., Merlivat, L., Millero, F. J., Monteiro, P. M. S., Munro, D. R., Murata, A., Newberger, T., Omar, A. M., Ono, T., Paterson,  
1120 K., Pearce, D., Pierrot, D., Robbins, L. L., Saito, S., Salisbury, J., Schlitzer, R., Schneider, B., Schweitzer, R., Sieger, R., Skjelvan, I.,  
Sullivan, K. F., Sutherland, S. C., Sutton, A. J., Tadokoro, K., Telszewski, M., Tuma, M., van Heuven, S. M. A. C., Vandemark, D., Ward,  
B., Watson, A. J., and Xu, S.: A multi-decade record of high-quality *f*CO<sub>2</sub> data in version 3 of the Surface Ocean CO<sub>2</sub> Atlas (SOCAT),  
*Earth System Science Data*, 8, 383–413, <https://doi.org/10.5194/essd-8-383-2016>, 2016.
- Ballantyne, A. P., Alden, C. B., Miller, J. B., Tans, P. P., and White, J. W. C.: Increase in observed net carbon dioxide uptake by land and  
1125 oceans during the past 50 years, *Nature*, 488, 70–72, <https://doi.org/10.1038/nature11299>, 2012.
- Behrenfeld, M. J. and Falkowski, P. G.: A consumer's guide to phytoplankton primary productivity models, *Limnology and Oceanography*,  
42, 1479–1491, <https://doi.org/10.1038/s41586-019-1098-2>, 1997.

- Berthet, S., S  ferian, R., Bricaud, C., Chevallier, M., Voldoire, A., and Eth  , C.: Evaluation of an Online Grid-Coarsening Algorithm in a Global Eddy-Admitting Ocean Biogeochemical Model, *Journal of Advances in Modeling Earth Systems*, 11, 1759–1783, <https://doi.org/https://doi.org/10.1029/2019MS001644>, 2019.
- Blondeau-Patissier, D., Gower, J. F., Dekker, A. G., Phinn, S. R., and Brando, V. E.: A review of ocean color remote sensing methods and statistical techniques for the detection, mapping and analysis of phytoplankton blooms in coastal and open oceans, *Progress in Oceanography*, 123, 123–144, <https://doi.org/https://doi.org/10.1016/j.pocean.2013.12.008>, 2014.
- Bopp, L., Resplandy, L., Orr, J. C., Doney, S. C., Dunne, J. P., Gehlen, M., Halloran, P., Heinze, C., Ilyina, T., S  ferian, R., Tjiputra, J., and Vichi, M.: Multiple stressors of ocean ecosystems in the 21st century: projections with CMIP5 models, *Biogeosciences*, 10, 6225–6245, <https://doi.org/10.5194/bg-10-6225-2013>, 2013.
- Bourgeois, T., Goris, N., Schwinger, J., and Tjiputra, J. F.: Stratification constrains future heat and carbon uptake in the Southern Ocean between 30  S and 55  S, *Nature Communications*, 13, 340, <https://doi.org/10.1038/s41467-022-27979-5>, 2022.
- Boyd, P., Claustre, H., Levy, M., Siegel, D. A., and Weber, T.: Multi-faceted particle pumps drive carbon sequestration in the ocean, *Nature*, 568, 327–335, <https://doi.org/10.1038/s41586-019-1098-2>, 2019.
- Boye, M., van den Berg, C. M., de Jong, J. T., Leach, H., Croot, P., and de Baar, H. J.: Organic complexation of iron in the Southern Ocean, *Deep Sea Research Part I: Oceanographic Research Papers*, 48, 1477–1497, [https://doi.org/https://doi.org/10.1016/S0967-0637\(00\)00099-6](https://doi.org/https://doi.org/10.1016/S0967-0637(00)00099-6), 2001.
- Buitenhuis, E., Rivkin, R. B., S  ailley, S., and Le Qu  r  , C.: Biogeochemical fluxes through microzooplankton, *Global Biogeochemical Cycles*, 24, <https://doi.org/10.1029/2009GB003601>, 2010.
- Bunsen, F.: Impact of recent climate variability on oceanic CO2 uptake in a global ocean biogeochemistry model, Master’s thesis, Kiel University Christian-Albrechts-Universit  t, Faculty of Mathematics and Natural Sciences, 2022.
- Bushinsky, S. M., Landsch  tzer, P., R  denbeck, C., Gray, A. R., Baker, D., Mazloff, M. R., Resplandy, L., Johnson, K. S., and Sarmiento, J. L.: Reassessing Southern Ocean air-sea CO2 flux estimates with the addition of biogeochemical float observations., *Global Biogeochemical Cycles*, 33, 1370–1388, <https://doi.org/10.1029/2019GB006176>, 2019.
- Butzin, M. and P  rtner, H. O.: Thermal growth potential of Atlantic cod by the end of the 21st century, *Global Change Biology*, 22, 4162–4168, <https://doi.org/10.1111/gcb.13375>, 2016.
- Campin, J.-M., Adcroft, A. J., Hill, C., and Marshall, J.: Conservation of properties in a free-surface model, *Ocean Modelling*, 6, 221–244, [https://doi.org/10.1016/S1463-5003\(03\)00009-X](https://doi.org/10.1016/S1463-5003(03)00009-X), 2004.
- Canadell, J., Monteiro, P., Costa, M., Cotrim da Cunha, L., Cox, P., Eliseev, A., Henson, S., Ishii, M., Jaccard, S., Koven, C., Lohila, A., Patra, P., Piao, S., Rogelj, J., Syampungani, S., Zaehle, S., and Zickfeld, K.: Global Carbon and other Biogeochemical Cycles and Feedbacks, p. 673–816, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/9781009157896.007>, 2021.
- Carr, M.-E., Friedrichs, M. A., Schmeltz, M., Noguchi Aita, M., Antoine, D., Arrigo, K. R., Asanuma, I., Aumont, O., Barber, R., Behrenfeld, M., Bidigare, R., Buitenhuis, E. T., Campbell, J., Ciotti, A., Dierssen, H., Dowell, M., Dunne, J., Esaias, W., Gentili, B., Gregg, W., Groom, S., Hoepffner, N., Ishizaka, J., Kameda, T., Le Qu  r  , C., Lohrenz, S., Marra, J., M  lin, F., Moore, K., Morel, A., Reddy, T. E., Ryan, J., Scardi, M., Smyth, T., Turpie, K., Tilstone, G., Waters, K., and Yamanaka, Y.: A comparison of global estimates of marine primary production from ocean color, *Deep Sea Research Part II: Topical Studies in Oceanography*, 53, 741–770, <https://doi.org/https://doi.org/10.1016/j.dsr2.2006.01.028>, the US JGOFS Synthesis and Modeling Project: Phase III, 2006.

- 1165 Chau, T. T. T., Gehlen, M., and Chevallier, F.: A seamless ensemble-based reconstruction of surface ocean  $p\text{CO}_2$  and air–sea  $\text{CO}_2$  fluxes over the global coastal and open oceans, *Biogeosciences*, 19, 1087–1109, <https://doi.org/10.5194/bg-19-1087-2022>, 2022.
- Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R., Galloway, J., Heimann, M., Le Quéré, C., Myneni, R., Piao, S., and Thornton, P.: Carbon and other biogeochemical cycles, in: *Climate change 2013: the physical science basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, pp. 465–570, Cambridge University Press, 2014.
- 1170 Claquin, P., Martin-Jézéquel, V., Kromkamp, J. C., Veldhuis, M. J. W., and Kraay, G. W.: Uncoupling of Silicon Compared With Carbon and Nitrogen Metabolisms and the Role of the Cell Cycle in Continuous Cultures of *Thalassiosira Pseudonana* (Bacillariophyceae) Under Light, Nitrogen, and Phosphorus Control, *Journal of Phycology*, 38, 922–930, <https://doi.org/10.1046/j.1529-8817.2002.t01-1-01220.x>, 2002.
- 1175 Cocco, V., Joos, F., Steinacher, M., Frölicher, T. L., Bopp, L., Dunne, J., Gehlen, M., Heinze, C., Orr, J., Oschlies, A., Schneider, B., Segschneider, J., and Tjiputra, J.: Oxygen and indicators of stress for marine life in multi-model global warming projections, *Biogeosciences*, 10, 1849–1868, <https://doi.org/10.5194/bg-10-1849-2013>, 2013.
- Cram, J. A., Weber, T., Leung, S. W., McDonnell, A. M. P., Liang, J.-H., and Deutsch, C.: The Role of Particle Size, Ballast, Temperature, and Oxygen in the Sinking Flux to the Deep Sea, *Global Biogeochemical Cycles*, 32, 858–876, <https://doi.org/https://doi.org/10.1029/2017GB005710>, 2018.
- 1180 Crisp, D., Dolman, H., Tanhua, T., McKinley, G. A., Hauck, J., Bastos, A., Sitch, S., Eggleston, S., and Aich, V.: How Well Do We Understand the Land-Ocean-Atmosphere Carbon Cycle?, *Reviews of Geophysics*, 60, e2021RG000736, <https://doi.org/https://doi.org/10.1029/2021RG000736>, e2021RG000736 2021RG000736, 2022.
- Danabasoglu, G., Yeager, S. G., Bailey, D., Behrens, E., Bentsen, M., Bi, D., Biastoch, A., Böning, C., Bozec, A., Canuto, V. M., Cassou, C., 1185 Chassignet, E., Coward, A. C., Danilov, S., Diansky, N., Drange, H., Farneti, R., Fernandez, E., Fogli, P. G., Forget, G., Fujii, Y., Griffies, S. M., Gusev, A., Heimbach, P., Howard, A., Jung, T., Kelley, M., Large, W. G., Leboissetier, A., Lu, J., Madec, G., Marsland, S. J., Masina, S., Navarra, A., George Nurser, A., Pirani, A., y Mélia, D. S., Samuels, B. L., Scheinert, M., Sidorenko, D., Treguier, A.-M., Tsujino, H., Uotila, P., Valcke, S., Voldoire, A., and Wang, Q.: North Atlantic simulations in Coordinated Ocean-ice Reference Experiments phase II (CORE-II). Part I: Mean states, *Ocean Modelling*, 73, 76–107, <https://doi.org/https://doi.org/10.1016/j.ocemod.2013.10.005>, 2014.
- 1190 Danilov, S.: Ocean modeling on unstructured meshes, *Ocean Modelling*, 69, 195–210, <https://doi.org/https://doi.org/10.1016/j.ocemod.2013.05.005>, 2013.
- Danilov, S., Wang, Q., Timmermann, R., Iakovlev, N., Sidorenko, D., Kimmritz, M., Jung, T., and Schröter, J.: Finite-Element Sea Ice Model (FESIM), version 2, *Geosci. Model Dev.*, 8, 1747–1761, <https://doi.org/10.5194/gmd-8-1747-2015>, 2015.
- Danilov, S., Sidorenko, D., Wang, Q., and Jung, T.: The Finite-volumE Sea ice–Ocean Model (FESOM2), *Geoscientific Model Development*, 10, 765–789, <https://doi.org/10.5194/gmd-10-765-2017>, 2017.
- 1195 de Baar, H. J., de Jong, J. T., Nolting, R. F., Timmermans, K. R., van Leeuwe, M. A., Bathmann, U., Rutgers van der Loeff, M., and Sildam, J.: Low dissolved Fe and the absence of diatom blooms in remote Pacific waters of the Southern Ocean, *Marine Chemistry*, 66, 1–34, [https://doi.org/https://doi.org/10.1016/S0304-4203\(99\)00022-5](https://doi.org/https://doi.org/10.1016/S0304-4203(99)00022-5), 1999.
- Denman, K. L., Brasseur, G., Chidthaisong, A., Ciais, P., Cox, P. M., Dickinson, R. E., Hauglustaine, D., Heinze, C., Holland, E., Jacob, D., 1200 Lohmann, U., Ramachandran, S., Leite da Silva Dias, P., Wofsy, S. C., and Zhang, X.: Couplings Between Changes in the Climate System and Biogeochemistry, in: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment*



- Report of the Intergovernmental Panel on Climate Change, edited by Solomon, S., Qin, D., Manning, M., Marquis, M., Averyt, K., Tignor, M. M. B., Miller, H. L., and Chen, Z. L., p. 499–587, Cambridge University Press, Cambridge, UK and New York, USA, 2007.
- 1205 DeVries, T., Le Quéré, C., Andrews, O., Berthet, S., Hauck, J., Ilyina, T., Landschützer, P., Lenton, A., Lima, I. D., Nowicki, M., Schwinger, J., and Séférian, R.: Decadal trends in the ocean carbon sink, *Proceedings of the National Academy of Sciences of the United States of America*, 116, 11 646–11 651, <https://doi.org/10.1073/pnas.1900371116>, 2019.
- Doney, S. C., Lindsay, K., Moore, J. K., Dutkiewicz, S., Friedrichs, M. A. M., and Matear, R. J.: Marine Biogeochemical Modeling: Recent Advances and Future Challenges, *Oceanography*, 14, 93–107, <https://doi.org/10.5670/oceanog.2001.10>, 2001.
- 1210 Doney, S. C., Lima, I., Feely, R. A., Glover, D. M., Lindsay, K., Mahowald, N., Moore, J. K., and Wanninkhof, R.: Mechanisms governing interannual variability in upper-ocean inorganic carbon system and air–sea CO<sub>2</sub> fluxes: Physical climate and atmospheric dust, *Deep Sea Research Part II: Topical Studies in Oceanography*, 56, 640–655, <https://doi.org/https://doi.org/10.1016/j.dsr2.2008.12.006>, surface Ocean CO<sub>2</sub> Variability and Vulnerabilities, 2009.
- Du, J., Ye, Y., Zhang, X., Völker, C., and Tian, J.: Southern Control of Interhemispheric Synergy on Glacial Marine Carbon Sequestration, *Geophysical Research Letters*, 49, e2022GL099048, <https://doi.org/https://doi.org/10.1029/2022GL099048>, e2022GL099048 2022GL099048, 2022.
- 1215 Dunne, J. P., Sarmiento, J. L., and Gnanadesikan, A.: A synthesis of global particle export from the surface ocean and cycling through the ocean interior and on the seafloor, *Global Biogeochem. Cycles*, 21, GB4006, <https://doi.org/10.1029/2006GB002907>, 2007.
- Elrod, V. A., Berelson, W. M., Coale, K. H., and Johnson, K. S.: The flux of iron from continental shelf sediments: a missing source for global budgets, *Geophysical Research Letters*, 31, L12 307, <https://doi.org/doi:10.1029/2004GL020216>, 2004.
- 1220 Fasham, M. J., Ducklow, H. W., and McKelvie, S. M.: A nitrogen-based model of plankton dynamics in the oceanic mixed layer, *Journal of Marine Research*, 48, 591–639, <https://doi.org/https://doi.org/10.1357/002224090784984678>, 1990.
- Fay, A. R. and McKinley, G. A.: Observed Regional Fluxes to Constrain Modeled Estimates of the Ocean Carbon Sink, *Geophysical Research Letters*, 48, <https://doi.org/10.1029/2021GL095325>, 2021.
- 1225 Fay, A. R., Gregor, L., Landschützer, P., McKinley, G. A., Gruber, N., Gehlen, M., Iida, Y., Laruelle, G. G., Rödenbeck, C., Roobaert, A., and Zeng, J.: SeaFlux: harmonization of air–sea CO<sub>2</sub> fluxes from surface *p*CO<sub>2</sub> data products using a standardized approach, *Earth System Science Data*, 13, 4693–4710, <https://doi.org/10.5194/essd-13-4693-2021>, 2021.
- Fennel, K., Mattern, J. P., Doney, S. C., Bopp, L., Moore, A. M., Wang, B., and Yu, L.: Ocean biogeochemical modelling, *Nature Reviews Methods Primers*, 2, 76, <https://doi.org/10.1038/s43586-022-00154-2>, 2022.
- 1230 Friedlingstein, P., Dufresne, J.-L., Cox, P. M., and Rayner, P. J.: How positive is the feedback between climate change and the carbon cycle?, *Tellus B*, 55, 692–700, <https://doi.org/10.1034/j.1600-0889.2003.01461.x>, 2003.
- Friedlingstein, P., Jones, M. W., O’Sullivan, M., Andrew, R. M., Bakker, D. C. E., Hauck, J., Le Quéré, C., Peters, G. P., Peters, W., Pongratz, J., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S. R., Anthoni, P., Bates, N. R., Becker, M., Bellouin, N., Bopp, L., Chau, T. T. T., Chevallier, F., Chini, L. P., Cronin, M., Currie, K. I., Decharme, B., Djeutchouang, L. M., Dou, X., Evans, W., Feely, R. A., Feng, L., Gasser, T., Gilfillan, D., Gkritzalis, T., Grassi, G., Gregor, L., Gruber, N., Gürses, Ö., Harris, I., Houghton, R. A., Hurtt, G. C., Iida, Y., Ilyina, T., Luijkx, I. T., Jain, A., Jones, S. D., Kato, E., Kennedy, D., Klein Goldewijk, K., Knauer, J., Korsbakken, J. I., Körtzinger, A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lienert, S., Liu, J., Marland, G., McGuire, P. C., Melton, J. R., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S.-I., Niwa, Y., Ono, T., Pierrot, D., Poulter, B., Rehder, G., Resplandy, L., Robertson, E., Rödenbeck, C., Rosan, T. M., Schwinger, J., Schwingshackl, C., Séférian, R., Sutton, A. J., Sweeney, C., Tanhua, T., Tans, P. P., Tian, H., Tilbrook, B., Tubiello, F., van der Werf, G. R., Vuichard, N., Wada, C., Wanninkhof, R., Watson, A. J., Willis, D., Wiltshire, A. J., Yuan, W., Yue, C., Yue, X.,

- 1240 Zaehle, S., and Zeng, J.: Global Carbon Budget 2021, *Earth System Science Data*, 14, 1917–2005, <https://doi.org/10.5194/essd-14-1917-2022>, 2022a.
- Friedlingstein, P., O’Sullivan, M., Jones, M. W., Andrew, R. M., Gregor, L., Hauck, J., Le Quéré, C., Luijkx, I. T., Olsen, A., Peters, G. P., Peters, W., Pongratz, J., Schwingshackl, C., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S. R., Alkama, R., Arneth, A., Arora, V. K., Bates, N. R., Becker, M., Bellouin, N., Bittig, H. C., Bopp, L., Chevallier, F., Chini, L. P., Cronin, M., Evans, W., Falk, S., Feely, 1245 R. A., Gasser, T., Gehlen, M., Gkritzalis, T., Gloege, L., Grassi, G., Gruber, N., Gürses, O., Harris, I., Hefner, M., Houghton, R. A., Hurtt, G. C., Iida, Y., Ilyina, T., Jain, A. K., Jersild, A., Kadono, K., Kato, E., Kennedy, D., Klein Goldewijk, K., Knauer, J., Korsbakken, J. I., Landschützer, P., Lefèvre, N., Lindsay, K., Liu, J., Liu, Z., Marland, G., Mayot, N., McGrath, M. J., Metzl, N., Monacci, N. M., Munro, D. R., Nakaoka, S.-I., Niwa, Y., O’Brien, K., Ono, T., Palmer, P. I., Pan, N., Pierrot, D., Pockock, K., Poulter, B., Resplandy, L., Robertson, E., Rödenbeck, C., Rodriguez, C., Rosan, T. M., Schwinger, J., Séférian, R., Shutler, J. D., Skjelvan, I., Steinhoff, T., Sun, Q., Sutton, 1250 A. J., Sweeney, C., Takao, S., Tanhua, T., Tans, P. P., Tian, X., Tian, H., Tilbrook, B., Tsujino, H., Tubiello, F., van der Werf, G. R., Walker, A. P., Wanninkhof, R., Whitehead, C., Willstrand Wranne, A., Wright, R., Yuan, W., Yue, C., Yue, X., Zaehle, S., Zeng, J., and Zheng, B.: Global Carbon Budget 2022, *Earth System Science Data*, 14, 4811–4900, <https://doi.org/10.5194/essd-14-4811-2022>, 2022b.
- Frölicher, T. L., Rodgers, K. B., Stock, C. A., and Cheung, W. W. L.: Sources of uncertainties in 21st century projections of potential ocean ecosystem stressors: UNCERTAINTIES IN STRESSOR PROJECTIONS, *Global Biogeochemical Cycles*, 30, 1224–1243, 1255 <https://doi.org/10.1002/2015GB005338>, 2016.
- Galbraith, E. D. and Skinner, L. C.: The Biological Pump During the Last Glacial Maximum, *Annual Review of Marine Science*, 12, 559–586, <https://doi.org/10.1146/annurev-marine-010419-010906>, PMID: 31899673, 2020.
- Garcia, H. E., Locarnini, R. A., Boyer, T. P., Antonov, J. I., Baranova, O. K., Zweng, M. M., Reagan, J. R., and Johnson, D. R.: World Ocean Atlas 2013, Volume 4: Dissolved Inorganic Nutrients (phosphate, nitrate, silicate). S. Levitus, Ed., A. Mishonov Technical Ed., Tech. rep., 1260 NOAA Atlas NESDIS 76, 2014.
- Garcia, H. E., Weathers, K. W., Paver, C. R., Smolyar, I., Boyer, T. P., Locarnini, R. A., Zweng, M. M., Mishonov, A. V., Baranova, O. K., Seidov, D., and Reagan, J. R.: World Ocean Atlas 2018, Volume 3: Dissolved Oxygen, Apparent Oxygen Utilization, and Oxygen Saturation. A. Mishonov Technical Editor, Tech. rep., NOAA Atlas NESDIS 83, 2019a.
- Garcia, H. E., Weathers, K. W., Paver, C. R., Smolyar, I., Boyer, T. P., Locarnini, R. A., Zweng, M. M., Mishonov, A. V., Baranova, 1265 O. K., Seidov, D., and Reagan, J. R.: World Ocean Atlas 2018, Volume 4: Dissolved Inorganic Nutrients (phosphate, nitrate, silicate). A. Mishonov Technical Editor, Tech. rep., NOAA Atlas NESDIS 84, 2019b.
- Geider, R. J. and La Roche, J.: The role of iron in phytoplankton photosynthesis, and the potential for iron-limitation of primary productivity in the sea, *Photosynthesis Research*, 39, 275–301, <https://doi.org/10.1007/BF00014588>, 1994.
- Geider, R. J., MacIntyre, H. L., and Kana, T. M.: A dynamic regulatory model of phytoplanktonic acclimation to light, nutrients, and 1270 temperature, *Limnol. Oceanogr.*, 43, 679–694, <https://doi.org/doi:10.4319/lo.1998.43.4.0679>, 1998.
- Gent, P. and McWilliams, J.: Isopycnal Mixing in Ocean Circulation Models, *J. Phys. Oceanogr.*, 20, 150–155, 1990.
- Gloege, L., McKinley, G. A., Landschützer, P., Fay, A. R., Frölicher, T. L., Fyfe, J. C., Ilyina, T., Jones, S., Lovenduski, N. S., Rodgers, K. B., Schlunegger, S., and Takano, Y.: Quantifying Errors in Observationally Based Estimates of Ocean Carbon Sink Variability, *Global Biogeochemical Cycles*, 35, e2020GB006788, <https://doi.org/https://doi.org/10.1029/2020GB006788>, e2020GB006788 2020GB006788, 1275 2021.
- Goris, N., Tjiputra, J. F., Olsen, A., Schwinger, J., Lauvset, S. K., and Jeansson, E.: Constraining Projection-Based Estimates of the Future North Atlantic Carbon Uptake, *Journal of Climate*, 31, 3959 – 3978, <https://doi.org/https://doi.org/10.1175/JCLI-D-17-0564.1>, 2018.

- Gray, A., Johnson, K. S., Bushinsky, S. M., Riser, S. C., Russell, J. L., Talley, L. D., Wanninkhof, R., Williams, N. L., and Sarmiento, J. L.: Autonomous biogeochemical floats detect significant carbon dioxide outgassing in the high-latitude Southern Ocean., *Geophysical Research Letters*, 45, 9049–9057, <https://doi.org/10.1029/2018GL078013>, 2018.
- 1280
- Gregor, L. and Gruber, N.: OceanSODA-ETHZ: a global gridded data set of the surface ocean carbonate system for seasonal to decadal studies of ocean acidification, *Earth System Science Data*, 13, 777–808, <https://doi.org/10.5194/essd-13-777-2021>, 2021.
- Gregor, L., Lebehot, A. D., Kok, S., and Scheel Monteiro, P. M.: A comparative assessment of the uncertainties of global surface ocean CO<sub>2</sub> estimates using a machine-learning ensemble (CSIR-ML6 version 2019a) – have we hit the wall?, *Geoscientific Model Development*, 12, 5113–5136, <https://doi.org/10.5194/gmd-12-5113-2019>, 2019.
- 1285
- Griffies, S.: The Gent-McWilliams Skew Flux, *J. Phys. Oceanogr.*, 28, 831–841, [https://doi.org/10.1175/1520-0485\(1998\)028<0831:TGMSF>2.0.CO;2](https://doi.org/10.1175/1520-0485(1998)028<0831:TGMSF>2.0.CO;2), 1998.
- Griffies, S. M., Biastoch, A., Böning, C., Bryan, F., Danabasoglu, G., Chassignet, E. P., England, M. H., Gerdes, R., Haak, H., Hallberg, R. W., Hazeleger, W., Jungclaus, J., Large, W. G., Madec, G., Pirani, A., Samuels, B. L., Scheinert, M., Gupta, A. S., Severijns, C. A., 1290 Simmons, H. L., Treguier, A. M., Winton, M., Yeager, S., and Yin, J.: Coordinated Ocean-ice Reference Experiments (COREs), *Ocean Modelling*, 26, 1–46, <https://doi.org/https://doi.org/10.1016/j.ocemod.2008.08.007>, 2009.
- Gruber, N., Gloor, M., Mikaloff Fletcher, S. E., Doney, S. C., Dutkiewicz, S., Follows, M. J., Gerber, M., Jacobson, A. R., Joos, F., Lindsay, K., Menemenlis, D., Mouchet, A., Müller, S. A., Sarmiento, J. L., and Takahashi, T.: Oceanic sources, sinks, and transport of atmospheric CO<sub>2</sub>, *Global Biogeochemical Cycles*, 23, GB1005, <https://doi.org/10.1029/2008GB003349>, 2009.
- 1295
- Gruber, N., Clement, D., Carter, B. R., Feely, R. A., van Heuven, S., Hoppema, M., Ishii, M., Key, R. M., Kozyr, A., Lauvset, S. K., Monaco, C. L., Mathis, J. T., Murata, A., Olsen, A., Perez, F. F., Sabine, C. L., Tanhua, T., and Wanninkhof, R.: The oceanic sink for anthropogenic CO<sub>2</sub> from 1994 to 2007, *Science*, 363, 1193–1199, <https://doi.org/10.1126/science.aau5153>, 2019.
- Hatta, M., Measures, C. I., Wu, J., Roshan, S., Fitzsimmons, J. N., Sedwick, P., and Morton, P.: An overview of dissolved Fe and Mn distributions during the 2010–2011 U.S. GEOTRACES north Atlantic cruises: GEOTRACES GA03, *Deep-Sea Research Part II: Topical Studies in Oceanography*, 116, 117–129, <https://doi.org/10.1016/j.dsr2.2014.07.005>, publisher: Elsevier ISBN: 1354-1013, 1354-1013, 2015.
- 1300
- Hauck, J., Völker, C., Wang, T., Hoppema, M., Losch, M., and Wolf Gladrow, D. A.: Seasonally different carbon flux changes in the Southern Ocean in response to the southern annular mode, *Global Biogeochemical Cycles*, 27, 1236–1245, <https://doi.org/10.1002/2013GB004600>, 2013.
- 1305
- Hauck, J., Völker, C., Wolf-Gladrow, D. A., Laufkötter, C., Vogt, M., Aumont, O., Bopp, L., Buitenhuis, E. T., Doney, S. C., Dunne, J., Gruber, N., Hashioka, T., John, J., Quéré, C. L., Lima, I. D., Nakano, H., Séférian, R., and Totterdell, I.: On the Southern Ocean CO<sub>2</sub> uptake and the role of the biological carbon pump in the 21st century, *Global Biogeochemical Cycles*, 29, 1451–1470, <https://doi.org/10.1002/2015GB005140>, 2015.
- Hauck, J., Köhler, P., Wolf-Gladrow, D., and Völker, C.: Iron fertilisation and century-scale effects of open ocean dissolution of olivine in a simulated CO<sub>2</sub> removal experiment, *Environmental Research Letters*, 11, 024 007, <https://doi.org/10.1088/1748-9326/11/2/024007>, 2016.
- 1310
- Hauck, J., Zeising, M., Le Quéré, C., Gruber, N., Bakker, D. C. E., Bopp, L., Chau, T. T. T., Gürses, Ö., Ilyina, T., Landschützer, P., Lenton, A., Resplandy, L., Rödenbeck, C., Schwinger, J., and Séférian, R.: Consistency and Challenges in the Ocean Carbon Sink Estimate for the Global Carbon Budget, *Frontiers in Marine Science*, 7, <https://doi.org/10.3389/fmars.2020.571720>, 2020.

- Hauck, J., Nissen, C., Landschützer, P., Rödenbeck, C., Bushinsky, S., and Olsen, A.: Sparse observations induce large biases in estimates  
1315 of the global ocean CO<sub>2</sub> sink: an ocean model subsampling experiment, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 381, 20220063, <https://doi.org/10.1098/rsta.2022.0063>, 2023.
- Hauck, J., Lenton, A., Langlais, C., and Matear, R.: The Fate of Carbon and Nutrients Exported Out of the Southern Ocean, *Global Biogeochemical Cycles*, 32, 1556–1573, <https://doi.org/10.1029/2018GB005977>, 2018.
- Henson, S. A., Sanders, R., Madsen, E., Morris, P. J., Le Moigne, F., and Quartly, G. D.: A reduced estimate of the strength of the ocean's  
1320 biological carbon pump, *Geophys. Res. Lett.*, 38, L04606, <https://doi.org/10.1029/2011GL046735>, 2011.
- Henson, S. A., Laufkötter, C., Leung, S., Giering, S. L. C., Palevsky, H. I., and Cavan, E. L.: Uncertain response of ocean biological carbon export in a changing world, *Nature Geoscience*, 15, 248–254, <https://doi.org/10.1038/s41561-022-00927-0>, 2022.
- Ho, D. T., Law, C. S., Smith, M. J., Schlosser, P., Harvey, M., and Hill, P.: Measurements of air-sea gas exchange at high  
1325 wind speeds in the Southern Ocean: Implications for global parameterizations, *Geophysical Research Letters*, 33, L16611, <https://doi.org/10.1029/2006GL026817>, 2006.
- Hohn, S.: Coupling and decoupling of biogeochemical cycles in marine ecosystems, Ph.D. thesis, Universität Bremen, Fachbereich Biologie, 2009.
- Huang, Y., Tagliabue, A., and Cassar, N.: Data-Driven Modeling of Dissolved Iron in the Global Ocean, *Frontiers in Marine Science*, 9, 1–14, <https://doi.org/10.3389/fmars.2022.837183>, 2022.
- 1330 Iida, Y., Kojima, A., Takatani, Y., Nakano, T., Sugimoto, H., Midorikawa, T., and Ishii, M.: Trends in pCO<sub>2</sub> and sea–air CO<sub>2</sub> flux over the global open oceans for the last two decades, *Journal of Oceanography*, 71, 637–661, <https://doi.org/10.1007/s10872-015-0306-4>, 2015.
- Johnson, R., Strutton, P. G., Wright, S. W., McMin, A., and Meiners, K. M.: Three improved Satellite Chlorophyll algorithms for the Southern Ocean, *Journal of Geophysical Research: Oceans*, 118, 3694 – 3703, <https://doi.org/doi:10.1002/jgrc.20270>, 2013.
- Joos, F. and Spahni, R.: Rates of change in natural and anthropogenic radiative forcing over the past 20,000 years, *Proceedings of the National  
1335 Academy of Sciences*, 105, 1425–1430, <https://doi.org/10.1073/pnas.0707386105>, 2008.
- Jungclaus, J. H., Fischer, N., Haak, H., Lohmann, K., Marotzke, J., Matei, D., Mikolajewicz, U., Notz, D., and von Storch, J. S.: Characteristics of the ocean simulations in the Max Planck Institute Ocean Model (MPIOM) the ocean component of the MPI-Earth system model, *Journal of Advances in Modeling Earth Systems*, 5, 422–446, <https://doi.org/https://doi.org/10.1002/jame.20023>, 2013.
- Juricke, S., Danilov, S., Koldunov, N., Oliver, M., and Sidorenko, D.: Ocean Kinetic Energy Backscatter Parametrization on  
1340 Unstructured Grids: Impact on Global Eddy-Permitting Simulations, *Journal of Advances in Modeling Earth Systems*, 12, <https://doi.org/10.1029/2019MS001855>, 2020.
- Karakuş, O., Völker, C., Iversen, M., Hagen, W., Wolf-Gladrow, D., Fach, B., and Hauck, J.: Modeling the Impact of Macrozooplankton on Carbon Export Production in the Southern Ocean, *Journal of Geophysical Research: Oceans*, 126, <https://doi.org/10.1029/2021JC017315>, 2021.
- 1345 Karakuş, O., Völker, C., Iversen, M., Hagen, W., and Hauck, J.: The Role of Zooplankton Grazing and Nutrient Recycling for Global Ocean Biogeochemistry and Phytoplankton Phenology, *Journal of Geophysical Research: Biogeosciences*, 127, e2022JG006798, <https://doi.org/https://doi.org/10.1029/2022JG006798>, e2022JG006798 2022JG006798, 2022.
- Keerthi, M. G., Prend, C. J., Aumont, O., and Lévy, M.: Annual variations in phytoplankton biomass driven by small-scale physical processes, *Nature Geoscience*, <https://doi.org/10.1038/s41561-022-01057-3>, 2022.

- 1350 Keppler, L., Landschützer, P., Gruber, N., Lauvset, S. K., and Stemmler, I.: Seasonal Carbon Dynamics in the Near-Global Ocean, *Global Biogeochemical Cycles*, 34, e2020GB006571, <https://doi.org/https://doi.org/10.1029/2020GB006571>, e2020GB006571 10.1029/2020GB006571, 2020.
- Koeve, W., Duteil, O., Oschlies, A., Kähler, P., and Segschneider, J.: Methods to evaluate CaCO<sub>3</sub> cycle modules in coupled global biogeochemical ocean models, *Geoscientific Model Development*, 7, 2393–2408, <https://doi.org/10.5194/gmd-7-2393-2014>, 2014.
- 1355 Koldunov, N. V., Aizinger, V., Rakowsky, N., Scholz, P., Sidorenko, D., Danilov, S., and Jung, T.: Scalability and some optimization of the Finite-volume Sea ice–Ocean Model, Version 2.0 (FESOM2), *Geoscientific Model Development*, 12, 3991–4012, <https://doi.org/10.5194/gmd-12-3991-2019>, 2019.
- Kriest, I. and Oschlies, A.: On the treatment of particulate organic matter sinking in large-scale models of marine biogeochemical cycles, *Biogeosciences*, 5, 55–72, <https://doi.org/10.5194/bg-5-55-2008>, 2008.
- 1360 Kulk, G., Platt, T., Dingle, J., Jackson, T., Jönsson, B. F., Bouman, H. A., Babin, M., Brewin, R. J. W., Doblin, M., Estrada, M., Figueiras, F. G., Furuya, K., González-Benítez, N., Gudfinnsson, H. G., Gudmundsson, K., Huang, B., Isada, T., Kovac, Z., Lutz, V. A., Marañón, E., Raman, M., Richardson, K., Rozema, P. D., Poll, W. H. v. d., Segura, V., Tilstone, G. H., Uitz, J., Dongen-Vogels, V. v., Yoshikawa, T., and Sathyendranath, S.: Primary Production, an Index of Climate Change in the Ocean: Satellite-Based Estimates over Two Decades, *Remote Sensing*, 12, <https://doi.org/10.3390/rs12050826>, 2020.
- 1365 Kwiatkowski, L., Torres, O., Bopp, L., Aumont, O., Chamberlain, M., Christian, J. R., Dunne, J. P., Gehlen, M., Ilyina, T., John, J. G., Lenton, A., Li, H., Lovenduski, N. S., Orr, J. C., Palmieri, J., Santana-Falcón, Y., Schwinger, J., Séférian, R., Stock, C. A., Tagliabue, A., Takano, Y., Tjiputra, J., Toyama, K., Tsujino, H., Watanabe, M., Yamamoto, A., Yool, A., and Ziehn, T.: Twenty-first century ocean warming, acidification, deoxygenation, and upper-ocean nutrient and primary production decline from CMIP6 model projections, *Biogeosciences*, 17, 3439–3470, <https://doi.org/10.5194/bg-17-3439-2020>, 2020.
- 1370 Kwon, E. Y., Primeau, F., and Sarmiento, J. L.: The impact of remineralization depth on the air–sea carbon balance, *Nature Geoscience*, 2, 630–635, <https://doi.org/110.1038/ngeo612>, 2009.
- Köhler, P., Abrams, J. F., Völker, C., Hauck, J., and Wolf-Gladrow, D. A.: Geoengineering impact of open ocean dissolution of olivine on atmospheric CO<sub>2</sub>, surface ocean pH and marine biology, *Environmental Research Letters*, 8, 014009, <https://doi.org/10.1088/1748-9326/8/1/014009>, 2013.
- 1375 Lacroix, F., Ilyina, T., and Hartmann, J.: Oceanic CO<sub>2</sub> outgassing and biological production hotspots induced by pre-industrial river loads of nutrients and carbon in a global modeling approach, *Biogeosciences*, 17, 55–88, <https://doi.org/10.5194/bg-17-55-2020>, 2020.
- Lacroix, F., Ilyina, T., Mathis, M., Laruelle, G. G., and Regnier, P.: Historical increases in land-derived nutrient inputs may alleviate effects of a changing physical climate on the oceanic carbon cycle, *Global Change Biology*, 27, 5491–5513, <https://doi.org/https://doi.org/10.1111/gcb.15822>, 2021.
- 1380 Landschützer, P., Gruber, N., and Bakker, D. C. E.: Decadal variations and trends of the global ocean carbon sink, *Global Biogeochemical Cycles*, 30, 1396–1417, <https://doi.org/https://doi.org/10.1002/2015GB005359>, 2016.
- Large, W. G. and Yeager, S. G.: Diurnal to decadal global forcing for ocean and sea-ice models : the datasets and flux climatologies., Tech. rep., CGD division of the National Center for Atmospheric Research, NCAR technical note : NCAR/TN-460+STR. Available on the GFDL CORE website., 2004.
- 1385 Large, W. G. and Yeager, S. G.: The global climatology of an interannually varying air-sea flux data set., *Clim Dyn*, 33, 341–364, <https://doi.org/10.1007/s00382-008-0441-3>, 2009.

- Large, W. G., McWilliams, J. C., and Doney, S. C.: Oceanic vertical mixing: A review and a model with a nonlocal boundary layer parameterization, *Reviews of Geophysics*, 32, <https://doi.org/10.1029/94RG01872>, 1994.
- 1390 Lauderdale, J. M. and Cael, B. B.: Impact of remineralization profile shape on the air-sea carbon balance, *Geophysical Research Letters*, 48, e2020GL091746., <https://doi.org/https://doi.org/10.1029/2020GL091746>, 2021.
- Laufkötter, C., Vogt, M., Gruber, N., Aita-Noguchi, M., Aumont, O., Bopp, L., Buitenhuis, E., Doney, S. C., Dunne, J., Hashioka, T., Hauck, J., Hirata, T., John, J., Le Quéré, C., Lima, I. D., Nakano, H., Seferian, R., Totterdell, I., Vichi, M., and Völker, C.: Drivers and uncertainties of future global marine primary production in marine ecosystem models, *Biogeosciences*, 12, 6955–6984, <https://doi.org/10.5194/bg-12-6955-2015>, 2015.
- 1395 Lauvset, S. K., Key, R. M., Olsen, A., van Heuven, S., Velo, A., Lin, X., Schirnick, C., Kozyr, A., Tanhua, T., Hoppema, M., Jutterström, S., Steinfeldt, R., Jeansson, E., Ishii, M., Perez, F. F., Suzuki, T., and Watelet, S.: A new global interior ocean mapped climatology: the  $1^\circ \times 1^\circ$  GLODAP version 2, *Earth System Science Data*, 8, 325–340, <https://doi.org/10.5194/essd-8-325-2016>, 2016.
- Le Quéré, C., Takahashi, T., Buitenhuis, E. T., Rödenbeck, C., and Sutherland, S. C.: Impact of climate change and variability on the global oceanic sink of CO<sub>2</sub>, *Global Biogeochemical Cycles*, 24, 1–10, <https://doi.org/10.1029/2009GB003599>, 2010.
- 1400 Lee, Z. and Marra, J. F.: The Use of VGPM to Estimate Oceanic Primary Production: A “Tango” Difficult to Dance, *Journal of Remote Sensing*, 2022, 9851 013, <https://doi.org/10.34133/2022/9851013>, 2022.
- Lenton, A., Tilbrook, B., Law, R. M., Bakker, D., Doney, S. C., Gruber, N., Ishii, M., Hoppema, M., Lovenduski, N. S., Matear, R. J., McNeil, B. I., Metzl, N., Mikaloff Fletcher, S. E., Monteiro, P. M. S., Rödenbeck, C., Sweeney, C., and Takahashi, T.: Sea–air CO<sub>2</sub> fluxes in the Southern Ocean for the period 1990–2009, *Biogeosciences*, 10, 4037–4054, <https://doi.org/10.5194/bg-10-4037-2013>, 2013.
- 1405 Lewis, K. M. and Arrigo, K. R.: Ocean Color Algorithms for Estimating Chlorophyll a, CDOM Absorption, and Particle Backscattering in the Arctic Ocean, *Journal of Geophysical Research: Oceans*, 125, e2019JC015706, <https://doi.org/https://doi.org/10.1029/2019JC015706>, e2019JC015706 2019JC015706, 2020.
- Lewis, K. M., van Dijken, G. L., and Arrigo, K. R.: Changes in phytoplankton concentration now drive increased Arctic Ocean primary production, *Science*, 369, 198–202, <https://doi.org/10.1126/science.aay8380>, 2020.
- 1410 Liao, E., Resplandy, L., Liu, J., and Bowman, K. W.: Amplification of the Ocean Carbon Sink During El Niños: Role of Poleward Ekman Transport and Influence on Atmospheric CO<sub>2</sub>, *Global Biogeochemical Cycles*, 34, e2020GB006574, <https://doi.org/https://doi.org/10.1029/2020GB006574>, e2020GB006574 2020GB006574, 2020.
- Long, M. C., Moore, J. K., Lindsay, K., Levy, M., Luo, J. Y., Krumhardt, K. M., Letscher, R. T., and Sylvester, Z. T.: Simulations with the Marine Biogeochemistry Library (MARBL), *Journal of Advances in Modeling Earth Systems*, 13, e2021MS002647, <https://doi.org/10.1029/2021MS002647>, 2021a.
- 1415 Long, M. C., Stephens, B. B., McKain, K., Sweeney, C., Keeling, R. F., Kort, E. A., Morgan, E. J., Bent, J. D., Chandra, N., Chevallier, F., Commane, R., Daube, B. C., Krummel, P. B., Loh, Z., Luijkx, I. T., Munro, D., Patra, P., Peters, W., Ramonet, M., Rödenbeck, C., Stavert, A., Tans, P., and Wofsy, S. C.: Strong Southern Ocean carbon uptake evident in airborne observations, *Science*, 374, 1275–1280, <https://doi.org/10.1126/science.abi4355>, 2021b.
- 1420 Losch, M., Strass, V., B. Cisewski, C. K., and Bellerby, R. G.: Ocean state estimation from hydrography and velocity observations during EIFEX with a regional biogeochemical ocean circulation model, *J. Mar. Sys.*, 129, 437–451, <https://doi.org/10.1016/j.jmarsys.2013.09.003>, 2014.
- Lévy, M., Franks, P. J. S., and Smith, K. S.: The role of submesoscale currents in structuring marine ecosystems, *Nature Communications*, 9, 47–58, <https://doi.org/10.1038/s41467-018-07059-3>, 2018.

- 1425 Maerz, J., Six, K. D., Stemmler, I., Ahmerkamp, S., and Ilyina, T.: Microstructure and composition of marine aggregates as co-determinants for vertical particulate organic carbon transfer in the global ocean, *Biogeosciences*, 17, 1765–1803, <https://doi.org/10.5194/bg-17-1765-2020>, 2020.
- Mahowald, N., Luo, C., del Corral, J., and Zender, C. S.: Interannual variability in atmospheric mineral aerosols from a 22-year model simulation and observational data, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/https://doi.org/10.1029/2002JD002821>, 1430 2003.
- Maier-Reimer, E., Mikolajewicz, U., and Winguth, A.: Future ocean uptake of CO<sub>2</sub>: interaction between ocean circulation and biology, *Climate Dynamics*, 12, 711–722, <https://doi.org/10.1007/s003820050138>, 1996.
- Maritorena, S., Siegel, D. A., and Peterson, A. R.: Optimization of a semianalytical ocean color model for global-scale applications, *Appl. Opt.*, 41, 2705–2714, <https://doi.org/10.1364/AO.41.002705>, 2002.
- 1435 Marshall, J., Adcroft, A., Hill, C., Perelman, L., and Heisey, C.: A finite-volume, incompressible Navier Stokes model for studies of the ocean on parallel computers, *Journal of Geophysical Research*, 102, 5753–5766, 1997.
- Matsuoka, A., Bricaud, A., Benner, R., Para, J., Sempéré, R., Prieur, L., Bélanger, S., and Babin, M.: Tracing the transport of colored dissolved organic matter in water masses of the Southern Beaufort Sea: relationship with hydrographic characteristics, *Biogeosciences*, 9, 925–940, <https://doi.org/10.5194/bg-9-925-2012>, 2012.
- 1440 McWilliams, J. C.: Submesoscale currents in the ocean, *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 472, 20160 117, <https://doi.org/10.1098/rspa.2016.0117>, 2016.
- Metzl, N., Brunet, C., Jabaud-Jan, A., Poisson, A., and Schauer, B.: Summer and winter air–sea CO<sub>2</sub> fluxes in the Southern Ocean, *Deep Sea Research Part I: Oceanographic Research Papers*, 53, 1548–1563, <https://doi.org/https://doi.org/10.1016/j.dsr.2006.07.006>, 2006.
- Misumi, K., Tsumune, D., Yoshida, Y., Uchimoto, K., Nakamura, T., Nishioka, J., Mitsudera, H., Bryan, F. O., Lindsay, K., Moore, J. K., and 1445 Doney, S. C.: Mechanisms controlling dissolved iron distribution in the North Pacific: A model study, *Journal of Geophysical Research*, 116, 1–17, <https://doi.org/10.1029/2010JG001541>, 2011.
- Mitchell, B.: Predictive bio-optical relationships for polar oceans and marginal ice zones, *Journal of Marine Systems*, 3, 91–105, [https://doi.org/https://doi.org/10.1016/0924-7963\(92\)90032-4](https://doi.org/https://doi.org/10.1016/0924-7963(92)90032-4), 1992.
- Mongwe, N. P., Vichi, M., and Monteiro, P. M. S.: The seasonal cycle of *p*CO<sub>2</sub> and CO<sub>2</sub> fluxes in the Southern Ocean: diagnosing anomalies 1450 in CMIP5 Earth system models, *Biogeosciences*, 15, 2851–2872, <https://doi.org/10.5194/bg-15-2851-2018>, 2018.
- Moore, C. M., Mills, M. M., Arrigo, K. R., Berman-Frank, I., Bopp, L., Boyd, P. W., Galbraith, E. D., Geider, R. J., Guieu, C., Jaccard, S. L., Jickells, T. D., La Roche, J., Lenton, T. M., Mahowald, N. M., Marañón, E., Marinov, I., Moore, J. K., Nakatsuka, T., Oschlies, A., Saito, M. A., Thingstad, T. F., Tsuda, A., and Ulloa, O.: Processes and patterns of oceanic nutrient limitation, *Nature Geoscience*, 6, 701–710, <https://doi.org/10.1038/ngeo1765>, 2013.
- 1455 Moriarty, R. and O’Brien, T. D.: Distribution of mesozooplankton biomass in the global ocean, *Earth System Science Data*, 5, 45–55, <https://doi.org/10.5194/essd-5-45-2013>, 2013.
- Moriarty, R., Buitenhuis, E. T., and Le Quééré, C.: Distribution of known macrozooplankton abundance and biomass in the global ocean, *Earth System Science Data*, 5, 241–257, <https://doi.org/10.5194/essd-5-241-2013>, 2013.
- Munhoven, G.: Mathematics of the total alkalinity–pH equation - pathway to robust and universal solution algorithms: the SolveSAPHE 1460 package v1.0.1, *Geoscientific Model Development*, 6, 1367–1388, <https://doi.org/10.5194/gmd-6-1367-2013>, 2013.

- Mustapha, S. B., Bélanger, S., and Larouche, P.: Evaluation of ocean color algorithms in the southeastern Beaufort Sea, Canadian Arctic: New parameterization using SeaWiFS, MODIS, and MERIS spectral bands, *Canadian Journal of Remote Sensing*, 38, 535–556, <https://doi.org/10.5589/m12-045>, 2012.
- 1465 Myriokefalitakis, S., Ito, A., Kanakidou, M., Nenes, A., Krol, M. C., Mahowald, N. M., Scanza, R. A., Hamilton, D. S., Johnson, M. S., Meskhidze, N., Kok, J. F., Guieu, C., Baker, A. R., Jickells, T. D., Sarin, M. M., Bikkina, S., Shelley, R., Bowie, A., Perron, M. M. G., and Duce, R. A.: Reviews and syntheses: the GESAMP atmospheric iron deposition model intercomparison study, *Biogeosciences*, 15, 6659–6684, <https://doi.org/10.5194/bg-15-6659-2018>, 2018.
- 1470 Nakano, H., Tsujino, H., Hirabara, M., Yasuda, T., Motoi, T., Ishii, M., and Yamanaka, G.: Uptake mechanism of anthropogenic CO<sub>2</sub> in the Kuroshio Extension region in an ocean general circulation model, *Journal of Oceanography*, 67, 765–783, <https://doi.org/10.1007/s10872-011-0075-7>, 2011.
- Nevison, C. D., Keeling, R. F., Kahru, M., Manizza, M., Mitchell, B. G., and Cassar, N.: Estimating net community production in the Southern Ocean based on atmospheric potential oxygen and satellite ocean color data, *Global Biogeochem. Cycles*, 26, GB1020, <https://doi.org/10.1029/2011GB004040>, 2012.
- 1475 Nielsdóttir, M. C., Moore, C. M., Sanders, R., Hinz, D. J., and Achterberg, E. P.: Iron limitation of the postbloom phytoplankton communities in the Iceland Basin, *Global Biogeochemical Cycles*, 23, 1–13, <https://doi.org/10.1029/2008GB003410>, 2009.
- Nissen, C., Timmermann, R., Hoppema, M., Gürses, Ö., and Hauck, J.: Abruptly attenuated carbon sequestration with Weddell Sea dense waters by 2100, *Nature Communications*, 13, 3402, <https://doi.org/10.1038/s41467-022-30671-3>, 2022.
- Orr, J. C.: On ocean carbon-cycle model comparison, *Tellus B: Chemical and Physical Meteorology*, 51, 509–510, <https://doi.org/10.3402/tellusb.v51i2.16334>, 1999.
- 1480 Orr, J. C. and Epitalon, J.-M.: Improved routines to model the ocean carbonate system: mocsy 2.0, *Geoscientific Model Development*, 8, 485–499, <https://doi.org/10.5194/gmd-8-485-2015>, 2015.
- Oziel, L., Schourup-Kristensen, V., Wekerle, C., and Hauck, J.: The pan-Arctic continental slope as an intensifying conveyor belt for nutrients in the central Arctic Ocean (1985–2015), *Global Biogeochemical Cycles*, p. e2021GB007268, <https://doi.org/10.1029/2021GB007268>, 2022.
- 1485 Pagnone, A., Völker, C., and Ye, Y.: Processes affecting dissolved iron across the Subtropical North Atlantic: a model study, *Ocean Dynamics*, 69, 989–1007, <https://doi.org/10.1007/s10236-019-01288-w>, 2019.
- Parekh, P., Follows, M. J., and E., B.: Modeling the global ocean iron cycle, *Global Biogeochem. Cycles*, 18, GB1002, <https://doi.org/doi:10.1029/2003GB002061>, 2004.
- 1490 Pradhan, H. K., Völker, C., Losa, S. N., Bracher, A., and Nerger, L.: Assimilation of Global Total Chlorophyll OC-CCI Data and Its Impact on Individual Phytoplankton Fields, *Journal of Geophysical Research: Oceans*, 124, 470–490, <https://doi.org/10.1029/2018JC014329>, 2019.
- Prowe, A. F., Pahlow, M., Dutkiewicz, S., Follows, M., and Oschlies, A.: Top-down control of marine phytoplankton diversity in a global ecosystem model, *Progress in Oceanography*, 101, 1–13, <https://doi.org/https://doi.org/10.1016/j.pocean.2011.11.016>, 2012.
- Raven, J. A.: The iron and molybdenum use efficiencies of plant growth with different energy, carbon and nitrogen sources, *New Phytologist*, 109, 279–287, 1988.
- 1495 Redfield, A., Ketchum, B., and Richards, F.: The influence of organisms on the composition of sea water, in: *The Sea*, edited by Hill, M., vol. 2, pp. 26–77, Interscience, 1963.
- Redi, M.: Oceanic Isopycnal Mixing by Coordinate Rotation, *J. Phys. Oceanogr.*, 12, 1154–1158, 1982.



- Regnier, P., Resplandy, L., Najjar, R. G., and Ciais, P.: The land-to-ocean loops of the global carbon cycle, *Nature*, 603, 401–410, <https://doi.org/10.1038/s41586-021-04339-9>, 2022.
- 1500 Resplandy, L., Hogikyan, A., Bange, H. W., Bianchi, D., Weber, T. S., Cai, W.-J., Doney, S. C., Fennel, K., Gehlen, M., Hauck, J., Lacroix, F., Landschützer, P., Quéré, C. L., Müller, J. D., Najjar, R. G., Roobaert, A., Berthet, S., Bopp, L., Chau, T. T.-T., Dai, M., Gruber, N., Ilyina, T., Kock, A., Manizza, M., Lachkar, Z., Laruelle, G. G., Liao, E., Lima, I. D., Nissen, C., Rödenbeck, C., Séférian, R., Schwinger, J., Toyama, K., Tsujino, H., and Regnier, P.: A Synthesis of Global Coastal Ocean Greenhouse Gas Fluxes, <https://doi.org/10.22541/essoar.168182303.39621839/v1>, 2023.
- 1505 Resplandy, L., Keeling, R. F., Roedenbeck, C., Stephens, B. B., Khatiwala, S., Rodgers, K. B., Long, M. C., Bopp, L., and Tans, P. P.: Revision of global carbon fluxes based on a reassessment of oceanic and riverine carbon transport, *Nature Geoscience*, 11, 504–509, <https://doi.org/10.1038/s41561-018-0151-3>, 2018.
- Rödenbeck, C., DeVries, T., Hauck, J., Le Quéré, C., and Keeling, R. F.: Data-based estimates of interannual sea–air CO<sub>2</sub> flux variations 1957–2020 and their relation to environmental drivers, *Biogeosciences*, 19, 2627–2652, <https://doi.org/10.5194/bg-19-2627-2022>, 2022.
- 1510 Rohr, T., Richardson, A. J., Lenton, A., and Shadwick, E.: Recommendations for the formulation of grazing in marine biogeochemical and ecosystem models, *Progress in Oceanography*, 208, 102 878, <https://doi.org/10.1016/j.pocean.2022.102878>, 2022.
- Sabine, C. L., Feely, R. A., Gruber, N., Key, R. M., Lee, K., Bullister, J. L., Wanninkhof, R., Wong, C. S., Wallace, D. W. R., Tilbrook, B., Millero, F. J., Peng, T.-H., Kozyr, A., Ono, T., and Rios, A. F.: The Oceanic Sink for Anthropogenic CO<sub>2</sub>, *Science*, 305, 367–371, <https://doi.org/10.1126/science.1097403>, 2004.
- 1515 Sallée, J.-B., Pellichero, V., Akhouldas, C., Pauthenet, E., Vignes, L., Schmidtko, S., Garabato, A. N., Sutherland, P., and Kuusela, M.: Summertime increases in upper-ocean stratification and mixed-layer depth, *Nature*, 591, 592–598, <https://doi.org/10.1038/s41586-021-03303-x>, 2021.
- Sarmiento, J. L. and Gruber, N.: *Ocean Biogeochemical Dynamics*, Princeton University Press, Princeton, NJ, 2006.
- Sathyendranath, S., Brewin, R. J., Brockmann, C., Brotas, V., Calton, B., Chuprin, A., Cipollini, P., Couto, A. B., Dingle, J., Doerffer, R., 1520 Donlon, C., Dowell, M., Farman, A., Grant, M., Groom, S., Horseman, A., Jackson, T., Krasemann, H., Lavender, S., Martinez-Vicente, V., Mazeran, C., Mélin, F., Moore, T. S., Müller, D., Regner, P., Roy, S., Steele, C. J., Steinmetz, F., Swinton, J., Taberner, M., Thompson, A., Valente, A., Zühlke, M., Brando, V. E., Feng, H., Feldman, G., Franz, B. A., Frouin, R., Gould, R. W., Hooker, S. B., Kahru, M., Kratzer, S., Mitchell, B. G., Muller-Karger, F. E., Sosik, H. M., Voss, K. J., Werdell, J., and Platt, T.: An Ocean-Colour Time Series for Use in Climate Studies: The Experience of the Ocean-Colour Climate Change Initiative (OC-CCI), *Sensors*, 19, <https://doi.org/10.3390/s19194285>, 2019.
- 1525 Scaife, A. A., Copsey, D., Gordon, C., Harris, C., Hinton, T., Keeley, S., O'Neill, A., Roberts, M., and Williams, K.: Improved Atlantic winter blocking in a climate model, *Geophysical Research Letters*, 38, <https://doi.org/https://doi.org/10.1029/2011GL049573>, 2011.
- Schartau, M., Engel, A., Schröter, J., Thoms, S., Völker, C., and Wolf-Gladrow, D.: Modelling carbon overconsumption and the formation of extracellular particulate organic carbon, *Biogeosciences*, 4, 433–454, <https://doi.org/10.5194/bg-4-433-2007>, 2007.
- Schlitzer, R.: Carbon export fluxes in the Southern Ocean: results from inverse modeling and comparison with satellite-based estimates, *Deep 1530 Sea Research Part II: Topical Studies in Oceanography*, 49, 1623–1644, [https://doi.org/https://doi.org/10.1016/S0967-0645\(02\)00004-8](https://doi.org/https://doi.org/10.1016/S0967-0645(02)00004-8), the Southern Ocean I: Climatic Changes in the Cycle of Carbon in the Southern Ocean, 2002.
- Schlitzer, R.: Export Production in the Equatorial and North Pacific Derived from Dissolved Oxygen, Nutrient and Carbon Data, *Journal of Oceanography*, 60, 53–62, <https://doi.org/10.1023/B:JOCE.0000038318.38916.e6>, 2004.
- Schlitzer, R., Anderson, R. F., Dodas, E. M., Lohan, M. C., Geibert, W., Tagliabue, A., Bowie, A. R., Jeandel, C., Maldonado, M. T., Landing, 1535 W. M., Cockwell, D., Abadie, C., Abouchami, W., Achterberg, E. P., Agather, A. M., Aguilar-Islas, A. M., van Aken, H. M., Andersen,

- M., Archer, C., Auro, M., de Baar, H. J. W., Baars, O., Baker, A. R., Bakker, K., Basak, C., Baskaran, M., Bates, N. R., Bauch, D., van Beek, P., Behrens, M. K., Black, E., Bluhm, K., Bopp, L., Bouman, H. A., Bowman, K., Bown, J., Boyd, P. W., Boye, M., Boyle, E. A., Branellec, P., Bridgestock, L., Brissebrat, G., Browning, T. J., Bruland, K. W., Brumsack, H. J., Brzezinski, M. A., Buck, C. S., Buck, K. N., Buesseler, K. O., Bull, A., Butler, E., Cai, P., Mor, P. C., Cardinal, D., Carlson, C., Carrasco, G., Casacuberta, N., Casciotti, K. L., 1540 Castrillejo, M., Chamizo, E., Chance, R., Charette, M. A., Chaves, J. E., Cheng, H., Chever, F., Christl, M., Church, T. M., Closset, I., Colman, A., Conway, T. M., Cossa, D., Croot, P. L., Cullen, J. T., Cutter, G. A., Daniels, C., Dehairs, F., Deng, F., Dieu, H. T., Duggan, B., Dulaquais, G., Dumousseaud, C., Echegoyen-Sanz, Y., Edwards, R. L., Ellwood, M., Fahrbach, E., Fitzsimmons, J. N., Russell Flegal, A., Fleisher, M. Q., van de Flierdt, T., Frank, M., Friedrich, J., Fripiat, F., Fröllje, H., Galer, S. J., Gamo, T., Ganeshram, R. S., Garcia-Orellana, J., Garcia-Solsona, E., Gault-Ringold, M., George, E., Gerringa, L. J., Gilbert, M., Godoy, J. M., Goldstein, S. L., Gonzalez, 1545 S. R., Grissom, K., Hammerschmidt, C., Hartman, A., Hassler, C. S., Hathorne, E. C., Hatta, M., Hawco, N., Hayes, C. T., Heimbürger, L. E., Helgoe, J., Heller, M., Henderson, G. M., Henderson, P. B., van Heuven, S. M. A. C., Ho, P., Horner, T. J., Hsieh, Y. T., Huang, K. F., Humphreys, M. P., Isshiki, K., Jacquot, J. E., Janssen, D. J., Jenkins, W. J., John, S., Jones, E. M., Jones, J. L., Kadko, D. C., Kayser, R., Kenna, T. C., Khondoker, R., Kim, T., Kipp, L., Klar, J. K., Klunder, M. B., Kretschmer, S., Kumamoto, Y., Laan, P., Labatut, M., Lacan, F., Lam, P. J., Lambelet, M., Lamborg, C. H., Le Moigne, F. A., Le Roy, E., Lechtenfeld, O. J., Lee, J. M., Lherminier, P., Little, 1550 S., López-Lora, M., Lu, Y., Masque, P., Mawji, E., McClain, C. R., Measures, C., Mehic, S., Barraqueta, J. L. M., van der Merwe, P., Middag, R., Mieruch, S., Milne, A., Minami, T., Moffett, J. W., Moncoiffe, G., Moore, W. S., Morris, P. J., Morton, P. L., Nakaguchi, Y., Nakayama, N., Niedermiller, J., Nishioka, J., Nishiuchi, A., Noble, A. E., Obata, H., Ober, S., Ohnemus, D. C., van Ooijen, J., O'Sullivan, J., Owens, S., Pahnke, K., Paul, M., Pavia, F., Pena, L. D., Peters, B. D., Planchon, F., Planquette, H. F., Pradoux, C., Puigcorbé, V., Quay, P., Queroue, F., Radic, A., Rauschenberg, S., Rehkämper, M., Rember, R., Remenyi, T. A., Resing, J. A., Rickli, J., Rigaud, S., 1555 Rijkenberg, M. J. A., Rintoul, S., Robinson, L. F., Roca-Martí, M., Rodellas, V., Roeske, T., Rolison, J. M., Rosenberg, M., Roshan, S., Rutgers van der Loeff, M. M., Ryabenko, E., Saito, M. A., Salt, L. A., Sanial, V., Sarthou, G., Schallenberg, C., Schauer, U., Scher, H., Schlosser, C., Schnetger, B., Scott, P., Sedwick, P. N., Semiletov, I., Shelley, R., Sherrell, R. M., Shiller, A. M., Sigman, D. M., Singh, S. K., Slagter, H. A., Slater, E., Smethie, W. M., Snaith, H., Sohrin, Y., Sohst, B., Sonke, J. E., Speich, S., Steinfeldt, R., Stewart, G., Stichel, T., Stirling, C. H., Stutsman, J., Swarr, G. J., Swift, J. H., Thomas, A., Thorne, K., Till, C. P., Till, R., Townsend, A. T., Townsend, 1560 E., Tuerena, R., Twining, B. S., Vance, D., Velazquez, S., Venchiarutti, C., Villa-Alfageme, M., Vivancos, S. M., Voelker, A. H., Wake, B. D., Warner, M. J., Watson, R., van Weerlee, E., Alexandra Weigand, M., Weinstein, Y., Weiss, D. J., Wisotzki, A., Woodward, E. M. S., Wu, J., Wu, Y., Wuttig, K., Wyatt, N. J., Xiang, Y., Xie, R. C., Xue, Z., Yoshikawa, H., Zhang, J., Zhang, P., Zhao, Y., Zheng, L., Zheng, X. Y., Zieringer, M., Zimmer, L. A., Ziveri, P., Zunino, P., and Zurbrick, C.: The GEOTRACES Intermediate Data Product 2017, *Chemical Geology*, 493, 210–223, <https://doi.org/10.1016/j.chemgeo.2018.05.040>, publisher: Elsevier, 2018.
- 1565 Schmidtko, S., Stramma, L., and Visbeck, M.: Decline in global oceanic oxygen content during the past five decades, *Nature*, 542, 335–339, <https://doi.org/10.1038/nature21399>, 2017.
- Schneider, B., Bopp, L., Gehlen, M., Segschneider, J., Frölicher, T. L., Cadule, P., Friedlingstein, P., Doney, S. C., Behrenfeld, M. J., and Joos, F.: Climate-induced interannual variability of marine primary and export production in three global coupled climate carbon cycle models, *Biogeosciences*, 5, 597–614, <https://doi.org/10.5194/bg-5-597-2008>, 2008.
- 1570 Scholz, P., Sidorenko, D., Gurses, O., Danilov, S., Koldunov, N., Wang, Q., Sein, D., Smolentseva, M., Rakowsky, N., and Jung, T.: Assessment of the Finite-volume Sea ice-Ocean Model (FESOM2.0) – Part 1: Description of selected key model elements and comparison to its predecessor version, *Geoscientific Model Development*, 12, 4875–4899, <https://doi.org/10.5194/gmd-12-4875-2019>, 2019.

- Scholz, P., Sidorenko, D., Danilov, S., Wang, Q., Koldunov, N., Sein, D., and Jung, T.: Assessment of the Finite Volume Sea Ice Ocean Model (FESOM2.0), Part II: Partial bottom cells, embedded sea ice and vertical mixing library CVMIX, *Geoscientific Model Development Discussions*, 2021, 1–42, <https://doi.org/10.5194/gmd-2021-94>, 2021.
- Schourup-Kristensen, V., Sidorenko, D., Wolf-Gladrow, D. A., and Völker, C.: A skill assessment of the biogeochemical model REcoM2 coupled to the Finite Element Sea-Ice Ocean Model (FESOM 1.3), *Geosci. Model Dev.*, 7, 2769–2802, <https://doi.org/10.5194/gmd-7-2769-2014>, 2014.
- Schourup-Kristensen, V., Wekerle, C., Wolf-Gladrow, D. A., and Völker, C.: Arctic Ocean biogeochemistry in the high resolution FESOM 1.4-REcoM2 model, *Progress in Oceanography*, 168, 65–81, <https://doi.org/10.1016/j.pocean.2018.09.006>, 2018.
- Schwinger, J., Goris, N., Tjiputra, J. F., Kriest, I., Bentsen, M., Bethke, I., Ilicak, M., Assmann, K. M., and Heinze, C.: Evaluation of NorESM-OC (versions 1 and 1.2), the ocean carbon-cycle stand-alone configuration of the Norwegian Earth System Model (NorESM1), *Geoscientific Model Development*, 9, 2589–2622, <https://doi.org/10.5194/gmd-9-2589-2016>, 2016.
- Séférián, R., Gehlen, M., Bopp, L., Resplandy, L., Orr, J. C., Marti, O., Dunne, J. P., Christian, J. R., Doney, S. C., Ilyina, T., Lindsay, K., Halloran, P. R., Heinze, C., Segsneider, J., Tjiputra, J., Aumont, O., and Romanou, A.: Inconsistent strategies to spin up models in CMIP5: implications for ocean biogeochemical model performance assessment, *Geoscientific Model Development*, 9, 1827–1851, <https://doi.org/10.5194/gmd-9-1827-2016>, 2016.
- Seifert, M., Nissen, C., Rost, B., and Hauck, J.: Cascading effects augment the direct impact of CO<sub>2</sub> on phytoplankton growth in a biogeochemical model, *Elementa: Science of the Anthropocene*, 10, <https://doi.org/10.1525/elementa.2021.00104>, 00104, 2022.
- Sidorenko, D., Wang, Q., Danilov, S., and Schröter, J.: FESOM under coordinated ocean-ice reference experiment forcing, *Ocean Dynamics*, 61, 881–890, <https://doi.org/10.1007/s10236-011-0406-7>, 2011.
- Sidorenko, D., Rackow, T., Jung, T., Semmler, T., Barbi, D., Danilov, S., Dethloff, K., Dorn, W., Fieg, K., Gößling, H. F., et al.: Towards multi-resolution global climate modeling with ECHAM6–FESOM. Part I: model formulation and mean climate, *Climate Dynamics*, 44, 757–780, <https://doi.org/10.1007/s00382-014-2290-6>, 2015.
- Siegel, D. A., Buesseler, K. O., Doney, S. C., Saille, S. F., Behrenfeld, M. J., and Boyd, P. W.: Global assessment of ocean carbon export by combining satellite observations and food-web models, *Global Biogeochem. Cycles*, 28, 181–196, <https://doi.org/doi:10.1002/2013GB004743>, 2014.
- Steele, M., Morley, R., and Ermold, W.: PHC: a global ocean hydrography with a high-quality Arctic Ocean, *J. Climate*, 14, 2079–2087, [https://doi.org/10.1175/1520-0442\(2001\)014<2079:PAGOHW>2.0.CO;2](https://doi.org/10.1175/1520-0442(2001)014<2079:PAGOHW>2.0.CO;2), 2001.
- Stewart, K., Kim, W., Urakawa, S., Hogg, A., Yeager, S., Tsujino, H., Nakano, H., Kiss, A., and Danabasoglu, G.: JRA55-do-based repeat year forcing datasets for driving ocean–sea-ice models, *Ocean Modelling*, 147, 101557, <https://doi.org/https://doi.org/10.1016/j.ocemod.2019.101557>, 2020.
- Storkey, D., Blaker, A. T., Mathiot, P., Megann, A., Aksenov, Y., Blockley, E. W., Calvert, D., Graham, T., Hewitt, H. T., Hyder, P., Kuhlbrodt, T., Rae, J. G. L., and Sinha, B.: UK Global Ocean GO6 and GO7: a traceable hierarchy of model resolutions, *Geoscientific Model Development*, 11, 3187–3213, <https://doi.org/10.5194/gmd-11-3187-2018>, 2018.
- Sundquist, E. T.: Geological Perspectives on Carbon Dioxide and the Carbon Cycle, pp. 55–59, American Geophysical Union (AGU), <https://doi.org/https://doi.org/10.1029/GM032p0005>, 1985.
- Sutton, A. J., Williams, N. L., and Tilbrook, B.: Constraining Southern Ocean CO<sub>2</sub> flux uncertainty using uncrewed surface vehicle observations., *Geophysical Research Letters*, 48, e2020GL091748, <https://doi.org/10.1029/2020GL091748>, 2021.

- 1610 Séférian, R., Berthet, S., Yool, A., Palmiéri, J., Bopp, L., Tagliabue, A., Kwiatkowski, L., Aumont, O., Christian, J., Dunne, J., Gehlen, M., Ilyina, T., John, J. G., Li, H., Long, M. C., Luo, J. Y., Nakano, H., Romanou, A., Schwinger, J., Stock, C., Santana-Falcón, Y., Takano, Y., Tjiputra, J., Tsujino, H., Watanabe, M., Wu, T., Wu, F., and Yamamoto, A.: Tracking Improvement in Simulated Marine Biogeochemistry Between CMIP5 and CMIP6, *Current Climate Change Reports*, 6, 95–119, <https://doi.org/10.1007/s40641-020-00160-0>, 2020.
- Tagliabue, A. and Völker, C.: Towards accounting for dissolved iron speciation in global ocean models, *Biogeosciences*, 8, 3025–3039, <https://doi.org/10.5194/bg-8-3025-2011>, 2011.
- 1615 Tagliabue, A., Mtshali, T., Aumont, O., Bowie, A. R., Klunder, M. B., Roychoudhury, A. N., and Swart, S.: A global compilation of dissolved iron measurements: focus on distributions and processes in the Southern Ocean, *Biogeosciences*, 9, 1–17, <https://doi.org/10.5194/bg-9-1-2012>, 2012.
- Tagliabue, A., Aumont, O., DeAth, R., Dunne, J. P., Dutkiewicz, S., Galbraith, E., Misumi, K., Moore, J. K., Ridgwell, A. J., Sherman, E., Stock, C. A., Vichi, M., Völker, C., and Yool, A.: How well do global ocean biogeochemistry models simulate dissolved iron distributions?, *Global Biogeochemical Cycles*, 30, <https://doi.org/10.1002/2015GB005289>, 2016.
- 1620 Tagliabue, A., Kwiatkowski, L., Bopp, L., Butenschön, M., Cheung, W., Lengaigne, M., and Vialard, J.: Persistent Uncertainties in Ocean Net Primary Production Climate Change Projections at Regional Scales Raise Challenges for Assessing Impacts on Ecosystem Services, *Frontiers in Climate*, 3, 738 224, <https://doi.org/10.3389/fclim.2021.738224>, 2021.
- 1625 Takahashi, T., Sutherland, S. C., Wanninkhof, R., Sweeney, C., Feely, R. A., Chipman, D. W., Hales, B., Friederich, G., Chavez, F., Sabine, C., Watson, A., Bakker, D. C., Schuster, U., Metzl, N., Yoshikawa-Inoue, H., Ishii, M., Midorikawa, T., Nojiri, Y., Körtzinger, A., Steinhoff, T., Hoppema, M., Olafsson, J., Arnarson, T. S., Tilbrook, B., Johannessen, T., Olsen, A., Bellerby, R., Wong, C., Delille, B., Bates, N., and de Baar, H. J.: Climatological mean and decadal change in surface ocean pCO<sub>2</sub>, and net sea–air CO<sub>2</sub> flux over the global oceans, *Deep Sea Research Part II: Topical Studies in Oceanography*, 56, 554–577, <https://doi.org/10.1016/j.dsr2.2008.12.009>, 2009.
- 1630 Taylor, M. H., Losch, M., and Bracher, A.: On the drivers of phytoplankton blooms in the Antarctic marginal ice zone: A modeling approach, *Journal of Geophysical Research*, 118, 63–75, <https://doi.org/doi:10.1029/2012JC008418>, 2013.
- Terhaar, J., Frölicher, T. L., and Joos, F.: Southern Ocean anthropogenic carbon sink constrained by sea surface salinity, *Science Advances*, 7, eabd5964, <https://doi.org/10.1126/sciadv.abd5964>, 2021.
- Terhaar, J., Frölicher, T. L., and Joos, F.: Observation-constrained estimates of the global ocean carbon sink from Earth system models, *Biogeosciences*, 19, 4431–4457, <https://doi.org/10.5194/bg-19-4431-2022>, 2022.
- 1635 Timmermann, R. and Beckmann, A.: Parameterization of vertical mixing in the Weddell Sea, *Ocean Modelling*, 6, 83–100, [https://doi.org/10.1016/S1463-5003\(02\)0061-6](https://doi.org/10.1016/S1463-5003(02)0061-6), 2004.
- Timmermann, R., Danilov, S., Schröter, J., Böning, C., Sidorenko, D., and Rollenhagen, K.: Ocean circulation and sea ice distribution in a finite element global sea ice–ocean model, *Ocean Modelling*, 27, 114–129, <https://doi.org/https://doi.org/10.1016/j.ocemod.2008.10.009>, 2009.
- 1640 Timmermann, R., Wang, Q., and Hellmer, H.: Ice-shelf basal melting in a global finite-element sea-ice/ice-shelf/ocean model, *Annals of Glaciology*, 53, 303–314, <https://doi.org/10.3189/2012AoG60A156>, 2012.
- Tjiputra, J. F., Schwinger, J., Bentsen, M., Morée, A. L., Gao, S., Bethke, I., Heinze, C., Goris, N., Gupta, A., He, Y.-C., Olivié, D., Seland, Ø., and Schulz, M.: Ocean biogeochemistry in the Norwegian Earth System Model version 2 (NorESM2), *Geoscientific Model Development*, 13, 2393–2431, <https://doi.org/10.5194/gmd-13-2393-2020>, 2020.
- 1645

- Tohjima, Y., Mukai, H., Machida, T., Hoshina, Y., and Nakaoka, S.-I.: Global carbon budgets estimated from atmospheric O<sub>2</sub>/N<sub>2</sub> and CO<sub>2</sub> observations in the western Pacific region over a 15-year period, *Atmospheric Chemistry and Physics*, 19, 9269–9285, <https://doi.org/10.5194/acp-19-9269-2019>, 2019.
- 1650 Tréguer, P. J., Sutton, J. N., Brzezinski, M., Charette, M. A., Devries, T., Dutkiewicz, S., Ehlert, C., Hawkings, J., Leynaert, A., Liu, S. M., Llopis Monferrer, N., López-Acosta, M., Maldonado, M., Rahman, S., Ran, L., and Rouxel, O.: Reviews and syntheses: The biogeochemical cycle of silicon in the modern ocean, *Biogeosciences*, 18, 1269–1289, <https://doi.org/10.5194/bg-18-1269-2021>, 2021.
- Tsujino, H., Urakawa, S., Nakano, H., Small, R. J., Kim, W. M., Yeager, S. G., Danabasoglu, G., Suzuki, T., Bamber, J. L., Bentsen, M., Böning, C. W., Bozec, A., Chassignet, E. P., Curchitser, E., Boeira Dias, F., Durack, P. J., Griffies, S. M., Harada, Y., Ilıcak, M., Josey, S. A., Kobayashi, C., Kobayashi, S., Komuro, Y., Large, W. G., Le Sommer, J., Marsland, S. J., Masina, S., Scheinert, M., Tomita, H., 1655 Valdivieso, M., and Yamazaki, D.: JRA-55 based surface dataset for driving ocean–sea-ice models (JRA55-do), *Ocean Modelling*, 130, 79–139, <https://doi.org/https://doi.org/10.1016/j.ocemod.2018.07.002>, 2018.
- Urakawa, L. S., Tsujino, H., Nakano, H., Sakamoto, K., Yamanaka, G., and Toyoda, T.: The sensitivity of a depth-coordinate model to diapycnal mixing induced by practical implementations of the isopycnal tracer diffusion scheme, *Ocean Modelling*, 154, 101 693, <https://doi.org/https://doi.org/10.1016/j.ocemod.2020.101693>, 2020.
- 1660 Vaittinada Ayar, P., Bopp, L., Christian, J. R., Ilyina, T., Krasting, J. P., Séférian, R., Tsujino, H., Watanabe, M., Yool, A., and Tjiputra, J.: Contrasting projections of the ENSO-driven CO<sub>2</sub> flux variability in the equatorial Pacific under high-warming scenario, *Earth System Dynamics*, 13, 1097–1118, <https://doi.org/10.5194/esd-13-1097-2022>, 2022.
- Völker, C. and Tagliabue, A.: Modeling organic iron-binding ligands in a three-dimensional biogeochemical ocean model, *Marine Chemistry*, 173, 67–77, <https://doi.org/10.1016/j.marchem.2014.11.008>, 2015.
- 1665 Völker, C. and Köhler, P.: Responses of ocean circulation and carbon cycle to changes in the position of the Southern Hemisphere westerlies at Last Glacial Maximum, *Paleoceanography*, 28, 726–739, <https://doi.org/https://doi.org/10.1002/2013PA002556>, 2013.
- Waite, A. M., Thompson, P. A., and Harrison, P. J.: Does energy control the sinking rates of marine diatoms?, *Limnology and Oceanography*, 37, 468–477, <https://doi.org/https://doi.org/10.4319/lo.1992.37.3.0468>, 1992.
- Wang, Q., Danilov, S., Sidorenko, D., Timmermann, R., Wekerle, C., Wang, X., Jung, T., and Schröter, J.: The Finite Element Sea Ice-Ocean 1670 Model (FESOM) v.1.4: formulation of an ocean general circulation model, *Geosci. Model Dev.*, 7, 663–693, <https://doi.org/10.5194/gmd-7-663-2014>, 2014.
- Wanninkhof, R.: Relationship between wind speed and gas exchange over the ocean revisited, *Limnology and Oceanography: Methods*, 12, 351–362, <https://doi.org/10.4319/lom.2014.12.351>, 2014.
- Wanninkhof, R., Park, G. H., Takahashi, T., Sweeney, C., Feely, R., Nojiri, Y., Gruber, N., Doney, S. C., McKinley, G. A., Lenton, A., 1675 Le Quéré, C., Heinze, C., Schwinger, J., Graven, H., and Khatiwala, S.: Global ocean carbon uptake: magnitude, variability and trends, *Biogeosciences*, 10, 1983–2000, <https://doi.org/10.5194/bg-10-1983-2013>, 2013.
- Wekerle, C., Wang, Q., Danilov, S., Schourup-Kristensen, V., von Appen, W.-J., and Jung, T.: Atlantic Water in the Nordic Seas: locally eddy-permitting ocean simulation in a global setup, *J. Geophys. Res. Oceans*, <https://doi.org/10.1002/2016JC012121>, 2017.
- Westberry, T., Behrenfeld, M. J., Siegel, D. A., and Boss, E.: Carbon-based primary productivity modeling with vertically resolved photoacclimation, *Global Biogeochemical Cycles*, 22, GB2024, <https://doi.org/doi:10.1029/2007GB003078>, 2008.
- 1680 Wolf-Gladrow, D. A., Zeebe, R. E., Klaas, C., Körtzinger, A., and Dickson, A. G.: Total alkalinity: The explicit conservative expression and its application to biogeochemical processes, *Marine Chemistry*, 106, 287–300,

- <https://doi.org/https://doi.org/10.1016/j.marchem.2007.01.006>, special issue: Dedicated to the memory of Professor Roland Wol-  
last, 2007.
- 1685 Wright, R. M., Le Quéré, C., Buitenhuis, E., Pitois, S., and Gibbons, M. J.: Role of jellyfish in the plankton ecosystem revealed using a  
global ocean biogeochemical model, *Biogeosciences*, 18, 1291–1320, <https://doi.org/10.5194/bg-18-1291-2021>, 2021.
- Yamanaka, Y. and Tajika, E.: The role of the vertical fluxes of particulate organic matter and calcite in the oceanic carbon cycle : Studies using  
an ocean biogeochemical general circulation model, *Global Biogeochemical Cycles*, 10, 361–382, <https://doi.org/10.1029/96GB00634>,  
1996.
- 1690 Ye, Y. and Völker, C.: On the Role of Dust-Deposited Lithogenic Particles for Iron Cycling in the Tropical and Subtropical Atlantic, *Global  
Biogeochemical Cycles*, 31, 1543–1558, <https://doi.org/10.1002/2017GB005663>, 2017.
- Ye, Y., Völker, C., and Gledhill, M.: Exploring the Iron-Binding Potential of the Ocean Using a Combined pH and DOC Parameteri-  
zation, *Global Biogeochemical Cycles*, 34, e2019GB006425, <https://doi.org/https://doi.org/10.1029/2019GB006425>, e2019GB006425  
10.1029/2019GB006425, 2020.
- 1695 Zeng, J., Nojiri, Y., Landschützer, P., Telszewski, M., and Nakaoka, S.: A Global Surface Ocean fCO<sub>2</sub> Climatology Based on a Feed-Forward  
Neural Network, *Journal of Atmospheric and Oceanic Technology*, 31, 1838 – 1849, <https://doi.org/https://doi.org/10.1175/JTECH-D-13-00137.1>, 2014.
- Zhang, R. and Vallis, G. K.: The Role of Bottom Vortex Stretching on the Path of the North Atlantic Western Boundary Current and on the  
Northern Recirculation Gyre, *Journal of Physical Oceanography*, 37, 2053 – 2080, <https://doi.org/10.1175/JPO3102.1>, 2007.