

ENSO statistics, teleconnections, and atmosphere-ocean coupling in the Taiwan Earth System Model version 1

Yi-Chi Wang¹, Wan-Ling Tseng^{2,*}, Yu-Luen Chen¹, [Shih-Yu Lee^{1*}](#), Huang-Hsiung Hsu¹, Hsin-Chien Liang¹

5 ¹Research Center for Environmental Changes, Academia Sinica, Taipei, Taiwan

²International Degree Program in Climate Change and Sustainable Development, National Taiwan University, Taipei, Taiwan

Correspondence to: [Shih-Yu Lee \(shihyu@gate.sinica.edu.tw\)](#), [Wan-Ling Tseng \(wtseng@ntu.edu.tw\)](#)

Abstract. This study provides an overview of the fundamental statistics and features of the El Niño Southern Oscillation (ENSO) in the historical simulations of the Taiwan Earth System Model version 1 (TaiESM1). Compared with observations, TaiESM1 can reproduce the fundamental features of observed ENSO signals, including seasonal phasing, thermocline coupling with winds, and atmospheric teleconnection during El Niño events. However, its ENSO response is approximately two times stronger than the observance in the spectrum, resulting in powerful teleconnection signals. The composite of El Niño events shows a strong westerly anomaly extending fast to the east Pacific in the initial stage in March, April, and May, initiating a warm sea surface temperature anomaly (SSTA) there. This warm SSTA maintains through September, October, and November (SON) and gradually diminishes after peaking in December. Analysis of wind stress-SST and heat flux-SST coupling proposes that biased positive SST-shortwave feedback contributes significantly to the strong warm anomaly over the eastern Pacific, especially in SON. Our analysis demonstrates TaiESM1's capability of simulating ENSO—a significant tropical climate variation on interannual scales with strong global impacts, and provides insights into mechanisms in TaiESM1 related to ENSO biases, laying the foundation for future model development to reduce uncertainties in TaiESM1 and climate models in general.

20 1 Introduction

The El Niño Southern Oscillation (ENSO) is the primary mode of interannual and decadal climate variability in the tropics ([Glantz, 2001; McPhaden et al., 2006](#)). It also affects climate variations in subtropical and mid-latitude regions across both hemispheres, as we, through teleconnection of Rossby waves ([Diaz et al. 2001; Yeh et al. 2018](#)). Therefore, its prediction is an essential part of global climate prediction on these scales ([Latif et al., 1998](#)). Additionally, ENSO is a crucial metric for climate model evaluation, especially for atmosphere–ocean coupling and associated physical feedbacks ([Planton et al., 2021](#)). Many studies have reported that coupled model intercomparison project (CMIP) models have successfully represented the basic features of observed ENSO, such as recognizable ENSO lifecycle and SST pattern over the tropical central and eastern Pacific ([Bellenger et al., 2014; Guilyardi et al., 2009, 2020; Lloyd et al., 2009, 2011](#)). However, many model biases are also

Deleted: .

Deleted: Shi

Deleted: Shi

Deleted: (shihyu@gate.sinica)

Deleted: .

Deleted: .

Deleted: .

Deleted: .

found in CMIP6 models (Beobide-Arsuaga et al., 2021; Capotondi et al., 2020; Chen & Jin, 2021), resulting in 30%–50% uncertainties in future ENSO projection (Beobide-Arsuaga et al. 2021).

The paradigm based on theory and observations depicts ENSO as a closely coupled oscillation system between the ocean and the atmosphere (Jin, 1997; Latif et al., 1998; McPhaden et al., 1998, 2006; Neelin et al., 1998; Philander, 1989; Wang, 2018; Wang & Picaut, 2013). An El Niño event begins from a westerly wind initiated over the west equatorial Pacific, typically in the spring of the first year (i.e., March, April, and May in year 0; MAM⁰). The westerly wind drives more warm water towards east and gradually warms the central Pacific. Around summer, eastward propagating oceanic Kelvin waves are triggered over the central Pacific, reducing the upwelling at the tropical east Pacific, deepening the thermocline, and warming the sea surface temperature (SST). Consequently, the zonal SST gradient and the easterly wind in the tropical Pacific are reduced. Such a retreat of the easterly wind further reduces the SST gradient, causing the so-called Bjerknes feedback between the easterly wind and the SST gradient (Bjerknes, 1969; Cane, 2005). Through this feedback, the warm SST increases and reaches a maximum around the following winter (i.e. December of the year 0 and January and February of the year 1; DJF⁺¹). Furthermore, following the warm SST anomaly (SSTA), the center of deep convection activity shifts toward the central Pacific, increasing latent heat flux and reducing shortwave heat flux into the ocean surface through deep cloud cover. Such seasonal phase-locking of an El Niño event is a crucial characteristic of the observed ENSO.

In contrast to the coupling nature of the atmosphere and ocean found in ENSO observations, the atmospheric feedback are found to dominate the modeled ENSO frequency and amplitude in CMIP models (Bellenger et al., 2014; Beobide-Arsuaga et al., 2021; Guilyardi et al., 2009; Lloyd et al., 2009, 2011). They noticed that the CMIP models tend to simulate a weaker Bjerknes feedback, namely, a weaker SST warming and westerly wind coupling. Furthermore, the heat flux-SST feedbacks are overemphasized in simulated ENSO dynamics, especially for the shortwave heat flux-SST feedback. Such overemphasized heat flux-SST feedback compensates for the weaker warming from the Bjerknes feedback, producing a seemingly realistic ENSO warming in CMIP models (Bayr et al., 2019). The biased Bjerknes and heat flux feedbacks are later found to be related to the biases of the seasonal variations of Walker circulation (Bayr et al., 2018, 2019). Such complexity resulting from intertwined atmospheric–ocean feedbacks makes it challenging for model developers to improve ENSO simulations without fully understanding how these mechanisms are represented in the coupled models.

Taiwan Earth System Model version 1 (TaiESM1; Lee et al., 2020) is the earth system model developed at the Research Center for Environmental Changes (RCEC), Academia Sinica. It participates in CMIP6 intercomparison activity and has been used in studying major climate variabilities and regional climate features (Chen & Jin, 2021; Park et al., 2020). While its overall performance of climate mean states and major variations has been evaluated and documented in Wang et al. (2021), in this study, we conducted more comprehensive investigation in ENSO's fundamental features and statistics in historical TaiESM1 simulations. We noticed that the ENSO amplitude increased significantly compared with the observations, with intense and prolonged warm SST from May to December. Especially over the eastern Pacific, the early onset of warming and sustained warming in September, October, and November (SON⁰) before peaks in December are the two primary prominent biases. We further analyzed the physical processes associated with ENSO's strong warming biases and found it is due to the

Deleted: .

Deleted: .

Deleted: .

Deleted: .

Deleted: .

Deleted: .

Deleted:)

Deleted: .

Deleted: .

80 biased positive feedback between SST-shortwave surface fluxes over the eastern equatorial Pacific. Such feedback is primarily attributed to the prevailing low clouds overlying the cold tongue region with large cold biases in SON⁰. Our results provide the baseline of ENSO performance of TaiESM1, and suggest the current ENSO biases in TaiESM1 is intertwined with biases of mean state and seasonal variation of the tropical climate system. The remainder of this study includes the following. Section 2 describes the TaiESM1 and observational dataset used for model evaluation and the methodology for analyzing ENSO. 85 Section 3 documents the basic characteristics of ENSO simulated in TaiESM1. More analysis focuses on seasonal variation of El Niño events in TaiESM1 and associated biases in Section 4. The study is concluded with a summary and discussion in Section 5.

Deleted: Niño

2 Data, Models, and Methodology

Based on the Community Earth System Model version 1.2.2 (CESM1.2.2; Hurrell et al., 2013) developed by the National Center for Atmospheric Research and sponsored by the National Science Foundation and the Department of Energy in the United States, TaiESM1 includes several physical schemes developed in-house in RCEC. These designs include convective triggering (Wang et al., 2015), radiation parameterization of 3-dimension topography (Lee et al., 2013), an aerosol scheme (Chen et al., 2013), and a probability density function-based cloud fraction scheme (Shiu et al., 2020). The ocean component is the same as CESM1 using the Parallel Ocean Program version 2 (POP2; Smith et al., 2010). An overall evaluation of climate variability in TaiESM1 shows that the simulated ENSO features stronger SST warming and atmospheric teleconnection compared with the base model CESM1 (Wang et al., 2021). In this study, we have conducted a more in-depth analysis of ENSO's fundamental features and statistics and identified the physical processes of ENSO biases within TaiESM1. The historical simulation of TaiESM1 from 1850 to 2014, driven by the forcing designed by the CMIP6, is analyzed. The historical run is initiated from the pre-industrial control run of TaiESM1. It utilized an atmospheric model with a horizontal resolution of 0.9° latitude × 1.25° longitude and 30 vertical layers. The community land model employed shares the same resolution as the atmospheric model. Additionally, the POP2 ocean model has a resolution of approximately 1.125° in longitude and 0.47° in latitude.

Deleted: .

Deleted: .

Deleted: .

Deleted: .

Deleted: .

Deleted: .

Deleted: .

We evaluated the model's performance using the atmospheric variables from the Collaborative Reanalysis Technical Environment Multireanalysis Ensemble version 2 (MRE2; Potter et al., 2018). The MRE2 is a product of the ensemble average of seven reanalysis products, including CFSR (Saha et al., 2010), ERA-Interim (Dee et al., 2011), MERRA (Rienecker et al., 2011), MERRA-2 (Gelaro et al., 2017), JRA-25 (Onogi et al., 2007), JRA-55 (Kobayashi et al., 2015), and 20CRv2c (Compo et al., 2011). Studies have found that the ensemble average can reduce the errors of individual reanalysis for selected atmospheric variables (Potter et al. 2018). For those variables not provided in MRE2, such as cloud cover, we used the ECMWF reanalysis version 5 (ERA5), the most up-to-date reanalysis produced by ECMWF (Hersbach et al., 2020). While previous studies have identified differences in air-sea feedbacks among reanalysis datasets, the ensemble mean of multiple reanalysis datasets can be used as the best estimate by reducing random errors through averaging (Kumar & Hu, 2012). We also obtained

Deleted: .

Deleted: .

Deleted: .

Deleted: .

Deleted: .

Deleted: .

Deleted: .

Deleted: .

Deleted: .

Deleted: .

130 precipitation data from the Global Precipitation Climatology Project (GPCP V2; [Adler et al., 2003](#); [Huffman et al., 2009](#)), SST
data from the Extended Reconstructed SST version 5 (ERSSTv5; [Huang et al., 2017](#)), and subsurface ocean data, such as sea
surface height (SSH) and potential subsurface temperature, from the Simple Ocean Data Assimilation version 3.3.2 (SODA
v3.2.2; [Carton et al., 2018](#)). The observational datasets used for this study span from 1980 to 2018, except for ERSST, which
covers the period from 1900 to 2018.

135 In this study, we employ regression and composite analyses as the primary tools to investigate ENSO features. We
represent El Niño using indices over crucial regions, including Niño 3 (5°N–5°S, 150°W–90°W), Niño 3.4 (5°N–5°S, 170°W–
120°W), and Niño 4 (5°N–5°S, 160°E–150°W), in the regression analysis. For the observational [Niño 3.4 index](#), we use a base
period between [1900 and 2014 from ERSSTv5](#), following the Niño index calculation of the Climate Prediction Center, NOAA.
[We also](#) use model data from 1900 to 2014 as the base period for TaiESM1's historic run. To avoid impacts of model bias on
140 longer timescales, such as interdecadal variation, we utilize the full length of available simulation data to obtain the most robust
statistics of ENSO feature simulated by TaiESM1.

We choose the composite method instead of the regression map to better identify teleconnection signals associated with
the El Niño events. In our preliminary analysis, the regressed maps of El Niño events show similar patterns with the composite
events in the tropics, but with much weaker signals in the midlatitudes (not shown). Furthermore, as the ENSO events simulated
145 in TaiESM1 shows very symmetric alternations between El Niño and La Niña events, our composite based on El [NinoNiño](#)
event is followed by La Niña event. To build the El Niño composite, we choose strong events with the Niño 3.4 index larger
than 1 standard deviation (i.e., 1.22°C) over the simulation period. The ERSSTv5 dataset includes eight El Niño events in
1982, 1986, 1987, 1991, 1994, 1997, 2002, and 2009. In comparison, TaiESM1 simulated 21 events throughout the entire
historical simulation. As the El Niño events simulated by TaiESM1 exhibit very strong amplitude, most of the composite fields
150 of these events passed the significance test at the 95% confidence level (not shown). Therefore, we will not denote regions
passing significant tests in the composite fields in the following analysis.

[In our analysis of TaiESM1's ability to simulate ENSO diversity, we examined its capability in distinguishing between
Eastern Pacific \(EP\) and Central Pacific \(CP\) El Niño events using the Niño3-Niño4 approach \(Kug et al., 2009\). During the
historical period of TaiESM1, we identified 23 CP events and 17 EP events. This higher frequency of CP events in TaiESM1
155 is consistent with previous findings in CMIP models \(Capotondi et al., 2020; Chen et al., 2017; McPhaden et al., 2011\). It is
worth noting that the total number of El Niño events exceeds 21 when using the El Niño 3.4 index, as some EP events transition
into CP events during the mature phase. In contrast, during the observation period from 1980 to 2014, we found a total of 4
EP events and 4 CP events. In the subsequent sections, we will also discuss the composites of EP and CP events in further
detail.](#)

160

Deleted: .

Deleted: ,.

Deleted: .

Deleted: data

Deleted: 1971

Deleted: 2000

Deleted: In contrast, we

Deleted: Nino

Deleted: To examine the ENSO diversity, Figure S1

170 **3 Basic statistics of ENSO in TaiESM1**

3.1 Niño 3.4 SST variability

Figure 1 shows the mean SST state (white contour) and monthly standard deviation (color shading) in ERSST and TaiESM1 over the tropical Pacific. The monthly standard deviation denotes the deviation of monthly SSTs from long-term monthly mean SST. TaiESM1 has a much more substantial equatorial Pacific SST variability, with an elongated region of high SST variation extending further into the warm pool region. Such an westward extension is collocated with the equatorial cold tongue, indicated by the 27°C isotherm thick white contour laying over the tropical eastern Pacific in the climatological SST mean field. TaiESM1 simulated a westward extension of the cold tongue compared with ERSSTv5. TaiESM1 also overestimated SST variation over other tropical oceans, including the Indian Ocean and warm pool.




Figure 1: Mean SST (white contours) and SST monthly standard deviation (color shading) for (a) detrended ERSSTv5 over 1958–2014 and (b) 1900–2014 of the TaiESM1’s historical simulations. The contour interval is 3°C, and the thick white contour indicates the 27°C isotherm.

180 Figure 2 shows the Niño3.4 SST index based on the ERSSTv5 and TaiESM1 historical simulations. TaiESM1 has more oscillatory ENSO signals with alternating cold and warm phases during 3–5 years compared with the observations. The Niño 3.4 index’s standard deviation of ERSSTv5 is 0.84°C, whereas that of TaiESM1 is 1.22°C. The simulated ENSO amplitude decreased after 1980 when the global temperature increased (Fig. 2a, b). [When comparing SST and surface wind fields between](#)

185 two 30-year periods, specifically 1950-1980 and 1984-2014, during which TaiESM1 exhibits distinct ENSO variability, Figure S1 demonstrates a notable shift in the background state towards a La Niña-like state. This shift is characterized by an increased zonal temperature gradient over the tropical Pacific and strengthened trade winds during the period of 1984-2014, in comparison to the period of 1950-1980. Previous studies investigating the ENSO response to changes in the observed mean state have indicated that such an increase in zonal wind stress can lead to a weakening of feedback mechanisms associated with El Niño (Fedorov et al., 2020; Zhao & Fedorov, 2020). This aligns with the observed decrease in ENSO variability in TaiESM1 during the period of 1984-2014.

190 _____ The power spectrum of Niño 3.4 confirms what is found in the time series of Niño 3.4, namely a much larger amplitude of ENSO in TaiESM1 (Fig. 2c). The amplitude of major peak between 3 and 4 years is around $250^{\circ}\text{C}^2/\text{month}$, while that of observed peak is around $75^{\circ}\text{C}^2/\text{month}$ in ERSSTv5. Similarly, around the secondary peak with period of 5 to 6 year in observations, TaiESM1 also shows two spectral peaks at 6 year and 8 year with stronger amplitude, respectively. Such model bias in representing ENSO-related spectral peaks have long noticed in CMIP models and still one of most challenging questions for climate models (Jha et al., 2014). Fig. 2d shows that the seasonal cycle of ENSO SST variance in ERSSTv5 and TaiESM1. Compared with the observations, the peak months simulated by TaiESM1 occurred in boreal winter, with one month delay than the observed, and a larger amplitude that was 1.5 times of the observed value.

Deleted: .

Deleted: .




Figure 2: Normalized time series of Niño3.4 index in (a) ERSST V5 (1900–2018) and (b) TaiESM1 historical run (1900–2014). The standard deviation of each dataset is noted on the upper-right side of the panels. The corresponding spectrums are shown in (c), with a black line for ERSSTv5 and a blue line for TaiESM1. (d) The seasonal cycle of SST variance in ERSSTv5 (black) and TaiESM1 (blue).

3.2 Atmosphere–ocean coupling of ENSO

As a coupled oscillation system, the coupling of atmosphere and ocean plays an important role in ENSO dynamics. To see how such coupling is simulated in TaiESM1, Figure 3 shows the regressed rainfall, wind stress, and SST to the Niño 3.4 index in the observations (Fig. 3a) and TaiESM1 (Fig. 3b). In Fig. 3a, the observation shows the west–east displacement of wind stress and warm SST over the tropical Pacific. The strong wind stress at 160°E is collocated with rainfall due to (mostly meridional) moisture convergence at the west and north edges of the warm SSTA, which was located in the central-eastern equatorial Pacific and was approximately 1° higher than the western equatorial Pacific. It is important to note that the major sea surface temperature anomaly (SSTA) did not occur in the eastern equatorial Pacific, where interannual variance was the highest. Instead, the SSTA occurred to the west of the region with the maximum variance. In TaiESM1 (as shown in Fig. 3b), the warm SST center is located approximately at 120°W, which is further east than in the observational data. Moreover, the magnitude of the SST anomaly is approximately 50% greater in TaiESM1 than in the observations. Compared with ERSST, the warm SSTA in TaiESM1 is meridionally narrower and more zonally elongated to the western Pacific around 155°E, causing stronger zonal and meridional SST gradients. Increase of the meridional SST gradient induces stronger meridional wind and moisture convergence over the equatorial Pacific. Zonally, the stronger westerly wind extends wider from 155°E to 120°W near the eastern edge of New Guinea Island. As a result, TaiESM1 produces strong wind stress and more deep convection (shown by rainfall; color shading in Fig. 3) to the north and west sides of the warm SSTA. Figure S2 shows the regressed magnitude of wind stress onto the Niño3.4 index and marks longitude center with dashed lines. While TaiESM1 reproduces the longitudinal center of wind stress response at 140°E as in observations, the response magnitude of wind stress to SST increase is weaker in TaiESM1 than in the observations.




Figure 3: Regression map of precipitation (mm/day, color shading), wind stress (1/s, vectors), and sea surface temperature (SST; °C, contours) on the normalized Niño-3.4 index for (a) Global Precipitation Climatology Project (GPCP), Simple Ocean Data Assimilation (SODA), Extended Reconstructed SST (ERSST), and (b) TaiESM1 historical simulation.

Deleted:

Deleted: S1

225 Figure 4 shows the regressed SSH to the wind stress averaged over the Niño-4 region (5°N–5°S, 150°E–90°E) to show
the thermocline response to the strengthening of equatorial wind stress over the western Pacific. In the observation, a west–
east dipole of thermocline response is found in Fig. 4a, showing thermocline deepening over the east Pacific (marked as black
square in Fig.4) and shallowing over the western subtropical Pacific. Compared with the observations, the TaiESM1 has
captured this west–east dipole of SSH response to equatorial wind stress, but with a much stronger magnitude over the eastern
equatorial Pacific (Fig. 4b). Such a strong response indicates that the SSH in TaiESM1 is more responsive than the observed
230 to the wind stress and can easily lead to an El Niño state through the Bjerknes feedback by reducing the zonal SST gradient
when the wind stress anomaly in the central equatorial Pacific initiates.




Figure 4: Regression map of sea surface height (SSH, cm/N/m², color shading) upon the normalized wind stress averaged over the Niño-4 region of 5°S–5°N and 160°E–160°W (a) from Simple Ocean Data Assimilation (SODA) and (b) the historical run of TaiESM1. The black square represents the Niño-4 region.

In addition to the two components of atmosphere-ocean coupling related to the wind stress, we also examine the heat
flux-SST coupling related to ENSO. Figures 5a and b show the shortwave radiation regressed to the Niño 3.4 index in MRE2
235 and TaiESM1. In the observations, the reduction in shortwave fluxes prevails in the tropics with the increased of warm SSTA
of the Niño 3.4 region because of emerging deep convection reflecting more shortwave radiation back (Fig. 5a). A zonal
gradient of shortwave fluxes is shown from $-160 \text{ W m}^{-2} \text{ K}^{-1}$ over the western Pacific (i.e. 170°E) to $-80 \text{ W m}^{-2} \text{ K}^{-1}$ over the
eastern Pacific (i.e. 120°W). Overall, TaiESM1 reproduced the negative feedback patterns in the deep tropics (3°S–3°N), but
with a much stronger shortwave reduction of $-200 \text{ W m}^{-2} \text{ K}^{-1}$ over the west Pacific in response to the warm SSTA of the Niño
240 3.4 region (Fig. 5b). Such pattern is consistent with stronger rainfall response of TaiESM1 in Fig. 3b, suggesting the stronger

deep convection response reflects more shortwave fluxes over the west Pacific. ~~In contrast, we observed an increase in~~ downwelling shortwave flux ~~over the subsidence regions adjacent to the JTCZ in both 10°N and 10°S over the east Pacific.~~ One notable feature in TaiESM1 is the shortwave fluxes can increase up to $60 \text{ Wm}^{-2} \text{ K}^{-1}$ over the tropical eastern Pacific (i.e. 120°W to 100°W), in contrast to a decrease in observations (Fig.5). Such difference suggests there is a biased cloud radiative response over the eastern Pacific when El Niño event occurs, which may induce biased heat flux-SST coupling in TaiESM1. We will further examine and discuss this bias in Section 4.

245

Deleted: On the other hand, increased

Deleted: Niño 3.4 SST is found near

Deleted: negative region's two fledges in TaiESM1 and the observations




Figure 5: Regression map of tropical surface downwelling shortwave radiation (RSDS; W/m^2 ; color) upon the normalized Niño-3.4 index for (a) MRE2 ensemble (1980–2017) and (b) TaiESM1 historical simulation (1900–2014).

3.3 Composite of El Niño structure and teleconnection

To evaluate the structure of El Niño events in TaiESM1, we compose the strong El Niño events with the Niño 3.4 index larger than 1 standard deviation of the entire time series (i.e., larger than 1.22°C). Under this definition, there are 21 Niño events in TaiESM1 historical run, and nine events in the MRE2 ensemble from 1980 to 2015.

Figures 6 and 7 present the seasonal variation of El Niño events in the tropic and its teleconnection pattern in the mid latitudes in the MRE2 ensemble and TaiESM1 by showing the 2m surface temperature (color shading; Fig. 6), sea-level pressure (SLP; contours; Fig. 6), precipitation (color shading; Fig. 7), and 300-hPa stream function (contours; Fig. 7). Overall, TaiESM1 reproduced the observed spatial structures and teleconnection patterns associated with El Niño; however, consistent with the over-simulated El Niño signals, TaiESM1 produces much stronger tropical SST warming and teleconnection in extratropical regions than the observations in all four seasons. As early as in June, July, and August in the first year (JJA⁰), TaiESM1 already simulates an SST anomaly with 2°C over the eastern Pacific (Fig. 6a, e) with clear rainfall response in central Pacific (Fig. 7a, e). A zonal dipole of surface temperature and rainfall between the eastern and western Pacific forms earlier than in the observations (color shading in Fig. 6a, e and Fig. 7a, e). In SON⁰, the warm SST anomaly grows even stronger and expands over the entire tropical Pacific in TaiESM1. As a result, very clear teleconnection similar to that of DJF⁺¹ can already be found in the north hemisphere, including horseshoe-shaped cooling in the western Pacific, and those over the Eurasia and United States (Fig.6b, f). In the meantime, TaiESM1 captures the responses in the south hemisphere in DJF⁺¹, including the warm and dry response over northern South America, the opposite responses over the southern South America, and the north–south dipole between Eurasia and South Asia (Fig. 6c, g). As for the rainfall response, TaiESM1 realistically simulates the shift of deep convection from the western Pacific to the central Pacific as the warm SST occurs in DJF⁺¹; however, the west shift of tropical SSTA causes the surface temperature response pattern also shift westward, resulting in enhanced stronger cooling in the East and Southeast Asia, The cooling further extends into the Indian ocean, causing an Indian Ocean Dipole-like response as depicted in Fig. 6g. In contrast to the weaker SSTA and surface temperature impacts in the observations (Fig. 6d) in MAM⁺¹, a strong teleconnection pattern in the surface temperature over the extratropical regions sustained into MAM⁺¹ in TaiESM1 (Fig. 6h). Furthermore, the overresponse of the rain band over the Indian Ocean could be due to the mean rainfall biases simulated in TaiESM1, as noticed by Wang et al. (2021). In terms of the atmospheric circulation anomaly, TaiESM1 successfully captures the southward Rossby wave propagation from the central Pacific to the southeastern Pacific (contours in Fig. 6a, e and Fig. 7a, e) in JJA⁰. From SON⁰ and into DJF⁺¹, teleconnection in TaiESM1 intensifies as the warm SST over the equatorial Pacific develops into the mature stage of El Niño (Fig. 6b, f; Fig. 7b, f). TaiESM1 reproduces the Rossby wave train response emitted from the equatorial Pacific into North America and the resulting dipole of surface temperature over North America during DJF⁺¹ (Fig. 6c, g). In line with the stronger temperature and rainfall responses observed during MAM+1, TaiESM1 exhibits El Niño-related circulation anomalies across the tropics (as shown in Figs. 6d, h and Figs. 7d, h).

Deleted: ¶

Deleted: , even

Deleted: Nino




Figure 6: The surface temperature (color shading) and SLP (contours; contour interval is 1 hPa; contours smaller than zero are dashed) of the El Niño composite in (a, e) JJA⁰, (b, f) SON⁰, (c, g) DJF⁺¹, and (d, h) MAM⁺¹ based on the MRE2 ensemble (left column) and TaiESM1 historical (right column).




Figure 7: As in Fig. 6 but showing precipitation (mm/day; color shading) and 300 hPa stream function (contours; contour interval of 2×10^6 m²/s anomalies; contour value smaller than zero is dashed).

Figure 8 shows the seasonal evolution of ocean subsurface potential temperature averaged over the selected El Niño events. The composites show four seasonal means from JJA⁰ when an El Niño event was identified to MAM⁺¹ in the following year. The green line shows the location of the 20°C subsurface isotherm (Z20) during El Niño events, and the gray line shows the climatological Z20. During JJA⁰ in the observations, Z20 deepens in the eastern Pacific and shallows in the western Pacific (Fig. 8a). While the TaiESM1 realistically simulated the climatological Z20 depth, it overestimated the response of subsurface temperatures and simulated a flatter Z20 profile in JJA⁰ (Fig. 8e). Such an overestimation bias in TaiESM1 continues from the beginning of the ENSO evolution through SON⁰ and DJF⁺¹ (Fig. 8b, c, f, and g). Accompanied by the warm bias over the eastern Pacific, the cold bias developed in SON⁰ when the cool water started to form in the tropical western Pacific (Fig. 8b–d, f–h). Such a zonal dipole of subsurface temperature bias in TaiESM1 manifests a basin-wide response of ocean circulations during El Niño events. Moreover, both warming and cooling from the surface to 100 m depth was much more pronounced in the model than in the observation, especially over the eastern Pacific. An unrealistic warming in the central-eastern equatorial Pacific is also notable, reflecting the unrealistic westward extension of positive SSTA. This bias led to an anomalously strong SST gradient up to 2°C between 180°W and 150°W, consistent with the wide-spread strong westerly anomalies. In the meantime, the composite of subsurface zonal currents corresponded to El Niño events simulated in TaiESM1 is shown in Figure S3. Strong westerly current anomaly suggests the zonal advection may also play a role in driving strong El Niño (Figure S3). More analysis is needed to determine which model components are more responsible for these biases.

Deleted: Fig.




Figure 8: Equatorial cross-section (5°S – 5°N) of the El Niño composite of the potential temperature anomaly (color shading) in (a, e) JJA⁰, (b, f) SON⁰, (c, g) DJF¹, and (d, h) MAM¹ based on SODA3.3.2 (left column) and TaiESM1 historical run (right column). The gray line shows the climatological 20°C isotherm (Z20), and the green dashed line shows the Z20 at the Niño state.

Deleted: ¶

310 **3.4 ENSO diversity and teleconnection**

Based on the EP and CP events identified by the Niño3-Niño4 approach, we make longitudinal profiles of SSTA for both EP events and CP events simulated in TaiESM1 compared to observations (Figure S4a). Our analysis reveals that TaiESM1 generally exhibits warmer SSTA over the tropical Pacific, particularly east of the 150°E line. Notably, during EP events, the model shows SSTA that can be as high as 1°C over the eastern Pacific region. To further analyze the impacts of EP and CP events, we constructed composites of surface temperature and SLP based on four EP events and four CP events during the observation period (1980-2014), as well as CP and EP events during the historical TaiESM1 period (1900-2014).

Regarding the EP composite, TaiESM1 successfully captures the observed features over the tropical Pacific, as illustrated in Figures S4b and S4c. However, the CP events identified in TaiESM1 exhibit elongated warm SSTA in the tropical region but with weaker teleconnections to the mid-latitudes in the northern hemisphere (Figures S4d and S4e). Additionally, the warming over North America is less pronounced and retreats towards the polar region in TaiESM1, whereas the observed cold surface temperature anomaly is replaced by a warm anomaly. These discrepancies suggest that biases in the model's mean state contribute to the model's biases in ENSO diversity and teleconnection patterns observed in TaiESM1, a common issue seen in other climate models as well (Ham & Kug, 2012).

325 **4 Linking the warm SST bias during El Niño events with simulated seasonal mean states**

In this section, we analyze the seasonal life cycle of the strong ENSO signals simulated in TaiESM1 to understand the intense tropical warming anomaly of El Niño events, and link this bias with seasonal mean states. Based on the El Niño events defined in the previous session, we construct the seasonal cycle of El Niño events by plotting the Hovmöller diagram averaging over the equator between 3°S to 3°N.

Figure 9 shows the Hovmöller diagram of SST anomalies and 1000 hPa zonal winds along the equator (3°S–3°N) based on the strong El Niño event selected from the ERSSTv5 and TaiESM1 historical run. In the observation in May, the warm SSTA occurs over the dateline and the westerly wind anomaly starts to propagate to 135°W. In JJA⁰, the warm SSTA slowly develops at 180°–135°W, progressively propagates to the eastern equatorial Pacific with westerly anomalies, and reaches maximum amplitude in November, December, and January (Fig. 9a). In contrast, the warm SSTA when initiated in May intensifies almost simultaneously in the basin east of 150°W in TaiESM1 (Fig. 9b). The warm anomaly quickly reaches 2°C over the entire eastern equatorial Pacific through JJA⁰ and into September. Such warming seems to be coupled with one branch quickly extending to the eastern equatorial Pacific after the initiation in May whereas the westerly anomaly's major branch is well coupled with SSTA over west Pacific most of time. The warming continues developing even when the westerly anomalies in the eastern equatorial Pacific weakened in JJA⁰ and reaches maximum in January as observations. This early development of a warm SSTA in the eastern equatorial Pacific in May and continuous warming in JJA⁰ and SON⁰ are two primary model bias that may contribute to strong El Niño events in TaiESM1.

Deleted: Fig.

Deleted: .




Figure 9: The Hovmöller diagram of composite SST anomalies (color) and zonal wind on 1000 hPa (contour) along the equator (3°S–3°N) for El Niño based on (a) observations (ERSST/MRE2 ensemble from 1980 to 2015) and (b) TaiESM1 historical run (1900–2014).

345

To understand the heat flux-SST coupling, we examined the similar composite of net surface heat flux (Figs. S5a, b) and latent heat fluxes (Figs. S5c, d), which are the two heat fluxes are prominent in observational El Niño events. Notably, in the observations, the seasonal variation of surface net heat fluxes in MRE2 is primarily controlled by latent heat fluxes, especially from May to November, as other heat fluxes play a minor role (Figs. S5c, d). However, the spatial patterns of net surface fluxes of TaiESM1 are primarily dominated by the shortwave surface fluxes (Figs. S5d) and amplified by the latent heat fluxes (Fig. 2b). Figure 10 shows the same Hovmöller diagram but for downwelling surface shortwave flux (color shading) and surface temperature (contours) composited over the El Niño events. Evolutions of shortwave radiation and SST during the lifecycle of El Niño events in observation and TaiESM1 are shown in Figs. 10a and b, respectively. In the observation, about 10 to 20 W/m² shortwave fluxes going upward (i.e. shown as negative on Fig.10) during the entire El Niño periods, which is well collocated with the warm SSTA, due to the increased shortwave reflection of deep convection triggered by warmer SST. Such reflection of shortwave fluxes by deep clouds induces negative feedback (i.e., higher SST, large reflected shortwave radiation) between shortwave radiation flux and SST in the tropics (Fig. 10a). The negative feedback intensified in September, reached the maximum in December, and continued to next February (Fig. 10a). In contrast to the observations, TaiESM1 produced unrealistically strong negative feedback over the western equatorial Pacific and Intertropical Convergence Zone (ITCZ) regions (Fig. 10b), and very strong positive shortwave radiation anomalies near the equatorial eastern Pacific after May in year 0 (Fig. 10a). The increase in shortwave influx starts from the eastern Pacific in May and gradually extends to 125°W in November, a feature that was not seen in observation. The westward progression of shortwave radiation increase contributes to erroneous strong SST warming, and its westward extension is seen in TaiESM1 from May to Jan in year 1 (Fig. 10b), instead of the observed simultaneous warming in the eastern Pacific.

350

355

360

Deleted: Fig. S2a

Deleted: Fig. S2c

Deleted: Fig

Deleted: 2

Deleted: Fig

Deleted: 2




Figure 10: A composite of El Niño events of RSDS (W/m^2 ; color) and surface temperature ($^{\circ}C$; contour) in (a) MRE2 and (b) TaiESM1. Both variables are averaged over the equator ($3^{\circ}S-3^{\circ}N$).

To understand the cause of the strong SSTA development from May to November identified in Fig. 9 and its relationship with shortwave radiation fluxes in Fig.10, we analyzed the seasonal cycle of low clouds (gray contours), vertical velocity (green contours), and SST (color shading) in the observations and TaiESM1 in Fig. 11a and b. Figure 11c shows the differences between TaiESM1 and MRE2. Figure 11d shows the climatological seasonal SST cycle at $100^{\circ}W$ to show the SST differences where cloud biases are prominent. On the seasonal timescale in observations, the SSTA is closely coupled with anomalies of vertical velocity and low-level clouds. Cold surface temperature is commonly collocated with excessive low-level clouds and subsidence (Fig. 11a). While TaiESM1 also exhibits a clear seasonal variation as the observations in the east tropical Pacific (Fig. 11b), the simulated seasonal cycle is rather asymmetric with colder bias during MAM⁰ and SON⁰ and warm biases during JJA⁰ and DJF⁺¹ (Fig. 11c). Compared with the observations, TaiESM1 warmed approximately one month later during MAM⁰ and cooled deeper over the eastern Pacific in SON⁰ (Fig. 11c). The warmer MAM⁰ sea surface in TaiESM1 provides a smaller zonal SST gradient and may lead to the earlier onset of the east propagating westerly anomaly found in MAM⁰ through the Bjerknes feedback (Fig. 9b). However, this cold surface temperature bias during SON⁰ provides a cold lower boundary for low stratus clouds to develop, leading to an environment for the strong positive feedback between shortwave fluxes and SST during El Niño events. Such impacts of cold tongue bias are also found in CESM1 and CESM2, which shares the same ocean model (i.e., POP2) as TaiESM1 (Wang et al., 2015; Wei et al., 2021).

Deleted: .




Figure 11: Climatological seasonal cycle of the surface temperature (shaded area; °C), low clouds (gray contours; %), and 500 hPa vertical velocity (green contours; hPa/s) of the tropical Pacific (3°S–3°N) in the (a) MRE2 ensemble and (b) TaiESM1, and (c) their differences. (d) The seasonal cycle of surface temperature at 100°W in MRE2 (black) and TaiESM1 (blue).

To understand further look into the bias in SON⁰, Fig. 12 shows the changes in cloud fraction and zonal circulations in
 390 SON⁰ during the composited El Niño events. TaiESM1 successfully reproduced the eastward shift of the upward branch related
 to deep convection in response to the warm SST during El Niño events, (Figs. 12a, b). However, TaiESM1 produced a much
 stronger response in circulations and cloud cover than the observations (Fig. 12b). Especially over the eastern Pacific, stronger
 upward motion anomaly occurs and about 10%–20% of low clouds is reduced in TaiESM1, indicating that a dramatic reduction
 of low-cloud regime (Fig. 11c). As a result, more shortwave heat influxes are allowed into the ocean surface in SON⁰ when El
 395 Niño events occur (Fig. 10b). Consistent with the seasonal variation of SSTA in Fig.9b, the increased shortwave fluxes thus
 help to keep the warm SSTA over the SON⁰ in the east Pacific after the warm SSTA initiates in MAM⁰ in TaiESM1 (Fig.10b).

Deleted: .




Figure 12: Longitudinal height cross-section of the El Niño composite of September, October, and November in year 0 (SON⁰) along the equator (5°S–5°N) based on (a) ERA5 and (b) TaiESM1. The cloud fraction (%) of the El Niño composite is shown in color shading, and the zonal and vertical winds (units: hPa/s) are shown as green vectors.

5 Summary

This study documented ENSO’s fundamental statistics and features in the TaiESM1, a CMIP6 participant. Compared with
 400 observational dataset, TaiESM1 has captured many prominent observed ENSO features, including a 3–5-year spectrum peak,
 seasonal phasing, the evolution of warm SST, deepening of the subsurface layer during the El Niño event, and teleconnection
 patterns in midlatitudes. However, the simulated El Niño signals in TaiESM1 are much stronger and more prolonged than the
 observed ENSO signals. Such strong signals are shown as intense warm SSTA over the tropical Pacific within the spatial
 structure of the composite of the El Niño event. In the meantime, in response to the tropical warm SSTA, the teleconnection
 405 wave activities associated with El Niño events are much stronger in TaiESM1, significantly affecting temperature and rainfall
 in high latitudinal regions, although with a similar spatial pattern to the observations.

To understand the cause of the warm El Niño biases, we investigated the seasonal cycle of strong El Niño events in
 TaiESM1. Compared with the observed delay propagation of SST after the wind anomaly’s initiation to the central Pacific in
 May, the simulated SST warming quickly propagates toward the eastern Pacific along with the westerly wind anomaly in
 410 TaiESM1. The SST warming continued through to November and reached a maximum in December, showing a stronger and
 more prolonged warm period than the observation. Moreover, in contrast to the negative feedback found in observations, our
 analysis shows that the strong El Niño warm anomalies in TaiESM1 is due to the strong SST-heat flux positive feedbacks,
 especially over the eastern Pacific in SON⁰. Further analysis shows that this biased feedback is due to the response of spurious
 low-cloud regime outside the west coast of South America simulated in TaiESM1. This biased cloud regime results from the
 415 seasonal variation of the cold tongue over the east Pacific, consistent with previous studies using the CESM family and CMIP
 models (Ham & Kug, 2012; Wei et al., 2021). During El Niño events, the stratus clouds over the eastern Pacific gradually
 diminished due to the warmer SST, allowing solar radiation to warm the ocean surface. This result leads to a positive feedback

Deleted: .

of downward solar radiation and SST over the eastern Pacific, an opposite sign of the observed relationship. This biased relationship is typical in the CMIP5 and CMIP6 models (Bayr et al. 2019; Beobide-Arsuaga et al. 2021).

In summary, TaiESM1 can reproduce many fundamental features of ENSO. However, it still possesses several biases shared by other CMIP6 models, including the lack of randomness of El Niño events, too strong El Niño magnitudes, and early SST warming in the early stage of El Niño events. The strong El Niño strength, our analysis found, is mainly resulted from the biased atmosphere-SST coupling accompanied by biases of the mean state and seasonal cycle of the cold tongue and Walker circulation, which is consistent with previous studies with CMIP5/CMIP6 models (Bayr et al., 2018; Chen et al., 2021). However, to resolve this bias (and others), more detailed analysis, including process-related metrics, and more model experiments, such as atmosphere-only and ocean-only experiments, is required to dissect the cause and effects of the seen biases. For TaiESM1, we plan to implement ocean-only experiments with the ocean component POP2, allowing us to quantify the ocean's response to biased winds and radiation fluxes. We will also conduct AMIP-type simulations to investigate the development of westerly wind anomalies under biased SST conditions. Combined with process-oriented diagnosis, these model experiments will help us to dissect and better comprehend the causes and effects of these observed biases. Also echoed with other studies of ENSO evaluations of CMIP6, our analysis suggests that a more process-based development strategy focusing on atmosphere–ocean coupling rather than a feature-based evaluation of ENSO is needed to reduce the uncertainty of ENSO simulations and future ENSO projection in climate models.

435 **Code and Data availability:**

All observational and analysis datasets used in this study are available online. The El Niño index can be downloaded from the NOAA Climate Prediction Center website (https://www.cpc.ncep.noaa.gov/products/analysis_monitoring/ensostuff/ensoyears_1971-2000_climo.shtml). The MRE2 ensemble can be downloaded from the website of the Collaborative REAnalysis Technical Environment – Intercomparison Project (<https://esgf-node.llnl.gov/projects/create-ip/>). The ERA5 monthly data can be downloaded from the Copernicus Climate Change Service Climate Data Store (doi: [10.24381/cds.f17050d7](https://doi.org/10.24381/cds.f17050d7); doi: [10.24381/cds.6860a573](https://doi.org/10.24381/cds.6860a573)). The Simple Ocean Data Assimilation (SODA) data v.2.3.3 can be downloaded from the SODA website (<http://www.soda.umd.edu/>). The Global Precipitation Climatology Project version 2.3 and Extended Reconstructed Sea Surface Temperature version 5 can be downloaded from the website of the NOAA PSL, Boulder, Colorado, USA (<https://psl.noaa.gov/data/gridded/data.gpcp.html>) and <https://psl.noaa.gov/data/gridded/data.noaa.ersst.v5.html>). The model code of TaiESM version 1 is available at <https://doi.org/10.5281/zenodo.3626654>. All post-process codes for producing figures presented in this paper are available at <https://doi.org/10.5281/zenodo.7740033>. The description and data of historical simulations of TaiESM1 can be found at <https://doi.org/10.22033/ESGF/CMIP6.9755>.

Deleted:

Deleted:

Author contribution

Yi-Chi Wang, Wan-Ling Tseng: Methodology, Investigation, Writing - Original draft, Writing – Review&Editing. **Yu-Luen Chen:** Software, Formal analysis, Visualization. **Shi-Yu Lee:** Conceptualization, Investigation, Writing – Review&Editing. **Huang-Hsiung Hsu:** Conceptualization, Methodology, Writing – Review&Editing, Supervision. **Hsin-Chien Liang:** Data curation.

Competing interests

The authors have no relevant financial or non-financial interests to disclose.

Acknowledgements

We would like to express our gratitude to the National Center for High-Performance Computing for providing the facilities for the CMIP6 participation of TaiESM1. This work was supported by the Taiwan National Science and Technology Council under grant numbers MOST 111-2111-M-001-013, MOST 111-2111-M-002-020, and MOST 110-2123-M-001-003.

References

- Adler, R. F., Huffman, G. J., Chang, A., Ferraro, R., Xie, P.-P., Janowiak, J., Rudolf, B., Schneider, U., Curtis, S., Bolvin, D., Gruber, A., Susskind, J., Arkin, P., & Nelkin, E. (2003). The Version-2 Global Precipitation Climatology Project (GPCP) Monthly Precipitation Analysis (1979–Present). *Journal of Hydrometeorology*, 4(6), 1147–1167. [https://doi.org/10.1175/1525-7541\(2003\)004<1147:TVGPCP>2.0.CO;2](https://doi.org/10.1175/1525-7541(2003)004<1147:TVGPCP>2.0.CO;2)
- Bayr, T., Latif, M., Dommenges, D., Wengel, C., Harlaß, J., & Park, W. (2018). Mean-state dependence of ENSO atmospheric feedbacks in climate models. *Climate Dynamics*, 50(9–10), 3171–3194. <https://doi.org/10.1007/s00382-017-3799-2>
- Bayr, T., Wengel, C., Latif, M., Dommenges, D., Lübbecke, J., & Park, W. (2019). Error compensation of ENSO atmospheric feedbacks in climate models and its influence on simulated ENSO dynamics. *Climate Dynamics*, 53(1–2), 155–172. <https://doi.org/10.1007/s00382-018-4575-7>
- Bellenger, H., Guilyardi, E., Leloup, J., Lengaigne, M., & Vialard, J. (2014). ENSO representation in climate models: from CMIP3 to CMIP5. *Climate Dynamics*, 42(7–8), 1999–2018. <https://doi.org/10.1007/s00382-013-1783-z>
- Beobide-Arsuaga, G., Bayr, T., Reintges, A., & Latif, M. (2021). Uncertainty of ENSO-amplitude projections in CMIP5 and CMIP6 models. *Climate Dynamics*, 56(11–12), 3875–3888. <https://doi.org/10.1007/s00382-021-05673-4>
- Bjerknes, J. (1969). Atmospheric Teleconnections from the Equatorial Pacific. *Monthly Weather Review*, 97(3), 163–172. [https://doi.org/10.1175/1520-0493\(1969\)097<0163:ATFTEP>2.3.CO;2](https://doi.org/10.1175/1520-0493(1969)097<0163:ATFTEP>2.3.CO;2)

Deleted: Ministry of

Deleted: MOST107

Deleted: 010

Deleted: 109

Deleted: 001-012-MY3

Deleted: 109

Deleted: 004

Deleted: ¶
¶

- Cane, M. A. (2005). The evolution of El Niño, past and future. *Earth and Planetary Science Letters*, 230(3–4), 227–240. <https://doi.org/10.1016/j.epsl.2004.12.003>
- 490 Capotondi, A., Deser, C., Phillips, A. S., Okumura, Y., & Larson, S. M. (2020). ENSO and Pacific Decadal Variability in the Community Earth System Model Version 2. *Journal of Advances in Modeling Earth Systems*, 12(12). <https://doi.org/10.1029/2019MS002022>
- Capotondi, A., Wittenberg, A. T., Kug, J., Takahashi, K., & McPhaden, M. J. (2020). ENSO Diversity. In *El Niño Southern Oscillation in a Changing Climate* (pp. 65–86). <https://doi.org/10.1002/9781119548164.ch4>
- 495 Carton, J. A., Chepurin, G. A., & Chen, L. (2018). SODA3: A New Ocean Climate Reanalysis. *Journal of Climate*, 31(17), 6967–6983. <https://doi.org/10.1175/JCLI-D-18-0149.1>
- Chen, C.-A., Hsu, H.-H., & Liang, H.-C. (2021). Evaluation and comparison of CMIP6 and CMIP5 model performance in simulating the seasonal extreme precipitation in the Western North Pacific and East Asia. *Weather and Climate Extremes*, 31, 100303. <https://doi.org/10.1016/j.wace.2021.100303>
- Chen, C., Cane, M. A., Wittenberg, A. T., & Chen, D. (2017). ENSO in the CMIP5 Simulations: Life Cycles, Diversity, and Responses to Climate Change. *Journal of Climate*, 30(2), 775–801. <https://doi.org/10.1175/JCLI-D-15-0901.1>
- 500 Chen, H.-C., & Jin, F.-F. (2021). Simulations of ENSO Phase-locking in CMIP5 and CMIP6. *Journal of Climate*, 1–42. <https://doi.org/10.1175/JCLI-D-20-0874.1>
- Chen, J.-P., Tsai, I.-C., & Lin, Y.-C. (2013). A statistical–numerical aerosol parameterization scheme. *Atmospheric Chemistry and Physics*, 13(20), 10483–10504. <https://doi.org/10.5194/acp-13-10483-2013>
- 505 Compo, G. P., Whitaker, J. S., Sardeshmukh, P. D., Matsui, N., Allan, R. J., Yin, X., Gleason, B. E., Vose, R. S., Rutledge, G., Bessemoulin, P., Brönnimann, S., Brunet, M., Crouthamel, R. I., Grant, A. N., Groisman, P. Y., Jones, P. D., Kruk, M. C., Kruger, A. C., Marshall, G. J., ... Worley, S. J. (2011). The Twentieth Century Reanalysis Project. *Quarterly Journal of the Royal Meteorological Society*, 137(654), 1–28. <https://doi.org/10.1002/qj.776>
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., 510 Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., ... Vitart, F. (2011). The ERA-Interim reanalysis: configuration and performance of the data assimilation system. *Quarterly Journal of the Royal Meteorological Society*, 137(656), 553–597. <https://doi.org/10.1002/qj.828>
- Fedorov, A. V., Hu, S., Wittenberg, A. T., Levine, A. F. Z., & Deser, C. (2020). *ENSO Low-Frequency Modulation and Mean State Interactions* (pp. 173–198). <https://doi.org/10.1002/9781119548164.ch8>
- 515 Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., da Silva, A. M., Gu, W., ... Zhao, B. (2017). The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2). *Journal of Climate*, 30(14), 5419–5454. <https://doi.org/10.1175/JCLI-D-16-0758.1>
- Glantz, M. H. (2001). *Currents of change: impacts of El Niño and La Niña on climate and society*. Cambridge University Press.
- 520

- Guilyardi, E., Capotondi, A., Lengaigne, M., Thual, S., & Wittenberg, A. T. (2020). ENSO Modeling: history, progress, and challenges. In M. J. McPhaden, A. Santoso, & W. Cai (Eds.), *El Niño Southern Oscillation in a Changing Climate* (pp. 199–226). American Geophysical Union. <https://doi.org/10.1002/9781119548164.ch9>
- 525 Guilyardi, E., Wittenberg, A., Fedorov, A., Collins, M., Wang, C., Capotondi, A., van Oldenborgh, G. J., & Stockdale, T. (2009). Understanding El Niño in Ocean–Atmosphere General Circulation Models: Progress and Challenges. *Bulletin of the American Meteorological Society*, *90*(3), 325–340. <https://doi.org/10.1175/2008BAMS2387.1>
- Ham, Y.-G., & Kug, J.-S. (2012). How well do current climate models simulate two types of El Niño? *Climate Dynamics*, *39*(1–2), 383–398. <https://doi.org/10.1007/s00382-011-1157-3>
- 530 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., ... Thépaut, J. (2020). The ERA5 global reanalysis. *Quarterly Journal of the Royal Meteorological Society*, *146*(730), 1999–2049. <https://doi.org/10.1002/qj.3803>
- Huang, B., Thorne, P. W., Banzon, V. F., Boyer, T., Chepurin, G., Lawrimore, J. H., Menne, M. J., Smith, T. M., Vose, R. S., & Zhang, H.-M. (2017). Extended Reconstructed Sea Surface Temperature, Version 5 (ERSSTv5): Upgrades, 535 Validations, and Intercomparisons. *Journal of Climate*, *30*(20), 8179–8205. <https://doi.org/10.1175/JCLI-D-16-0836.1>
- Huffman, G. J., Adler, R. F., Bolvin, D. T., & Gu, G. (2009). Improving the global precipitation record: GPCP Version 2.1. *Geophysical Research Letters*, *36*(17), L17808. <https://doi.org/10.1029/2009GL040000>
- 540 Hurrell, J. W., Holland, M. M., Gent, P. R., Ghan, S., Kay, J. E., Kushner, P. J., Lamarque, J.-F., Large, W. G., Lawrence, D., Lindsay, K., Lipscomb, W. H., Long, M. C., Mahowald, N., Marsh, D. R., Neale, R. B., Rasch, P., Vavrus, S., Vertenstein, M., Bader, D., ... Marshall, S. (2013). The Community Earth System Model: A Framework for Collaborative Research. *Bulletin of the American Meteorological Society*, *94*(9), 1339–1360. <https://doi.org/10.1175/BAMS-D-12-00121.1>
- Jha, B., Hu, Z.-Z., & Kumar, A. (2014). SST and ENSO variability and change simulated in historical experiments of CMIP5 models. *Climate Dynamics*, *42*(7–8), 2113–2124. <https://doi.org/10.1007/s00382-013-1803-z>
- 545 Jin, F.-F. (1997). An Equatorial Ocean Recharge Paradigm for ENSO. Part I: Conceptual Model. *Journal of the Atmospheric Sciences*, *54*(7), 811–829. [https://doi.org/10.1175/1520-0469\(1997\)054<0811:AEORPF>2.0.CO;2](https://doi.org/10.1175/1520-0469(1997)054<0811:AEORPF>2.0.CO;2)
- Kobayashi, S., Ota, Y., Harada, Y., Ebata, A., Moriya, M., Onoda, H., Onogi, K., Kamahori, H., Kobayashi, C., Endo, H., Miyaoaka, K., & Takahashi, K. (2015). The JRA-55 Reanalysis: General Specifications and Basic Characteristics. *Journal of the Meteorological Society of Japan. Ser. II*, *93*(1), 5–48. <https://doi.org/10.2151/jmsj.2015-001>
- 550 Kug, J.-S., Jin, F.-F., & An, S.-I. (2009). Two Types of El Niño Events: Cold Tongue El Niño and Warm Pool El Niño. *Journal of Climate*, *22*(6), 1499–1515. <https://doi.org/10.1175/2008JCLI2624.1>
- Kumar, A., & Hu, Z.-Z. (2012). Uncertainty in the ocean–atmosphere feedbacks associated with ENSO in the reanalysis products. *Climate Dynamics*, *39*(3–4), 575–588. <https://doi.org/10.1007/s00382-011-1104-3>

- 555 Latif, M., Anderson, D., Barnett, T., Cane, M., Kleeman, R., Leetmaa, A., O'Brien, J., Rosati, A., & Schneider, E. (1998). A review of the predictability and prediction of ENSO. *Journal of Geophysical Research: Oceans*, 103(C7), 14375–14393. <https://doi.org/10.1029/97JC03413>
- Lee, W.-L., Liou, K. N., & Wang, C. (2013). Impact of 3-D topography on surface radiation budget over the Tibetan Plateau. *Theoretical and Applied Climatology*, 113(1–2), 95–103. <https://doi.org/10.1007/s00704-012-0767-y>
- 560 Lee, W.-L., Wang, Y.-C., Shiu, C.-J., Tsai, I., Tu, C.-Y., Lan, Y.-Y., Chen, J.-P., Pan, H.-L., & Hsu, H.-H. (2020). Taiwan Earth System Model Version 1: description and evaluation of mean state. *Geoscientific Model Development*, 13(9), 3887–3904. <https://doi.org/10.5194/gmd-13-3887-2020>
- Lloyd, J., Guilyardi, E., & Weller, H. (2011). The role of atmosphere feedbacks during ENSO in the CMIP3 models. Part II: using AMIP runs to understand the heat flux feedback mechanisms. *Climate Dynamics*, 37(7–8), 1271–1292. <https://doi.org/10.1007/s00382-010-0895-y>
- 565 Lloyd, J., Guilyardi, E., Weller, H., & Slingo, J. (2009). The role of atmosphere feedbacks during ENSO in the CMIP3 models. *Atmospheric Science Letters*, 10(3), 170–176. <https://doi.org/10.1002/asl.227>
- McPhaden, M. J., Busalacchi, A. J., Cheney, R., Donguy, J.-R., Gage, K. S., Halpern, D., Ji, M., Julian, P., Meyers, G., Mitchum, G. T., Niiler, P. P., Picaut, J., Reynolds, R. W., Smith, N., & Takeuchi, K. (1998). The Tropical Ocean-Global Atmosphere observing system: A decade of progress. *Journal of Geophysical Research: Oceans*, 103(C7), 14169–14240. <https://doi.org/10.1029/97JC02906>
- 570 McPhaden, M. J., Lee, T., & McClurg, D. (2011). El Niño and its relationship to changing background conditions in the tropical Pacific Ocean. *Geophysical Research Letters*, 38(15). <https://doi.org/10.1029/2011GL048275>
- McPhaden, M. J., Zebiak, S. E., & Glantz, M. H. (2006). ENSO as an Integrating Concept in Earth Science. *Science*, 314(5806), 1740–1745. <https://doi.org/10.1126/science.1132588>
- 575 Neelin, J. D., Battisti, D. S., Hirst, A. C., Jin, F.-F., Wakata, Y., Yamagata, T., & Zebiak, S. E. (1998). ENSO theory. *Journal of Geophysical Research: Oceans*, 103(C7), 14261–14290. <https://doi.org/10.1029/97JC03424>
- Onogi, K., Tsutsui, J., Koide, H., Sakamoto, M., Kobayashi, S., Hatsushika, H., Matsumoto, T., Yamazaki, N., Kamahori, H., Takahashi, K., Kadokura, S., Wada, K., Kato, K., Oyama, R., Ose, T., Mannoji, N., & Taira, R. (2007). The JRA-25 Reanalysis. *Journal of the Meteorological Society of Japan. Ser. II*, 85(3), 369–432. <https://doi.org/10.2151/jmsj.85.369>
- 580 Park, J., Kim, H., Simon Wang, S.-Y., Jeong, J.-H., Lim, K.-S., LaPlante, M., & Yoon, J.-H. (2020). Intensification of the East Asian summer monsoon lifecycle based on observation and CMIP6. *Environmental Research Letters*, 15(9), 0940b9. <https://doi.org/10.1088/1748-9326/ab9b3f>
- Philander, S. (1989). *El Nino, La Nina, and the Southern Oscillation*. Academic Press.
- 585 Planton, Y. Y., Guilyardi, E., Wittenberg, A. T., Lee, J., Gleckler, P. J., Bayr, T., McGregor, S., McPhaden, M. J., Power, S., Roehrig, R., Vialard, J., & Voldoire, A. (2021). Evaluating Climate Models with the CLIVAR 2020 ENSO Metrics Package. *Bulletin of the American Meteorological Society*, 102(2), E193–E217. <https://doi.org/10.1175/BAMS-D-19-0337.1>

- 590 Potter, G. L., Carriere, L., Hertz, J., Bosilovich, M., Duffy, D., Lee, T., & Williams, D. N. (2018). Enabling Reanalysis Research Using the Collaborative Reanalysis Technical Environment (CREATE). *Bulletin of the American Meteorological Society*, 99(4), 677–687. <https://doi.org/10.1175/BAMS-D-17-0174.1>
- Rienecker, M. M., Suarez, M. J., Gelaro, R., Todling, R., Bacmeister, J., Liu, E., Bosilovich, M. G., Schubert, S. D., Takacs, L., Kim, G.-K., Bloom, S., Chen, J., Collins, D., Conaty, A., da Silva, A., Gu, W., Joiner, J., Koster, R. D., Lucchesi, R., ... Woollen, J. (2011). MERRA: NASA's Modern-Era Retrospective Analysis for Research and Applications. *Journal of Climate*, 24(14), 3624–3648. <https://doi.org/10.1175/JCLI-D-11-00015.1>
- 595 Saha, S., Moorthi, S., Pan, H.-L., Wu, X., Wang, J., Nadiga, S., Tripp, P., Kistler, R., Woollen, J., Behringer, D., Liu, H., Stokes, D., Grumbine, R., Gayno, G., Wang, J., Hou, Y.-T., Chuang, H., Juang, H.-M. H., Sela, J., ... Goldberg, M. (2010). The NCEP Climate Forecast System Reanalysis. *Bulletin of the American Meteorological Society*, 91(8), 1015–1058. <https://doi.org/10.1175/2010BAMS3001.1>
- 600 Shiu, C.-J., Wang, Y.-C., Hsu, H.-H., Chen, W.-T., Pan, H.-L., Sun, R., Chen, Y.-H., & Chen, C.-A. (2020). GTS v1.0: A Macrophysics Scheme for Climate Models Based on a Probability Density Function. *Geoscientific Model Development*. <https://doi.org/10.5194/gmd-2020-144>
- Smith, R., Jones, P., Briegleb, B., Bryan, F., Danabasoglu, G., Dennis, J., Dukowicz, J., Fox-Kemper, B., Eden, C., Gent, P., Hecht, M., Jayne, S., Jochum, M., Large, W., Lindsay, K., Maltrud, M., Norton, N., Peacock, S., Vertenstein, M., & Yeager, S. (2010). *The Parallel Ocean Program (POP) Reference Manual*.
- 605 Wang, C. (2018). A review of ENSO theories. *National Science Review*, 5(6), 813–825. <https://doi.org/10.1093/nsr/nwy104>
- Wang, C.-C., Lee, W.-L., Chen, Y.-L., & Hsu, H.-H. (2015). Processes Leading to Double Intertropical Convergence Zone Bias in CESM1/CAM5. *Journal of Climate*, 28(7), 2900–2915. <https://doi.org/10.1175/JCLI-D-14-00622.1>
- Wang, C., & Picaut, J. (2013). Understanding Enso Physics-A Review. In A. G. Union (Ed.), *Earth's Climate: The Ocean-Atmosphere Interaction* (pp. 21–48). <https://doi.org/10.1029/147GM02>
- 610 Wang, Y.-C., Pan, H. L., & Hsu, H. H. (2015). Impacts of the triggering function of cumulus parameterization on warm-season diurnal rainfall cycles at the Atmospheric Radiation Measurement Southern Great Plains Site. *Journal of Geophysical Research*, 120(20), 10,681-10,702. <https://doi.org/10.1002/2015JD023337>
- Wang, Y., Hsu, H., Chen, C., Tseng, W., Hsu, P., Lin, C., Chen, Y., Jiang, L., Lee, Y., Liang, H., Chang, W., Lee, W., & Shiu, C. (2021). Performance of the Taiwan Earth System Model in Simulating Climate Variability Compared With Observations and CMIP6 Model Simulations. *Journal of Advances in Modeling Earth Systems*, 13(7). <https://doi.org/10.1029/2020MS002353>
- 615 Wei, H., Subramanian, A. C., Karnauskas, K. B., DeMott, C. A., Mazloff, M. R., & Balmaseda, M. A. (2021). Tropical Pacific Air-Sea Interaction Processes and Biases in CESM2 and Their Relation to El Niño Development. *Journal of Geophysical Research: Oceans*, 126(6). <https://doi.org/10.1029/2020JC016967>
- 620 Zhao, B., & Fedorov, A. (2020). The Effects of Background Zonal and Meridional Winds on ENSO in a Coupled GCM. *Journal of Climate*, 33(6), 2075–2091. <https://doi.org/10.1175/JCLI-D-18-0822.1>

625

