In the following, author comments are in black and reviewer comments are in **blue**. When we present text passages from the revised manuscript, they appear in "*italics*".

There are many 1D models of blowing snow, several 2D models (operating on decametre grid scales as far back as the 1990s), and even 3D coupled snow-atmosphere models. An operational model that can be run on large scales is certainly of interest. The authors give a fair summary of previous work and a nearly complete description of the model being introduced.

We first would like to thank the reviewer for their constructive comments. In the following, we will give a detailed answer to each of them.

The paper is very long, however, for the amount of original material being presented; some of this is due to repetition and poor organization of material.

We understand the major concerns of both reviewers about the length and lack of clarity of the paper. In order to address this issue, we propose to extensively modify the structure of the paper to improve its readability. We will split Section 2 (Methodological choices) into two separate sections. The first one will be dedicated to a literature review largely reduced in length by focusing on the topics for which an added value is provided in SnowPappus compared with existing works. It includes discussions about the representation of saltation flux and lower boundary condition for suspension flux and terminal fall speed parameterizations. This will allow to make the logics of the manuscript clearer. For processes that are simply implemented in SnowPappus following existing models or previous literature, the literature overview will be considerably shortened, or even suppressed if it is not useful for the remaining part of the article, such as the discussion on sublimation (L449-452). Then, a second section will be dedicated to a concise model description without any theoretical interruptions between the description of the different implementations. We believe this organization will help the reader to understand more quickly our modelling choices.

In addition, several parts of the manuscript will be strongly shortened, primarily the description of numerical performance, various repetitions and unnecessary information or discussion will be withdrawn and a more concise writing style will be adopted when possible. Besides, the modification of the paper structure, which splits the section "Methodological choices" into 2 parts do not result in additional length, as the number of subsubsections does not changes a lot and repetitions will be avoided. Overall, the planned changes will allow to suppress Fig. 15 and move Fig. 14 in the appendix and to reduce overall text length by 20% (without taking appendix into account).

In the following, we will first list the text passages that will be strongly shortened, or withdrawn, in their order of appearance and identified by their section number and if necessary line number in the preprint. Then, we present a revised text outline to apply the described changes. New (sub)sections are written with their titles <u>underlined</u>, and there contains in plain text. Parts of the outline in which the organization remain identical with the preprint's outline are marked as UNCHANGED, although some length reductions have been also applied in these parts.

shortened text passages (with paragraph and line number from the initial manuscript):

2.1 Target, opportunities and constraints

2.2.2, wind profile

2.3.2 Blowing snow trajectories and transport modes

2.3.4 L257-263 discussion on the influence of particle size distribution on the suspension transport

2.3.3 L207-241 - Different types of transport models

2.3.7 L334-358 equations of simple saltation parameterizations P90 and S04

2.3.8 Influence of fetch distance

2.5 L502-508 Influence if snow transport on snow surface properties (state of the art)

3.2 L 560-570 Wind downscaling description

3.5 L 623-634 Methods for blowing snow occurrence measurements

4.6 L734-739, domain decomposition for parallel computing

5.1 L799-805 Discussion on the outlier in blowing snow fluxes evaluation withdrawn text passages :

2.4 L449-452, Sublimation

4.5 L712-717, Evaluation of blowing snow fluxes at *Col du Lac Blanc*, discussion on the outlier (repeated in the discussion)

new outline in the revised manuscript :

1. Introduction

General introduction followed by the target, opportunities and constraints of the development of SnowPappus (corresponds to Sect. 2.1 in the preprint)

2. Blowing snow flux computation : state of the art

2.1 Blowing snow occurrence

Useful theoretical background for blowing snow occurrence detection (Sect. 2.2.1 in the preprint)

2.2 Horizontal blowing snow fluxes

2.2.1 Notations and geometric considerations

2.2.2 Blowing snow particle trajectories and transport modes

State of the art on the trajectories of blowing snow particles and transport modes, focused on saltation and suspension (corresponds to Sect. 2.3.2 in the preprint)

2.2.3 Suspension transport modelling

-Existing types of suspension transport models (part of Sect. 2.3.3 in the preprint)

-Literature review on the effective terminal fall speed (mainly informations in Sect. 2.3.5 of the preprint)

- 2.2.4 Transition between saltation and suspension
 - The way it is treated in other models
 - State of the art on this transition zone
 - Problems of definition of the lower boundary condition for suspension transport
 - (corresponds mainly to Sect. 2.3.6 of the preprint)
- 2.2.5 Simple saltation models

-Short description of Sorensen et al., 2004 (S04) and Pomeroy et al., 1990 (P90) saltation parameterizations

-Discrepancies between S04 and P90 and discussion of the possible causes (informations in Sect. 2.3.7 of the preprint)

3. Model description

3.0 Crocus description

Very short description of Crocus snow model

3.1 Blowing snow occurrence

Equations of the different options implemented in SnowPappus for threshold wind speed for transport (Sect. 2.2.3 in the preprint)

3.2 Horizontal blowing snow flux

3.2.1 Suspension transport

- Reasons for the choice of the model type

- Logarithmic wind speed profile (Sect. 2.2.2 in the preprint)

- Equations for suspension transport in SnowPappus (corresponds to Sect. 2.3.4 in the preprint)

- Parameterization of terminal fall speed used in SnowPappus (informations in Sect.

2.3.5 of the preprint)

- Maximum height of suspension transport as a function of fetch distance (part of Sect. 2.3.8 of the preprint)

3.2.2 Saltation transport and transition with suspension

- Description of the two options (using S04 and P90) of flux computation in the saltation zone and the transition with suspension in SnowPappus (included in Sect. 2.3.7 of the preprint).

- Influence of fetch distance on saltation transport in SnowPappus (included in Sect.

2.3.8 of the preprint, with Fig. 3 appearing)

3.3 Sublimation

Sublimation options in SnowPappus

3.4 Mass balance

Mass balance implementation in SnowPappus

3.5 Influence of snow transport and deposition on snow surface properties

- Properties of deposited snow
- Blowing snow induced snow metamorphism

3.6 Implementation in SURFEX

How SnowPappus is included in SURFEX code (mainly Sect. 2.7 of the preprint, but includes the description of domain decomposition for parallel computing)

4. Evaluation : methods

UNCHANGED (with length reductions)

5. Results

mainly UNCHANGED but addition of 1 paragraph (see below) and Fig. 14 moved in appendix, as partly redundant with Fig. 13 .

5.1 Comparison of saltation parameterizations

Comparison of blowing snow fluxes obtained with S04 and P90 implementations in SnowPappus and implications for the comparison of S04 and P90 (corresponds to the end of Sect. 2.3.7 in the preprint).

6. Discussion

UNCHANGED (with length reductions)

7. Conclusion UNCHAGED Having said that "Blowing snow occurrence evaluation showed SnowPappus performs as well as a currently operational scheme" in the abstract, there needs to be a statement about the need and benefits for SnowPappus.

We agree with the reviewer that the needs and benefit of SnowPappus were insufficiently stressed out in the abstract. We thus propose to replace L 2-4 by "*Thus, the evolution of operational snow modelling systems towards 100-500 m resolutions requires representing this process at these resolutions, over large domains and entire snow seasons. We developed SnowPappus, a parsimonious blowing snow model coupled to the Crocus state-of-the-art snow model able to cope with these requirements.*"

Besides, the SYTRON operational scheme was indeed already available to simulate wind-induced snow transport coupled to Crocus, but it is designed to work on idealised geometries, where an altitudinal band of a mountain range is represented by 8 slopes with 8 orientations, where snow can be transported from one slope to the opposite one (Vionnet et al., 2018). Thus, this model is not suitable for distributed simulations. It is however used operationally for the detection of blowing snow occurrence. To clarify the added value of SnowPappus compared to this scheme, we propose to make less ambiguous the above-mentioned abstract phrase by changing "Blowing snow occurrence evaluation showed SnowPappus performs as well as a currently operational scheme" in "*Evaluations showed SnowPappus performs as well as currently operational scheme SYTRON in terms of blowing snow occurrence detection, while the latter does not give access to a spatialized information*".

Moreover, mention to SYTRON and its difference with SnowPappus will be discussed briefly in the end of the introduction, and the goal and strategy of the paper will be clarified. Given our reorganisation of the end of the introduction (see our response on manuscript length and organization), we present below the whole end of the introduction in the revised manuscript, and highlight in **bold** the sentences that address specifically these two issues.

"In the above-mentioned context of increasing resolution of snow modelling systems, the long-term project of CNRM aims at performing simulations with Crocus at the scale of the French Alps at 250m resolution in an operational purpose, associated with a data assimilation framework requiring ensemble runs of 50-100 members (Largeron et al., 2020; Cluzet et al., 2021). The 250 m resolution allows a trade-off between the need for precisely representing slopes and aspects, influencing mass and energy balance of the snowpack, and the expected computational cost. In this context, a numerically efficient representation of wind-induced snow transport that can be coupled to Crocus simulations is lacking while this is necessary to better account for its impact on avalanche forecasting over French mountains. Two blowing snow scheme coupled with Crocus exist yet : SYTRON (Vionnet et al., 2018) and Crocus-Meso-NH (Vionnet et al., 2014). However, both are unadapted to this geometry and resolution.

Thus, the goal of this paper is to describe and present first evaluations of a novel blowing snow scheme, SnowPappus, coupled to Crocus and able to be included in the aforementioned large-scale simulation system. Point-scale evaluation of blowing snow flux will be presented to discuss the modelling choices. In order to avoid a prohibitive computational cost, this scheme shall not be much more computationally intensive than the Crocus model itself, and it will be forced with 2D

wind fields downscaled from NWP systems rather than coupled with 3D high resolution atmospheric model."

There is no evaluation of the model in large-scale applications beyond demonstrating its computational feasibility.

This concern was also raised by Reviewer #1. In this model description paper, we considered that the evaluation of blowing snow fluxes was the main topic to address as this is the most direct observation of the newly simulated processes. These evaluations do not allow to determine if the spatial patterns of snow properties simulated are improved by the blowing snow module. This limitation is explicitly mentioned L858-860 "Snow redistribution in 2D simulations has not been evaluated in this article, and will be the subject of a future study expected to provide complementary insights to the following discussions". Such evaluations are definitely necessary but they raise major methodological difficulties which justify a dedicated and separated paper.

First of all, in large scale simulations, several sources of errors are superposed, making it hard to determine if differences between model and observations come from a misrepresentation of wind-induced snow transport or from other sources of errors. In particular, precipitation forcing at high altitudes suffer from high uncertainty partly due to a largely unexplained spatial variability in NWP precipitation outputs, a severe lack of observations to constrain meteorological analysis systems, and various issues in radar precipitation measurements over complex terrains. It usually leads forcing errors to prevail in snow simulation systems that are not forced by local observations are often not representative of the spatial scale of a simulation system (i.e. 250 m resolution in our case), evaluations have to rely on satellite observations which also involve complex retrieval methodologies and an appropriate consideration of associated uncertainties. Finally, methodological developments are required to compare simulated and observed snow maps as consistent spatial patterns may be simulated with slight localization inaccuracies, making traditional evaluation scores often unadapted (Gilleland et al., 2009).

Therefore, we also prepared a dedicated evaluation paper of the spatial distributions of snow height and Snow Melt Out Date against satellite stereo-imagery (Deschamps-Berger et al., 2020) and optical products (Gascoin et al., 2019). In this evaluation, the contributions of model and forcing uncertainties in simulation errors are quantified and compared. As expected, the forcing uncertainties prevail. This paper (Haddjeri et al.) will be submitted in September 2023 to The Cryosphere and will be complementary to the model description and first evaluations provided in this GMD model description paper.

Moreover, we would like to stress that we follow a strategy similar to other published windinduced snow transport models operating at similar or lower resolution. Due the same methodological challenges, the corresponding publications did neither describe evaluation of the simulated spatial patterns of snow depth or surface properties (Gallée et al., 2001; Amory et al. 2021; Sharma et al., 2021). These spatial evaluations, when performed, were sometimes conducted in a separate paper (Gerber et al., 2023). Smaller scale blowing snow models were often described along with a purely qualitative evaluation of snow depth patterns on areas ranging from hundreds of meters long transects to a few kilometre square test zones (Liston and Sturm (1998); Liston et al. (2007); Vionnet et al. (2014)), and sometimes without any spatial evaluation (Essery et al., 1999). Conversely, we provide in the present paper quantitative evaluations of the blowing snow flux and transport occurrence simulated with SnowPappus against a 10-years long observation time series at Col du Lac Blanc. As stressed in our introduction, such evaluations are very unusual in the currently available literature (except in Amory et al., 2021 in Antarctica), so concerning this direct variable, the evaluations provided in our paper are more advanced than similar literature on blowing snow models. These analyses allow to evaluate and discuss the parameterization choices done to compute blowing snow occurrence and fluxes and to question the interest of microstructure-based parameterizations for which the added value had never been assessed. This is also an important added value of our paper.

We propose to clarify the goal and strategy in the introduction. Given the introduction will be merged with Sect. 2.1 (see above our response on manuscript length and organization), we give here the full paragraphs including this clarification. The part of the text which is the most dedicated to it appears in bold.

"In the above-mentioned context of increasing resolution of snow modelling systems, the long-term project of CNRM aims at performing simulations with Crocus at the scale of the French Alps at 250m resolution in an operational purpose, associated with a data assimilation framework requiring ensemble runs of 50-100 members (Largeron et al., 2020; Cluzet et al., 2021). The 250 m resolution allows a trade-off between the need for precisely representing slopes and aspects, influencing mass and energy balance of the snowpack, and the expected computational cost. In this context, a numerically efficient representation of wind-induced snow transport that can be coupled to Crocus simulations is lacking while this is necessary to better account for its impact on avalanche forecasting over French mountains. Two blowing snow scheme coupled with Crocus exist yet : SYTRON (Vionnet et al., 2018) and Crocus-Meso-NH (Vionnet et al., 2014). However, both are unadapted to this geometry and resolution.

Thus, the goal of this paper is to describe and present first evaluations of a novel blowing snow scheme, SnowPappus, coupled to Crocus and able to be included in the above-mentioned large-scale simulation system. Point-scale evaluation of blowing snow flux will be presented to discuss the modelling choices"

In addition, L858-860 in the second part of the discussion will be replaced by :

"Snow redistribution in 2D simulations has not been evaluated in this article, due to several methodological challenges including dealing with the superposition of errors coming from the precipitation fields and finding relevant metrics. It will be the subject of a future study expected to provide complementary insights to the following discussions."

But the main problem is that this manuscript requires major copy editing for clarity and concision, beyond what can be achieved in review

As mentioned above, we propose a revised manuscript with very substantial changes, including a revised structure of the literature review and model description to improve the clarity of the manuscript. The length is also reduced by 20%.

I will expect to have more scientific comments on an easier to read revision, but here are a few corrections that I have noted:

There is, I think, a u* missing in equation 13

Indeed, the start of the equation will be corrected. The corrected one will start with : $Q_{susp}=c_r z_r u_*/(k (1 - gamma(u_*)))$

For reproducibility, quoting the values of F(T), A and B in equations 25 and 26 would save the reader some cross referencing. \mu_{satc} and Rh_i are essentially the same thing. Some signposting would help anyone hoping to find the SnowPappus code in the extensive SURFEX repository.

For reproducibility, we propose the following changes in the revised manuscript

- L 462, "A, B constants defined in (Gordon et al., 2006)" will be replaced by ", A = 0.0018 and B=3.6"
- in Eq. 26, 1-RH_i will be replaced by \mu_{satc}
- F(T) expression, which is quite complex, will be provided in appendix, to limit the increase of the size of the manuscript.
- -A Readme with path towards the most useful SURFEX routines for SnowPappus will be provided with the SURFEX repository.

Wind is measured at Huez and Chambon, so why no simulations forced by observed wind speed in figure 10?

We decided not to perform simulation with observed wind speed at Huez and Chambon as the wind data do not have the same quality as at the *Col du Lac Blanc* observatory. These date would required more treatment in order to get continuous wind time series. In particular, wind sensors are located at a fixed 3,5 m height above ground, so their height must be deduced from measured snow height. They are sometimes of very low quality, with numerous discontinuities and gaps.

Red points masked by the legend in figure 12a appear pink. Move the legend.

Will be fixed.

Vincent Vionnet is a co-author of this paper, so it should not be necessary to be so speculative about differences from Vionnet et al. (2018) in section 5.1.

We would like to apologize for this unclear formulation that did not correctly reflect our clear understanding of this issue. In fact we <u>were able to</u> retrieve the original simulation outputs from Vionnet et al. (2018) and found there was not any issue about it. As said L793 we "applied our evaluation process to these data[...] We obtained results very close to our own Sytron run with the original data". Regarding the dates, we were able to obtain the information on the precise temporal window used in Vionnet et al. (2018), which is from 01/11 to 15/04. Therefore, their temporal window is larger than ours, so it is not possible that the change in the temporal window by itself would cause a perfect detection in their case and not in ours. As a consequence, we are sure the reproducibility issue comes from an unreproducible data post-processing, that was applied when compiling the results at the daily time scale in Vionnet et al. (2018). This step was not applied in our evaluation dataset. To clarify this and taking into account the reviewer's call for concision, we propose to replace L789-794 by

"We were able to retrieve the original simulation outputs of Vionnet et al. (2018) and applied our evaluation process to these data (see code availability), obtaining results very close from ours. Thus, after discussion with the authors, it is clear that the issue comes from unreproducible data post-processing applied to the SPC data to compile results at the daily time scale." Of course, this unreproducible data processing is not satisfactory and we took a special care in this publication to respect the FAIR principles with all our data and provide all details in the Code and

REFERENCES

Data availability section to prevent such inconveniences.

Amory, C., Kittel, C., Le Toumelin, L., Agosta, C., Delhasse, A., Favier, V., and Fettweis, X.: Performance of MAR (v3. 11) in simulating the drifting-snow climate and surface mass balance of Adélie Land, East Antarctica, Geoscientific Model Development, 14, 3487–3510,830 <u>https://doi.org/0.5194/gmd-14-3487-2021</u>, 2021

Cluzet, B., Lafaysse, M., Deschamps-Berger, C., Vernay, M., and Dumont, M.: Propagating information from snow observations with CrocO ensemble data assimilation system: a 10-years case study over a snow depth observation network., Cryosphere Discussions, <u>https://doi.org/10.5194/tc-16-1281-2022</u>, 2021.

Deschamps-Berger, C., Gascoin, S., Berthier, E., Deems, J., Gutmann, E., Dehecq, A., ... & Dumont, M. (2020). Snow depth mapping from stereo satellite imagery in mountainous terrain: evaluation using airborne laser-scanning data. *The Cryosphere*, *14*(9), 2925-2940

Essery, R., Li, L., and Pomeroy, J.: A distributed model of blowing snow over complex terrain, Hydrological processes, 13, 2423–2438, https://doi.org/10.1002/(SICI)1099-1085(199910)13:14/15%3C2423::AID-HYP853%3E3.0.CO;2-U, 1999. Gallée, H., Guyomarc'h, G., and Brun, E (2001).: Impact of snow drift on the Antarctic ice sheet surface mass balance: possible sensitivity to snow-surface properties, Boundary-Layer Meteorology, 99, 1–19, <u>https://doi.org/10.1023/A:1018776422809</u> Gerber, F., Sharma, V., & Lehning, M. (2023). CRYOWRF-a validation and the effect of blowing

Gerber, F., Sharma, V., & Lehning, M. (2023). CRYOWRF-a validation and the effect of blowing snow on the Antarctic SMB. *Authorea Preprints*.

Gilleland, E., Ahijevych, D., Brown, B. G., Casati, B., & Ebert, E. E. (2009). Intercomparison of spatial forecast verification methods. *Weather and forecasting*, *24*(5), 1416-1430.

Günther, D., Marke, T., Essery, R. et Strasser, U. (2019). Uncertainties in Snowpack Simulations— Assessing the Impact of Model Structure, Parameter Choice, and Forcing Data Error on Point-Scale Energy Balance Snow Model Performance. Water Resour. Res., 55(4):2779–2800, <u>https://doi.org/10.1029/2018WR023403</u>.

Guyomarc'h, G. and Mérindol, L.: Validation of an application for forecasting blowing snow, Annals of Glaciology, 26, 138–143, <u>https://doi.org/10.3189/1998AoG26-1-138-143</u>, 1998

Largeron, C., Dumont, M., Morin, S., Boone, A., Lafaysse, M., Metref, S., Cosme, E., Jonas, T., Winstral, A., and Margulis, S. A.: Toward snow cover estimation in mountainous areas using modern data assimilation methods: a review, Frontiers in Earth Science, 8, 325, <u>https://doi.org/10.3389/feart.2020.00325</u>, 2020.

Lehning, M., Doorschot, J., and Bartelt, P.: A snowdrift index based on SNOWPACK model calculations, Annals of Glaciology, 31, 382–386,<u>https://doi.org/10.3189/172756400781819770</u>, 2000.

Liston, G. E. and Sturm, M.: A snow-transport model for complex terrain, Journal of Glaciology, 44, 498–516, https://doi.org/10.3189/S0022143000002021, 1998.

Liston, G. E., Haehnel, R. B., Sturm, M., Hiemstra, C. A., Berezovskaya, S., and Tabler, R. D.: Simulating complex snow distributions in windy environments using SnowTran-3D, Journal of Glaciology, 53, 241–256, <u>https://doi.org/10.3189/172756507782202865</u>, 2007.

Menard, C. B., Essery, R., Krinner, G., Arduini, G., Bartlett, P., Boone, A., ... & Yuan, H. (2021). Scientific and human errors in a snow model intercomparison. *Bulletin of the American Meteorological Society*, *102*(1), E61-E79.

Morin, S., Horton, S., Techel, F., Bavay, M., Coléou, C., Fierz, C., Gobiet, A., Hagenmuller, P., Lafaysse, M., Ližar, M., Mitterer, C., Monti, F., Müller, K., Olefs, M., Snook, J. S., van Herwijnen, A., and Vionnet, V.: Application of physical snowpack models in support of operational avalanche hazard forecasting: A status report on current implementations and prospects for the future, Cold Regions Science and Technology, 170, 102 910, <u>https://doi.org/10.1016/j.coldregions.2019.102910</u>, 2020b.

Raleigh, M. S., Lundquist, J. D. et Clark, M. P. (2015). Exploring the impact of forcing error characteristics on physically based snow simulations within a global sensitivity analysis framework. Hydrol. Earth Syst. Sci., 19(7):3153–3179. <u>https://doi.org/10.5194/hess-19-3153-2015</u>.

Schlögl, S., Marty, C., Bavay, M. et Lehning, M. (2016). Sensitivity of Alpine3D modeled snow cover to modifications in DEM resolution, station coverage and meteorological input quantities. Environmental Modelling & Software, 83:387–396. <u>https://doi.org/10.1016/j.envsoft.2016.02.017</u>

Schweizer, J., Jamieson, J. B., and Schneebeli, M.: Snow avalanche formation, Reviews of Geophysics, 41, <u>https://doi.org/10.1029/2002rg000123</u>, 2003.

Sharma, V., Gerber, F., and Lehning, M.: Introducing CRYOWRF v1. 0: Multiscale atmospheric flow simulations with advanced snow cover modelling, Geoscientific Model Development Discussions, pp. 1–46, <u>https://doi.org/10.5194/gmd-2021-231</u>, 2021

Viallon-Galinier, L., Hagenmuller, P., & Lafaysse, M. Forcing and evaluating detailed snow cover models with stratigraphy observations. *Cold Regions Science and Technology*, *180*, 103163, 2020

Vionnet, V., Guyomarc'h, G., Bouvet, F. N., Martin, E., Durand, Y., Bellot, H., Bel, C., and Puglièse, P.: Occurrence of blowing snow events at an alpine site over a 10-year period: Observations and modelling, Advances in water resources, 55, 53–63, <u>https://doi.org/10.1016/j.advwatres.2012.05.004</u>, 2013.

Vionnet, V., Martin, E., Masson, V., Guyomarc'h, G., Naaim-Bouvet, F., Prokop, A., Durand, Y., and Lac, C.: Simulation of wind-induced snow transport and sublimation in alpine terrain using a fully coupled snowpack/atmosphere model, The Cryosphere, 8, 395–415, <u>https://doi.org/10.5194/tc-8-395-2014</u>, 2014.

Vionnet, V., Guyomarc'h, G., Lafaysse, M., Naaim-Bouvet, F., Giraud, G., and Deliot, Y.: Operational implementation and evaluation of a blowing snow scheme for avalanche hazard forecasting, Cold Regions Science and Technology, 147, 1–10, <u>https://doi.org/10.1016/j.coldregions.2017.12.006</u>, 2018.

Walter, B., Weigel, H., Wahl, S., and Löwe, H.: Wind tunnel experiments to quantify the effect of aeolian snow transport on the surface snow microstructure, The Cryosphere Discuss. [preprint], <u>https://doi.org/10.5194/tc-2023-112</u>, in review, 2023.