

Supplement for

Application of the Multi-Scale Infrastructure for Chemistry and Aerosols version 0 (MUSICAv0) for air quality in Africa

Wenfu Tang¹, Louisa K. Emmons¹, Helen M. Worden¹, Rajesh Kumar², Cenlin He², Benjamin Gaubert¹, Zhonghua Zheng³, Simone Tilmes¹, Rebecca R. Buchholz¹, Sara-Eva Martinez-Alonso¹, Claire Granier^{4,5}, Antonin Soulie⁴, Kathryn McKain^{5,6}, Bruce C. Daube⁷, Jeff Peischl^{5,8}, Chelsea Thompson⁸, and Pieter Levelt^{1,9,10}

¹Atmospheric Chemistry Observations & Modeling Laboratory, National Center for Atmospheric Research, Boulder, CO, USA

²Research Applications Laboratory, National Center for Atmospheric Research, Boulder, CO, USA

³Department of Earth and Environmental Sciences, The University of Manchester, Manchester M13 9PL, United Kingdom

⁴Laboratoire d'Aérodologie, CNRS, Université de Toulouse, Toulouse, France

⁵Cooperative Institute for Research in Environmental Sciences (CIRES), University of Colorado, Boulder, CO, USA

⁶Global Monitoring Laboratory (GML), National Oceanic and Atmospheric Administration, Boulder, CO, USA

⁷Department of Earth and Planetary Sciences, Harvard University, Cambridge, MA, USA

⁸NOAA Chemical Sciences Laboratory, Boulder, CO, USA

⁹Royal Netherlands Meteorological Institute (KNMI), Utrechtseweg 297, 3730 AE De Bilt, the Netherlands

¹⁰University of Technology Delft, Mekelweg 5, 2628 CD Delft, the Netherlands




Figure S1. Comparisons of MUSICA v0 and WRF-Chem simulations and OMI tropospheric NO₂ column (molecules/cm²) in 2017. (a-d) Averaged OMI tropospheric NO₂ column in MAM (March, April, and May), JJA (June, July, and August), SON (September, October, and November), and DJF (December, January, and February). (e-h) MUSICA v0 model biases against OMI tropospheric NO₂ column in MAM, JJA, SON, and DJF. (i-l) is the same as (e-h) but for WRF-Chem. All data are gridded to 0.25 degree × 0.25 degree for plotting.




Figure S2. Comparisons of MUSICAv0 and WRF-Chem simulations and MODIS AOD in 2017. (a-d) Averaged MODIS AOD in MAM (March, April, and May), JJA (June, July, and August), SON (September, October, and November), and DJF (December, January, and February). (e-h) MUSICAv0 model biases against MODIS AOD in MAM, JJA, SON, and DJF. (i-l) is the same as (e-h) but for WRF-Chem. All data are gridded to $0.25 \text{ degree} \times 0.25 \text{ degree}$ for plotting.




Figure S3. Comparisons of MUSICA v0 and WRF-Chem simulations and OMI tropospheric CH_2O column ($\text{molecules}/\text{cm}^2$) in 2017. (a-d) Averaged OMI tropospheric CH_2O column in MAM (March, April, and May), JJA (June, July, and August), SON (September, October, and November), and DJF (December, January, and February). (e-h) MUSICA v0 model biases against OMI tropospheric CH_2O column in MAM, JJA, SON, and DJF. All data are gridded to 0.25 degree \times 0.25 degree for plotting.




Figure S4. Monthly time series of CO tracers of emissions from North Africa (green), West Africa (pink), East Africa (orange), Central Africa (blue), Southern Africa (yellow), and the rest of the world (grey) at (a) Assekrem, (b) Cape Point, (c) Izana, (d) Gobabeb, (e) Mare and (f) Ascension. The locations of the sites and the definition of the regions can be found in Figure 1.




Figure S5. Monthly time series of CO tracers of fire emissions (red), anthropogenic emissions (green), and waste burning emissions (yellow) at (a) Assekrem, (b) Cape Point, (c) Izana, (d) Gobabeb, (e) Mare and (f) Ascension. The locations of the sites can be found in Figure 1.




Figure S6. Modeled lightning NO emissions (g N/month/m^2) from the MUSICAv0 simulation in MAM (March, April, and May), JJA (June, July, and August), SON (September, October, and November), and DJF (December, January, and February).




Figure S7. Vertical profiles of air temperature (T; K) and water vapor mixing ratio (H_2O ; ppm) from the In-service Aircraft for a Global Observing System (IAGOS) measurements (black) and corresponding model output from MUSICA v0 (red), and WRF-Chem (blue) during 2017. Annual mean T profiles with the variation of the data in the pressure layer (25% quantile to 75% quantile) over (a) North Africa, (b) West Africa, (c) Central Africa, (d) East Africa, and (e) Southern Africa are shown. (f-j) are the same as (a-e) but for H_2O .




Figure S8. Spatial distribution of Taylor score of MUSICAv0 compared to satellite retrievals in each 5 degree \times 5 degree (latitude \times longitude) pixels. Taylor score of MUSICAv0 compared to MOPITT CO column retrievals for (a) 2017, (b) MAM (March, April, and May), (c) JJA (June, July, and August), (d) SON (September, October, and November), and (e) DJF (December, January, and February) are shown. (f-j) are the same as (a-e) but for Taylor score of MUSICAv0 compared to OMI tropospheric NO₂ column retrievals; (k-o) are the same as (a-e) but for Taylor score of MUSICAv0 compared to MODIS AOD; (p-t) are the same as (a-e) but for Taylor score of MUSICAv0 compared to OMI tropospheric HCHO column retrievals.