

# A modelling system for identification of maize ideotypes, optimal sowing dates and nitrogen fertilization under climate change – PREPCLIM-v1

Mihaela Caian<sup>1</sup>, Catalin Lazar<sup>2</sup>, Petru Neague<sup>1</sup>, Antoanelia Dobre<sup>1</sup>, Vlad Amihaesei<sup>1</sup>, Zenaida Chitu<sup>1</sup>,  
5 Adrian Irasoc<sup>1</sup>, Andreea Popescu<sup>1</sup>, George Cizmas<sup>2</sup>

<sup>1</sup> National Meteorological Administration Romania (NAM), Sos. Bucureşti-Ploieşti nr.97, Sector 1, 013686 Bucureşti România

<sup>2</sup> – National Agricultural Research and Development Institute (NARDI) Fundulea, 915200 Călăraşi, România

*Correspondence to:* Mihaela Caian (mihaela.caian@gmail.com)

10 **Abstract.** Climate change significantly threatens crop yields levels and stability. The complex interplay of factors at the local scale makes assessing these impacts difficult, requiring coupled climate-phenology models, which integrate climate data and crop information. Identifying suitable local management practices and crop varieties under future conditions becomes essential for developing effective adaptation strategies.

This study presents the implementation and application of an integrated climate-phenology adaptation support modelling 15 system. This is based on regional CORDEX climate models and the CERES Maize model from the DSSAT platform. Novel modules for optimal management and genotype identification under climate change have been developed in the system, employing a hybrid approach that combines deterministic modelling with machine learning (ML) techniques and genetic algorithms. This system was run as a regional pilot over Romania, operating in real-time in interaction with users, performing agro-climate projections (combination of fertilization, sowing date, genotype) and providing best crop 20 management simulated under climate change projections. Multi-model ensemble simulations were conducted for two radiative forcing scenarios RCP4.5 and RCP8.5 and twelve management scenarios, yielding novel results for the region. Results indicate a projected decrease in maize yields for the current genotype across all tested scenarios, primarily attributed to a shortened grain-filling period and reduced fertilization efficiency under warmer conditions. Soil initial conditions were found to significantly influence yield responses.

25 The analysis warns about a projected narrowing of the agro-management options for maintaining a high yield level. However, we find an added value from the impact of genotype selection in mitigating climate change impacts, even in extreme years. Genotype optimisation across six crossed cultivar dependent parameters revealed that while maximum yields declines, specific genotype windows exhibit increased intermediate yields under future climates compared to current 30 conditions. Sensitivity analysis identified the thermal time requirements during juvenile and maturity stages as the most critical factors influencing genotype performance under warmer climates.

This research demonstrates the added value of combining deterministic and data-driven modelling approaches within a coupled climate-crop system for developing effective adaptation strategies, including optimised fertilization pathways that contribute to climate change mitigation.

## 1. Introduction

According to the IPCC (2022), climate change is unequivocal, and its impacts appear more worrying today than decades ago. While research on the effects of climate change on crop yields and agricultural harvests has advanced (Arnell and Freeman, 2021; Hatfield et al., 2021; Rezaei et al., 2024), translating these findings into actionable solutions and scales 40 remains a challenge. This is primarily due to the high complexity of factors that intervene at the local scale of the crop (Malhi et al., 2021; Eyring et al., 2021) including sensitivities of the exchanges to variations in climate sub-components as atmosphere / soil/ biosphere's ecosystems under climate change, natural causes and human activities (Wheeler and Braun, 2013; Xie et al., 2023).

Given the projected global population increase estimated in scientific reports to over 9 billions by 2050 (Godfray and 45 Charles, 2010), global food production would have to increase by 70-100% to meet the growing demand (Smil, 2005; World Development Report, 2008; Selvaraju et al., 2011). This challenge is further compounded by the agro-climatic conditions expected to become vulnerable and gradually decline due to climate change, particularly impacting water availability (Stehr and von Storch, 2009; Villalobos et al., 2012; van Ittersum et al., 2013).

Another challenge of the problem comes from the need that approaches, and sustainable solutions must not only address the 50 needs of agricultural producers but also align with climate change mitigation goals for 2050, aiming for climate neutrality (Semenov and Strattonovitch, 2015; Dainelli et al., 2022; Mitchell et al., 2022).

Early studies investigating the impact of climate change on crop yields emphasized the necessity of high-resolution modelling approaches. These models should accurately represent management practices and the local effects of climate variables, such as temperature and precipitation (McKee et al., 1993; Trnka et al., 1995; Adams et al., 1998). These affect 55 thermal and water stress and plant physiological processes like stem water potential, stomatal opening, leaf transpiration efficiency (Espadafor et al., 2017). At the regional scale, the relationship between crop yield and water and thermal availability may exhibit strong dependencies on the crop type, geographical location, temporal scale, and plant developmental stage (Webber et al., 2018, 2020; Marcinkowski and Piniewski, 2018; Berti et al., 2019; Ceglar et al., 2020; Wu et al., 2021). For instance, simulations conducted by Kothari et al. (2022) in regions with arid climates, indicated for 60 future climate change a significant (~30%) decrease without adaptation, but a potential increase (15%) in corn yields under irrigated or under radiation-based genotype efficient use. These findings underscore the critical need for regional simulations that incorporate phenological characteristics with accurate soil moisture estimates to evaluate the effectiveness of various irrigation strategies under future climate scenarios.

In addition to atmospheric conditions, soil properties significantly influence plant growth. These influences occur through 65 physics-based interactions with climate and through alterations in soil chemical composition. Rising air temperatures have been shown to impact the soil carbon budget, with a decline in soil carbon potentially affecting plant and root processes, biochemical cycles, and species composition (Abhik Patra et al., 2021).

Crop modelling at local, regional and global scale has significantly advanced, enhancing our understanding of crop systems and enabling the simulation and projection of future yields. Studies (Tsvetsinskaya et al, 2001; Tao et al., 2009; Ganguly et al., 2013; Schuberger et al., 2020; Chen and Tao, 2022) consistently project global mean harvest reductions with differences in the regional pattern of climate change impact on crop and harvest (Asseng et al., 2015; Li et al. 2022). Not only projected spatial but also temporal variability of the climate change impact appears larger and accelerated, motivating intensified efforts on seasonal and multi-annual predictions of plant development and harvest (Baez-Gonzalez et al., 2005; Jin et al., 2022). Analysis of these simulations emphasized also the need to include crop uncertainty in climate scenarios assessments (Meehl et al., 2007, Rosenzweig et al. 2013, Basso Bruno et al., 2019; Chapagain et al., 2022).

Meanwhile, model simulations emerged as useful tool in plant breeding analysis (Bernardo, 2002; Banterng et al, 2004; Cooper and Messina et al., 2023; Mamassi et al., 2023), supporting the development of superior genotypes and breeding methods for maximizing crop performance. These simulations have proven effective in guiding cultivar selection through techniques such as parental selection and breeding by design (Peleman and van der Voort, 2003; Qiao et al., 2022).

In most recent years climate-crop modelling extended from deterministic crop models (Boogaard et al. 2013; Morell et al., 2016) to data-driven techniques approaches for assessing crop response to weather and climate change (Schwalbert et al., 2020; Meroni et al., 2021; Morales and Villalobos, 2023; Chang et al., 2023; Zhuang et al., 2024). Statistical methods as well as machine learning (ML) used for crop forecast and modelling were however shown to bring for now, limited benefits (Paudel et al., 2021), pointing to possibly hybrid techniques that include physical process in the modelling as a key approach for this challenging issue. On the other hand, breeding optimization techniques using fully deterministic model simulations require a huge number of simulations, analysis and inter-comparisons of predicted crop performance (Pfeiffer et al., 2007; Wang et al., 2023).

Here we present a novel hybrid approach developed in the frame of the PREPCLIM (“Preparing for climate change”) project in which we solve plant phenology using deterministic modelling and merge this technique with an on-line ML-genetic algorithms (GA) iteratively selecting along simulations the cross-range of optimal crop cultivar parameters, according to user-defined criteria for optimal target. The GA simulates the evolution of a population by applying in iterations, genetic operators (selection, crossover, mutation) to a set of candidate solutions (chromosomes). The chromosomes represent potential solutions to the problem and are encoded as strings of binary or symbolic values, with their fitness assessed by a problem-specific evaluation function here, user-required based. GAs have demonstrated success for optimizing agricultural practices using models like DSSAT for irrigation and fertilizer applications (Bai et al., 2021; Wang et al., 2023).

The hybrid approach implemented in this work focused on ideotype identification presents the advantage of physically treating the crop complex process involved along optimizing iterations, thus allowing specific inclusion and understanding of physical causes of responses and of optimal paths in various climate and management scenarios. Furthermore, it enhances the ability of choosing optimum conditions from continuous multi-dimensional intervals for gene parameters, as opposed to discrete sets. The continuum values approach is an important feature mainly for isolated extreme yield detection, or broader parameters’ range and high non-linearity, both aspects of increasing relevance in the context of climate change. Our findings

suggest a narrowing of agro-management adaptation opportunities under warmer climates, further emphasizing the importance of this hybrid genotype-agro-management approach to support finding solutions for the future.

The developed system aims to provide efficient and operational support for farmers and stakeholders. It leverages the state-of-the art DSSAT model, a widely used and extensively validated platform for agricultural modelling across diverse applications. The DSSAT model, incorporating complex parameterizations for soil processes, surface-atmosphere exchange, plant development stages, and their interactions with climate and management practices, undergoes continuous refinement through ongoing research and regional calibrations. For this study, the model was specifically adapted to the unique soil characteristics of the pilot region, including parameters such as porosity, composition per soil layers, and thermal properties.

The developed system exhibits portability to other regions with available soil and management data. Its functionality and user-friendliness are expected to improve through widespread adoption and the incorporation of advanced user requests and management options.

Section 2 presents the developed system and its data flow. Section 3a provides the motivation for system development, focusing on projected climate change impacts for the target region. Section 3b presents results obtained using the system to simulate projected changes in plant phenology and crop parameters for the target region, under various climate and management scenarios, for the current control genotype. Section 3c discusses results obtained using the system's genotype optimization package along agro-management scenarios. Finally, Section 4 presents perspectives and conclusions.

## 2. Data and methods

Projected changes in agro-climatic parameters for Romania were assessed under two Representative Concentration Pathways (RCPs): RCP4.5 and RCP8.5. These changes were computed as anomalies relative to historical simulations (Hist) using an ensemble of three CMIP5-CORDEX (Benestad et al., 2021, Karl et al., 2011) high resolution (11 km) climate models, based on the CNRM, EC-EARTH, and MPI global models coupled to the regional climate model RCA4. Subsequently, the DSSAT crop model (Jones et al., 2003; Hoogenboom et al., 2019) was employed to simulate projected changes in phenological and harvest parameters. The DSSAT model was driven by atmospheric conditions derived from each model of the ensemble for the historical period and for the two RCP emission scenarios.

A software package was developed for the DSSAT model that performs identification of optimal model parameters based on user-defined: criteria for optimum, climate-management scenario, region, and time horizon. Optimization goals include maximizing harvest, ensuring stable yields over time, and minimizing nitrogen leaching beyond the root zone (reducing water pollution risk). Management scenarios allow users to explore optimal cross-combinations of sowing dates, fertilization amounts, and genotypes.

Six main cultivar-specific parameters (P1 to P6) characterizing the maize genotype were analysed across wide ranges of physically realistic values, considering both current and extreme future climate projections for the target area. P1 represents the thermal time from seedling emergence to the end of the juvenile phase (ranging in these simulations from 100 to 500-

degree days above 8°C). It significantly influences crop flowering times (Liu et al., 2020), water availability, and ultimately, 135 yield. Studies have shown that utilizing longer-season maize cultivars (dependent also on P1) can lead to increased harvest in humid regions but decreased harvest in semi-humid regions (Mi et al., 2021).

P2, the photoperiod-development delay parameter (ranging in simulations here from 0.1 to 2.6 days) shows the extent to which development is delayed for each hour of photoperiod increase above the longest photoperiod of maximum 140 development rate (considered 12.5 hours). P2 influences the flowering time (Langworthy et al. 2018) and the rate of plant development, with long-day plants exhibiting faster development under longer day lengths (Angus et al., 1981). Related to these, studies have demonstrated the significant role of P2 in mitigating the negative impacts of waterlogging in warmer 145 climates (Liu et al., 2023). P3, the thermal time from silking to physiological maturity (tested here for values from 500 to 1500-degree days above 8°C), significantly influences maturity dates. It also has a main role in plant stress levels (longer-maturity hybrids increase harvest but under water stress it may provide lower yield (Su et al., 2021; Grewer et al, 2024)) and grain moisture at maturity (Tsimba et al., 2013). P4, representing the maximum number of kernels per plant, exhibits a 150 relatively predictable numerical response and is therefore held constant at the control value of 797.5 estimated for the region, in this analysis. P5, the kernel filling rate parameter (ranging from 6 to 12 mg/day), influences grain filling duration, desiccation, moisture at maturity and harvest (Chazarreta et al., 2021). P6, the phyllochron interval or the thermal time between successive leaves (ranging from 3 to 70 °C) is a critical parameter for estimating the duration of vegetative 155 development (Birch et al., 1998; Xu et al., 2023). P5 and P6 are important parameters of optimal plant adaptation to climate conditions, since they are drivers of the phenological response and yield formation, in conjunction with the temperature, radiation, humidity, water stress. These genotype (or cultivar specific) parameters are the primary ones considered in DSSAT model parameterizations for plant development processes (Hoogenboom et al., 2019).

The parameter ranges were rigorously tested in simulations to ensure their representatives for the target region, including an 155 analysis of extreme values. The control values for these cultivar-specific parameters  $P_i$  in the region are:  $P_1=200$ ,  $P_2=0.7$ ,  $P_3=800$ ,  $P_4=797.5$ ,  $P_5=8.60$ , and  $P_6=38.90$ . All the simulations for combinations of parameters values (cross- $P_i$  simulations) were performed under Hist, RCP4.5, and RCP8.5 emission scenarios. For each scenario, simulations were conducted for twelve agro-management scenarios consisting of sowing date changes and fertilization treatments, for each model of the ensemble.

By default, the twelve agro-management scenarios encompass four sowing dates (spaced five days apart) and three 160 fertilization levels (zero, then a regional average and its double). For each agro-management scenario, genotype optimization (finding the optimal set of  $P_i$  values under the given climate -agro-management and optimum criteria) was performed using two methods: 1) discretized parameter-space runs with subsequent post-processing ordering, and 2) continuum parameter-space search with iterative selection during simulations, employing genetic algorithms (GA).

The proposed GA-based method employs an iterative approach. It commences with an initial population of randomly 165 generated solutions (chromosomes) and undergoes iterative cycles (generations). In each generation, a selection process is performed to choose the fittest chromosomes for reproduction, based on their fitness scores. Subsequently, crossover (recombination) and mutation operators are applied to the selected chromosomes, generating offspring that inherit traits from


their parents. The new offspring replace some of the least fit individuals in the population, ensuring that the average fitness of the population improves over time. The convergence of the GA toward an optimal or near-optimal solution is achieved by 170 balancing exploration (searching the problem's space for diverse solutions exploiting promising regions) and exploitation (refining the best solutions found so far). Here GA has been newly applied to develop an innovative crop selection algorithm, optimizing genotypes across various agro-management scenarios. Steps along the algorithms are described in Annex.

The system generates output data (agro-climate and optimal paths) which is disseminated on two platforms (Fig.1). One is a 175 platform (Info-Platform, Fig.1a) providing one-way interactive (static) agro-climate information at local scale (NUTS3 level, aligned with the European Union's Nomenclature of Territorial Units for Statistics) over the region. It delivers pre-computed climate -agro-climate indicators, indices of agro-climate extremes based on observations, re-analysis and climate scenarios for future projections for the region.


The second platform (User-Platform, Fig.1b) is an operational, online, user-interactive (two-way) in real-time component, 180 where user requests are submitted, processed as input to the modelling chain and results delivered back to the user for a new, refined request.

The core of the modelling system integrates the DSSAT crop model (running on Linux OS) with regional climate models (Fig.2), with a pre-processing pack developed for coupling. This coupled system incorporates new features, that include the 185 ability of conducting parameter-varying cross-simulations and advanced algorithms for identifying optimal agro-management practices and genotype selections along simulations.

a)



b)



190 Fig.1 a): Info-Platform: Provides local-regional scale information derived from high-resolution regional climate models (CORDEX), e.g. climate, agro-climate data and indicators, indices of agro-climate extremes at the NUTS3 level. b): User-Platform for adaptation support: Processes in real time specific user requests, and simulates management scenarios, identifying optimal paths: Users input parameters (left, e.g: region, period (present / future climate scenarios), management options (e.g: sowing date, fertilization/irrigation time and amount, genotype); System Output (right, e.g: harvest, projected phenology dates, precipitation/evapotranspiration, Nitrogen and carbon balances, optimal management paths (dates and management actions), optimal genotype) estimated from ensemble simulations.

195




Fig.2: The PREPCLIM-v1 work schema: DSSAT-core and modelling components (middle), and data flow: input data (left), output information (right). Red modules were developed in PREPCLIM-v1.

200 The system was implemented and validated over Southern Romania, target agricultural area, for maize. Potential beneficiaries include researchers, farmers, policymakers, and maize breeders. The system can also assist maize breeders in adapting to climate change by enabling them to evaluate and select genotypes more resistant to challenging climatic conditions. Given the accelerating pace of climate change, such a system may provide valuable support in numerous ways.

205 **Table 1: The agro-management treatments (TR): each treatment is described in terms of the sowing date and fertilization amount, Nitrogen [kg/ha]. We denote two experiments: exper “1N” and exper “3N”, and fertilisations Fx0, Fx1, Fx2 have values dependent on the experiment: Fx0 is no fertilisation, Fx1 is the unit fertilisation of the experiment and Fx2 is the double unit fertilisation of the experiment. We define the unit fertilisation of the exper “1N” equal to 23 N/kg and the unit fertilisation of the exper “3N” as 60 kg/ha. Sowing date format is “DD.MM”.**

Treatment	TR1	TR2	TR3	TR4	TR5	TR6	TR7	TR8	TR9	TR10	TR11	TR12
Sowing date	1.04	15.04	1.05	15.05	1.04	15.04	1.05	15.05	1.04	15.04	1.05	15.05
Fertilization (exper “3N”)	Fx0 =0	Fx0 =0	Fx0 =0	Fx0 =0	Fx1=60 =60	Fx1=60 =60	Fx1=60 =60	Fx1=120 =120	Fx1=120 =120	Fx1=120 =120	Fx1=120 =120	Fx1=120 =120
Fertilization (exper “1N”)	Fx0 =0	Fx0 =0	Fx0 =0	Fx0 =0	Fx1=23 =23	Fx1=23 =23	Fx1=23 =23	Fx1=46 =46	Fx1=46 =46	Fx1=46 =46	Fx1=46 =46	Fx1=46 =46

### 3. Results

#### 210 3.1 Agro-climate changes in the region

##### 3.1.1 Climate changes in agro-climate indicators

Agro-climatic Indicators, derived from CORDEX models and available on the Info-Platform, provide time-series data for ensemble or individual model metrics at the NUTS3 level across Romania. Figure 3 illustrates projected changes in key agro-climatic characteristics.

215 The anticipated climate shift in the region is evidenced by changes in the Johansson Continentality Index (JCI, Fig.3a), calculated as  $JCI = 1.7 * dT / \sin(\phi) - 20.4$  (where  $dT$  is the annual maximum range of monthly mean temperatures and  $\phi$  is latitude; (Flocas, 1994; Baltas, 2007)). Changes in the JCI generally reveal robust evidence of large-scale changes influences on the regional climate. For this domain it shows a Southwards meridional gradient of the intra-annual variability (Arctic amplification remote impacts on Europe). Hence enhanced intra-annual variability (JCI) with much warmer summers than 220 winters over the main agricultural areas in South (and the opposite in the North), information useful for farmers to estimate changes in the sowing time.

In agreement with this, the Scorching Index (H32temp, Fig.3b, computed as the total degrees in summer days exceeding 32°C), used by farmers and agro-meteorologists to characterize the sub-regional drought conditions, projects severe drought conditions ( $H32temp \geq 51$ ), about doubling the Hist values and expanding significantly across the southern regions in 225 RCP8.5 with already high-level drought conditions ( $31 < H32temp < 51$ ) occurring in RCP4.5 (Fig.3b).

Accounting also for precipitation changes, the de Martonne Aridity Index (IM, the ratio of annual precipitation to a translation function (+10C) of the annual mean temperature), exhibits also significant projected changes. It shows

particularly increased aridity (low IM) in the south, southeast, and southwest regions, the major agricultural areas with an accelerating change up to 2100 (Fig.4, comparing projected differences to Hist for 2071-2100 versus 2021-2050).

230

a)





Fig.3: The JCI and the Scorching index H32temp indices. For each: (left): the index over Hist period 1971-2000 and changes (2021-2050) relative to it, under RCP4.5 (middle) and RCP8.5 (right). a) The JCI climate index classes are: marine for  $0 < k \leq 33$ , continental for  $33 < k \leq 66$  and exceptionally continental for  $66 < k \leq 100$ . b) The Scorching index H32temp classes are: reduced intensity drought for  $H32temp \in [0,10]$ , moderate intensity for  $H32temp \in (10,30]$ , high intensity for  $H32temp \in (30,50]$  and severe drought conditions for  $H32temp > 50$ .

235




240

Fig.4: The de Martonne aridity index (IM) for: Hist (left), RCP4.5 (middle) and RCP8.5 (right) for two horizons: 2021-2050 (top) and 2071-2100 (bottom). IM index classes are: arid for  $0 < IM \leq 10$  arid, semi-arid for  $10 < IM \leq 20$ , Mediterranean for  $20 < IM \leq 24$ , semi-humid for  $24 < IM \leq 28$ , wet for  $28 < IM \leq 36$ , very wet for  $36 < IM \leq 55$  very wet and extreme wet for  $IM > 55$ ; (all indices are time mean 30 years, ensemble mean).


245 3.1.2 Changes in agro-climate extremes

Projected changes in extremes for temperature and precipitation, highly useful information for agriculture, show important features in the region. A main aspect of interest is related to late-spring freezing days that may drastically affect the whole crop of the year. Fig.5a shows for South Romania (Călărași subregion) that in spite of the decreasing trend (5% p-level significance) of the total number of freezing days in spring, we still may have severe events with interestingly, a number of 250 freezing days in late-spring that is even higher in scenarios than in Hist, late-spring being one of the most vulnerable period to freezing for the plant already under development. Also note that successive extreme freezing years in late spring may occur.

a)



255 b)



c)




Fig.5: Extreme climate parameters (NUTS region 103032, representative for the target region), along historical (Hist) and RCP8.5 scenarios; a): FD, the number of frost days (minimum temperature  $< 0^{\circ}\text{C}$  in a 10-day period); b): RR10, the number of days with

260

heavy precipitation (>10 mm per day) in a 10-day period; c): RR, total precipitation (mm) per 10-day period; (left): the 10-day period is centred on April 5<sup>th</sup>; (right): the 10-day period is centred on April 25<sup>th</sup>. Values indicate the slope of the linear trend (black line) and the p-value of significance (p-values < 0.05 are statistically significant at the 5% level).

This late spring blizzard feature over the region, was analysed in a previous work and shown to be related to the combined context of Polar Jet instability meanwhile with warmer sea surface temperature in the Eastern Mediterranean (Caian and Andrei, 2019). Both these features are projected to enhance in a warmer climate (Lelieveld et al., 2012; Shaw and Miyawaki, 2024), indicating higher potential for severe late-spring blizzards in the region.

For precipitation, analysis of extreme precipitation events (RR10mm) and total precipitation (RR) reveals a notable shift in their temporal distribution within April. While a decreasing trend is observed in the first dekad, a positive trend emerges in the third (and second, not shown) dekades (Fig.5b). These suggest a time-shift tendency towards the end of April and into early May for the occurrence of intense and accumulated precipitation. Although statistically insignificant at the 5% level, this shift is consistently observed across all models within the CMIP5 ensemble. As for FD we note that higher extreme values of RR and RR10 are projected to occur under emission scenarios than Hist, mainly in RCP8.5 (Fig.5b,c), more often during late spring. Extreme daily precipitation is, in most cases detrimental for the crop, causing soil erosion and surface runoff mainly after drought periods.

### 3.2 Phenology and Harvest Projections for the Control Genotype

Projected changes in phenology for the control genotype (Pioneer 375) were simulated using the DSSAT model under historical (Hist) and multi-model climate projections of CRP4.5 and RCP8.5 scenarios. Further, multi-genotype simulations are discussed in Section 3.c.

Model validation was conducted using Control simulations (Ctrl) driven by ERA5 reanalysis data (Simmons et al., 2021) for each treatment outlined in Table 1 (experiment "3N").


These simulations, spanning the period 1976-2005, demonstrate the model's ability to capture inter-annual variability in harvest yields, including both high and low yield years, when compared to the measured available data for the region (suppl. S1). They also allowed model set-up improvements through sensitivity simulations and model calibration including soil parameters (suppl. S2) as soil water, nitrogen, and organic carbon content.

However, further improvements in model accuracy are to be achieved if incorporating factors such as inter-annual soil variability, the yearly impact of pests and diseases or the year-to-year variations in practices of fertilization and sowing dates. For example, simulations with fertilization specifications closer to the year's management practices (e.g. approximately 80-120 kg N/ha and sowing around April 15<sup>th</sup> for 1995) resulted in more accurate (reduced bias) predictions (TR6, TR10). These, together with the well simulated inter-annual variability, demonstrate the model's ability to capture the combined influence of climate and management practices on crop performance.

#### 3.2.1 Phenology dates - projected changes

Ensemble model simulations provide projected changes in phenology, for the control genotype (G0), under different 295 fertilization levels (0, 60, 120 kg/ha, Table 1, exper "3N") and sowing dates, averaged over 30-years, in scenarios (2021-2050, RCP4.5 and RCP8.5), versus Hist (experiment set-up "E\_3N\_G0"). Figure 6a,b illustrates the ensemble model changes, demonstrating an earlier anthesis date by up to ~6 days and an earlier maturity date by up to ~10 days across all 300 scenarios. These time-shifts result in a shortening of the grain-filling period by up to 10% across the ensemble, and are a consistent response observed in each individual model. Early sowing dates exhibit a more pronounced earlier shift in anthesis under warming scenarios, a response even more pronounced under RCP8.5.

Under warmer climates we note more frequent occurrences of critical situations with suboptimal grain filling and potential 305 crop failure, under fertilization. These were linked in previous studies to non-linear interactions between fertilization and temperature (Huang et al., 2024) with excessive fertilization during reproductive stages under elevated temperatures potentially inducing higher stress conditions. In our study premature ending of simulated vegetation season occurred more frequently in treatments with higher nitrogen fertilization, leading in average only small changes in maturity days. This may favour leaves development, enhanced transpiration and earlier depletion of the soil moisture leading later to water stress.



310 **Fig.6: Simulated a): anthesis dates ([dap], days after sowing) and b): maturity dates ([dap]), under historical conditions (black), RCP4.5 (blue), and RCP8.5 (red) scenarios, experiment setup E\_3N\_G0. Results are shown for the four sowing dates and nitrogen fertilization level of 60 kg/ha (Table 1, exper "3N").**

### 3.2.2 Harvest - projected changes

For harvest, the ensemble simulations (along E\_3N\_G0) project an overall decrease under both RCP4.5 and RCP8.5 scenarios and across all sowing dates and fertilization levels (Fig.7), compared to the historical period.

315 Harvest decline in climate scenarios is related to several factors: 1) reduced rainfall during the growing season (Fig.8), as evidenced by a strong correlation (0.5 in April to 0.8-0.9 in July-August, over 30 years) found between harvest (H) and accumulated precipitation in the Ctrl and in model simulations; 2) a shortened grain-filling period due to a projected earlier

flowering and an even earlier maturity across all the models (Fig.6), potentially limiting biomass accumulation; and 3) decreased fertilization efficiency under warming conditions, in the sense that H difference Hist minus scenario, increases 320 (non-linearly) with enhanced fertilisation (Fig.7). Hence for a same climate, the same increase in fertilisation brings less benefit in a warmer climate. This benefit for H is of about 10% in Hist versus 7.6% in RCP8.5 for early sowing and about 8% in Hist versus 4.3% in RCP8.5 for later sowing for doubling the N amount of nitrogen (Fig.7b,c). This efficiency decay feature underscores the primacy of reduced accumulated precipitation (Fig.8) and of higher temperature, that lead to a non-linear H response to fertilization (Huang et al, 2024). Their influence is noticed as well in the absence of fertilization 325 (Fig.7a), when H still declines in warmer climates, with a dominant control from precipitation. The correlation along sowing dates between H and accumulated precipitation until maturity (Pmat, Fig.8), is  $r(H, Pmat) > 0.96$  in both scenarios.





Fig.7: Simulated Harvest (kg/ha) under Hist (black), RCP4.5 (blue) and RCP8.5 (red) scenarios, for four sowing dates across three fertilization levels (Table 1, exper “3N”): 0 (a), 60 (b), and 120 (c) kg N/ha (from left to right), experiment setup E\_3N\_G0.



330

Fig.8: Precipitation accumulated until maturity (mm) in experiment E\_3N\_G0 (legend as in Fig. 7).

The role of the precipitation timing is emphasised: for late sowing, RCP8.5 shows more accumulated Pmat (and H) despite a shorter accumulation season (Fig.6) but having projected a precipitation increase towards late spring (Fig.5), that may  
335 significantly favour critical growth stages.

### 3.3 Optimal Genotype Identification

The system was further developed to extend the management scenarios for multi-genotype simulations and implement methods to identify ideotypes under each agro-climate scenario. The aim is to search for management scenarios that yield optimal outcomes defined by user-criteria such as maximizing harvest yield, stabilizing yield, or minimizing pollutant  
340 emissions. Two optimization methods are implemented: a discrete-parameter, purely deterministic technique, and a hybrid approach that combines deterministic modelling with continuous-parameter Machine Learning-based Genetic Algorithms for iterative genotype selection.

The deterministic method involves conducting multiple-genotype crop model simulations, with optimization performed as a post-processing step. Genotype parameters  $P_i$  are defined within pre-established limits and discretization. Multi-model  
345 simulations are then performed, where each parameter is individually varied while the remaining parameters are held constant. The total number of simulations in this case is determined by the chosen discretization level. In contrast, in the hybrid technique the  $P_i$  values are selected from a continuous range of values, identifying and iteratively improving the best sub-domains. This section presents the results of genotype optimization experiments (E\_1N\_Gn+w), built upon the E\_1N\_G0 and sets up initial (1st of January, yearly) soil moisture as best agreement with projections targeting near-term  
350 (2035 as centre of interval 2021-2050).

#### 3.3.1 Optimal genotype under climate change

##### i) harvest as a function of the genotype H(G) in scenarios versus current climate

We analyse the distribution of H obtained along multi-genotype simulations, ordered from maximum to minimum values and denote the genotypes corresponding to this ordering “H-ordered genotypes”, chain which is simulation (model, scenario)  
355 dependent. Comparing these H distributions for the two climate scenarios against Hist, indicates projected changes in the ensemble-model PDF (probability density function) of H under warmer climate.

A first outcome demonstrates in Fig.9a, b that for the H-ordered genotypes, a projected average decrease in Harvest (H) occurs within the range of maximum H values (genotypes in the upper H-percentile, interval GX (0%, 2.5%) of the H-ordered genotypes), under both scenarios, and mostly affecting the earlier sowing dates (Fig.9b). Across models of the  
360 ensemble, we note a strong modulation of this behaviour by precipitation, particularly for unfertilized scenarios. Precipitation exhibits high inter-model variability and significant regional-scale uncertainty, pointing to the need of ensemble modelling for reducing it. In contrast, the warming trend is a consistent feature across models in the region,

contributing other model-systematic responses such as earlier anthesis and maturity dates and shortening of the grain filling season.

365 The second note regards a different response projected in the intermediate H values (Fig.9a, c). Genotypes corresponding to the intermediate H values (genotypes of middle H-percentile, the interval GI (25%, 70%) of the H-ordered genotypes) show projected higher H values in GI in climate scenarios than in Hist (Fig.9c), affecting less the earlier sowing (Fig.9c).

These together lead to a narrowing of the H-values range of responses, in GX and GI, to the same managements applied, in scenarios compared to Hist. Same management spread would lead to closer H-responses, with enhancing the expectancy for

370 occurrence of intermediate values and decreasing the expectancy for highest H values (a third feature of projected changes).


Finally, we note that despite this narrowing, earlier sowings appear systematically as better timing options (Fig.9a), improving by up to 2-(4) % in scenarios (respectively Hist) unfertilized case and up to 8-(12) % in fertilised case (Fig.9a), with the lowest percentage for RCP8.5. Earlier sowing was reported in other recent studies as optimal for spring maize harvest (Djaman et al, 2022).

375 **ii) options for adaptation and mitigation using genotype analysis**



These three features of cross genotype-agro-management impact: - projected lower maxims of H in scenarios (mainly for early sowing), projected higher intermediate H (mainly mid-late sowing); - a narrowing of the range of H in GX and GI with higher/ lower expectancy of intermediate/ high values occurrence, have practical adaptation outcomes.

380 The first two points are equivalent to slopes' change of H as a function of the ordered genotype, as shown in (Supl. S3) in climate scenario versus Hist. Slope change information indicates the percentile (and genotype) threshold for improving the result in scenario compared to Hist, for a given agro management. Alternatively, for a given genotype one could find how a change in management practice could optimize the result. In this last case for example, one could choose a small shift in the



385 sowing, but using less fertilisation, less pollutant, meanwhile getting a same or even higher H, as shown for example in TR5 versus TR11 in Fig.9a, RCP4.5 (Fig.9).



b)



c)



390 **Fig.9 a): Harvest multi-model time mean, ordered from maximum to minimum value (left to right on x-axis, logarithmic scale).** The simulations are for: Hist (left), RCP4.5 (middle) and RCP8.5 (right), experiment setup E\_1N\_Gn+w, for cross-genotype changes in six  $P_i$  parameters (resulting 1890 simulations, x-axis); b) differences in projected harvest for a) RCP4.5 minus Hist (left) and RCP8.5 minus Hist (right), for the upper H percentile (the first 50 values, [1-50] on x axis) and intermediate in c), range [400-1400] on x-axis. (“Hmax left” indicates that increasing values of H are on leftward direction of the axis).

395 Apart from any comparison with Hist, it is important for long term adaptation, that one may find genetic combinations with high yield in specific target percentile under a given climate (e.g. first 50 values, as in Fig.9b).

At yearly level, the interest for some of these genotype parameters combinations may increase, providing that distinct weather favourable patterns will be identified, once with progress achieved in seasonal and annual weather forecasting (Dewitte et al., 2021).

#### 400 3.3.2 Optimal Genotype parameters $P_i$ under climate change

##### i) optimal genotype parameters

We further discriminate H response per genotype parameters ( $P_1$ - $P_6$ ), to understand the source of the changes in Fig.9 and the possible adaptation paths under climate and management scenarios.

Parameters’ analysis (Fig. 10) shows that in all simulations, higher harvest is obtained under: shorter thermal time from seedling to juvenile phase ( $P_1$ , Fig. 10 a), shorter photoperiod-delay ( $P_2$ , Fig.10b), slightly shorter thermal time between successive leaves appearance (phyllotchron,  $P_6$ , Fig. 10 e) in GI and longer in GX, but longer thermal time to maturity ( $P_3$ , Fig.10c) and higher grain filling rate ( $P_5$ , Fig. 10d). These results are in coherence with findings along recent works. Shorter  $P_1$  or lowering the seedling-juvenile thermal time for increasing H (Fig. 10a) is in agreement with Mi et al., (2021) for semi-humid areas, (the current class of this region, with semi-arid trends projected, Fig. 4), and the same for  $P_2$ , while slower 410 maturity ( $P_3$ ) and enhanced filling rate ( $P_5$ ) being linked to higher kernel weight and harvest in agreement with recent studies (Grewer et al., 2024).

##### ii) changes in optimal genotype parameters in climate scenarios

Comparing  $P_i$  in climate scenarios against Hist, reveals the new plant strategy put in place in the new climatic conditions, for maximizing the harvest. The ensemble simulations (Fig. 10) shows that highest harvests are reached with genotypes that

415 ensure a longer thermal time from seedling to juvenile phase and longer thermal time to maturity in scenarios compared to Hist. To a smaller extent this is also achieved by a longer photoperiod delay, higher grain filling rate and longer phyllochron interval, in scenarios, than for a same percentile of the Harvest in Hist. These show that under warmer climate it is essentially important to avoid too fast growth on main stages of the development.

Indeed, slower development phases are obtained in scenarios mainly by increasing P1 and P3 (Fig.10a, b) and related to 420 these, under longer photoperiod (P2 increases, Fig.10b). Other contributions come from ensuring a slower rate of appearance of successive leaves (P5 increase), while a higher grain filling rate (P6 increase) appears to partly compensate for the negative effect of higher temperature that decreases the seed-filling duration and seeds number and size and finally the harvest.

In other studies, this compensation was shown to be minor compared to the loss of seed-filling duration in warmer climate 425 (Singh et al., 2013) that points to P1 and P3 as main drivers for Harvest in climate scenarios. Percentages of the Pi changes in scenarios versus Hist for a given percentile of harvest (suppl. S4) confirm this main driving.




Fig.10: Pi values corresponding to ordered, decreasing harvest (on y axis, the number of the ordered simulation, y=1 is the highest harvest simulation). X-axis shows the Pi interval-number of discretization, increasing with increased values. Discretization here have used 5x7x6x1x3x3 intervals for P1xP2xP3xP4xP5xP6 (total 1890). Simulations are shown for two treatments (TR1 at top and TR12 at bottom), for: Hist (black), RCP4.5 (blue, only shown for the running mean) and RCP8.5 (red), ensemble time-mean; full lines show running means over 100 values window. The short arrows in a) and c) indicate, for a same harvest percentile (y=constant) the corresponding Pi intervals for Hist (black) and RCP8.5 (red); long arrows indicate the P0i values of the intersection of running-mean Pi for Hist with RCP8.5.

435

### iii) optimal genotype parameters in management and climate scenarios

Agro-treatments choice may significantly modulate the H response to genotype parameters. Delaying sowing, requires gradually decreasing Pi in order to maximize H (Fig.11, also in Fig.10), for both Hist and climate scenarios. For P1-P3 this

decrease reflects the priority in avoiding a too late end of the juvenile stage (and shift in climate conditions) and a too late

440 (autumn) maturity stage that is slowing the grain filling and leading crop failure.

However, Fig. 11 also shows that these Pi decreases cease or even reverse under extreme delay of sowing. For highest delays the development stage is getting too short under P1's too strong decrease while daily temperatures becoming higher, hampering the development. The same is seen for the maturity, with P3' too strong decrease favouring a too quick grain filling. Hence the plant strategy for adaptation after a threshold of sowing-delay is similar to the one already seen in its

445 adaptation to warmer climate, in scenarios. Higher harvest is then reached by gradually switching to only moderate decrease or even increases of Pi parameters along with gradual increasing delays in the sowing date.

This gradual switch in the mechanism of Pi performing high harvest, with sowing delay appears quite systematic for all Pi.

This crop adaptation mechanism, converging to the one projected for climate scenarios, shows that gradually under enhanced warming, the crucial priority in adaptation transfers, from the key issue of ensuring climatological conditions for the

450 development to the key issue of avoiding a too fast growth leading crop failure.

This gradual switch in the mechanism of Pi performing high harvest, with sowing delay appears quite systematic for all Pi.

This crop adaptation mechanism, converging to the one projected for climate scenarios, shows that gradually under enhanced warming, the crucial priority in adaptation transfers, from the key issue of ensuring climatological conditions for the

development to the key issue of avoiding a too fast growth leading crop failure.

455




Fig.11 As in Fig.10 but for all sowing dates, no fertilization Fx0 (TR1-4, top) and with fertilization Fx2 (TR9-12, bottom). Parameters Pi are shown for the top 200 highest harvest (y from 1 to 400). Grey colours are for Hist and yellow-red for RCP8.5 (light to dark from earlier to latest sowing).

460

#### iv) optimal genotype parameters in adaptation and mitigation strategy

For each agro-management and climate scenario one can identify threshold values of Pi ( $P_{0i}$  that depend on the Pi, the sowing date and the fertilization level, shown in Fig.10) of intersection between scenario and Hist. At this value, for the

genotype the two have the same H percentile. So P0i shows if we get an enhanced percentile or decreased from genotypes  
465 with higher or lower Pi in the scenario compared to Hist (Fig.10, shown by arrows).

Second remark is on the expectancy of an outcome. Since all the slopes of Pi, each as a function of H ordered-values are lower than in Hist (suppl. S3), there is a narrower Pi interval for all those Pi decreasing with H (e.g. P1) and a border one for those Pi increasing with H (P3, Fig.10c), in climate scenarios. P3 increases are broadening the interval for H-highest percentile, potentially presenting, in this sense, more expectancy (than P1, Fig.10a) on highest values outcome.

470 The genotyping results were found both in simulations involving deterministic and the hybrid deterministic-ML methods. The hybrid method involved the same cross-simulations, but the selection of Pi values for H optimization and ordering was no more following a pre-defined discretisation but instead a random picking up over a continuous interval of values with successively retrieving the best generation. It applies for optimization, classic Genetic Algorithms methods in which selection of pairs is based on the user-criteria (e.g. maximum harvest, stable harvest, etc.). Our results show that for the same  
475 physical intervals of the genotype parameters, the ML hybrid technique only after 20 generations shows at least 50% chances to get a better result than the deterministic model, while after 100 generations, it already increases at 80% chances to get better results with also computational efficiency. CPU time is reduced in this case by more than 30% using the hybrid technique compared to the fully deterministic model on a VM Linux platform. Hybrid method emerges as a better solution since it can identify improved optimums at lower computational prices.

#### 480 4. Discussions

The results found are in line with other results in recent studies, using different approaches and observational data, and offer an extended (continuum-parameter) assessment towards a more generalised frame, allowed by the implemented system. For the plant response under management treatment delaying sowing date, limiting elongations of the development phase was also found in other studies (Huang et al., 2020) to reduce the impact of temperature increase and, in some cases, precipitation  
485 decrease and water stress. This response was also found stronger under enhanced fertilization and delayed sowing (Fig. 10, 11). Also fertilization lowering P6 and enhancing leaf appearance rate (Fig. 10f), assessed in earlier studies mainly for warmer climates (Hokmalipour et al., 2011; Sardans et al., 2017; dos Santos et al., 2021) was recently put in relation to P2 decrease mainly along sensitive photoperiods (Hu et al., 2023) and to higher harvest, through enhanced evapo-transpiration maximizing the high N uptake (Lu et al., 2024). In warmer climate scenarios (Fig.10f, 11f), limitations in the expansion of  
490 new leaves (increase of P6, Fig.10) was shown to be an adaptive tolerance mechanism to drought and heat stress conditions (Fahad et al., 2017).

Further, for moderate sowing delay fertilisation was shown to require slower grain filling (P5, Fig.11d) under reduced P1, P2 and P3, controlling N stimulated growth under hydric stress conditions of current and projected climate for non-irrigated crop (Yang et al., 2024). Under high delay and warmer climate, a higher grain filling is required (Fig.11d). This increase for  
495 P5 under increased warming may reflect an adaptive strategy of plants to accelerate development under drought stress,

allowing plants to end their life cycle before impact of severe drought stress occurs (McKay et al., 2003; Roeber et al., 2022).

Simulations here emphasize and compare adaptation paths of gradual plant response to warming climate. These emphasise some reduction in the efficiency of adaptation through crop management in warmer climates. Meanwhile, genotyping shows 500 the possibility of identifying parameters still able to enhance efficiency of adaptation under climate and agro-management scenarios, hence suitable methods for an accelerating change. The ability of exploring continuum-parameter space not only offers a general picture of adaptation cross-solutions but identifies critical values of the parameters that for small perturbations may lead the system response into different states (threshold sowing-delays, or P0i for genotypes). Without an integrated modelling approach, estimating or emphasising these points meaningful for adaptation is hard, moreover since 505 these are simulation (climate-management scenario) dependent.

## 5. Conclusions

The main outcome of this study is that an agroclimatic real-time Interactive Service was implemented towards adaptation support, that allows performing real-time, user-requested, agro-management modelling scenarios for the region, under current and future climate. A novel feature of the system is the ability for identifying optimal management paths for the 510 user's request, along with multiple cross-cultivar parameters, such as cross-optimal sowing date, genotype parameters, amount and date of fertilization.

The system provides solutions and estimates the associated uncertainty by using multi-model ensembles for each agro-climate and management scenario. The crop optimization criteria are user-defined and can relate to high harvest, stable harvest, low pollution. The optimization module implemented uses a hybrid deterministic - ML methodology. It performs 515 multi-model simulations using physical models of climate and plant phenology and optimization is done either through discretizing the parameters' space and optimisation post-processing or using hybrid physical-ML Genetic Algorithms methods. ML methods are spanning continuous parameter's space and iteratively selecting along the simulations the best fit parameters, allowing to identify unprecedented optimal configurations (H maxims), not reachable under the discrete deterministic method. The overall system output information is layered and accessed from two interfaces: one static, for 520 information purpose (phenology, harvest, climate, extremes at high resolution NUTS3 level) and a second is real-time interactive online, through which the user places requests and receives the system-performed management simulations required (including uncertainty along multi-models) and identified optimal paths for adaptation. These platforms are operational for two emission scenarios RCP4.5 and RCP8.5 and twelve management scenarios (sowing dates and fertilization), for the time-horizon up to 2050, with open-source code (EERIS platform). The results of these were discussed 525 in this work for the pilot region South Romania.

For the current genotype, in both emission scenarios it is projected a mean decrease (14% in ensemble mean, with higher values per model) of the projected harvest, for all the management scenarios (sowing-dates and fertilization) tested.

This was linked to a projected shortening of the grain filling season (10% quicker with an earlier shift of both anthesis (5 day) and maturity (10 day) phases) and to a mean decrease of the fertilisation efficiency under climatic scenarios, stronger in 530 RCP 8.5 emissions.

The impact of genotype perturbations on crop parameters is analysed along six cross-genotype parameters, for agro-management-climate scenarios. The main questions: i) Can we identify optimal genotype parameters that lead to maximal harvest? How do these differ under projected climate change and/ or under agro-management options and can these enhance our understanding to guide our options? iii) Can be genotyping a (better) solution for adaptation under climate change in the 535 region?

These simulations showed that the maximal H values are projected to decline for all agro-management and breeding simulations performed, in emission scenarios compared to Hist, with a higher decline for earlier sowing. H-values then increase in the intermediate-percentile harvest in scenarios versus Hist and there is enhanced expectancy in scenarios to reach the historical values in this range through agro-management and breeding. These indicate a narrowing of the responses 540 range to same agro-managements, with less / more expectancy of reaching values in the highest / intermediate H-range of Hist, in climate scenarios. In practice, these express that we can identify the H-percentile (genotype), where agro-management choice will optimize the outcome compared to Hist, including finding solutions with lower fertilisation, less pollutant.

For effective support in adaptation applications, individual genotype parameters  $P_i$  were analysed in climate scenarios versus 545 Hist. This showed that the thermal times to juvenile ( $P_1$ ) and maturity ( $P_3$ ) are key genotype parameters driving harvest changes in the region, requiring increased values in climate scenarios compared to Hist for a same highest harvest-percentile range. This range is identified through critical values of the parameters ( $P_{0i}$ ), determined for each treatment and climate scenario. There is significant variability of  $P_{0i}$  under agro-management treatments. Moderate delayed sowing and enhanced 550 fertilisation may diminish the shifts in  $P_i$  in scenarios compared to Hist for a same H-percentile, in contrast to extreme managements.

These results show that Genetic approaches offer adaptation strategy support in helping plants to resist drought stress under warming climate. Moreover, it was shown that the optimization is improved by using a hybrid ML genetic algorithm method coupled to the deterministic model-output, leading to detecting better solutions, under a continuous-parameter space search. The system can be further used for searching paths along extreme drought years, along with 555 irrigation options investigation. Coupled with weather extended predictions (seasonal, year -decadal) this could provide near real-time adaptation support.

**Code and data availability:** The code is available in the Github repository at: <https://github.com/pneague/Genetic-Algorithm-for-Corn-Genotype-sowing-Date-Optimization> under a BSD 2-Clause Simplified License

560

**Author contribution:** MC: model implementation, code for optimal adaptation tool, pre and post-processing, model simulations, results analysis, development of the User-Platform, paper writing; LC: DSSAT model set-up, results analysis, paper review; PN: ML method implementation and runs, results analysis, paper writing; AD: model validation; VA: development of the Info-Platform; ZC and AI: platforms upload and update; AP: agro-meteorological station data providing; 565 GC: DSSAT model input for the target region.

**Competing interests:** The contact author has declared that none of the authors has any competing interests

**Acknowledgments:** The authors are grateful to UEFISCDI who provided the financial support of this work under the 570 Project Grant PREPCLIM PN-III-P2-2.1-PED-2019-5302.

## References

1. Adams, R. M., Hurd, B. H., Lenhart, S., & Leary, N. (1998): Effects of global climate change on agriculture: an interpretative review. *\*Climate Research\**, 11(1), 19–30.
- 575 2. Angus, J.F., Mackenzie, D.H., Morton, R., & Schafer, C.A. (1981): Phasic development in field crops II. Thermal and photoperiodic responses of spring wheat, *\*Field Crops Research\**, 4, 269-283, ISSN 0378-4290, [https://doi.org/10.1016/0378-4290\(81\)90078-2](https://doi.org/10.1016/0378-4290(81)90078-2).
3. Arnell, N.W., & Freeman, A. (2021): The effect of climate change on agro-climatic indicators in the UK. *\*Climatic Change\**, 165, 40, <https://doi.org/10.1007/s10584-021-03054-8>.
- 580 4. Asseng, S., Ewert, F., Martre, P., Rötter, R. P., Lobell, D., Cammarano, D., Kimball, B., et al. (2015): Rising temperatures reduce global wheat production. *\*Nature Climate Change\**, 5.
5. Baez-Gonzalez, A. D., Kiniry, J., Maas, S., et al. (2005): Large-Area Maize harvest Forecasting Using Leaf Area Index Based harvest Model. *\*Agronomy Journal\**, 97. 10.2134/agronj2005.0418.
6. Bai, Y., Yue, W., & Ding, C. (2021): Optimize the Irrigation and Fertilizer Schedules by Combining DSSAT and GA.
- 585 7. Baltas, E. (2007) Spatial Distribution of Climatic Indices in Northern Greece. *\*Meteorological Applications\**, 14, 69-78. [www.interscience.wiley.com <http://dx.doi.org/10.1002/met.7>](http://dx.doi.org/10.1002/met.7)
8. Banterng, P., A. Patanothai, K. Pannangpatch, S. Jogloy, and G. Hoogenboom. 2004. Determination of genetic coefficients of peanut lines for breeding applications, *European Journal of Agronomy* 21(3):297-310.
9. Basso, B., Shuai, G., Zhang, J., et al. (2019): harvest stability analysis reveals sources of large-scale nitrogen loss from the 590 US Midwest. *\*Sci Rep\**, 9, 5774, <https://doi.org/10.1038/s41598-019-42271-1>.
10. Benestad, R., Buonomo, E., Gutiérrez, J.M., et al. (2021): Guidance for EURO-CORDEX climate projections data use, Version 1.1.

11. Bernardo, R. (2002): \*Breeding for Quantitative Traits in Plants\*. Stemma Press, 9780972072403, <https://books.google.ro/books?id=3T2FQgAACAAJ>.

595 12. Berti, A., Maucieri, C., Bonamano, A., & Borin, M. (2019): Short-term climate change effects on maize phenological phases in northeast Italy. \*Italian Journal of Agronomy\*, DOI: 10.4081/ija.2019.1362.

13. Birch, C.J., Vos, J., Kiniry, J., Bos, H.J., & Elings, A. (1998): Phyllochron responds to acclimation to temperature and irradiance in maize, \*Field Crops Research\*, 59(3), 187-200, ISSN 0378-4290, [https://doi.org/10.1016/S0378-4290\(98\)00120-8](https://doi.org/10.1016/S0378-4290(98)00120-8).

600 14. Boogaard, H., Wolf, J., Supit, I., Niemeyer, S., & van Ittersum, M. (2013): A regional implementation of WOFOST for calculating harvest gaps of autumn-sown wheat across the European Union, \*Field Crops Research\*, 143, 130-142, ISSN 0378-4290, <https://doi.org/10.1016/j.fcr.2012.11.005>.

15. Caian, M., Georgescu, F., Pietrisi, M., & Catrina, O. (2021): Recent Changes in Storm Track over the Southeast Europe: A Mechanism for Changes in Extreme Cyclone Variability. \*Atmosphere\*, 12(10):1362.

605 <https://doi.org/10.3390/atmos12101362>

16. Caian, M.; Andrei, M.D. Late-Spring Severe Blizzard Events over Eastern Romania: A Conceptual Model of Development. \*Atmosphere\* 2019, \*10\*, 770. <https://doi.org/10.3390/atmos10120770>

17. Ceglar, A., Zampieri, M., Gonzalez-Reviriego, N., et al. (n.d.): Time-varying impact of climate on maize and wheat harvests in France since 1900. \*Environmental Research Letters\*, 15(9), <https://doi.org/10.1088/1748-9326/aba1be>.

610 18. Chang, Y., Latham, J., Licht, M. et al. A data-driven crop model for maize yield prediction. \*Commun Biol\* 6, 439 (2023). <https://doi.org/10.1038/s42003-023-04833-y>

19. Chapagain, R., Remenyi, T. A., Huth, N., Mohammed, C. L., & Ojeda, J. J. (2023): Investigating the effects of APSIM model configuration on model outputs across different environments. \*Frontiers in Agronomy\*, <https://www.frontiersin.org/articles/10.3389/fagro.2023.1213074>

615 20. Chazarreta, Y. D., Amas, J. I., & Otegui, M. E. (2021): Kernel filling and desiccation in temperate maize: Breeding and environmental effects, \*Field Crops Research\*, 271, 108243, ISSN 0378-4290, <https://doi.org/10.1016/j.fcr.2021.108243>.

21. Chen, Y., & Tao, F. (2022): Potential of remote sensing data-crop model assimilation and seasonal weather forecasts for early-season crop yield forecasting over a large area, \*Field Crops Research\*, 276, 108398, ISSN 0378-4290, <https://doi.org/10.1016/j.fcr.2021.108398>.

620 22. Cooper, M., & Messina, C. D. (2023): Breeding crops for drought-affected environments and improved climate resilience, \*The Plant Cell\*, 35(1), 162–186, <https://doi.org/10.1093/plcell/koac321>.

23. Dainelli, R., Calmanti, S., Pasqui, M., et al. (2022): Machine learning for regional crop harvest forecasting in Europe, \*Field Crops Research\*, 276, 108377, ISSN 0378-4290, <https://doi.org/10.1016/j.fcr.2021.108377>.

24. Djaman K., Samuel Allen, Dorlote S. Djaman, Komlan Koudahe, Suat Irmak, Naveen Puppala, Murali K. Darapuneni, 625 Sangamesh V. Angadi, Planting date and plant density effects on maize growth, yield and water use efficiency, \*Environmental Challenges\*, 6, 2022, 100417, ISSN 2667-0100, <https://doi.org/10.1016/j.envc.2021.100417>

25. dos Santos, C. L., et al. (2022): Maize Leaf Appearance Rates: A Synthesis From the United States Corn Belt. *Frontiers in Plant Science*, 13, <https://doi.org/10.3389/fpls.2022.872738>.

26. Espadafor, M., Orgaz, F., Testi, L., et al. (2017): Responses of transpiration and transpiration efficiency of almond trees to moderate water deficits, *Scientia Horticulturae*, 225, 6-14, ISSN 0304-4238, <https://doi.org/10.1016/j.scienta.2017.06.028>.

630 27. Eyring, V., Mishra, V., Griffith, G.P., et al. (2021): Reflections and projections on a decade of climate science. *Nature Climate Change*, 11(4), 279-285.

28. Fahad, et al. (2017): Crop Production under Drought and Heat Stress: Plant Responses and Management Options. *Frontiers in Plant Science*, 8, <https://doi.org/10.3389/fpls.2017.01147>.

635 29. Flocas, A. A. *Courses of meteorology and climatology*. Ziti Publications: Thessaloniki (1994)

30. Ganguly, 2013: Post harvest losses of agricultural produce. *International Journal of Agricultural Sciences*, 9(2), June, 2013 , 818-820

31. Godfray, H., Charles, J., et al. (2010): Food Security: The Challenge of Feeding 9 Billion People. *Science*, 327, 812-818.

640 32. Grewer, U., Kim, D.-H., & Waha, K. (2024): Too much, too soon? Early-maturing maize varieties as drought escape strategy in Malawi, *Food Policy*, 129, 102766, ISSN 0306-9192, <https://doi.org/10.1016/j.foodpol.2024.102766>.

33. Hatfield, J.L., et al. (2020): Indicators of climate change in agricultural systems. *Climatic Change*, 163, 1719–1732.

34. Hoogenboom, G., Porter, C.H., Sheila, V., Boote, K. J., Singh, U., White, J. W., Hunt, L.A., Ogoshi, R., Lizaso, J.I., Koo, J., Asseng, S., Singels, A., Moreno, L.P. and Jones, J. W. 2019. Decision Support System for Agrotechnology Transfer 645 (DSSAT) Version 4.7. DSSAT Foundation

35. Hokmalipour, S. (2011): The Study of Phyllochron and Leaf Appearance Rate in Three Cultivar of Maize (Zea mays L.) At Nitrogen Fertilizer Levels. *Agricultural and Food Sciences*.

36. Huang, M., et al. (2020): *Environmental Research Letters*, 15, 024015; DOI 10.1088/1748-9326/ab66ca

37. Huang N, Xiaomao Lin, Fei Lun, Ruiyun Zeng, Gretchen F. Sassenrath, Zhihua Pan, Nitrogen fertilizer use and climate 650 interactions: Implications for maize yields in Kansas, *Agricultural Systems*, Volume 220, 2024, 104079, ISSN 0308-521X, <https://doi.org/10.1016/j.agsy.2024.104079>

38. Hu, H., et al. (2023): The effects of photoperiod and temperature-related factors on maize leaf number and leaf positional distribution in the field. *Frontiers in Plant Science*, 14, <https://doi.org/10.3389/fpls.2023.1006245>.

39. IPCC, 2022 *Climate Change 2022: Impacts, Adaptation and Vulnerability*. IPCC Sixth Assessment Report

655 40. Jin, X., Jin, Y., Zhai, J., et al. (2022): Identification and prediction of crop Waterlogging Risk Areas under the impact of climate change. *Water*, 14, 1-21, 10.3390/w14121956.

41. Jones, J., Hoogenboom, G., Porter, C., et al. (2003): The DSSAT cropping system model. *European Journal of Agronomy*, 18, 235-265, [https://doi.org/10.1016/S1161-0301\(02\)00107-7](https://doi.org/10.1016/S1161-0301(02)00107-7).

42. Kakar, K.M., Khan, A., Khan, I., et al. (2014): Growth and yield response of maize (Zea mays L.) to foliar NPK- 660 fertilizers under moisture stress condition. *Soil and Environment*, 33(2),

43. Khan, F. , Khan, S. , Fahad, S. , Faisal, S. , Hussain, S. , Ali, S. and Ali, A. (2014) Effect of Different Levels of Nitrogen and Phosphorus on the Phenology and Yield of Maize Varieties. *American Journal of Plant Sciences*, 5, 2582-2590. doi: 10.4236/ajps.2014.517272.

44. Karl E. Taylor, Ronald J. Stouffer, Gerald A. Meehl, A summary of the CMIP5 Experiment Design, 2011

665 45. Kothari, K., Ale, S., Marek, G.W., Munster, C.L., Singh, V.P., Chen, Y., Marek, T.H., & Xue, Q. (2022): Simulating the climate change impacts and evaluating potential adaptation strategies for irrigated corn production in Northern High Plains of Texas, *Climate Risk Management*, 37, 100446, ISSN 2212-0963, <https://doi.org/10.1016/j.crm.2022.100446>

46. Langworthy, A. D., et al. (2018): Potential of summer-active temperate (C 3) perennial forages to mitigate the detrimental effects of supraoptimal temperatures on summer home-grown feed production in south-eastern Australian 670 dairying regions. *Crop Pasture Sci.*, 69, 808–820.

47. Lelieveld, J., Hadjinicolaou, P., Kostopoulou, E., et al. (2012): Climate change and impacts in the Eastern Mediterranean and the Middle East. *Climatic Change*, 114, 667–687, <https://doi.org/10.1007/s10584-012-0418-4>.

48. Li, M., & Tang, Y. (2022): Climate warming causes changes in wheat phenological development that benefit yield in the Sichuan Basin of China, *European Journal of Agronomy*, 139, 126574, ISSN 1161-0301, 675 <https://doi.org/10.1016/j.eja.2022.126574>

49. Liu, K., Harrison, M.T., Yan, H., et al. (2023): Silver lining to a climate crisis in multiple prospects for alleviating crop waterlogging under future climates. *Nature Communications*, 14(1), 765, 10.1038/s41467-023-36129-4.

50. Liu, K., et al. (2020): Identifying optimal sowing and flowering periods for barley in Australia: a modelling approach. *Agric. For. Meteorol.*, 282-283, 107871.

680 51. Lu, J., Stomph, T.J., Mi, G., et al. (2024): Identifying and quantifying the contribution of maize plant traits to nitrogen uptake and use through plant modelling, *in silico Plants*, 6(2), diae018, <https://doi.org/10.1093/insilicoplants/diae018>.

52. Malhi, G.S.; Kaur, M.; Kaushik, P. Impact of Climate Change on Agriculture and Its Mitigation Strategies: A Review. *Sustainability* 2021, 13, 1318. <https://doi.org/10.3390/su13031318>

53. Mamassi, A., Balaghi, R., Devkota, K.P., et al. (2023): Modelling genotype × environment × management interactions 685 for a sustainable intensification under rainfed wheat cropping system in Morocco. *Agriculture and Food Security*.

54. Marcinkowski, P., & Piniewski, M. (2018): Effect of climate change on sowing and harvest dates of spring barley and maize in Poland. *International Agrophysics*, 32(2), 265-27

55. Meehl, G.A., Stocker, T.F., Collins, W.D., et al. (2007): Global Climate Projections. In: Solomon, S., Qin, D., Manning, M., et al. (Eds.), Cambridge University Press.

690 56. Meroni M, François Waldner, Lorenzo Seguini, Hervé Kerdiles, Felix Rembold, Yield forecasting with machine learning and small data: What gains for grains?, *Agricultural and Forest Meteorology*, Volumes 308–309, 2021, 108555, ISSN 0168-1923, <https://doi.org/10.1016/j.agrformet.2021.108555>

57. Mi, N.; Cai, F.; Zhang, S.; et al. (2021): Thermal Time Requirements for Maize Growth in Northeast China and Their Effects on Yield and Water Supply under Climate Change Conditions. *Water*, 13, 2612. <https://doi.org/10.3390/w13192612>

695 58. Mitchell, R.J., Bellamy, P.E., Broome, A., et al.: Cumulative impact assessments of multiple host species loss from plant diseases show disproportionate reductions in associated biodiversity. *Journal of Ecology*, <https://doi.org/10.1111/1365-2745.13798>.

59. McKay, J. K., Richards, J. H., & Mitchell, T. (2003). Genetics of drought adaptation in *Arabidopsis thaliana*: I. Pleiotropy contributes to genetic correlations among ecological traits. *Molecular ecology*, 2003

700 60. McKee, T.B.; Doesken, N.J.; Kleist, J. (1993): The Relationship of Drought Frequency and Duration to Time Scales. In Proceedings of the Eighth Conference on Applied Climatology, Anaheim, CA, USA, 17–22 January 1993; pp. 179–184.

61. Morales, A., & Villalobos, F. (2023): Using machine learning for crop harvest prediction in the past or the future. *Frontiers in Plant Science*, 14, 1128388, 10.3389/fpls.2023.1128388.

62. Morell, F-J., Yang, H.S., Cassman, K.G., et al. (2016): Can crop simulation models be used to predict local to regional maize harvests and total production in the U.S. Corn Belt? *Field Crops Research*, 192, 1-12, ISSN 0378-4290, <https://doi.org/10.1016/j.fcr.2016.04.004>.

705 63. Patra Abhik, Vinod Kumar Sharma, Dhruba Jyoti Nath, Asik Dutta, Tapan Jyoti Purakayastha, Sarvendra Kumar, Mandira Barman, Kapil Atmaram Chobhe, Chaitanya Prasad Nath & Chiranjeev Kumawat (2022): Long-term impact of integrated nutrient management on sustainable harvest index of rice and soil quality under acidic inceptisol, \*Archives of 710 Agronomy and Soil Science\*, DOI: 10.1080/03650340.2022.2056597.

64. Paudel, Allard de Wit, Hendrik Boogaard, Diego Marcos, Sjoukje Osinga, Ioannis N. Athanasiadis, Interpretability of deep learning models for crop yield forecasting, *Computers and Electronics in Agriculture*, Volume 206, 2023, 107663, ISSN 0168-1699, <https://doi.org/10.1016/j.compag.2023.107663>

65. Peleman, J-D., & Rouppe van der Voort, J. (2003): Breeding by Design, *Trends in Plant Science*, 8(7), 330-334, ISSN 715 1360-1385, [https://doi.org/10.1016/S1360-1385\(03\)00134-1](https://doi.org/10.1016/S1360-1385(03)00134-1).

66. Pfeiffer, W., & McClafferty, B. (2007): HarvestPlus: Breeding Crops for Better Nutrition. *Crop Science*, 47. 10.2135/cropsci2007.09.0020IPBS.

67. Qiao, L., Xiaojun, Z., Li, X., et al. (2022): Genetic incorporation of genes for the optimal plant architecture in common wheat. *Molecular Breeding*, 42. 10.1007/s11032-022-01336-2.

720 68. Rezaei, E.E., Webber, H., Asseng, S., et al. (2023): Climate change impacts on crop harvests. *Nature Reviews Earth and Environment*.

69. Roeber, V., Schmulling T, Cortleven A., (2022): The Photoperiod: Handling and Causing Stress in Plants *Frontiers in Plant Science*, 12, [<https://doi.org/10.3389/fpls.2021.781988>](<https://doi.org/10.3389/fpls.2021.781>

725 70. Rosenzweig, C., Jones, J.W., Hatfield, J.L., et al. (2013): The Agricultural Model Intercomparison and Improvement Project (AgMIP): Protocols and pilot studies. *Agricultural and Forest Meteorology*, 170, 166-182, <https://doi.org/10.1016/j.agrformet.2012.09.011>.

71. Sardans, J., Grau, O., Chen, H.Y.H., et al.: Changes in nutrient concentrations of leaves and roots in response to global change factors. *Global Change Biology*, <https://doi.org/10.1111/gcb.13721>.

71. Schauberger, B., Jägermeyr, J., & Gornott, C. (2020): A systematic review of local to regional harvest forecasting  
730 approaches and frequently used data resources. European Journal of Agronomy, 120, 126153,  
<https://doi.org/10.1016/j.eja.2020.126153>.

72. Schwalbert , R., Amado , T.,Nieto , L. et al. (2020). Mid-season county-level corn yield forecast for US Corn Belt  
integrating satellite imagery and weather variables. *Crop Sci.* 60 : 739 – 750

73. Selvaraju, R., et al. (2011): Climate science in support of sustainable agriculture and food security. *Climatic Research*,  
735 47, 95–110.

74. Semenov, M., & Stratonovitch, P. (2015): Adapting wheat ideotypes for climate change: Accounting for uncertainties in  
CMIP5 climate projections. *Climate Research*, 65. 10.3354/cr01297.

75. Shaw, T.A., & Miyawaki, O. (2024): Fast upper-level jet stream winds get faster under climate change. *Nature Climate  
Change*, 14, 61–67. <https://doi.org/10.1038/s41558-023-01884-1>

740 76. Simmons, A. (2021): The ERA-Interim archive Version 2.0, ERA Report Series.

77. Singh R.P, P. V. Vara Prasad, K. Raja Reddy, Chapter Two - Impacts of Changing Climate and Climate Variability on  
Seed Production and Seed Industry,Editor(s): Donald L. Sparks,Advances in Agronomy,Academic Press,Volume  
118,2013,Pages 49-110,ISSN 0065-2113,ISBN 9780124059429,<https://doi.org/10.1016/B978-0-12-405942-9.00002-5>

78. Smil, V. 2005. The next 50 years: Unfolding trends. *Population and Development Review* 31: 605-643.

745 79. Stehr, N., & von Storch, H. (2000): Climate and society: climate as resource, climate as risk. *World Scientific*.

80. Su, Z., Liu, Z., Bai, F., et al. (2021): Cultivar selection can increase yield potential and resource use efficiency of spring  
maize to adapt to climate change in Northeast China. *Journal of Integrative Agriculture*, 20(2): 371–382.

81. Tao, F., Zhao, Z., Liu, J., & Yokozawa, M Modelling the impacts of weather and climate variability on crop productivity  
over a large area: A new super-ensemble-based probabilistic projection.

750 82. Tsvetinskaya, E.A., Mearns, L.O., & Easterling, W.E. (2002): Investigating the Effect of Seasonal Plant Growth and  
Development in Three-Dimensional Atmospheric Simulations. Part I: Simulation of Surface Fluxes over the Growing  
Season. [https://doi.org/10.1175/1520-0442\(2001\)014](https://doi.org/10.1175/1520-0442(2001)014)

83. Tsimba R, Gregory O. Edmeades, James P. Millner, & Peter D. Kemp (2013): The effect of planting date on maize:  
Phenology, thermal time durations and growth rates in a cool temperate climate, *Field Crops Research*, 150, 145-155, ISSN  
755 0378-4290, <https://doi.org/10.1016/j.fcr.2013.05.021>.

84. Trnka, M., et al. (2014): Adverse weather conditions for European wheat production will become more frequent with  
climate change. *Nature Climate Change*, 4, 637–643.

85. Villalobos, F., Pérez-Priego, O., Testi, L., et al. (2012): Effects of water supply on carbon and water exchange of olive  
trees. *European Journal of Agronomy*, 40, 1-7. 10.1016/j.eja.2012.02.004.

760 86. van Ittersum M., Kenneth G. Cassman, Patricio Grassini, Joost Wolf, Pablo Tittonell, Zvi Hochman,Yield gap analysis  
with local to global relevance—A review, *Field Crops Research*, Volume 143,2013,Pages 4-17,ISSN 0378-  
4290,<https://doi.org/10.1016/j.fcr.2012.09.009>

87. Yang Y. , Jiabo Yin, Shengyu Kang, Louise J. Slater, Xihui Gu, Aliaksandr Volchak, Quantifying the drivers of terrestrial drought and water stress impacts on carbon uptake in China, Agricultural and Forest Meteorology, Volume 344, 2024, 109817, ISSN 0168-1923, <https://doi.org/10.1016/j.agrformet.2023.109817>

765 88. Wang, Y., Jiang, K., Shen, H., et al. (2023): Decision-making method for maize irrigation in supplementary irrigation areas based on the DSSAT model and a genetic algorithm. Agricultural Water Management, 280, 108231.

89. Webber, H., Ewert, F., Olesen, J.E., et al. (2018a): Diverging importance of drought stress for maize and winter wheat in Europe. Nature Communications, 9(1), 1-11. 10.1038/s41467-018-06525-2.

770 90. Webber, H., et al. (2020): Pan-European multi-crop model ensemble simulations of wheat and grain maize under climate change scenarios, Open Data Journal for Agricultural Research, 6, 21-27.

91. Wheeler, T., & von Braun, J. (2013): Climate change impacts on global food security. Science, 341(6145), 508-13. 10.1126/science.1239402.

92. World Development Report 2008: Agriculture for Development.

775 93. Wu J-Z, Jing ZHANG, Zhang-ming GE, et al. (2021): Impact of climate change on maize harvest in China from 1979 to 2016, Journal of Integrative Agriculture, 20(1), 289-299, ISSN 2095-3119, [https://doi.org/10.1016/S2095-3119\(20\)63244-0](https://doi.org/10.1016/S2095-3119(20)63244-0).

94. Xie, W., Zhu, A., Ali, T., et al. (2023): Crop switching can enhance environmental sustainability and farmer incomes in China. Nature, 616, 300–305. <https://doi.org/10.1038/s>.

95. Xu Honggen, Bo Ming, Keru Wang, et al. (2023): Quantitative analysis of maize leaf collar appearance rates, Plant Physiology and Biochemistry, 196, 454-462, ISSN 0981-9428, <https://doi.org/10.1016/j.plaphy.2023.01.016>.

780 96. Zhuang H, Zhao Zhang, Fei Cheng, Jichong Han, Yuchuan Luo, Liangliang Zhang, Juan Cao, Jing Zhang, Bangke He, Jialu Xu, Fulu Tao, Integrating data assimilation, crop model, and machine learning for winter wheat yield forecasting in the North China Plain, Agricultural and Forest Meteorology, Volume 347, 2024, 109909, ISSN 0168-1923, <https://doi.org/10.1016/j.agrformet.2024.109909>

785

## Annex: Data and Methods: Steps in ML algorithm

Schema of steps in workflow of ML algorithms for optimal genotype identification:

Start with 10 randomly chosen solutions within the bounds of P1-P6;

Calculate the mean and std of harvest of each solution for the 30 years 1976-2005;

790 Calculate fitness = (Mean of harvest) – (Standard-deviation of Harvest/4);

Randomly choose 4 pairs of ‘parents’, with the probability being chosen weighted by the fitness;

For each pair of parents A and B, create identical children ‘a’ and ‘b’ to the parents, then choose a random number of P’s to be subjected to crossover, called x;

For each child, modify Px as follows:

795  $P_{xa} = \text{round} (B * P_{xa} + (1 - B) * P_{xb})$

$P_{xb} = \text{round} (1 - B) * P_{xa} + B * P_{xb}$

Where  $P_{xa}$  is the value of the x parameter of child “a” (initially identical to that of parent A), and B is the blending factor, set in this paper to 0.75. This technique is called blending, and it generates offspring chromosomes that inherit real-valued traits from both parents while exploring the search space between the parents' positions. The blending crossover promotes a

800 smoother and more gradual search for optimal solutions in continuous domains.

Then take each child, and with a probability of 0.5 perform a mutation on one of its chromosomes. This means setting one of the P’s to a random value between its allowed minimum and maximum.

At this point the children have been fully constructed. Discard the 8 parents with the lowest fitness and substitute them with the children.

805 Repeat.