



## The sensitivity of aerosol data assimilation to vertical profiles: case study of dust storm assimilation with LOTOS-EUROS v2.2

Mijie Pang<sup>1</sup>, Jianbing Jin<sup>1\*</sup>, Ting Yang<sup>2</sup>, Xi Chen<sup>1</sup>, Arjo Segers<sup>3</sup>, Hai Xiang Lin<sup>4,5</sup>, Hong Liao<sup>1</sup>, and Wei Han<sup>6\*</sup>

<sup>1</sup>Jiangsu Key Laboratory of Atmospheric Environment Monitoring and Pollution Control, Jiangsu Collaborative Innovation Center of Atmospheric Environment and Equipment Technology, School of Environmental Science and Engineering, Nanjing University of Information Science and Technology, Nanjing, Jiangsu, China

<sup>2</sup>State Key Laboratory of Atmospheric Boundary Layer Physics and Atmospheric Chemistry (LAPC), Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, China

<sup>3</sup>TNO, Department of Climate, Air and Sustainability, The Netherlands

<sup>4</sup>Delft Institute of Applied Mathematics, Delft University of Technology, Delft, The Netherlands

<sup>5</sup>Institute of Environment Sciences, Leiden University, The Netherlands

<sup>6</sup>CMA Earth System Modeling and Prediction Centre, Chinese Meteorological Administration, Beijing, China

**Correspondence:** Jianbing Jin (jianbing.jin@nuist.edu.cn) and Wei Han (hanwei@cma.gov.cn)

**Abstract.** Modeling and observational techniques are pivotal in aerosol research, yet each approach exhibits inherent limitations. Aerosol observation is constrained by its limited spatial and temporal coverage compared to models. On the other hand, models tend to possess higher uncertainties and biases compared to observations. Aerosol data assimilation has gained popularity as it combines the advantages of both methods. Despite numerous studies in this domain, few have addressed the challenges

5 faced in assimilating aerosol data with significant differences in magnitude and degree of freedom between the model state and observations, especially in the vertical direction. These challenges can lead to the preservation or even exacerbation of structural inaccuracies within the assimilation process. This study investigates the sensitivity of dust aerosol data assimilation to the vertical structure of the aerosol profile. We assimilate a variety of dust observations, encompassing ground-based particulate matter (PM<sub>10</sub>) measurements and satellite-derived dust optical depth (DOD) data, using the Ensemble Kalman Filter

10 (EnKF). The assimilation process is elucidated, detailing the assimilation of raw ground-based and satellite-based observations for an optimized three-dimensional (3D) posterior state. To demonstrate the impact of accurate versus erroneous prior aerosol vertical profiles on the assimilation result, we select three cases of super dust storms for analysis. Our findings reveal that the assimilation of ground observations would optimize the dust field at the ground in general. However, the vertical structure presents a more complex challenge. When the prior profile accurately reflects the true vertical structure, the assimilation process

15 can successfully preserve this structure. Conversely, if the prior profile introduces an incorrect structure, the assimilation can significantly deteriorate the integrity of the aerosol profile. This is also found in the assimilation of DOD, which exhibits a comparable pattern in its sensitivity to the initial aerosol profile's accuracy.



## 1 Introduction

In recent decades, atmospheric aerosol has garnered extensive attention due to its impact on the environment (Buseck and Pósfai, 1999; Zhao et al., 2020) and human health (Zhu et al., 2021). Numerous research studies have been conducted on the characterization (Mhawish et al., 2021), source apportionment (Wu et al., 2018), and model simulation (Lee et al., 2011) of aerosols. One crucial tool for understanding aerosol processes and forecasting is the chemical transport model (CTM), which can generate continuous three-dimensional aerosol fields on a large scale. However, despite significant efforts to develop and improve CTM parameter schemes, studies have pointed out that significant uncertainties still exist in CTM simulations (Vignati et al., 2010; Stier et al., 2013). These uncertainties primarily arise from emission sources (e.g., intensity, location), meteorological inputs (e.g., wind mixing, precipitation), and incomplete parameter schemes (e.g., chemical reactions) (Benedetti et al., 2018). At the same time, aerosol measurements have advanced owing to development of sensors and increased investment. These include chemical kinetics studies in laboratories (Kolb and Worsnop, 2012), a network of loosely-distributed ground monitoring stations (Gueymard and Yang, 2020), and regularly scanning satellites providing global coverage (Sogacheva et al., 2020). Although these measurements offer higher accuracy, challenges persist. So far, obtaining continuous three-dimensional aerosol measurements on a large scale remains impractical and costly. Measurements are limited to surface level concentrations of total particle matters, column-integrated aerosol optical depths (AOD), and vertical profiles of total aerosol extinction. These measurements have fewer degrees of freedom compared to the complexity of reality (Pappalardo et al., 2014; Qin et al., 2021).

Data assimilation incorporates model and measurements to generate a posterior that holds the strengths of both (Bannister, 2017). Various data assimilation techniques have been developed for different applications, including hydrology (Reichle et al., 2002), geology (Peng and Liu, 2022), and atmospheric science (Hamill, 2006). These methods are typically based on either variational or ensemble approaches (Whitaker et al., 2009; Law and Stuart, 2012). The variational method is particularly useful for emission inversion (Bergamaschi et al., 2022; Jin et al., 2023a). Ensemble-based filters also have demonstrated superior performance in sequential forecasting without the adjoint or tangent linear model that required by the variational method. By assimilating real-time observations, these filters generate optimized initial fields that enhance forecast accuracy (Houtekamer et al., 2005). Since the introduction of the classic Ensemble Kalman Filter (EnKF) (Evensen, 1994), a considerable amount of research has focused on its application and improvement. Several variations of the EnKF, such as EnSRF (Whitaker and Hamill, 2002) and LETKF (Hunt et al., 2007), have been proposed to address computational complexity, accuracy, and robustness issues. Although these advancements have been validated, challenges related to the model's background error covariance and observational data quality control can still impact the assimilation results significantly.

In data assimilation methodology, an observation operator is required for deriving the innovation vector that quantifies the difference between observation field and simulation field (Ma et al., 2020). These two usually have different degrees of freedom. With that, the optimized state then could be calculated considering the integration of background and observational penalties. In perspective of aerosol data assimilation, the main object is to reproduce the optimal aerosol states (Liu et al., 2011), concerning their spatial, vertical, aerosol species and size distribution features. While the available observations commonly measure their mixed state, such as AOD that is column-integrated optical extinction of all aerosols, and ground  $PM_{10}$



concentration measurement that is additive sum of all particles with diameter less than  $10\ \mu\text{m}$ . They are not directly comparable to the aerosol state. For ground measurements, the monitoring stations are sparsely distributed. High data vacancy is inevitable in AOD product as a result of retrieval failure. The order of the observations is therefore commonly much lower than the model state or the degree of freedom is limited compared to that of model state. Mathematically, the assimilation result highly depends on the background error covariance for transferring the increments from the low-degree observational space into the full model space. Given a partially erroneous prior, the assumptions about the prior states can be reserved or even exacerbated with the lack of observations. Only a few studies have focused on some of these aspects. We recently found that the fraud assumption of aerosol size distribution from prior model gives rise to worsen posterior (Jin et al., 2023b). In addition, the position error in the dust plume simulation can cause failure in assimilating valid observations (Jin et al., 2021). As to the vertical structure, most studies have focused on proposing relevant assimilation methods to assimilate observations like Cloud-Aerosol Lidar with Orthogonal Polarization (CALIPSO) (Sekiyama et al., 2010; Cheng et al., 2019; Ye et al., 2021), Light Detection and Ranging (LiDAR) (Wang et al., 2022). The sensitivities of assumptions in the prior simulations need to be explored.

Take dust aerosol data assimilation for instance, the most representative observation sources are direct observation like  $\text{PM}_{10}$ ,  $\text{PM}_{2.5}$  concentrations and indirect observation like dust optical depth (DOD), extinction coefficient (Jin et al., 2022). Since they all are a combination of different particles, it is necessary to perform the bias-correction before assimilation (Jin et al., 2019; Ma et al., 2020). Afterwards, through assimilation analysis, few observations can nudge the whole model state space. While in reality, the ground stations produce scattered concentrations on ground, satellite and LiDAR receive column-integrated information about the aerosol or a single profile once a day. None of them can provide a continuous vertical information about the aerosol from a comprehensive perspective in space. When there is fraud information about the dust aerosol structure in model prior, due to the considerably greater degree of freedom of model state, data assimilation algorithms may fail to correct it or even worse, further degrades the vertical dust aerosol loading.

There is limited research addressing the impact of flawed aerosol prior assumptions on the degradation of the assimilated posterior, particularly concerning vertical structures. Gwyther et al. (2023) delved into the implications of 4DVar data assimilation on mesoscale eddy representation in the vertical dimension, highlighting that inadequate prior knowledge of these structures significantly hinders improvement in the assimilated simulations. This issue extends to aerosol data assimilation, where erroneous assumptions regarding aerosol vertical distribution lead assimilation methods to maintain or exacerbate inaccuracies. This study focuses on several major dust storm events that occurred in the spring of 2021, utilizing them as case studies. We conduct separate data assimilation experiments involving ground-level  $\text{PM}_{10}$  measurements and satellite-based AOD data to demonstrate explicitly how misleading vertical prior information can distort the posterior results. The LOTOS-EUROS model facilitates the reproduction of dust dynamics and generates the initial dust load priors. For assimilation, we employ an EnKF algorithm, incorporating bias-corrected  $\text{PM}_{10}$  data from ground stations and DOD retrievals from the Himawari-8 satellite. To validate, we employ aerosol profile observations from CALIPSO and LiDAR systems. By presenting both favorable and detrimental scenarios, this work underscores the high sensitivity of data assimilation processes to the accuracy of assumed vertical structures, thereby emphasizing the criticality of precise vertical profiling in aerosol studies.



The remainder is organized as follows: Section 2 described the model used for dust aerosol simulation and the dust observations used in this study. Section 3 introduces the assimilation algorithm applied to assimilate the observations and the experiment settings. Section 4 discusses the assimilation results and validate the results with the CALIPSO and LiDAR observations. Finally, Section 5 summaries this study.

## 90 2 Dust aerosol simulation and observations

This section introduces the model used for simulating the dust aerosol. Then describe the observation methods applied for monitoring the dust including ground monitor station, Himawari-8, CALIPSO and LiDAR. Among these, ground PM<sub>10</sub> from ground monitor stations and DOD from Himawari-8 are used for assimilation. Extinction profile from CALIPSO and LiDAR are used to validate the posterior profile.

### 95 2.1 Dust simulation

#### 2.1.1 LOTOS-EUROS

The LOTOS-EUROS v2.2 is used to simulate the dust aerosol. The LOTOS-EUROS model is a 3D chemistry transport model aimed for air quality forecasting (Manders et al., 2017). It has also been applied in source apportionment and emission inversion worldwide. In this study, The modeling domain spans from 15° N to 50° N and from 70° E to 140° E with spatial resolution  
100 of 0.25° × 0.25°. Vertically, it comprises 21 layers with a top level at 10 km, which is adequate for recognizing the vertical structure. Dust aerosol processes including emission, advection, diffusion, deposition and sedimentation are considered in the model.

### 2.2 Dust observations for assimilation

#### 2.2.1 Ground PM<sub>10</sub>

105 There are over 1600 ground air quality monitor stations standing all over China. Their spatial distribution can be seen in Fig. 1 (blue scatters). They provide hourly air pollutant concentrations including PM<sub>2.5</sub>, PM<sub>10</sub>, O<sub>3</sub>, NO<sub>2</sub>, SO<sub>2</sub> and CO. PM<sub>10</sub> can appropriately represent the dust load for it is under the size definition of fine and coarse dust (Adebiyi et al., 2023). Its uncertainty is also relatively lower than some remote sensing instruments. However, PM<sub>10</sub> is a combination of various species. Assimilating such observations with weak or fraud representativeness can diverge the model. It is then necessary to perform  
110 bias-correction (BC) for removing the non-dust part before they are assimilated. Details concerning the method can be found in Jin et al. (2022). Afterwards, it can be used to correct the model simulation by data assimilation.

#### 2.2.2 Himawari-8 DOD

The Himawari-8 geostationary satellite, operated by the Japan Meteorology Agency (JMA), provides aerosol optical information covering East Asia and Western Pacific Region at a time-resolution of 10-min (Yumimoto et al., 2016). This allows for



115 monitoring the source and movement of dust at high spatial-temporal resolution. Its products, Ångström exponent (Å) and  
Aerosol Optical Thickness (AOT) at 550 nm, are selected to measure the dust optical depth (DOD). These data are merged  
to model grid ( $0.25^\circ \times 0.25^\circ$ ) and hourly resolution. A bias correction produce is applied before assimilation. Details can be  
found in Supplementary S1.

### 2.3 Dust observations for validation

120 Aerosol extinction coefficient from the CALIPSO and LiDAR is used to validate the vertical structure of dust aerosol.

#### 2.3.1 CALIPSO

The Cloud-Aerosol LiDAR with Orthogonal Polarization (CALIOP), onboard the CALIPSO satellite, provides vertical struc-  
tures and types of aerosols on a global scale (Winker et al., 2010). It has a spatial resolution of 333 m and a vertical resolution  
of 30 m. A normal extinction profile sample can be seen above the black dash line on map in Fig. 1 (a). It has been wildly used  
125 in investigating the aerosol distribution and evolution (Xu et al., 2020; Zhang et al., 2020; Han et al., 2022; Chen et al., 2023),  
data assimilation (Sekiya et al., 2010; Cheng et al., 2019; Escribano et al., 2022). Here, its product, extinction coefficient at  
532 nm, is taken as the aerosol profile validation set to identify the vertical distribution of the dust load.

#### 2.3.2 Polarization LiDAR


Polarization LiDAR deployed on the ground can provide continuous measurements about the vertical details of the atmo-  
spheric environment with high resolution (Hofer et al., 2017). Recently, Wang et al. (2021) collected the dust extinction coef-  
130 ficient data through a dual-wavelength aerosol LiDAR built in the tower of the Institute of Atmospheric Physics ( $39^\circ 58' 35''\text{N}$ ,  
 $116^\circ 22' 41''\text{E}$ , red star in Fig. 1 (a)). The device provided the dust information with a high vertical resolution of 30 m and  
temporal resolution of 15 min. The vertical profile development of the several super dust storms occur in spring, 2021 was  
captured thoroughly by the instrument, thus make it a crucial measurement source to learn about the vertical structure and  
135 its development during the dust storm. In this study, these data are employed to compare the vertical structure from model  
simulation and assimilation analysis.

## 3 Assimilation methodology and experiments

This Section first illustrates the methodology of the classic EnKF that were used in Section 3.1, then sensitivity of the aerosol  
posterior to the vertical structure were emphasised when assimilating the ground  $\text{PM}_{10}$  concentration in Section 3.2 and when  
140 assimilating the AOD in Section 3.3. Cases experiments for evaluating the sensitiveness are described in Section 3.4.

### 3.1 EnKF

The data assimilation algorithms used in this study is the stochastic EnKF formulated by Burgers et al. (1998). It is implemented  
in our self-developed PyFilter toolbox (Pang, last access: May. 2024). Its capability of improving dust forecasting had been



**Figure 1.** Spatial distribution of ground monitor stations (blue scatters), location of the LiDAR (red star) with upward scanning ray (orange cylinder) and aerosol extinction profile sample scanned by the CALIPSO satellite (trajectory is plotted as the gray line on ground).


proven in our recent research (Pang et al., 2023). We didn't adopt any other variant of EnKF for the reason that the problem we  
 145 are going to explain is contingent but universal in aerosol data assimilation applications.

EnKF is a Monte Carlo approach based on Kalman filter theory. EnKF maintains a set of model states to approximate the probability distribution of the model state or parameter. It includes the forecast step and the analysis step. In the forecast step, each ensemble member  $\mathbf{x}_{t-1}^{a,i}$  is integrated forward according to the model dynamics  $\mathcal{M}$  to generate a forecast for the system state  $\mathbf{x}_t^{f,i}$  at the next moment.

$$150 \quad \mathbf{x}_t^{f,i} = \mathcal{M}(\mathbf{x}_{t-1}^{a,i}) \quad (1)$$

In the analysis step, the states of each member is adjusted based on observational data to approach the true state. This adjustment process is implemented based on the error from model prior and observation. In EnKF, the model prior error covariance matrices  $\mathbf{P}^f$  is calculated by ensemble approximation method following:

$$\mathbf{P}^f = \frac{1}{N-1} (\mathbf{x}_i^f - \bar{\mathbf{x}}^f)(\mathbf{x}_i^f - \bar{\mathbf{x}}^f)^T \quad (2)$$



**Figure 2.** Schematic diagram of the sensitivity of data assimilation to aerosol vertical structure. Assimilation of ground and AOD observation under positive condition with AOD priors (gray hollow diamond), AOD observation (blue solid diamond) in upper box and true profile (red line), ensemble priors (gray dash line), prior mean (black line), ground observation (blue star) in lower box (a.1). Figure on the right are for the posteriors (a.2). Figures in the second row are prior (b.1) and posterior (b.2) of the ground assimilation under negative condition. Figures in the third row are prior (c.1) and posterior (c.2) of the AOD assimilation under negative condition.



155  $N$  is the number of ensembles.  $\mathbf{x}_i^f$  is the  $i$ th individual of the ensemble prior states and  $\bar{\mathbf{x}}^f$  means the average of ensemble members.

Then, a weight matrix  $\mathbf{K}$ , also referred to as Kalman Gain, can be obtained by

$$\mathbf{K} = \mathbf{P}^f \mathcal{H}^T (\mathcal{H} \mathbf{P}^f \mathcal{H}^T + \mathbf{R})^{-1} \quad (3)$$

The a posteriori ensemble individual  $\mathbf{x}_i^a$  is calculated by :

$$160 \quad \mathbf{x}_i^a = \mathbf{x}_i^f + \mathbf{K}(\mathbf{y} + \epsilon_i - \mathcal{H}\mathbf{x}_i^f) \quad (4)$$

$\epsilon_i$  is the extra perturbation and it's variance subjecting to the diagonal of observation error covariance  $\mathbf{R}$ . It serves to maintain the ensemble spread van Leeuwen (2020).

Meanwhile, the a posteriori  $\mathbf{x}^a$  can be updated via

$$\mathbf{x}^a = \bar{\mathbf{x}}^f + \mathbf{K}(\mathbf{y} - \mathcal{H}\bar{\mathbf{x}}^f) \quad (5)$$

165 From the equations above we can tell that the a posteriori is dependent on both prior and observations. These within different space are aligned by the observation operator  $\mathcal{H}$ .

Figure 2 briefly illustrates the process of EnKF applied in this paper. In the upper row of panel (a.1), we have ensemble AOD priors (gray hollow diamond) and AOD observation (blue solid diamond), and in below there are idealized aerosol vertical profiles including a true profile (red line), ensemble prior profiles (gray dash line) and an observation on ground level (blue star). It is noticeable that there are great differences between priors and observation both on ground and AOD in two conditions. Figures in panel (a) are in positive condition. Which both the AOD and 3-D mass priors has underestimated the loading while the structure is consistent with the true profile. Through assimilating ground or AOD observations, both the ground and vertical dust loading are nudged to be better consistent with the observations.

### 3.2 Ground $\text{PM}_{10}$ assimilation

175 In this section, impact of ground measurements data assimilation on the aerosol vertical structure is explained. In practice, each of the  $\text{PM}_{10}$  observations could be assimilated to nudge the 3D states that are correlated. The increments is transferred from the the observation space in the model space via the Kalman Gain  $\mathbf{K}$  shown above. To easily evaluate the posterior, we solely perform the assimilation analysis at the pixels where  $\text{PM}_{10}$  measurements are available. So here we confine the states only in vertical direction in one pixel to better identify the sensitivity, and assuming the prior states  $\mathbf{x}^f$  with total  $k$  levels:

$$180 \quad \mathbf{x}^f = [x_1^f, \dots, x_k^f]^T \quad (6)$$





In which case, the error covariance matrix  $\mathbf{P}^f$  is constructed by the variance of ensemble states and covariance in the vertical direction, and the observation operator  $\mathcal{H}$  only converts the single observation on ground level:

$$\mathbf{P}^f = \begin{bmatrix} C_{11} & C_{12} & \cdots & C_{1k} \\ C_{21} & C_{22} & \cdots & C_{2k} \\ \vdots & \vdots & \ddots & \vdots \\ C_{k1} & C_{k2} & \cdots & C_{kk} \end{bmatrix} \quad (7)$$

$$\mathcal{H} = [1 \quad 0 \quad \cdots \quad 0] \quad (8)$$

185 Calculation of  $\mathbf{K}$  follows the Eq. 3 and substitute Eq. 7 and Eq. 8 into it:

$$\mathbf{K} = \mathbf{P}^f \mathcal{H}^T (\mathcal{H} \mathbf{P}^f \mathcal{H}^T + \mathbf{R})^{-1} = [C_{11}, \dots, C_{k1}]^T (C_{11} + \sigma_y^2)^{-1} \quad (9)$$

$\sigma_y^2$  is the error of the ground observation.

Substitute  $\mathbf{K}$  into the update function:

$$[x_1^a, \dots, x_k^a]^T = [x_1^f, \dots, x_k^f]^T + [C_{11}, \dots, C_{k1}]^T (C_{11} + \sigma_y^2)^{-1} (y - x_1^f) \quad (10)$$

190 Here, we have the posterior state  $x_i^a$  on each level.

On the ground level, the posterior state is obtained as:

$$x_1^a = \frac{\sigma_y^2}{C_{11} + \sigma_y^2} x_1^f + \frac{C_{11}}{C_{11} + \sigma_y^2} y \quad (11)$$

If the observation error is much smaller than the prior error ( $\sigma_y^2 \ll C_{11}$ ). We can have:

$$x_0^a \approx y \quad (12)$$

195 which means that the posterior on ground level is close to the ground observation. The ground data assimilation has successfully tuned the ground prior state.

On the other hand, for the state on  $i$ th level, the posterior  $x_i^a$  is calculated by:

$$x_i^a = x_i^f + \frac{C_{k1}}{C_{11} + \sigma_y^2} (y - x_1^f) \quad (13)$$

To better identify the correctness of vertical structure, define a vertical ratio  $v_k$  here:

$$200 \quad v_k = \frac{x_i}{x_1} \quad (14)$$

which is the ratio of state on  $i$ th level to ground level.

Divide  $x_i^f$  on both sides of Eq. 13:

$$v_i^a = v_i^f + \frac{C_{k1}}{C_{11} + \sigma_y^2} \frac{y - x_1^f}{x_1^f} \quad (15)$$



When we have a strong vertical correlation and small observation error, we have  $\frac{C_{k1}}{C_{11} + \sigma_y^2} \approx 1$ . If the prior vertical structure  
 205 is far from the true structure  $v_i^f \ll v_i^{true}$  or  $v_i^f \gg v_i^{true}$ . Then, the vertical structure of posterior is high dependent on  $\frac{y - x_1^f}{x_1^f}$ . If  
 $x_1^f$  and  $y$  deviate far from each other, the increment can be large, thus further enhance the incorrect prior structure  $v_i^f$ .

Figure 2 (b) illustrates this kind of pattern. In panel (b.1), the priors have greatly underestimated the aerosol concentration  
 on the ground level. Meanwhile, over the upper space, it has overestimated the intensity and altitude of the dust loading to  
 some extent. After assimilating the ground observations, as shown in panel (b.2), the posterior aerosol loading (gray dash line)  
 210 is tuned to a large extent and much closer to the observation on ground layer. While for the vertical distribution, a greater  
 overestimation than the prior can be noticed. The original incorrect profile is significantly amplified.

### 3.3 AOD assimilation

Assimilation of AOD is carried out by connecting the 3-D aerosol priors  $x^f$  and AOD priors  $\tau^f$ . It is assumed that the vertical  
 structure of aerosol field remains the same for the AOD lacks the vertical information. Then, a 3-D mass concentration field  
 215 can be calculated by finding the optimal AOD field.

To begin with, value of AOD is related to the aerosol mass concentration, aerosol type, humidity, et al. Here, it's assumed to  
 be linearly related to the aerosol mass concentration field  $x^f$  solely. The column-integrated AOD is calculated by the summary  
 of aerosol mass concentration  $x_i^f$  on each level:

$$\tau^f = \sum_{i=1}^k \mathcal{M} x_i^f \quad (16)$$

220  $\mathcal{M}$  is the linear model operator that calculates the AOD from the mass concentration.

Calculation of posterior AOD  $\tau^a$  follows the Eq. 5:

$$\tau^a = \bar{\tau}^f + \mathbf{K}(y - \mathcal{H}\bar{\tau}^f) \quad (17)$$

Then, substitute Eq. 16 into Eq. 17. We can bridge the mass concentration field and AOD field:

$$\mathcal{M} \sum_{i=1}^k x_i^a = \mathcal{M} \sum_{i=1}^k x_i^f + \mathbf{K}(y - \mathcal{H}\bar{\tau}^f) \quad (18)$$

225 Same for AOD, let's confine the states in vertical direction in one pixel. We have one prior AOD  $\tau^f$  and aerosol mass prior  
 $x_i^f$  on total  $k$  levels. Define a vertical factor  $u_i$  which is the ratio of mass of  $i$ th level to the mass of all levels first:

$$u_i = \frac{x_i}{\sum_{i=1}^k x_i} \quad (19)$$

Substitute it into Eq. 18:

$$\mathcal{M} \frac{x_i^a}{u_i} = \mathcal{M} \frac{x_i^f}{u_i} + \mathbf{K}(y - \mathcal{H}\bar{\tau}^f) \quad (20)$$



230 Since we only have one observation and one AOD simulation, then the observation operator  $\mathcal{H}$ , error variance matrix  $\mathbf{P}^f$  and Kalman gain  $\mathbf{K}$  are:

$$\mathcal{H} = [1], \quad \mathbf{P}^f = [\sigma_\tau^2], \quad \mathbf{K} = \left[ \frac{\sigma_\tau^2}{\sigma_\tau^2 + \sigma_y^2} \right] \quad (21)$$

Substitute them into Eq. 20 and we have the mass posterior  $x_i^a$  on each layer:

$$\mathcal{M} \frac{x_i^a}{u_i^f} = \mathcal{M} \frac{x_i^f}{u_i^f} + \frac{\sigma_\tau^2}{\sigma_\tau^2 + \sigma_y^2} (y - \tau^f) \quad (22)$$

235

$$x_i^a = x_i^a + u_i^f \frac{\sigma_\tau^2}{\sigma_\tau^2 + \sigma_y^2} \frac{y - \tau^f}{\mathcal{M}} \quad (23)$$

This equation demonstrates that the analysis increment  $\frac{\sigma_\tau^2}{\sigma_\tau^2 + \sigma_y^2} \frac{y - \tau^f}{\mathcal{M}}$  from AOD assimilation is allocated to each level of mass by the vertical factor  $u_i$ . When we have an incorrect structure, we can notice that  $u_i \gg u_i^{true}$  or  $u_i \ll u_i^{true}$ . The analysis increment can be reallocated on the wrong level. Thus amplify the incorrect structure.

240 Figure 2 (c) demonstrates the negative pattern of assimilating AOD. In the upper row of panel (c.1), we have the ensemble AOD priors (gray hollow diamond) and AOD observation (blue solid diamond) that are combined to optimize the 3-D dust mass field (gray dash line). Similar to panel (b), assume we have an incorrect prior dust mass structure. By assimilating AOD observations, the AOD posteriors can be tuned to better fit the AOD observation as shown in the upper row in panel(c.2). While for the structure of mass field, this incorrect structure is overall amplified since the prior DOD simulation implies the model  
 245 underestimates the total dust column compared to the observation. Another pattern which the prior mass is incorrectly situated on ground level is also illustrated in Supplementary Fig. S1. It is notable that the sensitivity of AOD assimilation to the vertical structure is not as significant as the ground assimilation since the AOD contains information of total columns.

### 3.4 Case settings

To demonstrate the sensitivity of aerosol data assimilation on vertical structure, five assimilation analysis cases were selected  
 250 and name after *P-Ground-Calipso*, *N-Ground-Calipso*, *N-Ground-LiDAR*, *N-DOD-Calipso* and *NP-DOD-Calipso*, respectively. These cases were selected from the ensemble model run starting from 0:00 14th to 0:00 17th March and 0:00 14th to 0:00 17th April 2021. The background run consists of 32 ensembles and each of them are driven by perturbed emission inventories. Details concerning the generation of the ensembles can be found in Jin et al. (2022).

Table 1 lists information about the experiment cases. *P-Ground-Calipso* is the positive case which shows how the data  
 255 assimilation correct the surface concentrations and maintain the correct aerosol structure. *N-Ground-Calipso* and *N-Ground-LiDAR* are negative cases which shows how the assimilation deteriorates the profile when assimilating ground observations solely. The time of the former two cases is fixed, and their profiles follow the trajectory of CALIPSO. Hence, by comparison we can tell the impact of assimilation on vertical structure. The last case focuses on a fixed site. A continuous time-line is used, and assimilation analysis is performed on each time point. The profiles are compared with high-resolution LiDAR observations.



260 *N-DOD-Calipso* and *NP-DOD-Calipso* are negative cases that focus on assimilating DOD observations. Himawari-8 DOD observations are assimilated. Both of them are validated by the CALIPSO.

**Table 1.** Experiment configurations

Name	Time (CST)	Type	Assimilated observations	Validation data
<i>P-Ground-Calipso</i>	3:00 16th April 2021	positive	BC-PM <sub>10</sub>	CALIPSO
<i>N-Ground-Calipso</i>	3:00 16th March 2021	negative	BC-PM <sub>10</sub>	CALIPSO
<i>N-Ground-LiDAR</i>	15th March 2021	negative	BC-PM <sub>10</sub>	LiDAR
<i>N-DOD-Calipso</i>	14:00 16th March 2021	negative	Himawari-8 DOD	CALIPSO
<i>NP-DOD-Calipso</i>	14:00 29th March 2021	negative & positive	Himawari-8 DOD	CALIPSO

## 4 Results and discussions

This section delves into the sensitivities of data assimilation from ground and DOD observation, with a particular emphasis on the impact of prior vertical profile. Regarding ground observation assimilation, the discussion is enriched by presenting a positive case, alongside two illustrative negative examples. The validity of these scenarios is further reinforced from CALIPSO satellite observations and LiDAR measurements. This serves to underscore the substantial influence that ground assimilation can exert on the posterior. Then, the examination of DOD assimilation uncovers its intricacies by showcasing a pair of instances, highlighting favorable and unfavorable outcomes. These illustrations collectively emphasize the impact DOD assimilation has on the posterior, thereby underscoring the importance of understanding the vertical profile's role in the assimilation process for both ground and DOD observations.

### 4.1 Cases on ground assimilation

#### 4.1.1 Positive case

*P-Ground-Calipso* is aimed to show how the assimilation works properly. Figure 3 (a.1) is time series of ground PM<sub>10</sub> and BC-PM<sub>10</sub> from several ground stations that is close to CALIPSO trajectory ( $< 0.25^\circ$ ). The assimilation time point, 3:00 16th April 2021, is selected in this case (highlighted in green). Although it is on the later stage of the dust storm, the concentrations were as high as around  $400 \mu\text{g m}^{-3}$ . Through comparison between raw and bias-corrected PM<sub>10</sub> (solid and dash line), it is clear that the scanned region was still dominated by coarse dust aerosol. Figure 3 (c.1 and d.1) are snapshots of average of ensemble priors and posteriors. We can see that the priors had underestimated the dust load to the south of the domain and overestimated it to the east. After assimilation analysis, the dust load had been nudged that better fits the observations. The ground observations had been consistent with the posterior. RMSE has been reduced from  $254.83 \mu\text{g m}^{-3}$  to  $118.03 \mu\text{g m}^{-3}$ .

Take a further lookup to the vertical structure. Figure 3 (b) illustrates the extinction coefficient profile from CALIPSO. From the profile we can tell that the dust was concentrated on the ground over latitude of  $34^\circ$  to  $38^\circ$  (circled by dash line).




Vertically, it extended upwards accompanied with terrain from 1 km at 34.5° to 2.5 km at 37.5°. Figure 3 (c.1) shows the prior dust profile following the same trajectory of the CALIPSO. The color bar here is rescaled to show more detail. At around 41°  
285 to 42°, a heavy dust load was simulated (600-800  $\mu\text{gm}^{-3}$ ) while CALIPSO didn't capture the data there. Hence, it can't be validated, and we will focus on the dust at 34° to 38°. At 34° to 38°, the prior model had successfully reproduced the dust structure indicated by CALIPSO. The concentrations were about 100-300  $\mu\text{gm}^{-3}$ , which is underestimated as shown in Fig. 3 (d.2). After assimilation, the ground concentrations are increased to 400-500  $\mu\text{gm}^{-3}$  as shown in (c.1) and (d.1). The vertical structure is also reserved, which is identical to the structure shown by CALIPSO.

#### 290 4.1.2 Negative case validated by CALIPSO


*N-Ground-Calipso* is the negative case that illustrates how data assimilation degrades the aerosol vertical structure. Figure 4 (a.1) is time series of ground  $\text{PM}_{10}$  and  $\text{BC-PM}_{10}$  from several ground stations that is close to CALIPSO trajectory. As shown in panel (a), ground dust aerosol loading from ground stations in *N-Ground-Calipso* was under rapid increase since two hours before assimilation instance. Meaning that the region in the below rectangle in (c.1) was dust concentrated region.

295 Evidence from CALIPSO also indicated that there were heavy dust loading. As the Fig. 4 (b) shows, extinction coefficient profile from 37° to 42° (circled by black dash rectangle) exhibits extremely high values (maximum exceeded 2  $\text{km}^{-1}$ ). It spread upwards for 1 km at an altitude of 1 km. This region is highlighted by light blue dash rectangle in (c.1) and (d.1). Inferred from the farther ground stations (concentrations over 1000  $\mu\text{gm}^{-3}$ ), this region was dominated by dust aerosols and is situated on ground level.

300 The dust profile in (c.2) tells that the ensemble priors agreed that the aerosols were mainly concentrated upwards. The plume was floating up to 4 km with the latitude. Only a few dust aerosols over 36° to 37° on ground (300-600  $\mu\text{gm}^{-3}$ ) show consistency with the observations (shown in (c.1) and (c.2)). Great discrepancy exists between the prior and independent observations at 37° to 42°. After assimilating  $\text{BC-PM}_{10}$  concentrations, the overall loading are increased. Great underestimation in east was improved. However, The dust concentrated region circled in (d.1) are only improved to a small extent. This is due to lack  
305 of observations nearby and the much lower uncertainties exhibited by ensemble priors. In terms of the vertical structure, a disturbing profile is obtained. As Fig. 4 (d.2) shows, ground concentrations at 36° to 37° are aligned to over 600  $\mu\text{gm}^{-3}$ . This is consistent with surrounding stations. While for the reason mentioned earlier, dust loads over 37° to 42° are barely corrected. Yet, these trivial increments significantly enhance the erroneous vertical structure. On ground level, the dust concentrations rise from less than 100  $\mu\text{gm}^{-3}$  to nearly 300  $\mu\text{gm}^{-3}$  at 37° to 39°. The vertical loading are almost 3 times of loading from prior  
310 profile. What's more, the extremely low region at 39° to 42° represents a more intense amplification effect than that. Since the low values are more sensitive to variations, increments on these values can easily give rise to soaring of the vertical loading. In this case, this phenomenon can be vividly noticed through the comparison between prior, posterior and measurements for validation.



**Figure 3.** Time series of  $\text{PM}_{10}$  and  $\text{BC-PM}_{10}$  concentrations during the case from several ground stations that are close to the CALIPSO trajectory (a). Spatial distribution of ground dust concentrations from the average of the ensemble priors and the posteriors with scatters of ground  $\text{BC-PM}_{10}$  observations (c.1 and posterior d.1). The black line inside is where the CALIPSO scanned through. Figures on the right column is the extinction profile from CALIPSO (b), dust concentration profile following the CALIPSO scanning trajectory from the prior (c.2) and the posterior (d.2). The color bars in c.2 and d.3 are rescaled to show more detail. Black line at the bottom is the terrain altitude. The case time is at 3:00 on 16th April 2021 (CST).



**Figure 4.** Time series of PM<sub>10</sub> and BC-PM<sub>10</sub> concentrations during the case from several ground stations that are close to the CALIPSO trajectory (a). Spatial distribution of ground dust concentrations from the average of the ensemble priors and the posteriors with scatters of ground BC-PM<sub>10</sub> observations (c.1 and posterior d.1). The black line inside is where the CALIPSO scanned through. Figures on the right column is the extinction profile from CALIPSO (b), dust concentration profile following the CALIPSO scanning trajectory from the prior (c.2) and the posterior (d.2). Black line at the bottom is the terrain altitude. The case time is at 3:00 on 16th March 2021 (CST).



### 4.1.3 Negative case validated by LiDAR

315 *N-Ground-LiDAR* is another negative case. In this case, we don't focus on single time point. A series of assimilation analyses  
are conducted to investigate the effect. As Fig. 5 (a) shows, a high-resolution profile of dust extinction coefficient from LiDAR  
is given to demonstrate the changing structure of dust loading on a fix location. Entering at about 5:00 and then landing at  
7:00, this dust storm had lasted for about 20 hours in where the LiDAR located. Time range from 8:00 to 16:00 is selected as  
it comprises most of the dust period. During the whole period, the measured dust loading increased from 0.4 km to 1.4 km.  
320 Afterwards, the intensity descended gradually.

Figure 5 (b) illustrates the hourly prior and posterior dust profile during the selected period. The x-axis which represents  
concentrations is logarithmic rescaled here. In terms of prior profiles, dust loading have extended up to 5 km at 8:00 and  
declined to 3 km at 16:00. This structure is inconsistent with the LiDAR profile. The dust storm has been overestimated to a  
large extent in height. The ground aerosol concentrations are lower than  $200 \mu\text{gm}^{-3}$  throughout eight time points, which are  
325 much lower than the observations (over  $1000 \mu\text{gm}^{-3}$ ). After assimilation, this underestimation has been mitigated. The ground  
dust concentrations are amplified several times to better fit the observations. Meanwhile, the erroneous vertical structure is also  
intensified in the first six moments. Especially in 9:00, the dust loading above 2 km have been amplified to over  $1000 \mu\text{gm}^{-3}$ ,  
which will give out utterly incorrect information about the dust storm structure and impact the further forecasting.


## 4.2 Cases on DOD assimilation

330 In this section, the sensitivity of assimilating DOD on the aerosol vertical profile is presented. Starts with the *NP-DOD-Calipso*  
case. Figure 6 (a) show the spatial distribution of DOD observed by Himawari-8 at 14:00 29th March 2021. The dust plume then  
were approaching the east of China and reaching out to the East China Sea. The prior model has reproduced the distribution  
while the overall concentration is underestimated (less than  $300 \mu\text{gm}^{-3}$ ) as shown in panel (c.1). By assimilating the Himawari-  
8 DOD, the overall dust concentration field is increased to over  $1000 \mu\text{gm}^{-3}$ . The posterior dust plume is consistent with the  
335 DOD observations.


As the extinction profile in panel (b) shows, the dust was concentrated around  $32^\circ$  to  $38^\circ$  and extended upwards to altitude  
of 3 km. In terms of the prior, this structure is partly correct. The dust in red box is located on lower layer and aligns with  
the extinction profile. While the dust in blue box reports a higher position. By assimilating DOD, the total column dust field  
is optimized and the vertical structure is restored following the prior dust field. As a result, the underestimation is alleviated  
340 and the overall concentrations are increased. The correct dust structure in red box is maintained and the intensity is enhanced.  
However, the incorrect dust structure in blue box is also amplified, causing an obvious fraud dust loading which should located  
on lower level.

*N-DOD-Calipso* is another case that illustrates the negative aspects of assimilating DDO. Figure 7 comprises the same  
assimilation experiment results as in *NP-DOD-Calipso* except at 14:00 16th March 2021. The DOD observations report high  
345 values densely distributed in east and sparsely distributed in north. The prior model simulates the dust plume that concentrated





**Figure 5.** Time series of dust extinction coefficient profile obtained from LiDAR (a). The assimilation analysis is performed hourly from 8:00 to 16:00. Figure below is hourly dust profile line from prior (dash line) and posterior (solid line) (b). The profile data is extracted from the closest grid point to the LiDAR location. The x-axis is logarithmic rescaled. Note that instead of using the posterior in previous time to propagate the model, analysis here is separately conducted on the static background.




**Figure 6.** Time series of  $PM_{10}$  and  $BC-PM_{10}$  concentrations during the case from several ground stations that are close to the CALIPSO trajectory (a). Spatial distribution of ground dust concentrations from the average of the ensemble priors and the posteriors with scatters of ground  $BC-PM_{10}$  observations (c.1 and posterior d.1). The black line inside is where the CALIPSO scanned through. Figures on the right column is the extinction profile from CALIPSO (b), dust concentration profile following the CALIPSO scanning trajectory from the prior (c.2) and the posterior (d.2). Black line at the bottom is the terrain altitude. The case time is at 14:00 on 29th March 2021 (CST).



in north-west. As to the east, only rare dust is noticed. It is caused by instability of model to reproduce the long-term transport of dust. If we focus on where the CALIPSO scanned through, an increase of dust can be seen.

In terms of CALIPSO, it is certain that the dust is concentrated on lower level at  $43^{\circ}$  to  $45^{\circ}$ . There is also dust plume at  $39^{\circ}$  to  $42^{\circ}$  while can't be verified because of the missing of observation data. Similarly, the prior dust profiles show an incorrect dust structure as circled by black box in panel (c.2). After assimilation, this error is amplified from around  $100 \mu\text{g m}^{-3}$  to over  $600 \mu\text{g m}^{-3}$ .



**Figure 7.** Snapshot of Himawari-8 DOD at the assimilation instance (a). Spatial distribution of ground dust concentrations from the average of the ensemble priors (c.1 and posterior d.1). The black line inside is where the CALIPSO scanned through. Figures on the right column is the extinction profile from CALIPSO (b), dust concentration profile following the CALIPSO scanning trajectory from the prior (c.2) and the posterior (d.2). Black line at the bottom is the terrain altitude. The case time is at 14:00 on 16th March 2021 (CST).



## 5 Conclusions

Data assimilation has been widely used in construction of reanalysis dataset and improving the predictability of atmospheric model. The performance of assimilation algorithms relies not only on the methodology themselves but also on the observations. Nowadays, the complexity and dimensions of atmospheric model have been growing fast. For atmospheric aerosol model, high spatial and temporal resolution are becoming feasible. While in terms of observations, they are either outnumbered in space or discontinuous in time compared to the model. Particularly in vertical direction, the available aerosol measurements are still limited both in space and time. Through assimilating ground or satellite measurements, the aerosol field can be improved while in 3D space, it can also amplify the erroneous estimation about the aerosol structure.

In this paper, we explore the sensitivity of aerosol data assimilation to the vertical profile via carrying out five assimilation experiment cases. Dust aerosol, which is simulated by LOTOS-EUROS, is selected as the optimization target. The data assimilation methodology, EnKF, is adopted. Bias-corrected ground  $PM_{10}$  concentrations obtained from ground monitoring stations and DOD from the Himawari-8 are assimilated. Extinction coefficient profile from CALIPSO and Polarization LiDAR are used to validate the aerosol vertical structure. *P-Ground-Calipso* is aimed to show how ground data assimilation tune the vertical structure positively. After assimilation, the ground dust field has been nudged to better fit the observations and the vertical structure is enhanced, which is also consistent with the CALIPSO measurements. In contrast, *N-Ground-Calipso* exhibits drastic negative effect. In this case, even small increments on ground can greatly enhance the incorrect aerosol structure. *N-Ground-LiDAR* is another negative case which displays hourly prior and posterior profile. The continuous dust extinction profile from LiDAR is utilized to validate the structure. In most of the moments, the assimilation improves the ground aerosol field while significantly degrades the vertical dust loading. *N-DOD-Calipso* and *NP-DOD-Calipso* are aimed to show how DOD assimilation deteriorates the vertical structure. Different from the ground assimilation, the posterior vertical structure here is reallocated by the vertical ratio. When the vertical ratio is incorrect, assimilating DOD can also pass down this error.

In conclusion, integrating ground- and satellite-derived aerosol observations into models enhances both the analysis and forecasting accuracy of the model's state. However, challenges persist in reconciling these low-dimensional observations with the model's high-dimensional state. Specifically, the model's initial, potentially flawed, vertical aerosol structure could impair assimilation efforts. Alarmingly, such inaccuracies can be propagated and exacerbated within the model. This paper has only scratched the surface by presenting a handful of negative instances, yet it is evident that this issue is pervasive in aerosol data assimilation. The path forward entails establishing a complementary network of vertical observations and implementing advanced assimilation methodologies that are sensitive to vertical structures, thereby offering a promising avenue to surmount these challenges.

*Code and data availability.* PyFilter is archived on Zenodo (<https://zenodo.org/record/7611976>) (Pang, last access: May, 2024), also open at GitHub (<https://github.com/xxcvvv/open-PyFilter>). The source code and user guide of the LOTOS-EUROS is available at <https://airqualitymodeling.tno.nl/> (TNO, last access: May 2024). The CALIPSO data can be downloaded at <https://www-calipso.larc.nasa.gov/about/>



(NASA, last access: August 2023). The ground PM<sub>10</sub> observations can be obtained at <https://quotsoft.net/air/> (Wang, last access: May, 2024).  
385 The Himawari-8 aerosol product is available at <https://www.eorc.jaxa.jp/ptree>.

*Author contributions.* JJ conceived the study. MP carried out the experiments and data analysis. JJ and MP prepared the manuscript. YT, CX, AS, HXL, HL and WH provided useful comments on the paper.

*Acknowledgements.* This work is supported by the National Natural Science Foundation of China [grant No. 42105109 and 42205031] and Natural Science Foundation of Jiangsu Province (NO. BK20210664).

### 390 **Competing interests**

The authors declare that they have no conflict of interest.



## References

- Adebisi, A., Kok, J. F., Murray, B. J., Ryder, C. L., Stuu, J.-B. W., Kahn, R. A., Knippertz, P., Formenti, P., Mahowald, N. M., Pérez García-Pando, C., Klose, M., Ansmann, A., Samset, B. H., Ito, A., Balkanski, Y., Di Biagio, C., Romanias, M. N., Huang, Y., and Meng, J.: A  
395 Review of Coarse Mineral Dust in the Earth System, *Aeolian Res.*, 60, 100 849, <https://doi.org/10.1016/j.aeolia.2022.100849>, 2023.
- Bannister, R. N.: A Review of Operational Methods of Variational and Ensemble-Variational Data Assimilation, *Q. J. R. Meteorolog. Soc.*,  
143, 607–633, <https://doi.org/10.1002/qj.2982>, 2017.
- Benedetti, A., Reid, J. S., Knippertz, P., Marsham, J. H., Di Giuseppe, F., Rémy, S., Basart, S., Boucher, O., Brooks, I. M., Menut, L., Mona,  
L., Laj, P., Pappalardo, G., Wiedensohler, A., Baklanov, A., Brooks, M., Colarco, P. R., Cuevas, E., da Silva, A., Escribano, J., Flemming,  
400 J., Huneus, N., Jorba, O., Kazadzis, S., Kinne, S., Popp, T., Quinn, P. K., Sekiyama, T. T., Tanaka, T., and Terradellas, E.: Status and  
Future of Numerical Atmospheric Aerosol Prediction with a Focus on Data Requirements, *Atmos. Chem. Phys.*, 18, 10 615–10 643,  
<https://doi.org/10.5194/acp-18-10615-2018>, 2018.
- Bergamaschi, P., Segers, A., Brunner, D., Haussaire, J.-M., Henne, S., Ramonet, M., Arnold, T., Biermann, T., Chen, H., Conil, S., Delmotte,  
M., Forster, G., Frumau, A., Kubistin, D., Lan, X., Leuenberger, M., Lindauer, M., Lopez, M., Manca, G., Müller-Williams, J., O'Doherty,  
405 S., Scheeren, B., Steinbacher, M., Trisolino, P., Vítková, G., and Yver Kwok, C.: High-Resolution Inverse Modelling of European CH<sub>4</sub>  
Emissions Using the Novel FLEXPART-COSMO TM5 4DVAR Inverse Modelling System, *Atmos. Chem. Phys.*, 22, 13 243–13 268,  
<https://doi.org/10.5194/acp-22-13243-2022>, 2022.
- Burgers, G., Jan van Leeuwen, P., and Evensen, G.: Analysis Scheme in the Ensemble Kalman Filter, *Mon. Weather Rev.*, 126, 1719–1724,  
[https://doi.org/10.1175/1520-0493\(1998\)126<1719:ASITEK>2.0.CO;2](https://doi.org/10.1175/1520-0493(1998)126<1719:ASITEK>2.0.CO;2), 1998.
- 410 Buseck, P. R. and Pósfai, M.: Airborne Minerals and Related Aerosol Particles: Effects on Climate and the Environment, *PNAS*, 96, 3372–  
3379, <https://doi.org/10.1073/pnas.96.7.3372>, 1999.
- Chen, B., Dong, L., Huang, J., Wang, Y., Jing, Z., Yan, W., Wang, X., Song, Z., Huang, Z., Guan, X., Dong, X., and Huang, Y.: Analysis of  
Long-Term Trends in the Vertical Distribution and Transport Paths of Atmospheric Aerosols in Typical Regions of China Using 15 Years  
of CALIOP Data, *J. Geophys. Res. Atmos.*, 128, e2022JD038 066, <https://doi.org/10.1029/2022JD038066>, 2023.
- 415 Cheng, Y., Dai, T., Goto, D., Schutgens, N. A. J., Shi, G., and Nakajima, T.: Investigating the Assimilation of CALIPSO Global  
Aerosol Vertical Observations Using a Four-Dimensional Ensemble Kalman Filter, *Atmos. Chem. Phys.*, 19, 13 445–13 467,  
<https://doi.org/10.5194/acp-19-13445-2019>, 2019.
- Escribano, J., Di Tomaso, E., Jorba, O., Klose, M., Gonçalves Ageitos, M., Macchia, F., Amiridis, V., Baars, H., Marinou, E., Proestakis, E.,  
Urbanneck, C., Althausen, D., Bühl, J., Mamouri, R.-E., and Pérez García-Pando, C.: Assimilating Spaceborne Lidar Dust Extinction Can  
420 Improve Dust Forecasts, *Atmos. Chem. Phys.*, 22, 535–560, <https://doi.org/10.5194/acp-22-535-2022>, 2022.
- Evensen, G.: Sequential Data Assimilation with a Nonlinear Quasi-Geostrophic Model Using Monte Carlo Methods to Forecast Error Statis-  
tics, *J. Geophys. Res.*, 99, 10 143, <https://doi.org/10.1029/94JC00572>, 1994.
- Gueymard, C. A. and Yang, D.: Worldwide Validation of CAMS and MERRA-2 Reanalysis Aerosol Optical Depth Products Using 15 Years  
of AERONET Observations, *Atmos. Environ.*, 225, 117 216, <https://doi.org/10.1016/j.atmosenv.2019.117216>, 2020.
- 425 Gwyther, D. E., Keating, S. R., Kerry, C., and Roughan, M.: How Does 4DVar Data Assimilation Affect the Vertical Representation of  
Mesoscale Eddies? A Case Study with Observing System Simulation Experiments (OSSEs) Using ROMS v3.9, *Geosci. Model Dev.*, 16,  
157–178, <https://doi.org/10.5194/gmd-16-157-2023>, 2023.



- Hamill, T. M.: Ensemble-Based Atmospheric Data Assimilation, in: Predictability of Weather and Climate, edited by Palmer, T. and Hagedorn, R., pp. 124–156, Cambridge University Press, 1 edn., <https://doi.org/10.1017/CBO9780511617652.007>, 2006.
- 430 Han, Y., Wang, T., Tang, J., Wang, C., Jian, B., Huang, Z., and Huang, J.: New Insights into the Asian Dust Cycle Derived from CALIPSO Lidar Measurements, *Remote Sens. Environ.*, 272, 112 906, <https://doi.org/10.1016/j.rse.2022.112906>, 2022.
- Hofer, J., Althausen, D., Abdullaev, S. F., Makhmudov, A. N., Nazarov, B. I., Schettler, G., Engelmann, R., Baars, H., Fomba, K. W., Müller, K., Heinold, B., Kandler, K., and Ansmann, A.: Long-Term Profiling of Mineral Dust and Pollution Aerosol with Multiwavelength Polarization Raman Lidar at the Central Asian Site of Dushanbe, Tajikistan: Case Studies, *Atmos. Chem. Phys.*, 17, 14 559–14 577, 435 <https://doi.org/10.5194/acp-17-14559-2017>, 2017.
- Houtekamer, P. L., Mitchell, H. L., Pellerin, G., Buehner, M., Charron, M., Spacek, L., and Hansen, B.: Atmospheric Data Assimilation with an Ensemble Kalman Filter: Results with Real Observations, *Mon. Weather Rev.*, 133, 604–620, <https://doi.org/10.1175/MWR-2864.1>, 2005.
- Hunt, B. R., Kostelich, E. J., and Szunyogh, I.: Efficient Data Assimilation for Spatiotemporal Chaos: A Local Ensemble Transform Kalman 440 Filter, *Physica D*, 230, 112–126, <https://doi.org/10.1016/j.physd.2006.11.008>, 2007.
- Jin, J., Lin, H. X., Segers, A., Xie, Y., and Heemink, A.: Machine Learning for Observation Bias Correction with Application to Dust Storm Data Assimilation, *Atmos. Chem. Phys.*, 19, 10 009–10 026, <https://doi.org/10.5194/acp-19-10009-2019>, 2019.
- Jin, J., Segers, A., Lin, H. X., Henzing, B., Wang, X., Heemink, A., and Liao, H.: Position Correction in Dust Storm Forecasting Using LOTOS-EUROS v2.1: Grid-Distorted Data Assimilation v1.0, *Geosci. Model Dev.*, 14, 5607–5622, [https://doi.org/10.5194/gmd-14-5607-](https://doi.org/10.5194/gmd-14-5607-2021) 445 2021, 2021.
- Jin, J., Pang, M., Segers, A., Han, W., Fang, L., Li, B., Feng, H., Lin, H. X., and Liao, H.: Inverse Modeling of the 2021 Spring Super Dust Storms in East Asia, *Atmos. Chem. Phys.*, 22, 6393–6410, <https://doi.org/10.5194/acp-22-6393-2022>, 2022.
- Jin, J., Fang, L., Li, B., Liao, H., Wang, Y., Han, W., Li, K., Pang, M., Wu, X., and Lin, H. X.: 4DEnVar-based Inversion System for Ammonia Emission Estimation in China through Assimilating IASI Ammonia Retrievals, *Environ. Res. Lett.*, 18, 034 005, 450 <https://doi.org/10.1088/1748-9326/acb835>, 2023a.
- Jin, J., Henzing, B., and Segers, A.: How Aerosol Size Matters in Aerosol Optical Depth (AOD) Assimilation and the Optimization Using the Ångström Exponent, *Atmos. Chem. Phys.*, 23, 1641–1660, <https://doi.org/10.5194/acp-23-1641-2023>, 2023b.
- Kolb, C. E. and Worsnop, D. R.: Chemistry and Composition of Atmospheric Aerosol Particles, *Annu. Rev. Phys. Chem.*, 63, 471–491, <https://doi.org/10.1146/annurev-physchem-032511-143706>, 2012.
- 455 Law, K. J. H. and Stuart, A. M.: Evaluating Data Assimilation Algorithms, *Mon. Weather Rev.*, 140, 3757–3782, <https://doi.org/10.1175/MWR-D-11-00257.1>, 2012.
- Lee, L. A., Carslaw, K. S., Pringle, K. J., Mann, G. W., and Spracklen, D. V.: Emulation of a Complex Global Aerosol Model to Quantify Sensitivity to Uncertain Parameters, *Atmos. Chem. Phys.*, 11, 12 253–12 273, <https://doi.org/10.5194/acp-11-12253-2011>, 2011.
- Liu, Z., Liu, Q., Lin, H.-C., Schwartz, C. S., Lee, Y.-H., and Wang, T.: Three-Dimensional Variational Assimilation of MODIS Aerosol Optical Depth: Implementation and Application to a Dust Storm over East Asia, *J. Geophys. Res.*, 116, <https://doi.org/10.1029/2011JD016159>, 460 2011.
- Ma, C., Wang, T., Jiang, Z., Wu, H., Zhao, M., Zhuang, B., Li, S., Xie, M., Li, M., Liu, J., and Wu, R.: Importance of Bias Correction in Data Assimilation of Multiple Observations Over Eastern China Using WRF-Chem/DART, *J. Geophys. Res. Atmos.*, 125, e2019JD031 465, <https://doi.org/10.1029/2019JD031465>, 2020.



- 465 Manders, A. M. M., Builtjes, P. J. H., Curier, L., Denier van der Gon, H. A. C., Hendriks, C., Jonkers, S., Kranenburg, R., Kuenen, J. J. P., Segers, A. J., Timmermans, R. M. A., Visschedijk, A. J. H., Wichink Kruit, R. J., van Pul, W. A. J., Sauter, F. J., van der Swaluw, E., Swart, D. P. J., Douros, J., Eskes, H., van Meijgaard, E., van Ulft, B., van Velthoven, P., Banzhaf, S., Mues, A. C., Stern, R., Fu, G., Lu, S., Heemink, A., van Velzen, N., and Schaap, M.: Curriculum Vitae of the LOTOS–EUROS (v2.0) Chemistry Transport Model, *Geosci. Model Dev.*, 10, 4145–4173, <https://doi.org/10.5194/gmd-10-4145-2017>, 2017.
- 470 Mhawish, A., Sorek-Hamer, M., Chatfield, R., Banerjee, T., Bilal, M., Kumar, M., Sarangi, C., Franklin, M., Chau, K., Garay, M., and Kalashnikova, O.: Aerosol Characteristics from Earth Observation Systems: A Comprehensive Investigation over South Asia (2000–2019), *Remote Sens. Environ.*, 259, 112410, <https://doi.org/10.1016/j.rse.2021.112410>, 2021.  
NASA: Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observations, last access: August 2023.  
Pang, M.: Source Code of PyFilter, Zenodo, <https://doi.org/10.5281/zenodo.7611976>, last access: May 2024.
- 475 Pang, M., Jin, J., Segers, A., Jiang, H., Fang, L., Lin, H. X., and Liao, H.: Dust Storm Forecasting through Coupling LOTOS-EUROS with Localized Ensemble Kalman Filter, *Atmos. Environ.*, 306, 119831, <https://doi.org/10.1016/j.atmosenv.2023.119831>, 2023.  
Pappalardo, G., Amodeo, A., Apituley, A., Comeron, A., Freudenthaler, V., Linné, H., Ansmann, A., Bösenberg, J., D’Amico, G., Mattis, I., Mona, L., Wandinger, U., Amiridis, V., Alados-Arboledas, L., Nicolae, D., and Wiegner, M.: EARLINET: Towards an Advanced Sustainable European Aerosol Lidar Network, *Atmos. Meas. Tech.*, 7, 2389–2409, <https://doi.org/10.5194/amt-7-2389-2014>, 2014.
- 480 Peng, D. and Liu, L.: Quantifying Slab Sinking Rates Using Global Geodynamic Models with Data-Assimilation, *Earth Sci. Rev.*, 230, 104039, <https://doi.org/10.1016/j.earscirev.2022.104039>, 2022.  
Qin, W., Fang, H., Wang, L., Wei, J., Zhang, M., Su, X., Bilal, M., and Liang, X.: MODIS High-Resolution MAIAC Aerosol Product: Global Validation and Analysis, *Atmos. Environ.*, 264, 118684, <https://doi.org/10.1016/j.atmosenv.2021.118684>, 2021.  
Reichle, R. H., McLaughlin, D. B., and Entekhabi, D.: Hydrologic Data Assimilation with the Ensemble Kalman Filter, *Mon. Weather Rev.*, 130, 103–114, [https://doi.org/10.1175/1520-0493\(2002\)130<0103:HDAWTE>2.0.CO;2](https://doi.org/10.1175/1520-0493(2002)130<0103:HDAWTE>2.0.CO;2), 2002.
- 485 Sekiyama, T. T., Tanaka, T. Y., Shimizu, A., and Miyoshi, T.: Data Assimilation of CALIPSO Aerosol Observations, *Atmos. Chem. Phys.*, 10, 39–49, <https://doi.org/10.5194/acp-10-39-2010>, 2010.  
Sogacheva, L., Popp, T., Sayer, A. M., Dubovik, O., Garay, M. J., Heckel, A., Hsu, N. C., Jethva, H., Kahn, R. A., Kolmonen, P., Kosmale, M., de Leeuw, G., Levy, R. C., Litvinov, P., Lyapustin, A., North, P., Torres, O., and Arola, A.: Merging Regional and Global Aerosol Optical  
490 Depth Records from Major Available Satellite Products, *Atmos. Chem. Phys.*, 20, 2031–2056, <https://doi.org/10.5194/acp-20-2031-2020>, 2020.  
Stier, P., Schutgens, N. a. J., Bellouin, N., Bian, H., Boucher, O., Chin, M., Ghan, S., Huneeus, N., Kinne, S., Lin, G., Ma, X., Myhre, G., Penner, J. E., Randles, C. A., Samset, B., Schulz, M., Takemura, T., Yu, F., Yu, H., and Zhou, C.: Host Model Uncertainties in Aerosol Radiative Forcing Estimates: Results from the AeroCom Prescribed Intercomparison Study, *Atmos. Chem. Phys.*, 13, 3245–3270,  
495 <https://doi.org/10.5194/acp-13-3245-2013>, 2013.  
TNO: Source Code and User Guidance of LOTOS-EUROS, TNO, last access: May 2024.  
van Leeuwen, P. J.: A Consistent Interpretation of the Stochastic Version of the Ensemble Kalman Filter, *Q. J. R. Meteorolog. Soc.*, 146, 2815–2825, <https://doi.org/10.1002/qj.3819>, 2020.  
Vignati, E., Karl, M., Krol, M., Wilson, J., Stier, P., and Cavalli, F.: Sources of Uncertainties in Modelling Black Carbon at the Global Scale,  
500 *Atmos. Chem. Phys.*, 10, 2595–2611, <https://doi.org/10.5194/acp-10-2595-2010>, 2010.





- Wang, F., Yang, T., Wang, Z., Cao, J., Liu, B., Liu, J., Chen, S., Liu, S., and Jia, B.: A Comparison of the Different Stages of Dust Events over Beijing in March 2021: The Effects of the Vertical Structure on Near-Surface Particle Concentration, *Remote Sens.*, 13, 3580, <https://doi.org/10.3390/rs13183580>, 2021.
- Wang, H., Yang, T., Wang, Z., Li, J., Chai, W., Tang, G., Kong, L., and Chen, X.: An Aerosol Vertical Data Assimilation System (NAQPMS-PDAF v1.0): Development and Application, *Geosci. Model Dev.*, 15, 3555–3585, <https://doi.org/10.5194/gmd-15-3555-2022>, 2022.
- Wang, X.: Historical data on air quality in china, <https://quotsoft.net/air/>, last access: May. 2024.
- Whitaker, J. S. and Hamill, T. M.: Ensemble Data Assimilation without Perturbed Observations, *Mon. Weather Rev.*, 130, 1913–1924, [https://doi.org/10.1175/1520-0493\(2002\)130<1913:EDAWPO>2.0.CO;2](https://doi.org/10.1175/1520-0493(2002)130<1913:EDAWPO>2.0.CO;2), 2002.
- Whitaker, J. S., Compo, G. P., and Thépaut, J.-N.: A Comparison of Variational and Ensemble-Based Data Assimilation Systems for Reanalysis of Sparse Observations, *Mon. Weather Rev.*, 137, 1991–1999, <https://doi.org/10.1175/2008MWR2781.1>, 2009.
- Winker, D. M., Pelon, J., Coakley, J. A., Ackerman, S. A., Charlson, R. J., Colarco, P. R., Flamant, P., Fu, Q., Hoff, R. M., Kittaka, C., Kubar, T. L., Treut, H. L., McCormick, M. P., Mégie, G., Poole, L., Powell, K., Treppe, C., Vaughan, M. A., and Wielicki, B. A.: The CALIPSO Mission: A Global 3D View of Aerosols and Clouds, *B. Am. Meteorol. Soc.*, 91, 1211–1230, <https://doi.org/10.1175/2010BAMS3009.1>, 2010.
- Wu, X., Vu, T. V., Shi, Z., Harrison, R. M., Liu, D., and Cen, K.: Characterization and Source Apportionment of Carbonaceous PM<sub>2.5</sub> Particles in China - A Review, *Atmos. Environ.*, 189, 187–212, <https://doi.org/10.1016/j.atmosenv.2018.06.025>, 2018.
- Xu, X., Wu, H., Yang, X., and Xie, L.: Distribution and Transport Characteristics of Dust Aerosol over Tibetan Plateau and Taklimakan Desert in China Using MERRA-2 and CALIPSO Data, *Atmos. Environ.*, 237, 117 670, <https://doi.org/10.1016/j.atmosenv.2020.117670>, 2020.
- Ye, H., Pan, X., You, W., Zhu, X., Zang, Z., Wang, D., Zhang, X., Hu, Y., and Jin, S.: Impact of CALIPSO Profile Data Assimilation on 3-D Aerosol Improvement in a Size-Resolved Aerosol Model, *Atmos. Res.*, 264, 105 877, <https://doi.org/10.1016/j.atmosres.2021.105877>, 2021.
- Yumimoto, K., Nagao, T., Kikuchi, M., Sekiyama, T., Murakami, H., Tanaka, T., Ogi, A., Irie, H., Khatri, P., Okumura, H., Arai, K., Morino, I., Uchino, O., and Maki, T.: Aerosol Data Assimilation Using Data from Himawari-8, a Next-generation Geostationary Meteorological Satellite, *Geophys. Res. Lett.*, 43, 5886–5894, <https://doi.org/10.1002/2016GL069298>, 2016.
- Zhang, M., Su, B., Bilal, M., Atique, L., Usman, M., Qiu, Z., Ali, M. A., and Han, G.: An Investigation of Vertically Distributed Aerosol Optical Properties over Pakistan Using CALIPSO Satellite Data, *Remote Sens.*, 12, 2183, <https://doi.org/10.3390/rs12142183>, 2020.
- Zhao, C., Yang, Y., Fan, H., Huang, J., Fu, Y., Zhang, X., Kang, S., Cong, Z., Letu, H., and Menenti, M.: Aerosol Characteristics and Impacts on Weather and Climate over the Tibetan Plateau, *Natl Sci. Rev.*, 7, 492–495, <https://doi.org/10.1093/nsr/nwz184>, 2020.
- Zhu, C., Maharajan, K., Liu, K., and Zhang, Y.: Role of Atmospheric Particulate Matter Exposure in COVID-19 and Other Health Risks in Human: A Review, *Environ. Res.*, 198, 111 281, <https://doi.org/10.1016/j.envres.2021.111281>, 2021.