

Presentation, Calibration and Testing of the DCESS II Earth System Model of Intermediate Complexity (version 1.0)

Esteban Fernández¹, Gary Shaffer²

¹Graduate Program in Oceanography, Department of Oceanography, Faculty of Natural Sciences and Oceanography, University of Concepción, P.O. Box 160-C, Concepción, Chile

²Niels Bohr Institute, University of Copenhagen, 2100 Copenhagen Ø, Denmark

Correspondence to: Esteban Fernández (estfernandez@udec.cl)

Abstract. A new, Earth System Model of Intermediate Complexity, DCESS II, is presented that builds upon, improves and extends the Danish Center for Earth System Science (DCESS) Earth System model (DCESS I). DCESS II has considerably greater spatial resolution than DCESS I while retaining the fine, 100 m vertical resolution in the ocean. It contains modules for the atmosphere, ocean, ocean sediment, land biosphere and lithosphere and is designed to deal with global change simulations on scales of years to millions of years while using limited computational resources. Tracers of the atmospheric module are temperature, nitrous oxide, methane (^{12,13}C isotopes), carbon dioxide (^{12,13,14}C isotopes) and atmospheric oxygen. For the ocean module, tracers are conservative temperature, absolute salinity, water ¹⁸O, phosphate, dissolved inorganic carbon (^{12,13,14}C isotopes), alkalinity and dissolved oxygen. Furthermore, the ocean module considers simplified dynamical schemes for large-scale meridional circulation and sea-ice dynamics, stratification-dependent vertical diffusion, a gravity current approach to the formation of Antarctic Bottom Water and improvements in ocean biogeochemistry. DCESS II has two hemispheres with six zonal-averaged atmospheric boxes and twelve ocean ~~boxes~~ sectors distributed across the Indian-Pacific, the Atlantic, the Arctic and the Southern Oceans. A new, extended land biosphere scheme is implemented that considers three different vegetation types whereby net primary production depends on sunlight and atmospheric carbon dioxide. The ocean sediment and lithosphere model formulations are adopted from DCESS I but now applied to the multiple ocean and land regions of the new model.

A model calibration was carried out for the pre-industrial climate and model steady-state solutions were compared against available modern-day observations. For the most part, calibration results agree well with observed data, included excellent agreement with ocean carbon species. This serves to demonstrate model utility for dealing with the global carbon cycle. Finally, two idealized experiments were carried out in order to explore model performance. First, we forced the model by varying Ekman transport out of the model Southern Ocean, mimicking the effect of Southern Hemisphere westerly wind variations and second, we imposed freshwater melting pulses from the Antarctic ice sheet on to the model Southern Ocean shelf. Changes in ocean circulation and in the global carbon cycle found in these experiments are in line with ~~reasonable and agree with~~ results

~~from~~ much more complex models. Thus, we find DCESS II to be a useful and computational-friendly tool for simulations of past climates as well as for future Earth System projections.

1 Introduction

35 The carbon cycle is the backbone of the Earth's climate system since acts as a main regulator of global mean atmospheric temperature via atmospheric concentration of carbon dioxide ($p\text{CO}_2$). This cycle may be considered to be composed of two domains. One domain is a fast one with large exchange fluxes and relatively "rapid" reservoir turnovers (from years to thousands of years). This domain encompasses carbon in the atmosphere, ocean, superficial (bioturbated) ocean sediments and on land in vegetation, soil and freshwater. A second, slower domain (from hundreds of thousands to millions of years) consists of huge carbon stores in rocks and sediments that exchange carbon with the fast domain through volcanic emissions of CO_2 ,
40 chemical weathering, erosion and sediment formation on the sea floor. How this carbon is partitioned between the different Earth's reservoirs is what sets the $p\text{CO}_2$ in the atmosphere.

Earth system models (ESM) include one or both of these domains. They are thereby useful tools that can help us gain understanding of past climates as well as make future climate projections. Depending on their complexity and spatial
45 resolution, ESMs can take days, weeks or even months to run model simulations on the range of time scales mentioned above while using large computational resources. The Danish Center for Earth System Science (DCESS) model (DCESS I, Shaffer et al., 2008) is a low-order ESM with a simple geometry and ocean physics that deals with both domains of the global-scale carbon cycle and is thereby suitable for investigating Earth System changes on scales of years to millions of years while taking only minutes to days to run. The DCESS I model has proven to be useful tool in such studies as documented by many stand-
50 alone or intercomparison study publications (e.g. Eby et al., 2013; Harper et al., 2020; Joos et al., 2013; Macdougall et al., 2020; Shaffer, 2010; Shaffer and Lambert, 2018; Shaffer et al., 2009; Zickfeld et al., 2013). Nonetheless, DCESS I has serious limitations for addressing many important Earth System problems, like glacial-interglacial cycles for example, due to its one-hemisphere, two-sector horizontal resolution, lack of ocean dynamics, lack of seasonal cycles and simplified land vegetation scheme, among other factors. In order to address those deficiencies, ~~Here~~ here we present a new, Earth System Model, DCESS
55 II, that contains great improvements in model geometry and physical/biogeochemical processes. This new model is able to capture not only relevant environmental differences between major ocean basins, but also, for example, to produce synthetic ocean sediment cores from distinct ocean zones for more detailed comparison with data, and this while retaining much of the simplicity and the spirit of the DCESS I model. Thus, DCESS II is a simple, fast and highly flexible ESM of intermediate complexity well suited to run long-term experiments in a relatively simple way with no need for large computational resources.

60

This paper is organized as follows: in Sect. 2 we describe the modules of atmosphere, ocean, ocean sediment, land biosphere and lithosphere. In Sect. 3, we present the model solution and calibration procedure and show results for the model steady

state, pre-industrial simulation. In addition, we carry out two idealized experiments in order to explore and test model performance. Finally in Sect. 4 we discuss our results, outline future perspectives and present conclusions.

65 2 Model description

DCESS II is an intermediate complexity Earth System model containing atmosphere, ocean, land biosphere, lithosphere, and ocean sediment modules designed to deal with global climate simulations on scales of years to millions of years. It is an enhancement and extension of the original DCESS I model (Shaffer et al., 2008) and includes, for example, much improved horizontal and time resolution, as well as simplified ocean dynamics and seasonal cycles. Model geometry consists of two
70 hemispheres with a land/ocean area distribution and ocean depth distribution as shown in Fig. 1.




Figure 1. (a) Pre-industrial model land continental distribution and meridional boundaries of ocean boxes-sectors (dashed lines). There are twelve ocean boxes-sectors with their respective identifiers: Arc: Arctic Ocean, SOc: Southern Ocean, hn: high north, mn: mid north and en: equatorial north for Atlantic (A) and Indian-Pacific (P) Ocean. Box-Sector names in the Southern Hemisphere follow the same convention. High latitude North Pacific sector (hnP) and the Arctic Ocean are connected through the Bering Strait. Land south of 70° S is fully ice covered. Meridional sea-ice and snowline extents vary freely. (b) Observed present-day ocean area fraction profile of each ocean model box-sector calculated from Amante and Eakins (2009).
75

80 The atmospheric module considers three zonally averaged boxes per hemisphere as low, mid, and high latitude sectors divided at 35° (ϕ_{35}) and 55° (ϕ_{55}) North/South latitudes (Fig. 2a). In the ocean module we include an extra division at 65° N which allows us to consider four global ocean basins (Atlantic, Indian-Pacific, Arctic, and Southern Ocean; for simplicity, hereafter Indian-Pacific Ocean will be called only Pacific Ocean). The Southern Ocean is 360° wide and extends up to 69° S where it interacts with an ocean shelf extending to 70° S. In all there are 12 ocean sectors (compared to only 2 in DCESS I). Each ocean
85 sector is divided into 55 vertical layers with 100 m vertical resolution reaching 5500 m depth. An ocean sediment segment is

assigned to each of the layers (Fig. 2b). For each sector, ocean layer and ocean sediment areas are determined from observed ocean depth distributions (Fig. 1b).

90




Figure 2. (a) Ocean-atmosphere cross-section view depicting meridional distributions of ocean and atmospheric boxes. There are six zonal-averaged atmospheric boxes and twelve ocean [boxes-sectors](#) (see also Fig. 1a). The shaded vertical bar at 55° S extending from the surface to 2000 m depth is a virtual barrier against meridional geostrophic flow above the Drake Passage sill depth. The vertical barrier at 65° N extending from the bottom up to 1000 m depth is a physical barrier associated with Denmark Strait bathymetry. (b) Sketch of an idealized vertical area profile for an ocean [box-sector](#) (the actual profile for each [box-sector](#) is based on bathymetry observations; see Fig. 1b). Also shown is an idealized sediment segment contained in a 100 m thick box layer (the area of each specific segment is again based on bathymetric observations). The bioturbated layer (BL) is assumed to be 10 cm thick and is divided into seven sublayers as shown.

95

100 The land biosphere module considers three different types of vegetation per hemisphere whose latitudinal limits vary dynamically according to climate conditions. All model modules are described in detail in the following sections.

2.1 Atmosphere exchange, heat balance and ice/snow extent

We use a simple, zonally integrated energy balance model for the near surface atmospheric temperature, T_a (°C), forced by seasonally varying insolation, [heat exchange with the ocean](#)[air-sea-exchange](#), and meridional heat transport. In combination with simple sea ice and snow parameterizations, the model includes the ice-snow albedo feedback and the insulating effect of sea ice. Prognostic equations for mean T_a in the 0°-35°, 35°-55° and 55°-90° zones ($T_a^{l,m,h}$) are obtained by integrating the surface energy balance over the zones as,

105

$$\Lambda \frac{dT_a^{l,m,h}}{dt} = r^2 \int_0^{2\pi} \int_{\phi_{35}, \phi_{55}}^{\phi_{35}, \phi_{55}, \pi/2} (F^{\text{toa}} - F^T) \cos \phi \, d\phi \, d\xi \pm F^{\text{merid}} \quad (1)$$

110

where $\Lambda = A^{l,m,h} \rho_0 C_p b^{l,m,h}$, with $A^{l,m,h}$ the atmospheric box areas, ρ_0 the reference density of water, C_p is the specific heat capacity and $b^{l,m,h}$ are the thicknesses chosen to yield observed seasonal cycles of $T_a^{l,m,h}$ (partially based on Olsen et al., (2005)), r , ϕ and ξ are the Earth's radius, latitude and longitude respectively. Furthermore, F^{toa} and F^T are the vertical fluxes of heat through the top of atmosphere and the ocean surface, respectively, while F^{merid} correspond to the loss (equatorward box) or gain (poleward box) of heat due to meridional transport across ϕ_{35} or ϕ_{55} . A no flux boundary condition has been applied at the poles. ~~Meridional variations of $T_a(\phi)$ in the model are represented~~ At each point in time, we construct atmospheric temperature as a continuous function of latitude, $T_a(\phi)$, using ~~by~~ a fourth-order Legendre polynomial in sine of latitude, ϕ ,

$$120 \quad T_a(\phi) = \sum T_n P_n(\sin \phi); \quad n = 0, 2, 4 \quad (2)$$

where coefficients T_0 , T_2 and T_4 are determined by matching the area-weighted, zone mean values of $T_a(\phi)$ to the prognostic mean sector values of $T_a^{l,m,h}$ in each hemisphere. Observations show that eddy heat fluxes in the mid-latitude atmosphere are much greater than advective heat fluxes there (Oort and Peixoto, 1983). By neglecting the advective heat fluxes, Wang et al., (1999) developed suitable expressions for F^{merid} and the associated moisture flux, E^{merid} , in terms of T_a and $\partial T_a / \partial \phi$ at ϕ (for clarity we omit the index referencing low-mid and mid-high latitude boundaries, ϕ_{35} and ϕ_{55} respectively),

$$F^{\text{merid}} = -[K_t + L_v K_q \exp(-5420 T_a^{-1})] |\partial T_a / \partial \phi|^{m-1} \partial T_a / \partial \phi \quad (3)$$

$$E^{\text{merid}} = -K_q \exp(-5420 T_a^{-1}) |\partial T_a / \partial \phi|^{m-1} \partial T_a / \partial \phi \quad (4)$$

130

where K_t is a sensible heat exchange coefficient, K_q is a latent heat exchange coefficient and L_v is the latent heat of condensation ($2.25 \times 10^9 \text{ J m}^{-3}$). From observations, m is found to vary with latitude (Stone and Miller, 1980) and on this basis we take m to be 2.5 and 1.7 at ϕ_{35} and ϕ_{55} respectively. The temperatures and temperature gradients entering in Eqs. (3)-(4) above are obtained from Eq. (2). The poleward freshwater flux crossing a latitude ϕ is evaporated from an equatorward ocean ~~box-sector~~ and drains into the respective poleward Atlantic, Pacific, Arctic or Southern Ocean basin/~~box-sector~~ in accordance with observed land catchment areas (Rodriguez et al., 2011). In this way, each ocean ~~box-sector~~ receives a net freshwater flux from precipitation minus evaporation. The Arctic Ocean is an exception to this in as much as a fraction of E^{merid} crossing 55° N also drains there, emulating large northward flowing rivers in Canada and Siberia. Furthermore, an atmospheric zonal transport of freshwater of 0.15 Sv ($1 \text{ Sv} = 10^6 \text{ m}^3 \text{ s}^{-1}$) is prescribed from the tropical North Atlantic sector to the tropical North Pacific sector, emulating the effects of trade winds there (Dey and Döös, 2020; Lohmann and Lorenz, 2000). All these fluxes force respective ocean sectors (see Sect. 2.3).

140

Based on a calibration procedure that makes use of monthly observed atmospheric temperature, we derived the following expression for interhemispheric heat transport at the equator,

145

$$F^{eq} = K_t^{eq} \partial T_a / \partial \phi + 1.1 + [K_q^{eq} L_v \partial T_a / \partial \phi - 1.1] \exp \left[-5420 \left(\overline{T_a^{eq}}^{-1} - \overline{T_{a,PI}^{eq}}^{-1} \right) \right] \quad (5)$$

where K_t^{eq} and K_q^{eq} are sensible and latent heat coefficients and $\overline{T_a^{eq}}$ and $\overline{T_{a,PI}^{eq}}$ are the model and observed pre-industrial atmospheric mean temperature at the equator. For the $\overline{T_a^{eq}}$ calculation, Eq. (2) is used and evaluated at 0° latitude for Northern and Southern Hemispheres. The heat flux at the top of the atmosphere is taken as the balance between shortwave and longwave radiation as follows,

150

$$F^{toa} = (1 - \alpha)Q - (A + BT_a) \quad (6)$$

155

where α is the planetary albedo, equal to 0.62 for ice and snow-covered areas and equal to $\alpha_0 - \alpha_2(3 \sin^2 \phi - 1)$ otherwise. This formulation includes the effect of mean cloud cover and lower solar inclination at higher latitudes (Hartmann, 2016), and Q is the orbital, seasonal, and latitudinal (and if needed, paleo-time)-varying short-wave radiation (Berger and Loutre, 1991). The albedo of non-snow/ice-covered areas varies with vegetation type since forested areas have lower albedo than non-forested areas (Bonan, 2008). We adopt the approach in Eichinger et al., (2017) that include this effect by relating the α_0 factor to the vegetation type such as $\alpha_0 = 0.3 - \gamma (1 - f(\delta T_a)/f_0)$ where the factor 0.3 is the present-day value of α_0 , and γ is a multiplier equal to 0.02. $f(\delta T_a)$ is the ratio between the area of Grasslands/Savanna/Deserts zone and the total non-snow/ice covered area as a function of δT_a , the deviation of atmospheric temperature from the present-day value (f_0 is the ratio at $\delta T_a = 0$; see Sect. 2.5 for details about vegetation zones). Finally, α_2 is equal to 0.0875.

160

165

The second term on the right-hand side of Eq. (6) is the outgoing longwave radiation (Budyko, 1969), whereby A and BT_a are the flux at $T_a = 0$ and the deviation from this flux respectively. This simple formulation includes implicitly the radiative effects of changes in cloud cover and in atmospheric water vapor content. Greenhouse gas forcing is modelled by taking A to depend on deviations of (prognostic) atmospheric partial pressures of carbon dioxide, methane, and nitrous oxide ($p\text{CO}_2$, $p\text{CH}_4$ and $p\text{N}_2\text{O}$) from their pre-industrial values (PI), such that

170

$$A = A_{PI} - (A_{\text{CO}_2} + A_{\text{CH}_4} + A_{\text{N}_2\text{O}}) \quad (7)$$

where expressions for A_{CO_2} , A_{CH_4} and $A_{\text{N}_2\text{O}}$, ~~as~~ [are taken from Byrne and Goldblatt \(2014\); that are valid for atmospheric concentrations in the range of 200 – 10,000 ppm for CO₂ and 0.1 – 100 ppm for CH₄ and N₂O. Overlap of the absorption by](#)

175 [N₂O with CO₂ and CH₄ is included in these formulations.](#) ~~are well suited for low and high atmospheric greenhouse gas concentrations.~~ We take the year 1765 as ~~Our our~~ pre-industrial (PI) baseline [where](#) values of $p\text{CO}_{2,\text{PI}}$, $p\text{CH}_{4,\text{PI}}$ and $p\text{N}_2\text{O}_{\text{PI}}$ are 280, 0.72 and 0.27 ~~ppm_{#atm}~~, respectively, [as indicated by ice-core data \(IPCC, 2021 and citations therein\)](#). For simplicity and for the consideration of possible paleo-applications, we take constant values of A_{PI} and B for all atmospheric boxes (see Table 1). For each ocean sector, air-sea heat exchange is calculated according to Haney (1971) as,

180

$$F^T = -L_o - \lambda(T_a^{if} - T_o) \quad (8)$$

where L_o is the direct (solar) heating of the ocean surface layer, taken to be 40, 20, and 0 W m⁻² for the low, mid and high latitude sectors respectively, λ is a constant bulk transfer coefficient, as a good approximation taken to be 30 W m⁻² °C⁻¹ but
 185 set to zero for areas covered by sea ice (Haney, 1971), T_a^{if} is the ice-free mean atmospheric temperature for each sector, and T_o is the zone mean, ocean surface temperature of each ocean ~~box~~sector.

Table 1. Atmosphere module parameters.

Parameter	Symbol	Value
Sensible heat exchange coefficient	$K_t (\phi_{eq}/\phi_{lm}/\phi_{mh})$	$2.40 \times 10^{15}/4.21 \times 10^{11}/3.30 \times 10^{12} \text{ J s}^{-1} \text{ °C}^{-2.5}$
Latent heat exchange coefficient	$K_q (\phi_{eq}/\phi_{lm}/\phi_{mh})$	$-7.11 \times 10^5/1.56 \times 10^{10}/2.29 \times 10^{11} \text{ m}^3 \text{ s}^{-1} \text{ °C}^{-2.5}$
Pre-industrial, zero-degree, longwave radiation	A_{PI}	206.58 W m ⁻²
Temperature sensitivity of longwave radiation	B	2.21 W m ⁻² °C ⁻¹
Inverse timescale for sea-ice advance and retreat	Γ	$3 \times 10^{-2} \text{ s}^{-1}$
Ice-water heat exchange coefficient	κ_0	150 W m ⁻² °C ⁻¹

190 2.1.1 Sea ice and snow cover

Sea ice plays a pivotal role in the Earth's climate system, influencing radiative balance and thus global atmospheric temperature through its impact on surface albedo and air-sea heat exchange processes. For the sea ice cover, we take the same simple dynamical formulation as in Olsen et al., (2005) for the seasonal, equatorward sea ice extent. This formulation takes advantage of the meridional profile of T_a and assumes that: a) sea ice advance is proportional to the inverse timescale of cooling (τ^{adv})
 195 of the ocean mixed layer to the freezing temperature (T_f) by heat loss to the atmosphere and b) that the retreat is taken to be proportional to the inverse timescale of melting (τ^{ret}) of a seasonal sea ice cover. These inverse timescales are expressed as

$$\tau^{adv} = \frac{\lambda}{\rho_0 c_p d_u} \frac{T_a(\phi_i) - T_o}{T_o - T_f} \quad (9)$$

$$\tau^{ret} = \frac{k_i}{\delta_i^2 \rho_i L_i} [T_a(\phi_i) - T_f] + \frac{\kappa_0}{\delta_i \rho_i L_i} [T_o - T_f] \quad (10)$$

200

where d_u is the mixed layer depth (100 m), $T_a(\phi_i)$ is the atmospheric temperature at the sea ice edge, T_o is the ocean surface temperature, k_i , δ_i , ρ_i , L_i and κ_0 are the thermal conductivity of ice ($2.0 \text{ W m}^{-2} \text{ }^\circ\text{C}^{-1}$), the sea ice thickness (2 m), the density of ice (917 kg m^{-3}), the latent heat of fusion of ice ($3.34 \times 10^5 \text{ J kg}^{-1}$) and the heat transfer coefficient between ice and water ($150 \text{ W m}^{-2} \text{ }^\circ\text{C}^{-1}$; Bendtsen, 2002) respectively. Thus, changes in sea ice line position are determined individually for the

205 Arctic/Northern Atlantic, Northern Pacific and Southern Ocean as,

$$\frac{\partial \phi_i}{\partial t} = -\Gamma(\tau^{adv} - \tau^{ret}) \quad (11)$$

210 where Γ is a free parameter that together with κ_0 has been chosen to match the observed seasonal amplitude and the annual mean position of sea ice cover respectively. Land areas are covered by snow where $T_a(\phi) \leq 0 \text{ }^\circ\text{C}$ and taken to respond instantaneously to atmospheric temperature changes. Furthermore, present-day Northern Hemisphere ice sheets are prescribed as a constant area with ice albedo throughout the year independent of snow line position. All of Antarctica is covered by ice in the model.

2.2 Atmospheric chemistry and air-sea gas exchange

215 For each atmospheric box we consider partial pressures of $^{12,13,14}\text{CO}_2$, $^{12,13}\text{CH}_4$, N_2O and O_2 . The prognostic equation for the partial pressure of a gas χ is taken to be

$$\frac{dp(\chi)}{dt} = \frac{1}{v_a} [A_o^{if} \Psi_S(\chi) + \Psi_I(\chi) \pm \Psi_T(\chi)] \quad (12)$$

220 where v_a is an atmospheric mole volume, A_o^{if} an ice-free ocean surface area, Ψ_S is an air-sea gas exchange flux, Ψ_I are sources or sinks within the atmosphere or net transports to or from the atmosphere via weathering, volcanism, interaction with land biosphere and, for recent times, anthropogenic activities, and Ψ_T is the gas transport between adjacent atmospheric boxes.

Air-sea exchange for $^{12}\text{CO}_2$ for each atmospheric box is written as (for simplicity we omit atmospheric box index superscript)

225

$$\Psi_S = k_w \eta_{\text{CO}_2} (p\text{CO}_2 - p\text{CO}_{2,w}) \quad (13)$$

where the gas transfer velocity k_w is $0.39 u^2 (Sc/660)^{-0.5}$ with u the long-term annual mean wind speed at 10 m above ocean surface, Sc the CO_2 Schmidt number that depends on prognostic temperatures of the ocean surface layers (Wanninkhof, 1992),

230 and η_{CO_2} is the CO_2 solubility that is a function of temperature and salinity of the surface ocean layer (Weiss, 1974). $p\text{CO}_{2,w}$ is the prognostic CO_2 partial pressure at the ocean surface layer equal to $[\text{CO}_2]/\eta_{\text{CO}_2}$ whereby $[\text{CO}_2]$ is the prognostic dissolved CO_2 concentration of the ocean surface layer calculated from ocean carbonate carbon chemistry (see Sect. 2.4). Wind speeds are not calculated in our simplified atmosphere module, so we use observed values from the NOAA/CIRES/DOE 20th Century Reanalysis (V3) dataset for each ocean sector. For extremely warm climates experiments [\(annual mean sea-surface](#)
 235 [temperatures over 30 °C\)](#) we use the Schmidt number formulation from Gröger and Mikolajewicz (2011) who demonstrated that the Wanninkhof (1992) formulation underestimates Sc in such conditions.

Air-sea exchange for ${}^i\text{CO}_2$ ($i=13$ and 14) is given by

$$240 \quad \Psi_S^i = k_w \eta_{\text{CO}_2} {}^i\alpha_k \left({}^i\alpha_{aw} p^i\text{CO}_2 - {}^i\alpha_{wa} p\text{CO}_{2,w} \frac{[\text{DI}^i\text{C}]}{[\text{DIC}]} \right) \quad (14)$$

where ${}^i\alpha_k = 0.99912$ is the kinetic fractionation factor (Zhang et al., 1995), $[\text{DI}^i\text{C}]$ and $[\text{DIC}]$ are concentrations of dissolved inorganic carbon in the surface ocean layer, ${}^i\alpha_{aw}$ is the fractionation factor due to different solubilities in the equilibration process and ${}^i\alpha_{wa}$ is the fractionation factor in the dissociation reactions associated with ocean carbonate chemistry expressed
 245 as,

$${}^i\alpha_{wa} = \frac{[\text{CO}_2] + {}^i\alpha_{\text{HCO}_3}[\text{HCO}_3^-] + {}^i\alpha_{\text{CO}_3}[\text{CO}_3^{2-}]}{[\text{DIC}]} \quad (15)$$

where ${}^i\alpha_{\text{HCO}_3}$ and ${}^i\alpha_{\text{CO}_3}$ are individual fractionation factors for the carbon species $[\text{HCO}_3^-]$ and $[\text{CO}_3^{2-}]$ and surface layer values
 250 for these species are obtained from ocean carbonate chemistry calculations. Moreover, ${}^i\alpha_{\text{HCO}_3}$ and ${}^i\alpha_{\text{CO}_3}$ are functions of surface layer temperature (Zhang et al., 1995). As the rate at which different isotopes of a chemical element take part in a chemical reaction depends on their mass, we assume the ${}^{14}\text{C}/{}^{12}\text{C}$ fractionation to be twice as strong as for ${}^{13}\text{C}/{}^{12}\text{C}$, then ${}^{14}\alpha = 1 - 2(1 - {}^{13}\alpha)$.

255 For air-sea exchange of oxygen for each ocean surface sector

$$\Psi_S = k_w (\eta_{\text{O}_2} p\text{O}_2 - [\text{O}_2]) \quad (16)$$

with k_w as before but with O_2 Schmidt numbers that depend on prognostic temperatures of the ocean surface layers (Keeling
 260 et al., 1998). The O_2 solubility (η_{O_2}) was converted from Bunsen solubility coefficients that depend on prognostic temperatures and salinities (Weiss, 1970) to model units using the ideal gas mole volume, $[\text{O}_2]$ is the prognostic dissolved oxygen

concentration in the ocean surface layer (see Sect. 2.4). At present we do not include air-sea exchange of methane and nitrous oxide in this model version. However, this could be readily accomplished upon need, as this has been successfully implemented in the DCESS I model (Shaffer et al., 2017).

265

With regard to sources/sinks (the second term on the right-hand side of Eq. (12)), the model considers the following sources/sinks for each atmospheric tracer:

For carbon dioxide, there is net exchange with the land biosphere, oxidation of atmospheric methane, volcanic input, weathering of “old” organic carbon in rocks and weathering of carbonate and silicate rocks. As needed, anthropogenic CO₂ sources may be added associated with fossil fuel burning and/or land use change. All these sources/sinks are also considered for atmospheric ¹³CO₂. For ¹⁴CO₂ the same sources/sinks as above are included, except for old (and thus ¹⁴C-free) carbon sources which are inputs from volcanoes, organic carbon weathering and fossil fuel burning. In addition, ¹⁴C is produced naturally in the atmosphere via cosmic ray flux and, in recent times, by atomic bomb testing. For each atmospheric box, the cosmic ray source of ¹⁴CO₂ can be expressed as $A^{l,m,h} P_{14C}^{l,m,h} / A_{vg}$, where A_{vg} is the Avogadro number and $P_{14C}^{l,m,h}$ (in atoms m⁻² s⁻¹) is the magnitude of ¹⁴C production, chosen here to match the estimated pre-industrial atmospheric ¹⁴C concentration such that $\Delta^{14}C_{atm} \sim 0$ ‰. A small amount of atmospheric ¹⁴C enters the land biosphere and decays there radioactively. A smaller fraction decays directly in the atmosphere becoming an atmospheric sink of ¹⁴CO₂ with a decay rate λ_{14C} of 3.84×10^{-12} s⁻¹. By far, most of the ¹⁴C produced enters the ocean by air-sea exchange and by far most of this isotope decays within the ocean. Finally, a small amount of ¹⁴C enters the ocean sediment via sinking of biogenic particles and part of this returns to the ocean due to remineralization/dissolution in the ocean sediment.

For methane there is production within the land biosphere (see Sect. 2.56) and consumption associated with OH radicals in the troposphere. The latter leads to the net consumption of 2 O₂ molecules and production of 1 CO₂ molecule for every CH₄ molecule consumed. Since this reaction depletes the concentration of these radicals, atmospheric lifetime of CH₄ grows as methane concentration rises. We include this effect in the model by taking the atmospheric methane sink to be $\lambda_{CH_4} pCH_4$, with $\lambda_{CH_4} = v_a / \tau_{CH_4} M(pCH_4) / \tau_{CH_4,PI}$ where τ_{CH_4} is the (variable) atmospheric lifetime of methane. This lifetime is determined by fitting a function to the results from several modelling studies that consider a wide range of pCH₄ values. This fit yields a pre-industrial lifetime of 9.5 years, increasing for example to 10.8 and 15.1 years for 2 and 10 times the pre-industrial methane level, respectively (see Shaffer et al., 2017 for details). ~~is a fitted function from several modelling studies that consider a wide range of pCH₄ values and $\tau_{CH_4,PI}$ is the atmospheric pre industrial lifetime of methane equal to 9.5 years (Shaffer et al., 2017).~~ Potential sources like melting of methane hydrate in the arctic tundra and in ocean sediments and by human activities may be included in the model as needed. For ¹³CH₄ the same processes are considered but with their

285

290

295 respective fractionation factors (Sect. 2.5) and adding the $p^{13}\text{CH}_4/p^{12}\text{CH}_4$ ratio to the above OH-related atmospheric methane sink formulation.

For nitrous oxide there is production within the land biosphere and consumption in the atmosphere, here mainly due to photodissociation in the stratosphere. This is modelled as $\lambda_{\text{N}_2\text{O}}p\text{N}_2\text{O}$, with $\lambda_{\text{N}_2\text{O}} = v_a/\tau_{\text{N}_2\text{O}}$ where $\tau_{\text{N}_2\text{O}}$, the atmospheric lifetime of N_2O , is taken to be 150 years. Potential sources like N_2O flux from ocean denitrification and by human activities
300 may be included in the model as needed.

For oxygen, there is consumption associated with oxidation of atmospheric methane and reaction with OH radicals (see above), and sink (source) from organic matter remineralization (photosynthesis) on land. [Since methane is the end product of some of the remineralization on land \(see section 2.6\), the land biosphere is a net \$\text{O}_2\$ source in a steady state.](#) Furthermore, there are
305 long-term atmospheric oxygen sinks due to weathering of organic carbon in rocks and oxidation of reduced carbon emitted in lithosphere outgassing. A long-term, [quasi-steady state](#) of $p\text{O}_2$ is achieved in the model when these latter sinks balance net O_2 outgassing from the ocean from less O_2 consumption than production there due to burial of organic matter in the model ocean sediments. [However, for multi-million-year time scales, the global sulfur cycle would need to be included in the model to achieve “true” \$p\text{O}_2\$ steady states \(Berner, 2006\).](#) Additional sinks (sources) of atmospheric O_2 associated with recent land use
310 change and with burning of fossil fuels may be included in the model as needed.

The last term in Eq. (12) representing the gas transport between two adjacent atmospheric boxes is modelled as

$$\Psi_T(\chi) = -k_\chi \frac{dp(\chi)}{d\phi} \quad (17)$$

315

with $k_\chi = 6 \times 10^{-8} \text{ s}^{-1}$ based on data-model comparisons we made using the DCESS I model (Shaffer et al., 2008). For simplicity, we take the same value for all gases and for all atmospheric boundaries.

Evaporation and precipitation modify both salinity and oxygen isotopic composition in the surface ocean. Fractionation due to
320 evaporation enriches (depletes) the ^{18}O content in sea water (water vapor) from net evaporative model ocean (atmospheric) zones. Further depletion in water vapor takes place due to fractionation in the condensation process as the air mass cools in its poleward path. Given the importance of ^{18}O in paleoclimate studies, we incorporate atmospheric cycling of oxygen isotopes of water following Olsen et al., (2005), which gives the ^{18}O content in the well-known delta notation ($\delta^{18}\text{O}_w$) relative to Standard Mean Ocean Water (SMOW). This approach takes advantage of meridional atmospheric temperature profiles
325 estimating $\delta^{18}\text{O}_a$ at every atmospheric box division, where subindex “a” refers to isotopic excursion of ^{18}O in the atmosphere.

2.3 Ocean circulation and mixing

Simplified ocean dynamics in the model consist of a balance between the pressure gradient force and linear (Rayleigh) friction acting on the meridional velocity. Model flow is defined by this relation and hydrostatic and continuity equations,

$$330 \quad -\frac{1}{r\rho_0} \frac{\partial P}{\partial \phi} - r_f v = 0 \quad (18)$$

$$-\frac{1}{\rho_0} \frac{\partial P}{\partial z} - \frac{\rho}{\rho_0} g = 0 \quad (19)$$

$$\frac{1}{r \cos \phi} \frac{\partial v \cos \phi}{\partial \phi} + \frac{\partial w}{\partial z} = 0 \quad (20)$$

where r_f is a friction coefficient (Table 2), v and w are the meridional and vertical velocity components respectively, P is the pressure, g is gravity and ρ is the water density, calculated using the non-linear function of temperature, salinity, and pressure according to the TEOS-10 standard (International Thermodynamic Equation Of Seawater, IOC et al., (2010)) but modified for Boussinesq ocean models (Roquet et al., 2015).

340 **Table 2. Ocean module physical and biogeochemical parameters as well as applied Redfield ratios.**

Parameter	Symbol	Value
Ocean Physics		
Meridional flow friction coefficient	r_f	$1 \times 10^{-4} \text{ s}^{-1}$
Deep ocean horizontal diffusivity	K_h^d	$1 \times 10^3 \text{ m}^2 \text{ s}^{-1}$
Surface ocean horizontal diffusivity	K_h^s	$2.5 \times 10^4 \text{ m}^2 \text{ s}^{-1}$
e-folding depth for horizontal diffusivity scale	z_g	200 m
Glacial melt water flux to Southern Ocean shelf	F_g	0.07 Sv
Sea ice export flux from Southern Ocean shelf	F_i	0.13 Sv
Ocean Biochemistry		
Limitation factor (Southern Ocean/Arctic Ocean)	L_f	0.15/0.30 s^{-1}
Upper limit for rain ratio	$r_{Cal,max}$	0.35
Steepness factor for calcite production	μ	0.16
Nutrient remineralization length scale	ξ_N	833 m
Carbon remineralization length scale	ξ_C	769 m
Calcite dissolution length scale	ξ_{Cal}	2500 m
Redfield ratios		

Carbon to phosphate	r_{CP}	106
Alkalinity to phosphate	r_{AlkP}	16
Oxygen to phosphate (nutrient component)	r_{ONP}	32
Oxygen to phosphate (carbon component)	r_{OCP}	118

Since there are no meridional boundaries at the Drake Passage, a geostrophic flow cannot be maintained above the sill depth there. To account this feature in our simplified model, a virtual barrier against meridional flow is placed at 55° S from surface to 2000 m depth (Fig. 2). Equatorward Ekman transport there, forced by prevailing westerlies winds, is included by prescribing a net northward volume flux (E_k) in the surface layer taken to be 30 Sv. This transport, which carries Southern Ocean surface water properties, is injected into the “South Atlantic/South Pacific” sector on the corresponding density level, forming the model’s Antarctic Intermediate Water (AAIW). Moreover, a northward surface flow of 1 Sv at 65° N is prescribed to consider the net ocean exchange between the North Pacific and the Arctic Ocean across the Bering Strait. This flow carrying surface North Pacific properties enters on the corresponding density level in the Arctic Ocean. At all other latitudes, meridional flow is set to zero in the surface layer where exchanges rely entirely on horizontal and vertical mixing processes (Shaffer and Olsen, 2001). The zonally averaged velocity at each meridional zone boundary is related to the density field according to

$$v(z > d_u) = -\frac{g}{\alpha r_f \rho_0} \int_0^z \frac{\partial \rho}{\partial \phi} dz - \frac{g}{\alpha r_f} \frac{\partial \eta}{\partial \phi} \quad (21)$$

where η is the sea level and d_u is the depth of the mixed layer or, for Southern Ocean, the Drake Passage sill depth and for Arctic Ocean, the Denmark Strait sill depth. Sea levels of the model ocean zones are adjusted instantaneously to conserve mass and to form sea level gradients in Eq. (21) that lead to ocean meridional transports needed to balance the Ekman transport and atmospheric freshwater forcing at the ocean surface. Finally, the vertical flow is calculated from continuity equation, given meridional flow, Ekman transport, atmospheric freshwater forcing and shelf exchange (see below).

The effect of wind-driven gyre mixing is parameterized by a depth-dependent, horizontal diffusivity,

$$K_h(z) = K_h^d + K_h^s \exp\left(-\frac{z}{z_g}\right) \quad (22)$$

where K_h^d and K_h^s denote the deep and surface horizontal diffusivities and z_g is an e-folding, gyre depth (Table 2). At the Southern Ocean where wind-driven gyre mixing is not important, K_h^s is decreased by 90 %. The same approach is taken at the North Pacific sector where North Pacific Intermediate Water (NPIW) is formed with a reduction of 50 % in K_h^s . With this setting, the model can reproduce global-scale intermediate water masses (AAIW and NPIW) in a simple way as reported by

other model studies (England, 1993; Stocker et al., 1994). This is one of the few “regional tunings” in the model. The vertical
 370 diffusivity, K_v , is modelled using a stratification-dependent form, such that

$$K_v(z) = \min \left[K_v^{\max}, K_v^0 \left(\frac{N}{N_0} \right)^{-\gamma} \right]; \text{ for } N > 0 \quad (23)$$

where N is the Brunt-Väisälä frequency equal to $[(g/\rho_0)(\partial\rho/\partial z)]^{1/2}$. According to Gargett and Holloway (1984), we set
 375 $\gamma = 0.5$ indicating a relatively weak dependence on stratification associated with diapycnal mixing via breaking of internal
 waves. Furthermore, K_v^0 and N_0 are the respective scale values of K_v and N taken as global representative thermocline values
 equal to $3 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$ and 10^{-2} s^{-1} respectively (Kunze, 2017). For unstable stratification K_v is equated with K_v^{\max} ($8 \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$),
 accounting for model convective adjustment.

2.3.1 Southern Ocean shelf and Antarctic Bottom Water formation

380 Antarctic Bottom Water (AABW) constitutes a major ocean water mass filling and ventilating the abyssal ocean on a global
 scale (Johnson, 2008; Orsi et al., 1999). Also, AABW plays a crucial role through the global overturning circulation in the
 transport and redistribution of heat, salt, nutrients and carbon (among other tracers) and thus affects atmospheric CO_2
 concentrations and thereby global climate (Orsi, 2010; Purkey and Johnson, 2013).

385 AABW originates from dense shelf water, formed by air cooling and brine-rejection during sea ice formation, that can flow
 down the slope reaching abyssal ocean depths (Gordon, 2019). To address this process, we include a 500 m deep Southern
 Ocean shelf (Heywood et al., 2014) between 69° S and 70° S . Furthermore, we assume that there is enough air-sea heat
 exchanges on the shelf to maintain temperature there at freezing temperature T_f . Then, the prognostic equation for all shelf
 tracers but temperature (Ψ_{sh}) is,

$$\frac{\partial \Psi_{sh}}{\partial t} = \frac{1}{V_{sh}} \left[(F_g - F_c) \Psi_g - F_i \Psi_i + F_r \Psi_{SO} - F_{sl,0} \Psi_{sh} + F_{ge} A_{sh}^{if} \right] \quad (24)$$

where V_{sh} is the shelf volume, F_g is the freshwater flux due to ice sheet/calved iceberg melt and precipitation and F_c is the
 iceberg export from the shelf taken as 0.05 Sv and 0.02 Sv respectively. Furthermore, F_i is the sea-ice export to the Southern
 395 Ocean taken as 0.13 Sv (Haumann et al., 2016), F_r is the replacing flux from the Southern Ocean to the shelf and $F_{sl,0}$ is the
 volume flux leaving the shelf at the shelf break (see below). Each of these fluxes carry their own tracer values (Ψ_g , Ψ_i , Ψ_{SO}
 and Ψ_{sh}). Values of Ψ_g are zero for salinity, phosphate and alkalinity tracers, $\eta_{\text{CO}_2} P^{12,13,14} \text{CO}_2$ for $\text{DI}^{12,13,14}\text{C}$, $\eta_{\text{O}_2} \text{O}_2$ for
 dissolved oxygen and a constant value of -40 ‰ for $\delta^{18}\text{O}_w$. The latter is a rough estimate for the mean present day $\delta^{18}\text{O}_w$ value
 of ice exported from Antarctica based on snow input values and ice sheet flow (Masson-Delmotte et al., 2008). For Ψ_i , sea-ice

400 takes the shelf water value, except for salinity which is taken as constant equal to 5 g kg^{-1} . Furthermore, a fraction of the freshwater flux crossing 55° S falls directly into the Southern Ocean and the rest goes to Antarctica. In steady-state the amount going to Antarctica is equal to F_g . The last right-hand term corresponds to the air-sea gas exchange where F_{ge} is the gas flux and A_{sh}^{if} is the ice-free surface shelf area, meant mainly to represent coastal polynyas. Here we take A_{sh}^{if} to be 20 % of the total shelf area.

405

If it is denser than ambient Southern Ocean water, shelf water flows down along the continental slope. On a path to the deep ocean, downslope flow takes up ambient water via entrainment, typically increasing its volume more than twice its value at shelf break to finally fill the abyssal ocean as AABW (Orsi, 2002; Orsi et al., 2001). In order to include a comparable AABW formation process in our model, we address this using a simple formulation of entrainment following approaches described in
 410 Baines (2005, 2008). The governing equations for the downslope flux (F_{sl}) and the plume height (or thickness, H) normal to the slope are

$$\frac{dF_{sl}}{ds} = E \frac{F_{sl}}{H} \quad (25)$$

$$\frac{dH}{ds} = 2E + C_d - S_2 Ri \tan \theta \quad (26)$$

415

where s is the downslope distance from the shelf break depth obtained from observed bathymetry (Amante and Eakins, 2009) and E is the entrainment coefficient equal to $E_0(1 - Ri/Ri_c)$ for $0 \leq Ri \leq Ri_c$ and zero otherwise. Here Ri is the Richardson number, Ri_c is a critical value for Ri and E_0 is an amplitude parameter. Ri is defined as $GH^3 \cos \theta / F_{sl}^2$ whereby $G = g\Delta\rho(z)/\rho_0$ is the buoyancy, $\Delta\rho(z)$ is the difference between downflow density and the mean local ambient density and θ is
 420 the slope angle. Values for Ri_c and E_0 are taken to be 0.25 and 0.20 respectively (Xu et al., 2006). Finally, C_d and S_2 are the drag coefficient and a constant, respectively, with values taken from the literature cited above. Initial conditions at the shelf break [are](#) taken to be $H_0 = 100 \text{ m}$ and $F_{sl,0} = H_0(G_0 H_0)^{0.5}$ with G_0 calculated as above using the Southern Ocean density at the shelf break depth. We integrate Eq. (25) and Eq. (26) every 2 meters from the shelf break until the depth where the plume and the ambient water buoyancy are close enough as defined by a threshold. Then this downslope flow enters Southern Ocean
 425 model layers in accordance with the plume height. Although other overflow approaches have been proposed (Danabasoglu et al., 2010; Xu et al., 2006), we think that this simple and fast approach is well suited to capture the formation and insertion of AABW at depth and is a particular strength in our simplified model. In this context we note that, quite a few CMIP6 models form AABW incorrectly by deep, open-ocean convection and/or with lack of plume entrainment (Heuzé, 2021).

430 With the above physics and for any ocean zone, the general conservation equation of any ocean tracer ψ may be written as

$$\frac{\partial \psi}{\partial t} + \frac{1}{a \cos \phi} \frac{\partial(\cos \phi v \psi)}{\partial \phi} + \frac{\partial(w \psi)}{\partial z} = \frac{1}{a^2 \cos \phi} \frac{\partial}{\partial \phi} \left(\cos \phi K_h \frac{\partial \psi}{\partial \phi} \right) + \frac{\partial}{\partial z} \left(K_v \frac{\partial \psi}{\partial z} \right) + \Psi_S(\psi) + \Psi_B(\psi) + \Psi_I(\psi) \quad (27)$$

where Ψ_S is the air-sea exchange of heat and gases, Ψ_B is the exchange of dissolved substances with the ocean sediment and Ψ_I are internal source/sinks into the water column. Ocean tracers of temperature, salinity and $\delta^{18}\text{O}_w$ are forced only at the ocean surface via air-sea heat exchange and direct solar forcing for temperature and freshwater forcing for salinity and $\delta^{18}\text{O}_w$. Pacific and Atlantic Ocean model sectors between 35° S and 55° S south of Africa are connected via a zonal surface-intensified mixing scheme in order to emulate the role of Antarctic Circumpolar Current there. These specific terms are added to Eq. (27) for tracers of these [boxes-sectors](#) (msA and msP).

440 2.4 Ocean biogeochemical cycling

The ocean module considers the following biogeochemical ocean tracers: phosphate (PO_4), dissolved oxygen (O_2), dissolved inorganic carbon (DIC) in $^{12,13,14}\text{C}$ species, and alkalinity (ALK), which are all forced by new (export) production of organic matter and biogenic calcium carbonate shells in the lighted ocean surface layers. Furthermore, PO_4 , $\text{DI}^{12,13,14}\text{C}$ and ALK are forced by river inputs and concentration/dilution of the surface layer by evaporation/precipitation. Moreover, O_2 and $\text{DI}^{12,13,14}\text{C}$ are forced by air-sea exchange. In the ocean interior, all these tracers are influenced by remineralization of organic matter and dissolution of CaCO_3 shells in the water column as well as exchange with the ocean sediment. DI^{14}C is affected by radioactive decay in all ocean layers. For simplicity, we have neglected explicit nitrogen cycling and have assumed that all biogenic matter export from the surface layer is in the form of particle organic matter (POM) and that all CaCO_3 is in the form of calcite.

450 We take new (export) production of organic matter (NP) in each model surface layer to be a function of phosphorus (Maier-Reimer, 1993; Yamanaka and Tajika, 1996) and solar radiation as

$$\text{NP} = A_o^{if} z_{eu} L_f [\text{PO}_4] \frac{I}{I+I_{1/2}} \frac{[\text{PO}_4]}{[\text{PO}_4]+P_{1/2}} \quad (28)$$

455 where A_o^{if} is the ice-free ocean surface area, z_{eu} is the euphotic layer depth (100 m), $[\text{PO}_4]$ and I are phosphate content in the surface ocean and solar radiation there, respectively. $P_{1/2}$ and $I_{1/2}$ are their respective half saturation constants equal to $1 \mu\text{mol m}^{-3}$ for phosphate and 100 W m^{-2} for light (Mutshinda et al., 2017). L_f is an efficiency coefficient (units of s^{-1}) that estimates iron and/or other limitation factors on net primary production. We take the value of L_f to be equal to 1 for most ocean zones but set to some lower value for Southern and Arctic Oceans as determined by model fit to ocean data (Sect. 3.1.2). In the surface layer sources/sinks due to new production for PO_4 , DI^{12}C , ALK and O_2 are $-\text{NP}$, $-r_{\text{CP}}\text{NP}$, $r_{\text{AlkP}}\text{NP}$ and $(r_{\text{OCP}} + r_{\text{ONP}})\text{NP}$ respectively, where r_{CP} , r_{AlkP} , r_{OCP} and r_{ONP} are the Redfield ratios of C:P, ALK:P, $(\text{O}_2)_\text{C}:\text{P}$ and $(\text{O}_2)_\text{N}:\text{P}$ respectively. The subscripts C and N refer to a division of POM produced into “carbon” and “nutrient” parts respectively, as

explained below. We adopted the Redfield ratios used in Shaffer et al., (2008) and shown in Table 2. For $DI^{13,14}C$, the surface sink due to new production and associated isotope fractionation is $-^{13,14}\alpha_{Org}([DI^{13,14}C]/[DI^{12}C])_{eu} r_{CP}NP$. We take the
 465 fractionation factor $^{13}\alpha_{Org}$ to depend on surface ocean concentrations of dissolved carbon dioxide ($[CO_{2(aq)}]$) and phosphate according to Pagani et al., (1999):

$$^{13}\alpha_{Org} = 25 - \frac{116.96 L_f [PO_4] + 81.42}{[CO_{2(aq)}]} \quad (29)$$

470 where concentrations are in $\mu\text{mol kg}^{-1}$. As previously, $^{14}\alpha_{Org} = 1 - 2(1 - ^{13}\alpha_{Org})$.

Surface (export) production of biogenic calcite is related to new (export) production by $r_{Calc}r_{CP}NP$ where r_{Calc} is the “rain” ratio, the ratio between the production of $CaCO_3$ to the production of organic carbon. It is parameterized according to Maier-Reimer (1993) and Marchal et al., (1998) but with the addition of a dependence on the calcite saturation state of the ocean
 475 surface layer (Ω_S) following Shaffer et al., (2016):

$$r_{Calc} = r_{Calc,max} \frac{\exp(\mu(T_s - T_{ref}))}{1 + \exp(\mu(T_s - T_{ref}))} \frac{\Omega_S - 1}{v + (\Omega_S - 1)} \quad (30)$$

where $r_{Calc,max}$ is a rain ratio upper limit, μ is the steepness factor, T_s and T_{ref} are the surface temperature and the reference
 480 temperature (10 °C) respectively, and v is a half saturation constant taken to be 1 (Gangstø et al., 2011). Furthermore, $\Omega_S = [Ca^{2+}][CO_3^{2-}]/K_{sp}$ with calcium concentration given as $[Ca^{2+}] = [Ca^{2+}]_m(S/S_m)$ where $[Ca^{2+}]_m$ and S_m are the global ocean mean calcium and salinity values, taken as 10.57 mol m^{-3} and 35 for present day respectively, S is the ocean salinity and K_{sp} is the calcite solubility coefficient. There is no biogenic calcite production for subsaturated conditions ($r_{Calc} = 0$ for $\Omega_S \leq 1$). Values for $r_{Calc,max}$ and μ are determined by model fit to ocean and ocean sediment data. With this, surface sinks for $DI^{12}C$
 485 and ALK due to biogenic calcite production are $-r_{Calc}r_{CP}NP$ and $-2r_{Calc}r_{CP}NP$ respectively. For $DI^{13,14}C$, the surface sinks due to calcite production and associated isotope fractionation are $-^{13,14}\alpha_{Calc}([DI^{13,14}C]/[DI^{12}C])_{eu} r_{Calc}r_{CP}NP$, where $\alpha_{Calc} = 1$, assuming no carbon fractionation during biogenic calcite formation in the ocean surface layer.

Particles are assumed to sink out of the surface layer with settling speeds high enough to neglect advection and diffusion of
 490 them. This particulate flux decreases significantly with depth due to subsurface remineralization/dissolution with only a small fraction reaching the sea floor as shown by sediment trap data (Martin et al., 1987). To address this, we assume an exponential-type law for the vertical fraction of particulate organic matter (POM) “nutrient” and “carbon” component, each with a distinct e-folding length (ξ_N and ξ_C) motivated mainly by results of Shaffer et al., (1999). Additionally, we also include temperature

dependence (λ_Q) on remineralization rates as indicated by ocean data (Laufkotter et al., 2017; Marsay et al., 2015). Therefore, tracer sources in the water column due to remineralization of POM, “nutrient” (Φ_N) and “carbon” component (Φ_C), are expressed as the vertical gradient of POM,

$$\Phi_{N,C}(z) = \frac{\partial \text{POM}(z)_{N,C}}{\partial z} = \text{POM}(z)_{N,C} \frac{\lambda_Q(T_o(z))}{\xi_{N,C}} \quad (31)$$

where λ_Q is defined as $Q_{10}^{(T_o(z)-T_{o,ref})/10}$, whereby Q_{10} is the biotic activity increase for a 10 degree increase of T_o , $T_{o,ref}$ is a reference temperature taken as the present-day global area-weighted mean observed temperature from the World Ocean Atlas 2018 database (Boyer et al., 2018) for the upper 500 m (Komar and Zeebe, 2021) and $\xi_{N,C}$ are e-folding lengths for $T_o(z) = T_{o,ref} = 9.3$ °C. For the biogenic calcium carbonate particles flux (PCal), we take a simple exponential law with a constant e-folding length ξ_{Cal} and, as above, tracer sources in the water column due to dissolution of CaCO_3 produced in the euphotic zone (Φ_{Cal}) is expressed as

$$\Phi_{Cal}(z) = \frac{\partial \text{PCal}(z)}{\partial z} = \frac{\text{PCal}(z)}{\xi_{Cal}} \quad (32)$$

In Eqs. (31) and (32), POM_N , POM_C and PCal at $z = 0$ are the respective surface layer export productions, NP , $r_{CP}\text{NP}$ and $r_{CalC}r_{CP}\text{NP}$. Given the range of Q_{10} values shown by data and modelling studies (Laufkotter et al., 2017; Regaudie-de-Gioux and Duarte, 2012; Bendtsen et al., 2015) and to maintain model simplicity (see Sect. 2.5 and Sect. 2.6) we choose a constant value for Q_{10} of 2. Also for simplicity, ξ_N , ξ_C and ξ_{Cal} are taken to be constants for all ocean zones whose values (Table 2) has been chosen to fit ocean data (Sect. 3.1.2). Thus, the vertical source/sinks in the ocean interior for PO_4 , DI^{12}C , ALK and O_2 are Φ_N , $(\Phi_C + \Phi_{Cal})$, $(2\Phi_{Cal} - r_{AlkP}\Phi_N)$ and $(r_{ONP}\Phi_N + r_{OCP}\Phi_C)$ respectively. For $\text{DI}^{13,14}\text{C}$, the vertical distribution from remineralization and dissolution is $([\text{DI}^{13,14}\text{C}]/[\text{DI}^{12}\text{C}])_{eu} [^{13,14}\alpha_{Org}\Phi_C + ^{13,14}\alpha_{Cal}\Phi_{Cal}]$. All these vertical distributions are weighted by the ocean area profile $A_o(z)$ for each zone. In addition, the fluxes of P and $^{12,13,14}\text{C}$ that fall in the form of POM and/or biogenic calcite particles on the model ocean sediment surface at any depth of each zone are calculated as the product of $dA_o(z)/dz$ there and the difference between the particulate fluxes falling out of the ocean surface layer and the remineralization/dissolution taking place down to the depth of each zone.

520

Non-linear ocean carbonate chemistry is calculated using the recursive formulation of Antoine and Morel (1995) as explained in detail in Shaffer et al., (2008) in the context of a DCESS-model approach. This system yields ocean distributions of $\text{CO}_{2(aq)}$, CO_3^{2-} , HCO_3^- and hydrogen ion concentrations needed for calculations of air-sea exchange of carbon dioxide, carbon isotopic fractionation during air-sea exchange and in ocean new production, dissolution of calcite in the ocean sediment, and pH

525 (seawater scale) calculations. Profiles of carbonate saturation with respect to calcite are calculated as $K'_{CaCO_3}/([Ca^{2+}]_m S/S_m)$, where K'_{CaCO_3} is the apparent dissociation constant for calcite as function of T, S and pressure (Mucci, 1983).

2.5 Ocean Sediment

For the sediment module, we adopt the approach developed by Shaffer et al., (2008), the main features of which are summarized below.

530

Each model ocean layer of 100 m thickness is assigned a sediment segment composed of calcite, non-calcite mineral (NCM) and reactive organic matter. The segment is a bioturbated layer (BL) that is assumed to be 10 cm thick divided in 7 sublayers with highest resolution near the sediment surface such that sublayer boundaries are 0, 0.2, 0.5, 1, 1.8, 3.2, 6 and 10 cm. Sediment segment areas are determined by model topography from each ocean zone. POM and PCal rain fluxes, and ocean values of T, S, DIC, ALK, O₂ and PO₄ are taken from respective layers from ocean and ocean biogeochemistry modules. NCM fluxes (F_{NCM}) are parameterized as

535

$$F_{NCM} = NCF [1 + CAF \exp(-z/\lambda_{slope})] \quad (33)$$

540

where NCF is the open ocean non-calcite flux, CAF is an amplification factor at the coast (i.e. at $z = 0$) and λ_{slope} is the e-folding, water depth scale representing the effect of distance from the coast associated with continental slope topography. For simplicity, we apply the same value of NCF, CAF and λ_{slope} for all ocean sectors taken to be 0.3 g cm⁻² kyr⁻¹, 20 and 200 m respectively.

545

The sediment module is designed to address calcium carbonate (CaCO₃) dissolution and (oxic and anoxic) organic matter remineralization by calculating concentrations of reactive organic carbon (OrgC), pore-water O₂ and pore-water CO₃²⁻ for each sediment sublayer. To accomplish this, a key property is the sediment porosity (ϕ_S , not to be confused with the latitude symbol), which is parameterized as a function of calcite dry weight fraction, $(CaCO_3)_{dwf}$.

550

$$\phi_S(\zeta) = \phi_{S,min} + (1 - \phi_{S,min}) \exp(-\zeta/\alpha) \quad (34)$$

where ζ is the sediment vertical coordinate, $\phi_{(S,min)} = 1 - (0.483 + 0.45(CaCO_3)_{dwf})$ and $\alpha = 0.25(CaCO_3)_{dwf} + 3(1 - (CaCO_3)_{dwf})$ following Archer (1996). From sediment porosity, the sediment formation factor ($F_S = \phi_S^{-3}$) is determined in order to calculate bulk sediment diffusion coefficients of pore-water solutes.

555

To account for the role of benthic fauna, the bioturbation rate (D_b) is parameterized to depend on organic carbon rain rates but also to consider attenuation associated with very low dissolved oxygen concentrations. This is formulated as

$$D_b = D_b^0 \left(\frac{F_{OrgC}}{F_{OrgC}^0} \right)^{0.85} \frac{[O_{2,ocean}]}{[O_{2,ocean}] + O_{2,low}} \quad (35)$$

560

where $[O_{2,ocean}]$ is the ocean O_2 concentration at the sediment surface and $O_{2,low}$ is taken to be 20 mmol m^{-3} . Moreover, D_b^0 and F_{OrgC}^0 are the bioturbation rate scale and the organic carbon rain rate scale whose values are $1.38 \times 10^{-8} \text{ cm}^2 \text{ s}^{-1}$ and $1 \times 10^{-12} \text{ mol cm}^{-2} \text{ s}^{-1}$, based in part on Archer et al., (2002).

565 Oxygen remineralization rates in the BL are taken to scale as bioturbation rates (and thereby as organic carbon rain rates; Archer et al., (2002)), such as $\lambda_{ox} = \lambda_{ox}^0 D_b / D_b^0$. Anoxic remineralization rates in the BL are slower than oxic rates and will depend upon the specific remineralization reactions involved (e.g. denitrification faster than sulphate reduction). More organic rain would be associated with a more anoxic BL and a shift toward sulphate reduction. Therefore, we take $\lambda_{anox} = \beta \lambda_{ox}$ whereby β is taken to decrease for increasing organic carbon rain rate such that $\beta = \beta_0 (F_{OrgC} / F_{OrgC}^0)^\gamma$. As described in
 570 Shaffer et al., (2008), values for λ_{ox}^0 , β_0 and γ were constrained by organic carbon burial observations to be $1 \times 10^{-9} \text{ s}^{-1}$, 0.1 and -0.3 respectively.

Governing equations for OrgC, pore-water O_2 and CO_3^{2-} and $CaCO_3$ are second-order, non-linear coupled differential equations which are solved for each sediment segment using a semi-analytical iterative approach (steady state) or time stepping approach
 575 (time dependent) by imposing boundary conditions at the top and bottom of the BL and matching conditions at the sublayer boundaries. For simplicity we also apply the same calculated sediment remineralization rates to organic phosphorus raining on the sediment surface. Using a mass balance approach, sedimentation velocity is determined and used to calculate burial rates of phosphorus, organic carbon and carbonate carbon down out of the base of the BL. In this way the model produces synthetic sediment cores at every depth for each of the model ocean basins. Furthermore, model solutions provide fluxes
 580 between ocean and sediment layers of PO_4 , O_2 , DIC and ALK based on concentrations of these tracers in each respective adjacent ocean layer and sediment pore-water concentrations from organic matter remineralization and calcium carbonate dissolution. A detailed description of the sediment module is given in Appendix A of Shaffer et al., (2008).

2.6 Land Biosphere

585 [Eichinger et al., \(2017\) defined three different, dynamically varying vegetation zones to extend and improve the original DCESS I, one-zone land biosphere module. The vegetation zones - a grassland/desert zone bordered equatorward and poleward by tropical forest and extra tropical forest zones, respectively - were formulated by emulating the behavior of a complex land](#)

biosphere model (Gerber et al., 2004). With this approach, latitudinal boundaries of the zones could be defined as functions of global mean temperature alone encompassing implicit dependency on precipitation. In this way, the very different carbon distributions between, say, above-ground biomass and soil for each zone and the responses of these carbon reservoirs to changing climate and atmospheric CO₂ could be addressed. For example, while most of the carbon in tropical forests is found in above-ground, by far most of the carbon in extratropical forests is in the soil (Chapin III et al., 2011). With this new three-zone module, the size and timing of carbon exchanges between atmosphere and land were represented much more realistically in cooling and warming experiments than with the original, one-zone module (Eichinger et al., 2017). Furthermore, our three-zone approach allows for changing biosphere modulation of radiative forcing since albedo is higher for grasslands/deserts than for forests (see albedo formulations in section 2.1 above).

Here we use the same approach but now expanded to two hemispheres. The vegetation zones are ~~We consider a land biosphere model with three different, dynamically varying vegetation zones per hemisphere~~ (tropical forest, TF, grasslands, savanna and deserts, GSD, and extratropical forest, ET) that include carbon reservoirs for leaves (M_G), wood (M_W), litter (M_D) and soil (M_S) (Shaffer et al., 2008; Eichinger et al., 2017). ~~The meridional limits for~~ Latitudinal boundaries of each vegetation zone (ϕ_{TF-GSD} for TF-GSD boundary and ϕ_{GSD-EF} for GSD-EF boundary) are obtained using a fifth-order polynomial dependent on the deviation of hemispheric annual mean atmospheric temperature from the respective pre-industrial temperature such that $\phi(\delta T_a) = c_1 \delta T_a^5 + c_2 \delta T_a^4 + c_3 \delta T_a^3 + c_4 \delta T_a^2 + c_5 \delta T_a^1 + c_6$, with $\delta T_a = \bar{T}_a - \bar{T}_{a,PI}$. Polynomial coefficients are obtained by fitting data from Gerber et al., (2004) for Northern and Southern Hemisphere separately (Table 3). Poleward ET limits are the annual mean snowline for Northern Hemisphere and the fixed position at 55° S for Southern Hemisphere (there is no land south of 55° S, see Fig. 1a). We note that the above approach is only strictly valid for -10 °C < δT_a < 10 °C, the range considered in the original experiments of Gerber et al. (2004). For warming this corresponds to annual mean temperatures less than about 25 °C and corresponding atmospheric pCO₂ levels less than about 1000-1500 ppm. For more extreme warming situations in the distant past or future a re-evaluation of our land biosphere module would be necessary for use in model simulations.

Table 3: Coefficients for vegetation meridional limits, $\phi(\delta T_a)$, for Northern and Southern Hemispheres and global pre-industrial distribution of carbon storage and net primary production for all vegetation zones considered in the model.

		$c_1 (\times 10^{-5})$	$c_2 (\times 10^{-4})$	$c_3 (\times 10^{-3})$	$c_4 (\times 10^{-2})$	$c_5 (\times 10^{-1})$	c_6
ϕ_{TF-GSD}	NH	-1.803	-5.809	-5.168	4.970	10.920	11.280
	SH	8.413	7.339	-8.333	-8.764	-6.965	-19.660
ϕ_{GSD-EF}	NH	1.152	-1.785	-4.557	4.156	10.170	37.770
	SH	-0.651	2.131	1.857	-4.100	-6.615	-35.630
		Tropical forest		Grassland, Savanna and Deserts		Extratropical forests	

Leaves (Gt C)	30	50	20
Wood (Gt C)	270	50	180
Litter (Gt C)	16	40	64
Soil (Gt C)	200	500	800
NPP (Gt C yr ⁻¹)	25	20	15

615 Net primary production on land (NPP) takes up atmospheric CO₂ and is forced by seasonally varying solar radiation. For each vegetation type zone, land NPP is calculated according to

$$\text{NPP} = \text{NPP}_{\text{PI}} A_f f(I) \left[1 + f_{\text{CO}_2} \ln \left(\frac{p_{\text{CO}_2}}{p_{\text{CO}_2, \text{PI}}} \right) \right] \quad (36)$$

620 where NPP_{PI} is the pre-industrial net primary production (see Table 3), A_f is an area factor accounting for vegetation size change with respect to pre-industrial size, f_{CO_2} is the CO₂ fertilization factor equal to 0.37, a suitable value for the terrestrial biosphere (Eby et al., 2013; Zickfeld et al., 2013), p_{CO_2} is the model-calculated partial pressure of atmospheric carbon dioxide, $f(I) = f_0 + a \exp \left(-((I - b)/c)^2 \right)$ is a function fitted from model results (Hazarika et al., 2005) with coefficients a , b and c chosen to represent the seasonal cycle of solar radiation according to the specific orbital forcing parameters and f_0 is chosen
625 so that the annual mean value of $f(I)$ for each zone equal to one. [Note that each vegetation type has its own function \$f\(I\)\$ with its respective parameters.](#) With this formulation NPP responds in a seasonal cycle according to $f(I)$ and to atmospheric carbon dioxide on longer time scales.

630 With the descriptions above, [and the assumptions that NPP is distributed between leaves and wood in the fixed ratio 35:25, all leaf loss goes to litter, wood loss is divided between litter and soil in the fixed ratio 20:5, and litter loss is divided between the atmosphere \(as CO₂\) and the soil in the fixed ratio 45:10 \(Siegenthaler and Oeschger, 1987; Shaffer et al., 2008\).](#) the conservation equations for the land biosphere reservoirs of ¹²C for leaves (M_G), wood (M_W), litter (M_D) and soil (M_S) for each of the six vegetation zones are:

$$635 \quad \frac{dM_G}{dt} = \frac{35}{60} \text{NPP} - \frac{35}{60} \text{NPP}_{\text{PI}} \frac{M_G}{M_{G, \text{PI}}} \quad (37)$$

$$\frac{dM_W}{dt} = \frac{25}{60} \text{NPP} - \frac{25}{60} \text{NPP}_{\text{PI}} \frac{M_W}{M_{W, \text{PI}}} \quad (38)$$

$$\frac{dM_D}{dt} = \frac{35}{60} \text{NPP}_{\text{PI}} \frac{M_G}{M_{G, \text{PI}}} + \frac{20}{60} \text{NPP}_{\text{PI}} \frac{M_W}{M_{W, \text{PI}}} - \frac{55}{60} \text{NPP}_{\text{PI}} \frac{M_D}{M_{D, \text{PI}}} \lambda_Q \quad (39)$$

$$\frac{dM_S}{dt} = \frac{5}{60} \text{NPP}_{\text{PI}} \frac{M_W}{M_{W, \text{PI}}} + \frac{10}{60} \text{NPP}_{\text{PI}} \frac{M_D}{M_{D, \text{PI}}} \lambda_Q - \frac{15}{60} \text{NPP}_{\text{PI}} \frac{M_S}{M_{S, \text{PI}}} \lambda_Q \quad (40)$$

640 where $M_{G|W|D|S,PI}$ are the pre-industrial reservoir sizes (Table 3) and $\lambda_Q = Q_{10}^{(T_a - T_{a,PI})/10}$ with $Q_{10} = 2$ as above (Sect. 2.4).
 Atmosphere-land biosphere carbon dioxide flux from each vegetation zone is:

$$F_{CO_2} = -NPP + NPP_{PI} \left[\frac{45}{60} \frac{M_D}{M_{D,PI}} + \frac{15}{60} \frac{M_S}{M_{S,PI}} \right] \lambda_Q \quad (41)$$

645 For isotopes ^{13}C and ^{14}C , Eqs. (37)-(41) are extended considering fractionation factors for photosynthesis and, for ^{14}C , radioactive decay as described in Shaffer et al., (2008).

Finally, land biosphere methane and nitrous oxide productions (F_{CH_4} and F_{N_2O} respectively) take place in soil and are proportional to the reservoir size and temperature dependent according to λ_Q where again $Q_{10} = 2$ for both. In addition, we
 650 assume methane emissions only from wet areas (zones TF and ET). Thus, for each vegetation zone fluxes of these two greenhouse gases are given as

$$F_{CH_4|N_2O} = \tau_{CH_4|N_2O,PI} (pCH_4|pN_2O)_{PI} \frac{M_S}{M_{S,PI}} \lambda_Q \quad (42)$$

655 where $\tau_{CH_4|N_2O,PI} = \tau_{CH_4|N_2O,PI}$ and $\tau_{CH_4|N_2O,PI}$ is the pre-industrial atmospheric life time of CH_4 and N_2O (9.5 and 150 years, respectively). $^{13,14}CH_4$ fluxes are $^{13,14}\alpha_M F_{CH_4,PI} (M_S^{13,14}/M_{S,PI}) \lambda_Q$ with $^{13}\alpha_M = 0.97$ the fractionation factor for CH_4 production. As above, we assume the $^{14}C/^{12}C$ fractionation to be twice as strong as for $^{13}C/^{12}C$ (see Sect. 2.2). Fluxes to specific atmospheric boxes are equal to the biosphere fluxes within the latitudinal boundaries of the boxes.

2.7 Lithosphere module (rock weathering, volcanism, and river input)

660 We follow the same approach as in the DCESS I model (Sect. 2.7 in Shaffer et al., (2008)) by considering river inputs of phosphorus and carbon species, climate-dependent carbonate and silicate weathering rates and lithosphere outgassing. However here we extend this approach to consider distributions of continents, river mouths (Dai and Trenberth, 2002) and volcanoes (NCEI Volcano Location Database). In the following, we restrict ourselves to presenting the main features of this module (see Shaffer et al., (2008) again for more details).

665

Weathering rates of rocks containing phosphorus (W_P), as well as carbonate and silicate weathering rates (W_{Sil} and W_{Cal}) are taken to depend on deviation of mean atmospheric temperature from its pre-industrial value in the form,

$$W_{P|Sil|Cal} = \lambda_Q W_{P|Sil|Cal,PI} = Q_{10}^{(T_a - T_{a,PI})/10} W_{P|Sil|Cal,PI} \quad (43)$$

670

where $W_{P|Sil|Cal,PI}$ are the pre-industrial weathering rates for phosphorus, silicate and carbonate and $Q_{10} = 2$ as for the other model components. [Vegetation affects weathering rates by modifying surface pH through production of CO₂ or organic acids or by altering the physical properties of soil such as erosion of exposed mineral areas and by water cycling content \(Drever, 1994; Berner, 1995\). Our model does not explicitly include these and other factors like a direct dependency of atmospheric pCO₂ levels \(Krissansen-Totton and Catling, 2017\). Such factors would add extra tuneable complexity and be beyond the scope and balance of our simplified model.](#) Silicate weathering consumes 2 moles of atmospheric CO₂ per mole of silicate mineral weathered while the carbonate weathering consumes only 1 mole of atmospheric CO₂ per mole of carbonate mineral weathered. Both types of weathering supply bicarbonate ion to ocean surface layers, modifying dissolved inorganic carbon and alkalinity concentrations. The phosphorus supply is equal to W_P . Therefore, expressions for total river inputs for PO₄, DIC and ALK tracers are

$$R_P = W_P = \lambda_Q W_{P,PI} \quad (44)$$

$$R_{DIC} = 2(W_{Sil} + W_{Cal}) = 2\lambda_Q (W_{Sil,PI} + W_{Cal,PI}) \quad (45)$$

$$R_{ALK} = 2(W_{Sil} + W_{Cal}) - r_{ALKP} W_P = \lambda_Q [2(W_{Sil,PI} + W_{Cal,PI}) - r_{ALKP} W_{P,PI}] \quad (46)$$

Values of $W_{P,PI}$, $W_{Cal,PI}$ and $W_{Sil,PI}$ are obtained from the assumed pre-industrial steady state equal to the global ocean burial rate of phosphate (B_{OrgP}) and carbonate (B_{Cal}), respectively, and the assumption that $W_{Sil,PI}$ can be taken to be a fixed ratio of carbonate weathering: $W_{Sil,PI} = \gamma_{Sil} W_{Cal,PI}$ with $\gamma_{Sil} = 0.85$. These total river inputs are distributed among the twelve ocean sectors according to river mouth distributions mentioned above. For example, this leads to no river input to the model Southern Ocean sector.

Sources of carbon to the atmosphere are weathering of rocks containing old organic carbon (W_{OrgC}) and lithosphere outgassing (Vol). As above, $W_{OrgC} = \lambda_Q W_{OrgC,PI}$ and Vol may either be taken as a constant and equal to its pre-industrial value (Vol_{PI}) or may be prescribed as an external forcing of the Earth System. Given the above together with assigned or calculated ¹³C content for the different model inputs and outputs (including for example for organic carbon burial), overall steady state conservation equations can be formulated for both ¹²C and ¹³C. These conservation equations can then be used to derive expressions for $W_{OrgC,PI}$ and Vol_{PI} as given in Shaffer et al., (2008) (their Eqs. (41) and (42)). These total atmospheric carbon inputs are distributed among the 6 atmosphere sectors according to the same meridional distribution as carbonate and silicate weathering and volcano distribution mentioned above.

From the above, the pre-industrial steady state equations for phosphorus, carbon-12 and carbon-13 are given by

$$W_P - B_{OrgP} = 0 \quad (47)$$

$$W_{OrgC} + Vol - \frac{\gamma_{Sil}}{1+\gamma_{Sil}} B_{Cal} - B_{OrgC} = 0 \quad (48)$$

$$705 \quad W_{OrgC} \delta^{13}C_{OrgC} + Vol \delta^{13}C_{Vol} - \frac{\gamma_{Sil}}{1+\gamma_{Sil}} B_{Cal} \delta^{13}C_{Cal} - B_{OrgC} \delta^{13}C_{OrgC} = 0 \quad (49)$$

For total oxygen content in the ocean-atmosphere system, the pre-industrial steady state equation is

$$\frac{r_{OCP}}{r_{CP}} [B_{OrgC} - W_{OrgC} - f_{old,OM} Vol] + r_{ONP} (B_P - W_P) = 0 \quad (50)$$

710

where r_{ONP} , r_{OCP} and r_{CP} are as mentioned in Sect. 2.4, and $f_{old,OM}$ is the fraction of Vol originating from old organic matter that results from the above conservation calculations for ^{12}C and ^{13}C .

3 Model solution, calibration and validation

3.1 Pre-industrial steady state solution

715 3.1.1 Solution procedure

The ocean module equations are discretized on a staggered grid type, with tracer values defined at the centre of boxes and velocities and diffusivities determined at box edges. Centered differences are used for derivatives, diffusion and vertical advection, whereas an upwind scheme is used for the coarsely resolved, meridional advection. Prognostic equations for the atmosphere (including snow and sea ice cover), land biosphere, lithosphere and the ocean modules are solved simultaneously using a fourth order Runge-Kutta algorithm with a two week time step. Prognostic equations for the ocean sediment are solved by simple time stepping with a one year time step. The complete, coupled model is written in Fortran language and runs at a speed of about 10 kyr of simulation per 30 minutes of computer time on a high-end personal computer.

720

3.1.2 Calibration procedure

For model calibration we used a similar approach as in Shaffer et al., (2008) but here consisting of six steps. For the first step we considered the atmospheric module with atmospheric pCO_2 , pCH_4 and pN_2O set to their pre-industrial values (280, 0.72 and $0.27 \mu atm$ respectively) and a slab ocean for air-sea heat exchange. We adjusted the free parameters listed in Table 1 to give a steady state global annual mean atmospheric temperature of $15 \text{ }^\circ C$ and a climate sensitivity of $3 \text{ }^\circ C$ per doubling CO_2 ; as indicated by several lines of evidence and model estimates (Meehl et al., 2020; Zelinka et al., 2020; IPCC, 2021), a poleward transport of heat and water vapor consistent with observations and annual mean latitudes and seasonal cycle amplitudes of sea ice lines in accordance with observed data. In the second step we couple the physics part of the ocean module to the physics part of atmosphere module and physical ocean free parameters (Table 2) were adjusted in order to get the best fit to observed mass and heat transport as well as temperature and salinity distributions. In the third step we couple the land biosphere to the

730

previously calibrated “physics model” version, and we adjust free parameters of the set of functions $f(I)$ in Eq. (36) to give observed annual mean NPP values for each vegetation zone as well as their annual cycle amplitudes ~~carbon reservoirs for each~~
735 ~~vegetation zones as well as an atmospheric carbon dioxide seasonal cycle amplitude~~ from consistent with observations. This largely sets the modelled annual cycle of atmospheric carbon dioxide of each atmospheric box. In the fourth step we incorporate ocean biogeochemical tracers PO_4 , $\text{DI}^{12,13,14}\text{C}$, ALK and O_2 to the model version of step three. For calibration we start with homogeneous vertical values of 2.17×10^{-3} , 2.32 and 2.43 mol m^{-3} for PO_4 , DIC and ALK respectively (in the following for simplicity DIC will be used to mean DI^{12}C). Furthermore, in this step we use a fixed atmospheric O_2 equal to 0.2095 #atm , an
740 atmospheric $\delta^{13}\text{C}$ equal to -6.5 ‰ and an atmospheric ^{14}C production chosen to keep $\Delta^{14}\text{C}_{\text{atm}} \sim 0 \text{ ‰}$ (Sarmiento and Gruber, 2006; Shaffer et al., 2008). At this stage we assume that all biogenic particles falling to the ocean bottom remineralize completely there. We then made initial guesses for the values of the biogeochemical free parameters of the ocean module listed in Table 2. These choices were partially based on DCESS I model (Shaffer et al., 2008). The atmosphere-land biosphere-ocean
745 model was spun up with uniform atmosphere and ocean tracer distributions to a steady state after about 10,000 model years and results were compared with atmosphere, land biosphere and ocean data. Then all parameter values were adjusted by trial and error in order to get steady state solutions that better satisfied requirements described in the previous steps as well as observed global annual mean distributions of T, S, PO_4 , DIC, ALK and dissolved O_2 .

In the fifth calibration step we coupled the sediment module to the step four calibrated model. For conservation, the total burial
750 rate of a tracer was added to the ocean surface layer of each ~~box~~ sector under consideration of present-day rivers distribution. After solving this new closed-system for a new, pre-industrial steady state, we adjusted all free parameters in order to obtain steady state solutions that better satisfied the data-based constrains of the previous calibration steps. In the sixth and final calibration step, we coupled the lithosphere module to the step five calibrated model whereby river inputs are equated with tracer burial fluxes from the fifth calibration step (tracer burial fluxes are now leaving the system). In addition, weathering
755 rates and lithosphere outgassing were calculated from the tracer burial fluxes and the assumption of the pre-industrial steady state for phosphorus and $^{12,13}\text{C}$ (Sect. 2.7). A last slight trial and error adjustment is made to satisfy the data-based requirements from all previous calibration steps. For this final calibration, we made a long run until steady state is achieved such that all model components vary less than a 0.001 % during 1000 model years. Resulting global annual means for atmospheric temperature and CO_2 , CH_4 and N_2O atmospheric partial pressures are $15.12 \text{ }^\circ\text{C}$, 279.96, 0.72 and $0.27 \text{ } \mu\text{atm}$ respectively. For
760 atmospheric isotopes, this final calibration gives a global mean atmospheric $\delta^{13}\text{C}$ of -6.53 ‰ and a ^{14}C atmospheric production of $1.66 \times 10^4 \text{ atom m}^{-2} \text{ s}^{-1}$, respectively. Global mean ocean values are 2.16 and $145.07 \text{ mmol m}^{-3}$ for PO_4 and dissolved O_2 and 2.30 and 2.41 mol m^{-3} for DIC and ALK respectively.


3.1.3 Atmosphere tracers and transport results

The steady-state, pre-industrial solution gives the annual mean atmospheric temperatures of $15.4 \text{ }^\circ\text{C}$ for Northern Hemisphere
765 and $14.8 \text{ }^\circ\text{C}$ Southern Hemisphere, whereas annual mean sea ice extensions for Northern (Arctic and North Pacific sectors)

and Southern Hemispheres are 66.2° N, 64.9° N and 63.2° S respectively. While Southern Ocean model result is close to the observed value (~64° S), our Arctic Ocean sea ice line position is about 7° too far south with respect to observational estimates (Fetterer et al., 2017). This could be attributable to ocean and sea ice dynamics that are not captured in our simplified, zonal-averaged model. The Northern Hemisphere snow line is found at 58.8° N. Annual mean poleward atmospheric heat transport across 35° N/S and 55° N/S are 4.4/4.8 and 3.1/3.4 PW, and poleward water vapor transports in the atmosphere at the same latitudes are 0.76/0.82 and 0.43/0.45 Sv respectively. These model results agree well observational estimates (Trenberth and Caron, 2001). The seasonal cycle of sea ice is relatively well represented into the model, with a maximum (minimum) extension at the end of winter (summer) and with amplitudes of 4.7 and 9.0 degrees for Arctic and Southern Ocean respectively with differences of 0.4 and 0.9 degrees respective to observed estimates (Fetterer et al., 2017). We believe that model-data disagreements in sea ice are due to the simplicity of parameterization, but also due to anthropogenic signal in the modern-day sea-ice data. Annual mean model difference in atmospheric $p\text{CO}_2$ between Northern and Southern Hemispheres is only 0.7 μatm , a somewhat lower value than observations. However, much of this difference could be explained by the effect on the observations of NH anthropogenic CO_2 emissions. On the other hand, the atmospheric $p\text{CO}_2$ seasonal cycle amplitude of 4.6 and 0.9 ppm for Northern and Southern Hemisphere respectively agrees rather well with monthly-mean observations from Mauna Loa and Cape Grim observatories. This annual cycle amplitude responds strongly to the annual land vegetation dynamics as well as the ocean-land distribution between Northern and Southern Hemispheres. The Southern Ocean plays a role here as well, by dampening Southern Hemisphere cycle amplitude.

3.1.4 Ocean circulation and heat transport results

The steady state, pre-industrial model has the large-scale ocean circulation shown in Fig. 3. There are two meridional cells in the model Atlantic Ocean. The upper, clockwise cell is an Atlantic Meridional Overturning Circulation (AMOC) with a maximum transport of 19 Sv near 1000 m depth in the northern North Atlantic and a maximum depth of about 3500 m. Below 2000 m depth the flow returns southward to the Southern Ocean where upwells below the Drake Passage sill depth (2000 m), largely in response to the surface northward Ekman transport at 55° S. Both the intensity and depth penetration of modelled AMOC are in line with observed data and complex model results (Talley et al., 2003; Hirschi et al., 2020). The lower, counterclockwise cell carries mainly AABW, which fills the whole abyssal Atlantic. The upper, southward branch of this cell returns to the Southern Ocean and upwells there.



795 **Figure 3.** Pre-industrial, steady state model representation of the meridional overturning circulation (S_v) for: **(a)** the Atlantic Ocean, **(b)** the Pacific Ocean and **(c)** the global ocean. Red contours (positive values) represent clockwise circulation and blue contours (negative values) are for counterclockwise circulation as indicated schematically by arrows.

The Pacific Ocean overturning circulation is dominated by a counterclockwise cell carrying AABW which fills the whole deep ocean. The lower branch of the northward flow in this cell upwells in the North Pacific and returns southward in a near surface flow. The upper branch of the northward flow in this cell upwells and return southward to the Southern Ocean below 2000 m without crossing the equator. There is also a second counterclockwise cell carrying Antarctic Intermediate Water (AAIW) north of the equator above 2000 m depth. The upper, southward branch of this cell joins the southward, near surface flow of the deep, counterclockwise cell described above. There is a weak clockwise cell confined above 500 m depth in the North Pacific related in part to the imposed model outflow to the Arctic Ocean through Bering Strait.

805

The mediterranean Arctic Ocean presents an estuarine-type meridional circulation of 1 Sv intensity entering at 700 m depth and flowing out at 300 m depth. This counterclockwise circulation is maintained mainly by the zonal-averaged geometry and freshwater inputs from precipitation and runoff. The Southern Ocean shelf produces 5.4 Sv of overflow water which after entrainment increases its volume to ~16 Sv upon outflow in the deep Southern Ocean. This agrees quite well with observed estimates (Orsi, 2002; Gordon, 2019). Such realistic entrainment and subsequent AABW formation, as achieved in the model


810

with the prescribed, present day Antarctic continental slope (Amante and Eakins, 2009), underlines the usefulness of our simplified, gravity current approach to this problem.

815 Northward ocean heat transport is found throughout the Atlantic Ocean, peaking at 35° N with 0.83 PW and falling
subsequently to 0.75 and 0.19 PW at 55° N and 65° N respectively. The high value at 55° N is related to the maximum ocean
circulation intensity there. The South Atlantic carries 0.37 and 0.34 PW northward at 35° S and 55° S respectively. The North
Pacific Ocean transports heat northwards, with values of 0.41 and 0.29 PW at 35° N and 55° N respectively. The South Pacific
Ocean transports heat southwards, peaking at 35° S with 1.21 PW and 0.84 PW at 55° S. At the equator, the Atlantic and the
820 modelled ocean transports agree well with data-based estimates (Trenberth and Caron, 2001), except at 55° N where the
modelled value is greater than observations.

3.1.5 Ocean tracer and biological production results

Model ocean profiles of T, S and $\delta^{18}\text{O}_w$, are plotted together with observational data in Fig. 4 (all observed data shown in Figs.
4-7 are from the Global Ocean Data Analysis Project version2.2022 (GLODAPv2, Lauvset et al., (2022)), except for those of
825 $\delta^{18}\text{O}_w$ where data from the Global Seawater Oxygen-18 Database – v1.22 (Schmidt, 1999; Bigg and Rohling, 2000) have been
used. In general, there is a good model-data agreement especially at mid-depth and in the deep and abyssal ocean. In the upper
ocean, the model profiles are well within the observed mean range values. In particular for temperature, the model captures
quite well surface-to-deep ocean transitions in most of the model ocean sectors. Modelled temperatures in the Arctic Ocean
below 1000 m are about 3.5 °C warmer than observations, likely reflecting the extensive ocean area covered by sea-ice and/or
830 the lack of local deepwater formation mechanisms like wintertime coastal polynyas in our simplified model. South of 35° S
model temperature fall at the warm end of the observed data.




835 **Figure 4.** Pre-industrial, steady state model ocean vertical profiles (dark blue lines) of conservative temperature (T), absolute salinity (S) and water ^{18}O isotopic excursion ($\delta^{18}\text{O}_w$) compared to observed data (light blue dots) for each ocean model ~~box~~-sector shown in Fig. 1a. T and S are compared with data from the GLODAPv2 database (Lauvset et al., 2022), $\delta^{18}\text{O}_w$ values are compared with data from Global Seawater Oxygen-18 Database – v1.22 (Schmidt, 1999; Bigg and Rohling, 2000, <https://data.giss.nasa.gov/o18data/>).

840 Model salinity is quite well represented and captures prominent features of global ocean, such as the salinity minimum at 650 m and 1050 m depth for msA and msP ~~boxes~~-sectors representing the AAIW, the relatively surface minimum in mnP and hnP sectors associate with North Pacific Intermediate Water (NPIW; Sverdrup et al., 1942), and the vertical salinity structure in the Arctic Ocean with a surface salinity minimum, associated with freshwater inputs from runoff and sea-ice and snow melt and the strong halocline as found in observations (Aagaard et al., 1981). The lack of equatorial upwelling in our simplified model may explain the relatively warm and salty waters in the north and south tropical sectors. For $\delta^{18}\text{O}_w$, the model captures 845 well the vertical distribution as well as the meridional gradient in each model sector, reflecting the global evaporation/precipitation distribution associated with isotopic fractionation of water.

850 Model ocean profiles of PO_4 and O_2 are plotted together with observational data in Fig. 5. The model achieves a good fit to the observed values in almost all ocean model ~~boxes~~-sectors although the model results for O_2 are slightly lower than observed for the mid-depth Atlantic Ocean. Surface phosphate is strongly controlled by ocean new production. Low surface values are

found in all ocean ~~boxes~~sectors, except at the Southern Ocean where the highest surface values are found due to intense upwelling and to the low biological-production efficiency as expected from the low value of L_f in our model. Furthermore, the model captures well differences between Pacific and Atlantic Ocean basins. These differences are largely a consequence of the large-scale ocean circulation as described above. The oldest waters are found at the North Pacific Ocean (model sectors mnP and hnP, see $\Delta^{14}\text{C}$ model profiles in Fig. 6), and consequently, the highest (lowest) values of phosphate (dissolved oxygen) in the ocean interior are found there. High dissolved O_2 values at both surface and abyssal ocean are relatively well represented in the model in response to air-sea gas exchange and AABW ventilation. At the ocean interior, O_2 distributions respond mainly to organic matter remineralization and, for msA and msP sector for example, ventilation from AAIW. Misrepresentation of PO_4 and dissolved O_2 above 1000 m depth of hnP sector is related to the way NPIW is treated in the model. The PO_4 (dissolved O_2) excess (deficit) in Arctic model sector is related to the overestimation in ocean temperature there which influences organic matter remineralization.



865 **Figure 5.** Pre-industrial, steady state model ocean vertical profiles (dark blue lines) of phosphate (PO_4) and dissolved oxygen (O_2) compared to observed data (light blue dots) for each ocean model ~~box~~-sector shown in Fig. 1a. Observed values are from the GLODAPv2 database (Lauvset et al., 2022).

The ocean carbon cycle model tracers DIC and ALK, are plotted together with observed DIC and ALK in Fig. 6. The comparison with observational data shows excellent model-data agreement for both DIC and ALK. Both the vertical structure and the spatial differences among ocean model ~~boxes~~-sectors are well captured by the model. Furthermore, both the depths of the DIC maximum and the slightly shallower depths of the PO₄ maximum (Fig. 5) in well-stratified ocean zones agree well with observations indicating good e-folding length choices used for model remineralization rates. These distributions reflect the interplay between the cycling of organic matter and calcite as well as the air-sea exchange of CO₂.




Figure 6. Pre-industrial, steady state model ocean vertical profiles (dark blue lines) of total dissolved inorganic carbon (DIC) and alkalinity (ALK) compared to observed data (light blue dots) for each ocean model ~~box~~-sector shown in Fig. 1a. Observed values are from the GLODAPv2 database (Lauvset et al., 2022).

Vertical distributions of carbonate ion (CO_3^{2-}), CO_3^{2-} saturation with calcite ($\text{CO}_{3,\text{sat}}^{2-}$) and water CO_2 are plotted in Fig. 7. The crossing point between CO_3^{2-} and $\text{CO}_{3,\text{sat}}^{2-}$ is the calcite saturation depth (CSD) which is found to be around 4000 m depth for the entire Atlantic Ocean and which reaches a minimum depth of around 700 m in the North Pacific sector where the maximum water CO_2 concentration even exceeds that of CO_3^{2-} . These results showing undersaturation throughout most of the deep North Pacific Ocean would imply low values of CaCO_3 in ocean sediments there. This is confirmed by the results for modelled and observed ocean sediments presented below.





Figure 7. Pre-industrial, steady state model ocean vertical profiles of carbonate ion (CO_3^{2-}), carbonate ion saturation with calcite ($\text{CO}_3^{2-}, \text{sat}$), and water CO_2 for each ocean model ~~box~~-sector shown in Fig. 1a.

890 Ocean carbon isotopes ($\delta^{13}\text{C}$ and $\Delta^{14}\text{C}$) are plotted together with observational data in Fig. 8. As for DIC and ALK, there is excellent model-data agreement in nearly all model ocean zones. As light carbon is preferentially taken up during photosynthesis, export of organic matter leaves the euphotic zone enriched in ^{13}C (higher values of $\delta^{13}\text{C}$). Upon remineralization of this organic matter at depth, light carbon is released into the water column leaving the water there depleted in ^{13}C (lower values of $\delta^{13}\text{C}$). As shown by the agreement of model $\delta^{13}\text{C}$ results with ocean as well as atmosphere data, the

895 model deals well with this fractionation process as well as with the temperature-dependent fractionation associated with air-sea exchange of CO_2 . As for PO_4 and O_2 , meridional gradients and Pacific-Atlantic Ocean differences reflect ocean circulation as seen clearly in the $\Delta^{14}\text{C}$ profiles. The oldest waters are found in the North Pacific and exhibit the most negative values of $\delta^{13}\text{C}$ and $\Delta^{14}\text{C}$ in the ocean interior due to a longer time for receiving light carbon from organic matter remineralization and due to a longer time for radioactive decay to act, respectively. Modelled surface ocean $\delta^{13}\text{C}$ values are well within the data-

900 based estimates with a global surface mean model result of 2.15 ‰. As before, the model-data disagreement of $\delta^{13}\text{C}$ in the upper 1000 m depth in hnP sector is related to NPIW formation in the model.



905 **Figure 8.** Pre-industrial, steady state model ocean vertical profiles (dark blue lines) of isotopic excursion of ^{13}C ($\delta^{13}\text{C}$) and isotopic excursion of ^{14}C ($\Delta^{14}\text{C}$) compared to observed data (light blue dots) for each ocean model ~~box-sector~~ shown in Fig. 1a. Observed values are from the GLODAPv2 database (Lauvset et al., 2022). For $\Delta^{14}\text{C}$ only values below 1000 m depth have been plotted as shallower depths are strongly affected by the anthropogenic signal including atomic bomb ^{14}C inputs. Note that bomb ^{14}C also affects deeper levels in the North Atlantic and Arctic ~~boxes-sectors~~ (mnA, hnA and Arc).

910 With all the above, and despite minor model-data disagreements, the model reproduces quite well the main global features of both physical and biogeochemical tracers. Better ocean modelled distributions of S, $\delta^{18}\text{O}_w$, DIC, ALK, $\delta^{13}\text{C}$ and $\Delta^{14}\text{C}$ are key improvements that have been made in comparison to the much-simpler DCESS I model where distributions of those tracers had shortcomings, in particular for S, $\delta^{18}\text{O}_w$ and $\delta^{13}\text{C}$ (Shaffer et al., 2008).

915 Model global ocean new (exported) production (NP) is $5.84 \text{ Gt C yr}^{-1}$, slightly higher than the 5.4 Gt C yr^{-1} of the DCESS I model (Shaffer et al., 2008). This new estimate is at the lower end of the estimates from Dunne et al., (2007) ($9.6 \pm 3.6 \text{ Gt C yr}^{-1}$) but almost matches a modern and possibly more robust estimate of Siegel et al., (2014) of $5.90 \text{ Gt C yr}^{-1}$. By far, the tropical ocean model zones (35° S - 35° N) are the most productive model regions, accounting for more than 66 % of global exported production partitioned into a 53 % and 13 % for the Pacific and Atlantic Ocean respectively. The Arctic Ocean and high north
 920 latitude zone of Pacific Ocean have the lowest values of exported NP with 0.2 % and 1.1 % of the total respectively. The Southern Ocean also has a rather low value with 5 % of total export production. Northern and southern mid-latitude model

sectors (35°-55° N/S) show around 22 % of total NP, with 5 % and 7 % for mnP and mnA and 8 % and 2 % for msP and msA ~~boxes-sectors~~ respectively. This meridional distribution is also observed in the publications cited above. We believe that lack of equatorial upwelling of rich-nutrient waters in the model probably contributed to the relatively low value of ocean NP in our model.

Global biogenic calcite production in the model is 1.25 Gt C yr⁻¹ where more than 65 % is produced in tropical model zones of the Pacific Ocean. Tropical zones of the Atlantic Ocean account only for 16 % of global calcium carbonate production. Extratropical ocean model zones from Northern and Southern hemispheres and combining Pacific and Atlantic ~~boxes-sectors~~ represent only a 15 % of total biogenic calcite production. Table 4 shows values for organic matter and calcite production for each ocean model ~~boxsector~~.

Table 4: Pre-industrial organic matter and biogenic calcite ocean production estimates. All values are given in Gt C yr⁻¹.

Ocean limits	Atlantic Ocean		Pacific Ocean	
	Organic matter	Biogenic Calcite	Organic matter	Biogenic Calcite
65° N-55° N	0.20	0.01	0.07	3.52×10 ⁻³
55° N-35° N	0.38	0.06	0.30	0.04
35° N-0°	0.43	0.12	1.34	0.35
0°-35° S	0.35	0.09	1.80	0.47
35° S-55° S	0.15	0.02	0.47	0.07
	Southern Ocean (55° S-70° S)		Arctic Ocean (90° N-65° N)	
	0.31	0.01	0.02	5.43×10 ⁻⁴

Our estimates are well into the observed range of 0.5-1.6 Gt C yr⁻¹ of Berelson et al., (2007) but slightly higher than more recent estimates of 0.91±0.14 Gt C yr⁻¹ (Sulpis et al., 2021). Overall, our model result of new (exported) production is well into the range of CMIP6 models (Planchat et al., 2023), but our biogenic calcite production is at the upper end of those model estimates. However, we note that CMIP6 ensemble results tend to globally underestimate the exported inorganic carbon compared to observational estimates (Planchat et al., 2023).

The calcite carbon to organic carbon flux ratio is similar for the Pacific/Atlantic Oceans, with ratios of 0.6/0.5, 0.9/0.7 and 1.2/1.1 in tropical model sectors at 1000, 2000 and 3000 m depth respectively, and ratios of 0.2, 0.3 and 0.5 for northern latitudes sectors of those basins at these depths. These ratios are somewhat lower than those estimated from sediment trap data (Berelson et al., 2007). The longer e-folding scale ξ_{Cal} than ξ_C and the temperature effect on organic matter remineralization and on the CaCO₃ rain ratio (see Sect. 2.4) help explain these results.

3.1.6 Sediment results

In the model pre-industrial steady state, 0.11 Gt C yr⁻¹ of organic carbon is buried in ocean sediments. This represents 12 % of total organic carbon falling on sediment surface; the remaining 88 % is remineralized to DIC that returns to the water column. [This modelled value is well within the range of data-based estimations \(Bernier, 1982; Hedges and Keil, 1995; Cartapanis et al., 2018\).](#) However, the higher ends of some of these estimates may include contributions from estuaries and delta environments that our model does not consider. Other modelled results are in the range of 0.02 – 0.09 Gt C yr⁻¹ (Bernier 1991; [Munhoven, 2007; Willeit et al., 2023](#)). More than 90 % of ~~this burial~~ [our modelled burial](#) takes place at water depths shallower than 1000 m [such that 0.011 Gt C yr⁻¹ is buried below this depth, in agreement with deep sea burial estimates \(Hayes et al., 2021\)](#). The highest modelled burial rates for organic carbon per ocean sector are found in the tropical Pacific Ocean and the North Atlantic Ocean (35° N-65° N) sectors with over 40 % and 27 % of the total organic carbon burial, respectively. The global inventory of bioturbated layer (BL) organic carbon is 122 Gt C, somewhat higher than found in the pre-industrial, DCESS I model simulation (92 Gt C).

For calcite, the global annual mean burial rate is 0.21 Gt C yr⁻¹ where more than 57 % takes place in the tropical Pacific sectors and 23 % is buried at the combined north-tropical and mid-latitude Atlantic Ocean sectors. [Our global result compares well with data-based estimations and model results in the range of 0.13 – 0.45 Gt C yr⁻¹ \(Milliman and Droxler, 1995; Cartapanis et al., 2018; Hayes et al., 2021; Willeit et al., 2023\).](#) Of the total annual mean burial rate of CaCO₃ in the Pacific and Atlantic Ocean (0.14 and 0.07 Gt C yr⁻¹ respectively) 49 % and 45 % of calcite is buried at depths greater than 1000 m in these basins, respectively. [Other estimates also show such a nearly half-to-half, shelf-slope vs deep ocean division \(Milliman, 1993\).](#) Modelled global ocean mean calcite dry-weight fraction (CaCO₃)_{dwf} is 0.349, whereas Pacific and Atlantic oceans present mean values of (CaCO₃)_{dwf} of 0.353 and 0.440 respectively, where highest values are found in south tropical Pacific and north Atlantic sectors with values of 0.459 and 0.553 respectively. Model global inventory of bioturbated layer calcite carbon is 767 Gt C, somewhat lower than the DCESS I pre-industrial estimate (1010 Gt C) but in good agreement with the 800 Gt C data-based estimate from Archer (1996). Of this, 503 Gt C is found in the Pacific Ocean BL and 249 Gt C in the Atlantic Ocean one, equivalent to 66 % and 32 % of the total, respectively. The results above highlight how the model captures ocean carbon cycle differences between Pacific and Atlantic Ocean.

Figure 9 shows the dry-weight fractions of organic matter (OM)_{dwf} and calcite (CaCO₃)_{dwf} as well as the sediment velocity out of the bottom of the BL (w_{sed}) for each ocean model ~~box~~ [sector](#). In general, relatively high (OM)_{dwf}, relatively low (CaCO₃)_{dwf} and rapid sedimentation rates are found at shallow depths. This can be explained mainly by high prescribed flux of non-calcite minerals at such depths that 1. fill the BL mainly with such minerals and 2. rapidly flush the BL thereby promoting high (OM)_{dwf} burial since relatively little organic matter is remineralized in the BL on such short time scales. At intermediate depths above the model's CSD, (CaCO₃)_{dwf} is higher, (OM)_{dwf} is lower and sedimentation rates are more moderate. This results from

enhanced CaCO_3 accumulation in sediments bounded by ocean layers supersaturated with carbonate ion (see Fig. 7), but also, from much lower non-calcite mineral and organic matter rain rates leading to slower BL flushing that allows more complete organic matter remineralization. At those intermediate depths, the modelled effect of dissolved oxygen on organic matter remineralization can be seen in the North Pacific model sectors where this remineralization is slowed down considerably as ocean layers there become nearly suboxic/anoxic. This also leads to a local increase in sediment rates. In the deep ocean below the model's CSD, rapid decreases in CaCO_3 content, very low organic matter contents and slower sedimentation rates are the result of calcite dissolution combined with low constant non-calcite rain rates and still lower organic matter rain rates. The Arctic Ocean model sediment is almost completely composed of non-calcite mineral as the sediment flux of biogenic material falling on sediment surface is very small due to the extremely limited surface new production. In the Southern Ocean, model sediment is also composed mainly of non-calcite-mineral but contains larger fractions of $(\text{OM})_{dwf}$ and $(\text{CaCO}_3)_{dwf}$ than in the Arctic Ocean. The water depth of the calcite compensation depth (CCD), defined operationally where $(\text{CaCO}_3)_{dwf}$ equals 0.1, is found around 4800 m for both tropical Pacific and tropical Atlantic Ocean sectors. In contrast, there are major differences between North Atlantic and North Pacific sectors with CCD depths around 4400 m and 2100 m depth, respectively.




Figure 9. Pre-industrial, steady state model sediment vertical profiles of dry-weight fractions of organic matter, $(\text{OM})_{dwf}$, and calcite, $(\text{CaCO}_3)_{dwf}$, as well as sedimentation velocity (W_{sed}), all at the base of the bioturbated layer of each ocean model ~~box~~ sector shown in Fig. 1a.

3.1.7 Lithosphere results

1000 Table 5 lists model weathering and burial rates for the pre-industrial steady state. From weathering-burial steady-state balances
outlined in Sect. 2.7 and global mean values of ^{13}C isotope excursions for organic carbon ($\delta^{13}\text{OrgC} = -22.06 \text{ ‰}$) and carbonate
($\delta^{13}\text{Carb} = 2.22 \text{ ‰}$) obtained from the fifth calibration step, modelled outgassing from volcanoes is found to be $0.140 \text{ Gt C yr}^{-1}$,
1005 in good agreement with to modern-day data-based estimates (Masson-Delmotte et al., 2021; IPCC, 2021). Results for
carbonate and silicate weathering rates are similar to those found with the DCESS I model, 0.109 and $0.092 \text{ Gt C yr}^{-1}$,
respectively (Shaffer et al., 2008). Our carbonate weathering rate is well into the range of other observational and model
estimates, $0.088 - 0.241 \text{ Gt C yr}^{-1}$, but our silicate weathering rates are somewhat lower than other corresponding estimates,
 $0.122 - 0.236 \text{ Gt C yr}^{-1}$ (Gaillardet et al., 1999; Ludwig et al., 1999; Munhoven, 2002; Hartmann et al., 2009; Lenton et al.,
2018; Willeit et al., 2023). Phosphorus weathering rates are higher than those found in the DCESS I model, $1.663 \times 10^3 \text{ mol P}$
 s^{-1} but are in better agreement with observational estimates, $2.019 - 3.071 \times 10^3 \text{ mol P s}^{-1}$ (Filippelli, 2002; Paytan and
1010 Mclaughlin, 2007). Organic carbon weathering rates agree well with recent estimations of $0.068 \text{ Gt C yr}^{-1}$ (Zondervan et al.,
2023). Based on these results, although simple, our continental weathering-climate relationship gives reasonable results when
compared against models having more complicated relationships.

Table 5: Weathering, lithosphere outgassing and burial estimates for pre-industrial, steady state balances.

Property	Symbol	Estimated value
Phosphorus weathering	$W_{\text{P,PI}}$	$2.507 \times 10^3 \text{ mol P s}^{-1}$
Organic carbon weathering	$W_{\text{OrgC,PI}}$	$0.064 \text{ Gt C yr}^{-1}$
Carbonate weathering	$W_{\text{Cal,PI}}$	$0.115 \text{ Gt C yr}^{-1}$
Silicate weathering	$W_{\text{Sil,PI}}$	$0.098 \text{ Gt C yr}^{-1}$
Lithosphere outgassing	V_{olPI}	$0.140 \text{ Gt C yr}^{-1}$
Organic carbon burial	$B_{\text{OrgC,PI}}$	$0.106 \text{ Gt C yr}^{-1}$
Carbonate burial	$B_{\text{Cal,PI}}$	$0.213 \text{ Gt C yr}^{-1}$

1015 The modelled river input of total inorganic carbon is $0.43 \text{ Gt C yr}^{-1}$, a value close to that found in the DCESS I model (0.40 Gt
 C yr^{-1}) and close to the $0.41 \text{ Gt C yr}^{-1}$ from data-based estimates (Li et al., 2017). More than 51 % of this river input flux, is
delivered to the Atlantic Ocean, while 40 % drains to the Pacific Ocean and the rest to the Arctic Ocean. The Southern Ocean
does not receive inorganic carbon from rivers. With the results above, modelled steady state pre-industrial global ocean

outgassing of CO₂ is 0.11 Gt C yr⁻¹ which is balanced by net uptake of atmospheric CO₂ that is expressed as $2W_{Sil,PI} + W_{Cal,PI} - (Vol_{PI} + W_{OrgC,PI})$.

3.2 Testing the model

We carry out two simple experiments to test model performance. In the first experiment (Exp 1) we force the model with an imposed, time-dependent function of northward Ekman transport out of the Southern Ocean at 55° S. We considered two cases for this experiment, doubling (Exp 1a) and halving (Exp 1b) this transport value on a timescale of 1500 years. With this, we try to emulate the role of Southern Ocean westerly winds on large-scale ocean circulation as it has been hypothesized that intensity and position shifts of these winds may drive global-scale climate changes on glacial-interglacial timescales (Gray et al., 2023; Lee et al., 2011; Toggweiler et al., 2006). In the second experiment (Exp 2) we force the model with an imposed, time-dependent function of freshwater input to the Southern Ocean shelf to emulate a melting pulse of the Antarctic ice sheet. We “melt” the equivalent of $2.5 \times 10^{15} \text{ m}^3$ freshwater (equivalent to 5 m sea level rise) over two different timescales: a fast pulse to get a sea level rise of 1 m (Exp 2a) and a slow pulse to get a 0.1 m rise (Exp 2b) after 250 years of model simulation. This is consistent with modelled assessments about Antarctic ice sheet retreats in the present climate context (Ruckert et al., 2017; Deconto and Pollard, 2016). In both experiments we focus on changes on large scale ocean circulation and its impact on ocean tracer distributions as well as on global climate.

3.2.1 Southern Ocean Ekman transport forcing

The forcing and model results are presented in Fig. 10. With the increase of northward Ekman transport (Exp 1a) across 55° S, there is more upwelling in the Southern Ocean sector (Fig. 10b). However, this Southern Hemisphere forcing does not significantly impact AMOC intensity in the North Atlantic Ocean, as shown by the streamline evolution at 55° N at 1000 m depth (Fig. 10e). Rather, this forcing leads to the appearance of a clockwise recirculation cell between 55° S and 40° S confined between 1000 m and 2000 m depth for both Atlantic and Pacific Oceans, as indicated by the vertical flux (defined positive upwards) in msA and msP sectors at 1500 m depth showed in Fig. 10c-d. In Exp 1b, a decrease in northward Ekman transport across 55° S causes a decrease in the Southern Ocean upwelling (Fig. 10b). In this experiment there is an increase in the upward flux above 2000 m depth in msA and a decrease in the downward flux below 1000 m depth in msP, fluxes that subsequently join the upper ocean northward flow and the deep southward flow in the Atlantic and Pacific Oceans respectively. Insensitivity of North Atlantic AMOC intensity as well as intermediate-depth Southern Hemisphere recirculation for increased westerlies forcing has also been reported in other modelled experiments (Jochum and Eden, 2015; Rahmstorf and England, 1997).




Figure 10. Results for 3000-year simulations for two different choices of northward Ekman transport across 55° S. **(a)** Applied Ekman transport evolutions, **(b)-(d)** vertical transport (Vert transport, defined positive upwards) at 1500 m depth for Southern Ocean, msA and msP sectors respectively, **(e)** northward transport across 55° N above 1000 m depth in the Atlantic Ocean, **(f)** AABW transport reaching the abyssal Southern Ocean, **(g)-(h)** Southern Ocean new production evolution and Southern Ocean surface $\delta^{13}\text{C}$ change and **(i)-(j)** changes in atmospheric CO_2 and atmospheric temperature (solid lines are for Southern Hemisphere mean T_a and dashed lines are for T_a for atmospheric southern polar zone, 55° S-90° S).

1050

1055

1060

These ocean interior circulation changes lead to a denser (lighter) water being upwelled in the Southern Ocean for Exp_1a and Exp_2_1b respectively, leading to less (more) entrainment and consequently decreasing (increasing) amounts of AABW reaching the abyssal ocean for these experiments (Fig. 10f). More (less) upwelling together with this Southern Hemisphere recirculation, enhance (reduce) ocean new production in the Southern Ocean, as more (less) nutrients are carried to the surface there (Fig. 10g). These changes in Southern Ocean NP are not reflected in surface ocean $\delta^{13}\text{C}$ values (Fig. 10h, as well as in msA and msP) in this zone of low model new production efficiency. Rather, the decrease (increase) in $\delta^{13}\text{C}$ values in Exp 1a (Exp 1b) is associated with lighter (heavier) signal of ^{13}C from more (less) upwelling of deeper water originating from northern

latitudes of Atlantic and Pacific model zones. Air-sea exchange of CO₂ with the Southern Ocean reflects changes in the upwelling rates there as more or less carbon-rich deeper water is carried to surface layers there. More (less) outgassing there leads to an increase (decrease) in atmospheric CO₂ concentration (Fig. 10i) and a resulting rise (fall) in atmospheric temperature (Fig. 10j). [Such a Southern Hemisphere westerly winds intensity-atmospheric CO₂ relationship has also been reported in another modelled experiment \(d'Orgeville et al., 2010\).](#) Changes in atmospheric CO₂ content is nearly uniform in all atmospheric boxes due to the rapid turnover time of the atmosphere. The temperature change in southern polar atmospheric box is around 2.5 °C and -1.8 °C for Exp 1a and Exp 1b respectively. However, this warming (cooling) signal decreases rapidly farther northward such that mean Southern Hemisphere atmospheric temperature changes are around 0.7 and -0.5 °C for Exp 1a and 1b respectively. [Finally, our Exp1a results are in line with the importance of Southern Hemisphere westerly winds on deglacial atmospheric CO₂ rise as found in models and proxy-data reconstructions \(Mayr et al., 2013; Menviel et al., 2018\).](#)

3.2.2 Antarctic ice sheet melt freshwater forcing


Forcing and model results for this experiment are shown in Fig. 11. The rapid freshwater pulse to the Southern Ocean shelf (Exp 2a) produces a greater and more abrupt decrease in AABW fluxes flowing down off the shelf break ($F_{sl,0}$), as well as in the abyssal ocean, F_{sl} , (Fig. 11b). For Exp 2a, $F_{sl,0}$ and F_{sl} reach minimum values of 3.5 and 11 Sv respectively, i.e. around 30 % less than the pre-industrial values. In Exp 2b, modest declines of 0.7 and 2 Sv are obtained for $F_{sl,0}$ and F_{sl} , representing a decrease of only 13 % compared to modern-day values (Gordon, 2019). [Such effect on AABW production rates due to Antarctic meltwater on Southern Ocean shelf has been observed in similar Antarctic-meltwater experiments using more complex models \(Fogwill et al., 2015; Silvano et al., 2018; Li et al., 2023\).](#) Consequently, the abyssal counterclockwise circulation decreases its intensity by more than 11 and 2 Sv for fast and slow freshwater pulses respectively, as shown by the northward streamline evolution at 4500 m depth crossing 55° S (Fig. 11c). [Such behavior has been reported for numerous model experiments including a multi-model ensemble of CMIP6 model experiments \(Park and Latif, 2019; Mackie et al., 2020; Chen et al., 2023\).](#) This corresponds to a 75 % decrease compared to only a 14 % decrease for the fast and slow freshwater pulses, respectively, underlining the sensitivity of model AABW circulation to the speed and not only the size of the Antarctic Ice Sheet melt water pulse. Such lower AABW ventilation rates lead to the decline in dissolved O₂ content of the abyssal northern sectors of the Pacific and Atlantic Ocean (Fig. 11d-e). [This suggests ageing of the abyssal ocean, also reported in Li et al., \(2023\).](#) A greater AMOC penetration depth, as a consequence of weakened AABW abyssal circulation, may inhibit a further decrease in the abyssal O₂ content of the North Atlantic Ocean as more atmospheric oxygen is carried to those depths.

Opposite behaviors are observed in the abyssal ocean $\Delta^{14}\text{C}$ content for northernmost model sectors of North Atlantic and North Pacific Ocean (hnA and hnP, respectively). The increase in the hnA sector (Fig. 11f) is caused by deeper AMOC penetration depth as AABW cell gets weaker leading to increased transport of near surface ^{14}C to the abyssal ocean. In the hnP sector, the decrease in ^{14}C content stems from longer residence times due to the reduced AABW ventilation there leading to more

1095 radioactive decay. The oxygen-18 signature in the abyssal ocean is quite sensitive to the freshwater forcing to the Southern Ocean shelf as water melt from the Antarctica ice sheet comes with an isotopic signal of -40‰ in the model. This is shown in Fig. 11 h-i, where similar changes are observed at-in the boxes-sectors directly north of the Southern Ocean (msA and msP) at 4500 m depth, with changes around of -0.17 and -0.09 per mil for fast and slow forcing experiments, respectively. This demonstrates the model's ability to capture the $\delta^{18}\text{O}_w$ signal from Antarctic ice-sheet melting, shown to be a better indicator of such melting than ocean salinity changes (Kim and Timmermann, 2024). Despite all these ocean interior changes, Exp 2a and Exp 2b forcings lead only to small changes in atmospheric CO_2 of around 1 ppm or less in both experiments. As a consequence, there is nearly no change in global atmospheric temperature although small, sea-ice dynamics-driven oscillations of 20 year-period are found in southern polar atmospheric temperature (not shown). We stress that this simplified experiment was carried out mainly to test the model performance to isolated fresh water forcing. A comprehensive study of the global climate system response to Antarctic ice sheet melting would also require consideration of climatic effects leading to the melting in the first place, i.e. a coupling of the climate system and the ice sheet.

1100

1105



1110 **Figure 11.** Results for 3000-year simulations for two different choices of Antarctic icesheet melt rate. **(a)** Applied melt rates evolutions, **(b)**
AABW transport flowing off the Southern Ocean shelf ($F_{sl,0}$) and reaching abyssal depths (F_{sl}), **(c)** total northward transport (combined
Atlantic and Pacific) crossing 55° S below 4500 m depth, **(d)-(e)** changes in dissolved O₂ content in the ocean at 4500 m depth for northern
sectors of the Atlantic and the Pacific Ocean, respectively (mnA and mnP), **(f)-(g)** ocean $\Delta^{14}\text{C}$ change at 4500 m depth for the northernmost
sectors of the Atlantic and the Pacific Ocean, respectively (55° N-65° N, hnA and hnP), **(h)-(i)** $\delta^{18}\text{O}_w$ change at 4500 m depth for ocean
~~boxes-sectors~~ directly north of the Southern Ocean (msA and msP) and **(j)** global mean atmospheric $p\text{CO}_2$ change.

1115 **4 Discussion and conclusions**

We have described and tested a new Earth System Model of Intermediate Complexity, DCESS II, that has been calibrated to
the pre-industrial Earth System. For this relatively simple model, efforts were made to limit the number of model free
parameters and to constrain their values as much as possible using observations. At the end of this process, [despite model
limitations](#), we [generally](#) find ~~generally~~ excellent agreement with available modern observations. [When compared with results
from the DCESS I model, a number of improvements with our new model stand out as described below.](#)

Some key features of the physical part of the model like horizontal resolution, a simplified dynamical scheme for large-scale
ocean circulation, stratification-dependent vertical diffusion, a gravity current approach to the formation of Antarctic Bottom
Water as well as a dynamical formulation for sea-ice, have helped us to achieve good agreement to observed data like
atmospheric heat and water vapor transport and ocean temperature and salinity distributions (Lauvset et al., 2022)- [with space
and time resolution not possible using the original DCESS I model](#). These improvements ~~with respect to the original DCESS
I-model~~, together with updates in ocean biogeochemistry such as a light-dependence on surface new (exported) production of
organic matter, dependence on calcite saturation state for the biogenic carbonate production, as well as local temperature
dependence on organic matter remineralization, result in accurate model representations of ocean biogeochemical tracers,
highlighted by an excellent model-data agreement with regard to marine carbon cycle species, [well beyond that achieved with
the DCESS I model in particular with respect to carbon 13 \(Shaffer et al., 2008\)](#). Furthermore, the incorporation of a new land
biosphere module ~~considering with~~ three different [vegetation](#) types ~~of vegetation~~ and a light-atmospheric CO₂-dependence on
net primary production on land, as well as ocean sediment and lithosphere modules applied to multiple ocean and land sectors
make [results of the new model much more amenable to comparison with data than is the case for the DCESS I model. All the
above improvements help make](#) the DCESS II model an excellent while economical tool for studying and gaining short as well
as long-term understanding of the global carbon cycle.

Two experiments were conducted in order to explore and test model performance. In the first experiment (Exp 1) we forced
the model in a way that emulates the role of the Southern Hemisphere westerly winds on the large-scale ocean circulation. In
the second experiment (Exp 2) we introduced (at two different rates) a freshwater “melt” volume equivalent to a global sea
level rise of 5 m into the shelf surrounding Antarctica. In both experiments, model results show important physical and
biogeochemical changes in the ocean driving perturbations in the global carbon cycle. This is reflected by responses in

atmospheric CO₂ as well as in abyssal ocean $\Delta^{14}\text{C}$ and dissolved O₂ content. These experiments serve to demonstrate how the model captures the global role that Southern Hemisphere westerly winds and Antarctic Bottom Water play in the Earth System.

1145 [Our Exp1 supports the role of the Southern Hemisphere westerly winds in modulating past glacial-interglacial, atmospheric carbon dioxide variations, while our Exp2 points toward the impact that future Antarctic ice sheet melting would have on the formation of Antarctic Bottom water and on the deep global ocean in general.](#)

The development and calibration of the DCESS II model as described here has also set the stage for future work. For example, coupling of the model to an Antarctic Ice Sheet (AIS) model will provide a more realistic Earth System simulation when compared to our idealized experiment 2 described above. For this work an extended and improved version of the simple, well tested DAIS model (Shaffer, 2014) will be used. In another example, planned incorporation of methane ~~and~~, nitrogen and sulfur cycles into [the next DCESS II update](#) will significantly improve its ability to deal realistically with deep-time global warming events associated with massive carbon inputs to the Earth System. [Under such conditions, suboxic/anoxic ocean conditions may arise leading, for example, to denitrification and sulfate reduction that must be addressed in such an update.](#)

1155 Groundwork for this step has already been done in as much as these global biogeochemical cycles [and suboxic/anoxic processes have been](#) ~~were~~ incorporated into the DCESS I model (Shaffer et al., 2017). In addition, other potential model applications include the study of different global climate events like Dansgaard-Oeschger oscillations, longer climatic events like the Eocene-Oligocene Transition or even the assessment of deep-time mass extinction events by taken advantage of the relative simplicity of the model for setting proper boundary conditions like orbital forcing and continental distribution among others. [In this regard, when such deep-time model applications with quite different boundary conditions from today are to be made, careful consideration should be made in choosing the few \(only two\) regional tunings of the model.](#)

1160

In conclusion, we have presented, validated and tested a simple and fast new Earth System Model of intermediate complexity intended to be a flexible, comprehensive and economical Earth System modelling platform. Despite its limitations like relatively low horizontal resolution and multiple parameterizations, the model represents quite well most Earth System components, especially the global-scale carbon cycle. Due to its simplicity, it could be easily modified in terms of boundary conditions to address specific past epochs. Thus, we find DCESS II to be a useful tool for studies of past, present and future global change on time scales of years to millions of years while not needing large computational resources.

1165

1170 *Code availability.* The code for the DCESS II model [v1.0 is archived on Zenodo \(https://doi.org/10.5281/zenodo.13738105, Fernández, 2024\)](#) ~~and others DCESS model versions are freely available at~~ [as well as in https://www.dcess.dk/](#).

Data availability. Ocean observed data used in Figs. 4-6 and in Fig. 8 are from the Global Ocean Data Analysis Project version 2.2022 (GLODAP-v2.2022, Lauvset et al., 2022) database and can be accessed at the National Oceanic and Atmospheric Administration (NOAA) National Centers for Environmental Information (NCEI) under [https://doi.org/10.25921/1f4w-0t92](#). Observed values of seawater ¹⁸O showed in Fig. 4 are from the Global Seawater Oxygen-18 Database - v1.22 and can be

1175

accessed at <https://data.giss.nasa.gov/o18data/> (Schmidt, 1999; Bigg and Rohling, 1999). Sea-ice observations are from the Sea Ice Index, Version 3 of the National Snow and Ice Data Center, Boulder, Colorado, USA (Fetterer et al., 2017) and can be accessed at <https://doi.org/10.7265/N5K072F8>. World Ocean Atlas database (Boyer et al., 2018) used for calculation of reference temperature for organic matter remineralization can be accessed at NOAA NCEI under
1180 <https://www.ncei.noaa.gov/archive/accession/NCEI-WOA18>. Southern Hemisphere atmospheric CO₂ data were obtained from the Cape Grim Baseline Air Pollution Station belonging to the Commonwealth Scientific and Industrial Research Organisation (CSIRO) Oceans & Atmosphere and the Australian Bureau of Meteorology. This data can be freely downloaded at <https://capegrim.csiro.au/>. Northern Hemisphere atmospheric CO₂ data were obtained from the Mauna Loa Observatory
1185 <https://gml.noaa.gov/ccgg/trends/>. Present-day volcano distributions were obtained from the Volcano Location Database belonging to the NOAA NCEI. These data can be freely downloaded from <https://www.ngdc.noaa.gov/hazel/view/hazards/volcano/loc-data>. Reference wind values for air-sea exchange are from the NOAA/CIRES/DOE 20th Century Reanalysis (V3) dataset provided by the NOAA PSL, Boulder, Colorado, USA, from their website at <https://psl.noaa.gov>.

1190 *Author contributions.* EF and GS designed the work, EF led the development of the new model and code to which GS contributed. EF wrote the original draft, and together with GS discussed model concepts and results and wrote the final manuscript.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. We thank to the Chilean National Agency for Research and Development (ANID) for financial support of
1195 this work which was largely supported through the ANID/CONICYT-PFCHA/Doctorado Nacional/2017-21171747 and FONDECYT (Chile) 1190230 grants. We also received support from FONDECYT (Chile) grant 1230534 and the ANID Millennium Science Initiative Program (Millennium Institute of Oceanography ICN12_019). In addition, we thank Roberto Rondanelli who helped us with equatorial-cross heat transport parameterization. Finally, we would like to extend special thanks to Donald Canfield who invited EF to spend time at his lab at the University of Southern Denmark and whose valuable
1200 comments and discussions helped us with the formulation of the ocean biogeochemistry module.

References

- Aagaard, K., Coachman, L. K., and Carmack, E.: On the halocline of the Arctic Ocean, Deep Sea Research Part A. Oceanographic Research Papers, 28, 529-545, [https://doi.org/10.1016/0198-0149\(81\)90115-1](https://doi.org/10.1016/0198-0149(81)90115-1), 1981.
- Amante, C. and Eakins, B.W.: ETOPO1 1 Arc-Minute Global Relief Model: Procedures, Data Sources and Analysis, NOAA
1205 Technical Memorandum NESDIS NGDC-24. National Geophysical Data Center, NOAA, <https://www.ncei.noaa.gov/products/etopo-global-relief-model>, 2009.
- Antoine, D. and Morel, A.: Modelling the seasonal course of the upper ocean pCO₂ (I). Development of a one-dimensional model, Tellus B, 47, 103-121, <https://doi.org/10.1034/j.1600-0889.47.issue1.11.x>, 1995.
- Archer, D. E.: An atlas of the distribution of calcium carbonate in sediments of the deep sea, Global Biogeochem. Cy., 10,
1210 159-174, <https://doi.org/10.1029/95GB03016>, 1996.

- Archer, D. E., Morford, J. L., and Emerson, S. R.: A model of suboxic sedimentary diagenesis suitable for automatic tuning and gridded global domains, *Global Biogeochem. Cy.*, 16, 17-1-17-21, <https://doi.org/10.1029/2000GB001288>, 2002.
- Baines, P. G.: Mixing regimes for the flow of dense fluid down slopes into stratified environments, *J. Fluid Mech.*, 538, 245-267, <https://doi.org/10.1017/S0022112005005434>, 2005.
- 1215 Baines, P. G.: Mixing in Downslope Flows in the Ocean - Plumes versus Gravity Currents, *Atmos-Ocean*, 46, 405-419, <https://doi.org/10.3137/ao.460402>, 2008.
- Bendtsen, J.: Climate sensitivity to changes in solar insolation in a simple coupled climate model, *Clim. Dynam.*, 18, 595-609, <https://doi.org/10.1007/s00382-001-0198-4>, 2002.
- Bendtsen, J., Hilligsoe, K. M., Hansen, J. L. S., and Richardson, K.: Analysis of remineralisation, lability, temperature
1220 sensitivity and structural composition of organic matter from the upper ocean, *Prog. Oceanogr.*, 130, 125-145, <https://doi.org/10.1016/j.pocean.2014.10.009>, 2015.
- Berelson, W. M., Balch, W. M., Najjar, R., Feely, R. A., Sabine, C., and Lee, K.: Relating estimates of CaCO₃ production, export, and dissolution in the water column to measurements of CaCO₃ rain into sediment traps and dissolution on the sea floor: A revised global carbonate budget, *Global Biogeochem. Cy.*, 21, GB1024, <https://doi.org/10.1029/2006gb002803>, 2007.
- 1225 Berger, A. and Loutre, M. F.: Insolation values for the climate of the last 10 million years, *Quaternary Sci. Rev.*, 10, 297-317, [https://doi.org/10.1016/0277-3791\(91\)90033-Q](https://doi.org/10.1016/0277-3791(91)90033-Q), 1991.
- [Berner, R. A.: Burial of organic carbon and pyrite sulfur in the modern ocean: its geochemical and environmental significance. *Am. J. Sci.*, 282, 451-473, <https://doi.org/10.2475/ajs.282.4.451>, 1982.](#)
- [Berner, R. A.: A model for atmospheric CO₂ over Phanerozoic time, *Am. J. Sci.*, 291, 339-376, <https://doi.org/10.2475/ajs.291.4.339>, 1991.](#)
- 1230 [Berner, R. A.: Chemical weathering and its effect on atmospheric CO₂ and climate, in: *Chemical Weathering Rates of Silicate Minerals*, edited by: White, A. F., and Brantley, S. L., De Gruyter, Berlin, Boston, 565-584, <https://doi.org/10.1515/9781501509650-015>, 1995.](#)
- [Berner, R. A.: GEOCARBSULF: A combined model for Phanerozoic atmospheric O₂ and CO₂, *Geochim. Cosmochim. Ac.*,
1235 70, 5653-5664, <https://doi.org/10.1016/j.gca.2005.11.032>, 2006.](#)
- Bonan, G. B.: Forests and Climate Change: Forcings, Feedbacks, and the Climate Benefits of Forests, *Science*, 320, 1444-1449, <https://doi.org/10.1126/science.1155121>, 2008.
- Boyer, T. P., García, H. E., Locarnini, R. A., Zweng, M. M., Mishonov, A. V., Reagan, J. R., Weathers, K. A., Baranova, O. K., Paver, C. R., Seidov, D., and Smolyar, I. V.: World Ocean Atlas 2018, NOAA National Centers for Environmental
1240 Information [data set], <https://www.ncei.noaa.gov/archive/accession/NCEI-WOA18>, 2018.
- Budyko, M. I.: Effect of Solar Radiation Variations on Climate of Earth, *Tellus*, 21, 611-619, <https://doi.org/10.3402/tellusa.v21i5.10109>, 1969.
- Byrne, B. and Goldblatt, C.: Radiative forcing at high concentrations of well- mixed greenhouse gases, *Geophys. Res. Lett.*, 41, 152-160, <https://doi.org/10.1002/2013gl058456>, 2014.

- 1245 [Cartapanis, O., Galbraith, E. D., Bianchi, D., and Jaccard, S. L.: Carbon burial in deep-sea sediment and implications for oceanic inventories of carbon and alkalinity over the last glacial cycle, *Clim Past*, 14, 1819-1850, <https://doi.org/10.5194/cp-14-1819-2018>, 2018.](#)
- [Chapin III, S. F., Matson, P. A., and Vitousek, P.: Principles of terrestrial ecosystem ecology, Springer, New York, NY, USA, <https://doi.org/10.1007/978-1-4419-9504-9>, 2011.](#)
- 1250 [Chen, J. J., Swart, N. C., Beadling, R., Cheng, X., Hattermann, T., Jüling, A., Li, Q., Marshall, J., Martin, T., Muilwijk, M., Pauling, A. G., Purich, A., Smith, I. J., and Thomas, M.: Reduced deep convection and bottom water formation due to Antarctic meltwater in a multi-model ensemble, *Geophys. Res. Lett.*, 50, e2023GL106492, <https://doi.org/10.1029/2023gl106492>, 2023.](#)
- Dai, A. and Trenberth, K. E.: Estimates of Freshwater Discharge from Continents: Latitudinal and Seasonal Variations, *J. Hydrometeorol*, 3, 660-687, [https://doi.org/10.1175/1525-7541\(2002\)003<0660:EOFDfC>2.0.CO;2](https://doi.org/10.1175/1525-7541(2002)003<0660:EOFDfC>2.0.CO;2), 2002.
- 1255 [Danabasoglu, G., Large, W. G., and Briegleb, B. P.: Climate impacts of parameterized Nordic Sea overflows, *J. Geophys. Res.*, 115, C11005, <https://doi.org/10.1029/2010JC006243>, 2010.](#)
- [DeConto, R. M. and Pollard, D.: Contribution of Antarctica to past and future sea-level rise, *Nature*, 531, 591-597, <https://doi.org/10.1038/nature17145>, 2016.](#)
- [Dey, D. and Döös, K.: Atmospheric Freshwater Transport From the Atlantic to the Pacific Ocean: A Lagrangian Analysis, *Geophys. Res. Lett.*, 47, e2019GL086176, <https://doi.org/10.1029/2019GL086176>, 2020.](#)
- 1260 [d'Orgeville, M., Sijp, W. P., England, M. H., and Meissner, K. J.: On the control of glacial-interglacial atmospheric CO₂ variations by the Southern Hemisphere westerlies, *Geophys. Res. Lett.*, 37, L21703, <https://doi.org/10.1029/2010gl045261>, 2010.](#)
- [Drever, J. I.: The effect of land plants on weathering rates of silicate minerals, *Geochim. Cosmochim. Ac.*, 58, 2325-2332, \[https://doi.org/10.1016/0016-7037\\(94\\)90013-2\]\(https://doi.org/10.1016/0016-7037\(94\)90013-2\), 1994.](#)
- 1265 [Dunne, J. P., Sarmiento, J. L., and Gnanadesikan, A.: A synthesis of global particle export from the surface ocean and cycling through the ocean interior and on the seafloor, *Global Biogeochem. Cy.*, 21, GB4006, <https://doi.org/10.1029/2006gb002907>, 2007.](#)
- [Eby, M., Weaver, A. J., Alexander, K., Zickfeld, K., Abe-Ouchi, A., Cimadoribus, A. A., Crespin, E., Drijfhout, S. S., Edwards, N. R., Eliseev, A. V., Feulner, G., Fichefet, T., Forest, C. E., Goosse, H., Holden, P. B., Joos, F., Kawamiya, M., Kicklighter, D., Kienert, H., Matsumoto, K., Mokhov, I. I., Monier, E., Olsen, S. M., Pedersen, J. O. P., Perrette, M., Philippon-Berthier, G., Ridgwell, A., Schlosser, A., Schneider Von Deimling, T., Shaffer, G., Smith, R. S., Spahni, R., Sokolov, A. P., Steinacher, M., Tachiiri, K., Tokos, K., Yoshimori, M., Zeng, N., and Zhao, F.: Historical and idealized climate model experiments: an intercomparison of Earth system models of intermediate complexity, *Clim. Past*, 9, 1111-1140, \[https://doi.org/10.5194/cp-9-\]\(https://doi.org/10.5194/cp-9-1111-2013\)](#)
- 1275 1111-2013, 2013.
- [Eichinger, R., Shaffer, G., Albarran, N., Rojas, M., and Lambert, F.: An improved land biosphere module for use in the DCESS Earth system model \(version 1.1\) with application to the last glacial termination, *Geosci. Model Dev.*, 10, 3481-3498, <https://doi.org/10.5194/gmd-10-3481-2017>, 2017.](#)

- England, M. H.: Representing the Global-Scale Water Masses in Ocean General Circulation Models, *J. Phys. Oceanogr.*, 23, 1523-1552, [https://doi.org/10.1175/1520-0485\(1993\)023<1523:RTGSWM>2.0.CO;2](https://doi.org/10.1175/1520-0485(1993)023<1523:RTGSWM>2.0.CO;2), 1993.
- Fetterer, F., Knowles, K., Meier, W. N., Savoie, M., and Windnagel, A. K.: Sea Ice Index, Version 3, National Snow and Ice Data Center, Boulder, Colorado USA [data set], <https://doi.org/10.7265/N5K072F8>, 2017.
- [Filippelli, G. M.: The Global Phosphorus Cycle, *Reviews in Mineralogy and Geochemistry*, 48, 391-425, <https://doi.org/10.2138/rmg.2002.48.10>, 2002.](https://doi.org/10.2138/rmg.2002.48.10)
- 1285 [Fogwill, C. J., Phipps, S. J., Turney, C. S. M., and Golledge, N. R.: Sensitivity of the Southern Ocean to enhanced regional Antarctic ice sheet meltwater input, *Earth's Future*, 3, 317-329, <https://doi.org/10.1002/2015ef000306>, 2015.](https://doi.org/10.1002/2015ef000306)
- [Gaillardet, J., Dupré, B., Louvat, P., and Allègre, C. J.: Global silicate weathering and CO₂ consumption rates deduced from the chemistry of large rivers, *Chem. Geol.*, 159, 3-30, \[https://doi.org/10.1016/S0009-2541\\(99\\)00031-5\]\(https://doi.org/10.1016/S0009-2541\(99\)00031-5\), 1999.](https://doi.org/10.1016/S0009-2541(99)00031-5)
- Gangstør, R., Joos, F., and Gehlen, M.: Sensitivity of pelagic calcification to ocean acidification, *Biogeosciences*, 8, 433-458, <https://doi.org/10.5194/bg-8-433-2011>, 2011.
- 1290 Gargett, A. E. and Holloway, G.: Dissipation and diffusion by internal wave breaking, *J. Mar. Res.*, 42, 15-27, 1984.
- Gerber, S., Joos, F., and Prentice, I. C.: Sensitivity of a dynamic global vegetation model to climate and atmospheric CO₂, *Glob. Change Biol.*, 10, 1223-1239, <https://doi.org/10.1111/j.1529-8817.2003.00807.x>, 2004.
- Gordon, A. L.: Bottom Water Formation, in: *Encyclopedia of Ocean Sciences (Third Edition)*, edited by: Cochran, J. K., 1295 Bokuniewicz, H. J., and Yager, P. L., Academic Press, Oxford, 120-126, <https://doi.org/10.1016/B978-0-12-409548-9.04019-7>, 2019.
- Gray, W. R., De Lavergne, C., Jnglin Wills, R. C., Menviel, L., Spence, P., Holzer, M., Kageyama, M., and Michel, E.: Poleward Shift in the Southern Hemisphere Westerly Winds Synchronous With the Deglacial Rise in CO₂, *Paleoceanography and Paleoclimatology*, 38, e2023PA004666, <https://doi.org/10.1029/2023pa004666>, 2023.
- 1300 Gröger, M. and Mikolajewicz, U.: Note on the CO₂ air-sea gas exchange at high temperatures, *Ocean Model.*, 39, 284-290, <https://doi.org/10.1016/j.ocemod.2011.05.003>, 2011.
- Haney, R. L.: Surface Thermal Boundary Condition for Ocean Circulation Models, *J. Phys. Oceanogr.*, 1, 241-248, [https://doi.org/10.1175/1520-0485\(1971\)001<0241:Stbcfo>2.0.Co;2](https://doi.org/10.1175/1520-0485(1971)001<0241:Stbcfo>2.0.Co;2), 1971.
- 1305 Harper, D. T., Hönisch, B., Zeebe, R. E., Shaffer, G., Haynes, L. L., Thomas, E., and Zachos, J. C.: The Magnitude of Surface Ocean Acidification and Carbon Release During Eocene Thermal Maximum 2 (ETM-2) and the Paleocene-Eocene Thermal Maximum (PETM), *Paleoceanography and Paleoclimatology*, 35, e2019PA003699, <https://doi.org/10.1029/2019pa003699>, 2020.
- Hartmann, D. L.: *Global Physical Climatology (2nd ed.)*, Elsevier Press, Amsterdam, Netherlands, <https://doi.org/10.1016/C2009-0-00030-0>, 2016.
- 310 [Hartmann, J., Jansen, N., Dürr, H. H., Kempe, S., and Köhler, P.: Global CO₂-consumption by chemical weathering: What is the contribution of highly active weathering regions?, *Global Planet. Change*, 69, 185-194, <https://doi.org/10.1016/j.gloplacha.2009.07.007>, 2009.](https://doi.org/10.1016/j.gloplacha.2009.07.007)

- Haumann, F. A., Gruber, N., Münnich, M., Frenger, I., and Kern, S.: Sea-ice transport driving Southern Ocean salinity and its recent trends, *Nature*, 537, 89-92, <https://doi.org/10.1038/nature19101>, 2016.
- 1315 [Hayes, C. T., Costa, K. M., Anderson, R. F., Calvo, E., Chase, Z., Demina, L. L., Dutay, J. C., German, C. R., Heimbürger-Boavida, L. E., Jaccard, S. L., Jacobel, A., Kohfeld, K. E., Kravchishina, M. D., Lippold, J., Mekik, F., Missiaen, L., Pavia, F. J., Paytan, A., Pedrosa-Pamies, R., Petrova, M. V., Rahman, S., Robinson, L. F., Roy-Barman, M., Sanchez-Vidal, A., Shiller, A., Tagliabue, A., Tessin, A. C., Van Hulten, M., and Zhang, J.: Global ocean sediment composition and burial flux in the deep sea, *Global Biogeochem. Cy.*, 35, e2020GB006769, <https://doi.org/10.1029/2020gb006769>, 2021.](#)
- 1320 Hazarika, M. K., Yasuoka, Y., Ito, A., and Dye, D.: Estimation of net primary productivity by integrating remote sensing data with an ecosystem model, *Remote Sens. of Environ.*, 94, 298-310, <https://doi.org/10.1016/j.rse.2004.10.004>, 2005.
- [Hedges, J. I. and Keil, R. G.: Sedimentary organic matter preservation: an assessment and speculative synthesis, *Mar. Chem.*, 49, 81-115, \[https://doi.org/10.1016/0304-4203\\(95\\)00008-F\]\(https://doi.org/10.1016/0304-4203\(95\)00008-F\), 1995.](#)
- Heuzé, C.: Antarctic Bottom Water and North Atlantic Deep Water in CMIP6 models, *Ocean Sci.*, 17, 59-90, <https://doi.org/10.5194/os-17-59-2021>, 2021.
- 1325 Heywood, K. J., Schmidtko, S., Heuzé, C., Kaiser, J., Jickells, T. D., Queste, B. Y., Stevens, D. P., Wadley, M., Thompson, A. F., Fielding, S., Guihen, D., Creed, E., Ridley, J. K., and Smith, W.: Ocean processes at the Antarctic continental slope, *Philos. T. R. Soc. A*, 372, 20130047, <https://doi.org/10.1098/rsta.2013.0047>, 2014.
- Hirschi, J. J. M., Barnier, B., Böning, C., Biastoch, A., Blaker, A. T., Coward, A., Danilov, S., Drijfhout, S., Getzlaff, K., 1330 Griffies, S. M., Hasumi, H., Hewitt, H., Iovino, D., Kawasaki, T., Kiss, A. E., Koldunov, N., Marzocchi, A., Mecking, J. V., Moat, B., Molines, J. M., Myers, P. G., Penduff, T., Roberts, M., Treguier, A. M., Sein, D. V., Sidorenko, D., Small, J., Spence, P., Thompson, L., Weijer, W., and Xu, X.: The Atlantic Meridional Overturning Circulation in High-Resolution Models, *J. Geophys. Res.-Oceans*, 125, e2019JC015522, <https://doi.org/10.1029/2019jc015522>, 2020.
- IOC, SCOR, IAPSO: The international thermodynamic equation of seawater - 2010: Calculation and use of thermodynamic 1335 properties. Intergovernmental Oceanographic Commission, Manuals and Guides No. 56, UNESCO (English), 196 pp., 2010.
- Jochum, M. and Eden, C.: The Connection between Southern Ocean Winds, the Atlantic Meridional Overturning Circulation, and Indo-Pacific Upwelling, *J. Climate*, 28, 9250-9257, <https://doi.org/10.1175/jcli-d-15-0263.1>, 2015.
- Johnson, G. C.: Quantifying Antarctic Bottom Water and North Atlantic Deep Water volumes, *J. Geophys. Res.*, 113, C05027, <https://doi.org/10.1029/2007jc004477>, 2008.
- 1340 [IPCC: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, in press, <https://doi.org/10.1017/9781009157896>, 2021.](#)
- 1345 ~~Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J., Maycock, T., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B. (Eds.):~~

~~Summary for Policymakers, in: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, Tech. rep., Intergovernmental Panel on Climate Change, <https://doi.org/10.1017/9781009157896>, 2021.~~

1350 Joos, F., Roth, R., Fuglestedt, J. S., Peters, G. P., Enting, I. G., Von Bloh, W., Brovkin, V., Burke, E. J., Eby, M., Edwards, N. R., Friedrich, T., Frölicher, T. L., Halloran, P. R., Holden, P. B., Jones, C., Kleinen, T., Mackenzie, F. T., Matsumoto, K., Meinshausen, M., Plattner, G.-K., Reisinger, A., Segschneider, J., Shaffer, G., Steinacher, M., Strassmann, K., Tanaka, K., Timmermann, A., and Weaver, A. J.: Carbon dioxide and climate impulse response functions for the computation of greenhouse gas metrics: a multi-model analysis, *Atmos. Chem. Phys.*, 13, 2793-2825, [https://doi.org/10.5194/acp-13-2793-](https://doi.org/10.5194/acp-13-2793-2013)
1355 2013, 2013.

Keeling, R. F., Stephens, B. B., Najjar, R. G., Doney, S. C., Archer, D., and Heimann, M.: Seasonal variations in the atmospheric O₂/N₂ ratio in relation to the kinetics of air-sea gas exchange, *Global Biogeochem. Cy.*, 12, 141-163, <https://doi.org/10.1029/97gb02339>, 1998.

[Kim, H. and Timmermann, A.: Seawater oxygen isotopes as a tool for monitoring future meltwater from the Antarctic ice-sheet, *Communications Earth & Environment*, 5, <https://doi.org/10.1038/s43247-024-01514-4>, 2024.](https://doi.org/10.1038/s43247-024-01514-4)

1360 Komar, N. and Zeebe, R. E.: Reconciling atmospheric CO₂, weathering, and calcite compensation depth across the Cenozoic, *Science Advances*, 7, eabd4876, <https://doi.org/10.1126/sciadv.abd4876>, 2021.

[Krissansen-Totton, J. and Catling, D. C.: Constraining climate sensitivity and continental versus seafloor weathering using an inverse geological carbon cycle model, *Nat. Commun.*, 8, 15423, <https://doi.org/10.1038/ncomms15423>, 2017.](https://doi.org/10.1038/ncomms15423)

1365 Kunze, E.: Internal-Wave-Driven Mixing: Global Geography and Budgets, *J. Phys. Oceanogr.*, 47, 1325-1345, <https://doi.org/10.1175/jpo-d-16-0141.1>, 2017.

Laufkotter, C., John, J. G., Stock, C. A., and Dunne, J. P.: Temperature and oxygen dependence of the remineralization of organic matter, *Global Biogeochem. Cy.*, 31, 1038-1050, <https://doi.org/10.1002/2017gb005643>, 2017.

1370 Lauvset, S. K., Lange, N., Tanhua, T., Bittig, H. C., Olsen, A., Kozyr, A., Alin, S. R., Álvarez, M., Azetsu-Scott, K., Barbero, L., Becker, S., Brown, P. J., Carter, B. R., Cotrim da Cunha, L., Feely, R. A., Hoppema, M., Humphreys, M. P., Ishii, M., Jeansson, E., Jiang, L. Q., Jones, S. D., Lo Monaco, C., Murata, A., Müller, J. D., Pérez, F. F., Pfeil, B., Schirnick, C., Steinfeldt, R., Suzuki, T., Tilbrook, B., Ulfso, A., Velo, A., Woosley, R. J., and Key, R. M.: Global Ocean Data Analysis Project version 2.2022 (GLODAPv2.2022), NOAA National Centers for Environmental Information [data set], <https://doi.org/10.25921/1f4w-0t92>, 2022.

1375 Lee, S. Y., Chiang, J. C. H., Matsumoto, K., and Tokos, K. S.: Southern Ocean wind response to North Atlantic cooling and the rise in atmospheric CO₂: Modeling perspective and paleoceanographic implications, *Paleoceanography*, 26, PA1214, <https://doi.org/10.1029/2010pa002004>, 2011.

[Lenton, T. M., Daines, S. J., and Mills, B. J. W.: COPSE reloaded: An improved model of biogeochemical cycling over Phanerozoic time, *Earth-Sci. Rev.*, 178, 1-28, <https://doi.org/10.1016/j.earscirev.2017.12.004>, 2018.](https://doi.org/10.1016/j.earscirev.2017.12.004)

1380

- Li, M., Peng, C., Wang, M., Xue, W., Zhang, K., Wang, K., Shi, G., and Zhu, Q.: The carbon flux of global rivers: A re-evaluation of amount and spatial patterns, *Ecol. Indic.*, 80, 40-51, <https://doi.org/10.1016/j.ecolind.2017.04.049>, 2017.
- [Li, Q., England, M. H., Hogg, A. M., Rintoul, S. R., and Morrison, A. K.: Abyssal ocean overturning slowdown and warming driven by Antarctic meltwater, *Nature*, 615, 841-847, <https://doi.org/10.1038/s41586-023-05762-w>, 2023.](#)
- 1385 Lohmann, G. and Lorenz, S.: On the hydrological cycle under paleoclimatic conditions as derived from AGCM simulations, *J. Geophys. Res.*, 105, 17417-17436, <https://doi.org/10.1029/2000JD900189>, 2000.
- [Ludwig, W., Amiotte-Suchet, P., and Probst, J.-L.: Enhanced chemical weathering of rocks during the last glacial maximum: a sink for atmospheric CO₂?, *Chem. Geol.*, 159, 147-161, \[https://doi.org/10.1016/S0009-2541\\(99\\)00038-8\]\(https://doi.org/10.1016/S0009-2541\(99\)00038-8\), 1999.](#)
- MacDougall, A. H., Frölicher, T. L., Jones, C. D., Rogelj, J., Matthews, H. D., Zickfeld, K., Arora, V. K., Barrett, N. J., Brovkin, V., Burger, F. A., Eby, M., Eliseev, A. V., Hajima, T., Holden, P. B., Jeltsch-Thömmes, A., Koven, C., Mengis, N., Menviel, L., Michou, M., Mokhov, I. I., Oka, A., Schwinger, J., Séférian, R., Shaffer, G., Sokolov, A., Tachiiri, K., Tjiputra, J., Wiltshire, A., and Ziehn, T.: Is there warming in the pipeline? A multi-model analysis of the Zero Emissions Commitment from CO₂, *Biogeosciences*, 17, 2987-3016, <https://doi.org/10.5194/bg-17-2987-2020>, 2020.
- 1395 [Mackie, S., Smith, I. J., Ridley, J. K., Stevens, D. P., and Langhorne, P. J.: Climate response to increasing Antarctic iceberg and ice shelf melt, *J Climate*, 33, 8917-8938, <https://doi.org/10.1175/jcli-d-19-0881.1>, 2020.](#)
- Maier-Reimer, E.: Geochemical cycles in an ocean general circulation model. Preindustrial tracer distributions, *Global Biogeochem. Cy.*, 7, 645-677, <https://doi.org/10.1029/93gb01355>, 1993.
- Marchal, O., Stocker, T. F., and Joos, F.: A latitude-depth, circulation-biogeochemical ocean model for paleoclimate studies. Development and sensitivities, *Tellus B*, 50, 290-316, <https://doi.org/10.3402/tellusb.v50i3.16130>, 1998.
- 1400 Marsay, C. M., Sanders, R. J., Henson, S. A., Pabortsava, K., Achterberg, E. P., and Lampitt, R. S.: Attenuation of sinking particulate organic carbon flux through the mesopelagic ocean, *P. Natl. Acad. Sci. USA*, 112, 1089-1094, <https://doi.org/10.1073/pnas.1415311112>, 2015.
- Martin, J. H., Knauer, G. A., Karl, D. M., and Broenkow, W. W.: VERTEX: carbon cycling in the northeast Pacific, *Deep Sea Research Part A. Oceanographic Research Papers*, 34, 267-285, [https://doi.org/10.1016/0198-0149\(87\)90086-0](https://doi.org/10.1016/0198-0149(87)90086-0), 1987.
- 1405 Masson-Delmotte, V., Hou, S., Ekaykin, A., Jouzel, J., Aristarain, A., Bernardo, R. T., Bromwich, D., Cattani, O., Delmotte, M., Falourd, S., Frezzotti, M., Gallée, H., Genoni, L., Isaksson, E., Landais, A., Helsen, M. M., Hoffmann, G., Lopez, J., Morgan, V., Motoyama, H., Noone, D., Oerter, H., Petit, J. R., Royer, A., Uemura, R., Schmidt, G. A., Schlosser, E., Simões, J. C., Steig, E. J., Stenni, B., Stievenard, M., Van Den Broeke, M. R., Van De Wal, R. S. W., Van De Berg, W. J., Vimeux, F., and White, J. W. C.: A Review of Antarctic Surface Snow Isotopic Composition: Observations, Atmospheric Circulation, and
- 1410 Isotopic Modeling, *J. Climate*, 21, 3359-3387, <https://doi.org/10.1175/2007jcli2139.1>, 2008.
- [Mayr, C., Lücke, A., Wagner, S., Wissel, H., Ohlendorf, C., Haberzettl, T., Oehlerich, M., Schäbitz, F., Wille, M., Zhu, J., and Zolitschka, B.: Intensified Southern Hemisphere westerlies regulated atmospheric CO₂ during the last deglaciation, *Geology*, 41, 831-834, <https://doi.org/10.1130/G34335.1>, 2013.](#)

- 1415 [Meehl, G. A., Senior, C. A., Eyring, V., Flato, G., Lamarque, J.-F., Stouffer, R. J., Taylor, K. E., and Schlund, M.: Context for interpreting equilibrium climate sensitivity and transient climate response from the CMIP6 Earth system models, *Science Advances*, 6, eaba1981, <https://doi.org/10.1126/sciadv.aba1981>, 2020.](#)
- [Menviel, L., Spence, P., Yu, J., Chamberlain, M. A., Matear, R. J., Meissner, K. J., and England, M. H.: Southern Hemisphere westerlies as a driver of the early deglacial atmospheric CO₂ rise, *Nat. Commun.*, 9, <https://doi.org/10.1038/s41467-018-04876-4>, 2018.](#)
- 1420 [Milliman, J. D.: Production and accumulation of calcium carbonate in the ocean: Budget of a nonsteady state, *Global Biogeochem. Cy.*, 7, 927-957, <https://doi.org/10.1029/93GB02524>, 1993.](#)
- [Milliman, J. and Droxler, A.: Calcium carbonate sedimentation in the global ocean: Linkages between the neritic and pelagic environments, *Oceanography*, 8, 92-94, <https://doi.org/10.5670/oceanog.1995.04>, 1995.](#)
- Mucci, A.: The solubility of calcite and aragonite in seawater at various salinities, temperatures, and one atmosphere total pressure, *Am. J. Sci.*, 283, 780-799, <https://doi.org/10.2475/ajs.283.7.780>, 1983.
- 1425 [Munhoven, G.: Glacial–interglacial changes of continental weathering: estimates of the related CO₂ and HCO₃⁻ flux variations and their uncertainties, *Global Planet. Change*, 33, 155-176, \[https://doi.org/10.1016/S0921-8181\\(02\\)00068-1\]\(https://doi.org/10.1016/S0921-8181\(02\)00068-1\), 2002.](#)
- [Munhoven, G.: Glacial–interglacial rain ratio changes: Implications for atmospheric CO₂ and ocean–sediment interaction, *Deep Sea Research Part II: Topical Studies in Oceanography*, 54, 722-746, <https://doi.org/10.1016/j.dsr2.2007.01.008>, 2007.](#)
- 1430 [Mutshinda, C., Finkel, Z., Widdicombe, C., and Irwin, A.: Phytoplankton traits from long-term oceanographic time-series, *Mar. Ecol. Prog. Ser.*, 576, 11-25, <https://doi.org/10.3354/meps12220>, 2017.](#)
- Olsen, S. M., Shaffer, G., and Bjerrum, C. J.: Ocean oxygen isotope constraints on mechanisms for millennial-scale climate variability, *Paleoceanography and Paleoclimatology*, 20, PA1014, <https://doi.org/10.1029/2004PA001063>, 2005.
- Oort, A. H. and Peixoto, J. P.: Global Angular-Momentum and Energy-Balance Requirements from Observations, *Adv. Geophys.*, 25, 355-490, [https://doi.org/10.1016/S0065-2687\(08\)60177-6](https://doi.org/10.1016/S0065-2687(08)60177-6), 1983.
- 1435 [Orsi, A. H.: Recycling bottom waters, *Nat. Geosci.*, 3, 307-309, <https://doi.org/10.1038/ngeo854>, 2010.](#)
- [Orsi, A. H., Johnson, G. C., and Bullister, J. L.: Circulation, mixing, and production of Antarctic Bottom Water, *Prog. Oceanogr.*, 43, 55-109, \[https://doi.org/10.1016/S0079-6611\\(99\\)00004-X\]\(https://doi.org/10.1016/S0079-6611\(99\)00004-X\), 1999.](#)
- [Orsi, A. H., Jacobs, S. S., Gordon, A. L., and Visbeck, M.: Cooling and ventilating the Abyssal Ocean, *Geophys. Res. Lett.*, 28, 2923-2926, <https://doi.org/10.1029/2001gl012830>, 2001.](#)
- 1440 [Orsi, A. H., Smethie, W. M. J., and Bullister, J., L.: On the total input of Antarctic waters to the deep ocean: A preliminary estimate from chlorofluorocarbon measurements, *J. Geophys. Res.*, 107, 31-1-31-14, <https://doi.org/10.1029/2001JC000976>, 2002.](#)
- [Pagani, M., Arthur, M. A., and Freeman, K. H.: Miocene evolution of atmospheric carbon dioxide, *Paleoceanography*, 14, 273-292, <https://doi.org/10.1029/1999pa900006>, 1999.](#)
- 1445 [Park, W. and Latif, M.: Ensemble global warming simulations with idealized Antarctic meltwater input, *Clim Dynam.*, 52, 3223-3239, <https://doi.org/10.1007/s00382-018-4319-8>, 2019.](#)

[Paytan, A. and McLaughlin, K.: The oceanic phosphorus cycle, Chem. Rev., 107, 563-576, https://doi.org/10.1021/cr0503613, 2007.](https://doi.org/10.1021/cr0503613)

- 1450 Planchat, A., Kwiatkowski, L., Bopp, L., Torres, O., Christian, J. R., Butenschön, M., Lovato, T., Séférian, R., Chamberlain, M. A., Aumont, O., Watanabe, M., Yamamoto, A., Yool, A., Ilyina, T., Tsujino, H., Krumhardt, K. M., Schwinger, J., Tjiputra, J., Dunne, J. P., and Stock, C.: The representation of alkalinity and the carbonate pump from CMIP5 to CMIP6 Earth system models and implications for the carbon cycle, *Biogeosciences*, 20, 1195-1257, <https://doi.org/10.5194/bg-20-1195-2023>, 2023.
- 1455 Purkey, S. G. and Johnson, G. C.: Antarctic Bottom Water Warming and Freshening: Contributions to Sea Level Rise, Ocean Freshwater Budgets, and Global Heat Gain, *J. Climate*, 26, 6105-6122, <https://doi.org/10.1175/JCLI-D-12-00834.1>, 2013.
- Rahmstorf, S. and England, M. H.: Influence of Southern Hemisphere Winds on North Atlantic Deep Water Flow, *J. Phys. Oceanogr.*, 27, 2040-2054, [https://doi.org/10.1175/1520-0485\(1997\)027<2040:IOSHWO>2.0.CO;2](https://doi.org/10.1175/1520-0485(1997)027<2040:IOSHWO>2.0.CO;2), 1997.
- Regaudie-de-Gioux, A. and Duarte, C. M.: Temperature dependence of planktonic metabolism in the ocean, *Global Biogeochem. Cy.*, 26, GB1015, <https://doi.org/10.1029/2010gb003907>, 2012.
- 1460 Rodriguez, J. M., Johns, T. C., Thorpe, R. B., and Wiltshire, A.: Using moisture conservation to evaluate oceanic surface freshwater fluxes in climate models, *Clim. Dynam.*, 37, 205-219, <https://doi.org/10.1007/s00382-010-0899-7>, 2011.
- Roquet, F., Madec, G., McDougall, T. J., and Barker, P. M.: Accurate polynomial expressions for the density and specific volume of seawater using the TEOS-10 standard, *Ocean Model.*, 90, 29-43, <https://doi.org/10.1016/j.ocemod.2015.04.002>, 2015.
- 1465 Ruckert, K. L., Shaffer, G., Pollard, D., Guan, Y., Wong, T. E., Forest, C. E., and Keller, K.: Assessing the Impact of Retreat Mechanisms in a Simple Antarctic Ice Sheet Model Using Bayesian Calibration, *PLOS ONE*, 12, e0170052, <https://doi.org/10.1371/journal.pone.0170052>, 2017.
- Sarmiento, J. L., and Gruber, N.: *Ocean Biogeochemical Dynamics*, Princeton University Press, Princeton, New Jersey, USA, <https://doi.org/10.2307/j.ctt3fgxqx>, 2006.
- 1470 Schmidt, G. A., Bigg, G. R., and Rohling, E. J.: *Global Seawater Oxygen-18 Database - v1.22*, <https://data.giss.nasa.gov/o18data/>, 1999.
- Shaffer, G.: Long-term effectiveness and consequences of carbon dioxide sequestration, *Nat. Geosci.*, 3, 464-467, <https://doi.org/10.1038/ngeo896>, 2010.
- Shaffer, G.: Formulation, calibration and validation of the DAIS model (version 1), a simple Antarctic ice sheet model sensitive to variations of sea level and ocean subsurface temperature, *Geosci. Model Dev.*, 7, 1803-1818, <https://doi.org/10.5194/gmd-7-1803-2014>, 2014.
- 1475 Shaffer, G. and Lambert, F.: In and out of glacial extremes by way of dust-climate feedbacks, *P. Natl. Acad. Sci. USA*, 115, 2026-2031, <https://doi.org/10.1073/pnas.1708174115>, 2018.
- Shaffer, G. and Olsen, S. M.: Sensitivity of the thermohaline circulation and climate to ocean exchanges in a simple coupled
- 1480 model, *Clim. Dynam.*, 17, 433-444, <https://doi.org/10.1007/pl00013739>, 2001.

- Shaffer, G., Bendtsen, J., and Ulloa, O.: Fractionation during remineralization of organic matter in the ocean, *Deep Sea Research Part I: Oceanographic Research Papers*, 46, 185-204, [https://doi.org/10.1016/S0967-0637\(98\)00061-2](https://doi.org/10.1016/S0967-0637(98)00061-2), 1999.
- Shaffer, G., Malskær Olsen, S., and Pepke Pedersen, J. O.: Presentation, calibration and validation of the low-order, DCESS Earth System Model (Version 1), *Geosci. Model Dev.*, 1, 17-51, <https://doi.org/10.5194/gmd-1-17-2008>, 2008.
- 1485 Shaffer, G., Olsen, S. M., and Pedersen, J. O. P.: Long-term ocean oxygen depletion in response to carbon dioxide emissions from fossil fuels, *Nat. Geosci.*, 2, 105-109, <https://doi.org/10.1038/ngeo420>, 2009.
- Shaffer, G., Huber, M., Rondanelli, R., and Pepke Pedersen, J. O.: Deep time evidence for climate sensitivity increase with warming, *Geophys. Res. Lett.*, 43, 6538-6545, <https://doi.org/10.1002/2016gl069243>, 2016.
- Shaffer, G., Fernández Villanueva, E., Rondanelli, R., Pedersen, J. O. P., Olsen, S. M., and Huber, M.: Implementation of methane cycling for deep-time global warming simulations with the DCESS Earth system model (version 1.2), *Geosci. Model Dev.*, 10, 4081-4103, <https://doi.org/10.5194/gmd-10-4081-2017>, 2017.
- 1490 Siegel, D. A., Buesseler, K. O., Doney, S. C., Saille, S. F., Behrenfeld, M. J., and Boyd, P. W.: Global assessment of ocean carbon export by combining satellite observations and food-web models, *Global Biogeochem. Cy.*, 28, 181-196, <https://doi.org/10.1002/2013gb004743>, 2014.
- 1495 [Siegenthaler, U. and Oeschger, H.: Biospheric CO₂ emissions during the past 200 years reconstructed by deconvolution of ice core data, *Tellus B*, 39B, 140-154, <https://doi.org/10.1111/j.1600-0889.1987.tb00278.x>, 1987.](https://doi.org/10.1111/j.1600-0889.1987.tb00278.x)
- [Silvano, A., Rintoul, S. R., Peña-Molino, B., Hobbs, W. R., van Wijk, E., Aoki, S., Tamura, T., and Williams, G. D.: Freshening by glacial meltwater enhances melting of ice shelves and reduces formation of Antarctic Bottom Water, *Science Advances*, 4, \[eap9467\]\(https://doi.org/10.1126/sciadv.aap9467\), <https://doi.org/10.1126/sciadv.aap9467>, 2018.](https://doi.org/10.1126/sciadv.aap9467)
- 1500 Stocker, T. F., Broecker, W. S., and Wright, D. G.: Carbon uptake experiments with a zonally-averaged global ocean circulation model, *Tellus B*, 46, 103-122, <https://doi.org/10.3402/tellusb.v46i2.15756>, 1994.
- Stone, P. H. and Miller, D. A.: Empirical relations between seasonal changes in meridional temperature gradients and meridional fluxes of heat, *J. Atmos. Sci.*, 37, 1708-1721, [https://doi.org/10.1175/1520-0469\(1980\)037<1708:ERBSCI>2.0.CO;2](https://doi.org/10.1175/1520-0469(1980)037<1708:ERBSCI>2.0.CO;2), 1980.
- 1505 Sulpis, O., Jeansson, E., Dinuer, A., Lauvset, S. K., and Middelburg, J. J.: Calcium carbonate dissolution patterns in the ocean, *Nat. Geosci.*, 14, 423-428, <https://doi.org/10.1038/s41561-021-00743-y>, 2021.
- Sverdrup, H. U., Johnson, M. W., and Fleming, R. H.: *The Oceans: Their physics, chemistry, and general biology*, Prentice Hall, Inc., New York, USA, 1942.
- Talley, L. D., Reid, J. L., and Robbins, P. E.: Data-based meridional overturning streamfunctions for the global ocean, *J. Climate*, 16, 3213-3226, [https://doi.org/10.1175/1520-0442\(2003\)016<3213:Dmosft>2.0.Co;2](https://doi.org/10.1175/1520-0442(2003)016<3213:Dmosft>2.0.Co;2), 2003.
- 1510 Toggweiler, J. R., Russell, J. L., and Carson, S. R.: Midlatitude westerlies, atmospheric CO₂, and climate change during the ice ages, *Paleoceanography*, 21, PA2005, <https://doi.org/10.1029/2005pa001154>, 2006.
- Trenberth, K. E. and Caron, J. M.: Estimates of Meridional Atmosphere and Ocean Heat Transports, *J. Climate*, 14, 3433-3443, [https://doi.org/10.1175/1520-0442\(2001\)014<3433:eomaa>2.0.co;2](https://doi.org/10.1175/1520-0442(2001)014<3433:eomaa>2.0.co;2), 2001.

- 1515 Wang, X. L., Stone, P. H., and Marotzke, J.: Global thermohaline circulation. Part II: Sensitivity with interactive atmospheric transports, *J. Climate*, 12, 83-91, [https://doi.org/10.1175/1520-0442\(1999\)012<0083:GTCPIS>2.0.CO;2](https://doi.org/10.1175/1520-0442(1999)012<0083:GTCPIS>2.0.CO;2), 1999.
- Wanninkhof, R.: Relationship between wind speed and gas exchange over the ocean, *J. Geophys. Res.*, 97, 7373-7382, <https://doi.org/10.1029/92JC00188>, 1992.
- Weiss, R. F.: The solubility of nitrogen, oxygen and argon in water and seawater, *Deep Sea Research and Oceanographic Abstracts*, 17, 721-735, [https://doi.org/10.1016/0011-7471\(70\)90037-9](https://doi.org/10.1016/0011-7471(70)90037-9), 1970.
- 1520 Weiss, R. F.: Carbon dioxide in water and seawater: the solubility of a non-ideal gas, *Marine Chemistry*, 2, 203-215, [https://doi.org/10.1016/0304-4203\(74\)90015-2](https://doi.org/10.1016/0304-4203(74)90015-2), 1974.
- [Willeit, M., Ilyina, T., Liu, B., Heinze, C., Perrette, M., Heinemann, M., Dalmonech, D., Brovkin, V., Munhoven, G., Börker, J., Hartmann, J., Romero-Mujalli, G., and Ganopolski, A.: The Earth system model CLIMBER-X v1.0 – Part 2: The global carbon cycle, *Geosci. Model Dev.*, 16, 3501-3534, <https://doi.org/10.5194/gmd-16-3501-2023>, 2023.](#)
- 1525 Xu, X. B., Chang, Y. S., Peters, H., Ozgokmen, T. M., and Chassignet, E. P.: Parameterization of gravity current entrainment for ocean circulation models using a high-order 3D nonhydrostatic spectral element model, *Ocean Model.*, 14, 19-44, <https://doi.org/10.1016/j.ocemod.2006.02.006>, 2006.
- Yamanaka, Y. and Tajika, E.: The role of the vertical fluxes of particulate organic matter and calcite in the oceanic carbon cycle: Studies using an ocean biogeochemical general circulation model, *Global Biogeochem. Cy.*, 10, 361-382, <https://doi.org/10.1029/96gb00634>, 1996.
- 1530 Zhang, J., Quay, P. D., and Wilbur, D. O.: Carbon isotope fractionation during gas-water exchange and dissolution of CO₂, *Geochim. Cosmochim. Ac.*, 59, 107-114, [https://doi.org/10.1016/0016-7037\(95\)91550-D](https://doi.org/10.1016/0016-7037(95)91550-D), 1995.
- [Zelinka, M. D., Myers, T. A., McCoy, D. T., Po-Chedley, S., Caldwell, P. M., Ceppi, P., Klein, S. A., and Taylor, K. E.: Causes of higher climate sensitivity in CMIP6 models, *Geophys. Res. Lett.*, 47, e2019GL085782, <https://doi.org/10.1029/2019gl085782>, 2020.](#)
- 1535 Zickfeld, K., Eby, M., Weaver, A. J., Alexander, K., Cresspin, E., Edwards, N. R., Eliseev, A. V., Feulner, G., Fichefet, T., Forest, C. E., Friedlingstein, P., Goosse, H., Holden, P. B., Joos, F., Kawamiya, M., Kicklighter, D., Kienert, H., Matsumoto, K., Mokhov, I. I., Monier, E., Olsen, S. M., Pedersen, J. O. P., Perrette, M., Philippon-Berthier, G., Ridgwell, A., Schlosser, A., Schneider Von Deimling, T., Shaffer, G., Sokolov, A., Spahni, R., Steinacher, M., Tachiiri, K., Tokos, K. S., Yoshimori, M., Zeng, N., and Zhao, F.: Long-Term Climate Change Commitment and Reversibility: An EMIC Intercomparison, *J. Climate*, 26, 5782-5809, <https://doi.org/10.1175/jcli-d-12-00584.1>, 2013.
- [Zondervan, J. R., Hilton, R. G., Dellinger, M., Clubb, F. J., Roylands, T., and Ogrič, M.: Rock organic carbon oxidation CO₂ release offsets silicate weathering sink, *Nature*, 623, 329-333, <https://doi.org/10.1038/s41586-023-06581-9>, 2023.](#)
- 1545