

**The Ross Sea and Amundsen Sea Ice-Sea Model (RAISE v1.0): a high-resolution ocean-sea ice-ice shelf coupling model for simulating the Dense Shelf Water and Antarctic Bottom Water in the Ross Sea, Antarctica**

5 Zhaoru Zhang<sup>1,2,3,4</sup>, Chuan Xie<sup>1</sup>, Chuning Wang<sup>1</sup>, Yuanjie Chen<sup>1</sup>, Heng Hu<sup>1</sup>, Xiaoqiao Wang<sup>1,5</sup>

<sup>1</sup>Key Laboratory of Polar Ecosystem and Climate Change, Ministry of Education and School of Oceanography, Shanghai Jiao Tong University, 1954 Huashan Road, Shanghai, 200030, China

<sup>2</sup>Shanghai Key Laboratory of Polar Life and Environment Sciences, Shanghai Jiao Tong University,  
10 Shanghai, China

<sup>3</sup>Shanghai Frontiers Science Center of Polar Science, Shanghai Jiao Tong University, 1954 Huashan Road, Shanghai, 200030, China

<sup>4</sup>Key Laboratory for Polar Science, Polar Research Institute of China, Ministry of Natural Resources, Shanghai, 200136, China

15 <sup>5</sup>High Impact Weather Key Laboratory of China Meteorological Administration (CMA), Changsha, 410073, China

*Correspondence to:* Zhaoru Zhang (zrzhang@sjtu.edu.cn)

## 20 **Abstract**

The Ross Sea in the Southern Ocean is a key region for the formation of the Antarctic Bottom Water (AABW) that supplies the lower limb of the global overturning circulation, and contributes to 20–40% of the total AABW production. AABW primarily originates from polynyas characterized by strong sea ice production and ocean convection that lead to the formation of Dense Shelf Water (DSW), the precursor of the AABW. The production and characteristics of DSW in the Ross Sea and AABW in the surrounding ocean are significantly affected by ice shelf meltwater transported from the nearby Amundsen Sea. The scarcity of long-term observations in the Ross Sea hinders the understanding of DSW and AABW variability, and numerical models are needed to explore the multi-scale variations of these water masses and the forcing mechanisms. In this work, a coupled high-resolution ocean-sea ice-ice shelf model is developed for the Ross Sea and Amundsen Sea, named RAISE (**R**oss-**A**mundsen Sea **I**ce-**S**ea Model). Detailed descriptions of the model configurations are provided. This study represents an attempt to thoroughly evaluate the DSW properties and associated ocean-sea ice-ice shelf coupling processes among modelling studies in the Southern Ocean, using multiple datasets including satellite-based observations and hydrographic measurements from the World Ocean Database, Argo profilers and seal-tag sensors. In particular, the modelled temporal variations of DSW properties in polynyas and its key export passages are compared with long-term mooring observations, which are rarely seen in studies of the DSW temporal variability before. RAISE demonstrates high skills in simulating the observed sea ice production rates in the Ross Sea polynyas, and the modelled spatial and temporal variability of DSW are significantly and strongly correlated with observations. RAISE can also effectively capture the observed long-term freshening trend of DSW prior to 2014 and the rebounding of DSW salinity after 2014. RAISE


shows an overestimate of DSW density in the Ross Sea, which is associated with an underestimate of ice shelf melting rates in the Amundsen Sea, missing ice shelf calving processes and subglacial discharge in the model. A sensitivity experiment simulating increased freshwater discharge from these processes can significantly improve the simulation of DSW properties.

## 1 Introduction

The Southern Ocean is the production site of bottom water mass in the global ocean — the Antarctic Bottom Water (AABW), which supplies the lower limb of the global thermohaline circulation. AABW originates predominantly from polynyas on the Antarctic continental shelves or in open ocean regions, defined as low sea ice concentration areas that allow large ocean-atmosphere heat fluxes in the cold seasons and thus significant new ice production. Brine rejection during ice formation drives deep ocean convection and the formation of the Dense Shelf Water (DSW) in the polynya regions. DSW is subsequently transported across the continental slope and into the open ocean, entraining other water masses such as the Ice Shelf Water (ISW) and the Circumpolar Deep Water (CDW) and finally forming AABW.

The Ross Sea is a key region for the formation of DSW and contributes to 20–40% of the global AABW production (Meredith et al., 2013; Solodoch et al., 2022). DSW is primarily formed in two coastal polynyas, the Terra Nova Bay polynya (TNBP) off the Victoria Land and the Ross Ice Shelf polynya (RISP, also called the Ross Sea polynya) off the largest ice shelf in Antarctica (Fig. 1). Intense katabatic winds blowing from the Transantarctic Mountains or the ice shelf toward the ocean drive the formation of these polynyas, enhancing sea ice production and the DSW formation. DSW is then transported along three deep troughs on the Ross Sea shelf — the Drygalski Trough, the Joides Trough and the Glomar Challenger Trough (Fig. 1) to the slope, where it sinks down the slope to the ocean bottom or is carried to East Antarctic regions by the westward Antarctic Slope Current, forming AABW in the Pacific sector of the Southern Ocean. The formation and properties of DSW in the Ross Sea are significantly affected by freshwater input from the Amundsen Sea, originating from the melting of ice shelves in the Amundson

Sea (Kusahara and Hasumi, 2013; Nakayama et al., 2014). Under surface warming and enhanced on-shelf intrusion of the warm CDW, there has been accelerated melting of ice shelves in the Amundsen Sea, and the increased meltwater transport into the Ross Sea causes a freshening trend of DSW in the Ross Sea over the past few decades (Jacobs et al., 2022). Such trend is found to be reversed in recent years due to interactions between major climate modes, leading to changes in winds and sea ice exchange between these seas (Castagno et al., 2019; Silvano et al., 2020; Guo et al., 2020).



**Figure 1. (a)** Map of the Ross Sea and the Amundsen Sea. The bathymetric contours are shown as thin grey lines. Grey shading indicates ice shelves. The movement of dense shelf water (DSW) and Antarctic Bottom Water (AABW) are illustrated by blue arrows, and the movement of circumpolar deep water (CDW) is illustrated by red arrows. The Antarctic Slope Current and coastal current are illustrated by thick grey arrows. TNBP denotes the Terra Nova Bay polynya, and RISP denotes the Ross Ice Shelf polynya. Color indicates the climatological annual-accumulative sea ice production from satellite estimates based on the AMSR-E data by Nakata et al. (2021). **(b)** The RAISE model domain and grid. Grey color indicates ice shelves including the Ross Ice Shelf, the Getz Ice Shelf (Getz), the Dotson Ice Shelf (Dot), the Crosson Ice Shelf (Cro), the Thwaites Ice Shelf (Thw), the Cosgrove Ice Shelf (Cos) and the Pine Island Ice Shelf (Pine). The yellow line denotes the cross-shore transect from the MEOP dataset. The red dots indicate mooring locations in the Terra Nova Bay and on the Ross Sea slope near Cape Adare, the magenta dot indicates the long-term observational site near the Ross Island mentioned in Jacobs et al. (2022), and the blue dots indicate locations in the three troughs of the Ross Sea for examining the long-term variations of DSW.

In recent years, several ocean-sea ice-ice shelf coupling models have been developed for the Ross Sea. Dinniman et al. (2018) developed a 5-km coupled ocean-sea ice-ice shelf model for the Ross Sea using the Regional Ocean Modelling System (ROMS), on the basis of an ocean circulation model (Dinniman et al., 2004) and a coupled ocean-ice shelf model (Dinniman et al., 2007; Dinniman et al., 2011). This model is employed to study the future changes of atmospheric forcings and freshwater inflow on the formation of DSW, the on-shelf intrusion of CDW and basal melting of the Ross Ice Shelf; changes in the freshwater inflow from the Amundsen Sea are simulated by reducing salinity at the eastern and western boundaries of the Ross Sea. Yan et al. (2023) developed a coupled ocean-sea ice-ice shelf model for the Ross Sea based on the Massachusetts Institute of Technology General Circulation Model (MITgcm), also with a horizontal resolution of approximately 5 km. This model is used to analyze the seasonality of salinity budget to understand the controlling mechanisms for the bottom water variations.

As DSW formation is significantly affected by the meltwater inflow from the Amundsen Sea ice shelves,  
100 to achieve realistic simulations of the DSW features, it would be necessary to include processes in the  
Amundsen Sea and directly simulate the melting of ice shelves and the impacts of these meltwater on the  
Ross Sea shelf and slope environments.

Given the concerns above, in this work, we developed a coupled ocean-sea ice-ice shelf high-  
resolution model covering both the Ross Sea and the Amundsen Sea, named as RAISE (**R**oss-**A**mundsen  
105 **S**ea **I**ce-**S**ea Model). Compared to earlier modelling work, this study for the first time provides  
comprehensive evaluations of the coupled model performance on simulating the DSW properties,  
particularly its temporal variability, in the Ross Sea with cruise and mooring observations; the associated  
sea ice production rates are also assessed using satellite-retrieved datasets. It is found that the model can  
well capture the variations of sea ice production rates in the Ross Sea polynyas, the long-term freshening  
110 trend of DSW as well as the rebounding of salinity after 2014 that is revealed by observations. DSW  
variations at higher frequencies in the DSW formation sites and major export passages on the slope are  
also well represented by the model compared to mooring observations.

## **2 Model setup**

### 115 **2.1 The ocean model**


The ocean model of RAISE is an implementation of the Regional Ocean Modelling System  
(ROMS), which is a primitive-equation, finite-volume model with a terrain-following vertical coordinate  
system (Haidvogel et al. 2008; Shchepetkin and McWilliams 2009). The model domain covers the Ross  
Sea, the Amundsen Sea and the adjacent open ocean in the Pacific sector of the Southern Ocean (Fig. 1b).

120 This model is an updated version of the one used in Xie et al. (2024) and Zhang et al. (2024a), and the major differences among these models are the application of nudging for temperature and salinity, as will be explained later in this section. The model horizontal resolution varies from ~2 km in the coastal areas to ~6 km in the open ocean. This model includes 32 vertical layers, with variable thicknesses that depend on water column depth and are smaller at the surface and bottom. On the Ross Sea shelf and slope, the  
125 thickness of the bottom layer varies from 10 m over banks to 60 m in the Drygalski Trough. In the open ocean, the bottom layer thickness varies from 100 m to 200 m. The model bathymetry and ice shelf draft are interpolated from BedMachine-Antarctica-v2.0 (Morlighem et al. 2020), which has a spatial resolution of 500 m on the Antarctic Polar Stereographic projection. Vertical mixing of momentum and tracers are computed using the K-profile parameterization (KPP) mixing scheme (Large et al. 1994).

130 Initial conditions of temperature and salinity come from simulations from a circum-Antarctic ocean-sea ice-ice shelf model with a horizontal resolution of 10 km (Dinniman et al. 2015). Alternative initial conditions from the World Ocean Atlas 2018 (WOA18) are also employed for this model, and we found that after a 5-year spin-up period, these conditions yield quite similar model simulations to those initialized by the model results from Dinniman et al. (2015). Temperature, salinity, sea surface height and  
135 depth-averaged velocity for the open boundaries are derived from daily mean data of the Met Office Global Seasonal forecasting system version 5 (GloSea5) (Maclachlan et al. 2015). Hydrographic simulations from a couple of global ocean-sea ice reanalysis products are compared with the EN4 dataset for the Ross Sea, the Amundsen Sea and the nearby open ocean, including C-GLORS, GECCO3, GLORYS12V1, ORAS5 and GloSea5, and it is found that GloSea5 has the overall best performance in  
140 simulating temperature and salinity in this region (Fig. 2). Below 1000 m (the isobath at the shelf break),



the average root mean square errors of temperature and salinity over the model domain relative to the EN4 dataset (Good et al., 2013) are 0.165 °C and 0.054 psu, respectively. Tidal forcing is derived from the global tidal solution TPXO9 (Egbert and Erofeeva 2002), including 15 major tidal constituents (K1, S2, M4, P1, O1, Q1, S1, MS4, MN4, MF, 2N2, M2, K2, MM and N2) forced at the open boundaries via  
145 sea surface height and barotropic velocity. Atmospheric forcing fields for the model are obtained from the ERA5 reanalysis product, including 3-hourly data for surface wind and air temperature, and daily data for sea level pressure, humidity, cloudiness and precipitation. Compared to the model used in Xie et al. (2024) and Zhang et al. (2024a), surface temperature and salinity are nudged to a monthly mean climatology in this model, provided by the WOA18 dataset (Locarnini et al., 2018). Due to the limited  
150 observational data in the Antarctic shelf regions in WOA18, which are mostly collected during the summer, surface nudging is only applied to the off-shelf regions. Such nudging results in improved simulations of sea ice production and DSW properties. (Figs. S2 and S3 in the Supplementary Information)



**Figure 2.** Vertical profiles of root mean square errors of **(a)** temperature and **(b)** salinity from C-GLORS, GECCO3, 155 GLORYS12V1, GloSea5 and ORAS5 relative to the EN4 dataset.

Numerical dyes are released in the model to trace the movement and distributions of major water masses, including the DSW, CDW and ISW. The dyes are released continuously during the simulation periods of the experiments. DSW dyes are released at model grid points in the polynya areas where sea 160 ice production occurs, and the dye values are proportional to the ice production rates. CDW dyes are released at grid points in the open ocean (offshore of the 1000-m isobath) where temperatures are greater than 0°C, with the initial dye values set as 100. ISW dyes are released at grid points where ice shelf exists, and the dye values are proportional to the ice shelf basal melting rates.

## 2.2 The sea ice module

165 The sea ice module (Budgell 2005) of RAISE is based on two-layer ice thermodynamics and a molecular sublayer beneath the sea ice described by Mellor and Kantha (1989) and Häkkinen and Mellor (1992), and elastic-viscous-plastic rheology for ice dynamics (Hunke and Dukowicz, 1997; Hunke, 2001). A snow layer is included, and snow is converted into ice when the snow-ice interface is below sea level. The sea-ice model also includes a simple estimate of frazil ice production (Steele et al. 1989). Boundary 170 conditions of sea ice concentration are obtained from daily data of the Advanced Microwave Scanning Radiometer-Earth Observing System (AMSR-E) and the Advanced Microwave Scanning Radiometer 2 (AMSR2) datasets, provided by the University of Bremen using the ARTIST sea ice algorithm (Spreen et al. 2008).

## 2.3 The ice shelf module

175 The ice shelves in the model are static, and there are no thickness or extent changes of an ice shelf over time. Configurations of the ice shelf module follow those in Dinniman et al. (2007) and Dinniman et al. (2011). The hydrostatic pressure at the base of the ice shelf is computed based on the assumption that ice is in isostatic equilibrium. Friction between the ice shelf and the water is computed as a quadratic stress, and is applied as a body force over the top three ocean layers beneath the ice shelf. At the interface  
 180 between the ocean and ice shelf, a parameterization scheme with a viscous sublayer model is used with three equations representing the conservation of heat across the ocean-ice shelf boundary, the conservation of salt and a linearized version of the freezing point of sea water as a function of salinity and pressure (Holland and Jenkins, 1999). The conservation of heat across the ocean-ice shelf boundary is expressed as:

$$185 \quad \frac{\rho_I w_B L_f}{C_{pw}} = -\rho_W \gamma_T (T_B - T_W), \quad (1)$$

where  $\rho_I$  is the ice density and specified as  $930 \text{ kg m}^{-3}$ ,  $L_f$  is the latent heat of fusion of ice specified as  $3.34 \times 10^5 \text{ J kg}^{-1}$ ,  $C_{pw}$  is the specific heat capacity of seawater specified as  $4000 \text{ J kg}^{-1} \text{ }^\circ\text{C}^{-1}$ ,  $\rho_W$  is the seawater density of the uppermost ocean layer ( $\text{kg m}^{-3}$ ),  $w_B$  denotes the rate of ice melting ( $> 0$ ) or freezing ( $< 0$ ) ( $\text{m s}^{-1}$ ),  $T_B$  is the interface temperature (freezing point),  $T_W$  is the temperature of seawater  
 190 at a certain distance from the ocean-ice shelf interface, and  $\gamma_T$  is the heat transfer coefficient ( $\text{m s}^{-1}$ ) representing the molecular and turbulent mixing coefficient of heat within the ocean boundary layer adjacent to the ice shelf. The conservation of salt across the ocean-ice shelf boundary is expressed as:

$$\rho_I w_B S_B = \rho_W \gamma_S (S_B - S_W), \quad (2)$$

where  $S_B$  is the salinity at the ocean-ice shelf interface,  $S_W$  represents the salinity of the uppermost ocean  
 195 grid cell in the model and  $\gamma_S$  is the salt transfer coefficient ( $\text{m s}^{-1}$ ).  $\gamma_T$  and  $\gamma_S$  are specified following

McPhee et al. (1987) that assumes a viscous molecular sub-layer adjacent to the ice–ocean boundary. The last equation is a linearized version of the equation for the freezing point of sea water, which is written as:

$$T_B = aS_B + bP_B + c, \quad (3)$$

200 where the salinity coefficient  $a$  is specified as  $-5.7 \times 10^{-2} \text{ }^\circ\text{C}$ , the pressure coefficient  $b$  is specified as  $-7.61 \times 10^{-4} \text{ }^\circ\text{C dbar}^{-1}$ , and  $c$  the depth of the ice shelf base. The variables  $w_B$ ,  $T_B$ , and  $S_B$  can be solved by simultaneously solving Equations 1–3.

Using the configurations described in Section 2.1–2.3, the model is integrated from 2003 to 2019 starting from a 5-year spin-up simulation, and the model simulation is referred to as the CTRL simulation. 205 A sensitivity experiment Melt+ is conducted in which the basal melting rates of ice shelves in the Amundsen Sea are increased by tuning the heat and salt transfer coefficients ( $\gamma_T$  and  $\gamma_S$ ) in Equations 1 and 2 (see the details in Section 4.5), in order to explore the effects of underrepresented ice shelf melting and unrepresented ice shelf calving on freshwater fluxes and thus the DSW formation; detail information of this experiment is provided in Section 4.5.

210

### 3 Validation datasets and methodology

#### 3.1 Ocean

For the validations of hydrographic properties and water masses in the study region including the DSW, we use hydrography data of the World Ocean Database (WOD) available at the National Centers 215 for Environmental Information (<https://www.ncei.noaa.gov/products/world-ocean-database>) and Argo data provided by the International Argo Program and the national programs that contribute to it

(<https://argo.ucsd.edu>). Hydrographic measurements along a cross-shelf transect (Fig. 1b) from the Marine Mammals Exploring the Oceans Pole to Pole (MEOP) elephant-seal data (Treasure et al. 2017) are also used to assess the simulated water masses in the Ross Sea. Salinity data collected near the Ross  
220 Island during summer (December to February) from Jacobs et al. (2022) are employed to evaluate the interannual variation and trend of the DSW salinity from 2003 to 2019. High-frequency variability of DSW formed in the Terra Nova Bay polynya is evaluated by hydrography measurements from a mooring (Fig. 1b) deployed by the MORSea Project of the Italian National Research Antarctic Program for 2008–  
2016. High-frequency variability of DSW on the slope is evaluated by measurements from two moorings  
225 (Fig. 1b) deployed by the U.S. Cape Adare Long-term Mooring (CALM) program for 2008–2011.

### 3.2 Sea ice and ice shelf


Model simulations of sea ice concentration are compared with the AMSR-E and AMSR2 datasets provided by the University of Bremen, available as daily data with a horizontal resolution of 6.25 km (Spren et al., 2008). The simulations of sea ice production (SIP) are evaluated against the satellite-  
230 retrieved SIP dataset provided by the Institute of Low Temperature Science at the Hokkaido University ([http://www.lowtem.hokudai.ac.jp/wwwod/polar-seaflux/southern\\_ocean\\_new/AMSR-POLAR/](http://www.lowtem.hokudai.ac.jp/wwwod/polar-seaflux/southern_ocean_new/AMSR-POLAR/)), which are calculated using the AMSR-E data for sea ice concentration and the ERA5 data for heat fluxes. This dataset includes estimates of frazil ice production (Nakata et al., 2019; Nakata et al., 2021). The data are provided monthly on the polar stereographic grid with a spatial resolution of 6.25 km, and are available  
235 in the period of 2003–2010. For the assessment of temporal variations of SIP, we use a SIP product spanning a longer period, i.e. from 1992 to 2013 as described in Tamura et al. (2016). For this product, sea ice production is estimated by the heat flux calculation using the SSM/I passive microwave data and

atmospheric reanalysis products including ERA-40, ERA-interim and NCEP24; frazil ice is not included in this dataset. Melting rates of ice shelves in the Amundsen Sea are compared with the estimates from  
 240 Rignot et al. (2013), Depoorter et al. (2013) and Liu et al. (2015) based on satellite observations.

## 4 Simulation results

### 4.1 Sea ice concentration and production


The modelled spatial distributions of SIC during the ice freezing seasons are shown in Fig. 3,  
 245 assessed against SIC distributions derived from the AMSR-E/AMSR2 products. Compared with the satellite data (Fig. 3a–c), the RAISE model overall underestimates SIC over most of the model domain, which is attributed to high temperature bias in the surface ocean of the model as will be presented in Section 4.2. Correspondingly, the model underestimates the sea ice extent, but demonstrates good skill in capturing the temporal variability of the ice extent (Fig. 3g). Correlation between the modelled and  
 250 satellite-derived temporal variations of sea ice extent anomalies reaches 0.68 ( $P < 0.001$ ).



**Figure 3.** Seasonal mean sea ice concentration from (a–c) the AMSR product and (d–f) the RAISE model simulation for austral (a, d) autumn, (b, e) winter and (c, f) spring averaged over 2003–2019. (g) Time series of sea ice area anomalies from the AMSR-E/AMSR2 and from the RAISE simulation. The correlation coefficient (R) and p-value (P) between these time series are provided.

Sea ice production is the determinant factor for the DSW formation in the Southern Ocean. Comparisons between the modelled and satellite-estimated annual accumulative SIP rates (over the freezing seasons March to October) averaged over 2003–2010 are shown in Fig. 4. It can be seen that the model can well simulate locations and shape of the two major formation sites of DSW in the Ross Sea, the TNBP and RISP, while the simulated SIP rates are higher than satellite estimates by 14.9 km<sup>3</sup> and 236 km<sup>3</sup> for the TNBP and RISP area-mean values, respectively. Such differences are on the one hand due to inadequate representations of sea ice thermodynamic and dynamic processes in the model, which lead to an overestimate of sea ice thickness in the model in the polynya areas (Fig. S1 in the Supplementary Information). On the other hand, there are also estimation errors in the satellite products. For example, satellite estimation does not include oceanic heat fluxes, and based on observed vertical temperature profiles in the Terra Nova Bay by Thompson et al. (2020), temperature in the subsurface layer in this region is lower than the surface layer, and hence there would be more sea ice production if the vertical oceanic heat fluxes are considered in the satellite retrieval algorithms. Interannual variations of the modelled annual accumulative SIP rates are significantly correlated with those from satellite estimates for both the TNBP (R=0.62, P=0.04) and RISP (R=0.55, P=0.08), demonstrating that the model can well simulate the temporal variability of ice production in the Ross Sea polynyas. Previous studies demonstrated that DSW primarily exists in the western portion of the RISP and not in the eastern portion (Orsi and Wiederwhol, 2009; Wang et al., 2021), as a result of more ice shelf meltwater from the

Amundsen Sea transported to the eastern portion and more local meltwater from the Ross Ice Shelf in the  
 275 eastern portion. Therefore, sea ice production in the western portion of the RISP contributes most to the  
 DSW production in this polynya. We compared the interannual variability of modelled sea ice production  
 in the western RISP (west of 186°E) to that from the satellite estimate, and the correlation is significantly  
 improved ( $R=0.76$ ,  $P=0.01$ ).




280 **Figure 4.** (a) Cumulative sea ice production in March–October averaged over 2003–2010 from satellite estimates.  
 (b) The simulated cumulative sea ice production in March–October averaged over 2003–2019. (c) The scatter plot  
 of cumulative sea ice production (unit:  $\text{km}^3$ ) averaged over the Terra Nova Bay polynya from the model simulation  
 versus that from satellite estimate. (d) Same as (c) but for the Ross Ice Shelf polynya. (e) Same as (d) but for the  
 western Ross Ice Shelf polynya. The correlation coefficients ( $R$ ) and  $p$ -values ( $P$ ) between modelled and satellite-  
 285 retrieved data are provided for (c), (d) and (e).



## 4.2 Hydrography and water masses

The temperature-salinity diagrams of Ross Sea waters below the depth of 100 m from WOD and RAISE are shown in Fig. 5. The model can well depict the distributions of major water masses in the subsurface and bottom layers of the Ross Sea, including the salty DSW, the warm CDW and the cold ISW that are all components of the AABW. Compared to WOD, there is some deficiency for the model to capture the high-salinity ends (34.9–35 g kg<sup>-1</sup>) of DSW and high-temperature ends (1.5–1.7 °C) of




**Figure 5.** Temperature-salinity diagrams from (a) the WOD dataset and (b) the RAISE model simulation over the model domain. The major water masses are labeled, including the Dense Shelf Water (DSW), Circumpolar Deep Water (CDW), Ice Shelf Water (ISW) and Winter Water (WW).

CDW. Model simulations of potential temperature and salinity are also compared with seal-tag CTD measurements from MEOP on a transect across the Ross Sea and the adjacent open ocean (Fig. 1b). The observed spatial structures of temperature and salinity are well captured by the model (Fig. 6), and the model also performs well in simulating the on-shelf intrusion of warm CDW and the distribution of dense DSW (defined as neutral density  $\gamma^n > 28.27 \text{ kg m}^{-3}$ ). Compared to observations, the model slightly

overestimates temperature and underestimates salinity in the surface layer. In the subsurface layer, the model has lower temperature in the open ocean and higher temperature on the shelf, indicating stronger CDW intrusion in the model relative to the observational data; the subsurface salinity is underestimated

305 in the model. The warm bias in the surface layer of the model is responsible for the underestimation of sea ice concentration in the model as shown in Fig. 3.




**Figure 6.** Vertical sections of (a, c) potential temperature and (b, d) salinity from (a, b) the MEOP data and (c, d) the RAISE model simulation along the cross-shore transect in the Ross Sea and open ocean (shown in Fig. 1b).

310 Contours indicate isolines of neutral density.

The climatological (2003–2019 average) model simulations of potential temperature and salinity in the bottom 100-m layer (the layer mainly composed of DSW or AABW) on the Ross Sea shelf and slope and the adjacent open ocean are evaluated against climatology from the WOD and Argo data (Fig.

315 7). The spatial distributions of modelled temperature and salinity compare well with those from observations (Fig. 7a, c). Linear regression results reveal that the spatial correlation between the modelled and observed potential temperature can reach 0.91 (Fig. 7b), with a slope value of 0.86. The correlation between modelled and observed salinity is 0.80 (Fig. 7d), with a slope value slightly lower than that for temperature and approaching 0.7. These results suggest that the climatological spatial patterns of DSW  
320 hydrography in the Ross Sea and AABW hydrography in the open ocean can be well represented by the model. It is noted that the model overestimates salinity in the lower-salinity range and underestimates salinity in the higher-salinity range (Fig. 7d). This means while the model produces larger salinity in the polynyas compared to observations, in other regions featured by high salinities there can be underestimates of salinity by the model.




325

**Figure 7.** Comparisons of modelled climatological fields with historical hydrographic observations. **(a, c)** Modelled climatological potential temperature and salinity fields in the bottom 100-m layer, overlaid with historical observations (circled points). Historical data include all CTD, XBT, MBT, drifting buoy, glider and ocean station profiling measurements from WOD and Argo data. **(b, d)** Scatter plots of modelled and observed potential temperature and salinity in the Ross Sea. Black dashed lines denote 1:1 ratio lines, and gray dashed lines denote linear regression fits. Colors of points denote the location of measurements; the three regions (shelf/shelf break/ocean) are separated by the 700-m and 3000-m isolines on the shelf break (marked by dashed lines in a, c).

The intrusion of warm CDW is a major mechanism for causing ice shelf basal melting and generating the ISW, which subsequently affects the DSW characteristics. The CDW dyes are initially released at model grid points in the open ocean where water temperature is above 0°C. As seen in Fig. 8a that shows the CDW dye values 5 years after release at the 15<sup>th</sup> model level (200–400 m), CDW mainly intrudes onto the continental shelves via troughs and spreads over the shelf regions. High dye values are also present beneath the Ross Ice Shelf north of 80°S in the eastern Ross Sea, while low dye values reach much further south (to 82°S) beneath the ice shelf in the western Ross Sea, which could result from a southward flow that has been reported in earlier studies (Budillon et al., 2003; Jendersie et al., 2018; Stewart et al., 2019). The presence of CDW under the ice shelf is essential for its basal melting rate. Figure 8b and 8c show the distributions of vertically integrated values of ISW dye originating from ice shelves in the Ross Sea and Amundsen Sea, respectively. Compared to the western portion of the RIS, there is more ISW beneath the eastern portion, indicating stronger influence on the hydrography over the Ross Sea shelf. The ISW dye values are much higher in the Amundsen Sea and decrease dramatically towards the Ross Sea, indicating strong basal melting of ice shelves in the former region that provides fresh meltwater input to the Ross Sea, which plays a more important role in modulating the salinity and

stratification on the Ross Sea shelf compared to the meltwater released from the RIS. The transport time of ISW dyes from the Amundsen Sea to the Ross Sea is about 2 years.




350

**Figure 8.** Values of (a) CDW dyes, (b) ISW dyes originating from the Ross Ice Shelf, and (c) ISW dyes from the Amundsen Sea ice shelves 5 years after the release time of these dyes in the model simulation.

In Fig. 9, DSW dyes are released separately for the TNBP and RISP. We can see that DSW formed  
 355 in the TNBP (Fig. 9a) is mainly transported to the slope via the Drygalski Trough, and DSW formed in  
 the RISP (Fig. 9b) is transported mainly via the Joides Trough and the Glamor Challenger Trough, while  
 a portion also flows to the slope via the Drygalski Trough. Exports of DSW through the Drygalski Trough,  
 Joides Trough and Clamor Challenger Trough contribute to the total DSW export by 41%, 14% and 45%,


respectively. Once crossing the slope and reaching the deep ocean, DSW turns into AABW and is mainly  
360 transported westward toward the Indian sector of the Southern Ocean by the Antarctic Slope Current.  
DSW dyes released in the RISP cover a larger area in the open ocean than those released in the TNBP.  
DSW formed in the western portion of RISP is also carried southward beneath the Ross Ice Shelf, and  
can reach as far as 84°S near the grounding line of the ice shelf, which can be associated with the  
southward flow as mentioned above as well as the role of tidal currents (Arzeno et al., 2014). Such  
365 intrusion can be important for the basal melting of the Ross Ice Shelf, which is categorized as “cold-water  
cavity” where the DSW acts as the main thermal forcing (Rignot et al., 2013; Adusumilli et al., 2020).



**Figure 9.** Vertically integrated values of DSW dyes originating from the (a) Terra Nova Bay polynya (TNBP)  
and (b) the Ross Ice Shelf polynya (RISP) 5 years after the release time in the model simulation. DT, JT and  
370 GCT denote the Drygalski Trough, Joides Trough and Glamor Challenger Trough, respectively.

### 4.3 Temporal variability of DSW

Temporal variations of neutral density in the middle and bottom layer of the TNBP are compared to measurements from mooring observations conducted by the Italian MORSea project (Fig. 10). Both the variations of neutral density at middle (500 m) and bottom depths (1060 m) are examined. While the



375


**Figure 10.** (a) Time series of 5-day-average neutral density at 500 m from the CTRL simulation, the Melt+ simulation and mooring observations in the Terra Nova Bay polynya (TNBP, see the mooring location in Fig. 1b) during 2010–2016. (b) Same as (a) but for neutral density at 1060 m during 2008–2016. (c) Time series of neutral density anomalies at 500 m from the CTRL simulation and mooring observations in the TNBP during 2010–2016. (d) Same as (c) but for neutral density at 1060 m during 2008–2016. The coefficient of correlation (R) between neutral density (anomaly) from the CTRL simulation and mooring and the corresponding p-value (P) are provided for each panel.

380


model has an overestimate of density in both layers compared to the mooring observations, which might be related to the model overestimate of sea ice production and inadequate representations of other processes (see the discussions in Section 4.4), the model can well capture the temporal variability of DSW density. The correlations between variations of modelled and observed neutral density at middle and bottom depths reach 0.76 and 0.55, respectively, both of which are significant ( $P < 0.001$ ). Removing the seasonal cycles, correlations between variations of modelled and observed neutral density anomalies at middle and bottom depths are 0.65 and 0.53, respectively ( $P < 0.001$ ). This demonstrates the model can reasonably simulate the temporal variations of ocean-sea ice processes forming DSW at its originating sites. The annual average DSW production rate in the TNBP estimated from the RAISE model simulation is 0.33 Sv, which is between the estimate of 0.28 Sv from Jendersie et al. (2018) using a coupled ocean-sea ice model for the Ross Sea and the estimate of 0.43 Sv from Miller et al. (2024) using observations from a mooring in the TNBP. The estimated annual average DSW production rate in the RISP from the model is 1.23 Sv. The model also performs well in simulating the temporal variations of DSW properties in its key outflow passage on the slope, as shown in Fig. 11. comparisons with observations from two moorings deployed by the U.S. CALM project near Cape Adare (CA1 and CA2 in Fig. 1b) show that correlations between simulated and observed variations of DSW neutral density near the ocean bottom reach 0.75 and above ( $P < 0.001$ ), though there is also overestimate of density in the model, which could be associated with the model bias in the polynya areas. Removing the seasonal cycles, correlations between the simulated and observed variations of DSW neutral density reach 0.70 at 1735 m and 0.63 at 1929 m, both of which are significant ( $P < 0.001$ ). Time series of the estimated DSW outflow fluxes at the exits of the three troughs (across the transects as shown in Fig. 9) for different years in 2003–2019 and



for the multi-year average are provided in Fig. 12. For the Drygalski Trough, the Joides Trough and the  
 405 Glamor Challenger Trough, the transports are relatively strong during February–May, April–July and  
 September–December, respectively; the annual mean outflow fluxes for the three troughs are 0.62 Sv,  
 0.19 Sv and 0.81 Sv, respectively.




**Figure 11.** (a) Time series of 5-day-average neutral density at 1735 m from the CTRL simulation, the Melt+  
 410 simulation, and the CA1 mooring observations at the slope near Cape Adare (see Fig. 1b) during 2008–2011. (b)  
 Same as (a) but for neutral density at 1929 m, and the observations are from the CA2 mooring. (c) Time series of  
 neutral density anomalies at 1735 m from the CTRL simulation and mooring observations at CA1 during 2008–  
 2011. (d) Same as (c) but for neutral density at 1929 m at the CA2 mooring location during 2008–2011. The  
 coefficient of correlation (R) between neutral density from the CTRL simulation and mooring and the  
 415 corresponding p-value (P) are provided for each panel.



**Figure 12.** Time series of the DSW outflow fluxes at the exits of **(a)** the Drygalski Trough, **(b)** the Joides Trough and **(c)** the Glamor Challenger Trough for different years in 2003–2019 (grey thin lines) and the multi-year average (black thick line).

420            The decadal variation of DSW has been the focus of DSW studies in the past ten years, which reveal that DSW in the Ross Sea has shown a freshening trend based on observations (Jacobs et al., 2010; Jacobs et al., 2022), attributed to increased transport of ice shelf meltwater from the Amundsen Sea into the Ross Sea (Nakayama et al., 2014). Recent work found that such trend is reversed since 2014 (Castagno et al., 2019), and explained the reversal by the combined effects of positive phase of the Southern Annular  
425            Mode and extreme El Niño conditions (Silvano et al., 2020) and reduced input of meltwater from the Amundsen Sea (Guo et al., 2020). Comparing the model simulations of DSW salinity from 2003 to 2019 with observational data near the Ross Island from Jacobs et al. (2022), we found that the RAISE model can capture the freshening trend prior to 2014 as well as the reversal of freshening after 2014 quite well

(Fig. 13a); the interannual variation of modelled and observed DSW salinity is significantly correlated (R=0.66, and P=0.004). The decadal variations of bottom water salinity are also examined for four locations in the three troughs for DSW exports and the TNBP, which all show freshening trend of DSW before 2014 and rebounding of salinity after 2014 (Fig. 13b–e). The estimated freshening trend prior to 2014 varies from -0.008 PSU yr<sup>-1</sup> to -0.004 PSU yr<sup>-1</sup> at the four locations, which falls in the range of -0.08 to -0.01 PSU per decade as estimated by Castagno et al. (2019) based on long-term cruise observations.




**Figure 13.** (a) Time series of summer bottom water salinity near the Ross Island from the CTRL simulation, the Melt+ simulation and CTD observations from Jacobs et al. (2022) during 2003–2019. (b–e) Time series of simulated summer bottom water salinity in CTRL at the long-term observation locations in (b) the Drygalski Trough, (c) the Joides Trough, (d) the Glamor Challenger Trough and (e) the Terra Nova Bay during 2003–2019. See the locations in Fig. 1b. The trends for the periods prior to 2014 and after 2014 with the significance test results are labeled in panels b–e.

#### 4.4 Ice shelf melting rates

As the melting of ice shelves in the Amundsen Sea and the Ross Sea can have significant impacts on the salinity and stratification in the Ross Sea, and thus the formation of DSW, the simulated melting rates of ice shelves are evaluated against estimates based on satellite data (Fig. 14). In the Ross Sea, the model simulated melting rate for the Ross Ice Shelf averaged over 2003–2019 is about  $79 \text{ Gt yr}^{-1}$ , which is in line with  $47.7 \pm 34 \text{ Gt yr}^{-1}$  as estimated by Rignot et al. (2013) from remote sensing, while higher than the estimates of  $34 \pm 25 \text{ Gt yr}^{-1}$  by Depoorter et al. (2013) and  $27 \pm 22 \text{ Gt yr}^{-1}$  by Liu et al. (2015). In the Amundsen Sea, the simulated melting rate of the Getz Ice Shelf is higher than the satellite estimates from the studies above by  $58\text{--}71 \text{ Gt yr}^{-1}$ , while for the Dotson, Crosson, Thwaites and Pine Island ice shelves, the simulated melting rates are significantly lower than all satellite estimates. In total, in the Amundsen Sea the RAISE model underestimates the ice shelf melting rates by  $107\text{--}172 \text{ Gt yr}^{-1}$  compared with the satellite-retrieved values. Such underestimates are largely attributed to the absence of subglacial runoff in the RAISE model, which is demonstrated to impose dramatic effects on basal melting of Antarctic ice shelves (Nakayama et al., 2021; Goldberg et al., 2023; Gwyther et al., 2023). In addition to the underestimated ice shelf melting rates, the model does not include ice shelf calving as a static ice-shelf module is employed in RAISE; ice shelf calving is not included in the satellite estimates by Rignot

460 et al. (2013) and Depoorter et al. (2013) as well, and is separately considered in Liu et al. (2015) apart from the basal melting process. Liu et al. (2015) suggests that ice calving can contribute to a mass loss of ice shelves of  $270 \pm 22$  Gt yr<sup>-1</sup> in the Amundsen Sea. The inadequate representation of freshwater input related to underestimation of basal melting rates and missing calving processes of ice shelves in the Amundsen Sea could have significant influence on freshwater volume in the Ross Sea, via the westward  
 465 transport of meltwater by the Antarctic Slope Current and coastal currents, and could be one reason for the overestimate of DSW salinity in the Ross Sea by the model. In the next section, a sensitivity experiment is conducted to evaluate the effects of missing freshwater discharge associated with these processes on the Ross Sea DSW characteristics.




470 **Figure 14.** Basal melting rates of ice shelves in the Ross Sea and Amundsen Sea from the RAISE simulation and satellite estimates of earlier studies.

## 4.5 A sensitivity experiment of increasing freshwater discharge associated with ice shelf basal melting and calving

As shown in Figs. 10 and 11, the RAISE simulation shows overestimate of DSW neutral density compared to mooring observations in the Ross Sea polynya and slope regions. A possible reason for such bias is the inadequate representation of freshwater input from underrepresented basal melting rates and missing calving processes of ice shelves in the RAISE model, as discussed in Section 4.4. To test the role of these processes on the Ross Sea DSW properties, we conducted a sensitivity experiment Melt+, in which the basal melting rates of ice shelves in the Amundsen Sea are artificially increased to compensate for the missing freshwater discharge associated with underestimated melting rates, absence of ice shelf calving as well as subglacial runoff. Contributions of these processes to the freshwater discharge sum up to  $\sim 450 \text{ Gt yr}^{-1}$  based on the estimates in Section 4.4, including the contribution from subglacial runoff in the Amundsen Sea estimated as  $\sim 10 \text{ Gt yr}^{-1}$  by Goldberg et al. (2023). Increases of the basal melting rates are achieved by modulating the heat and salt transfer coefficients at the ocean-ice shelf interface following Nakayama et al. (2020). The Melt+ experiment spans the period of 2003 to 2019. Time series of the DSW neutral density in the TNBP and on the Ross Sea slope in Melt+ are presented in Fig. 10 and Fig. 11, respectively, which are substantially lower than those from the CTRL simulation and much closer to the observations. In Melt+, the root-mean-square error of DSW neutral density is reduced by  $0.028 \text{ kg m}^{-3}$  in the TNBP, and reduced by  $0.019 \text{ kg m}^{-3}$  on the Ross Sea slope. The simulation of salinity in the area near the Ross Island in Melt+ is also notably reduced (by  $0.032 \text{ PSU}$ ) compared to that from CTRL (Fig. 13a). There is substantial decrease of salinity over the Ross Sea shelf by  $0.02\text{--}0.1 \text{ PSU}$  (Fig. 15), followed by reduction of DSW thicknesses on the Ross Sea shelf and slope that can reach 100 m and reduction of

AABW thicknesses over the majority of the open ocean by over 100 m. These results suggest that accurate representation of ice shelf melting, calving and subglacial runoff (note that it can also cause ice shelf melting) processes are crucial for accurate simulations of DSW formation and properties in the Ross Sea, which will then affect the simulation of AABW production in the open ocean.



**Figure 15.** (a) Spatial distributions of salinity in the model bottom layer of the Ross Sea averaged over the simulation period. (b) Changes of salinity in the model bottom layer in Melt+ relative to CTRL. (c) Spatial distributions of DSW thicknesses on the Ross Sea shelf and slope and AABW thicknesses in the open ocean in the CTRL simulation. (d) Changes of DSW and AABW thicknesses in the Melt+ simulation relative to the CTRL simulation.

## 5 Conclusions and prospects

In this work, a high-resolution coupled ocean-sea ice-ice shelf model (RAISE) is developed for  
505 the Ross Sea and Amundsen Sea in the Southern Ocean. A major function of this model is to simulate the  
formation of Dense Shelf Water in the Ross Sea and the Antarctic Bottom Water in the open ocean in the  
Pacific sector, which is controlled by sea ice production in coastal polynyas and on the continental shelf  
as well as the discharge of freshwater released from ice shelf melting, which is in turn modulated by the  
intrusion of the warm Circumpolar Deep Water. The RAISE model demonstrates its ability in effectively  
510 simulating the spatial distributions and temporal variations of sea ice production rates compared with  
satellite estimates. The modelled temperature and salinity distributions of DSW in the Ross Sea are in  
fairly good agreement with observations from the combined WOD and Argo data. The simulated temporal  
variations of DSW hydrography in both the Terra Nova Bay polynya and slope region of the Ross Sea  
are significantly and highly correlated with those obtained from mooring measurements. The RAISE  
515 model can also well capture the freshening trend of DSW prior to 2014 and the rebounding of salinity  
after 2014. Compared with satellite estimates, the RAISE model significantly underestimates the melting  
rates of ice shelves in the Amundsen Sea, which may be an important reason for the overestimate of DSW  
density in the Ross Sea. In a sensitivity experiment in which the basal melting rates of ice shelves are  
increased to compensate for the underrepresented ice shelf melting rates and the absence of ice shelf  
520 calving and subglacial runoff processes, the DSW density is notably reduced compared to the CTRL  
simulation and is in better agreement with observations. Such results demonstrate the importance of  
accurate representation of freshwater released from ice shelves for accurately simulating the DSW  
formation and hydrography.



In the future, the model configurations can be further optimized to improve the simulations for  
525 DSW. First, the modelled sea ice production rates in the polynyas are higher than estimates based on  
satellite data. While satellite products cannot be treated as ground truth as a couple of physical processes  
are missing in the retrieval algorithms, such as oceanic heat fluxes, the model may also have incomplete  
or misrepresented processes that lead to an overestimate of sea ice production. The ice-ocean drag or ice-  
atmosphere drag parameterization schemes can be tuned to yield better simulations for sea ice production.  
530 Second, the RAISE model resolution is about 5 km in the slope area, coarser than the baroclinic Rossby  
deformation radius in this region that is suggested to be  $\sim 1$  km (Mack et al. 2019; Stewart and Thompson  
2015). This will result in inadequate representation of CDW on-shelf intrusion associated with eddy  
activities, and may be one of the reasons for inadequate representation of ice shelf basal melting rates in  
the model. The model horizontal resolution can be further enhanced to improve the CDW intrusion  
535 processes. Moreover, subglacial runoff is not included in the RAISE model, and such runoff can on the  
one hand contribute directly to the freshwater fluxes, and on the other hand contribute to ice shelf melting  
and consequently freshwater discharge. The subglacial discharge needs to be included in the model for  
more accurate simulation of freshwater volume and stratification in the Ross Sea. Finally, the RAISE  
model uses a static ice shelf module, and in the future a dynamic ice shelf module should be considered  
540 to include the ice shelf calving processes and their contributions to freshwater fluxes into the Amundsen  
Sea and the Ross Sea.

*Code and data availability.* The source code used for the simulations described here are archived at  
<https://doi.org/10.5281/zenodo.12735787> (Zhang, 2024b). The scripts used to generate the grid and

545 forcing files as well as scripts and data used to generate the figures included in this paper are archived at [https://doi.org/ 10.5281/zenodo.14472621](https://doi.org/10.5281/zenodo.14472621) (Zhang, 2024c). The model output can be obtained from the authors upon request. The ERA5 atmospheric reanalysis data, used for atmospheric forcing files, were collected from the Climate Data Store, available at <https://cds.climate.copernicus.eu>. The GloSea5 reanalysis product, used for the model boundary conditions, were collected from the Copernicus Marine  
550 Data Store, available at <https://data.marine.copernicus.eu>. The BedMachine Antarctica v2 topography data used for the model grid file were collected from the National Snow and Ice Data Center and available at <https://nsidc.org/data/nsidc-0756/versions/2>.

*Author contributions.* ZZ conceptualized the idea of this work, designed the numerical configurations of the RAISE model and wrote the manuscript draft. CX contributed to the implementation of the model  
555 development, numerical simulations and model result validations. CW, YC and XW contributed to the model development. HH performed part of the CTRL simulation. All authors contributed to the writing and editing of this manuscript.

*Competing interests.* The corresponding author has declared that none of the authors has any competing interests.

560 *Acknowledgements.* This work is funded by Key Research & Development Program of the Ministry of Science and Technology of China (Grant No 2022YFC2807601), the National Natural Science Foundation of China (Grant No 42476271), the Shanghai Pilot Program for Basic Research of Shanghai Jiao Tong University (Grant No TQ1400201), the Impact and Response of Antarctic Seas to Climate Change (Grant No IRASCC 1-02-01B), and the Shenlan Program funded by Shanghai Jiao Tong

565 University (Grant No SL2020MS021). We thank Michael S. Dinniman from the Old Dominion University for the support on the model development, and Pasquale Castagno and Giorgio Budillon on providing the mooring data from the Italian MORSea project used for model validation, which was financially and logistically supported by the Italian National Programme for Antarctic Research (PNRA).

## References

- 570 Adusumilli, S., Fricker, H. A., Medley, B., Padman, L., and Siegfried, M. R.: Interannual variations in meltwater input to the Southern Ocean from Antarctic ice shelves, *Nat. Geosci.*, 13, 616–620, <https://doi.org/10.1038/s41561-020-0616-z>, 2020.
- Arzeno, I. B., Beardsley, R. C., Limeburner, R., Owens, B., Padman, L., Springer, S. R., Stewart, C. L., and Williams, M. J. M.: Ocean variability contributing to basal melt rate near the ice front of Ross Ice  
575 Shelf, Antarctica, *Journal of Geophysical Research: Oceans*, 119, 4214–4233, <https://doi.org/10.1002/2014JC009792>, 2014.
- Budgell, W. P.: Numerical simulation of ice-ocean variability in the Barents Sea region: Towards dynamical downscaling, *Ocean Dynamics*, 55, 370–387, <https://doi.org/10.1007/s10236-005-0008-3>, 2005.
- 580 Budillon, G., Pacciaroni, M., Cozzi, S., Rivaro, P., Catalano, G., Ianni, C., and Cantoni, C.: An optimum multiparameter mixing analysis of the shelf waters in the Ross Sea, *Antarctic Science*, 15, 105–118, <https://doi.org/10.1017/S095410200300110X>, 2003.

- Castagno, P., Capozzi, V., DiTullio, G. R., Falco, P., Fusco, G., Rintoul, S. R., Spezie, G., and Budillon, G.: Rebound of shelf water salinity in the Ross Sea, *Nat Commun*, 10, 5441, 585 <https://doi.org/10.1038/s41467-019-13083-8>, 2019.
- Depoorter, M. A., Bamber, J. L., Griggs, J. A., Lenaerts, J. T. M., Ligtenberg, S. R. M., van den Broeke, M. R., and Moholdt, G.: Calving fluxes and basal melt rates of Antarctic ice shelves, *Nature*, 502, 89–92, <https://doi.org/10.1038/nature12567>, 2013.
- Dinniman, M. S. and Klinck, J. M.: A model study of circulation and cross-shelf exchange on the west 590 Antarctic Peninsula continental shelf, *Deep Sea Research Part II: Topical Studies in Oceanography*, 51, 2003–2022, <https://doi.org/10.1016/j.dsr2.2004.07.030>, 2004.
- Dinniman, M. S., Klinck, J. M., and Smith, W. O.: Influence of sea ice cover and icebergs on circulation and water mass formation in a numerical circulation model of the Ross Sea, *Antarctica, J. Geophys. Res.*, 112, 2006JC004036, <https://doi.org/10.1029/2006JC004036>, 2007.
- 595 Dinniman, M. S., Klinck, J. M., and Smith, W. O.: A model study of Circumpolar Deep Water on the West Antarctic Peninsula and Ross Sea continental shelves, *Deep Sea Research Part II: Topical Studies in Oceanography*, 58, 1508–1523, <https://doi.org/10.1016/j.dsr2.2010.11.013>, 2011.
- Dinniman, M. S., Klinck, J. M., Bai, L.-S., Bromwich, D. H., Hines, K. M., and Holland, D. M.: The 600 Effect of Atmospheric Forcing Resolution on Delivery of Ocean Heat to the Antarctic Floating Ice Shelves, *Journal of Climate*, 28, 6067–6085, <https://doi.org/10.1175/JCLI-D-14-00374.1>, 2015.
- Dinniman, M. S., Klinck, J. M., Hofmann, E. E., and Smith, W. O.: Effects of Projected Changes in Wind, Atmospheric Temperature, and Freshwater Inflow on the Ross Sea, *Journal of Climate*, 31, 1619–1635, <https://doi.org/10.1175/JCLI-D-17-0351.1>, 2018.

- Good, S. A., Martin, M. J., and Rayner, N. A.: EN4: quality controlled ocean temperature and salinity  
605 profiles and monthly objective analyses with uncertainty estimates, *Journal of Geophysical Research: Oceans*, 118, 6704–6716, <https://doi.org/10.1002/2013JC009067>, 2013.
- Egbert, G. D. and Erofeeva, S. Y.: Efficient Inverse Modeling of Barotropic Ocean Tides, *Journal of Atmospheric and Oceanic Technology*, 19, 183–204, [https://doi.org/10.1175/1520-0426\(2002\)019<0183:EIMOBO>2.0.CO;2](https://doi.org/10.1175/1520-0426(2002)019<0183:EIMOBO>2.0.CO;2), 2002.
- 610 Goldberg, D. N., Twelves, A. G., Holland, P. R., and Wearing, M. G.: The Non-Local Impacts of Antarctic Subglacial Runoff, *JGR Oceans*, 128, e2023JC019823, <https://doi.org/10.1029/2023JC019823>, 2023.
- Guo, G., Gao, L., and Shi, J.: Modulation of dense shelf water salinity variability in the western Ross Sea associated with the Amundsen Sea Low, *Environ. Res. Lett.*, 16, 014004, <https://doi.org/10.1088/1748-9326/abc995>, 2020.  
615
- Gwyther, D. E., Dow, C. F., Jendersie, S., Gourmelen, N., and Galton-Fenzi, B. K.: Subglacial Freshwater Drainage Increases Simulated Basal Melt of the Totten Ice Shelf, *Geophysical Research Letters*, 50, e2023GL103765, <https://doi.org/10.1029/2023GL103765>, 2023.
- Häkkinen, S. and Mellor, G. L.: Modeling the seasonal variability of a coupled Arctic ice-ocean system,  
620 *Journal of Geophysical Research: Oceans*, 97, 20285–20304, <https://doi.org/10.1029/92JC02037>, 1992.
- Haidvogel, D. B., Arango, H., Budgell, W. P., Cornuelle, B. D., Curchitser, E., Di Lorenzo, E., Fennel, K., Geyer, W. R., Hermann, A. J., Lanerolle, L., Levin, J., McWilliams, J. C., Miller, A. J., Moore, A. M., Powell, T. M., Shchepetkin, A. F., Sherwood, C. R., Signell, R. P., Warner, J. C., and Wilkin, J.:

- 625 Ocean forecasting in terrain-following coordinates: Formulation and skill assessment of the Regional  
Ocean Modeling System, *Journal of Computational Physics*, 227, 3595–3624,  
<https://doi.org/10.1016/j.jcp.2007.06.016>, 2008.
- Holland, D. M. and Jenkins, A.: Modelling thermodynamic ice-ocean interactions at the base of an ice  
shelf, *Journal of Physical Oceanography*, 29, 1787–1800, [https://doi.org/10.1175/1520-](https://doi.org/10.1175/1520-0485(1999)029<1787:MTIOIA>2.0.CO;2)  
630 0485(1999)029<1787:MTIOIA>2.0.CO;2, 1999.
- Hunke, E. C. and Dukowicz, J. K.: An Elastic–Viscous–Plastic Model for Sea Ice Dynamics, *Journal of  
Physical Oceanography*, 27, 1849–1867, [https://doi.org/10.1175/1520-](https://doi.org/10.1175/1520-0485(1997)027<1849:AEVPMF>2.0.CO;2)  
0485(1997)027<1849:AEVPMF>2.0.CO;2, 1997.
- Hunke, E. C.: Viscous–Plastic Sea Ice Dynamics with the EVP Model: Linearization Issues, *Journal of  
635 Computational Physics*, 170, 18–38, <https://doi.org/10.1006/jcph.2001.6710>, 2001.
- Jacobs, S. S. and Giulivi, C. F.: Large Multidecadal Salinity Trends near the Pacific–Antarctic Continental  
Margin, *Journal of Climate*, 23, 4508–4524, <https://doi.org/10.1175/2010JCLI3284.1>, 2010.
- Jacobs, S. S., Giulivi, C. F., and Dutrieux, P.: Persistent Ross Sea Freshening From Imbalance West  
Antarctic Ice Shelf Melting, *JGR Oceans*, 127, e2021JC017808,  
640 <https://doi.org/10.1029/2021JC017808>, 2022.
- Jendersie, S., Williams, M. J. M., Langhorne, P. J., and Robertson, R.: The Density-Driven Winter  
Intensification of the Ross Sea Circulation, *Journal of Geophysical Research: Oceans*, 123, 7702–7724,  
10.1029/2018JC013965, 2018.

- Kusahara, K. and Hasumi, H.: Modeling Antarctic ice shelf responses to future climate changes and  
645 impacts on the ocean, *Journal of Geophysical Research: Oceans*, 118(5), 2054–2475,  
<https://doi.org/10.1002/jgrc.20166>, 2013.
- Large, W. G., McWilliams, J. C., and Doney, S. C.: Oceanic vertical mixing: A review and a model with  
a nonlocal boundary layer parameterization, *Reviews of Geophysics*, 32, 363–403,  
<https://doi.org/10.1029/94RG01872>, 1994.
- 650 Liu, Y., Moore, J. C., Cheng, X., Gladstone, R. M., Bassis, J. N., Liu, H., Wen, J., and Hui, F.: Ocean-  
driven thinning enhances iceberg calving and retreat of Antarctic ice shelves, *Proc. Natl. Acad. Sci.*  
*U.S.A.*, 112, 3263–3268, <https://doi.org/10.1073/pnas.1415137112>, 2015.
- Mack, S. L., Dinniman, M. S., Klinck, J. M., McGillicuddy Jr., D. J., and Padman, L.: Modeling Ocean  
Eddies on Antarctica’s Cold Water Continental Shelves and Their Effects on Ice Shelf Basal Melting,  
655 *Journal of Geophysical Research: Oceans*, 124, 5067–5084, <https://doi.org/10.1029/2018JC014688>,  
2019.
- MacLachlan, C., Arribas, A., Peterson, K. A., Maidens, A., Fereday, D., Scaife, A. A., Gordon, M.,  
Vellinga, M., Williams, A., Comer, R. E., Camp, J., Xavier, P., and Madec, G.: Global Seasonal  
forecast system version 5 (GloSea5): a high-resolution seasonal forecast system, *Quarterly Journal of*  
660 *the Royal Meteorological Society*, 141, 1072–1084, <https://doi.org/10.1002/qj.2396>, 2015.
- McPhee, M. G., Maykut, G. A., and Morison, J. H.: Dynamics and thermodynamics of the ice/upper ocean  
system in the marginal ice zone of the Greenland Sea, *Journal of Geophysical Research: Oceans*, 92,  
7017-7031, <https://doi.org/10.1029/JC092iC07p07017>, 1987.

- Mellor, G. L. and Kantha, L.: An ice-ocean coupled model, *Journal of Geophysical Research: Oceans*, 665 94, 10937–10954, <https://doi.org/10.1029/JC094iC08p10937>, 1989.
- Meredith, M. P.: Replenishing the abyss, *Nature Geosci*, 6, 166–167, <https://doi.org/10.1038/ngeo1743>, 2013.
- Miller, U. K., Zappa, C. J., Gordon, A. L., Yoon, S. T., Stevens, C., and Lee, W. S.: High Salinity Shelf Water production rates in Terra Nova Bay, Ross Sea from high-resolution salinity observations, *Nature* 670 *Communications*, 15, 373, <https://doi.org/10.1038/s41467-023-43880-1>, 2024.
- Morlighem, M., Rignot, E., Binder, T., Blankenship, D., Drews, R., Eagles, G., Eisen, O., Ferraccioli, F., Forsberg, R., Fretwell, P., Goel, V., Greenbaum, J. S., Gudmundsson, H., Guo, J., Helm, V., Hofstede, C., Howat, I., Humbert, A., Jokat, W., Karlsson, N. B., Lee, W. S., Matsuoka, K., Millan, R., Mouginot, J., Paden, J., Pattyn, F., Roberts, J., Rosier, S., Ruppel, A., Seroussi, H., Smith, E. C., Steinhage, D., 675 Sun, B., Broeke, M. R. van den, Ommen, T. D. van, Wessem, M. van, and Young, D. A.: Deep glacial troughs and stabilizing ridges unveiled beneath the margins of the Antarctic ice sheet, *Nature Geoscience*, 13, 132–137, <https://doi.org/10.1038/s41561-019-0510-8>, 2020.
- Nakata, K., Ohshima, K. I., and Nihashi, S.: Estimation of Thin-Ice Thickness and Discrimination of Ice Type From AMSR-E Passive Microwave Data, *IEEE Transactions on Geoscience and Remote Sensing*, 680 57, 263–276, <https://doi.org/10.1109/TGRS.2018.2853590>, 2019.
- Nakata, K., Ohshima, K. I., and Nihashi, S.: Mapping of Active Frazil for Antarctic Coastal Polynyas, With an Estimation of Sea-Ice Production, *Geophysical Research Letters*, 48, e2020GL091353, <https://doi.org/10.1029/2020GL091353>, 2021.



- Nakayama, Y., Cai, C., and Seroussi, H.: Impact of subglacial freshwater discharge on Pine Island Ice Shelf, *Geophysical Research Letters*, 48(18): e2021GL093923, <https://doi.org/10.1029/2021GL093923>, 2021.
- Nakayama, Y., Timmermann, R., Rodehacke, C. B., Schröder, M., and Hellmer, H. H.: Modeling the spreading of glacial meltwater from the Amundsen and Bellingshausen Seas, *Geophysical Research Letters*, 41, 7942–7949, <https://doi.org/10.1002/2014GL061600>, 2014.
- 685 Nakayama, Y., Timmermann, R., and H. Hellmer, H.: Impact of West Antarctic ice shelf melting on Southern Ocean hydrography, *The Cryosphere*, 14, 2205–2216, <https://doi.org/10.5194/tc-14-2205-2020>, 2020.
- Orsi, A. H., and Wiederwohl, C. L.: A recount of Ross Sea water, *Deep-Sea Research Part II*, 56(13), 778–795, <https://doi.org/10.1016/j.dsr2.2008.10.033>, 2009.
- 695 Rignot, E., Jacobs, S., Mouginot, J., and Scheuchl, B.: Ice-Shelf Melting Around Antarctica, *Science*, 341, 266–270, <https://doi.org/10.1126/science.1235798>, 2013.
- Shchepetkin, A., and McWilliams, J.: Correction and commentary for “ocean forecasting in terrain-following coordinate: Formulation and skill assessment of the regional ocean modeling system” by Haidvogel et al., 3595–3634, *Journal of Computational Physics*, 228, 8985–9000, 2009.
- 700 Silvano, A., Foppert, A., Rintoul, S. R., Holland, P. R., Tamura, T., Kimura, N., Castagno, P., Falco, P., Budillon, G., Haumann, F. A., Naveira Garabato, A. C., and Macdonald, A. M.: Recent recovery of Antarctic Bottom Water formation in the Ross Sea driven by climate anomalies, *Nat. Geosci.*, 13, 780–786, <https://doi.org/10.1038/s41561-020-00655-3>, 2020.

- Solodoch, A., Stewart, A. L., Hogg, A. McC., Morrison, A. K., Kiss, A. E., Thompson, A. F., Purkey, S.  
705 G., and Cimoli, L.: How Does Antarctic Bottom Water Cross the Southern Ocean?, *Geophysical  
Research Letters*, 49, e2021GL097211, <https://doi.org/10.1029/2021GL097211>, 2022.
- Spren, G., Kaleschke, L., and Heygster, G.: Sea ice remote sensing using AMSR-E 89-GHz channels,  
*Journal of Geophysical Research: Oceans*, 113, <https://doi.org/10.1029/2005JC003384>, 2008.
- Steele, M., Mellor, G. L., and Mcphee, M. G.: Role of the molecular sublayer in the melting or freezing  
710 of sea ice, *Journal of Physical Oceanography*, 19, 139–147, [https://doi.org/10.1175/1520-0485\(1989\)019<0139:ROTMSI>2.0.CO;2](https://doi.org/10.1175/1520-0485(1989)019<0139:ROTMSI>2.0.CO;2), 1989.
- Stewart, C. L., Christoffersen, P., Nicholls, K. W., Williams, M. J. M., and Dowdeswell, J. A.: Basal  
melting of Ross Ice Shelf from solar heat absorption in an ice-front polynya, *Nat. Geosci.*, 12, 435–  
440, <https://doi.org/10.1038/s41561-019-0356-0>, 2019.
- 715 Stewart, A. L. and Thompson, A. F.: Eddy-mediated transport of warm Circumpolar Deep Water across  
the Antarctic Shelf Break, *Geophysical Research Letters*, 42, 432–440,  
<https://doi.org/10.1002/2014GL062281>, 2015.
- Tamura, T., Ohshima, K. I., Fraser, A. D., and Williams, G. D.: Sea ice production variability in Antarctic  
coastal polynyas, *JGR Oceans*, 121, 2967–2979, <https://doi.org/10.1002/2015JC011537>, 2016.
- 720 Thompson, L., Smith, M., Thomson, J., Stammerjohn, S., Ackley, S., and Loose, B.: Frazil ice growth  
and production during katabatic wind events in the Ross Sea, Antarctica, *The Cryosphere*, 14, 3329–  
3347, <https://doi.org/10.5194/tc-14-3329-2020>, 2020.
- Treasure, A., Roquet, F., Ansorge, I., Bester, M., Boehme, L., Bornemann, H., Charrassin, J., Chevallier,  
D., Costa, D., Fedak, M., Guinet, C., Hammill, M., Harcourt, R., Hindell, M., Kovacs, K., Lea, M.,

- 725 Lovell, P., Lowther, A., Lydersen, C., McIntyre, T., McMahon, C., Muelbert, M., Nicholls, K., Picard, B., Reverdin, G., Trites, A., Williams, G., and P.J.N. de Bruyn.: Marine mammals exploring the oceans pole to pole: A review of the MEOP consortium, *Oceanography*, 30(2), 132–138, <https://doi.org/10.5670/oceanog.2017.234>, 2017.
- Wang, X., Zhang, Z., Dinniman, M. S., Uotila, P., Li, X., and Zhou, M.: The response of sea ice and high-  
730 salinity shelf water in the Ross Ice Shelf Polynya to cyclonic atmosphere circulations. *The Cryosphere*, 17, 1107–1126, *The Cryosphere*, 17, 1107–1126, <https://doi.org/10.5194/tc-17-1107-2023>, 2023.
- Xie, C., Zhang, Z., Wang, C., and Zhou, M.: The response of Ross Sea shelf water properties to enhanced Amundsen Sea ice shelf melting, *Journal of Geophysical Research: Oceans*, 129, e2024JC020919, <https://doi.org/10.1029/2024JC020919>, 2024.
- 735 Yan, L., Wang, Z., Liu, C., Wu, Y., Qin, Q., Sun, C., Qian, J., and Zhang, L.: The Salinity Budget of the Ross Sea Continental Shelf, Antarctica, *JGR Oceans*, 128, e2022JC018979, <https://doi.org/10.1029/2022JC018979>, 2023.
- Zhang, Z., Xie, C., Castagno, P., England, M., Wang, X., Dinniman, M. S., Silvano, A., Wang, C., Zhou, L., Li, X., Zhou, M., and Budillon, G.: Evidence for large-scale climate forcing of dense shelf water  
740 variability in the Ross Sea, *Nature Communications*, 15, 8190, <https://doi.org/10.1038/s41467-024-52524-x>, 2024a.
- Zhang, Z., Xie, C., Wang, C., Chen, Y., Hu, H., and Wang, X.: The Ross Sea and the Amundsen Sea high resolution ocean-sea ice-ice shelf coupled model (RAISE v1.0) [Software]. Zenodo. <https://doi.org/10.5281/zenodo.12735787>, 2024b.

745 Zhang, Z., Xie, C., Wang, C., Chen, Y., Hu, H., and Wang, X.: The Ross Sea and Amundsen Sea Ice-Sea Model (RAISE v1.0): a high-resolution ocean-sea ice-ice shelf coupling model for simulating the Dense Shelf Water and Antarctic Bottom Water in the Ross Sea, Antarctica [Software]. Zenodo. <https://doi.org/10.5281/zenodo.14472621>, 2024c.