

ISWNM-NSCS v2.0: advancing the internal solitary wave numerical model with background currents and horizontally inhomogeneous stratifications

Yankun Gong¹, Xueen Chen², Jiexin Xu¹, Zhiwu Chen¹, Qingyou He^{1,3}, Ruixiang Zhao⁴, Xiao-Hua Zhu⁴, Shuqun Cai^{1,5}

¹State Key Laboratory of Tropical Oceanography, South China Sea Institute of Oceanology, Chinese Academy of Sciences, Guangzhou 510301, China

²College of Oceanic and Atmospheric Sciences, Ocean University of China, Qingdao, 266100, China

³Guangdong Key Lab of Ocean Remote Sensing, South China Sea Institute of Oceanology, Chinese Academy of Sciences, Guangzhou 510301, China

⁴State Key Laboratory of Satellite Ocean Environment Dynamics, Second Institute of Oceanography, Ministry of Natural Resources, Hangzhou, 310012, China

⁵University of Chinese Academy of Sciences, Beijing, 100049, China

Correspondence to: Shuqun Cai (caisq@scsio.ac.cn)

Abstract. A new version of an internal solitary wave (ISW) model, the Internal Solitary Wave Numerical Model-Northern South China Sea version 2.0 (ISWNM-NSCS v2.0), is presented. The background currents and horizontally inhomogeneous stratifications are implemented in ISWNM-NSCS v2.0 to better reproduce ISW properties, including arrival time, mode-1 wave amplitude, wave-induced velocity, characteristic half-width and propagation direction. Optimized viscosity and diffusivity coefficients (i.e., $1.0 \times 10^{-2} \text{ m}^2 \text{ s}^{-1}$ in horizontal and $1.0 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$ in vertical) are also introduced to maintain stable stratifications within the model domain, thereby prolonging the model's valid forecasting period. A mooring station around the Dongsha Atoll is used for model evaluation and numbers of sensitivity experiments are implemented to illustrate the individual effect of the major updates. In comparison with ISWFM-NSCS v1.0, ISWNM-NSCS v2.0 significantly enhances model accuracy in forecasting ISW characteristics, with a 37% improvement in arrival time, a 34% improvement in mode-1 wave amplitude, a 25% improvement in wave-induced velocity, and an 85% improvement in half-width.

1 Introduction

For decades, numerical modelling has been essential for investigating internal solitary waves (ISWs), particularly in dynamically complex regions like the northern South China Sea (NSCS), where large-amplitude ISWs with observed displacements exceeding 200 m and wavelengths of ~ 3 km (Huang et al., 2017; Alford et al., 2015). Theoretical foundations established through frameworks like the Korteweg-de Vries (KdV) equation (Benney, 1966) and its extensions (Grimshaw et al., 2010) have progressively evolved alongside computational advances. By the early 21st century, limitations of two-layer analytical approximations (Holloway et al., 1997) and two-dimensional hydrostatic models (Du et al., 2008) became

apparent for simulating the NSCS's full-depth nonlinear wave dynamics. This spurred development of high-resolution 3D non-hydrostatic solvers capable of resolving critical processes including generation, propagation, and dissipation of ISWs (Simmons et al., 2011). Contemporary open-source frameworks like SUNTANS (Zhang et al., 2011), MITgcm (Vlasenko et al., 2005; Alford et al., 2015), and FVCOM (Lai et al., 2019) now enable realistic simulations of ISW generation, propagation, and dissipation through advanced numerical schemes validated against modern observational arrays. These advances form the foundation for our ISWFM-NSCS model, which extends this capability toward operational forecasting in the NSCS basin.

An ISW forecasting model in the NSCS (Gong et al., 2023), called ISWFM-NSCS v1.0 hereafter, is one of the 3D realistic ISW models developed by a primitive equation ocean solver (MITgcm, Marshall et al., 1997). ISWFM-NSCS v1.0 employs a high horizontal resolution of 500 m and 90 vertical layers, designed to resolve large amplitudes and short wavelengths of ISWs observed in the NSCS. ISWFM-NSCS v1.0 has demonstrated robust performance in characterizing small-scale dynamics within 15 modal days, as evidenced by comparisons to field observations and satellite images. However, as discussed in Gong et al. (2023), there remains potential for enhancement in the coefficient configurations, initial conditions and boundary conditions.

Firstly, the valid forecasting period in ISWFM-NSCS v1.0 is limited to 15 days, as the stratifications significantly weaken beyond this time frame. Viscosity and diffusivity parameters determine the extent of mixing and dissipation of energy in the model. Higher coefficients increase the damping of internal waves, leading to smoother wave fields and reduced wave amplitudes. This can help in preventing numerical instabilities but might underestimate the wave energy and dynamics (Legg and Huijts, 2006). In contrast, low-valued coefficients might lead to numerical noise and spurious oscillations (Álvarez et al., 2019). Therefore, optimizing the settings for viscosity and diffusivity coefficients is essential for maintaining numerical stability in high-resolution models (Nagai and Hibiya, 2015; Stewart et al., 2017) and extending the valid forecasting period (e.g., 30 days or more).

Secondly, initial stratification profiles (temperature and salinity) in ISWFM-NSCS v1.0 are horizontally homogeneous. However, previous studies (e.g., Centurioni et al., 2004; Chao et al., 2007) have observed that the thermocline in the Luzon Strait rises in the west due to the influence of the northward-flowing Kuroshio. Zheng et al. (2007) proposed that the deepening of the thermocline towards the east could potentially hinder the development of eastward-moving solitons in the Pacific. Additionally, 2D idealized model simulations by Shaw et al. (2009) and Buijsman et al. (2010) demonstrated that a sharper and shallower thermocline on the western side of the Luzon Strait, compared to the eastern side, leads to the generation of larger westward-propagating solitons. While these studies shed light on how horizontally inhomogeneous stratification affects the east-west asymmetry of internal solitary waves, their findings have yet to be verified using a 3D realistic model.

Thirdly, ISWFM-NSCS v1.0 does not account for background currents, which are a crucial factor in ISW dynamics. While it is well-established that background stratification and currents significantly affect the characteristics and behaviour of ISWs (DeCarlo et al., 2015; Li et al., 2016), the effects of more complex dynamical motions, such as oceanic currents and

mesoscale eddies, remain inadequately explored. Specifically, the effects of the Kuroshio (Caruso et al., 2006) and mesoscale eddies (Xie et al., 2015) on ISWs have not yet been examined using a 3D realistic model. The absence of such considerations in ISWFM-NSCS v1.0 suggests a gap in our understanding, as incorporating these dynamic elements could provide deeper insights into how ISWs interact with background currents.

In this work, we present an updated version of a high-performance model for predicting ISWs in the NSCS (called ISWNM-NSCS v2.0 hereafter) and evaluate the roles of optimized turbulence configurations, horizontally inhomogeneous stratifications, and background currents in precisely forecasting ISWs through sensitivity numerical experiments. The structure of the manuscript is outlined as follows. A description of the model is given and the major updates made from ISWFM-NSCS v1.0 to ISWNM-NSCS v2.0 are summarized in Sect. 2. The updated model results and corresponding calibrations are detailed in Sect. 3. Moreover, we provide a quantitative analysis to demonstrate the roles of optimized viscosity and diffusivity, horizontally inhomogeneous stratifications, and background currents in the ISWNM-NSCS v2.0 in Sect. 4. Conclusions follow in Sect. 5.

2 Description of the model and major updates

As ISWFM-NSCS v1.0 has been described in Gong et al. (2023), only the main concepts are reviewed here. In ISWNM-NSCS, a 3D realistic non-hydrostatic oceanic solver MITgcm (Marshall et al., 1997, Adcroft et al., 2008) is employed to reproduce generation, propagation and dissipation processes of ISWs in the NSCS. As discussed in ISWFM-NSCS v1.0, the horizontal cell size is set to 500 m, providing approximately 6-8 grid points per characteristic half-width for typical ISWs in the NSCS which aligns with established resolution standards for internal wave process studies (e.g., Zhang et al., 2011; Lai et al., 20219), with 90 vertical layers (i.e., from 5 m at the sea surface to 120 m at the bottom). The configuration represents an optimal balance between computational efficiency and dynamical precision for regional-scale forecasting. More detailed sensitivity analyses of resolution thresholds are available in Gong et al. (2023).

A time step of 10 sec ensures compliance with the Courant-Friedrichs-Lewy (CFL) conditions. Horizontally homogeneous temperature and salinity profiles are initialized by using the climatology WOA 2018 dataset (<https://www.ncei.noaa.gov/access/world-ocean-atlas-2018/>, last access: 10 July 2024) and the model bathymetry is obtained by interpolating the GEBCO dataset (https://www.gebco.net/data_and_products/gridded_bathymetry_data, last access: 10 July 2024). Eight primary barotropic tidal constituents, extracted from TPXO8-atlas dataset (Egbert and Erofeeva, 2002), are applied at each lateral boundary, with a 25 km wide sponge layer absorbing internal wave energy (Zhang et al., 2011). However, background currents and eddies have not been taken into account in ISWFM-NSCS v1.0. To mitigate grid-scale instability, constant turbulent parameters (including viscosity and diffusivity) are applied as follows: $A_h = 0.5 \text{ m}^2 \text{ s}^{-1}$; $A_v = 5.0 \times 10^{-3} \text{ m}^2 \text{ s}^{-1}$; $K_h = 0.5 \text{ m}^2 \text{ s}^{-1}$; $K_v = 5.0 \times 10^{-3} \text{ m}^2 \text{ s}^{-1}$.

In comparison to field observations and satellite imagery, a test case applying the ISWFM-NSCS v1.0 shows well performance in reproducing ISW properties within first 10 model days. Specifically, the root mean square deviations (RMSD)

of arrival time, maximum vertical wave amplitude and baroclinic velocity are 0.71 h, 37.27 m and, 0.41 m s^{-1} , respectively. However, the stratification profiles gradually weaken with descending thermocline depth after the 10th model days, which limit the effective duration of the forecasting model. As discussed in ISWFM-NSCS v1.0, the prediction accuracy may be improved to take into account background currents (Xie et al., 2015) and horizontally inhomogeneous stratifications (Buijsman et al., 2010). Then, this section describes the major updates in comparison with ISWFM-NSCS v1.0 (marked in red in Fig. 1).

2.1 Optimizations of viscosity and diffusivity

In ISWFM-NSCS v1.0, high values of eddy viscosity and diffusivity coefficients (i.e., $0.5 \text{ m}^2 \text{ s}^{-1}$ in horizontal and $5.0 \times 10^{-3} \text{ m}^2 \text{ s}^{-1}$ in vertical) are empirically selected to be sufficient to eliminate grid-scale noise in the velocity and mixing fields (Legg and Huijts, 2006). However, these values generally weaken the background stratifications within the entire model domain, in particular after two weeks, potentially dampening ISW amplitudes and underestimating the wave nonlinearity. Hence, the valid forecasting duration of ISWFM-NSCS v1.0 is less than 15 days, due to the weakening stratification. Therefore, we optimize the eddy viscosity and diffusivity coefficients to extend the valid forecasting duration. The specific parameter updates are detailed below.

Vertical diffusivity in ISWFM-NSCS v2.0 is set to $1.0 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$ (Fig. 1), consistent with microstructure measurements of background diapycnal mixing in summer NSCS (Shang et al., 2017). Horizontal tracer diffusivity adopts $1.0 \times 10^{-2} \text{ m}^2 \text{ s}^{-1}$, following established subgrid-scale parameterizations for mesoscale-resolving models (Large et al., 1994). While transient events (e.g., internal tide breaking) elevate diffusivity to $O(10^{-5} \sim 10^{-4} \text{ m}^2 \text{ s}^{-1})$ (Sun et al., 2016; Yang et al., 2016), our constant coefficients represent baseline values. Sensitivity analyses confirmed that such diffusivity magnitudes preserved large-scale internal wave energetics despite minor impacts on short-wavelength features (Jachec, 2007; Vlasenko et al., 2010). For momentum closure, horizontal eddy viscosity ($A_h = 1.0 \times 10^{-2} \text{ m}^2 \text{ s}^{-1}$) parameterizes unresolved lateral dissipation from inertial ranges and mesoscale processes (Smagorinsky, 1963), while vertical eddy viscosity ($A_v = 1.0 \times 10^{-2} \text{ m}^2 \text{ s}^{-1}$) represents turbulence from shear instabilities and internal wave breaking (Mellor and Yamada, 1982). This configuration aligns with implementations for marginal China Seas using MITgcm (Min et al., 2023; Vlasenko et al., 2018).

To establish physical robustness, we further validate these values through systematic sensitivity experiments (Section 5.1), demonstrating that horizontal and vertical viscosities/diffusivities of $1.0 \times 10^{-2} \text{ m}^2 \text{ s}^{-1}$ and $1.0 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$ optimally reproduce observed ISW properties in the NSCS while maintaining numerical stability.

2.2 Horizontally inhomogeneous stratifications

Numerous satellite images concluded the east-west asymmetric characteristics of ISWs in the Luzon Strait in the NSCS (e.g., Jackson and Apel, 2004; Alford et al., 2015). In previous literatures, the asymmetry of ISWs is a multifaceted phenomenon influenced by asymmetric barotropic tides, water depth differences between the NSCS and Pacific Ocean,

westward thermocline shoaling related to the Kuroshio current, and internal tide resonance in a double ridge configuration.

130 In ISWFM-NSCS v1.0, three factors have been considered except for the east–west gradients in the thermocline.

The westward shoaling of the thermocline is associated with the northward-flowing Kuroshio current, which is centered between the west and east ridges, as evidenced by drifter observations (Centurioni et al., 2004) and model results (Chao et al., 2007). Based on a nonhydrostatic ROMS model, Buijsman et al. (2010) demonstrated that westward-propagating solitons are 28% larger than eastward-propagating solitons, due to the inhomogeneous thermocline, which is secondary to the effect of a
135 deeper Pacific Ocean. In addition, Zheng et al. (2007) argued that the eastward deepening of the thermocline might inhibit the formation of eastward-propagating solitons in the Pacific.

Aforementioned evidences highlight the significant roles of thermocline structure in shaping the east-west asymmetric characteristics of ISWs in the Luzon Strait. Therefore, horizontally inhomogeneous stratifications are conducted as the initial conditions (Figs. 2a and 2b) in ISWNM-NSCS v2.0. Temperature and salinity profiles are both extracted from the global
140 HYCOM re-analysis dataset (<https://www.hycom.org/>, last access: 12 July 2024).

2.3 Background currents and eddies

In the NSCS, background circulations such as the Kuroshio Current play a crucial role in influencing behaviour and characteristics of ISWs (DeCarlo et al., 2015; Li et al., 2016). The Kuroshio, a major western boundary current, brings warm and salty waters into the NSCS, thereby significantly affecting the local hydrography and stratification (Hu et al., 2020).
145 When ISWs encounter the Kuroshio, the interaction can alter the propagation speed and direction of ISWs (Alford et al., 2010). Specifically, a strong north-westward flow of the Kuroshio (looping or leaking type) can accelerate ISWs traveling in the same direction, enhancing their amplitude and nonlinearity. Conversely, ISWs moving against the Kuroshio current can experience deceleration and reduced amplitude, affecting their arrival time and energy. This dynamic interaction affects temporal and spatial characteristics of ISWs, impacting the mixing processes and energy distribution in the NSCS (Xie et al.,
150 2021).

Mesoscale eddies are another ubiquitous phenomenon (Chelton et al., 2011), which are dynamically important in modulating currents and temperature in the NSCS. Their interaction with ISWs can be expected to happen frequently in the deep basin (Liu et al., 2000; Liu et al., 2008; Huang et al., 2017). Numerical simulations by Xie et al. (2015) demonstrated that mesoscale eddies can redistribute the energy of ISWs along their wave fronts. In regions where energy is focused, the
155 amplitudes of ISWs tend to increase, while in spreading regions, they decrease. Previous observational studies (Park and Farmer, 2013; Li et al., 2016) also showed that mesoscale structures can substantially distort the propagation paths of ISWs in the NSCS, leading to dramatic changes in wave amplitude at fixed locations.

Given the importance of background circulations and mesoscale eddies to ISW properties, background currents are not only added as the initial condition (Fig. 2c), but also continuously imposed at the four lateral boundaries (Fig. 3) over time in
160 ISWNM-NSCS v2.0. The background zonal and meridional velocity fields, associated with the corresponding temperature and salinity fields, are directly derived from the global HYCOM re-analysis dataset (<https://www.hycom.org/>, last access: 12

July 2024). These three-dimensional datasets are linearly interpolated onto the model grid to initialize the baseline dynamic conditions, while the time-varying velocity fields from the HYCOM dataset are imposed as lateral boundary forcing across all four domain edges, thereby continuously driving the internal circulation patterns through dynamic coupling.

165 **3 Model results and calibrations**

Following the aforementioned updates, a reference test case (i.e., control run, EXP. 1) is launched since 05 August 2014 and run for 30 days, including two spring-neap cycles. The model runs with a sampling rate of 1 h in the whole domain, and also a sampling rate of 1 min at a targeted location for comparing to field observations.

Then, the model performance is evaluated in three stages: first, by comparing the background current field with the
170 HYCOM reanalysis dataset; second, by comparing the spatial characteristics of ISWs with satellite imagery; and third, by comparing five wave properties (including arrival time, maximum vertical amplitudes, baroclinic velocities, propagation direction and characteristic half-widths) of 28 ISWs with field observational data from the targeted mooring DS.

3.1 Comparison with HYCOM reanalysis dataset

To evaluate the model accuracy in reproducing the correct background current field, we run an extra 3D model (EXP. 0)
175 with the same configurations as EXP. 1, but exclude the surface tide forcing at four lateral boundaries (see details in Table 1). Furthermore, both the horizontal resolution (reduced from 500 m to $1/12^\circ$) and the vertical resolution (reduced from 90 layers to 40 layers) are adjusted downward in EXP. 0 to maintain consistency with the HYCOM reanalysis dataset.

As shown in Fig. 4, we depict four snapshots with a 10-day interval from 05 August to 04 September for the HYCOM reanalysis dataset and the EXP. 0 model results, respectively. Note that main flow patterns are marked with red arrows. It is
180 clear that in both HYCOM reanalysis dataset and EXP. 0 results, the Kuroshio flows northward from the east side of Philippine to the east side of Taiwan Island (Figs. 4a and 4b) with a leaking-pattern intrusion after 20 days (Figs. 4c and 4g). In addition, an anticyclonic eddy is also reproduced in the model at the east side of Luzon Strait after 10 days (Figs. 4b and 4f). However, the model (EXP. 0) might omit a small eddy at the west side of Taiwan Island near the north boundary at the end (Fig. 4h), which is likely to generate remotely and propagate into the model domain. But overall, the model can correctly
185 reproduce background current field in the NSCS, including the Kuroshio and mesoscale eddies.

3.2 Comparison with satellite images

In addition to validating the model's ability in background current regime, we subsequently examine the control run (EXP. 1) in ISW field via a comparison between the model with MODIS imagery (available in the NASA Worldview website, <https://worldview.earthdata.nasa.gov>, last access: 19 July 2024). Given the model's one-hour sampling rate, we choice four
190 closest snapshots of sea surface height gradients for comparison with MODIS imagery (Fig. 5). Note that detailed approaches to compute sea surface height gradients can be found in Gong et al. (2023).

Figs. 5a (05:00 UTC, 14 August) and 5b (05:15 UTC, 14 August) both illustrate two consecutive ISWs (labelled as IW1 and IW2) separated by approximately 120 km. The predicted curvatures, lengths, and positions of IW1 and IW2 exhibit a high degree of consistency with the corresponding features observed in the satellite imagery. Nevertheless, the numerical simulations also reveal two additional ISWs on the continental slope (Fig. 5a), which are obscured in the MODIS-Aqua image from 14 August due to cloud cover (Fig. 5b). On 15 August, the cloud cover cleared, allowing the MODIS-Terra image to capture a clearer depiction of the ISWs (Fig. 5d). The three ISWs shown in Figs. 5c and 5d are located in close proximity and exhibit similar wave crestline lengths extending from the Luzon Strait to the continental slope. Additionally, in shallower waters, the simulated IW1 shows an ISW packet with secondary waves, a feature also observed in the satellite imagery.

Throughout the extended 15-day forecast period in ISWNM-NSCS v2.0, EXP. 1 continues to demonstrate strong performance in depicting spatial distributions of ISWs (Figs. 5e – 5h). Specifically, satellite-observed shallowing and diffracting processes around the Dongsha Atoll at 05:25 UTC on 28 August (Fig. 5f) are clearly captured by the model at 05:00 UTC on 28 August (Fig. 5e). After ISWs impact the Dongsha Atoll (IW4), their wave crests are divided into two branches (IW2 and IW3). The lengths of the wave crests shorten as they bypass the atoll and continue to propagate westward, eventually reconverging behind the island (IW1). It is worth mentioning that the model and the satellite observations remain consistent even on the 25th day of the forecast period (31 August, Figs. 5g and 5h).

Given that the control run does not account for wind effects above the sea surface, some subtle differences in wave characteristics remain. Nevertheless, the model effectively illustrates the spatial features of ISWs in the NSCS, as evidenced by comparisons with the MODIS imagery.

3.3 Comparison with field observations

To conduct a more detailed evaluation of the model's accuracy in predicting ISWs, we incorporate field observations from the Dongsha (hereafter DS) mooring station (117°44.7'E, 20°44.2'N; deployed from 1 August to 6 September, 2014). The mooring included ADCPs (2-min sampling; 16/8-m vertical bins) and distributed temperature, CTD, and CT sensors (10–15 sec sampling). More details can be found in Gong et al. (2023). We examine the vertical structure and arrival time of ISWs after their passage through the deep basin by plotting temperature and baroclinic velocities (or wave-induced velocities) for the periods of 08 to 14 August and 25 to 31 August, respectively. For clarity, Fig. 6 only displays the comparison for the upper 900 m, encompassing the primary wave-induced temperature fluctuations.

Throughout the initial 15-day period, both the forecasting model and in-situ observations capture individual solitons and ISW packets (Figs. 6a and 6b). From 08 to 14 August, there is a notable increase in wave amplitude and nonlinearity, reflecting the transition from neap to spring barotropic tides. The model's predictions show consistent arrival times, baroclinic velocities (as indicated by color shades in Fig. 6), and maximum amplitudes (represented by contours in Fig. 6) with those observed in the in-situ data. Although the model omits some small trailing waves (indicated by green arrows in Fig. 6a) in the observations, it nonetheless demonstrates strong performance in forecasting ISWs at the initial 15 days.

225 Throughout the extended 15-day period, Figs. 6c and 6d continue to exhibit high consistency in predicting arrival times of ISWs. However, the rates of false positives (simulated ISWs that are not observed, indicated by blue arrows in Fig. 6) and false negatives (observed ISWs that are not reproduced, indicated by green arrows in Fig. 6) are relatively higher compared to those observed during the initial 15 days. By considering all ISWs captured at the DS station in EXP. 1, the false positive rate is 8.6% (3 out of 35) and the false negative rate is 11.4% (4 out of 35).

230 To assess the model's accuracy more quantitatively, we have identified the 28 well-predicted ISWs (indicated by red arrows in Fig. 6). We extract their ISW characteristic parameters, such as arrival time, maximum vertical amplitudes, baroclinic velocities, and wave direction, and compare them with field data (see red circles and green triangles in Fig. 7). Note that detailed approaches to extract wave properties can be found in Gong et al. (2023). To evaluate the significance of difference between in-situ observational wave properties and those in ISWNW-NSCS v2.0, we further conduct independent

235 two-sample t-tests. Firstly, the arrival times of ISWs are displayed on the top and bottom of Fig. 7. The discrepancy between the model results and observations is consistently less than one hour and a half, with a RMSD of 0.64 h ($p=0.11$), suggesting that the control run (EXP. 1) accurately captures arrival times of ISWs. Secondly, the model's average of maximum vertical amplitude (~ 88 m) is comparable to the observed value (~ 95 m), though the RMSD for the amplitude is 26.51 m ($p=0.21$). Thirdly, the average maximum baroclinic velocities are 1.34 m s^{-1} in the model and 1.23 m s^{-1} in the observations,

240 respectively, with an RMSD of 0.39 m s^{-1} ($p=0.17$). Finally, the average wave propagation directions are approximately 298° in the model and 288° in the observations, with an RMSD of 13.74° ($p=0.03$). Overall, EXP. 1 successfully reproduces the four key wave features of ISWs observed near the Dongsha Atoll.

4 Sensitivity experiments to evaluate model updates

Building on the standard experiment (EXP. 1, CTRL), we modify its initial and boundary conditions, to individually

245 assess the impact of turbulence parameter optimization, horizontally inhomogeneous stratifications and background currents on the prediction accuracy of ISW model in the NSCS. Details in configuration modifications are as follows (also see Table 1).

1) EXP. 2: Compared with EXP. 1, initial stratification profiles are horizontally homogeneous, derived from the seasonal-averaged WOA18 dataset. Moreover, background currents are excluded at both the initial conditions and boundary

250 conditions.

2) EXP. 3: Compared with EXP. 2, mixing coefficients (i.e., viscosity and diffusivity) are imposed as $A_h = 0.5 \text{ m}^2 \text{ s}^{-1}$; $A_v = 5.0 \times 10^{-3} \text{ m}^2 \text{ s}^{-1}$; $K_h = 0.5 \text{ m}^2 \text{ s}^{-1}$; $K_v = 5.0 \times 10^{-3} \text{ m}^2 \text{ s}^{-1}$ in horizontal and vertical, respectively. Note that the setups in EXP. 3 are identical to those in ISWFM-NSCS v1.0 with the exception of the extended forecasting time (30 days).

3) EXP. 4: Compared with EXP. 1, background currents are only configured as initial conditions but no longer

255 continuously imposed at four lateral boundaries.

4.1 Roles of optimized viscosity and diffusivity

Various three-dimensional models employing different viscosity and diffusivity configurations have been applied to simulate internal solitary waves (ISWs) in the northern South China Sea (e.g., Alford et al., 2015; Lai et al., 2019). However, determining optimal turbulence coefficients for accurate extended-range ISW prediction remains challenging. To address this, we conduct sensitivity experiments EXP.2 and EXP.3, differing by two orders of magnitude in viscosity/diffusivity values.

Analysis of horizontally averaged buoyancy frequency profiles (Fig. 8) reveals fundamental differences in stratification stability. EXP.3 (higher coefficients) exhibits progressive weakening of stratification over 30 days, with thermocline depth descending and maximum buoyancy frequency halving from 0.018 s^{-1} to 0.01 s^{-1} . Conversely, EXP.2 (lower coefficients) maintains stable stratification, showing only a modest reduction from 0.018 s^{-1} to 0.015 s^{-1} . This degradation in EXP.3 directly impacts forecast capability, as evidenced by ISW evolution patterns (Fig. 9). While both experiments accurately reproduce westward-propagating ISWs (individual solitons and wave packets) during the first 15 days, significant deviations emerge thereafter. Beyond 15 days, EXP.3 develops spurious small-scale eddies due to accumulated boundary energy flux, which is unable to accurately forecast ISW properties. Consequently, the valid forecast period in ISWFM-NSCS v1.0 is 15 days rather than 30 days.

Validation against DS station observations (Fig. 10) confirms superior vertical structure representation in EXP.2. Over the extended 15-day period, EXP.2 maintains narrower characteristic half-widths (indicating higher nonlinearity) and stronger stratification compared to EXP.3. Statistical analysis reveals comparable initial error rates (5.9-17.7% false positives/negatives), though EXP.3 degrades more rapidly over time. Quantitative assessment of 28 ISWs (Fig. 11, Table 2) further demonstrates EXP.2's advantages: arrival time biases remain below 1.5 h (RMSD=0.77h, $p<0.01$) versus growing >2 h errors in EXP.3 (RMSD=1.01h, $p<0.01$). While both experiments show amplitude overestimation initially and underestimation later, EXP.2 achieves better baroclinic velocity accuracy (RMSD=0.40 m/s vs. 0.52 m/s) and significantly superior half-width reproduction (RMSD=0.28km, $p<0.01$ vs. 1.13km, $p=0.01$). Performance degradation correlates strongly with stratification loss in both experiments.

In summary, EXP.3 (ISWFM-NSCS v1.0) achieves reliable 15-day forecasts of ISWs in the NSCS. However, EXP.2's reduced viscosity and diffusivity extend the valid forecasting window to 30 days while better preserving wave nonlinearity and vertical structure, establishing optimized turbulence parameters for extended-range prediction. Building on these findings, we further systematically examine ISWFM-NSCS v2.0's sensitivity to mixing parameters through a series of sensitivity coefficient values (from $0.01\times\text{CTRL}$ to $100\times\text{CTRL}$) in the hereinafter, and compare constant turbulent coefficients against dynamic closure schemes to evaluate ISW model performance.

4.2 Roles of horizontally inhomogeneous stratification

The westward shoaling of thermocline, driven the northward-flowing Kuroshio, has been identified as a critical factor contributing to the west-east asymmetry of ISWs in the Luzon Strait (e.g., Zheng et al., 2007; Buijsman et al., 2010). However, the significant role of horizontally inhomogeneous stratification in this process has yet to be validated by a realistic ISW model. Here, we compare two sensitivity experiments (EXP. 2 and EXP. 4) with different initial conditions, namely EXP. 2 with horizontally homogeneous stratifications from WOA18 dataset and EXP. 4 with horizontally inhomogeneous stratifications from HYCOM reanalysis dataset.

First, we examine the effects of horizontally inhomogeneous stratifications on the spatial characteristics of ISWs and review Fig. 9. In EXP. 4, ISW crestlines are longer and more prone to distortion by background processes (Figs. 9d and 9h), which closely replicates the ISW scenario simulated in the control run (EXP. 1). Additionally, spurious eastward-propagating ISWs from the Luzon Strait appear in EXP. 2 (Fig. 9b), which are not reproduced in EXP. 4 (Fig. 9d). This discrepancy arises because the control run, initialized with the HYCOM reanalysis dataset, accounts for the east-west asymmetric thermocline associated with the Kuroshio. Moreover, the southern portion of ISW crestline is much more distinct in EXP. 4 (Fig. 9h) in comparison with EXP. 2 (Fig. 9f), especially as the ISWs approach the Dongsha Atoll and bifurcate into two branches.

Next, we analyse the differences in ISW vertical structures between EXP. 2 and EXP. 4 using data from the selected transect and the DS station. During the initial 15 days, successive westward-propagating internal solitons and ISW packets are captured along the transect both in EXP. 2 and EXP. 4 (Figs. 9b and 9d), as the stratification remain stable. In contrast, during the extended 15 days, EXP. 2 tends to underestimate the ISW nonlinearity, consequently missing an ISW in the deep basin (Fig. 9f), whereas EXP. 4 continues to reproduce it, albeit with a less significant amplitude (Fig. 9h). Given that the primary differences manifest during the final 15 days, we further compare the single-point outputs in EXP. 2 and EXP. 4. It is evident that EXP. 4 (Fig. 10d) captures more ISWs with narrower characteristic half-widths than EXP. 2 (Fig. 10b) at the DS station.

Finally, we conduct a quantitative assessment of the sensitivity models' ability to replicate ISWs by calculating the biases and RMSDs for five key wave properties (Fig. 11 and Table 2) in the cases with and without horizontally inhomogeneous stratification. Fig. 11f illustrates that bias of arrival time in EXP. 4 significantly exceed that in EXP. 2, resulting in a RMSD of 1.20 h ($p < 0.01$) in EXP. 4, compared to 0.77 h ($p < 0.01$) in EXP. 2. This may be due to the omission of the lateral boundary forcing in EXP. 4, resulting in the inability to continuously maintain horizontally inhomogeneous stratification. The RMSDs of maximum wave-induced velocities are very close in the two experiments (see Table 2), namely 0.40 m s^{-1} ($p = 0.20$) in EXP. 2 versus 0.44 m s^{-1} ($p = 0.10$) in EXP. 4. Nonetheless, EXP. 4 demonstrates superior performance in reproducing mode-1 wave amplitude, with a RMSD of 31.94 m ($p = 0.27$) versus 39.17 m ($p = 0.29$) in EXP. 2, as well as in accurately capturing propagation direction, with a RMSD of 9.66° ($p = 0.97$) versus 10.76° ($p = 0.04$) in EXP. 2.

In summary, horizontally inhomogeneous stratification is essential in the initial conditions of ISWNM-NSCS v2.0 particularly during the first 15 days. However, to sustain the west-east asymmetric stratification within the model domain, it is necessary to impose time-variable background currents at the four lateral boundaries.

4.3 Roles of background currents

As inferred from Sec. 4.2, time-variable boundary conditions are crucial for maintaining the horizontally inhomogeneous stratification within the model domain. Here, we extract the background currents (including temperature, salinity and velocities) from the HYCOM reanalysis dataset and impose them at the four lateral boundaries in the control run (EXP. 1). Then, we compare the model results from EXP. 1 with those from EXP. 4, which is solely initialized with 3D stratification and currents but does not include continuous lateral boundary forcing.

Regarding the spatial distribution of ISWs, horizontal gradients of sea surface heights in EXP. 1 (Fig. 9a) exhibit a pattern analogous to that observed in EXP. 4 (Fig. 9d) at 12:00 UTC on 11 August. However, a notable difference emerges at 02:00 UTC on 27 August, wherein the ISW crests in EXP. 1 are longer and more susceptible to distortion by background processes compared to those in EXP. 4. We subsequently examine the differences in the vertical structures of ISWs between two cases along the selected transect and over a 15-day time series at the DS station (Figs. 10a and 10d) during the extended 15 days. Although both cases successfully reproduce distinct vertical structures of ISWs along the transect, ISWs in EXP. 1 (Fig. 9e) exhibit greater nonlinearity compared to those in EXP. 4 (Fig. 9h), particularly within the deep basin. Fig. 10 illustrates that the rate of false positives is 8.6% (3 out of 35) both in EXP. 1 and EXP. 4, but the rate of false negatives (11.4%, 4 out of 35) in EXP. 1 is lower than that (17.1%, 6 out of 35) in EXP. 4.

From a quantitative perspective, EXP. 1 demonstrates superior precision (47%) in predicting the arrival time of ISWs, as evidenced by a RMSD of 0.64 h ($p=0.11$), compared to a RMSD of 1.20 h ($p<0.01$) in EXP. 4. This improved precision is attributed to the presence of time-variable boundary conditions in EXP. 1, which results in a stable stratification. Conversely, in EXP. 4, the bias in arrival time progressively exceeds 1.5 h during the final 15 days (Fig. 11f). Additionally, the control run (EXP. 1) exhibits superior performance in reproducing maximum amplitudes, baroclinic velocities and half-widths than EXP. 4 (see black circles and green stars in Figs. 11b, 11c and 11e). Specifically, the RMSDs in EXP. 1 are 26.51 m ($p=0.21$), 0.39 m s^{-1} ($p=0.17$), and 0.17 km ($p<0.01$), whereas in EXP. 4 the RMSDs are 31.94 m ($p=0.27$), 0.44 m s^{-1} ($p=0.10$), and 0.50 km ($p=0.04$), respectively.

To sum up, by incorporating time-variable background currents at the lateral boundaries, the effects of background flows and mesoscale eddies on the propagation processes of ISWs are more accurately represented. This improvement enhances the model's accuracy in forecasting key wave features, such as arrival time, baroclinic velocities, maximum vertical amplitudes, and characteristic half-widths.

5 Discussion on the turbulence configurations

5.1 Sensitivity of ISWNM-NSCS v2.0 to viscosity and diffusivity

350 In order to examine the sensitivity of the SWNM-NSCS v2.0 results to the horizontal and vertical eddy viscosity and diffusivity coefficients, a set of numerical experiments (CTRL, EXP. A1–A8, K1-K8) are carried out in the NSCS for 30 days. For each experiment, we compare RMSDs for the five ISW properties with the field observations at the DS mooring station. The configurations and results are summarized in Table 3.

355 In the EXPs. A1-A4, the horizontal viscosity coefficient (A_h) is ranging from $100\times\text{CTRL}$ ($1.0\times 10^0 \text{ m}^2 \text{ s}^{-1}$) to $0.01\times\text{CTRL}$ ($1.0\times 10^{-4} \text{ m}^2 \text{ s}^{-1}$). As a result, the EXPs. A1-A4 reveal exceptional stability in four key ISW properties across different scaling factors. Specifically, the arrival time (0.64 h), baroclinic velocity ($\sim 0.40 \text{ m s}^{-1}$), and propagation direction (13.67° – 13.74°) show negligible RMSD variations in EXP. A1–A4. Only maximum mode-1 amplitude exhibits mild degradation (i.e., 27.53–28.59 m vs. 26.51 m in CTRL), while characteristic half-width shows marginal improvement in EXP. A2 (i.e., 0.15 km vs. 0.17 km in CTRL). Conversely, the EXPs. A5-A8 show greater sensitivity for vertical eddy viscosity. In details, EXP. 360 A5 significantly degrades propagation direction (16.05° vs. 13.74° in CTRL), and most sensitivity experiments for A_v worse reproduce maximum wave amplitudes. Although EXP. A6 slightly improves arrival time (0.58 h vs. 0.64 h in CTRL), no sensitivity experiments achieve $>5\%$ improvement across multiple ISW properties.

However, in the EXPs. K1-K4, modifications to horizontal diffusivity (K_h) yield mixed results. While arrival time and baroclinic velocity remain stable ($\pm 0.04 \text{ h}$ and $\pm 0.02 \text{ m s}^{-1}$), EXP. K3 ($K_h=0.1\times\text{CTRL}$) substantially degrades maximum 365 wave amplitude predictions (31.85 m vs. 26.51 m in CTRL). Characteristic half-width consistently worsens (0.18–0.22 km vs. 0.17 km in CTRL), though EXP. K4 slightly improves prediction of propagation direction (13.16° vs. 13.74° in CTRL). For vertical diffusivity (EXPs. K5-K8), extreme scaling causes pronounced effects. Specifically, although EXP. K5 ($K_v=100\times\text{CTRL}$) shows a slight improvement in arrival time (0.55 h vs. 0.64 in CTRL), it significantly degrades four other wave properties simultaneously, namely maximum amplitude (31.92 m vs. 26.51 m in CTRL), baroclinic velocity (0.45 m 370 s^{-1} vs. 0.39 m s^{-1} in CTRL), propagation direction (15.26° vs. 13.74° in CTRL), and characteristic half-width (0.28 km vs. 0.17 km in CTRL). Conversely, EXP. K7 ($K_v=0.1\times\text{CTRL}$) improves maximum amplitude (27.10 m vs. 26.51 m in CTRL) and baroclinic velocity (0.38 m s^{-1} vs. 0.39 m s^{-1} in CTRL), but this is offset by half-width degradation (0.21 km vs. 0.17 km in CTRL).

Overall, no sensitivity experiment outperforms CTRL across all five ISW properties, but only isolated cases (e.g., arrival 375 time in the EXP. A6) show $>5\%$ improvement in single metrics. The CTRL run maintains the most balanced performance, with all RMSDs within intermediate ranges. The fluctuations in ISW properties across all 16 sensitivity experiments confirm that viscosity and diffusivity configurations appear robust in the CTRL run (EXP. 1).

5.2 Application of the K-Profile Parameterization (KPP) scheme in ISWNM-NSCS v2.0

Strong dissipation was observed in the northern region west of the Luzon Strait, extending across the NSCS, driven by wave-wave interactions, direct breaking of internal tides during shoaling, and scattering by abyssal hills (Jiang et al., 2025). However, the control simulation (EXP. 1), which employs constant eddy viscosity and diffusivity values, likely introduces artifacts in the stability of the upper thermocline, thereby distorting the representation of ISW dynamics. While adaptive turbulent closure schemes, such as the Mellor-Yamada hierarchy (Mellor and Yamada, 1982) and the K-Profile Parameterization (KPP; Large et al., 1994), have proven effective in simulating ISW dynamics in stratified shelf seas (e.g., Vlasenko et al., 2005), recent work by Thakur et al. (2022) challenges this paradigm. Their findings suggest that excluding KPP background mixing better preserves vertical velocity gradients and aligns kinetic energy spectra with observations. To reconcile these insights with regional NSCS dynamics, we integrate the KPP scheme into ISWNM-NSCS v2.0 in EXP. 5 to evaluate its influence on ISW dynamics. Results are depicted by green histograms in Fig. 12, revealing the comparative performance of this approach.

Qualitatively, both EXP. 1 (control run) and EXP. 5 (KPP scheme) clearly reproduce the spatial evolution of internal solitons and wave packets from generation to dissipation (not shown). Quantitatively, however, their performance diverges. EXP. 5 shows marginally better precision in predicting ISW arrival times (RMSD = 0.58 h, $p = 0.11$ vs. 0.63 h in EXP. 1, $p = 0.11$), likely due to the KPP scheme's enhanced representation of stable stratification. Conversely, EXP. 1 outperforms EXP. 5 in capturing maximum wave amplitudes (RMSD = 26.51 m vs. 37.22 m, $p = 0.21$), baroclinic velocities (0.39 m s^{-1} vs. 0.52 m s^{-1} , $p = 0.17$), propagation direction (13.74° vs. 14.46° , $p = 0.03\text{--}0.04$), and half-widths (0.17 km vs. 0.25 km , $p < 0.01$), respectively, as shown in Fig. 12 (black lines: EXP. 1; green histograms: EXP. 5). These results suggest that while the KPP scheme improves arrival time accuracy, it degrades predictions of other critical ISW properties. Thus, integrating vertical turbulence parameterizations like KPP into ISWNM-NSCS v2.0 may not universally enhance model skill in the NSCS. Future work should explore hybrid or dynamically adaptive closure schemes to better balance stratification effects with ISW-specific dynamics, ensuring robust predictions across all wave properties.

5.3 Comparison to the General Estuarine Transport Model-General Ocean Turbulence Model (GETM-GOTM)

While constant eddy viscosities and diffusivities demonstrate robust ISW simulation capability in basin-scale applications, this simplified approach fundamentally differs from advanced turbulence closures in coastal models like GETM-GOTM (Burchard et al., 2004). Notably, our choice of constant coefficients represents a pragmatic operational trade-off rather than a physical optimum. While the ISWNM-NSCS v2.0 effectively captures ISW dynamics in the NSCS, it cannot resolve scale-dependent turbulence interactions like dynamic schemes. In comparison, the GETM-GOTM employs a hydrostatic solver optimized for shallow systems ($<100 \text{ m}$ depth), incorporating scale-aware turbulence closures ($k\text{-}\epsilon/k\text{-}\omega$), sediment-coupled biogeochemistry, and wetting-drying schemes essential for coastal processes. These capabilities enable it to resolve stratified shear flows with small error in regions like the Baltic Sea and Wadden Sea (Stips et al., 2004, 2008; Tiessen et al., 2012).

410 Conversely, ISWNM-NSCS v2.0 leverages the non-hydrostatic core of MITgcm to capture vertical accelerations governing large-amplitude ISWs in deep basins (>2000 m), where non-hydrostatic pressure gradients dominate the local nonlinear wave evolution (Vlasenko et al., 2005). Consequently, GETM-GOTM's turbulence closures address sediment-induced mixing absent in our deep-water domain, while our simplified viscosity scheme aligns with operational constraints when validated against observed ISW decay rates. Crucially, this functional specialization highlights how turbulence
415 parameterizations must correspond to hydrodynamic solvers and target phenomena, where GETM-GOTM excels in coastal ecological processes whereas ISWNM-NSCS v2.0 prioritizes abyssal wave dynamics.

In future work, we will implement and benchmark k - ϵ and k - ω closures within ISWNM-NSCS to quantify performance trade-offs across scales. Systematic comparisons against GETM-GOTM across shelf-break zones (e.g., continental shelf and slope in the NSCS) will evaluate whether dynamic schemes improve representation of wave-sediment interactions and
420 turbulent dissipation while maintaining computational viability for forecasting.

6 Conclusions

A robust 3D non-hydrostatic model ISWNM-NSCS v2.0 for forecasting ISWs in the NSCS has been presented. A reference test case was launched from 05 August 2014 and ran for 30 days, during which in-situ observations are available. Various wave properties are better characterized with ISWNM-NSCS v2.0 compared with ISWFM-NSCS v1.0. The major
425 updates and findings are as follows.

1) Optimized viscosity and diffusivity coefficients (i.e., $A_h = K_h = 1.0 \times 10^{-2} \text{ m}^2 \text{ s}^{-1}$, $A_v = K_v = 1.0 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$) contribute to the stabilization of stratification profiles within the model domain, thereby extending the valid forecasting period to 30 days. By comparing the biases between sensitivity model results with in-situ observations at the DS station, we found that ISWFM-NSCS v1.0 gradually loses forecast precision regarding wave nonlinearities after 15 days. Specifically, the RMSD
430 of ISW characteristic half-widths is 1.13 km in SWFM-NSCS v1.0 (EXP. 3), compared to 0.28 km with the optimized turbulence coefficients (EXP. 2).

2) Horizontally inhomogeneous stratifications are implemented as the initial conditions in ISWNM-NSCS v2.0, resulting in west-east asymmetric thermoclines on either side of the Luzon Strait. Considering these horizontally inhomogeneous stratifications, mode-1 wave amplitudes are more accurately reproduced, with a RMSD of 31.94 m in EXP. 4 versus 39.17 m
435 in EXP. 2. Similarly, the propagation direction is better represented, with an RMSD of 9.66° in EXP. 4 versus 10.76° in EXP. 2.

3) Time-variable background currents at four lateral boundaries are essential for maintaining the horizontally inhomogeneous stratification within the model domain. Additionally, background circulations, such as the Kuroshio Current and mesoscale eddies, have been shown to significantly impact the behaviours and characteristics of ISWs in the NSCS. As
440 compared with EXP. 4 (with RMSDs of 31.94 m, 0.44 m s^{-1} , and 0.50 km), applying the background currents could enhance

the performance of ISWNM-NSCS v2.0 in reproducing maximum vertical amplitudes, baroclinic velocities, and characteristic half-widths, resulting in improved RMSDs of 26.51 m, 0.39 m s⁻¹, and 0.17 km.

In summary, ISWNM-NSCS v2.0 incorporates optimized turbulence coefficients, horizontally inhomogeneous stratifications, and background currents, compared with ISWFM-NSCS v1.0. As a result, ISWNM-NSCS v2.0 demonstrates considerable improvements in the model's ability to accurately predict a range of wave properties, achieving a 37% improvement in arrival time, a 34% improvement in mode-1 wave amplitude, a 25% improvement in wave-induced velocity, and an 85% improvement in characteristic half-width.

Code and data availability. The MODIS satellite imagery can be freely downloaded from the NASA Worldview website (<https://worldview.earthdata.nasa.gov>, last access: 11 July 2024, Plato et al., 2019). The code of Massachusetts Institute of Technology general circulation model for ISWNM-NSCS v2.0 can be accessed at <https://doi.org/10.5281/zenodo.14847454> (last access: 11 February 2025). The input files, including initial and boundary conditions, as well as the corresponding output data for ISWNM-NSCS v2.0, are freely accessible through an open-access data repository available at <https://doi.org/10.5281/zenodo.14842090> (Gong, 2025, last access: 10 February 2025).

Author contributions. YG, XC, ZC, QH, and SC developed the software. JX, RZ, and XZ provided guidance on the processing of observational data. YG conducted the numerical model. YG wrote the draft of the manuscript with the help of all the co-authors. All authors reviewed the final manuscript. YG, XC, ZC, and SC were responsible for conceptualisation, and for the workshop and training in the use of the software.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Financial support. This work was jointly supported by the National Natural Science Foundation of China (NSFC) under contract Nos. 42130404, 42206012, 42276022, 42176025, 42276011 and U23A2032; Natural Science Foundation of Guangdong Province (2024A1515012703, 2024A1515012549); State Key Laboratory of Satellite Ocean Environment Dynamics, SIO, MNR (QNHX2308); the Science and Technology Projects of Guangzhou (2024A04J3587, 2024A04J9022); South China Sea Institute of Oceanology (SCSIO2023QY02 and LTOZZ2205); and Strategic Priority Research Program of the Chinese Academy of Sciences (XDA0370304). Yankun Gong is supported by CAS Cultivating Young Scientists Foundation. The numerical simulations were supported by the High Performance Computing Division, with assistance from HPC managers Wei Zhou and Dandan Sui at the South China Sea Institute of Oceanology.

References

- Adcroft, A., Campin, J. M., Dutkiewicz, S., Evangelinos, C., Ferreira, D., Forget, G., ... and Molod, A.: MITgcm user manual. Massachusetts Institute of Technology, <http://hdl.handle.net/1721.1/117188>, 2008.
- 475 Alford, M. H., Lien, R. C., Simmons, H., Klymak, J., Ramp, S., Yang, Y. J., ... and Chang, M. H.: Speed and evolution of nonlinear internal waves transiting the South China Sea. *Journal of Physical Oceanography*, 40, 1338-1355, <https://doi.org/10.1175/2010JPO4388.1>, 2010.
- Alford, M. H., Peacock, T., MacKinnon, J. A., Nash, J. D., Buijsman, M. C., Centurioni, L. R., ... and Paluszkiwicz, T.: The formation and fate of internal waves in the South China Sea, *Nature*, 521, 65-69, <https://doi.org/10.1038/nature14399>,
 480 2015.
- Álvarez, Ó., Izquierdo, A., González, C. J., Bruno, M., and Mañanes, R.: Some considerations about non-hydrostatic vs. hydrostatic simulation of short-period internal waves. A case study: The Strait of Gibraltar. *Continental Shelf Research*, 181, 174-186, <https://doi.org/10.1016/j.csr.2019.05.016>, 2019.
- Benney, D. J.: Long non-linear waves in fluid flows, *Journal of Mathematics and Physics*, 45, 52-63,
 485 <https://doi.org/10.1002/sapm196645152>, 1966.
- Buijsman, M. C., McWilliams, J. C., and Jackson, C. R.: East-west asymmetry in nonlinear internal waves from Luzon Strait, *Journal of Geophysical Research: Oceans*, 115, <https://doi.org/10.1029/2009JC006004>, 2010.
- Burchard, H., Bolding, K., and Villarreal, M. R.: Three-dimensional modelling of estuarine turbidity maxima in a tidal estuary. *Ocean Dynamics*, 54, 250-265, <https://doi.org/10.1007/s10236-003-0073-4>, 2004.**
- 490 Caruso, M., Gawarkiewicz, G. G., and Beardsley, R.: Interannual variability of the Kuroshio Current intrusion in the South China Sea, *Journal of Oceanography*, 62, 559–575. <https://doi.org/10.1007/s10872-006-0076-0>, 2006.
- Centurioni, L. R., Niiler, P. P., and Lee, D. K.: Observations of inflow of Philippine Sea surface water into the South China Sea through the Luzon Strait, *Journal of Physical Oceanography*, 34, 113-121, [https://doi.org/10.1175/1520-0485\(2004\)034<0113:OOIOPS>2.0.CO;2](https://doi.org/10.1175/1520-0485(2004)034<0113:OOIOPS>2.0.CO;2), 2004.
- 495 Chao, S. Y., Ko, D. S., Lien, R. C., and Shaw, P. T.: Assessing the west ridge of Luzon Strait as an internal wave mediator, *Journal of Oceanography*, 63, 897-911, <https://doi.org/10.1007/s10872-007-0076-8>, 2007.
- Chelton, D. B., Schlax, M. G., & Samelson, R. M.: Global observations of nonlinear mesoscale eddies, *Progress in oceanography*, 91, 167-216, <https://doi.org/10.1016/j.pocean.2011.01.002>, 2011.
- DeCarlo, T. M., Karnauskas, K. B., Davis, K. A., and Wong, G. T. F.: Climate modulates internal wave activity in the
 500 Northern South China Sea, *Geophysical Research Letters*, 42, 831–838. <https://doi.org/10.1002/2014gl062522>, 2015.
- Du, T., Tseng, Y. H., and Yan, X. H.: Impacts of tidal currents and Kuroshio intrusion on the generation of nonlinear internal waves in Luzon Strait, *Journal of Geophysical Research: Oceans*, 113, <https://doi.org/10.1029/2007JC004294>, 2008.
- Egbert, G. D., and Erofeeva, S. Y.: Efficient inverse modeling of barotropic ocean tides, *Journal of Atmospheric and Oceanic technology*, 19, 183-204, [https://doi.org/10.1175/1520-0426\(2002\)019<0183:EIMOBO>2.0.CO;2](https://doi.org/10.1175/1520-0426(2002)019<0183:EIMOBO>2.0.CO;2), 2002.

- 505 Grimshaw, R., Pelinovsky, E., Talipova, T., and Kurkina, O.: Internal solitary waves: propagation, deformation and disintegration. *Nonlinear Processes in Geophysics*, 17(6), 633-649, <https://doi.org/10.5194/npg-17-633-2010>, 2010.
- Gong, Y., Chen, X., Xu, J., Xie, J., Chen, Z., He, Y., and Cai, S.: An internal solitary wave forecasting model in the northern South China Sea (ISWFM-NSCS), *Geosci. Model Dev.*, 16, 2851–2871, <https://doi.org/10.5194/gmd-16-2851-2023>, 2023.
- Holloway, P. E., Pelinovsky, E., Talipova, T., and Barnes, B.: A nonlinear model of internal tide transformation on the
510 Australian North West Shelf. *Journal of Physical Oceanography*, 27(6), 871-896, [https://doi.org/10.1175/1520-0485\(1997\)027<0871:ANMOIT>2.0.CO;2](https://doi.org/10.1175/1520-0485(1997)027<0871:ANMOIT>2.0.CO;2), 1997.
- Hu, S., Sprintall, J., Guan, C., McPhaden, M. J., Wang, F., Hu, D., and Cai, W.: Deep-reaching acceleration of global mean ocean circulation over the past two decades. *Science advances*, 6, eaax7727, <https://doi.org/10.1126/sciadv.aax7727>, 2020.
- Huang, X., Zhang, Z., Zhang, X., Qian, H., Zhao, W., and Tian, J.: Impacts of a mesoscale eddy pair on internal solitary
515 waves in the northern South China Sea revealed by mooring array observations, *Journal of Physical Oceanography*, 47, 1539-1554, <https://doi.org/10.1175/JPO-D-16-0111.1>, 2017.
- Jachec, S. M.: *Understanding the evolution and energetics of internal tides within Monterey Bay via numerical simulations (Doctoral dissertation, Stanford University)*, 2007.
- Jackson, C. R., and Apel, J.: An atlas of internal solitary-like waves and their properties. *Contract*, 14(03-C), 0176,
520 https://www.internalwaveatlas.com/Atlas2_PDF/IWAtlas2_FrontMatter.pdf, 2004.
- Jiang, S., Dai, D., Wang, D., Wang, S., Li, Y., Guo, J., and Qiao, F.: Inferring diapycnal mixing using the internal wave continuum from the high resolution ocean model. *Ocean Modelling*, 195, 102525, <https://doi.org/10.1016/j.ocemod.2025.102525>, 2025.
- Lai, Z., Jin, G., Huang, Y., Chen, H., Shang, X., and Xiong, X.: The generation of nonlinear internal waves in the South
525 China Sea: A three-dimensional, nonhydrostatic numerical study, *Journal of Geophysical Research: Oceans*, 124, 8949-8968, <https://doi.org/10.1029/2019JC015283>, 2019.
- Large, W. G., McWilliams, J. C., and Doney, S. C.: Oceanic vertical mixing: A review and a model with a nonlocal boundary layer parameterization. *Reviews of geophysics*, 32, 363-403, <https://doi.org/10.1029/94RG01872>, 1994.
- Legg, S., and Huijts, K. M.: Preliminary simulations of internal waves and mixing generated by finite amplitude tidal flow
530 over isolated topography, *Deep Sea Research Part II: Topical Studies in Oceanography*, 53, 140-156, <https://doi.org/10.1016/j.dsr2.2005.09.014>, 2006.
- Li, Q., Wang, B., Chen, X., Chen, X., and Park, J.-H.: Variability of nonlinear internal waves in the South China Sea affected by the Kuroshio and mesoscale eddies. *Journal of Geophysical Research: Oceans*, 121, 2098–2118. <https://doi.org/10.1002/2015jc011134>, 2016.
- 535 Liu, Y., Yuan, Y., Su, J., and Jiang, J.: Circulation in the South China Sea in summer of 1998. *Chinese Science Bulletin*, 45, 1648-1655, <https://doi.org/10.1007/BF02898979>, 2000.

- Liu, Y., Weisberg, R. H., and Yuan, Y.: Patterns of upper layer circulation variability in the South China Sea from satellite altimetry using the self-organizing map. *Acta Oceanologica Sinica*, 27(Suppl.), 129, https://digitalcommons.usf.edu/msc_facpub/332/, 2008.
- 540 Marshall, J., Hill, C., Perelman, L., and Adcroft, A.: Hydrostatic, quasi-hydrostatic, and nonhydrostatic ocean modelling, *Journal of Geophysical Research: Oceans*, 102, 5733-5752, <https://doi.org/10.1029/96JC02776>, 1997.
- Mellor, G. L., and Yamada, T.: Development of a turbulence closure model for geophysical fluid problems, *Reviews of Geophysics*, 20(4), 851-875, <https://doi.org/10.1029/RG020i004p00851>, 1982.
- 545 Min, W., Li, Q., Xu, Z., Wang, Y., Li, D., Zhang, P., ... and Yin, B.: High-resolution, non-hydrostatic simulation of internal tides and solitary waves in the southern East China Sea. *Ocean Modelling*, 181, 102141, <https://doi.org/10.1016/j.ocemod.2022.102141>, 2023.
- Nagai, T., and Hibiya, T.: Internal tides and associated vertical mixing in the Indonesian Archipelago. *Journal of Geophysical Research: Oceans*, 120, 3373-3390, <https://doi.org/10.1002/2014JC010592>, 2015.
- Park, J.-H., and Farmer, D.: Effects of Kuroshio intrusions on nonlinear internal waves in the South China Sea during winter, *Journal of Geophysical Research: Oceans*, 118, 7081-7094, <https://doi.org/10.1002/2013jc008983>, 2013.
- 550 Simmons, H., Chang, M. H., Chang, Y. T., Chao, S. Y., Fringer, O., Jackson, C. R., and Ko, D. S.: Modeling and prediction of internal waves in the South China Sea, *Oceanography*, 24, 88-99, 2011.
- Shang, X. D., Liang, C. R., & Chen, G. Y.: Spatial distribution of turbulent mixing in the upper ocean of the South China Sea, *Ocean Science*, 13(3), 503-519, <https://doi.org/10.5194/os-13-503-2017>, 2017.
- 555 Shaw, P. T., Ko, D. S., and Chao, S. Y.: Internal solitary waves induced by flow over a ridge: With applications to the northern South China Sea, *Journal of Geophysical Research: Oceans*, 114, <https://doi.org/10.1029/2008JC005007>, 2009.
- Smagorinsky, J.: General circulation experiments with the primitive equations: I. The basic experiment. *Monthly weather review*, 91(3), 99-164, [https://doi.org/10.1175/1520-0493\(1963\)091<0099:GCEWTP>2.3.CO;2](https://doi.org/10.1175/1520-0493(1963)091<0099:GCEWTP>2.3.CO;2), 1963.
- 560 Stewart, K. D., Hogg, A. M., Griffies, S. M., Heerdegen, A. P., Ward, M. L., Spence, P., and England, M. H.: Vertical resolution of baroclinic modes in global ocean models, *Ocean Modelling*, 113, 50-65, <https://doi.org/10.1016/j.ocemod.2017.03.012>, 2017.
- Stips, A., Bolding, K., Pohlmann, T., and Burchard, H.: Simulating the temporal and spatial dynamics of the North Sea using the new model GETM (general estuarine transport model). *Ocean Dynamics*, 54, 266-283, <https://doi.org/10.1007/s10236-003-0077-0>, 2004.
- 565 Stips, A. K., Bolding, K., and Lilover, M.: Scenario simulations of recent Baltic Sea inflows using the hydrodynamic transport model GETM. In 2008 IEEE/OES US/EU-Baltic International Symposium (pp. 1-6). IEEE, <https://doi.org/10.1109/BALTIC.2008.4625527>, 2008.
- Sun, H., Yang, Q., Zhao, W., Liang, X., and Tian, J.: Temporal variability of diapycnal mixing in the northern South China Sea, *Journal of Geophysical Research: Oceans*, 121, 8840-8848, <https://doi.org/10.1002/2016JC012044>, 2016.

- 570 Thakur, R., Arbic, B. K., Menemenlis, D., Momeni, K., Pan, Y., Peltier, W. R., ... and Ma, Y.: Impact of vertical mixing parameterizations on internal gravity wave spectra in regional ocean models. *Geophysical Research Letters*, 49(16), e2022GL099614, <https://doi.org/10.1029/2022GL099614>, 2022.
- Tiessen, M., Nauw, J., Ruudij, P., and Gerkema, T.: Numerical modeling of physical processes in the North Sea and Wadden Sea with GETM/GOTM, <https://doi.org/10.3990/2.197>, 2012.
- 575 Vlasenko, V., Stashchuk, N., and Hutter, K.: Baroclinic tides: theoretical modeling and observational evidence. Cambridge University Press, 2005.
- Vlasenko, V., Stashchuk, N., Guo, C., and Chen, X.: Multimodal structure of baroclinic tides in the South China Sea, *Nonlinear Processes in Geophysics*, 17, 529-543, <https://doi.org/10.5194/npg-17-529-2010>, 2010.
- Vlasenko, V., Stashchuk, N., and Nimmo-Smith, W. A. M.: Three-dimensional dynamics of baroclinic tides over a seamount. *Journal of Geophysical Research: Oceans*, 123(2), 1263-1285, <https://doi.org/10.1002/2017JC013287>, 2018.
- 580 Xie, J., He, Y., Chen, Z., Xu, J., and Cai, S.: Simulations of internal solitary wave interactions with mesoscale eddies in the northeastern South China Sea, *Journal of Physical Oceanography*, 45, 2959–2978. <https://doi.org/10.1175/jpo-d-15-0029.1>, 2015.
- Xie, J., Fang, W., He, Y., Chen, Z., Liu, G., Gong, Y., and Cai, S.: Variation of internal solitary wave propagation induced by the typical oceanic circulation patterns in the northern South China Sea deep basin, *Geophysical Research Letters*, 48, e2021GL093969, <https://doi.org/10.1029/2021GL093969>, 2021.
- 585 Yang, Q., Zhao, W., Liang, X., and Tian, J.: Three-dimensional distribution of turbulent mixing in the South China Sea. *Journal of Physical Oceanography*, 46(3), 769-788, <https://doi.org/10.1175/JPO-D-14-0220.1>, 2016.
- Zhang, Z., Fringer, O. B., and Ramp, S. R.: Three-dimensional, nonhydrostatic numerical simulation of nonlinear internal wave generation and propagation in the South China Sea, *Journal of Geophysical Research: Oceans*, 116, <https://doi.org/10.1029/2010JC006424>, 2011.
- 590 Zheng, Q., Susanto, R. D., Ho, C. R., Song, Y. T., and Xu, Q.: Statistical and dynamical analyses of generation mechanisms of solitary internal waves in the northern South China Sea, *Journal of Geophysical Research: Oceans*, 112, <https://doi.org/10.1029/2006JC003551>, 2007.

595




Figure 1. (a) The schematic of configuration and implementation of ISWFM-NSCS v1.0, which includes initial and boundary conditions. (b) Same as (a) but for ISWNM-NSCS v2.0. Note that the major model updates are marked in red in (b).




Figure 2. (a) Horizontally inhomogeneous temperature near the sea surface at the initial conditions (00:00 UTC 05 August 2014), which is derived from HYCOM reanalysis dataset. (b) Meridionally-averaged temperature profile through the entire model domain, showing west-eastward asymmetry of thermoclines. (c) Background velocity near the sea surface at the initial conditions, which is derived from HYCOM reanalysis dataset. (d) Model bathymetry, obtained by GEBCO dataset.




Figure 3. (a) Background currents and tidal forcing at the west lateral boundary, including x-t diagram of background zonal velocities at the sea surface (top panel), latitudinal averaged zonal velocity in the upper 1000 m (front panel), and tidal amplitudes of four primary constituents (right panel). Note that color ranges are $-1.5 \sim 1.5 \text{ m s}^{-1}$ and $-0.2 \sim 0.2 \text{ m s}^{-1}$ in the top and front panels, respectively. (b-d) Same as (a) but for east, north and south lateral boundaries, respectively.




Figure 4. (a-d) Background currents and eddies at the sea surface from 05 August to 04 September in 2014 with a time interval of 10 days, derived from the global HYCOM reanalysis dataset (<https://www.hycom.org/>). (e-h) Same as (a-d) but
615 derived from the model results of EXP. 0.




Figure 5. (a) Horizontal gradients of sea surface heights induced by internal solitary waves (ISWs) at 05:00 UTC on 14 August 2014, and (b) corresponding MODIS-Aqua image captured at 05:15 UTC on 14 August 2014. (c), (e), and (g) Same as (a) but at 03:00 UTC on 15 August, 05:00 UTC on 28 August, and 03:00 UTC on 31 August 2014, respectively. (d), (f), and (h) Same as (b) but with MODIS-Terra imagery at 02:50 UTC on 15 August, MODIS-Aqua imagery at 05:25 UTC on 28 August, and MODIS-Terra imagery at 02:50 UTC on 31 August 2014, respectively. Note that the MODIS images were freely accessible at the NASA Worldview website (<https://worldview.earthdata.nasa.gov>, open source).




Figure 6. (a) Temperature and wave-induced (or baroclinic) velocities along the main propagation direction of ISWs from 08 August to 15 August based on field observations at the DS station. (b) Same as (a) but for the standard experiment (EXP. 1). (c-d) Same as (a-b) but from 25 August to 31 September. Red arrows highlight ISWs correctly captured by the model, while blue and green arrows demote the false positive and false negative results, respectively.





Figure 7. Maximum mode-1 wave amplitudes (a), baroclinic velocities (b), and propagation directions (c) of twenty-eight ISWs at the DS station. Field observations are represented by red circles, while numerical model results are indicated by green triangles. Note that averaged values are shown as solid lines.



635 **Figure 8.** (a) Time series of horizontally-averaged background buoyancy frequency in the upper 500 m through the entire model domain in EXP. 2. (b) Black and red lines represent the averaged buoyancy frequency profiles at the beginning and at the ending of the model, respectively. (c-d) Same as (a-b) but in EXP. 3.




Figure 9. Sea surface height gradients, and temperature and baroclinic velocities along the transect (marked as dashed line in Fig. 8a) at 12:00 UTC on 11 August 2014 in the cases (a) EXP. 1, (b) EXP. 2, (c) EXP. 3, and (d) EXP. 4, respectively. (e-h) Same as (a-d) but at 02:00 UTC on 27 August 2014. Small panels in the bottom left display the zonal barotropic velocity (in m s^{-1}) in the Luzon Strait. Solid lines represent the barotropic tidal conditions at the specified times.




Figure 10. Time series of temperature and baroclinic velocities at station DS from 25 August to 01 September 2014 for the
645 model runs: (a) EXP. 1, (b) EXP. 2, (c) EXP. 3, and (d) EXP. 4. Red arrows highlight ISWs detected by the model, while
blue and green arrows denote the false positive and false negative results, respectively.




Figure 11. Bias in arrival time (a), maximum mode-1 amplitudes (b), wave-induced velocities (c), propagation directions (d), and half-widths (e) for 14 ISWs at the first 15 predicting days at station DS. (f)-(j) Same as (a)-(e) but for the remaining 14 ISWs during the final 15 predicting days. The control run (EXP. 1) is depicted by black lines and dark shades, while the three sensitivity experiments are shown as histograms in different colours (EXP. 2 – EXP. 4).




Figure 12. Same as Fig. 11 but for the sensitivity experiment (EXP. 5) considering the K-Profile Parameterization (KPP) scheme.

Table 1. Summary of all experimental configurations.

No.	A_h, K_h	A_v, K_v	Initial conditions	Boundary conditions
EXP. 0	$1.0 \times 10^{-2} \text{ m}^2 \text{ s}^{-1}$	$1.0 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$	3D currents & Stratifications (HYCOM)	Background currents (HYCOM)
EXP. 1	$1.0 \times 10^{-2} \text{ m}^2 \text{ s}^{-1}$	$1.0 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$	3D currents & Stratifications (HYCOM)	Surface tides (TPXO8) & Background currents (HYCOM)
EXP. 2	$1.0 \times 10^{-2} \text{ m}^2 \text{ s}^{-1}$	$1.0 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$	Horizontally homogeneous stratifications (WOA18)	Surface tides (TPXO8)
EXP. 3	$5.0 \times 10^{-1} \text{ m}^2 \text{ s}^{-1}$	$5.0 \times 10^{-3} \text{ m}^2 \text{ s}^{-1}$	Horizontally homogeneous stratifications (WOA18)	Surface tides (TPXO8)
EXP. 4	$1.0 \times 10^{-2} \text{ m}^2 \text{ s}^{-1}$	$1.0 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$	3D currents & Stratifications (HYCOM)	Surface tides (TPXO8)

Table 2. Root mean square deviation (RMSD) of five wave properties via comparing different sensitivity experiments with field observations at the DS station. Note that the values in parentheses denote the RMSD of the ISWs during the initial and final 15-day periods, respectively.

No.	RMSD of arrival time [h]	RMSD of maximum mode-1 amplitude [m]	RMSD of baroclinic velocity [m s ⁻¹]	RMSD of propagation direction [°]	RMSD of characteristic half- width [km]
EXP. 1	0.64 (0.64, 0.64)	26.51 (24.17, 29.01)	0.39 (0.40, 0.39)	13.74 (8.75, 17.89)	0.17 (0.19, 0.14)
EXP. 2	0.77 (0.74, 0.81)	39.17 (46.54, 28.22)	0.40 (0.42, 0.38)	10.76 (8.79, 12.67)	0.28 (0.26, 0.30)
EXP. 3	1.01 (0.69, 1.28)	40.39 (45.44, 33.56)	0.52 (0.40, 0.63)	10.09 (8.67, 11.54)	1.13 (0.16, 1.65)
EXP. 4	1.20 (0.70, 1.60)	31.94 (29.79, 34.28)	0.44 (0.49, 0.38)	9.66 (7.79, 11.47)	0.50 (0.18, 0.71)

660 **Table 3.** Same as Table 2 but for sensitivity experiments with varying viscosities and diffusivities. Note that bold red (blue) values indicate experiments performing >5% better (worse) than CTRL (EXP. 1) in the RMSD metrics. Percentages in brackets indicate the magnitude of RMSD change relative to CTRL, with $\uparrow$ ($\downarrow$) denoting degradation (improvement).

No.	RMSD of arrival time [h]	RMSD of maximum mode-1 amplitude [m]	RMSD of baroclinic velocity [m s ⁻¹]	RMSD of propagation direction [°]	RMSD of characteristic half- width [km]
EXP. 1 (CTRL)	0.64	26.51	0.39	13.74	0.17
EXP. A1 ($A_h=100\times\text{CTRL}$)	0.64	27.53	0.40	13.70	0.17
EXP. A2 ($A_h=10\times\text{CTRL}$)	0.64	27.61	0.40	13.70	0.15 ($\uparrow 12\%$)
EXP. A3 ($A_h=0.1\times\text{CTRL}$)	0.64	28.59 ($\downarrow 8\%$)	0.40	13.71	0.22 ($\downarrow 29\%$)
EXP. A4 ($A_h=0.01\times\text{CTRL}$)	0.64	28.08 ($\downarrow 6\%$)	0.40	13.67	0.23 ($\downarrow 35\%$)
EXP. A5 ($A_v=100\times\text{CTRL}$)	0.64	28.11 ($\downarrow 6\%$)	0.41 ($\downarrow 5\%$)	16.05 ($\downarrow 17\%$)	0.21 ($\downarrow 24\%$)
EXP. A6 ($A_v=10\times\text{CTRL}$)	0.58 ($\uparrow 9\%$)	31.41 ($\downarrow 18\%$)	0.41 ($\downarrow 5\%$)	14.50 ($\downarrow 6\%$)	0.17
EXP. A7 ($A_v=0.1\times\text{CTRL}$)	0.62	28.47 ($\downarrow 7\%$)	0.41 ($\downarrow 5\%$)	15.09 ($\downarrow 10\%$)	0.20 ($\downarrow 18\%$)
EXP. A8 ($A_v=0.01\times\text{CTRL}$)	0.62	31.64 ($\downarrow 19\%$)	0.42 ($\downarrow 8\%$)	13.89	0.24 ($\downarrow 41\%$)
EXP. K1 ($K_h=100\times\text{CTRL}$)	0.61 ($\uparrow 5\%$)	27.71	0.39	13.74	0.20 ($\downarrow 18\%$)
EXP. K2 ($K_h=10\times\text{CTRL}$)	0.63	29.31 ($\downarrow 11\%$)	0.41 ($\downarrow 5\%$)	14.17	0.18 ($\downarrow 6\%$)
EXP. K3 ($K_h=0.1\times\text{CTRL}$)	0.60 ($\uparrow 6\%$)	31.85 ($\downarrow 20\%$)	0.40	14.17	0.22 ($\downarrow 29\%$)
EXP. K4 ($K_h=0.01\times\text{CTRL}$)	0.64	28.78 ($\downarrow 9\%$)	0.39	13.16	0.22 ($\downarrow 29\%$)
EXP. K5 ($K_v=100\times\text{CTRL}$)	0.55 ($\uparrow 14\%$)	31.92 ($\downarrow 20\%$)	0.45 ($\downarrow 15\%$)	15.26 ($\downarrow 11\%$)	0.28 ($\downarrow 65\%$)
EXP. K6 ($K_v=10\times\text{CTRL}$)	0.64	30.26 ($\downarrow 14\%$)	0.41 ($\downarrow 5\%$)	14.59 ($\downarrow 6\%$)	0.21 ($\downarrow 24\%$)
EXP. K7 ($K_v=0.1\times\text{CTRL}$)	0.62	27.10	0.38	13.45	0.21 ($\downarrow 24\%$)
EXP. K8 ($K_v=0.01\times\text{CTRL}$)	0.64	29.88 ($\downarrow 13\%$)	0.42 ($\downarrow 8\%$)	14.28	0.19 ($\downarrow 12\%$)