



Data-driven rolling model for global wave height

Xinxin Wang³, Jiuke Wang⁴, Wenfang Lu⁵, Changming Dong⁶, Hao Qin³, Haoyu Jiang^{*1,2,3}

¹ College of Life Sciences and Oceanography, Shenzhen University, Shenzhen, China

² Laboratory for Regional Oceanography and Numerical Modeling, Qingdao Marine Science and Technology Center

5 ³ Shenzhen Research Institute, China University of Geosciences, Shenzhen, China

⁴ School of Artificial Intelligence, Sun Yat-Sen University, Zhuhai, China

⁵ School of Marine Sciences, Sun Yat-Sen University, Zhuhai, China

⁶ School of Marine Sciences, Nanjing University of Information Science & Technology, Nanjing, China

10 Corresponding author: Haoyu Jiang (Haoyujiang@szu.edu.cn)

Abstract. Significant Wave Height (SWH) is crucial for many human activities, such as marine navigation, offshore operations, and coastal management. Traditionally, SWH is modeled using numerical wave models, which, while accurate, are computationally intensive and constrained by incomplete physical representations of wave spectral evolution. This study
15 introduces a simple global deep learning-based model for SWH, which uses the current SWH field and the wind field at the next time step as inputs to predict the SWH field at the next time step. This approach mirrors the rolling prediction strategy of numerical wave models. After training on a re-analysis dataset, the errors of the model diverge lightly with time when given a good initial field because no spectral information is used. However, after diverging for ~200 hours, the errors stabilize, remaining comparable to those of state-of-the-art numerical wave models. Additionally, the error divergence can be mitigated
20 through the assimilation of altimeter measurements. This deep learning model can not only serve as an efficient surrogate for traditional numerical wave models but also provide a baseline for statistical modeling of global SWH due to its simplicity in inputs and outputs.

1 Introduction

Wind-generated surface gravity waves (hereafter, waves) are one of the most common physical phenomena on the sea surface.
25 These waves impact nearly all human activities in the ocean, including ocean engineering, maritime navigation, fisheries, and port operations. Moreover, ocean waves play a crucial role in many geophysical processes at the sea surface, such as the exchange of mass, momentum, and energy within the wave boundary layer. Thus, it is essential to keep improving our ability to model ocean waves.

30 Numerical Wave Models (NWMs) are the most widely used tool for forecasting and hindcasting waves. These models apply numerical methods to solve wave action balance equations, thereby representing the evolution of wave spectra. Over years of development, widely used NWMs like WAVEWATCH III (WW3) (WW3DG, 2019) and SWAN (Simulating Waves Nearshore) (Booij et al., 1999) have demonstrated the capability to provide spatio-temporal distributions of wave parameters, such as Significant Wave Height (SWH), given reliable wind forcing fields (e.g., Alday et al., 2021; Liu et al., 2021).



35

However, NWMs have notable limitations. First, their computational cost is high, particularly with high-resolution models, making them difficult to use in time-sensitive or resource-constrained scenarios. It is noted that the evolution of wave spectra in NWMs occurs within a five-dimensional space (two spatial dimensions, time, frequency, and direction), adding complexity to the numerical computations. Second, the accuracy of NWMs is constrained by incomplete physical representations and numerical effects.

40

The rapid development of artificial intelligence (AI) offers potential solutions to the limitations of traditional NWMs. Recent advancements in AI weather forecasting have demonstrated that AI-based models can achieve better accuracy than numerical models with much lower computational costs (e.g., Lam et al. 2023, Bi et al. 2023), providing the confidence for developing AI-based wave models. Consequently, some studies have already explored AI applications in wave modeling. Some have attempted to replicate the AI weather forecasting approach by treating wave modeling as a purely nonlinear auto-regression problem of spatio-temporal series (e.g., Zhou et al., 2021; Ouyang et al., 2023). However, according to our understanding, this method is simply wrong because phase-averaged wave modeling is more of a forcing problem than an initial value problem.

45

50

Recent studies have adopted a rolling SWH prediction strategy similar to NWMs, utilizing both initial SWH fields (past and present) and forcing wind fields (future winds) as inputs, with future SWHs as outputs. However, most of these studies have focused on wind-sea-dominated nearshore areas, where swell propagation is not a dominant factor in wave modeling (e.g., Cao et al. 2023; Gao et al. 2023). These studies have found that the error in these AI models increases over time compared to NWM hindcasts. This is not surprising because the models do not account for spectral information, and different spectra with the same SWH respond differently to the same forcing. If such an error divergence is too large, the AI model will not be able to run independently without the initial SWH field from NWMs. Conversely, if the divergence is minor, the model may still be valuable for various applications. However, to the best of our knowledge, no study has yet discussed whether such a model combined wind and SWH inputs can operate effectively using a rolling strategy without relying on NWM data.

55

60

One potential solution to solving this problem in AI wave modeling is straightforward, that is, to incorporate the full directional wave spectrum, allowing the AI to approximate the solution of the wave action balance equation. However, applying this method to global SWH modeling presents significant challenges. The global directional wave spectra at any given moment form a very large 4-D matrix. When using these 4-D matrices as inputs and outputs of an AI model, the training will require an enormous dataset and a complex model architecture.

65

From an engineering perspective, the divergence in model simulation errors can be mitigated through data assimilation. In NWMs, the assimilation of altimeter-measured SWH does not always yield positive outcomes because altimeters provide only



wave height information without detailed wave spectra (e.g., Ardhuin et al. 2019, Jiang et al., 2022). However, it is worth investigating whether the assimilation of altimeter data can enhance the accuracy of AI-based SWH modeling.

70

In this study, we propose a global-scale deep learning-based model for SWH. The model utilizes a rolling prediction strategy, similar to NWMs, by taking the current SWH field and the wind field at the next time step as inputs and predicting the SWH field at that next time step. This model is designed to address two key questions: 1) How does a simplified global AI wave model, using an input-output framework similar to NWMs but without incorporating spectral data, handle error divergence?
75 2) Can the assimilation of altimeter data help reduce error divergence and improve the reliability of SWH modeling?

After training the model on a re-analysis dataset, it was observed that, as expected, the AI model experiences a slight divergence in error over time when provided with a good initial field. However, after approximately 200 hours, the error stabilizes, and the stabilized errors are not significantly larger than those of state-of-the-art NWMs, which is somewhat surprising.
80 Additionally, we demonstrate that the issue of error divergence can be partially mitigated through the assimilation of altimeter measurements. The remainder of this paper is organized as follows: Section 2 describes the data and methodologies employed in this study. Section 3 presents the results from the AI model and their evaluation, followed by discussions and conclusions in Section 4.

2 Materials and Methods

85 2.1 Data

2.1.1 ERA5 Wind and Wave Data

The ERA5 is a comprehensive global climate reanalysis dataset, covering the period from 1950 to the present, with hourly data on a wide range of atmospheric and wave parameters (Hersbach et al. 2020). This dataset is based on state-of-the-art modeling technology and has assimilated global historical observations to produce global estimates of these parameters. The
90 wave data in ERA5 is derived from the Wave Model (WAM) hindcast and has assimilated SWH data from various altimeters, including ERS-1/2, ENVISAT, JASON-1/2, CRYOSAT-2, and SARAL, using an optimal interpolation scheme. This assimilation enhances the accuracy of SWH data, particularly in the open ocean, making ERA5 more reliable compared to other NWM hindcasts. Due to its accuracy and consistency, ERA5 data products have been widely utilized in wave-related research (e.g., Jiang and Mu 2019, Jiang 2020). The dataset is available through the Climate Data Store, with pre-interpolated
95 resolutions up to $0.25^\circ \times 0.25^\circ$ for atmospheric parameters and $0.5^\circ \times 0.5^\circ$ for wave parameters.

In this study, we utilized the global SWH and 10-meter longitudinal and latitudinal components of neutral wind (U10 and V10) from the ERA5 dataset for the period 2000-2017 to train the global AI SWH model. The corresponding data in the year 2022



100 was used for validation to prevent over-fitting, while the model testing was conducted with data in the year 2020. Both the wind and wave data used here are at a $0.5^\circ \times 0.5^\circ \times 1\text{h}$ spatio-temporal resolution.

2.1.2 CCI-Sea State Dataset

105 The altimeter dataset used in this study for data assimilation experiment and model evaluation is the Climate Change Initiative (CCI)-Sea State dataset version 3 (Dodet et al., 2020). This dataset provides accurate and consistent global SWH data. The SWH data have undergone rigorous quality control and joint calibration to minimize systematic errors across altimeters. Additionally, a non-parametric empirical mode decomposition technique has been employed for data de-noising, effectively reducing random measurement errors. As shown by Jiang (2023), after reducing random noise, the typical error of SWH from the CCI-Sea State is only ~ 0.15 m in the open ocean, making the dataset well-suited for calibrating and evaluating SWHs from NWMs. To minimize land contamination, altimeter measurements within 50 km offshore were excluded from the dataset.

110

2.1.3 WAVEWATCH-III Hindcast

115 The SWH data from the WAVEWATCH-III model hindcast with the physical parameterizations by Liu (2021), hereafter referred to as WW3-ST6, were utilized as a benchmark to evaluate the performance of the AI model. This hindcast is driven by ERA5 10-m surface winds and has a spatial-temporal resolution of $0.25^\circ \times 0.25^\circ \times 3\text{h}$. Although it does not assimilate wave observations, the WW3-ST6 hindcast shows good agreement with observational data, achieving an overall RMSE of approximately 0.35 m (or 5%–15% of SWH) compared to altimeter data in the open ocean. Detailed information and access to the dataset can be found in Liu (2021).

2.2 Deep Learning Model

2.2.1 Model inputs and outputs

120 The deep learning model for SWH in this study employs an input-output structure similar to NWMs. The SWH field at any time point T_i (initial SWH field) and the wind field (U10 and V10) at the next time point (one hour later in this case) T_{i+1} are used to predict the SWH field at T_{i+1} . The model can then further predict the SWH field at T_{i+2} using the SWH field at T_{i+1} and the wind field at T_{i+2} , which is a rolling prediction strategy. We understand that adding historical wind information might enhance the accuracy of the AI SWH model. Particularly, if a long series of wind fields are used as inputs, the model can work in a different way that the initial SWH field is not needed (e.g., Song and Jiang, 2023). However, one of our aims at this stage is to maintain the model's similarity to NWMs to test the effectiveness of this straightforward and simple input-output structure.

125



2.2.2 Model Structure

This study employs a U-Net architecture for the AI modeling of global-scale SWH. U-Net is a convolutional neural network (CNN) originally designed for biomedical image segmentation. It is characterized by its U-shaped structure, which combines an encoder and a decoder through skip connections (Ronneberger et al. 2015). The encoder progressively extracts features from the input through convolution and pooling, while the decoder reconstructs spatial resolution using de-convolution and up-sampling. Skip connections link corresponding layers of the encoder and decoder, preserving high-resolution details. We believe such a CNN-based deep learning model is well-suited for wave statistical modeling using our input-output structure. The processes of both local wave generation by wind and wave propagation in space can be captured by convolutional kernels at different scales.

Figure 1 presents a schematic of the U-Net architecture used in this study. The input matrix consists of three channels: the global SWH field at T_i and the U10 and V10 fields at T_{i+1} . To handle the wraparound at the -180° and 180° longitude boundary, we used an engineering trick of extending the input fields from -180° to 180° (720 longitudes) to -190° to 190° (760 longitudes), to simplify the propagation of signals near the two boundaries. The final output, the SWH field at T_{i+1} , also spans -190° to 190° , but only data from -180° to 180° were retained in the computation of the cost function for training.




Figure 1: An illustration of the U-Net architecture used in this study. Each cube represents a feature map, with the numbers on the sides indicating the number of channels. The legend on the lower-right panel explains the meaning of the different arrows used in the schematic diagram.

2.2.3 Model Training

For model training, the training set was randomly shuffled, and the model was then trained to minimize the Mean Squared Error (MSE) between the model output and the target output:



$$Loss = \frac{1}{n} \sum_{j=1}^{Lat} \sum_{i=1}^{Lon} [(y_{i,j} - x_{i,j}) \cos \theta_j]^2 \quad (1)$$

where x and y denote the SWH from the AI SWH model and ERA5, respectively; the subscripts i and j denote the i -th longitudinal and j -th latitudinal grid point; θ_j denotes the latitude of the j -th latitudinal grid point. This cosine term was introduced to account for the area change of grid points with latitudes. We used six batches for training and trained the model for 30 epochs at a learning rate of 0.0001 using the AdamW optimizer. Training took approximately one hour per epoch on an NVIDIA GeForce 4090 GPU. Once trained, the model requires less than 10 minutes to compute (infer) the global SWH for one year at a spatio-temporal resolution of $0.5^\circ \times 0.5^\circ \times 1h$ on an NVIDIA 3060 GPU.

2.2.4 Epoch Ensemble Method

To further improve the accuracy and stability of the model predictions, this study employs the epoch ensemble method. This approach mitigates potential issues like over-fitting or under-fitting, which can arise from relying on a single model, by leveraging the diversity of models trained across different epochs. The simplest way of using this method is to retain several models obtained in different epochs during the training process and average their outputs during inference. This straightforward yet effective strategy enhances model performance without requiring additional training. In this study, the ensemble size was set to four, and the ensemble mean reduced the Root-MSE (RMSE) by ~30% compared to individual models. The final AI models established in this study are available from the Git-hub repository at: <https://github.com/YulKeal/AI-Rolling-Wave-Height-Model>.

2.3 Error Metrics

The Bias, RMSE, Correlation Coefficient (CC), and Scatter Index (SI) are used as the error metrics to evaluate the performance of the AI SWH model, which are defined as:

$$Bias = \frac{1}{n} \sum_{i=1}^n (y_i - x_i) \quad (2)$$

$$RMSE = \sqrt{\frac{1}{n} \sum_{i=1}^n (y_i - x_i)^2} \quad (3)$$

$$CC = \frac{\sum_{i=1}^n (y_i - \bar{y})(x_i - \bar{x})}{\sqrt{\sum_{i=1}^n (y_i - \bar{y})^2} \sqrt{\sum_{i=1}^n (x_i - \bar{x})^2}} \quad (4)$$

$$SI = RMSE / \bar{y} \quad (5)$$

where x and y denote the SWH from the AI models and reference data (which can be either ERA5 or CCI-sea state altimeter data), respectively; n is the sample size, and the bars over x and y denote their mean values. These error metrics were also used to monitor the training process of the model.



2.4 Data Assimilation

To reduce error divergence in the long-term operation of the rolling prediction, we tried to incorporate data assimilation techniques by integrating altimeter measurements to correct the model’s “initial” SWH field. In the assimilation of NWMs, spectral information is used so that it requires a method to transform observations of SWH or other integrated wave parameters into wave spectra. However, in our case, only SWH fields and wind fields are used as the model inputs, without involving spectral information. Thus, the assimilation of this model also does not need to involve the spectral information, which simplifies the assimilation.

Here, we employed a simple objective analysis method for data assimilation:

$$A(i, j, t) = M(i, j, t) + \sum_{k=1}^N w(i, j, t)_k g(O_k - M_k) \quad (6)$$

$$w(i, j, t)_k = \exp\left[-d_k^2(i, j, t) / 2R(i, j, t)^2\right] / \sum_{k=1}^N \exp\left[-d_k^2(i, j, t) / 2R(i, j, t)^2\right] \quad (7)$$

$$R(i, j, t) = \min[d_k(i, j, t)] \quad (k = 1, 2, 3, \dots, N) \quad (8)$$

$$d_k(i, j, t) = \sqrt{(S_k / S_1)^2 + (T_k / T_1)^2} \quad (S_k < 1500 \text{ km}, T_k < 48 \text{ h}) \quad (9)$$


where i, j , and t represent longitude, latitude, and time, respectively; M (model) and A (assimilated) represent the model outputs before and after assimilation, respectively; $k \in \{1, 2, \dots, N\}$ represents the number of observations to be assimilated at a given time; O_k and M_k represent the values of observed and corresponding modeled SWH at the spatio-temporal location of the k -th observation; w_k represents the weight factor for correction at location (i, j, t) for the k -th observation; d_k represent the spatio-temporal distance from location (i, j, t) to the k -th observation; $R(i, j, t)$ represent the distance from location (i, j, t) to its nearest observation. S_k and T_k are the spatial and temporal differences between the location (i, j, t) and the k -th observation, respectively, and S_1 and T_1 are tuning coefficients to combine spatial and temporal distances. Previous studies often used a 30-min-50-km window to collocate SWH from altimeters and other sources (e.g., Jiang 2020), thus, S_1 and T_1 are set to 50 km and 0.5 h, respectively.

In our data assimilation experiment, assimilation was conducted every 6 hours, beginning after the first 24 hours of the model run. During each assimilation, the SWH data from the CCI-Sea State dataset were used to correct the AI model’s hindcasts using Equations 6-9. It is noted that in Equation 9, the upper limits of S_k and T_k mean that only observations within 1500 km can influence the value of the target grid point. Only observations from the past 48 hours were used to correct the current SWH field. After assimilation, the prediction for the next time step used the assimilated SWH field as inputs for the AI model.



3 Results

205 The performance of the proposed rolling AI model for SWH was evaluated on the 2020 test dataset. We selected initial SWH fields every 36 hours from 00:00:00 2020-Jan-1 (i.e., the 0th, 36th, 72nd, ..., 8460th hours of 2020, totaling 236 sets of experiments). For each initial SWH field, a 300-hour rolling modeling was conducted. Figure 2 shows the variation of global overall error metrics compared to ERA5 SWH with simulation time. The orange and blue lines represent the mean values of the error metrics for the 236 experiments before and after assimilation, respectively, with the shaded areas indicating the range of these metrics across different starting times.



210 **Figure 2:** The variation of global overall error metrics between the AI SWH model outputs and ERA5 with simulation time: (a) CC, (b) bias, (c) RMSE, and (d) SI. The orange and blue lines represent the mean values of the error metrics for the 236 experiments starting from different initial SWH fields, before and after assimilation, respectively. The shaded areas around the lines indicate the range of error metrics across different experiments with varying initial SWH fields.

215 For the condition without assimilation, the curves for all four error metrics show that the errors of the AI SWH model increase rapidly with simulation time initially. As mentioned in the introduction, this trend is expected given the absence of spectral information. However, as simulation time progresses, the rate of error growth diminishes, and the model stabilizes after ~240 hours. This means the model can still capture some aspects of SWH evolution over time. Remarkably, the global overall mean values for CC, bias, RMSE, and SI are around 0.985, 0.06 m, 0.23 m, and 0.09, respectively, comparable to state-of-the-art
220 NWMs. This suggests that the simple AI model can function independently, at least, in certain scenarios.

To further test whether such a model can operate independently, we conducted a “cold start” experiment using an initial SWH field of zero. The results, shown in Fig. 3, are compared with the “hot start” results from Fig. 2. Although the initial SWH fields are the same (zeros) in the “cold start” experiment, varying wind fields at different starting times lead to differences in



error metrics (depicted by blue shadows in Fig. 3). As expected, the cold start experiment shows larger errors initially, but
225 these errors diminish over time, converging to values similar to those from the “hot start” after approximately 240 hours. This
convergence demonstrates the robustness of the model.




Figure 3: The variation of global overall error metrics between the AI SWH model outputs and ERA5 with simulation time:
(a) CC, (b) bias, (c) RMSE, and (d) SI. The orange lines and shaded areas are the same as those in Fig. 2, but no epoch ensemble
230 is used. The blue lines and shaded area are the corresponding results for the cold start with an initial field of zero SWH.

Go back to Fig. 2, when data assimilation is applied, the errors are significantly reduced across all metrics, except for bias
before the ~100th hour. The increase in bias is likely due to minor inconsistencies between ERA5 and CCI-Sea State, and the
bias remains less than 0.06 m. For the other error metrics, assimilation reduces the time required for error stabilization to ~72
235 hours while also lowering the final converged errors of the AI model. When stabilized, the global overall CC, bias, RMSE,
and SI reach 0.992, 0.05 m, 0.17 m, and 0.07, respectively. Although these metrics are calculated relative to ERA5 data rather
than direct observations, these values seem to be completely acceptable for most operational wave modeling applications.

To further understand the model's performance, we plotted the scatter plot and geographical distributions of the four
240 aforementioned error metrics for both results with and without data assimilation. The results for 6-h, 24-h, 72-h, and 240-h
simulations without data assimilation are shown in Fig. S1, S2, and S3 of the Supporting Information and Fig. 4, respectively.
Although errors increase with simulation time, as shown in Fig. 2, these results indicate that the spatial error patterns remain
consistent across different simulation time. Our primary focus is on the 240-h hindcast results in Fig. 4, where the errors have
stabilized. This is because high-quality SWH initial field like ERA5 is not always easily available, and when the AI SWH
245 model is run independently, the stabilized error is more typical and meaningful as the reference.




Figure 4: Comparison of SWHs from the AI model at 240-h hindcast time (when the errors are stable) with ERA5 for the year 2020. (a) Scatter plot between the SWHs from the two datasets. (b-e) Global spatial distributions of CC, bias, RMSE, and SI, respectively.

250

The scatter plot in Fig. 4a shows a good overall agreement between the SWHs generated by the AI model and those from ERA5, with most points closely aligning with the 1:1 line. The SI of 0.093, the CC of 0.986, and the RMSE of 0.23 m are already better than those typically observed between contemporary global NWM hindcasts and altimeter data. Although such a direct comparison might not be entirely fair or reasonable, these values indicate that this simple AI SWH model is capable of effectively modeling the distribution and variability of global SWHs.

255

Regarding the spatial distributions of errors, the CCs (Fig. 4b) are close to 0.99 in the westerlies of both hemispheres and in marginal and (semi-)enclosed seas where wind-seas occur frequently dominated. However, in the tropical oceans, especially along their eastern coasts where swells are predominant (“swell pools”, Chen et al. (2002)), the CCs are below 0.9 (~0.85 in the Indian Ocean, ~0.8 in the Atlantic Ocean, and ~0.7 in the Pacific Ocean).

260

Three main factors contribute to these lower CC values in the swell pools. First, the wind-sea growing process can be regarded as a forcing problem while swell propagation is more of an initial value problem. This difference is evident in the CCs observed



over different simulation time. For example, in regions of westerlies, the CCs remain stable at around 0.99 across 6-h, 24-h,
265 72-h, and 240-h hindcasts (Fig. S1b-S3b and Fig. 4b). Conversely, in the Pacific swell pool, CCs decrease significantly with
simulation time: from 0.98 at 6 hours, to 0.92 at 24 hours, to 0.8 at 72 hours, and finally to 0.7 at 240 hours. Compared to the
wind-sea growth, it is far more challenging, if not physically impossible, for the AI model to accurately learn the swell
propagation process using only the evolution of SWH spatial patterns without directional wave spectra. Despite this limitation,
the AI model still manages to capture some rough characteristics of swell energy propagation from the SWH data, which is
270 why its performance is still reasonable in these swell-dominated regions.

Second, SWHs in swell pools typically vary within a narrow range of approximately 0.5~3.5 m. This limited variability means
that even if absolute RMSEs are relatively low, the CC values may still be low, making it challenging to achieve high CC
values. These regions also exhibit the lowest CCs in the comparisons between other global SWH data, i.e., NWM hindcast
275 versus altimeter observations. For reference, comparisons between SWHs from WW3-ST6 and CCI-Sea State are shown in
Fig. S4, where CCs in these swell pools are also lower than 0.8.

Third, the Garden Sprinkle Effect (GSE), a numerical error associated with swell propagation, can introduce “random” errors
into SWHs when swells have propagated over large distances. Such swells are very common in swell pools and it is probably
280 impossible for the AI model to learn how these numerical errors evolve using the ERA5 SWH data.

Regarding the bias, the values vary in the range of ± 0.15 m in most parts of the ocean but can reach 0.3 m to the Southwest of
South America and 0.2 m to the Southeast of Africa. These relatively large biases are related to the accumulation of error with
simulation time. It is not clearly known why the bias has such a distribution. We also plotted the distribution of bias in other
285 years and found that the regions with the largest bias are slightly different in different years but the overall patterns are similar.
For example, the results in the year 2000 are shown in Fig. S5 where the error maps look similar to those in Fig. 4. It is noted
that although the data from the year 2000 is used in the model training, the training is only based on 1-h simulations without
rolling so the AI model has never “seen” the exact input for the simulation time of more than two hours. It is not a wrong way
to use the data in the training set to do a long-term rolling test (of course, using an independent testing set should be better). In
290 all years, these biases are not significant compared to the typical annual mean SWH in the corresponding regions. Besides,
these biases can largely be corrected by some simple post-process methods such as point-by-point linear regression.

The RMSE pattern in Fig. 4d shares some similarities with the bias pattern, with the largest RMSE values also found southwest
of South America, indicating that bias significantly contributes to the overall error. However, the swell pools with relatively
295 low CCs are not prominently visible in the maps of bias, RMSE, and SI. In terms of SI, apart from the regions with relatively
large RMSE, high SI values are also observed near small islands and archipelagos such as Indonesia. The annual mean SWHs
in these areas are lower than in the open ocean, so even a small RMSE can lead to a relatively large SI. Moreover, these areas




represent only a small portion of the global ocean, so their contributions to the overall loss function are minimal. However, wave behavior in these regions differs significantly from that in the open ocean, leading to a different input-output relationship for the model, which complicates the training process. Additionally, NWMs also encounter numerical errors when handling these small islands. These factors make it more challenging for the AI model to effectively "learn" from data near small islands.

The AI model does not directly incorporate ice information, treating ice-covered regions simply as land. As a result, higher errors are expected in polar regions. However, this is not clearly evident in Fig. 4. In contrast, Fig. S1 shows that errors in polar regions increase rapidly at first but then stabilize with simulation time. This pattern may be due to the variability of sea ice, which leads the model to primarily learn the rapid response of SWHs to wind forcing in polar regions. Consequently, SWHs in polar regions are less sensitive to the initial field compared to other areas. It is also noted that only the data outside the marginal ice zone is used, meaning waves that propagate into these regions are not considered. For waves generated in the marginal ice zone and propagating out, they contribute minimally to the overall SWH energy so they are neglected by the AI model.

The distributions of these error metrics suggest that the AI model performs well across global oceans in general, both in wind-sea- and swell-dominated regions. To provide a more intuitive understanding of the AI SWH model's performance, an animation comparing the global SWH distributions from ERA5 and our AI model is presented in Movie S1 in the Supporting Information. Slightly different from the 240-h hindcast results in Fig. 4, the results in Movie S1 are generated by continuously rolling the AI model from 01-Jan-2020 00:00:00. A simple visual inspection of the movie indicates that the AI model effectively captures SWH evolution, suggesting that the AI model could serve as an effective surrogate for NWMs, at least for some wind-sea-dominated regions.

To further evaluate the model's performance, we compared the SWHs from the 240-h~272-h hindcasts of the AI model with those from the CCI-Sea State, with the results shown in Fig. 5. This direct comparison with altimeter-measured SWHs provides a more independent and commonly used method for wave model evaluation. To ensure sufficient collocation between the altimeter and model data, we extended the simulation period by 36 hours, making sure that every altimeter data record in the open ocean can be collocated with a "model grid point".

The comparison shows that the AI model also aligns well with the altimeter data. In Fig. 5a, most data points lie along the 1:1 line, with a bias close to zero, an RMSE of 0.336 m, a CC of 0.968, and an SI of 0.123. These overall error metrics are comparable to those observed between WW3-ST6 and CCI-Sea State in Fig. S4 where the bias, RMSE, CC, and SI are 0.032 m, 0.326 m, 0.972, and 0.119, respectively.



330

Figure 5: The same as Fig. 4, but the comparison is between the 240-h SWH hindcasts of the AI model and the CCI-Sea State dataset.

Regarding the spatial patterns of errors, Fig. 5b-5e are similar to Fig. 4b-4e though the magnitudes of errors are generally larger in Fig. 5. The CCs in Fig. 5b are ~ 0.98 in the westerlies but are only ~ 0.6 in the Pacific swell pool. In contrast, the two corresponding CCs are ~ 0.97 and ~ 0.7 , respectively, in Fig. S4. For other error metrics, the differences between the two models are even smaller. The biases vary in a similar range, and the RMSEs and SIs show similar patterns in Fig. 5 and Fig. S4. Notably, in the westerlies the RMSE and SI values from the AI model are even slightly lower than those from WW3-ST6, a state-of-the-art NWM hindcast. These findings further demonstrate the strong performance of the AI SWH model, particularly in open ocean regions that are not always predominated by swells.

340

For the 240-h rolling hindcast results of the AI model after data assimilation every six hours, the corresponding comparisons with ERA5 and CCI-Sea State are shown in Fig. 6 and 7, respectively. Compared to the results without assimilation in Fig. 4, all the error metrics of the model improve significantly after data assimilation in Fig. 6. Specifically, the CCs in the Pacific, Atlantic, and Indian Ocean swell pools increase from ~ 0.7 , ~ 0.8 , and ~ 0.85 in Fig. 4 to ~ 0.88 , ~ 0.92 , and 0.95 in Fig. 6, respectively. The magnitudes of bias, RMSE, and SI also decrease across the oceans after assimilation, although the bias and RMSE remain relatively high in regions to the southwest of South America and southeast of Africa. The comparison between

345



350 Fig. 5 and Fig. 7 shows a similar result: the overall errors become significantly smaller, particularly in the swell-dominated regions, after assimilation. Similar to Supplementary Movie S1, the comparison animation of the results after assimilation is placed in Supplementary Movie S2, where the AI model better captured the SWH evolution.






Figure 6: The same as Fig. 4, but the AI model has assimilated the data from CCI-Sea State every six hours.



355 **Figure 7:** The same as Fig. 6, but the comparison is with the CCI-Sea State dataset.

4 Discussion

The results demonstrate that it is feasible to develop a usable AI SWH model using only an initial SWH field and the wind field at the next time step as inputs. These are likely the minimum requirements for the inputs of an AI SWH model. As noted
360 in the introduction, relying solely on SWH fields as inputs is insufficient since wind-seas cannot be accurately modeled without wind information. Similarly, one can expect that if the initial SWH field is excluded from the inputs, the AI model would struggle to simulate ocean swells using only the input of the current wind field. To confirm this, we trained an AI model using only the wind field as input, with SWH at the same time step as the output, with the results shown in Fig. 8. The model performance in Fig. 8 is significantly worse than that in Fig. 4, in both wind-sea- and swell-dominated regions. In areas with frequent wind-seas, such as the westerlies, although the CCs exceed 0.95, the RMSEs can also surpass 0.4 m in both hemispheres, much higher than the values in Fig. 4. In the Pacific and Atlantic swell pools, the CCs are even lower than 0.4.
365 These results underscore the critical importance of including both of the initial SWH field and the forcing wind field as inputs for the AI SWH model.



370 **Figure 8:** The same as Fig. 4, but the AI model is trained only using the wind field at the corresponding time as the input.

From a physical perspective, the SWH at a given location is influenced by the wind speed, fetch, and duration in wind-sea conditions. The wind input provides information on both wind speed and fetch. Meanwhile, duration (or historical wind) information is partially and implicitly conveyed by the SWH input, as it is computed in a rolling simulation using a recursive method that incorporates past wind data. Although the implicit information provided by global SWHs is not as comprehensive as that from global directional wave spectra, the spatial distribution of SWHs still contains significant historical wind information. This explains why including the SWH input is beneficial for modeling wind-sea-dominated regions and why the AI model can slightly outperform the NWM in these areas.

375
380 In swell-dominated regions, where local wind speeds remain low almost all years, using only the wind input fails to provide any meaningful information about the SWH, as illustrated in Fig. 8. However, as previously mentioned, the AI model can still learn some rough statistical characteristics of swell energy propagation from the data, especially in regions like swell pools. This is also demonstrated in Movie S1, where the propagation of swells generated by extra-tropical storms into tropical regions is distinctly observable.

385



Although the above analysis underscores the importance of including SWH input for the AI model, the quality of the initial SWH is not important if the model is run in a rolling way for relatively long time. The "cold start" experiment has demonstrated that the model error can stabilize within 240 hours, even without an initial SWH field. However, we do not recommend using such a "cold start" in practice because a better initial field or data assimilation can greatly accelerate the speed of error convergence and such a better initial field is almost always available (e.g., using the output of the model in Fig. 8).

Regarding data assimilation, the assimilation of altimeter SWH measurements is sometimes believed not to be always helpful in NWMs, and may even have negative effects in some cases. This is because there are different approaches to using SWH data to correct directional wave spectra, and improper corrections can adversely affect the model results. However, in this AI model, the spectral information is encapsulated within the SWH, and both the computation and assimilation are directly based on the SWH. Consequently, if the assimilated SWH data is more accurate than the output of the AI model, the assimilation will positively impact the results.

It is not surprising that data assimilation can significantly improve the performance of the AI model, but it is noted that the computational cost of assimilation in this AI model is low. In the assimilation of NWMs, SWH observations are used to correct the spectral densities of directional wave spectra, a four-dimensional array (latitude, longitude, frequency, direction) at a given time step, using empirical relations. In contrast, the assimilation process in the AI model bypasses the need for wave spectral information, requiring corrections only to a two-dimensional SWH array at a given time, also significantly reducing the complexity of the model.

5 Concluding Remarks

In this study, a global-scale AI model for SWH is proposed. The model takes the current SWH field and the wind field at the next time step as inputs, and outputs the SWH field at the subsequent time step. Such a rolling computation method is similar to that used in NWMs, but the spectral information is not used.

As expected, the lack of spectral data leads to an increase in model error during the early stages of the rolling simulation when given a good-quality initial SWH field. However, the rate of error growth slows as the simulation progresses, nearly halting after ~200 hours. More surprisingly, once the error stabilizes, its overall magnitude is not significantly larger than that of state-of-the-art NWMs, particularly under wind-sea-dominated conditions. Although the performance of the AI model in swell-dominated regions is somewhat inferior to that of NWMs, it still produces meaningful outputs, with a correlation coefficient (CC) exceeding 0.7. This suggests that a simple AI model, using only the current SWH field and the wind field at the next time step as inputs, can be practical for many applications, including operational forecasting, at least in regions outside of swell pools.



420 Additionally, this study demonstrates that the issue of error divergence can be partially mitigated through the assimilation of
altimeter measurements. By using a simple objective analysis method, the assimilation helps the error of the model to stabilize
more rapidly and reduces the magnitude of the stabilized error, resulting in a more reliable AI SWH model.

425 An important advantage of the AI SWH model proposed here is its low computational cost compared to traditional NWMs.
This makes the AI model particularly valuable in time-sensitive and resource-constrained scenarios, where it can be used as a
surrogate for the NWMs. One potential application of this model is ensemble modeling, both in operational wave forecasting
and wave climate studies. In these applications, it is challenging to run NWMs multiple times using wind fields from different
ensemble members of weather forecast models (for wave forecasting) or of various climate scenarios for long-term projection
(for wave climate projection). In contrast, these tasks can be efficiently completed using the AI model, even on a standard
laptop.

430 There are many directions for future work. At this stage, the AI model is trained only on SWH data, limiting its applicability
to other wave parameters, such as mean wave periods. Developing an AI model for these additional wave parameters would
require training from scratch with the relevant data. Whether the current model framework, using the corresponding wave
parameter at the current time step and the wind field at the next time step as inputs, can be extended to these parameters remains
to be tested, which can be one future direction. While we acknowledge the potential for a more refined deep learning
architecture to marginally improve model performance, we believe the bottleneck of the current AI model lies in the physics
of the input-output relationship. Therefore, it is difficult to further improve the model performance without changing the model
inputs.

440 We have demonstrated that the current SWH field and the wind field at the next time step are minimum requirements for the
inputs of an AI SWH model. Such simplicity of model inputs and outputs makes this model a potential baseline for AI-based
modeling of global SWH. A promising future direction of this work involves incorporating additional inputs, such as ocean
currents and sea ice, into the model. The most ambitious version of this approach, as mentioned in the introduction, would
involve using global directional wave spectra at the current time step and the wind field at the next time step as inputs, with
the global directional wave spectra at the next time step as the output. While training such a model would be challenging, it is
not an impossible task, and the rapid advancements in AI may make this goal more achievable in the future.

Author contributions

450 Conceptualization: HJ, JW, WL, CD, HQ
Methodology: XW, HJ



Investigation: XW

Visualization: XW

Supervision: HJ

Writing—original draft: HJ, XW

455 Writing—review & editing: HJ, JW, WL, CD, HQ

Competing interests

All authors declare that they have no competing interests.

460 Acknowledgments

This work was jointly supported by the National Key Research and Development Program of China (2023YFC3008203), the National Natural Science Foundation of China (42376172), and the Guangdong Basic and Applied Basic Research Foundation (2022A1515240069, 2024A1515012032, 2023A1515240047).

465 Open Research

The ERA5 data is downloaded from Copernicus Climate Data (<https://cds.climate.copernicus.eu/>). The CCI-Sea State dataset is downloaded from Centre for Environmental Data Analysis (<https://archive.ceda.ac.uk/>). The WW3-ST6 dataset is available from Liu et al (2021). The AI models established in this study are available from the git-hub repository (<https://github.com/YulKeal/AI-Rolling-Wave-Height-Model>).

470

References

- Alday, M., Accensi, M., Arduin, F., and Dodet, G.: A global wave parameter database for geophysical applications. Part 3: Improved forcing and spectral resolution, *Ocean Modelling*, 166, 101848, <https://doi.org/10.1016/j.ocemod.2021.101848>, 2021.
- 475 Arduin, F., Stopa, J. E., Chapron, B., Collard, F., Husson, R., Jensen, R. E., Johannessen, J., Mouche, A., Passaro, M., Quartly, G. D., Swail, V., and Young, I.: Observing Sea States, *Front. Mar. Sci.*, 6, 124, <https://doi.org/10.3389/fmars.2019.00124>, 2019.
- Bi, K., Xie, L., Zhang, H., Chen, X., Gu, X., and Tian, Q.: Accurate medium-range global weather forecasting with 3D neural networks, *Nature*, 619, 533–538, <https://doi.org/10.1038/s41586-023-06185-3>, 2023.
- 480 Booij, N., Ris, R. C., and Holthuijsen, L. H.: A third-generation wave model for coastal regions: 1. Model description and validation, *J. Geophys. Res.*, 104, 7649–7666, <https://doi.org/10.1029/98JC02622>, 1999.



- Cao, H., Liu, G., Huo, J., Gong, X., Wang, Y., Zhao, Z., and Xu, D.: Multi factors-PredRNN based significant wave height prediction in the Bohai, Yellow, and East China Seas, *Front. Mar. Sci.*, 10, 1197145, <https://doi.org/10.3389/fmars.2023.1197145>, 2023.
- 485 Chen, G., Chapron, B., Ezraty, R., and Vandemark, D.: A Global View of Swell and Wind Sea Climate in the Ocean by Satellite Altimeter and Scatterometer, *J. Atmos. Oceanic Technol.*, 19, 1849–1859, [https://doi.org/10.1175/1520-0426\(2002\)019<1849:AGVOSA>2.0.CO;2](https://doi.org/10.1175/1520-0426(2002)019<1849:AGVOSA>2.0.CO;2), 2002.
- Dodet, G., Piolle, J.-F., Quilfen, Y., Abdalla, S., Accensi, M., Arduin, F., Ash, E., Bidlot, J.-R., Gommenginger, C., Marechal, G., Passaro, M., Quartly, G., Stopa, J., Timmermans, B., Young, I., Cipollini, P., and Donlon, C.: The Sea State CCI dataset
490 v1: towards a sea state climate data record based on satellite observations, *Earth Syst. Sci. Data*, 12, 1929–1951, <https://doi.org/10.5194/essd-12-1929-2020>, 2020.
- Gao, Z., Liu, X., Yv, F., Wang, J., and Xing, C.: Learning wave fields evolution in North West Pacific with deep neural networks, *Applied Ocean Research*, 130, 103393, <https://doi.org/10.1016/j.apor.2022.103393>, 2023.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers,
495 D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., De Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.: The ERA5 global reanalysis, *Quart J Royal Meteorol Soc*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- 500 Jiang, H.: Evaluation of altimeter undersampling in estimating global wind and wave climate using virtual observation, *Remote Sensing of Environment*, 245, 111840, <https://doi.org/10.1016/j.rse.2020.111840>, 2020.
- Jiang, H.: Random, Environmental, and Representativeness Errors in Ocean Remote Sensing Versus In Situ Data: An Example of Wave Heights From Altimeters, *IEEE Trans. Geosci. Remote Sensing*, 61, 1–13, <https://doi.org/10.1109/TGRS.2023.3285348>, 2023.
- 505 Jiang, H. and Mu, L.: Wave Climate from Spectra and Its Connections with Local and Remote Wind Climate, *Journal of Physical Oceanography*, 49, 543–559, <https://doi.org/10.1175/JPO-D-18-0149.1>, 2019.
- Jiang, H., Mironov, A., Ren, L., Babanin, A. V., Wang, J., and Mu, L.: Validation of Wave Spectral Partitions From SWIM Instrument On-Board CFOSAT Against In Situ Data, *IEEE Trans. Geosci. Remote Sensing*, 60, 1–13, <https://doi.org/10.1109/TGRS.2021.3110952>, 2022.
- 510 Lam, R., Sanchez-Gonzalez, A., Willson, M., Wirnsberger, P., Fortunato, M., Alet, F., Ravuri, S., Ewalds, T., Eaton-Rosen, Z., Hu, W., Merose, A., Hoyer, S., Holland, G., Vinyals, O., Stott, J., Pritzel, A., Mohamed, S., and Battaglia, P.: Learning skillful medium-range global weather forecasting, *Science*, 382, 1416–1421, <https://doi.org/10.1126/science.adi2336>, 2023.
- Liu, Q., Babanin, A. V., Rogers, W. E., Zieger, S., Young, I. R., Bidlot, J., Durrant, T., Ewans, K., Guan, C., Kirezci, C., Lemos, G., MacHutchon, K., Moon, I., Rapizo, H., Ribal, A., Semedo, A., and Wang, J.: Global Wave Hindcasts Using the



- 515 Observation-Based Source Terms: Description and Validation, *J Adv Model Earth Syst*, 13, e2021MS002493, <https://doi.org/10.1029/2021MS002493>, 2021.
- Ouyang, L., Ling, F., Li, Y., Bai, L., and Luo, J.-J.: Wave forecast in the Atlantic Ocean using a double-stage ConvLSTM network, *Atmospheric and Oceanic Science Letters*, 16, 100347, <https://doi.org/10.1016/j.aosl.2023.100347>, 2023.
- Ronneberger, O., Fischer, P., and Brox, T.: U-Net: Convolutional Networks for Biomedical Image Segmentation, in: *Medical Image Computing and Computer-Assisted Intervention – MICCAI 2015*, vol. 9351, edited by: Navab, N., Hornegger, J., Wells, W. M., and Frangi, A. F., Springer International Publishing, Cham, 234–241, https://doi.org/10.1007/978-3-319-24574-4_28, 2015.
- 520 Song, Y. and Jiang, H.: A Deep Learning–Based Approach for Empirical Modeling of Single-Point Wave Spectra in Open Oceans, *Journal of Physical Oceanography*, 53, 2089–2103, <https://doi.org/10.1175/JPO-D-22-0198.1>, 2023.
- 525 WAVEWATCH III® Development Group: User Manual and System Documentation of WAVEWATCH III® Version 5.16, NOAA/NWS/NCEP/MMAB, Coll. Park, MD, USA, 2016.
- Zhou, S., Xie, W., Lu, Y., Wang, Y., Zhou, Y., Hui, N., and Dong, C.: ConvLSTM-Based Wave Forecasts in the South and East China Seas, *Front. Mar. Sci.*, 8, 680079, <https://doi.org/10.3389/fmars.2021.680079>, 2021.