

A new parameterisation for homogeneous ice nucleation driven by highly variable dynamical forcings

Alena Kosareva¹, Stamen Dolaptchiev¹, Peter Spichtinger², and Ulrich Achatz¹

¹Institut für Atmosphäre und Umwelt, Goethe-Universität Frankfurt, Frankfurt am Main, Germany

²Johannes Gutenberg-Universität Mainz, Mainz, Germany

Correspondence: Alena Kosareva (kosareva@iau.uni-frankfurt.de)

Abstract. The present work aims to extend the parameterisation of homogeneous ice nucleation introduced in Dolaptchiev et al. (2023) by incorporating variable ice mean mass and generalizing the approach under different conditions. The proposed method involves introducing an empirically derived correction to the parameterisation based on a large data set of parcel model simulations to account for the mean ice particle mass variations. The method is validated against ensemble simulations using time-resolved double-moment ice microphysics, showing a mean deviation of less than 16% ~~from the reference solution~~, with robust performance across a range of conditions. The uncertainty of the ~~extended~~-proposed parameterisation is evaluated for ~~the~~ increasing integration time steps. The method remains computationally efficient and produces sufficiently accurate results, even with larger time steps, making it suitable for integration into numerical weather prediction and climate models. It is shown that the generalized approach not only provides a good representation of individual nucleation events but also effectively captures the statistics across the ensemble data. The prediction of the ice mixing ratio is also assessed against the reference ~~full~~-double-moment system results. ~~Despite a significant error in the initial prediction, it~~ It is demonstrated that ~~the integration of the system over several time steps equilibrates the inconsistencies~~, following the nucleation event and re-initialization of the ice mixing ratio, the system equilibrates toward the reference solution within a few time steps. This refined parameterisation offers a more accurate prediction of ice number concentration and ice mixing ratio and is not limited to gravity ~~wave-induced~~-wave-induced perturbations and can be supplemented by other relevant dynamical effects, such as large scale motions or even turbulence.

1 Introduction

Cirrus clouds correspond to up to 30% of the total cloud cover, ~~however~~; however, they are still not well understood and their overall impact on climate in particular on the radiation forcing remains highly uncertain (e.g., Gasparini et al. (2018); Joos et al. (2014); Boucher et al. (2013)). The albedo effect and the greenhouse effect of these clouds are similar in magnitude, meaning that the microphysical details ~~such as~~, such as the number and shape of the ice crystals, can significantly influence their net radiative effect (Krämer et al. (2020); Matus and L'Ecuyer (2017); Wang et al. (2020); Zhang et al. (1999)) (Krämer et al., 2020; Matus and L'Ecuyer, 2017; Wang et al., 2020; Zhang et al., 1999). Some studies also indicate the indirect effects of cirrus clouds on future climate (e.g. Gettelman et al. (2012)) (e.g. Gettelman et al., 2012), which occurs mostly due to increases in ice crystal concentrations due to homogeneous nucleation from anthropogenic sulfur emissions.

25 These microphysical characteristics of cirrus are affected by a complex interaction of the small-scale cloud processes and ~~various interactions with the~~ the dynamics of the surrounding atmosphere. Observations suggest that gravity wave (GW) dynamics significantly impact the properties and life cycle of cirrus clouds (~~Kärcher and Ström (2003); Atlas and Bretherton (2023)~~) (Kärcher and Ström, 2003; Atlas and Bretherton, 2023).

Vertical velocity perturbations, associated with gravity waves, turbulence, or other processes, lead to the homogeneous freezing of aqueous solution droplets (~~Koop et al. (2000); Baumgartner et al. (2022)~~) at lower (Koop et al., 2000; Baumgartner et al., 2022; Kärcher et al. (2002)) at low temperature conditions in the tropopause region. The effect of ~~the~~ gravity waves and other local dynamical processes on the nucleation has been widely investigated ~~utilising different approaches for description of~~ utilizing different approaches to describe the updrafts. Those interactions are studied, for example ~~in~~ Dinh et al. (2016); Kärcher and Podglajen (2019) utilising, in Dinh et al. (2016) and Kärcher and Podglajen (2019) utilizing a stochastic approach, in ~~Joos et al. (2008)~~ Joos et al. (2008) using a GW linear theory, and using the large-scale grid-resolved updraft velocity alongside ~~with sub-grid-scale turbulence induced~~ subgrid-scale turbulence-induced updrafts in Zhou et al. (2016).

Based on the results from Baumgartner and Spichtinger (2019) studying the homogeneous nucleation due to constant updraft velocities, Dolaptchiev et al. (2023) ~~is~~ has extended the asymptotic approach to incorporate GW dynamics. The present work complements Dolaptchiev et al. (2023) and has been designed ~~to for~~ further usage in connection with GW parameterisations for a more realistic representation of GW-cirrus interactions in coarse-grid models. Recently, transient and lateral-propagation effects have been introduced into GW parameterisations (~~Kim et al. (2021); Achatz et al. (2023); Voelker et al. (2024)~~). ~~Corresponding parameterisations bear the potential to provide~~ (Kim et al., 2021; Achatz et al., 2023; Voelker et al., 2024), offering further potential for a more realistic ~~representation of GW-cirrus interactions~~ simulation of these interactions.

~~It was previously shown (Gierens et al. (2003)) that the deposition coefficient largely affects the nucleation process, while at the same time being dependent on the radius or mass of ice crystals. Mean mass variability have~~ The mean mass variability of the ice particles has a significant effect on the cloud ice number concentration prediction, which is also observed in numerical parcel model simulations in Dolaptchiev et al. (2023). The influence of a cloud ice particle size uncertainty in a climate model was assessed in Wang et al. (2020) ~~and~~ illustrates the sensitivity of the surface precipitation to the change in particle effective radius ~~due to the larger energy imbalance,~~ as well as spatial variability of short and long wave radiation. The effects of variable mass become more prominent in ~~the conditions of~~ conditions of a higher number of pre-existing ice particles. Finally, it was previously shown (Gierens et al., 2003) that the deposition (or accommodation) coefficient, controlling the growth of ice crystals in the kinetic regime, largely affects the nucleation process, while at the same time being dependent on the radius or mass of ice crystals.

The current work focuses on generalizing the approach introduced in Dolaptchiev et al. (2023) to include mean mass variability for a wide range of initial conditions, e.g., dynamical forcing and environmental conditions. The asymptotic approach is supplemented by an empirical correction ~~which that~~ can be used for various forcing representations. The proposed method is not restricted to the ~~GW-induced~~ GW-induced dynamics and can be used for any kind of air motion, as e.g., large-scale updrafts or even turbulent fluctuations. This work is also relevant for a further implementation into a global numerical weather prediction (NWP) model, where it can be coupled to the dynamical parameterisations (e.g., GW and turbulence). However, due

60 to the generalized formulation of the approach and robustness for larger time steps, it can also be applied to climate models with large time steps.

The article is organized as follows: ~~The~~ Sect. 2.1 provides a brief overview of the approach used and the motivation for its generalization. Sections 2.2 and 2.3 describe the chosen approach for constructing the parameterisation and technical details of the used ~~data-set~~dataset. In the Sect. 3, the proposed parameterisation with ~~corrections~~correction is verified against reference
65 calculations, and the range of applicability is tested based on ensemble calculations. Concluding remarks can be found in Sect. 4.

2 Theory and methodology

2.1 Double-moment approach to ice physics representation coupled to GW

In the present work, cirrus clouds are modeled using a double-moment bulk microphysics scheme (~~Spichtinger et al. (2023b)~~
70 ~~) that assumes a (based on Spichtinger et al., 2023a) that assumes an~~ unimodal ice mass distribution function, where ~~first and second the zeroth and first~~ moments of the distribution function are ice ~~mixing-ratio q_i and ice~~ number concentration n_i and ice mixing ratio q_i , respectively. Ice crystals are assumed to have a spherical shape, which simplifies the description of the properties. The mentioned assumption is reasonable for very small, newly nucleated ice particles formed through homogeneous nucleation. However, as ice particles grow through deposition, their shape becomes more complex, introducing
75 some uncertainty in the calculations. For consistency, we maintain the same shape assumption in deposition calculations, acknowledging that this introduces a certain error. Although a more sophisticated shape approximation would be beneficial for studying ice growth dynamics, this does not affect the developed correction and goes beyond the topic of the current study; former nucleation parameterisations also assume spherical ice particles at their early stage of life (see, e.g., Kärcher and Lohmann, 2002; Re
The ice physics scheme used here is described in ~~Spichtinger and Gierens (2009); Spreitzer et al. (2017)~~ Spichtinger and Gierens (2009); S
80 with the addition of ~~sedimentational~~sedimentation sinks. The influence of Lagrangian pressure and temperature variations, e.g., due to GWs, on homogeneous nucleation and as well as mean mass variability, are incorporated in Dolaptchiev et al. (2023). The resulting system of equations for n_i, q_i and saturation ratio over ice S reads:

$$\frac{Dn}{Dt} \frac{Dn_i}{Dt} = J \exp(B(S - S_c)) + \frac{1}{\rho} \frac{\partial(\rho n v_n)}{\partial z}, \quad (1)$$

$$\frac{Dq}{Dt} \frac{Dq_i}{Dt} = \frac{p_{si}}{p} D \left(\frac{q}{n} \frac{q_i}{n_i} \right)^{1/3} (S - 1) T n_i + \frac{J \exp B(S - S_c) \hat{m}}{J \exp(B(S - S_c))} + \frac{1}{\rho} \frac{\partial(\rho q v_q)}{\partial z}, \quad (2)$$

$$85 \quad \frac{DS}{Dt} = -D \left(\frac{q}{n} \frac{q_i}{n_i} \right)^{1/3} (S - 1) T n_i - \frac{p}{p_{si}} J \exp(B(S - S_c)) - S \frac{1}{\pi} \frac{D\pi}{Dt} \left(\frac{L_i}{T R_v} - \frac{c_p}{R} \right). \quad (3)$$

Here ~~saturation-ratio~~the saturation ratio over ice is described as $S = \frac{p_v}{p_{si}}$, p_v is the water vapor pressure, p_{si} is the saturation pressure over ice, p is the pressure, \hat{m} is the mean mass of newly nucleated ice particles, J is a homogeneous nucleation rate, including also the number concentration of solution droplets, B is a nucleation parameter, S_c is a critical saturation

ratio, D is a ~~deposition-coefficient~~, growth coefficient, including the diffusion coefficient and the Howell factor, v_n, v_q are sedimentation velocities, T is the temperature, ρ is the density, π is the ~~Exner-pressure~~ Exner pressure, L_i denotes the latent heat of sublimation, c_p is the specific heat capacity of air at constant pressure, R_v is the ideal gas constant for water vapor, and R is the ideal gas constant for dry air, respectively. The values of the used parameters are presented in Table A1. In general formulation, the critical value of the saturation ratio over ice is temperature dependent; in the current work, the value is kept constant while other studies (such as Spichtinger et al. (2023a)) show an insignificant influence of S_c on nucleation processes. Equations (1)-(3) are written in terms of material derivative $\frac{D}{Dt}$, i.e., all quantities are described following the motion of air parcels. ~~Right-hand~~ The right-hand parts of the system of equations describe the ice crystal growth due to the deposition of water vapor, the formation of new ice crystals through homogeneous nucleation ~~and~~, and the sedimentation of the ice crystals due to gravity. The ~~later-latter~~ is not included ~~into-in~~ the consideration further because of the much larger sedimentation time scale (see Dolaptchiev et al. (2023)).

In Baumgartner and Spichtinger (2019) ~~realistic-~~, detailed air parcel simulations are conducted using a box model and a bulk microphysics scheme (based on Spichtinger and Gierens, 2009), showing that for various environmental conditions, assuming a constant mean mass is generally valid ~~during long periods over the duration~~ of the nucleation. However, in case of pre-existing ice particles just before nucleation, when the saturation ratio exceeds one, the ice crystals' mean mass increases, leading to a higher deposition rate. This increased deposition affects the saturation ratio over ice (as a sink), which in turn influences the number of nucleated ice crystals, ~~reducing the actual nucleation rate~~. Given that it is crucial to include these effects in the current model, where the mean mass of ice crystals can be determined from the relation $m_{mean} = q_i/n_i q_i/n_i$. This is achieved by incorporating the evolution equation for q_i and substituting $m(t) = q/n$ $m(t) = q_i/n_i$ in the diffusional growth term.

Time scale separation is employed in Dolaptchiev et al. (2023) to assess the different processes and link the ice physics equations with the effects of GWs. The details of the coupling between the processes' time scales and assumptions to the GW theory are presented in the Appendix.

Further study is done considering the ice physics of a single air parcel influenced by GW dynamics, adopting a Lagrangian framework. In this case, material derivatives in the system of equations are replaced by time derivatives $\frac{D}{Dt} = \frac{d}{dt}$.

As shown in Dolaptchiev et al. (2023), considering only dominant processes, the system of Eqs. (1)-(3) can be reduced to the following form:

$$\frac{dn}{dt} \frac{dn_i}{dt} = J \exp(B(S - S_c)), \quad (4)$$

$$\frac{dq}{dt} \frac{dq_i}{dt} = \frac{p}{p_{si}} \frac{p_{si}}{p} D \left(\frac{q}{n} \frac{q_i}{n_i} \right)^{1/3} (S - 1) T n_i, \quad (5)$$

$$\frac{dS}{dt} = -D \left(\frac{q}{n} \frac{q_i}{n_i} \right)^{1/3} (S - 1) T n_i + S F(t). \quad (6)$$

Where ~~A forcing term~~ $F(t)$ ~~is a forcing term~~ incorporating constant background updraft and GWs dynamics is written in the following form

$$F(t) = \frac{gL_i}{c_p R_v T^2} \left(w_{00} + \sum_j \hat{w}_j \cos(\omega_j t + \phi_j) \right),$$

where g is the acceleration due to gravity, w_{00} is a constant background updraft, \hat{w}_j , Ω_j , ϕ_j are the vertical wind amplitude, frequency, and phase of GW number j . It is worth mentioning that the application of the full system of Eqs. (1)-(3) is not a standard approach, whereas the reduced system (4)-(6) is close to the reduced model in Baumgartner and Spichtinger (2019), and variations of it are commonly used in NWP and climate applications, however including the ice mass concentration.

Referring to the previous results from Dolaptchiev et al. (2023), in the case of constant mass, the asymptotic technique can be applied ~~in order~~ to obtain the prototype ~~parameterisation~~. ~~The nucleation event is~~ parameterization of the homogeneous nucleation process. By employing asymptotic expansions, the authors systematically decompose the governing equations into different scale regimes corresponding to the key microphysical processes: nucleation and deposition. In the nucleation regime, where supersaturation rapidly increases due to gravity wave-induced cooling, they derive an analytical solution for the evolution of ice crystal number concentration. In the growth (or deposition) regime, where ice crystals grow primarily by vapor deposition, the governing equations are solved separately under the assumption of gradual saturation adjustment. The solutions from both regimes are then matched using asymptotic matching techniques. This approach results in a reduced analytical expression for the number of nucleated ice crystals per nucleation event, explicitly linking it to wave parameters (such as frequency and amplitude) and background atmospheric conditions (such as temperature and humidity). The nucleation event is roughly determined where the saturation ~~ration ratio~~ reaches the critical value S_c , and the predicted value for the ice number concentration is found from:

$$N_{post} = \begin{cases} 2 \frac{S_c F(t_0)}{D m_c^{1/3} T (S_c - 1)} - n_{pre}, & \text{if } n_{pre} < \frac{S_c F(t_0)}{D m_c^{1/3} T (S_c - 1)}; \\ n_{pre}, & \text{otherwise.} \end{cases} \quad (7)$$

Here $D^{(*)} = D m_c^{1/3}$ ~~is a deposition coefficient in case of constant mean mass with reference value~~ $m_c = 10^{-12}$ kg ~~is the reference mass the value~~, n_{pre} , n_{post} are the ice number concentrations before and after the nucleation event, t_0 is the time of a nucleation event when $S = S_c$. Note that nucleation already occurs for values of S close to S_c , see asymptotic analysis in Spichtinger et al. (2023a).

In Dolaptchiev et al. (2023), the asymptotic approach was extended to account for the variable mass in the ~~deposition~~ ~~coefficient~~, ~~in depositional growth coefficient in (5), (6) under~~ the regime of slowly varying ice mixing ratio q_i during the nucleation event. In such a case, the different processes can be decoupled, and only n_i undergoes a quick change during the nucleation regime.

Numerical studies of the reduced system (4)-(6) based on a simplified parcel model showed several possible regimes relevant ~~for to~~ the nucleation process (see Fig. 1). One regime ~~under the conditions of at~~ high values of the pre-existing ice crystal number concentration (shown in Fig. 1 in black) follows the same tendency as a constant mass case. The evolution of parameters can be separated into three phases (see Baumgartner and Spichtinger (2019)), where the nucleation part ~~is describing only the~~




Figure 1. Evolution of the ice number concentration n_i , ice mixing ratio q_i , saturation ratio over ice S and mean mass of ice particles $m = q_i/n_i$ calculated based on system (4)-(6). Two cases of initial conditions: low pre-existing ice $n_{init} = 10^2 \text{ kg}^{-1}$ and higher values of pre-existing ice $n_{init} = 10^6 \text{ kg}^{-1}$. The blue line corresponds to the critical saturation ratio, dashed lines mark the time of the nucleation event, where $S = S_c$ in red for the case $n_{init} = 10^2 \text{ kg}^{-1}$, in black for the case $n_{init} = 10^6 \text{ kg}^{-1}$.

describes only the phase of rapid change of ice number concentration. Evolution of the ice number concentration n_i , ice mixing ratio q_i , saturation ratio over ice S and mean mass of ice particles $m = q_i/n_i$ calculated based on system (4)-(6). Two cases of initial conditions: low pre-existing ice $n_{init} = 10^2$ and higher values of pre-existing ice $n_{init} = 10^6$. Whereas In contrast, the saturation ratio and ice mixing ratio change slowly during this short time, and therefore can be assumed to not vary during this phase. However, conditions of low-to-zero pre-existing ice number concentrations (shown in red) lead to a different behaviour due to the deposition-growth coefficient change. Assumptions of the time separation for the nucleation regime do not hold, since the ice mixing ratio is changing on the same time scale as the ice number concentration. The extension to be proposed here aims to cover the whole range of initial conditions allowing to use, allowing the use of the parameterisation accounting for mass variability independently from the regime.

2.2 Correction of the deposition-growth coefficient for variable mean mass

In the present study we are utilising, we are utilizing the previously obtained result for the constant mass case (7). This approach would allow for a clear physical relation between N_{post} and n_{post} , the initial number concentration n_{init} , and the forcing term $F(t_0)$ at the point t_0 of the nucleation event. It would also account for different background conditions for temperature by design. Since mass variability changing the deposition changes the growth coefficient directly, as one can see from (6), the straightforward solution is to correct the deposition-growth coefficient in the constant mass parameterisation to a factor, representing the mass at the point of the nucleation event. The corrected parameterisation has the following form:

$$Nn_{post} = \begin{cases} 2 \frac{S_c F(t_0)}{D(m_{eff})^{1/3} T(S_c - 1)} - n_{pre}, & \text{if } n_{pre} < \frac{S_c F(t_0)}{D(m_{eff})^{1/3} T(S_c - 1)}; \\ n_{pre}, & \text{otherwise,} \end{cases} \quad (8)$$




Figure 2. Evolution of n_i, q_i, S, m in the reduced system (4)-(6) case and in the pre-nucleation regime, solving (5), (6) with assumption of $n = n_{init} = const, n_i = n_{init} = const$. The blue line corresponds to the critical saturation ratio, the dashed lines mark the time of the nucleation event, where $S = S_c$.

where $D^{(**)} = D\left(\frac{q_0}{n_0}\right)^{1/3}$ is a deposition coefficient corrected by a factor $(m_{eff})^{1/3}$ is a growth coefficient correction factor representing the effective mean mass of the ice crystals at the start of $\left(\frac{q_0}{n_0}\right)$, t_0 is the time of the nucleation event and $q_0 = q(t_0)$, $n_0 = n(t_0)$.

In order to find the ~~the desired deposition coefficient~~ desired growth coefficient, one needs to define the parameters ~~which~~ that can be used for the correction. Figure 2 shows the evolution of the parameters calculated from the reduced system (4)-(6) and known pre-nucleation regime parameters, which can be used for deriving the correction and parameterisation. Here a pre-nucleation regime is defined as solution of the (4)-(6) system in the regime, where initial ice number concentration stays constant, which leads to solution of Eqs. (5),(6) with $n = n_{init} = const$ decoupled from Eq. 4. Sharp changes in the ice mixing ratio and mass in ~~case of the case of a~~ reduced system make it impossible to approximate the necessary ~~q_0 or $\frac{q_0}{n_0}$~~ quantities $q_i(t_0)$ or $\frac{q_i(t_0)}{n_i(t_0)}$ from pre-nucleation regime values itself. Therefore ~~the correction to the deposition coefficient~~, the introduced effective mass is approximated based on a large data set. Variables ~~which that~~ which that can be used in the approximation include the forcing term at the point of the nucleation event $F(t_0)$, initial number the initial number concentration n_{init} and, and the mass m^* predicted from the pre-nucleation regime, assuming no change in ice number concentration before the nucleation event.

The exact ~~correction m_0~~ effective mass m_{eff} is determined from the results of the full system, such that the constant-mass approach leads to the observed ~~$N_{post} n_{post}$~~ . Assuming a known evolution of the parameters during the nucleation event, Eq. (8), under the conditions allowing for further nucleation, can be solved for ~~m_0~~ m_{eff} :

$$180 \quad m_{0eff} = \left(\frac{2S_c F(t_0)}{(N_{post} + N_{pre})D(S_c - 1)} \frac{2S_c F(t_0)}{(n_{post} + n_{pre})DT(S_c - 1)} \right)^3, \quad (9)$$

where $t_0, N_{pre}, N_{post}, n_{pre}, n_{post}$ are known from the resolved full system (1)-(3). A similar approach was used in Baumgartner and Spichti, where the increase of the ice mass during the nucleation was found to be linear with a slope proportional to some constant mean mass.

Next, we discuss the construction of the data set from ~~the ensemble calculations based on ensemble simulations based on~~
 185 ~~Eqs. (1)-(3) disregarding~~, disregarding the sedimentation process. ~~This simplification can be justified by the small sizes of ice~~
~~particles during and shortly after the nucleation event.~~ The variability during the nucleation event depends on the initial condi-
 tions, such as ice number concentration, ice mixing ratio, or mean mass of ice particles, and on the ~~gravity waves~~ GW forcing.
 Since we want to represent the nucleation process itself, the initial saturation ratio is set to a value of $S(t=0) = 1.4$, close to
 the critical value. Other ~~initial~~ conditions are varied randomly using a uniform distribution in ~~logarithmic space, i.e., $\log(n_{init})$~~
 190 ~~in~~ physically meaningful ranges motivated ~~from the paper (Krämer et al. (2016))~~ by observations (Krämer et al., 2016): we
 choose $10^{-4} \text{ kg}^{-1} < n_{init}(t=0) < 10^7 \text{ kg}^{-1}$, and $10^{-16} \text{ kg} < m_{mean}(t=0) < 10^{-12} \text{ kg}$. ~~The lower bound for the initial~~
~~number concentration is chosen to represent the conditions of the low-to-no pre-existing ice, which can occur in applications~~
~~to the global NWP models.~~

Construction of the forcing is done based on the output from the global ICON ~~model (Zängl et al. (2015)) coupled with~~
 195 ~~(ICOsahedral Non-hydrostatic) model (Zängl et al., 2015) coupled with the~~ GW parameterisation Multi-Scale Gravity Wave
 Model (MS-GWaM) (~~Bölöni et al. (2021); Kim et al. (2021); Achatz et al. (2023); Kim et al. (2023); Voelker et al. (2024))~~ (Bölöni et al., 2021).
 This parameterisation is based on ~~WKB theory (Bölöni et al. (2021)) and~~ Wentzel–Kramers–Brillouin (WKB) theory and is
 implemented using Lagrangian ray volumes, which are considered as carriers of the GW fields’ wave-action density (Bölöni et al., 2021).
 The current version of ~~the~~ MS-GWaM allows to account for convectively generated GWs and GWs generated from processes
 200 other than convection or flow over mountains. ~~Influence~~ ~~The influence~~ of the GWs generated by the mountains is incorpo-
 rated by using ~~original scheme (Lott and Miller (1997)) included in the original scheme (Lott and Miller, 1997) included in the~~
 ICON model. For constructing the forcing term, we utilize ICON version 2.6.5-nwp1, with 120 vertical levels and at horizontal
 resolution R2B5 ($\sim 80 \text{ km}$); ~~the~~ km). The initial conditions are taken for ~~the~~ 1st of May 2010, and the model is integrated
 forward for 3 weeks, accounting for 2 weeks of spin-up. Information on subgrid-scale gravity wave related vertical wind per-
 205 turbations is retrieved from MS-GWaM using the ray volume with ~~the~~ maximum vertical wind amplitude per cell and ~~the~~
 corresponding frequency. In order to cover the variety of possible forcing term configurations, the following cases are included
 in the data set construction:

1. Large-scale vertical velocity from ICON run w_{00} , in such a case the forcing term reads: $F(t) = \frac{gL_i}{c_p R_v T^2} w_{00}$;
2. ~~Single~~ ~~A single~~ GW from MS-GWaM with maximum wind amplitude \hat{w} within the cell, the corresponding frequency ω
 210 and ~~initial~~ phase ϕ which is picked randomly: $F(t) = \frac{gL_i \hat{w}}{c_p R_v T^2} \cos(\omega t + \phi)$;
3. ~~Superposition~~ ~~A superposition~~ of GWs, where the main GW is taken from ~~the~~ MS-GWaM and other GWs have ran-
 domly varying amplitudes A_j , frequencies ω_j and phases ϕ_j independent from ~~the~~ main GW and rescaled in ~~the a~~
 way, that the total momentum flux added is ~~1mPa~~ 1mPa, ~~similar to approach used in Dolaptchiev et al. (2023)~~: $F(t) =$
 $\frac{gL_i \hat{w}}{c_p R_v T^2} \cos(\omega t + \phi) + \sum_j \frac{gL_i A_j}{c_p R_v T^2} \cos(\omega_j t + \phi_j)$;
- 215 4. As in case 2 in the presence of ~~(large scale)~~ background updraft w_{00} :

$$F(t) = \frac{gL_i}{c_p R_v T^2} \left(w_{00} + \hat{w} \cos(\omega t + \phi) \right);$$

5. As in case 3 in the presence of (large scale) background updraft w_{00} :

$$F(t) = \frac{gL_i}{c_p R_v T^2} \left(w_{00} + \hat{w} \cos(\omega t + \phi) + \sum_j \frac{gL_i A_j}{c_p R_v T^2} \cos(\omega_j t + \phi_j) \right).$$

It is worth mentioning that the GW diagnosed from the MS-GWaM model dominates in cases 3 and 5 due to the larger amplitudes of the vertical velocity perturbations.

Distributions of the large-scale vertical winds and vertical winds from the GWs used for constructing the forcing are presented in Fig. 3 (a, b). The probabilities for the corresponding frequency ω are also presented in Fig. 3, as well as those for the calculated GW momentum flux $\rho w w$ for the considered GW (see Fig. 3 (c, d)). The data is taken from the instantaneous output of the global ICON simulation at altitudes from 8 to 14 km, most relevant for cirrus formation. While this range does not explicitly cover the full tropical tropopause layer (TTL), including altitudes up to 18 km, we note that the selected range captures the majority of ice formation conditions observed in the Upper Troposphere/Lower Stratosphere (UTLS). Moreover, the vertical velocity distributions in this range are comparable to those at higher altitudes, and the proposed parameterization is additionally tested under conditions representative of stronger forcing (see Appendix C). These tests suggest that the approach remains robust and physically consistent even beyond the original range of sampled data.

It is seen in Fig. 3 that large-scale updraft is varied the large-scale updraft varies in range from -0.2 ms^{-1} up to 0.4 ms^{-1} , where in the majority of the cells, vertical velocity is close to zero. Therefore, we include cases 2) and 3) without the large-scale updraft when creating the dataset. Vertical wind perturbations \hat{w} obtained from the GW parameterisation show stronger vertical velocity, with a rare maximum above 2 ms^{-1} . Corresponding frequencies ω are sampled in a range between Coriolis parameter $f = 10^{-4} \text{ s}^{-1}$ and buoyancy frequency $N = 2 \cdot 10^{-2} \text{ s}^{-1}$ in the tropopause region and the form of the ω spectrum resembles the Desaubies form (Orr et al., 2010) (VanZandt, 1982). It is worth noting that the resulting momentum flux based on the highest-amplitude ray volume is aligned with the observational statistics obtained from the balloon measurements (Hertzog et al., 2012) (Hertzog et al., 2012) and previously calculated momentum fluxes from the MS-GWaM parameterisation (Kim et al., 2021). Therefore, the total momentum flux considered lies in the physical range, and the forcing term applied in the parcel simulations is not overestimated realistic to some degree. Further support for the realism of the applied conditions comes from Fig. 3 b, where the PDF of gravity wave-induced vertical velocity perturbations \hat{w} . The spread and frequency of occurrence show a good agreement with available observations from balloon measurements (Podglajen et al., 2016; Köhler et al., 2023).

The variation of vertical velocities used in the construction of the forcing is sampled from the wide range of conditions, including the large updrafts up to 10 ms^{-1} . Because of the general model formulation, one can add the vertical velocity contributions to the forcing term based on other information provided by the model, for example, by the turbulence scheme. However, such an extension of the forcings is not tested in the current work.

The exact correction m_0 value of the effective mass m_{eff} is found from Eq. 9 (9) based on the ensemble calculations for $\sim 2 \cdot 10^5$ nucleation events; the results are shown in Fig. 4. The data set is created using random initial conditions for n_i, q_i and various GW forcing configurations as discussed above. The internal variability of the data is quite high, especially depending




Figure 3. Number Frequencies of occurrences of (a) the large-scale vertical velocity w , (b) the vertical velocity amplitude \hat{w} , (c) the frequency of the GWs ω defined from the MS-GWaM parameterisation, and (d) the momentum flux corresponding to the detected GW.




Figure 4. Exact m_0 The exact m_{eff} correction depending on the initial number concentration (a) and on the forcing term value at the nucleation event point (b). Blue markers correspond to calculations with approximately same initial n_i constant n_{init} , black markers correspond to calculation-calculations made with approximately same-constant values of $F(t_0)$.

on the forcing term at the point of the nucleation event. However, by fixing certain parameters (blue and black markers), one can see the-a clear dependence of m_0 from m_{eff} on the chosen parameters. Variability-over-The variability with the forcing term is much larger, whereas under the conditions of smaller initial number concentrations m_0 , m_{eff} stays nearly constant. The dependence on the forcing term is decaying with large variability for smaller values of the forcing at the point of the nucleation event. The increase of the mass with an increase of the initial number concentration is connected to the fact that pre-existing ice suppresses the nucleation process, hence, the number of newly nucleated ice crystals is smaller. Because the initial ice mean mass is quite high, and the new particles are four orders of magnitude lighter, the recalculated mean mass

would increase if more pre-existing ice is present. The observed behavior of the mass in dependence on the forcing term is the opposite, it decreases with increasing $F(t_0)$. A larger forcing term at the point of the nucleation is associated with the larger updrafts leading to a faster change in the saturation ratio over ice. The depletion of the water vapor via deposition in such conditions is ineffective, leading to smaller mass values.

2.3 Mean mass fitting for parameterisation

In the present section, we consider options for approximating the dependence of m_0 on m_{eff} based on the parameters known from the pre-nucleation regime. Since the evolution of the ice mixing ratio q_i is not captured in this regime (due to fixed deposition coefficient and n), one can fit the behaviour of the correction growth coefficient and n_i , we propose to fit a function depending on the known parameters, such as initial ice number concentration n_{init} and forcing term at the point of the nucleation event $F(t_0)$, to the ensemble simulations results. These parameters are selected based on the definition of the correction given in Eq. (9), and their relevance is further supported by the clear dependence observed in Fig. 4, where only a slight spread is visible in cases with fixed n_{init} and $F(t_0)$.

Based on the created data set and relative behaviour of m_0 the relative behavior of m_{eff} in dependence on n_{init} and $F(t_0)$, an approximation to the data is found. The non-linear least squares method is applied to fit an assumed function to the known data from the ensemble simulations. The method requires to suggest a certain functional dependency on the parameters. Given the data set behaviour, the chosen simplified function is:

$$m_{0,eff}(F(t_0), n_{init}) = \exp(a_1 + a_2 F(t_0)^{1/3} + a_3 n_{init}^{1/3} + a_4 n_{init} F(t_0)^{1/3}). \quad (10)$$

The mean error for m_{eff} during the target data set reproduction is around 30% for, which translates into a mean error of $\sim 10\%$ in radius, for the set of $\sim 2 \cdot 10^5$ nucleation events with different GW forcing representations.

As an extension, a more complex dependency on n_{init} , $F(t_0)$, and involving one more parameter m^* , defined from the pre-nucleation regime, is introduced and tested. The results showed an improvement compared to the fit-ease previously introduced fit (10); see the corresponding figures and discussion in the Appendix B. Mean error for While the use of an advanced fit improves the prediction of the effective mass (m_{eff}), the ice number concentration prediction is reduced only by after nucleation (n_{post}), which is the primary target of the parameterization, is already well captured by the simple fit. The more complex approach yields only a marginal improvement of about 2% in the mean error. Besides, the advanced fit adds more complexity to the calculation, requires the use of an additional parameter and because of the additional terms used can lead to diverging results in cases out of sample range with regard to the reference data used for the parameter optimization. Therefore results presented below are obtained with the simplified fit approach only, given the efficiency and robustness of the original simple fit, we recommend its use and present all further analysis based on this version.

3 Results. Evaluation of the Model

Further results are shown for an independent data set consisting of 490 nucleation events ~~in~~ in order to validate the chosen fit. Initial conditions for n_i and m_i were created randomly, and the forcing term is based on ICON simulation ~~is~~ different from the one used for the construction of the fit. In the following, we will first asses the proposed fit. This will be followed by stability and accuracy tests, depending on the time step chosen for the integration of the Eqs. (5)-(6) in the parameterised case.

We acknowledge that developing a parameterization based on a limited dataset is inherently challenging and introduces certain limitations. The considered conditions are representative of the UTLS region and are specific to the processes as captured by the ICON model. As a result, extreme or anomalous cases may be underrepresented. Nevertheless, we argue that the physically grounded formulation of the proposed parameterization, along with its robustness to variations in forcing, allows it to provide reasonable predictions even outside the original fitting range. To support this claim, we validated the parameterization on an independent dataset and included results for selected test cases with unusually strong vertical velocities in the Appendix C.

3.1 Extended parameterisation

~~Comparison~~ A comparison of the application of the chosen fit (10) and the exact ~~effective~~ mass calculated from the nucleation event simulations is shown in Fig. 5. ~~Direct~~ A direct comparison of predictions with the exact values for ~~m_0~~ m_{eff} shows a good agreement ~~of for~~ for the major part of the data points. Nonetheless, the fit is not able to capture the higher possible ~~m_0~~ m_{eff} values at higher ~~initial n values of n_{init}~~ n_{init} . Fig. 6 shows the exact post-nucleation ice number concentration and the one ~~obtained~~ obtained with the use of the extended parameterisation (8). The underestimation of the ~~m_0~~ m_{eff} values at higher values of pre-existing ice ~~number concentrations~~ leads to an overestimation of the ice number concentration as shown in Fig. 6.

Referring to the Fig. 5, the most frequently occurring cases of ~~m_0~~ m_{eff} in the range between 10^{-13} kg and 10^{-12} kg, and the corresponding final ~~N_{post}~~ n_{post} in Fig. 6, in the range between 10^5 and 10^6 , are successfully represented by the chosen fit (10) and ~~the~~ corrected parameterisation (8). However, the outlying extreme cases are poorly captured due to ~~the~~ simplicity of the fit. The overall ~~behaviour~~ behavior of the fit stays similar to the exact data, ~~therefore; therefore,~~ with decreasing ~~n_{init}~~ n_{init} , the prediction for the ~~m_0~~ m_{eff} stays nearly independent from n_{init} (see Fig. 5 (a)). The cases with an increased forcing term correspond to the smaller values of ~~m_0~~ m_{eff} , increasing the final prediction for ~~N_{post}~~ n_{post} in the exact data, and the same ~~behaviour~~ behavior is captured by the fit.

To assess how well the chosen approach represents individual cases, the deviations of ~~m_0 and N_{post}~~ m_{eff} , ~~effective ice~~ particle radius R_{eff} and n_{post} from their exact values are calculated. The ~~effective radius is calculated based on the effective~~ mass as follows: $R_{eff} = \left(\frac{3m_{eff}}{4\pi\rho_i} \right)^{\frac{1}{3}}$. The probability density function (PDF) of the deviations is presented in Fig. 7. Here the deviations are calculated as $\Delta M = (M_{predicted} - M_{exact})/M_{exact} \cdot 100 \%$, with ~~$M = m_0, N_{post}$~~ $M = m_{eff}, R_{eff}, n_{post}$. The mean errors for ~~m_0 and N_{post}~~ m_{eff} and n_{post} predictions are 24% and 16% ~~correspondingly,~~ respectively. Despite the substantial deviation in effective mass, the prediction for the effective radius shows a good agreement and the mean error is only about 10%. More importantly, 90% of the error for ~~the~~ prediction of the ice number concentration lies below 37%.




Figure 5. Exact m_0 - m_{eff} correction depending on the initial number concentration and m_0 - m_{eff} prediction from the constructed fit (10) for independent ensemble calculations.




Figure 6. Post-nucleation ice number concentrations defined from the full system and prediction from the constructed parameterisation with correction (10).

320 **Majority** The majority of the individual cases are represented sufficiently well, whereas outliers and poorly captured events with deviation in N_{post} a deviation in n_{post} above 50% correspond to less than 10% of the distribution.

For a better understanding of the prediction deviations, we also consider the PDF of the exact parameters and the approximated ones in order to assess the ability of our fit to capture the statistical properties of the data set. Previous ensemble simulations using constant mass parameterisation (Dolapchiev et al. (2023)) by Dolapchiev et al. (2023) showed




Figure 7. Probability density functions for deviations of the \bar{m}_0 , m_{eff} , R_{eff} and N_{post} , n_{post} predictions from the exact solutions. Vertical lines correspond to the 90th, 50th and 10th percentiles of the distribution.

that while individual events can vary significantly, or even be missed, the statistical quantities for the ensemble still align well.

Comparisons of the PDFs for the mass \bar{m}_0 correction, effective mass m_{eff} and the predicted ice number concentration and the ice mixing ratio after the nucleation event are shown in Fig. 8. Here the prediction for the ice mixing ratio q_i is made based on the estimated mass \bar{m}_0 and predicted by the corrected parameterisation ice number concentration N_{post} via $q_{i,post} = \bar{m}_0 N_{post}$. Such approach contains some physical inconsistencies, because the approximated ice mean mass \bar{m}_0 differs from the mean mass after the nucleation event, which is supposed to be used. However such estimation gives a suitable approximation for the further simulation of q_i . Further details are shown in Fig. 13 and 14 and discussed later in Sect. 3.2.

The PDF of the predicted mass from Fig. 8 at the point of the nucleation exhibits a slight underestimation of the mean value, due to the inability to correctly predict larger \bar{m}_0 , m_{eff} (also shown in Fig. 5). For the same reason, the tail of the distribution is shorter, and the maximum value has a mismatch of approximately one order of magnitude. The lower values of \bar{m}_0 , m_{eff} are represented better, and the minimum values are in the same range. Comparison of the PDFs of the ice number concentration shows a good correspondence of major statistical indicators, such as mean, maximum and minimum values. However, the extended parameterisation gives occasional overestimation of N_{post} , n_{post} values for larger forcing term values of the forcing term and underestimation of the N_{post} , n_{post} in cases of small forcing (see the right panel in Fig. 6). The comparison of

Apart from the values of n_i , the prediction for the ice mixing ratio PDFs can be considered as rough estimation of the q_i ; which then is integrated over time with the larger time step. Therefore consistent underestimation of the q_i refers to a physical inconsistency discussed above is made based on the estimated mass m_{eff} and the ice number concentration n_{post} , calculated using the corrected parameterisation, via $q_{i,post} = m_{eff} n_{post}$. Such an approach contains some physical inconsistencies because the approximated ice mean mass m_{eff} differs from the mean mass after the nucleation event, which is supposed to be used.




Figure 8. Probability density functions for m_0, N_{post}, m_{eff} and q_i, n_{post} exact ensemble solutions and predictions. Vertical lines correspond to the mean of the distribution.

However, even with a roughly one-order-of-magnitude underestimation, this prediction can still be used for further integration, minimizing the risk of numerical issues related to the rapid increase in the estimation gives a suitable approximation for the further simulation of q_i . Further, the evolution of the parameters for two isolated cases is presented in Figs. 13 and 14 to show the q_i prediction relevance.

3.2 Influence of the time step

The results shown in Sect. 3.1 are conducted using the same time step for the solution of the full system (1)-(3) and coupled system (5),(6) employed in the parameterisation. Such a decision is made in-order to keep the results comparable without introducing further uncertainties. However, the major goal of the parameterisation is to increase the numerical performance, especially by using larger time steps as in coarse resolution models, e.g., NWP and climate models. Therefore, the following section is devoted to the depiction of the uncertainties and evaluation of the results under different used time steps.

The investigation of the stability of numerical integration and applicability of the defined approach is done based on a data set with 6 GWs and a constant background updraft w_{00} . One of the sources of uncertainty is the accuracy of the detection of the forcing term at the point-time of the nucleation where $S = S_c$. The nucleation event can be misplaced, changing the defined time of the nucleation event t_0 , depending on the time step used for the calculation of the saturation ratio evolution (6). The calculation of the forcing term is sensitive to the changes in t_0 , which leads to further errors in the estimation of the $F(t_0)$ and other parameters.

Further, the deviations of the predictions from the exact values ΔM are calculated as mentioned before in Sect. 3.1 for the considered parameters $M = F(t_0), m_0, N_{post}, M = F(t_0), m_{eff}, n_{post}$. First, in Fig. 9 we compare deviations in the forcing $F(t_0)$ when solving slow-scale processes for S, q_i based on system (5) and (6) with five different time steps choices. In Figs.




Figure 9. Probability density function of the deviation of ~~detected from the parameterised approach~~ deviation between the parameterized forcing term $F(t_0)$ and the corresponding value detected from the exact ~~full-system~~ full-system solution in %. Horizontal bars correspond to ~~maximum, the 90, 50, and 10 percentiles and minimum~~ of the distribution.

9-11, horizontal bars represent percentiles of the distribution to illustrate data variability. Note that in several cases, the 10th percentile lies near zero and may not be easily distinguishable in the plots due to scale limitations. The smallest chosen time step of one second gives the exact solution, whereas longer time steps lead to deviations from the exact prediction. Such deviations may influence the estimated $m_0 m_{eff}$ via errors introduced in t_0 and $F(t_0)$ determination. However, the crucial impact it would have on the $N_{post} n_{post}$ prediction by directly changing $F(t_0)$ in the formula (8). Maximum differences above 50% occur rarely and are associated with outliers, for instance cases where, some cases where the nucleation event is missed with the usage of the larger time step. One of such cases such case is illustrated further in this section in Fig. 13. Mean deviations in the forcing term corresponding to the case of 30 seconds time step is second time steps are 2.6%, and the 90th percentile is 3.5%. The mean and the 90th percentile of the PDF are doubled with doubling the time step and rise up to 15% and 31% for the largest considered time step of two minutes. Considering the exact differences instead of relative errors, it is found that larger time steps mostly lead to an underestimation of the forcing. Hence, because of the direct relation to the ice number concentration, one would expect underestimation of the N_{post} with increasing the n_{post} with increasing time step.

Uncertainties from the detection of the forcing term then lead to a variation in $m_0 m_{eff}$, considered in Fig. 10. The overall error we introduce by using the fit is illustrated by the first PDF when the same time step as in the full system is used. Additional The additional deviation introduced by differences in $F(t_0)$ leads to a variation of about 2-3% in the mean error of $m_0 m_{eff}$, which is negligible compared to the overall error introduced by the fit.

Comparing the error of the ice number concentration prediction from the exact solution in Fig. 11, one observes the increase in the errors when going to higher larger time steps. The mean and 90th percentile of the distribution stay nearly the same when increasing the time step from 1 second to 60 seconds. The increase in mean error becomes larger for a time step $dt = 120$ s and at, and at a time step $dt = 240$ s it becomes closer to 50%. We conclude that one can use a time step of 60 seconds without a significant error increase. To estimate the quality of the new parameterisation, one can compare the mean errors to the relative accuracy of measurements (Krämer et al. (2016, 2020)). Mean error achieved with using (see, e.g., Krämer et al., 2016, 2020).

The resulting mean error using a time step of 60 seconds is of the same order as the accuracy of measurements, which is about 20% (Luebke et al. (2016)). Therefore 15% (c.f. Luebke et al., 2016). Therefore, the proposed approach and chosen time step is sufficient are well-suited for the representation of the process.

390 In order to check the dependence of the statistics on the time step used in integrating equations (5),(6), the comparison of the PDFs of m_0 , N_{post} and q_i , m_{eff} and n_{post} is shown in Fig. 12. The overall shape of the approximated PDF of mass m_0 agrees with the exact solution. However, the case with the largest time step of two minutes shows underestimated m_0 an underestimated m_{eff} minimum for several orders of magnitude. This leads to poor representation of the cases with higher N_{post} and consequent overestimation of the N_{post} maximum (see Fig. 12 middle-lower panel). Predictions of the ice number concentrations are not far from the exact full system results if one compares the center of the PDF. The peak of the predicted N_{post} PDFs stays in the range of 2% difference compared to the full system for all the parameterisation cases. However, the prediction made with using the largest time step considered substantially overestimates the maximum N_{post} value. It is worth to mention, mentioning that the ice number concentrations obtained with the parameterisation and full system are in the range of the observed quantities for the homogeneous nucleation (Krämer et al. (2020)) (Krämer et al., 2020).

400 Concerning the ice mixing ratio prediction, only the parameterisation with the finest time step gives underestimated q_i for the full PDF. The PDF of q_i is consistently increasing with increase of the time step, and peaks of the PDFs differ for about 50% for cases with time step of 30 and 60 seconds, and by the factor of 2 when using the 2 and 4 minutes time steps.

Further, we present two nucleation events cases in order to directly compare the evolution and parameterisation prediction with different time step steps (see Figs. 13 and 14). By the means of such comparison, one can make an additional conclusion on the correctness and possibility of use of q_i as mentioned earlier and assess numerical stability of the method under different time steps. Initial conditions for the presented cases are picked from the ensemble calculation with 6 superposed GWs with the presence of a background updraft.

Both cases do not show any difference between the reduced system (4)-(6) and the full system (1)-(3). The corrected parameterisation (8) is tested for larger time step steps of 30, 120, 240 s, omitting the case with the smallest time step. The first case in Fig. 13 illustrates that parameterisation even with larger time steps the parameterization gives a good agreement with the full system, even with larger time steps. The evolution of the saturation ratio over ice, ice mixing ratio and mass are, and mass is aligned with the full system results after some time following the nucleation event. Prediction The prediction of the ice number concentration is not that exact for cases with larger time steps, but the difference is still in the range of 30%. After the prediction of q_i using the m_0 and N_{post} , m_{eff} and n_{post} from the parameterisation one can see that integration is done without numerical problems. Evolution of The evolution of the ice mixing ratio q_i shows a jump due to the nucleation process, followed by a short relaxation period, after which the q_i prediction is adjusted and overlaps with the full system result.

415

The second test case presented in Fig. 14 has been picked on purpose to show several possible outcomes of using the parameterisation and the consequences of usage of larger time steps. The evolution of the slowly-varying parameters such as S , q_i , m is captured relatively well by the parameterisation with the time steps of 30, 120 seconds. The largest considered time step, however, leads to smaller variations in the saturation ratio and hence the nucleation event is not captured by the parameterisation.

420

The parameterisation using a time step of 30 seconds gives a sufficiently good agreement with a deviation from the exact solution not larger than 22%. However, even though the parameterisation with ~~2 minutes~~ a 2-minute time step predicts the nucleation event, the difference in the final ice number concentration is increased to 46%. Other variables and the evolution of the slowly varying part ~~is~~ are described well in cases where the nucleation is captured as such. Considering different cases and referring to the representation of the PDFs ~~the~~ a time step of 60 seconds can be suggested for parameterisation for the optimal result. Further increase in time step leads to ~~the decrease of accuracy~~ a decrease in accuracy, and in cases of ~~a~~ the largest time step of 4 minutes, even to a wrong representation of the process, where some events are not captured. Such behaviour is observed due to the fact that the pre-existing ice grows too fast, depleting the available water vapor and thus preventing the saturation ratio ~~to increase to~~ from increasing to values triggering the nucleation.

4 Conclusions

~~A~~ This study aimed to extend the applicability of the nucleation parameterization Dolaptchiev et al. (2023) to a broader range of atmospheric conditions. Specifically, it addressed the regime of low pre-existing ice, where the rapid evolution of the ice mixing ratio limits the validity of the asymptotic approach used in earlier work. To overcome this limitation, a generalisation of the parameterisation derived in Dolaptchiev et al. (2023), including the effect of variable ice mean mass, is proposed in the article. ~~The employed approach has several advantages compared to~~ Compared to the constant mean mass assumption ~~as the introduced~~ the proposed approach has the advantage of incorporating a physically grounded correction to the ~~deposition coefficient~~ growth coefficient, reflecting the actual variation in ice particle growth under different conditions. The considered method allows ~~to include the~~ for the inclusion of the mean mass variation, which leads to a better representation of the predicted ice number concentration distribution.

The new approach is validated against ensemble simulations based on the full double-moment scheme, showing a good agreement with the exact solution. The representation of individual nucleation events is ~~sufficient~~ done with reasonable accuracy, and the mean deviation from the full system is less than 16% for the ice number concentration, which is similar to the usual accuracy of measurements. A vast ~~amount of cases is represented well~~ number of cases are represented accurately, but the substantial differences remain for the extreme cases of the high forcing term values. The probability density function of the occurring ~~n_i~~ n_i determined from independent ensemble simulations captures the major statistical characteristics ~~such as~~ such as the mean and the center of the PDF.

Because of the computational efficiency demand for ~~the NWP's application~~ NWPs and climate applications, the effect of uncertainties resulting from large ice microphysics time steps in the proposed extended parameterisation is assessed as well. The optimal representation of individual nucleation events combined with the sufficiently well captured statistics of ~~N_{post}~~ n_{post} occurrence is achieved with the $dt = 60$ seconds. This recommendation is comparable ~~with~~ to the fast physics time step of about 4 minutes ~~used in large-scale NWP model~~ typically used in NWP models. Evaluation of the numerical effectiveness shows that parameterisation is several times faster than the solution of the full system with ~~the~~ a time step of 1 second.

Further research involves implementation Given the strong agreement of the proposed approach for describing homogeneous
 455 ice nucleation in a global Numerical Weather Prediction (NWP) model. Besides the advanced description of the physical
process by accounting for the mass variability, it is planned to employ an approach coupled to the dynamical parameterisations.
The method with resolved physics, its robustness, and sufficient accuracy, the approach is now ready for further application
in NWP models, such as ICON. The generalization of the method allows for direct coupling with forcings from dynamical
parameterizations, thereby enhancing its integration into operational systems. For instance, the detailed representation of local
 460 subgrid-scale GWs dynamical fields impacting cirrus clouds is planned to can be incorporated using the Multi-Scale Gravity
 Wave parameterization (MS-GWaM) (Böläni et al., 2021; Voelker et al., 2024) coupled to the described ice nucleation param-
 eterisation. Additionally, this coupling approach can be supplemented by the extended to other relevant dynamical effects, e.g.,
 turbulence.

Code and data availability. The air parcel model source code and Python scripts for processing and plotting the data are available on Zenodo
 465 at <https://doi.org/10.5281/zenodo.14244119>. Data set from ICON simulation used in the construction of the initial conditions for the parcel
 model, as well as all output data used for plots, can be found on Zenodo at <https://doi.org/10.5281/zenodo.13819511>.

Appendix A

Coupling of the ice physics processes to the GW theory is done under the assumptions of mid-frequency and high-frequency
 GWs, where the GW time scale T_w is tied to the Brunt-Vaisala frequency $N = \sqrt{\frac{g}{\theta} \frac{d\theta}{dz}}$ by the following relation:

$$T_w = \frac{1}{\varepsilon^\beta N}, \quad \beta = 0, 1; \quad \varepsilon = 0.1,$$

where $\beta = 0$ characterizes the high-frequency and $\beta = 1$ the mid-frequency GW time scale, ε is an introduced small parameter
used for defining the distinguished limit in the asymptotic analysis. For instance, the characteristic time scales in the condi-
 470 tions of the troposphere, where $N = 10^{-2} \text{ s}^{-1}$, are $T_w = 100$ and $T_w = 1000$ seconds for the high and mid-frequency waves,
 respectively. The diffusional growth time scale is estimated from $T_d = \left(D m_c^{1/3} \frac{p_{si,c}}{p_{00} q_{v,c}} T_{00} n_c \right)^{-1}$ and has a value of about
 ~ 340 seconds. Here, n_c is the reference ice number concentration, m_c is the reference mean mass of ice particles. Given the
 characteristic time scales and expanding the Exner pressure π in terms of a small parameter ε one can write the following
 expression for the derivative of π (~~Dolaptchiev et al. (2023)~~) (Dolaptchiev et al., 2023):

$$475 \quad \frac{D\pi}{Dt} = \varepsilon^{2+\alpha} \frac{D\pi'}{Dt} + \varepsilon w \frac{d\bar{\pi}}{dz_s}, \quad (\text{A1})$$

where $\pi = \left(\frac{p}{p_{00}} \right)^{\frac{R}{c_p}}$, $\alpha = 0$ corresponds to the strong and $\alpha = 1$ to the weak stratification case (~~Achatz et al. (2017, 2023)~~)
 (Achatz et al., 2017, 2023), $\bar{\pi}$ is the ~~Extner-Exner~~ Exner pressure of the hydrostatically balanced reference atmosphere, R is the gas
constant for dry air, and π' describes the fluctuations due to the wave field.

As shown in Dolaptchiev et al. (2023), applying the scaling analysis, one can neglect the first term on the ~~right-hand~~ right-hand side in Eq. (A1). In ~~case, the case~~ where the leading order vertical velocity is solely due to a single GW, w in the second term of Eq. (A1) can be written as

$$w = |\hat{w}|^{(0)} \cos(\omega t_w + \phi).$$

For a superposition of j gravity waves, the following expression applies:

$$w = \sum_j |\hat{w}_j| \cos(\underline{\Omega} \omega_j t_w + \phi_j).$$

Here \hat{w}_j , $\underline{\Omega} \omega_j$, ϕ_j are the vertical wind amplitude, frequency, and phase of GW number j .

480 Substituting the following expressions in the Eq.(6) and also taking into account constant background updraft w_{00} , the last term can be written in the following form:

$$F(t) = \frac{gL_i}{\underline{c_p R_v T_{00}^2}} \frac{gL_i}{c_p R_v T^2} \left(\sum_j \hat{w}_j \cos(\Omega_j t + \phi_j) + w_{00} \right), \quad (\text{A2})$$

where L_i is the latent heat of sublimation, g is the acceleration due to gravity, c_p is the heat capacity, R_v is the gas constant for dry air, $\underline{T_{00}} T$ is the background reference temperature.

485 Appendix B

The parameter selection and limitations of the proposed fit are further evaluated using additional results obtained with a more advanced fit formulation for m_{eff} . This version incorporates a third parameter, m^* , and introduces a more complex functional dependency, expanding the simple fit to a third-order polynomial and increasing the number of fitting coefficients to 17. This extension is designed to better capture the spread observed in the dataset shown before in Fig. 4.

490 The PDFs of deviations of the m_{eff} , mean radius of ice particle R_0 , and n_{post} , predicted with the use of the more complex fit, from the reference are shown in Fig. B1. As mentioned in section 2.3, the improvement in the target parameter n_{post} is minor. The PDFs of predicted m_{eff} and n_{post} (see Fig. B2) indicate a better representation of the maximum and right tail of the PDF of ice number concentration. Despite those improvements, the use of a more comprehensive fit is not considered further, due to efficiency reasons and divergent behavior of the polynomials in the extreme cases.

495 Appendix C

In order to further quantify the proposed parameterization and test it for the larger vertical velocity perturbations, a set of additional calculations was performed using the constant updraft and extreme amplitudes of GW-induced vertical velocity. The results are compared to the commonly used homogeneous nucleation parameterizations such as Kärcher and Lohmann (2002) and a reference system (1)-(3) with resolved ice physics. The ice number concentration prediction is presented depending on

500 constant vertical updraft and \hat{w} vertical updraft created by the GW (see Fig.C1). The proposed parameterization shows a good

Table A1. Values or ranges for the physical quantities within the relevant ranges for [the UTLS region](#) ([Baumgartner and Spichtinger \(2019\)](#); [Dolaptchiev et al. \(2023\)](#)), see also ([Baumgartner and Spichtinger, 2019](#); [Dolaptchiev et al., 2023](#))

Parameter	Description	Value or range
T_{00}	Reference background temperature	210 K
ρ_{00}	Reference background density	0.5 kg m^{-3}
ρ_i	Ice particles density	920 kg m^{-3} to 929 kg m^{-3}
p_{00}	Reference background pressure	300 hPa
J	Parameter for the nucleation rate	$4.9 \cdot 10^4 \text{ kg}^{-1} \text{ s}^{-1}$
B	Parameter for the nucleation rate	337
S_c	Critical saturation ratio over ice	1.43 to 1.64 <u>1.5</u>
D	Deposition <u>Growth</u> parameter	$4.3 \cdot 10^{-8} \text{ kg}^{2/3} \text{ s}^{-1} \text{ K}^{-1}$
L_i	Latent heat of sublimation	$2.8 \cdot 10^6 \text{ J kg}^{-1}$
c_p	Specific heat of dry air	$1005 \text{ J kg}^{-1} \text{ K}^{-1}$
R	Gas constant for dry air	$287 \text{ J kg}^{-1} \text{ K}^{-1}$
<u>R_v</u>	<u>Gas constant for water vapor</u>	<u>$416 \text{ J kg}^{-1} \text{ K}^{-1}$</u>
$p_{si,c}$	Reference saturation pressure over ice	1 Pa
n_c	characteristic <u>Characteristic</u> ice number concentration	$2 \cdot 10^6 \text{ kg}^{-1}$
m_c	characteristic <u>Characteristic</u> mean ice crystal mass	10^{-12} kg
<u>\hat{m}</u>	<u>Mean mass of newly nucleated ice particles</u>	<u>10^{-16} kg</u>

agreement with the traditional approaches and captures the larger n_i for the larger vertical updraft, which was also found by Kärcher and Lohmann (2002) using a bin model. Cases where vertical updraft was forced by the GWs also show the same behavior as the constant updraft, however, the validation is done against the resolved version. The extended parameterization overestimates n_i compared to the resolved physics, but the final predictions stay in the same order of magnitude and follow the expected physical trend. Notably, the larger deviations occur primarily in scenarios with extreme vertical velocity perturbations, up to 15 ms^{-1} , which represent rare and highly energetic events in the UTLS region. Despite this, the parameterization successfully captures the qualitative and quantitative response of the system under such conditions. This demonstrates both the robustness and the flexibility of the approach, making it a reliable and efficient choice for representing a wide range of atmospheric scenarios, including those with strong dynamic forcing.

Author contributions. AK and SD proposed the concept of generalization of the parameterisation. The extension was built based on the collaborative work of SD, PS, and UA, utilizing data produced from the ICON model with the gravity waves parameterisation developed within the group, under the leadership of UA. AK developed the methodology and conducted the investigation, defining the extended parameterisation with input from colleagues. The source code of the parcel model was developed by AK with the assistance of SD, scripts for

data analysis and outputs were written and documented by AK. Data interpretation was carried out collectively by all authors. AK wrote the
515 manuscript, incorporating contributions and revisions from all co-authors. SD, UA, and PS provided supervision and guidance throughout
the project.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Acknowledgements. All authors thank the German Research Foundation (DFG) for support through the CRC 301 "TPChange" (Project-ID
428312742, Projects B06 "Impact of small-scale dynamics on UTLS transport and mixing", B07 "Impact of cirrus clouds on tropopause
520 structure", and Z03 "Joint model development").

References

- Achatz, U., Ribstein, B., Senf, F., and Klein, R.: The interaction between synoptic-scale balanced flow and a finite-amplitude mesoscale wave field throughout all atmospheric layers: weak and moderately strong stratification, *Quarterly Journal of the Royal Meteorological Society*, 143, 342–361, <https://doi.org/10.1002/qj.2926>, 2017.
- 525 Achatz, U., Kim, Y.-H., and Voelker, G. S.: Multi-scale dynamics of the interaction between waves and mean flows: From non-linear WKB theory to gravity-wave parameterizations in weather and climate models, *Journal of Mathematical Physics*, 64, <https://doi.org/https://doi.org/10.1063/5.0165180>, 2023.
- Atlas, R. and Bretherton, C. S.: Aircraft observations of gravity wave activity and turbulence in the tropical tropopause layer: prevalence, influence on cirrus clouds, and comparison with global storm-resolving models, *Atmospheric Chemistry and Physics*, 23, 4009–4030, <https://doi.org/10.5194/acp-23-4009-2023>, 2023.
- 530 Barahona, D. and Nenes, A.: Parameterization of cirrus cloud formation in large-scale models: Homogeneous nucleation, *Journal of Geophysical Research-Atmospheres*, 113, <https://doi.org/10.1029/2007JD009355>, 2008.
- Baumgartner, M. and Spichtinger, P.: Homogeneous nucleation from an asymptotic point of view, *Theoretical and Computational Fluid Dynamics*, 33, 83–106, <https://doi.org/10.1007/s00162-019-00484-0>, 2019.
- 535 Baumgartner, M., Rolf, C., Groß, J.-U., Schneider, J., Schorr, T., Möhler, O., Spichtinger, P., and Krämer, M.: New investigations on homogeneous ice nucleation: the effects of water activity and water saturation formulations, *Atmospheric Chemistry and Physics*, 22, 65–91, <https://doi.org/10.5194/acp-22-65-2022>, 2022.
- Böloni, G., Kim, Y.-H., Borchert, S., and Achatz, U.: Toward transient subgrid-scale gravity wave representation in atmospheric models. Part I: Propagation model including nondissipative wave–mean-flow interactions, *Journal of the Atmospheric Sciences*, 78, 1317–1338, <https://doi.org/10.1175/JAS-D-20-0065.1>, 2021.
- 540 Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M., Kondo, Y., Liao, H., Lohmann, U., et al.: Clouds and aerosols, in: *Climate change 2013: The physical science basis. Contribution of working group I to the fifth assessment report of the intergovernmental panel on climate change*, pp. 571–657, Cambridge University Press, 2013.
- Dinh, T., Podglajen, A., Hertzog, A., Legras, B., and Plougonven, R.: Effect of gravity wave temperature fluctuations on homogeneous ice nucleation in the tropical tropopause layer, *Atmospheric Chemistry and Physics*, 16, 35–46, <https://doi.org/10.5194/acp-16-35-2016>, 2016.
- 545 Dolaptchiev, S. I., Spichtinger, P., Baumgartner, M., and Achatz, U.: Interactions between Gravity Waves and Cirrus Clouds: Asymptotic Modeling of Wave-Induced Ice Nucleation, *Journal of the Atmospheric Sciences*, 80, 2861 – 2879, <https://doi.org/https://doi.org/10.1175/JAS-D-22-0234.1>, 2023.
- Gasparini, B., Meyer, A., Neubauer, D., Münch, S., and Lohmann, U.: Cirrus cloud properties as seen by the CALIPSO satellite and ECHAM-HAM global climate model, *Journal of Climate*, 31, 1983–2003, <https://doi.org/10.1175/JCLI-D-16-0608.1>, 2018.
- 550 Gettelman, A., Liu, X., Barahona, D., Lohmann, U., and Chen, C.: Climate impacts of ice nucleation, *Journal of geophysical research: Atmospheres*, 117, <https://doi.org/10.1029/2012JD017950>, 2012.
- Gierens, K. M., Monier, M., and Gayet, J.-F.: The deposition coefficient and its role for cirrus clouds, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/10.1029/2001JD001558>, 2003.
- 555 Hertzog, A., Alexander, M. J., and Plougonven, R.: On the intermittency of gravity wave momentum flux in the stratosphere, *Journal of the Atmospheric Sciences*, 69, 3433–3448, <https://doi.org/10.1175/JAS-D-12-09.1>, 2012.

- Joos, H., Spichtinger, P., Lohmann, U., Gayet, J.-F., and Minikin, A.: Orographic cirrus in the global climate model ECHAM5, *Journal of Geophysical Research: Atmospheres*, 113, <https://doi.org/10.1029/2007JD009605>, 2008.
- Joos, H., Spichtinger, P., Reutter, P., and Fusina, F.: Influence of heterogeneous freezing on the microphysical and radiative properties of orographic cirrus clouds, *Atmospheric Chemistry and Physics*, 14, 6835–6852, <https://doi.org/10.5194/acp-14-6835-2014>, 2014.
- Kärcher, B. and Ström, J.: The roles of dynamical variability and aerosols in cirrus cloud formation, *Atmospheric Chemistry and Physics*, 3, 823–838, <https://doi.org/10.5194/acp-3-823-2003>, 2003.
- Kärcher, B., Hoffmann, F., Podglajen, A., Hertzog, A., Pluogonven, R., Atlas, R., Corcos, M., Grabowski, W. W., and Gasparini, B.: Effects of Turbulence on Upper-Tropospheric Ice Supersaturation, *Journal of the Atmospheric Sciences*, 81, 1589–1604, <https://doi.org/10.1175/JAS-D-23-0217.1>, 2024.
- Kim, Y.-H., Bölöni, G., Borchert, S., Chun, H.-Y., and Achatz, U.: Toward transient subgrid-scale gravity wave representation in atmospheric models. Part II: Wave intermittency simulated with convective sources, *Journal of the Atmospheric Sciences*, 78, 1339–1357, <https://doi.org/10.1175/JAS-D-20-0066.1>, 2021.
- Kim, Y.-H., Voelker, G. S., Bölöni, G., Zängl, G., and Achatz, U.: Crucial role of obliquely propagating gravity waves in the quasi-biennial oscillation dynamics, *EGUspere*, 2023, 1–18, <https://doi.org/10.5194/acp-24-3297-2024>, 2023.
- Köhler, L., Green, B., and Stephan, C. C.: Comparing loon superpressure balloon observations of gravity waves in the tropics with global storm-resolving models, *Journal of Geophysical Research: Atmospheres*, 128, e2023JD038 549, <https://doi.org/10.1029/2023JD038549>, 2023.
- Koop, T., Luo, B., Tsias, A., and Peter, T.: Water activity as the determinant for homogeneous ice nucleation in aqueous solutions, *Nature*, 406, 611–614, <https://doi.org/10.1038/35020537>, 2000.
- Krämer, M., Rolf, C., Luebke, A., Afchine, A., Spelten, N., Costa, A., Meyer, J., Zoeger, M., Smith, J., Herman, R. L., et al.: A microphysics guide to cirrus clouds–Part 1: Cirrus types, *Atmospheric Chemistry and Physics*, 16, 3463–3483, <https://doi.org/10.5194/acp-16-3463-2016>, 2016.
- Krämer, M., Rolf, C., Spelten, N., Afchine, A., Fahey, D., Jensen, E., Khaykin, S., Kuhn, T., Lawson, P., Lykov, A., et al.: A microphysics guide to cirrus–Part 2: Climatologies of clouds and humidity from observations, *Atmospheric Chemistry and Physics*, 20, 12 569–12 608, <https://doi.org/10.5194/acp-20-12569-2020>, 2020.
- Kärcher, B. and Lohmann, U.: A parameterization of cirrus cloud formation: Homogeneous freezing of supercooled aerosols, *Journal of Geophysical Research: Atmospheres*, 107, AAC 4–1–AAC 4–10, <https://doi.org/10.1029/2001JD000470>, 2002.
- Kärcher, B. and Podglajen, A.: A Stochastic Representation of Temperature Fluctuations Induced by Mesoscale Gravity Waves, *Journal of Geophysical Research: Atmospheres*, 124, 11 506–11 529, <https://doi.org/10.1029/2019JD030680>, 2019.
- Lott, F. and Miller, M. J.: A new subgrid-scale orographic drag parametrization: Its formulation and testing, *Quarterly Journal of the Royal Meteorological Society*, 123, 101–127, <https://doi.org/10.1002/qj.49712353704>, 1997.
- Luebke, A. E., Afchine, A., Costa, A., Groß, J.-U., Meyer, J., Rolf, C., Spelten, N., Avallone, L. M., Baumgardner, D., and Krämer, M.: The origin of midlatitude ice clouds and the resulting influence on their microphysical properties, *Atmospheric Chemistry and Physics*, 16, 5793–5809, <https://doi.org/10.5194/acp-16-5793-2016>, 2016.
- Matus, A. V. and L’Ecuyer, T. S.: The role of cloud phase in Earth’s radiation budget, *Journal of Geophysical Research: Atmospheres*, 122, 2559–2578, <https://doi.org/10.1002/2016JD025951>, 2017.

- Orr, A., Bechtold, P., Scinocca, J., Ern, M., and Janiskova, M.: Improved middle atmosphere climate and forecasts in the ECMWF model through a nonorographic gravity wave drag parameterization, *Journal of Climate*, 23, 5905–5926, <https://doi.org/10.1175/2010JCLI3490.1>, 2010.
- Podglajen, A., Hertzog, A., Plougonven, R., and Legras, B.: Lagrangian temperature and vertical velocity fluctuations due to gravity waves in the lower stratosphere, *Geophysical Research Letters*, 43, 3543–3553, <https://doi.org/10.1002/2016GL068148>, 2016.
- Ren, C. and Mackenzie, A. R.: Cirrus parametrization and the role of ice nuclei, *Quarterly Journal of the Royal Meteorological Society*, 131, 1585–1605, <https://doi.org/10.1256/qj.04.126>, 2005.
- Spichtinger, P. and Gierens, K. M.: Modelling of cirrus clouds—Part 1a: Model description and validation, *Atmospheric Chemistry and Physics*, 9, 685–706, <https://doi.org/10.5194/acp-9-685-2009>, 2009.
- Spichtinger, P., Marschalik, P., and Baumgartner, M.: Impact of formulations of the homogeneous nucleation rate on ice nucleation events in cirrus, *Atmospheric Chemistry and Physics*, 23, 2035–2060, <https://doi.org/10.5194/acp-23-2035-2023>, 2023a.
- Spichtinger, P., Marschalik, P., and Baumgartner, M.: Impact of formulations of the homogeneous nucleation rate on ice nucleation events in cirrus, *Atmospheric Chemistry and Physics*, 23, 2035–2060, <https://doi.org/10.5194/acp-23-2035-2023>, 2023b.
- Spreitzer, E. J., Marschalik, M. P., and Spichtinger, P.: Subvisible cirrus clouds—a dynamical system approach, *Nonlinear Processes in Geophysics*, 24, 307–328, <https://doi.org/10.5194/npg-24-307-2017>, 2017.
- VanZandt, T.: A universal spectrum of buoyancy waves in the atmosphere, *Geophysical Research Letters*, 9, 575–578, <https://doi.org/10.1029/GL009i005p00575>, 1982.
- Voelker, G. S., Bölöni, G., Kim, Y.-H., Zängl, G., and Achatz, U.: MS-GWaM: A Three-Dimensional Transient Gravity Wave Parametrization for Atmospheric Models, *Journal of the Atmospheric Sciences*, 81, 1181–1200, <https://doi.org/10.1175/JAS-D-23-0153.1>, 2024.
- Wang, Y., Su, H., Jiang, J. H., Xu, F., and Yung, Y. L.: Impact of cloud ice particle size uncertainty in a climate model and implications for future satellite missions, *Journal of Geophysical Research: Atmospheres*, 125, e2019JD032119, <https://doi.org/10.1029/2019JD032119>, 2020.
- Zängl, G., Reinert, D., Rípodas, P., and Baldauf, M.: The ICON (ICOsahedral Non-hydrostatic) modelling framework of DWD and MPI-M: Description of the non-hydrostatic dynamical core, *Quarterly Journal of the Royal Meteorological Society*, 141, 563–579, <https://doi.org/10.1002/qj.2378>, 2015.
- Zhang, Y., Macke, A., and Albers, F.: Effect of crystal size spectrum and crystal shape on stratiform cirrus radiative forcing, *Atmospheric Research*, 52, 59–75, [https://doi.org/10.1016/S0169-8095\(99\)00026-5](https://doi.org/10.1016/S0169-8095(99)00026-5), 1999.
- Zhou, C., Penner, J. E., Lin, G., Liu, X., and Wang, M.: What controls the low ice number concentration in the upper troposphere?, *Atmospheric Chemistry and Physics*, 16, 12411–12424, <https://doi.org/10.5194/acp-16-12411-2016>, 2016.




Figure 10. Probability density function of the deviation of $m_0 - m_{eff}$ predicted by fit (10) from exact mass calculated from the full system using (9) in %. Horizontal bars correspond to the 90, 50, and 10 percentiles of the distribution.




Figure 11. Probability density function of the deviation of $N_{post} - n_{post}$ predicted using fit (10) by (8) from exact mass calculated from the full system in %. Horizontal bars correspond to the 90, 50, and 10 percentiles of the distribution.




Figure 12. Probability density functions of $m_0, N_{post}, m_{eff}, n_{post}$ and q_i from the full model and predicted by the parameterisation (8) with different used time steps.




Figure 13. Comparison of the parameters evolution in full system, reduced system, and parameterisation with proposed correction. Initial conditions: $S = 1.4$, $n = 50$ $n_i = 50 \text{ kg}^{-1}$, $q = 2.6 \cdot 10^{-13}$ $q_i = 2.6 \cdot 10^{-13} \text{ kg kg}^{-1}$.




Figure 14. Comparison of the parameters evolution in full system, reduced system, and parameterisation with proposed correction. Initial conditions: $S = 1.4$, $n = 0.018$ $n_i = 0.018$ kg^{-1} , $q = 6 \cdot 10^{-16}$ $q_i = 6 \cdot 10^{-16}$ kg kg^{-1} .




Figure B1. Probability density functions for deviations of the m_{eff} , R_{eff} , and n_{post} predictions made with the advanced fit from the exact solutions. Vertical lines correspond to the 90th, 50th, and 10th percentiles of the distribution.




Figure B2. Probability density functions for m_{eff} , n_{post} , and q_i exact ensemble solutions and predictions made with the advanced fit. Vertical lines correspond to the mean of the distribution.




Figure C1. The ice number concentration calculated based on the proposed approach with the conditions of $T = 220\text{K}$, time step of 10s and with the prescribed constant updraft (param.) and single GW forcing (param. GW forcing) is shown depending on the w or \hat{w} . For comparisons, the results based on the reference system (1)-(3) without sedimentation are shown for the GW forcing (ref. GW forcing). As a reference for the constant updraft, the values from detailed microphysics calculations from Kärcher and Lohmann (2002) are shown, where the upper/middle/lower symbols correspond to an initialization of narrower/baseline/broader aerosol size distributions.