

SICNet_{season} V1.0: a transformer-based deep learning model for seasonal Arctic sea ice prediction by incorporating sea ice thickness data

Yibin Ren¹, Xiaofeng Li¹, Yunhe Wang¹

¹Key Laboratory of Ocean Observation and Forecasting, Key Laboratory of Ocean Circulation and Waves, Institute of Oceanology, Chinese Academy of Sciences, Qingdao, China

Correspondence to: Xiaofeng Li (lixf@qdio.ac.cn)

Abstract. The Arctic sea ice ~~is suffering~~suffers dramatic retreating in summer and fall, which has far-reaching consequences on the global climate and commercial activities. Accurate seasonal sea ice predictions ~~are significant in inferring~~significantly ~~infer~~ climate change and planning commercial activities. However, seasonally predicting the summer sea ice encounters a significant obstacle known as the spring predictability barrier (SPB): ~~predictions made later than the date of melt onset (roughly May)~~predictions made later than May demonstrate good skill in predicting summer sea ice, while predictions made on or earlier than May exhibit considerably lower skill. This study develops a transformer-based deep-learning model, SICNet_{season} (V1.0), to predict the Arctic sea ice concentration on a seasonal scale. Including spring sea ice thickness (SIT) data in the model significantly improves the prediction skill at the SPB point. A 20-year (2000-2019) testing demonstrates that the detrended anomaly correlation coefficient (ACC) of Sep. sea ice extent (sea ice concentration > 15%) predicted by our model at May/Apr. is improved by 7.7%/10.61% over the ACC predicted by the state-of-the-art dynamic model [SEAS5](#) from the European Centre for Medium-Range Weather Forecasts (ECMWF). Compared with the anomaly persistence benchmark, the mentioned improvement is 41.02%/36.33%. Our deep learning model significantly reduces prediction errors of Sep.'s sea ice concentration on seasonal scales compared to [SEAS5](#) ~~ECMWF~~ and Persistence. The spring SIT data is key in ~~optimizing the SPB~~optimizing the predictions around the SPB, contributing to a more than 20% ACC enhancement in Sep.'s SIE at four to five months lead predictions. Our model achieves good generalization in predicting the Sep. SIE of 2020-2023.

1 Introduction

Arctic sea ice plays a significant role in the global climate because it modulates the thermal and dynamic exchanges between the ocean and the atmosphere (Ding et al., 2017; Kapsch et al., 2013; Liu et al., 2021a; Olonscheck et al., 2019). In recent decades, global warming has resulted in a dramatic retreat in Arctic sea ice during the summer and fall (Cao et al., 2017; Shu et al., 2022). This decline triggers a system-positive feedback mechanism that causes the Arctic's surface air temperature to increase 2-4 times faster than the global mean state, known as the Arctic amplification (AA) (England et al., 2021; Pithan

30 and Mauritsen, 2014; Screen et al., 2013; Screen and Simmonds, 2010). AA accelerates sea ice decline, strengthening
positive feedback (Jenkins and Dai, 2021; Kumar et al., 2010). If the situation is unchanged, climate models project that the
Arctic will become ice-free during summer by the 2050s (Jahn et al., 2024; Kim et al., 2023; Andersson et al., 2024). The
dramatic Arctic sea ice loss has led to consequences for global climate (Francis and Vavrus, 2012) and commercial activities
(Min et al., 2022). For example, it ~~it may weaken the stratospheric polar vortex~~ weakens the stratospheric polar vortex in the
35 winter, increasing extreme cold events in the Northern Hemisphere (Blackport et al., 2019; Cohen et al., 2014). Furthermore,
the lower sea ice area during summer extends the navigability of the Arctic Passage to seasonal scales (Cao et al., 2022).

Sea ice predictions are helpful for better understanding global climate change and support human activities in the Arctic
(Lindsay et al., 2008; Merryfield et al., 2013). Therefore, sea ice prediction, commonly represented by parameters such as
sea ice concentration (SIC) or sea ice extent (SIE, defined as the sum of grid cell area where SIC>15%), has always attracted
40 substantial efforts (Guemas et al., 2016; Stroeve and Notz, 2015). Various prediction systems are proposed, such as
numerical (Chevallier et al., 2013; Liang et al., 2020; Mu et al., 2020; Wang et al., 2013; Yang et al., 2019; Zhang et al.,
2008, 2022), statistical (Gregory et al., 2020; Wang et al., 2016, 2022; Yuan et al., 2016), and deep learning models (Jun
Kim et al., 2020; Ren et al., 2022; Ren and Li, 2023). However, accurate sea ice prediction for Arctic summer remains
challenging, particularly at seasonal or even longer scales (Zampieri et al., 2018; Blanchard-Wrigglesworth et al., 2015,
45 2023). One of the biggest challenges is the spring predictability barrier (SPB): predictions for summer sea ice made ~~before or~~
~~at the timing of melt onset before or on May~~ show significantly lower skill than predictions made after ~~the timing of melt~~
~~onset May~~ (Bonan et al., 2019; Bushuk et al., 2020; Day et al., 2014; Zeng et al., 2023). ~~The Sea Ice Prediction Network~~
~~(SIPN) for the Sep.' SIE in the Arctic showed that individual dynamical and statistical predictions could not beat the anomaly~~
~~persistence benchmark when predictions are made from early June to early August, with a root mean squared error (RMSE)~~
50 ~~of 0.5–0.7 million km² (Blanchard-Wrigglesworth et al., 2015, 2023).~~ Studies show that SPB is evident in nearly all the
fully coupled global climate models (GCMs) in Phase 5 of the Coupled Model Intercomparison Project (CMIP5), a crucial
initiative providing climate projections to support essential climate research worldwide (Blanchard-Wrigglesworth et al.,
2011; Tietsche et al., 2014). So, ~~optimizing the SPB~~ optimizing the predictions around the SPB is an urgent task for accurate
summer sea ice predictions.

55 Experiments based on ensemble simulations reveal that the predictability of summer SIE is limited before spring due to the
ice motion and growth in winter (Bushuk et al., 2020). However, the predictability increases rapidly after the melting processes
in the spring (Bushuk et al., 2020). The satellite observations show that the spring sea ice thickness (SIT) correlates more with
the summer SIE than the spring SIE (Landy et al., 2022). These findings indicate that the spring SIT may be a key factor in
~~optimizing the SPB~~ optimizing the predictions around the SPB (Bushuk et al., 2020, n.d.). Recently, researchers have
60 assimilated the CryoSat-2 observed SIT data, the first summer SIT observations, into the Geophysical Fluid Dynamics
Laboratory (GFDL) ocean–sea ice model and found that the prediction skill of Sep.'s SIC is improved significantly when the
model is initialized with SIT anomaly in Jul. and Aug. (Zhang et al., 2023). This study further proves that the summer SIT data
contributes to Sep.'s sea ice prediction. However, as the SPB flag is May for most studies, whether the ~~predictions around the~~

带格式的: 字体: (默认)+西文标题 (Times New Roman)

带格式的: 字体: (默认) Times New Roman, (中文) Times New Roman

SPB could be optimized ~~or even overcome~~ by including SIT data remains largely unknown.

65 Currently, numerical models ~~account for the mainstream~~ are widely used in operationally sea ice prediction, but they are inflexible and have been limited by the SPB (Msadek et al., 2014; Sigmond et al., 2013). Statistical models are good at long-term prediction but cannot model complex nonlinear relationships and face SPB challenges. Deep learning models are more flexible than numerical models and more ~~powerful-potent~~ than traditional statistical ones, and they have been successfully used in Earth prediction problems (Li et al., 2021; Reichstein et al., 2019). Researchers have successfully developed deep-learning
70 models to predict polar sea ice state from synoptic to sub-seasonal scales (Andersson et al., 2021; Dong et al., 2024; Li et al., 2024; Mu et al., 2023; Palerme et al., 2024; Ren et al., 2022; Ren and Li, 2023; Song et al., 2024; Wu et al., 2022; Yang et al., ~~in 2024~~; Zhu et al., 2023), bringing the new potential to solve the SPB problem to improve the seasonal prediction skill from a data-driven perspective.

This work develops a seasonal sea ice prediction model named SICNet_{season} (V1.0) to ~~optimize the SPB~~ optimize the predictions around the SPB. SICNet_{season} is a transformer-based deep learning model with a physically constrained loss function based on SIC morphology. It takes historical SIC and SIT data as predictors and predicts the SIC of the following six months. The SIC data is the satellite-observed data from the National Snow and Ice Data Center (NSIDC) (Cavalieri et al., 1996). The SICNet_{season} model is trained on data from 1979-2019 and tested with data from 2000-2019 by a leave-one-year-out strategy. Data from the recent four years, 2020-2023, is employed to verify the model's generalization. Experiments demonstrate that our
80 model significantly optimizes the SPB with a higher detrend correlation coefficient (ACC) compared with anomaly persistence (Persistence) and the state-of-the-art dynamic ~~predictions-model~~ SEAS5 from the European Centre for Medium-Range Weather Forecasts (ECMWF) (Johnson et al., 2019). Our model significantly reduces the errors ~~of Sep in September~~. SIC/SIE in four to five months lead predictions ~~than compared to~~ the two compared models. The spring SIT data is key in optimizing the SPB optimizing the predictions around the SPB. Our model generalized well in predicting the Sep. SIE of 2020-2023. Finally,
85 we compare our SICNet_{season} model with the an IceNet-inspired ~~published deep learning~~ U-Net model-IceNet. Finally, a comparison between our and the published deep learning model IceNet (Andersson et al., 2021) is discussed. IceNet is a probability prediction model for Arctic SIE based on convolutional neural network (CNN) units and the U-Net architecture. It achieved state-of-the-art performance in predicting the probability of SIE for six months (Andersson et al., 2021). Therefore,
~~Therefore, we construct an IceNet-inspired U-Net model-chose IceNet as a comparison model. The IceNet is a probability prediction model for Arctic SIE based on convolutional neural networks (CNN) units and the U-Net architecture. It achieved the state-of-the-art performance in predicting the SIE of six months (Andersson et al., 2021). Therefore, we choose the IceNet as a comparison model.~~
90 prediction model for Arctic SIE based on convolutional neural networks (CNN) units and the U-Net architecture. It achieved the state-of-the-art performance in predicting the SIE of six months (Andersson et al., 2021). Therefore, we choose the IceNet as a comparison model.

带格式的: 下标

2 Data

1.1 Sea ice concentration data

95 The SIC data of 1979-2023 are experiment data. The SIC is the satellite-observed data obtained from the NSIDC. It is a daily observation derived from the Nimbus-7 Scanning Multichannel Microwave Radiometer (SMMR) and the Defense Meteorological Satellite Program (DMSP) Special Sensor Microwave Imager (SSM/I and SSMIS) (DiGirolamo-Cavaleri et al., 1996, 2022). The projection of the SIC data is the north-polar stereographic with a 25 km spatial resolution.

带格式的: 非突出显示

1.2 Sea ice thickness data

100 The SIT data is the reanalysis SIT from the Pan-Arctic Ice Ocean Modeling and Assimilation System (PIOMAS). PIOMAS is a numerical model with sea ice and ocean components, and it assimilates SIC and sea surface temperature (Zhang and Rothrock, 2003). PIOMAS SIT agrees well with in situ, airborne, and satellite measurements (Schweiger et al., 2011). It is daily data with an 18 km spatial resolution. Although the PIOMAS generally overestimates thin ice and underestimates thick ice regions, it is widely adopted by Arctic studies (Collow et al., 2015; Kwok et al., 2020; Nakanowatari et al., 2022).


105 The SIC and PIOMAS SIT data are converted to a Northern Polar Stereographic Grid with 80 km resolution. The temporal resolution is one month.

3 Method

3.1 Framework of SICNet_{season}

The SICNet_{season} model is derived from a transformed-based U-Net deep learning model, SwinUNet (Cao Hu and Wang, 2023)(Cao Hu and Wang, 2023). It accepts a three-dimensional sea ice data sequence and predicts a three-dimensional SIC sequence of the future, Fig. 1. The inputs to the model are monthly mean fields. The predicted target is the SIC of the next six months. For example, if we make predictions in May, the six months' predictions will cover the full summer and fall, the months from Jun. to Nov.

带格式的: 字体: (默认)+西文标题 (Times New Roman)



115 Figure 1. Framework of model SICNet_{season}. (a) Input consists of SIT of the last three months, SIC of the last six months, SIC
 120 anomaly of the last three months, and SIC climatology of six target months, 96×96×18. (b) The encoder comprises four swin-
 transformer blocks and three patch-merging operators. (c) The decoder contains three swin-transformer blocks and three patch-
 expanding operators. (d) Concatenations connect the feature maps from the encoder and the decoder module. (e) A CNN layer
 with sigmoid activation transforms the feature map to the predicted SIC of six-month leads. (f) Model training procedure. The loss
 function is combined with the normalized integrated ice-edge error (NIIEE) and the mean square error (MSE).

The input for SICNet_{season} is a 96×96×18 SIC and SIT sequence, composed of SIT of the ~~last last~~ three months, SIC of the
~~last last~~ six months, SIC anomaly of the ~~last last~~ three months, and SIC climatology of the six target months (Fig. 1a). ~~The~~
~~lengths of the input factors are determined by lots of experiments. We determine the length of input factors by combining~~
 125 ~~domain knowledge and manual tuning experiments. The primary domain knowledge we considered is the spring-fall~~
~~reemergence mechanism. It occurs between pairs of months where the ice edge is in the same position, such as in May and~~
~~December (Blanchard-Wrigglesworth et al., 2011; Day et al., 2014). The spring sea ice anomaly is positively correlated with~~
~~fall sea ice anomalies, and there is also a weaker reemergence between fall sea ice anomalies and anomalies the following~~
 130 ~~spring (Bushuk et al., 2015). Therefore, we set the initial input length of the SIC/SIT/SIC anomaly as six months. We change~~
~~the input length manually (from six to one in step one) to fine-tune the deep learning model to find the best-matched length~~
~~for each factor. The SIC climatology of the target months provides an essential mean state of the prediction SIC. It represents~~
~~the monthly cycle signal that IceNet has considered (Andersson et al., 2021).~~

The input is fed into the encoder to capture spatiotemporal correlations among SIC/SIT data sequences at different levels
 to form multi-scale correlation maps. The encoder comprises four swin-transformer (Liu et al., 2021b) blocks and three patch
 135 merging operators (Fig. 1b). A swin-transformer block is a transformer unit integrated with shifted windows (Liu et al.,
 2021b). A transformer operator captures global dependencies through an attention mechanism. The shifted windows help the
 transformer operator capture local dependencies like the convolution operator. Therefore, local and global spatiotemporal

带格式的: 字体: (默认) Times New Roman, (中文) Times New Roman

dependencies among sea ice sequences can be captured. The patch merging operator downscales the captured feature maps like the pooling layer in CNN models. The decoder upscales the feature maps through the patch expanding operator and swin-transformer blocks (Fig. 1c). The extracted correlation maps of the encoder and decoder are stacked to form fused spatiotemporal maps (Fig. 1d). A CNN layer transforms the decoded feature maps to the same shape as the target SIC sequence. Here, it is a $96 \times 96 \times 6$ array (Fig. 1e). As the range of SIC is 0-1, we employ the sigmoid function to activate the last feature map to transform the predicted values to 0-1.

During the training procedure, the loss is calculated between the predicted and ground values. Then, the model's parameters are trained by minimizing the loss value iteratively. We will explain the loss function in the following section.

3.2 Integrated ice-edge-constrained loss function

For a deep-learning model, the loss function is crucial during the training procedure as it guides the optimization of the model's parameters. Here, the loss is the difference between the predicted values and the ground ones from NSIDC. Generally, the mean square error (MSE) is a basic fundamental loss function for prediction tasks. The MSE measures the mean state for all predicted values and cannot reflect the spatial differences between 2-dimensional SIC patterns. To address the issue, we proposed a normalized integrated ice-edge error (NIIEE) loss function that considers the spatial distribution of SIC to constrain the model's optimization (Fig. 1g).

The NIIEE loss is based on the integrated ice-edge error (IIEE), a common professional metric for sea ice predictions. The IIEE represents the error regions ~~that~~ the prediction model overestimated and underestimated (Goessling et al., 2016). It measures the spatial similarity between two 2-dimension SIC patterns. Initially, the IIEE binarizes the SIC by 15% to describe the SIE. For the SIC prediction here, we do not perform binarization. Let P_{SIC}/G_{SIC} represent the predicted/ground SIC; the IIEE is calculated by Eq. (1). We normalize IIEE to the range of 0-1 to form the NIIEE loss by Eq. (2). If the NIIEE loss is 0, the predicted SIC and the ground SIC will match in spatial and numerical. The fundamental MSE loss has been demonstrated to be effective adequate for prediction tasks. If the number of all predicted values is N , the MSE is calculated by Eq. (3). We combine the NIIEE with MSE as the loss function of the SICNet_{season}. A constant scale factor, 0.01, is multiplied by NIIEE to balance its range with that of MSE, Eq. (4).

$$IIEE = (P_{SIC} \cup G_{SIC}) - (P_{SIC} \cap G_{SIC}) \quad (1)$$

$$NIIEE = \frac{IIEE}{P_{SIC} \cup G_{SIC}} = 1 - \frac{P_{SIC} \cap G_{SIC}}{P_{SIC} \cup G_{SIC}} \quad (2)$$

$$MSE = \frac{\sum (P_{SIC} - G_{SIC})^2}{N} \quad (3)$$

$$Loss = 0.01 \times NIIEE + MSE \quad (4)$$

4 Experiments

4.1 Model training

The model is trained on a computer station with an NVIDIA Tesla V100 32-GB card. The training and test samples are constructed by step-by-step sliding. The testing period is 2000-2019. The leave-one-year-out strategy is adopted to train/evaluate our SICNet_{season} model. For example, if the testing year is 2000, the training set is from 1979-1999 and 2001-2019. [The leave-one-year-out strategy is widely adopted by statistical models to maximize the sample volume while obtaining a multi-year evaluation \(Wang et al., 2022; Yuan et al., 2016\).](#) The validation set is split 20% from the training set. We set the batch size as eight and the initial learning rate as 0.0001. We employ the early stopping strategy to break the training procedure when the validation loss does not decrease. The model is trained three times to eliminate random errors. The testing set is run on three trained models, and the mean values are adopted as the final predictions. Data from the recent four years, 2020-2023, is employed to verify the model's generalization. Data from these four years did not participate in the training stage. They are fed into the trained models obtained by the leave-one-year-out strategy to get the predictions. The predictions are the mean values of the 20 trained models.

4.2 Evaluation metrics

The mean absolute error (MAE), Binary Accuracy (BACC), and detrend ACC are evaluation metrics. The MAE is for SIC, and the other two metrics are for SIE. To accurately calculate the metrics, we use the maximum observed monthly SIE since 1979 to mask the predictions. Assuming the predicted/truth value of the i_{th} grid is p_i/g_i , the number of validation grids is N . The MAE values are calculated by (5). The BACC of time t is obtained by using one to subtract the ratio of IIEE to the area of the activated grid cell region (the maximum observed SIE during 1979-2019) of t by equation (6). The detrend ACC of SIE is the anomaly correlation coefficient of two detrend SIE series. Each SIE series has 20 elements, from 2000 to 2019.

$$MAE = \frac{\sum_i^N |p_i - g_i|}{N} \quad (5)$$

$$BACC = (1 - \frac{IIEE}{\text{area of the activated grid cell region}}) \times 100\% \quad (6)$$

4.3 Model skill in seasonal predictions


We compare the SICNet_{season} with the Persistence and the [ECMWF dynamical model-SEAS5](#) to validate our model's ability to [optimize the SPB](#)[optimize the predictions around the SPB](#). [Blanchard Wrigglesworth](#)[The Persistence model estimates the target SIC values by adding the current anomaly to the climate mean state at the target time, widely adopted as a benchmark for sea ice prediction \(Wang et al., 2016\)\(Wang et al., 2016Blanchard Wrigglesworth et al., 2023\).](#) [The Persistence is the anomaly persistence model. It assumes the anomaly constant in time and estimates the target SIC values by adding the current anomaly to the climate mean state at the target time, widely adopted as a benchmark for sea ice prediction \(Wang et al., 2016\).](#) The SEAS5 is a new seasonal forecast system from the ECMWF that shows excellent sea ice prediction skills

(Johnson et al., 2019). A BACC value of 100% indicates that the predicted SIE matches the observed SIE 100% ~~in~~ ~~spatially~~ spatially. The metrics are calculated for 20 testing years, 2000-2019, in a leave-one-year-out training/testing strategy. As the SPB occurred in the target summer month, we focus on the four summer months, June to September.


200 Fig. 2 shows the detrended ACC and BACC of target months, Jun.-Sep., on six lead months' predictions. As shown in Figs. 2(a) and (b), the predictions of Persistence and [SEAS5](#) ~~ECMWF~~ show an apparent SPB: the detrended ACC drops sharply when the predictions are made earlier than May, with a maximum ACC gap between two adjacent lead months marked by black lines. Taking September for example, the detrend ACC is 56.39% for Persistence when the prediction is made in Jun. (three months lead). Then, it decreases to 26.59% in May's prediction (four months lead), Fig. 2(a). For [ECMWFSEAS5](#), the 205 ACC of Jun.'s prediction is 83.94%, then drops to 59.91% at May's prediction, forming a 24.03% ACC gap, Fig. 2(b). Although the SICNet_{season}'s prediction also shows an SPB feature, the black line in Fig.2 (c), the ACC at May's prediction is improved to 67.61%, and the ACC gap is reduced to 15.6%. Further, the ACC difference is calculated between SICNet_{season} and Persistence/[ECMWFSEAS5](#). Compared with Persistence, SICNet_{season} improves the ACC in most predictions, Fig. 2d. The ACC improvements along the SPB flag are more than 30% on average (the lead months right to the black line in Fig. 210 2d). Compared with the [ECMWFSEAS5](#), SICNet_{season} also improves the prediction skill of the SPB. When the target months are Jun. and Jul., SICNet_{season} shows a much higher prediction skill than [ECMWFSEAS5](#) in four to six months lead predictions, Fig. 2e. For the target month Sep., the SICNet_{season} improves the ACC by 7.70%/10.61% than [ECMWFSEAS5](#) when prediction is made in May/Apr. (four/five months lead in Fig. 2e). For the target month Aug./Sep., the SICNet_{season} shows lower ACCs than the [ECMWFSEAS5](#) when prediction is made on or before Mar. (five/six months lead for Aug./Sep.). 215 However, for the predictions made adjacent to the SPB flag line, the SICNet_{season} achieves larger ACCs than the [ECMWFSEAS5](#) (values right to the black line in Fig. 2e). Therefore, the SICNet_{season} optimizes the SPBs significantly compared to the well-known numerical model.

The BACC of [ECMWFSEAS5](#) also shows a similar SPB characteristic to the ACC. A sharp BACC drop occurred when the prediction was made on and before May, the black line in Fig. 2(g). The maximum BACC gaps of Persistence and 220 SICNet_{season} occurred in the second lead month. However, the maximum BACC gap of SICNet_{season} is about 2%, much lower than the 10% gap of Persistence and [ECMWFSEAS5](#). Compared with the Persistence and [ECMWFSEAS5](#), the SICNet_{season} improves the BACC by more than 10% in predicting SIE of Aug. and Sep. three to six months lead, Figs. 2(i) and (j).


Detrended ACC




Detrended ACC Difference




BACC





BACC Difference





带格式的: 居中



Detrended ACC



BACC







Figure 2. Detrend ACC of SIE, BACC of SIE, and their differences of Persistence, ECMWFSEASS, and SICNet_{season} from Jun. to Sep., averaged by 2000-2019. (a)-(c) Detrend ACC of three models. Two detrend SIE series (predicted and observed) calculate each value. (d)-(e) Detrend ACC differences between SICNet_{season} and Persistence/ECMWFSEASS. (f)-(h) BACC of three models. Each BACC is a mean value during 20 testing years. (i)-(j) BACC differences of SICNet_{season} and Persistence/ECMWFSEASS. The

black line indicates the SPB: a maximum decrease between two adjacent lead months. The red signifies a high/improvement in ACC/BACC, and the blue signifies a decrease.

4.4 Performance in predicting SIC of Sep.

As September's sea ice draws wider attention than other months, we calculate the MAE of SIC of Sep. predicted by three models. Fig. 3 shows the spatial MAE of Persistence, [ECMWFSEAS5](#), and SICNet_{season} on six lead months. The MAE in the three models is not much different for the first two lead months. When the lead month is one, the MAE of [ECMWFSEAS5](#) is slightly better than that of Persistence and SICNet_{season}, indicating that the [dynamic SEAS5](#) model performs well in monthly predicting. This result may be due to the good atmospheric initialization in SEAS5, which beat many machine learning and dynamical models in sub-seasonal scale SIC prediction (Bushuk et al., 2024). However, when the lead month is longer than three, the [ECMWFSEAS5](#)'s MAEs are much more than 45% in the Pacific sector, mainly containing the Beaufort Sea, the Chukchi Sea, the East Siberian Sea, and the Laptev Sea, Figs. 3(j)-(l). The SICNet_{season} reduces the MAEs to 20-30% for most regions in the Pacific Arctic, Figs. 3(q)-(r). Compared with Persistence, SICNet_{season} also reduces MAEs by 5-10% in the mentioned four local seas, Figs. 3(d)-(f). Therefore, the SICNet_{season} significantly reduces the SIC errors of Sep. in seasonal scale predictions (three to six months lead).



245




Figure 3. The MAEs of September's predictions are based on three compared models: each value is averaged by 20 testing years. (a)-(g) MAEs of Persistence. (h)-(m) MAEs of [ECMWF-SEAS5](#). (n)-(s) MAEs of SICNet_{season}.

4.5 SIT's contributions to seasonal predictions

250 We further conduct a comparison experiment to validate the role of SIT data in seasonal predictions based on SICNet_{season}. The model without SIT as an input is named SICNet_{season_nosit}. The other settings for SICNet_{season_nosit} are the same as those for SICNet_{season}. The detrended ACC and BACC are shown in Fig. 4.

255 Without the SIT data as input, the model's prediction skill drops apparently in three to six months lead predictions, Fig. 4(a). For the target month Sep., the detrend ACC is 76.41% when the prediction is made in Jun. (three months' lead). Then, the ACC drops to 26.43% at May's prediction (four months' lead). By including SIT data as input, the ACC of May's prediction is improved by 41.18% in model SICNet_{season}, Fig. 4(c). For the target month of August, the ACC improvement at May's prediction (three months' lead) by including SIT data is 42.44%. Therefore, the SIT data is important to improve the model's prediction skill on SPB.

For target months Aug. and Sep., the BACCs of SICNet_{season_nosit} show an apparent drop in three to six months lead predictions. By including SIT data as the model's input, the BACC improvement is 0.95%/2.02% for the target month, Aug./Sep. at May's predictions, Fig. 4(f). Then, we calculate the MAE of the target month Sep., Fig. 5. The MAEs of the first two lead months are similar for the two models. When the lead month is larger than three, the MAEs of SICNet_{season_nosit} in the Beaufort Sea, the East Siberian Sea, and the Laptev Sea are 30-45%, with red circles ~~eyes~~ in Figs. 5(d)-(f). By including SIT data, the MAEs in the ~~mentioned three regions are reduced to 20-35% by SICNet_{season}, three mentioned regions~~ are reduced to 20-35% by SICNet_{season}, as shown in the red circles ~~eyes~~ in Figs. 5(j)-(l). Therefore, including SIT data reduces the errors of Sep.'s SIC by more than 10% in seasonal scale predictions.

域代码已更改




Figure 4. Detrended ACC of SICNet_{season_nosit} (a) and SICNet_{season} (b). (c) ACC difference obtained by SICNet_{season} minus SICNet_{season_nosit}. BACC of SICNet_{season_nosit} (d) and SICNet_{season} (e). (f) BACC difference like (c). The red signifies a high/improvement in ACC/BACC, and the blue signifies a decrease.




Figure 5. The MAEs of Sep.'s SIC predicted by $SICNet_{season}$ and $SICNet_{season_nosit}$. Each value is averaged by 20 testing years. (a)-(g) MAE of $SICNet_{season_nosit}$. (h)-(m) MAE of $SICNet_{season}$. The red cycles marked the regions where the MAE is reduced typically by including SIT data.

4.6 Generalization in predicting the SIEs of 2020-2023


280 To verify our model's generalization in predicting the SIEs of recent years, we employed the twenty trained models to predict the SIE of the recent four years, 2020-2023. The twenty models are trained for 2000-2019, as mentioned in the earlier sections. The data from 2020-2023 is "blind" for the models. The mean values of the twenty models' predictions are the final predictions. As the temporal span of four years is too short for calculating ACC, we use the BACC as the metric (Fig. 6). Compared with Persistence and [ECMWFSEAS5](#), the SICNet_{season} achieves higher BACCs in predicting SIEs of Aug. and
285 Sep. For the target month Sep., the BACC of SICNet_{season} is 10% higher than those of the other two models in three to six months lead predictions.

We draw the observed and predicted [Sep.'s SIEs of 2020, 2022, and 2023](#) in Fig. 7. [The Sep.'s SIE in 2020/2023 \(4.0/4.37 milkm²\) is the second/sixth lowest record in the Arctic since 1979. The SIE in 2022 Sep. \(4.90 milkm²\) has been large since 2015. We focus on the seasonal scale predictions with four to six lead months. The ECMWF model achieves good accuracy when the lead month is one, with the largest BACC values in 2020-2022, Fig. 7\(a\)-\(g\)\(m\). This result proves that the dynamic model is good at short-term predictions. However, for the lead month longer than two, our SICNet_{season} data-driven model shows obvious advantages over the \[ECMWFSEAS5\]\(#\) and Persistence. For predictions made on or before May, lead months of four to six, the BACCs of SICNet_{season} are much higher than those of Persistence and \[ECMWFSEAS5\]\(#\), Fig. 7. ~~The SIE in 2020/2023 is the second/sixth lowest record in the Arctic since 1979.~~ At May's prediction, our model achieved a BACC of 82.25%/90.39%/82.08% in 2020/2022/2023, about 12%/10% more than 10% higher than over the Persistence and \[ECMWFSEAS5\]\(#\), Fig. 7\(a\)\(d\)\(w-g\). Therefore, the SICNet_{season} model achieves good generalization ability in predicting the SIEs of 2020-2023.](#)

带格式的: 字体: (默认) Times New Roman, (中文) Times New Roman

带格式的: 字体: (默认) Times New Roman, (中文) Times New Roman


带格式的: 上标



300 Figure 6. BACC of 2020-2023. (a) Persistence, (b) ~~ECMWF~~SEAS5, and (c) SICNet_{season}. Each value is a mean value of the four testing years. The horizontal axis represents the six lead months, and the vertical axis represents the target months, Jun. to Sep. The red signifies a high/improvement in ACC/BACC, and the blue signifies a decrease.

域代码已更改

带格式的: 字体: (中文)+中文正文 (宋体), 10 磅, 非加粗, (中文) 中文(中国)






Figure 7. Predicted Sep. SIEs and their BACCs of 2020-/2022/2023 in four to six months lead by Persistence, ECMWFSEAS5, and SICNet_{season}. (a)-(c) 2020, (d)-(f) 2022, (g)-(i) 2023.

带格式的: 字体: (默认) Times New Roman, (中文) Times New Roman

4.7 Comparison with the representative deep learning model

We compare the SICNet_{season} against the representative deep learning sea ice prediction model, the U-Net (IceNet-inspired) ~~IceNet~~ model. ~~The IceNet is a seasonal sea ice prediction model that performs state-of-the-art SIE probability prediction.~~ ~~The IceNet is a seasonal sea ice prediction model and achieves state-of-the-art performance in SIE prediction~~ (Andersson et al., 2021). It is a ~~CNN-based CNN-U-Net-based model for classification tasks-model~~, and it outputs the probability of three classes: open water (SIC≤15%), marginal ice (15% < SIC < 80%), and full ice (SIC≥80%). Differently, our SICNet_{season} outputs the 0-100% range SIC values. The IceNet's inputs consist of 50 monthly mean variables, including SIC, 11 climate variables, statistical SIC forecasts, and metadata.


~~The original IceNet model has some unique designs in inputs and training strategy. As we focus on the differences in model structures, we construct a U-Net (IceNet-inspired) model for comparison.~~

~~To make a fair comparison, we~~ We set the inputs (including SIT data) of the ~~IceNet-U-Net (IceNet-inspired)~~ to the same ones as SICNet_{season}. The loss function is also set as the NIIEE+MSE. We ~~reconstruct IceNet's~~ set the output layer of U-Net (IceNet-inspired) ~~by replacing the original softmax with theas~~ a sigmoid activation function to output. ~~The sigmoid function outputs continuous values of 0-100%. transform the original IceNet's output to the continuous values of 0-100%.~~ We also change the number of CNN filters to make the number of training parameters in U-Net (IceNet-inspired) ~~IceNet~~ equal to that in SICNet_{season}, about 140 million. The training and testing settings of U-Net (IceNet-inspired) ~~IceNet~~ are the same as those of SICNet_{season}. ~~The U-Net (IceNet-inspired) IceNet is trained using the same leave-one-year-out strategy as the SICNet_{season}.~~ ~~For example, if the testing year is 2019, the training set is data from 1979-2018, and the testing data is 2019. Then, the testing data moves to 2018; the remaining data (1979-2017, 2019) is the training set. For each training/testing pair, the model is trained three times to eliminate randomness, and the final prediction for testing data is the mean value of the three models.~~

带格式的: 缩进: 首行缩进: 0 厘米

带格式的: 缩进: 首行缩进: 1 字符

带格式的: 下标



域代码已更改

330 Figure 8. Detrended ACC of IceNet (a) and SICNet_{season} (b). (c) ACC difference obtained by SICNet_{season} minus U-Net (IceNet-inspired). BACC of U-Net (IceNet-inspired) (d) and SICNet_{season} (e). (f) BACC difference like (c). The red signifies a high/improvement in ACC/BACC, and the blue signifies a decrease.

带格式的: 字体: (中文)+中文正文 (宋体), 10 磅, 非加粗, (中文) 中文(中国)




Figure 9. The predicted Sep. SIEs of U-Net (IceNet-inspired) and SICNet_{season} in six months' lead: (a) 2012, (b) 2017, (c) 2018, and (d) 2019.

Fig. 8 shows the detrend ACC, BACC, and the differences between the two models. Compared with the U-Net (IceNet-inspired) model, our SICNet_{season} model significantly improves the ACC at most predictions, Fig. 8(c). For the target month, Aug./Sep., the SPB feature is obvious in the U-Net (IceNet-inspired): the maximum ACC gap is about 40%/30% at predictions made in May and Jun., Fig. 8(a). Our SICNet_{season} model optimizes the ACC gap with an improvement of 31.8%/20.8% at May's predictions, Fig. 8(c). The ACC improvements are also larger than 15% for predictions made before May. Therefore, compared with the state-of-the-art deep learning model U-Net (IceNet-inspired), our model achieves more skillful seasonal predictions by optimizing the SPB.

Unlike the ACC values, the BACC values of U-Net (IceNet-inspired) are more significant than those of SICNet_{season} on most predictions, Fig. 8(f). This result implies that U-Net (IceNet-inspired) is more dependent on SIE trends than SICNet_{season}. This difference can be attributed to the distinct fundamental units employed by the two models. The U-Net (IceNet-inspired) is a CNN-based model, and the weight-sharing mechanism of convolutional kernels forces the model to

350 capture the most "common" local dependencies in spatial. Though representative, these "common" local dependencies tend to yield smoother model outputs. The SICNet_{season} is a transformer-based model. The attention mechanism of the transformer can capture global dependencies without weight-sharing. As a result, "personalized" global dependencies are extracted, and the output is not smooth like the output of a CNN-based model. The "common" local dependencies have more apparent trend features than the "personalized" global dependencies. Fig. 9 shows the Sep.'s SIEs predicted by U-Net (IceNet-inspired) and
355 SICNet_{season} in the sixth-month lead. The SIEs of U-Net (IceNet-inspired) are smoother than those of SICNet_{season}. For 2012 and 2017, the SIEs' locations of the two models are very similar. For the other two years, the SIEs of U-Net (IceNet-inspired) match the observed SIEs better than those of SICNet_{season}. However, the SIEs of U-Net (IceNet-inspired) are over-smoothed and fail to characterize some abnormal characteristics, such as the SIE in Sep. 2018, Fig. 9(c).

Therefore, our transformer-based SICNet_{season} is more skillful than the representative CNN-based model U-Net (IceNet-inspired) in [optimizing the SPB](#)~~optimizing the predictions around the SPB~~. The SICNet_{season} exhibits a lower dependency on SIE trends and lower smooth results than the CNN-based model.

5 Conclusion

This study develops a deep-learning model, SICNet_{season}, to predict the Arctic SIC on a seasonal scale. The model is derived from a SwinUNET architecture. It inputs the historical SIC, SIT, and SIC climatology of target moths and predicts the SIC of
365 the next six months. A spatially constrained loss function NIIEE is employed to train the model considering sea ice distribution. We employ a 20-year (2000-2019) testing set to validate the model's performance. The summer season, Jun. to Sep., is the target period. The detrend ACC, BACC, and MAE are metrics. Comparison experiments with Persistence and seasonal predictions of [ECMWFSEAS5](#) are made to validate our model's performance. In particular, an ablation experiment is carried out to investigate the role of SIT data in [optimizing the SPB](#)~~optimizing the predictions around the SPB~~. A
370 generalization experiment with data from the last four years, 2020-2023, is carried out—the seasonal predictions of Sep. SIEs are analyzed. Finally, we discuss the advantages and disadvantages of our model and the typical CNN-based model, U-Net (IceNet-inspired). Given the mentioned efforts, our study draws the following conclusions.

First, our deep learning model, SICNet_{season}, is skillful in predicting the Arctic sea ice seasonally. Compared with the dynamic model [ECMWFSEAS5](#), SICNet_{season} optimizes the SPB significantly. The detrended ACC of Sep. SIE predicted by
375 SICNet_{season} in May/Apr. is improved by 7.7%/10.61% over the ACC predicted by the [ECMWFSEAS5](#). Compared with the anomaly persistence benchmark, the mentioned improvement is 41.02%/36.33%. Our deep learning model significantly reduces prediction errors of Sep.'s SIC on seasonal scales compared to [ECMWFSEAS5](#) and Persistence, a 20-30% reduction measured by MAE.

Second, the spring SIT data is key in [optimizing the SPB](#)~~optimizing the predictions around the SPB~~, contributing to a more
380 than 20% ACC enhancement in Sep.'s SIE at four to five months lead predictions. By including SIT data, the MAEs in the Beaufort Sea, the East Siberian Sea, and the Laptev Sea are reduced by more than 10% compared with those without SIT

data.

Third, our model achieves good generalization in predicting the Sep. SIEs of 2020-2023. When predicting the Sep.'s SIE in 2020/2023 (second/sixth lowest record) in May, SICNet_{season} achieved a BACC of 82.25%/82.08%, about 12%/10% higher than Persistence and [ECMWFSEAS5](#).

Fourth, our transformer-based SICNet_{season} is more skillful than the CNN-based ~~model~~ [U-Net \(-IceNet-inspired\) model](#) in seasonal sea ice predictions. Our SICNet_{season} model optimizes the ACC gap with an improvement of 31.8%/20.8% at May's predictions over the [U-Net \(IceNet-inspired\)](#) ~~IceNet~~. The SICNet_{season} exhibits a lower dependency on SIE trends and lower smooth results than the CNN-based model. This is due to the attention mechanism of the transformer operator extracting "personalized" global dependencies, while the CNN operator captures the most "common" local dependencies globally. The "common" local dependencies smooth the map and depend more on the trend than "personalized" ones.

Code and data availability

The code, the exact input/output data, and the saved well-trained weights of the developed model SICNet_{season} are available at <https://doi.org/10.5281/zenodo.14561423>.

395 Acknowledgment

This work was supported by the National Science Foundation of China (42206202), (U2006211), (42221005), Laoshan Laboratory Innovation Project (LSKJ202202302), (LSKJ202204302), in part by the China-Portugal Xinghai "Belt and Road" (2022YFE0204600).

Author contribution

400 All authors designed the experiments and carried them out. Yibin Ren developed and evaluated the model. Xiaofeng Li designed experiments and revised the manuscript. Yunhe Wang analyzed the experimental results.

Competing interests

The authors declare no competing interests.

405 **References**

- Andersson, T. R., Hosking, J. S., Pérez-Ortiz, M., Paige, B., Elliott, A., Russell, C., Law, S., Jones, D. C., Wilkinson, J., Phillips, T., Byrne, J., Tietsche, S., Sarojini, B. B., Blanchard-Wrigglesworth, E., Aksenov, Y., Downie, R., and Shuckburgh, E.: Seasonal Arctic sea ice forecasting with probabilistic deep learning, *Nat Commun*, 12, <https://doi.org/10.1038/s41467-021-25257-4>, 2021.
- 410 Blackport, R., Screen, J. A., van der Wiel, K., and Bintanja, R.: Minimal influence of reduced Arctic sea ice on coincident cold winters in mid-latitudes, *Nat Clim Chang*, 9, 697–704, <https://doi.org/10.1038/s41558-019-0551-4>, 2019.
- Blanchard-Wrigglesworth, E., Armour, K. C., Bitz, C. M., and Deweaver, E.: Persistence and inherent predictability of arctic sea ice in a GCM ensemble and observations, *J Clim*, 24, 231–250, <https://doi.org/10.1175/2010JCLI3775.1>, 2011.
- Blanchard-Wrigglesworth, E., Cullather, R. I., Wang, W., Zhang, J., and Bitz, C. M.: Model forecast skill and sensitivity to initial conditions in the seasonal Sea Ice Outlook, *Geophys Res Lett*, 42, 8042–8048, <https://doi.org/10.1002/2015GL065860>, 2015.
- 415 Blanchard-Wrigglesworth, E., Bushuk, M., Massonnet, F., Hamilton, L. C., Bitz, C. M., Meier, W. N., and Bhatt, U. S.: Forecast Skill of the Arctic Sea Ice Outlook 2008–2022, *Geophys Res Lett*, 50, <https://doi.org/10.1029/2022GL102531>, 2023.
- 420 Bonan, D. B., Bushuk, M., and Winton, M.: A spring barrier for regional predictions of summer Arctic sea ice, *Geophys Res Lett*, 46, 5937–5947, <https://doi.org/10.1029/2019GL082947>, 2019.
- [Bushuk, M., Ali, S., Bailey, D. A., Bao, Q., Batté, L., Bhatt, U. S., Blanchard-Wrigglesworth, E., Blockley, E., Cawley, G., Chi, J., Counillon, F., Coulombe, P. G., Cullather, R. I., Diebold, F. X., Dirkson, A., Exarchou, E., Göbel, M., Gregory, W., Guemas, V., Hamilton, L., He, B., Horvath, S., Ionita, M., Kay, J. E., Kim, E., Kimura, N., Kondrashov, D., Labe, Z. M., Lee, W. S., Lee, Y. J., Li, C., Li, X., Lin, Y., Liu, Y., Maslowski, W., Massonnet, F., Meier, W. N., Merryfield, W. J., Myint, H., Acosta Navarro, J. C., Petty, A., Qiao, F., Schröder, D., Schweiger, A., Shu, Q., Sigmond, M., Steele, M., Stroeve, J., Sun, N., Tietsche, S., Tsamados, M., Wang, K., Wang, J., Wang, W., Wang, Y., Wang, Y., Williams, J., Yang, Q., Yuan, X., Zhang, J., and Zhang, Y.: Predicting September Arctic Sea Ice: A Multimodel Seasonal Skill Comparison, *Bull Am Meteorol Soc*, 105, E1170–E1203, <https://doi.org/10.1175/BAMS-D-23-0163.1>, 2024.](https://doi.org/10.1175/BAMS-D-23-0163.1)
- 425 [Bushuk, M., Giannakis, D., and Majda, A. J.: Arctic Sea Ice Reemergence: The Role of Large-Scale Oceanic and Atmospheric Variability, *J Clim*, 28, 5477–5509, <https://doi.org/10.1175/JCLI-D-14-00354.1>, 2015.](https://doi.org/10.1175/JCLI-D-14-00354.1)
- 430 Bushuk, M., Winton, M., Bonan, D. B., Blanchard-Wrigglesworth, E., and Delworth, T. L.: A mechanism for the Arctic sea ice spring predictability barrier, *Geophys Res Lett*, 47, <https://doi.org/10.1029/2020GL088335>, 2020.
- 435 Bushuk, M., Zhang, Y., Winton, M., Hurlin, B., Delworth, T., Lu, F., Jia, L., Zhang, L., Cooke, W., Harrison, M., Johnson, N. C., Kapnick, S., Mchugh, C., Murakami, H., Rosati, A., Tseng, K.-C., Wittenberg, A. T., Yang, X., and Zeng, A. F.:

- Mechanisms of Regional Arctic Sea Ice Predictability in Two Dynamical Seasonal Forecast Systems, <https://doi.org/10.1175/JCLI-D-21, n.d.>
- Cao Hu and Wang, Y. and C. J. and J. D. and Z. X. and T. Q. and W. M.: Swin-Unet: Unet-Like Pure Transformer
440 for Medical Image Segmentation, in: Computer Vision – ECCV 2022 Workshops, 205–218, 2023.
- Cao, Y., Liang, S., Chen, X., He, T., Wang, D., and Cheng, X.: Enhanced wintertime greenhouse effect reinforcing Arctic amplification and initial sea-ice melting, *Sci Rep*, 7, <https://doi.org/10.1038/s41598-017-08545-2>, 2017.
- Cao, Y., Liang, S., Sun, L., Liu, J., Cheng, X., Wang, D., Chen, Y., Yu, M., and Feng, K.: Trans-Arctic shipping routes expanding faster than the model projections, *Global Environmental Change*, 73,
445 <https://doi.org/10.1016/j.gloenvcha.2022.102488>, 2022.
- ~~DiGirolamo, N. E., Parkinson, C. L., Cavalieri, D. J., Gloersen, P., and Zwally, H. J.: Sea ice concentrations from Nimbus-7 SMMR and DMSP SSM/I-SSMIS passive microwave data., Boulder, Colorado USA: NASA National Snow and Ice Data Center Distributed Active Archive Center, <https://doi.org/10.5067/MPYG15WAA4WX>, 2022. Cavalieri, D. J., Parkinson, C. L., Gloersen, P., and Zwally, H. J.: Sea ice concentrations from Nimbus-7 SMMR and DMSP SSM/I-SSMIS passive
450 ~~microwave data, Boulder, Colorado USA: NASA DAAC at the National Snow and Ice Data Center, 25, 1996.~~~~
- Chevallier, M., Méliá, D. S. Y., Voldoire, A., Déqué, M., and Garric, G.: Seasonal forecasts of the pan-arctic sea ice extent using a GCM-based seasonal prediction system, *J Clim*, 26, 6092–6104, <https://doi.org/10.1175/JCLI-D-12-00612.1>, 2013.
- Cohen, J., Screen, J. A., Furtado, J. C., Barlow, M., Whittleston, D., Coumou, D., Francis, J., Dethloff, K., Entekhabi, D., Overland, J., and Jones, J.: Recent Arctic amplification and extreme mid-latitude weather, <https://doi.org/10.1038/ngeo2234>,
455 2014.
- Collow, T. W., Wang, W., Kumar, A., and Zhang, J.: Improving arctic sea ice prediction using PIOMAS initial sea ice thickness in a coupled ocean-atmosphere model, *Mon Weather Rev*, 143, 4618–4630, <https://doi.org/10.1175/MWR-D-15-0097.1>, 2015.
- Day, J. J., Tietsche, S., and Hawkins, E.: Pan-arctic and regional sea ice predictability: Initialization month dependence, *J
460 Clim*, 27, 4371–4390, <https://doi.org/10.1175/JCLI-D-13-00614.1>, 2014.
- Ding, Q., Schweiger, A., L'Heureux, M., Battisti, D. S., Po-Chedley, S., Johnson, N. C., Blanchard-Wrigglesworth, E., Harnos, K., Zhang, Q., Eastman, R., and Steig, E. J.: Influence of high-latitude atmospheric circulation changes on summertime Arctic sea ice, *Nat Clim Chang*, 7, 289–295, <https://doi.org/10.1038/nclimate3241>, 2017.
- Dong, X., Yang, Q., Nie, Y., Zampieri, L., Wang, J., Liu, J., and Chen, D.: Antarctic sea ice prediction with A convolutional
465 long short-term memory network, <https://doi.org/10.1016/j.ocemod.2024.102386>, 1 August 2024.
- England, M. R., Eisenman, I., Lutsko, N. J., and Wagner, T. J. W.: The Recent Emergence of Arctic Amplification, *Geophys Res Lett*, 48, <https://doi.org/10.1029/2021GL094086>, 2021.
- Francis, J. A. and Vavrus, S. J.: Evidence linking Arctic amplification to extreme weather in mid-latitudes, *Geophys Res Lett*, 39, <https://doi.org/10.1029/2012GL051000>, 2012.

- 470 Goessling, H. F., Tietsche, S., Day, J. J., Hawkins, E., and Jung, T.: Predictability of the Arctic sea ice edge, *Geophys Res Lett*, 43, 1642–1650, <https://doi.org/10.1002/2015GL067232>, 2016.
- Gregory, W., Tsamados, M., Stroeve, J., and Sollich, P.: Regional september sea ice forecasting with complex networks and gaussian processes, *Weather Forecast*, 35, 793–806, <https://doi.org/10.1175/WAF-D-19-0107.1>, 2020.
- Guemas, V., Blanchard-Wrigglesworth, E., Chevallier, M., Day, J. J., Déqué, M., Doblas-Reyes, F. J., Fučkar, N. S., Germe, A., Hawkins, E., Keeley, S., Koenigk, T., Salas y Méliá, D., and Tietsche, S.: A review on Arctic sea-ice predictability and prediction on seasonal to decadal time-scales, *Quarterly Journal of the Royal Meteorological Society*, 142, 546–561, <https://doi.org/10.1002/qj.2401>, 2016.
- Jahn, A., Holland, M. M., and Kay, J. E.: [Projections of an ice-free Arctic Ocean](https://doi.org/10.1038/s43017-023-00515-9), <https://doi.org/10.1038/s43017-023-00515-9>, 1 March 2024.
- 480 Jenkins, M. and Dai, A.: The Impact of Sea-Ice Loss on Arctic Climate Feedbacks and Their Role for Arctic Amplification, *Geophys Res Lett*, 48, <https://doi.org/10.1029/2021GL094599>, 2021.
- Johnson, S. J., Stockdale, T. N., Ferranti, L., Balmaseda, M. A., Molteni, F., Magnusson, L., Tietsche, S., Decremier, D., Weisheimer, A., Balsamo, G., Keeley, S. P. E., Mogensen, K., Zuo, H., and Monge-Sanz, B. M.: SEAS5: The new ECMWF seasonal forecast system, *Geosci Model Dev*, 12, 1087–1117, <https://doi.org/10.5194/gmd-12-1087-2019>, 2019.
- 485 Jun Kim, Y., Kim, H. C., Han, D., Lee, S., and Im, J.: Prediction of monthly Arctic sea ice concentrations using satellite and reanalysis data based on convolutional neural networks, *Cryosphere*, 14, 1083–1104, <https://doi.org/10.5194/tc-14-1083-2020>, 2020.
- Kapsch, M. L., Graversen, R. G., and Tjernström, M.: Springtime atmospheric energy transport and the control of Arctic summer sea-ice extent, *Nat Clim Chang*, 3, 744–748, <https://doi.org/10.1038/nclimate1884>, 2013.
- 490 Kim, Y. H., Min, S. K., Gillett, N. P., Notz, D., and Malinina, E.: [Observationally-constrained projections of an ice-free Arctic even under a low emission scenario](https://doi.org/10.1038/s41467-023-38511-8), *Nat Commun*, 14, <https://doi.org/10.1038/s41467-023-38511-8>, 2023.
- Kumar, A., Perlwitz, J., Eischeid, J., Quan, X., Xu, T., Zhang, T., Hoerling, M., Jha, B., and Wang, W.: Contribution of sea ice loss to Arctic amplification, *Geophys Res Lett*, 37, <https://doi.org/10.1029/2010GL045022>, 2010.
- Kwok, R., Cunningham, G. F., Kacimi, S., Webster, M. A., Kurtz, N. T., and Petty, A. A.: Decay of the Snow Cover Over Arctic Sea Ice From ICESat-2 Acquisitions During Summer Melt in 2019, *Geophys Res Lett*, 47, <https://doi.org/10.1029/2020GL088209>, 2020.
- Landy, J. C., Dawson, G. J., Tsamados, M., Bushuk, M., Stroeve, J. C., Howell, S. E. L., Krumpen, T., Babb, D. G., Komarov, A. S., Heorton, H. D. B. S., Belter, H. J., and Aksenov, Y.: A year-round satellite sea-ice thickness record from CryoSat-2, *Nature*, 609, 517–522, <https://doi.org/10.1038/s41586-022-05058-5>, 2022.
- 500 Liang, X., Zhao, F., Li, C., Zhang, L., and Li, B.: Evaluation of ArcIOPS sea ice forecasting products during the ninth chinare-arctic in summer 2018, *Adv Polar Sci*, 31, 14–25, <https://doi.org/10.13679/j.advps.2019.0019>, 2020.
- Lindsay, R. W., Zhang, J., Schweiger, A. J., and Steele, M. A.: Seasonal predictions of ice extent in the Arctic Ocean, *J Geophys Res Oceans*, 113, <https://doi.org/10.1029/2007JC004259>, 2008.

- Liu, Z., Risi, C., Codron, F., He, X., Poulsen, C. J., Wei, Z., Chen, D., Li, S., and Bowen, G. J.: Acceleration of western Arctic sea ice loss linked to the Pacific North American pattern, *Nat Commun*, 12, <https://doi.org/10.1038/s41467-021-21830-z>, 2021a.
- Liu, Z., Lin, Y., Cao, Y., Hu, H., Wei, Y., Zhang, Z., Lin, S., and Guo, B.: Swin Transformer: Hierarchical Vision Transformer using Shifted Windows, in: 2021 IEEE/CVF International Conference on Computer Vision (ICCV), 9992–10002, <https://doi.org/10.1109/ICCV48922.2021.00986>, 2021b.
- 510 Li, X., Liu, B., Zheng, G., Ren, Y., Zhang, S., Liu, Y., Gao, L., Liu, Y., Zhang, B., and Wang, F.: Deep-learning-based information mining from ocean remote-sensing imagery, *Natl Sci Rev*, 7, <https://doi.org/10.1093/NSR/NWAA047>, 2021.
- Li, Y., Qiu, Y., Jia, G., Yu, S., Zhang, Y., Huang, L., and Lepparanta, M.: An Explainable Deep Learning Model for Daily Sea Ice Concentration Forecast, *IEEE Transactions on Geoscience and Remote Sensing*, 62, 1–17, <https://doi.org/10.1109/TGRS.2024.3386930>, 2024.
- 515 Merryfield, W. J., Lee, W. S., Wang, W., Chen, M., and Kumar, A.: Multi-system seasonal predictions of Arctic sea ice, *Geophys Res Lett*, 40, 1551–1556, <https://doi.org/10.1002/grl.50317>, 2013.
- Min, C., Yang, Q., Chen, D., Yang, Y., Zhou, X., Shu, Q., and Liu, J.: The Emerging Arctic Shipping Corridors, *Geophys Res Lett*, 49, <https://doi.org/10.1029/2022GL099157>, 2022.
- Msadek, R., Vecchi, G. A., Winton, M., and Gudgel, R. G.: Importance of initial conditions in seasonal predictions of Arctic sea ice extent, *Geophys Res Lett*, 41, 5208–5215, <https://doi.org/10.1002/2014GL060799>, 2014.
- 520 Mu, B., Luo, X., Yuan, S., and Liang, X.: IceTFT v1.0.0: interpretable long-term prediction of Arctic sea ice extent with deep learning, *Geosci Model Dev*, 16, 4677–4697, <https://doi.org/10.5194/gmd-16-4677-2023>, 2023.
- Mu, L., Nerger, L., Tang, Q., Loza, S. N., Sidorenko, D., Wang, Q., Semmler, T., Zampieri, L., Losch, M., and Goessling, H. F.: Toward a data assimilation system for seamless sea ice prediction based on the AWI climate model, *J Adv Model Earth Syst*, 12, <https://doi.org/10.1029/2019MS001937>, 2020.
- 525 Nakanowatari, T., Inoue, J., Zhang, J., Watanabe, E., and Kuroda, H.: A New Norm for Seasonal Sea Ice Advance Predictability in the Chukchi Sea: Rising Influence of Ocean Heat Advection, *J Clim*, 35, 2723–2740, <https://doi.org/10.1175/JCLI-D-21-0425.1>, 2022.
- Olonscheck, D., Mauritsen, T., and Notz, D.: Arctic sea-ice variability is primarily driven by atmospheric temperature fluctuations, *Nat Geosci*, 12, 430–434, <https://doi.org/10.1038/s41561-019-0363-1>, 2019.
- 530 Palerme, C., Lavergne, T., Rusin, J., Melsom, A., Brajard, J., Kvanum, A. F., MacDonald Sørensen, A., Bertino, L., and Müller, M.: Improving short-term sea ice concentration forecasts using deep learning, *Cryosphere*, 18, 2161–2176, <https://doi.org/10.5194/tc-18-2161-2024>, 2024.
- Pithan, F. and Mauritsen, T.: Arctic amplification dominated by temperature feedbacks in contemporary climate models, *Nat Geosci*, 7, 181–184, <https://doi.org/10.1038/ngeo2071>, 2014.
- 535 Reichstein, M., Camps-Valls, G., Stevens, B., Jung, M., Denzler, J., Carvalhais, N., and Prabhat: Deep learning and process understanding for data-driven Earth system science, *Nature*, 566, 195–204, <https://doi.org/10.1038/s41586-019-0912-1>, 2019.

- Ren, Y. and Li, X.: Predicting the Daily Sea Ice Concentration on a Subseasonal Scale of the Pan-Arctic During the Melting Season by a Deep Learning Model, *IEEE Transactions on Geoscience and Remote Sensing*, 61, <https://doi.org/10.1109/TGRS.2023.3279089>, 2023.
- 540 Ren, Y., Li, X., and Zhang, W.: A data-driven deep learning model for weekly sea ice concentration prediction of the Pan-Arctic during the melting season, *IEEE Transactions on Geoscience and Remote Sensing*, 60, <https://doi.org/10.1109/TGRS.2022.3177600>, 2022.
- Schweiger, A., Lindsay, R., Zhang, J., Steele, M., Stern, H., and Kwok, R.: Uncertainty in modeled Arctic sea ice volume, *J Geophys Res Oceans*, 116, <https://doi.org/10.1029/2011JC007084>, 2011.
- 545 Screen, J. A. and Simmonds, I.: The central role of diminishing sea ice in recent Arctic temperature amplification, *Nature*, 464, 1334–1337, 2010.
- Screen, J. A., Simmonds, I., Deser, C., and Tomas, R.: The atmospheric response to three decades of observed arctic sea ice loss, *J Clim*, 26, 1230–1248, <https://doi.org/10.1175/JCLI-D-12-00063.1>, 2013.
- 550 Shu, Q., Wang, Q., Arthun, M., Wang, S., Song, Z., Zhang, M., and Qiao, F.: Arctic Ocean Amplification in a warming climate in CMIP6 models, *Sci Adv*, 8, eabn9755, <https://doi.org/10.1126/sciadv.abn9755>, 2022.
- Sigmond, M., Fyfe, J. C., Flato, G. M., Kharin, V. V., and Merryfield, W. J.: Seasonal forecast skill of Arctic sea ice area in a dynamical forecast system, *Geophys Res Lett*, 40, 529–534, <https://doi.org/10.1002/grl.50129>, 2013.
- Song, C., Zhu, J., and Li, X.: Assessments of Data-Driven Deep Learning Models on One-Month Predictions of Pan-Arctic Sea Ice Thickness, *Adv Atmos Sci*, 41, 1379–1390, <https://doi.org/10.1007/s00376-023-3259-3>, 2024.
- 555 Stroeve, J. and Notz, D.: Insights on past and future sea-ice evolution from combining observations and models, *Glob Planet Change*, 135, 119–132, <https://doi.org/https://doi.org/10.1016/j.gloplacha.2015.10.011>, 2015.
- Tietsche, S., Day, J. J., Guemas, V., Hurlin, W. J., Keeley, S. P. E., Matei, D., Msadek, R., Collins, M., and Hawkins, E.: Seasonal to interannual Arctic sea ice predictability in current global climate models, *Geophys Res Lett*, 41, 1035–1043, <https://doi.org/10.1002/2013GL058755>, 2014.
- 560 Wang, L., Yuan, X., Ting, M., and Li, C.: Predicting summer arctic sea ice concentration intraseasonal variability using a vector autoregressive model, *J Clim*, 29, 1529–1543, <https://doi.org/10.1175/JCLI-D-15-0313.1>, 2016.
- Wang, W., Chen, M., and Kumar, A.: Seasonal prediction of arctic sea ice extent from a coupled dynamical forecast system, *Mon Weather Rev*, 141, 1375–1394, <https://doi.org/10.1175/MWR-D-12-00057.1>, 2013.
- 565 Wang, Y., Yuan, X., Bi, H., Bushuk, M., Liang, Y., Li, C., and Huang, H.: Reassessing seasonal sea ice predictability of the Pacific-Arctic sector using a Markov model, *Cryosphere*, 16, 1141–1156, <https://doi.org/10.5194/tc-16-1141-2022>, 2022.
- Wu, A., Che, T., Li, X., and Zhu, X.: Routeview: an intelligent route planning system for ships sailing through Arctic ice zones based on big Earth data, *Int J Digit Earth*, 15, 1588–1613, <https://doi.org/10.1080/17538947.2022.2126016>, 2022.
- 570 Yang, Q., Mu, L., Wu, X., Liu, J., Zheng, F., Zhang, J., and Li, C.: Improving Arctic sea ice seasonal outlook by ensemble prediction using an ice-ocean model, *Atmos Res*, 227, 14–23, <https://doi.org/10.1016/j.atmosres.2019.04.021>, 2019.

[Yang, Z., Liu, J., Song, M., Hu, Y., Yang, Q., and Fan, K.: Extended seasonal prediction of Antarctic sea ice using ANTSIC-UNet, EGU sphere, 2024, 1–25, <https://doi.org/10.5194/egusphere-2024-1001>, 2024.](https://doi.org/10.5194/egusphere-2024-1001)
~~Yang, Z., Liu, J., Song, M., Hu, Y., Yang, Q., and Fan, K.: Extended seasonal prediction of Antarctic sea ice using ANTSIC UNet, <https://doi.org/10.5194/egusphere-2024-1001>, n.d.~~

575

Yuan, X., Chen, D., Li, C., Wang, L., and Wang, W.: Arctic sea ice seasonal prediction by a linear markov model, *J Clim*, 29, 8151–8173, <https://doi.org/10.1175/JCLI-D-15-0858.s1>, 2016.

Zampieri, L., Goessling, H. F., and Jung, T.: Bright Prospects for Arctic Sea Ice Prediction on Subseasonal Time Scales, *Geophys Res Lett*, 45, 9731–9738, <https://doi.org/10.1029/2018GL079394>, 2018.

580 Zeng, J., Yang, Q., Li, X., Yuan, X., Bushuk, M., and Chen, D.: Reducing the Spring Barrier in Predicting Summer Arctic Sea Ice Concentration, *Geophys Res Lett*, 50, <https://doi.org/10.1029/2022GL102115>, 2023.

Zhang, J. and Rothrock, D. A.: Modeling Global Sea Ice with a Thickness and Enthalpy Distribution Model in Generalized Curvilinear Coordinates, *Mon Weather Rev*, 131, 845–861, [https://doi.org/10.1175/1520-0493\(2003\)131<0845:MGSIIWA>2.0.CO;2](https://doi.org/10.1175/1520-0493(2003)131<0845:MGSIIWA>2.0.CO;2), 2003.

585 Zhang, J., Steele, M., Lindsay, R., Schweiger, A., and Morison, J.: Ensemble 1-Year predictions of Arctic sea ice for the spring and summer of 2008, *Geophys Res Lett*, 35, <https://doi.org/10.1029/2008GL033244>, 2008.

Zhang, Y.-F., Bushuk, M., Winton, M., Hurlin, B., Delworth, T., Harrison, M., Jia, L., Lu, F., Rosati, A., and Yang, X.: Subseasonal-to-Seasonal Arctic Sea Ice Forecast Skill Improvement from Sea Ice Concentration Assimilation, *J Clim*, 35, 4233–4252, <https://doi.org/10.1175/JCLI-D-21-0548.1>, 2022.

590 Zhang, Y. F., Bushuk, M., Winton, M., Hurlin, B., Gregory, W., Landy, J., and Jia, L.: Improvements in September Arctic Sea Ice Predictions Via Assimilation of Summer CryoSat-2 Sea Ice Thickness Observations, *Geophys Res Lett*, 50, <https://doi.org/10.1029/2023GL105672>, 2023.

Zhu, Y., Qin, M., Dai, P., Wu, S., Fu, Z., Chen, Z., Zhang, L., Wang, Y., and Du, Z.: Deep Learning-Based Seasonal Forecast of Sea Ice Considering Atmospheric Conditions, *Journal of Geophysical Research: Atmospheres*, 128,

595 <https://doi.org/10.1029/2023JD039521>, 2023.