

Combining empirical and mechanistic understanding of spruce bark beetle outbreak dynamics in the LPJ-GUESS (v4.1, r13130) vegetation model

Fredrik Lagergren¹, Anna Maria Jönsson¹, Mats Lindeskog¹, and Thomas A. M. Pugh^{1,2,3}

¹Department of Physical Geography and Ecosystem Science, Lund University, Lund, Sweden.

²School of Geography, Earth and Environmental Sciences, University of Birmingham, Birmingham, UK.

³Birmingham Institute of Forest Research, University of Birmingham, Birmingham, UK.

Correspondence to: Fredrik Lagergren (Fredrik.lagergren@nateko.lu.se)

Abstract. For evaluating the forests' performance in the future, dynamic vegetation models are important tools.

Tree mortality is an important function in such models and, especially for needle leaved forest in the temperate and boreal zones, bark beetles are important for the mortality pattern. The European spruce bark beetle (SBB, *Ips typographus*) has in the most recent years replaced wind as the most important disturbance agent in European forests. Historically, SBB damage is typically triggered by wind storms as they create breeding material with no defences to overcome for the beetles. Drought can contribute to increased damage and prolonged outbreaks by lowering the defence of the trees, but has been the main driver of some of the European forest damage in the last decade. In this study we implemented an SBB damage module in a dynamic vegetation model (LPJ-GUESS) that includes representation of wind damage and forest management, with the aim to present a general concept that can be used also for other bark beetle species. The module was calibrated against observations of storm and SBB damage in Sweden, Switzerland, Austria and France. An index of the SBB population size that changed over time driven by phenology, drought, storm felled spruce trees and density of the beetle population, was used to scale modelled damage. The model was able to catch the start and duration of outbreaks triggered by storm damage reasonably well but there were discrepancies in levels, which partly can be related to salvage logging of storm felled forest and sanitary cutting of infested trees. The model showed increased damage in most recent years with warm and dry conditions, although below the level reported, which may suggest that the drought stress response of spruce in LPJ-GUESS is underestimated. The new model forms a basis to explore vulnerability of European forests to spruce bark beetle infestations.

1. Introduction

Intensified forest management, reforestation and fertilization effects from increased CO₂ concentration and nitrogen deposition have promoted the growth and stock of European forests over recent decades (Ciais et al., 2008; Scheel et al., 2022). However, over the most recent years this trend no longer exists, due to increased harvest and disturbances (Palahi et al., 2021; Patacca et al., 2022; Wernick et al., 2021). The disturbances and associated tree mortality can be related to weather and climate, directly or in combination with a biological agent, and the forest's structure and species composition is important as most epidemic species are selective in host preference (Balla et al., 2021). The prevailing paradigm of European forestry has increased the vulnerability of forests to disturbances by promoting monocultures (Forzieri et al., 2021). Forest policy in Europe is determined both at the national and European Union levels. Tools that can provide projections of how disturbances are likely to change

under different climate conditions and management actions at these national-to-continental scales are important to be able to develop appropriate forest policy.

40 Bark beetles are a group of insect species that in most cases colonize dead or stressed trees. A few species are, however, also able to kill healthy trees, potentially leading to outbreaks that can cause severe mortality, especially in temperate and boreal needle-leaved forests (Kautz et al., 2017; Lindgren and Raffa, 2013). In Europe, the most damaging bark beetle is the spruce bark beetle (*Ips typographus*, hereafter referred to as SBB), causing tree death corresponding to tens of million m³ wood per year (Patacca et al., 2022). The most recent years have seen accelerating losses of trees from SBB outbreaks, mainly related to droughts reducing the capacity of trees to defend themselves and climate warming accelerating the lifecycle of the beetles (Hlásny et al., 2021). Historically, 45 however, outbreaks have mainly been triggered by storm damage, which creates breeding material in which the beetles are not required to overcome the tree defence (Schroeder, 2001). Norway spruce (*Picea abies*), the host tree, is favoured by forestry and is often managed with thinning and clear-cuts that creates dense even-aged stands, that over time becomes increasingly vulnerable to both storm damage and SBB attacks. The interaction of these multiple drivers makes SBB damage challenging to model at any scale. To further complicate the situation, 50 counter measures such as salvage and sanitary cutting (SSC), which are effective in preventing and shorten outbreaks of SBB, have a high variation in intensity over time and space. Salvage logging of storm felled trees reduce the material where SBB can have very efficient regeneration and can even reduce the SBB population if the harvest is done between infection and emergence of the new generation. Sanitary cutting of infected living 55 trees will take away the new generation, which can be a significant factor for ending an outbreak situation. Whilst landscape- and national-level models for SBB exist (De Bruijn et al., 2014; Jönsson et al., 2012; Marie et al., 2024; Seidl and Rammer, 2017; Seidl et al., 2014; Temperli et al., 2013), the capability to explicitly model historical forest damage from SBB has not yet been demonstrated at European-scale.

The life cycle and temperature dependent phenology of SBB are well studied (Wermelinger and Seifert, 1998). 60 However, to be able to evaluate model simulations on spruce bark beetle population dynamics and the shift between endemic and epidemic conditions, not only phenological understanding is needed, but also detailed knowledge about forest conditions. This includes stand composition and structure, timing and magnitude of storm damage, and subsequent management actions such as SSC (Jönsson et al., 2012). So far, empirical approaches have been applied in SBB modelling at continental (Marini et al., 2017) and regional scale (Soukhovolsky et al., 65 2022). Most empirical models include response functions of storm felling, temperature and precipitation deficit, but require prior knowledge of the SBB population, as the output from such models is a relative change in SBB population or damage, limiting the simulation of full outbreak cycles. In detailed mechanistic models, the timing of SSC counter measures is an important component (Jönsson et al., 2012), which makes data availability a limitation to large-scale model calibration. The dynamical vegetation model (DVM) ORCHIDEE has recently 70 been updated with a mechanistic SBB functionality (Marie et al., 2024), but lacks SSC functionality. The iLand landscape simulator has mechanistic SBB components and can simulate salvage logging, but has been evaluated in protected areas to rule out the influence of SSC (Seidl and Rammer, 2017). A model without mechanistic responses of climate and forest structure may, however, not be able to project future conditions and the effect of adaption in proactive and reactive forest management strategies. Conversely, the SBB lifecycle has several 75 temperature dependent stages for reproduction and survival, which makes representation by a continuous response

function, as typically employed by empirical models, difficult. Yet, when working over large regions with gridded climate datasets, the weather variability within a grid cell related to altitude, aspect and forest edge effects smooth out some of the non-linearity. This means that an empirical approach can be efficient in applications over large regions, also when considering the highest resolution of gridded climate data available today (typically 3 km, e.g. Lind et al., 2020). An approach that blends the insight of mechanistic modelling with the ability of large-scale empirical assessments to capture the net effects of unrepresented smaller-scale processes is needed.

Here, we seek to develop and evaluate a model for forest damage from SBB outbreaks within the LPJ-GUESS dynamic vegetation model (Smith et al., 2001; Smith et al., 2014) that are able to simulate the magnitude and frequency of large outbreak events, as well as the typical background rates. This is important when the model is applied at relatively large scales with a focus on carbon cycling, forest productivity and resilience. It is also important to be able to simulate the impacts of management interventions such as SSC, and systematic changes in forest structure. Furthermore, because SBB is only one of many bark beetles that have large impacts on forest dynamics across northern forests (Kautz et al., 2017), there is also a need to develop a model structure that is flexible to simulate a category of pest with multiple bark beetle species with different characteristics across a wide variety of forest across the temperate and boreal forest biomes. The aim of this study was therefore to develop a model with the following characteristics:

1. To be able to catch outbreak dynamics, triggered by storm damage, drought stress and temperature-driven changes in beetle phenology, without detailed accounting of SBB population.
2. To utilise simple empirical relationships where available, but also make use of suitable mechanistic knowledge that is relevant at the scale of interest.
3. To generate fractions of killed individuals in vegetation in tree size/age cohorts for feedback to modelled vegetation dynamics
4. To provide a generalised concept to use for different types of bark beetles by changing the underlying functions related to weather and insect – host tree interactions.

To achieve this, a semi-empirical SBB damage model, including SSC functionality, with components of negative feedback from a dense SBB population, amplification of damage after storm felling, where also warm and dry weather can trigger and contribute to sustained outbreaks, will be calibrated based on storm and SBB damage statistics from four countries in Europe. In this model, we seek to represent key aspects of the interaction of SBB with climate change and management, specifically phenology, SSC, interaction with storm and drought, whilst minimising process complexity by adopting more empirical elements wherever possible.

2. Material and Methods

2.1. General description of LPJ-GUESS

LPJ-GUESS is a dynamic vegetation model adapted to both global and European simulation domains, which simulates the development of vegetation cohorts belonging to different plant functional types (PFT) in replicate patches (Smith et al., 2001; Smith et al., 2014). The cohorts compete for water, nutrients and light within the patches, driven by climate, nitrogen deposition and atmospheric CO₂ concentration. The PFTs differ in parameters

related to physiological response functions, allometry and bioclimatic restrictions. Replicate patches are subject to the same climatic and edaphic conditions, but differ due to stochastic elements in the processes of cohort establishment and individual tree mortality and disturbance. A range of different forest management classes can be simulated in each model grid cell, for instance clear-cut or continuous harvesting following a particular regime or unmanaged vegetation (Lindeskog et al., 2021). Within these classes, there may be patches of different ages since the last patch-destroying (i.e. stand-replacing) natural disturbance or clear-cut harvest event. Processes in the model includes e.g. light absorption, photosynthesis, auto- and heterotrophic respiration, allocation and different types of mortality. Patch destroying disturbances apart from fire have typically been simulated as random events with a fixed average return time.

In the present study LPJ-GUESS version 4.1, subversion revision 13130 was used. In addition to the standard trunk version of the model, this revision also included the storm damage module from Lagergren et al. (2012). In this module, the simulated storm-damaged fraction of a cohorts is the product of a cohort's sensitivity index (SI), the triggering wind load (WL) and a calibration factor (CF).

$$DF_{\text{cohort}} = SI_{\text{cohort}} \times WL \times CF \quad (1)$$

The SI is based on tree shape (height to diameter ratio), neighbouring cohorts' and stands' height and time since thinning.

2.2. Setup of simulations representing the current state of spruce forest

The global set of PFTs were used in the simulations, as the aim was to produce a biotic model that could be applied routinely in the standard version of LPJ-GUESS. This set has seven boreal and temperate tree PFTs, of which spruce is represented by the shade tolerant boreal needle leaved evergreen tree type (BNE). The spruce forest for each region/country was represented in the simulations by nine age classes of clear-cut forestry with a 90-year rotation period established at 10-year intervals after year 1859, continuous-cover forestry (CCF) with short cutting interval (12 years, 15% of biomass removed at each occasion) and long (25 years, 30%) and unmanaged forest. For the clear-cut rotations thinning was done at age 9, 27 and 45 years with strength of 10%, 30% and 25% of biomass removed. Planting or establishment in the managed forest types were set to BNE only. The results were weighted by weights based on the age-class distribution from Poulter et al. (2019) for year 2010. The age-class data had a regional resolution for France and a national for Austria and Switzerland. For Sweden national inventory data for 2008-2012 with a county resolution were used instead of the Poulter et al. (2019) data (Skogsdata 2013, <https://www.slu.se/en/Collaborative-Centres-and-Projects/the-swedish-national-forest-inventory/foreststatistics/skogsdata/>). To get also weights for CCF and potential natural vegetation (PNV) from the datasets, the short and long CCF classes were used to represent the fraction of the 91-110 and 111-140 years age classes in the inventory data respectively, and PNV was used to represent the fraction of forest older than 140 years. The weights, related to the age in 2010, were used as input to the landcover functionality in LPJ-GUESS (Lindeskog et al., 2021) and were held constant over the simulations. Each age class, CCF type and natural forest was run for five replicate patches in each grid cell. Random patch-destroying disturbances were initially set to a return time of 500 years, but after the introduction of management in a patch, these were turned off to avoid resetting the age structure. Fire disturbance was simulated with the BLAZE module and it was also turned off for

managed patches. In Europe, BNE also represents silver fir (*Abies alba*), which is not attacked by SBB, this could potentially lead to overestimation of SBB damage in stands simulated as unmanaged, as BNE in those stands would represent both species. In Switzerland that has the highest fraction of PNV (25%), silver fir only makes up 6.6% of the total spruce and fir standing volume. In the French regions there are 0-12% PNV, but here silver fir makes up 49% of total fir and spruce. Austria has only 7% PNV and in Sweden there are no silver fir. Since PNV multiplied by fir fraction was low (<6%) in all regions/countries, we ignored this problem. Furthermore, as *Picea abies* is the primary species constituting the BNE PFT within Europe, the parameterisation here can also be considered applicable in the European version of the model (Pugh et al., Manuscript).

2.3. Implementation of a bark beetle damage module in LPJ-GUESS

European storm and bark beetle damage statistics were compiled by Marini et al. (2017), and they used the dataset to derive empirical models describing the increase rate (R) of forest volume loss due to bark beetles (D_{SBB}) to one year (t) from the previous year ($t-1$):

$$R = \log_e(D_{SBB\ t}/D_{SBB\ t-1}) \quad (2)$$

The top-rated model was:

$$R = -0.099 + 0.223T_t + 0.265D_{storm\ t-1} - 0.351D_{SBB\ t-1} - 0.151W_{t-1} - 0.052W_t - 0.233T_t D_{storm\ t-1} + 0.153T_t W_{t-1} + 0.233D_{storm\ t-1} W_t, \quad (3)$$

depending on the thermal sum between 1 May and 30 July (T) with a threshold of $+5^\circ\text{C}$, storm felled volume (D_{storm}), D_{SBB} , and cumulative rainfall between 1 March and 31 July (W). All variables were standardized to mean 0 and standard deviation 1, D_{storm} was log-transformed before standardization.

Using this concept, we implemented an additive model for bark beetle damage in LPJ-GUESS, taking advantage of the existing LPJ-GUESS formulations for water stress and mortality of trees (L , stem litter from spruces larger than a limit (d_{lim}) killed by other agents than bark beetles last year), and the bark-beetle phenology of Jönsson et al. (2007).

$$R = f(P_{gridcell\ t-1}) + f(P_{patch\ t-1}/L) + f(\text{water stress}) + f(\text{phenology}) \quad (4)$$

$$P_{patch} = e^R P_{patch\ t-1} \quad (5)$$

$$M = P_{patch} k_0 = e^R P_{patch\ t-1} k_0 \quad (6)$$

In which an index of the population size at the start of the year ($P_{patch\ t-1}$), calculated both at patch and grid-cell level, determines the mass of bark-beetle killed trees (M) together with R and a calibration factor (k_0). At start of a simulation P_{patch} was initiated with a value of 10 for all patches. As there is a linear dependency of R for M (Eq. 5), R representing increase rate of damage (Eq. 2) corresponds to R representing increase rate of P_{patch} . Of the components of R (Fig. 1, Eq. 4), $f(P_{patch}/L)$ represents the negative feedback from a denser population relative to the amount of substrate with no defence (L , typically with a high value after storm damage) for a group of trees (patch), $f(P_{gridcell})$ represent the negative feedback when a high population in the landscape leads to lower patch level R because of swarming induced competition, $f(\text{water stress})$ has a positive impact on R as the defence in

healthy trees is reduced with water limitation, and $f(\text{phenology})$ has a positive impact resulting from faster phenological development of the SBB.

185 The fraction of the different age and management classes, from the landcover functionality in LPJ-GUESS, were used to calculate P_{gridcell} weighted over the classes. With all variables in the (Marini et al., 2017) model at ± 2 standard deviation from the mean, R has a range of $-4.66 - 3.36$, but interactions between variables prevent it to reach higher numbers. R calculated from the observation data used in the present study (see 2.4 below) has a range of $-2.2 - 2.9$, but initial high numbers in the start of an outbreak were often missing as inventories only began
190 when already under an outbreak situation. The total range of R (Fig. 1a) in the presented model was set to $-3.8 - 6.0$, where the possible outcome range of the different parts of the model (Eq. 4, Fig. 1a) were given weights of the similar magnitude as in the (Marini et al., 2017) model. The maximum R can also be translated to an extreme case of population increase rate with two successful generations in a year with 21 female offspring per mother ($e^{6.1} = 21.1^2$). To enable that an outbreak can also be sustained at the highest population levels, the lowest possible
195 total negative feedback from population size is just below the highest possible positive impact of water stress and phenology.




Figure 1: The different components of the increase rate of the bark beetle population index (R). (a) Ranges of the components and the total (Eq. 4). For $f(P_{\text{gridcell}})$ and $f(P_{\text{patch}}/L)$, the light shaded areas show the part of the ranges that were varied in the parameter optimization, where the sum of the minimum of $f(P_{\text{gridcell}})$ and $f(P_{\text{patch}}/L)$ were kept constant to have the possible total negative feedback from the population index constant (see section 2.6). (b-e) Shape of the functions for the components of the increase rate of the bark beetle population index (R), b Eq. 7, c Eq. 8, d Eq. 10, e Eq. 12. The default parameter setting (Table 7) is shown by thick grey lines (b-d). The functions are also shown in colour for the min and max value of parameters included in the calibration and sensitivity analysis (section 2.6), using the default setting for the other parameters. For $f(\text{phenology})$, (e), no parameters were tested but the response depends on the grid-cell's 30-year running mean of the length of the autumn swarming period (ASP, ASP30) and function are shown for ASP30 from 2 to 75 days.

The response function for the P_{gridcell} component of R (Fig. 1b, Eq. 4) was:

$$f(P_{\text{gridcell}}) = \text{MIN}(k_{\text{gc_max}}, \text{MAX}(k_{\text{gc_min}}, -\log_e(P_{\text{gridcell}} \times k_1) \times k_2)), \quad (7)$$

where $k_{\text{gc_max}}$ and $k_{\text{gc_min}}$ determine the range, k_1 the intercept and k_2 the slope of the expression. The same type of function was used for the combined response of P_{patch} and L (Fig. 1c, Eq. 4):

$$f(P_{\text{patch } t-1}/L) = \text{MIN}(k_{p_max}, \text{MAX}(k_{p_min}, -\log_e(P_{\text{patch}}/L \times k_3) \times k_4)) , \quad (8)$$

215 where the min and max are set by k_{p_max} and k_{p_min} respectively, and k_3 and k_4 set the intercept and slope. To avoid division by zero, a fixed background level (k_{base_bm} , set to 0.0001 kg C m⁻²) was added to the available brood material (L):

$$L = k_{base_bm} + L_{mortality} , \quad (9)$$

220 where $L_{mortality}$ is C mass of stem mortality of spruce trees above a diameter threshold (d_{lim}) for previous year caused by other reasons than bark beetles (including storm), d_{lim} was set to 15 cm (Jönsson et al., 2012). The dependency of $P_{gridcell}$, P_{patch} and L corresponds to D_{storm} and D_{SBB} in the Marini et al. (2017) model. To take advantage of LPJ-GUESS' ability to model drought impact, the Marini et al. (2017) dependency of rainfall was replaced with a dependency of the ratio between water supply to the canopy and canopy water demand ($wscal$), as calculated by LPJ-GUESS. The value goes from zero at complete shutdown of photosynthesis and transpiration to one at no stress, and it was used to assess the dependency of drought ($1 - wscal$, Fig. 1d, Eq. 4):

$$f(\text{water stress}) = \text{MIN}(k_{drought_max}, (1 - wscal_{mean}) \times k_5) , \quad (10)$$

230 where k_5 is the slope, determining the point of full effect on R . The mean $wscal$ calculated over the month May-July for both previous year ($wscal_{t-1}$) and current year ($wscal_t$) for the BNE PFT were used. Based on the (Marini et al., 2017) model, the data from the previous year were given a three times higher weight in the default setting, but in the calibration and sensitivity analysis (see section 2.4 below) the previous year weight (k_{pyw}) was varied between 1/4 to 4:

$$wscal_{mean} = \frac{wscal_{t-1}k_{pyw} + wscal_t}{k_{pyw} + 1} \quad (11)$$

235 For a more mechanistic approach of taking phenology into account, the dependency of T in the (Marini et al., 2017) model was replaced with a dependency on the length of the autumn swarming period (ASP, Eq. 4) in comparison with the grid-cell specific 30-year average as:

$$f(\text{phenology}) = \text{ASP} \frac{k_6}{\text{ASP}_{30} + k_7} , \quad (12)$$

240 where ASP is the number of flight days of the first new generation according to Jönsson et al. (2011). The slope depends on the grid-cell-level 30-year running mean of ASP (ASP_{30}) and two parameters k_6 (slope) and k_7 (dampening). Calculated ASP was capped at 90 days (ASP_{max}). Normalizing with ASP_{30} instead of a direct relationship ($R_{phen} = \text{ASP} \times 2 / \text{ASP}_{max}$) gives a more responsive function at lower ASP_{30} (Fig. 1e). The dependency of ASP instead of, e.g., the number of generations per year was chosen as ASP is a continuous variable which better catch the average when there is a variability in the climate, such in mountainous regions, then a discrete variable.

245 In the managed forest of Europe, countermeasures against outbreaks are often performed. We included functionality of salvage cutting of storm-felled trees and sanitary cutting of infested trees (SSC). The salvage cutting part was done by reducing $L_{mortality}$ by 90% for larger storm events ($>5 \text{ m}^3 \text{ ha}^{-1}$ wood at patch level). In Sweden forest owners are not allowed to leave more than 5 m³ of damaged spruce wood with $d > 10$ cm per hectare after

a storm, as regulated by the Swedish Forestry Act (Swedish Forest Agency, <https://www.skogsstyrelsen.se/en/laws-and-regulations/skogsvarsdlagen/>, last access: 12 May 2025). The total grid-cell maximum salvage cutting capacity ($salv_{max}$) was set to 50% of the 10-year average harvest rate, as it has to be done before the new generation of bark beetles emerge (which occurs approximately 6 months into the year) in order to have an effect. If the 90% of the storm damage ($damage_{available}$) was $> salv_{max}$, L_{mort} was reduced by $salv_{max}/damage_{available}$ instead of 90%. Sanitary cutting was applied by reducing P_{patch} by 25% if the fraction of available spruce volume that would be killed was $>1\%$. This reduction was done before Eq. (6) was applied. At this stage of the model development the effect of the SSC was just applied in the bark beetle accounting, the damaged trees were not removed in the main carbon accounting of the model. This can cause some underestimation of the heterotrophic respiration, but it was considered as insignificant for the present study which only focuses on SBB outbreak dynamics.

The bark beetle accounting and application of damage is placed in the “mortality_guess” function within the vegdynam.cpp code together with the wind damage application, and it is called at the end of each simulated year (Lagergren et al., 2024a).

2.4. Data of storm and bark beetle damage to forest in Europe

Data of damaged volume of spruce forest were combined with statistics of standing spruce volume to assess the fractions in a country/region damaged by storm (DF_{storm}) or killed by SBB (DF_{SBB}), which were calculated and used for further analysis.

European storm and bark beetle damage statistics were compiled by Marini et al. (2017), for some countries separated into administrative or topographical units. From that data set we used damage statistics ($m^3 yr^{-1}$) from South Sweden (data separated into 10 counties), North-East France (five former administrative units), Switzerland (lowland and mountains) and Austria (whole country), to cover regions with large interannual variability in ASP. As storms in Europe mainly occur in autumn and winter, and as the vegetation and bark beetle effect, e.g., will be the same for a storm event in October or in February the next year, and the amount of damage caused by the storm is much more important than its timing during this period when considering its impact on bark beetle dynamics during the following growing season, the storm damage statistics for a specific year were compiled for a storm season of 12 months from July until June the next year when building the dataset used for the calibration.

Data of spruce standing volume were available by year (1961-2010) for Sweden (<https://www.slu.se/en/Collaborative-Centres-and-Projects/the-swedish-national-forest-inventory/long-time-series/time-series-from-1953/>), for year 2008 for France (<https://inventaire-forestier.ign.fr/spip.php?rubrique250>), for year 1985, 1995 and 2006 for Switzerland (<https://www.lfi.ch/publikationen/publikation-en.php>) and for 2008 in Austria (<https://www.bfw.gv.at/en/departments-en/forest-inventory/>). For the other countries than Sweden with data only for one or three years, the data were interpolated between the inventory years and kept constant before the first and after the last year.

In recent years Europe has faced several SBB outbreaks that have been driven by warm and dry conditions rather than triggered by storm events (Nardi et al., 2023; Trubin et al., 2022). To test if this new situation in driving

factors was important for the model parameterization, national level storm and bark beetle damage statistics from the Standardized Disturbance Index (SDI) dataset (Patacca et al., 2022) in years 2011-2019 for Switzerland and Austria were used. For Switzerland the damage was split between Lowland and Mountain assuming that the proportion of the country totals for Lowland and Mountain parts were the same as in the 1990-2010 data described above.

Table 1: Summary of available storm and SBB damage data from Marini et al. (2017), data used in the primary calibration and additional data from SDI (Patacca et al., 2022) used in the validation as well as the number of simulated climate grid-cells in the different parts.

Country or part of country	Regions	Gridcells per region	Primary calibration years (Marini et al., 2017)	Additional calibration years (Patacca et al., 2022)
South Sweden	10	3-9	1990-2010	
Switzerland	2	8-9	1990-2010	2011-2019
Austria	1	39	1990-2010	2011-2019
East France	5	4-17	2000-2010	

In order to focus on bark beetle outbreak dynamics without introducing additional uncertainties associated with wind data and wind damage modelling, we prescribed wind damage from observed data. To still take advantage of the wind module's capacity to distribute wind damage among patches and cohorts' sensitivity (Eq. 1), a calibration was done to adjust WL so that modelled damage followed DF_{storm} (denoted WL_{stat}). As a common linear scaling was used to go from DF_{storm} to WL_{stat} , the exact DF_{storm} time series will not be reproduced by the model. In a first step, a factor of 2 was found to approximately generate the same average level of WL_{stat} calculated from DF_{storm} as WL calculated from wind (see section 2.5 below) data for years 1990-2010.

$$WL_{stat} = DF_{storm} \times 2 \quad (13)$$

After evaluating the quotient between preliminary LPJ-GUESS simulation results and inventoried DF_{storm} at regional level for all available years (Table 1), we concluded to use a separate function for northern Europe (Sweden) depending also on latitude (LAT):

$$WL_{stat} = DF_{storm} \times 2/f(LAT) \quad (14)$$

$$f(LAT) = -0.00412 \times LAT^2 + 0.425 \times LAT - 9.94, \quad (15)$$

where Eq. (15) was fitted to DF_{storm} quotients in S Sweden 1965-2010. The scaling with LAT, as a proxy for productivity (LAT explains 82% of the variation in county average site quality class in Southern Sweden and 92% for all counties in Sweden (Tab. 3.11a in Roberge et al., 2023), is reasonable as a higher WL is needed to trigger a certain level of damage when the cohorts have a lower SI (as they have with lower productivity, see Eq. 1).

2.5. Climate data and area delimitation




Figure 2: The climate gridcells simulated for the ten counties in South Sweden, the five counties in North-East France, the two larger regions in Switzerland and in Austria.

The simulations were driven by daily weather data 1901-2019 at 0.5° resolution from the CRU-JRA V2.1 dataset (<https://catalogue.ceda.ac.uk/uuid/10d2c73e5a7d46f4ada08b0a26302ef7>). For nitrogen deposition, monthly data from Lamarque et al. (2010) were used. The model was run with forest vegetation for all grid cells within each region/country (Fig. 2). For the years with no wind damage statistics, WL was calculated from CRU-JRA windspeed from the cubed exceedance of the 99.5 percentile of daily wind speed accumulated over storm season as in Lagergren et al. (2012). Wind damage was applied from 1951 and forward in the simulations.

2.6. Calibration and sensitivity test

For calibrating and testing the parameter sensitivity of the model, data for 1990-2010 from the 10 southern counties in South Sweden (excluding the Gotland island, which has a low fraction of spruce forest and non-typical soils), 5 counties in North-East France (only data for 1999-2010), Austria (whole country) and the lowland and mountain regions of Switzerland were used. For South Sweden, North-East France and Switzerland the modelling results and calibration data of DF_{SBB} were averaged over the counties/regions for use in the calibration and sensitivity test. The parameters were first adjusted based on expert judgement, giving a default set of parameters (Table 2).

Table 2: Parameters in the bark beetle model and reference to the equation they were used in. Min and max value shown for parameters used in the calibration and sensitivity analysis.

Parameter	Part of model	Default	Min	Max	Eq.
k_{gc_min}	$f(P_{gridcell})$	-2.0	-2.5	-1.0	7
k_{gc_max}	$f(P_{gridcell})$	1.0			7
k_1	$f(P_{gridcell})$	0.05	0.005	0.5	7

k_2	$f(P_{gridcell})$	0.25	0.15	0.5	7
k_{base_bm}	$f(P_{patch} / L)$	0.0001			9
k_{p_min}	$f(P_{patch} / L)$	-1.8	-2.8	-1.3	8
k_{p_max}	$f(P_{patch} / L)$	1.0			8
k_3	$f(P_{patch} / L)$	0.03	0.005	0.15	8
k_4	$f(P_{patch} / L)$	0.4	0.25	0.8	8
k_{pyw}	$f(\text{water stress})$	3	0.25	4	11
$k_{drought_max}$	$f(\text{water stress})$	2.0			10
k_5	$f(\text{water stress})$	8	5	27	10
k_6	$f(\text{phenology})$	3.33			12
k_7	$f(\text{phenology})$	75			12

Eight of the 14 parameters were then selected for a calibration procedure and a sensitivity test. These selected parameters were mainly related to the shape of the functions; other parameters were not included as we wanted to keep the range of the response functions at approximately the same magnitude as the weight of the original Marini et al. (2017) model (Eq. 3, see section 2.3). For this reason, the max parameters were excluded from the calibration. As previously stated, the k_{base_bm} parameter was set to a low value to avoid division by zero. To have a linear response within the wanted range (Fig 1a) of the $f(\text{phenology})$ function for the space of ASP (Fig. 1e) there was little room to adjust the parameters of Eq. 12 and they were therefore set fixed. To further reduce the number of calculations and to keep the total weight of the population size dependency constant, the sum of the minimum of the ranges $f(P_{gridcell})$ and $f(P_{patch} / L)$ was kept constant ($k_{gc_min} + k_{p_min} = -3.8$). This meant that the number of tested parameters could be reduced to 7 as k_{p_min} could be replaced with $-3.8 - k_{gc_min}$. For each tested parameter 7 discrete numbers were used, evenly spread between the ranges in Table 2. LPJ-GUESS was first run with the default parameter setting. Output at cohort level of all variables used in the bark beetle module was produced for this simulation. These data were then applied in a stand-alone version of the bark beetle module implemented in Matlab (R2018b) for all 7^7 parameter combinations. This approach misses the feedback from bark-beetle mortality to the vegetation state that is simulated within LPJ-GUESS, but greatly reduces the calculation time. Each parameter combination was first run with the default k_0 , k_0 was then iteratively adjusted until the mean quotient between simulated and observed maximum DF_{SBB} for the calibration period over the four regions/countries equalled 1. The results were sorted by R^2 , root mean square error (RMSE) and absolute bias both at country level (arithmetic mean over the 1-10 regions, Table 1) and by arithmetic mean over the four countries' mean values, and the top ranked parameter combination was selected based on highest R^2 , lowest RMSE and lowest absolute Bias. A combined statistic measure (CSM) was also calculated for all 7^7 models (X) by summing normalized $1 - R^2$, RMSE and absolute Bias as:

$$CSM_X = \frac{(1-R_X^2)-(1-R_{mean}^2)}{(1-R_{max}^2)-(1-R_{min}^2)} + \frac{RMSE_X - RMSE_{mean}}{RMSE_{max} - RMSE_{min}} + \frac{|Bias_X| - |Bias|_{mean}}{|Bias|_{max} - |Bias|_{min}}, \quad (15)$$

where the normalization is based on deviation from mean relative to the range (max-min). The CSM_X values were sorted in ascendant order and we present parameters and statistics for the highest rated model as well as the range of parameters for the 50 highest rated models.

LPJ-GUESS was run both with and without SSC, then the parameter testing and calibration procedure was repeated both with and without inclusion of the 2011-2019 calibration data for Switzerland and Austria. As a final test of the impact of SSC for the results, the stand-alone implementation was also run with inclusion of SSC for the parameter set obtained without SSC and vice versa. The setting including SSC using only the 1990-2010 data was considered as the main base run, results from the other runs are in most cases presented in the Supplement.

2.7. Robustness test and exploration of the climate change signal

To test the robustness of the approach to calibrate the model for different parameter combinations with structure and L_{mort} prescribed from an LPJ-GUESS simulation with default parameters, LPJ-GUESS was finally run with the optimized parameter set, which generated the right corresponding feedback of the damage associated with that setting to the simulated vegetation. A simple test of climate sensitivity was also done for this setting by applying +2 °C to the climate data throughout the simulation.

3. Results

3.1. Model optimization

The top ranked set of model parameters differed depending on the country or region assessed (Table 3), the calibration period included and whether SSC was included or not (Table S1). Most of the parameters of the model common for all four regions in the main base-run optimization (with SSC and not including calibration data for Switzerland and Austria 2011-2019) had a large span within the 50 highest ranked models (Table 3) but only one parameter value or a narrow span was dominant (Fig. S1). It should be noted that all calibrations were based on data from all countries, then the optimum model among the 7^7 -parameter space for the regions countries, or all together, was selected, which explains why there is a difference for S Sweden and NE France when including calibration data 2011-2019 for Austria and Switzerland (Table 3 vs Table S1a and Table S1b vs Table S1c).

Table 3: Parameters in the main base run setting (with SSC and not including calibration data for Switzerland and Austria 2011-2019), for the top-ranked model in terms of combined statistics of bias, RMSE and R^2 for all four regions/countries together and for each region/country separately. For all four regions/countries together, also the parameter range for the 50 highest ranked combinations is shown (min_50, max_50). The k_0 values are the results of the calibration, LPJ-GUESS was run with a k_0 of 0.003. The numbers in parenthesis is the order number (from smallest to largest) of the seven values tested within the full parameter range (Table 2).

	k_1	k_2	k_{gc_min}	k_{pyw}	k_3	k_4	k_5	$k_0 \times 1000$
Default	0.05 (4)	0.25 (3)	-2 (3)	3 (6)	0.03 (4)	0.4 (4)	8 (3)	2.89
All four	0.005 (1)	0.5 (7)	-1.75 (4)	0.25 (1)	0.15 (7)	0.8 (7)	5 (1)	5.12
min_50	0.005 (1)	0.15 (1)	-1.75 (4)	0.25 (1)	0.005 (1)	0.5 (5)	5 (1)	
max_50	0.5 (7)	0.5 (7)	-1 (7)	4 (7)	0.15 (7)	0.8 (7)	20 (6)	
S Sweden	0.5 (7)	0.15 (1)	-2 (3)	0.5 (3)	0.009 (2)	0.8 (7)	27 (7)	0.0989
Switzerland	0.005 (1)	0.15 (1)	-2.25 (2)	1 (4)	0.005 (1)	0.25 (1)	6 (2)	0.192
Austria	0.12 (5)	0.15 (1)	-2.5 (1)	1 (4)	0.15 (7)	0.5 (5)	27 (7)	0.00272
NE France	0.005 (1)	0.35 (4)	-2.5 (1)	4 (7)	0.15 (7)	0.4 (4)	20 (6)	0.0428

3.2. Model performance

The optimization procedure resulted in reduced bias and RMSE and increased R^2 compared to the default setting (Table 4, Fig. 3, Table S2, Fig. S2-4).

Table 4: Statistics for different parameter settings; default (Table 2), the top-ranked model in terms of combined statistics of bias, RMSE and R^2 for all four regions/countries together and for each region/country separately, in the main base run.

	Default			Combined all four			Region/country		
	R^2	RMSE	Bias	R^2	RMSE	Bias	R^2	RMSE	Bias
All four	0.30	0.29%	0.10%	0.40	0.22%	0.04%			
S Sweden	0.54	0.28%	0.21%	0.76	0.12%	0.07%	0.72	0.08%	0.01%
Switzerland	0.34	0.31%	-0.02%	0.43	0.38%	0.07%	0.45	0.34%	0.01%
Austria	0.03	0.20%	0.10%	0.19	0.18%	0.08%	0.26	0.15%	0.07%
NE France	0.29	0.59%	-0.30%	0.21	0.64%	-0.28%	0.44	0.60%	-0.08%

The model captured the outbreak dynamics well for S Sweden. In Switzerland R^2 was rather low, mainly because the peak in damage after the 1999 storm in 2001 was not captured. In North-East France there was a large spread in the observed outbreak after the 1999 storm, with Alsace and Lorraine having a large peak in 2001, while the other counties had a more slowly evolving progress. In Austria the outbreak starting in 2003 after the 2002 storm, with significant storm damage also in 2008 and 2008, lasted almost a decade though the response from the functions of phenology and drought did not indicate that these factors supported the outbreak, resulting in low R^2 . Including years 2011-2019 data from Austria and Switzerland in the calibration resulted in similar outbreak pattern (Fig. 3, Fig. S2).




Figure 3: Observed and modelled fraction of spruce forest damaged by storm (left y-axis) and SBB (right y-axis) in four regions/countries, with modelled SBB damage from different parameter settings (Table 3) in the main base run. For Sweden ($n = 10$), Switzerland ($n = 2$) and France ($n = 5$) $\pm$ standard deviation between regions in observed SBB damage is shown with dotted lines.

410





Figure 4: Accumulated fraction of simulated and observed spruce forest damaged by SBB over the period with observations. Calibrations are shown using only data up until 2010 (10) and also including data for Switzerland and Austria 2011-2019 (19) as well as with (W) or without (WO) salvage and sanitary cutting. The main base run setting, cW10-rW, is showed by bold line.

Accumulating the SBB damage over time show that level of damage during outbreak situations was generally well captured for all four region/countries (Fig. 4). The difference depending on setting for the calibration was quite large, and in the main base run the total damage for SE France was underestimated. During non-outbreak

420 situation in Sweden the accumulated damage is higher than observed while it agrees better in Austria and Switzerland but this may to a large extent depend on the way damage is reported, as discussed in Sect. 4.

3.3. Model sensitivity



425 **Figure 5:** Test of the sensitivity of including salvage cutting of storm felled trees and sanitary cutting of infected trees (SSC) for the fraction of SBB killed trees. The calibrations with (cW) and without (cWO) SSC using calibration data for 1990-2010 (10) were run both with (rW) and without (rWO) SSC. In Fig. S5, the calibrations also including 2011-2019 data for Austria and Switzerland are shown.

Including the 2011-2019 data for Austria and Switzerland in the calibration resulted in higher accumulated SBB damage in Austria and NE France but lower level in Switzerland with SSC (Fig. 4). Without SSC it resulted in substantially higher accumulated damage in Switzerland and NE France as the outbreaks continued at a high level for more years (Fig. S3-4).




Figure 6: Sensitivity test of modelled storm and spruce bark beetle damage for temperature and model environment. Model output from the calibrated stand-alone version with tree structure and storm damage input from the default LPJ-GUESS run (def, “All4 opt” in Fig. 3) and the calibrated model run with LPJ-GUESS with normal climate (SBB sim) and with +2 ° temperature (sim+2) compared to observations (obs).

Applying the parameter set obtained with SSC without SSC resulted in 1.7 (Austria) to seven (Sweden) times higher simulated damage levels and the calibration without SSC run with SSC showed reductions of similar magnitude (Fig. 5, Fig. S5). It should though be noted that the negative feedback on forest vulnerability when trees are killed is not included in these stand-alone implementations, giving a higher background damage level in the run without SSC.

Up to this point all results are for the stand-alone implementation based on the vegetation from the default LPJ-GUESS run, i.e. with feedback from SBB damage from the default run instead of from the stand-alone. Applying the calibrated parameter set of the main base run in LPJ-GUESS showed a very good match for S Sweden, a minor underestimation in NE France and underestimations by 35% and 70% in Switzerland and Austria, respectively (Fig. 6). An in-depth analysis for Austria showed that the difference was due to one grid cell where the missing negative feedback from spruce killing resulted in accumulated modelled damage that widely exceeded the availability of spruce trees in the stand-alone version. Removing this grid cell gave similarly low levels as for the LPJ-GUESS run with calibrated parameters, showing that the applied stand-alone optimization methodology resulted in an underestimation in this case. It can, therefore, be concluded that this calibration process resulted in a calibrated model with conservative damage estimates.

The simple climate sensitivity test of increasing the temperature by 2 °C resulted in an expected increase in SBB damage in S Sweden and Austria (Fig. 6). In Switzerland and NE France, the warming resulted in significantly reduced biomass of the boreal BNE PFT and consequently a reduction in SBB damage as this PFT was close to its environmental limits and outside Norway spruce' native distribution in these regions (Caudullo et al., 2016), already in the present climate.

4. Discussion

The course of a of an insect outbreak can be boiled down to predisposing, triggering and contributing factors (Saxe, 1993). A summary of these factors by country are shown in Fig. 7, and will be the base for our discussion.

In a remote-sensing based modelling study, tree height, soil moisture and nearby clear-cuts were together with high spruce volume identified as the most important factors predisposing forests to SBB damage (Müller et al., 2022). In this study, damage was assessed for the spruce fraction of the forest simulated for stands planted with the BNE PFT. The predisposition is, therefore, mainly determined by the amount of spruce stem volume with a diameter larger than 15 cm (d_{lim}), which depends on tree allometry, age-class distribution, growth rate and tree density. The diameter limit has no direct physiological relationship to SBB preference but is related to bark thickness, which is directly related to the possibility to breed (Schlyter and Anderbrant, 1993), with more bark beetle offspring per unit bark area in larger trees (Weslien and Regnander, 1990). In thinner trees there may be parts of the stem that has thick enough bark for successful breeding, but the beetles need a dense cover of beetle larvae galleries over a major part of the stem to overcome the resin defence. In larger trees the bark can be too thick at the base and the beetles attack higher up on the tree where a lower water potential in the stem (Tyree and Ewers, 1991) also contributes to the success of the attack. The tree density (trees per ha) determines how quickly a tree reaches d_{lim} , as reduced density means increased diameter growth, and, in turn, depends on plant number, thinning and mortality. Age- and competition-related mortality also contributes to the background endemic SBB

population, which is important for the level of damage when an outbreak is triggered. In this study, the size of age classes was determined for one occasion at country or county scale, which is a simplification. Likewise, the availability of spruce trees with $d > d_{\text{lim}}$ in a grid cell is based on age-class data for the year 2010. Currently available large-scale datasets do not intersect stand age and species composition, which induces a big uncertainty when attempting to model outbreaks that are strongly dependent on the availability of trees of a particular species and size. A step forward in this respect could be to base the current stage on national forest inventories using plot level information on individual trees and use harvest and damage statistics for damage, thinning and clear-cut levels, which then in turn determine the tree composition and size distribution (Pugh et al., Manuscript).




Figure 7: Sensitivity for SBB outbreak divided in predisposing (biomass of spruce trees with a diameter larger than 15 cm), triggering (fraction of spruce volume damaged by storm) and contributing (the sum of the phenology and water stress related components for the increase rate of the bark beetle population index, R Eq. 4) factors. For the phenology and water stress R components sum the running 3-year mean and the trend over time (dashed line) are also shown.

Bark beetles commonly develop epidemic levels during the growing season following a storm event ramping up the populations in the defenceless damaged trees. This then leads to killing of trees the second season after the storm. The timing of such outbreaks is quite well captured by the presented model. In managed forest, however, the occurrence and timing of counter measures such as salvage logging of storm-felled trees, sanitary cutting of

495 infected trees and insect traps have a very big impact on the outcome (Jönsson et al., 2012), factors that vary greatly in time and by region over Europe, as well as at finer scale (Wichmann and Ravn, 2001). By timely salvage logging, a large fraction of the beetles can be removed from the forest, substantially dampening the damage done. European-wide models calibrated over a longer time span (e.g. Marini et al., 2017) should not therefore be expected to have a high explanation of particular events in comparison to more local model applications (e.g. 500 Soukhovolsky et al., 2022). However, they should be able to capture the general characteristics of these events well. One possible route forward to improved evaluation of large-scale models is assembling local-scale datasets that combine high-resolution information on species composition and forest structure, with detailed information on bark beetle detection and SSC. With on-going developments in remote sensing of forest structure and composition combined with datasets on forest harvesting, this may be increasingly feasible in the near future (e.g. 505 Jamali et al., 2024).

After a major storm, the remaining trees may have disturbed root systems making them more vulnerable to drought which may enhance SBB outbreaks, but also making them unstable – predisposing them to further storm damage. As SBB prefers defenceless trees even in an outbreak situation, such wind damage can serve as traps for the beetles if the wood is salvaged in time. This situation occurred in Southern Sweden when the 2006-2007 storm 510 dampened the outbreak following the major 2004-2005 storm. This relatively subtle mechanism is not captured in our model, which overestimated SBB damage in this outbreak. The outcome after a trigger depends on the initial level of the SBB population. Many SBB models require an initial population or damage level, which means that they work in relative terms (Marini et al., 2017; Soukhovolsky et al., 2022), but a dynamic forest model needs to operate with absolute damage levels making the modelling more challenging.

515 The way damage is inventoried and reported may also be different between countries and over time. For example, the neighbouring countries Austria and Switzerland have very different reported patterns, in which Austria has a relatively high damage level for almost all years for both wind and SBB damage but relatively low peaks, while Switzerland has large storm damage followed by large SBB damage after the 1999 storm, but low or absent storm damage except from that. The periods of zero bark beetle damage recorded indicate periods with low monitoring activity caused by low bark beetle activity. E.g., the country level compilation and gap filling of the DFDE 520 database (Patacca et al., 2022) show seven times higher expert gap filled SBB damage than reported 1990-2005 for Sweden, and in France the machine learning gap filling resulted in 2.4 times higher damage 1991-2000.

Warm weather accompanied with drought are often seen as a contributing factor (e.g. Bakke, 1983), but in recent year it has also triggered as well as contributed to sustained outbreaks of SBB in Europe (Nardi et al., 2023; Trubin 525 et al., 2022). There was no county-level data of SBB damage available for Sweden and France later than 2010, but country totals (Patacca et al., 2022) show levels that greatly exceed our small peak in modelled damage during 2018-2019. Drought has generally been seen as the main driver of these events (George et al., 2022; Kärvelo et al., 2023). With the data used in our calibration and the additive form of the response function we might have underestimated this response, and for Austria and Switzerland the model also underestimated the drought-induced 530 peak even when data for those years were included. Part of the reason for underestimating the damage could be that LPJ-GUESS may be failing to simulate sufficiently increased water stress in the vegetation during this time period. For instance, in Austria and Switzerland, there were no abnormally low values of the water stress scaler wscal (indicating water stress) post 2015. This may be a failing of the model parameterization for water stress or

of the input climate forcing dataset (Steinkamp and Hickler, 2015). Also for the most recent application of SBB damage with the ORCHIDEE vegetation model, a shortcoming when it comes to reflect last years' damage levels associated with extreme drought has been concluded (Marie et al., 2024). We note, however, that the 2018 drought generates a fairly strong water stress response in the simulations for southern Sweden (Fig. 7) and this is reflected in some increase in beetle damage in the years 2018 and 2019 (Fig. 3). An additional contributing factor that not has been explicitly accounted for in the present study is root rot, which can significantly increase the severity of an outbreak (Honkaniemi et al., 2018).

The simple test with a 2° C higher temperature gave an expected increase in modelled damage in S Sweden and Austria. Similar tests resulting in a strong increase in modelled SBB damage has also been done by (Jönsson et al., 2012) and (Seidl and Rammer, 2017). In Switzerland and NE France, however, a reduction was simulated, mainly related to a change of the predisposing spruce biomass $>d_{lim}$ as simulated growth of the boreal-parameterized BNE PFT was reduced in the warmer climate. The SBB phenology response, which is a function of the length of the first generation's swarming period, also reaches a plateau when the climate is warm enough to allow a complete second generation to emerge every year. A third generation can be completed with very hot conditions, but the consequences for SBB population dynamics depends on host tree availability.

Whilst predicting absolute damage levels is challenging, our model is generally effective at indicating when there are elevated periods of damage risk due to SBB (Fig. 7). The most obvious case is year 2003 with a strong drought (Granier et al., 2007) that contributed to prolonged outbreaks in Switzerland, Austria and NE France, but also in the recent years (2017-2019) the indicators also show increased risks coinciding with observed damage for all assessed countries (Patacca et al., 2022). Combined with the process detail related to forest structure and management, this means that the model can be a powerful tool to explore how different forest structures and climate and management scenarios might interact to shift forests towards increased or decreased vulnerability.

The model that we have developed here is parameterised for SBB. However, many different species of bark beetles that have the potential to cause large outbreaks and tree mortality, meaning that effectively accounting for the impact of these species on large-scale forest dynamics requires that we can develop methods to generalise responses across species to some degree. These other species of bark beetles can be specific in temperature sums for evolving and dormancy periods and for density dependent defence overcome and competition (Bentz et al., 2019). A common pattern for many species is increased risk for outbreaks related to drought (Reed and Hood, 2021), which is positive for a general concept of bark beetle damage modelling. The implementation of the SBB outbreak dynamics in LPJ-GUESS has introduced a concept of insect functional types (IFTs), which can be modelled in parallel with different response functions and specific PFT hosts. With this concept the model could also be applied in other parts of the world, if sufficient data on bark beetle phenology and damage exist to parameterise and calibrate the model.

6. Conclusions

Nature and human interactions drive and control SBB outbreaks in the intensively managed forest in Europe, which makes modelling challenging. The modelling concept we present here was able to catch the main timing and duration of observed damage but there was a substantial spread in absolute agreement, as also reflected in

uncertainty in parameter estimates. With more detailed information of the human factors, mainly in form of salvage logging and sanitary cutting, it should be possible to improve the explanatory power of the model. The SBB module is sensitive to the climate change signal, though the magnitude of temperature driven SBB damage seen in recent years was underestimated. In the modelling framework of the powerful LPJ-GUESS dynamical vegetation model it can, therefore, be a useful tool for exploring the SBB vulnerability of future climate and management scenarios.

Code availability

The underlying LPJ-GUESS code subversion revision 13130 and simulation settings are archived at Zenodo (Lagergren et al., 2024a). Analysis code is available via Github ([https://github.com/LPJ-GUESS/spruce bark beetle](https://github.com/LPJ-GUESS/spruce_bark_beetle); doi on acceptance).

Data availability

Model simulations underlying the results herein are archived at Zenodo (Lagergren et al., 2024b).

Author contribution

TP initiated the study. FL developed the SBB damage modelling concept with interaction from AMJ and TP. ML invented and implemented the insect functional type concept and coded the phenology of SBB into LPJ-GUESS. FL conducted all simulations and analysis of the results with feedback from all authors. FL lead the writing of the manuscript with contribution from all authors.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

The model developments described in this paper have been funded under the European Union's Horizon Europe research and innovation programme (grant agreement numbers 101059888, CLIMB-Forest; 101056755, ForestPaths; 101084481) and the Horizon 2020 programme (grant agreement no. 758873, TreeMort) as well as from the ForestValue programme, the European Commission, Vinnova, the Swedish Energy Agency and Formas for the project FORECO. This study is a contribution to the Swedish government's strategic research areas BECC and MERGE and the Nature-based Future Solutions profile area at Lund University. We thank Arjan Meddens, Rupert Seidl, Nikica Ogris and Cornelius Senf for discussions on the early stages of this work and Markus Kautz for providing phenology data.

References

- 600 Bakke, A.: Host tree and bark beetle interactions during a mass outbreak of *Ips typographus* in Norway, Journal of Applied Entomology, 96, 118-125, <https://doi.org/10.1111/j.1439-0418.1983.tb03651.x>, 1983.
- Balla, A., Silini, A., Cherif-Silini, H., Bouket, A. C., Moser, W. K., Nowakowska, J. A., Oszako, T., Benia, F., and Belbahri, L.: The Threat of Pests and Pathogens and the Potential for Biological Control in Forest Ecosystems, Forests, 12, <https://doi.org/10.3390/f12111579>, 2021.
- 605 Bentz, B. J., Jönsson, A. M., Schroeder, M., Weed, A., Wilcke, R. A. I., and Larsson, K.: *Ips typographus* and *Dendroctonus ponderosae* Models Project Thermal Suitability for Intra- and Inter-Continental Establishment in a Changing Climate, Frontiers in Forests and Global Change, 2, <https://doi.org/10.3389/ffgc.2019.00001>, 2019.
- Caudullo, G., Tinner, W., and de Rigo, D.: *Picea abies* in Europe: distribution, habitat, usage and threats, in: European Atlas of Forest Tree Species, edited by: San-Miguel-Ayaz, J., de Rigo, D., Caudullo, G., Houston
- 610 Durrant, T., and Mauri, A., Publ. Off. EU, Luxembourg, e012300+, 2016.
- Ciais, P., Schelhaas, M. J., Zaehle, S., Piao, S. L., Cescatti, A., Liski, J., Luyssaert, S., Le-Maire, G., Schulze, E. D., Bouriaud, O., Freibauer, A., Valentini, R., and Nabuurs, G. J.: Carbon accumulation in European forests, Nature Geoscience, 1, 425-429, <https://doi.org/10.1038/ngeo233>, 2008.
- de Bruijn, A., Gustafson, E. J., Sturtevant, B. R., Foster, J. R., Miranda, B. R., Lichti, N. I., and Jacobs, D. F.:
- 615 Toward more robust projections of forest landscape dynamics under novel environmental conditions: Embedding PnET within LANDIS-II, Ecological Modelling, 287, 44-57, <https://doi.org/10.1016/j.ecolmodel.2014.05.004>, 2014.
- Forzieri, G., Girardello, M., Ceccherini, G., Spinoni, J., Feyen, L., Hartmann, H., Beck, P. S. A., Camps-Valls, G., Chirici, G., Mauri, A., and Cescatti, A.: Emergent vulnerability to climate-driven disturbances in European
- 620 forests, Nature Communications, 12, <https://doi.org/10.1038/s41467-021-21399-7>, 2021.
- George, J. P., Bürkner, P. C., Sanders, T. G. M., Neumann, M., Cammalleri, C., Vogt, J. V., and Lang, M.: Long-term forest monitoring reveals constant mortality rise in European forests, Plant Biology, 24, 1108-1119, <https://doi.org/10.1111/plb.13469>, 2022.
- Granier, A., Reichstein, M., Bréda, N., Janssens, I. A., Falge, E., Ciais, P., Grünwald, T., Aubinet, M.,
- 625 Berbigier, P., Bernhofer, C., Buchmann, N., Facini, O., Grassi, G., Heinesch, B., Ilvesniemi, H., Keronen, P., Knohl, A., Köstner, B., Lagergren, F., Lindroth, A., Longdoz, B., Loustau, D., Mateus, J., Montagnani, L., Nys, C., Moors, E., Papale, D., Peiffer, M., Pilegaard, K., Pita, G., Pumpanen, P., Rambal, S., Rebmann, C., Rodrigues, A., Seufert, G., Tenhunen, J., Vesala, T., and Wang, Q.: Evidence for soil water control on carbon and water dynamics in European forests during the extremely dry year: 2003, Agricultural and Forest
- 630 Meteorology, 143, 123-145, <https://doi.org/10.1016/j.agrformet.2006.12.004>, 2007.
- Hlásny, T., König, L., Krokene, P., Lindner, M., Montagné-Huck, C., Müller, J., Qin, H., Raffa, K. F., Schelhaas, M. J., Svoboda, M., Viiri, H., and Seidl, R.: Bark Beetle Outbreaks in Europe: State of Knowledge and Ways Forward for Management, Current Forestry Reports, 7, 138-165, <https://doi.org/10.1007/s40725-021-00142-x>, 2021.

- 635 Honkaniemi, J., Ojansuu, R., Kasanen, R., and Heliövaara, K.: Interaction of disturbance agents on Norway spruce: A mechanistic model of bark beetle dynamics integrated in simulation framework WINDROT, Ecological Modelling, 388, 45-60, <https://doi.org/10.1016/j.ecolmodel.2018.09.014>, 2018.
- Jamali, S., Olsson, P. O., Müller, M., and Ghorbanian, A.: Kernel-Based Early Detection of Forest Bark Beetle Attack Using Vegetation Indices Time Series of Sentinel-2, Ieee Journal of Selected Topics in Applied Earth
640 Observations and Remote Sensing, 17, 12868-12877, <https://doi.org/10.1109/jstars.2024.3425795>, 2024.
- Jönsson, A. M., Harding, S., Bärning, L., and Ravn, H. P.: Impact of climate change on the population dynamics of *Ips typographus* in southern Sweden, Agricultural and Forest Meteorology, 146, 70-81, <https://doi.org/10.1016/j.agrformet.2007.05.006>, 2007.
- Jönsson, A. M., Schroeder, L. M., Lagergren, F., Anderbrant, O., and Smith, B.: Guess the impact of *Ips*
645 *typographus* - An ecosystem modelling approach for simulating spruce bark beetle outbreaks, Agricultural and Forest Meteorology, 166-167, 188-200, <https://doi.org/10.1016/j.agrformet.2012.07.012>, 2012.
- Jönsson, A. M., Harding, S., Krokene, P., Lange, H., Lindelöw, Å., Okland, B., Ravn, H. P., and Schroeder, L. M.: Modelling the potential impact of global warming on *Ips typographus* voltinism and reproductive diapause, Climatic Change, 109, 695-718, <https://doi.org/10.1007/s10584-011-0038-4>, 2011.
- 650 Kärvmö, S., Huo, L., Öhrn, P., Lindberg, E., and Persson, H. J.: Different triggers, different stories: Bark-beetle infestation patterns after storm and drought-induced outbreaks, Forest Ecology and Management, 545, <https://doi.org/10.1016/j.foreco.2023.121255>, 2023.
- Kautz, M., Meddens, A. J. H., Hall, R. J., and Arneeth, A.: Biotic disturbances in Northern Hemisphere forests - a synthesis of recent data, uncertainties and implications for forest monitoring and modelling, Global Ecology and
655 Biogeography, 26, 533-552, <https://doi.org/10.1111/geb.12558>, 2017.
- Lagergren, F., Jönsson, A. M., Blennow, K., and Smith, B.: Implementing storm damage in a dynamic vegetation model for regional applications in Sweden, Ecological Modelling, 247, 71-82, <https://doi.org/10.1016/j.ecolmodel.2012.08.011>, 2012.
- Lagergren, F., Jönsson, A. M., Lindeskog, M., and Pugh, T. A. M.: Code and simulation setting underlying:
660 Combining empirical and mechanistic understanding of spruce bark beetle outbreak dynamics in the LPJ-GUESS (v4.1, r13130) vegetation model (Version 1), Zenodo, <https://doi.org/10.5281/zenodo.14411974>, 2024a.
- Lagergren, F., Jönsson, A. M., Lindeskog, M., and Pugh, T. A. M.: Simulation results underlying: Combining empirical and mechanistic understanding of spruce bark beetle outbreak dynamics in the LPJ-GUESS (v4.1, r13130) vegetation model (Version 1) [Data set], Zenodo, <https://doi.org/10.5281/zenodo.14415079>, 2024b.
- 665 Lamarque, J. F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Lioussé, C., Mieville, A., Owen, B., Schultz, M. G., Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne, J., Cooper, O. R., Kainuma, M., Mahowald, N., McConnell, J. R., Naik, V., Riahi, K., and van Vuuren, D. P.: Historical (1850-2000) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols: methodology and application, Atmospheric Chemistry and Physics, 10, 7017-7039, <https://doi.org/10.5194/acp-10-7017-2010>,
670 2010.

- Lind, P., Belušić, D., Christensen, O. B., Dobler, A., Kjellström, E., Landgren, O., Lindstedt, D., Matte, D., Pedersen, R. A., Toivonen, E., and Wang, F. X.: Benefits and added value of convection-permitting climate modeling over Fenno-Scandinavia, *Climate Dynamics*, 55, 1893-1912, <https://doi.org/10.1007/s00382-020-05359-3>, 2020.
- 675 Lindeskog, M., Smith, B., Lagergren, F., Sycheva, E., Ficko, A., Pretzsch, H., and Rammig, A.: Accounting for forest management in the estimation of forest carbon balance using the dynamic vegetation model LPJ-GUESS (v4.0, r9710): implementation and evaluation of simulations for Europe, *Geoscientific Model Development*, 14, 6071-6112, <https://doi.org/10.5194/gmd-14-6071-2021>, 2021.
- Lindgren, B. S. and Raffa, K. F.: Evolution of tree killing in bark beetles (Coleoptera: Curculionidae): trade-offs
680 between the maddening crowds and a sticky situation, *Canadian Entomologist*, 145, 471-495, <https://doi.org/10.4039/tce.2013.27>, 2013.
- Marie, G., Jeong, J., Jactel, H., Petter, G., Cailleret, M., McGrath, M. J., Bastrikov, V., Ghattas, J., Guenet, B., Lansø, A. S., Naudts, K., Valade, A., Yue, C., and Luyssaert, S.: Simulating *Ips typographus* L. outbreak dynamics and their influence on carbon balance estimates with ORCHIDEE r8627, *Geoscientific Model Development*, 17, 8023-8047, <https://doi.org/10.5194/gmd-17-8023-2024>, 2024.
- 685 Marini, L., Økland, B., Jönsson, A. M., Bentz, B., Carroll, A., Forster, B., Grégoire, J.-C., Hurling, R., Nageleisen, L. M., Netherer, S., Ravn, H. P., Weed, A., and Schroeder, M.: Climate drivers of bark beetle outbreak dynamics in Norway spruce forests, *Ecography*, 40, 1426-1435, <https://doi.org/10.1111/ecog.02769>, 2017.
- 690 Müller, M., Olsson, P.-O., Eklundh, L., Jamali, S., and Ardö, J.: Features predisposing forest to bark beetle outbreaks and their dynamics during drought, *Forest Ecology and Management*, 523, <https://doi.org/10.1016/j.foreco.2022.120480>, 2022.
- Nardi, D., Jactel, H., Pagot, E., Samalens, J. C., and Marini, L.: Drought and stand susceptibility to attacks by the European spruce bark beetle: A remote sensing approach, *Agricultural and Forest Entomology*, 25, 119-129,
695 <https://doi.org/10.1111/afe.12536>, 2023.
- Palahi, M., Valbuena, R., Senf, C., Acil, N., Pugh, T. A. M., Sadler, J., Seidl, R., Potapov, P., Gardiner, B., Hetemäki, L., Chirici, G., Francini, S., Hlásny, T., Lerink, B. J. W., Olsson, H., Olabarria, J. R. G., Ascoli, D., Asikainen, A., Bauhus, J., Berndes, G., Donis, J., Fridman, J., Hanewinkel, M., Jactel, H., Lindner, M., Marchetti, M., Marušák, R., Sheil, D., Tomé, M., Trasobares, A., Verkerk, P. J., Korhonen, M., and Nabuurs, G.
700 J.: Concerns about reported harvests in European forests, *Nature*, 592, E15-E17, <https://doi.org/10.1038/s41586-021-03292-x>, 2021.
- Patacca, M., Lindner, M., Lucas-Borja, M. E., Cordonnier, T., Fidej, G., Gardiner, B., Hauf, Y., Jasinevičius, G., Labonne, S., Linkevičius, E., Mahnken, M., Milanovic, S., Nabuurs, G. J., Nagel, T. A., Nikinmaa, L., Panyatov, M., Bercak, R., Seidl, R., Ostrogović Sever, M. Z., Socha, J., Thom, D., Vuletic, D., Zudin, S., and
705 Schelhaas, M. J.: Significant increase in natural disturbance impacts on European forests since 1950, *Global Change Biology*, <https://doi.org/10.1111/gcb.16531>, 2022.

- Poulter, B., Aragão, L., Andela, N., Bellassen, V., Ciais, P., Kato, T., Lin, X., Nachin, B., Luyssaert, S., Pederson, N., Peylin, P., Piao, S., Pugh, T., Saatchi, S., Schepaschenko, D., Schelhaas, M., and Shvidenko, A.: The global forest age dataset and its uncertainties (GFADv1.1), PANGAEA [dataset], <https://doi.org/10.1594/PANGAEA.897392>, 2019.
- Pugh, T. A. M., Arneth, A., Eckes-Shephard, A. H., Jönsson, A. M., Lagergren, F., Lindeskog, M., Miller, P. A., Nieradzik, L., Olin, S., Piltz, K., Rammig, A., Suvanto, S., Wittenbrink, M., Zhong, H., Astigarraga, J., Cienciala, E., Esquivel-Muelbert, A., Fridman, J., Lehtonen, A., Ruiz Benito, P. R., Peltoniemi, M., Schelhaas, M.-J., Senf, C., Viana-Soto, A., Stadelmann, G., Tagesson, T., Talarczyk, A., and Zavala, M. A.: Dynamic vegetation modelling for the past and future of Europe's forests, Manuscript.
- Reed, C. C. and Hood, S. M.: Few generalizable patterns of tree-level mortality during extreme drought and concurrent bark beetle outbreaks, *Science of the Total Environment*, 750, <https://doi.org/10.1016/j.scitotenv.2020.141306>, 2021.
- Roberge, C., Nilsson, P., Wikberg, P.-E., and Fridman, J.: Forest statistics 2023 - Official Statistics of Sweden, Swedish University of Agricultural Sciences, Department of Forest Resource Management, Umeå, 168 pp.2023.
- Saxe, H.: Triggering and predisposing factors in the "Red" decline syndrome of Norway spruce (*Picea abies*), *Trees-Structure and Function*, 8, 39-48, 1993.
- Scheel, M., Lindeskog, M., Smith, B., Suvanto, S., and Pugh, T. A. M.: Increased Central European forest mortality explained by higher harvest rates driven by enhanced productivity, *Environmental Research Letters*, 17, <https://doi.org/10.1088/1748-9326/ac9635>, 2022.
- Schlyter, F. and Anderbrant, O.: Competition and niche separation between 2 bark beetles - existence and mechanism, *Oikos*, 68, 437-447, <https://doi.org/10.2307/3544911>, 1993.
- Schroeder, L. M.: Tree Mortality by the Bark Beetle *Ips typographus* (L.) in storm-disturbed stands, *Integrated Pest Management Reviews*, 6, 169-175, <https://doi.org/10.1023/A:1025771318285>, 2001.
- Seidl, R. and Rammer, W.: Climate change amplifies the interactions between wind and bark beetle disturbances in forest landscapes, *Landscape Ecology*, 32, 1485-1498, <https://doi.org/10.1007/s10980-016-0396-4>, 2017.
- Seidl, R., Rammer, W., and Spies, T. A.: Disturbance legacies increase the resilience of forest ecosystem structure, composition, and functioning, *Ecological Applications*, 24, 2063-2077, <https://doi.org/10.1890/14-0255.1>, 2014.
- Smith, B., Prentice, I. C., and Sykes, M. T.: Representation of vegetation dynamics in the modelling of terrestrial ecosystems: comparing two contrasting approaches within European climate space, *Global Ecology and Biogeography*, 10, 621-637, <https://doi.org/10.1046/j.1466-822X.2001.t01-1-00256.x>, 2001.
- Smith, B., Wårlind, D., Arneth, A., Hickler, T., Leadley, P., Siltberg, J., and Zaehle, S.: Implications of incorporating N cycling and N limitations on primary production in an individual-based dynamic vegetation model, *Biogeosciences*, 11, 2027-2054, <https://doi.org/10.5194/bg-11-2027-2014>, 2014.
- Soukhovolsky, V., Kovalev, A., Tarasova, O., Modlinger, R., Křenová, Z., Mezei, P., Škvarenina, J., Rožnovský, J., Korolyova, N., Majdák, A., and Jakuš, R.: Wind Damage and Temperature Effect on Tree

- Mortality Caused by *Ips typographus* L.: Phase Transition Model, *Forests*, 13, <https://doi.org/10.3390/f13020180>, 2022.
- 745 Steinkamp, J. and Hickler, T.: Is drought-induced forest dieback globally increasing?, *Journal of Ecology*, 103, 31-43, <https://doi.org/10.1111/1365-2745.12335>, 2015.
- Temperli, C., Bugmann, H., and Elkin, C.: Cross-scale interactions among bark beetles, climate change, and wind disturbances: a landscape modeling approach, *Ecological Monographs*, 83, 383-402, <https://doi.org/10.1890/12-1503.1>, 2013.
- 750 Trubin, A., Mezei, P., Zabihi, K., Surový, P., and Jakuš, R.: Northernmost European spruce bark beetle *Ips typographus* outbreak: Modelling tree mortality using remote sensing and climate data, *Forest Ecology and Management*, 505, <https://doi.org/10.1016/j.foreco.2021.119829>, 2022.
- Tyree, M. T. and Ewers, F. W.: The hydraulic architecture of trees and other woody plants, *New Phytologist*, 119, 345-360, <https://doi.org/10.1111/j.1469-8137.1991.tb00035.x>, 1991.
- 755 Wermelinger, B. and Seifert, M.: Analysis of the temperature dependent development of the spruce bark beetle *Ips typographus* (L) (Col, Scolytidae), *Journal of Applied Entomology*, 122, 185-191, <https://doi.org/10.1111/j.1439-0418.1998.tb01482.x>, 1998.
- Wernick, I. K., Ciais, P., Fridman, J., Höglberg, P., Korhonen, K. T., Nordin, A., and Kauppi, P. E.: Quantifying forest change in the European Union, *Nature*, 592, E13-E14, <https://doi.org/10.1038/s41586-021-03293-w>, 2021.
- 760 Weslien, J. and Regnander, J.: Colonization densities and offspring production in the bark beetle *Ips typographus* (L.) in standing spruce trees, *Journal of Applied Entomology*, 109, 358-366, <https://doi.org/10.1111/j.1439-0418.1990.tb00064.x>, 1990.
- Wichmann, L. and Ravn, H. P.: The spread of *Ips typographus* (L.) (Coleoptera, Scolytidae) attacks following heavy windthrow in Denmark, analysed using GIS, *Forest Ecology and Management*, 148, 31-39, [https://doi.org/10.1016/s0378-1127\(00\)00477-1](https://doi.org/10.1016/s0378-1127(00)00477-1), 2001.