

# Lagrangian tracking of sea ice in Community Ice Code (CICE; version 5)

Chenhui Ning<sup>1</sup>, Shiming Xu<sup>1,2</sup>, Yan Zhang<sup>3,1</sup>, Xuantong Wang<sup>1</sup>, Zhihao Fan<sup>1</sup>, and Jiping Liu<sup>4,5</sup>

<sup>1</sup>Ministry of Education Key Laboratory for Earth System Modeling, Department of Earth System Science (DESS), Tsinghua University, Beijing, China

<sup>2</sup>University Corporation for Polar Research (UCPR), Beijing, China

<sup>3</sup>Institute of Applied Physics and Computational Mathematics (IAPCM), Beijing, China

<sup>4</sup>School of Atmospheric Sciences, Sun Yat-sen University, Zhuhai, China

<sup>5</sup>Southern Marine Science and Engineering Guangdong Laboratory (Zhuhai), Zhuhai, China

**Correspondence:** Shiming Xu (xusm@tsinghua.edu.cn) and Jiping Liu (liujp63@mail.sysu.edu.cn)

**Abstract.** Sea ice models are essential tools for simulating the thermodynamic and dynamic processes of the sea ice and the coupling with the polar atmosphere and ocean. Popular models such as the Community Ice Code (CICE) are usually based on non-moving, locally orthogonal Eulerian grids. However, the various in-situ observations such as ice tethered buoys and drift stations, are subjected to sea ice drift and hence by nature Lagrangian. Furthermore, the statistical analysis of sea ice kinematics requires the Lagrangian perspective. As a result, the offline sea ice tracking with model output is usually carried out for many scientific and validation practices. Certain limitations exist, such as the need of *REVLang: high frequency high-frequency* model outputs, as well *REVLang: as* unaccountable tracking errors. In order to facilitate Lagrangian diagnostics in current sea ice models, we design and implement an online Lagrangian tracking module in CICE under the coupled model system of CESM (Community Earth System Model). In this work, we introduce its design and implementation in detail, as well as the numerical experiments for the validation and the analysis of sea ice *REV2: deformations deformation*. In particular, the sea ice model is forced with historical atmospheric reanalysis data, and the Lagrangian tracking results are compared with the observed buoys' tracks for the years from 1979 to 2001. Moreover, high-resolution simulations are carried out with the Lagrangian tracking to study the multi-scale sea ice *REV2: deformations deformation* modeled by CICE. Through scaling analysis, we show that CICE simulates multi-fractal sea ice *REV2: deformations deformation fields* in both the spatial and the temporal domain, as well as the spatial-temporal coupling characteristics. The analysis with model output on the Eulerian grid shows systematic difference with the Lagrangian tracking-based results, highlighting the importance of the Lagrangian perspective for scaling analysis. Related topics, including the *REVLang: subdaily sub-daily* sea ice kinematics and the potential application of the Lagrangian tracking module, are also discussed.

*Copyright statement.* TEXT

## 20 1 Introduction

Sea ice floes are inherently Lagrangian points that undergo thermodynamic and dynamic changes throughout <sup>REVLing:</sup> ~~the~~ their lifetime. Under the dynamic forcings from the atmosphere and the ocean, <sup>REVLing:</sup> ~~the sea ice drifts and with the build-up of internal stress, deforms and undergoes plastic failures.~~ sea ice drifts and, as internal stress accumulates, it deforms and undergoes plastic failures. The drift of sea ice floes is associated with constant thermodynamic growth and melt of the sea ice, hence fundamental to the energy and ice/water balance in the polar regions (Haas, 2009). Furthermore, highly nonlinear and anisotropic linear kinematic features manifest with the sea ice <sup>REV2:</sup> ~~deformations~~ deformation fields, which <sup>REVLing:</sup> ~~are prevalent~~ prevail from meters to the geophysical scales (Marsan et al., 2004). The accurate observation and modeling of the sea ice drift and <sup>REV2:</sup> ~~deformations~~ deformation are key to both our scientific understanding of the climate system and human activities in the polar region.

Due to the harsh conditions of the polar environment, the long-term direct measurements of the sea ice are usually carried out through in-situ deployments of buoys. These autonomous systems, which are usually attached to the sea ice, <sup>REVLing:</sup> ~~report back their locations, the sea ice conditions, as well as the associated atmospheric and ocean conditions.~~ relay information about their locations, sea ice conditions, and associated atmospheric and oceanic conditions. Since they drift with the sea ice, their locations are also representative of the sea ice floe <sup>REV2:</sup> ~~they are attached to~~ to which they are attached. For example, the International Arctic Buoy Programme (IABP: <https://iabp.apl.uw.edu>) compiles historical and real-time buoy measurements in the Arctic region. The data are widely used in the study of sea ice <sup>REV2:</sup> ~~dynamic~~ dynamics (Rigor et al., 2002) and thermodynamic processes (Perovich et al., 2014), as well as the data assimilation for numerical weather forecasts (Inoue et al., 2009).

The drift and the deformation of sea ice are the result of its dynamic response to the atmospheric and oceanic forcings. <sup>REVLing:</sup> ~~Unlike the Newtonian fluids of the air and the water, the sea ice patches undergo deformations with distinctive characteristics of multi-fractal, plastic faults.~~ Unlike the Newtonian fluids of air and water, sea ice patches undergo multi-fractal deformations characterized by localized plastic faults. <sup>REVLing:</sup> ~~These deformations are highly localized and anisotropic, which usually correspond to sea ice leads and ridges and manifest as linear kinematic features.~~ These highly localized and anisotropic deformation typically correspond to sea ice leads and ridges, manifesting as linear kinematic features [LKFs, (Kwok et al., 1998)]. While sea ice ridging is the major way of forming thick sea ice, <sup>REVLing:</sup> ~~leads are important source of~~ leads serve as important sources of heat and moisture for the polar regions especially during winter (Rothrock, 1975; Andreas and Cash, 1999). Therefore, sea ice <sup>REV2:</sup> ~~deformations~~ deformation are crucially important for the coupled climate system of the polar region. In order to study how the sea ice deforms, we usually carry out multi-scale analysis through the statistics of the deformation rates (i.e., the speed at which the sea ice deforms). Specifically, the deformation rates, denoted <sup>REV2:</sup>  ~~$\dot{\epsilon}$ 's~~  $\dot{\epsilon}$ 's, can be computed for individual sea ice patches at various temporal and spatial scales. Since the sea ice is constantly drifting, the computing of  $\dot{\epsilon}$ 's and the related analysis should also be carried out from a Lagrangian perspective. Satellite remote sensing based datasets such as RGPS (Lindsay and Stern, 2003) utilize <sup>REVLing:</sup> ~~SAR~~ Synthetic Aperture Radar (SAR) images collected at <sup>REVLing:</sup> ~~different~~ various times to produce large-scale, high-resolution maps of the sea ice drift and deformation. In particular, both the correlation-based and the feature tracking approaches when processing the SAR images ensure that the analysis of the drift/deformation is Lagrangian by nature. These datasets are widely used for the study <sup>REVLing:</sup> ~~the~~ of

multi-scale sea ice kinematics (Marsan et al., 2004) and the validation of numerical simulations (Kwok et al., 2008; Rampal et al., 2019).

55 For the simulation of sea ice and its kinematics, we construct numerical models which <sup>REVLang:</sup>~~involves~~involve a layered structure with: (1) the mathematical modeling of the physical processes, (2) the numerical treatments for the spatial-temporal discretization and the integration, and (3) the code implementation and the simulation on parallel computers. Popular sea ice models, such as CICE (Community ICE code, <https://github.com/CICE-Consortium/CICE>) and SI<sup>3</sup> (Vancoppenolle et al., 2023), are usually based on the spatial discretization <sup>REVLang:</sup>~~on~~using locally orthogonal structured grids. With the rheology model such as Viscous-Plastic [VP, see also Elasto-Viscous-Plastic (Hunke and Lipscomb, 2008)], these models (Bouchat et al., 2022; Hutter and Losch, 2020) are capable <sup>REV2:</sup>~~to reproduce~~of reproducing certain statistics of the observed multi-scale sea ice kinematics (Kwok et al., 2008). However, the model's output, including the instantaneous and the average model status at different temporal scales, are typically defined on the model's native, Eulerian grid. One notable exception is the neXtSIM model, which is based on Lagrangian moving mesh and <sup>REVLang:</sup>~~natively~~inherently supports the scaling analysis (Rampal et al., 65 2019). But for CICE and many widely adopted sea ice models, the model output is <sup>REVLang:</sup>~~insufficient~~insufficient especially for the scaling analysis at large temporal scales, since it inherently requires a Lagrangian perspective. A typical practice to overcome this limitation is to reconstruct Lagrangian tracks with the model's output, such as <sup>REVLang:</sup>~~that based on the models'~~those derived from daily velocity fields in Bouchat et al. (2022). Certain limitations are still present, however, especially given the ever-growing resolution of current models (Xu et al., 2021; Zhang et al., 2023). High-frequency model output is needed to reconstruct realistic 70 Lagrangian tracks, <sup>REVLang:</sup>~~which requires large amount for~~entailing substantial data storage and off-line computation. Furthermore, the analysis of small-scale sea ice <sup>REV2:</sup>~~deformations~~deformation [i.e., minute-scale as in Oikkonen et al. (2017)] requires even finer spatial and temporal model output and larger overhead with the offline tracking analysis. Therefore, more flexible Lagrangian diagnostic tools are needed for the scaling analysis of sea ice kinematics and future development of sea ice models.

In this paper we introduce the online Lagrangian tracking of sea ice and its model integration in the model of CICE 75 (version 5). The tracking of sea ice is carried out along with the model's numerical integration, and it supports very high-frequency tracking (at the model's time step) and large numbers of Lagrangian points. The model integration is carried out and further validated through the numerical experiments in the coupled framework of Community Earth System Model (CESM, version 2: <https://www.cesm.ucar.edu/models/cesm2>). Specifically, the comparison with observed buoy tracks is carried out with atmospherically forced historical simulations. Furthermore, high-resolution experiment with 7km-resolution in the Arctic 80 region is carried out, and we evaluate the spatial-temporal scaling of wintertime sea ice <sup>REV2:</sup>~~deformations~~deformation. In Section 2 we introduce in detail the tracking algorithm and the integration in CICE. Section 3 includes the numerical experiments and detailed analysis of simulation results. Finally in Section 4, we summarize the article and discuss related topics, including potential applications of the sea ice Lagrangian tracking and the high-resolution simulation of sea ice kinematics.

## 2 Lagrangian tracking in CICE

85 The Lagrangian tracking of sea ice is tightly integrated with the dynamics processes of CICE (version 5). The model grid of CICE is a two-dimensional, logically rectangular, structured grid with the size of  $(nx\_global, ny\_global)$  and indexed by  $(i, j)$ , respectively. Typical lateral boundary conditions are supported for global configurations, including the east-west periodic boundary and the tripolar grid boundary (Murray, 1996). For the Lagrangian tracking, each active point has a specific logical location of  $(x, y)$  in the two-dimensional continuous space, satisfying:  $0 \leq x < nx\_global$  and  $0 \leq y < ny\_global$ .  
90 Furthermore, there is a bi-projection between <sup>REVLang: the point's</sup>each point's logical location and <sup>REVLang: the corresponding geolocation</sup>its corresponding geolocation. The Lagrangian tracking is carried out within the model grid's domain following the tracking algorithm, while maintaining the grid's lateral boundary conditions and the land/ocean distribution.


In this section, the Lagrangian tracking algorithm is introduced in detail in Sec. 2.1, which is based on the existing advection framework of transport remapping (Dukowicz and Baumgardner, 2000). Regarding the commonly used functionality of CICE,  
95 the Lagrangian tracking also supports parallel computing, tripolar grids, and a simple logging system for the record of the tracking results. <sup>REVLang: Details</sup>Detailed implementation and the integration in CICE <sup>REVLang: is</sup>are covered in Sec. 2.2.

### 2.1 Lagrangian tracking algorithm

The Lagrangian tracking algorithm is based on the transport remapping advection scheme, which is available in CICE (Dukowicz and Baumgardner, 2000). As a conservative, two-dimensional, semi-Lagrangian scheme, transport remapping <sup>REVLang: is</sup>based on the Arakawa-B staggered grid. At each dynamics step of CICE, the backward tracking of the corner points is  
100 carried out for the advection of tracers onto the Eulerian grid cells (corners of the hollow quadrilateral in Fig. <sup>REVI: 7.a1.a</sup>7.a1.a). For the Lagrangian tracking, we utilize the backward tracking vectors and compute the forward tracking for all 4 corner points (filled grey arrows in Fig. <sup>REVI: 7.a1.a</sup>7.a1.a), and they form the new quadrilateral (outlined in blue). Then based on the previous location of the Lagrangian point (black circle), we can compute its new location (blue circle) as follows.

105 The tracking of the Lagrangian point is based on the local, normalized coordinate of the grid cell <sup>REVLang: that the point is in</sup>containing the point. Under the local coordinate, the cell's four corners <sup>REVLang: corresponding</sup>correspond to  $(0.0, 0.0)$ ,  $(0.0, 1.0)$ ,  $(1.0, 0.0)$  and  $(1.0, 1.0)$  respectively. The previous location of the Lagrangian point under this coordinate is  $(x_{local}, y_{local})$ , satisfying:  $0 \leq x_{local} < 1$  and  $0 \leq y_{local} < 1$ . Note that given the point's logical position in the model grid as  $(x, y)$ , the following hold: (1) the cell <sup>REVI: the where</sup>point is present has the index of  $(\lfloor x \rfloor + 1, \lfloor y \rfloor + 1)$ ; and (2)  $x_{local} = x - \lfloor x \rfloor$  and  $y_{local} = y - \lfloor y \rfloor$ . After  
110 the forward tracking, the four corners have the new locations of:  $(x_{i,j}, y_{i,j})$  for  $i \in \{0, 1\}$  and  $j \in \{0, 1\}$ . Then, the new location of the Lagrangian point, denoted  $(x_{local}^*, y_{local}^*)$  is computed through the bilinear interpolations with  $x_{i,j}$ 's and  $y_{i,j}$ 's, assuming that its relative location within the new quadrilateral remains  $(x_{local}, y_{local})$ . In the case that  $x_{local}^*$  or  $y_{local}^*$  is larger than 1 or smaller than 0, the Lagrangian point has drifted out of the current grid cell, which we denote as the migration of the point. Since CICE utilizes domain decomposition for parallel computing, the migration potentially is between blocks (sub-domains after  
115 decomposition) and entails communication between parallel processes. The detailed support is introduced further in Section 2.2.






**Figure 1.** Local Lagrangian tracking scheme. Panel (a) shows the backward tracking (the double-lined quadrilateral) and the forward tracking (the quadrilateral in blue line) of the corners of the current T-cell at  $(i, j)$  (thick black quadrilateral). In panel (b), the tracking of a Lagrangian point (black circle) at normalized relative position  $(x, y)$  within the current T-cell is based on linear interpolation of the four corner points [i.e.,  $(x_i, y_j)$ , for  $i = \{0, 1\}$  and  $j = \{0, 1\}$ ].

Based on the geolocations of the four cell corners and the Lagrangian point's relative location within the cell, we can compute the geolocation of the Lagrangian point. Specifically, first, the three-dimensional locations of the cell corners are computed with their latitudes and longitudes. Through linear interpolations, we locate the three-dimensional location of the Lagrangian point. Finally, the latitude and longitude of the Lagrangian point can then be determined.

## 2.2 Implementation in CICE

### 2.2.1 ~~REV2: Basic support~~ Software implementation

For the Lagrangian tracking in CICE, we define the data structure ~~REVLang: is~~ as `lagr_point` in the FORTRAN module of `ice_transport_driver`. It contains necessary fields of information for the Lagrangian point, including its current status, ~~REVLang: its~~ lifetime, ~~REVLang: its~~ current location, and other essential information. Furthermore, each of the parallel CICE processes ~~REVLang: contains~~ maintains a large, pre-allocated pool of available Lagrangian points (i.e., instances of `lagr_point`). When a new Lagrangian point is created in the current process or migrated from another process, an unused slot is claimed from this



**Figure 2.** Typical events for Lagrangian points, including: the creation, the melt (i.e., ice concentration below 5%), drifting within a block or between blocks (i.e., migration), and migration onto lands.

pool. Similarly, when the point is dead (i.e., due to melting) or *REVLag: migrates* migrates out of the block, the slot is reclaimed and recycled in the pool.

130 The life cycle of a Lagrangian point consists of several stages (Fig. 2) and *REVLag: the transition* transitions between the stages, called events. Upon its creation (type-I event), the Lagrangian point is assigned to a specific geolocation, and consequently, a specific block and a specific processor. The Lagrangian point drifts (type-II event), until the sea ice melts (sea ice concentration lower than 5%, type-V event) or it is automatically deactivated (e.g., exceeding the prescribed maximum lifetime). When the Lagrangian points *REVLag: migrates* migrate outside the current block, *REVLag: it they* will be delivered to the corresponding adjacent  
 135 block (type-III event, also see below). This process is carried out through the built-in boundary exchange operations of CICE. A special *landing* event of Lagrangian points (type-IV) is supported, but not possible in the current implementation.

### 2.2.2 Overall model integration

As shown in Figure 3, the tracking of all Lagrangian points occurs within each dynamics time step (i.e., *step\_dynamics*) after the computation of the prognostic velocities and the advection process (which computes the backward tracking vectors).

140 To ensure numerically stable integration for the advection, there is strict limitation over the time step and hence the backward/forward tracking vectors. As a consequence, Lagrangian points cannot migrate beyond one grid cell in either direction. When a specific point migrates beyond the current block's boundary (i.e., thick rectangular boundary in Fig. 2), it will be strictly within the outer boundary of the block (i.e., within the thin rectangular boundary in Fig. 2).

The Lagrangian tracking contains 4 major steps (shown below). For each Lagrangian point, its status is maintained, tracked  
145 with the forward drifting vectors, and <sup>REVLang: theits</sup> location information updated. In the case of the Lagrangian point <sup>REVLang: mi-</sup>gratesmigrating outside the current block, it is recorded for further boundary exchange. Then the Lagrangian points are exchanged between blocks, with newly migrated Lagrangian points recorded for the current block.

1. For each active Lagrangian point:

(a) Check for deactivation (due to melt, life-span, etc.), with necessary management of the Lagrangian point slots

150 (b) Aging of the Lagrangian point

(c) Retrieve the local advection information

(d) Do Lagrangian tracking, and update the logical and the geophysical positions

(e) Check for potential migration out of the current block

(f) For a migrating point:

155 i. Record its information for boundary exchange

ii. Carry out management of the Lagrangian point slot

2. Boundary exchange for Lagrangian points


3. Activate newly migrated Lagrangian points

4. Deactivate Lagrangian points (due to landing, etc.) with necessary managements

### 160 2.2.3 Miscellaneous

#### **Boundary exchanges for migrating Lagrangian points:**

Each dynamics time step involves a single FORTRAN function call to the boundary exchange, carrying out the migration of Lagrangian points between blocks. The maximum exchangeable Lagrangian point count per boundary cell can be changed in the code through the FORTRAN parameter LAGR\_BNDY\_SIZE\_PARAM. It is worth <sup>REV2: to note</sup>noting that the single boundary  
165 exchange due to the migration of Lagrangian points only incurs a very small computational overhead. Within the same dynamics time step, the EVP solver usually contains over 100 <sup>REVLang: subcycles</sup>sub-cycles of the elastic waves and the corresponding boundary exchanges (Lemieux et al., 2010; Xu et al., 2021). Furthermore, the Lagrangian tracking utilizes the transport remapping scheme's backward tracking. In general, the EVP solver dominates the overall simulation time of the sea ice dynamics, with the Lagrangian tracking consuming less than 3% of the time for all the numerical experiments in Section 3.



**Figure 3.** Overall integration of Lagrangian tracking in the time step of CICE.


#### 170 **Support for tripolar grids:**

Tripolar grids are commonly used in high-resolution global configurations of CICE and CESM (Small et al., 2014). For example, the  $0.1^\circ$  grid of TX0.1 is used for the global ocean mesoscale-resolving simulations of CESM. The commonly used U-fold tripolar grids are supported for our implementation of the Lagrangian tracking in CICE, with the schematics shown in Fig. 4.

175 When the Lagrangian point drifts beyond the northern boundary of the blocks on the tripolar folding line, it is passed to the corresponding block and grid cell through the boundary exchange process. This corresponds to the case with  $y$  larger than  $ny\_global$ , where  $ny\_global$  is the grid cell count in the  $j$  direction. Then, the  $x$  and  $y$  position of the point are modified accordingly to the correct values, following the topology on the tripolar boundary.

#### **Logging system:**

180 A simple text-based logging system is implemented to report the results of the Lagrangian tracking. Each parallel processor generates a log file that contains the records of all active Lagrangian points on the processor. The records include the major events, including the creation of the Lagrangian points, the melt events, the migration events, as well as the Lagrangian points' status at regular time intervals. The time step count for reporting Lagrangian points' information is a user-prescribed compile-time parameter (see Appendix C for details).



**Figure 4.** Support of Lagrangian point migration for tripolar grids. The model’s global grid is of size  $nx\_global$  by  $ny\_global$ , with the northern tripolar boundary shown. U-fold tripolar grids are supported (i.e., sea ice velocities are defined on the corner points of the Arakawa-B grid and along the northern boundary).

### 185 3 Numerical experiments and analysis

We carry out numerical experiments of the Lagrangian tracking with CICE (version 5) and CESM (version 2). CICE is configured with 5 ice thickness categories with multiple vertical layers, as well as full thermodynamic and dynamic processes. Key model configurations include the mushy-layer vertical thermodynamics, the Elasto-Viscous-Plastic (EVP) rheology model, and the Delta-Eddington radiation scheme. Detailed parameterizations schemes and major parameters of CICE are further covered in Appendix A. Two model resolutions of CICE are used: the nominally  $1^\circ$  grid of GX1V6 which is built-in in CESM, and the nominally  $0.15^\circ$  grid of TS015 [previously implemented in CESM (Xu et al., 2021)]. Notably, the TS015 grid has the horizontal dimension of  $2400 \times 1680$  globally, and the mean grid resolution is about  $7km$  in the Arctic region. Prominent, multi-fractal sea ice <sup>REV2:</sup>deformations deformation can be simulated at this resolution (Xu et al., 2021).

For all the experiments, CICE is coupled to the Slab Ocean Model (SOM) and forced by the CORE-II dataset through the coupling framework in CESM. The CORE-II dataset is based on NCEP atmospheric reanalysis and further used in the Ocean Model Intercomparison Projects (Griffies et al., 2016). It contains two separate forcing datasets: the Normal Year Forcing (NYF) which is annually repeating, and the Inter-Annual Forcing (IAF) which is for the years between 1948 and 2007. While the NYF dataset is usually used for the long-term simulations and the <sup>REVLan:</sup>spin-up spin-up of the ocean/sea-ice coupled model,

the IAF dataset can be used for the hindcast of the historical states of the ocean and the sea ice (Wang et al., 2016). Both the  
200 NYF and the IAF datasets are used for the numerical experiments, covered in the following of <sup>REVLang:</sup>the this section.


The simulated Arctic sea ice climatology under NYF is shown in Fig. 5. The seasonality of the sea ice extent at both  $1^\circ$   
and  $0.15^\circ$  (not shown) is consistent with the observational climatology (NSIDC). The overestimation of SIE during winter  
months mainly manifests in the peripheral seas of lower latitudes. In particular, in the <sup>REVLang:</sup>Atlantic Atlantic Arctic region, the  
overestimation of SIE may be due to the absence of ocean heat transport of the SOM model. During summer, the SIE agrees  
205 well with the observation. In terms of the sea ice volume, there is general underestimation compared with PIOMAS (Schweiger  
et al., 2011). Thick, multi-year sea ice manifests north to the Greenland and the Canadian Arctic Archipelago (CAA), with  
the clockwise circulation in the Canadian Basin as controlled by the Beaufort High. With the generally good agreement of the  
modeled sea ice states to the observational datasets, we consider it sufficient for further analysis of the Lagrangian tracking.  
Moreover, further improvements of the model's simulation results are planned for future work, including the coupling with a  
210 dynamic ocean model as well as the tuning of parameterization schemes in CICE.

### 3.1 Climatology of Lagrangian points under NYF

Under the annually-repeating NYF dataset, we carry out the basic test of the Lagrangian tracking with the low-resolution,  $1^\circ$   
grid. After the model has reached quasi-equilibrium after 20-year's <sup>REVLang:</sup>spin-up spin-up, we deploy one Lagrangian <sup>REVLang:</sup>points point  
at the center of every grid cell with sea ice (SIC > 5%) in the Northern Hemisphere. The initial locations of the points are shown  
215 in Figure 6.a. With the summer melt and the transport to lower latitudes, the count of active Lagrangian points decreases with  
each year. After a full year since the deployment, the Lagrangian points outside the Arctic Basin are all lost due to melting  
(Fig. 6.b). In Figure 7 <sup>REV2:</sup>we show the two-month tracks of the points for the first whole winter after deployment. The overall  
distribution of remaining points after a melt season, as well as their tracks in the winter, shows: (1) the anti-clockwise drift in  
the Beaufort Gyre at distinctive stages throughout the winter; (2) the convergence of points to the north of Greenland/CAA;  
220 and (3) the transpolar drift from the eastern part of the basin and the outflow in the Fram Strait. After 5 years and 10 years  
(Fig. 6.c and d), the Lagrangian points are generally lost. The surviving points are mainly retained due to the accumulation in  
the Beaufort Gyre. Among all of the 3599 points that are originally in the basin ( $80^\circ N$  and north), about 2063 (57.3%) points  
are lost through melting in the basin within the first 20 years, while 1083 (30.1%) due to export from the Fram Strait (FS).  
In later years, the surviving points are mainly lost due to the circulation to the transpolar drift (hence the FS export) or to the  
225 eastern part of the basin with <sup>REVLang:</sup>a more prominent summer melt. The daily locations of the Lagrangian points are shown by  
the animation in the supplementary to the manuscript. It is worth <sup>REV2:</sup>to note noting that the NYF forcing dataset does not sup-  
port direct comparison with observations. Hence the experiment results, including the simulated circulation of the Lagrangian  
points and the loss due to melt process, <sup>REVLang:</sup>provides provide us with basic test of the Lagrangian tracking functionalities.


### 3.2 Validation of Lagrangian tracking under IAF

230 We further <sup>REVLang:</sup>carry out the evaluate the Lagrangian tracking through the IAF-based simulation and <sup>REVLang:</sup>the comparison COM-  
pare it with the buoy measurements. Based on the  $1^\circ$  grid (GX1V6) and the model's <sup>REVLang:</sup>spin-up spin-up status under NYF, we




**Figure 5.** March (a) and September (b) sea ice thickness simulated by CESM and GX1V6 grid. The last *REVLans: 10-year's 10 years'* results are used to compute the thickness fields (filled contour) and the ice edge with sea ice concentration (SIC) at 15% (red line). The climatological sea ice concentration from NSIDC (based on passive microwave sensors) of the corresponding months are computed for the years between 1979 and 2000, with the sea ice edge shown by the yellow line (SIC=15%). The seasonal cycle of the Arctic sea ice extent (SIE) and the volume (SIV) are shown in panel (c) and (d), respectively. The simulation results from both GX1V6 and TS015 are included. For comparison, the SIE is based on NSIDC dataset, and the SIV is computed from the PIOMAS dataset for the same period (i.e., 1979 to 2000).*REV2:* [Colormap changed for ice thickness.]





**Figure 6.** Lagrangian points (dots) at the initial deployment on Jan-1st (a), after 1 year (b), after 5 years (c) and 10 years (d). The previous locations of each point *REVLang:isare* also shown of up to 2 weeks (lines). *REV2:One in every two Lagrangian points is shown in (b,c,d).*



**Figure 7.** <sup>REV2:</sup>Sea-level pressure (SLP, filled contour) and the tracks of Lagrangian points for two bi-monthly periods: November-December (left) and February-March (right) for the NYF-based simulation with the GX1V6 grid. The tracking results during the first whole winter after the deployment of the Lagrangian points are shown. One in every five Lagrangian points, along with its track throughout the two periods, are indicated by the dark blue dot and the thin blue lines respectively.

carry out the historical simulation for the years between 1979 and 2001. For every day during the winter (December to March) at 0:00 (UTC), we deploy a Lagrangian point at every grid cell <sup>REVLag:</sup>with the presence of sea ice where sea ice is present. For each Lagrangian point, we track its location <sup>REVLag:</sup>effor up to 90 days. The daily instantaneous locations of the points are recorded  
235 and further compared with the observations.

The observational dataset from the International Arctic Buoy Programme (Rigor et al., 2008, IABP) during the same years is used for validation (data available at: <https://iabp.apl.uw.edu/data.html>, last access: 10 May 2023). Since we deploy Lagrangian points to the center of the grid cells, we use the following criteria to match <sup>REVLag:</sup>to the physical buoys' locations of IABP. In total, 621 buoys with 49,004 hourly locations are available for comparison. We further split each buoy's continuous track into  
240 sub-tracks of 14 days for further comparisons. For each buoy's (sub-)track, we screen over the newly deployed Lagrangian points on its starting date. The Lagrangian point nearest to the track's starting location is located, and the simulated track is then compared against the observation. Leap years are ignored in the IAF dataset and the simulations, hence the buoys' tracks starting on February 29th are matched with the Lagrangian points on March 1st.

Figure 8 shows the matching between the buoys' and the corresponding Lagrangian points' locations during the winter of  
245 1993–1994 (December to March). All of the sub-tracks, each covering <sup>REVLag:</sup>14 day's 14-day buoy's track, are shown. The two major sea ice drift regimes include the clockwise circulation in the Canadian basin and the clockwise drift and export of the sea

ice in the region east to  $60^{\circ}W$  and west to  $130^{\circ}E$ . In both regimes, the modeled Lagrangian points' tracks are highly consistent with the observations.


Furthermore, Figure 9 shows for all the sub-tracks the distance between the locations of the buoys and the corresponding Lagrangian points. In total there are 2849 sub-tracks for comparison, each corresponding to a buoy's drift within 14 days. For the starting locations of the buoys' sub-tracks, the average distances to the matching virtual buoys are all within  $50km$ . The average grid size within the Arctic Basin is about  $50km$ , with finer (coarser) resolution in the area near (far from) the Greenland. For over 75% of the matching buoy pairs, the initial distance is less than  $27km$ . After 14-day's tracking, the distance gradually grows, but for 75% of all pairs, the distance is within  $60km$ . Besides, the median (mean) distance is  $38km$  ( $43km$ ). On average, the average drift distance of buoys is  $87km$  at 14-day scale. For fast-moving buoys such as those entering the Fram Strait, the distances between the Lagrangian points and the matching physical buoys are larger, but the relative error always remains low within the two-week tracking period. In general, based on the Lagrangian tracking in CICE, the sea ice drift as simulated by the model matches well with buoys' observations. The tracking uncertainties may arise from the limited spatial resolution of the model, the uncertainty of the atmospheric forcing dataset, as well as the sea ice model's dynamics in simulating the observed drifts. Besides, due to the regular deployment of the Lagrangian points, there is no exact match of the buoys' initial locations. Further attribution of the tracking error to the various contributing factors, including the model and the data's uncertainty, as well as the initial location errors, is planned for future study.

### 3.3 Scaling analysis of sea ice kinematics in the $7km$ -resolution simulation


We further carry out high-resolution simulations of sea ice kinematics and related analysis of the sea ice deformations. In particular, with the  $7km$  resolution of the TS015 grid in the Arctic region, the model is capable of simulating the multi-fractal sea ice deformation (Xu et al., 2021). The experiment is based on the NYF dataset and the model's quasi-equilibrium status after the spin-up process. The Lagrangian tracking is utilized for the diagnostics of the wintertime sea ice deformation and related statistics over multiple spatial and temporal scales. We first introduce the basic framework of scaling analysis and the use of Lagrangian tracking in Sec. 3.3.1. Furthermore, the detailed analysis of the spatial scaling (Sec. 3.3.2) and the temporal scaling and the spatial-temporal coupling (Sec. 3.3.3) are covered.

#### 3.3.1 Lagrangian tracking and scaling analysis

The sea ice deformation rates ( $\dot{\epsilon}$ 's) are defined with the simulated velocities. Since CICE is based on the locally orthogonal grid and Arakawa-B staggering, the model's velocity as well as the derived velocity of Lagrangian points are defined as fields of  $u$ 's and  $v$ 's. The grid invariant deformation rates, including the divergence rate ( $\dot{\epsilon}_{div}$ ) and the shear rate ( $\dot{\epsilon}_{shear}$ ) are then defined as follows. The total deformation rate ( $\dot{\epsilon}_{total}$ ) which is subjected to further scaling analysis, can be further defined with  $\dot{\epsilon}_{div}$  and  $\dot{\epsilon}_{shear}$ .



**Figure 8.** Simulated Lagrangian points (blue) and those of the corresponding IABP buoys (red) during the winter of 1993–1994. The last location (+) of each record and the track (line) of up to 14 days are shown for each buoy/Lagrangian point. The wintertime mean (DJFM) sea ice concentration is also shown in the background. <sup>REV1</sup>: [Increased the size of '+' markers on the map.]



**Figure 9.** Distance between the modeled Lagrangian points and the corresponding buoys. The yellow line is the median, the lower and upper edges of the green shade are the 25th and the 75th percentiles, respectively. The horizontal bar (grey dashed line) marks half of the mean grid size ( $\sqrt{dx \cdot dy}/2$ ) of the GXIV6 grid in the Arctic Basin.

$$\dot{\epsilon}_{\text{div}} = u_x + v_y \quad (1)$$

$$\dot{\epsilon}_{\text{shear}} = \sqrt{(u_x - v_y)^2 + (u_y + v_x)^2} \quad (2)$$

$$280 \quad \dot{\epsilon}_{\text{total}} = \sqrt{\dot{\epsilon}_{\text{div}}^2 + \dot{\epsilon}_{\text{shear}}^2} \quad (3)$$

In these equations,  $u_x$ ,  $u_y$ ,  $v_x$  and  $v_y$  are the spatial derivatives of  $u$  and  $v$  in the two orthogonal directions. The direct model output of the instantaneous and/or the temporal mean of the velocity fields, as well as the various deformation rates, are defined on the Eulerian model grid. In Figure 10 we show the maps of the daily-mean  $\dot{\epsilon}_{\text{total}}$  in the Arctic region for two typical days of the experiment with TS015. LKFs manifest with the highly localized, anisotropic sea ice <sup>REV2: deformations</sup> deformation. It is



285 worth <sup>REV2:</sup>to note noting that the daily-mean velocity and the deformation fields (shown in Fig. 10) are inherent Eulerian, hence not proper for the scaling analysis. For example, the offline tracking (at hourly scale) should be carried out for further analysis [e.g., Bouchat et al. (2022)].

The velocity derivatives at various temporal/spatial scales (i.e.,  $\{u, v\}_{\{x, y\}}$ 's) are further computed over the various Lagrangian patches. A set of adjacent Lagrangian points form an enclosed sea ice patch and start with the position on regular Eulerian grid points. Their locations change with the Lagrangian tracking, due to the sea ice drift and deformation. At certain time (delayed by  $T$ ), their new locations are recorded and the displacement from their original locations are computed as  $\Delta_x$ 's and  $\Delta_y$ 's. Then the average velocity  $\bar{u}$  and  $\bar{v}$  for each Lagrangian point can be computed as  $\bar{u} = \frac{\Delta_x}{\Delta_t}$  and  $\bar{v} = \frac{\Delta_y}{\Delta_t}$  respectively. The spatial velocity derivatives of the Lagrangian patch are computed with the <sup>REVLanq:</sup>the patch's area ( $A$ ) and the line integral of velocity over its outer boundary, following Kwok et al. (2008) and Rampal et al. (2019). Details of the computation are further covered in Appendix B.

Given the total deformation rates of all sea ice patches, we compute the average value of  $\dot{\epsilon}_{total}$ 's within the similar spatial scale ( $L = \sqrt{A}$ ). In particular, the  $q$ -th order of  $\dot{\epsilon}_{total}$ 's, computed as  $\dot{\epsilon}_{total}^q$ 's, along with their average value are also computed for  $q \in \{0.5, 1, 1.5, 2, 2.5, 3\}$ . By the scaling law (Marsan et al., 2004), under the spatial scale of  $L$  and the temporal scale of  $T$ , we have:

$$300 \quad \langle \dot{\epsilon}^q(T, L) \rangle \sim T^{-\alpha(q)} \quad (4)$$


$$\langle \dot{\epsilon}^q(T, L) \rangle \sim L^{-\beta(q)} \quad (5)$$

Where  $\alpha(q)$  and  $\beta(q)$  are the structure functions of the temporal and the spatial scaling. For a specific value of  $q$ , we carry out the linear fitting of  $\langle \dot{\epsilon}^q(T, L) \rangle$  with respect to  $T$  (or  $L$ ) to estimate the value of  $\alpha$  (or  $\beta$ ). For the observed multi-fractal <sup>REV2:</sup>deformations deformation of the sea ice (Marsan et al., 2004), we witness convex structure functions for both  $\alpha(q)$  and  $\beta(q)$  with respect to  $q$ . <sup>REVI:</sup>[A generalized analysis framework with non-fixed degree of multifractality for the sea ice formation is also available](#) (Weiss, 2008; Bouchat et al., 2022). <sup>REVI:</sup>[For comparison, the forms in Eqs. 6 and 7 also assume the underlying multi-fractal, long-normal multiplicative model. In this study, they are adopted because their quadratic form is sufficient in capturing the convex shape of the structure functions](#) (Marsan et al., 2004; Rampal et al., 2019). Correspondingly,  $\alpha(q)$  and  $\beta(q)$  can be fitted as follows, with the fitted <sup>REVLanq:</sup>parameter parameters of  $a$  and  $c$  both larger than 0.

$$310 \quad \beta(q) = a \cdot q^2 + b \cdot q \quad (6)$$

$$\alpha(q) = c \cdot q^2 + d \cdot q \quad (7)$$

In order to evaluate the simulated sea ice <sup>REV2:</sup>deformations deformation, we deploy a Lagrangian point at the center of each grid cell for every model day at the time of 0:00. The maximum allowed life span of each Lagrangian point is 30 days. As a consequence, the temporal scaling of up to 30-day can be studied (i.e., the maximum value of  $T$  at 30-day). The model regularly reports the location of every Lagrangian point every 6 hours (i.e., the minimum value of  $T$  at 6-hour). Both parameters can be



**Figure 10.** Daily-mean sea ice total deformation on Dec-20 (left) and Feb-6 (right) simulated by CESM and TS015. The region subjected to further scaling analysis is outlined by the dashed blue line in each panel.

configured at the compile time of the CICE model. The model grid's native resolution in the Arctic is about  $7.3\text{km}$ . Given that the effective resolution is coarser (Xu et al., 2021), we evaluate the  $REV2$ :deformationsdeformation at the spatial scale from  $30\text{km}$  to  $480\text{km}$  (4 times to 64 times the grid's resolution). For the scaling analysis, we limit the initial locations of all Lagrangian points to be at least  $400\text{km}$  away from land (i.e., within the outlined region in Fig. 10). The regions near the coast and the continental shelf break, where constant deformation features persist, are therefore excluded from the scaling analysis.

After the model's  $REVL$ ang:spin-upspin-up, we output a whole winter's Lagrangian tracking results, together with the daily mean Eulerian fields of the sea ice velocity and the deformation. Furthermore, we also study the difference between the Lagrangian tracking-based scaling analysis and that based on the temporal/spatial mean fields of the model output on Eulerian locations.

### 3.3.2 Spatial scaling

We carry out spatial scaling analysis for two typical periods during the winter, focusing on 4 typical temporal scales: 1-day, 3-day, 10-day, and 30-day. Figure 11 and 12 show the results around Dec. 20th and Feb. 6th, respectively. At  $q = 1$  and 1-day scale, the spatial  $REVL$ ang:scalingscalings of sea ice deformation are both very shallow:  $\beta = 0.0623$  and  $0.121$ , respectively. These two days correspond to different strengths and depths of the Beaufort High and slightly different Arctic Oscillation indices, and consequently the drift and deformation fields are distinct [see Fig. 6 and 11 of Xu et al. (2021)]. At different orders of momentum, the mean deformation rates all follow the power-law scaling. On higher orders ( $q > 1$ ), large deformation events are more dominant in the mean deformation rate, corresponding to steeper scaling (i.e., larger values of  $\beta$ ). Furthermore, the value of  $\beta$  grows nonlinearly with  $q$ . With the increase of the moment order, the deformation rate decreases faster with respect  $REVL$ ang:to the increase of the spatial scale ( $L$ ). Correspondingly, the convex shape of  $\beta(q)$  indicates that the model simulates multi-fractal sea ice  $REV2$ :deformationsdeformation.




335 For the period of study surrounding Dec. 20th, at larger temporal scales from 1-day to 30-day, the average deformation rates decrease at all spatial scales and moments of order. More importantly, the value and the shape of  $\beta(q)$  also change significantly with the temporal scale. At  $q = 1$  and the 1-day temporal scale, the value of  $\beta$  is 0.0623. For larger temporal scales, the mean deformation rate gradually decreases, which also applies to the structure function of  $\beta$ . In particular, the convexity of  $\beta(q)$ , indicated by the fitted value of  $a$ , decreases drastically from 0.137 (1-day) to 0.064 (30-day). The change in  $a$  indicates that  
340 there is <sup>REVLang: a</sup> strong coupling between the spatial and temporal domain for the sea ice deformation. Similar results hold for the period around Feb. 6th, with the two following major differences: (1) more convex  $\beta(q)$  around Feb. 6th than Dec. 20th; (2) the faster decrease of  $a$  with respect to  $T$  (i.e., the temporal scale). The most significant drop in  $a$  occurred from the 3-day to the 10-day scale for Feb. 6th (from 0.151 to 0.101). For comparison, that for Dec. 20th is between 10-day to 30-day (from 0.093 to 0.064). Regarding the differences between the two periods, we conjecture that the process-dependent deformations  
345 are the major cause, including the strength of the spatial-temporal coupling. Further analysis is needed for the attribution of these differences to various factors, including the sea ice status, its deformation history, as well as the forcings.

For comparison, we also <sup>REVLang: shows</sup> show the results based on Eulerian deformation fields in both Figure 11 and 12, including the power-law fittings and the structure functions of  $\beta(q)$ . The method details are introduced in Appendix B. For both periods of study, the scaling analysis with Lagrangian tracking results show steeper  $\beta$  functions with higher convexity across all  
350 temporal scales (i.e., larger values of  $a$ 's). For Feb. 6th, the difference between the two is more pronounced. Note that the scaling analysis should be carried out based on the Lagrangian perspective. The objective <sup>REVLang: for</sup> of the comparisons is to demonstrate that there exist systematic differences <sup>REVLang: of</sup> in the scaling analysis when using the Eulerian model outputs. The quantitative differences <sup>REVLang: of</sup> in the deformation rates may arise from: (1) that the deformation events are misaligned between the Eulerian and the Lagrangian <sup>REVLang: perspective</sup> perspectives, and (2) the secondary effect of changing shapes and scales when  
355 the sea ice deforms within the Lagrangian perspective, which is not captured by the Eulerian framework.

### 3.3.3 Temporal scaling and spatial-temporal coupling

For the temporal scaling analysis, we focus on the spatial scale of the model's effective resolution ( $L = 22km$ ). Figure 13 shows the results for the outlined region in Figure 10 in the month of December (left) and February (right). The analyzed temporal scale is in the range between 1-day and 30-day. Similar to the spatial scaling results, the power-law scaling is witnessed for the  
360 temporal scaling (top panels of Fig. 13). Also, convex structure functions of  $\alpha(q)$  are present for both months, with the fitted value of  $c$  as 0.063 and 0.059, respectively. <sup>REVLang: This</sup> These results indicate that the model also simulates multi-fractal sea ice deformation in the temporal domain.


Evidently, the power-law scaling at sub-daily scales is much shallower than that between 1-day and 30-day. Similar behavior of the sea ice is observed in Oikkonen et al. (2017), which covers the temporal scale from about 10 minutes. We consider the  
365 shallower scaling in the simulations <sup>REVLang: is to be</sup> qualitatively consistent with the observation. However, although the model's dynamics timestep is sufficiently short (see Appendix A), the temporal resolution of the forcing dataset is much coarser, at 6-hourly. In order to fully study the simulation of the sub-daily sea ice <sup>REV2: deformations</sup> deformation, we need high-frequency forcing datasets or the coupled simulation with the high-resolution interactive atmospheric component (Zhang et al., 2023).



**Figure 11.** Spatial scaling around Dec. 20th. Four temporal *REVLing*: scalescales are evaluated: 1-day (first column), 3-day (second column), 10-day (third column), and 30-day (last column). Results with Lagrangian tracking (triangles) are compared with *REVLing*: thatthose based on temporally mean Eulerian deformation fields (dots). Black lines in each panel represent the fittings with the spatial scale of 30km and coarser, and red dashed lines *REVLing*: represent the fittings with the whole resolution range of TS015 (7.3km and coarser). The fitted parameter of  $a$  for the structure function  $\beta(q)$  is shown in the legend.

Similar to the analysis in Section 3.3.2, we also compute the temporal mean Eulerian deformation fields and the equivalent  
 370 scaling analysis. Since we only output daily model fields, the analysis is limited to the temporal scale between 1-day and 30-day  
 (middle row of Fig. 13). Apparent power-law scaling is witnessed for both months. However, the structure function of  $\alpha(q)$   
 with the Lagrangian tracking is systematically less convex than that based on Eulerian fields, with lower values of  $c$  for both  
 months (bottom row of Fig. 13). This result, together with that in Section 3.3.2, show that the scaling analysis should be carried  
 out in the Lagrangian framework. Although the scaling analysis with the model's Eulerian mean fields yields apparent multi-  
 375 fractal sea ice *REV2*: deformationsdeformation, the results are systematically different from *REVLing*: the those based on the Lagrangian  
 analysis.


Furthermore, we conduct the initial analysis of the spatial-temporal coupling during the whole winter (December to March).  
 For the region of study (outlined in Fig. 10), we utilize all of the Lagrangian tracking results by forming and tracking Lagrangian



**Figure 12.** Same as Fig. 11, but for Feb. 6th.


patches at various spatial and temporal scales. Specifically, for each combination of the spatial and the temporal scale (i.e.,  $L$  and  $T$ ), we form Lagrangian patches that satisfy the following criteria: (1) they start at interleaved Eulerian grid locations separated by  $L/2$  in both directions, and (2) the time difference between their starting time and 0:00 of Dec. 1st separated by  $n \cdot T/2$ , where  $n \in \{0, 1, 2, 3, \dots\}$ . With these two criteria, we in effect split the Lagrangian tracking results into sea ice Lagrangian patches with 50% temporal and spatial overlapping. As a result, we not only attain full coverage of the study area/period, but also avoid potential sampling issues associated with the changing weather regimes.

The relationship between the curvature parameter of the spatial (temporal) scaling structure function and the temporal (spatial) scale is shown in Figure 14.a (b). There is evident coupling between the spatial domain and the temporal domain, with decreased curvature of the spatial (temporal) structure function at larger temporal (spatial) scales. In particular, for the temporal scaling, there is a good fit of the curvature parameter  $c$  to the Power-Law (Fig. 14.b), which is consistent with various estimations based on in-situ and satellite-based remote sensing observations (Rampal et al., 2008; Marsan and Weiss, 2010). However, for the spatial scaling, the relationship of  $a$  to the temporal scale is much flatter and in particular, less convex for a Power-Law fit (Fig. 14.a). We note that during different periods of the winter, the sea ice drift and deformation patterns are



**Figure 13.** Temporal scaling for December (left) and February (right) at the spatial scale of  $22km$  (i.e.,  $3\times$  the original grid resolution). Similar to Fig. 11 and Fig. 12, the results with Lagrangian tracking and Eulerian mean deformation fields are shown by triangles and dots, respectively. Note that the *REVLang*:fitting fittings are only computed for 1-day or coarser temporal scales (indicated by the abscissa range of the fitted lines).

highly heterogeneous (due to changing *REVLang*:weathers weather), with the sea ice condition undergoing significant changes. For example, the spatial scaling exponent shows large temporal variability (Rampal et al., 2019) and is likely linked to the atmospheric forcing patterns (Xu et al., 2021). Therefore, we adopt another fitting with the *log*-quadratic form:  $y = p \cdot \log(x)^2 + q$ .  
 395 This new form yields *REVLang*:a much higher fitting to the relationship between the curvature parameter  $a$  and the temporal scale  $T$  ( $R^2$  from 0.882 to 0.989). The efficacy of the Power-Law fitting, as well as the full analysis of the spatial-temporal coupling as simulated by CICE, is beyond the scope of this study. Especially, the sensitivity to sea ice rheology model and other dynamic processes is planned for both CICE standalone and coupled experiments with CESM.



**Figure 14.** Temporal-spatial coupling (Dec.-Mar.) analysis of Lagrangian tracking deformation fields: The curvature of the structure function of the spatial(temporal) scaling exponent as a function of temporal(spatial) scale is shown in panel a(b). The solid and dashed lines represent the logarithm-quadratic fitting of  $y = p \cdot (\log(x))^2 + q$  and the power-law fitting of  $y = p \cdot x^q$ , respectively. The red (black) line in the first panel shows the fitting with all the spatial scales (the scales with the model’s effective resolutions). The  $R^2$  of each fitting in top panels are shown, respectively. Scaling analysis with both time scale and spatial scale as independent variables are shown in (c), correspondent structure functions are shown in (d).

#### 4 Summary and discussion

400 We design and implement the online Lagrangian tracking in the sea ice model of CICE. It is incorporated in the dynamics process of CICE, and fully supports the domain decomposition and parallel computing in CICE. Compared with the common practice of offline sea-ice tracking (Bouchat et al., 2022; Sumata et al., 2023), the online tracking has several advantages. First, the offline tracking usually requires the storage of high-frequency sea ice drift data, which is not needed for the online tracking. Second, since the online tracking is carried out per dynamics step, the tracking error is minimal. For offline tracking, 405 the tracking uncertainty arises from the *REVLang: relatively* relatively coarse temporal mean (e.g., daily) sea ice drift fields, as well as the variable tracking frequency [see examples in Bouchat et al. (2022)]. Moreover, the study of sea ice dynamics at small temporal scales (such as sub-daily motions) with offline tracking requires *REVLang: an* an even finer resolution of the stored sea ice drift. The study of these processes is readily supported by the online tracking [see Fig. 13 and also Oikkonen et al. (2017)]. Computationally, the Lagrangian tracking only induces a small overhead to the overall simulation. For example, for the daily 410 deployment of Lagrangian points in the *7km* experiment in Section 3, there are on average 30 to 35 million active Lagrangian points, and the extra computational overhead is less than 3%. Therefore, the current implementation is capable *REVLang: to support* Of supporting high-resolution simulations with a large quantity of Lagrangian points.

In Section 3.3 we carried out the spatial and the temporal scaling analysis with the Lagrangian tracking based on high-resolution simulation during winter. As shown, the CICE model simulates multi-fractal sea ice *REV2: deformations* deformation 415 fields both spatially and temporally, as well as the tight coupling between the spatial and the temporal domain. The scaling

properties of the simulations are consistent with observed statistics based on high-resolution synthetic aperture radar satellite payloads. In particular, we compare the Lagrangian-based scaling statistics with the counterparts based on Eulerian model outputs. Results highlight the importance of using Lagrangian-based diagnosis for the scaling analysis: although the analysis based on Eulerian output also yields convex structure functions, the fitted convexity parameters are significantly different from those based on the Lagrangian perspective.

In order to compare with the observed sea ice drift and deformation (e.g., RGPS) as well as the scaling statistics, we plan to carry out high-resolution, atmospherically-forced historical simulations with the CICE/CESM model assisted with the Lagrangian tracking. Consequently, fine tuning of the sea ice model parameters *REVLang: are* is needed for: (1) the optimization of the simulation of various sea ice parameters such as sea ice thickness, and (2) improved comparability to the observed sea ice *REV2: deformations* deformation. Especially, in Bouchat and Tremblay (2020) the authors proposed *REVLang: the* a new estimation of deformation rates regarding their uncertainties. Better consistency with model simulations is attained with further incorporation of the model simulation and tracking uncertainties (Bouchat et al., 2022). We plan to incorporate the method and carry out *REVLang: a* systematic evaluation of the model's capability to *REVLang: simulated* simulate the observed multi-scale sea ice *REV2: deforma-* tions deformation. In particular, the sensitivity to the sea ice rheology is a key aspect of the planned future work (Tsamados et al., 2013; Bouchat et al., 2022).

In this first version of the Lagrangian tracking in CICE, virtual buoys are supported, which can be deployed at regular time intervals and on regular locations. Section 3.2 shows that with the pairing between the physical buoys and the respective virtual buoys, the simulated tracks are generally consistent with observations. For future development, we plan to support the deployment of physical buoys in the simulation at prescribed time and locations, so that we can better compare with observational datasets such as IABP. Moreover, key Lagrangian tracking related parameters (in Appendix B) *REVLang: is* are planned to be configured through a subset of the namelist in the future development plan. Finally, the current design of Lagrangian tracking is based on the Arakawa-B grid sea ice dynamics and the transport remapping scheme. For the compatibility with ongoing development in CICE6 (<https://github.com/CICE-Consortium/CICE/releases>), we also plan to incorporate flux-form based Lagrangian tracking which is suitable for Arakawa-C grid sea ice dynamics (Lemieux et al., 2024).

The Lagrangian tracking can be used in a variety of science questions and applications related to sea ice. Large-scale Lagrangian survey can be carried out in the simulations, complementary to buoys' measurements which are limited in terms of spatial and temporal coverage. For example, virtual buoys can be deployed in historical simulations, which enables systematic study of the thermodynamic and dynamic history of the sea ice (Sumata et al., 2023). Furthermore, CICE is widely adopted for the high-resolution sea ice operational systems (Smith et al., 2016; Yang et al., 2020). The online tracking of sea ice in key regions can be carried out for sea ice forecasts, in order to support operations and ship navigation in polar waters.

*Acknowledgements.* The authors would like to thank the editors and referees for their invaluable efforts in improving the manuscript. This work is mainly supported by the joint project of INTERAAC co-funded by the National Key R&D Program of China (grant no.: 2022YFE0106700) and the Research Council of Norway (grant no.: 328957). JL is supported by the National Key R&D Program of China

(grant no.: 2018YFA0605900). SX is also partially supported by the National Natural Science Foundation of China (grant no.: 42030602),  
450 the International Partnership Program of Chinese Academy of Sciences (grant no.: 183311KY5B20200015), and the Research Council of  
Norway (grant no.: 328957). The authors would also like to acknowledge the computational and technical support from the National Super-  
computing Center at Wuxi for the numerical experiments.

*Code and data availability.* The original codebase of the CESM model (version 2, <https://www.cesm.ucar.edu/models/cesm2>) and the asso-  
ciated CICE model (version 5) is available through: <https://github.com/ESCOMP/CESM/tree/release-cesm2.2.2> (last access: 2024-Jan-20).  
455 <sup>REV2:</sup>The CESM code is also archived at: <https://zenodo.org/records/12616647> (last access: 2024-Jun-20). The sea ice concentration data  
is available at NSIDC through: <https://noaadata.apps.nsidc.org/NOAA/G02135/> (last access: 2024-Feb-21). The sea ice thickness dataset  
of PIOMAS is downloaded from: <https://psc.apl.uw.edu/research/projects/arctic-sea-ice-volume-anomaly/data/> (last access: 2024-Feb-21).  
The buoy tracks from IABP program are downloaded from: [https://iabp.apl.uw.edu/Data\\_Products/BUOY\\_DATA/3HOURLY\\_DATA/](https://iabp.apl.uw.edu/Data_Products/BUOY_DATA/3HOURLY_DATA/) (last  
access: 2024-Feb-21).

460 The Lagrangian tracking in CICE (version 5) and the sample Lagrangian tracking output are available at: [https://zenodo.org/records/](https://zenodo.org/records/12616647)  
[12616647](https://zenodo.org/records/12616647) <sup>REV2:</sup>(last access: 2024-Mar-7)(last access: 2024-Jun-20). <sup>REV2:</sup>The CICE namelist used in the numerical simulations are also included.

The animation produced with the Lagrangian tracking with the  $1^\circ$  grid under NYF dataset is accessible as supplementary material of this  
article.

## Appendix A: Model configuration and parameters of CICE

465 The CICE (version 5.0) is configured with 5 discretized thickness categories for the Ice Thickness Distribution (ITD), 8 and 3  
layers for sea ice and its snow cover, respectively. The Elasto-Viscous-Plastic (EVP) rheology model is used for all experiments.  
The ice strength parameterization follows that in Rothrock (1975), which relates the ice strength to the gain of the potential  
energy during the ridging process. The ridging/rafting parameterization uses an exponential distribution form of the ridged  
ice in the ITD. For sea ice thermodynamics, the mushy layer physics parameterization is adopted (Turner et al., 2013). The  
470 Delta-Eddington (D-E) scheme is used for the radiation processes (Holland et al., 2012). Table A1 lists the key parameters of  
the model. The parameters are the same across all numerical experiments by default, except the dynamics time step and the  
EVP subcycling count (NDTE). These two parameters are set differently between the experiments with GX1V6 and those with  
TS015. The shorter time step is adopted for TS015 to ensure numerical stability, and the value of NDTE is further enlarged for  
the numerical convergence of the EVP rheology (Lemieux et al., 2010; Xu et al., 2021).

## 475 Appendix B: Scaling analysis with Lagrangian tracking and Eulerian fields

We carry out the scaling analysis of the sea ice <sup>REV2:</sup>~~deformations~~deformation at various spatial and temporal scales (Rampal et al.,  
2019; Bouchat and Tremblay, 2020). The Lagrangian tracking results, as well as the model outputs on Eulerian grids, are



**Table A1.** Major model parameters of CICE<sup>REV2</sup>: [Model parameters are added for the sake of completeness]

Parameter	Value	Notes
Time stepping		
DT	3600s (GX1V6) and 1800s (TS015)	Time step for sea ice dynamics
NDTE	120 (GX1V6) and 960 (TS015)	EVP subcycling count
Parameterization: ridging and redistribution		
$a^*$	0.5	$e$ -folding scale for participation function during ridging
$\mu_{rdg}$	$4\sqrt{m}$	$e$ -folding scale for redistribution
Ice strength parameterization (R75)		
$C_f$	17	Parameter for potential/frictional energy during ridging
Albedo parameterization (CCSM3, or default)		
albice_i	0.45	Albedo at infra-red spectral band for bare ice
albice_v	0.75	Albedo at visible spectral band for bare ice
albsnow_i	0.73	Albedo at infra-red spectral band for snow surface
albsnow_v	0.98	Albedo at visible spectral band for snow surface
Shortwave parameterization (Delta-Eddington)		
$\rho_s$	$330\text{ kg/m}^3$	Snow density
$R_{fresh}$	$100\text{ }\mu\text{m}$	Freshly-fallen snow grain radius
$R_{nonmelt}$	$500\text{ }\mu\text{m}$	Seasoned snow grain radius
$R_{melt}$	$1000\text{ }\mu\text{m}$	Melting snow grain radius
Thermodynamics parameterization (Mushy Layer & MU71)		
a_rapid_mode	$0.5 \times 10^{-3}\text{ m}$	Brine channel diameter
aspect_rapid_mode	1	Brine convection aspect ratio
dSdt_slow_mode	$-1.5 \times 10^{-7}\text{ m/s/K}$	Brine drainage strength
phi_c_slow_mode	0.05	Critical liquid fraction
phi_i_mushy	0.85	Solid fraction at ice bottom

used to compute the mean sea ice deformation rates at the corresponding scales. A set of Lagrangian points form an enclosed Lagrangian patch, for which we compute the spatial derivatives of velocities (i.e.,  $u_x$ ,  $u_y$ ,  $v_x$  and  $v_y$ ) as follows:

$$\begin{aligned}
 u_x &= \frac{1}{A} \oint u dy \\
 u_y &= -\frac{1}{A} \oint u dx \\
 v_x &= \frac{1}{A} \oint v dy \\
 v_y &= -\frac{1}{A} \oint v dx
 \end{aligned}
 \tag{B1}$$

These derivatives are used to compute the deformation rates as in Eqs. 2 to 3. The line integral in Eqs. B1 is carried out over the Lagrangian patch that originally covers a regular rectangular domain. Then each point on the patch's outer boundary, called a vertex, has its location of  $(x_i^j, y_i^j)$  at the time of  $t_j$ , and the new location of  $(x_i^{j+1}, y_i^{j+1})$  at the time of  $t_{j+1}$ . Figure B1 shows the schematics for a Lagrangian patch with 4 corner points. We can estimate its mean velocity  $(u_i, v_i)$  as the ratios of the displacements to the time difference:  $(x_i^{j+1} - x_i^j)/(t_{j+1} - t_j)$  and  $(y_i^{j+1} - y_i^j)/(t_{j+1} - t_j)$ . Furthermore, in order to compute the line integrals at time  $t$ , we compute the mean velocity on each edge as follows (for the example in Fig. B1):


$$u_x = \frac{1}{A} \sum_{i=1}^4 \frac{1}{2} (u_{i+1} + u_i) (y_{i+1} - y_i),
 \tag{B2}$$

where  $A$  is the area the set of Lagrangian points cover:

$$A = \frac{1}{2} \sum_{i=1}^4 (x_i y_{i+1} - x_{i+1} y_i),
 \tag{B3}$$

In order to complete the line integral, we set  $x_5 = x_1$ , and similarly the corresponding values of  $y_5$ ,  $u_5$ , and  $v_5$ . For larger spatial scales, we start with a larger set of Lagrangian points that have their original locations on the Eulerian grid locations and cover a locally rectangular area. Then the velocities, their spatial derivatives, and the line integrals are computed, followed by the computation of the deformation rates. This is equivalent to the aggregation of all the smallest rectangular units of 4 Lagrangian points that constitute the whole rectangular area. Note that at the beginning of each model day, we deploy the Lagrangian points on all Eulerian grid locations, so that we attain full spatial and temporal coverage, from the daily scale and above. <sup>REVLang:</sup> Besides Additionally, we further exclude the cases with the size change by <sup>REVLang:</sup> the factor of 2 or larger.

For comparison, we also compute the equivalent deformation rates based on Eulerian fields. We directly obtain the temporally mean velocities on each Eulerian grid location, and compute the velocity gradients correspondingly (Fig. B1.b). Different spatial scales correspond to the various areas that <sup>REVLang:</sup> corresponds to are associated with different rectangular grid cells, which do not change <sup>REVLang:</sup> throughout the simulation.



**Figure B1.** Schematics of the scaling analysis with Lagrangian tracking and the equivalent Eulerian model outputs. The locations of Eulerian and Lagrangian points at the initial time  $t_0$  are shown in panel (a), and those at time  $t_1$  are shown in panel (b) and (c), respectively. The spatial scale  $L$  is defined as the square root  $A$  which is the area of the region covered by the four Lagrangian/Eulerian points. Note that in our analysis, the initial locations of the Lagrangian points are the regular, Eulerian grid points. For comparison, on the same time scale (i.e.,  $T = t_1 - t_0$ ), we compute the model-output mean velocities on the Eulerian grid points, as well as the displacement-based velocity estimations for Lagrangian points.

For both Lagrangian statistics and the Eulerian counterparts, we only compute the deformation field that is contained by the areas covered at the coarsest spatial scale. This practice ensures that there is no preferential sampling of the deformation events.

A wide range of both spatial and temporal scales are adopted for the computation of the deformation rates. The statistics of  $\dot{\epsilon}(T, L)$ , which is the deformation rate at the spatial scale of  $L$  and the temporal scale of  $T$ , include its  $q$ -th order [i.e.,  $\dot{\epsilon}^q(T, L)$ ] and the mean value are computed over the studied area. It is worth <sup>REV2: to note</sup>noting that the spatial scale of  $L$  varies within the studied domain, as well as with time for the Lagrangian tracking. Therefore, both the mean value of  $\dot{\epsilon}^q(T, L)$ 's and that of the spatial scales of  $L$ 's are used for the scaling analysis.

### Appendix C: Major parameters of Lagrangian tracking

510 Tab. C1 shows the major user-specified parameters for running the Lagrangian tracking in CICE. In the current implementation, they are configured at the compile-time of the model. In the future, we plan to implement certain parameters to be configurable through the namelist.

*Author contributions.* JL and SX conceived the work. SX carried out the design and the implementation of the Lagrangian tracking in CICE. SX, CN, and YZ carried out the numerical experiments. CN and SX analyzed the results. All the authors contributed to the writing of the manuscript.

**Table C1.** User-prescribed parameters of Lagrangian tracking

Parameter	Type	Default value	Description
LAGR_BUFFER_SIZE_PARAM	Integer	100000	Size of the pool of Lagrangian points
LAGR_BNDY_SIZE_PARAM	Integer	400	Per-cell maximum count of migrating Lagrangian points
lagr_aice_thres	Real	0.05	Threshold value of SIC for detecting melt events
LAGR_REPORT_INTERVAL	Real	18000 <i>s</i>	The interval of reporting Lagrangian points' status
LAGR_ACTIVATION_INTERVAL	Real	86400 <i>s</i>	Virtual buoy activation interval (in seconds)
LAGR_ACTIVATION_DNSTY	Real	1.0	Deployment density of virtual buoys (1 per cell)
LAGR_VIRTUAL_BUOY_MAX_LIFE_DURATION	Real	2592000 <i>s</i>	Maximum lifetime of virtual buoys (in seconds)

*Competing interests.* The authors declare that they have no conflict of interest.

## References

- Andreas, E. L. and Cash, B. A.: Convective heat transfer over wintertime leads and polynyas, *Journal of Geophysical Research: Oceans*, 104, 25 721–25 734, <https://doi.org/10.1029/1999JC900241>, 1999.
- 520 Bouchat, A. and Tremblay, B.: Reassessing the Quality of Sea-Ice Deformation Estimates Derived From the RADARSAT Geophysical Processor System and Its Impact on the Spatiotemporal Scaling Statistics, *Journal of Geophysical Research: Oceans*, 125, e2019JC015 944, <https://doi.org/https://doi.org/10.1029/2019JC015944>, 2020.
- Bouchat, A., Hutter, N., Chanut, J., Dupont, F., Dukhovskoy, D., Garric, G., Lee, Y. J., Lemieux, J.-F., Lique, C., Losch, M., Maslowski, W., Myers, P. G., Olason, E., Rampal, P., Rasmussen, T., Talandier, C., Tremblay, B., and Wang, Q.: Sea Ice Rheology Experiment (SIREx):
- 525 1. Scaling and Statistical Properties of Sea-Ice Deformation Fields, *Journal of Geophysical Research: Oceans*, 127, e2021JC017 667, <https://doi.org/10.1029/2021JC017667>, 2022.
- Dukowicz, J. K. and Baumgardner, J. R.: Incremental Remapping as a Transport/Advection Algorithm, *Journal of Computational Physics*, 160, 318 – 335, <https://doi.org/10.1006/jcph.2000.6465>, 2000.
- Griffies, S. M., Danabasoglu, G., Durack, P. J., Adcroft, A. J., Balaji, V., Böning, C. W., Chassignet, E. P., Curchitser, E., Deshayes, J.,
- 530 Drange, H., Fox-Kemper, B., Gleckler, P. J., Gregory, J. M., Haak, H., Hallberg, R. W., Heimbach, P., Hewitt, H. T., Holland, D. M., Ilyina, T., Jungclaus, J. H., Komuro, Y., Krasting, J. P., Large, W. G., Marsland, S. J., Masina, S., McDougall, T. J., Nurser, A. J. G., Orr, J. C., Pirani, A., Qiao, F., Stouffer, R. J., Taylor, K. E., Treguier, A. M., Tsujino, H., Uotila, P., Valdivieso, M., Wang, Q., Winton, M., and Yeager, S. G.: OMIP contribution to CMIP6: experimental and diagnostic protocol for the physical component of the Ocean Model Intercomparison Project, *Geoscientific Model Development*, 9, 3231–3296, <https://doi.org/10.5194/gmd-9-3231-2016>, 2016.
- 535 Haas, C.: Dynamics Versus Thermodynamics: The Sea Ice Thickness Distribution, chap. 4, pp. 113–151, John Wiley & Sons, Ltd, <https://doi.org/10.1002/9781444317145.ch4>, 2009.
- Holland, M. M., Bailey, D. A., Briegleb, B. P., Light, B., and Hunke, E.: Improved Sea Ice Shortwave Radiation Physics in CCSM4: The Impact of Melt Ponds and Aerosols on Arctic Sea Ice, *Journal of Climate*, 25, 1413 – 1430, <https://doi.org/h10.1175/JCLI-D-11-00078.1>, 2012.
- 540 Hunke, E. C. and Lipscomb, W. H.: CICE: the Los Alamos Sea Ice Model Documentation and Software User’s Manual Version 4.0 LA-CC-06-012, Tech. rep., Los Alamos National Laboratory, 2008.
- Hutter, N. and Losch, M.: Feature-based comparison of sea ice deformation in lead-permitting sea ice simulations, *The Cryosphere*, 14, 93–113, <https://doi.org/10.5194/tc-14-93-2020>, 2020.
- Inoue, J., Enomoto, T., Miyoshi, T., and Yamane, S.: Impact of observations from Arctic drifting buoys on the reanalysis of surface fields,
- 545 *Geophysical Research Letters*, 36, <https://doi.org/10.1029/2009GL037380>, 2009.
- Kwok, R., Schweiger, A., Rothrock, D. A., Pang, S., and Kottmeier, C.: Sea ice motion from satellite passive microwave imagery assessed with ERS SAR and buoy motions, *Journal of Geophysical Research: Oceans*, 103, 8191–8214, <https://doi.org/10.1029/97JC03334>, 1998.
- Kwok, R., Hunke, E. C., Maslowski, W., Menemenlis, D., and Zhang, J.: Variability of sea ice simulations assessed with RGPS kinematics, *Journal of Geophysical Research*, 113, <https://doi.org/10.1029/2008JC004783>, 2008.
- 550 Lemieux, J.-F., Tremblay, B., Sedláček, J., Tupper, P., Thomas, S., Huard, D., and Auclair, J.-P.: Improving the numerical convergence of viscous-plastic sea ice models with the Jacobian-free Newton-Krylov method, *Journal of Computational Physics*, 229, 2840 – 2852, <https://doi.org/10.1016/j.jcp.2009.12.011>, 2010.

- Lemieux, J.-F., Lipscomb, W., Craig, A., Bailey, D. A., Hunke, E., Blain, P., Rasmussen, T. A. S., Bentsen, M., Dupont, F., Hebert, D., and Allard, R.: CICE on a C-grid: new momentum, stress, and transport schemes for CICEv6.5, *Geoscientific Model Development Discussions*, 2024, 1–42, <https://doi.org/10.5194/gmd-2023-239>, 2024.
- Lindsay, R. W. and Stern, H. L.: The RADARSAT Geophysical Processor System: Quality of Sea Ice Trajectory and Deformation Estimates, *Journal of Atmospheric and Oceanic Technology*, 20, 1333–1347, [https://doi.org/10.1175/1520-0426\(2003\)020<1333:TRGPSQ>2.0.CO;2](https://doi.org/10.1175/1520-0426(2003)020<1333:TRGPSQ>2.0.CO;2), 2003.
- Marsan, D. and Weiss, J.: Space/time coupling in brittle deformation at geophysical scales, *Earth and Planetary Science Letters*, 296, 353–359, <https://doi.org/10.1016/j.epsl.2010.05.019>, 2010.
- Marsan, D., Stern, H., Lindsay, R., and Weiss, J.: Scale Dependence and Localization of the Deformation of Arctic Sea Ice, *Phys. Rev. Lett.*, 93, 178 501, <https://doi.org/10.1103/PhysRevLett.93.178501>, 2004.
- Murray, R. J.: Explicit Generation of Orthogonal Grids for Ocean Models, *Journal of Computational Physics*, 126, 251–273, <https://doi.org/10.1006/jcph.1996.0136>, 1996.
- Oikkonen, A., Haapala, J., Lensu, M., Karvonen, J., and Itkin, P.: Small-scale sea ice deformation during N-ICE2015: From compact pack ice to marginal ice zone, *Journal of Geophysical Research: Oceans*, 122, 5105–5120, <https://doi.org/10.1002/2016JC012387>, 2017.
- Perovich, D., Richter-Menge, J., Polashenski, C., Elder, B., Arbetter, T., and Brennick, O.: Sea ice mass balance observations from the North Pole Environmental Observatory, *Geophysical Research Letters*, 41, 2019–2025, <https://doi.org/10.1002/2014GL059356>, 2014.
- Rampal, P., Weiss, J., Marsan, D., Lindsay, R., and Stern, H.: Scaling properties of sea ice deformation from buoy dispersion analysis, *Journal of Geophysical Research: Oceans*, 113, <https://doi.org/10.1029/2007JC004143>, 2008.
- Rampal, P., Dansereau, V., Olason, E., Bouillon, S., Williams, T., Korosov, A., and Samaké, A.: On the multi-fractal scaling properties of sea ice deformation, *The Cryosphere*, 13, 2457–2474, <https://doi.org/10.5194/tc-13-2457-2019>, 2019.
- Rigor, I., Clemente-Colon, P., and Hudson, E.: The international Arctic buoy programme (IABP): A cornerstone of the Arctic observing network, *OCEANS 2008*, pp. 1–3, <https://doi.org/10.1109/OCEANS.2008.5152136>, 2008.
- Rigor, I. G., Wallace, J. M., and Colony, R. L.: Response of Sea Ice to the Arctic Oscillation, *Journal of Climate*, 15, 2648–2663, [https://doi.org/10.1175/1520-0442\(2002\)015<2648:ROSITT>2.0.CO;2](https://doi.org/10.1175/1520-0442(2002)015<2648:ROSITT>2.0.CO;2), 2002.
- Rothrock, D. A.: The energetics of the plastic deformation of pack ice by ridging, *Journal of Geophysical Research (1896-1977)*, 80, 4514–4519, <https://doi.org/10.1029/JC080i033p04514>, 1975.
- Schweiger, A., Lindsay, R., Zhang, J., Steele, M., Stern, H., and Kwok, R.: Uncertainty in modeled Arctic sea ice volume, *Journal of Geophysical Research: Oceans*, 116, <https://doi.org/10.1029/2011JC007084>, 2011.
- Small, R. J., Bacmeister, J., Bailey, D., Baker, A., Bishop, S., Bryan, F., Caron, J., Dennis, J., Gent, P., Hsu, H.-m., Jochum, M., Lawrence, D., Munoz, E., diNezio, P., Scheitlin, T., Tomas, R., Tribbia, J., Tseng, Y.-h., and Vertenstein, M.: A new synoptic scale resolving global climate simulation using the Community Earth System Model, *Journal of Advances in Modeling Earth Systems*, 6, 1065–1094, <https://doi.org/10.1002/2014MS000363>, 2014.
- Smith, G. C., Roy, F., Reszka, M., Surcel Colan, D., He, Z., Deacu, D., Belanger, J.-M., Skachko, S., Liu, Y., Dupont, F., Lemieux, J.-F., Beaudoin, C., Tranchant, B., Drevillon, M., Garric, G., Testut, C.-E., Lellouche, J.-M., Pellerin, P., Ritchie, H., Lu, Y., Davidson, F., Buehner, M., Caya, A., and Lajoie, M.: Sea ice forecast verification in the Canadian Global Ice Ocean Prediction System, *Quarterly Journal of the Royal Meteorological Society*, 142, 659–671, <https://doi.org/10.1002/qj.2555>, 2016.
- Sumata, H., de Steur, L., Divine, D. V., Granskog, M. A., and Gerland, S.: Regime shift in Arctic Ocean sea ice thickness, *Nature*, 615, 443–449, <https://doi.org/10.1038/s41586-022-05686-x>, 2023.

- Tsamados, M., Feltham, D. L., and Wilchinsky, A. V.: Impact of a new anisotropic rheology on simulations of Arctic sea ice, *Journal of Geophysical Research: Oceans*, 118, 91–107, <https://doi.org/10.1029/2012JC007990>, 2013.
- Turner, A. K., Hunke, E. C., and Bitz, C. M.: Two modes of sea-ice gravity drainage: A parameterization for large-scale modeling, *Journal of Geophysical Research: Oceans*, 118, 2279–2294, <https://doi.org/10.1002/jgrc.20171>, 2013.
- 595 Vancoppenolle, M., Rousset, C., Blockley, E., Aksenov, Y., Feltham, D., Fichefet, T., Garric, G., Guemas, V., Iovino, D., Keeley, S., Madec, G., Massonnet, F., Ridley, J., Schroeder, D., and Tietsche, S.: SI3, the NEMO Sea Ice Engine, <https://doi.org/10.5281/zenodo.7534900>, 2023.
- Wang, Q., Ilicak, M., Gerdes, R., Drange, H., Aksenov, Y., Bailey, D. A., Bentsen, M., Biastoch, A., Bozec, A., Boning, C., Cassou, C., Chassignet, E., Coward, A. C., Curry, B., Danabasoglu, G., Danilov, S., Fernandez, E., Fogli, P. G., Fujii, Y., Griffies, S. M., Iovino, D.,  
600 Jahn, A., Jung, T., Large, W. G., Lee, C., Lique, C., Lu, J., Masina, S., Nurser, A. G., Rabe, B., Roth, C., Salas y Melia, D., Samuels, B. L., Spence, P., Tsujino, H., Valcke, S., Voldoire, A., Wang, X., and Yeager, S. G.: An assessment of the Arctic Ocean in a suite of interannual CORE-II simulations. Part I: Sea ice and solid freshwater, *Ocean Modelling*, 99, 110–132, <https://doi.org/h10.1016/j.ocemod.2015.12.008>, 2016.
- Weiss, J.: Intermittency of principal stress directions within Arctic sea ice, *Phys. Rev. E*, 77, 056106,  
605 <https://doi.org/10.1103/PhysRevE.77.056106>, 2008.
- Xu, S., Ma, J., Zhou, L., Zhang, Y., Liu, J., and Wang, B.: Comparison of sea ice kinematics at different resolutions modeled with a grid hierarchy in the Community Earth System Model (version 1.2.1), *Geoscientific Model Development*, 14, 603–628, <https://doi.org/10.5194/gmd-14-603-2021>, 2021.
- Yang, C.-Y., Liu, J., and Xu, S.: Seasonal Arctic Sea Ice Prediction Using a Newly Developed Fully Coupled Regional Model  
610 With the Assimilation of Satellite Sea Ice Observations, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS001938, <https://doi.org/10.1029/2019MS001938>, 2020.
- Zhang, S., Xu, S., Fu, H., Wu, L., Liu, Z., Gao, Y., Wan, W., Xie, J., Wan, L., Lu, H., Duan, J., Liu, Y., Fei, Y., Guo, X., Wang, Z., Wang, X., Wang, Z., Qu, B., Li, M., Zhao, H., Jiang, Y., Yang, G., Wang, H., An, H., Zhang, X., Zhang, Y., Ma, W., Yu, F., Xu, J., and Shen, X.:  
615 Toward Earth System Modeling with Resolved Clouds and Ocean Submesoscales on Heterogeneous Many-Core HPCs, *National Science Review*, <https://doi.org/10.1093/nsr/nwad069>, 2023.