



AI-NAOS: An AI-Based Nonspherical Aerosol Optical Scheme for Chemical Weather Model GRAPES_Meso5.1/CUACE

Xuan Wang¹, Lei Bi¹, Hong Wang², Yaqiang Wang², Wei Han³, Xueshun Shen³, Xiaoye Zhang²

¹Key Laboratory of Geoscience Big Data and Deep Resource of Zhejiang Province, School of Earth Sciences, Zhejiang University, Hangzhou, China

²Key Laboratory of Atmospheric Chemistry of CMA, Institute of Atmospheric Composition, Chinese Academy of Meteorological Sciences, Beijing, China

³CMA Earth System Modeling and Prediction Centre (CEMC), China Meteorological Administration, Beijing, China

Correspondence to: Lei Bi (bilei@zju.edu.cn)

Abstract The AI-based Nonspherical Aerosol Optical Scheme (AI-NAOS) is a newly developed aerosol optical module that improves the representation of aerosol optical properties for radiative transfer simulations in atmospheric models. It incorporates the nonsphericity and inhomogeneity (NSIH) of internally mixed aerosol particles through a deep learning method. Specifically, the AI-NAOS considers black carbon (BC) as fractal aggregates and models soil dust (SD) as superspheroids, encapsulated partially or completely with hygroscopic aerosols such as sulfate, nitrate, and aerosol water. To obtain AI-NAOS, a database of the optical properties for the models was constructed using the invariant imbedding T-matrix method (IITM), and deep neural networks (DNN) were trained based on this database. In this study, the AI-NAOS was integrated into the mesoscale version 5.1 of Global/Regional Assimilation and Prediction System with Chinese Unified Atmospheric Chemistry Environment (GRAPES_Meso5.1/CUACE). Real-case simulations were conducted during a winter with high pollution, comparing BC aerosols evaluated using three schemes with spherical aerosol models (external-mixing, core-shell, and volume-mixing) and the AI-NAOS scheme. The results showed that NSIH effect led to a moderate estimation of absorbing aerosol optical depth (AAOD) and obvious changes in aerosol radiative effects, short-wave heating rates, temperature profiles, and boundary layer height. The AAOD values based on three spherical schemes were 70.4%, 125.3%, and 129.3% over Sichuan Basin, benchmarked to the AI-NAOS results. Compared to the external-mixing scheme, the direct radiative effect (DRE) induced by the NSIH effect reached +1.6 W/m² at the top-of-atmosphere (TOA) and -2.9 W/m² at surface. The NSIH effect could enhance the short-wave heating rate, reaching 20%. Thus, the warming effect at 700 hPa and the cooling effect on the ground were strengthened by 21% and 13%, reaching +0.04 and -0.10 K, which led to a reduction in the height of the Planetary Boundary Layer (PBL) by -11 meters. In addition, the precipitation was inhibited by the NSIH effect, causing a 15% further decrease. Therefore, the NSIH effects demonstrated their non-negligible impacts and highlighted the importance of incorporating them into chemical weather models.



30 1 Introduction

Aerosols play a significant role in the Earth's climate system through various pathways, including direct scattering and absorbing radiation, and also indirectly, by affecting cloud formation and dimming snow, hence influencing the energy budget and climate processes. However, larger uncertainties regarding aerosol radiative forcing still exist due to challenges in identifying their distribution, transportation, and physical properties (Yu et al., 2006). The direct radiative effect (DRE) of aerosols, which refers to the radiation changes caused by aerosol absorption and scattering, is the primary driver of aerosol–climate interactions. In the weather scale, the aerosol-radiation interaction (ARI) is quite important for predicting air quality, as it can significantly alter the meteorology, affecting visibility and PM_{2.5} concentrations (Wang et al., 2022a; Peng et al., 2022). Including ARI in mesoscale weather forecasts is essential for more accurate radiation flux estimation, particularly during dust storms or haze episodes (Wang et al., 2010, 2015a, b).

Aerosol optical properties, namely, the extinction coefficient (q_{ext}), single scattering albedo (ω), and asymmetry factor (g), are key parameters for understanding direct aerosol–radiation interactions in weather/climate models. These optical properties of aerosols vary due to multiple factors, including wavelength, complex refractive index, and particle size and morphology. Among these factors, morphological features are particularly challenging to address and can be divided into two aspects: (1) nonsphericity and (2) inhomogeneity. In general, simplified spherical models are commonly used to represent hygroscopic aerosol particles such as sulfate, nitrate, and ammonium, which is consistent with observations in both the field and laboratory (Buseck and Pósfai, 1999; Wise et al., 2007). However, insoluble aerosols like BC and mineral dust aerosols are nonspherical (Adachi et al., 2007; Pósfai et al., 2013). Additionally, BC and mineral dust aerosols are not always homogeneous, as they can mix with other particles during transport, resulting in embedded aerosols within coating materials (Adachi et al., 2010).

Several electromagnetic scattering algorithms are available for calculating the optical properties of aerosol particles with different morphological features. The Lorenz–Mie theory is widely applied for calculating optical properties of homogenous spherical particles (Bohren and Huffman, 1998). The multiple sphere T-matrix method (MSTM) is an implementation of a generalized Mie theory and is suitable for analyzing multiple sphere domains, such as coated spheres and fractal aggregates with spherical monomers (Mackowski, 2014). The discrete dipole approximation (DDA) method is a general scattering algorithm for particles of arbitrary geometry and composition, in which the scatterer is divided into small cubical subvolumes (Yurkin and Hoekstra, 2011). The invariant imbedding T-matrix method (IITM) is a more efficient algorithm for analyzing randomly oriented nonspherical and inhomogeneous particles, which are discretized in terms of multiple inhomogeneous spherical layers (Bi and Yang, 2014; Bi et al., 2013, 2022; Wang et al., 2023c).

The aerosol optical parameterization scheme in climate/weather models is closely tied to the capabilities of electromagnetic scattering algorithms. Spherical aerosol models are commonly used in climate/weather models due to the relatively low computational time required by the Lorenz–Mie theory. Examples of spherical aerosol models include (1) the external-mixing model, where each kind of aerosols is assumed to be a homogeneous sphere, (2) the volume-mixing model,



where all species are homogeneously mixed, (3) the Maxwell-Garnet model, where isolated spherules are suspended in an embedding sphere, and (4) the core-shell model for concentrically coated particles (Bond and Bergstrom, 2006; Fast et al., 2006; Kotchenova and Vermote, 2007). For example, in Weather Research and Forecasting model coupled with Chemistry (WRF/Chem), multiple aerosol optical schemes are included, such as the volume-mixing, the Maxwell-Garnet, and the core-shell models (Barnard et al., 2010). In the mesoscale version 5.1 of Global/Regional Assimilation and Prediction System coupled with Chinese Unified Atmospheric Chemistry Environment model (GRAPES_Meso5.1/CUACE), only the external-mixing model is included (Gong and Zhang, 2008; Wang et al., 2018). As a climate model, the Community Atmosphere Model version 5 (CAM5) incorporates both volume-mixing and core-shell treatments in Aerosol Two-dimensional bin module for formation and Aging Simulation (ATRAS) model using look-up tables (Matsui, 2017).

Implementing electromagnetic scattering algorithms in climate and weather models is unrealistic due to the prohibitively long computational time. Therefore, look-up tables with pre-computed optical properties are typically used to incorporate morphologically realistic aerosol models in optical parameterization, reducing computational costs. For example, bare fractal aggregates model and core-grey-shell model have been applied to address the nonsphericity and inhomogeneity of BC using look-up tables in the Multi-scale Atmospheric Transport and Chemistry model (MATCH) and the Sectional Aerosol module for Large Scale Applications (SALSA) (Andersson and Kahnert, 2016). Recently, deep learning has emerged as a new method to handle the optical properties of complex internal mixing models. Wang et al. (2022b) incorporated coated super-spheroid models for mineral aerosols into the WRF/chem model using a fully connected neural network (Wang et al., 2022b).

In this study, we developed a comprehensive AI-based nonspherical aerosol optical scheme (AI-NAOS) to investigate the direct aerosol radiation effects by employing morphologically realistic models based on the deep learning method. In this scheme, insoluble aerosols, such as soil dust (SD) and BC, were both treated as nonspherical cores that are partially or fully embedded in a spherical or nonspherical shell consisting of hygroscopic aerosols. Specifically, we modelled dust as super-spheroids, which have been proved successful in matching the optical properties of dust samples and the polarized radiance observations from Polarisation and Anisotropy of Reflectances for Atmospheric Science coupled with Observations from a Lidar (PARASOL) (Lin et al., 2018, 2021). BC was assumed to exist as fractal aggregates that could accurately reproduce the measured linear backscattering depolarization ratio (Luo et al., 2019; Wang et al., 2023b). We developed optical property databases of these two models using the IITM method. Subsequently, several deep neural networks (DNN) were trained based on the aforementioned databases and then integrated into the AI-NAOS module. The AI-NAOS scheme was online-coupled with the chemical weather models, including WRF/chem and GRAPES_Meso5.1/CUACE.

In the remaining sections of this paper, we will present the details of the AI-NAOS module in Section 2. We will analyze real-case evaluation performed by the GRAPES_Meso5.1/CUACE model in Section 3. Finally, in Section 4, we will summarize and discuss the conclusions.



2 Methods

2.1 Aerosol optical modelling

In the aerosol module of CUACE model, there are seven aerosol species, including two insoluble aerosols, black carbon and soil dust, and five hygroscopic aerosols, such as organic carbon, sulfate, sea salt, nitrate, and ammonium. The optical properties of these aerosols are accessed through a series of look-up tables, where complex refractive indices are implicitly represented. In the CUACE model, the refractive indices of SD were obtained from the Aeolian dust experiment on climate (ADEC), while the high-resolution transmission molecular absorption database (HITRAN) were applied to represent refractive indices of other aerosols (Rothman et al., 2005; Wang et al., 2006). In the aerosol optical modelling process, aerosol particles of each species were simplified as spheres and completely dissolved in aerosol water, which undergo hygroscopic growth, and then mix externally. This aerosol scheme is a typical external-mixing scheme in which aerosol particles are always assumed to be spherical and homogeneous.

As an update, the AI-NAOS module incorporates improvements related to nonsphericity and inhomogeneity into the aerosol scheme. Additionally, two widely used internal-mixing schemes, the volume-mixing and core-shell methods, have also been integrated into the aerosol module to facilitate a better understanding of the differences between aerosol schemes with various optical models.

Two nonspherical models, fractal aggregates and super-spheroids, were incorporated to represent insoluble aerosols, while hygroscopic aerosols were treated as a coating shell to introduce inhomogeneity. Specifically, the fractal aggregate model composed of identical spheres was found to be a good fit for the morphology of BC particles. The fractal law was used to characterize aggregates, shown below (Sorensen, 2001; Teng et al., 2019):

$$N_s = k_0 \left(\frac{R_g}{R} \right)^{D_f}, \quad (1)$$

In Equation 1, D_f is the fractal dimension, k_0 is the scaling prefactor, N_s is the number of spherical monomer, R is the radius of monomer, and R_g is the radius of gyration (Sorensen, 2011). The fractal dimension and the scaling prefactor are major factors influencing the compactness of fractal structure, which is connected with the aging progress of BC (Forrest and Witten, 1979; Li et al., 2016). Tuning a smaller scaling prefactor or fractal dimension allows for the creation of a looser and more linear structure, which shows better fitness for newly emitted BC. Conversely, a larger scaling prefactor and fractal dimension is more suitable for aged BC after transportation. Generally, the fractal dimension ranges from 1.8 to 2.8 (Wang et al., 2017; Wu et al., 2023). In this study, the fractal dimension was set at 2.1, and the scaling prefactor was set at 1.2. A tuneable algorithm called FracVAL was used to generate this fractal aggregates (Morán et al., 2019). The refractive index of BC was assumed to be $1.95 + 0.79i$ (Bond and Bergstrom, 2006); however, this value can be flexibly adjusted in the module. All hygroscopic aerosols were treated as a homogenous spherical coating shell using the volume-mixing method. The center




of the spherical coating shell was located at the midpoint of the major axis of the fractal aggregates, which was defined based on two spherical monomers with the greatest separations, partially embedding the fractal framework as shown in Figure 1a.

A simplified equation of super-ellipsoid (referred to as super-spheroid) was used for geometric modelling (Barr, 1981; Bi et al., 2018a):

$$130 \quad \left(\frac{x}{a}\right)^{\frac{2}{n}} + \left(\frac{y}{a}\right)^{\frac{2}{n}} + \left(\frac{z}{c}\right)^{\frac{2}{n}} = 1, \quad (2)$$

in which n is the roundness parameter and a/c is the aspect ratio. In this case, the roundness parameter was fixed at 2.6, which was suitable for optical modelling of SD (Lin et al., 2018). The coating shell was also modelled as a super-spheroid with the roundness parameter ranging from 2.6 to 1.0, for which the core was fully embedded as shown in Figure 1b.



135 **Figure 1. Optical modelling for (a) fractal aggregates framework of black carbon (BC) partially encapsulated with spherical coating of hygroscopic aerosols and (b) super-spheroid framework of soil dust (SD) fully coated with another super-spheroid of hygroscopic aerosols with various volume fractions.**

2.2 Database of optical properties

140 Next, we discuss optical property databases that have been constructed for nonspherical and inhomogeneous particles, specifically encapsulated BC fractal aggregates and coated SD super-spheroids. These databases utilized six parameters to describe the microphysics of a certain particle, including size parameter, volume fraction (vf), and complex refractive indices of both insoluble aerosols and hygroscopic aerosols. Three optical properties, namely, extinction efficiency (q_{ext}), single scattering albedo (ω), and asymmetry factor (g) were determined using these parameters. The volume fraction is a key parameter that describes the mixing status and is defined as shown below:

$$145 \quad \text{vf} = \frac{V_{\text{insol}}}{V_{\text{hygro}} + V_{\text{insol}}}, \quad (3)$$



in which V_{insol} is the volume of insoluble aerosols, and V_{hygro} is the volume of hygroscopic aerosols in a single particle. The size parameter (x) is a size defined relative to wavelength and is given by:

$$x = \frac{\pi D}{\lambda}, \quad (4)$$

150 in which λ is the wavelength, and D is the diameter. For nonspherical particles, the actual diameter is the length of the major axis, which can be determined by two monomers with the longest distance in fractal aggregates and the longest dimension in super-spheroids. Accounting for the convenience of matching with spherical models, the diameter defined in Equation 4 was calculated based on the equi-volume sphere. As shown in Table 1, the size parameter exceeds 20, which is sufficient to handle particles with diameters less than 4 μm even in photosynthetically active radiative bands. The complex refractive indices for hygroscopic aerosols and two insoluble aerosols, BC and SD, are denoted by mr_{hygro} and mr_{insol} for the real
 155 part and mi_{hygro} and mi_{insol} for the imaginary part. The volume fraction of insoluble aerosols ranges from 0.04 to 1.0 for BC aerosols and from 0.15 to 1.0 for SD aerosols. These parameters are summarized in Table 1.

Table 1. Optical property database of nonspherical and inhomogeneous (NSIH) particles. Black carbon (BC) was modelled as encapsulated fractal aggregates while soil dust (SD) was assumed to be a coated super-spheroid.

	coated fractal aggregates	coated super-spheroids
mr_{hygro}	1.20, 1.25, 1.30, 1.35, 1.40, 1.45, 1.50, 1.55, 1.60	1.2, 1.3, 1.4, 1.5, 1.6, 1.8
mi_{hygro}	0, 0.001, 0.005, 0.01, 0.05, 0.1	10^{-5} , 10^{-4} , 0.001, 0.005, 0.01, 0.05, 0.1, 0.5
vf(%)	3.66, 5.42, 8.46, 11.79, 16.77, 24.3, 35.82, 43.28, 52.03, 61.74, 71.57, 80.78, 88.89, 94.99, 100	14.68, 18.06, 22.51, 28.35, 36.02, 46.11, 59.38, 76.88, 100
mr_{insol}	1.65, 1.75, 1.85, 1.95	1.2, 1.4, 1.5, 1.6
mi_{insol}	0.5, 0.6, 0.7, 0.8	0.0005, 0.001, 0.005, 0.01, 0.05, 0.1, 0.5
x	0.1-21.0	0.05-26.4

2.3 Bulk optical properties

160 In the aerosol module of the CUACE, aerosol particles are categorized into 12 size bins based on their diameter, ranging from 0.01 μm to 40.96 μm . When utilizing this model, it is necessary to integrate the optical properties over the diameter within each size bin to obtain bulk optical properties that are more representative, as the oscillation of optical properties along with the particle size occurs. To achieve this, we used the log-normal size distribution for each size bin, ranging from the lower bound D_{min} to the upper bound D_{max} . The probability density function (PDF) of log-normal
 165 distribution is defined as shown below:

$$n(D) = \frac{1}{D\sqrt{2\pi}\sigma} e^{-\frac{(\ln D - \ln D_m)^2}{2\sigma^2}}, \quad (5)$$



in which σ is the standard deviation, and D_m is the mean diameter. Based on the PDF, bulk optical properties can be calculated by integration as shown below:

$$Q_{\text{ext}} = \frac{\int_{D_{\text{min}}}^{D_{\text{max}}} q_{\text{ext}} D^2 n(D) dD}{\int_{D_{\text{min}}}^{D_{\text{max}}} D^2 n(D) dD}, \quad (6)$$

$$170 \quad \langle \text{SSA} \rangle = \frac{\int_{D_{\text{min}}}^{D_{\text{max}}} \omega q_{\text{ext}} D^2 n(D) dD}{\int_{D_{\text{min}}}^{D_{\text{max}}} q_{\text{ext}} D^2 n(D) dD}, \quad (7)$$

$$\langle G \rangle = \frac{\int_{D_{\text{min}}}^{D_{\text{max}}} g \omega q_{\text{ext}} D^2 n(D) dD}{\int_{D_{\text{min}}}^{D_{\text{max}}} \omega q_{\text{ext}} D^2 n(D) dD}, \quad (8)$$

in which Q_{ext} , $\langle \text{SSA} \rangle$ and $\langle G \rangle$ are the bulk extinction efficiency, bulk single scattering albedo and bulk asymmetry factor, respectively.

The bulk optical properties were compared and are shown in Figure 2 for different aerosol optical schemes, including the external-mixing, volume-mixing, core-shell, and AI-NAOS modules. The AI-NAOS module contained encapsulated fractal aggregates model of BC and coated super-spheroids of SD. The calculations of bulk optical properties were performed considering the presence of hygroscopic aerosols and either BC or SD as the insoluble aerosol species. The volume fraction of insoluble aerosols was assumed to be 0.33. The complex refractive indices of BC, SD, and hygroscopic aerosols were $1.95+0.79i$, $1.5+0.01i$, and $1.4+0.0001i$, respectively. Generally, BC aerosols are small particles with diameters around $0.2 \mu\text{m}$, and size parameters around 0.9–1.6 for photosynthetically active radiative bands (Wu et al., 2023). As shown in the top row of Figure 2, the extinction efficiency of the AI-NAOS scheme was smaller compared to the core-shell scheme and the volume-mixing scheme, but larger compared to the external-mixing scheme. As for SSA, it could be observed that the external-mixing scheme yielded the largest results, while the other three schemes were close. It is likely that the AI-NAOS scheme led to improvements in the estimation of BC absorption, as the external-mixing scheme underestimated BC absorption and the volume-mixing scheme overestimated this parameter (Kahnert et al., 2012). The asymmetry factor calculated by the AI-NAOS was slightly higher than other schemes, leading to a smaller backscattering fraction. With respect to SD aerosol, the aerosols' diameters were much larger, so we focus on the part where the size parameter was larger than 10.0, as shown in the bottom row in Figure 2. The extinction efficiency based on the AI-NAOS scheme was larger, and the SSA was similar compared to other schemes for large SD particles, indicating stronger absorption due to the NSIH effect of the SD. The asymmetry factor of the AI-NAOS scheme was the smallest.




Figure 2. Optical properties of (a-c) black carbon (BC) and (d-f) soil dust (SD) with four optical modelling schemes (external-mixing, AI-NAOS, core-shell and volume-mixing scheme).

2.4 Deep Neural Networks

195 Aerosol optical parameterizing is a complex task that involves various species and multiple processes, such as aerosol
mixture and hygroscopic growth, that need to be addressed. In the past, the look-up table method was widely used for this
purpose, but it had limitations in terms of simplifying some processes and consuming a relatively large amount of storage
space. To overcome these challenges, a deep learning method can be applied. A deep neural network (DNN) can efficiently
and accurately provide continuous predictions while requiring less storage space compared to a look-up table (Yu et al.,
200 2022a, b). Additionally, the DNN method eliminates the need for multiple interpolations among various dimensions, making
it easier to compute the optical properties at the particle size, volume fraction, and complex refraction indices based on the
real-time status of aerosols. A multiple-target DNN model has been applied to infer single scattering properties of
encapsulated fractal aggregates of BC (Wang et al., 2023b). In this study, four DNNs have been trained based on bulk optical
property databases of spherical and NSIH particles using this DNN architecture, including spheres, core-shell spheres,
205 encapsulated fractal aggregates and coated super-spheroids. Note that, the present DNNs are different from those in previous



studies, in which DNNs are obtained in terms of the single-scattering properties for general application purposes. More details regarding the DNNs are provided in the Appendix.

Figure 3 illustrates the performance of DNNs for these two NSIH models, and three statistical metrics, namely, mean absolute error (MAE), root mean square error (RMSE), and coefficient of determination (R-squared), were used to evaluate the DNN. Approximately 100,000 samples were randomly selected from the database for each optical property of each NSIH model. The scatter points in Figure 3 represent the evaluation results of the DNN predictions and IITM calculations, with the colour denoting the probability density scaled from 0 to 1. For coated fractal aggregates of BC, the RMSE of bulk extinction efficiency, bulk SSA, and bulk asymmetry factor were 0.0114, 0.0017, and 0.0013, respectively. For coated super-spheroids of SD, the RMSE values were 0.0240, 0.0062, and 0.0052, respectively. The R-squared values for all three optical properties were above 0.99, indicating a good fit between the true values from the IITM calculations and the predicted values from the DNNs. Overall, the optical properties of the NSIH models can be accurately predicted using the DNN models.




Figure 3. Comparison of optical properties calculated using the invariant imbedding T-matrix method (IITM) and predictions from deep neural networks (DNNs). (a–c) encapsulated fractal aggregate model for black carbon. (d–f) coated super-spheroid model for soil dust. The probability density is scaled from 0 to 1.

2.5 Nonspherical Aerosol Optical Scheme

A basic process in the AI-NAOS module is the determination of particles states, including the volume of each kind of particle, volume fraction of mixed particles, and the morphology of particles. Coupled with a size-segregated aerosol module, the particle states are independent among every size bin. In a specific bin, all hygroscopic aerosols are first



225 internally mixed with aerosol water based on a volume-mixing assumption and then encapsulated other two insoluble
aerosol, respectively. The key issue is to determine the amount of hygroscopic aerosols coating, i.e., the volume fraction.
The simplest method is to make all hygroscopic aerosols be the coating, like the core-shell scheme, which means the volume
fraction ranges from zero to one. A more appropriate method is to consider a lower limit of the volume fraction, based on
worldwide field observations (Wang et al., 2023a). Specifically, the lower limit of the volume fraction was set to be 0.3 for
230 BC and 0.6 for SD. Thus, not all hygroscopic aerosols were internally mixed with insoluble aerosols when the volume of
hygroscopic aerosols was large enough. These extra mixed hygroscopic aerosols were assumed to be homogenous spheres.
Due to hygroscopic growth, the diameters of different hygroscopic aerosol species were also different. Similar to the core-
shell method, the diameter of internally mixed particle was determined using the volume-weighted method. In this way, all
seven aerosol species could be treated as three kinds of internally-mixed particles, including two nonspherical and
235 inhomogeneous particles (partially encapsulated fractal aggregates of BC, and fully coated super-spheroid of SD) and the
homogeneous and spherical particle of hygroscopic aerosols.

Once the state of aerosols was determined, the optical properties of three internally-mixed aerosol particles were
inferred by the DNNs integrated in the AI-NAOS module. The total optical properties were calculated based on the external-
mixing assumption, in which extinction and scattering coefficients were the sum of each species. This implementation of the
240 AI-NAOS module provided an efficient estimation of optical properties for nonspherical and inhomogeneous particles.
Generally, three optical properties (extinction coefficient, single-scattering albedo, and asymmetry factor) were considered
sufficient for radiation transfer calculation in the chemical weather model.

3 Model evaluation

3.1 Model configuration

245 The AI-NAOS module was on-line coupled with chemical weather model GRAPES_Meso5.1/CUACE. In a previous
study, the NSIH effect of SD had been evaluated using the WRF/chem model, revealing that the nonsphericity of SD could
cause more surface dimming and more solar heating (Wang et al., 2022b). At that time, BC was still assumed to be spherical
and treated using the core-shell model. Thus, this study primarily focuses on the NSIH effects of BC, where various models
of BC were compared while SD was always assumed to be super-spheroids. Real-case simulations were conducted using
250 GRAPES_Meso5.1/CUACE to gain a better understanding of the impact of aerosols on the weather. Five different aerosol
treatments were applied in the simulations, including a control experiment without aerosol optical effects, the AI-NAOS
scheme, the core-shell scheme with spherical but inhomogeneous model, and the external-mixing and the volume-mixing
schemes for the spherical and homogeneous model. To assess the NSIH effect from the BC aerosol, different models were
applied to only the BC aerosols, while the SD aerosols were modelled as super-spheroids among all schemes to avoid the
255 influence due to SD. The AI-NAOS scheme was directly connected with the shortwave radiation scheme (Goddard Space



Flight Center Scheme); namely, the primary driver of NSIH effect on aerosol–weather interactions was a perturbation in the solar radiation.

260 Real-scene studies were conducted to investigate the NSIH effect on the thermodynamic structure of the atmosphere under a heavy pollution scenario. The case was carried out on January 12, 2018, running for 72 hours, with the last 48 hours used for evaluation. The simulation domain covered East China (95–125°E, 20–50°N) with a horizontal resolution of $0.1^\circ \times 0.1^\circ$. The evaluations mainly focused on three regions with high emissions: (1) Sichuan Basin (103.5–108.5°E, 28.5–32.5°N), (2) Middle Yangtze Plain (110.5–114.5°E, 27.5–32.5°N), and (3) North China Plain (114–120°E, 33–39°N). The initial meteorological field and lateral boundary conditions were constructed using the NCEP FNL data on 0.25×0.25 grids (National Centers For Environmental Prediction/National Weather Service/NOAA/U.S. Department Of Commerce, 2015).
265 The Multi-resolution Emission Inventory model for Climate and air pollution research (MEIC) was used for chemical emissions (Li et al., 2017).

The configurations of physical and chemical processes in GRAPES_Meso5.1/CUACE simulations are summarized in Table 2.

Table 2. Physical and chemical processes in the GRAPES_Meso5.1/CUACE.

Process	Scheme
Microphysics	Thompson (Thompson et al., 2004)
Shortwave radiation	Goddard (Chou et al., 1998)
Longwave radiation	RRTM (Mlawer et al., 1997)
Surface layer	SFCLAY (Pleim, 2007)
Land surface	Noah (Chen et al., 1997)
Planetary boundary layer	MRF (Hong and Pan, 1996)
Cumulus	KFETA (Kain and Fritsch, 1993)
Gas-phase chemistry	RADM2 (Stockwell et al., 1990)
Aerosol	CUACE (Zhou et al., 2012)
Aerosol optics	<ol style="list-style-type: none">1. NA (no aerosol optical effect)2. external-mixing3. AI-NAOS4. core-shell5. volume -mixing

270

3.2 Aerosol optical properties

It is essential to evaluate three bulk aerosol optical properties (Q_{ext} , $\langle \text{SSA} \rangle$, and $\langle G \rangle$), which are refined directly in the AI-NAOS module. To effectively display these properties of the vertical atmospheric column, the column aerosol optical



properties, which include aerosol optical depth (AOD), absorbing aerosol optical depth (AAOD), and column asymmetry
275 factor (G), are defined as follows (Taylor, 2012):

$$b_{\text{ext}} = \sum_{i=1}^{12} \sum_{j=1}^3 Q_{\text{ext}ij} \frac{\pi}{4} D_{ij}^2 N_{ij}, \quad (9)$$


$$\text{AOD} = \int_{z=0}^{H_{\text{TOA}}} b_{\text{ext}} dz, \quad (10)$$

$$\text{AAOD} = \int_{z=0}^{H_{\text{TOA}}} b_{\text{ext}} (1 - \langle \text{SSA} \rangle) dz, \quad (11)$$

$$G = \frac{\int_{z=0}^{H_{\text{TOA}}} b_{\text{ext}} \langle \text{SSA} \rangle \langle G \rangle dz}{\int_{z=0}^{H_{\text{TOA}}} b_{\text{ext}} \langle \text{SSA} \rangle dz}, \quad (12)$$

280 where i is the index of size bin, j is the index of aerosol particle, N is the number concentration of aerosol particles, b_{ext} is the extinction coefficient, z is the height above ground, and H_{TOA} is the height of TOA.

Figure 4 illustrates three column optical properties at $0.55 \mu\text{m}$ based on the AI-NAOS scheme, as well as anomalies based on the other three aerosol optical schemes. All properties were averaged over 48 hours from the 25th to the 72th hour to yield a general picture. Three distinct centers with extreme AOD values, Sichuan Basin, Middle Yangtze Plain, and North
285 China Plain, are marked with black rectangles from west to east. These three regions also stood out as major sources of aerosol emissions. The North China Plain was characterized by intense industrial manufacturing while residential activities and transportation played prominent roles in the Sichuan Basin and Middle Yangtze Plain compared to the surrounding areas. Within the AI-NAOS module, the spatial average of AOD was 0.29, 0.37, and 0.36 over the Sichuan Basin, Middle
290 Yangtze Plain and North China Plain, respectively. The AOD values were larger compared to the external-mixing scheme but slightly smaller compared to the core-shell and volume mixing schemes. With respect to AAOD, the values were found to be 0.06, 0.08, and 0.08 averaged over the three marked regions, from west to east, respectively. Taking the results under the AI-NAOS module as a benchmark, the AAOD values based on three spherical schemes (the external-mixing, core-shell and volume-mixing scheme) were 70.4%, 125.3%, and 129.3% over Sichuan Basin, 70.0%, 119.6%, and 126.1% over the Middle Yangtze Plain and 70.4%, 116.0%, and 124.4% over the North China Plain, respectively. The column asymmetry
295 factor is illustrated in the bottom panel of Figure 4. The values based on the AI-NAOS module was 0.66, 0.64, and 0.63 averaged over the three marked regions, which was roughly +0.03, +0.03, and +0.01 larger compared to the three spherical schemes.



300 **Figure 4. Aerosol column optical properties based on the AI-NAOS scheme and anomalies based on three spherical schemes (external-mixing, core-shell, and volume-mixing).**

The spatially averaged column optical properties of the three spherical schemes, all of which were benchmarked to values under the AI-NAOS module, are summarized in Table 3. It is clear that there was minimal variation in the asymmetry factor, whereas a significant disparity existed in the AAOD. The AAOD, indicating the aerosol absorption, is greatly affected by aerosol optical modelling. Specifically, the external-mixing scheme, lacking both nonspherical and inhomogeneous effects, estimated notably weaker absorption, while the core-shell model, only considering inhomogeneous

305




effects, estimated stronger absorption. The AI-NAOS module is anticipated to correctly deal with the aerosol absorption, yielding a moderate estimation.

Table 3. Aerosol optical properties of three spherical schemes averaged over three marked regions, benchmarked to values under the AI-NAOS module.

Optical Properties	Regions	External-mixing	Core-shell	Volume-mixing
AOD	Sichuan Basin	91.7%	104.9%	107.0%
	Middle Yangtze Plain	91.7%	105.1%	108.0%
	North China Plain	92.3%	104.8%	108.4%
AAOD	Sichuan Basin	70.4%	125.3%	129.3%
	Middle Yangtze Plain	70.0%	119.6%	126.1%
	North China Plain	71.4%	116.0%	124.4%
G	Sichuan Basin	96.2%	94.9%	98.8%
	Middle Yangtze Plain	96.0%	94.8%	98.3%
	North China Plain	96.2%	95.4%	98.0%

310 The absorption enhancement (E_{abs}), defined as the ratio of absorption section of coated and bare BC particles, is an important metric of aerosol absorption. A significant issue identified is that field observations typically reveal smaller E_{abs} values than those predicted by model simulations. To mitigate this discrepancy, a series of modifications aimed at managing the E_{abs} value were conducted, drawing on microphysical complexities such as coating thickness and morphology (Chen et al., 2023; Huang et al., 2023). However, these strategies were rooted in the core-shell model, utilizing look-up tables or
 315 empirical formulas without accounting for aerosol nonsphericity. This limitation potentially leads to inaccuracies in estimations of aerosol absorption.

320 Figure 5 illustrates the distribution of absorption enhancement and single scattering albedo over the Sichuan Basin. The E_{abs} value was the ratio of AAOD under the external-mixing scheme and other internally-mixed schemes. The median values of E_{abs} under the AI-NAOS, core-shell and volume-mixing schemes were 1.42, 1.79, and 1.86, respectively. It is clear that the encapsulated fractal models in the AI-NAOS module could induce less absorption enhancement as the lensing effect was weak, as the BC core was only partially coated by non-absorptive hygroscopic aerosols. The median values of single scattering albedo were 0.81, 0.78, and 0.78 under the three schemes, which could also be attributed to the lensing effect. The E_{abs} value could be extremely large (1.80 ± 0.29) when the BC was heavily coated and dropped down to 1.22 ± 0.09 with thin coating (Zhai et al., 2022). In most cases, the mean value of E_{abs} observed in China was quite small, ranging from 1.17
 325 to 1.50 (Wu et al., 2018; Ma et al., 2020; Sun et al., 2021; Zheng et al., 2022). Taking all factors into consideration, the AI-NAOS results, which provided a moderate estimation of AAOD while maintaining a lower E_{abs} value, appeared to be a favorable choice.



330 **Figure 5. Probability density function of absorption enhancement and single scattering albedo over the Sichuan Basin within the AI-NAOS, core-shell, and volume-mixing schemes.**

3.3 Shortwave direct radiation effect

The shortwave radiative flux is directly influenced by the optical properties of the atmospheric column. Figure 6 depicts the aerosol shortwave DRE at the TOA, within the atmosphere, and at the surface, based on four aerosol optical schemes (external-mixing, AI-NAOS, core-shell and volume-mixing). Similar to Section 3.1, these DREs were averaged over a 48-hour period.

335 At the TOA, the spatially averaged DREs based on the AI-NAOS module were +1.9, -2.8, and -1.2 W/m² over the three marked regions, respectively. The external-mixing scheme yielded values of +0.3, -4.6, and -3.0 W/m², while the core-shell scheme showed values of +2.4, -2.8, and -1.2 W/m², and the volume-mixing scheme showed values of +3.4, -1.8, and -0.4 W/m². It is observed that aerosols exhibited a warming impact on the Earth's climate system over the Sichuan Basin, whereas a cooling effect was discovered over the Middle Yangtze Plain and North China Plain. In comparison to the AI-NAOS scheme, the warming effect was much weaker with the external-mixing scheme but more pronounced with the core-shell and volume-mixing schemes.

345 At the surface, the spatially averaged DREs based on the AI-NAOS module were -21.5, -37.8 and -31.2 W/m² over the three regions, respectively, indicating a cooling effect of aerosols at the surface. This cooling effect was weaker with the external-mixing scheme and stronger with the core-shell scheme and volume-mixing schemes. This finding can be attributed to atmospheric absorption, which was determined by the difference between net solar fluxes of TOA and the surface. It is evident that the changes in atmosphere absorption based on the AI-NAOS scheme were extremely large over the three regions, with values of +23.4, +35.0 and +30.0 W/m², respectively. As shown in the second panel of Figure 6, the atmosphere absorption based on the three spherical schemes were also benchmarked to the AI-NAOS scheme, with values of

350



80.7%, 112.5%, and 118.9% over Sichuan Basin, 79.6%, 109.2%, and 115.7% over Middle Yangtze Plain, and 80.2%, 105.8%, and 114.3% over North China Plain.

Taking into consideration of the NSIH effect, a warming effect on the Earth's climate system and a cooling effect on the surface were found compared to the external-mixing scheme. Specifically, the DRE induced by the NSIH effect was
 355 +1.6, +1.8, and +1.8 W/m² at TOA and -2.9, -5.3, and -4.1 W/m² at surface, over the three regions, respectively. Furthermore, these effects can be strengthened using the volume-mixing scheme, which may be too strong and lead to an overestimation.




Figure 6. Aerosol shortwave direct radiation effect based on four aerosol optical schemes (external-mixing, AI-NAOS, core-shell, and volume-mixing) (a) at the top of atmosphere (TOA), (b) within the atmosphere, and (c) at the surface.

360 3.4 Thermodynamic structure


Solar radiation directly heats the Earth's system and influences the thermodynamic structure of the atmosphere. Scattering aerosols, such as sulfate, can partially offset the warming effects from greenhouse gases, while absorptive aerosols, such as BC, can lead to strong warming effects (Kaufmann et al., 2011; Ramanathan and Carmichael, 2008). The overall aerosol DRE could result in a spatially averaged surface temperature decrease of -0.487K over China (Persad and
 365 Caldeira, 2018), with cooling reaching up to -0.7K in specific regions like the Sichuan Basin (Giorgi, 2002).

Figure 7 illustrates anomalies in temperature at 700 hPa, the height of the planetary boundary layer (H_{PBL}), and the 2-meter temperature. As shown in the first column of Figure 7, the aerosol DRE induced a warming effect at 700 hPa over the planetary boundary layer (PBL) and a cooling effect at 2 meters above surface, leading to a more stable atmosphere layer and



370 a decrease in H_{PBL} . Within the AI-NAOS module, anomalies in temperature at 700 hPa were +0.17, +0.12, and +0.14 K, while anomalies in the 2-meter temperature were -0.83, -1.11, and -1.37K over the three regions, respectively. The values of decreases in H_{PBL} were -75, -108 and -90 meters over the three regions, respectively.

375 The warming effect at 700 hPa and the cooling effect on the ground were less prominent based on the external-mixing scheme. The temperature anomalies between the two schemes were +0.04, +0.03, and +0.02 K at 700 hPa and -0.10, -0.14, and -0.17 K on the ground, over the three regions, respectively. The NSIH effect is observed to enhance the temperature anomalies by at least 21% at 700 hPa and 13% on the ground. Similarly, the H_{PBL} could be further decreased by NSIH effect, with values of -11, -15 and -12 meters over the three regions, respectively. Generally, the anomalies in temperature and H_{PBL} could be more intensified by the core-shell and volume-mixing schemes, which exhibited larger DRE anomalies.




380 **Figure 7. Anomalies in temperature at 700 hPa, height of planetary boundary layer, and 2-meter temperature caused by the direct radiative effect (DRE). The anomalies are calculated as the difference between the results under the AI-NAOS module and the control scheme without aerosol effects (first column), as well as the difference between the results under three spherical schemes and the AI-NAOS module (second to fourth columns).**

Figure 8 illustrates the vertical profiles of perturbations in temperature and aerosol short-wave radiation heating rate (HR) based on the four optical aerosol schemes, averaged over three specific areas. The aerosol heating effect decreases with



385 the height as aerosol concentrations are higher near the emission source on lower levels. The HR values within the AI-NAOS
module were 0.64, 0.72, and 0.66 K/day at the surface, and 0.29, 0.34, and 0.31 K/day at 700 hPa, over the three specific
areas, respectively. When vertically averaged, the values within the AI-NAOS module were 0.54, 0.61, and 0.56 K/day,
which were +0.09, +0.11, and +0.10 K/day higher than the values within the external-mixing scheme, over the three specific
390 areas, respectively. Therefore, the NSIH effect could enhance the heating rate by at least 20%. This enhancement could be
even stronger within the core-shell and volume-mixing schemes, reaching 28% and 36%, respectively, over the Sichuan
Basin.

Temperature perturbations increased with height, indicating a cooling effect at the surface and a warming effect above
the PBL. Vertically averaged temperature perturbations under the AI-NAOS module were -0.53, -0.45, and -0.40 K across
the three regions, respectively, indicating an overall cooling effect. This cooling effect of the three spherical schemes was
395 92%, 113%, and 111% compared to the AI-NAOS module over the Sichuan Basin, respectively. In general, NSIH effects
induce direct changes in the heating rate based on anomalies of atmosphere absorption, further influencing temperature and
thermodynamic structures.



400 **Figure 8. Vertical profiles of (a-c) temperature and (d-f) short-wave heating rate anomalies based on four aerosol optical schemes (external-mixing, AI-NAOS, core-shell and volume-mixing). The solid lines represent the median value, while the shaded areas cover the range from 25 to 75 percentage.**

3.4 Precipitation

We then examined the impact of aerosol-induced changes in thermodynamic structure on precipitation anomalies. The
405 direct radiation effect plays a role in reducing solar radiation flux, temperature and moisture fluxes at the surface, leading to
the inhibition of convection and a decrease in precipitation (Nabat et al., 2015). Additionally, the semi-direct effect, caused
by aerosol absorption heating the lower atmosphere, results in increased atmospheric stability and suppression of convection



410 within the PBL, while inducing stronger convection above this boundary layer (Allen et al., 2019). Collectively, these two
effects contribute to reduced precipitation in East Asia. However, anomalous circulations driven by direct radiation effect
can impact moisture transport, leading to enhanced precipitation in South China but reduced precipitation in North China
(Chen et al., 2018; Huang et al., 2007).

415 Figure 9 depicts the anomalies in accumulated precipitation of the last 48 hours between the AI-NAOS module and a
control scheme with zero AOD. The analysis focused on a specific region within the precipitation center (101.5-112.5°E,
29.5-34.5°N), encompassing parts of the Sichuan Basin and Middle Yangtze Plain. Overall, there was a suppression effect
over this region due to the aerosol-induced stability with weaker convections in the atmosphere. The spatially averaged
anomalies were -0.24 mm under the AI-NAOS module, while they were -0.21, -0.26 and -0.27 mm corresponding to the
external-mixing, core-shell, and volume-mixing schemes. Notably, the suppression effect was less pronounced in the
external-mixing scheme. The NSIH effect amplified the suppression effect by 15%. In the core-shell and volume-mixing
schemes, this effect could be even more pronounced, similar to the thermodynamic effect.

420




Figure 9. Anomalies in accumulative precipitation between the AI-NAOS module and the control scheme over a specific region within the precipitation center (101.5–112.5°E, 29.5–34.5°N).

425 4 Conclusion

A new aerosol optical module, referred to as AI-NAOS, was developed to incorporate the nonsphericity and inhomogeneity of aerosol particles. Using integrated DNNs, aerosol optical properties were effectively calculated based on



NSIH aerosol models. This newly developed aerosol optical module was on-line coupled with the chemical weather model, GRAPES_Meso5.1/CUACE.

430 In the AI-NAOS scheme, BC was assumed to be fractal aggregates while soil dust was modelled as super-spheroids. Both of these insoluble aerosols were either fully or partially encapsulated with hygroscopic aerosols, which were treated with the volume-mixing method. The aerosol optical properties were computed using the IITM algorithm. Two databases of optical properties for BC and SD were established, including more than 700, 000 records. Direct incorporating the two optical property databases into the aerosol parameterization scheme using the look-up table method is inconvenient. This
435 method consumes a significant amount of storage and introduces additional errors from interpolation. To address this issue, two DNNs were trained based on the two optical property databases. A unified interface was implemented in the AI-NAOS scheme to calculate the optical properties of BC and SD using the DNNs.

Real-scenario simulations were conducted to assess the NSIH effect of BC aerosols using the GRAPES_Meso5.1/CUACE model. In the simulation, five aerosol treatments were applied, including the control experiment
440 without aerosol optical effects, the AI-NAOS module, and three spherical schemes.

The AI-NAOS module provided a moderate estimation of AAOD, falling between the underestimation from the external-mixing scheme and the overestimation from the core-shell and the volume-mixing schemes. For example, the AAOD values based on the three spherical schemes were 70.4%, 125.3%, and 129.3% over Sichuan Basin, benchmarked to the AI-NAOS results. The absorption enhancements were 1.42, 1.79, and 1.86 based on the AI-NAOS, core-shell and the
445 volume-mixing schemes. It was clear that the lowest absorption enhancement value was induced under the AI-NAOS module, indicating a favorable choice that matched low E_{abs} measurements. It is worth noting that the asymmetry factor based on the NSIH scheme was the largest among all schemes, indicating the weakest backscattering fraction.

The aerosol shortwave DRE is directly influenced by the NSIH effect. Compared to the external-mixing scheme, the DRE induced by the NSIH effect was +1.6, +1.8, and +1.8 W/m^2 at TOA and -2.9, -5.3, and -4.1 W/m^2 at the surface over
450 three regions, respectively. This effect was stronger within the core-shell and volume-mixing schemes, which could mainly be attributed to the difference in aerosol absorption. Compared to the AI-NAOS scheme, the DREs in the atmosphere were 80.7%, 112.5%, and 118.9% over the Sichuan Basin based on the three spherical schemes, respectively.

The NSIH effect also resulted in changes in the thermal structure. The heating rates within the AI-NAOS module were 0.64, 0.72, and 0.66 K/day on the ground and 0.29, 0.34, and 0.31 K/day at 700 hPa. Compared to the external-mixing
455 scheme, the NSIH effect could enhance the heating rate, reaching at least 20%. This enhancement could be stronger within the core-shell and volume-mixing schemes, reaching 28% and 36%. The temperature anomalies under the AI-NAOS scheme were +0.17, +0.12, and +0.14 K at 700 hPa and -0.83, -1.11, and -1.37 K on the ground over three regions, respectively. Compared to the external-mixing scheme, the NSIH effect could enhance the temperature anomalies by at least 21% at 700 hPa and 13% on the ground. This anomaly could be larger within the core-shell and volume-mixing schemes. The vertically
460 averaged temperature anomalies of the three spherical schemes were 92%, 113%, and 111% benchmarked to the AI-NAOS scheme over the Sichuan Basin. As a result, the PBL becomes more stable, with its height decreasing by at least -11 m due to



the NSIH effect. The precipitation was suppressed with weaker convection in the stable atmosphere, being -0.24 mm under the AI-NAOS module. Compared to the external-mixing scheme, the NSIH effect enhanced the suppression effect by 15%.

465 In this study, a comprehensive assessment was conducted to analyze the NSIH effect of BC aerosols. The findings indicated that the NSIH effect cannot be disregarded in weather forecast models. A crucial aspect is that the estimation of aerosol absorption can be improved by incorporating the NSIH effect. However, additional real-case studies, encompassing weather and climate scales, should be undertaken in the future.

Temporarily, the refractive indices and fractal dimension of BC were fixed. As the AI-NAOS module is driven by DNNs, it is convenient to adjust refractive indices without rebuilding the DNN. This flexibility proves advantageous for
470 investigating the absorption ability of BC. Additionally, there are plans to train a new DNN that incorporates BC particles with various fractal dimensions, aiming to support more detailed research in this area. As for SD, the database of coated super-spheroid for large particles will be updated when a computational program based on the ray tracing technique is available. In the current AI-NAOS module, the nonsphericity and inhomogeneity are only considered with insoluble aerosol particles. Although hygroscopic aerosols, such as sea salt, may be completely dissolved in aerosol water after deliquescence,
475 a solid core coated with liquid saline still exists until the relative humidity reaches 97% (Zeng et al., 2013). The NSIH effect has a significant impact on the single-scattering properties of sea salt, especially with respect to depolarization (Bi et al., 2018b; Kanngießner and Kahnert, 2021; Lin and Bi, 2024). Additionally, it was found that the NSIH effect on optical properties could induce a cooling effect, weaken vertical velocity and reduce the accumulated rainfall of Typhoon Fitow (Zhu et al., 2022). Note that the NSIH effect of sea salt aerosol is not important in this study, because we focused on the land
480 area. In the future, we will consider the NSIH features of sea salt particles, developing a more comprehensive nonspherical and inhomogeneous aerosol optical scheme. Furthermore, it is anticipated that the AI-NAOS scheme will be routinely updated and implemented in various chemical weather, climate, and radiation transfer models, as well as utilized for data assimilation purposes.

485

Appendix. Deep neural networks of bulk optical properties.


As shown in Table A1, there were three input parameters (x , $m_{r_{\text{hygro}}}$, $m_{i_{\text{hygro}}}$) for spheres in the volume-mixing scheme and six parameters (x , $m_{r_{\text{hygro}}}$, $m_{i_{\text{hygro}}}$, v_f , $m_{r_{\text{insol}}}$, $m_{i_{\text{insol}}}$) for the core-shell model and two NSIH models of BC and SD. The architecture of the DNNs included fully connected layers and duplicated residual blocks, as depicted in Figure A1.
490 Specifically, the first two hidden layers consisted of fully-connected layers with 40 and 20 nodes, respectively. The following setup included three branches corresponding to the extinction coefficient, single-scattering albedo, and asymmetry factor. In each branch, two residual blocks were included and each block was composed by two fully-connected layers with 20 nodes and a shortcut connection. The architecture is similar to our previous work (Wang et al., 2023b). However, in this



495 study, the DNNs were applied to bulk optical properties in various size bins, and the scattering matrices were excluded from the database. Additionally, we followed the method proposed in Wang et al. (2022b) to constrain the extrapolation of the DNN for large coated super-spheroids by using the asymptotic values of the core-shell model.

Table A1. DNNs of bulk optical properties (used in the AI-NAOS module).

Aerosol	Particle Model	DNN input	DNN output
Hygroscopic aerosols	volume-mixing (sphere)	$(x, m_{r_{\text{hygro}}}, m_{i_{\text{hygro}}})$	$(Q_{\text{ext}}, \langle \text{SSA} \rangle, \langle G \rangle)$
BC	core-shell	$(x, m_{r_{\text{hygro}}}, m_{i_{\text{hygro}}}, v_f, m_{r_{\text{insol}}}, m_{i_{\text{insol}}})$	
	encapsulated fractal soot		
SD	coated super-spheroid		



500

Figure A1. The architecture of DNNs in this study.

Data availability. The data presented in this paper are available on Zenodo (<https://doi.org/10.5281/zenodo.11208003>).

505 *Code availability.* The AI-NAOS aerosol optical module codes developed in this study are available on Zenodo (<https://doi.org/10.5281/zenodo.11181275>). We used the GRAPES_Meso5.1/CUACE code provided by the CMA Earth System Modeling and Prediction Center (<https://cemc.cma.cn/>) together with the Chinese Academy of Meteorological Sciences (<https://www.cma.gov.cn/>). We cannot distribute the GRAPES_Meso5.1/CUACE code due to its license.

510 *Acknowledgement.* We used the computing facilities of the China HPC Cloud Computing Center. This work was supported by the National Key Scientific and Technological Infrastructure project Earth System Numerical Simulation Facility (EarthLab).



Financial support. This research was supported by the NSFC Major Project (42090030).

515

Competing interests. The contact author has declared that none of the authors has any competing interests.

References

- Adachi, K., Chung, S. H., Friedrich, H., and Buseck, P. R.: Fractal parameters of individual soot particles determined using electron tomography: Implications for optical properties, *J. Geophys. Res.*, 112, D14202, <https://doi.org/10.1029/2006JD008296>, 2007.
- Adachi, K., Chung, S. H., and Buseck, P. R.: Shapes of soot aerosol particles and implications for their effects on climate, *J. Geophys. Res.*, 115, D15206, <https://doi.org/10.1029/2009JD012868>, 2010.
- Allen, R. J., Amiri-Farahani, A., Lamarque, J.-F., Smith, C., Shindell, D., Hassan, T., and Chung, C. E.: Observationally constrained aerosol–cloud semi-direct effects, *npj Clim Atmos Sci*, 2, 16, <https://doi.org/10.1038/s41612-019-0073-9>, 2019.
- Andersson, E. and Kahnert, M.: Coupling aerosol optics to the MATCH (v5.5.0) chemical transport model and the SALSA (v1) aerosol microphysics module, *Geosci. Model Dev.*, 9, 1803–1826, <https://doi.org/10.5194/gmd-9-1803-2016>, 2016.
- Barnard, J. C., Fast, J. D., Paredes-Miranda, G., Arnott, W. P., and Laskin, A.: Technical Note: Evaluation of the WRF-Chem “Aerosol Chemical to Aerosol Optical Properties” Module using data from the MILAGRO campaign, *Atmos. Chem. Phys.*, 10, 7325–7340, <https://doi.org/10.5194/acp-10-7325-2010>, 2010.
- Barr: Superquadrics and Angle-Preserving Transformations, *IEEE Comput. Grap. Appl.*, 1, 11–23, <https://doi.org/10.1109/MCG.1981.1673799>, 1981.
- Bi, L. and Yang, P.: Accurate simulation of the optical properties of atmospheric ice crystals with the invariant imbedding T-matrix method, *J. Quant. Spectrosc. Radiat. Transfer*, 138, 17–35, <https://doi.org/10.1016/j.jqsrt.2014.01.013>, 2014.
- Bi, L., Yang, P., Kattawar, G. W., and Mishchenko, M. I.: Efficient implementation of the invariant imbedding T-matrix method and the separation of variables method applied to large nonspherical inhomogeneous particles, *J. Quant. Spectrosc. Radiat. Transfer*, 116, 169–183, <https://doi.org/10.1016/j.jqsrt.2012.11.014>, 2013.
- Bi, L., Lin, W., Liu, D., and Zhang, K.: Assessing the depolarization capabilities of nonspherical particles in a super-ellipsoidal shape space, *Opt. Express*, 26, 1726, <https://doi.org/10.1364/OE.26.001726>, 2018a.
- Bi, L., Lin, W., Wang, Z., Tang, X., Zhang, X., and Yi, B.: Optical Modeling of Sea Salt Aerosols: The Effects of Nonsphericity and Inhomogeneity, *J. Geophys. Res. Atmos.*, 123, 543–558, <https://doi.org/10.1002/2017JD027869>, 2018b.
- Bi, L., Wang, Z., Han, W., Li, W., and Zhang, X.: Computation of Optical Properties of Core-Shell Super-Spheroids Using a GPU Implementation of the Invariant Imbedding T-Matrix Method, *Front. Remote Sens.*, 3, 903312, <https://doi.org/10.3389/frsen.2022.903312>, 2022.
- Bohren, C. F. and Huffman, D. R.: *Absorption and Scattering of Light by Small Particles*, 1st ed., Wiley, 1998.
- Bond, T. C. and Bergstrom, R. W.: Light Absorption by Carbonaceous Particles: An Investigative Review, *Aerosol Sci. Technol.*, 40, 27–67, <https://doi.org/10.1080/02786820500421521>, 2006.
- Buseck, P. R. and Pósfai, M.: Airborne minerals and related aerosol particles: Effects on climate and the environment, *Proc. Natl. Acad. Sci. U.S.A.*, 96, 3372–3379, <https://doi.org/10.1073/pnas.96.7.3372>, 1999.
- Chen, F., Janjić, Z., and Mitchell, K.: Impact of Atmospheric Surface-layer Parameterizations in the new Land-surface Scheme of the NCEP Mesoscale Eta Model, *Bound.-Layer Meteorol.*, 85, 391–421, <https://doi.org/10.1023/A:1000531001463>, 1997.
- Chen, G., Wang, W.-C., and Chen, J.-P.: Circulation responses to regional aerosol climate forcing in summer over East Asia, *Clim Dyn*, 51, 3973–3984, <https://doi.org/10.1007/s00382-018-4267-3>, 2018.
- Chen, G., Wang, J., Wang, Y., Wang, J., Jin, Y., Cheng, Y., Yin, Y., Liao, H., Ding, A., Wang, S., Hao, J., and Liu, C.: An Aerosol Optical Module With Observation-Constrained Black Carbon Properties for Global Climate Models, *J Adv Model Earth Syst*, 15, e2022MS003501, <https://doi.org/10.1029/2022MS003501>, 2023.



- Chou, M.-D., Suarez, M. J., Ho, C.-H., Yan, M. M.-H., and Lee, K.-T.: Parameterizations for Cloud Overlapping and Shortwave Single-Scattering Properties for Use in General Circulation and Cloud Ensemble Models, *J. Climate*, 11, 202–214, [https://doi.org/10.1175/1520-0442\(1998\)011<0202:PFCOAS>2.0.CO;2](https://doi.org/10.1175/1520-0442(1998)011<0202:PFCOAS>2.0.CO;2), 1998.
- 560 Fast, J. D., Gustafson, W. I., Easter, R. C., Zaveri, R. A., Barnard, J. C., Chapman, E. G., Grell, G. A., and Peckham, S. E.: Evolution of ozone, particulates, and aerosol direct radiative forcing in the vicinity of Houston using a fully coupled meteorology-chemistry-aerosol model, *J. Geophys. Res.*, 111, D21305, <https://doi.org/10.1029/2005JD006721>, 2006.
- Forrest, S. R. and Witten, T. A.: Long-range correlations in smoke-particle aggregates, *J. Phys. A: Math. Gen.*, 12, L109–L117, <https://doi.org/10.1088/0305-4470/12/5/008>, 1979.
- 565 Giorgi, F.: Direct radiative forcing and regional climatic effects of anthropogenic aerosols over East Asia: A regional coupled climate-chemistry/aerosol model study, *J. Geophys. Res.*, 107, 4439, <https://doi.org/10.1029/2001JD001066>, 2002.
- Gong, S. L. and Zhang, X. Y.: CUACE/Dust – an integrated system of observation and modeling systems for operational dust forecasting in Asia, *Atmos. Chem. Phys.*, 8, 2333–2340, <https://doi.org/10.5194/acp-8-2333-2008>, 2008.
- Hong, S.-Y. and Pan, H.-L.: Nonlocal Boundary Layer Vertical Diffusion in a Medium-Range Forecast Model, *Mon. Wea. Rev.*, 124, 2322–2339, [https://doi.org/10.1175/1520-0493\(1996\)124<2322:NBLVDI>2.0.CO;2](https://doi.org/10.1175/1520-0493(1996)124<2322:NBLVDI>2.0.CO;2), 1996.
- 570 Huang, X.-F., Peng, Y., Wei, J., Peng, J., Lin, X.-Y., Tang, M.-X., Cheng, Y., Men, Z., Fang, T., Zhang, J., He, L.-Y., Cao, L.-M., Liu, C., Zhang, C., Mao, H., Seinfeld, J. H., and Wang, Y.: Microphysical complexity of black carbon particles restricts their warming potential, *One Earth*, S2590332223005596, <https://doi.org/10.1016/j.oneear.2023.12.004>, 2023.
- Huang, Y., Chameides, W. L., and Dickinson, R. E.: Direct and indirect effects of anthropogenic aerosols on regional precipitation over east Asia, *J. Geophys. Res.*, 112, D03212, <https://doi.org/10.1029/2006JD007114>, 2007.
- 575 Kahnert, M., Nousiainen, T., Lindqvist, H., and Ebert, M.: Optical properties of light absorbing carbon aggregates mixed with sulfate: assessment of different model geometries for climate forcing calculations, *Opt. Express*, 20, 10042, <https://doi.org/10.1364/OE.20.010042>, 2012.
- Kain, J. S. and Fritsch, J. M.: Convective Parameterization for Mesoscale Models: The Kain-Fritsch Scheme, in: *The Representation of Cumulus Convection in Numerical Models*, edited by: Emanuel, K. A. and Raymond, D. J., American Meteorological Society, Boston, MA, 165–170, 1993.
- 580 Kanngießer, F. and Kahnert, M.: Optical properties of water-coated sea salt model particles, *Opt. Express*, 29, 34926, <https://doi.org/10.1364/OE.437680>, 2021.
- Kaufmann, R. K., Kauppi, H., Mann, M. L., and Stock, J. H.: Reconciling anthropogenic climate change with observed temperature 1998–2008, *Proc. Natl. Acad. Sci. U.S.A.*, 108, 11790–11793, <https://doi.org/10.1073/pnas.1102467108>, 2011.
- 585 Kotchenova, S. Y. and Vermote, E. F.: Validation of a vector version of the 6S radiative transfer code for atmospheric correction of satellite data Part II Homogeneous Lambertian and anisotropic surfaces, *Appl. Opt.*, 46, 4455, <https://doi.org/10.1364/AO.46.004455>, 2007.
- Li, W., Shao, L., Zhang, D., Ro, C.-U., Hu, M., Bi, X., Geng, H., Matsuki, A., Niu, H., and Chen, J.: A review of single aerosol particle studies in the atmosphere of East Asia: morphology, mixing state, source, and heterogeneous reactions, *J. Cleaner Prod.*, 112, 1330–1349, <https://doi.org/10.1016/j.jclepro.2015.04.050>, 2016.
- 590 Lin, W. and Bi, L.: Optical modeling of sea salt aerosols using in situ measured size distributions and the impact of larger size particles, *Adv. Atmos. Sci.*, <https://doi.org/10.1007/s00376-024-3351-3>, 2024.
- Lin, W., Bi, L., and Dubovik, O.: Assessing Superspheroids in Modeling the Scattering Matrices of Dust Aerosols, *J. Geophys. Res. Atmos.*, 123, <https://doi.org/10.1029/2018JD029464>, 2018.
- 595 Lin, W., Bi, L., Weng, F., Li, Z., and Dubovik, O.: Capability of Superspheroids for Modeling PARASOL Observations Under Dusty-Sky Conditions, *J. Geophys. Res. Atmos.*, 126, <https://doi.org/10.1029/2020JD033310>, 2021.
- Luo, J., Zhang, Q., Luo, J., Liu, J., Huo, Y., and Zhang, Y.: Optical Modeling of Black Carbon With Different Coating Materials: The Effect of Coating Configurations, *J. Geophys. Res. Atmos.*, 124, 13230–13253, <https://doi.org/10.1029/2019JD031701>, 2019.
- 600 Ma, Y., Huang, C., Jabbour, H., Zheng, Z., Wang, Y., Jiang, Y., Zhu, W., Ge, X., Collier, S., and Zheng, J.: Mixing state and light absorption enhancement of black carbon aerosols in summertime Nanjing, China, *Atmos. Environ.*, 222, 117141, <https://doi.org/10.1016/j.atmosenv.2019.117141>, 2020.
- Mackowski, D. W.: A general superposition solution for electromagnetic scattering by multiple spherical domains of optically active media, *J. Quant. Spectrosc. Radiat. Transfer*, 133, 264–270, <https://doi.org/10.1016/j.jqsrt.2013.08.012>, 2014.
- 605



- Matsui, H.: Development of a global aerosol model using a two-dimensional sectional method: 1. Model design, *J. Adv. Model. Earth Syst.*, 9, 1921–1947, <https://doi.org/10.1002/2017MS000936>, 2017.
- 610 Mlawer, E. J., Taubman, S. J., Brown, P. D., Iacono, M. J., and Clough, S. A.: Radiative transfer for inhomogeneous atmospheres: RRTM, a validated correlated-k model for the longwave, *J. Geophys. Res.*, 102, 16663–16682, <https://doi.org/10.1029/97JD00237>, 1997.
- Morán, J., Fuentes, A., Liu, F., and Yon, J.: FracVAL: An improved tunable algorithm of cluster–cluster aggregation for generation of fractal structures formed by polydisperse primary particles, *Comput. Phys. Commun.*, 239, 225–237, <https://doi.org/10.1016/j.cpc.2019.01.015>, 2019.
- 615 Nabat, P., Somot, S., Mallet, M., Sevault, F., Chiacchio, M., and Wild, M.: Direct and semi-direct aerosol radiative effect on the Mediterranean climate variability using a coupled regional climate system model, *Clim Dyn*, 44, 1127–1155, <https://doi.org/10.1007/s00382-014-2205-6>, 2015.
- National Centers For Environmental Prediction/National Weather Service/NOAA/U.S. Department Of Commerce: NCEP GDAS/FNL 0.25 Degree Global Tropospheric Analyses and Forecast Grids, <https://doi.org/10.5065/D65Q4T4Z>, 2015.
- 620 Peng, Y., Wang, H., Zhang, X., Zheng, Y., Zhang, X., Zhang, W., Liu, Z., Gui, K., Liu, H., Wang, Y., and Che, H.: Aerosol-radiation interaction in the operational atmospheric chemistry model GRAPES_Meso5.1/CUACE and its impacts on mesoscale NWP in Beijing-Tianjin-Hebei, China, *Atmos. Res.*, 280, 106402, <https://doi.org/10.1016/j.atmosres.2022.106402>, 2022.
- Persad, G. G. and Caldeira, K.: Divergent global-scale temperature effects from identical aerosols emitted in different regions, *Nat Commun*, 9, 3289, <https://doi.org/10.1038/s41467-018-05838-6>, 2018.
- Pleim, J. E.: A Combined Local and Nonlocal Closure Model for the Atmospheric Boundary Layer. Part I: Model Description and Testing, *J. Appl. Meteorol. Climatol.*, 46, 1383–1395, <https://doi.org/10.1175/JAM2539.1>, 2007.
- Pósfai, M., Axisa, D., Tompa, É., Freney, E., Bruintjes, R., and Buseck, P. R.: Interactions of mineral dust with pollution and clouds: An individual-particle TEM study of atmospheric aerosol from Saudi Arabia, *Atmos. Res.*, 122, 347–361, <https://doi.org/10.1016/j.atmosres.2012.12.001>, 2013.
- 630 Ramanathan, V. and Carmichael, G.: Global and regional climate changes due to black carbon, *Nat. Geosci.*, 1, 221–227, <https://doi.org/10.1038/ngeo156>, 2008.
- Rothman, L. S., Jacquemart, D., Barbe, A., Chris Benner, D., Birk, M., Brown, L. R., Carleer, M. R., Chackerian, C., Chance, K., Coudert, L. H., Dana, V., Devi, V. M., Flaud, J.-M., Gamache, R. R., Goldman, A., Hartmann, J.-M., Jucks, K. W., Maki, A. G., Mandin, J.-Y., Massie, S. T., Orphal, J., Perrin, A., Rinsland, C. P., Smith, M. A. H., Tennyson, J., Tolchenov, R. N., Toth, R. A., Vander Auwera, J., Varanasi, P., and Wagner, G.: The HITRAN 2004 molecular spectroscopic database, *J. Quant. Spectrosc. Radiat. Transfer*, 96, 139–204, <https://doi.org/10.1016/j.jqsrt.2004.10.008>, 2005.
- Sorensen, C. M.: Light Scattering by Fractal Aggregates: A Review, *Aerosol Sci. Technol.*, 35, 648–687, <https://doi.org/10.1080/02786820117868>, 2001.
- 640 Sorensen, C. M.: The Mobility of Fractal Aggregates: A Review, *Aerosol Sci. Technol.*, 45, 765–779, <https://doi.org/10.1080/02786826.2011.560909>, 2011.
- Stockwell, W. R., Middleton, P., Chang, J. S., and Tang, X.: The second generation regional acid deposition model chemical mechanism for regional air quality modeling, *J. Geophys. Res.*, 95, 16343, <https://doi.org/10.1029/JD095iD10p16343>, 1990.
- 645 Sun, J., Xie, C., Xu, W., Chen, C., Ma, N., Xu, W., Lei, L., Li, Z., He, Y., Qiu, Y., Wang, Q., Pan, X., Su, H., Cheng, Y., Wu, C., Fu, P., Wang, Z., and Sun, Y.: Light absorption of black carbon and brown carbon in winter in North China Plain: comparisons between urban and rural sites, *Sci. Total Environ.*, 770, 144821, <https://doi.org/10.1016/j.scitotenv.2020.144821>, 2021.
- Taylor, F. W.: *Theory of Atmospheric Radiative Transfer: A Comprehensive Introduction*, by Manfred Wendisch and Ping Yang: Scope: textbook. Level: advanced undergraduate, *Contemp. Phys.*, 53, 506–506, <https://doi.org/10.1080/00107514.2012.732967>, 2012.
- 650 Teng, S., Liu, C., Schnaiter, M., Chakrabarty, R. K., and Liu, F.: Accounting for the effects of nonideal minor structures on the optical properties of black carbon aerosols, *Atmos. Chem. Phys.*, 19, 2917–2931, <https://doi.org/10.5194/acp-19-2917-2019>, 2019.
- Thompson, G., Rasmussen, R. M., and Manning, K.: Explicit Forecasts of Winter Precipitation Using an Improved Bulk Microphysics Scheme. Part I: Description and Sensitivity Analysis, *Mon. Wea. Rev.*, 132, 519–542, [https://doi.org/10.1175/1520-0493\(2004\)132<0519:EFOWPU>2.0.CO;2](https://doi.org/10.1175/1520-0493(2004)132<0519:EFOWPU>2.0.CO;2), 2004.



- Wang, H., Shi, G., Li, S., Li, W., Wang, B., and Huang, Y.: The impacts of optical properties on radiative forcing due to dust aerosol, *Adv. Atmos. Sci.*, 23, 431–441, <https://doi.org/10.1007/s00376-006-0431-5>, 2006.
- 660 Wang, H., Zhang, X., Gong, S., Chen, Y., Shi, G., and Li, W.: Radiative feedback of dust aerosols on the East Asian dust storms, *J. Geophys. Res. Atmos.*, 115, <https://doi.org/10.1029/2009JD013430>, 2010.
- Wang, H., Shi, G. Y., Zhang, X. Y., Gong, S. L., Tan, S. C., Chen, B., Che, H. Z., and Li, T.: Mesoscale modelling study of the interactions between aerosols and PBL meteorology during a haze episode in China Jing–Jin–Ji and its near surrounding region – Part 2: Aerosols’ radiative feedback effects, *Atmos. Chem. Phys.*, 15, 3277–3287, <https://doi.org/10.5194/acp-15-3277-2015>, 2015a.
- 665 Wang, H., Shi, G. Y., Zhang, X. Y., Gong, S. L., Tan, S. C., Chen, B., Che, H. Z., and Li, T.: Mesoscale modelling study of the interactions between aerosols and PBL meteorology during a haze episode in China Jing–Jin–Ji and its near surrounding region – Part 2: Aerosols’ radiative feedback effects, *Atmos. Chem. Phys.*, 15, 3277–3287, <https://doi.org/10.5194/acp-15-3277-2015>, 2015b.
- Wang, H., Peng, Y., Zhang, X., Liu, H., Zhang, M., Che, H., Cheng, Y., and Zheng, Y.: Contributions to the explosive growth of PM_{2.5} mass due to aerosol–radiation feedback and decrease in turbulent diffusion during a red alert heavy haze in Beijing–Tianjin–Hebei, China, *Atmos. Chem. Phys.*, 18, 17717–17733, <https://doi.org/10.5194/acp-18-17717-2018>, 2018.
- 670 Wang, H., Zhang, X. Y., Wang, P., Peng, Y., Zhang, W. J., Liu, Z. D., Han, C., Li, S. T., Wang, Y. Q., Che, H. Z., Huang, L. P., Liu, H. L., Zhang, L., Zhou, C. H., Ma, Z. S., Chen, F. F., Ma, X., Wu, X. J., Zhang, B. H., and Shen, X. S.: Chemistry–Weather Interacted Model System GRAPES_Meso5.1/CUACE CW V1.0: Development, Evaluation and Application in Better Haze/Fog Prediction in China, *J. Adv. Model. Earth Syst.*, 14, e2022MS003222, <https://doi.org/10.1029/2022MS003222>, 2022a.
- Wang, J., Wang, J., Cai, R., Liu, C., Jiang, J., Nie, W., Wang, J., Moteki, N., Zaveri, R. A., Huang, X., Ma, N., Chen, G., Wang, Z., Jin, Y., Cai, J., Zhang, Y., Chi, X., Holanda, B. A., Xing, J., Liu, T., Qi, X., Wang, Q., Pöhlker, C., Su, H., Cheng, Y., Wang, S., Hao, J., Andreae, M. O., and Ding, A.: Unified theoretical framework for black carbon mixing state allows greater accuracy of climate effect estimation, *Nat. Commun.*, 14, 2703, <https://doi.org/10.1038/s41467-023-38330-x>, 2023a.
- 680 Wang, X., Bi, L., Han, W., and Zhang, X.: Single-Scattering Properties of Encapsulated Fractal Black Carbon Particles Computed Using the Invariant Imbedding T-Matrix Method and Deep Learning Approaches, *J. Geophys. Res. Atmos.*, 128, e2023JD039568, <https://doi.org/10.1029/2023JD039568>, 2023b.
- Wang, Y., Liu, F., He, C., Bi, L., Cheng, T., Wang, Z., Zhang, H., Zhang, X., Shi, Z., and Li, W.: Fractal Dimensions and Mixing Structures of Soot Particles during Atmospheric Processing, *Environ. Sci. Technol. Lett.*, 4, 487–493, <https://doi.org/10.1021/acs.estlett.7b00418>, 2017.
- 685 Wang, Z., Bi, L., Wang, H., Wang, Y., Han, W., and Zhang, X.: Evaluation of a new internally-mixed aerosol optics scheme in the weather research and forecasting model, *J. Quant. Spectrosc. Radiat. Transfer*, 283, 108147, <https://doi.org/10.1016/j.jqsrt.2022.108147>, 2022b.
- 690 Wang, Z., Bi, L., and Kong, S.: Flexible implementation of the particle shape and internal inhomogeneity in the invariant imbedding T-matrix method, *Opt. Express*, 31, 29427, <https://doi.org/10.1364/OE.498190>, 2023c.
- Wise, M. E., Semeniuk, T. A., Brientjes, R., Martin, S. T., Russell, L. M., and Buseck, P. R.: Hygroscopic behavior of NaCl-bearing natural aerosol particles using environmental transmission electron microscopy, *J. Geophys. Res.*, 112, <https://doi.org/10.1029/2006JD007678>, 2007.
- 695 Wu, C., Wu, D., and Yu, J. Z.: Quantifying black carbon light absorption enhancement with a novel statistical approach, *Atmos. Chem. Phys.*, 18, 289–309, <https://doi.org/10.5194/acp-18-289-2018>, 2018.
- Wu, Y., Cheng, T., Zheng, L., Zhang, Y., and Zhang, L.: Particle size amplification of black carbon by scattering measurement due to morphology diversity, *Environ. Res. Lett.*, 18, 024011, <https://doi.org/10.1088/1748-9326/acaede>, 2023.
- 700 Yu, H., Kaufman, Y. J., Chin, M., Feingold, G., Remer, L. A., Anderson, T. L., Balkanski, Y., Bellouin, N., Boucher, O., Christopher, S., DeCola, P., Kahn, R., Koch, D., Loeb, N., Reddy, M. S., Schulz, M., Takemura, T., and Zhou, M.: A review of measurement-based assessments of the aerosol direct radiative effect and forcing, *Atmos. Chem. Phys.*, 6, 613–666, <https://doi.org/10.5194/acp-6-613-2006>, 2006.
- Yu, J., Bi, L., Han, W., and Zhang, X.: Application of a Neural Network to Store and Compute the Optical Properties of Non-Spherical Particles, *Adv. Atmos. Sci.*, <https://doi.org/10.1007/s00376-021-1375-5>, 2022a.
- 705 Yu, J., Bi, L., Han, W., Wang, D., and Zhang, X.: Jacobians of single-scattering optical properties of super-spheroids computed using neural networks, *Opt. Express*, 30, 38513, <https://doi.org/10.1364/OE.471821>, 2022b.



- Yurkin, M. A. and Hoekstra, A. G.: The discrete-dipole-approximation code ADDA: Capabilities and known limitations, *J. Quant. Spectrosc. Radiat. Transfer*, 112, 2234–2247, <https://doi.org/10.1016/j.jqsrt.2011.01.031>, 2011.
- 710 Zeng, J., Zhang, G., Long, S., Liu, K., Cao, L., Bao, L., and Li, Y.: Sea salt deliquescence and crystallization in atmosphere: an in situ investigation using x-ray phase contrast imaging: *In situ* imaging of sea salt deliquescence and crystallization, *Surf. Interface Anal.*, 45, 930–936, <https://doi.org/10.1002/sia.5184>, 2013.
- Zhai, J., Yang, X., Li, L., Bai, B., Liu, P., Huang, Y., Fu, T.-M., Zhu, L., Zeng, Z., Tao, S., Lu, X., Ye, X., Wang, X., Wang, L., and Chen, J.: Absorption Enhancement of Black Carbon Aerosols Constrained by Mixing-State Heterogeneity, *Environ. Sci. Technol.*, 56, 1586–1593, <https://doi.org/10.1021/acs.est.1c06180>, 2022.
- 715 Zheng, H., Kong, S., Chen, N., and Wu, C.: Secondary inorganic aerosol dominated the light absorption enhancement of black carbon aerosol in Wuhan, Central China, *Atmos. Environ.*, 287, 119288, <https://doi.org/10.1016/j.atmosenv.2022.119288>, 2022.
- Zhou, C.-H., Gong, S., Zhang, X.-Y., Liu, H.-L., Xue, M., Cao, G.-L., An, X.-Q., Che, H.-Z., Zhang, Y.-M., and Niu, T.: Towards the improvements of simulating the chemical and optical properties of Chinese aerosols using an online coupled model – CUACE/Aero, *Tellus B*, 64, 18965, <https://doi.org/10.3402/tellusb.v64i0.18965>, 2012.
- 720 Zhu, L., Shu, S., Wang, Z., and Bi, L.: More or Less: How Do Inhomogeneous Sea-Salt Aerosols Affect the Precipitation of Landfalling Tropical Cyclones?, *Geophys. Res. Lett.*, 49, <https://doi.org/10.1029/2021GL097023>, 2022.