

PyEt v1.3.1: a Python package for the estimation of potential evapotranspiration

Matevz Vremec^{1,*}, Raoul A. Collenteur^{2,*}, and Steffen Birk¹

*These authors contributed equally to this work.

¹Department of Earth Sciences, NAWI Graz Geocenter, University of Graz, Graz, Austria

²Eawag, Swiss Federal Institute of Aquatic Science and Technology, Department Water Resources and Drinking Water, Dübendorf, Switzerland

Correspondence: Matevz Vremec (matevz.vremec@uni-graz.at)

Abstract.

Evapotranspiration (ET) is a crucial flux of the hydrological water balance, commonly estimated using (semi-)empirical formulas. The estimated flux may strongly depend on the formula used, adding uncertainty to the outcomes of environmental studies using ET. Climate change may cause additional uncertainty, as the ET estimated by each formula may respond differently to changes in meteorological input data. To include the effects of model uncertainty and climate change, and facilitate the use of these formulas in a consistent, tested, and reproducible workflow, we present *PyEt*. *PyEt* is an open-source Python package for the estimation of daily potential evapotranspiration (PET) using available meteorological data. It allows the application of twenty different PET methods on both time series and gridded datasets. The majority of the implemented methods are benchmarked against literature values and tested with continuous integration to ensure the correctness of the implementation. This article provides an overview of *PyEt*'s capabilities, including the estimation of PET with twenty PET methods for station, and gridded data, a simple procedure for calibrating the empirical coefficients in the alternative PET methods, and estimation of PET under warming and elevated atmospheric CO₂ concentration. Further discussion on the advantages of using *PyEt* estimates as input for hydrological models, sensitivity/uncertainty analyses, and hind/forecasting studies, especially in data-scarce regions, is provided.

15 1 Introduction

Evaporation — the process by which water is converted from its liquid to vapor phase — is a central component of the global hydrological cycle (Katul et al., 2012). Evaporation has far-reaching impacts on both human societies and ecosystems (Oki and Kanae, 2006; Fisher et al., 2011). In the remainder of this paper, the term evapotranspiration (ET) is used to refer to the total evaporation flux from soil and water bodies (evaporation) and vegetated surfaces (transpiration; Allen et al., 1998; Dingman, 2015). Information about the magnitude of the ET flux is important across different geoscience disciplines: it assists in predicting irrigation demands and crop water requirements in agriculture, supports efficient water resources management, guides operational strategies in hydropower and meteorological studies, and plays a crucial role in ecological research and climate-change impact assessments. Given that climate change — through warming and elevated CO₂ concentrations — is

set to alter evapotranspiration, the need for its accurate estimation is of paramount importance, as it affects our understanding and assessment of past, present, and potential future impacts on ecosystem functioning (Milly and Dunne, 2016; Yang et al., 2019; Caretta et al., 2022).

Evapotranspiration (~~ET~~) can hardly be measured directly (Wang and Dickinson, 2012; Jensen and Allen, 2016), and is therefore commonly estimated using (semi-)empirical formulas from other, more easily obtained meteorological variables such as temperature, wind speed, and radiation. Over time, dozens of methods have been proposed and applied. Each of these methods generally results in slightly different estimates of evapotranspiration, depending on the methods and data used (Oudin et al., 2005; McMahon et al., 2013; Xu and Singh, 2000, 2001; Lemaitre-Basset et al., 2022). Most of these formulas estimate either the reference crop evapotranspiration (ET_0), which is ET from a reference surface or crop that is not short of water (Allen et al., 1998), or the potential evapotranspiration (PET), which is the maximum rate of ET that would occur given a sufficient water supply (Xiang et al., 2020). Potential evapotranspiration is determined by meteorological conditions, whereas water availability determines if actual evapotranspiration occurs at its potential rate (Jensen and Allen, 2016). Differences in the potential evapotranspiration estimate may cascade through a modeling chain and ultimately impact the results of a study. For example, ~~Prudhomme and Williamson (2013); Lemaitre-Basset et al. (2022); Bormann (2010)~~ Prudhomme and Williamson (2013), Lemaitre-Basset et al. (2022), and Bormann (2010) showed that the method used affects the results from hydrological climate change impact studies. Similarly, the estimation of water demand for efficient crop and irrigation management depends on potential evapotranspiration, and may thus be impacted by the methods used (Kumar et al., 2012).

To account for the structural uncertainty of the different PET models, it has been recommended to use multiple methods (Seiller and Anctil, 2016; Beven and Freer, 2001; Velázquez et al., 2013). Such an approach can help improve the understanding of the effect of model uncertainty on PET estimates in, for example, historical climate studies (Zhou et al., 2020; Dakhlaoui et al., 2020; Yang et al., 2019) and climate change impact studies (Bormann, 2010; Seiller and Anctil, 2016; Gharbia et al., 2018; Shi et al., 2020). Climate change impact studies often rely on climate projection data or global observational datasets, which are generally available in a gridded format (i.e., netCDF, GRIB), thus requiring tools that can efficiently process such data. Some of these datasets also contain only a limited set of observed or projected meteorological variables, requiring PET estimation methods that use fewer inputs. It may also be necessary to account for environmental variables that change over time and impact the evapotranspiration, such as vegetation changes and increases in atmospheric CO_2 concentrations (Fatichi et al., 2016; Ainsworth and Rogers, 2007; Vremec et al., 2022). ~~To efficiently account for the structural uncertainty, the software used to compute PET should ideally have multiple methods available, and be flexible enough to deal with such changing environmental variables.~~ Studies like those mentioned above, require software programs that 1) include multiple PET estimation methods, 2) be flexible in adjusting input parameters (e.g., empirical coefficients, crop data, and meteorological inputs), and 3) be applicable to both time series and gridded data, given the spatial nature of many of these studies.

~~The FAIR principles of findability, accessibility, interoperability, and reusability are of crucial importance for the credibility and reproducibility of scientific studies (Barker et al., 2022). Tools~~ Existing tools for calculating evapotranspiration, such as 'Evapotranspiration' in R (Guo et al., 2016), and "PyETo" (Richards, 2019) and 'pyfao56' in Python (Thorpe, 2022), ~~have significantly advanced the field by offering valuable computational solutions. While these tools represent important advances~~

in research software development, there is an opportunity to enhance their compliance are primarily designed for station-based
60 time-series data. This limits their applicability with gridded datasets. While Peterson et al. (2020) extended 'Evapotranspiration'
to 'AWAPer' to process gridded data, its use is limited to Australia. For the large community of geoscientists working with
Python, the number of available PET methods from existing packages is limited (3 for PyETo and 1 for pyfao56), compared to
21 methods in the R package. This highlights a gap in the availability of a software for the estimation of multiple PET methods
for both time-series and gridded data, with the input parameter flexibility required for advanced studies on PET. Given the
65 increasing need to understand and predict environmental changes accurately across the globe, the availability of such software
is of paramount importance for the geoscience community.

Opportunities also exist to further align these tools with the FAIR standards of findability, accessibility, interoperability, and
reusability for research software, crucial importance for the credibility and reproducibility of scientific studies (Barker et al., 2022)
. This involves improving methodological testing through continuous integration, ~~expanding compatibility with various data~~
70 ~~formats like NetCDF to support regional and global ET estimation, inclusion of~~ inclusion of additional alternative PET meth-
ods, and enabling more flexibility in adjusting internal empirical coefficients. Such enhancements not only adhere to best
practices in software development, but also broaden the scope and applicability of these tools in diverse geoscientific contexts.
The refinement and development of evapotranspiration estimation tools that fully embrace the FAIR principles are therefore
crucial steps toward advancing the field, ensuring more reliable and comprehensive research outcomes in the face of evolving
75 scientific needs (Wood et al., 1998; DeJonge and Thorp, 2017).

In this paper we introduce *PyEt*, an open-source Python package for the estimation of potential evapotranspiration. The aim
of *PyEt* is to provide researchers and practitioners with a wide variety of tested, documented, and flexible Python functions ~~for~~
~~the estimation of potential evapotranspiration~~ that support multiple PET methods for both station and gridded data. All methods
have a common application programming interface, allowing users to easily test different PET models for their application and,
80 if desired, address structural uncertainty and changing conditions. The majority of the implemented methods are benchmarked
against literature values and tested with continuous integration to ensure the correctness of the implementation. Allowing
different types of input data, *PyEt* is also applicable in regions with sparsely distributed measurement stations, where standard
meteorological data (e.g., wind, relative humidity) are often unavailable. The software is available under MIT-license from the
Python Package Index (PyPI) (Vremec and Collenteur, 2022), and developed as a community project on GitHub (www.github.
85 com/pyet-org/PyEt).

The remainder of this paper is structured as follows. In the next section, the software design, capabilities, and benchmarking
tests are described. The third section introduces the software through four examples, showing potential future users how to
apply *PyEt* in real-world applications. These examples concentrate on addressing practical problems commonly faced by
geoscientists in their daily work. The fourth section discusses future potential applications of *PyEt*, and how we think it can
90 help the scientific community improve the estimation of potential evapotranspiration. In the fifth and final section, conclusions
and future plans are outlined.

2 PyEt Python Package

2.1 Software design

The basic design principle for *PyEt* was to build a software that is intuitive and easy-to-use by novice users with little programming experience, yet flexible enough to allow advanced users to perform more complex analyses. The software uses a modular design, with formulas shared by different PET methods implemented as a single function. This reduces the amount of code and makes it easier to maintain the software and implement new methods. All the PET methods are intended to work with the minimum input data required by the PET models (e.g., radiation, temperature), but also allow more user input if such data is available and allowed by the PET method (e.g., humidity, surface resistance in the Penman-Monteith model). Utility functions are available to the user or are called internally to compute unavailable variables (e.g., solar radiation from latitude value). Moreover, the constants in the empirical PET formulas (e.g., the Stefan Boltzmann constant) are function arguments with default values from the literature, which may also be changed by the user to adapt the empirical relationship to another region. Finally, the available methods should work for both station (1D) and gridded data (2D/3D).

PyEt is part of the wider Python ecosystem, and depends on three widely used and well-developed Python packages from the scientific Python stack: Numpy (Harris et al., 2020), Pandas (McKinney, 201), and Xarray (Hoyer and Hamman, 2017). The input and output data of *PyEt* are formatted as time series data in `Pandas.Series` or `Xarray.DataArrays`, which allows using all the Pandas and/or Xarray functions on the data (Harris et al., 2020; McKinney, 201; Hoyer and Hamman, 2017). These functions include gap-filling and selection functions for interpolation, resampling, clustering, and many more. Being part of a wider ecosystem, users can leverage other Python packages for visualization (e.g., Matplotlib (Hunter, 2007), MetPy (May et al., 2022)) and optimization and uncertainty analyses (Scipy (Virtanen et al., 2020), SpotPy (Houska et al., 2015)).

The software is hosted and developed on the GitHub platform, and distributed under MIT-license through the Python Packaging Index (PyPI). Documentation and example applications are available on a dedicated ReadTheDocs website (<http://pyet.readthedocs.io>). The documentation for individual methods is also directly available in Python from the documentation strings. Each release of *PyEt* is automatically stored in the Zenodo repository and assigned a Digital Object Identifier (DOI). As such, *PyEt* complies with many of the recommendations for good research software development as given in, for example, Hutton et al. (2016) and the FAIR4RS (FAIR for Research Software) principles (Barker et al., 2022). The scripts or the Jupyter notebooks used to apply *PyEt* improve the reproducibility and provide a transparent report of the entire calculation process (Kluyver et al., 2016).

2.2 Implemented methods and benchmarking

Twenty methods are currently implemented in *PyEt* for the estimation of daily potential evapotranspiration. ~~In this paper, Apart from the Penman-Monteith method is considered the standard for estimating potential evapotranspiration due to its widespread use and acceptance. All other methods implemented~~, which is considered as the standard by the Food and Agriculture Organization (FAO) (Allen et al., 1998) and the World Meteorological Organization (WMO), multiple alternative methods are




Figure 1. Scatter plots showing estimated PET with *PyEt* against PET values estimated with the R package *Evapotranspiration* from Guo et al. (2016), and literature values from Allen et al. (1998), McMahon et al. (2013), and Schrödter (1985).

125 [also available](#) in *PyEt* will be referred to as alternative methods. An overview of these methods and the required input data is provided in Table 1. Depending on the method, different (amounts of) input data are required to compute the potential evapotranspiration. It is often also possible to provide different input data to the same method (e.g., the average or the minimum and maximum daily temperatures), or even that some input data is optional (as described in the footnotes of Table 1). In the case of optional input data, utility functions are used internally to estimate that data. In the example of the Penman-Monteith method, 130 solar radiation does not necessarily need to be provided by the user and can be estimated from the latitude and actual duration of sunshine hours instead.

The *PyEt* project is intended to be used by a wide community, and any errors in the code may have consequences for other studies applying *PyEt* to obtain PET estimates. Special attention was therefore paid to benchmark the available methods to published literature values and data from well-known research and meteorological institutes (Allen et al., 1998; McMahon 135 et al., 2013; Schrödter, 1985; Walter et al., 2000). These benchmarks are also implemented in the continuous integration and tested using the *unittest* testing framework (unittest, 2022). This ensures that the benchmarks are satisfied each time the software is updated in the future. New methods added to *PyEt* will be required to be accompanied by the appropriate benchmark data and tests. Figure 1 shows the results for each benchmarked method, indicating that the PET estimates from all these methods are equal to the benchmark values (i.e., all values are on the 1:1 line). Despite our best efforts, we acknowledge here that 4 140 methods have not (yet) been benchmarked due to a lack of appropriate data.

2.3 The Penman-Monteith equation

To illustrate how *PyEt* might be helpful in several applications, one of the most used and most versatile PET methods is discussed in more detail, the Penman-Monteith method (Monteith, 1965). In its different forms, the Penman-Monteith equation can be used to estimate reference crop evapotranspiration (Allen et al., 1998; Walter et al., 2000), potential evapotranspiration 145 (Monteith, 1965), and potential crop evapotranspiration. The latter is defined as "the maximum value of ET from a specific crop

Table 1. Data requirements for different PET or ET_0 models, the corresponding *PyEt* function, and if benchmarking of the method was performed. The literature-references include both the original publications of the models and the manuscripts from which the equations were taken: [1] McMahon et al. (2013), [2] Oudin et al. (2005) [3] Xu and Singh (2001), [4] Ansoorge and Beran (2019); Rosenberry et al. (2004), [5] Schrödter (1985), [6] Schiff (1975), [7] Jensen and Allen (2016), [8] Xu and Singh (2000)

Method name ⁰	<i>PyEt</i> function	Climate data				Location		Bench.	Literature
		T	RH	R	u_2	Lat.	El.		
Penman-Penman-Monteith	<i>penmanpm</i>	✓ ^a	✓ ^{b,c}	✓ ^d	✓	✓ ^d	✓ ^e	✓	Penman (1948) Monteith (1965)
Penman-Monteith-ET₀: ASCE-PM	<i>pm_asce</i>	✓ ^a	✓ ^{b,c}	✓ ^d	✓	✓ ^d	✓ ^e	✓	Monteith (1965) Walter et al. (2000)
ET₀: ASCE-PM-FAO-56	<i>pm_ascefao56</i>	✓ ^a	✓ ^{b,c}	✓ ^d	✓	✓ ^d	✓ ^e	✓	Walter et al. (2000) Allen et al. (1998)
ET₀: FAO-56-Penman	<i>pm_fao56penman</i>	✓ ^a	✓ ^{b,c}	✓ ^d	✓	✓ ^d	✓ ^e	✓	Allen et al. (1998) Penman (1948)
Priestley-Taylor	<i>priestley_taylor</i>	✓	✓ ^h	✓ ^h	-	✓ ^h	✓ ^e	✓	Priestley and Taylor (1972)
Kimberly-Penman	<i>kimberly_penman</i>	✓ ^a	✓ ^{b,c}	✓ ^d	✓	✓ ^d	✓ ^e	-	Wright (1982)
Thom-Oliver	<i>thom_oliver</i>	✓ ^a	✓ ^{b,c}	✓ ^d	✓	✓ ^d	✓ ^e	-	Thom and Oliver (1977)
Blaney-Criddle	<i>blaney_criddle</i>	✓	- ⁱ	- ⁱ	- ⁱ	✓	-	✓	Blaney and others (1952), [1, 3, 5]
Hamon	<i>hamon</i>	✓	-	-	-	✓	-	✓	Hamon (1963), [2, 4]
Romanenko	<i>romanenko</i>	✓	✓	-	-	-	-	✓	Romanenko (1961), [3]
Linacre	<i>linacre</i>	✓ ^j	-	-	-	-	✓	✓	Linacre (1977), [3]
Haude	<i>haude</i>	✓	✓ ^k	-	-	-	-	✓	Haude (1955), [6]
Turc	<i>turc</i>	✓	✓	✓	-	-	-	✓	Turc (1961), [8]
Jensen-Haise	<i>jensen_haise</i>	✓	-	✓ ^l	-	✓ ^l	-	✓	Jensen and Haise (1963), [2, 7]
McGuinness-Bordne	<i>mcguinness_bordne</i>	✓	-	-	-	✓	-	✓	McGuinness and Bordne (1972), [8]
Hargreaves	<i>hargreaves</i>	✓ ^m	-	-	-	✓	-	✓	Hargreaves and Samani (1982), [1, 7]
ET ₀ : FAO-24	<i>fao_24</i>	✓	✓	✓	✓	-	✓ ^e	-	Jensen et al. (1990)
ET ₀ : Abtew	<i>abtew</i>	✓	-	✓	-	-	-	✓	Abtew (1996), [8]
Makkink	<i>makkink</i>	✓	-	✓	-	-	✓ ^e	✓	Makkink (1957), [1]
Oudin	<i>oudin</i>	✓	-	-	-	✓	-	-	Oudin et al. (2005)

⁰ The corresponding literature to each method is provided in Table A1, in Appendix. ^a T_{max} and T_{min} can also be provided. ^b RH_{max} and RH_{min} can also be provided. ^c If actual vapor pressure is provided, RH is not needed. ^d Input for radiation can be (1) Net radiation, (2) solar radiation or (3) sunshine hours. If (1), then latitude is not needed. If (1, 3) latitude and elevation is needed. ^e One must provide either the atmospheric pressure or elevation. ^f The PM method can be used to estimate potential crop evapotranspiration, if leaf area index or crop height data is available. ^g The effect of CO₂ on stomatal resistance can be included using the formulation of Yang et al. 2019. (Yang et al., 2019). ^h If net radiation is provided, RH and Lat are not needed. ⁱ If method==2, u_2 , RH_{min} and sunshine hours are required. ^j Additional input of T_{max} and T_{min} , or T_{dew} . ^k Input can be RH or actual vapor pressure. ^l If method==1, latitude is needed instead of R_s . ^m T_{max} and T_{min} also needed.

type having specific properties under conditions of full soil water supply, but not necessarily having a saturated surface" by Jensen and Allen (2016). Through the introduction of the Penman-Monteith equation by Monteith (1965), a broad applicability to different surfaces and vegetation types was achieved (Jensen and Allen, 2016). This was done by implementing the plant aerodynamic resistance (r_a) and the surface resistance (r_s) in the PET formula.

150 Users of *PyEt* can include leaf/canopy cover measurements (Leaf Area Index – LAI) to calculate surface resistance (r_s), thereby accounting for the effects of crop management and phenology on ET. A modified stomatal resistance model also allows for the inclusion of the sensitivity of the stomatal resistance (r_l) to the atmospheric CO₂ concentration (as shown in, for example, Yang et al. 2016):

$$r_s = \frac{r_l(\text{CO}_2)}{0.5\text{LAI}} = \frac{r_{l-300} \times \{1 + S_{r_l-\text{CO}_2} \times (\text{CO}_2 - 300)\}}{0.5\text{LAI}}$$

155 where $S_{r_l-[\text{CO}_2]}$ ppm⁻¹ is the relative sensitivity of r_l to Δ CO₂ and r_{l-300} s m⁻¹ is the reference stomatal resistance when atmospheric CO₂ concentration is 300 ppm. The relative sensitivity of r_l to Δ CO₂ represents the change in r_l per ppm increase in CO₂ concentration.

If measurements of crop height exist, these data can be used to calculate the aerodynamic resistance to vapor and heat transfer (r_a) to represent the effects of crop phenology on PET:

$$160 \quad r_a = \frac{\ln \left[\frac{z_m - d}{z_{om}} \right] \ln \left[\frac{z_h - d}{z_{oh}} \right]}{k^2 u_z}$$

where z_m is the reference level at which the wind speed is measured; z_h is the height of the temperature and humidity measurements; k is the von Karman constant ($= 0.41$), u_z is the measured wind speed (Allen et al., 1998) and d is the zero plane displacement height, taken as $0.67h_c$ m; z_{om} is the roughness parameter for momentum ($= 0.123h_c$) m and z_{oh} is the roughness parameter for heat and water vapor ($= 0.1z_{om}$) m (Jensen and Allen, 2016).

165 2.3 Performance

The computational efficiency of various PET methods was assessed by examining the computation times in relation to the time series length and the number of cells in an Xarray dataset. Computation time was evaluated by running all models on a benchmark configuration (with time series of varying lengths) using a 12th Gen Intel Core i7-1255U processor with 10 cores and 12 logical processors. All Xarray.DataArrays cover a period of 1 year, while the spatial resolution changes. This comparison highlights the trade-offs between computational complexity and data size, but also demonstrates the performance of the methods.

Figure 2 demonstrates that all *PyEt* methods maintain computational times below 1 second for time series data with lengths ranging from 10 to 10,000 days. For multi-dimensional data, the computation time does not exceed 10 seconds for larger Xarray.DataArrays up to 100,000 cells. Moreover, the results in Figure 2 show that the problem scales well, and does not take proportionally more time for larger data sets. Notably, the Penman-Monteith and Priestley-Taylor methods exhibited the largest processing times, whereas methods like Jensen-Haise, Turc, Makkink, and Romanenko were faster. Future improvements will aim to increase this efficiency, in particular to support faster calculations in large-scale global studies.




Figure 2. Computational efficiency of different PET methods: This figure shows the comparative processing times of different PET estimation methods with respect to regarding the length of the time series (a) and the size of the Xarray data (b).

3 Example use cases

180 Below, four **example**-use cases of *PyEt* are presented to illustrate how the software can be used. The first example shows how to efficiently compute different potential evapotranspiration estimates using 20 various methods for station data. This example also illustrates how to use *PyEt* in general. The second example illustrates how to provide 3D estimates of PET using 3 different methods and gridded Xarray data. The third example shows how to calibrate different PET methods to local conditions and use the calibrated formula for hindcasting. The fourth and final example illustrates a workflow to account for the effects of

185 warming and elevated CO₂ in climate change impact studies. The source code for these and other examples can be found in a Zenodo repository related to this paper (Matevz Vremec and Raoul Collenteur, 2024).

3.1 Example 1: Estimation of PET from station data

In this example, potential evapotranspiration is estimated for the town of De Bilt in The Netherlands using data provided by the Royal Netherlands Meteorological Institute (KNMI). The reference method used by the KNMI for the estimation of

190 potential evapotranspiration is the Makkink method, also implemented in *PyEt*. The PET computed with the Makkink method is compared to the PET values from all other methods in *PyEt*. A number of Several steps are taken in a Python script to estimate PET. The code implementing these steps is shown in the code example below. *PyEt* provides a convenient way to compute the PET with all available methods, `pyet.calculate_all()`:

1. Import the necessary Python packages.

195 `import pandas as pd`

```
import pyet
```

2. Load the meteorological data.

```
meteo = pd.read_csv("meteo.csv",  
                    index_col=0, parse_dates=True)
```

200 3. Determine the necessary input data for the PET model.

```
tmean, tmax, tmin, rh, rs, wind, \  
pet_knmi = (meteo[col] for col in  
            meteo.columns)  
lat = 0.91 # latitude  
205 elev = 4 # elevation
```

4. Estimate the potential evapotranspiration with all methods or the method of choice.

```
pet_df = pyet.calculate_all(tmean,  
                             wind, rs, elev, lat, tmax, tmin, rh)  
pet_mak = pyet.makkink(tmean, rs,  
210 elevation=elev)
```

5. Visualize and analyze the results.

```
pet_df.plot()  
pet_df.boxplot()  
pet_df.cumsum().plot()
```

215 The results from this analysis are shown in Figure 3. From these visualizations, it is clear that the potential evapotranspiration depends on the chosen method. This can accumulate up to a 35% deviation of the estimated annual flux from the mean in this example. Such substantial differences between the estimated fluxes motivate the use of multiple methods (ensemble modelling) (Beven and Freer, 2001; Krueger et al., 2010; Shi et al., 2020; Oudin et al., 2005). This example showed how *PyEt* can be used efficiently for this task.

220 **3.2 Example 2: Estimate PET for gridded data**

Gridded 3-dimensional data (x, y, t) obtained from satellites, radar imagery, or post-processed products is rapidly becoming widely available. More and more public data sets exist with global PET estimates at 0.1 degree resolution (e.g., Martens et al., 2017; Xie et al., 2022). *PyEt* also supports such gridded data, as illustrated here for the E-OBS gridded data set (Cornes et al.,




Figure 3. Potential evapotranspiration estimates for the year 2018 computed with all available PET methods, plotted as (a) time series, (b) box plots (b), (c) cumulative PET.

2018) for Europe. The application of *PyEt* on gridded datasets is illustrated for the FAO-56, Makkink, and Hargreaves method. `Xarray.DataArrays` are used as input data instead of `Pandas.Series`. *PyEt* methods will return the same data type, a `Xarray.DataArray`. The workflow is comparable to that in the first example, except that now the individual PET methods were used.

The results for the three methods and three time steps are shown in Figure 4. These again show that, depending on the PET method, results may differ, also spatially. Looking more closely at Figure 4, we can observe that the FAO-56 and Makkink method do not compute PET in eastern parts of Europe. The data do not include relative humidity and solar radiation for these areas, and thus PET cannot be computed using the FAO-56 or Makkink method. If NaN (not-a-number) values are present in the required input data for a *PyEt* method, the method also returns a NaN value. The Hargreaves method, on the other hand, does not require solar radiation or relative humidity data. It can therefore be used to compute PET in the eastern parts of Europe. This example showed how *PyEt* can be applied to estimate PET using gridded data, and demonstrated the benefits of using alternative PET methods when data such as radiation or relative humidity are missing.

3.3 Example 3: Calibration of PET models

The available input data often does not suffice to compute potential evapotranspiration with the Penman-Monteith equation. This can be the case in data-scarce regions or time periods, or when using historical data or data from climate models. In such cases, alternative PET methods can be calibrated to the estimates obtained from the Penman-Monteith equation for a period when sufficient data is available. The calibrated method can then be used to estimate PET in periods of data scarcity. As concluded by several authors (Jensen and Allen, 2016; Valipour, 2015; Yang et al., 2021; Dlouhá et al., 2021), calibration




Figure 4. Daily PET estimates for Europe from 2018-6-6 to 2018-6-8 using meteorological data obtained from the E-OBS dataset (Cornes et al., 2018).

of alternative models is often crucial to ensure that the model fits the regional climate. In this example, it is shown how the calibration of temperature-based PET models affects the model uncertainty for studies focusing on current and past climates.

The approach is illustrated for the town of Graz, Austria, where the input data required for Penman-Monteith are only available from 2000 to 2021. Imagine, however, that for our study we also need potential evapotranspiration data for the period 1961 to 2021, but only temperature data is available (e.g., from the Spartacus temperature dataset (Hiebl and Frei, 2016)). Several steps are taken to calibrate the following five temperature-based methods: Oudin, Hargreaves, McGuinness-Bordne, Hamon, and Blaney-Criddle. First, the PET for the period 2000-2021 is computed using the Penman-Monteith equation. In the second step, the coefficients of the temperature-based PET equations are estimated by calibrating the estimated PET from temperature-based methods to the Penman-Monteith PET. Calibration is done by minimizing the sum of the squared residuals between these two PET estimates, using SciPy's (Virtanen et al., 2020) *least_squares* method. In the third and final step, these calibrated coefficients are used to estimate the PET for the period 1961-2021.

Figure 5 shows the computed PET with default (row 1) and the calibrated coefficients (row 2). The model bias (mm/day) and the coefficient of determination (R^2) between simulated and observed (Penman-Monteith) PET show an improved model fit for all methods after calibration. The use of calibrated methods reduces the model bias, which is visually illustrated by the




Figure 5. Density scatter plots comparing simulated and observed (FAO-56) PET for uncalibrated (row 1) and calibrated models (row 2). The last column shows the annual PET sums for the period 1961-2021 using uncalibrated (row 1) and calibrated models (row 2).

annual PET flux (composed of daily values) in the last column of Figure 5. Using the Spartacus temperature dataset (Hiebl and Frei, 2016), PET can now be estimated back to 1961 using the calibrated alternative PET methods.

3.4 Example 4: The effect of CO₂ on future PET estimates

In this example, it is shown how to account for changing environmental conditions affecting the PET flux when modelling the effects of climate change. Under a warmer and CO₂ richer future (Caretta et al., 2022), potential evapotranspiration tends to increase with increasing temperature (and vapor pressure deficit). A reduction in PET is expected under elevated CO₂ due to an increased stomatal resistance (Field et al., 1995; Ainsworth and Rogers, 2007). The increase in CO₂ is still commonly ignored in PET models employed for climate change studies, although excluding its stomatal effect may lead to an overestimation of PET (Kingston et al., 2009; Milly and Dunne, 2016; Vremec et al., 2022; Riedel et al., 2023). The effect of temperature increases on PET can be easily modelled with all available PET methods, as temperature is an input for all methods. The CO₂ stomatal effect, however, can only be directly accounted for with the Penman-Monteith method (Liu et al., 2022). Using a CO₂-dependent stomatal resistance model implemented in *PyEt* (Yang et al., 2019), the effect of elevated CO₂ on stomatal resistance can be considered (see Eq. A1A2). When calculating PET with alternative methods, Kruijt et al. (2008) and Trnka et al. (2014) argued that an adjustment factor for the atmospheric CO₂ concentration (f_{CO_2}) can be used to account for the effect of elevated CO₂ concentrations on PET. The scaling factor can be obtained from literature values (Kruijt et al., 2008; Trnka et al., 2014). Alternatively, the factor can be calibrated using the Penman-Monteith equation together with the CO₂-dependent stomatal resistance model (Eq. A1) to match the local climate and vegetation:

$$\begin{aligned}
PET_{CO_2} &= f_{CO_2} PET_{300} \\
&= (1 + S_{PET_{CO_2}} (CO_2 - 300)) PET_{300}
\end{aligned}
\tag{1}$$

where $S_{PET_{CO_2}}$ is the relative sensitivity of PET to CO_2 , PET_{300} is the computed Penman-Monteith estimate at 300ppm [CO_2] (preindustrial concentration), while PET_{CO_2} is the computed Penman-Monteith estimate under elevated CO_2 concentration (Yang et al., 2019). Such relationships can be easily implemented in *PyEt*, and f_{CO_2} can be obtained by calculating PET_{300} and PET_{CO_2} with the Penman-Monteith equation (Eq. A1) at ambient and elevated CO_2 concentration, respectively.

Building on the previous example, the Graz study area served as a practical example to demonstrate the application of the calibrated models in assessing the impact of warming and elevated CO_2 concentration on PET based on the projected increase in temperature and CO_2 concentration from the representative concentration pathways (RCPs) (Van Vuuren et al., 2011). Daily PET was calculated for each RCP scenario (2.6, 4.5, 6.0, and 8.5) by adding the projected increase in temperature and CO_2 concentration to the existing data for 2020-2021. Figure 6 shows the increase in the average annual PET (aggregated from daily values) under warming and elevated CO_2 concentrations according to the RCP scenarios. In figure 6c, the effects of elevated CO_2 concentration on PET were neglected, and only increases in temperature were considered. Similar to Milly and Dunne (2016), Yang et al. (2019), and Vremec et al. (2022), this example shows that neglecting the effect of elevated CO_2 on PET (Fig. 6-c) can lead to overestimation of PET under future conditions.




Figure 6. Projected increase in temperature (6-a) and atmospheric CO_2 concentration (6-b) under the RCP scenarios, and calculated increase in the average annual PET with warming (6-c), and PET with warming and elevated CO_2 concentration (6-d). The uncertainty bounds represent the 5th-95th percentile of the PET model ensemble.

4 Discussion

4.1 Improved handling of PET in scientific studies

Evapotranspiration data from lysimeter or eddy correlation measurements (Pastorello et al., 2020) are rare and, if available
290 at all, only locally available for relatively short time periods. Thus, there is a widespread need to estimate evapotranspiration
from more readily available meteorological data using (semi-)empirical approaches. In general, these approaches follow three
steps, as outlined for example by Allen et al. (1998). Firstly, the potential evapotranspiration of a reference surface (hence
reference evapotranspiration) is estimated using meteorological data. Secondly, a crop coefficient may be applied to transform
the reference evapotranspiration into the potential crop evapotranspiration. Thirdly, a soil-water balance approach is used to
295 account for reduced actual evapotranspiration if the soil-water storage is depleted. *PyEt* is designed to perform the first two
steps. It can be easily complemented by soil-water balance approaches to calculate actual evapotranspiration. Hydrological
models, however, often use PET directly as input.

Rainfall-runoff models represent one type of hydrological model where PET is commonly used as an input, either as gridded
data in distributive models, or as a spatially aggregated values in lumped-parameter models. Some studies (e.g., Andréassian
300 et al., 2004; Oudin et al., 2005; Sperna Weiland et al., 2012) found that PET had little impact on the performance of such
models, and thus advocated the use of simplistic PET models. However, Jayathilake and Smith (2021) found that model
performance was clearly sensitive to PET at sites where evapotranspiration was water limited. More importantly, the choice of
the PET model has been shown to affect the results of hydrological projections in climate change impact assessments (Kay and
Davies, 2008; Seiller and Anctil, 2016; Dallaire et al., 2021; Lemaitre-Basset et al., 2022)). PET is expected to be even more
305 influential in the assessment of groundwater recharge (e.g., Bakundukize et al., 2011) and crop water demands (e.g., Webber
et al., 2016)), which – compared to runoff – are more directly linked to evapotranspiration. Thus, the selection of appropriate
PET models needs to account for the research context and variable of interest.

Bormann (2010) found that PET models that are based on the same or similar climate variables exhibit different sensitivity
to observed climate change. This finding suggests that appropriate PET models need to be specifically selected for the given
310 region of interest. Guo et al. (2017) provides pointers to examining which variables are likely to be the most important for a
particular location. For more detailed insights into the application of individual PET methods across various climates, refer to
Table A2 and the studies by Allen et al. (1998); McMahon et al. (2013); Jensen and Allen (2016); Yang et al. (2021); Pimentel
et al. (2023). The comparison of PET estimates for Europe shown in Figure 4 illustrates the spatial variability of differences
in PET estimates obtained from different methods; as can be seen, the magnitude and pattern of PET estimates are similar in
315 some regions (e.g., Scandinavia) but differ more strongly in others (e.g., Southeast Europe).

As indicated above, the performance of PET models may vary depending on the region considered. Approaches that were
found applicable in one region may perform less well in other regions. In this case, PET models can be calibrated to a reference
data set by adjustment of the coefficients in the model equation, as shown in the third example. The reference data set can
either be observed evapotranspiration (e.g., from lysimeters) or PET obtained from a model considered to be reliable. This
320 has been illustrated by Example 3, where the coefficients of temperature-based models were adjusted to achieve the best fit

to the Penman-Monteith model. This approach can also be used to obtain consistent spatial distributions of PET. As shown in Example 2 (Figure 4), the limited data availability for Eastern Europe did not allow the application of the FAO-56 or Makkink method, while sufficient data was available for the Hargreaves method. Thus, one may consider calibrating the latter to one of the former methods where these are applicable, and only then apply it to obtain estimates for the entire region. For a more advanced calibration procedure, see for example Haslinger and Bartsch (2016).

~~In many cases~~ Often, the range of PET models that can potentially be employed is pre-determined by data availability. This may be the case if historical records of climate data are to be used for the PET estimation, for example, as many weather stations do not measure all climate variables included in the Penman-Monteith equation. Yet, this is also often the case in assessments of hydrological impacts of climate change if projected climate variables have high uncertainty. Lai et al. (2022), for example, concluded that the high uncertainty of wind speed projected in complex terrain may increase the uncertainty in PET, whereas air temperature and solar radiation have low uncertainty and thus should be the parameters preferred in the PET model. Given the climate variables for which data is available, Table 1 can be used to identify the PET models that come into consideration. However, it is advised to evaluate the assumptions and limitations of the individual methods regarding their applicability in the given case. Please refer to the comments in Table A2 and the references in Table 1 for this purpose. We generally recommend applying all models that have been identified as suitable (PET model ensemble), but the purpose and specific implementation of such a multi-model approach will depend on the research context. Example 4 (section 3.4) illustrated how PET model ensembles can be used to include model uncertainties in PET projections under warming and elevated atmospheric CO₂ concentration. Since the latter effect is frequently excluded in hydrological projections, Milly and Dunne (2016) and Yang et al. (2019) advocate the inclusion of the effect of elevated CO₂ on stomatal resistance when estimating PET under warming and elevated atmospheric CO₂ concentrations.

To improve reliability and efficiency in estimating PET, it is crucial to use a reproducible workflow. Scripts provide an efficient way to document the modelling process and are an important step towards full reproducibility. As shown in the examples, Jupyter Notebooks (Kluyver et al., 2016) provide a solution for publishing code, results, and explanations in a single document. As such, the presented package and its application in this paper are in line with the steps suggested by Hutton et al. (2016) to improve reproducibility in hydrological studies. To speed up adaptation of the methods and allow a faster transfer between research teams, formal procedures such as benchmarking (e.g., Maxwell et al., 2014) can help to ensure confidence in key complex codes.

4.2 Building the PyEt community and outlook

As a community project, the success of *PyEt* depends on the uptake from and interaction with the community. This, in turn, depends on the ease of use and the trust in the project. Emphasis was put on designing a user-friendly, well-documented software, including various user examples, and extensive benchmark testing using continuous integration. Since the initial launch of *PyEt*, the package has already seen a good community uptake. Apart from applications of the software in projects related to the Authors, which include estimating PET under conditions of warming and elevated CO₂ concentrations, assessing potential crop evapotranspiration, and providing inputs for hydrological models (e.g., Vremec et al., 2022; Forstner et al.,

355 2022; Collenteur et al., 2023; Jemeljanova et al., 2023), *PyEt* has also been independently used by other researchers. These studies have ~~utilized~~used *PyEt* for ~~a variety of~~various purposes: integrating PET estimates with machine learning models for enhanced analytical capabilities (Vaz et al., 2022), and combining them with software for computing groundwater recharge and the water balance (Hassanzadeh et al., 2023). The software has also played a role in evaluating hydroclimatic changes and generating regional and global PET estimates (e.g., Tercini and Mello Júnior, 2023; Aguayo et al., 2024). The quick uptake of
360 the software by the community ~~shows~~confirms the need for this software.

The primary channel for communication with the *PyEt* community is GitHub, which provides several options for discussions, tracking code issues, and code development. Users are encouraged to ask questions in GitHub discussions and to report potential issues, suggest improvements, and feature requests via the GitHub issue tracker. As a community project, we plan to continue to improve the existing code and develop new capabilities based on feedback and with help from the community. An
365 example of developments that are currently underway is the adaptation of the current methods to also work for hourly data, allowing the estimation of hourly PET. Other future work will focus on improvements in usability and the inclusion of other alternative methods.

5 Conclusions

This paper introduced *PyEt*, a Python package for the estimation of daily potential evapotranspiration (PET). The package
370 enables the inclusion of model uncertainty and climate change in the PET estimation in a consistent, tested, and reproducible
environment. With *PyEt*, PET can be estimated using 20 different methods with just a few lines of Python code ~~not only for~~.
Unlike existing tools for PET calculation, which are designed for station-based time series, *PyEt* can also be applied to gridded
(1D) time series but also for (3D) gridded data sets, which goes beyond existing tools for PET calculation data sets. This is of
great practical relevance, particularly in climate impact studies, where gridded data sets are often used. The examples described
375 in this paper illustrate how *PyEt* can be used in geoscientific studies to (1) facilitate the characterization of model uncertainty
using a ~~multi-model~~ multimodel approach (model ensembles); (2) calibrate PET models and apply them in data-scarce regions
and time periods; (3) include the effects of warming and elevated atmospheric CO₂ concentrations. The use of Python scripts
and Jupyter Notebooks ensure reproducibility and provides a transparent report of the PET computation process. We believe
that *PyEt* will help improve the handling of PET and allow a more sophisticated and comprehensive consideration of PET in
380 environmental studies, particularly those related to climate change.

Code and data availability. The Jupyter Notebook and data used in this study are available in the "examples" folder of the GitHub repository
and also available on Zenodo (version v.1.3.1, DOI: 10.5281/zenodo.5896799). The authors welcome code contributions, bug reports, and
feedback from the community to further improve the software. *PyEt* is free and open-source software available under the MIT license. Source
code is available at the project's home page on GitHub. Full documentation is available on ReadTheDocs. *PyEt* is meant as a community
385 project, and the Authors welcome contributions and feedback to continue to improve and develop the project are welcome.

Appendix: PET Methods - Climate suitability and applications

Table A1: List of variables and symbols used in the paper and supplementary section

<u>Variable</u>	<u>Description</u>	<u>Units*</u>
<u>PET</u>	<u>Potential evapotranspiration</u>	<u>mm day⁻¹</u>
<u>T</u>	<u>Mean daily temperature</u>	<u>°C</u>
<u>T_{max}</u>	<u>Maximum daily temperature</u>	<u>°C</u>
<u>T_{min}</u>	<u>Minimum daily temperature</u>	<u>°C</u>
<u>T_{dew}</u>	<u>Mean daily dew-point temperature</u>	<u>°C</u>
<u>RH</u>	<u>Mean daily relative humidity</u>	<u>%</u>
<u>RH_{max}</u>	<u>Maximum daily relative humidity</u>	<u>%</u>
<u>RH_{min}</u>	<u>Minimum daily relative humidity</u>	<u>%</u>
<u>u₂</u>	<u>Wind speed measured at 2 m</u>	<u>m s⁻¹</u>
<u>R_n</u>	<u>Net radiation</u>	<u>MJ m⁻² d⁻¹</u>
<u>R_s</u>	<u>Incoming solar radiation</u>	<u>MJ m⁻² d⁻¹</u>
<u>R_a</u>	<u>Extraterrestrial daily radiation</u>	<u>MJ m⁻² d⁻¹</u>
<u>G</u>	<u>Soil heat flux</u>	<u>MJ m⁻² d⁻¹</u>
<u>n</u>	<u>Actual duration of sunshine</u>	<u>hour</u>
<u>N</u>	<u>Maximum possible duration of sunshine or daylight hours</u>	<u>hour</u>
<u>Elev</u>	<u>Elevation above sea level</u>	<u>m</u>
<u>lat</u>	<u>Latitude</u>	<u>radians</u>
<u>lat_{deg}</u>	<u>Latitude</u>	<u>degrees</u>
<u>p</u>	<u>Atmospheric pressure</u>	<u>kPa</u>
<u>λ</u>	<u>Latent heat of vaporization</u>	<u>MJ kg⁻¹</u>
<u>Δ</u>	<u>Slope of the saturation vapor pressure curve</u>	<u>kPa K⁻¹</u>
<u>ρ_a</u>	<u>Air density</u>	<u>kg m⁻³</u>
<u>ρ_w</u>	<u>Water density (= 1)</u>	<u>Mg m⁻³</u>
<u>c_p</u>	<u>Specific heat of dry air</u>	<u>MJ kg⁻¹ K⁻¹</u>
<u>e₀</u>	<u>Saturation vapor pressure of the air at T</u>	<u>kPa</u>
<u>e_s</u>	<u>Saturation vapor pressure of the air</u>	<u>kPa</u>
<u>e_a</u>	<u>Actual vapor pressure of the air</u>	<u>kPa</u>
<u>γ</u>	<u>Psychrometric constant</u>	<u>kPa K⁻¹</u>
<u>r_s</u>	<u>Bulk surface resistance</u>	<u>s m⁻¹</u>

Table A1: (continued)

<u>Variable</u>	<u>Description</u>	<u>Units*</u>
r_L	Bulk stomatal resistance	$s\ m^{-1}$
$S_{r_L=[CO_2]}$	Relative sensitivity of r_L to $\Delta [CO_2]$	ppm^{-1}
r_a	Bulk aerodynamic resistance	$s\ m^{-1}$
h_c	Crop height	m
LAI	Leaf area index	-
CO_2	Atmospheric CO_2 concentration	ppm
K_u	Unit conversion factor (=86400)	$s\ d^{-1}$
a_w	Penman wind coefficient	-
b_w	Penman wind coefficient	-
C_n	Numerator constant that changes with reference type	$K\ mm\ s^3\ Mg^{-1}\ d^{-1}$
C_d	Denominator constant that changes with reference type	$s\ m^{-1}$
α	Surface albedo	-
α_L	Priestley-Taylor coefficient	-
P_{y-h}	Percentage of actual day-light hours for the day compared to the number of day-light hour during the entire year	-
as1	Empirical coefficient for extraterrestrial radiation	-
bs1	Empirical coefficient for extraterrestrial radiation	-
a	Empirical coefficient for Net Long-Wave radiation	-
b	Empirical coefficient for Net Long-Wave radiation	-
k	Empirical/calibration coefficient	-

Table A2. Overview of PET methods: climate suitability, applications, and limitations. ET_0 - reference crop/surface ET, PET_C - potential crop/surface ET, PET_{OW} - potential ET for open water, PET_{SL} - potential ET for shallow lakes, PET_{RR} - potential ET for rainfall-runoff modelling. Based on Allen et al. (1998); McMahon et al. (2013); Jensen and Allen (2016); Yang et al. (2021); Pimentel et al. (2023)

Method name	Application	Climate	Limitations/Comments
Penman <u>Penman-Monteith</u>	PET_{OW}, PET_{SL} ET_0 , PET_C , PET_{RR}	All climates	High input data requirement
Penman-Monteith <u>FAO-56</u>	ET_0 , PET_C, PET_{RR}	All climates	High input data requirement
FAO-56 <u>Penman</u>	ET_0 PET_{OW} , PET_{SL} , PET_{RR}	All climates	High input data requirement
Priestley-Taylor	PET_C , PET_{RR}	Temperate/polar	Calibration recommended for semi-arid and arid regions; often underestimates in high vapor pressure deficit areas.
Kimberly-Penman	ET_0 -alfalfa	Temperate/continental	High input data requirement
Thom-Oliver	ET_0	Temperate/continental	High input data requirement
Blaney-Criddle	ET_0 , PET_{RR}	Temperate	Overestimates in calm, moist, shaded areas; underestimates in windy, dry, sunny ones.
Hamon	PET_C	All climates	Recommended regional calibration.
Romanenko	PET_C , PET_{RR}	All climates	Best recommended model for PET in China.
Linacre	PET_C	All climates	Recommended regional calibration.
Haude	PET_C	All climates	Recommended regional calibration.
Turc	ET_0 , PET_C	Humid	Underestimates in areas with large daily vapor pressure deficits.
Jensen-Haise	PET_0 , PET_{RR}	Continental	Recommended regional calibration.
McGuinness-Bordne	PET_C , PET_{RR}	All climates	Recommended regional calibration.

Appendix: PET Methods - equations

Penman-Monteith (*pm*)

390 Through the introduction of the Penman-Monteith equation by Monteith (1965), a broad applicability of PET estimation to different surfaces and vegetation types was achieved (Jensen and Allen, 2016). This was done by implementing the plant aerodynamic resistance (r_a) and the surface resistance (r_s) in the PET formula:

$$PET = \frac{1}{\lambda \rho_w} \left[\frac{\Delta(R_n - G) + \rho_a c_p K_u (e_s - e_a) / r_a}{\Delta + \gamma \left(1 + \frac{r_s}{r_a}\right)} \right] \quad (A1)$$

Users of *PET* can include leaf/canopy cover measurements (Leaf Area Index - LAI) to calculate surface resistance (r_s), thereby accounting for the effects of crop management and phenology on PET. A modified stomatal resistance model also allows for the inclusion of the sensitivity of the stomatal resistance (r_l) to the atmospheric CO₂ concentration (as shown in, for example, Yar
395
:

$$r_s = \frac{r_l(\text{CO}_2)}{0.5\text{LAI}} = \frac{r_{l-300} \{1 + S_{r_l-\text{CO}_2} (\text{CO}_2 - 300)\}}{0.5\text{LAI}} \quad (A2)$$

where $S_{r_l-\text{CO}_2}$ [ppm⁻¹] is the relative sensitivity of r_l to Δ [CO₂] and r_{l-300} [s m⁻¹] is the reference stomatal resistance when atmospheric CO₂ concentration is 300 ppm. The relative sensitivity of r_l to Δ [CO₂] represents the change in r_l per ppm
400 increase in CO₂ concentration.

If measurements of crop height exist, these data can be used to calculate the aerodynamic resistance to vapor and heat transfer (r_a) to represent the effects of crop phenology on PET:

$$r_a = \frac{\ln \left[\frac{z_m - d}{z_{om}} \right] \ln \left[\frac{z_h - d}{z_{oh}} \right]}{k^2 u_z} \quad (A3)$$

where z_m is the reference level at which the wind speed is measured; z_h is the height of the temperature and humidity
405 measurements; k is the von Karman constant ($= 0.41$), u_z is the measured wind speed (Allen et al., 1998) and d is the zero plane displacement height, taken as $0.67h_c$ [m]; z_{om} is the roughness parameter for momentum ($= 0.123h_c$) [m] and z_{oh} is the roughness parameter for heat and water vapor ($= 0.1z_{om}$) [m] (Jensen and Allen, 2016).

Free parameters in the Penman-Monteith equation, available for calibration, include the bulk stomatal $r_l = 100$ (ranging between 40 - 150) and surface resistance (ranging between 50 - 200), with values for specific surfaces/crops found in (Jensen and Allen, 201
410 . Additionally, one can adjust the surface albedo ($\alpha = 0.23$, ranges between 0.04 for water surfaces and 0.9 for snow) (Jensen and Allen, 2016) to estimate net shortwave radiation (equation 38 in Allen et al. (1998)), or the empirical coefficients for net long-wave radiation $a = 1.35$ and $b = -0.35$ (equation 39 in Allen et al. (1998)), or the empirical coefficients for

clear-sky radiation $as1 = 0.25$ and $bs1 = 0.5$ (equation 36 in Allen et al. (1998)). Optional meteorological inputs include G , T_{max} , T_{min} , RH_{max} , RH_{min} , p , N .

415 **ASCE-PM (*pm_asce*)**

The ASCE Penman-Monteith equation for is computed after Walter et al. (2000) [equation 1]:

$$PET = \frac{0.408\Delta(R_n - G) + \gamma \frac{C_n}{T+273} u_2 (e_s - e_a)}{\Delta + \gamma(C_d u_2)} \quad (A4)$$

where $C_n = 900$ and $C_d = 0.34$ are empirical coefficients for short reference vegetation (grass), while for tall reference vegetation (alfalfa) $C_n = 1600$ and $C_d = 0.38$ can be specified. The free parameters a , b , $as1$, $bs1$ and α are consistent with those specified for the Penman-Monteith method, while the optional meteorological inputs remain the same.

420

FAO-56 (*pm_fao56*)

The FAO-56 Penman-Monteith equation for reference crop evapotranspiration is computed after Allen et al. (1998) [equation 6]:

$$PET = \frac{0.408\Delta(R_n - G) + \gamma \frac{900}{T+273} u_2 (e_s - e_a)}{\Delta + \gamma(1 + 0.34u_2)} \quad (A5)$$

425 The free parameters a , b , $as1$, $bs1$ and α are consistent with those specified for the Penman-Monteith method, while the optional meteorological inputs remain the same.

Penman (*penman*)

The Penman's PET formulation is computed after Penman (1948):

$$PET = \frac{1}{\lambda \rho_w} \left[\frac{\Delta(R_n - G) + \gamma(e_s - e_a)(a_w + b_w u_2)}{\Delta + \gamma} \right] \quad (A6)$$

430 where free parameters for the Penman's wind function include $a_w = 1$ and $b_w = 0.537$ (Valiantzas, 2006), while Penman (1948) suggested values of $a_w = 2.626$ and $b_w = 1.381$. The free parameters a , b , $as1$, $bs1$ and α are consistent with those specified for the Penman-Monteith method, while the optional meteorological inputs remain the same.

Priestley-Taylor (*priestley_taylor*)

Priestley-Taylor's PET formulation is computed after Priestley and Taylor (1972):

$$435 \quad \underline{PET = \alpha_L \frac{\Delta(R_n - G)}{\lambda \rho_w (\Delta + \gamma)}} \quad (A7)$$

where $\alpha_L = 1.26$ is an empirical coefficient. The free parameters a , b , $as1$, $bs1$ and α are consistent with those specified for the Penman-Monteith method, while the optional meteorological inputs remain the same.

The Kimberly-Penman equation (Wright, 1982) is computed after Oudin et al. (2005):

$$PET = \frac{\Delta(R_n - G) + \gamma(e_s - e_a)w}{\lambda \rho_w (\Delta + \gamma)} \quad (A8)$$

$$440 \quad \text{where } w = u_2 \left[0.4 + 0.14 \exp\left(-\left(\frac{j-173}{58}\right)^2\right) \right] + \left[0.605 + 0.345 \exp\left(-\left(\frac{j-243}{80}\right)^2\right) \right].$$

The free parameters a , b , $as1$, $bs1$ and α are consistent with those specified for the Penman-Monteith method, while the optional meteorological inputs remain the same.

Thom-Oliver (*thom_oliver*)

Thom-Oliver's PET formulation is computed Thom and Oliver (1977), as used in Oudin et al. (2005):

$$445 \quad \underline{PET = \frac{\Delta(R_n - G) + 2.5\gamma(e_s - e_a) \cdot a_w(1 + b_w u_2)}{\lambda rh_o_w (\Delta + \gamma(1 + \frac{r_s}{r_a}))}} \quad (A9)$$

where $a_w = 2.6$ and $b_w = 0.536$. The free parameters a , b , $as1$, $bs1$, α , r_l , r_s and $S_{x_l-[CO_2]}$ are consistent with those specified for the Penman-Monteith method, while the optional meteorological inputs remain the same.

Blaney-Criddle (*blaney_criddle*)

Three different approaches can be taken to estimate the Blaney-Criddle PET, depending on the selected method:

$$450 \quad \underline{PET = \begin{cases} a + bP_y(0.46T + 8.13) & \text{if } Method = 0 \text{ (after Schrödter (1985)),} \\ k \cdot P_y(0.457T + 8.128) & \text{if } Method = 1 \text{ (after Xu and Singh (2001) [equation 6]),} \\ k1 + b_{var} \cdot P_y(0.46T + 8.13) & \text{if } Method = 2 \text{ (after McMahon et al. (2013) [equation S9.7 and S9.8]),} \end{cases}} \quad (A10)$$

where $k = 0.65$, $a = -1.55$ and $b = 0.96$ are empirical coefficients, and $k1 = 0.0043 \cdot RH_{min} - \frac{n}{N} - 1.41$, and $b_{var} = 0.81917 - 0.00409$

Hamon (*hamon*)

The PET formulation after (Hamon, 1963), as used in (Oudin et al., 2005):

$$455 \quad PET = k \cdot \left(\frac{N}{12}\right)^2 \cdot \exp\left(\frac{T}{16}\right) \quad (A11)$$

where $k = 1$ is a calibration coefficient.

Romanenko (*romanenko*)

Romanenko's PET formulation Romanenko (1961) as used in Oudin et al. (2005):

$$PET = k \left(1 + \frac{T}{25}\right)^2 \cdot \left(1 - \frac{e_a}{e_s}\right) \quad (A12)$$

460 where $k = 4.5$ is an empirical coefficient (Oudin et al., 2005). Optional meteorological inputs include T_{\max} , T_{\min} , RH_{\max} and RH_{\min} .

Linacre (*linacre*)

Linacre's PET formula Linacre (1977), as used in Oudin et al. (2005):

$$PET = \frac{\frac{500 \cdot T_m}{100 - \text{lat}_{\text{deg}}} + 15 \cdot (T - T_{\text{dew}})}{80 - T} \quad (A13)$$

465 where $T_m = T + 0.006 \cdot \text{Elev}$.

Haude (*haude*)

Haude's PET formulation Haude (1955), as used in Schiff (1975) is computed as:

$$PET = k \cdot FK \cdot (e_0 - e_a) \cdot 10 \quad (A14)$$

where $k = 1$ is a calibration coefficient and FK represents Haude's monthly coefficients, as adapted by Schiff (1975).

470 Turc (*turc*)

The PET formula, as derived from Turc (1961) and used in McMahon et al. (2013) (equations S9.10 and S9.11), is computed as:

$$PET = k \cdot c \cdot \frac{T}{T + 15} (23.88R_s + 50) \quad (A15)$$

where $k = 0.013$ is an empirical coefficient and c , dependent on the relative humidity (RH), is defined as:

$$c = \begin{cases} 1 + \frac{50 - rh}{70} & \text{if } rh < 50, \\ 1 & \text{otherwise.} \end{cases} \quad (A16)$$

Jensen-Haise (*jensen_haise*)

The PET according to the Jensen-Haise model (Jensen and Haise, 1963), varies depending on the chosen method:

$$PET = \begin{cases} k \frac{R_s}{\lambda \rho_w} (T - T_x) & \text{if } Method = 0 \text{ (after Jensen and Allen (2016))}, \\ k \frac{R_a T}{\lambda \rho_w} & \text{if } Method = 1 \text{ (after Oudin et al. (2005))}, \end{cases} \quad (A17)$$

where $cr = 0.025$ is an empirical coefficient and $T_x = -3$ as used in Jensen and Allen (2016).

480 **McGuinness-Bordne (*mcguinness_bordne*)**

McGuinness-Bordne's PET equation (McGuinness and Bordne, 1972), as used in Oudin et al. (2005):

$$PET = k \frac{R_a (T + 5)}{\lambda \rho_w} \quad (A18)$$

where $k = 0.0147$ is an empirical coefficient as suggested by Xu and Singh (2000).

Hargreaves (*hargreaves*)

485 The Hargreaves PET equation (Hargreaves and Samani, 1982) is computed as:

$$PET = k_{CHS} \frac{R_a}{\lambda \rho_w} \sqrt{T_{\max} - T_{\min}} (T + 17.8) \quad (A19)$$

- If *method* = 0 (after Jensen and Allen (2016)), the empirical coefficient $k = 0.0135$, and $c_{CHS} = 0$.
- If *method* = 1 (after McMahon et al. (2013)), $k = 0.0135$. The coefficient c_{CHS} is calculated as $0.00185 \cdot (T_{\max} - T_{\min})^2 - 0.0433 \cdot (T_{\max} - T_{\min}) + 0.0001$.

490 **FAO-24 (fao_24)**

The FAO-24 PET equation from Doorenbos (1977); Jensen et al. (1990), as used in Xu and Singh (2000) (equation 11 and 12), is given by:

$$PET = a + \frac{\Delta}{\Delta + \gamma} \frac{R_s}{\lambda \rho_w} (1 - \alpha) b \quad (\text{A20})$$

where $a = -0.3$ and $b = 1.066 - 0.13 \cdot \frac{RH}{100} + 0.045u_2 - 0.02 \cdot \frac{RH}{100} u_2 - 0.315 \cdot (\frac{RH}{100})^2 - 0.0011u_2$. Free parameters include the surface albedo α , consistent with those used in the Penman-Monteith method.

Abtew (abtew)

Abtew's PET equation (Abtew, 1996), as used in Xu and Singh (2000) (equation 14):

$$PET = k \frac{R_s}{\lambda \rho_w} \quad (\text{A21})$$

where $k = 0.53$ is an empirical coefficient as suggested by Xu and Singh (2000).

500 **Makkink (makink)**

Makkink's PET equation (Makkink, 1957):

$$PET = \frac{\Delta}{\Delta + \gamma} \frac{R_s}{\lambda \rho_w} \quad (\text{A22})$$

where $k = 0.65$ is the empirical coefficients recommended by Hiemstra and Sluiter (2011) ranging between (0.61-0.77) Jensen and Allen (2002).

505 **Makkink-KNMI (makink_knmi)**

The Royal Netherlands Meteorological Institute (KNMI) employs a slightly modified version of the Makkink equation, tailored specifically for conditions in the Netherlands, as described by Hiemstra and Sluiter (2011):

$$PET = k \frac{\Delta}{\Delta + \gamma} \frac{R_s}{\lambda \rho_w} \quad (\text{A23})$$

where $k = 0.65$. The calculations for s , e_s , γ , and λ are specified as follows:

510 $-\Delta = \frac{7.5 \cdot 237.3}{(237.3 + T)^2} \ln(10) \cdot e_{s_i}$

- $e_s = 0.6107 \cdot 10^{\frac{7.5T}{237.3+T}}$,
- $\gamma = 0.0646 + 0.00006T$,
- $\lambda = (2501 - 2.375T)1000$.

Oudin (oudin)

515 According to Oudin et al. (2005) (equation 2), the potential evapotranspiration (PET) can be expressed as:

$$PET = \begin{cases} R_a \frac{(T+k_2)}{\lambda \rho_w k_1} & \text{if } T + k_2 > 0 \\ 0 & \text{otherwise,} \end{cases} \quad (\text{A24})$$

where $k_2 = 5$ and $k_1 = 100$ (ranging between 75-100) are empirical coefficients recommended by Oudin et al. (2005).

Author contributions. Conceptualization, M.V., S.B. and R.C.; software, M.V. and R.C.; investigation, M.V.; writing—original draft preparation, M.V. and R.C. ; writing—review and editing, S.B. and R.C.; supervision, S.B.. All authors have read and agreed to the published
520 version of the manuscript.

Competing interests. The authors declare no conflict of interest.

Acknowledgements. We acknowledge the financial support by the University of Graz and the funding of the Earth System Sciences research program of ~~the~~ the Austrian Academy of Sciences (ÖAW project ClimGrassHydro). We acknowledge the ZAMG dataset (<https://data.hub.zamg.ac.at>), KNMI dataset (<https://www.knmi.nl/home>), and the E-OBS dataset from the EU-FP6 project UERRA (<http://www.uerra.eu>)
525 and the Copernicus Climate Change Service, and the data providers in the ECA&D project (<https://www.ecad.eu>).

References

- Abtew, W.: Evapotranspiration measurements and modeling for three wetland systems in South Florida 1, *JAWRA Journal of the American Water Resources Association*, 32, 465–473, publisher: Wiley Online Library, 1996.
- 530 Aguayo, R., León-Muñoz, J., Aguayo, M., Baez-Villanueva, O. M., Zambrano-Bigiarini, M., Fernández, A., and Jacques-Coper, M.: PatagoniaMet: A multi-source hydrometeorological dataset for Western Patagonia, *Scientific Data*, 11, 6, <https://doi.org/10.1038/s41597-023-02828-2>, 2024.
- Ainsworth, E. A. and Rogers, A.: The response of photosynthesis and stomatal conductance to rising [CO₂]: mechanisms and environmental interactions, *Plant, cell & environment*, 30, 258–270, <https://doi.org/10.1111/j.1365-3040.2007.01641.x>, publisher: Wiley Online Library, 2007.
- 535 Allen, R. G., Pereira, L. S., Raes, D., Smith, M., and others: Crop evapotranspiration-Guidelines for computing crop water requirements-FAO Irrigation and drainage paper 56, Fao, Rome, 300, D05 109, 1998.
- Andréassian, V., Perrin, C., and Michel, C.: Impact of imperfect potential evapotranspiration knowledge on the efficiency and parameters of watershed models, *Journal of Hydrology*, 286, 19–35, <https://doi.org/10.1016/j.jhydrol.2003.09.030>, 2004.
- Ansorge, L. and Beran, A.: Performance of simple temperature-based evaporation methods compared with a time series of pan evaporation 540 measures from a standard 20 m² tank, *Journal of Water and Land Development*, 41, 1–11, <https://doi.org/10.2478/jwld-2019-0021>, 2019.
- Bakundukize, C., Van Camp, M., and Walraevens, K.: Estimation of groundwater recharge in Bugesera region (Burundi) using soil moisture budget approach, *GEOLOGICA BELGICA*, 14, 85–102, <http://hdl.handle.net/1854/LU-1204652>, 2011.
- Barker, M., Chue Hong, N., Katz, D. S., Lamprecht, A.-L., Martinez Ortiz, C., Psomopoulos, F., Harrow, J., Castro, L., Gruenpeter, M., Martinez, P., and Honeyman, T.: Introducing the FAIR Principles for research software, *Scientific Data*, 9, <https://doi.org/10.1038/s41597-022-01710-x>, 2022.
- 545 Beven, K. and Freer, J.: *Journal of Hydrology*, 249, 11–29, [https://doi.org/https://doi.org/10.1016/S0022-1694\(01\)00421-8](https://doi.org/https://doi.org/10.1016/S0022-1694(01)00421-8), 2001.
- Blaney, H. F. and others: Determining water requirements in irrigated areas from climatological and irrigation data, Tech. rep., UNITED STATES DEPARTMENT OF AGRICULTURE, 1952.
- Bormann, H.: Sensitivity analysis of 18 different potential evapotranspiration models to observed climatic change at German climate stations, 550 *Climatic Change*, 104, 729–753, <https://doi.org/10.1007/s10584-010-9869-7>, 2010.
- Caretta, M. A., Mukherji, A., Arfanuzzaman, M., Betts, R. A., Gelfan, A., Hirabayashi, Y., Lissner, T. K., Liu, J., Gunn, E. L., Morgan, R., Mwangi, S., and Supratid, S.: Water. In: *Climate Change 2022: Impacts, Adaptation, and Vulnerability, Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [H.-O. Pörtner, D.C. Roberts, M. Tignor, E.S. Poloczanska, K. Mintenbeck, A. Alegría, M. Craig, S. Langsdorf, S. Lösche, V. Möller, A. Okem, B. Rama (eds.)], publisher: Cambridge 555 University Press, 2022.
- Collenteur, R. A., Moeck, C., Schirmer, M., and Birk, S.: Analysis of nationwide groundwater monitoring networks using lumped-parameter models, *Journal of Hydrology*, 626, 130 120, <https://doi.org/10.1016/j.jhydrol.2023.130120>, 2023.
- Cornes, R. C., van der Schrier, G., van den Besselaar, E. J. M., and Jones, P. D.: An Ensemble Version of the E-OBS Temperature and Precipitation Data Sets, *Journal of Geophysical Research: Atmospheres*, 123, 9391–9409, <https://doi.org/https://doi.org/10.1029/2017JD028200>, 560 2018.
- Dakhlaoui, H., Seibert, J., and Hakala Assendelft, K.: Sensitivity of discharge projections to potential evapotranspiration estimation in Northern Tunisia, *Regional Environmental Change*, 20, <https://doi.org/10.1007/s10113-020-01615-8>, 2020.

- Dallaire, G., Poulin, A., Arsenault, R., and Brissette, F.: Uncertainty of potential evapotranspiration modelling in climate change impact studies on low flows in North America, *Hydrological Sciences Journal*, 66, 689–702, <https://doi.org/10.1080/02626667.2021.1888955>, 2021.
- DeJonge, K. C. and Thorp, K. R.: Implementing Standardized Reference Evapotranspiration and Dual Crop Coefficient Approach in the DSSAT Cropping System Model, *Transactions of the ASABE*, 60, 1965–1981, <https://doi.org/10.13031/trans.12321>, 2017.
- Dingman, S. L.: *Physical hydrology*, Waveland Press, Inc, Long Grove, Illinois, third edition edn., 2015.
- Dlouhá, D., Dubovský, V., and Pospíšil, L.: Optimal Calibration of Evaporation Models against Penman–Monteith Equation, *Water*, 13, <https://doi.org/10.3390/w13111484>, 2021.
- Doorenbos, J.: Guidelines for predicting crop water requirements, *FAO irrigation and drainage paper*, 24, 1–179, publisher: FAO, 1977.
- Fatichi, S., Leuzinger, S., Paschalis, A., Langley, J. A., Barraclough, A. D., and Hovenden, M. J.: Partitioning direct and indirect effects reveals the response of water-limited ecosystems to elevated CO₂, *Proceedings of the National Academy of Sciences*, 113, 12 757–12 762, <https://doi.org/10.1073/pnas.1605036113>, publisher: National Acad Sciences, 2016.
- Field, C. B., Jackson, R. B., and Mooney, H. A.: Stomatal responses to increased CO₂: implications from the plant to the global scale, *Plant, Cell and Environment*, 18, 1214–1225, <https://doi.org/10.1111/j.1365-3040.1995.tb00630.x>, 1995.
- Fisher, J. B., Whittaker, R. J., and Malhi, Y.: ET come home: potential evapotranspiration in geographical ecology, *Global Ecology and Biogeography*, 20, 1–18, <https://doi.org/https://doi.org/10.1111/j.1466-8238.2010.00578.x>, 2011.
- Forstner, V., Vremec, M., Herndl, M., and Birk, S.: Effects of dry spells on soil moisture and yield anomalies at a montane managed grassland site: A lysimeter climate experiment, *Ecohydrology*, <https://doi.org/10.1002/eco.2518>, 2022.
- Gharbia, S. S., Smullen, T., Gill, L., Johnston, P., and Pilla, F.: Spatially distributed potential evapotranspiration modeling and climate projections, *Science of The Total Environment*, 633, 571–592, <https://doi.org/10.1016/j.scitotenv.2018.03.208>, 2018.
- Guo, D., Westra, S., and Maier, H. R.: An R package for modelling actual, potential and reference evapotranspiration, *Environmental Modelling & Software*, 78, 216–224, <https://doi.org/https://doi.org/10.1016/j.envsoft.2015.12.019>, 2016.
- Guo, D., Westra, S., and Maier, H. R.: Sensitivity of potential evapotranspiration to changes in climate variables for different Australian climatic zones, *Hydrology and Earth System Sciences*, 21, 2107–2126, <https://doi.org/10.5194/hess-21-2107-2017>, 2017.
- Hamon, W. R.: Estimating potential evapotranspiration, *Transactions of the American Society of Civil Engineers*, 128, 324–338, publisher: American Society of Civil Engineers, 1963.
- Hargreaves, G. H. and Samani, Z. A.: Estimating potential evapotranspiration, *Journal of the irrigation and Drainage Division*, 108, 225–230, [https://doi.org/10.1061/\(ASCE\)0733-9437\(1983\)109:3\(341\)](https://doi.org/10.1061/(ASCE)0733-9437(1983)109:3(341)), publisher: American Society of Civil Engineers, 1982.
- Harris, C. R., Millman, K. J., Van Der Walt, S. J., Gommers, R., Virtanen, P., Cournapeau, D., Wieser, E., Taylor, J., Berg, S., Smith, N. J., and others: Array programming with NumPy, *Nature*, 585, 357–362, <https://doi.org/https://doi.org/10.1038/s41586-020-2649-2>, publisher: Nature Publishing Group, 2020.
- Haslinger, K. and Bartsch, A.: Creating long-term gridded fields of reference evapotranspiration in Alpine terrain based on a recalibrated Hargreaves method, *Hydrology and Earth System Sciences*, 20, 1211–1223, <https://doi.org/10.5194/hess-20-1211-2016>, 2016.
- Hassanzadeh, A., Vázquez-Suñé, E., Valdivielso, S., and Corbella, M.: An Open-Source Python Library for Water Balance Modeling, <https://doi.org/10.2139/ssrn.4450673>, 2023.
- Haude, W.: Determination of evapotranspiration by an approach as simple as possible, *Mitt Dt Wetterdienst*, 2, 1955.
- Hiebl, J. and Frei, C.: Daily temperature grids for Austria since 1961—concept, creation and applicability, *Theoretical and applied climatology*, 124, 161–178, <https://doi.org/https://doi.org/10.1007/s00704-015-1411-4>, publisher: Springer, 2016.

- Hiemstra, P. and Sluiter: Interpolation of Makkink evaporation in the Netherlands, Royal Netherlands Meteorological Institute (KNMI), 2011.
- Houska, T., Kraft, P., Chamorro-Chavez, A., and Breuer, L.: SPOTting Model Parameters Using a Ready-Made Python Package, PLOS ONE, 10, 1–22, <https://doi.org/10.1371/journal.pone.0145180>, publisher: Public Library of Science, 2015.
- 605 Hoyer, S. and Hamman, J.: xarray: ND labeled arrays and datasets in Python, Journal of Open Research Software, 5, <https://doi.org/10.1038/s41592-019-0686-2>, publisher: Ubiquity Press, 2017.
- Hunter, J. D.: Matplotlib: A 2D graphics environment, Computing in Science & Engineering, 9, 90–95, <https://doi.org/10.1109/MCSE.2007.55>, publisher: IEEE COMPUTER SOC, 2007.
- Hutton, C., Wagener, T., Freer, J., Han, D., Duffy, C., and Arheimer, B.: Most computational hydrology is not reproducible, so is it really science?, Water Resources Research, 52, 7548–7555, <https://doi.org/https://doi.org/10.1002/2016WR019285>, 2016.
- 610 Jayathilake, D. I. and Smith, T.: Assessing the impact of PET estimation methods on hydrologic model performance, Hydrology Research, 52, 373–388, <https://doi.org/10.2166/nh.2020.066>, 2021.
- Jemeljanova, M., Collenteur, R. A., Kmoch, A., Bikše, J., Popovs, K., and Kalvāns, A.: Modeling hydraulic heads with impulse response functions in different environmental settings of the Baltic countries, Journal of Hydrology: Regional Studies, 47, 101416, <https://doi.org/10.1016/j.ejrh.2023.101416>, 2023.
- 615 Jensen, M. E. and Allen, R. G.: Evaporation, Evapotranspiration, and Irrigation Water Requirements, American Society of Civil Engineers, second edition edn., <https://doi.org/10.1061/9780784414057>, eprint: <https://ascelibrary.org/doi/pdf/10.1061/9780784414057>, 2016.
- Jensen, M. E. and Haise, H. R.: Estimating evapotranspiration from solar radiation, Journal of the Irrigation and Drainage Division, 89, 15–41, <https://doi.org/10.1061/JRCEA4.0000287>, publisher: American Society of Civil Engineers, 1963.
- 620 Jensen, M. E., Burman, R. D., Allen, R. G., and others: Evapotranspiration and irrigation water requirements, ASCE, New York, 1990.
- Katul, G. G., Oren, R., Manzoni, S., Higgins, C., and Parlange, M. B.: Evapotranspiration: A process driving mass transport and energy exchange in the soil-plant-atmosphere-climate system, Reviews of Geophysics, 50, <https://doi.org/https://doi.org/10.1029/2011RG000366>, 2012.
- Kay, A. and Davies, H.: Calculating potential evaporation from climate model data: A source of uncertainty for hydrological climate change impacts, Journal of Hydrology, 358, 221–239, <https://doi.org/10.1016/j.jhydrol.2008.06.005>, 2008.
- 625 Kingston, D. G., Todd, M. C., Taylor, R. G., Thompson, J. R., and Arnell, N. W.: Uncertainty in the estimation of potential evapotranspiration under climate change, Geophysical Research Letters, 36, <https://doi.org/https://doi.org/10.1029/2009GL040267>, 2009.
- Kluyver, T., Ragan-Kelley, B., Pé Rez, F., Granger, B., Bussonnier, M., Frederic, J., Kelley, K., Hamrick, J., Grout, J., Corlay, S., Ivanov, P., Avila, D., n, Abdalla, S., Willing, C., and Team, J. D.: Jupyter Notebooks – a publishing format for reproducible computational workflows, Positioning and Power in Academic Publishing: Players, Agents and Agendas, pp. 87–90, <https://doi.org/10.3233/978-1-61499-649-1-87>, 2016.
- 630 Krueger, T., Freer, J., Quinton, J. N., Macleod, C. J. A., Bilotta, G. S., Brazier, R. E., Butler, P., and Haygarth, P. M.: Ensemble evaluation of hydrological model hypotheses, Water Resources Research, 46, <https://doi.org/https://doi.org/10.1029/2009WR007845>, 2010.
- Kruijt, B., Witte, J.-P. M., Jacobs, C. M. J., and Kroon, T.: Effects of rising atmospheric CO₂ on evapotranspiration and soil moisture: A practical approach for the Netherlands, Journal of Hydrology, 349, 257–267, <https://doi.org/10.1016/j.jhydrol.2007.10.052>, 2008.
- Kumar, R., Jat, M. K., and Shankar, V.: Methods to estimate irrigated reference crop evapotranspiration – a review, Water Science and Technology, 66, 525–535, <https://doi.org/10.2166/wst.2012.191>, 2012.

- Lai, C., Chen, X., Zhong, R., and Wang, Z.: Implication of climate variable selections on the uncertainty of reference crop evapotranspiration projections propagated from climate variables projections under climate change, *Agricultural Water Management*, 259, 107 273, 640 <https://doi.org/10.1016/j.agwat.2021.107273>, 2022.
- Lemaitre-Basset, T., Oudin, L., Thirel, G., and Collet, L.: Unraveling the contribution of potential evaporation formulation to uncertainty under climate change, *Hydrology and Earth System Sciences*, 26, 2147–2159, <https://doi.org/10.5194/hess-26-2147-2022>, 2022.
- Linacre, E. T.: A simple formula for estimating evaporation rates in various climates, using temperature data alone, *Agricultural Meteorology*, 18, 409–424, [https://doi.org/https://doi.org/10.1016/0002-1571\(77\)90007-3](https://doi.org/https://doi.org/10.1016/0002-1571(77)90007-3), 1977.
- 645 Liu, Z., Han, J., and Yang, H.: Assessing the ability of potential evaporation models to capture the sensitivity to temperature, *Agricultural and Forest Meteorology*, 317, 108 886, <https://doi.org/10.1016/j.agrformet.2022.108886>, 2022.
- Makkink, G. F.: Testing the Penman formula by means of lysimeters, *Journal of the Institution of Water Engineers*, 11, 277–288, 1957.
- Martens, B., Miralles, D. G., Lievens, H., van der Schalie, R., de Jeu, R. A. M., Fernández-Prieto, D., Beck, H. E., Dorigo, W. A., and Verhoest, N. E. C.: GLEAM v3: satellite-based land evaporation and \hack\newlineroor-zone soil moisture, *Geoscientific Model Development*, 650 10, 1903–1925, <https://doi.org/10.5194/gmd-10-1903-2017>, 2017.
- Matevz Vremec and Raoul Collenteur: PyEt: A Python package for estimating potential evapotranspiration, <https://doi.org/10.5281/ZENODO.5896799>, 2024.
- Maxwell, R. M., Putti, M., Meyerhoff, S., Delfs, J., Ferguson, I. M., Ivanov, V., Kim, J., Kolditz, O., Kollet, S. J., Kumar, M., Lopez, S., Niu, J., Paniconi, C., Park, Y., Phanikumar, M. S., Shen, C., Sudicky, E. A., and Sulis, M.: Surface-subsurface model intercom- 655 parison: A first set of benchmark results to diagnose integrated hydrology and feedbacks, *Water Resources Research*, 50, 1531–1549, <https://doi.org/10.1002/2013WR013725>, 2014.
- May, R. M., Goebbert, K. H., Thielen, J. E., Leeman, J. R., Camron, M. D., Bruick, Z., Bruning, E. C., Manser, R. P., Arms, S. C., and Marsh, P. T.: MetPy: A meteorological Python library for data analysis and visualization, *Bulletin of the American Meteorological Society*, <https://doi.org/10.1175/BAMS-D-21-0125.1>, place: Boston MA, USA Publisher: American Meteorological Society, 2022.
- 660 McGuinness, J. and Bordne, E.: A comparison of lysimeter derived potential evapotranspiration with computed values, *Tech. Bull.*, 1452, *Agric. Res. Serv.*, US Dep. of Agric., Washington, DC, <https://doi.org/10.22004/ag.econ.171893>, 1972.
- McKinney, W.: Data Structures for Statistical Computing in Python, in: *Proceedings of the 9th Python in Science Conference*, edited by Walt, S. v. d. and Millman, J., pp. 56 – 61, <https://doi.org/10.25080/Majora-92bf1922-00a>, 201.
- McMahon, T. A., Peel, M. C., Lowe, L., Srikanthan, R., and McVicar, T. R.: Estimating actual, potential, reference crop and pan 665 evaporation using standard meteorological data: a pragmatic synthesis, *Hydrology and Earth System Sciences*, 17, 1331–1363, <https://doi.org/10.5194/hess-17-1331-2013>, 2013.
- Milly, P. and Dunne, K.: Potential evapotranspiration and continental drying, *Nature Climate Change*, 6, <https://doi.org/10.1038/nclimate3046>, 2016.
- Monteith, J. L.: Evaporation and environment, in: *Symposia of the society for experimental biology*, vol. 19, pp. 205–234, Cambridge 670 University Press (CUP) Cambridge, 1965.
- Oki, T. and Kanae, S.: Global Hydrological Cycles and World Water Resources, *Science*, 313, 1068–1072, <https://doi.org/10.1126/science.1128845>, 2006.
- Oudin, L., Michel, C., and Anctil, F.: Which potential evapotranspiration input for a lumped rainfall-runoff model?, *Journal of Hydrology*, 303, 275–289, <https://doi.org/10.1016/j.jhydrol.2004.08.025>, 2005.

- 675 Pastorello, G., Trotta, C., Canfora, E., Chu, H., Christianson, D., Cheah, Y.-W., Poindexter, C., Chen, J., Elbashandy, A., Humphrey, M., Isaac, P., Polidori, D., Reichstein, M., Ribeca, A., van Ingen, C., Vuichard, N., Zhang, L., Amiro, B., Ammann, C., Arain, M. A., Ardö, J., Arkebauer, T., Arndt, S. K., Arriga, N., Aubinet, M., Aurela, M., Baldocchi, D., Barr, A., Beamesderfer, E., Marchesini, L. B., Bergeron, O., Beringer, J., Bernhofer, C., Berveiller, D., Billesbach, D., Black, T. A., Blanken, P. D., Bohrer, G., Boike, J., Bolstad, P. V., Bonal, D., Bonnefond, J.-M., Bowling, D. R., Bracho, R., Brodeur, J., Brümmner, C., Buchmann, N., Burban, B., Burns, S. P., Buysse, P., Cale, P.,
- 680 Cavagna, M., Cellier, P., Chen, S., Chini, I., Christensen, T. R., Cleverly, J., Collalti, A., Consalvo, C., Cook, B. D., Cook, D., Coursolle, C., Cremonese, E., Curtis, P. S., D'Andrea, E., da Rocha, H., Dai, X., Davis, K. J., Cinti, B. D., Grandcourt, A. d., Ligne, A. D., De Oliveira, R. C., Delpierre, N., Desai, A. R., Di Bella, C. M., Tommasi, P. d., Dolman, H., Domingo, F., Dong, G., Dore, S., Duce, P., Dufrêne, E., Dunn, A., Dušek, J., Eamus, D., Eichelmann, U., ElKhidir, H. A. M., Eugster, W., Ewenz, C. M., Ewers, B., Famulari, D., Fares, S., Feigenwinter, I., Feitz, A., Fensholt, R., Filippa, G., Fischer, M., Frank, J., Galvagno, M., Gharun, M., Gianelle, D., Gielen, B., Gioli, B.,
- 685 Gitelson, A., Goded, I., Goeckede, M., Goldstein, A. H., Gough, C. M., Goulden, M. L., Graf, A., Griebel, A., Gruening, C., Grünwald, T., Hammerle, A., Han, S., Han, X., Hansen, B. U., Hanson, C., Hatakka, J., He, Y., Hehn, M., Heinesch, B., Hinko-Najera, N., Hörtnagl, L., Hutley, L., Ibrom, A., Ikawa, H., Jackowicz-Korczynski, M., Janouš, D., Jans, W., Jassal, R., Jiang, S., Kato, T., Khomik, M., Klatt, J., Knohl, A., Knox, S., Kobayashi, H., Koerber, G., Kolle, O., Kosugi, Y., Kotani, A., Kowalski, A., Kruijt, B., Kurbatova, J., Kutsch, W. L., Kwon, H., Launiainen, S., Laurila, T., Law, B., Leuning, R., Li, Y., Liddell, M., Limousin, J.-M., Lion, M., Liska, A. J., Lohila, A., López-Ballesteros, A., López-Blanco, E., Loubet, B., Loustau, D., Lucas-Moffat, A., Lüers, J., Ma, S., Macfarlane, C., Magliulo, V., Maier, R., Mammarella, I., Manca, G., Marcolla, B., Margolis, H. A., Marras, S., Massman, W., Mastepanov, M., Matamala, R., Matthes, J. H., Mazzenga, F., McCaughey, H., McHugh, I., McMillan, A. M. S., Merbold, L., Meyer, W., Meyers, T., Miller, S. D., Minerbi, S., Moderow, U., Monson, R. K., Montagnani, L., Moore, C. E., Moors, E., Moreaux, V., Moureaux, C., Munger, J. W., Nakai, T., Neiryneck, J., Nesic, Z., Nicolini, G., Noormets, A., Northwood, M., Noretto, M., Nouvellon, Y., Novick, K., Oechel, W., Olesen, J. E., Ourcival, J.-M., Papuga, S. A., Parmentier, F.-J., Paul-Limoges, E., Pavelka, M., Peichl, M., Pendall, E., Phillips, R. P., Pilegaard, K., Pirk, N., Posse, G., Powell, T., Prasse, H., Prober, S. M., Rambal, S., Rannik, , Raz-Yaseef, N., Rebmann, C., Reed, D., Dios, V. R. d., Restrepo-Coupe, N., Reverter, B. R., Roland, M., Sabbatini, S., Sachs, T., Saleska, S. R., Sánchez-Cañete, E. P., Sanchez-Mejia, Z. M., Schmid, H. P., Schmidt, M., Schneider, K., Schrader, F., Schroder, I., Scott, R. L., Sedláč, P., Serrano-Ortiz, P., Shao, C., Shi, P., Shironya, I., Siebicke, L., Šigut, L., Silberstein, R., Sirca, C., Spano, D., Steinbrecher, R., Stevens, R. M., Sturtevant, C., Suyker, A., Tagesson, T.,
- 700 Takanashi, S., Tang, Y., Tapper, N., Thom, J., Tomassucci, M., Tuovinen, J.-P., Urbanski, S., Valentini, R., van der Molen, M., van Gorsel, E., van Huissteden, K., Varlagin, A., Verfaillie, J., Vesala, T., Vincke, C., Vitale, D., Vygodskaya, N., Walker, J. P., Walter-Shea, E., Wang, H., Weber, R., Westermann, S., Wille, C., Wofsy, S., Wohlfahrt, G., Wolf, S., Woodgate, W., Li, Y., Zampedri, R., Zhang, J., Zhou, G., Zona, D., Agarwal, D., Biraud, S., Torn, M., and Papale, D.: The FLUXNET2015 dataset and the ONEFlux processing pipeline for eddy covariance data, *Scientific Data*, 7, 225, <https://doi.org/10.1038/s41597-020-0534-3>, 2020.
- 705 Penman, H. L.: Natural evaporation from open water, bare soil and grass, *Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences*, 193, 120–145, publisher: The Royal Society London, 1948.
- Peterson, T. J., Wasko, C., Saft, M., and Peel, M. C.: AWAPer: An R package for area weighted catchment daily meteorological data anywhere within Australia, *Hydrological Processes*, 34, 1301–1306, <https://doi.org/10.1002/hyp.13637>, 2020.
- Pimentel, R., Arheimer, B., Crochemore, L., Andersson, J. C. M., Pechlivanidis, I. G., and Gustafsson, D.: Which Potential Evapotranspiration Formula to Use in Hydrological Modeling World-Wide?, *Water Resources Research*, 59, e2022WR033447, <https://doi.org/10.1029/2022WR033447>, 2023.

- Priestley, C. H. B. and Taylor, R. J.: On the assessment of surface heat flux and evaporation using large-scale parameters, *Monthly weather review*, 100, 81–92, 1972.
- Prudhomme, C. and Williamson, J.: Derivation of RCM-driven potential evapotranspiration for hydrological climate change impact analysis
715 in Great Britain: a comparison of methods and associated uncertainty in future projections, *Hydrology and Earth System Sciences*, 17, 1365–1377, <https://doi.org/10.5194/hess-17-1365-2013>, 2013.
- Richards, M.: PyETo, <https://github.com/woodcrafty/PyETo>, 2019.
- Riedel, T., Weber, T. K. D., and Bergmann, A.: Near constant groundwater recharge efficiency under global change in a central European catchment, *Hydrological Processes*, 37, e14 805, <https://doi.org/10.1002/hyp.14805>, 2023.
- 720 Romanenko, V.: Computation of the autumn soil moisture using a universal relationship for a large area, *Proc. of Ukrainian Hydrometeorological Research Institute*, 3, 12–25, 1961.
- Rosenberry, D. O., Stannard, D. I., Winter, T. C., and Martinez, M. L.: Comparison of 13 equations for determining evapotranspiration from a prairie wetland, Cottonwood Lake Area, North Dakota, USA, *Wetlands*, 24, 483–497, [https://doi.org/10.1672/0277-5212\(2004\)024\[0483:COEFDE\]2.0.CO;2](https://doi.org/10.1672/0277-5212(2004)024[0483:COEFDE]2.0.CO;2), 2004.
- 725 Schiff, H.: Berechnung der potentiellen Verdunstung und deren Vergleich mit aktuellen Verdunstungswerten von Lysimetern, *Archiv für Meteorologie, Geophysik und Bioklimatologie, Serie B*, 23, 331–342, <https://doi.org/https://doi.org/10.1007/BF02242689>, publisher: Springer, 1975.
- Schrödter, H.: Hinweise Für den Einsatz Anwendungsorientierter Bestimmungsverfahren, Springer Berlin Heidelberg, Berlin, Heidelberg, https://doi.org/10.1007/978-3-642-70434-5_8, publication Title: Verdunstung: Anwendungsorientierte Meßverfahren und Bestimmungsmethoden, 1985.
- 730 Seiller, G. and Anctil, F.: How do potential evapotranspiration formulas influence hydrological projections?, *Hydrological Sciences Journal*, 61, 2249–2266, <https://doi.org/10.1080/02626667.2015.1100302>, 2016.
- Shi, L., Feng, P., Wang, B., Liu, D. L., Cleverly, J., Fang, Q., and Yu, Q.: Projecting potential evapotranspiration change and quantifying its uncertainty under future climate scenarios: A case study in southeastern Australia, *Journal of Hydrology*, 584, 124756, <https://doi.org/https://doi.org/10.1016/j.jhydrol.2020.124756>, 2020.
- 735 Sperna Weiland, F. C., Tisseuil, C., Dürr, H. H., Vrac, M., and van Beek, L. P. H.: Selecting the optimal method to calculate daily global reference potential evaporation from CFSR reanalysis data for application in a hydrological model study, *Hydrology and Earth System Sciences*, 16, 983–1000, <https://doi.org/10.5194/hess-16-983-2012>, 2012.
- Tercini, J. R. B. and Mello Júnior, A. V.: Impact of Hydroclimatic Changes on Water Security in the Cantareira Water Production System, Brazil, *Atmosphere*, 14, 1836, <https://doi.org/10.3390/atmos14121836>, 2023.
- 740 Thom, A. and Oliver, H.: On Penman’s equation for estimating regional evaporation, *Quarterly Journal of the Royal Meteorological Society*, 103, 345–357, publisher: Wiley Online Library, 1977.
- Thorp, K. R.: pyfao56: FAO-56 evapotranspiration in Python, *SoftwareX*, 19, 101208, <https://doi.org/10.1016/j.softx.2022.101208>, 2022.
- Trnka, M., Rötter, R. P., Ruiz-Ramos, M., Kersebaum, K. C., Olesen, J. E., Žalud, Z., and Semenov, M. A.: Adverse weather conditions for European wheat production will become more frequent with climate change, *Nature Climate Change*, 4, 637–643, <https://doi.org/10.1038/nclimate2242>, 2014.
- 745 Turc, L.: Estimation of irrigation water requirements, potential evapotranspiration: a simple climatic formula evolved up to date, *Ann. Agron*, 12, 13–49, 1961.
- unittest: unittest, <https://docs.python.org/3/library/unittest.html>, 2022.

- 750 Valiantzas, J. D.: Simplified versions for the Penman evaporation equation using routine weather data, *Journal of Hydrology*, 331, 690–702, <https://doi.org/10.1016/j.jhydrol.2006.06.012>, 2006.
- Valipour, M.: Evaluation of radiation methods to study potential evapotranspiration of 31 provinces, *Meteorology and Atmospheric Physics*, 127, 289–303, <https://doi.org/10.1007/s00703-014-0351-3>, 2015.
- Van Vuuren, D. P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., Hurtt, G. C., Kram, T., Krey, V., Lamarque, J.-F., and
 755 others: The representative concentration pathways: an overview, *Climatic change*, 109, 5–31, <https://doi.org/10.1007/s10584-011-0148-z>, publisher: Springer, 2011.
- Vaz, P. J., Schütz, G., Guerrero, C., and Cardoso, P. J. S.: A Study on the Prediction of Evapotranspiration Using Freely Available Meteorological Data, in: *Computational Science – ICCS 2022*, edited by Groen, D., de Mulatier, C., Paszynski, M., Krzhizhanovskaya, V. V., Dongarra, J. J., and Sloot, P. M. A., pp. 436–450, Springer International Publishing, Cham, 2022.
- 760 Velázquez, J. A., Schmid, J., Ricard, S., Muerth, M. J., Gauvin St-Denis, B., Minville, M., Chaumont, D., Caya, D., Ludwig, R., and Turcotte, R.: An ensemble approach to assess hydrological models’ contribution to uncertainties in the analysis of climate change impact on water resources, *Hydrology and Earth System Sciences*, 17, 565–578, <https://doi.org/10.5194/hess-17-565-2013>, 2013.
- Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D., Burovski, E., Peterson, P., Weckesser, W., Bright, J., van der Walt, S. J., Brett, M., Wilson, J., Millman, K. J., Mayorov, N., Nelson, A. R. J., Jones, E., Kern, R., Larson, E., Carey, C. J., Polat,
 765 , Feng, Y., Moore, E. W., VanderPlas, J., Laxalde, D., Perktold, J., Cimrman, R., Henriksen, I., Quintero, E. A., Harris, C. R., Archibald, A. M., Ribeiro, A. H., Pedregosa, F., van Mulbregt, P., and SciPy 1.0 Contributors: SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python, *Nature Methods*, 17, 261–272, <https://doi.org/10.1038/s41592-019-0686-2>, 2020.
- Vremec, M. and Collenteur, R.: `pyet-pypi`, <https://pypi.org/project/pyet/>, 2022.
- Vremec, M., Forstner, V., Herndl, M., Collenteur, R., Schaumberger, A., and Birk, S.: Sensitivity of evapotranspiration and seepage to elevated atmospheric C O 2 from lysimeter experiments in a montane grassland, *Journal of Hydrology*, p. 128875, <https://doi.org/10.1016/j.jhydrol.2022.128875>, 2022.
- 770 Walter, I. A., Allen, R. G., Elliott, R., Jensen, M., Itenfisu, D., Mecham, B., Howell, T., Snyder, R., Brown, P., Echings, S., and others: ASCE’s standardized reference evapotranspiration equation, in: *Watershed management and operations management 2000*, pp. 1–11, 2000.
- Wang, K. and Dickinson, R. E.: A review of global terrestrial evapotranspiration: Observation, modeling, climatology, and climatic variability, *Reviews of Geophysics*, 50, <https://doi.org/https://doi.org/10.1029/2011RG000373>, 2012.
- 775 Webber, H., Gaiser, T., Oomen, R., Teixeira, E., Zhao, G., Wallach, D., Zimmermann, A., and Ewert, F.: Uncertainty in future irrigation water demand and risk of crop failure for maize in Europe, *Environmental Research Letters*, 11, 074 007, <https://doi.org/10.1088/1748-9326/11/7/074007>, 2016.
- Wood, E. F., Lettenmaier, D. P., Liang, X., Lohmann, D., Boone, A., Chang, S., Chen, F., Dai, Y., Dickinson, R. E., Duan, Q., Ek, M.,
 780 Gusev, Y. M., Habets, F., Irannejad, P., Koster, R., Mitchel, K. E., Nasonova, O. N., Noilhan, J., Schaake, J., Schlosser, A., Shao, Y., Shmakin, A. B., Verseghy, D., Warrach, K., Wetzel, P., Xue, Y., Yang, Z.-L., and Zeng, Q.-c.: The Project for Intercomparison of Land-surface Parameterization Schemes (PILPS) Phase 2(c) Red–Arkansas River basin experiment:, *Global and Planetary Change*, 19, 115–135, [https://doi.org/10.1016/S0921-8181\(98\)00044-7](https://doi.org/10.1016/S0921-8181(98)00044-7), 1998.
- Wright, J. L.: New evapotranspiration crop coefficients, *Proceedings of the American Society of Civil Engineers, Journal of the Irrigation and Drainage Division*, 108, 57–74, 1982.
- 785 Xiang, K., Li, Y., Horton, R., and Feng, H.: Similarity and difference of potential evapotranspiration and reference crop evapotranspiration – a review, *Agricultural Water Management*, 232, 106 043, <https://doi.org/https://doi.org/10.1016/j.agwat.2020.106043>, 2020.

- Xie, Z., Yao, Y., Zhang, X., Liang, S., Fisher, J. B., Chen, J., Jia, K., Shang, K., Yang, J., Yu, R., Guo, X., Liu, L., Ning, J., and Zhang, L.:
The Global LAnd Surface Satellite (GLASS) evapotranspiration product Version 5.0: Algorithm development and preliminary validation,
790 *Journal of Hydrology*, 610, 127 990, <https://doi.org/https://doi.org/10.1016/j.jhydrol.2022.127990>, 2022.
- Xu, C.-Y. and Singh, V. P.: Evaluation and generalization of radiation-based methods for calculating evaporation, *Hydrological Processes*,
14, 339–349, 2000.
- Xu, C.-Y. and Singh, V. P.: Evaluation and generalization of temperature-based methods for calculating evaporation, *Hydrological Processes*,
15, 305–319, <https://doi.org/https://doi.org/10.1002/hyp.119>, 2001.
- 795 Yang, Y., Roderick, M., Zhang, S., McVicar, T., and Donohue, R.: Hydrologic implications of vegetation response to elevated CO₂ in climate
projections, *Nature Climate Change*, 9, <https://doi.org/10.1038/s41558-018-0361-0>, 2019.
- Yang, Y., Chen, R., Han, C., and Liu, Z.: Evaluation of 18 models for calculating potential evapotranspiration in different climatic zones of
China, *Agricultural Water Management*, 244, 106 545, <https://doi.org/10.1016/j.agwat.2020.106545>, 2021.
- 800 Zhou, J., Wang, Y., Su, B., Wang, A., Tao, H., Zhai, J., Kundzewicz, Z. W., and Jiang, T.: Choice of potential
evapotranspiration formulas influences drought assessment: A case study in China, *Atmospheric Research*, 242, 104 979,
<https://doi.org/https://doi.org/10.1016/j.atmosres.2020.104979>, 2020.